

UNIFORM BOUND ON THE NUMBER OF PARTITIONS FOR OPTIMAL CONFIGURATIONS OF THE OHTA-KAWASAKI ENERGY IN 3D

XIN YANG LU AND JUN-CHENG WEI

ABSTRACT. We study a 3D ternary system derived as a sharp-interface limit of the Nakazawa-Ohta density functional theory of triblock copolymers, which combines an interface energy with a long range interaction term. Although both the binary case in 2D and 3D, and the ternary case in 2D, are quite well studied, very little is known about the ternary case in 3D. In particular, it is even unclear whether minimizers are made of finitely many components. In this paper we provide a positive answer to this, by proving that the number of components in a minimizer is bounded from above by some quantity depending only on the total masses and the interaction coefficients. One key difficulty is that the 3D structure prevents us from uncoupling the Coulomb-like long range interaction from the perimeter term, hence the actual shape of minimizers is unknown, not even for small masses. This is due to the lack of a quantitative isoperimetric inequality with two mass constraints in 3D, and it makes the construction of competitors significantly more delicate.

Keywords. Pattern formation, small volume-fraction limit, triblock copolymers.

AMS Subject Classification. 49S05, 35K30, 35K55

1. INTRODUCTION

Energy functionals entailing a direct competition between an attractive short-range force and a repulsive Coulombic long-range force are mathematically studied intensively in recent years to understand physical problems such as Gamow's liquid drop problem and self-assembly of block copolymers.

In Gamow's liquid drop model [10], the volume of the nucleus $\Omega \subset \mathbb{R}^3$ is fixed, that is, $|\Omega| = m$. Here m is referred to as "mass". The binding energy is given by

$$\mathcal{E}_{\text{liquid}}(\Omega) := \text{Per}(\Omega) + \frac{1}{8\pi} \iint_{\Omega \times \Omega} \frac{1}{|x - y|} dx dy,$$

where the first term is the perimeter term which is $\mathcal{H}^2(\Omega)$, the surface area of Ω , and it arises because of lower nucleon density near the nucleus boundary; the second term is the Coulombic term which is introduced due to the presence of positively charged protons [3].

In Ohta and Kawasaki's diblock copolymer model [21], the free energy is represented by

$$\mathcal{E}_{\text{diblock}}(\Omega) := \text{Per}(\Omega) + \gamma \iint_{\Omega \times \Omega} G(x, y) dx dy,$$

where the first term is the perimeter term $\mathcal{H}^2(\Omega)$ which favors a large ball; the second term prefers splitting and models long-range interaction between monomers due to the connectivity of different subchains in

copolymer molecules. Here

$$G(x, y) = \frac{1}{4\pi|x - y|} + R(x, y)$$

is the Green's function of $-\Delta$ operator in \mathbb{R}^3 , $R(x, y)$ is the regular part of $G(x, y)$, and γ is the long-range interaction coefficient which is determined by the percentage of each type monomer, the number of all monomers in a chain molecule, the repulsion between unlike monomers, and the average distance between two adjacent monomers [6]. During each experiment, the total mass of each type monomer is fixed. So the energy is minimized under the mass constraint $|\Omega| = m$.

The diblock copolymer model is a model in binary systems. In this paper we study a counter model in ternary systems which was introduced by Nakazawa and Ohta to study triblock copolymers [20]. A triblock copolymer is a chain molecule consisting of three types of subchains, a subchain of type A monomers connected to a subchain of type B monomers and then connected to a subchain of type C monomers. Block copolymers can be used as a material in artificial organ technology and controlled drug delivery.

The free energy functional of triblock copolymers used here is a sharp interface model, derived as the Γ -limit of Nakazawa and Ohta's diffuse interface model, by Ren and Wei in [24, 23]

$$\mathcal{E}_{\text{triblock}}(\Omega_1, \Omega_2) := \frac{1}{2} \sum_{i=0}^2 \text{Per}(\Omega_i) + \sum_{i,j=1}^2 \gamma_{ij} \iint_{\Omega_i \times \Omega_j} G(x, y) \, dx \, dy.$$

Here $\Omega_0 = (\Omega_1 \cup \Omega_2)^c$, the perimeter term is defined by

$$\frac{1}{2} \sum_{i=0}^2 \text{Per}(\Omega_i) = \sum_{0 \leq i < j \leq 2} \mathcal{H}^2(\partial\Omega_i \cap \partial\Omega_j),$$

and the long-range interaction coefficients γ_{ij} form a 2×2 symmetric matrix. Using a "droplet" scaling argument, as done by Choksi and Peletier in [4, 5], and by Alama, Bronsard, the first author, and Wang in [1], it can be shown that the leading order of the free energy takes the form

$$E_0(\Omega_1, \Omega_2) = \sum_k e_0(|\Omega_{1,k}|, |\Omega_{2,k}|), \quad \Omega_i = \cup_k \Omega_{i,k}, \quad i = 1, 2, \quad (1)$$

where

$$\begin{aligned} e_0 : [0, +\infty) \times [0, +\infty) &\longrightarrow \mathbb{R}, \\ e_0(m_1, m_2) &:= \inf \left\{ \sum_{0 \leq i < j \leq 2} \mathcal{H}^2(\partial\Omega_i \cap \partial\Omega_j) \right. \\ &\quad \left. + \sum_{i,j=1}^2 \frac{\Gamma_{ij}}{4\pi} \int_{\Omega_i \times \Omega_j} \frac{1}{|x - y|} \, dx \, dy : |\Omega_i| = m_i, \, i = 1, 2 \right\}, \end{aligned}$$

where Γ_{ij} is a suitable scaling of γ_{ij} , which we will present in Section 2. That is, E_0 seeks the optimal partition $\Omega_i = \bigcup_k \Omega_{i,k}$, with each couple $(\Omega_{1,k}, \Omega_{2,k})$ minimizing e_0 .

2. SETTING UP THE PROBLEM

The aim of this section is to introduce the main energy of this paper. Choksi and Peletier showed in [4, Theorem 4.2] that, with the domain being the unit torus \mathbb{T}^3 , in the small mass volume fraction regime, the first order Γ -limit of the energies (see [4, Equation (1.8)])

$$E_\eta^{3d}(v) := \begin{cases} \eta \int_{\mathbb{T}^3} |\nabla v| dx + \eta \|v - \frac{1}{|\mathbb{T}^3|} \int_{\mathbb{T}^3} v dx\|_{\mathcal{H}^{-1}(\mathbb{T}^3)}^2 & \text{if } v \in BV(\mathbb{T}^3; \{0, \eta^{-3}\}), \\ +\infty & \text{otherwise,} \end{cases}$$

is of the form

perimeter + long range interaction,

i.e.,

$$E_0^{3d}(v) := \begin{cases} \sum_{k=0}^{\infty} e_0(m_k) & \text{if } v = \sum_{k=0}^{\infty} m_k \delta_{x_k}, \sum_{i=0}^{\infty} m_k = M = \text{total mass,} \\ +\infty & \text{otherwise,} \end{cases}$$

with

$$e_0^{3d}(m) = \inf \left\{ \int_{\mathbb{R}^3} |\nabla z| dx + \Gamma \|z\|_{H^{-1}(\mathbb{R}^3)}^2 : z \in BV(\mathbb{R}^3; \{0, 1\}), \|z\|_{L^1(\mathbb{R}^3)} = M \right\}.$$

The H^{-1} norm can be made explicit:

$$\|z\|_{H^{-1}(\mathbb{R}^3)}^2 = \int_{\mathbb{R}^3 \times \mathbb{R}^3} G(|x - y|) z(x) z(y) dx dy,$$

where G denotes the Green's function of \mathbb{R}^3 . That is, the minima seeks the optimal partition, in which each component minimizes the energy e_0^{3d} . An analogous result, but for ternary systems on the two dimensional torus, was obtained in [1, Theorem 3.2].

With the same arguments as in [4, 1], we can show that again, with the domain being the unit torus \mathbb{T}^3 , in the small mass volume fraction regime, the first order Γ -limit of the energies (which are the analogue of [1, Equation (1.8)] for ternary systems in 3D)

$$E_{\text{ternary}, \eta}^{3d}(v_{1, \eta}, v_{2, \eta}) := \begin{cases} \frac{\eta}{2} \sum_{i=0}^2 \int_{\mathbb{T}^3} |\nabla v_{i, \eta}| dx \\ + \sum_{i, j=1}^2 \eta^4 \gamma_{ij} \int_{\mathbb{T}^3 \times \mathbb{T}^3} G_{\mathbb{T}^3}(|x - y|) v_{i, \eta}(x) v_{j, \eta}(y) dx dy & \text{if } v_{1, \eta}, v_{2, \eta} \in BV(\mathbb{T}^3; \{0, \frac{1}{\eta^3}\}), \\ +\infty & \text{otherwise,} \end{cases}$$

$G_{\mathbb{T}^3} := \text{Green's function of } \mathbb{T}^3 \text{ with zero average,}$

can be again written in the form

$$E_{\text{ternary}, 0}^{3d}(v_1, v_2) := \begin{cases} \sum_{k=0}^{\infty} e_0(m_{1, k}, m_{2, k}) & \text{if } v_i = \sum_{k=0}^{\infty} m_{i, k} \delta_{x_{i, k}}, \sum_{k=0}^{\infty} m_{i, k} = M_i, \\ +\infty & \text{otherwise,} \end{cases} \quad (2)$$

$M_i = \text{total mass of type } i \text{ constituent,} \quad i = 1, 2,$

where

$$e_0(m_1, m_2) = \inf \left\{ \sum_{0 \leq i < j \leq 2} \mathcal{H}^2(\partial\Omega_i \cap \partial\Omega_j) + \sum_{i,j=1}^2 \Gamma_{ij} \|z_i\|_{H^{-1}(\mathbb{R}^3)}^2 : \right. \\ \left. z_i \in BV(\mathbb{R}^3; \{0, 1\}), \|z_i\|_{L^1(\mathbb{R}^3)} = m_i, \right. \\ \left. \Omega_i = \text{supp } z_i, i = 1, 2, |\Omega_1 \cap \Omega_2| = 0 \right\}, \quad \Omega_0 = (\Omega_1 \cup \Omega_2)^c,$$

and $\Gamma_{ij}\eta^{-3} = \gamma_{ij} \geq 0$ are coefficients penalizing the Coulomb interaction. Observe that the problem of minimizing $E_{\text{ternary},0}^{3d}$ is determined once we fix the total masses M_i and the interaction coefficients Γ_{ij} . Each couple of sets (Ω_1, Ω_2) , with the appropriate masses and minimizing e_0 , is referred to as a “cluster”. Similarly to [1], it is not restrictive to require $x_{i,k} \neq x_{i,j}$ whenever $k \neq j$, $i = 1, 2$, but allowing for $x_{1,k} \neq x_{2,j}$ for some k, j . Thus, it is possible to impose $x_{1,k} = x_{2,k} = x_k$ for all k , and allow for some masses $m_{i,k} = 0$, but still requiring $m_{1,k} + m_{2,k} > 0$ for all k . This is due to the ternary nature of the system, to account for the fact that at some x_k we might have both types of constituents (hence both $m_{1,k}$ and $m_{2,k}$ are positive), while at another x_j we might have only one type of constituent (hence only one between $m_{1,j}$ and $m_{2,j}$ is positive).

Next, we introduce the main energy of this paper: given connected sets Ω_1, Ω_2 , with $\mathbf{1}_{\Omega_i} \in BV(\mathbb{R}^3; \{0, 1\})$ and $|\Omega_1 \cap \Omega_2| = 0$, define the energy

$$E(\Omega_1, \Omega_2) := \sum_{0 \leq i < j \leq 2} \mathcal{H}^2(\partial\Omega_i \cap \partial\Omega_j) + \sum_{i,j=1}^2 \gamma_{ij} \int_{\Omega_i \times \Omega_j} |x - y|^{-1} dx dy, \quad (3)$$

where $\Omega_0 = (\Omega_1 \cup \Omega_2)^c$. Here γ_{ij} denote the interaction strengths, and are positive, of order $O(1)$. Then, given disjoint unions

$$\left(\bigsqcup_k \Omega_{1,k}, \bigsqcup_k \Omega_{2,k} \right),$$

with $\Omega_{i,k}$ being connected, the total energy of this configuration is defined by

$$\mathcal{E} \left(\bigsqcup_k \Omega_{1,k}, \bigsqcup_k \Omega_{2,k} \right) := \sum_k E(\Omega_{1,k}, \Omega_{2,k}).$$

Note that our main energy is quite similar to [1, Equation (1.8)], [1, Equation (1.8)], and (2), with the main difference being in the Green’s function in the interaction term; and also to $\mathcal{E}_{\text{liquid}}$, $\mathcal{E}_{\text{diblock}}$, and $\mathcal{E}_{\text{triblock}}$, with the main difference being that the interaction between different components are suppressed.

In the following, when we say “optimal configuration”, unless otherwise specified, we mean a configuration $(\bigsqcup_k \Omega_{1,k}, \bigsqcup_k \Omega_{2,k})$ minimizing \mathcal{E} .

In 2D, due to the fact that the Green’s function is a logarithmic term, the interaction was simply the product of the masses, hence it was equivalent to minimize the perimeter, subject to two mass constraints. It is well known that the double bubble is the unique such minimizer (see e.g. [8, 18] for the 2D case, and [12] for the 3D case, and also [22, 7, 16, 17]). In the ternary 3D case, however, such simplification is not available, and the shape of the minimizers is unclear, even for small masses. This is a significant hurdle, and studying the shape of minimizers is hindered by the lack of a quantitative isoperimetric inequality with two mass constraints in 3D

Therefore, a priori, it is even unclear whether optimal configurations have finitely many clusters, as we cannot exclude the presence of infinitely many components with very small masses. Our main result is to show that this is not the case:

Theorem 1. *There exists a computable constant $K = K(M_1, M_2, \gamma_{11}, \gamma_{22})$ such that any optimal configuration has at most K clusters.*

Notation. Since the position of the clusters is rarely relevant, in this paper we denote by B_m a ball of mass m .

3. UNIFORM UPPER BOUND ON THE NUMBER OF CLUSTERS

The proof of Theorem 1 will be split over several lemmas. Throughout the entire section, M_i , $i = 1, 2$, will denote the total masses of type i constituent, and γ_{ij} , $i, j = 1, 2$ will denote the interaction coefficients. These parameters completely determine the minimization problem for $\mathcal{E}_{\text{triblock}}$ in 3D. All the M_i and γ_{ij} will assumed to be given, and do not change throughout the section. Our proof will proceed as follows.

- (1) First, in Lemma 2, we bound from above the number of clusters made purely of one constituent type. Such upper bound will depend only on M_i , γ_{ii} , $i = 1, 2$.
- (2) Then, in Lemma 3, we show that the largest cluster's mass cannot be too small. Such lower bound will depend only on M_i , γ_{ii} , $i = 1, 2$.
- (3) Finally, in Lemmas 4 and 5 we show that the total mass of each cluster is bounded from below by a constant depending only on M_i , γ_{ii} , $i = 1, 2$. Since there is only so much total mass (i.e., $M_1 + M_2$), this allows us to infer Theorem 1.

Lemma 2. *Consider an optimal configuration, made of clusters $\Omega_{i,k}$, $i = 1, 2$, $k \geq 1$. Then*

$$\#\{k : |\Omega_{1,k}| |\Omega_{2,k}| = 0\}$$

is bounded from above by a constant depending only on M_i , γ_{ii} , $i = 1, 2$.

Proof. It is well known (see e.g. [13, 2, 19, 9, 14, 15], and references therein) that there exist $m_{i,B} = m_{i,B}(\gamma_{ii})$, $i = 1, 2$, such that, for all $m \leq m_{i,B}(\gamma_{ii})$, the minimizer of

$$\inf_{|X|=m} \left\{ \mathcal{H}^2(\partial X) + \gamma_{ii} \int_{X \times X} |x - y|^{-1} dx dy \right\}$$

is given by B_m . Since $\mathcal{H}^2(\partial B_m)$ (resp. $\int_{X \times X} |x - y|^{-1} dx dy$) scales like $m^{2/3}$ (resp. $m^{5/3}$), the perimeter term is dominating for all sufficiently small masses. Thus there exist geometric constants $m_{i,S} = m_{i,S}(\gamma_{ii}) \leq m_{i,B}(\gamma_{ii})$ such that

$$\begin{aligned} & \mathcal{H}^2(\partial B_{m_1}) + \gamma_{ii} \int_{B_{m_1} \times B_{m_1}} |x - y|^{-1} dx dy \\ & + \mathcal{H}^2(\partial B_{m_2}) + \gamma_{ii} \int_{B_{m_2} \times B_{m_2}} |x - y|^{-1} dx dy \\ & > \mathcal{H}^2(\partial B_{m_1+m_2}) + \gamma_{ii} \int_{B_{m_1+m_2} \times B_{m_1+m_2}} |x - y|^{-1} dx dy, \end{aligned}$$

for all $m_1, m_2 \leq m_{i,S}(\gamma_{ii})$, i.e. combining the two balls is energetically favorable whenever $m_1, m_2 \leq m_{i,S}(\gamma_{ii})$. Thus we cannot have two balls of the type i constituent, both with masses less than $m_{i,S}(\gamma_{ii})$. Since the total mass is $M_1 + M_2 < +\infty$, the proof is complete. \square

Lemma 3. *Consider an optimal configuration, made of clusters $\Omega_{i,k}$, $i = 1, 2$, $k \geq 1$. Then*

$$m_i^+ := \sup_k m_{i,k}, \quad m_{i,k} := |\Omega_{i,k}|,$$

is bounded from below by

$$\min \left\{ \frac{M_i}{2}, \left(\frac{c_i M_i}{2 \sum_{i=1}^2 [\sqrt[3]{36\pi} M_i^{2/3} + \gamma_{ii} \int_{B_{M_i} \times B_{M_i}} |x - y|^{-1} dx dy]} \right)^3 \right\}.$$

Note that, curiously, this lower bound is independent of γ_{12} . As it will be clear from the proof, this is due to the fact that an upper bound for the energy of an optimal configuration is given by the energy of two balls of masses M_1 and M_2 respectively. Such upper bound is clearly independent of γ_{12} .

Proof. The idea is that, for very small masses, the perimeter term is sub-additive and dominating. Assume $m_i^+ \leq M_i/2$, as otherwise $M_i/2$ is already a lower bound. Note that

$$E(\Omega_{1,k}, \Omega_{2,k}) \geq \mathcal{S}(m_{1,k}, m_{2,k}) \quad \forall k \geq 1,$$

where

$$\mathcal{S}(m_1, m_2) = \text{perimeter of the standard double bubble with masses } m_1 \text{ and } m_2,$$

and, by [11, Theorem 4.2] (applied with $v_1 = m_1$, $x = v_2 = m_2$, $n = 3$)

$$\mathcal{S}(m_1, m_2) \geq \sum_{i=1}^2 c_i m_i^{2/3}, \quad c_1 = c_2 = \frac{\sqrt[3]{36\pi}}{2}.$$

Thus the total energy of our optimal configuration satisfies

$$\sum_{k \geq 1} E(\Omega_{1,k}, \Omega_{2,k}) \geq \sum_{i=1}^2 c_i \sum_{k \geq 1} m_{i,k}^{2/3}.$$

By the concavity of the function $t \mapsto t^{2/3}$, the sum $\sum_{k \geq 1} m_{i,k}^{2/3}$ is minimum when $m_{i,k} \in \{0, m_i^+\}$ for all k . Since $\sum_{k \geq 1} m_{i,k} = M_i$, there are at least $\lfloor \frac{M_i}{m_i^+} \rfloor$ many clusters containing type i constituents, thus

$$\begin{aligned} \sum_{k \geq 1} E(\Omega_{1,k}, \Omega_{2,k}) &\geq \sum_{i=1}^2 c_i \sum_{k \geq 1} m_{i,k}^{2/3} \geq \sum_{i=1}^2 c_i \left\lfloor \frac{M_i}{m_i^+} \right\rfloor (m_i^+)^{2/3} \\ &\geq \sum_{i=1}^2 c_i \frac{M_i - m_i^+}{(m_i^+)^{1/3}} \geq \sum_{i=1}^2 \frac{c_i}{2} \frac{M_i}{(m_i^+)^{1/3}}. \end{aligned}$$

Since our configuration was an optimal one, its energy does not exceed that of two balls, which we denote by B_{M_1} and B_{M_2} , of masses M_1 and M_2 , respectively. Thus the above line continues as

$$\begin{aligned} \sum_{i=1}^2 \frac{c_i}{2} \frac{M_i}{(m_i^+)^{1/3}} &\leq \sum_{k \geq 1} E(\Omega_{1,k}, \Omega_{2,k}) \\ &\leq \sum_{i=1}^2 \left[\sqrt[3]{36\pi} M_i^{2/3} + \gamma_{ii} \int_{B_{M_i} \times B_{M_i}} |x - y|^{-1} dx dy \right], \end{aligned}$$

hence

$$(m_i^+)^{1/3} \geq \frac{c_i M_i}{2 \sum_{i=1}^2 [\sqrt[3]{36\pi} M_i^{2/3} + \gamma_{ii} \int_{B_{M_i} \times B_{M_i}} |x - y|^{-1} dx dy]},$$

and the proof is complete. \square

Lemma 4. *Consider an optimal configuration, made of clusters $\Omega_{i,k}$, $i = 1, 2$, $k \geq 1$. Assume $\sup_k |\Omega_{1,k}|$ and $\sup_k |\Omega_{2,k}|$ are achieved on different clusters, i.e., without loss of generality,*

$$|\Omega_{1,1}| = m_1^+ = \sup_k |\Omega_{1,k}|, \quad |\Omega_{2,2}| = n_2^+ = \sup_k |\Omega_{2,k}|.$$

Then

$$\inf_k \sum_{i=1}^2 |\Omega_{i,k}|$$

is bounded from below by a constant depending only on M_i , γ_{ii} , $i = 1, 2$.

Proof. Consider a cluster $(\Omega_{1,k}, \Omega_{2,k})$, with $k \geq 3$, and let

$$|\Omega_{2,1}| =: m_2, \quad |\Omega_{1,2}| = n_1, \quad \varepsilon_i := \Omega_{i,k} > 0, \quad i = 1, 2.$$

Note that $m_1^+ \geq n_1$, $n_2^+ \geq m_2$. The construction will be slightly different depending on the values of $\frac{m_1^+}{m_2}$, $\frac{m_2}{n_2^+}$, and $\frac{\varepsilon_1}{\varepsilon_2}$.

Case 1: $\frac{m_1^+}{m_2} \geq \frac{\varepsilon_1}{\varepsilon_2}$. Consider the competitor constructed in the following way (see Figure 1).

- Move mass ε_1 (resp. $r m_2$, with $r := \frac{\varepsilon_1}{m_1^+} \leq 1$) of type I (resp. type II) constituent from the cluster $(\Omega_{1,k}, \Omega_{2,k})$ to $(\Omega_{1,1}, \Omega_{2,1})$. This is possible since we are discussing the case $\frac{m_1^+}{m_2} \geq \frac{\varepsilon_1}{\varepsilon_2}$, i.e. $r m_2 = \varepsilon_1 \frac{m_2}{m_1^+} \leq \varepsilon_2$.
- Replace $(\Omega_{1,k}, \Omega_{2,k})$ and $(\Omega_{1,1}, \Omega_{2,1})$ with $B_{\varepsilon_2 - r m_2}$ (of type II constituent) and $(\tilde{\Omega}_{1,1}, \tilde{\Omega}_{2,1}) := (1 + r)^{1/3} (\Omega_{1,1}, \Omega_{2,1})$ respectively, while every other cluster remains unaltered.

Now we estimate the change in energy. Since our initial configuration was optimal,

$$\begin{aligned} 0 &\leq E((1 + r)^{1/3} (\Omega_{1,1}, \Omega_{2,1})) + E(\emptyset, B_{\varepsilon_2 - r m_2}) \\ &\quad - E(\Omega_{1,1}, \Omega_{2,1}) - E(\Omega_{1,k}, \Omega_{2,k}). \end{aligned} \tag{4}$$




FIGURE 1. Schematic representation of the construction of the competitor: original clusters (top), and modified clusters (bottom). Though the objects in question are three dimensional, for better clarity, we represented the construction in two dimensions. Only the affected clusters are represented here. The clusters are drawn deliberately deformed, to emphasize the fact that we do not know the clusters' precise shapes.

By a straightforward scaling argument,

$$\begin{aligned}
& E((1+r)^{1/3}(\Omega_{1,1}, \Omega_{2,1})) \\
&= (1+r)^{2/3} \sum_{0 \leq i < j \leq 2} \mathcal{H}^2(\partial\Omega_{i,1} \cap \partial\Omega_{j,1}), \quad \Omega_{0,1} := (\Omega_{1,1} \cup \Omega_{2,1})^c, \\
&\quad + (1+r)^{5/3} \sum_{i,j=1}^2 \gamma_{ij} \int_{\Omega_{i,1} \times \Omega_{j,1}} |x-y|^{-1} dx dy \\
&\leq (1+r) \sum_{0 \leq i < j \leq 2} \mathcal{H}^2(\partial\Omega_{i,1} \cap \partial\Omega_{j,1}) + (1+3r) \sum_{i,j=1}^2 \gamma_{ij} \int_{\Omega_{i,1} \times \Omega_{j,1}} |x-y|^{-1} dx dy \\
&\leq (1+3r) \underbrace{\left[\sum_{0 \leq i < j \leq 2} \mathcal{H}^2(\partial\Omega_{i,1} \cap \partial\Omega_{j,1}) + \sum_{i,j=1}^2 \gamma_{ij} \int_{\Omega_{i,1} \times \Omega_{j,1}} |x-y|^{-1} dx dy \right]}_{=E(\Omega_{1,1}, \Omega_{2,1})},
\end{aligned}$$

where we used the estimates

$$(1+r)^{2/3} \leq 1+r \leq 1+3r, \quad (1+r)^{5/3} \leq (1+r)^2 \stackrel{(r \leq 1)}{\leq} 1+3r.$$

Thus, in view of Lemma 3,

$$\begin{aligned} E((1+r)^{1/3}(\Omega_{1,1}, \Omega_{2,1})) - E(\Omega_{1,1}, \Omega_{2,1}) &\leq 3rE(\Omega_{1,1}, \Omega_{2,1}) \leq \varepsilon_1 H_1(M_1, M_2, \gamma_{11}, \gamma_{22}), \\ H_1(M_1, M_2, \gamma_{11}, \gamma_{22}) &:= \sum_{i=1}^2 \frac{3}{m_1^+} \left[\sqrt[3]{36\pi} M_i^{2/3} + \gamma_{ii} \int_{B_{M_i} \times B_{M_i}} |x-y|^{-1} dx dy \right]. \end{aligned} \quad (5)$$

Now we estimate $E(\emptyset, B_{\varepsilon_2 - rm_2}) - E(\Omega_{1,k}, \Omega_{2,k})$:

$$\begin{aligned} E(\emptyset, B_{\varepsilon_2 - rm_2}) - E(\Omega_{1,k}, \Omega_{2,k}) &\leq \mathcal{S}(0, \varepsilon_2 - rm_2) - \mathcal{S}(\varepsilon_1, \varepsilon_2) \\ &= \mathcal{S}(0, \varepsilon_2 - rm_2) - \mathcal{S}(\varepsilon_1, \varepsilon_2 - rm_2) + \mathcal{S}(\varepsilon_1, \varepsilon_2 - rm_2) - \mathcal{S}(\varepsilon_1, \varepsilon_2) \\ &\leq -c_1 \varepsilon_1^{2/3}, \quad c_1 := \frac{\sqrt[3]{36\pi}}{2}, \end{aligned}$$

where the last line is due to [11, Theorem 3.2], which gives

$$\mathcal{S}(\varepsilon_1, \varepsilon_2 - rm_2) - \mathcal{S}(\varepsilon_1, \varepsilon_2) \leq 0,$$

and [11, Theorem 4.2] (applied with $v_1 = \varepsilon_1$, $x = v_2 = \varepsilon_2 - rm_2$, $n = 3$), which gives

$$\begin{aligned} \mathcal{S}(\varepsilon_1, \varepsilon_2 - rm_2) &\geq \frac{\sqrt[3]{36\pi}}{2} [\varepsilon_1^{2/3} + (\varepsilon_2 - rm_2)^{2/3} + (\varepsilon_1 + \varepsilon_2 - rm_2)^{2/3}] \\ &\geq \frac{\sqrt[3]{36\pi}}{2} [\varepsilon_1^{2/3} + 2(\varepsilon_2 - rm_2)^{2/3}] = \frac{\sqrt[3]{36\pi}}{2} \varepsilon_1^{2/3} + \underbrace{\sqrt[3]{36\pi}(\varepsilon_2 - rm_2)^{2/3}}_{=\mathcal{S}(0, \varepsilon_2 - rm_2)}. \end{aligned}$$

Combining with (4) and (5) gives the necessary condition

$$\begin{aligned} 0 &\leq E((1+r)^{1/3}(\Omega_{1,1}, \Omega_{2,1})) + E(\emptyset, B_{\varepsilon_2 - rm_2}) - E(\Omega_{1,1}, \Omega_{2,1}) - E(\Omega_{1,k}, \Omega_{2,k}) \\ &\leq \varepsilon_1 H_1(M_1, M_2, \gamma_{11}, \gamma_{22}) - c_1 \varepsilon_1^{2/3}, \end{aligned} \quad (6)$$

hence

$$\varepsilon_1^{1/3} \geq H_1(M_1, M_2, \gamma_{11}, \gamma_{22}) c_1^{-1},$$

thus completing the proof for this case.

Case 2: $\frac{n_2^+}{n_1} \geq \frac{\varepsilon_2}{\varepsilon_1}$. The competitor constructed in a way similar to the previous case.

- Move mass ε_2 (resp. rn_1 , with $r := \frac{\varepsilon_2}{n_2^+} \leq 1$) of type II (resp. type I) constituent from the cluster $(\Omega_{1,k}, \Omega_{2,k})$ to $(\Omega_{1,2}, \Omega_{2,2})$. This is possible since we are discussing the case $\frac{n_2^+}{n_1} \geq \frac{\varepsilon_2}{\varepsilon_1}$, i.e. $rn_1 = \varepsilon_2 \frac{n_1}{n_2^+} \leq \varepsilon_1$.
- Replace $(\Omega_{1,k}, \Omega_{2,k})$ and $(\Omega_{1,2}, \Omega_{2,2})$ with $B_{\varepsilon_1 - rn_1}$ (of type I constituent) and $(1+r)^{1/3}(\Omega_{1,2}, \Omega_{2,2})$ respectively, while every other cluster remains unaltered.

Then the proof proceeds like in the previous case. With the same arguments from Case 1, we obtain

$$\begin{aligned} E((1+r)^{1/3}(\Omega_{1,2}, \Omega_{2,2})) - E(\Omega_{1,2}, \Omega_{2,2}) &\leq 3rE(\Omega_{1,2}, \Omega_{2,2}) \leq \varepsilon_2 H_2(M_1, M_2, \gamma_{11}, \gamma_{22}), \\ H_2(M_1, M_2, \gamma_{11}, \gamma_{22}) &:= \sum_{i=1}^2 \frac{3}{n_2^+} \left[\sqrt[3]{36\pi} M_i^{2/3} + \gamma_{ii} \int_{B_{M_i} \times B_{M_i}} |x-y|^{-1} dx dy \right], \end{aligned}$$

which is the analogue of (5), and

$$\begin{aligned} 0 &\leq E((1+r)^{1/3}(\Omega_{1,2}, \Omega_{2,2})) + E(\emptyset, B_{\varepsilon_1 - rn_1}) - E(\Omega_{1,2}, \Omega_{2,2}) - E(\Omega_{1,k}, \Omega_{2,k}) \\ &\leq \varepsilon_2 H_2(M_1, M_2, \gamma_{11}, \gamma_{22}) - c_2 \varepsilon_2^{2/3}, \end{aligned}$$

for some computable, purely geometric constant $c_2 > 0$, which is the analogue of (6). Thus

$$\varepsilon_2^{1/3} \geq H_2(M_1, M_2, \gamma_{11}, \gamma_{22}) c_2^{-1},$$

concluding the proof for this case.

Finally, note that the above two cases are exhaustive: if Case 1 does not hold, i.e. $\frac{\varepsilon_2}{\varepsilon_1} < \frac{m_2}{m_1^+}$, using $m_1^+ \geq n_1$, $n_2^+ \geq m_2$, we get

$$\frac{\varepsilon_2}{\varepsilon_1} < \frac{m_2}{m_1^+} \leq \frac{n_2^+}{n_1},$$

i.e. Case 2 holds. The proof is thus complete. \square

Lemma 5. *Consider an optimal configuration, made of clusters $\Omega_{i,k}$, $i = 1, 2$, $k \geq 1$. Assume $\sup_k |\Omega_{1,k}|$ and $\sup_k |\Omega_{2,k}|$ are achieved on the same clusters, i.e., without loss of generality,*

$$|\Omega_{i,1}| = m_i^+ = \sup_k |\Omega_{i,k}|, \quad i = 1, 2.$$

Then

$$\inf_k \sum_{i=1}^2 |\Omega_{i,k}|$$

is again bounded from below by a constant depending only on M_i , γ_{ii} , $i = 1, 2$.

Proof. We rely on Lemma 4: Consider another cluster $(\Omega_{1,k}, \Omega_{2,k})$, $k \geq 2$. Let $|\Omega_{1,k}| = \varepsilon_1 > 0$, $|\Omega_{2,k}| = \varepsilon_2 > 0$, and note that one of the following cases must hold.

- (1) If $\frac{m_1^+}{m_2^+} \geq \frac{\varepsilon_1}{\varepsilon_2}$, then we can use the construction from Case 1 of Lemma 4.
- (2) If $\frac{m_1^+}{m_2^+} \leq \frac{\varepsilon_1}{\varepsilon_2}$, i.e. $\frac{m_2^+}{m_1^+} \geq \frac{\varepsilon_2}{\varepsilon_1}$, then we can use the construction from Case 2 of Lemma 4.

The proof is thus complete. \square

ACKNOWLEDGMENTS

XYL acknowledges the support of NSERC. The research of J. Wei is partially supported by NSERC. We are grateful to Chong Wang for useful discussions and suggestions.

REFERENCES

- [1] S. ALAMA, L. BRONSARD, X. LU, AND C. WANG, *Periodic minimizers of a ternary non-local isoperimetric problem*, Indiana Univ. Math. J., (2021).
- [2] M. BONACINI AND R. CRISTOFERI, *Local and global minimality results for a nonlocal isoperimetric problem on R^N* , SIAM Journal on Mathematical Analysis, 46 (2014), pp. 2310–2349.
- [3] R. CHOKSI, C. B. MURATOV, AND I. TOPALOGLU, *An old problem resurfaces nonlocally: Gamow's liquid drops inspire today's research and applications*, Notices of the AMS, 64 (2017), pp. 1275–1283.
- [4] R. CHOKSI AND M. A. PELETIER, *Small volume fraction limit of the diblock copolymer problem: I. sharp-interface functional*, SIAM Journal on Mathematical Analysis, 42 (2010), pp. 1334–1370.

- [5] ———, *Small volume-fraction limit of the diblock copolymer problem: II. Diffuse-interface functional*, SIAM Journal on Mathematical Analysis, 43 (2011), pp. 739–763.
- [6] R. CHOKSI AND X. REN, *On the derivation of a density functional theory for microphase separation of diblock copolymers*, Journal of Statistical Physics, 113 (2003), pp. 151–176.
- [7] R. DORFF, G. LAWLOR, D. SAMPSON, AND B. WILSON, *Proof of the planar double bubble conjecture using metacalibration methods*, Involve, a Journal of Mathematics, 2 (2010), pp. 611–628.
- [8] J. FOISY, M. ALFARO GARCIA, J. BROCK, N. HODGES, AND J. ZIMBA, *The standard double soap bubble in R^2 uniquely minimizes perimeter*, Pacific journal of mathematics, 159 (1993), pp. 47–59.
- [9] R. L. FRANK AND E. H. LIEB, *A compactness lemma and its application to the existence of minimizers for the liquid drop model*, SIAM Journal on Mathematical Analysis, 47 (2015), pp. 4436–4450.
- [10] G. GAMOW, *Mass defect curve and nuclear constitution*, Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character, 126 (1930), pp. 632–644.
- [11] M. HUTCHINGS, *The structure of area-minimizing double bubbles*, The Journal of Geometric Analysis, 7 (1997), pp. 285–304.
- [12] M. HUTCHINGS, F. MORGAN, M. RITORÉ, AND A. ROS, *Proof of the double bubble conjecture*, Annals of Mathematics, (2002), pp. 459–489.
- [13] H. KNÜPFER AND C. B. MURATOV, *On an isoperimetric problem with a competing nonlocal term I: The planar case*, Communications on Pure and Applied Mathematics, 66 (2013), pp. 1129–1162.
- [14] H. KNÜPFER, C. B. MURATOV, AND M. NOVAGA, *Low density phases in a uniformly charged liquid*, Communications in Mathematical Physics, 345 (2016), pp. 141–183.
- [15] ———, *Emergence of nontrivial minimizers for the three-dimensional Ohta–Kawasaki energy*, Pure and Applied Analysis, 2 (2019), pp. 1–21.
- [16] E. MILMAN AND J. NEEMAN, *The Gaussian double-bubble conjecture*, arXiv preprint arXiv:1801.09296, (2018).
- [17] F. MORGAN AND M. RITORÉ, *Geometric measure theory and the proof of the double bubble conjecture*, in Proc. Clay Research Institution Summer School, MSRI, 2001.
- [18] F. MORGAN AND W. WICHIRAMALA, *The standard double bubble is the unique stable double bubble in R^2* , Proceedings of the American Mathematical Society, 130 (2002), pp. 2745–2751.
- [19] C. MURATOV AND H. KNÜPFER, *On an isoperimetric problem with a competing nonlocal term II: The general case*, Communications on Pure and Applied Mathematics, 67 (2014), pp. 1974–1994.
- [20] H. NAKAZAWA AND T. OHTA, *Microphase separation of ABC-type triblock copolymers*, Macromolecules, 26 (1993), pp. 5503–5511.
- [21] T. OHTA AND K. KAWASAKI, *Equilibrium morphology of block copolymer melts*, Macromolecules, 19 (1986), pp. 2621–2632.
- [22] B. W. REICHARDT, C. HEILMANN, Y. Y. LAI, AND A. SPIELMAN, *Proof of the double bubble conjecture in R^4 and certain higher dimensional cases*, Pacific journal of mathematics, 208 (2003), pp. 347–366.
- [23] X. REN AND J. WEI, *Triblock copolymer theory: Free energy, disordered phase and weak segregation*, Physica D: Nonlinear Phenomena, 178 (2003), pp. 103–117.
- [24] X. REN AND J. WEI, *Triblock copolymer theory: Ordered ABC lamellar phase*, Journal of Nonlinear Science, 13 (2003), pp. 175–208.

DEPARTMENT OF MATHEMATICAL SCIENCES, LAKEHEAD UNIVERSITY, THUNDER BAY, ON, CANADA
 Email address: XLU8@LAKEHEADU.CA

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC, CANADA
 Email address: JCWEI@MATH.UBC.CA