

ON THE ANTI-CANONICAL GEOMETRY OF WEAK \mathbb{Q} -FANO THREEFOLDS, III

CHEN JIANG AND YU ZOU

ABSTRACT. For a terminal weak \mathbb{Q} -Fano 3-fold X , we show that the m -th anti-canonical map defined by $|-mK_X|$ is birational for all $m \geq 59$.

1. INTRODUCTION

Throughout this paper, we work over an algebraically closed field of characteristic 0 (for instance, the complex number field \mathbb{C}). We adopt standard notation in [14].

A normal projective variety X is called a *weak \mathbb{Q} -Fano variety* (resp. *\mathbb{Q} -Fano variety*) if $-K_X$ is nef and big (resp. ample). According to the minimal model program, (weak) \mathbb{Q} -Fano varieties form a fundamental class in birational geometry. Motivated by the classification theory of 3-dimensional algebraic varieties, we are interested in the study of explicit geometry of (weak) \mathbb{Q} -Fano varieties with terminal or canonical singularities. In this direction there are a lot of works in the literature, see for example [16, 17, 3, 18, 4, 19, 5, 6, 12, 10, 11].

Given a terminal weak \mathbb{Q} -Fano 3-fold X , the m -th *anti-canonical map* $\varphi_{-m,X}$ (or simply φ_{-m}) is the rational map induced by the linear system $|-mK_X|$. We are interested in the fundamental question of finding an optimal integer c_3 such that φ_{-m} is birational for all $m \geq c_3$. The existence of such c_3 follows from the boundedness result in [13]. More generally, Birkar [1] showed that, fix a positive integer d , there exists a positive integer c_d such that φ_{-m} is birational for all $m \geq c_d$ and for all terminal weak \mathbb{Q} -Fano d -folds, which is one important step towards the solution of the Borisov–Alexeev–Borisov conjecture. The following example shows that $c_3 \geq 33$.

Example 1.1 ([8, List 16.6, No. 95]). A general weighted hypersurface $X_{66} \subset \mathbb{P}(1, 5, 6, 22, 33)$ is a \mathbb{Q} -factorial terminal \mathbb{Q} -Fano 3-fold of Picard number 1 with φ_{-m} birational for $m \geq 33$ but φ_{-32} not birational.

In [5], it was showed that for a terminal weak \mathbb{Q} -Fano 3-fold X , φ_{-m} is birational for all $m \geq 97$, which seems far from being optimal comparing to Example 1.1. Later in [6], it was showed that any terminal weak \mathbb{Q} -Fano 3-fold is birational to some terminal weak \mathbb{Q} -Fano 3-fold Y such that $\varphi_{-m,Y}$ is birational for all $m \geq 52$. Moreover, in recent works [10, 11], we can make use of the behavior of the pluri-anti-canonical maps studied in [5] in the classification of terminal \mathbb{Q} -Fano 3-folds. So we believe that a better

Date: August 1, 2023.

2020 Mathematics Subject Classification. 14J45, 14J30, 14C20, 14E05.

Key words and phrases. weak \mathbb{Q} -Fano 3-fold, anti-pluricanonical map, birationality.

understanding of the behavior of the pluri-anti-canonical maps (including new methods developed during the approach) will help us understand the classification of terminal \mathbb{Q} -Fano 3-folds better.

The main goal of this paper is to give an improvement of [5, 6] without passing to a birational model. The main theorem of this paper is the following.

Theorem 1.2. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. Then the m -th anti-canonical map φ_{-m} defined by $|-mK_X|$ is birational for all $m \geq 59$.*

Remark 1.3. Theorem 1.2 holds for canonical weak \mathbb{Q} -Fano 3-folds by taking a \mathbb{Q} -factorial terminalization by [14, Theorem 6.23, Theorem 6.25].

For terminal \mathbb{Q} -Fano 3-folds, we have a slightly better bound.

Theorem 1.4. *Let X be a terminal \mathbb{Q} -Fano 3-fold. Then the m -th anti-canonical map φ_{-m} defined by $|-mK_X|$ is birational for all $m \geq 58$.*

To prove the main theorem, we already have several criteria to determine the birationality in [5, 6], which are optimal in many cases (cf. [5, Example 5.12]). In order to study the birationality of $|-mK_X|$, as indicated in [4, 5, 6], it is crucial to study when $|-mK_X|$ is not composed with a pencil. In fact, finding a criterion for $|-mK_X|$ not composed with a pencil is one of the central problems in [5, 6] ([5, Problem 1.3], [6, Problem 1.5]). Comparing to the birationality criteria, the non-pencil criteria in [5, 6] are not satisfactory. As one of the main ingredients of this paper, we give a new criterion for $|-mK_X|$ not composed with a pencil.

Theorem 1.5 (=Theorem 4.2). *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. If*

$$h^0(X, -mK_X) > 12m + 1$$

for some positive integer m , then $|-mK_X|$ is not composed with a pencil.

The following special case is already interesting for the study of anti-canonical systems of terminal weak \mathbb{Q} -Fano 3-folds, and might have applications on upper bounds of degrees of terminal weak \mathbb{Q} -Fano 3-folds (cf. [16, 17]).

Corollary 1.6. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. If $h^0(X, -K_X) > 13$, then $|-K_X|$ is not composed with a pencil.*

The paper is organized as the following. In Section 2, we recall basic knowledge. In Section 3, we recall the birationality criteria of terminal weak \mathbb{Q} -Fano 3-folds in [5, 6] with some generalizations. In Section 4, we prove the new criterion Theorem 4.2 and give an effective method to apply it. In Section 5, we prove the main theorem.

Notation

For the convenience of readers, we list here the notation that will be frequently used in this paper. Let X be a terminal weak \mathbb{Q} -Fano 3-fold.

| | |
|--------------------------|---------------------------------------|
| φ_{-m} | the rational map defined by $ -mK_X $ |
| $P_{-m} = h^0(X, -mK_X)$ | the m -th anti-plurigenus of X |

| | |
|--|--|
| $B_X = \{(b_i, r_i)\}$ | the Reid basket of orbifold points of X |
| $\mathcal{R}_X = \{r_i\}$ | the collection of local indices of X |
| $r_X = \text{lcm}\{r_i \mid r_i \in \mathcal{R}_X\}$ | the Cartier index of K_X |
| $r_{\max} = \max\{r_i \mid r_i \in \mathcal{R}_X\}$ | the maximal local index of X |
| $\sigma(B_X) = \sum_i b_i$ | an invariant of B_X contributing to the Riemann-Roch formula |
| $\sigma'(B_X) = \sum_i \frac{b_i^2}{r_i}$ | an invariant of B_X contributing to the Riemann-Roch formula |
| $\gamma(B_X) = \sum_i \frac{1}{r_i} - \sum_i r_i + 24$ | an invariant of B_X from the Miyaoka inequality |
| $B_X^{(0)} = \{n_{1,r}^0 \times (1, r)\}$ | the initial basket of B_X |

2. PRELIMINARIES

Let X be a terminal weak \mathbb{Q} -Fano 3-fold. Denote by r_X the Cartier index of K_X . For any positive integer m , the number $P_{-m} = h^0(X, \mathcal{O}_X(-mK_X))$ is called the m -th *anti-plurigenus* of X and φ_{-m} denotes the m -th *anti-canonical map* defined by $|-mK_X|$.

2.1. The fibration induced by $|D|$. Let X be a terminal weak \mathbb{Q} -Fano 3-fold. Consider a \mathbb{Q} -Cartier Weil divisor D on X with $h^0(X, D) \geq 2$. Then there is a rational map defined by $|D|$:

$$\Phi_{|D|} : X \dashrightarrow \mathbb{P}^{h^0(X, D)-1}.$$

By Hironaka's desingularization theorem, we can take a projective birational morphism $\pi : W \rightarrow X$ such that:

- (i) W is smooth;
- (ii) the movable part $|M|$ of $|\pi^*(D)|$ is base point free and, consequently, $\gamma := \Phi_{|D|} \circ \pi$ is a morphism;
- (iii) the sum of $\pi_*^{-1}(D)$ and the exceptional divisors of π has simple normal crossing support.

Let $W \xrightarrow{f} \Gamma \xrightarrow{s} Z$ be the Stein factorization of γ with $Z := \gamma(W) \subset \mathbb{P}^{h^0(X, D)-1}$. We have the following commutative diagram:

$$\begin{array}{ccc}
 W & \xrightarrow{f} & \Gamma \\
 \pi \downarrow & \searrow \gamma & \downarrow s \\
 X & \xrightarrow{\Phi_{|D|}} & Z.
 \end{array}$$

If $\dim(\Gamma) \geq 2$, then a general member S of $|M|$ is a smooth projective surface by Bertini's theorem. In this case, $|D|$ is said to be *not composed with a pencil of surfaces* (not composed with a pencil, for short).

If $\dim(\Gamma) = 1$, then $\Gamma \cong \mathbb{P}^1$ as $h^1(\Gamma, \mathcal{O}_\Gamma) \leq h^1(W, \mathcal{O}_W) = h^1(X, \mathcal{O}_X) = 0$. Furthermore, a general fiber S of f is a smooth projective surface by Bertini's theorem. In this case, $|D|$ is said to be *composed with a (rational) pencil of surfaces* (composed with a pencil, for short).

In each case, S is called a *generic irreducible element* of $|M|$. We can also define a generic irreducible element of a moving linear system on a surface in the similar way.

Definition 2.1. Keep the same notation as above. Let D' be another \mathbb{Q} -Cartier Weil divisor on X with $h^0(X, D') \geq 2$. We say that $|D|$ and $|D'|$ are *composed with the same pencil*, if both of them are composed with pencils and they define the same fibration structure $W \rightarrow \mathbb{P}^1$. In particular, $|D|$ and $|D'|$ are not composed with the same pencil if one of them is not composed with a pencil.

2.2. Reid's Riemann–Roch formula and Chen–Chen's method. A basket B is a collection of pairs of coprime integers where a pair is allowed to appear several times, say

$$\{(b_i, r_i) \mid i = 1, \dots, s; b_i \text{ is coprime to } r_i\}.$$

For simplicity, we will alternatively write a basket as a set of pairs with weights, say for example,

$$B = \{2 \times (1, 2), (1, 3), (3, 7), (5, 11)\}.$$

Let X be a terminal weak \mathbb{Q} -Fano 3-fold. According to Reid [20], there is a basket of (virtual) orbifold points

$$B_X = \{(b_i, r_i) \mid i = 1, \dots, s; 0 < b_i \leq \frac{r_i}{2}; b_i \text{ is coprime to } r_i\}$$

associated to X , where a pair (b_i, r_i) corresponds to an orbifold point Q_i of type $\frac{1}{r_i}(1, -1, b_i)$. Denote by \mathcal{R}_X the collection of r_i (counted with multiplicities) appearing in B_X , and $r_{\max} = \max\{r_i \mid r_i \in \mathcal{R}_X\}$. Note that the Cartier index r_X of K_X is just $\text{lcm}\{r_i \mid r_i \in \mathcal{R}_X\}$.

According to Reid [20], for any positive integer n ,

$$P_{-n} = \frac{1}{12}n(n+1)(2n+1)(-K_X^3) + (2n+1) - l(n+1) \quad (2.1)$$

where $l(n+1) = \sum_i \sum_{j=1}^n \frac{\overline{jb_i}(r_i - \overline{jb_i})}{2r_i}$ and the first sum runs over Reid's basket of orbifold points. Here $\overline{jb_i}$ means the smallest non-negative residue of $jb_i \bmod r_i$.

Set $\sigma(B_X) = \sum_i b_i$ and $\sigma'(B_X) = \sum_i \frac{b_i^2}{r_i}$. From (2.1) for $n = 1, 2$,

$$-K_X^3 = 2P_{-1} + \sigma(B_X) - \sigma'(B_X) - 6; \quad (2.2)$$

$$\sigma(B_X) = 10 - 5P_{-1} + P_{-2}. \quad (2.3)$$

Denote

$$\gamma(B_X) := \sum_i \frac{1}{r_i} - \sum_i r_i + 24.$$

By [13] and [20, 10.3],

$$\gamma(B_X) \geq 0. \quad (2.4)$$

We recall Chen–Chen's method on basket packing from [3]. Given a basket

$$B = \{(b_i, r_i) \mid i = 1, \dots, s; 0 < b_i \leq \frac{r_i}{2}; b_i \text{ is coprime to } r_i\}$$

and assume that $b_1 r_2 - b_2 r_1 = 1$, then the new basket

$$B' = \{(b_1 + b_2, r_1 + r_2), (b_3, r_3), \dots, (b_s, r_s)\}$$

is called a *prime packing* of B . We say that a basket B' is *dominated by* B , denoted by $B \succeq B'$, if B' can be achieved from B by a sequence of prime packings (including the case $B = B'$).

By [3, §2.5], there is a unique basket $B_X^{(0)}$, called *the initial basket of* B_X , of the form $B_X^{(0)} = \{n_{1,r}^0 \times (1, r) \mid r \geq 2\}$ such that $B_X^{(0)} \succeq B_X$. By [3, §2.7], we have

$$n_{1,2}^0 = 5 - 6P_{-1} + 4P_{-2} - P_{-3}; \quad (2.5)$$

$$n_{1,3}^0 = 4 - 2P_{-1} - 2P_{-2} + 3P_{-3} - P_{-4}; \quad (2.6)$$

$$n_{1,4}^0 = 1 + 3P_{-1} - P_{-2} - 2P_{-3} + P_{-4} - \sigma_5, \quad (2.7)$$

where $\sigma_5 = \sum_{r \geq 5} n_{1,r}^0$. We refer to [3] for more details.

2.3. Auxiliary results. We list here some useful results on terminal weak \mathbb{Q} -Fano 3-folds.

Proposition 2.2. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. Then*

- (1) $r_X = 840$ or $r_X \leq 660$ ([5, Proposition 2.4]);
- (2) $P_{-8} \geq 2$ ([3, Theorem 1.1]); moreover, if $P_{-1} = 0$ and $P_{-2} > 0$, then $P_{-6} \geq 2$ ([3, Case 1 of Proof of Proposition 3.10]);
- (3) $-K_X^3 \geq \frac{1}{330}$ ([3, Theorem 1.1]); moreover, if $P_{-1} = 0$ and $P_{-2} > 0$, then $-K_X^3 \geq \frac{1}{70}$, and if in addition $P_{-4} \geq 2$, then $-K_X^3 \geq \frac{1}{30}$ ([3, (4.1), Lemma 4.2, Case I of Proof of Theorem 4.4]);
- (4) if $P_{-1} = 0$, then $2 \in \mathcal{R}_X$ ([5, Proof of Theorem 1.8, Page 106]).

Lemma 2.3. *Suppose that $\{(b_i, r_i) \mid 1 \leq i \leq k\}$ is a collection of pairs of integers with $0 < 2b_i \leq r_i$ for $1 \leq i \leq k$. Then $\sum_{i=1}^k (r_i - \frac{1}{r_i}) \geq \frac{3}{2} \sum_{i=1}^k b_i$.*

Proof. $r_i \geq 2b_i$ implies that $r_i - \frac{1}{r_i} \geq \frac{3}{2}b_i$. \square

3. THE CRITERIA FOR BIRATIONALITY

In this section we recall the birationality criteria of terminal weak \mathbb{Q} -Fano 3-folds in [5, 6]. Here we remark that all birationality criteria in this section are from [5, 6] except for Theorem 3.5 and Corollary 3.7 (which are minor generalizations of [6, Theorem 5.9]). Also we provide Lemma 3.3 in order to apply Corollary 3.7 efficiently. In fact, in [6], [6, Theorem 5.9] is only used for very special cases, but in this paper, thanks to Lemma 3.3, we make use of Corollary 3.7 in many cases.

3.1. General settings. We recall numerical invariants needed in the birationality criteria, namely, $\nu_0, m_0, a(m_0), m_1, \mu'_0, N_0$.

Notation 3.1. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold.*

Let ν_0 be a positive integer such that $P_{-\nu_0} > 0$.

Take a positive integer m_0 such that $P_{-m_0} \geq 2$. Set

$$a(m_0) = \begin{cases} 6, & \text{if } m_0 \geq 2; \\ 1, & \text{if } m_0 = 1. \end{cases}$$

Take $m_1 \geq m_0$ to be an integer with $P_{-m_1} \geq 2$ such that $|-m_0 K_X|$ and $|-m_1 K_X|$ are not composed with the same pencil.

Set $D := -m_0 K_X$ and keep the same notation as in Subsection 2.1. Denote S to be a generic irreducible element of $|M_{-m_0}| = \text{Mov } |\pi^*(-m_0 K_X)|$. Choose a positive rational number μ'_0 such that

$$\mu'_0 \pi^*(-K_X) - S \sim_{\mathbb{Q}} \text{effective } \mathbb{Q}\text{-divisor}.$$

Set $N_0 = r_X(\pi^*(-K_X)^2 \cdot S)$.

Remark 3.2 ([6, Remark 5.8]). Here we explain how to choose μ'_0 . In general, by assumption, we can always take $\mu'_0 = m_0$. On the other hand, if $|-m_0 K_X|$ and $|-k K_X|$ are composed with the same pencil for some positive integer k , and $\frac{k}{P_{-k}-1} < m_0$, then we can take $\mu'_0 = \frac{k}{P_{-k}-1}$ as

$$k\pi^*(-K_X) \sim_{\mathbb{Q}} (P_{-k} - 1)S + \text{effective } \mathbb{Q}\text{-divisor}.$$

Lemma 3.3. *In Notation 3.1, $N_0 \geq \lceil \frac{r_X}{m_1 \nu_0 r_{\max}} \rceil$.*

Proof. We may modify π such that $|M_{-m_1}| = \text{Mov } |\pi^*(-m_1 K_X)|$ is base point free. Pick a generic irreducible element C of the base point free linear system $|M_{-m_1}|_S$. Since $\pi^*(-m_1 K_X) \geq M_{-m_1}$, $\pi^*(-m_1 K_X)|_S \geq C$. Set

$$\zeta := (\pi^*(-K_X) \cdot C) = (\pi^*(-K_X)|_S \cdot C)_S.$$

By [5, Proposition 5.7(v)], $\zeta \geq \frac{1}{\nu_0 r_{\max}}$. Since $\pi^*(-K_X)|_S$ is nef,

$$\pi^*(-K_X)^2 \cdot S \geq \pi^*(-K_X)|_S \cdot \frac{1}{m_1} C \geq \frac{1}{m_1 \nu_0 r_{\max}}.$$

Hence $N_0 \geq \lceil \frac{r_X}{m_1 \nu_0 r_{\max}} \rceil$ as N_0 is an integer by [5, Lemma 4.1]. \square

3.2. Birationality criteria. We recall the birationality criteria of terminal weak \mathbb{Q} -Fano 3-folds.

Theorem 3.4 ([5, Theorem 5.11]). *Keep the setting in Notation 3.1. Then the m -th anti-canonical map φ_{-m} is birational if one of the following conditions holds:*

- (1) $m \geq \max\{m_0 + m_1 + a(m_0), \lfloor 3\mu'_0 \rfloor + 3m_1\}$;
- (2) $m \geq \max\{m_0 + m_1 + a(m_0), \lfloor \frac{5}{3}\mu'_0 + \frac{5}{3}m_1 \rfloor, \lfloor \mu'_0 \rfloor + m_1 + 2r_{\max}\}$;
- (3) $m \geq \max\{m_0 + m_1 + a(m_0), \lfloor \mu'_0 \rfloor + m_1 + 2\nu_0 r_{\max}\}$.

As another criterion, we have the following modification of [6, Theorem 5.9].

Theorem 3.5. *Keep the setting in Notation 3.1. Fix a real number $\beta \geq 8$. Then the m -th anti-canonical map φ_{-m} is birational if*

$$m \geq \max \left\{ m_0 + a(m_0), \left\lceil \mu'_0 + \frac{4\nu_0 r_{\max}}{1 + \sqrt{1 - \frac{8}{\beta}}} \right\rceil - 1, \lfloor \mu'_0 + \sqrt{\beta r_X / N_0} \rfloor \right\}.$$

Proof. The proof is the same as [6, Theorem 5.9] by replacing [6, Lemma 5.10] with Lemma 3.6. \square

Lemma 3.6 ([2, Theorem 2.8]). *Let S be a smooth projective surface and L be a nef and big \mathbb{Q} -divisor on S satisfying the following conditions:*

- (1) $L^2 > \beta$, for some real number $\beta \geq 8$, and

- (2) $(L \cdot C_P) \geq \frac{4}{1+\sqrt{1-\frac{8}{\beta}}}$ for all irreducible curves C_P passing through any very general point $P \in S$.

Then the linear system $|K_S + \lceil L \rceil|$ separates two distinct points in very general positions. Consequently, $|K_S + \lceil L \rceil|$ gives a birational map.

We will use the following version of Theorem 3.5.

Corollary 3.7. *Keep the setting in Notation 3.1. Then the m -th anti-canonical map φ_{-m} is birational if one of the following conditions holds:*

- (1) $m \geq \max\{m_0 + a(m_0), \lceil \mu'_0 \rceil + 4\nu_0 r_{\max} - 1, \lfloor \mu'_0 + \sqrt{8r_X/N_0} \rfloor\}$;
(2) $\nu_0 r_{\max} \geq \sqrt{\frac{r_X}{2N_0}}$ and

$$m \geq \max \left\{ m_0 + a(m_0), \left\lfloor \mu'_0 + 2\nu_0 r_{\max} + \frac{r_X}{N_0 \nu_0 r_{\max}} \right\rfloor \right\}.$$

Proof. (1) follows directly from [6, Theorem 5.9] or Theorem 3.5 with $\beta = 8$. For (2), take

$$\beta = \frac{N_0}{r_X} \left(2\nu_0 r_{\max} + \frac{r_X}{N_0 \nu_0 r_{\max}} \right)^2 \geq 8$$

in Theorem 3.5. Then

$$\begin{aligned} \frac{4\nu_0 r_{\max}}{1 + \sqrt{1 - \frac{8}{\beta}}} &= \frac{4\nu_0 r_{\max} \sqrt{\beta}}{\sqrt{\beta} + \sqrt{\beta - 8}} \\ &= \frac{4\nu_0 r_{\max} \sqrt{\beta}}{\sqrt{\frac{N_0}{r_X} \left(2\nu_0 r_{\max} + \frac{r_X}{N_0 \nu_0 r_{\max}} \right)^2} + \left| 2\nu_0 r_{\max} - \frac{r_X}{N_0 \nu_0 r_{\max}} \right|} \\ &= \sqrt{\beta r_X / N_0}. \end{aligned}$$

So the conclusion follows from Theorem 3.5. \square

Finally we explain the strategy to apply the birationality criteria to assert the birationality. It is clear that in order to apply Theorem 3.4 and Corollary 3.7, we need to control the values of (some of) ν_0 , m_0 , m_1 , μ'_0 , N_0 and r_X, r_{\max} . To be more precise, we need to give upper bounds of ν_0 , m_0 , m_1 , μ'_0 , r_X , r_{\max} and lower bounds of N_0 . Here m_0 and ν_0 can be controlled by Proposition 2.2 (in particular we can always take $m_0 = 8$), μ'_0 can be controlled by Remark 3.2, N_0 can be controlled by Lemma 3.3 (in most cases, we use the trivial lower bound $N_0 \geq 1$), and r_X, r_{\max} can be controlled by (2.4). So the most important and difficult part is to bound m_1 . We will deal with this issue in the next section.

4. A NEW CRITERION FOR $|-mK|$ NOT COMPOSED WITH A PENCIL

In this section, we give a new criterion on when $|-mK_X|$ is not composed with a pencil for a terminal weak \mathbb{Q} -Fano 3-fold X . Such a criterion is essential in order to apply criteria for birationality in Section 3 (see also [4, 5, 6]). In [5], the following proposition is used to determine when $|-mK|$ is not composed with a pencil.

Proposition 4.1 ([5, Corollary 4.2]). *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. If*

$$P_{-m} > r_X(-K_X^3)m + 1$$

for some positive integer m , then $|-mK_X|$ is not composed with a pencil.

However, Proposition 4.1 is too weak for application, especially when $r_X(-K_X^3)$ is large (see Example 4.10). In [6], there is a modification of this inequality (cf. [6, Lemma 4.2, Proposition 5.2]) but one has to replace X with a birational model. In this paper, by technique developed recently in [12], we give a new criterion.

Theorem 4.2. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. If*

$$P_{-m} > 12m + 1$$

for some positive integer m , then $|-mK_X|$ is not composed with a pencil.

4.1. A structure theorem of terminal weak \mathbb{Q} -Fano 3-folds. We recall the following structure theorem of terminal weak \mathbb{Q} -Fano 3-folds from [12]. It plays the role of Fano–Mori triples as in [6]. Unlike [6], we do not need to replace by a birational model (cf. [6, Proposition 3.9]).

Proposition 4.3 ([12, Proposition 4.1]). *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. Then there exists a normal projective 3-fold Y birational to X satisfying the following properties:*

- (1) *Y is \mathbb{Q} -factorial terminal;*
- (2) *$-K_Y$ is big;*
- (3) *for any sufficiently large and divisible positive integer n , $|-nK_Y|$ is movable;*
- (4) *for a general member $M \in |-nK_Y|$, M is irreducible and $(Y, \frac{1}{n}M)$ is canonical;*
- (5) *there exists a projective morphism $g : Y \rightarrow S$ with connected fibers where F is a general fiber of g , such that one of the following conditions holds:*
 - (a) *S is a point and Y is a \mathbb{Q} -Fano 3-fold with $\rho(Y) = 1$;*
 - (b) *$S = \mathbb{P}^1$ and F is a smooth weak del Pezzo surface;*
 - (c) *S is a del Pezzo surface with at worst Du Val singularities and $\rho(S) = 1$, and $F \simeq \mathbb{P}^1$.*

Here we remark that in the proof of [12, Proposition 4.1], Y is obtained by running a K -MMP on a \mathbb{Q} -factorialization of X , so the induced map $X \dashrightarrow Y$ is a contraction, that is, it does not extract any divisor.

4.2. Bounding coefficients of anti-canonical divisors. In this subsection, we discuss coefficients of certain divisors in the \mathbb{Q} -linear system of the anti-canonical divisor in several cases.

Lemma 4.4. *Let S be a smooth weak del Pezzo surface and C a non-zero effective integral divisor on S which is movable. If $-K_S \sim_{\mathbb{Q}} aC + B$ for some positive rational number a and some effective \mathbb{Q} -divisor B , then $a \leq 4$.*

Proof. By classical surface theory, it is well-known that there is a birational map from S to \mathbb{P}^2 or the Hirzebruch surface \mathbb{F}_0 or \mathbb{F}_2 . So by taking push-forward, we may replace S by \mathbb{P}^2 or \mathbb{F}_0 or \mathbb{F}_2 . Here C is not contracted by the pushforward as it is movable.

If $S = \mathbb{P}^2$, then intersecting with a general line L , we get $a \leq a(C \cdot L) \leq (-K_S \cdot L) = 3$.

If $S = \mathbb{F}_0$, then we may find a ruling structure $\phi : \mathbb{F}_0 \rightarrow \mathbb{P}^1$ such that C is not vertical. Then we get $a \leq 2$ by intersecting with a fiber of ϕ .

If $S = \mathbb{F}_2$, then we consider the natural ruling structure $\phi : \mathbb{F}_2 \rightarrow \mathbb{P}^1$. If C is not vertical, then intersecting with a fiber of ϕ , we get $a \leq 2$. If C is vertical, then intersecting with $-K_S$ we get $a(-K_S \cdot C) \leq K_S^2$ which implies that $a \leq 4$. □

Lemma 4.5. *Let Y be a \mathbb{Q} -factorial terminal \mathbb{Q} -Fano 3-fold with $\rho(Y) = 1$ and D an integral divisor with $h^0(D) \geq 2$. If $-K_Y \sim_{\mathbb{Q}} aD + B$ for some positive rational number a and some effective \mathbb{Q} -divisor B , then $a \leq 7$. Moreover, the equality holds if and only if $Y \simeq \mathbb{P}(1, 1, 2, 3)$, $B = 0$, and $\mathcal{O}_Y(D) \simeq \mathcal{O}_Y(1)$.*

Proof. Suppose that $a \geq 7$. As $\rho(Y) = 1$, we have $-K_Y \sim_{\mathbb{Q}} tD$ for some rational number $t \geq a \geq 7$. Recall that the \mathbb{Q} -Fano index of Y is defined by

$$q\mathbb{Q}(Y) = \max\{q \mid -K_Y \sim_{\mathbb{Q}} qA, A \text{ is a Weil divisor}\}.$$

By [18, Corollary 3.4(ii)], $t = q\mathbb{Q}(Y) \geq 7$. As $h^0(D) \geq 2$, there are two different effective divisors $D_1, D_2 \in |D|$ such that $-K_Y \sim_{\mathbb{Q}} tD_1 \sim_{\mathbb{Q}} tD_2$, which implies that $Y \simeq \mathbb{P}(1, 1, 2, 3)$ by [18, Theorem 1.4(vi)]. But in this case $t = q\mathbb{Q}(Y) = 7$. Hence $a = 7$, $B = 0$, and $\mathcal{O}_Y(D) \simeq \mathcal{O}_Y(1)$. □

Lemma 4.6. *Keep the setting in Proposition 4.3, suppose that $S = \mathbb{P}^1$. If $-K_Y \sim_{\mathbb{Q}} \omega F + E$ for some positive rational number ω and some effective \mathbb{Q} -divisor E , then $\omega \leq 12$.*

This lemma is from the proof of [12, Proposition 4.2]. For the reader's convenience, we recall the proof here.

Proof. We may assume that $\omega > 2$. By Proposition 4.3(3)(4), for a sufficiently large and divisible integer n , $|-nK_Y|$ is movable, and there exists an effective \mathbb{Q} -divisor $M \sim -nK_Y$ such that $(Y, \frac{1}{n}M)$ is canonical. Since $-K_Y$ is big, we can write $-K_Y \sim_{\mathbb{Q}} A + N$, where A is an ample \mathbb{Q} -divisor and N is an effective \mathbb{Q} -divisor. Set $B_\epsilon = \frac{1-\epsilon}{n}M + \epsilon N$ for a rational number $0 < \epsilon < 1$. Take two general fibers F_1, F_2 of g . Denote

$$\Delta = (1 - \frac{2}{\omega})B_\epsilon + \frac{2}{\omega}E + F_1 + F_2.$$

Then

$$-(K_Y + \Delta) \sim_{\mathbb{Q}} -(1 - \frac{2}{\omega})(K_Y + B_\epsilon) \sim_{\mathbb{Q}} (1 - \frac{2}{\omega})\epsilon A$$

is ample as $\omega > 2$. Hence by the connectedness lemma ([12, Lemma 2.6]), $\text{Nklt}(Y, \Delta)$ is connected. By construction, $F_1 \cup F_2 \subset \text{Nklt}(Y, \Delta)$, then $\text{Nklt}(Y, \Delta)$ dominates \mathbb{P}^1 . By the inversion of adjunction ([14, Lemma 5.50]), $(F, (1 - \frac{2}{\omega})B_\epsilon|_F + \frac{2}{\omega}E|_F)$ is not klt for a general fiber F of g . As being klt is an open condition on the coefficients, by the arbitrariness of ϵ , it follows that $(F, (1 - \frac{2}{\omega})\frac{1}{n}M|_F + \frac{2}{\omega}E|_F)$ is not klt for a very general fiber F of g .

On the other hand, as $(Y, \frac{1}{n}M)$ is canonical, $(F, \frac{1}{n}M|_F)$ is canonical by Bertini's theorem (see [14, Lemma 5.17]). Since M is a general member

of a movable linear system by assumption, $M|_F$ is a general member of a movable linear system on F . So each irreducible component of $M|_F$ is nef. Also we can take M such that $M|_F$ and $E|_F$ have no common irreducible component. By construction, $\frac{1}{n}M|_F \sim_{\mathbb{Q}} E|_F \sim_{\mathbb{Q}} -K_F$. So we can apply [12, Theorem 3.3] to $F, \frac{1}{n}M|_F, E|_F$, which implies that $\frac{2}{\omega} \geq \frac{1}{6}$. Hence $\omega \leq 12$. \square

Lemma 4.7. *Keep the setting in Proposition 4.3, suppose that S is a del Pezzo surface. Suppose that D is a non-zero effective integral divisor on Y which is movable. If $-K_Y \sim_{\mathbb{Q}} \omega D + E$ for some positive rational number ω and some effective \mathbb{Q} -divisor E , then $\omega \leq 12$.*

Proof. As S is a del Pezzo surface with at worst Du Val singularities and $\rho(S) = 1$, there are 3 cases (see [15], [17, Remark 3.4(ii)]):

- (1) $K_S^2 = 9$ and $S \simeq \mathbb{P}^2$;
- (2) $K_S^2 = 8$ and $S \simeq \mathbb{P}(1, 1, 2)$;
- (3) $1 \leq K_S^2 \leq 6$.

Consider the linear system \mathcal{H} on S defined by

$$\mathcal{H} := \begin{cases} |\mathcal{O}_{\mathbb{P}^2}(1)| & \text{if } S \simeq \mathbb{P}^2; \\ |\mathcal{O}_{\mathbb{P}(1,1,2)}(2)| & \text{if } S \simeq \mathbb{P}(1, 1, 2); \\ |-K_S| & \text{if } 2 \leq K_S^2 \leq 6; \\ |-2K_S| & \text{if } K_S^2 = 1. \end{cases}$$

Then \mathcal{H} is base point free and defines a generically finite map (cf. [7, Theorem 8.3.2]). By Bertini's theorem, we can take a general element $H \in \mathcal{H}$ such that H and $G = g^{-1}(H) = g^*H$ are smooth. Note that for a general fiber C of $g|_G$, $C \simeq \mathbb{P}^1$, $(-K_G \cdot C) = 2$, and $G|_G \sim (H^2) \cdot C$.

Note that $g|_G$ is factored through by a ruled surface over H , so $K_G^2 \leq 8 - 8g(H)$. Then

$$\begin{aligned} (-K_Y|_G)^2 &= (-K_G + G|_G)^2 \\ &= K_G^2 + 4H^2 \\ &\leq 8 - 8g(H) + 4H^2 \\ &= -4(K_S \cdot H) \leq 24. \end{aligned}$$

By construction, as $|G|$ defines a morphism from Y to a surface and D is movable, $D|_G$ is an effective non-zero integral divisor for a general G . So we may write

$$-K_Y|_G \sim_{\mathbb{Q}} \omega D|_G + E|_G. \quad (4.1)$$

Take a general fiber C of $g|_G$. If $(D|_G \cdot C) \neq 0$, then by (4.1) intersecting with C , $\omega \leq 2$. If $(D|_G \cdot C) = 0$, then $D|_G$ is vertical over H and thus $D|_G$ is numerically equivalent to a multiple of C . By Proposition 4.3(3), $-K_Y|_G$ is nef. Then by (4.1) intersecting with $-K_Y|_G$,

$$24 \geq (-K_Y|_G)^2 \geq \omega(-K_Y|_G \cdot D|_G) \geq \omega(-K_Y|_G \cdot C) = 2\omega,$$

which implies that $\omega \leq 12$. \square

4.3. A new geometric inequality. Now we are prepared to prove Theorem 4.2.

Proof of Theorem 4.2. It suffices to show that, if $|-mK_X|$ is composed with a pencil, then $P_{-m} \leq 12m + 1$.

Take $g : Y \rightarrow S$ to be the morphism in Proposition 4.3. Take a common resolution $\pi : W \rightarrow X$, $q : W \rightarrow Y$. We may modify π such that $f : W \rightarrow \mathbb{P}^1$ is the fibration induced by $|-mK_X|$ as in Subsection 2.1. See the following diagram:

$$\begin{array}{ccccc} & & W & \xrightarrow{f} & \mathbb{P}^1 \\ & \swarrow \pi & & \searrow q & \\ X & \text{---} & & \text{---} & Y \xrightarrow{g} S. \end{array}$$

Denote by F_W a general fiber of f . Then

$$\pi^*(-mK_X) \sim (P_{-m} - 1)F_W + E, \quad (4.2)$$

where E is an effective \mathbb{Q} -divisor on W . Set $\omega = \frac{P_{-m}-1}{m}$. Pushing forward (4.2) to Y , we have

$$-K_Y \sim_{\mathbb{Q}} \omega q_* F_W + E_Y, \quad (4.3)$$

where E_Y is an effective \mathbb{Q} -divisor on Y . Note that $q_* F_W$ is a general member of a movable linear system.

Case 1. S is a point.

In this case, $\omega \leq 7$ by (4.3) and Lemma 4.5.

Case 2. $S = \mathbb{P}^1$.

If $S = \mathbb{P}^1$ and $q_* F_W|_F = 0$, then $q_* F_W \sim F$ and $-K_Y \sim_{\mathbb{Q}} \omega F + E_Y$. By Lemma 4.6, $\omega \leq 12$.

If $S = \mathbb{P}^1$ and $q_* F_W|_F \neq 0$, then $q_* F_W|_F$ is a movable effective non-zero integral divisor on F . Restricting (4.3) on F , we have $-K_F \sim_{\mathbb{Q}} \omega(q_* F_W|_F) + E_Y|_F$. By Lemma 4.4, $\omega \leq 4$.

Case 3. S is a del Pezzo surface.

In this case, $\omega \leq 12$ by (4.3) and Lemma 4.7.

Combining all above cases, we proved that $\frac{P_{-m}-1}{m} = \omega \leq 12$ as long as $|-mK_X|$ is composed with a pencil. \square

Applying Proposition 4.1 and Theorem 4.2, we have the following criteria for $|-mK|$ not composed with a pencil (cf. [6, Proposition 5.4]).

Proposition 4.8. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. Let t be a positive real number and m a positive integer. If $m \geq t, m \geq \frac{r_{\max} t}{3}$, and one of the following conditions holds:*

- (1) $m > -\frac{3}{4} + \sqrt{\frac{12}{t(-K_X^3)} + 6r_X + \frac{1}{16}};$
- (2) $m > -\frac{3}{4} + \sqrt{\frac{12}{t(-K_X^3)} + \frac{72}{-K_X^3} + \frac{1}{16}},$

then $|-mK_X|$ is not composed with a pencil.

Proof. By [6, Proposition 5.3],

$$P_{-m} \geq \frac{1}{12}m(m+1)(2m+1)(-K_X^3) + 1 - \frac{2m}{t}.$$

The assumption implies that either $P_{-m} > r_X(-K_X^3)m + 1$ or $P_{-m} > 12m + 1$. Hence $|-mK_X|$ is not composed with a pencil by Proposition 4.1 and Theorem 4.2. \square

By the same method we have the following corollary.

Corollary 4.9. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. Let t be a positive real number and m a positive integer. If $m \geq t, m \geq \frac{r_{\max}t}{3}$, and $m > -\frac{3}{4} + \sqrt{\frac{12}{t(-K_X^3)} + \frac{6}{t(-K_X^3)} + \frac{1}{16}}$ for some positive real number l , then $P_{-m} - 1 > \frac{m}{t}$.*

We illustrate in the following example on how efficient Proposition 4.8 is comparing to [5, Corollary 4.2].

Example 4.10. Suppose that X is a terminal weak \mathbb{Q} -Fano 3-fold with $P_{-1} = 0$, $B_X = \{2 \times (1, 2), (2, 5), (3, 7), (4, 9)\}$, and $-K_X^3 = \frac{43}{315}$. Then [5, Corollary 4.2] implies that $|-mK_X|$ is not composed with a pencil for all $m \geq 61$ (see the last paragraph of [5, Page 106]). On the other hand, by Proposition 4.8, $|-mK_X|$ is not composed with a pencil for all $m \geq 23$ (see Case 4 of Proof of Theorem 5.6), which significantly improves the previous result. Also it can be computed directly by (2.1) to get $P_{-22} = 260 < 12 \times 22 + 1$, which tells that the estimates in Proposition 4.8 is efficient enough comparing to directly using the Riemann–Roch formula.

4.4. A remark on [5, Corollary 4.2]. In this subsection, we discuss the equality case of [5, Corollary 4.2] for terminal \mathbb{Q} -Fano 3-folds.

Proposition 4.11. *Let X be a terminal \mathbb{Q} -Fano 3-fold and let m be a positive integer. If*

$$P_{-m} = r_X(-K_X^3)m + 1$$

and $|-mK_X|$ is composed with a pencil, then

- (1) $r_X(-K_X^3) = 1$;
- (2) *if moreover the Weil divisor class group of X has no m -torsion element, then $h^0(X, -kK_X) = k + 1$ for all $1 \leq k \leq m$.*

Proof. We recall the proof of [5, Corollary 4.2]. As $|-mK_X|$ is composed with a pencil, take $D = -mK_X$ and keep the notation in Subsection 2.1, we have

$$\pi^*(-mK_X) \sim (P_{-m} - 1)S + F \tag{4.4}$$

where S is a generic irreducible element of $\text{Mov}[\pi^*(-mK_X)]$ and F is an effective \mathbb{Q} -divisor. Then

$$m(-K_X^3) \geq (P_{-m} - 1)(\pi^*(-K_X)^2 \cdot S) \geq \frac{1}{r_X}(P_{-m} - 1)$$

by [5, Lemma 4.1].

Now by assumption, the equality holds. So $(\pi^*(-K_X)^2 \cdot F) = 0$. This implies that F is π -exceptional as $-K_X$ is ample. So (4.4) implies that

$$-mK_X \sim (P_{-m} - 1)\pi_*S.$$

Then

$$-K_X \sim_{\mathbb{Q}} r_X(-K_X^3)\pi_*S. \quad (4.5)$$

By [9, Lemma 2.3], $(\pi_*S)^3 \geq \frac{1}{r_X}$. Then (4.5) implies that

$$-K_X^3 \geq \frac{(r_X(-K_X^3))^3}{r_X},$$

which implies that $r_X(-K_X^3) = 1$ as it is a positive integer. Under the assumption that the Weil divisor class group of X has no m -torsion element, (4.5) implies that $-K_X \sim \pi_*S$. So the conclusion follows as $-kK_X \sim k\pi_*S$ is composed with a pencil for any $1 \leq k \leq m$ (see [5, Page 63, Case (f_p)]). \square

The following example shows that Proposition 4.11 is non-empty.

Example 4.12 ([8, List 16.6, No. 88]). A general weighted hypersurface $X_{42} \subset \mathbb{P}(1, 1, 6, 14, 21)$ is a terminal \mathbb{Q} -Fano 3-fold with $r_X(-K_X^3) = 1$ and $B_X = \{(1, 2), (1, 3), (1, 7)\}$. By [18, Proposition 2.9], the Weil divisor class group of X is torsion free. Certainly, $P_{-k} = k + 1$ and $|-kK_X|$ is composed with a pencil for $1 \leq k \leq 5$.

5. PROOF OF MAIN RESULTS

In this section, we apply the birationality criteria (Theorem 3.4 and Corollary 3.7) and the non-pencil criteria (Proposition 4.8) to prove the main theorem. The proof will be divided into several cases:

1. $r_X = 840$;
2. $P_{-2} = 0$;
3. $P_{-2} > 0$, $P_{-1} = 0$, and $r_{\max} \geq 14$;
4. $P_{-2} > 0$, $P_{-1} = 0$, and $r_{\max} \leq 13$;
5. $P_{-1} > 0$ and $r_{\max} \geq 14$;
6. $P_{-1} > 0$ and $r_{\max} \leq 13$.

Here recall that $r_X = \text{lcm}\{r_i \mid r_i \in \mathcal{R}_X\}$ is the Cartier index of K_X , and $r_{\max} = \max\{r_i \mid r_i \in \mathcal{R}_X\}$ is the maximal local index.

5.1. The case $r_X = 840$.

Theorem 5.1. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold with $r_X = 840$. Then φ_{-m} is birational for all $m \geq 48$.*

Proof. Keep the setting in Notation 3.1. By [6, Lemma 6.5] and the first line of its proof, we know that $r_{\max} = 8$, $P_{-1} \geq 1$, and $-K_X^3 \geq \frac{47}{840}$. Take $m_0 = 8$ and $\nu_0 = 1$. By Corollary 4.9 (with $l = 1$, $t = 4.5$, $-K_X^3 \geq \frac{47}{840}$), we have $P_{-12} - 1 > 12$.

If $|-12K_X|$ and $|-8K_X|$ are composed with the same pencil, then take $\mu'_0 = \frac{12}{P_{-12}-1} < 1$ by Remark 3.2. By Proposition 4.8(2) (with $t = 13.5$, $-K_X^3 \geq \frac{47}{840}$), we can take $m_1 = 36$. By Lemma 3.3, $N_0 \geq \lceil \frac{840}{8m_1} \rceil = 3$. Then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 48$.

If $|-12K_X|$ and $|-8K_X|$ are not composed with the same pencil, then take $m_1 = 12$ and $\mu'_0 = m_0 = 8$. Then by Theorem 3.4(3), φ_{-m} is birational for all $m \geq 36$. \square

5.2. The case $P_{-2} = 0$.

Theorem 5.2. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. If $P_{-2} = 0$, then φ_{-m} is birational for all $m \geq 51$.*

Proof. Keep the setting in Notation 3.1. In this case, the possible baskets are classified in [3, Theorem 3.5] with 23 cases in total (see Table 2 in Appendix). Here we refer to the numbering in Table 2.

For No. 1-5 of Table 2, $r_X \leq 84$, $-K_X^3 \geq \frac{1}{84}$, $P_{-8} \geq 2$, and $r_{\max} \leq 11$. So we can take $m_0 = 8$. By Corollary 4.9 (with $l = 2, t = 5.7$), $P_{-21} - 1 > \frac{21}{2}$. If $|-21K_X|$ and $|-8K_X|$ are composed with the same pencil, then take $\mu'_0 = \frac{21}{P_{-21}-1} < 2$ by Remark 3.2. By Proposition 4.8(1) (with $t = 6.6$), we can take $m_1 = 25$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 48$. If $|-21K_X|$ and $|-8K_X|$ are not composed with the same pencil, then take $m_1 = 21$ and $\mu'_0 = m_0 = 8$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 51$.

For No. 6-13 and No. 16-23 of Table 2, $r_X \leq 78$, $-K_X^3 \geq \frac{1}{30}$, $P_{-6} \geq 2$, and $r_{\max} \leq 14$. So we can take $m_0 = 6$. By Corollary 4.9 (with $l = 1, t = 3.6$), $P_{-17} - 1 > 17$. If $|-17K_X|$ and $|-6K_X|$ are composed with the same pencil, then take $\mu'_0 = \frac{17}{P_{-17}-1} < 1$ by Remark 3.2. By Proposition 4.8(1) (with $t = 4.9$), we can take $m_1 = 23$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 51$. If $|-17K_X|$ and $|-6K_X|$ are not composed with the same pencil, then take $m_1 = 17$ and $\mu'_0 = m_0 = 6$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 51$.

For No. 14 of Table 2, $r_X = 210$, $-K_X^3 = \frac{17}{210}$, $P_{-5} = 2$, and $r_{\max} = 7$. So we can take $m_0 = 5$. By Corollary 4.9 (with $l = 1, t = 4.2$), $P_{-10} - 1 > 10$. If $|-10K_X|$ and $|-5K_X|$ are composed with the same pencil, then take $\mu'_0 = \frac{10}{P_{-10}-1} < 1$ by Remark 3.2. By Proposition 4.8(2) (with $t = 12$), we can take $m_1 = 30$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 51$. If $|-10K_X|$ and $|-5K_X|$ are not composed with the same pencil, then take $m_1 = 10$ and $\mu'_0 = m_0 = 5$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 29$.

For No. 15 of Table 2, $r_X = 120$, $-K_X^3 = \frac{3}{40}$, $P_{-5} = 2$, and $r_{\max} = 8$. So we can take $m_0 = 5$. By Corollary 4.9 (with $l = 1, t = 4$), $P_{-11} - 1 > 11$. If $|-11K_X|$ and $|-5K_X|$ are composed with the same pencil, then take $\mu'_0 = \frac{11}{P_{-11}-1} < 1$ by Remark 3.2. By Proposition 4.8(1) (with $t = 10$), we can take $m_1 = 27$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 46$. If $|-11K_X|$ and $|-5K_X|$ are not composed with the same pencil, then take $m_1 = 11$ and $\mu'_0 = m_0 = 5$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 32$. \square

5.3. The case $P_{-2} > 0$ and $P_{-1} = 0$.

Lemma 5.3. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. If $P_{-1} = 0$ and $P_{-2} > 0$, then*

$$\gamma(B_X) \geq 0, \quad 2 \in \mathcal{R}_X, \quad \sigma(B_X) \geq 11. \quad (5.1)$$

Proof. By (2.3), we have $\sigma(B_X) = 10 - 5P_{-1} + P_{-2} = 10 + P_{-2} \geq 11$. Other statements follow from (2.4) and Proposition 2.2(4). \square

Theorem 5.4. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. If $P_{-2} > 0$, $P_{-1} = 0$, and $r_{\max} \geq 14$, then φ_{-m} is birational for all $m \geq 59$. Moreover, φ_{-58} may not be birational only if $B_X = \{(1, 2), 2 \times (1, 3), (8, 17)\}$ and $|-24K_X|$ is composed with a pencil.*

Proof. Keep the setting in Notation 3.1. By [5, Case II of Proof of Theorem 3.12] (especially the last paragraph of Subsubcase II-3-iii) or the second paragraph of [5, Case IV of Proof of Theorem 1.8] (see Table 3 in Appendix), we can see that $r_{\max} \leq 13$ provided $P_{-4} = 1$. Hence by assumption, $P_{-4} \geq 2$ and we can always take $m_0 = 4$.

Case 1. $r_{\max} \geq 16$.

It is not hard to search by hands or with the help of a computer program to get all possible B_X satisfying (5.1) and $r_{\max} \geq 16$. Here note that $\sigma(B_X) \geq 11$ implies that $\sum_i r_i > 2\sigma(B_X) \geq 22$.

If $22 \leq r_{\max} \leq 24$, then there is no B_X satisfying (5.1). If $16 \leq r_{\max} \leq 21$, then all possible B_X satisfying (5.1) are listed in Table 1.

Here we explain briefly how to get Table 1. The algorithm is the following: first we can list all possible \mathcal{R}_X satisfying $2 \in \mathcal{R}_X$ and $\gamma(B_X) \geq 0$; then we find all possible b_i for those \mathcal{R}_X such that $\sigma(B_X) \geq 11$. For example, let us consider the case $r_{\max} = 17$. As $2 \in \mathcal{R}_X$, $\{2, 17\} \subset \mathcal{R}_X$. So we can list all possible \mathcal{R}_X with $\gamma(B_X) \geq 0$ by enumeration method by considering the second largest r_i :

$$\begin{aligned} &\{2, 5, 17\}; && \{2, 2, 4, 17\}; && \{2, 4, 17\}; \\ &\{2, 3, 3, 17\}; && \{2, 2, 3, 17\}; && \{2, 3, 17\}; \\ &\{2, 2, 2, 2, 17\}; && \{2, 2, 2, 17\}; && \{2, 2, 17\}; \\ &\{2, 17\}. \end{aligned}$$

Then all possible B_X with $\sigma(B_X) \geq 11$ are listed in Table 1; for instance, there is no such basket B_X with $\mathcal{R}_X = \{2, 4, 17\}$ because in this case $\sigma(B_X) \leq 1 + 1 + 8 = 10$.

TABLE 1. Baskets satisfying Lemma 5.3 with $r_{\max} \geq 16$

| No. | B_X | $-K^3$ |
|-----|--|---------|
| 1 | $\{(1, 2), (10, 21)\}$ | < 0 |
| 2 | $\{2 \times (1, 2), (10, 21)\}$ | $5/21$ |
| 3 | $\{2 \times (1, 2), (9, 20)\}$ | < 0 |
| 4 | $\{2 \times (1, 2), (9, 19)\}$ | < 0 |
| 5 | $\{(1, 2), (1, 3), (9, 19)\}$ | < 0 |
| 6 | $\{3 \times (1, 2), (9, 19)\}$ | $9/38$ |
| 7 | $\{3 \times (1, 2), (8, 19)\}$ | $5/38$ |
| 8 | $\{4 \times (1, 2), (7, 18)\}$ | $5/18$ |
| 9 | $\{(1, 2), (2, 5), (8, 17)\}$ | < 0 |
| 10 | $\{3 \times (1, 2), (8, 17)\}$ | < 0 |
| 11 | $\{2 \times (1, 2), (1, 3), (8, 17)\}$ | < 0 |
| 12 | $\{2 \times (1, 2), (1, 4), (8, 17)\}$ | < 0 |
| 13 | $\{(1, 2), 2 \times (1, 3), (8, 17)\}$ | $7/102$ |
| 14 | $\{4 \times (1, 2), (8, 17)\}$ | $4/17$ |
| 15 | $\{4 \times (1, 2), (7, 17)\}$ | $2/17$ |
| 16 | $\{2 \times (1, 2), (2, 5), (7, 16)\}$ | $11/80$ |
| 17 | $\{4 \times (1, 2), (7, 16)\}$ | < 0 |

Continued on next page

Table 1 – continued from previous page

| No. | B_X | $-K^3$ |
|-----|--|--------|
| 18 | $\{3 \times (1, 2), (1, 3), (7, 16)\}$ | 5/48 |
| 19 | $\{5 \times (1, 2), (7, 16)\}$ | 7/16 |

For No. 2 of Table 1, $-K_X^3 = \frac{5}{21}$ and in this case $P_{-2} = 2$, $r_X = 42$, $r_{\max} = 21$. We can take $\mu'_0 = m_0 = 2$. By Proposition 4.8(1) (with $t = 2.28$), we can take $m_1 = 16$. Then by Theorem 3.4(1), φ_{-m} is birational for all $m \geq 54$.

For other cases with $-K_X^3 > 0$, we have $-K_X^3 \geq \frac{7}{102}$ and $r_{\max} \leq 19$. By Corollary 4.9 (with $l = 1$, $t = 2$), $P_{-13} - 1 > 13$. If $|-13K_X|$ and $|-4K_X|$ are not composed with the same pencil, then take $m_1 = 13$ and $\mu'_0 = m_0 = 4$. Then by Theorem 3.4(1), φ_{-m} is birational for all $m \geq 51$.

So we may assume that $|-13K_X|$ and $|-4K_X|$ are composed with the same pencil. Then in the following we can take $\mu'_0 = \frac{13}{P_{-13}-1} < 1$ by Remark 3.2 and $m_0 = 4$.

For No. 6-8 of Table 1, $-K_X^3 \geq \frac{5}{38}$, $r_X \leq 38$, $r_{\max} \leq 19$. By Proposition 4.8(1) (with $t = 2.4$), we can take $m_1 = 16$. Then by Theorem 3.4(1), φ_{-m} is birational for all $m \geq 50$.

For No. 14-16 and No. 18-19 of Table 1, $-K_X^3 \geq \frac{5}{48}$, $r_X \leq 80$, $r_{\max} \leq 17$. By Proposition 4.8(1) (with $t = 3.88$), we can take $m_1 = 22$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 56$.

For No. 13 of Table 1, $-K_X^3 = \frac{7}{102}$, $r_X = 102$, $r_{\max} = 17$. By Proposition 4.8(1) (with $t = 3.6$), we can take $m_1 = 25$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 59$. Moreover, if $|-24K_X|$ is not composed with a pencil, then take $m_1 = 24$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 58$.

Case 2. $14 \leq r_{\max} \leq 15$.

For the remaining cases $14 \leq r_{\max} \leq 15$, we have $-K_X^3 \geq \frac{1}{30}$ by Proposition 2.2(3) as $P_{-4} \geq 2$. By Corollary 4.9 (with $l = 1$, $t = 3.4$), $P_{-17} - 1 > 17$.

If $|-17K_X|$ and $|-4K_X|$ are not composed with the same pencil, then take $m_1 = 17$ and $\mu'_0 = m_0 = 4$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 51$.

If $|-17K_X|$ and $|-4K_X|$ are composed with the same pencil, then take $\mu'_0 = \frac{17}{P_{-17}-1} < 1$ by Remark 3.2 and $m_0 = 4$.

If $r_{\max} = 15$, then we claim that $r_X \leq 60$. In fact, as $\{15, 2\} \subset \mathcal{R}_X$, by (2.4), $s \notin \mathcal{R}_X$ for all $s > 7$. If $7 \notin \mathcal{R}_X$, then r_X divides 60. If $7 \in \mathcal{R}_X$, then $\mathcal{R}_X = \{15, 7, 2\}$ by (2.4), and moreover $B_X = \{(1, 2), (3, 7), (7, 15)\}$ by $\sigma(B_X) \geq 11$. But this basket has $-K_X^3 < 0$ by (2.2), which is absurd. Hence $r_X \leq 60$. By Proposition 4.8(1) (with $t = 4$, $-K_X^3 \geq \frac{1}{30}$), we can take $m_1 = 21$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 51$.

If $r_{\max} = 14$, then we claim that $B_X = \{6 \times (1, 2), (5, 14)\}$. Suppose that $B_X = \{(b_1, r_1), \dots, (b_k, r_k), (b, 14)\}$ with $b \in \{1, 3, 5\}$. Then $\sigma(B_X) \geq 11$ implies that $\sum_{i=1}^k b_i \geq 6$. If there exists some $r_i > 2$, then $\sum_{i=1}^k r_i > 2 \sum_{i=1}^k b_i \geq 12$, that is, $\sum_{i=1}^k r_i \geq 13$. So (2.4) implies that $\sum_{i=1}^k \frac{1}{r_i} \geq 3 - \frac{1}{14}$. On the other hand, (2.4) implies that $\frac{3}{2}k + 14 - \frac{1}{14} \leq 24$, which says that $k \leq 6$. So $\sum_{i=1}^k \frac{1}{r_i} \leq 5 \times \frac{1}{2} + \frac{1}{3} < 3 - \frac{1}{14}$, a contradiction. So all $r_i = 2$ and

$k \geq 6$. Then $\gamma(B_X) \geq 0$ implies that $k = 6$, and $\sigma(B_X) \geq 11$ implies that $b = 5$. We conclude that $B_X = \{6 \times (1, 2), (5, 14)\}$. In this case, $-K_X^3 = \frac{3}{14}$, $r_X = r_{\max} = 14$. By Proposition 4.8(1) (with $t = 2$), we can take $m_1 = 10$. Then by Theorem 3.4(1), φ_{-m} is birational for all $m \geq 32$.

Combining all above cases, we have proved the theorem. \square

Lemma 5.5 (cf. [3, Case I of Proof of Theorem 4.4]). *Let X be a terminal weak \mathbb{Q} -Fano 3-fold with $P_{-1} = 0$ and $P_{-2} > 0$. If $P_{-4} \geq 2$ and $-K_X^3 < \frac{1}{12}$, then B_X is dominated by one of the following initial baskets:*

$$\begin{aligned} &\{8 \times (1, 2), 3 \times (1, 3)\}; \\ &\{9 \times (1, 2), (1, 4), (1, 5)\}; \\ &\{9 \times (1, 2), (1, 4), (1, 6)\}; \\ &\{9 \times (1, 2), 2 \times (1, 5)\}. \end{aligned}$$

Note that in the latter 3 cases, all possible packings have $r_{\max} \leq 9$.

Proof. Following [3, Case I of Proof of Theorem 4.4], we only need to consider the cases $(P_{-3}, P_{-4}) = (1, 2)$ or $(0, 2)$ in [3, Subcase I-3 of Proof of Theorem 4.4].

If $(P_{-3}, P_{-4}) = (1, 2)$, then [3, Subcase I-3 of Proof of Theorem 4.4] shows that B_X is dominated by $\{8 \times (1, 2), 3 \times (1, 3)\}$. (Actually it shows moreover that B_X is dominated by $\{7 \times (1, 2), (2, 5), 2 \times (1, 3)\}$.)

If $(P_{-3}, P_{-4}) = (0, 2)$, then $P_{-1} = 0$ and $P_{-2} = 1$. Then by (2.5)–(2.7), $n_{1,2}^0 = 9$, $n_{1,3}^0 = 0$, $n_{1,4}^0 + \sigma_5 = 2$. So B_X is dominated by $\{9 \times (1, 2), (1, s_1), (1, s_2)\}$ for some $s_2 \geq s_1 \geq 4$. The case $(s_1, s_2) = (4, 4)$ is ruled out by [3, Subcase I-3 of Proof of Theorem 4.4]. Hence we get the conclusion by (2.4) and [3, Lemma 3.1]. \square

Theorem 5.6. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. If $P_{-1} = 0$, $P_{-2} > 0$, and $r_{\max} \leq 13$, then φ_{-m} is birational for all $m \geq 56$.*

Proof. Keep the setting in Notation 3.1. By Proposition 2.2, $P_{-6} \geq 2$ and $-K_X^3 \geq \frac{1}{70}$. We always take $\nu_0 = 2$.

If $P_{-4} = 1$, then $P_{-2} = 1$. Following the second paragraph of [5, Case IV of Proof of Theorem 1.8] (see Table 3 in Appendix), we have $r_X \leq 130$. By Corollary 4.9 (with $l = 1, t = 5.5$, $-K_X^3 \geq \frac{1}{70}$), $P_{-24} - 1 > 24$. If $|-24K_X|$ and $|-6K_X|$ are not composed with the same pencil, then take $m_1 = 24$ and $\mu'_0 = m_0 = 6$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 56$. If $|-24K_X|$ and $|-6K_X|$ are composed with the same pencil, then take $m_0 = 6$ and $\mu'_0 = \frac{24}{P_{-24}-1} < 1$ by Remark 3.2. By Proposition 4.8(1) (with $t = 6.9$), we can take $m_1 = 30$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 56$.

From now on, we assume that $P_{-4} \geq 2$. Note that $-K_X^3 \geq \frac{1}{30}$ by Proposition 2.2(3). By Corollary 4.9 (with $l = 1, t = 3.6$), $P_{-16} - 1 > 16$. If $|-16K_X|$ and $|-4K_X|$ are not composed with the same pencil, then take $m_1 = 16$ and $\mu'_0 = m_0 = 4$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 46$.

In the following discussions, we assume that $|-16K_X|$ and $|-4K_X|$ are composed with the same pencil. We can always take $m_0 = 4$ and $\mu'_0 = \frac{16}{P_{-16}-1} < 1$ by Remark 3.2.

Case 1. $r_{\max} \leq 6$ or $r_{\max} \in \{10, 12\}$.

If $r_{\max} \leq 6$, then $r_X \leq 60$. If $r_{\max} = 10$ (resp. $r_{\max} = 12$), then $r_X \leq 210$ (resp. $r_X \leq 84$) by [5, Page 107]. Then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 52$.

Case 2. $r_{\max} = 7$.

If $r_{\max} = 7$, then r_X divides $\text{lcm}(2, 3, 4, 5, 6, 7) = 420$. Hence either $r_X = 420$ or $r_X \leq 210$. If $r_X = 420$, then $\{4, 5, 7\} \subset \mathcal{R}_X$ and one element of $\{3, 6\}$ is in \mathcal{R}_X . Suppose that

$$B_X = \{(b_1, r_1), \dots, (b_k, r_k), (1, r), (1, 4), (a_5, 5), (a_7, 7)\}$$

where $r \in \{3, 6\}$, $a_5 \leq 2$, $a_7 \leq 3$. Then $\sigma(B_X) \geq 11$ implies that $\sum_{i=1}^k b_i \geq 4$. Lemma 2.3 implies that

$$\gamma(B_X) \leq 24 - (7 - \frac{1}{7} + 5 - \frac{1}{5} + 4 - \frac{1}{4} + 3 - \frac{1}{3} + 4 \times \frac{3}{2}) < 0, \quad (5.2)$$

a contradiction. Hence $r_X \leq 210$. Then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 43$.

Case 3. $r_{\max} = 8$.

If $r_{\max} = 8$, then we claim that $r_X \leq 168$. In fact, if $r_X > 168$, then $\{5, 7\} \subset \mathcal{R}_X$. Suppose that

$$B_X = \{(b_1, r_1), \dots, (b_k, r_k), (a_5, 5), (a_7, 7), (a_8, 8)\}$$

where $a_5 \leq 2$, $a_7 \leq 3$, $a_8 \leq 3$. Then $\sigma(B_X) \geq 11$ implies that $\sum_{i=1}^k b_i \geq 3$. Similar to (5.2), Lemma 2.3 implies that $\gamma(B_X) < 0$, a contradiction. Hence $r_X \leq 168$.

Then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 43$.

Case 4. $r_{\max} = 9$.

If $r_{\max} = 9$, then we claim that $r_X \leq 252$ or $B_X = \{2 \times (1, 2), (2, 5), (3, 7), (4, 9)\}$.

If 7 and 8 are not in \mathcal{R}_X , then $r_X \leq 180$.

If $8 \in \mathcal{R}_X$, then as $\{2, 8, 9\} \subset \mathcal{R}_X$, we know that 6 and 7 are not in \mathcal{R}_X by $\gamma(B_X) \geq 0$. If $5 \notin \mathcal{R}_X$, then $r_X = 72$. If $5 \in \mathcal{R}_X$, then $\mathcal{R}_X = \{2, 5, 8, 9\}$ as $\gamma(B_X) \geq 0$, but in this case $\sigma(B_X) \leq 10$, a contradiction.

If $7 \in \mathcal{R}_X$, then as $\{2, 7, 9\} \subset \mathcal{R}_X$, we know that at most one element of $\{4, 5, 6\}$ is in \mathcal{R}_X by $\gamma(B_X) \geq 0$. If $5 \notin \mathcal{R}_X$, then $r_X \leq 252$. If $5 \in \mathcal{R}_X$, then by $\sigma(B_X) \geq 11$ and $\gamma(B_X) \geq 0$, it is not hard to check that the only possible basket is $B_X = \{2 \times (1, 2), (2, 5), (3, 7), (4, 9)\}$. This concludes the claim.

If $r_X \leq 252$, then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 50$.

If $B_X = \{2 \times (1, 2), (2, 5), (3, 7), (4, 9)\}$, then $-K_X^3 = \frac{43}{315}$. By Proposition 4.8(2) (with $t = 7.6$), we can take $m_1 = 23$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 41$.

Case 5. $r_{\max} = 11$.

If $r_{\max} = 11$, then we claim that $r_X \leq 264$ or $B_X = \{3 \times (1, 2), (1, 3), (2, 5), (5, 11)\}$ or $B_X = \{2 \times (1, 2), (1, 3), (3, 7), (5, 11)\}$.

As $\{2, 11\} \subset \mathcal{R}_X$, we know that at most one element of $\{6, 7, 8, 9, 10\}$ is in \mathcal{R}_X by $\gamma(B_X) \geq 0$. If $10 \in \mathcal{R}_X$, then $r_X = 110$ by [5, Page 107]. If $9 \in \mathcal{R}_X$ or $8 \in \mathcal{R}_X$, then $r_X \leq 264$ by [5, Page 107]. If $7 \in \mathcal{R}_X$, then $5 \notin \mathcal{R}_X$ by

$\gamma(B_X) \geq 0$. So either $r_X = 154$ or at least one element of $\{3, 4\}$ is in \mathcal{R}_X . For the latter case, it is not hard to check that the only basket satisfying $\sigma(B_X) \geq 11$ and $\gamma(B_X) \geq 0$ is $B_X = \{2 \times (1, 2), (1, 3), (3, 7), (5, 11)\}$. If $6 \in \mathcal{R}_X$, then we get a contradiction by Lemma 2.3 as (5.2).

If none element of $\{6, 7, 8, 9, 10\}$ is in \mathcal{R}_X , then r_X divides 660 and $r_X < 660$ by [5, Page 107]. So either $r_X \leq 220$ or $r_X = 330$. Moreover, if $r_X = 330$, then $\{2, 3, 5, 11\} \subset \mathcal{R}_X$ and $4 \notin \mathcal{R}_X$, and it is not hard to check that the only basket satisfying $\sigma(B_X) \geq 11$ and $\gamma(B_X) \geq 0$ is $B_X = \{3 \times (1, 2), (1, 3), (2, 5), (5, 11)\}$. This concludes the claim.

If $r_X \leq 264$, then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 56$.

If $B_X = \{3 \times (1, 2), (1, 3), (2, 5), (5, 11)\}$, then $-K_X^3 = \frac{31}{330}$. By Proposition 4.8(2) (with $t = 7.6$), we can take $m_1 = 28$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 50$.

If $B_X = \{2 \times (1, 2), (1, 3), (3, 7), (5, 11)\}$, then $-K_X^3 = \frac{50}{462}$. By Proposition 4.8(2) (with $t = 7$), we can take $m_1 = 26$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 48$.

Case 6. $r_{\max} = 13$.

If $r_{\max} = 13$, then $r_X \leq 390$ or $r_X = 546$ by [5, Page 107].

If $r_X = 546$, then again by [5, Page 107], $B_X = \{(1, 2), (1, 3), (3, 7), (6, 13)\}$ and $-K_X^3 = \frac{61}{546}$. By Proposition 4.8(2) (with $t = 6$), we can take $m_1 = 26$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 52$.

If $r_X \leq 390$ and $-K_X^3 \geq \frac{1}{12}$, then by Proposition 4.8(2) (with $t = 6.9$), we can take $m_1 = 30$. Then by Theorem 3.4(2), φ_{-m} is birational for all $m \geq 56$.

If $r_X \leq 390$ and $-K_X^3 < \frac{1}{12}$, then by Lemma 5.5, B_X is dominated by $\{8 \times (1, 2), 3 \times (1, 3)\}$. As $r_{\max} = 13$, this implies that B_X is dominated by either $\{3 \times (1, 2), (6, 13), 2 \times (1, 3)\}$ or $\{6 \times (1, 2), (5, 13)\}$. So we get the following possibilities of B_X by $\gamma(B_X) \geq 0$:

| | |
|---|------------------|
| $\{3 \times (1, 2), (6, 13), 2 \times (1, 3)\}$ | $-K^3 = 5/78;$ |
| $\{2 \times (1, 2), (6, 13), (2, 5), (1, 3)\}$ | $-K^3 = 19/195;$ |
| $\{2 \times (1, 2), (6, 13), (3, 8)\}$ | $-K^3 = 11/104;$ |
| $\{(1, 2), (6, 13), (3, 7), (1, 3)\}$ | $-K^3 = 61/546;$ |
| $\{6 \times (1, 2), (5, 13)\}$ | $-K^3 = 1/13.$ |

In the above list, only the first and the last have $-K_X^3 < \frac{1}{12}$. In particular, in these cases, $r_X \leq 78$. Then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 55$.

Combining all above cases, we have proved the theorem. \square

5.4. The case $P_{-1} > 0$.

Lemma 5.7. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. If $P_{-1} > 0$ and $r_{\max} \geq 16$, then $P_{-4} \geq 2$.*

Proof. If $P_{-4} = 1$, then $P_{-2} = P_{-3} = 1$. Since $r_{\max} \geq 16$, by the classification in [3, Subsubcase II-4f of Proof of Theorem 4.4], B_X is dominated by $\{2 \times (1, 2), 2 \times (1, 3), (1, s_1), (1, s_2)\}$ with $s_2 \geq s_1 \geq 5$, which means that

$s_1 + s_2 = r_{\max} \geq 16$. But in this case $2 \times \frac{3}{2} + 2 \times \frac{8}{3} + 16 - \frac{1}{16} > 24$, contradicting (2.4) and [3, Lemma 3.1]. So $P_{-4} \geq 2$. \square

Theorem 5.8. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. If $P_{-1} > 0$ and $r_{\max} \geq 14$, then φ_{-m} is birational for all $m \geq 52$.*

Proof. Keep the setting in Notation 3.1. We always take $\nu_0 = 1$.

If $14 \leq r_{\max} \leq 15$, then $r_X \leq 210$ by [5, Page 104]. By Proposition 2.2(2), we can take $\mu'_0 = m_0 = 8$. Then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 52$.

If $r_{\max} = 24$, then $B_X = \{(b, 24)\}$ with $b \in \{1, 5, 7, 11\}$. If $P_{-1} = 1$, then $b = \sigma(B_X) = 5 + P_{-2} \geq 6$ by (2.3), hence $b \geq 7$ and $P_{-2} \geq 2$. By (2.2), we have $-K_X^3 \geq \frac{23}{24}$. Similarly, if $P_{-1} = 2$, then $P_{-2} \geq 2P_{-1} - 1 = 3$, and thus $b \geq 5$. Hence by (2.2), $-K_X^3 \geq \frac{47}{24}$. If $P_{-1} \geq 3$, then by (2.2), $-K_X^3 \geq b - \frac{b^2}{24} \geq \frac{23}{24}$. In summary, $-K_X^3 \geq \frac{23}{24}$ and $P_{-2} \geq 2$. We can take $\mu'_0 = m_0 = 2$. By Proposition 4.8(2) (with $t = 1$), we can take $m_1 = 9$. Then by Theorem 3.4(1), φ_{-m} is birational for all $m \geq 33$.

In the following, we consider $16 \leq r_{\max} \leq 23$. By Lemma 5.7, we always take $\mu'_0 = m_0 = 4$ and $\nu_0 = 1$.

If $r_{\max} = 23$, then $B_X = \{(b, 23)\}$ with $1 \leq b \leq 11$. If $P_{-1} = 1$, then $b = 5 + P_{-2} \geq 6$ and thus by (2.2) $-K_X^3 \geq \frac{10}{23}$. If $P_{-1} = 2$, then $b = P_{-2} \geq 2P_{-1} - 1 = 3$, hence by (2.2) $-K_X^3 \geq \frac{14}{23}$. If $P_{-1} \geq 3$, then $-K_X^3 \geq b - \frac{b^2}{23} \geq \frac{22}{23}$. In summary, $-K_X^3 \geq \frac{10}{23}$. By Proposition 4.8(1) (with $t = \frac{36}{23}$), we can take $m_1 = 12$. Then by Theorem 3.4(1), φ_{-m} is birational for all $m \geq 48$.

If $20 \leq r_{\max} \leq 22$, then by (2.4) we have $r_X \leq 60$. Then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 50$.

If $18 \leq r_{\max} \leq 19$, then $r_X \leq 190$ by [5, Page 104]. Then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 52$.

If $16 \leq r_{\max} \leq 17$, then $r_X \leq 240$ by [5, Page 104]. Then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 52$. \square

Theorem 5.9. *Let X be a terminal weak \mathbb{Q} -Fano 3-fold. If $P_{-1} > 0$ and $r_{\max} \leq 13$, then φ_{-m} is birational for all $m \geq 58$.*

Proof. Keep the setting in Notation 3.1. By Theorem 5.1 and Proposition 2.2(1), we may assume that $r_X \leq 660$. By Proposition 2.2(2), we can take $m_0 = 8$ and $\nu_0 = 1$. We take $\mu'_0 = 8$ unless stated otherwise.

Case 1. $r_{\max} \leq 8$.

If $r_{\max} \leq 8$, then r_X divides $\text{lcm}(8, 7, 6, 5) = 840$. As $r_X \leq 660$, $r_X = 420$ or $r_X \leq 280$.

If $r_X = 420$, then by Proposition 4.8(1) (with $t = 19.5$, $-K_X^3 \geq \frac{1}{330}$), we can take $m_1 = 52$. By Lemma 3.3, $N_0 \geq \lceil \frac{420}{8m_1} \rceil = 2$. Then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 48$.

If $r_X \leq 280$, then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 55$.

Case 2. $r_{\max} = 9$.

If $r_{\max} = 9$, then r_X divides 2520. As $r_X \leq 660$ and 9 divides r_X , we have $r_X \leq 360$ or $r_X \in \{504, 630\}$.

If $r_X \leq 360$, then by Corollary 4.9 (with $l = 4, t = 10$, $-K_X^3 \geq \frac{1}{330}$), we have $P_{-30} - 1 > \frac{30}{4}$. If $|-30K_X|$ and $|-8K_X|$ are composed with

the same pencil, then take $\mu'_0 = \frac{30}{P_{-30}-1} < 4$ by Remark 3.2. Then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 57$. If $|-30K_X|$ and $|-8K_X|$ are not composed with the same pencil, then take $m_1 = 30$ and $\mu'_0 = m_0 = 8$. Then by Theorem 3.4(3), φ_{-m} is birational for all $m \geq 56$.

If $r_X = 630$, then $-K_X^3 \geq \frac{1}{315}$ (note that $-K_X^3 \geq \frac{1}{330}$ and $r_X(-K_X^3)$ is an integer). By Corollary 4.9 (with $l = 2.8, t = 11$), we have $P_{-33} - 1 > \frac{33}{2.8}$. If $|-33K_X|$ and $|-8K_X|$ are composed with the same pencil, then take $\mu'_0 = \frac{33}{P_{-33}-1} < 2.8$ by Remark 3.2. By Proposition 4.8(1) (with $t = 20$, $-K_X^3 \geq \frac{1}{315}$), we can take $m_1 = 63$. By Lemma 3.3, $N_0 \geq \lceil \frac{630}{9m_1} \rceil = 2$. Then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 52$. If $|-33K_X|$ and $|-8K_X|$ are not composed with the same pencil, then take $m_1 = 33$ and $\mu'_0 = m_0 = 8$. By Lemma 3.3, $N_0 \geq \lceil \frac{630}{9m_1} \rceil = 3$. Then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 48$.

If $r_X = 504$, then $\mathcal{R}_X = \{9, 8, 7\}$ by (2.4). Write $B_X = \{(a, 7), (b, 8), (c, 9)\}$, where $a \leq 3, b \in \{1, 3\}$ and $c \in \{1, 2, 4\}$. If $P_{-1} \geq 2$, then by (2.2),

$$\begin{aligned} -K_X^3 &= 2P_{-1} + \frac{a(7-a)}{7} + \frac{b(8-b)}{8} + \frac{c(9-c)}{9} - 6 \\ &\geq \frac{6}{7} + \frac{7}{8} + \frac{8}{9} - 2 > 0.6. \end{aligned} \quad (5.3)$$

If $P_{-1} = 1$, then by (2.2) and (2.3),

$$\begin{aligned} -K_X^3 &= \frac{a(7-a)}{7} + \frac{b(8-b)}{8} + \frac{c(9-c)}{9} - 4 > 0, \\ \sigma(B_X) &= a + b + c = 5 + P_{-2} \geq 6. \end{aligned} \quad (5.4)$$

So it is easy to check that $-K_X^3 \geq \frac{73}{504}$ by considering all possible values of (a, b, c) . By Proposition 4.8(2) (with $t = 7.3$, $-K_X^3 \geq \frac{73}{504}$), we can take $m_1 = 22$. By Lemma 3.3, $N_0 \geq \lceil \frac{504}{9m_1} \rceil = 3$. Then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 44$.

Case 3. $r_{\max} = 10$.

If $r_{\max} = 10$, then we claim that $r_X \leq 210$ or $r_X = 420$.

By (2.4), at most one element of $\{7, 8, 9\}$ is in \mathcal{R}_X . If $7 \notin \mathcal{R}_X$, then r_X divides either $120 = \text{lcm}(10, 8, 60)$ or $180 = \text{lcm}(10, 9, 60)$. If $7 \in \mathcal{R}_X$, then r_X divides $420 = \text{lcm}(10, 7, 60)$, so either $r_X \leq 210$ or $r_X = 420$. This concludes the claim.

If $r_X \leq 210$, then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 48$.

If $r_X = 420$, then by (2.4), $\mathcal{R}_X = \{10, 7, 4, 3\}$. Write $B_X = \{(1, 3), (1, 4), (a, 7), (b, 10)\}$, where $a \leq 3$ and $b \in \{1, 3\}$. If $P_{-1} \geq 2$, then by (2.2),

$$\begin{aligned} -K_X^3 &= 2P_{-1} + \frac{2}{3} + \frac{3}{4} + \frac{a(7-a)}{7} + \frac{b(10-b)}{10} - 6 \\ &\geq \frac{2}{3} + \frac{3}{4} + \frac{6}{7} + \frac{9}{10} - 2 > 1. \end{aligned} \quad (5.5)$$

If $P_{-1} = 1$, then by (2.2) and (2.3),

$$\begin{aligned} -K_X^3 &= \frac{2}{3} + \frac{3}{4} + \frac{a(7-a)}{7} + \frac{b(10-b)}{10} - 4 > 0, \\ \sigma(B_X) &= 2 + a + b = 5 + P_{-2} \geq 6. \end{aligned} \quad (5.6)$$

So it is easy to check that $-K_X^3 \geq \frac{13}{420}$ by considering all possible values of (a, b) . By Corollary 4.9 (with $l = 1$, $t = 4.8$), we have $P_{-16} - 1 > 16$. If $|-16K_X|$ and $|-8K_X|$ are composed with the same pencil, then take $\mu'_0 = \frac{16}{P_{-16}-1} < 1$ by Remark 3.2. Then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 58$. If $|-16K_X|$ and $|-8K_X|$ are not composed with the same pencil, then take $m_1 = 16$ and $\mu'_0 = m_0 = 8$. Then by Theorem 3.4(3), φ_{-m} is birational for all $m \geq 44$.

Case 4. $r_{\max} = 11$.

If $r_{\max} = 11$, then we claim that $r_X \leq 330$ or $r_X \in \{385, 396, 440, 462, 660\}$.

By (2.4), at most one element of $\{7, 8, 9, 10\}$ is in \mathcal{R}_X . So r_X divides one element of $\{1980, 1320, 4620\}$. As $r_X \leq 660$ and 11 divides r_X , it is clear that $r_X \leq 330$ or $r_X \in \{385, 396, 440, 462, 495, 660\}$. Moreover, if $r_X = 495$, then $\{11, 9, 5\} \subset \mathcal{R}_X$, which contradicts (2.4). This concludes the claim.

If $r_X \leq 330$, then by Corollary 4.9 (with $l = 7.6$, $t = 7.6$, $-K_X^3 \geq \frac{1}{330}$), $P_{-28} - 1 > \frac{28}{7.6}$. If $|-28K_X|$ and $|-8K_X|$ are composed with the same pencil, then take $\mu'_0 = \frac{28}{P_{-28}-1} < 7.6$ by Remark 3.2. Then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 58$. If $|-28K_X|$ and $|-8K_X|$ are not composed with the same pencil, then take $m_1 = 28$ and $\mu'_0 = m_0 = 8$. Then by Theorem 3.4(3), φ_{-m} is birational for all $m \geq 58$.

If $r_X = 385$ (resp. 396), then $-K_X^3 \geq \frac{2}{385}$ (resp. $\geq \frac{2}{396}$). By Corollary 4.9 (with $l = 1.5$, $t = 9$), $P_{-33} - 1 > \frac{33}{1.5}$. If $|-33K_X|$ and $|-8K_X|$ are composed with the same pencil, then take $\mu'_0 = \frac{33}{P_{-33}-1} < 1.5$ by Remark 3.2. Then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 56$ (resp. ≥ 57). If $|-33K_X|$ and $|-8K_X|$ are not composed with the same pencil, then take $m_1 = 33$ and $\mu'_0 = m_0 = 8$. By Lemma 3.3, $N_0 \geq \lceil \frac{385}{11m_1} \rceil = 2$. Then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 47$ (resp. ≥ 48).

We claim that if $r_X \in \{440, 462, 660\}$, then $-K_X^3 \geq \frac{74}{462}$ or $B_X = \{(1, 2), (2, 5), (1, 3), (1, 4), (1, 11)\}$ with $-K_X^3 = \frac{17}{660}$.

If $r_X = 440$, then by (2.4), $\mathcal{R}_X = \{11, 8, 5\}$. Arguing similarly as (5.3), we get $-K_X^3 > 0.5$ when $P_{-1} \geq 2$. Arguing similarly as (5.4), we get constrains for B_X when $P_{-1} = 1$. By (2.2) and considering all the possible baskets when $P_{-1} = 1$, we can see that $-K_X^3 \geq \frac{97}{440}$.

If $r_X = 462$, then by (2.4), $\mathcal{R}_X = \{11, 7, 6\}$ or $\{11, 7, 3, 2\}$ or $\{11, 7, 3, 2, 2\}$. Arguing similarly as (5.5), we get $-K_X^3 > 0.5$ or $-K_X^3 > 0.9$ or $-K_X^3 > 1$ when $P_{-1} \geq 2$. Arguing similarly as (5.6), we get constrains for B_X when $P_{-1} = 1$. By (2.2) and considering all the possible baskets when $P_{-1} = 1$, we can see that $-K_X^3 \geq \frac{85}{462}$ or $-K_X^3 \geq \frac{95}{462}$ or $-K_X^3 \geq \frac{74}{462}$ unless $B_X = \{2 \times (1, 2), (1, 3), (2, 7), (1, 11)\}$ with $-K_X^3 = \frac{1}{231}$. But the last basket has $P_{-5} = 0$, which is absurd.

If $r_X = 660$, then by (2.4), $\mathcal{R}_X = \{11, 5, 4, 3\}$ or $\{11, 5, 4, 3, 2\}$. Arguing similarly as (5.5), we get $-K_X^3 > 1$ or $-K_X^3 > 1.5$ when $P_{-1} \geq 2$. Arguing similarly as (5.6), we get constrains for B_X when $P_{-1} = 1$. By (2.2) and considering all the possible baskets when $P_{-1} = 1$, we can see that $-K_X^3 \geq \frac{167}{660}$ or $-K_X^3 \geq \frac{233}{660}$ unless $B_X = \{(1, 2), (2, 5), (1, 3), (1, 4), (1, 11)\}$ with $-K_X^3 = \frac{17}{660}$.

To summarize, we conclude the claim that $-K_X^3 \geq \frac{74}{462}$ or $B_X = \{(1, 2), (2, 5), (1, 3), (1, 4), (1, 11)\}$ with $-K_X^3 = \frac{17}{660}$.

If $-K_X^3 \geq \frac{74}{462}$, then by Proposition 4.8(2) (with $t = 5.7$), we can take $m_1 = 21$. Then by Theorem 3.4(3), φ_{-m} is birational for all $m \geq 51$.

Now we consider the case $B_X = \{(1, 2), (2, 5), (1, 3), (1, 4), (1, 11)\}$ with $-K_X^3 = \frac{17}{660}$. By Corollary 4.9 (with $l = 1, t = 4.8$), $P_{-18} - 1 > 18$. If $|-18K_X|$ and $|-8K_X|$ are composed with the same pencil, then take $\mu'_0 = \frac{18}{P_{-18}-1} < 1$ by Remark 3.2. By Proposition 4.8(2) (with $t = 14.4$), we can take $m_1 = 53$. By Lemma 3.3, $N_0 \geq \lceil \frac{660}{11m_1} \rceil = 2$. Then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 52$. If $|-18K_X|$ and $|-8K_X|$ are not composed with the same pencil, then take $m_1 = 18$ and $\mu'_0 = m_0 = 8$. By Lemma 3.3, $N_0 \geq \lceil \frac{660}{11m_1} \rceil = 4$. Then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 45$.

Case 5. $r_{\max} = 12$.

If $r_{\max} = 12$, then we claim that $r_X \leq 132$ or $r_X = 420$.

By (2.4), at most one element of $\{6, 7, 8, 9, 10, 11\}$ is in \mathcal{R}_X . So r_X divides one element of $\{120, 180, 420, 660\}$. Recall that 12 divides r_X , it is clear that $r_X \leq 132$ or $r_X \in \{180, 420, 660\}$. Moreover, if $r_X \in \{180, 660\}$, then $\{12, 9, 5\} \subset \mathcal{R}_X$ or $\{12, 11, 5\} \subset \mathcal{R}_X$, which contradicts (2.4). This concludes the claim.

If $r_X \leq 132$, then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 43$.

If $r_X = 420$, then by (2.4), $\mathcal{R}_X = \{12, 7, 5\}$. Arguing similarly as (5.3), we get $-K_X^3 \geq \frac{241}{420}$ when $P_{-1} \geq 2$. Arguing similarly as (5.4), we get constraints for B_X when $P_{-1} = 1$. By (2.2) and considering all the possible baskets when $P_{-1} = 1$, we can see that $-K_X^3 \geq \frac{241}{420}$. By Proposition 4.8(2) (with $t = 2.75$), we can take $m_1 = 11$. Hence by Lemma 3.3, $N_0 \geq \lceil \frac{420}{12m_1} \rceil = 4$. Then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 40$.

Case 6. $r_{\max} = 13$.

If $r_{\max} = 13$, then we claim that $r_X \leq 364$ or $r_X = 390$ or $r_X = 546$.

By (2.4), at most one element of $\{6, 7, 8, 9, 10, 11, 12\}$ is in \mathcal{R}_X . So r_X divides one element of $\{5460, 1560, 2340, 8580\}$. Recall that 13 divides r_X and $r_X \leq 660$, it is clear that $r_X \leq 364$ or $r_X \in \{390, 429, 455, 468, 520, 546, 572, 585\}$. Moreover, if $r_X \in \{429, 455, 468, 520, 572, 585\}$, then we can see that \mathcal{R}_X violates (2.4) by discussing the factors. For example, if $r_X = 455$, then $\{13, 7, 5\} \subset \mathcal{R}_X$ which violates (2.4). This concludes the claim.

If $P_{-4} = 1$, then $P_{-k} = 1$ for $1 \leq k \leq 4$. By [3, Subsubcase II-4f of Proof of Theorem 4.4], B_X is dominated by

$$\{2 \times (1, 2), 2 \times (1, 3), (1, s_1), (1, s_2)\}$$

for some $s_2 \geq s_1 \geq 4$. As $r_{\max} = 13$, $(s_1, s_2) = (6, 7)$. Considering all possible packings, we get the following possibilities of B_X :

$$\begin{aligned} &\{2 \times (1, 2), 2 \times (1, 3), (2, 13)\}; \\ &\{(1, 2), (2, 5), (1, 3), (2, 13)\}; \\ &\{(3, 7), (1, 3), (2, 13)\}; \\ &\{(1, 2), (3, 8), (2, 13)\}; \\ &\{2 \times (2, 5), (2, 13)\}. \end{aligned}$$

Then $r_X \leq 273$ or $B_X = \{(1, 2), (2, 5), (1, 3), (2, 13)\}$.

If $r_X \leq 273$, then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 55$.

If $B_X = \{(1, 2), (2, 5), (1, 3), (2, 13)\}$, then $-K_X^3 = \frac{23}{390}$. By Corollary 4.9 (with $l = 1, t = 3$), $P_{-13} - 1 > 13$. If $|-13K_X|$ and $|-8K_X|$ are composed with the same pencil, then take $\mu'_0 = \frac{13}{P_{-13}-1} < 1$ by Remark 3.2. Then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 56$. If $|-13K_X|$ and $|-8K_X|$ are not composed with the same pencil, then take $m_1 = 13$ and $\mu'_0 = m_0 = 8$. Then by Lemma 3.3, $N_0 \geq \lceil \frac{390}{13m_1} \rceil = 3$. Then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 44$.

So from now on, we assume that $P_{-4} \geq 2$ and take $m_0 = 4$. We take $\mu'_0 = 4$ unless stated otherwise.

If $r_X \leq 364$, then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 57$.

If $r_X = 546$, then by (2.4), $\mathcal{R}_X = \{13, 7, 3, 2\}$. Arguing similarly as (5.5), we get $-K_X^3 > 0.9$ when $P_{-1} \geq 2$. Arguing similarly as (5.6), we get constrains for B_X when $P_{-1} = 1$. By (2.2) and considering all the possible baskets when $P_{-1} = 1$, we can see that $-K_X^3 \geq \frac{157}{546}$. By Proposition 4.8(2) (with $t = 3.6$), we can take $m_1 = 16$. Hence by Lemma 3.3, $N_0 \geq \lceil \frac{546}{13m_1} \rceil = 3$. Then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 44$.

If $r_X = 390$, then by (2.4), $\mathcal{R}_X = \{13, 6, 5\}$ or $\{13, 5, 3, 2\}$ or $\{13, 5, 3, 2, 2\}$. Arguing similarly as (5.5), we get $-K_X^3 > 0.5$ or $-K_X^3 > 0.8$ or $-K_X^3 > 1$ when $P_{-1} \geq 2$. Arguing similarly as (5.6), we get constrains for B_X when $P_{-1} = 1$. By (2.2) and considering all the possible baskets when $P_{-1} = 1$, we can see that $-K_X^3 \geq \frac{133}{390}$ or $-K_X^3 \geq \frac{23}{390}$ or $-K_X^3 \geq \frac{62}{390}$. By Corollary 4.9 (with $l = 1, t = 3$, $-K_X^3 \geq \frac{23}{390}$), $P_{-13} - 1 > 13$. If $|-13K_X|$ and $|-4K_X|$ are composed with the same pencil, then take $\mu'_0 = \frac{13}{P_{-13}-1} < 1$ by Remark 3.2. Then by Corollary 3.7(1), φ_{-m} is birational for all $m \geq 56$. If $|-13K_X|$ and $|-4K_X|$ are not composed with the same pencil, then take $m_1 = 13$ and $\mu'_0 = m_0 = 4$. Then by Lemma 3.3, $N_0 \geq \lceil \frac{390}{13m_1} \rceil = 3$. Then by Corollary 3.7(2), φ_{-m} is birational for all $m \geq 40$.

Combining all above cases, we have proved the theorem. \square

5.5. Proof of main results.

Proof of Theorem 1.2. It follows from Theorems 5.2, 5.4, 5.6, 5.8, 5.9. \square

Proof of Theorem 1.4. From the proof of Theorem 1.2, φ_{-58} may not be birational only if $B_X = \{(1, 2), 2 \times (1, 3), (8, 17)\}$, $P_{-1} = 0$, and $|-24K_X|$ is composed with a pencil. In this case, $r_X(-K_X^3) = 7$ and $P_{-24} = 169 = 7 \times 24 + 1$ by (2.1). But this contradicts Proposition 4.11. \square

6. APPENDIX

The possible baskets with $P_{-2} = 0$ are the following (cf. [3, Theorem 3.5]).

TABLE 2. Baskets with $P_{-2} = 0$

| No. | B_X | $-K^3$ | P_{-3} | P_{-4} | P_{-5} | P_{-6} | P_{-7} | P_{-8} |
|-----|--|--------|----------|----------|----------|----------|----------|----------|
| 1 | $\{2 \times (1, 2), 3 \times (2, 5), (1, 3), (1, 4)\}$ | 1/60 | 0 | 0 | 1 | 1 | 1 | 2 |
| 2 | $\{5 \times (1, 2), 2 \times (1, 3), (2, 7), (1, 4)\}$ | 1/84 | 0 | 1 | 0 | 1 | 1 | 2 |
| 3 | $\{5 \times (1, 2), 2 \times (1, 3), (3, 11)\}$ | 1/66 | 0 | 1 | 0 | 1 | 1 | 2 |

Continued on next page

Table 2 – continued from previous page

| No. | B_X | $-K^3$ | P_{-3} | P_{-4} | P_{-5} | P_{-6} | P_{-7} | P_{-8} |
|-----|---|--------|----------|----------|----------|----------|----------|----------|
| 4 | $\{5 \times (1, 2), (1, 3), (3, 10), (1, 4)\}$ | 1/60 | 0 | 1 | 0 | 1 | 1 | 2 |
| 5 | $\{5 \times (1, 2), (1, 3), 2 \times (2, 7)\}$ | 1/42 | 0 | 1 | 0 | 1 | 2 | 3 |
| 6 | $\{4 \times (1, 2), (2, 5), 2 \times (1, 3), 2 \times (1, 4)\}$ | 1/30 | 0 | 1 | 1 | 2 | 2 | 4 |
| 7 | $\{3 \times (1, 2), (2, 5), 5 \times (1, 3)\}$ | 1/30 | 1 | 1 | 1 | 3 | 3 | 4 |
| 8 | $\{2 \times (1, 2), (3, 7), 5 \times (1, 3)\}$ | 1/21 | 1 | 1 | 1 | 3 | 4 | 5 |
| 9 | $\{(1, 2), (4, 9), 5 \times (1, 3)\}$ | 1/18 | 1 | 1 | 1 | 3 | 4 | 5 |
| 10 | $\{3 \times (1, 2), (3, 8), 4 \times (1, 3)\}$ | 1/24 | 1 | 1 | 1 | 3 | 3 | 5 |
| 11 | $\{3 \times (1, 2), (4, 11), 3 \times (1, 3)\}$ | 1/22 | 1 | 1 | 1 | 3 | 3 | 5 |
| 12 | $\{3 \times (1, 2), (5, 14), 2 \times (1, 3)\}$ | 1/21 | 1 | 1 | 1 | 3 | 3 | 5 |
| 13 | $\{2 \times (1, 2), 2 \times (2, 5), 4 \times (1, 3)\}$ | 1/15 | 1 | 1 | 2 | 4 | 5 | 7 |
| 14 | $\{(1, 2), (3, 7), (2, 5), 4 \times (1, 3)\}$ | 17/210 | 1 | 1 | 2 | 4 | 6 | 8 |
| 15 | $\{2 \times (1, 2), (2, 5), (3, 8), 3 \times (1, 3)\}$ | 3/40 | 1 | 1 | 2 | 4 | 5 | 8 |
| 16 | $\{2 \times (1, 2), (5, 13), 3 \times (1, 3)\}$ | 1/13 | 1 | 1 | 2 | 4 | 5 | 8 |
| 17 | $\{(1, 2), 3 \times (2, 5), 3 \times (1, 3)\}$ | 1/10 | 1 | 1 | 3 | 5 | 7 | 10 |
| 18 | $\{4 \times (1, 2), 5 \times (1, 3), (1, 4)\}$ | 1/12 | 1 | 2 | 2 | 5 | 6 | 9 |
| 19 | $\{4 \times (1, 2), 4 \times (1, 3), (2, 7)\}$ | 2/21 | 1 | 2 | 2 | 5 | 7 | 10 |
| 20 | $\{4 \times (1, 2), 3 \times (1, 3), (3, 10)\}$ | 1/10 | 1 | 2 | 2 | 5 | 7 | 10 |
| 21 | $\{3 \times (1, 2), (2, 5), 4 \times (1, 3), (1, 4)\}$ | 7/60 | 1 | 2 | 3 | 6 | 8 | 12 |
| 22 | $\{3 \times (1, 2), 7 \times (1, 3)\}$ | 1/6 | 2 | 3 | 4 | 9 | 12 | 17 |
| 23 | $\{2 \times (1, 2), (2, 5), 6 \times (1, 3)\}$ | 1/5 | 2 | 3 | 5 | 10 | 14 | 20 |

The possible baskets with $P_{-1} = 0$ and $P_{-2} = P_{-4} = 1$ are the following (cf. the second paragraph of [5, Case IV of Proof of Theorem 1.8]).

TABLE 3. Baskets with $P_{-1} = 0$ and $P_{-2} = P_{-4} = 1$

| No. | B_X | r_X |
|-----|--|-------|
| 1 | $\{9 \times (1, 2), (1, 3), (1, 7)\}$ | 42 |
| 2 | $\{8 \times (1, 2), (2, 5), (1, 7)\}$ | 70 |
| 3 | $\{8 \times (1, 2), (2, 5), (1, 6)\}$ | 30 |
| 4 | $\{7 \times (1, 2), (3, 7), (1, 6)\}$ | 42 |
| 5 | $\{6 \times (1, 2), (4, 9), (1, 6)\}$ | 18 |
| 6 | $\{7 \times (1, 2), (3, 7), (1, 5)\}$ | 70 |
| 7 | $\{6 \times (1, 2), (4, 9), (1, 5)\}$ | 90 |
| 8 | $\{5 \times (1, 2), (5, 11), (1, 5)\}$ | 110 |
| 9 | $\{4 \times (1, 2), (6, 13), (1, 5)\}$ | 130 |

ACKNOWLEDGMENTS

The second named author would like to thank her advisor, Professor Meng Chen, for his support and encouragement. The authors are grateful to Meng Chen for discussions and suggestions. This work was supported by NSFC for Innovative Research Groups (Grant No. 12121001) and National Key Research and Development Program of China (Grant No. 2020YFA0713200). The first author is a member of LMNS, Fudan University. We would like to thank the referee for useful suggestions.

REFERENCES

- [1] C. Birkar, *Anti-pluricanonical systems on Fano varieties*, Ann. of Math. (2) 190 (2019), no. 2, 345–463.
- [2] J.-J. Chen, J. A. Chen, M. Chen, Z. Jiang, *On quint-canonical birationality of irregular threefolds*, Proc. Lond. Math. Soc. (3) 122 (2021), no. 2, 234–258.

- [3] J. A. Chen, M. Chen, *An optimal boundedness on weak \mathbb{Q} -Fano 3-folds*, Adv. Math. 219 (2008), no. 6, 2086–2104.
- [4] M. Chen, *On anti-pluricanonical systems of \mathbb{Q} -Fano 3-folds*, Sci. China Math. 54 (2011), no. 8, 1547–1560.
- [5] M. Chen, C. Jiang, *On the anti-canonical geometry of \mathbb{Q} -Fano threefolds*, J. Differential Geom. 104 (2016), no. 1, 59–109.
- [6] M. Chen, C. Jiang, *On the anti-canonical geometry of weak \mathbb{Q} -Fano threefolds II*, Ann. Inst. Fourier (Grenoble) 70 (2020), no. 6, 2473–2542.
- [7] I. V. Dolgachev, *Classical algebraic geometry: A modern view*, Cambridge University Press, Cambridge, 2012.
- [8] A. R. Iano-Fletcher, *Working with weighted complete intersections*, Explicit birational geometry of 3-folds, pp. 101–173, London Math. Soc. Lecture Note Ser., 281, Cambridge Univ. Press, Cambridge, 2000.
- [9] C. Jiang, *On birational geometry of minimal threefolds with numerically trivial canonical divisors*, Math. Ann. 365 (2016), no. 1-2, 49–76.
- [10] C. Jiang, *Characterizing terminal Fano threefolds with the smallest anti-canonical volume*, arXiv:2109.03453v1, to appear in Birational geometry, Kähler–Einstein metrics and Degenerations, Moscow, Shanghai and Pohang, June–November 2019, Springer Proceedings in Mathematics & Statistics (PROMS, volume 409).
- [11] C. Jiang, *Characterizing terminal Fano threefolds with the smallest anti-canonical volume, II*, arXiv:2207.03832v1, contributing to the special volume in honor of Professor Shokurov’s seventieth birthday.
- [12] C. Jiang, Y. Zou, *An effective upper bound for anti-canonical volumes of canonical \mathbb{Q} -Fano threefolds*, Int. Math. Res. Not., rnac100, <https://doi.org/10.1093/imrn/rnac100>.
- [13] J. Kollár, Y. Miyaoka, S. Mori, H. Takagi, *Boundedness of canonical \mathbb{Q} -Fano 3-folds*, Proc. Japan Acad. Ser. A Math. Sci. 76 (2000), no. 5, 73–77.
- [14] J. Kollár, S. Mori, *Birational geometry of algebraic varieties*, Cambridge tracts in mathematics, 134, Cambridge University Press, Cambridge, 1998.
- [15] M. Miyanishi, D.-Q. Zhang, *Gorenstein log del Pezzo surfaces of rank one*, J. Algebra 118 (1988), no. 1, 63–84.
- [16] Y. G. Prokhorov, *The degree of Fano threefolds with canonical Gorenstein singularities*. Mat. Sb. 196 (2005), no. 1, 81–122; translation in Sb. Math. 196 (2005), no. 1-2, 77–114.
- [17] Y. G. Prokhorov, *The degree of \mathbb{Q} -Fano threefolds*, Mat. Sb. 198 (2007), no. 11, 153–174; translation in Sb. Math. 198 (2007), no. 11-12, 1683–1702.
- [18] Y. G. Prokhorov, *\mathbb{Q} -Fano threefolds of large Fano index, I*, Doc. Math. 15 (2010), 843–872.
- [19] Y. G. Prokhorov, *On Fano threefolds of large Fano index and large degree*, Mat. Sb. 204 (2013), no. 3, 43–78; translation in Sb. Math. 204 (2013), no. 3-4, 347–382.
- [20] M. Reid, *Young person’s guide to canonical singularities*, Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), pp. 345–414, Proc. Sympos. Pure Math., 46, Part 1, Amer. Math. Soc., Providence, RI, 1987.

CHEN JIANG, SHANGHAI CENTER FOR MATHEMATICAL SCIENCES & SCHOOL OF MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, SHANGHAI, 200438, CHINA

Email address: chenjiang@fudan.edu.cn

YU ZOU, YAU MATHEMATICAL SCIENCES CENTER, TSINGHUA UNIVERSITY, BEIJING, 100084, CHINA

Email address: fishlinazy@tsinghua.edu.cn