

Closed Vortex Filament in a Cylindrical Domain: Circulation Quantization

S.V. TALALOV

Department of Applied Mathematics, Togliatti State University,
14 Belorusskaya str., Tolyatti, Samara region, 445020 Russia.
svt_19@mail.ru

Abstract

This article investigates quantum oscillations of a vortex ring with zero thickness that evolves in a cylindrical domain $V = D \times [0, L]$. The symbol D denotes the planar domain which is bounded by some closed connected curve S . The quantization scheme of this dynamical system is based on the approach proposed by the author earlier. As result, we find the discrete values Γ_n for circulation Γ . In contrast to the traditional approach, where such quantities are usually postulated, the values Γ_n are deduced rigorously as the consequence of the conventional scheme of quantum theory. The model demonstrates the splitting of levels also. In particular, the levels correction values depend on the domain V : both the cylinder height L and the form of the curve S affect the final formula for the quantities Γ_n . Moreover, we prove that the basic circulation levels demonstrate a "fine structure". These anomalous terms, which are proportional to the value \hbar^2 , are calculated in the article as well. The conclusions are compared with some results of numerical simulations by other authors.

keywords: vortex ring, constrained Hamiltonian systems, circulation quantization

PACS numbers: 47.10.Df 47.32.C

As it seems, the theory of quantum fluids cannot be considered complete. Without attempting to make a review of the literature on this topic, we will mention only the Landau two-fluids model (see, for example, [1]) which marks the beginning of a whole new direction of research in this field. Probably, the quantum description of vortex filaments arising in fluids is one of the most interesting areas for research. Thus, a number of authors (see, for example, [2]) assume that the topological defects – vortices – are the key to understanding the turbulence nature in quantum fluids. In the recent paper [3] it was emphasized that the investigation of individual quantum vortices

at certain scales is very important in this regard. In the work [3] the numerical investigations were performed within the framework of a generalized Gross-Pitaevskii model. The determining role of circulation in the study of turbulence was also pointed out in article [4].

The problem of circulation quantization is inextricably linked with these studies. So far, the quantization was postulated [1]:

$$\Gamma_n \equiv \oint_{\gamma} \mathbf{u}(\ell) d\ell = \frac{n\hbar}{\mu} \quad n = 0, 1, 2, \dots, \quad (1)$$

where γ is some closed curve in space E_3 and $\mathbf{u}(\ell) \equiv \mathbf{u}(\mathbf{r}(\ell))$ – fluid velocity in the point $\mathbf{r}(\ell) \in \gamma$. Parameter μ is some parameter with a dimension of mass. As it has been repeatedly stated in the literature, such quantization rules are similar to the quantization rules in the old Bohr quantum theory. In contradiction with the contemporary quantum mechanics, quantum levels in the old theory do not depend on boundary conditions. Moreover, the quantization of each observable should be deduced from the general principles of quantum theory, and not be postulated separately.

In this article, we suggest the scheme of the quantum description of the vortex ring moving in the cylindrical domain. The approach under consideration is a further development of the ideas stated in the work [5]. The suggested method leads to circulation quantization naturally.

We consider the following dynamical system. A closed vortex filament with zero thickness described by the curve $\mathbf{r} = \mathbf{r}(t, \ell)$, moves parallel to the axis of the cylindrical fluid-filled domain $V = D \times [0, L]$. Note that the quantum vortex structures in a similar domain were investigated in the article [6] recently.

The dynamics of our object is described by the Local Induction Equation. Naturally, we assume that the Z-axis coincides with the axis of the considered cylinder. The theory has three dimensional constants: the fluid's density ϱ_0 , the speed of sound in this fluid v_0 and the natural scale length

$$R_0 \simeq \max \|\mathbf{r}_1 - \mathbf{r}_2\|, \quad \mathbf{r}_1, \mathbf{r}_2 \in S.$$

For convenience, we will use the auxiliary constants $t_0 = R_0/v_0$, μ_0 and $\mathcal{E}_0 = \mu_0 v_0^2/2$. These constants define the scales of time, masses and energy correspondingly. In the work [5], parameter μ_0 appeared as the central charge for the centrally extended Galilei group. Despite the fact that there is a natural mass parameter $\tilde{\mu}_0 = \varrho_0 R_0^2 L$ in this work, we will use the mass μ_0 too.

Therefore, we have an additional dimensionless parameter here: $\alpha = \mu_0/\tilde{\mu}_0$. We will discuss its significance in theory later.

For convenience, we introduce the dimensionless parameter $\xi = \ell/R$ and the "dimensionless time" $\tau = t/t_0$ for all evolving closed curves $\mathbf{r}(t, \ell)$ with length $2\pi R$. With these denotations, the vortex filament we are considering is given by the formula

$$\mathbf{r}(\tau, \xi) = \mathbf{q} + R \int_0^{2\pi} [\xi - \eta] \mathbf{j}(\tau, \eta) d\eta, \quad R \ll R_0, \quad (2)$$

where the notation $[x]$ means the integer part of the number $x/2\pi$ and the variables $\mathbf{q} = \mathbf{q}(\tau)$ may be τ -dependent. 2π -periodical vector function $\mathbf{j}(\xi) \in E_3$ defines the unit tangent vector. Because our vortex filament is closed, the following equalities are fulfilled too:

$$\int_0^{2\pi} j_k(\xi) d\xi = 0 \quad (k = x, y, z), \quad (3)$$

The function $\mathbf{r}(\tau, \xi)$ satisfies the LIE equation

$$\partial_\tau \mathbf{r}(\tau, \xi) = \frac{1}{R_0} \partial_\xi \mathbf{r}(\tau, \xi) \times \partial_\xi^2 \mathbf{r}(\tau, \xi). \quad (4)$$

In addition to the equation (4) that describes the evolution of the curve $\mathbf{r}(\cdot, \xi)$, we postulate the standard hydrodynamic formula for the momentum [7]:

$$\tilde{\mathbf{p}} = \frac{\varrho_0}{2} \int \mathbf{r} \times \boldsymbol{\omega}(\mathbf{r}) dV. \quad (5)$$

The vector $\boldsymbol{\omega}(\mathbf{r})$ means vorticity. As well-known (see, for example [8]), the vorticity of the closed vortex filament is calculated by means of the formula

$$\boldsymbol{\omega}(\mathbf{r}) = \Gamma \int_0^{2\pi} \delta(\mathbf{r} - \mathbf{r}(\xi)) \partial_\xi \mathbf{r}(\xi) d\xi, \quad (6)$$

where symbol Γ means circulation. Taking into account the formulae (2), (3) and (6), the following expression for the canonical momentum is deduced by

direct calculations:

$$\begin{aligned}\tilde{\mathbf{p}} &= \varrho_0 R^2 \Gamma \mathbf{f}, \\ \mathbf{f} &= \frac{1}{2} \int_0^{2\pi} \int_0^{2\pi} [\xi - \eta] \mathbf{j}(\eta) \times \mathbf{j}(\xi) d\xi d\eta.\end{aligned}\tag{7}$$

In the work [5] the author investigated the small perturbations of the vortex ring described by the formula (2) with tangent vector

$$\mathbf{j}_0(\xi) = \{-\sin \xi, \cos \xi, 0\}\tag{8}$$

and coordinates $\mathbf{q}_0 = q_x^0 \mathbf{e}_x + q_y^0 \mathbf{e}_y + (q_z^0 + c\tau) \mathbf{e}_z$, where $q_{x,y,z}^0 = \text{const}$. This means that we consider the quantities

$$\mathbf{q} = \mathbf{q}_0 + \varepsilon \mathbf{q}_{\text{pert}}, \quad \mathbf{j}(\tau, \xi) = \mathbf{j}_0(\xi) + \varepsilon \mathbf{j}_{\text{pert}}(\tau, \xi),\tag{9}$$

where small parameter $\varepsilon < 1$ and the values $|\mathbf{q}_{\text{pert}}|$ and $|\mathbf{j}_{\text{pert}}|$ are restricted. Let us note here the work [9], where, in particular, the system of small-perturbed straightforward vortex filaments with certain interactions were investigated.

Next, we will not write the "pert" index explicitly, hoping that this will not lead to misunderstandings. The symmetry of the original object allows the introduction of cylindrical coordinates (ρ, ϕ, z) . Thus, the function $\mathbf{j}(\tau, \xi)$ can be written in form $\mathbf{j}(\tau, \xi) = j_\rho(\tau, \xi) \mathbf{e}_\rho + j_\phi^0(\tau, \xi) \mathbf{e}_\phi + j_z(\tau, \xi) \mathbf{e}_z$ and the linearized equation (4) takes the following form in the cylindrical basis:

$$\begin{aligned}\partial_\tau \mathbf{j}(\tau, \xi) &= \left(j_z(\tau, \xi) + \partial_\xi^2 j_z(\tau, \xi) \right) \mathbf{e}_\rho - \\ &- \left(\partial_\xi^2 j_\rho(\tau, \xi) - 2 \partial_\xi j_\phi(\tau, \xi) \right) \mathbf{e}_z.\end{aligned}$$

This equation demonstrates that $\partial_\tau j_\phi^0(\tau, \xi) \equiv 0$.

Further we will consider the case $j_\phi^0(\tau, \xi) \equiv 0$ only and the complex-valued function $\mathbf{j} = j_\rho + i j_z$. Here it is appropriate to introduce the separate notation \mathbf{j} for the complex perturbation amplitude to avoid any ambiguity. Linearized equation for the amplitude $\mathbf{j}(\tau, \xi)$ was deduced in the article [5]:

$$\partial_\tau \mathbf{j} = -i \partial_\xi^2 \mathbf{j} - \frac{i}{2} (\mathbf{j} - \bar{\mathbf{j}}).$$

After solving this equation, we have the following representation for the function $j(\tau, \xi)$:

$$j(\tau, \xi) = \sum_n j_n e^{i[n\xi + n\sqrt{n^2-1}\tau]}, \quad (10)$$

where $j_n \equiv \text{const}$ and the coefficients \bar{j}_{-n} and j_n are connected to each other as follows:

$$\bar{j}_{-n} = 2 \left[n\sqrt{n^2-1} - n^2 + \frac{1}{2} \right] j_n. \quad (11)$$

Restrictions (3) lead to the restrictions for the coefficients j_0 and $j_{\pm 1}$:

$$j_0 = \bar{j}_0, \quad j_1 = -\bar{j}_{-1}.$$

Although these restrictions are deduced from the constraint (3), they are consistent with the formula (11).

The formulas (7), (10) and (3) reveal the following connection between the perturbation $\mathbf{p} = p_x + ip_y$ of the momentum (5) and the amplitude j :

$$\tilde{\mathbf{p}} = 2\pi\varrho_0 R^2 \Gamma j_{-1}. \quad (12)$$

The projection p_z of the momentum on the Z axis turns out to be unperturbed and is written as follows:

$$\tilde{p}_z = \pi\varrho_0 R^2 \Gamma. \quad (13)$$

Proofs and details can be found in the work [5].

It is also necessary to pay attention to the following important point. The local variable "fluid velocity" is absent in our theory. However, we intend take the movement of the fluid that surrounds the vortex filament into account, in a certain way. To do this, we declare the value Γ as a dynamic variable here, in addition to the variables $\mathbf{j}(\xi)$ and \mathbf{q} . We believe that this assumption allows us to take into account the dynamics of the surrounding fluid in a minimal way. We denote as \mathcal{A} the set of the dynamical variables $\{\mathbf{q}, \mathbf{j}(\xi), \Gamma\}$. Despite the fact that the variable Γ is a conserved quantity, such an extension of the original dynamical system (4) turns out to be nontrivial. In order to make sure of this, we now extend the set \mathcal{A} . Let

us denote as \mathcal{A}' the set of the independent variables $(\mathbf{q}; \mathbf{p}; \mathbf{j}(\xi))$. Formula $\mathbf{p} = \alpha \tilde{\mathbf{p}}$ and formula (7) perform the injection \mathcal{F}

$$\mathcal{F}: \mathcal{A} \xrightarrow{\mathcal{F}} \mathcal{A}', \quad \text{Ran } \mathcal{F} \equiv \Omega \subset \mathcal{A}'$$

for every constant α . Next, we will demonstrate that the set \mathcal{A}' is more appropriate for subsequent quantization of our dynamical system.

It is clear that the formula (7), which defines the momentum through the variables of the set \mathcal{A} , is not executed on the set \mathcal{A}' in general. Indeed, the vectors \mathbf{p} and \mathbf{f} are independent vectors here. But we can state that $\forall \mathbf{p} \in \mathcal{A}', \forall \mathbf{j} \in \mathcal{A}'$ the relation

$$\mathbf{p} = \alpha \varrho_0 R^2 \Gamma \mathbf{B} \mathbf{f}, \quad \mathbf{f} = \mathbf{f}[\mathbf{j}] \quad (14)$$

will be fulfilled for some real number Γ and some matrix $\mathbf{B} = \mathbf{B}(\theta) \in SO(3)$. The angle θ is some angle of rotation around the vector $\mathbf{n} = \mathbf{p} \times \mathbf{f}$. We can introduce certain constraints which define subset $\Omega \subset \mathcal{A}'$ so that one-to-one correspondence $\mathcal{A} \longleftrightarrow \Omega$ is established. For example, we can postulate the equality

$$(\mathbf{p}\mathbf{f})^2 = \mathbf{p}^2 \mathbf{f}^2, \quad \mathbf{f} = \mathbf{f}[\mathbf{j}], \quad \mathbf{p}, \mathbf{j} \in \mathcal{A}'. \quad (15)$$

Indeed, constraint (15) is fulfilled identical if and only if when the proportionality $\mathbf{p} \propto \mathbf{f}$ takes place. Let us note that in this paper we are investigating the circulation only. Despite the fact that the value Γ is defined initially only on the set $\mathcal{A} \approx \Omega$, we can assume that it is also defined everywhere on the set \mathcal{A}' . Formula (14) defines the appropriate expansion $\Omega \rightarrow \mathcal{A}'$. Moreover, the value Γ is same both on the set Ω and the set \mathcal{A}'/Ω because the identity $\mathbf{B}^T \mathbf{B} = I, \forall \mathbf{B} \in SO(3)$. Indeed, the following equality is fulfilled on the set \mathcal{A}' :

$$\exists \Gamma \in \mathbb{R} : \quad |\mathbf{p}|^2 = \alpha^2 \varrho_0^2 R^4 \Gamma^2 |\mathbf{f}|^2.$$

Taking into account the equalities (12) and (13), let us write this formula in the following form:

$$\exists \Gamma \in \mathbb{R} : \quad \Phi_\Gamma \equiv |\mathbf{p}|^2 - \alpha^2 \pi^2 \varrho_0^2 \Gamma^2 R^4 \left(1 + 4\varepsilon^2 |\mathbf{j}_{-1}|^2\right) = 0. \quad (16)$$

The value $|\mathbf{p}|^2 = p_z^2 + \varepsilon^2 |\mathbf{p}|^2$ here. In fact, equality $\Phi_\Gamma = 0$ defines the function $\Gamma = \Gamma(\mathbf{p}, \mathbf{j})$ implicitly. This form will be useful for our subsequent studies.

In order to obtain possible discrete values of circulation Γ_n , we need a quantum version of the model under consideration. Both the hamiltonian structure of the considered dynamical system and subsequent quantization scheme have been considered in the work [5] in detail¹. Without going into details in this paper, we define the fundamental hamiltonian variables here as $p_i, q_j, \mathbf{j}_m, \bar{\mathbf{j}}_n$ and their Poisson brackets:

$$\begin{aligned} \{p_i, q_j\} &= \delta_{ij}, & i, j &= x, y, z, \\ \{\mathbf{j}_m, \bar{\mathbf{j}}_n\} &= (i/\mathcal{E}_0 t_0) \delta_{mn}, & m, n &= -1, -2, \dots \end{aligned}$$

As a consequence, the structure of the Hilbert space \mathbf{H} of quantum states of our dynamical system can be defined as follows:

$$\mathbf{H} = \mathbf{H}_3 \otimes \mathbf{H}_F,$$

where the symbol \mathbf{H}_3 denotes the Hilbert space of a free structureless particle in the domain V (the space $\mathbf{L}^2(V) = \mathbf{L}^2([0, L]) \otimes \mathbf{L}^2(D)$ in our case) and the symbol \mathbf{H}_F denotes the Fock space for the infinite number of the harmonic oscillators. The creation and annihilation operators which are defined in the space \mathbf{H}_F , have standard commutation relations

$$[\hat{a}_m, \hat{a}_n^+] = \hat{I}_F, \quad \hat{a}_m |0\rangle = 0, \quad m, n = 1, 2, \dots, \quad |0\rangle \in \mathbf{H}_F,$$

The variables \mathbf{q}, \mathbf{p} and $\mathbf{j}(\xi)$ are quantized in accordance with the rule

$$q_{x,y,z} \rightarrow q_{x,y,z} \otimes \hat{I}_F, \quad p_{x,y,z} \rightarrow -i\hbar \frac{\partial}{\partial q_{x,y,z}} \otimes \hat{I}_F,$$

$$\mathbf{j}_{-n} \rightarrow \sqrt{\frac{\hbar}{t_0 \mathcal{E}_0}} (\hat{I}_3 \otimes \hat{a}_n),$$

where $n = 1, 2, \dots$. The symbols \hat{I}_3 and \hat{I}_F denote the identical operators in spaces \mathbf{H}_3 and \mathbf{H}_F correspondingly. Next, we will not write both the index $n = 1$ and the constructions $(\dots \otimes \hat{I}_F)$, $(\hat{I}_3 \otimes \dots)$ explicitly: $\hat{a}_1 = \hat{a}$, $(\hat{I}_3 \otimes \hat{a}_n) = \hat{a}_n$ and so on.

As discussed above, in classical theory there should be constraints on the set \mathcal{A}' , to ensure the parallelism of vectors \mathbf{p} and \mathbf{f} . Does this requirement

¹The case $V = \mathbf{R}_3$ was considered in the work [5] only. As a consequence, both energy of the system and circulation take continuous values for this case.

make sense for the quantum version of the system in question? First, there are no quantum states in the "box" V which correspond to the definite momentum \mathbf{p} . Moreover, the average square distance δp^2 of any component $p_{x,y,z}$ is limited from below. Indeed, the following inequalities take place for obvious geometrical reasons:

$$\delta q_{x,y} \leq R_0, \quad \delta q_z \leq L,$$

where $\delta q = \sqrt{(q - \langle q \rangle)^2}$. Taking into account Heisenberg's uncertainty principle, we can conclude that the following inequalities take place for our quantum system:

$$\delta p_{x,y} \geq \frac{\hbar}{2R_0}, \quad \delta p_z \geq \frac{\hbar}{2L}.$$

On the contrary, operator $\hat{\mathbf{f}}$ has states with certain eigenvalues \mathbf{f} . These conclusions make possible to consider space \mathbf{H} completely, not restricted to any subspace which is the consequence of classical constraints. In fact, we quantize the extended classical system \mathcal{A}' instead of the initial system \mathcal{A} . We have demonstrated above that the value Γ we are interested in is the same for the set \mathcal{A}' and the constraint surface $\Omega \subset \mathcal{A}'$.

To find permissible circulation values, we consider the equation

$$\hat{\Phi}_\Gamma |\psi_\Gamma\rangle = 0,$$

where the operator $\hat{\Phi}_\Gamma$ is the quantized function $\Phi_\Gamma(\mathbf{p}, \mathbf{j})$ (see (16)). Explicitly, this equation is written as:

$$\left[-\hbar^2 \left(\frac{\partial^2}{\partial z^2} + \varepsilon^2 \Delta_2 \right) - \alpha^2 \pi^2 \varrho_0^2 \Gamma^2 R^4 \left(1 + \varepsilon^2 \frac{8\hbar}{\mu_0 v_0 R_0} \hat{a}^+ \hat{a} \right) \right] |\psi_\Gamma\rangle = 0. \quad (17)$$

We will look for solutions to this equation in the form

$$|\psi_\Gamma\rangle = |\Psi_1\rangle |\Psi_D\rangle |j\rangle,$$

where

$$|\Psi_1\rangle \in \mathbf{L}^2([0, L]), \quad |\Psi_D\rangle \in \mathbf{L}^2(D), \quad |j\rangle \in \mathbf{H}_F.$$

Let us consider the vectors $|\Psi_1\rangle = |\Psi_1^n\rangle$ and $|\Psi_D\rangle = |\Psi_D^m\rangle$ such that

$$\langle q_z | \Psi_1^n \rangle \equiv \Psi_n(q_z) = \sin\left(\frac{\pi n}{L}\right), \quad \langle q_x, q_y | \Psi_D^m \rangle \equiv \Psi_m(q_x, q_y),$$

in the coordinate representation. Functions $\Psi_m(q_x, q_y)$ are eigenfunctions for the Laplace operator Δ_2 in the planar domain D . These functions satisfy the boundary conditions $\Psi_m(q_x, q_y) = 0$ on the curve $S = \partial D$. Next, we consider the vectors

$$|j\rangle = |j_k\rangle \equiv \frac{1}{\sqrt{k}} (\hat{a}^+)^k |j_0\rangle, \quad |j_0\rangle = \hat{a}_{n_1}^+ \hat{a}_{n_2}^+ \dots \hat{a}_{n_\ell}^+ |0\rangle, \quad n_j > 1.$$

Choosing the solution of the equation (17) in the form

$$|\psi_\Gamma\rangle = |n, m, k\rangle \equiv |\Psi_1^n\rangle |\Psi_D^m\rangle |j_k\rangle,$$

we come to equality

$$\hbar^2 \left[\left(\frac{\pi n}{L} \right)^2 + \varepsilon^2 \lambda_m^2 \right] - \pi^2 \alpha^2 \varrho_0^2 \Gamma^2 R^4 \left(1 + \varepsilon^2 \frac{8 \hbar k}{\mu_0 v_0 R_0} \right) = 0. \quad (18)$$

Numbers n, m, k are natural numbers here and symbols λ_m denote the eigenvalues of the Laplace operator Δ_2 in the domain D . The equality (18) allows us to find the circulation $\Gamma = \Gamma_{n,m,k}$ explicitly. In accordance with our initial supposition (9), we can consider only such quantum numbers n, m, k that ensure the fulfillment of the relations

$$\lambda_m \leq (\pi n / L), \quad 8 \hbar k \leq \mu_0 v_0 R_0.$$

As a consequence, we can write an expression for circulation $\Gamma_{n,m,k}$ as a series by constants ε and \hbar :

$$\Gamma_{n,m,k} = \frac{\hbar n}{\mu_v} \left[1 + \varepsilon^2 \left(\frac{1}{2} \left(\frac{L \lambda_m}{\pi n} \right)^2 - \frac{4 \hbar k}{\mu_0 v_0 R_0} \right) \right] + \mathcal{O}(\varepsilon^4, \hbar^3), \quad (19)$$

where the notation $\mu_v = \mu_0 (R/R_0)^2$ was introduced. The vectors $|n, m, k\rangle$ form the spaces $\mathbf{H}_{n,m,k} \in \mathbf{H}$ in a way that the following decomposition is fulfilled:

$$\mathbf{H} = \bigoplus_{m,n,k} \mathbf{H}_{n,m,k}.$$

The theory under consideration should be enhanced by the "superselection rules": any pure state of the system that is a linear combination of vectors from different spaces $\mathbf{H}_{n,m,k}$ must be forbidden.

In the limit $\varepsilon \rightarrow 0$, provided the quantum numbers m and k are limited, we have the expression

$$\Gamma_{n,m,k} \longrightarrow \Gamma_n = \frac{\hbar n}{\mu_v}, \quad n = 0, 1, 2, \dots$$

which coincides with the standard formula (1). We emphasize that this formula is a consequence of the first principles of quantum theory here. Apparently, constant μ_0 can be chosen based on agreement with a possible experiments.

Let us consider terms of the formula (19) in square brackets that are proportional to the value ε^2 . The first term can be considered as a certain "form - factor", because it is determined only by the form of the domain V . The second term demonstrates the "fine structure" of the levels $\Gamma_{n,m,k}$ that was found. Indeed, it is proportional to the value \hbar^2 . Note also that the second term does not depend on the form of the domain V but depends on the excitations of the initial vortex ring. What does formula (19) predict for the value $\mu_v \Gamma / \hbar$ in its experimental determination? It is clear that we have the "large" peaks in the integer values and certain "small" peaks at the non-integer values. It will be interesting to note that in the work [3] the numerical simulations for circulation confirms this distribution of the value of $\mu_v \Gamma / \hbar$. Although this was explained by errors there, the model considered here gives a different explanation of this fact.

References

- [1] S.J. Putterman. *Superfluid Hydrodynamics*. Elsevier, NY 1974.
- [2] M. Tsubota, K. Fujimoto and S. Yui. J. Low. Temp. Phys. 188, 119 (2017).
- [3] N.P. Müller, J.I. Polanco, G. Krstulovic, Phys. Rev. X 11, 011053 (2021).
- [4] : K.P. Iyer, K.R. Sreenivasan, P. K. Yeung, Phys. Rev. X 9, 041006 (2019).
- [5] S.V. Talalov. Eur. Journ. Mech B/Fluids. **92**. pp. 100 - 106. (2022). arXiv: math-ph/2112.04859v1.
- [6] S. Inui, T. Nakagawa, M. Tsubota, Phys. Rev. B 102, 224511 (2020).

- [7] G.K. Batchelor, *An Introducton to Fluid Dynamics*. Cambrige Univ. Press. (1970).
- [8] S.V. Alekseenko, P.A. Kuibin, V.L. Okulov, *Theory of concentrated vortices*. Springer-Verlag, Berlin Heidelberg (2007).
- [9] A.J. Majda, A.L. Bertozzi, *Vorticity and Incompressible Flow*. Cambridge Univ. Press. (2002).