

The road problem and homomorphisms of directed graphs

Sophie MacDonald

July 17, 2023

Abstract

We make progress on a generalization of the road (colouring) problem. The road problem was posed by Adler-Goodwyn-Weiss and solved by Trahtman. The generalization was posed, and solved in certain special cases, by Ashley-Marcus-Tuncel. We resolve two new families of cases, of which one generalizes the road problem and follows Trahtman’s solution, and the other generalizes a result of Ashley-Marcus-Tuncel with a proof quite different from theirs. Along the way, we prove a universal property for the fiber product of certain graph homomorphisms, which may be of independent interest. We provide polynomial-time algorithms for relevant constructions and decision problems.

1 Introduction

1.1 Background: from toral automorphisms to finite automata

In this paper, we are concerned with a generalization of the road (colouring) problem. That problem, posed by Adler-Goodwyn-Weiss in [1] and solved by Trahtman in [22], asked whether every strongly connected, aperiodic directed graph of constant out-degree is the underlying graph of a synchronizing deterministic finite automaton (DFA). Trahtman’s road colouring theorem (Theorem 4.1 below) gives an affirmative answer, and [11, 12] give a generalization to periodic graphs (Theorem 4.2).

The motivation for the road problem comes from ergodic theory. Specifically, a weak form of the road colouring theorem was used to prove the main theorem of [1], which gives a criterion for measure-theoretic isomorphism of certain Markov chains. This was the first time the road problem was explicitly posed, although the real origin of the problem is the earlier paper [2], which concerns the analogous isomorphism problem for hyperbolic automorphisms of the two-dimensional torus.

The present paper concerns a generalization of the road problem motivated by graph-theoretic invariants for a different, but related, isomorphism relation in ergodic theory. Specifically, Ashley-Marcus-Tuncel [5] identified a graph-theoretic criterion for isomorphism of one-sided stationary Markov chains, implicit in [10], and gave a complete, effectively computable set of isomorphism invariants. They observed that a certain conjectural uniqueness result (the $O(G)$ conjecture, below in §3.1) would, if proven, simplify the set of invariants, and proved the conjecture in a family of special cases. The conjecture reduces to a generalization of the road problem involving certain right-resolving graph homomorphisms (which we call *right-resolvers*; see §2.1). For graphs of constant out-degree, these homomorphisms coincide with road colourings.

1.2 Main ideas and contributions

The main purpose of this paper is to present new results toward the $O(G)$ conjecture. There are two main ideas in the paper. The first idea concerns the *stability relation* of a right-resolver, which was introduced in [16] for DFAs or road colourings. For context, Kari [17] solved the road problem in the Eulerian case by finding, for a given graph, a road colouring with nontrivial (i.e. not merely diagonal) stability relation, then recursively finding a synchronizing road colouring of the strictly smaller quotient

graph (in which the states are stability classes), and lifting it to the original graph. Trahtman’s solution of the full road problem uses the same inductive strategy, paired with a more sophisticated technique for obtaining a colouring with a nontrivial stability relation.

By determining how the stability relation behaves with respect to composition of right-resolvers, we are able to study the aforementioned recursive lifting constructions more systematically, allowing us to apply them toward the $O(G)$ conjecture. In particular, we adapt Trahtman’s proof of the road colouring theorem to cover a larger family of cases (Theorem 4.3), in which the out-degrees of states are allowed to vary cyclically. To do so, we generalize, to the setting of right-resolvers, a sufficient condition for a road colouring to have a nontrivial stable pair, based on the idea of a function graph with a unique tallest tree. This condition is at the heart of all proofs of the road colouring theorem to date, and its importance has motivated detailed analysis [8].

The second main idea in this paper is a graph property that we call *bunchiness*, along with a weaker property called *almost bunchiness*. Bunchy and almost bunchy graphs are characterized by the property that the right-resolvers they admit are unique up to automorphisms in a certain sense (Proposition 5.2). We highlight the implicit role of bunchiness both in [5] and in the road colouring literature, and prove the $O(G)$ conjecture for bunchy and almost bunchy graphs (Theorem 5.4). Furthermore, we show that the fiber product of right-resolving homomorphisms satisfies a universal property (Theorem 5.7) that further highlights the essential role of bunchy graphs.

Motivated by these results, we introduce a new conjecture, which we call the bunchy factor conjecture (see §6), asserting essentially that the $O(G)$ conjecture can be proved using the stability approach that Trahtman used to prove the road colouring theorem, with bunchy graphs as the base of the recursion. Another way of articulating this conjecture is that the barrier to proving the $O(G)$ conjecture is our lack of a sufficiently general method of producing homomorphisms with nontrivial stability relation. The fact that the bunchy factor conjecture implies the $O(G)$ conjecture is made explicit in Proposition 6.2, which relies primarily on Proposition 5.9, a uniqueness result that uses the universal property of the fiber product in an essential way.

1.3 Organization of the paper

In §2, we recall and adapt standard material on graphs and homomorphisms. In §3, we define the stability relation for a right-resolver, give its main structural properties, and relate it to synchronization. In §4, we recall from [5] the connection between the road problem and the $O(G)$ conjecture, and state our generalization of the road colouring theorem.

In §5, we introduce the concepts of bunchiness and almost bunchiness and present results involving them, including the $O(G)$ conjecture for bunchy and almost bunchy graphs and the universal property of the fiber product. In §6, we pose the bunchy factor conjecture, which has several equivalent formulations, and discuss its relation to the $O(G)$ conjecture and the road problem. In §7, we give polynomial-time algorithms for construction and decision problems involving right-resolvers, and discuss the algorithmic implications of the $O(G)$ and bunchy factor conjectures.

The proofs of many results in §§2–7, comprising the structural properties of graphs and right-resolvers, are deferred to §8. The proof of our generalization of the road colouring theorem is deferred to §9.

2 Graphs and graph homomorphisms

2.1 Basic definitions

We take all graphs to be finite and directed. A graph G consists of a set $V(G)$ of states, or vertices, and a set $E(G)$ of edges, together with a pair of maps $s, t : E(G) \rightarrow V(G)$ giving the source and target of each edge. Loops (edges e with $s(e) = t(e)$) and parallel edges (distinct edges e, e' with $s(e) = s(e')$, $t(e) = t(e')$) are allowed. For $I \in V(G)$, we write $E_I(G) = s^{-1}(I)$ for the set of outgoing edges from I . We write $F(I) = t(E_I(G))$ for the set of follower states of I , and we write $E_{IJ}(G) = s^{-1}(I) \cap t^{-1}(J)$

for the set of edges from I to J . We write $L(G)$ for the language of G , i.e. the set of finite edge paths in G , i.e. $e_1 e_2 \dots e_n$ where $t(e_i) = s(e_{i+1})$, $1 \leq i \leq n-1$. We also refer to elements of $L(G)$ as *words*. The maps s, t extend to $s, t : L(G) \rightarrow V(G)$ by $s(e_1 \dots e_n) = s(e_1)$, $t(e_1 \dots e_n) = t(e_n)$. We define $L_I(G) = \{u \in L(G) \mid s(u) = I\}$, and $L_{IJ}(G) = \{u \in L_I(G) \mid t(u) = J\}$. A *cycle* in G is a path $u \in L(G)$ with $s(u) = t(u)$, i.e. an element of $L_{II}(G)$ for some $I \in V(G)$.

A *graph homomorphism* $\Phi : G \rightarrow H$ is a pair of maps $\Phi : E(G) \rightarrow E(H)$, $\partial\Phi : V(G) \rightarrow V(H)$ such that $s \circ \Phi = \partial\Phi \circ s$ and $t \circ \Phi = \partial\Phi \circ t$. If there is a surjective homomorphism from G to H , then we say that H is a *factor* of G , and that G is an *extension* of H . Observe that every factor of a strongly connected graph is strongly connected. A graph homomorphism $\Phi : G \rightarrow H$ induces a map $L(G) \rightarrow L(H)$ (also written Φ) in the obvious way.

A graph isomorphism is a homomorphism that is injective and surjective (i.e. on both edges and states), and an automorphism of a graph G is an isomorphism from G to itself. We denote the group of automorphisms of G by $\text{Aut}(G)$, and by $P(G)$ the (normal) subgroup of $\text{Aut}(G)$ which acts trivially on states and permutes parallel edges. We generally identify isomorphic graphs, and use the symbol $=$ to denote isomorphism, except when discussing algorithms for deciding isomorphism, or confirming that the automorphism group of a given graph is trivial.

We now introduce the class of homomorphisms with which we are concerned.

Definition 2.1 (right-resolver). Let G, H be graphs. Let $\Phi : G \rightarrow H$ be a surjective graph homomorphism. We say that Φ is *right-resolving*, or is a *right-resolver*, if, for each $I \in V(G)$, the restriction $\Phi|_{E_I(G)} : E_I(G) \rightarrow E_{\partial\Phi(I)}(H)$ is a bijection. We denote the set of right-resolvers $G \rightarrow H$ by $\text{hom}_R(G, H)$, and we write $H \leq_R G$ if $\text{hom}_R(G, H) \neq \emptyset$.

Remark 2.1. The term “right-resolving” comes from symbolic dynamics, where the words are of primary importance and the actual graph is secondary. A graph homomorphism $\Phi : G \rightarrow H$ is right-resolving if and only if the associated map $\Phi : L(G) \rightarrow L(H)$ satisfies a certain condition on the symbols (edges) appearing to the right of a given symbol in a word. See [5], §8.2 for details if interested.

Remark 2.2. The class of right-resolving graph homomorphisms is closed under composition. This reduces to the fact, applied to the outgoing edges from each state, that a composition of bijections is a bijection. This means that the relation \leq_R is transitive, and since the graphs are finite, it is clearly antisymmetric, so it is indeed a partial order on the set of all graphs (really on the set of equivalence classes of graphs up to isomorphism).

The following lemma is evident but we state it explicitly for future reference.

Lemma 2.1. *The image of a right-resolver is determined up to graph isomorphism by the partition of the domain into fibers. That is, if $H_1, H_2 \leq_R G$ via $\Phi_i \in \text{hom}_R(G, H_i)$, and for any $I_1, I_2 \in V(G)$ we have $\partial\Phi_1(I_1) = \partial\Phi_1(I_2)$ if and only if $\partial\Phi_2(I_1) = \partial\Phi_2(I_2)$, then in fact the H_i are isomorphic.*

Note that the converse is not true: for a given G, H with $H \leq_R G$, there may exist $\Phi, \Phi' \in \text{hom}_R(G, H)$ with distinct partitions $\{(\partial\Phi)^{-1}(I) \mid I \in V(H)\}$, $\{(\partial\Phi')^{-1}(I) \mid I \in V(H)\}$ of $V(G)$. However, this cannot occur when H is \leq_R -minimal:

Theorem 2.2 ([5], Theorem 3.2 and Corollary 3.3(a)). *For any graph G , there exist a unique \leq_R -minimal graph $M(G) \leq_R G$ and a unique map $\Sigma_G : V(G) \rightarrow V(M(G))$ such that $\partial\Phi = \Sigma_G$ for any $\Phi \in \text{hom}_R(G, M(G))$.*

The construction of $M(G)$ was first given in [14], though not in this notation. We discuss the proof of Theorem 2.2 in §8.1. The notion of $M(G)$, for a graph G , provides context for road colourings:

Definition 2.2 (M_D and road colourings). For $D \geq 1$, let M_D be the graph with a single state and D self-loops. For a graph G of constant out-degree D , a *road colouring* of G is a right-resolver $G \rightarrow M_D$.

Note that each M_D is \leq_R -minimal, and that $\text{hom}_R(G, M_D)$ is nonempty if and only if G has constant out-degree D , in which case $M(G) = M_D$.

2.2 Subgraphs and connectedness

A *sink* in a graph G is a state $I \in V(G)$ is a state with no outgoing edges, i.e. $F(I) = \emptyset$. We assume throughout that all graphs are sink-free; this is purely for convenience, as all of the results that do not require strong connectedness can be proved for graphs with sinks, with routine but tedious modifications to the proofs. We say that a graph G is *strongly connected*, or *irreducible*, if for any ordered pair $I, J \in V(G)$, there is a (directed) edge path in G from I to J , i.e. $L_{IJ}(G) \neq \emptyset$. Note that strongly connected graphs are sink-free. The *period* $\text{per}(G)$ of a strongly connected graph G is the gcd of its cycle lengths.

A graph H is a *subgraph* of a graph G if $E(H) \subseteq E(G)$, $V(H) \subseteq V(G)$, and the maps s, t with respect to H agree with their counterparts on G , restricted to H . An *induced subgraph* of a graph G is a subgraph H such that $E_{IJ}(H) = E_{IJ}(G)$ for every $I, J \in V(H)$. A *strong component* of a graph is a maximal strongly connected subgraph, i.e. a strongly connected subgraph that is not a proper subgraph of another strongly connected graph.

A *principal subgraph* of a graph G is a subgraph H such that $E_I(H) = E_I(G)$ for every $I \in V(H)$. Note that every principal subgraph is induced. Note also that if H is a principal subgraph of G , and K is a principal subgraph of H , then K is a principal subgraph of G . A *principal component* is a strongly connected principal subgraph; note that the principal components are precisely the minimal principal subgraphs. In particular, any two principal components of a given graph have disjoint sets of states. The principal components of G correspond to the sink states in the *condensation* of G , which is the directed acyclic graph in which the states are the strong components, or maximal strongly connected subgraphs, of G , and with an edge $C_1 \rightarrow C_2$ in the condensation if there is an edge $I_1 \rightarrow I_2$ for $I_i \in V(C_i)$ in G .

Let G, H be graphs with $H \leq_R G$, and let $\Phi \in \text{hom}_R(G, H)$. Let K be a subgraph of G . Note, by the right-resolving property, that in order for $\Phi|_K : K \rightarrow H$ to be surjective, it is necessary and sufficient that $\partial\Phi|_{V(K)} : V(K) \rightarrow V(H)$ be surjective and that K be a principal subgraph of G .

Remark 2.3. The road problem and the $O(G)$ problem were both originally raised for strongly connected graphs, which is a natural restriction given the origins of both problems in the ergodic theory of stationary Markov chains. Moreover, strong connectedness is used in an important way in a lemma used to prove both the road colouring theorem and the almost bunchy case of the bunchy factor conjecture. This is why the $O(G)$ conjecture and the bunchy factor conjecture are stated only for strongly connected graphs.

However, it is quite natural from an automata-theoretic perspective, especially concerning computational complexity, to consider graphs that are not strongly connected. For instance, Eppstein [15] shows that it is NP-complete to determine whether the minimum length of a synchronizing word for a given synchronizing DFA is at most some given value. Eppstein's examples are not strongly connected. There are also several graph problems from symbolic dynamics [13] that are NP-complete in general but have polynomial-time algorithms in the strongly connected case.

3 Stability and synchronization

3.1 Transitions, stability, and synchronization

A right-resolver on a graph G induces transition maps on $V(G)$ in the standard way:

Definition 3.1 (transition map). Let G, H be graphs with $H \leq_R G$. Let $\Phi \in \text{hom}_R(G, H)$. For $I \in V(G)$ and $u \in L_{\partial\Phi(I)}(H)$, we write $I \cdot u$ for the terminal state $t(\gamma)$ of the unique $\gamma \in L_I(G)$ with $\Phi(\gamma) = u$. That is, $I \cdot u = t((\Phi|_{L_I(G)})^{-1}(u))$. We denote by S_Φ the set of maps of the form $I \mapsto I \cdot u$ with respect to Φ .

We now introduce the notion of a *congruence* (see [9], Chapter 1, or [17], §3), of which we will see two important examples. The main example will be the stability relation, but we will also use a congruence in Proposition 8.7 to construct the maximal bunchy factor $B(G)$ of a given graph G .

Definition 3.2 (congruences and quotients). Let G, H be graphs with $H \leq_R G$, let $\Phi \in \text{hom}_R(G, H)$, and let \sim be an equivalence relation on $V(G)$. We say that \sim is a *congruence* with respect to Φ if it is invariant under transitions, i.e. for all $I \in V(H)$, all $u \in L_I(H)$, and all $I_1, I_2 \in (\partial\Phi)^{-1}(I)$ with $I_1 \sim I_2$, we have $I_1 \cdot u \sim I_2 \cdot u$. We “overload” a congruence \sim by defining it also on paths (in particular, edges), by saying that $\gamma_1 \sim \gamma_2$, for $\gamma_1, \gamma_2 \in L(G)$, if $\Phi(\gamma_1) = \Phi(\gamma_2)$ and $s(\gamma_1) \sim s(\gamma_2)$. Define the *quotient graph* G/\sim by $V(G/\sim) = V(G)/\sim$, $E(G/\sim) = E(G)/\sim$, $s([e]_\sim) = [s(e)]_\sim$, and $t([e]_\sim) = [t(e)]_\sim$.

Remark 3.1. Let G, H be graphs with $H \leq_R G$, let $\Phi \in \text{hom}_R(G, H)$, and let \sim be a congruence on G with respect to Φ . Observe that there are right-resolvers $G \rightarrow G/\sim$, $G/\sim \rightarrow H$ which compose to Φ , where the right-resolver $G \rightarrow G/\sim$ is the quotient map, and the right-resolver $G/\sim \rightarrow H$ takes a \sim class to the image in H of any of its representatives.

Remark 3.2. The coarsest congruence, with respect to a right-resolver $\Phi \in \text{hom}_R(G, H)$, is the total relation on the fibers, i.e. the relation $\bigsqcup_{I \in V(H)} ((\partial\Phi)^{-1}(I))^2$. The quotient of G by this relation is simply H , with Φ as the quotient map.

Definition 3.3 (stability relation for a right-resolver). Let G, H be graphs with $H \leq_R G$ and let $\Phi \in \text{hom}_R(G, H)$. The *stability relation* for Φ , written \sim_Φ , is the equivalence relation on $V(G)$ defined as follows: for $I \in V(H)$ and $I_1, I_2 \in (\partial\Phi)^{-1}(I)$, $I_1 \sim_\Phi I_2$ if and only if for all $u \in L_I(H)$, there exists $v \in L_{t(u)}(H)$ such that $I_1 \cdot uv = I_2 \cdot uv$.

Lemma 3.1. *Let G, H be graphs with $H \leq_R G$ and let $\Phi \in \text{hom}_R(G, H)$. The stability relation \sim_Φ is a congruence with respect to Φ .*

Proof. Let $I \in V(H)$ and $I_1, I_2 \in (\partial\Phi)^{-1}(I)$ with $I_1 \sim_\Phi I_2$. Let $u \in L_I(H)$ and let $v \in L_{t(u)}(H)$. Since $I_1 \sim_\Phi I_2$, there exists $w \in L_{t(v)}(H)$ such that $I_1 \cdot uvw = I_2 \cdot uvw$. Therefore $I_1 \cdot u \sim_\Phi I_2 \cdot u$, so \sim_Φ is indeed a congruence. \square

We now define *synchronizing* right-resolvers in terms of the stability relation, then show in Proposition 3.2 that, at least in the strongly connected case, this definition is equivalent to a more obvious notion of synchronization for a right-resolver.

Definition 3.4 (synchronizer). Let G, H be graphs with $H \leq_R G$ and let $\Phi \in \text{hom}_R(G, H)$. We say that Φ is *synchronizing*, or is a *synchronizer*, if each fiber $(\partial\Phi)^{-1}(I)$, $I \in V(H)$, is a \sim_Φ class. We denote the set of synchronizers $G \rightarrow H$ by $\text{hom}_S(G, H)$, and we write $H \leq_S G$ if $\text{hom}_S(G, H) \neq \emptyset$.

Note that \sim_Φ depends on Φ only through S_Φ . That is, if Φ, Φ' are such that $S_\Phi = S_{\Phi'}$, then $\sim_\Phi = \sim_{\Phi'}$.

Proposition 3.2. *Let G, H be graphs with $H \leq_R G$ and let $\Phi \in \text{hom}_R(G, H)$. Then Φ is synchronizing if and only if for every $I \in V(H)$, there exists $u \in L_I(H)$ with $|(\partial\Phi)^{-1}(I) \cdot u| = 1$.*

Proof. First, suppose that Φ is synchronizing and let $I \in V(H)$. If $|(\partial\Phi)^{-1}(I)| = 1$, then we are done. Otherwise, there exist at least two distinct states $I_1, I_2 \in (\partial\Phi)^{-1}(I)$, and $I_1 \sim_\Phi I_2$ since $(\partial\Phi)^{-1}(I)$ is a \sim_Φ class by assumption. Therefore, there exists $u_1 \in L_I(H)$ with $I_1 \cdot u_1 = I_2 \cdot u_1$. In particular, $|(\partial\Phi)^{-1}(I) \cdot u_1| < |(\partial\Phi)^{-1}(I)|$. Continuing inductively, we can produce a sequence of words u_1, \dots, u_n such that $t(u_i) = s(u_{i+1})$ and $|(\partial\Phi)^{-1}(I) \cdot u_1 \cdots u_n| = 1$. This proves the first claim.

For the converse, let $I \in V(H)$ and $I_1, I_2 \in (\partial\Phi)^{-1}(I)$. We will show that $I_1 \sim_\Phi I_2$. Let $v \in L_I(H)$ be arbitrary. By the assumption that each fiber can be collapsed to a single state, let $w \in L_{t(v)}(H)$ be such that $|(\partial\Phi)^{-1}(t(v)) \cdot w| = 1$. Then $|(\partial\Phi)^{-1}(I) \cdot vw| = 1$; in particular, $I_1 \cdot vw \sim_\Phi I_2 \cdot vw$. Therefore $(\partial\Phi)^{-1}(I)$ is a \sim_Φ class, and Φ is synchronizing. \square

We now summarize the structure of stability, in the sense of its behaviour with respect to composition of right-resolvers. For the proof of Theorem 3.3, see §8.2.

Theorem 3.3. *Let G, K, H be graphs with $H \leq_R K \leq_R G$. Let $\Psi \in \text{hom}_R(G, K)$, $\Delta \in \text{hom}_R(K, H)$, and let $\Phi = \Delta \circ \Psi$.*

1. *The \sim_Ψ classes in $V(G)$ are the intersections of \sim_Φ classes with $\partial\Psi$ fibers. In particular, Ψ is synchronizing if and only if every $\partial\Psi$ fiber is contained in a \sim_Φ class.*
2. *If $K = G / \sim_\Phi$ and Ψ is the quotient map for \sim_Φ , then Ψ is synchronizing and \sim_Δ is trivial.*
3. *If \sim_Δ is trivial, then $\sim_\Phi = \sim_\Psi$.*
4. *Φ is synchronizing if and only if both Ψ and Δ are synchronizing.*

The following observation follows immediately from Theorem 3.3(4).

Corollary 3.4. *The relation \leq_S is transitive, and is thus a partial order on the class of graphs (again, really isomorphism classes of graphs), refining the partial order \leq_R .*

Conjecture (*O(G) conjecture, Question 4.6 in [5]*). *Let G be a strongly connected graph. Then the set of graphs H with $H \leq_S G$ has a unique \leq_S -minimal element $O(G)$.*

Remark 3.3. This remark is intended for readers interested in algebraic or categorical perspectives on automata theory. Recall that for a complete DFA, or road colouring $\Phi \in \text{hom}_R(G, M_D)$, where G is a graph of constant out-degree D , the set S_Φ of transition maps forms a transformation semigroup under composition. Indeed, a complete DFA is essentially a finite transformation semigroup together with a choice of generators; this perspective is taken explicitly in [4, 11] and mentioned in [20], the first paper on the road problem after [1].

For a general right-resolver $\Phi \in \text{hom}_R(G, H)$, one could see S_Φ as the semigroup of transitions of a partial finite automaton (PFA), where a given transition is defined only on a single fiber. However, it is more helpful to see S_Φ as a *semigroupoid* (equivalently, if the empty word is included, a small category). One reason is that, as we show in §7, it can be decided in polynomial time whether Φ is synchronizing, and the length of a word synchronizing a given fiber is bounded by a polynomial in $|V(G)|$. This is in contrast to the high level of complexity typical of related problems in subset synchronization and synchronization of PFAs [7, 23].

The reader may verify as an exercise, generalizing Cayley's theorem or specializing the Yoneda lemma, that every finite semigroupoid is isomorphic to S_Φ for some graphs G, H (although possibly with sinks) and some $\Phi \in \text{hom}_R(G, H)$, with appropriate generalizations to the infinite case. Moreover, just as every group is a quotient of a free group, S_Φ is a quotient of the free semigroupoid $L(H)$.

3.2 Sufficient conditions for stability

We now give a pair of sufficient conditions for nontrivial stability, both of which are used in the proof of the road colouring theorem and one of which is also used in §5 to obtain a right-resolver with nontrivial stability on an almost bunchy graph.

The first condition involves a special case of the operation known as in-amalgamation ([18], §2.4):

Lemma 3.5. *Let G, H be graphs with $H \leq_R G$ and let $\Phi \in \text{hom}_R(G, H)$ be a right-resolver. Let $I \in V(H)$. Suppose that there exist $I_1, I_2 \in (\partial\Phi)^{-1}(I)$ such that $|E_{I_1 J}(G)| = |E_{I_2 J}(G)|$ for all $J \in V(G)$. Then there exists $\Phi' \in \text{hom}_R(G, H)$ such that $I_1 \sim_{\Phi'} I_2$.*

Proof. We first claim that $F(I_1) = F(I_2)$, where we recall from §2 the notation $F(\cdot)$ for the set of follower states of a given state. Indeed, the assumption that $|E_{I_1 J}(G)| = |E_{I_2 J}(G)|$ for all $J \in V(G)$ implies in particular that, for any $J \in V(G)$, we have $|E_{I_1 J}(G)| > 0$, equivalently $J \in F(I_1)$, if and only $|E_{I_2 J}(G)| > 0$, equivalently $J \in F(I_2)$.

Let $F = F(I_1) = F(I_2)$. For each $J \in F$, choose a bijection $\Theta_J : E_{I_2 J}(G) \rightarrow E_{I_1 J}(G)$. Define Φ' as follows: $\partial\Phi' = \partial\Phi$, $\Phi'|_{E(G) \setminus E_{I_2}(G)} = \Phi|_{E(G) \setminus E_{I_2}(G)}$, and, for each $J \in F$, $\Phi'|_{E_{I_2 J}(G)} = \Phi|_{E_{I_1 J}(G)} \circ \Theta_J$.

That is, Φ' agrees with Φ on states and on all edges with initial state other than I_2 , but may disagree with Φ on the outgoing edges from I_2 , specifically by permutations of parallel edges. Since $\Phi'|_{E_{I_2}(G)} : E_{I_2}(G) \rightarrow E_I(G)$ is a bijection (being a composition of bijections), Φ' is indeed right-resolving.

By the construction of Φ' , for any $a \in E_I(G)$, we have $I_1 \cdot a = I_2 \cdot a$. Any $w \in L_I(H)$ is of the form $w = au$ with $a \in E_I(G)$, so $I_1 \cdot w = I_2 \cdot w$. Therefore $I_1 \sim_{\Phi'} I_2$. \square

In Lemma 3.5, the states I_1, I_2 are said to be *in-amalgamated* by the operation $G \rightarrow G / \sim_{\Phi'}$; the inverse operation is known as *in-splitting*. The lemma shows in particular that no fiber of a \leq_S -minimal graph G over $M(G)$ has two states that can be in-amalgamated. Trahtman applies a special case of this fact to graphs of constant out-degree, and we follow his line of application; see the first paragraph of the proof of Theorem 4.3, found in §9.2.

The second sufficient condition, given in Proposition 3.8, concerns *minimal images*:

Definition 3.5 (minimal image). Let G, H be graphs with $H \leq_R G$, and let $\Phi \in \text{hom}_R(G, H)$. A *minimal image* is a set of the form $U = (\partial\Phi)^{-1}(I) \cdot u$ for some $I \in V(H)$ and $u \in L_I(H)$, such that $|U \cdot v| = |U|$ for any $v \in L_{t(u)}(H)$.

Remark 3.4. Let G, H be graphs with $H \leq_R G$, and let $\Phi \in \text{hom}_R(G, H)$. For any $I \in V(H)$, any $u \in L_I(H)$, and any $v \in L_{t(u)}(H)$, if $U = (\partial\Phi)^{-1}(I) \cdot u$, we clearly have $|U \cdot v| \leq |U|$. If there exists $v \in L_{t(u)}(H)$ such that this inequality is strict, then $|U|$ is not minimal, i.e. U is not a minimal image. This is the reason for the term.

For a right-resolver Φ on a strongly connected graph, all minimal images have the same size, which is called the *degree* of Φ , and a word with minimal image is called a *magic word*. See [5], §9.1 for a treatment of degrees, using symbolic dynamics. In this paper, we only need a small fragment of the theory of degree, which we establish in a self-contained way with no connectedness assumptions, using the properties of stability.

We use minimal images to give a criterion for stability that can be seen as a pairwise version of Proposition 3.2.

Lemma 3.6. *Let G, H be graphs with $H \leq_R G$, and let $\Phi \in \text{hom}_R(G, H)$. For $I \in V(H)$ and $I_1, I_2 \in (\partial\Phi)^{-1}(I)$, we have $I_1 \sim_{\Phi} I_2$ if and only if $I_1 \cdot u = I_2 \cdot u$ for every word $u \in L_I(H)$ such that $(\partial\Phi)^{-1}(I) \cdot u$ is a minimal image.*

Proof. Let $I \in V(H)$ and let $I_1, I_2 \in (\partial\Phi)^{-1}(I)$. First suppose that $I_1 \sim_{\Phi} I_2$, and let $u \in L_I(H)$. If $I_1 \cdot u \neq I_2 \cdot u$, then let $v \in L_{t(u)}(H)$ be such that $I_1 \cdot uv = I_2 \cdot uv$. Then $|(\partial\Phi)^{-1}(I) \cdot uv| < |(\partial\Phi)^{-1}(I) \cdot u|$, so $(\partial\Phi)^{-1}(I) \cdot u$ is not a minimal image. Contrapositively, if $(\partial\Phi)^{-1}(I) \cdot u$ is a minimal image, then $I_1 \cdot u = I_2 \cdot u$.

Conversely, suppose that $I_1 \cdot u = I_2 \cdot u$ for every word $u \in L_I(H)$ such that $(\partial\Phi)^{-1}(I) \cdot u$ is a minimal image. Let $r = \min_{u \in L_I(H)} |(\partial\Phi)^{-1}(I) \cdot u|$ and let $u \in L_I(H)$. We claim that there exists $w \in L_{t(u)}(H)$ with $I_1 \cdot uw = I_2 \cdot uw$. Indeed, if $I_1 \cdot u \neq I_2 \cdot u$, then $(\partial\Phi)^{-1}(I) \cdot u$ is not a minimal image, so there exists $v \in L_{t(u)}(H)$ such that $|(\partial\Phi)^{-1}(I) \cdot uv| < |(\partial\Phi)^{-1}(I) \cdot u|$. We can thus inductively extend v to obtain the desired w , so indeed $I_1 \sim_{\Phi} I_2$. \square

The following easy observation about minimal images is the main reason that our results toward the bunched factor conjecture (the generalized road colouring theorem and the related result for almost bunched graphs) require strong connectedness.

Lemma 3.7. *Let G, H be strongly connected graphs with $H \leq_R G$, and let $\Phi \in \text{hom}_R(G, H)$. Every minimal image for Φ has the same cardinality, and for every $I' \in V(G)$, there exists a minimal image U with $I' \in U$.*

In the proof of the generalized road colouring theorem, we apply Lemma 3.7 both directly and via Proposition 3.8. The proof of Proposition 3.8 is adapted from the proof of Lemma 10.4.4 in [9].

Proposition 3.8. *Let G, H be strongly connected graphs with $H \leq_R G$ and let $\Phi \in \text{hom}_R(G, H)$. Let $I, J \in V(H)$ and let $u_1, u_2 \in L_{I, J}(H)$ be such that $U_i = (\partial\Phi)^{-1}(I) \cdot u_i$ are minimal images. If $|U_1 \Delta U_2| = 2$, say $U_1 \Delta U_2 = \{J_1, J_2\}$ (where Δ denotes the symmetric difference), then $J_1 \sim_\Phi J_2$.*

Proof. Let $r = |U_1| = |U_2|$. Suppose without loss of generality that $J_i \in U_i$, and let $U_0 = U_1 \cap U_2$, so that $U_i = U_0 \cup \{J_i\}$. For any $v \in L_J(H)$, we have $(U_1 \cup U_2) \cdot v = (U_0 \cdot v) \cup (\{J_1, J_2\} \cdot v)$. We must have $|U_0 \cdot v| = |U_0| = r - 1$ and $J_i \cdot v \notin U_0 \cdot v$, since otherwise we would have $|(\partial\Phi)^{-1}(I) \cdot u_i v| = |U_i \cdot v| < r$, contradicting the minimality assumption. Therefore $r - 1 + |\{J_1, J_2\} \cdot v| = |(U_1 \cup U_2) \cdot v|$. Note that $|(U_1 \cup U_2) \cdot v| \in \{r, r + 1\}$.

Let $v \in L_J(H)$ be such that $(\partial\Phi)^{-1}(J) \cdot v$ is a minimal image. By Lemma 3.6, to show that $J_1 \sim_\Phi J_2$, we need to show that $|\{J_1, J_2\} \cdot v| = 1$, or equivalently $|(U_1 \cup U_2) \cdot v| \leq r$. By strong connectedness and Lemma 3.7, we have $|(\partial\Phi)^{-1}(J) \cdot v| = r$. But $(U_1 \cup U_2) \cdot v \subseteq (\partial\Phi)^{-1}(J) \cdot v$, so indeed $|(U_1 \cup U_2) \cdot v| \leq r$. \square

4 The $O(G)$ conjecture and the road problem

4.1 Generalization of the road colouring theorem

We first introduce the class of graphs involved in the theorem. A *bunch* in a graph G is a state $I \in V(G)$ with $|F(I)| = 1$. (This terminology, introduced in [22] and used also in [9], is the origin of our term *bunchy*, introduced in §5.) A strongly connected graph in which every state is a bunch is a *cycle of bunches*. Let M be a cycle of bunches with $V(M) = \{I_0, \dots, I_{p-1}\}$, where $F(I_i) = \{I_{i+1}\}$ (subscripts indexing states in a cycle of length p should be read modulo p throughout), and let $D_i = |E_{I_i}(M)|$. Note that M is \leq_R -minimal if and only if the sequence of out-degrees D_0, \dots, D_{p-1} is not a cyclic shift of a sequence obtained by concatenating a strictly shorter sequence with itself more than once.

Let M be a \leq_R -minimal cycle of bunches. Let $O_{M, q}$ be the cycle of bunches in which the sequence of out-degrees consists of q cyclic repetitions of D_0, \dots, D_{p-1} . Note that $O_{M, 1} = M$. Observe that, for a strongly connected graph G with $M = M(G)$ a cycle of bunches, if $q = \text{per}(G)/\text{per}(M)$ and H is a cycle of bunches with $H \leq_S G$, then $H = O_{M, q}$. Let $O_{D, p} = O_{M_D, p}$ be the cycle of bunches of period p and constant out-degree D . Note that $O_{D, 1} = M_D$. For a strongly connected, aperiodic graph G of constant out-degree D , a synchronizer $G \rightarrow M_D$ is precisely a synchronizing road colouring of G .

Theorem 4.1 (Trahtman, [22]). *Let G be a strongly connected, aperiodic graph of constant out-degree D . Then $M_D \leq_S G$.*

Theorem 4.2 (Béal-Perrin [12], Budzban-Feinsilver [11]). *Let G be a strongly connected graph of constant out-degree D and period p . Then $O_{D, p} \leq_S G$.*

We prove the following generalization:

Theorem 4.3. *Let G be a strongly connected graph such that $M(G)$ is a cycle of bunches. Let $q = \text{per}(G)/\text{per}(M(G))$. Then $O_{M(G), q} \leq_S G$.*

The proof (see §9) follows that of Theorems 4.1 and 4.2. The strategy is to show that if G is not itself a cycle of bunches, then there exists $\Phi \in \text{hom}_R(G, M(G))$ with \sim_Φ nontrivial, by constructing two minimal images that differ by a pair and applying Proposition 3.8. A very similar strategy is used to prove the bunchy factor conjecture for almost bunchy graphs (Corollary 5.6), the (substantial) difference being the different techniques used to obtain the requisite pair of minimal images.

4.2 The $O(G)$ conjecture implies the road colouring theorem

We now recall the sense in which the $O(G)$ conjecture was first understood to relate to the road problem. Although the road colouring theorem clearly implies the $O(G)$ conjecture for strongly connected, aperiodic graphs of constant out-degree, the converse implication may not be apparent. Indeed, the

$O(G)$ conjecture asserts that $O(G)$ is well-defined for every strongly connected graph, but does not immediately say how to compute $O(G)$, whereas the road colouring theorem explicitly specifies the form of $O(G)$ for the graphs G to which it applies. However, the $O(G)$ conjecture does imply the road colouring theorem, via a key result from [1], for which we require a definition.

Definition 4.1 (higher edge graph). Let G be a graph. For $k \geq 2$, the k -th *higher edge graph* of G is the graph $G^{[k]}$ with edge set consisting of edge paths $e_1e_2 \cdots e_{k-1}e_k$ of length k in G , and states given by $s(e_1e_2 \cdots e_{k-1}e_k) = e_1e_2 \cdots e_{k-1}$, $t(e_1e_2 \cdots e_{k-1}e_k) = e_2 \cdots e_{k-1}e_k$. We define $G^{[1]} = G$.

It is a standard result ([18], Chapter 2) that $G \leq_S G^{[k]}$ for any strongly connected graph G and any $k \geq 1$. In our terminology, Adler-Goodwyn-Weiss showed the following:

Lemma 4.4 ([1], Lemma 4). *Let G be a strongly connected, aperiodic graph of constant out-degree D . Then for all sufficiently large k , we have $M_D \leq_S G^{[k]}$.*

Together with an easy observation about partially ordered sets, the Adler-Goodwyn-Weiss result shows that the $O(G)$ conjecture implies the road colouring theorem.

Lemma 4.5. *Let (\mathcal{P}, \preceq) be a partially ordered set such that, for any $y \in \mathcal{P}$, there exists a unique \preceq -minimal element $O(y) \preceq y$. If $x \preceq y$, then $O(x) = O(y)$.*

Proposition 4.6 ([5]). *Suppose that the $O(G)$ conjecture is true. Let G be a strongly connected, aperiodic graph of constant out-degree D . Then $M_D \leq_S G$.*

Proof. Let k be large enough that $M_D \leq_S G^{[k]}$, by Lemma 4.4. Then in fact $M_D = O(G^{[k]})$. Since $G \leq_S G^{[k]}$ as well, the result follows by Lemma 4.5. \square

5 Bunchiness

In this section, we define and characterize the classes of bunchy and almost bunchy graphs, and demonstrate the importance of bunchy graphs to the structural properties of right-resolvers.

5.1 Bunchy and almost bunchy graphs

We recall from Theorem 2.2, for a graph G , the notation $\Sigma_G : V(G) \rightarrow V(M(G))$ for the unique state map among right-resolvers $G \rightarrow M(G)$.

Definition 5.1 (bunchy states and graphs). Let G be a graph. We say that a state $I \in V(G)$ is *bunchy* if $\Sigma_G|_{F(I)} : F(I) \rightarrow F(\Sigma_G(I)) \subseteq V(M(G))$ is a bijection. We say that G is bunchy if every $I \in V(G)$ is bunchy. We say that G is *almost bunchy* if for each $I, J \in V(M)$, there exists at most one $I' \in \Sigma_G^{-1}(I)$ such that $|F(I') \cap \Sigma_G^{-1}(J)| \geq 2$.

Remark 5.1. The definition of almost bunchiness means that for every ordered pair of Σ_G fibers in G , say the fibers of states $I, J \in V(M)$, there is at most one state in the fiber of I that does not “look bunchy”, in the sense that it has non-parallel outgoing edges into the fiber of J . In other words, an almost bunchy graph almost satisfies the conditions for bunchiness, but an exception is allowed for each ordered pair of fibers.

The following is evident but we state it explicitly for reference:

Lemma 5.1. *The classes of bunchy and almost bunchy graphs are closed under right-resolvers. Moreover, if G is a bunchy graph and C is a principal subgraph of G with $M(C) = M(G)$, then C is also bunchy.*

Remark 5.2. We briefly discuss examples of bunched and almost bunched graphs. The only strongly connected bunched graphs of constant out-degree are the cycles of bunches. For a given \leq_R -minimal cycle of bunches M with sequence of out-degrees D_0, \dots, D_{p-1} , the only strongly connected bunched graphs G with $M(G) = M$ are the graphs $O_{M,q}$ introduced in the previous section.

A strongly connected almost bunched (but not bunched) graph of constant out-degree is a graph with a unique non-bunched state, i.e. a state I with $|F(I)| \geq 2$, together with a path from each element of $F(I)$ back to I . One example that has been considered in the literature is the graph W_n studied in [3], first discussed in [24], and of interest due to its slow synchronization.

An almost bunched graph G can have at most $|V(M(G))|^2$ non-bunched states, one for each ordered pair of Σ_G fibers. One way to obtain an almost bunched graph is to start with a bunched graph and perform a sequence of in-splittings (recall Lemma 3.5, and see also [18], §2.4), but not all in-splittings will preserve almost bunchedness, and not all almost bunched graphs arise this way.

Bunched and almost bunched graphs are characterized in terms of automorphisms, with an important uniqueness consequence for the sets of transition maps induced on them by right-resolvers:

Proposition 5.2. *A graph G is almost bunched if and only if there is a unique right-resolver $G \rightarrow M(G)$ up to permutations of parallel edges: that is, if and only if, for any $\Phi_1, \Phi_2 \in \text{hom}_R(G, M(G))$, there exist $\sigma \in P(G), \tau \in P(M(G))$ such that $\Phi_1 = \tau \circ \Phi_2 \circ \sigma$. Moreover, G is bunched if and only if we can take $\tau = \text{id}$ regardless of Φ_1, Φ_2 .*

Proposition 5.3. *Let G be an almost bunched graph. Let $\Phi_1, \Phi_2 \in \text{hom}_R(G, M(G))$. Then $S_{\Phi_1} = S_{\Phi_2}$. In particular, $\sim_{\Phi_1} = \sim_{\Phi_2}$.*

For the proofs of Propositions 5.2 and 5.3, see §8.3. The following definition is now justified.

Definition 5.2. Let G be an almost bunched graph. We denote by \sim_G the unique relation on $V(G)$ with $\sim_G = \sim_\Phi$ for any $\Phi \in \text{hom}_R(G, M(G))$.

5.2 Proof of the $O(G)$ conjecture in the bunched case

We now resolve the $O(G)$ conjecture in the almost bunched case (which includes the bunched case). This extends Corollary 4.3 in [5], which resolves the conjecture for graphs G such that $M(G)$ has no parallel edges (so G is trivially bunched). The proof here is quite different from the proof in the no-parallel-edges case, and yields a polynomial-time algorithm (Algorithm 7.2) for constructing $O(G)$.

Theorem 5.4. *Let G be an almost bunched graph and let $H \leq_S G$. If H is \leq_S -minimal, then $H = G / \sim_G$. In particular, the set $\{K \mid K \leq_S G\}$ has a unique \leq_S -minimal element $O = G / \sim_G$.*

Proof. Let $\Psi \in \text{hom}_S(G, H)$ and $\Delta \in \text{hom}_R(H, M)$. We can naturally identify $V(H) = V(G) / \sim_\Psi$ by Lemma 2.1. The hypothesis that H is \leq_S -minimal implies that \sim_Δ is trivial, so $\sim_\Psi = \sim_{\Delta \circ \Psi}$ by Theorem 3.3(3). By Proposition 5.3, we have $\sim_{\Delta \circ \Psi} = \sim_G$. We can thus identify $V(H)$ with $V(G) / \sim_G = V(G / \sim_G)$. Thus $H = G / \sim_G$ by a second application of Lemma 2.1. \square

In the strongly connected case, we can apply Proposition 3.8, which is also used in the proof of the road colouring theorem, to say more.

Proposition 5.5. *Let G be a strongly connected almost bunched graph. If G is not bunched, then \sim_G is nontrivial.*

Proof. Let $I, J \in V(M(G))$ and $I' \in \Sigma_G^{-1}(I)$ with $|F(I') \cap \Sigma_G^{-1}(J)| \geq 2$. Let $J_1, J_2 \in F(I') \cap \Sigma_G^{-1}(J)$, $J_1 \neq J_2$, and let $e_i \in E_{I', J_i}(G)$. Let $\Phi \in \text{hom}_R(G, M(G))$ and let $a_i = \Phi(e_i)$. By strong connectedness and Lemma 3.7, there exists a minimal image $U \subseteq \Sigma_G^{-1}(I)$ with $I' \in U$. Let $U_0 = U \setminus \{I'\}$. Then $U_0 \cdot a_1 = U_0 \cdot a_2$ since G is almost bunched. Moreover, $J_i = I' \cdot a_i \notin U_0 \cdot a_i$ (otherwise, minimality would be contradicted). Thus $(U \cdot a_1)\Delta(U \cdot a_2) = \{J_1, J_2\}$. By Proposition 3.8, $J_1 \sim_\Phi J_2$. \square

Corollary 5.6. *Let G be a strongly connected almost bunched graph. Then $O(G)$ is bunched.*

Proof. If $|V(G)| = 1$ then the claim is clearly true. Suppose that the conclusion is true for all almost bunched G with $|V(G)| \leq_R N$, and let G be almost bunched with $|V(G)| = N + 1$. If G is bunched, then $O(G)$ is clearly bunched by Lemma 5.1. If G is not bunched, then \sim_G is nontrivial by Proposition 5.5, so $|V(G/\sim_G)| \leq_R N$. Moreover, since $G/\sim_G \leq_S G$, it follows that $O(G) = O(G/\sim_G)$ is bunched by the inductive hypothesis and Lemma 5.1. \square

5.3 Universal property of the fiber product

We recall a standard construction known as the fiber product, and derive several new properties. Chief among these is the one exhibited in Theorem 5.7, which is an analogue of the universal property often enjoyed by the fiber product, or pullback, in other categories (see e.g. [21], Definition 3.1.15 and subsequent discussion).

Definition 5.3 (fiber product). Let H_1, H_2, K be graphs and let $\Psi_i : H_i \rightarrow K$ be graph homomorphisms. The *fiber product* of Ψ_1, Ψ_2 is the graph $P = H_1 \times_{\Psi_1, \Psi_2} H_2$ where

$$V(P) = \bigsqcup_{I \in V(K)} (\partial\Psi_1)^{-1}(I) \times (\partial\Psi_2)^{-1}(I)$$

$$E(P) = \{(e_1, e_2) \mid e_i \in E(H_i), \Psi_1(e_1) = \Psi_2(e_2)\}$$

together with the coordinate projections $\hat{\Psi}_i : P \rightarrow H_i$. We write $\Psi_P = \Psi_i \circ \hat{\Psi}_i : P \rightarrow K$.

Remark 5.3. To see that $\Psi_1 \circ \hat{\Psi}_1 = \Psi_2 \circ \hat{\Psi}_2$ and thus that Ψ_P is well-defined, note that for every $(I_1, I_2) \in V(P)$ and every $(e_1, e_2) \in E_{(I_1, I_2)}(P)$, we have, by the definition of P ,

$$\Psi_1 \circ \hat{\Psi}_1(e_1, e_2) = \Psi_1(e_1) = \Psi_2(e_2) = \Psi_2 \circ \hat{\Psi}_2(e_1, e_2)$$

Remark 5.4. Observe that the $\hat{\Psi}_i$ are surjective (respectively, right-resolving) when the Ψ_i are surjective (respectively, right-resolving). Moreover, if C is a principal subgraph of P such that the restricted state maps $\partial\hat{\Psi}_i|_{V(C)} : V(C) \rightarrow V(H_i)$ are surjective, then $H_i \leq_R C$, indeed $\hat{\Psi}_i|_C \in \text{hom}_R(C, H_i)$. In particular, this condition is satisfied if the H_i are strongly connected and C is a principal component of P .

Remark 5.5. Often the convention is taken that $V(P) = V(H_1) \times V(H_2)$. However, all of the elements of the full Cartesian product that are not elements of $V(P)$, as we have defined it, would be isolated states, and in particular would be sinks. Our definition has the feature that the fiber product of two sink-free graphs (or rather, of two right-resolvers defined on such graphs) is also sink-free.

We now state the universal property of the fiber product. Compare with a similar diagram in [5] (p. 289). See §8.4 for the proof.

Theorem 5.7. *Let H_1, H_2 be bunched graphs with $M(H_1) = M(H_2) = M$. Let $\Psi_i \in \text{hom}_R(H_i, M)$ be right-resolvers, and let $P = H_1 \times_{\Psi_1, \Psi_2} H_2$. Let G be a common right-resolving extension of H_1, H_2 via $\Phi_i \in \text{hom}_R(G, H_i)$. Then there exist a principal subgraph C of P and right-resolvers $\Delta_i \in \text{hom}_R(G, C)$ such that $\Phi_i = \hat{\Psi}_i \circ \Delta_i$ and $\partial\Delta_1 = \partial\Delta_2$. In particular, $H_i \leq_R C \leq_R G$, with $\hat{\Psi}_i|_C \in \text{hom}_R(C, H_i)$.*

Remark 5.6. The bunchedness hypothesis on the H_i cannot be dropped, as the following construction illustrates. Let G be a graph and let $\Phi_1, \Phi_2 \in \text{Aut}(G)$. In the notation of the theorem, we will take $H_1 = H_2 = G$. (Recall that any automorphism is right-resolving.) Let $M = M(G)$. Let $\Psi_i \in \text{hom}_R(G, M)$ and $P = G \times_{\Psi_1, \Psi_2} G$. Let C be a principal subgraph of P with $\hat{\Psi}_i|_{V(C)}$ surjective, and let $\Delta_i \in \text{hom}_R(G, C)$ with $\Phi_i = \hat{\Psi}_i|_C \circ \Delta_i$. Then $\hat{\Psi}_i|_C$ and Δ_i are isomorphisms, since they compose to an isomorphism. In particular, since $\Psi_1 \circ \hat{\Psi}_1|_C = \Psi_2 \circ \hat{\Psi}_2|_C$, we have $\Psi_1 = \Psi_2 \circ \tau$ where $\tau = \hat{\Psi}_2|_C \circ (\hat{\Psi}_1|_C)^{-1}$ is an isomorphism. In other words, any two elements of $\text{hom}_R(G, M)$ agree up to an automorphism of G . That condition always holds when G is bunched (see Proposition 5.2), but fails in general.

We now give two applications of the universal property. The first, Proposition 5.9, is applied in Proposition 6.2, which is the main motivation for the bunchy factor conjecture. See §8.5 for the proof of Lemma 5.8.

Lemma 5.8. *Let H_1, H_2 be bunchy graphs with $M(H_1) = M(H_2) = M$. Let $\Psi_i \in \text{hom}_R(H_i, M)$, and let $P = H_1 \times_{\Psi_1, \Psi_2} H_2$. Then P is bunchy. In particular, if C is a principal subgraph of P such that the restrictions $\partial \hat{\Psi}_i|_{V(C)} : V(C) \rightarrow V(H_i)$ are surjective, then C is bunchy.*

Proposition 5.9. *Let G be a graph. Let $H_1, H_2 \leq_S G$ be bunchy. Then $O(H_1) = O(H_2)$, i.e. G has at most one \leq_S -minimal bunchy synchronizing factor.*

Proof. Let $M = M(G)$, let $\Phi_i \in \text{hom}_S(G, H_i)$, and let $\Psi_i \in \text{hom}_R(H_i, M)$. Let $P = H_1 \times_{\Psi_1, \Psi_2} H_2$. By Theorem 5.7, there is a principal subgraph C of P admitting $\Delta_i \in \text{hom}_R(G, C)$ such that $\Phi_i = \hat{\Psi}_i|_{C \circ \Delta_i}$. Since each Φ_i is synchronizing, each restriction $\hat{\Psi}_i|_C$ is synchronizing as well, so $H_i \leq_S C$. Since C is bunchy by Lemma 5.8, we know that $O(C)$ is well-defined, and thus, by Lemma 4.5, we have $O(H_1) = O(C) = O(H_2)$ as claimed. \square

For the second application of the universal property, recall that the only strongly connected bunchy graphs of constant out-degree are the cycles of bunches. In particular, by the periodic road colouring theorem, for any strongly connected graph G of constant out-degree D and period p , the unique maximal bunchy right-resolving factor of G , namely $O_{D,p}$, is a synchronizing factor of G (and is indeed equal to $O(G)$). We now show that every graph G has a unique maximal bunchy right-resolving factor $B(G)$. The construction is similar to that of the auxiliary graph \tilde{G} in [5], §5. See §8.5 for the proof, as well as an explicit construction of $B(G)$ yielding a polynomial-time algorithm (Algorithm 7.3).

Proposition 5.10. *Let G be a graph.*

- (1) *The set $\{H \leq_R G \mid H \text{ is bunchy}\}$ has a unique \leq_R -maximal element $B = B(G)$.*
- (2) *Let $H \leq_R G$ be bunchy and $\Phi \in \text{hom}_R(G, H)$. Then Φ factors through B , i.e. there exist $\Delta \in \text{hom}_R(G, B)$, $\Theta \in \text{hom}_R(B, H)$ such that $\Phi = \Theta \circ \Delta$.*

6 The $O(G)$ conjecture and bunchy synchronizing factors

As we have seen, the $O(G)$ conjecture holds for strongly connected graphs G such that $M(G)$ is a cycle of bunches, and for almost bunchy graphs (including the bunchy graphs) by Theorem 5.4. Moreover, for strongly connected almost bunchy graphs, and strongly connected graphs that factor onto cycles of bunches, we know that there is a bunchy synchronizing factor, which we show inductively by assuming non-bunchiness and obtaining a right-resolver with a nontrivial stability relation. It seems plausible that, if the $O(G)$ conjecture is true, then it can be proven by a similar approach: assume non-bunchiness, find a right-resolver with nontrivial stability relation, recursively find a bunchy synchronizing factor, and apply Proposition 5.9. The next proposition gives several equivalent formulations of the hypothesis that this approach can be made to work. See §8.6 for the proof.

Proposition 6.1. *The following statements are equivalent.*

- (1) *Any strongly connected \leq_S -minimal graph is bunchy.*
- (2) *For any strongly connected graph G , there exists some bunchy $H \leq_S G$.*
- (3) *For any non-bunchy strongly connected graph G , there exists some $\Phi \in \text{hom}_R(G, M(G))$ with \sim_Φ nontrivial.*
- (4) *For any strongly connected graph G , $B(G) \leq_S G$.*

Conjecture (bunchy factor conjecture). *The assertions in Proposition 6.1 are true.*

Proposition 6.2. *The bunchy factor conjecture implies the $O(G)$ conjecture.*

Proof. Let G be a strongly connected graph and let $A = \{H \mid H \leq_S G, H \text{ is } \leq_S\text{-minimal}\}$. Clearly $|A| \geq 1$. By hypothesis, every element of A is bunchy. By Proposition 5.9, $|A| \leq 1$, so A has a single element, namely $O(G)$. \square

Observe that the bunchy factor conjecture is a straightforward generalization of the road problem. As discussed above, the $O(G)$ conjecture was already known to imply the road colouring theorem, via the higher-edge result from [1]. By contrast, the bunchy factor conjecture implies the road colouring theorem more directly, without reference to [1].

7 Computing with right-resolvers

We now discuss the computational problems of constructing $O(G_1), O(G_2)$, for input graphs G_1, G_2 such that the $O(G_i)$ are known to exist, and deciding whether the $O(G_i)$ are isomorphic. Although one could apply generic graph isomorphism algorithms, which are efficient in practice (Theorem 1 in [19] gives a polynomial-time reduction from directed to undirected graph isomorphism, and see [6] for a survey of the state of the art), it is desirable to have a polynomial-time algorithm, in particular one that only uses constructions involved in the theory of right-resolvers and synchronization. We do not attempt detailed complexity analyses, noting only that all of the procedures we give can be easily seen to run in polynomial time.

7.1 Basic routines

In [5], a polynomial-time algorithm is given for computing $M(G)$, along with Σ_G and a total ordering of $V(M(G))$ such that any graph isomorphism $M(G) \rightarrow M(H)$ must preserve the order of the states. Deciding whether $M(G), M(H)$ are isomorphic is therefore no harder than constructing them. Moreover, we can use Σ_G to construct right-resolvers, as follows.

Algorithm 7.1. Construct a right-resolver from a graph to its minimal right-resolving factor.

1. Input: a graph G .
2. Construct $M(G)$ and Σ_G .
3. For each $I, J \in V(M(G))$:
 1. Choose a total ordering on $E_{IJ}(M)$.
 2. For each $I' \in \Sigma_G^{-1}(I)$:
 1. Choose a total ordering on $\bigcup_{J' \in \Sigma_G^{-1}(J)} E_{I'J'}(G)$, i.e. the edges $e \in E_{I'}(G)$ with $\Sigma_G(t(e)) = J$.
 2. For each $e \in E_{I'}(G)$ with $\Sigma_G(t(e)) = J$, record as $\Phi(e)$ the edge in $E_{IJ}(M)$ with the same position in the total ordering of $E_{IJ}(M)$ as e has in $\bigcup_{J' \in \Sigma_G^{-1}(J)} E_{I'J'}(G)$.
4. Return: Φ .

There are also obvious polynomial-time procedures for constructing the fiber product of two right-resolvers, and for determining whether there is a path from one given state to another (in a graph that may not be strongly connected). With these basic routines, we can construct the stability relation of a right-resolver in polynomial time, as follows.

Algorithm 7.2. Construct the stability relation of a right-resolver.

1. Input: graphs G, H , and $\Phi \in \text{hom}_R(G, H)$.

2. Construct $P = G \times_{\Phi, \Phi} G$.
3. Populate the set U of states $(I_1, I_2) \in V(P)$ with no outgoing path to the diagonal in $V(P)$.
4. Populate and output: the set \sim_{Φ} of states $(I_1, I_2) \in V(P)$ with no outgoing path to U .

Recall that, by definition, $\Phi \in \text{hom}_S(G, H)$ if and only if $H = G / \sim_{\Phi}$, i.e. the $\partial\Phi$ -fibers are precisely the \sim_{Φ} classes. Since this is easy to check, Algorithm 7.2 can be used to decide whether Φ is synchronizing.

A similar procedure can be used to construct the maximum bunchy factor $B(G)$, following the construction in Proposition 8.7:

Algorithm 7.3. Construct the maximum bunchy right-resolving factor of a graph.

1. Input: a graph G .
2. Construct $M(G)$, along with the quotient map $\Sigma_G : V(G) \rightarrow V(M)$.
3. Construct a graph H with the following data:

$$V(H) = V(G) \times V(G)$$

$$|E_{(I_1, I_2)(J_1, J_2)}(H)| = \begin{cases} 1, & \Sigma_G(I_1) = \Sigma_G(I_2), \Sigma_G(J_1) = \Sigma_G(J_2), \text{ and } J_i \in F(I_i) \\ 0, & \text{otherwise} \end{cases}$$

4. Populate the set \approx_0 of pairs (I_1, I_2) with a path in H from the diagonal to (I_1, I_2) .
5. Construct and output: the transitive closure \approx of the relation \approx_0 .

Regarding step 4: referring to the description of \approx_0 in Proposition 8.7, a path in H from the diagonal to (I_1, I_2) corresponds to a pair of paths $\gamma, \delta \in L(G)$ witnessing $I_1 \approx_0 I_2$.

7.2 Decision procedures for common synchronizing factors and extensions

See §8.6 for the proof of the following Proposition, which collects several similar statements relating common synchronizing factors and common synchronizing extensions. These results allow us to use fiber products of two graphs to decide questions about common factors of the graphs.

Proposition 7.1. *Let G_1, G_2 be strongly connected graphs with $M(G_1) = M(G_2) = M$.*

1. *If G_1, G_2 have a common synchronizing factor, then there exist $\Phi_i \in \text{hom}_R(G_i, M)$ and a principal component C of $Q = G_1 \times_{\Phi_1, \Phi_2} G_2$ such that $\hat{\Phi}_i|_C \in \text{hom}_S(C, G_i)$.*
2. *Assume the $O(G)$ conjecture. Then the converse holds in (1). That is, suppose that there exist $\Phi_i \in \text{hom}_R(G_i, M)$ and a principal component C of $Q = G_1 \times_{\Phi_1, \Phi_2} G_2$, such that $\hat{\Phi}_i|_C \in \text{hom}_S(C, G_i)$. Then G_1, G_2 have a common synchronizing factor, specifically $O(G_1) = O(C) = O(G_2)$.*
3. *If the G_i are bunchy, then the equivalence described in (1) and (2) holds unconditionally (i.e. without assuming any unproven conjectures).*
4. *Assume the bunchy factor conjecture. Then we have $O(G_1) = O(G_2)$ if and only if, for the essentially unique $\Phi_i \in \text{hom}_R(B(G_i), M)$, there is a principal component C of $P = B(G_1) \times_{\Phi_1, \Phi_2} B(G_2)$ such that $\hat{\Phi}_i|_C \in \text{hom}_S(C, B(G_i))$.*

Ashley-Marcus-Tuncel give a polynomial-time algorithm for deciding whether two strongly connected graphs have a common strongly connected synchronizing extension. Their algorithm relies on the construction of a graph that they call \tilde{G} , from an input graph G ; see Theorem 5.2 and Remark 5.10 in [5]. If the $O(G)$ conjecture is true, then by Proposition 7.1(2), the Ashley-Marcus-Tuncel algorithm also decides whether $O(G_1) = O(G_2)$ for G_1, G_2 strongly connected. Without assuming any unproven conjectures, a negative result from the algorithm shows, by Proposition 7.1(1), that G_1, G_2 have no common synchronizing factor, while a positive result is inconclusive.

For bunchy G_1, G_2 , however, Theorem 5.4 and Proposition 7.1(3) show that it can be decided in polynomial time whether $O(G_1), O(G_2)$ are isomorphic without the Ashley-Marcus-Tuncel algorithm:

Algorithm 7.4. Decide whether $O(G_1), O(G_2)$ are isomorphic for G_1, G_2 strongly connected and bunchy.

1. Input: strongly connected bunchy graphs G_1, G_2 such that $M(G_1) = M(G_2) = M$.
2. Choose $\Phi_i \in \text{hom}_R(G_i, M)$, using Algorithm 7.1.
3. Construct $Q = G_1 \times_{\Phi_1, \Phi_2} G_2$.
4. For each principal component C , decide whether $\hat{\Phi}_i|_C \in \text{hom}_S(C, G_i)$, using Algorithm 7.2. If so, halt, and return the result that $O(G_1) = O(G_2)$.
5. If no affirmative result is returned in step 4, then halt and return the result that $O(G_1) \neq O(G_2)$.

Regarding steps 4–5: Proposition 7.1(3) shows that we have $O(G_1) = O(G_2)$ if and only if $\hat{\Phi}_i|_C \in \text{hom}_S(C, G_i)$ for some principal component C of Q . The same algorithm would work even if G_1, G_2 are not strongly connected but only bunchy, with “principal component C ” replaced by “principal subgraph C such that each $\partial\hat{\Phi}_i|_{V(C)}$ is surjective”, but it is not clear that the number of such subgraphs is bounded by a polynomial in $|V(G)|$.

Furthermore, if the bunchy factor conjecture is true, then isomorphism of $O(G_1), O(G_2)$ is equivalent, by Proposition 6.1, to isomorphism of $O(B(G_1)), O(B(G_2))$, which can be decided without the use of the Ashley-Marcus-Tuncel \tilde{G} algorithm, using the following procedure. Note that this works even if we do not have an efficient way to find an element of $\text{hom}_S(G_i, B(G_i))$.

Algorithm 7.5. Decide isomorphism of $O(G_1), O(G_2)$ (assuming the bunchy factor conjecture).

1. Input: strongly connected graphs G_1, G_2 .
2. Construct $B(G_1)$ and $B(G_2)$, using Algorithm 7.3.
3. Decide whether $O(B(G_1)) = O(B(G_2))$, using Algorithm 7.4.
4. Output: the Boolean value of “ $O(B(G_1)) = O(B(G_2))$ ”.

Observe that the procedure of Algorithm 7.5 is essentially what is described in Proposition 7.1(4).

8 Proofs of structural results and additional details

8.1 Remarks on the proof of Theorem 2.2

In this subsection, we revisit the first proof of Theorem 2.2 given in [5]. The proof of the uniqueness of $M(G)$, for a given graph G , is without issue, but the proof of the uniqueness of Σ_G is not quite complete. That proof seems to assume that, for graphs G, H with $H \leq_R G$, two right-resolvers $G \rightarrow H$ with distinct state maps must partition $V(G)$ differently. In general, this is false: per Lemma 8.3 below, counterexamples arise precisely when $\text{Aut}(H)$ acts nontrivially on $V(H)$.

The second proof given in [5], constructing $V(M(G))$ by successive refinements of an initial state partition, can be made complete by observing that all of the state maps corresponding to the successive refinements are invariant under $\text{Aut}(G)$.

Proposition 8.1 (Lemma 3.1 in [5]). *Let G, H_1, H_2 be graphs with $H_i \leq_R G$ via $\Psi_i \in \text{hom}_R(G, H_i)$, with partitions $\alpha_i = V(H_i)$ of $V(G)$. Then there is a graph $K \leq_R H_1, H_2$, with $V(K)$ equal to the finest common coarsening of α_1, α_2 .*

Corollary 8.2 (Theorem 3.2 in [5]). *Let G be a graph. Then there exists a unique \leq_R -minimal graph $M(G) \leq_R G$, with $V(M(G))$ given by the finest common coarsening of the partitions $\alpha = V(H)$ of $V(G)$, where $H \leq_R G$.*

We will show further that for any $\Phi_1, \Phi_2 \in \text{hom}_R(G, M(G))$, we have $\partial\Phi_1 = \partial\Phi_2$. Note that $\partial\Phi_1, \partial\Phi_2$ at least have the same sets of fibers, since, if they did not, then by Proposition 8.1, we could take their finest common coarsening, contradicting the minimality of $M(G)$. It is therefore enough to show that $\text{Aut}(M(G))$ acts trivially on $M(G)$. For this, we need a lemma.

Lemma 8.3. *Let G, H be graphs with $H \leq_R G$, and let $\Phi_1, \Phi_2 \in \text{hom}_R(G, H)$. Suppose that the $\partial\Phi_i$ have the same fibers, i.e. for any $I_1, I_2 \in V(G)$, we have $\partial\Phi_1(I_1) = \partial\Phi_1(I_2)$ if and only if $\partial\Phi_2(I_1) = \partial\Phi_2(I_2)$. Then there exists an automorphism $\tau \in \text{Aut}(H)$ such that $\partial\Phi_2 = \partial(\tau \circ \Phi_1)$.*

Proof. By the assumption of equal fibers, there exists a (unique) bijection $T : V(H) \rightarrow V(H)$ with $\partial\Phi_2 = T \circ \partial\Phi_1$. We need to find $\tau \in \text{Aut}(H)$ with $\partial\tau = T$. Let $I \in V(H)$. For $J \in F(I)$, note, by the right-resolving hypothesis on the Φ_i and the choice of T , that $|E_{IJ}(H)| = |E_{T(I)T(J)}(H)|$. Letting $\tau|_{E_{IJ}(H)} : E_{IJ}(H) \rightarrow E_{T(I)T(J)}(H)$ be any bijection, we have $\tau \in \text{Aut}(H)$ with $T = \partial\tau$ and $\partial\Phi_2 = \partial(\tau \circ \Phi_1)$. \square

We now discuss the quotient of a graph by its automorphism group. Let G be a graph. We will construct a graph $K = G/\text{Aut}(G) \leq_R G$ as follows. Let $V(K)$ consist of the orbits in $V(G)$ under $\text{Aut}(G)$. Let $I, J \in V(K)$ and $I_1, I_2 \in I$. We need to specify $|E_{IJ}(K)|$. We claim that $\sum_{J' \in J} |E_{I_1 J'}(G)| = \sum_{J' \in J} |E_{I_2 J'}(G)|$. Indeed, let $\tau \in \text{Aut}(G)$ such that $I_2 = \partial\tau(I_1)$. Then $\partial\tau(F(I_1)) = \partial\tau(F(I_2))$, and $\tau(J) = J$, so for any $J' \in J \cap F(I_1)$, we have $\partial\tau(J') \in J \cap F(I_2)$. Therefore $\sum_{J' \in J} |E_{I_1 J'}(G)| \leq_R \sum_{J' \in J} |E_{I_2 J'}(G)|$. Replacing τ with τ^{-1} , we obtain equality. Define K by specifying that $|E_{IJ}(K)| = \sum_{J' \in J} |E_{I_1 J'}(G)|$. This edge count ensures that $K \leq_R G$.

In particular, if $\text{Aut}(G)$ acts nontrivially on $V(G)$, then $K \neq G$, so G is not \leq_R -minimal. It follows that, for a \leq_R -minimal graph M , $\text{Aut}(M)$ acts trivially on $V(M)$. This observation, together with Lemma 8.3, shows that for any $\Phi_1, \Phi_2 \in \text{hom}_R(G, M(G))$, we have $\partial\Phi_1 = \partial\Phi_2$.

8.2 Proof of Theorem 3.3

We prove Theorem 3.3 in several stages. First, in Lemma 8.4, we determine how stability classes intersect with fibers in compositions of right-resolvers.

Lemma 8.4. *Let G, K, H be graphs with $H \leq_R K \leq_R G$. Let $\Psi \in \text{hom}_R(G, K)$, $\Delta \in \text{hom}_R(K, H)$, and let $\Phi = \Delta \circ \Psi$. Then \sim_Ψ -classes are intersections of \sim_Φ -classes with $\partial\Psi$ -fibers. That is, for $I_1, I_2 \in V(G)$, if $I_1 \sim_\Psi I_2$, then $I_1 \sim_\Phi I_2$; conversely, if $I_1 \sim_\Phi I_2$ and moreover $\partial\Psi(I_1) = \partial\Psi(I_2)$, then $I_1 \sim_\Psi I_2$.*

Proof. First suppose that $I_1 \sim_\Psi I_2$. Then in particular $\partial\Psi(I_1) = \partial\Psi(I_2) = \hat{I}$. Let $I = \partial\Delta(\hat{I})$ and $u \in L_I(H)$, and consider the unique $\lambda \in L_{\hat{I}}(K)$ such that $\Delta(\lambda) = u$. Since $I_1 \sim_\Psi I_2$, there exists $\mu \in L_{t(\lambda)}(K)$ such that $I_1 \cdot \lambda\mu = I_2 \cdot \lambda\mu$. Letting $v = \Delta(\mu)$, we have $I_1 \cdot uv = I_2 \cdot uv$, so $I_1 \sim_\Phi I_2$.

For the converse, suppose that $I_1 \sim_\Phi I_2$ and $\partial\Psi(I_1) = \partial\Psi(I_2) = \hat{I}$. Let $I = \Delta(\hat{I})$, let $\lambda \in L_I(H)$, let $u = \Delta(\lambda)$, and let $v \in L_{t(u)}(H)$ be such that $I_1 \cdot uv = I_2 \cdot uv$. Consider the unique $\mu \in L_{t(\lambda)}(K)$ such that $\Delta(\mu) = v$. Then $I_1 \cdot \lambda\mu = I_2 \cdot \lambda\mu$, so indeed $I_1 \sim_\Psi I_2$. \square

Corollary 8.5 follows immediately from Lemma 8.4, and together they comprise Theorem 3.3(1).

Corollary 8.5. *If $\Phi = \Delta \circ \Psi$ is a composition of right-resolvers, then Ψ is synchronizing if and only if every $\partial\Psi$ -fiber is contained in a \sim_Φ -class.*

Proof of Theorem 3.3(2). By assumption, the \sim_Φ classes are precisely the $\partial\Psi$ -fibers, so by Lemma 8.4, they are also the \sim_Ψ classes. Thus indeed $\Psi \in \text{hom}_S(G, K)$.

Moreover, let $I \in V(H)$, let $I'_1, I'_2 \in (\partial\Delta)^{-1}(I)$, and suppose that $I'_1 \sim_\Delta I'_2$. We claim that $I'_1 = I'_2$. Toward this end, we claim that $(\partial\Psi)^{-1}(\{I'_1, I'_2\})$ is a subset of a \sim_Φ class. Indeed, let $u \in L_I(H)$, and let $I_i \in (\partial\Psi)^{-1}(I'_i)$. Let $v \in L_{t(u)}(H)$ be such that $I'_1 \cdot uv = I'_2 \cdot uv = J'$. Let $J_i = I_i \cdot uv$. Then $J_1, J_2 \in (\partial\Psi)^{-1}(J')$. Since Ψ is synchronizing, there exists $\gamma \in L_{J'}(K)$ with $J_1 \cdot \gamma = J_2 \cdot \gamma$. Let $w = \Delta(\gamma)$. Then $I_1 \cdot uvw = I_2 \cdot uvw$. Since $u \in L_I(H)$ was arbitrary, it follows that $I_1 \sim_\Phi I_2$ as claimed. Since, by assumption, a \sim_Φ class is precisely a $\partial\Psi$ -fiber of a single state of K , we must have $I'_1 = I'_2$ as claimed. Therefore \sim_Δ is indeed trivial. \square

Lemma 8.6. *Let G, K, H be graphs with $H \leq_R K \leq_R G$. Let $\Psi \in \text{hom}_R(G, K)$, $\Delta \in \text{hom}_R(K, H)$, and let $\Phi = \Delta \circ \Psi$. Let $I \in V(H)$, let $I'_1, I'_2 \in (\partial\Delta)^{-1}(I)$, and let $I_i \in (\partial\Psi)^{-1}(I'_i)$. If $I_1 \sim_\Phi I_2$, then $I'_1 \sim_\Delta I'_2$.*

Proof. Suppose that $I_1 \sim_\Phi I_2$. Let $u \in L_I(H)$. Let $v \in L_{t(u)}(H)$ be such that $I_1 \cdot uv = I_2 \cdot uv$. Then, since $I'_i \cdot uv = \partial\Phi(I_i \cdot uv)$, we have $I'_1 \cdot uv = I'_2 \cdot uv$. Therefore $I'_1 \sim_\Delta I'_2$. \square

Proof of Theorem 3.3(3). Suppose that \sim_Δ is trivial. Let $I \in V(H)$, let $I'_1, I'_2 \in (\partial\Delta)^{-1}(I)$ with $I'_1 \neq I'_2$, and let $I_i \in (\partial\Psi)^{-1}(I'_i)$. Since \sim_Δ is trivial, there exists $u \in L_I(H)$ such that, for every $v \in L_{t(u)}(H)$, we have $I'_1 \cdot uv \neq I'_2 \cdot uv$. This implies that $I_1 \cdot uv \neq I_2 \cdot uv$, so $I_1 \not\sim_\Phi I_2$. It follows that each \sim_Φ class is contained inside a $\partial\Psi$ fiber. By Lemma 8.4, this shows that $\sim_\Phi = \sim_\Psi$. \square

The final part of Theorem 3.3 can be proved in the strongly connected case using symbolic dynamics, via the multiplicativity of degree under composition of right-resolvers; see [18], §9.1. Using the theory developed so far, we give a self-contained proof without the assumption of strong connectedness.

Proof of Theorem 3.3(4). Suppose that Ψ, Δ are synchronizing. Let $I \in V(H)$. Let $I_1, I_2 \in (\partial\Phi)^{-1}(I)$. We need to show that $I_1 \sim_\Phi I_2$. Let $u \in L_I(K)$. We need to find $v \in L_{t(u)}(H)$ such that $I_1 \cdot uv = I_2 \cdot uv$. Let $I'_i = \partial\Psi(I_i)$. Then $\partial\Delta(I'_1) = \partial\Delta(I'_2)$, so $I'_1 \sim_\Delta I'_2$ since Δ is synchronizing. Let $v_1 \in L_{t(u)}(H)$ be such that $I'_1 \cdot uv_1 = I'_2 \cdot uv_1$. Note that $I'_i \cdot uv_1 = \partial\Psi(I_i \cdot uv_1)$. Let $J' = I'_i \cdot uv_1$ and $J_i = I_i \cdot uv_1$. Then $\partial\Psi(J_1) = J' = \partial\Psi(J_2)$, so $J_1 \sim_\Psi J_2$ since Ψ is synchronizing. Let $\gamma \in L_{J'}(K)$ be such that $J_1 \cdot \gamma = J_2 \cdot \gamma$. Let $v_2 = \Delta(\gamma)$. Then $I_i \cdot uv_1 v_2 = J_i \cdot \gamma$, so taking $v = v_1 v_2$, we have $I_1 \cdot uv = I_2 \cdot uv$. Thus $I_1 \sim_\Phi I_2$, so Φ is indeed synchronizing.

We prove the converse by the contrapositive. Suppose that Δ is not synchronizing. Then there exist $I \in V(H)$ and $I'_1, I'_2 \in (\partial\Delta)^{-1}(I)$ such that $I'_1 \not\sim_\Delta I'_2$. By Lemma 8.6, there exist $I_i \in (\partial\Psi)^{-1}(I'_i)$ such that $I_1 \not\sim_\Phi I_2$. Thus Φ is not synchronizing. Similarly, suppose that Ψ is not synchronizing. Then there exist $I' \in V(K)$ and $I_1, I_2 \in (\partial\Psi)^{-1}(I')$ such that $I_1 \not\sim_\Psi I_2$. By Lemma 8.4, $I_1 \not\sim_\Phi I_2$, so Φ is not synchronizing. \square

8.3 Proofs of Propositions 5.2 and 5.3

Proof of Proposition 5.2. Let $M = M(G)$. First, suppose that G is almost bunchy. Let $\Phi_1, \Phi_2 \in \text{hom}_R(G, M)$. For each $I, J \in V(M)$, and each $I' \in \Sigma_G^{-1}(I)$, let $A_{I', J} = \{e \in E_{I'}(G) \mid \Sigma_G(t(e)) = J\}$. If there exists $I' \in \Sigma_G^{-1}(I)$ with $|F(I') \cap \Sigma_G^{-1}(J)| \geq 2$ (by almost bunchiness, there is at most one such I' for any given I, J), then let $\tau_{IJ} \in P(M)$ be the permutation of parallel edges in M given by

$$\begin{aligned} \tau_{IJ}|_{E(M) \setminus E_{IJ}(M)} &= \text{id}|_{E(M) \setminus E_{IJ}(M)} \\ \tau_{IJ}|_{E_{IJ}(M)} &= \Phi_1|_{A_{I', J}} \circ (\Phi_2|_{A_{I', J}})^{-1} \end{aligned}$$

If there is no $I' \in \Sigma_G^{-1}(I)$ with $|F(I') \cap \Sigma_G^{-1}(J)| \geq 2$, then let $\tau_{IJ} = \text{id}$. Distinct τ_{IJ} are permutations of disjoint sets and therefore commute. Let $\tau = \prod_{I, J \in V(M)} \tau_{IJ}$. Note that if G is bunchy, then $\tau_{IJ} = \text{id}$ for all I, J , so $\tau = \text{id}$.

Now, for each $I, J \in V(M)$ and each $I' \in \Sigma_G^{-1}(I)$, let $\sigma_{IJ, I'} \in P(G)$ be given by

$$\begin{aligned}\sigma_{IJ, I'}|_{E(G) \setminus A_{I', J}} &= \text{id}|_{E(G) \setminus A_{I', J}} \\ \sigma_{IJ, I'}|_{A_{I', J}} &= \Phi_2|_{A_{I', J}}^{-1} \circ \tau^{-1} \circ \Phi_1|_{A_{I', J}}\end{aligned}$$

All of the $\sigma_{IJ, I'}$ commute. Let $\sigma = \prod_{I, J \in V(M), I' \in \Sigma_G^{-1}(I)} \sigma_{IJ, I'}$. Then for $I \in V(M)$, $J \in F(I)$, and $I' \in \Sigma_G^{-1}(I)$,

$$\tau \circ \Phi_2 \circ \sigma|_{A_{I', J}} = \tau \circ \Phi_2 \circ (\Phi_2|_{A_{I', J}}^{-1} \circ \tau^{-1} \circ \Phi_1|_{A_{I', J}}) = \Phi_1|_{A_{I', J}}$$

This concludes the proof in the ‘‘only if’’ direction.

For the ‘‘if’’ direction, which we prove in the contrapositive, suppose that G is not almost bunchy. Let $I, J \in V(M)$ and $I_1, I_2 \in \Sigma_G^{-1}(I)$ such that $|F(I_i) \cap \Sigma_G^{-1}(J)| \geq 2$ for $i = 1, 2$. Let $e_{i,1}, e_{i,2} \in A_{I_i, J}$ be such that $t(e_{i,1}) \neq t(e_{i,2})$. Let $a_1, a_2 \in E_{IJ}(M)$, and let $\Phi_1, \Phi_2 \in \text{hom}_R(G, M)$ be such that $\Phi_1(e_{1,j}) = a_j$ but $\Phi_2(e_{2,1}) = a_2$ and $\Phi_2(e_{2,2}) = a_1$. (The behaviour of Φ_i on $E(G) \setminus \{e_{i,j}\}_{i,j=1,2}$ is irrelevant.) Then there do not exist $\sigma \in P(G)$, $\tau \in P(M)$ such that $\Phi_1 = \tau \circ \Phi_2 \circ \sigma$.

Finally, suppose that G is almost bunchy, but not bunchy. Let $I, J \in V(M)$ and $I' \in \Sigma_G^{-1}(I)$ such that $|F(I') \cap \Sigma_G^{-1}(J)| \geq 2$. Let $e_1, e_2 \in A_{I', J}$ be such that $t(e_1) \neq t(e_2)$. Let $a_1, a_2 \in E_{IJ}(M)$, and let $\Phi_1, \Phi_2 \in \text{hom}_R(G, M)$ be such that $\Phi_1(e_i) = a_i$, $\Phi_2(e_1) = a_2$, and $\Phi_2(e_2) = a_1$. Then there does not exist $\sigma \in P(G)$ such that $\Phi_1 = \Phi_2 \circ \sigma$. \square

Proof of Proposition 5.3. Let $M = M(G)$. It is enough to show that S_{Φ_1}, S_{Φ_2} share a generating set T , which we now construct and examine. For $I, J \in V(M)$, if $J \notin F(I)$, let $T_{I, J, 1} = T_{I, J, 2} = \emptyset$. If $J \in F(I)$, then let $T_{I, J, i}$ be the set of maps $f_{a, i} : \Sigma_G^{-1}(I) \rightarrow \Sigma_G^{-1}(J)$ of the form $I' \mapsto t \circ (\Phi_i|_{E_{I', J}})^{-1}(a)$, i.e. $I' \cdot f_{a, i} = I' \cdot a$ with respect to Φ_i , where $a \in E_{IJ}(M)$. Clearly $T_{I, J, i}$ generates S_{Φ_i} , in the sense that S_{Φ_i} is the smallest collection of maps closed under composition and containing the $T_{I, J, i}$ as I, J range over $V(M)$.

We claim that $T_{I, J, 1} = T_{I, J, 2}$. Indeed, let $a \in E_{IJ}(M)$ and let $I' \in \Sigma_G^{-1}(I)$ such that $|F(I') \cap \Sigma_G^{-1}(J)| \geq 2$. It is enough to show that $f_{a, 1} \in T_{I, J, 2}$, as this will show that $T_{I, J, 1} \subseteq T_{I, J, 2}$, from which equality follows by symmetry. By almost bunchiness, there is at most one state $I' \in \Sigma_G^{-1}(I)$ with $|F(I') \cap \Sigma_G^{-1}(J)| \geq 2$. If there is no such state, then clearly $f_{a, 1} = f_{a, 2}$, so assume that such a state I' exists. Observe that $I' \cdot f_{b, 2} = I' \cdot f_{a, 1}$ where $b = \Phi_2 \circ (\Phi_1|_{E_{I', (G)}})^{-1}(a)$. Moreover, for every $I'' \in \Sigma_G^{-1}(I)$ with $I'' \neq I'$, we also have, by almost bunchiness, that $I'' \cdot f_{b, 2} = I'' \cdot f_{a, 1}$. Therefore $f_{a, 1} = f_{b, 2} \in T_{I, J, 2}$, so indeed $T_{I, J, 1} = T_{I, J, 2}$ as claimed. Let $T_{I, J} = T_{I, J, i}$ and let $T = \bigcup_{I, J} T_{I, J}$. Then T generates both S_{Φ_1}, S_{Φ_2} , so indeed $S_{\Phi_1} = S_{\Phi_2}$. \square

8.4 Proof of Theorem 5.7

Proof of Theorem 5.7. Define $T : V(G) \rightarrow V(P)$ as follows: for $I' \in V(G)$, let $T(I') = (\partial\Phi_1(I'), \partial\Phi_2(I'))$. Note that $\partial\Psi_P \circ T = \partial(\Psi_i \circ \Phi_i) = \Sigma_G$ for each i . Let C be the subgraph of P induced by $T(V(G))$. Let $I' \in V(G)$ and $J' \in F(I')$; $I = \Sigma_G(I')$ and $J = \Sigma_G(J')$; and $I_i = \partial\Phi_i(I')$ and $J_i = \partial\Phi_i(J')$. Observe that $(I_1, I_2) = T(I')$ and $(J_1, J_2) = T(J')$.

As in the proof of Proposition 5.2, let $A_{I', J} = \{e \in E_{I', (G)} \mid \Sigma_G(t(e)) = J\}$. Since H_i is bunchy, we have $F(I_i) \cap \Sigma_{H_i}^{-1}(J) = \{J_i\}$, so $\Phi_i|_{A_{I', J}} = E_{I_i, J_i}(H_i)$. It therefore makes sense to define

$$\Delta_i|_{A_{I', J}} = (\hat{\Psi}_i|_{E_{(I_1, I_2)(J_1, J_2)}(P)})^{-1} \circ \Phi_i|_{A_{I', J}} : A_{I', J} \rightarrow E_{(I_1, I_2)(J_1, J_2)}(P)$$

Gluing these together, we obtain maps $\Delta_i : E(G) \rightarrow E(P)$. For $e \in A_{I', J}$, we have $s(\Delta_i(e)) = T(s(e))$ and $t(\Delta_i(e)) = T(t(e))$, so the $\Delta_i : G \rightarrow P$ are graph homomorphisms with $\partial\Delta_i = T$.

We now claim that $\Delta_i(E(G)) = E(C)$. Enumerate $E_{IJ}(M) = \{a^{(1)}, \dots, a^{(k)}\}$, and let $e_i^{(j)} = (\Psi_i|_{E_{I_i, J_i}(H_i)})^{-1}(a^{(j)})$. Then

$$(e_1^{(j)}, e_2^{(j)}) \in E_{(I_1, I_2)(J_1, J_2)}(P) = E_{(I_1, I_2)(J_1, J_2)}(C)$$

where equality holds because $(I_1, I_2), (J_1, J_2) \in V(C)$ and C is an induced subgraph of P . Fixing (I_1, I_2) and varying J' , thus varying (J_1, J_2) , the sets $E_{(I_1, I_2)(J_1, J_2)}(P)$ exhaust $E_{(I_1, I_2)}(P)$, so $E_{(I_1, I_2)}(C) = E_{(I_1, I_2)}(P)$. Since $\Delta_i|_{E_{I'}(G)} : E_{I'}(G) \rightarrow E_{(I_1, I_2)}(P)$ is surjective, we have $E_{(I_1, I_2)}(C) = \Delta_i(E_{I'}(G))$. Thus indeed $E(C) = \Delta_i(E(G))$.

Since we already know that $E_{(I_1, I_2)}(C) = E_{(I_1, I_2)}(P)$ for each $(I_1, I_2) \in V(C)$, it follows that C is indeed a principal subgraph of P . Finally, we have

$$\hat{\Psi}_i \circ \Delta_i|_{A_{I', J}} = \hat{\Psi}_i \circ (\hat{\Psi}_i|_{E_{(I_1, I_2)(J_1, J_2)}(P)})^{-1} \circ \Phi_i|_{A_{I', J}} = \Phi_i|_{A_{I', J}}$$

as claimed. This shows that $\hat{\Psi}_i|_C : C \rightarrow H_i$ is surjective, so $H_i \leq_R C$. \square

8.5 Proofs of Lemma 5.8 and Proposition 5.10, and construction of $B(G)$

Proof of Lemma 5.8. Let $(I_1, I_2) \in V(P)$ with $I_i \in V(H_i)$. Let $e = (e_1, e_2), e' = (e'_1, e'_2) \in E_{(I_1, I_2)}(P)$ such that $t(\partial\Psi_P(e)) = t(\partial\Psi_P(e'))$. We must show that $t(e) = t(e')$. Toward that goal, note that for each $i = 1, 2$, we have

$$\begin{aligned} t(\partial\Psi_P(e)) &= \partial\Psi_P(t(e)) = \partial\Psi_i(t(e_i)) \\ t(\partial\Psi_P(e')) &= \partial\Psi_P(t(e')) = \partial\Psi_i(t(e'_i)) \end{aligned}$$

Therefore $\partial\Psi_i(t(e_i)) = \partial\Psi_i(t(e'_i))$ for each i , since $t(\partial\Psi_P(e)) = t(\partial\Psi_P(e'))$ by hypothesis. But since the H_i are bunchy, we in fact have $t(e_i) = t(e'_i)$ for each i , so indeed $t(e) = t(e')$ as required. This shows that $\partial\Psi_P|_{F((I_1, I_2))}$ is a bijection, so P is bunchy.

Let C be a principal subgraph of P such that $\partial\hat{\Psi}_i|_{V(C)} : V(C) \rightarrow V(H_i)$ are surjective. Then $M(C) = M$. For each $J \in V(M)$, we have $\Sigma_C^{-1}(J) = \Sigma_P^{-1}(J) \cap V(C)$. Thus C is bunchy. \square

Proof of Proposition 5.10. To prove (1), let $H_1, H_2 \leq_R G$ be \leq_R -maximal among the bunchy right-resolving factors of G . Let $\Phi_i \in \text{hom}_R(G, H_i)$ and $\Psi_i \in \text{hom}_R(H_i, M)$. Let $P = H_1 \times_{\Psi_1, \Psi_2} H_2$. By Theorem 5.7, there exist a principal subgraph C of P , and $\Delta_i \in \text{hom}_R(G, C)$, such that $\Phi_i = \hat{\Psi}_i|_C \circ \Delta_i$, so $H_i \leq_R C$. By Lemma 5.8, C is bunchy, so by the maximality of the H_i , we have $H_1 = C = H_2$. This proves uniqueness, so we can take $B = H_i = C$.

To prove (2), let $H \leq_R G$ be bunchy and let $\Phi \in \text{hom}_R(G, H)$. Let $\Phi' \in \text{hom}(G, B)$, $\Psi \in \text{hom}_R(H, M)$, $\Psi' \in \text{hom}(B, M)$. Let $P = H \times_{\Phi, \Phi'} B$. By the universal property (Theorem 5.7), there exist a principal subgraph C of P and $\Delta, \Delta' \in \text{hom}_R(G, C)$ with $\Phi = \hat{\Psi} \circ \Delta$ and $\Phi' = \hat{\Psi}' \circ \Delta'$. Again by Lemma 5.8, C is bunchy, so by (1) and the fact that $B \leq_R C$, we have $B = C$. This proves (2) with $\Theta = \hat{\Psi}'|_C$. \square

We now present the construction of $B(G)$ described in Algorithm 7.3.

Proposition 8.7. *Let G be a graph. For $J_1, J_2 \in V(G)$ with $\Sigma_G(J_1) = \Sigma_G(J_2)$, write $J_1 \approx_0 J_2$ if there exist paths $\gamma = \gamma_1 \cdots \gamma_n, \delta = \delta_1 \cdots \delta_n \in L(G)$ where $\gamma_i, \delta_i \in E(G)$, such that $s(\gamma_1) = s(\delta_1)$, $t(\gamma_n) = J_1$, $t(\delta_n) = J_2$, and $\Sigma_G(t(\gamma_i)) = \Sigma_G(t(\delta_i))$ for each i . Let \approx denote the transitive closure of \approx_0 and let $\Phi \in \text{hom}_R(G, M(G))$. Then \approx is a congruence with respect to Φ , and $B(G) = G/\approx$.*

Proof. Let $M = M(G)$. Let $I \in V(M)$, $I_1, I_2 \in \Sigma_G^{-1}(I)$, and $J \in F(I)$. Suppose that $I_1 \approx I_2$. Let $J_i \in F(I_i) \cap \Sigma_G^{-1}(J)$. We need to show that $J_1 \approx J_2$. Let $I_1 = I^{(0)}, I^{(1)}, \dots, I^{(n)} = I_2 \in \Sigma_G^{-1}(I)$ with $I^{(j)} \approx_0 I^{(j+1)}$. Let $\gamma^{(j)}, \delta^{(j)} \in L(G)$ witness the relation $I^{(j)} \approx_0 I^{(j+1)}$. Choose $J^{(j)} \in F(I^{(j)}) \cap \Sigma_G^{-1}(J)$, with $J^{(0)} = J_1$ and $J^{(n)} = J_2$. Let $e^{(j)} \in E_{I^{(j)}J^{(j)}}(G)$. Then $\gamma^{(j)}e^{(j)}, \delta^{(j)}e^{(j+1)}$ witness $J^{(j)} \approx_0 J^{(j+1)}$. This shows that $J_1 \approx J_2$.

Let $\Phi \in \text{hom}_R(G, M(G))$. As in the previous paragraph, let $I \in V(M)$, let $J \in F(I)$, and let $I_1, I_2 \in \Sigma_G^{-1}(I)$. Suppose that $I_1 \approx I_2$. Let $a \in E_{IJ}(M)$, let $J_i = I_i \cdot a$, and let $e_i = (\Phi|_{E_{I_i}(G)})^{-1}(a)$. Then $t(e_i) = J_i$, so $J_1 \approx J_2$. Therefore \approx is indeed a congruence for Φ . It follows that $F([I_i]_{\approx}) \cap \Sigma_{G/\approx}^{-1}(J) = \{[J_i]_{\approx}\}$, so indeed G/\approx is bunchy.

To see that G/\approx is \leq_R -maximal among the bunchy factors of G , let $H \leq_R G$ be bunchy and $\Psi \in \text{hom}_R(G, H)$. If $I_1 \approx_0 I_2$ and H is bunchy, then we must have $\partial\Psi(I_1) = \partial\Psi(I_2)$. Therefore the partition into $\partial\Psi$ -fibers corresponds to an equivalence relation that coarsens the symmetric, reflexive relation \approx_0 , and thus also coarsens the transitive closure \approx . Considering $V(H)$ as a partition of $V(G)$, we must therefore have $V(H) \preceq V(G)/\approx$, so $(G/\approx) = B(G)$ by the maximality of $B(G)$. \square

8.6 Proofs of Propositions 6.1 and 7.1

Lemma 8.8. *Let G_1, G_2, H, K be graphs with $K \leq_R H \leq_R G_i$. Let $\Delta \in \text{hom}_R(H, K)$. Let $\Psi_i \in \text{hom}_R(G_i, H)$ and $P = G_1 \times_{\Psi_1, \Psi_2} G_2$. Let $\Phi_i = \Delta \circ \Psi_i$ and $Q = G_1 \times_{\Phi_1, \Phi_2} G_2$. Then, noting that $V(P) = V(Q)$ and $E(P) \subseteq E(Q)$, we have $\hat{\Phi}_i|_P = \hat{\Psi}_i$, and every principal subgraph of P is a principal subgraph of Q .*

Proof. Let C be a principal subgraph of P . By the definition of the fiber product, we have $V(P) = V(Q) = V(G_1) \times V(G_2)$. Let $I \in V(H)$ and let $I_i \in (\partial\Psi_i)^{-1}(I)$. Suppose that $(I_1, I_2) \in V(C)$. We need to show that $E_{(I_1, I_2)}(C) = E_{(I_1, I_2)}(Q)$.

Since $E_{(I_1, I_2)}(C) = E_{(I_1, I_2)}(P)$ by the definition of a principal subgraph, it is enough to show that $E_{(I_1, I_2)}(P) = E_{(I_1, I_2)}(Q)$ for any $(I_1, I_2) \in V(P)$. Clearly $E_{(I_1, I_2)}(P) \subseteq E_{(I_1, I_2)}(Q)$, so it is enough to show that $|E_{(I_1, I_2)}(P)| = |E_{(I_1, I_2)}(Q)|$. To see this equality, note that since $\Delta \circ \Psi_P = \Phi_Q$, we have

$$|E_{(I_1, I_2)}(P)| = |E_{\partial\Psi_P(I_1, I_2)}(H)| = |E_{\partial\Phi_Q(I_1, I_2)}(K)| = |E_{(I_1, I_2)}(Q)|$$

where the equalities follow from the facts that Ψ_P, Δ , and Φ_Q respectively are right-resolving. Therefore $E_{(I_1, I_2)}(P) = E_{(I_1, I_2)}(Q)$. This shows that C is indeed a principal subgraph of Q . Moreover, for $e = (e_1, e_2) \in E(P)$, we have $\hat{\Phi}_i(e) = e_i = \hat{\Psi}_i(e)$, so indeed $\hat{\Phi}_i|_P = \hat{\Psi}_i$. \square

Lemma 8.9. *Let G_1, G_2, K be graphs with $K \leq_S G_i$ via $\Phi_i \in \text{hom}_S(G_i, K)$. Let C be a principal subgraph of $P = G_1 \times_{\Phi_1, \Phi_2} G_2$ such that the $\partial\hat{\Phi}_i|_{V(C)} : V(C) \rightarrow V(G_i)$ are surjective. Then $\hat{\Phi}_i|_C \in \text{hom}_S(C, G_i)$. In particular, C is a common synchronizing extension of the G_i .*

Proof. Let $I \in V(K)$ and let $I_i, I'_i \in V(G_i)$ with $\partial\Phi_i(I_i) = \partial\Phi_i(I'_i) = I$. Since the Φ_i are synchronizing, we have $I_i \sim_{\Phi_i} I'_i$. Suppose that $(I_1, I_2), (I'_1, I'_2) \in V(C)$. In order to show that $\hat{\Phi}_i|_C \in \text{hom}_S(C, G_i)$, we claim that $(I_1, I_2) \sim_{\Phi_P} (I'_1, I'_2)$. This will show that $\Phi_P|_C \in \text{hom}_S(C, K)$. Since $\Phi_P|_C = \Phi_i \circ \hat{\Phi}_i|_C$, it will then follow that $\hat{\Phi}_i|_C \in \text{hom}_S(C, G_i)$ by Theorem 3.3(4).

To prove this claim, let $u \in L_I(K)$. Since $I_1 \sim_{\Phi_1} I'_1$, there exists $v_1 \in L_{t(u)}(K)$ such that $I_1 \cdot uv_1 = I'_1 \cdot uv_1$. Similarly, since $I_2 \sim_{\Phi_2} I'_2$, there exists $v_2 \in L_{t(v_1)}(K)$ such that $I_2 \cdot uv_1v_2 = I'_2 \cdot uv_1v_2$. Then in particular $(I_1, I_2) \cdot uv_1v_2 = (I'_1, I'_2) \cdot uv_1v_2$. Since $u \in L_I(K)$ was arbitrary, we have $(I_1, I_2) \sim_{\Phi_P} (I'_1, I'_2)$ as claimed. \square

Proof of Proposition 7.1(1). Let $K \leq_S G_i$ and $\Psi_i \in \text{hom}_S(G_i, K)$. Let $P = G_1 \times_{\Psi_1, \Psi_2} G_2$ and let C be a principal subgraph of P such that $\partial\hat{\Psi}_i|_{V(C)} : V(C) \rightarrow V(G_i)$ are surjective. By Lemma 8.9, we have $\hat{\Psi}_i|_C \in \text{hom}_S(C, G_i)$. Let $\Delta \in \text{hom}_R(K, M)$, let $\Phi_i = \Delta \circ \Psi_i$, and let $Q = G_1 \times_{\Phi_1, \Phi_2} G_2$. Then, by Lemma 8.8, C is a principal subgraph of Q , with $\hat{\Phi}_i|_C = \hat{\Psi}_i|_C$. In particular, $\hat{\Phi}_i|_C \in \text{hom}_S(C, G_i)$. \square

We now give an equivalent form of the $O(G)$ conjecture. Fragments of this result appear in Proposition 7.1. The logical structure of Proposition 8.10 (one statement is equivalent to the equivalence of four other statements) is unusual.

Proposition 8.10. *Let \mathcal{F} be a family of graphs satisfying the following conditions:*

- (i) *If $G \in \mathcal{F}$ and $H \leq_R G$, then $H \in \mathcal{F}$.*
- (ii) *Let $G_1, G_2, K \in \mathcal{F}$ with $K \leq_R G_i$. Let $\Phi_i \in \text{hom}_R(G_i, K)$, let $P = G_1 \times_{\Phi_1, \Phi_2} G_2$, and let C be a principal subgraph of P such that the $\partial\hat{\Phi}_i|_{V(C)}$ are surjective. Then $C \in \mathcal{F}$.*

Then the following assertions are equivalent.

- (1) For any $G \in \mathcal{F}$, there exists a unique \leq_S -minimal graph $O(G) \leq_S G$.
- (2) For any $G_1, G_2 \in \mathcal{F}$, the following assertions are equivalent.
 - (a) $O(G_1), O(G_2)$ exist and are equal.
 - (b) G_1, G_2 have a common synchronizing factor.
 - (c) $M(G_1) = M(G_2) = M$ and there exist $\Phi_i \in \text{hom}_R(G_i, M)$ such that $\hat{\Phi}_i \in \text{hom}_S(C, G_i)$ for some principal subgraph C of $Q = G_1 \times_{\Phi_1, \Phi_2} G_2$.
 - (d) G_1, G_2 have a common synchronizing extension $K \in \mathcal{F}$.

Remark 8.1. The proof of Proposition 8.10 in fact shows that both (1) and (2) are equivalent to the assertion that (d) implies (b). Note that for $G_1 = G_2 = G$, (a) states that $O(G)$ is well-defined, while (b)–(d) are trivial.

Remark 8.2. The $O(G)$ conjecture states that the equivalent statements (1) and (2) in Proposition 8.10 hold with \mathcal{F} equal to the class of all strongly connected graphs.

Proof of Proposition 8.10. First, assume (2). Let G be a graph and let $H_1, H_2 \leq_S G$ be \leq_S -minimal. Since H_1, H_2 have the common synchronizing extension G , they satisfy condition (d), so they also satisfy condition (b), i.e. there exists a common synchronizing factor $K \leq_S H_i$. But since the H_i were assumed minimal, we must have $H_1 = K = H_2 = O(G)$.

Now, assume (1) and deduce (2) as follows. Trivially, (a) implies (b) and (c) implies (d). Moreover, (b) implies (c) by Proposition 7.1(1). Finally, assume (d). Suppose that G_1, G_2 have a common synchronizing extension K . Then $O(G_1) = O(K) = O(G_2)$, so (d) implies (a). \square

Proof of Proposition 7.1(2). This is immediate from Proposition 8.10, specifically, the equivalence of 2(b) and 2(c), with \mathcal{F} taken to be the class of all strongly connected graphs. \square

Proof of Proposition 7.1(3). By Theorem 5.4, this is immediate from Proposition 8.10, again via the equivalence of 2(b) and 2(c), but with \mathcal{F} taken to be the class of strongly connected almost bunchy graphs. \square

Proposition 8.11. *Let \mathcal{F} be a family of graphs such that, if $G \in \mathcal{F}$ and $H \leq_R G$, then $H \in \mathcal{F}$. Then the following assertions are equivalent.*

- (1) Any \leq_S -minimal graph $H \in \mathcal{F}$ is bunchy.
- (2) For any $G \in \mathcal{F}$, there exists some bunchy $H \leq_S G$.
- (3) For any non-bunchy $G \in \mathcal{F}$, there exists some $\Phi \in \text{hom}_R(G, M(G))$ with \sim_Φ nontrivial.
- (4) For any $G \in \mathcal{F}$, $B(G) \leq_S G$.

Remark 8.3. Note that Proposition 8.11 is a more detailed version of Proposition 6.1.

Proof. To see that (1) implies (2), let $G \in \mathcal{F}$ and consider the set $\{H \mid H \leq_S G\}$. Being a finite partially ordered set, this set must have at least one minimal element. If (1) holds, then this minimal element is bunchy. Thus (2) holds.

To see that (2) implies (3), let $G \in \mathcal{F}$ be non-bunchy. There exists at least one \leq_S -minimal graph $H \leq_S G$. If (2) holds, then G is not \leq_S -minimal, so $H \neq G$. Let $\Psi \in \text{hom}_S(G, H)$. The $\partial\Psi$ -fibers are precisely the \sim_Ψ -classes. Since $H \neq G$, the $\partial\Psi$ -fibers are not merely singletons, so \sim_Ψ is nontrivial. Let $M = M(G)$ and $\Delta \in \text{hom}_R(H, M)$, and let $\Phi = \Delta \circ \Psi$. By Lemma 8.4, the \sim_Φ -classes are unions of \sim_Ψ -classes, so in particular, \sim_Φ is nontrivial.

To see that (3) implies (1), let $G \in \mathcal{F}$ be non-bunchy. If (3) holds, then there exists $\Phi \in \text{hom}_R(G, M(G))$ with \sim_Φ nontrivial. Then $G / \sim_\Phi \leq_S G$ and $G / \sim_\Phi \neq G$; in particular, G is not \leq_S -minimal. This proves (1) in the contrapositive.

Finally, we show that (2) and (4) are equivalent. If (4) holds, then (2) holds since $B(G)$ is bunchy. Conversely, assume (2) and let $G \in \mathcal{F}$. Then there exists $H \leq_S G$ bunchy. Let $\Phi \in \text{hom}_S(G, H)$ and let $B = B(G)$. By Proposition 5.10(2) and Theorem 3.3(4), we have $\Phi = \Delta \circ \Theta$ for some $\Theta \in \text{hom}_S(G, B)$, $\Delta \in \text{hom}_S(B, H)$. Therefore (4) holds. \square

Corollary 8.12. *Let \mathcal{F} be a family of graphs such that, if $G \in \mathcal{F}$ and $H \leq_R G$, then $H \in \mathcal{F}$. Suppose that any \leq_S -minimal element of \mathcal{F} is bunchy. Then $O(G)$ exists for any $G \in \mathcal{F}$.*

The proof of Corollary 8.12 is a trivial adaptation of the proof of Proposition 6.2.

Proof of Proposition 7.1(4). Assuming the bunchy factor conjecture, the $O(G_i)$ are well-defined. By the equivalence of conditions (a),(c) in Proposition 8.11, the hypothesis on the $B(G_i)$ is equivalent to the equality $O(B(G_1)) = O(B(G_2))$, which in turn is equivalent to the equality $O(G_1) = O(G_2)$ by Proposition 8.11. \square

9 Proof of Theorem 4.3, following Trahtman

In this section, we recall Trahtman's proof of the road colouring theorem and reformulate it in a form applicable to Theorem 4.3.

9.1 Systems of maps with unique tallest trees

Trahtman's proof uses an idea well known in the literature on the combinatorics of transformations of finite sets—namely, that, the graph of a transformation f of a set V is a directed graph of constant out-degree 1 with V for its set of states where, for each state $I \in V$, the unique edge with source I has target $I \cdot f$. When V is finite, the graph consists of a set of state-disjoint directed cycles together with trees rooted on the cycles, directed toward their roots. Trahtman's proof is based on the construction of a transformation with a unique tallest tree (the height of a tree being the maximum distance from the root to a leaf). We need a few definitions in order to extend the notion of a unique tallest tree to a system of maps taking one set to another, cyclically, as opposed to a transformation of a single set.

Let $p \in \mathbb{N}$ and let $\{V_i\}_{0 \leq i \leq p-1}$ be disjoint finite sets. Let $a_i : V_i \rightarrow V_{i+1}$ be maps, with subscripts read modulo p . For $k \geq 0$, let $b_k = a_k a_{k+1} \cdots a_{p-1} a_0 \cdots a_{k-1} : V_k \rightarrow V_k$. As noted above in the general discussion of graphs of transformations, since each b_k is a transformation of a finite set, it is eventually periodic—that is, for each k , there exist $m \geq 0$, $z \geq 1$ such that $b_k^{m+z} = b_k^m$. Moreover, the orbits of individual elements $I \in V_k$ may vary in their eventually periodic behaviour. Specifically, for a given $I \in V_k$, consider the lexicographically minimal (ℓ, m, z) with $0 \leq \ell \leq p-1$, $m \geq 0$, $z \geq 1$, such that $I \cdot a_k \cdots a_{k+\ell-1} b_{k+\ell}^{m+z} = I \cdot a_k \cdots a_{k+\ell-1} b_{k+\ell}^m$. Define the *height* $h(I) = mp + \ell$ and the *root* $\rho(I) = I \cdot a_k \cdots a_{k+\ell-1} b_{k+\ell}^{m+z}$. The idea is that $h(I)$ is the number of steps required until the orbit of I reaches the root $\rho(I)$ and becomes periodic, with $z(I) = z$ being the period as a multiple of p .

To talk about unique tallest trees, let $h_{\max, k} = \max\{h(I) \mid I \in V_k\}$, and let $h_k(J) = \max\{h(I) \mid I \in V_k, \rho(I) = J\}$. Let $z_k = \text{lcm}\{z(I) \mid I \in V_k\}$. We say that the system $(V_i, a_i)_{0 \leq i \leq p-1}$ has a *unique tallest tree* at V_k if there is a unique J with $h_k(J) = h_{\max, k}$. Note that the terms we have defined here still make sense even if the V_i are not all pairwise disjoint, as long as any two are either equal or disjoint, since we can make them disjoint by replacing V_i with $V_i \times \{i\}$.

We now present our interpretation of a key step in Trahtman's proof, applying Proposition 3.8 and closely following [9].

Lemma 9.1. *Let G, H be strongly connected graphs with $H \leq_R G$. Let $\Phi \in \text{hom}_R(G, H)$. Let $I_0, \dots, I_{p-1} \in V(H)$, not necessarily distinct, be such that $I_{i+1} \in F(I_i) \neq \emptyset$, and let $a_i \in E_{I_i I_{i+1}}(H)$*

(with subscripts read modulo p). Suppose that the system $((\partial\Phi)^{-1}(I_i), a_i)_{0 \leq i \leq p-1}$ has a unique tallest tree, where we write a_i for the map $I \mapsto I \cdot a_i$, $I \in (\partial\Phi)^{-1}(I_i)$. Then \sim_Φ is nontrivial.

Proof. For $0 \leq i \leq p-1$, let $V_i = (\partial\Phi)^{-1}(I_i)$. Suppose without loss of generality that the system has a unique tallest tree at V_0 . Let $I \in V_0$ be a state of maximal height $h(I) = h_{\max,0}$. Let $m \geq 0$, $0 \leq p-1$ be such that $h_{\max,0} = mp + \ell$, and let $z = z(I)$ and let $R = \rho(I)$. Note that $R \in V_\ell$. That is, upon cyclic application of the maps a_i , I is eventually mapped to R , then returns to R every z cycles around the graph.

By strong connectedness and Lemma 3.7, let $U \subseteq I_0$ be a minimal image such that $I \in U$. We claim that there is no other state $I' \in U$ with $\rho(I') = R$ and $I' \neq I$. Indeed, suppose that there is such a state I' . Then $I \cdot a_0 \cdots a_{\ell-1} b_\ell^m = I' \cdot a_0 \cdots a_{\ell-1} b_\ell^m = R$, so $|U \cdot a_0 \cdots a_{\ell-1} b_\ell^m| < |U|$, contradicting the minimality of U . This proves the claim. Let $U_0 = U \setminus \{I\}$. Then every element of U_0 has height strictly less than $h_{\max,0}$.

Let $u_1 = a_0 \cdots a_{\ell-1} b_\ell^{m-1} a_\ell \cdots a_{\ell-2} : V_0 \rightarrow V_{\ell-1}$, where the tail $a_\ell a_{\ell+1} \cdots a_{\ell-2}$ includes each a_i exactly once, other than $a_{\ell-1}$, and the subscripts are read modulo p . The effect of applying u_1 to V_0 is to bring I to one step before its first encounter its root R ; since I has maximal height, every other $I' \in V_0$ has already reached its root and is in the periodic part of its orbit after application of u_1 .

Let $u_2 = u_1 b_{\ell-1}^{z\ell} = a_0 \cdots a_{\ell-1} b_{\ell-1}^{m+z\ell} a_\ell \cdots a_{\ell-2} : V_0 \rightarrow V_{\ell-1}$. Let $J_i = I \cdot u_i$ and $U_i = U \cdot u_i$. As observed in the previous paragraph, after application of u_1 , I is not yet in the periodic part of its orbit (i.e. the orbit under cyclic application of the maps a_i), but every other element of U_1 is in the periodic part of its orbit.

Observe that $J_1 \neq J_2$ by the assumed value of $h(I)$. However, since I is the unique element of U with this maximal height, we have $U_1 \Delta U_2 = \{J_1, J_2\}$. Since the U_i are minimal images, we have $J_1 \sim_\Phi J_2$ by Proposition 3.8. \square

9.2 Obtaining a right-resolver with a unique tallest tree

Let G be a graph. We define a *total order colouring* to be a total ordering of each edge set $E_I(H)$, i.e. a labeling of the edges of G such that, if $|E_I(G)| = k$, then the edges in $E_I(G)$ are labeled bijectively by $\{0, \dots, k-1\}$. Suppose that $M = M(G)$ is a cycle of bunches. Then, once a total order colouring of M is fixed, total order colourings of G correspond bijectively with right-resolvers $\Phi \in \text{hom}_R(G, M)$.

Letting $V(M) = \{I_0, \dots, I_{p-1}\}$, there is exactly one edge $a_i \in E_{I_i}(H)$ labeled 0 for each i . This yields a subgraph W of G consisting of edges labeled 0, which is a spanning subgraph of G of constant out-degree 1. Every graph of constant out-degree 1 consists of a set of state-disjoint cycles, together with trees rooted on the cycles, directed toward their roots. The height of a tree is the maximum path length from a state in the tree to its root. Observe that the system $((\partial\Phi)^{-1}(I_i), a_i)_{i=0}^{p-1}$ has a unique tallest tree, in the sense of mappings, if and only if there is a unique tallest tree in W .

We now present our interpretation of the main technical lemma in the proof of the road colouring theorem (Lemma 10.4.6 in [9]). The following is not how the lemma is stated in [9], but one can follow the proof and observe that it is equivalent.

Lemma 9.2 (Trahtman). *Let G be a strongly connected graph such that $M(G)$ is a cycle of bunches. At least one of the following is true:*

1. G is itself a cycle of bunches.
2. G has two distinct bunches whose outgoing edges have the same target.
3. G admits a total order colouring with a unique tallest tree.

With this result, we can prove our generalization of the road colouring theorem.

Proof of Theorem 4.3. Let $M = M(G)$. The claim is trivially true if $|V(G)| = |V(M)|$, in which case $G = M$. Suppose that it is true for all H with $|V(H)| \leq_R N$ and $M(H) = M$. Suppose that $|V(G)| = N + 1$. If G is bunched, then we are done. If G is not bunched, but has two states that can

be in-amalgamated, then they are stable for some $\Phi \in \text{hom}_R(G, M)$ by Lemma 3.5. Let $G' = G / \sim_\Phi$. Then $|V(G / \sim_\Phi)| < |V(G)|$, so by the inductive hypothesis, there is some bunchy $H \leq_S G / \sim_\Phi \leq_S G$.

Now, suppose that G does not have two states that can be in-amalgamated—in particular, G does not have two distinct bunches whose outgoing edges have the same target. Since G is not bunchy, it is in particular not a cycle of bunches, so by Lemma 9.2, it admits a total order colouring with unique tallest tree. As remarked above, this total order colouring corresponds to some right-resolver $\Phi \in \text{hom}_R(G, M)$, which has \sim_Φ nontrivial by Lemma 9.1. Then once more G / \sim_Φ is strictly smaller than G . We can then apply Proposition 8.11 to conclude that any \leq_S -minimal O with $M = M(O)$ is a cycle of bunches. If $O \leq_S G$, then $O = O_{M,p}$ for $p = \text{per}(G)/\text{per}(M)$. \square

Acknowledgments

The author thanks Brian Marcus and Tom Meyerovitch for their patient, generous advice and supervision; Theo Morrison for a number of insightful questions; the anonymous referees for very careful reading and suggestions; and Mike Boyle for suggesting (to Brian Marcus) a renewed attack on the $O(G)$ problem, shortly after the road problem was solved in 2007. The author was supported through NSERC Discovery Grant EJYR GR010163 (PI: Brian Marcus).

References

- [1] R.L. Adler, L.W. Goodwyn, and B. Weiss. Equivalence of topological Markov shifts. *Isr. J. Math.*, 27(1):49–63, 1977.
- [2] R.L. Adler and B. Weiss. Entropy, a complete metric invariant for automorphisms of the torus. *Proc. Nat. Acad. Sci.*, 57(6):1573–1576, 1967.
- [3] D. Ananichev, V. Gusev, and M. Volkov. Slowly synchronizing automata and digraphs. In Petr Hlinený and Antonín Kucera, editors, *MFCS 2010*, volume 6281 of *Lect. Notes Comput. Sci.*, pages 55–65. Springer-Verlag Berlin Heidelberg, 2010.
- [4] J. Araújo, P. Cameron, and B. Steinberg. Between primitive and 2-transitive: synchronization and its friends. *EMS Surv. Math. Sci.*, 40(2):101–184, 2017.
- [5] J. Ashley, B. Marcus, and S. Tuncel. The classification of one-sided Markov chains. *Ergod. Theor. Dynam. Syst.*, 17(2):269–295, 1997.
- [6] L. Babai. Groups, graphs, algorithms: the graph isomorphism problem. In B. Sirakov, P. Ney de Souza, and M. Viana, editors, *ICM 2018*, volume 3, pages 3303–3320. World Scientific, 2018.
- [7] M.V. Berlinkov. On two algorithmic problems about synchronizing automata. In A.M. Shur and M.V. Volkov, editors, *DLT 2014*, volume 8633 of *LNCS*, pages 61–67, 2014.
- [8] M.V. Berlinkov. On the probability of being synchronizable. In S. Govindarajan and A. Maheshwari, editors, *CALDAM 2016*, volume 9602 of *Lect. Notes Comput. Sci.*, pages 73–84. Springer International Publishing Switzerland, 2016.
- [9] J. Berstel, D. Perrin, and C. Reutenauer. *Codes and Automata*, volume 129 of *Encycl. Math. Appl.* Cambridge University Press, 2009.
- [10] M. Boyle and S. Tuncel. Regular isomorphism of Markov chains is almost topological. *Ergod. Theor. Dynam. Syst.*, 10(1):89–100, 1990.
- [11] G. Budzban and P. Feinsilver. The generalized road coloring problem and periodic digraphs. *Appl. Algebra Engrg. Comm. Comput.*, 22(1):21–35, 2011.

- [12] M.-P. Béal and D. Perrin. A quadratic algorithm for road coloring. *Discrete Appl. Math.*, 169:15–29, 2014.
- [13] J. Cai and R. Frongillo. Computational complexity of problems for deterministic presentations of sofic shifts, 2021. Online preprint, arXiv:2112.03484v1.
- [14] D.G. Corneil and C.C. Gotlieb. An efficient algorithm for graph isomorphism. *J. Assoc. Comput. Mach.*, 17(1):51–64, 1970.
- [15] D. Eppstein. Reset sequences for monotonic automata. *SIAM J. Comput.*, 19(3):500–510, 1990.
- [16] K. Culik II, J. Karhumäki, and J. Kari. A note on synchronized automata and road coloring problem. In W. Kuich, G. Rozenberg, and A. Salomaa, editors, *DLT 2001*, volume 2295 of *Lect. Notes Comput. Sci.*, pages 175–185. Springer-Verlag Berlin Heidelberg, 2002.
- [17] J. Kari. Synchronizing finite automata on Eulerian digraphs. *Theor. Comput. Sci.*, 295:223–232, 2003.
- [18] D. Lind and B. Marcus. *An Introduction to Symbolic Dynamics and Coding*. Cambridge University Press, 2nd edition, 2020.
- [19] G.L. Miller. Graph isomorphism, general remarks. *J. Comput. Syst. Sci.*, 18:128–142, 1979.
- [20] G.L. O’Brien. The road-colouring problem. *Isr. J. Math.*, 39(1–2):145–154, 1981.
- [21] E. Riehl. *Category Theory in Context*. Aurora. Dover, 2017.
- [22] A.N. Trahtman. The road coloring problem. *Isr. J. Math.*, 172(1):51–60, 2009.
- [23] V. Vorel. Subset synchronization and careful synchronization of binary finite automata. *Intern. J. Found. Comput. Sci.*, 27(5):557–577, 2016.
- [24] H.W. Wielandt. Unzerlegbare, nicht negative Matrizen (German). *Math. Z.*, 52:642–648, 1950.