

Error analysis of a fully discrete scheme for the Cahn-Hilliard-Magneto-hydrodynamics problem

Hailong Qiu

Received: date / Accepted: date

Abstract In this paper we analyze a fully discrete scheme for a general Cahn-Hilliard equation coupled with a nonsteady Magneto-hydrodynamics flow, which describes two immiscible, incompressible and electrically conducting fluids with different mobilities, fluid viscosities and magnetic diffusivities. A typical fully discrete scheme, which is comprised of conforming finite element method and the Euler semi-implicit discretization based on a convex splitting of the energy of the equation is considered in detail. We prove that our scheme is unconditionally energy stability and obtain some optimal error estimates for the concentration field, the chemical potential, the velocity field, the magnetic field and the pressure. The results of numerical tests are presented to validate the rates of convergence.

Keywords Nonstationary Magneto-hydrodynamics flow · Cahn-Hilliard equation · Fully discrete scheme · Conforming finite element · Energy stability · Optimal error estimates

Mathematics Subject Classification (2010) 65N30 · 76M10 · 76W05

1 Introduction

In the paper, we derive error estimates for a fully discrete, first order in time, finite element method for the Cahn-Hilliard-Magneto-hydrodynamics problem for two phase flow. Let $\Omega \subset \mathbf{R}^d$ ($d = 2, 3$) stands for an open convex polygonal or polyhedral domain with Lipschitz continuous boundary $\partial\Omega$. For all $\phi \in$

Hailong Qiu
School of Mathematics and Physics, Yancheng Institute of Technology, Yancheng, 224051, China.
This work is supported by the Natural Science Foundation of China (No. 11701498).
E-mail: qhllf@163.com

$H^1(\Omega)$, $\mathbf{u} \in \mathbf{L}^2(\Omega)$ and $\mathbf{B} \in \mathbf{L}^2(\Omega)$, consider the following energy:

$$\mathbf{E}(\phi, \mathbf{u}, \mathbf{B}) = \int_{\Omega} \left(\frac{1}{2} \|\mathbf{u}\|^2 + \frac{S_c}{2} \|\mathbf{B}\|^2 + \frac{\lambda \epsilon}{2} \|\nabla \phi\|^2 + \frac{\lambda}{4\epsilon} (1 - \phi^2)^2 \right) dx, \quad (1)$$

where ϕ , \mathbf{u} and \mathbf{B} denote respectively the concentration field, the velocity field and the magnetic field, and the parameter $\epsilon > 0$ stands for the interfacial thickness between the two phases. The Cahn-Hilliard-Magneto-hydrodynamics system is a gradient flows of this energy:

$$\partial_t \phi - \epsilon \operatorname{div}(\kappa(\phi) \nabla \mu) - \nabla \phi \cdot \mathbf{u} = 0, \quad \text{in } \Omega \times T, \quad (2a)$$

$$-\epsilon \Delta \phi + \epsilon^{-1} (\phi^3 - \phi) = \mu, \quad \text{in } \Omega \times T, \quad (2b)$$

$$\partial_t \mathbf{u} - \operatorname{div}(2\nu(\phi) \mathbb{D}(\mathbf{u})) + (\mathbf{u} \cdot \nabla) \mathbf{u} + S_c \mathbf{B} \times \operatorname{curl} \mathbf{B} \quad (2c)$$

$$+ \nabla p = \lambda \mu \nabla \phi, \quad \text{in } \Omega \times T,$$

$$\partial_t \mathbf{B} + \operatorname{curl}(\eta(\phi) \operatorname{curl} \mathbf{B}) - \operatorname{curl}(\mathbf{u} \times \mathbf{B}) = \mathbf{0}, \quad \text{in } \Omega \times T, \quad (2d)$$

$$\operatorname{div} \mathbf{u} = 0, \quad \text{in } \Omega \times T, \quad (2e)$$

$$\operatorname{div} \mathbf{B} = 0, \quad \text{in } \Omega \times T. \quad (2f)$$

This problem is considered in conjunction with natural and no-flux/no-flow boundary conditions:

$$\frac{\partial \phi}{\partial \mathbf{n}} = \frac{\partial \phi}{\partial \mathbf{n}} = 0, \quad \text{on } \partial \Omega \times T, \quad (3a)$$

$$\mathbf{u} = \mathbf{0}, \quad \text{on } \partial \Omega \times T, \quad (3b)$$

$$\mathbf{B} \cdot \mathbf{n} = 0, \quad \text{on } \partial \Omega \times T, \quad (3c)$$

$$\operatorname{curl} \mathbf{B} \times \mathbf{n} = \mathbf{0}, \quad \text{on } \partial \Omega \times T, \quad (3d)$$

and initial conditions:

$$\phi(\mathbf{x}, 0) = \phi_0(\mathbf{x}), \quad \mathbf{u}(\mathbf{x}, 0) = \mathbf{u}_0(\mathbf{x}), \quad \mathbf{B}(\mathbf{x}, 0) = \mathbf{B}_0(\mathbf{x}), \quad \text{in } \Omega, \quad (4)$$

where $T > 0$ is time, $\phi \approx \pm 1$ represents two different fluids, μ and p are respectively the chemical potential and the pressure. $\mathbb{D}(\mathbf{u}) := \frac{1}{2}(\nabla \mathbf{u} + \nabla \mathbf{u}^T)$ is the strain-rate tensor. The given functions $\kappa(\phi) > 0$, $\nu(\phi) > 0$ and $\eta(\phi) = \frac{1}{\eta_0 \sigma(\phi)} > 0$ stand for respectively the mobility, the fluid viscous, the magnetic diffusivity with σ the electrical conductivity. η_0 is the magnetic permeability. $S_c := \frac{1}{\eta_0 \rho_0}$ denotes the coupling coefficient, here ρ_0 is the reference density. λ denotes the mixing energy density.

The (2)-(4) system satisfies the following energy laws:

$$\begin{aligned} \mathbf{E}(\phi(t), \mathbf{u}(t), \mathbf{B}(t)) + \int_{\Omega} \left(\nu(\phi) \|\nabla \mathbf{u}\|^2 + S_c \eta(\phi) \|\nabla \mathbf{B}\|^2 + \lambda \epsilon \kappa(\phi) \|\nabla \mu\|^2 \right) dx \\ = \mathbf{E}(\phi_0, \mathbf{u}_0, \mathbf{B}_0). \end{aligned} \quad (5)$$

In the last decades, phase field approaches for two phases incompressible flows have been widely developed to model and numerically solve the topological transitions of interfaces (cf. [6, 11, 1, 14, 19, 21, 32, 7] and references therein).

Recently, the research of interaction of electromagnetic fields with two immiscible, incompressible and electrically conducting fluids has become more and more important for the design and analysis of engineering field, such as fusion reactors, pump accelerators, metallurgical industry and Magneto-hydrodynamics generators [26, 23, 24, 4]. Very recently, in the work [33], two phase Magneto-hydrodynamics problem about the diffuse interface between two different incompressible fluids is considered.

Though many error estimates are available for fully discrete scheme to the Magneto-hydrodynamics equations [31, 25, 30, 22, 18] and for the Cahn-Hilliard/Navier-Stokes equations [6, 7, 8, 9, 5, 27, 10, 3], it is not trivial to cope with the system which couple Magneto-hydrodynamics with Cahn-Hilliard, since the phase field dependent coefficients affects the whole system. The major difficulties cause from the phase field dependent coefficients and from the coupling nonlinear terms. To the best of our knowledge, error estimates for fully discrete scheme of problem (2)-(4) are not yet set up [33]. However, the analysis in [7, 5, 3] cannot be easily developed to the fully discrete form of problem (2)-(4), as the time-space discretization raises another difficulty, particularly in deriving error estimates for the pressure.

In this paper, applying some useful techniques given in references [29, 5, 3], we analyze a fully discrete scheme for problem (2)-(4), which is comprised of conforming mixed finite element method in space and the Euler semi-implicit discretization with a convex splitting method in time. The main purpose of this paper is to derive some optimal error estimates for the concentration field, the chemical potential, the velocity field, the magnetic field and the pressure for the Cahn-Hilliard-Magneto-hydrodynamics system in fully discrete form. Furthermore, the error estimates are established do not need any CFL conditions. The highlight of this paper is to set up the error estimates for the fully discrete solution $(\phi_h^n, \mu_h^n, \mathbf{u}_h^n, p_h^n, \mathbf{B}_h^n)$ as follows:

$$\begin{aligned} \max_{1 \leq n \leq N} \|\nabla(\phi(t_n) - \phi_h^n)\| + \left(\Delta t \sum_{n=1}^N \|\nabla(\mu(t_n) - \mu_h^n)\|^2 \right)^{\frac{1}{2}} &\leq C(\Delta t + h^{k+1}), \\ \max_{1 \leq n \leq N} \|\mathbf{u}(t_n) - \mathbf{u}_h^n\| + \max_{1 \leq n \leq N} \|\mathbf{B}(t_n) - \mathbf{B}_h^n\| &\leq C(\Delta t + h^{k+1}), \\ \left(\Delta t \sum_{n=1}^N \|p(t_n) - p_h^n\|^2 \right)^{\frac{1}{2}} &\leq C(\Delta t + h^{k+1}). \end{aligned}$$

Numerical tests are given to confirm the theoretical rates of convergence.

The paper is organized as follows. In section 2 we present some preliminary results and the well-posedness of weak solution of Cahn-Hilliard-Magneto-hydrodynamics system. In section 3 we give fully discrete scheme and obtain its unconditionally energy stability. In section 4 we prove some optimal error estimates for the concentration field, the chemical potential, the velocity field, the magnetic field and the pressure. In section 5 we present some numerical tests to checked the theoretical results of the our scheme.

2 Continuous problem

2.1 Preliminaries

In this subsection we give some standard notations. Let $C^m(\Omega)$ ($m \in \mathbb{N}$) be the space of functions with up to m times continuously differentiable in Ω , and let $C^{m,1}(\Omega)$ be the space of functions in $C^m(\Omega)$ that are Lipschitz continuous in Ω . let $(L^p(\Omega), \|\cdot\|_{L^p})$ and $(W^{k,p}(\Omega), \|\cdot\|_{k,p})$ denote respectively the Lebesgue spaces and Sobolev spaces. For simplicity, we denote $\|\cdot\| := \|\cdot\|_{L^2}$, and denote by $H^k(\Omega)$ the Sobolev space $W^{k,2}(\Omega)$.

To define a weak formulation of problem (2)-(4), we introduce the following Sobolev spaces

$$\begin{aligned}\mathcal{X} &:= H_0^1(\Omega)^d = \{\mathbf{v} \in H^1(\Omega)^d : \mathbf{v}|_{\partial\Omega} = 0\}, \\ \mathcal{X}_0 &:= \{\mathbf{v} \in L^2(\Omega)^d : \nabla \cdot \mathbf{v} = 0, \mathbf{v}|_{\partial\Omega} = 0\}, \\ \mathcal{W} &:= H_n^1(\Omega)^d := \{\mathbf{w} \in H^1(\Omega)^d : \mathbf{w} \cdot \mathbf{n}|_{\partial\Omega} = 0\}, \\ \mathcal{W}_0 &:= \{\mathbf{w} \in H^1(\Omega)^d : \text{curl } \mathbf{w} \times \mathbf{n}|_{\partial\Omega} = 0\}, \\ \mathcal{X}_0 &:= \{\mathbf{v} \in \mathcal{X} : \text{div } \mathbf{v} = 0\}, \\ \mathcal{W}_n &:= \{\mathbf{w} \in \mathcal{W} : \text{div } \mathbf{w} = 0\}, \\ \mathcal{Q} &:= \{\varphi \in H^1(\Omega) : \frac{\partial \varphi}{\partial \mathbf{n}}|_{\partial\Omega} = 0\}, \\ \mathcal{M} &:= L_0^2(\Omega) = \{q \in L^2(\Omega), \int_{\Omega} q d\mathbf{x} = 0\}.\end{aligned}$$

The following useful inequalities hold [28, 12, 13, 15, 9, 18]:

$$\|\mathbf{v}\| \leq C_{\Omega} \|\nabla \mathbf{v}\|, \quad \forall \mathbf{v} \in \mathcal{X}, \quad (6a)$$

$$\|\mathbf{v}\|_{L^6} \leq C_{\Omega} \|\nabla \mathbf{v}\|, \quad \forall \mathbf{v} \in \mathcal{X}, \quad (6b)$$

$$\|\mathbb{D}\mathbf{v}\| \geq c_0 \|\nabla \mathbf{v}\|, \quad \forall \mathbf{v} \in \mathcal{X}, \quad (6c)$$

$$\|\mathbf{v}\|_{L^3} \leq C_{\Omega} \|\mathbf{v}\|^{\frac{6-d}{6}} \|\nabla \mathbf{v}\|^{\frac{d}{6}}, \quad \forall \mathbf{v} \in \mathcal{X}, \quad (6d)$$

$$\|\mathbf{v}\|_{L^4} \leq C_{\Omega} \|\mathbf{v}\|^{\frac{4-d}{4}} \|\nabla \mathbf{v}\|^{\frac{d}{4}}, \quad \forall \mathbf{v} \in \mathcal{X}, \quad (6e)$$

$$\|\mathbf{v}\|_{L^\infty} \leq C_{\Omega} \|\mathbf{v}\|_{H^1}^{1/2} \|\mathbf{v}\|_{H^2}^{1/2}, \quad \forall \mathbf{v} \in H^2(\Omega)^d, \quad (6f)$$

$$c_1 \|\nabla \mathbf{B}\|^2 \leq \|\text{curl } \mathbf{B}\|^2 + \|\text{div } \mathbf{B}\|^2, \quad \forall \mathbf{B} \in \mathcal{W}, \quad (6g)$$

$$\|\text{curl } \mathbf{B}\| \leq \sqrt{2} \|\nabla \mathbf{B}\|, \|\text{div } \mathbf{B}\| \leq \sqrt{d} \|\nabla \mathbf{B}\|, \quad \forall \mathbf{B} \in H^1(\Omega)^d, \quad (6h)$$

$$\|\phi\|_{L^p} \leq C_{\Omega} \|\phi\|_{H^1}, \quad (2 \leq p \leq 6) \quad \forall \mathbf{v} \in \mathcal{Q}, \quad (6i)$$

$$\|\phi\|_{L^3} \leq C_{\Omega} \|\phi\|^{\frac{6-d}{6}} \|\nabla \phi\|^{\frac{d}{6}} + C_{\Omega} \|\phi\|, \quad \forall \mathbf{v} \in \mathcal{Q}, \quad (6j)$$

$$\|\phi\|_{L^\infty} \leq C_{\Omega} \|\Delta \phi\|^{\frac{d}{2(6-d)}} \|\phi\|_{L^6}^{\frac{3(4-d)}{2(6-d)}} + C_{\Omega} \|\phi\|_{L^6}, \quad \forall \mathbf{v} \in \mathcal{Q}, \quad (6k)$$

where c_0 , c_1 and C_{Ω} are positive constants depending on Ω .

We define the following bilinear terms:

$$\begin{aligned} a_\phi(\varphi; \phi, \theta) &= \int_{\Omega} \kappa(\varphi) \nabla \phi \cdot \nabla \theta d\mathbf{x}, & a_f(\varphi; \mathbf{u}, \mathbf{v}) &= \int_{\Omega} 2\nu(\varphi) \mathbb{D}(\mathbf{u}) : \mathbb{D}(\mathbf{v}) d\mathbf{x}, \\ a_B(\varphi; \mathbf{B}, \mathbf{H}) &= \int_{\Omega} \eta(\varphi) \operatorname{curl} \mathbf{B} \cdot \operatorname{curl} \mathbf{H} d\mathbf{x} & + \int_{\Omega} \eta(\varphi) \operatorname{div} \mathbf{B} \cdot \operatorname{div} \mathbf{H} d\mathbf{x}, \\ d(\mathbf{v}, q) &= \int_{\Omega} q \operatorname{div} \mathbf{v} d\mathbf{x}, \end{aligned}$$

and trilinear terms:

$$\begin{aligned} b(\mathbf{w}, \mathbf{u}, \mathbf{v}) &= \frac{1}{2} \int_{\Omega} [(\mathbf{w} \cdot \nabla) \mathbf{u}] \cdot \mathbf{v} - [(\mathbf{w} \cdot \nabla) \mathbf{v}] \cdot \mathbf{u} d\mathbf{x} = \int_{\Omega} [(\mathbf{w} \cdot \nabla) \mathbf{u}] \cdot \mathbf{v} + \frac{1}{2} [(\nabla \cdot \mathbf{w}) \mathbf{u}] \cdot \mathbf{v} d\mathbf{x}, \\ c_{\widehat{B}}(\mathbf{H}, \mathbf{B}, \mathbf{v}) &= \int_{\Omega} \mathbf{H} \times \operatorname{curl} \mathbf{B} \cdot \mathbf{v} d\mathbf{x}, & c_{\widehat{B}}(\mathbf{u}, \mathbf{B}, \mathbf{H}) &= \int_{\Omega} (\mathbf{u} \times \mathbf{B}) \cdot \operatorname{curl} \mathbf{H} d\mathbf{x}. \end{aligned}$$

In addition, using the definition of $b(\cdot, \cdot, \cdot)$, it follows that

$$b(\mathbf{u}, \mathbf{v}, \mathbf{v}) = 0, \quad \mathbf{u} \in \mathcal{X}, \mathbf{v} \in H^1(\Omega)^d. \quad (7)$$

Applying $(\mathbf{B} \times \operatorname{curl} \mathbf{H}, \mathbf{v}) = (\mathbf{v} \times \mathbf{H}, \operatorname{curl} \mathbf{B})$, one finds that

$$c_{\widehat{B}}(\mathbf{B}, \mathbf{B}, \mathbf{u}) - c_{\widehat{B}}(\mathbf{u}, \mathbf{B}, \mathbf{B}) = 0, \quad \mathbf{u} \in \mathcal{X}, \mathbf{B} \in \mathcal{W}. \quad (8)$$

The bilinear term $d(\cdot, \cdot)$ satisfies the LBB condition [12, 28]:

$$\sup_{\mathbf{v} \in \mathcal{X}, \mathbf{v} \neq \mathbf{0}} \frac{d(\mathbf{v}, q)}{\|\mathbf{v}\|_1} \geq \widehat{\beta} \|q\|, \quad \forall q \in \mathcal{M}, \quad (9)$$

where $\widehat{\beta} > 0$ is constant depend on Ω .

A weak formulation for (2)-(4) may be written as follows: find $(\phi, \mu, \mathbf{u}, p, \mathbf{B})$ such that

$$(\partial_t \phi, \psi) + \epsilon a_\phi(\phi; \mu, \psi) + (\nabla \phi \cdot \mathbf{u}, \psi) = 0, \quad (10a)$$

$$\epsilon^{-1}(\phi^3 - \phi, \theta) + \epsilon(\nabla \phi, \nabla \theta) = (\mu, \theta), \quad (10b)$$

$$(\partial_t \mathbf{u}, \mathbf{v}) + a_f(\phi; \mathbf{u}, \mathbf{v}) + b(\mathbf{u}, \mathbf{u}, \mathbf{v}) + S_c c_{\widehat{B}}(\mathbf{B}, \mathbf{B}, \mathbf{v}) \quad (10c)$$

$$-d(\mathbf{v}, p) = \lambda(\mu \nabla \phi, \mathbf{v}),$$

$$d(\mathbf{u}, q) = 0, \quad (10d)$$

$$(\partial_t \mathbf{B}, \mathbf{H}) + a_B(\phi; \mathbf{B}, \mathbf{H}) - c_{\widehat{B}}(\mathbf{u}, \mathbf{B}, \mathbf{H}) = 0, \quad (10e)$$

for all $(\psi, \theta, \mathbf{v}, q, \mathbf{H}) \in \mathcal{Q} \times \mathcal{Q} \times \mathcal{X} \times \mathcal{M} \times \mathcal{W}$.

2.2 Wellposedness of solution

This subsection builds a well-posedness result of (10). For simplicity, we consider the following problem: find $(\phi, \mu, \mathbf{u}, \mathbf{B}) \in (\mathcal{Q}, \mathcal{Q}, \mathcal{X}_0, \mathcal{W})$, such that

$$(\partial_t \phi, \psi) + \epsilon a_\phi(\phi; \mu, \psi) + (\nabla \phi \cdot \mathbf{u}, \psi) = 0, \quad (11a)$$

$$\epsilon^{-1}(\phi^3 - \phi, \theta) + \epsilon(\nabla \phi, \nabla \theta) = (\mu, \theta), \quad (11b)$$

$$(\partial_t \mathbf{u}, \mathbf{v}) + a_f(\phi; \mathbf{u}, \mathbf{v}) + b(\mathbf{u}, \mathbf{u}, \mathbf{v}) + S_{cc\hat{B}}(\mathbf{B}, \mathbf{B}, \mathbf{v}) = \lambda(\mu \nabla \phi, \mathbf{v}), \quad (11c)$$

$$(\partial_t \mathbf{B}, \mathbf{H}) + a_B(\phi; \mathbf{B}, \mathbf{H}) - c_{\hat{B}}(\mathbf{u}, \mathbf{B}, \mathbf{H}) = 0, \quad (11d)$$

for all $(\psi, \theta, \mathbf{v}, \mathbf{H}) \in \mathcal{Q} \times \mathcal{Q} \times \mathcal{X}_0 \times \mathcal{W}$.

The next theorems give a well-posedness result for weak solution of problem (11). They were proved by the similar lines as in [20, 2, 17]. Thus, we skip the proofs of the following theorems.

Theorem 1 *Suppose that the initial conditions $\phi_0, \mathbf{u}_0, \mathbf{B}_0$ satisfy*

$$\phi_0 \in H^1(\Omega), \quad \mathbf{u}_0 \in L^2(\Omega)^d, \quad \mathbf{B}_0 \in L^2(\Omega)^d. \quad (12)$$

Furthermore, assume that the given functions κ, ν and η satisfy

$$\kappa, \nu, \eta \in C(\overline{\Omega} \times [0, T] \times R; R^+). \quad (13)$$

Then problem (11) has at least one solution $(\phi, \mu, \mathbf{u}, \mathbf{B})$ such that

$$\begin{aligned} \phi &\in L^\infty(0, T, H^1(\Omega)) \cap L^2(0, T, H^3), \\ \mathbf{u} &\in L^\infty(0, T, L^2(\Omega)^d) \cap L^2(0, T, \mathcal{X}), \\ \mathbf{B} &\in L^\infty(0, T, L^2(\Omega)^d) \cap L^2(0, T, \mathcal{W}) \\ \mu &\in L^2(0, T, \mathcal{Q}). \end{aligned} \quad (14)$$

Theorem 2 *Taking the place of (13) by*

$$\nu, \eta, \kappa \in C^{0,1}(\overline{\Omega} \times [0, T] \times R; R^+), \quad (15)$$

and retaining the other assumptions in Theorem 1. In addition, we suppose $(\phi, \mu, \mathbf{u}, \mathbf{B})$ satisfy

$$\phi \in L^2(0, T, W^{1,\infty}), \quad \mathbf{u} \in L^2(0, T, W^{1,\infty}(\Omega)^d), \quad \mathbf{B} \in L^2(0, T, W^{1,\infty}(\Omega)^d). \quad (16)$$

Then problem (11) has a unique solution $(\phi, \mu, \mathbf{u}, \mathbf{B})$.

3 Fully discrete scheme

In this section, we give a fully discrete scheme based on applying conforming finite element method in space and Euler semi-implicit discretization with a convex splitting method in time for (10) and obtain some unconditionally energy stability.

Let \mathbf{K}_h be a conforming, quasi-uniform family of triangulations of Ω into triangles when $d=2$ and tetrahedra when $d=3$, respectively. Furthermore, we introduce four finite element spaces $\mathcal{X}_h, \mathcal{M}_h, \mathcal{W}_h, \mathcal{Q}_h$ with $\mathcal{X}_h \subset \mathcal{X}, \mathcal{M}_h \subset \mathcal{M}, \mathcal{W}_h \subset \mathcal{W}, \mathcal{Q}_h \subset \mathcal{Q}$ as follows.

$$\begin{aligned}\mathcal{X}_h &= \{\mathbf{v} \in \mathbf{C}^0(\overline{\Omega})^d \cap \mathcal{X} : \mathbf{v}|_{\mathbb{K}} \in P_{r+1}(\mathbb{K})^d, \forall \mathbf{K} \in \mathbf{K}_h\}, \\ \mathcal{M}_h &= \{q \in C^0(\overline{\Omega}) \cap \mathcal{M} : q|_{\mathbb{K}} \in P_r(\mathbb{K}), \forall \mathbf{K} \in \mathbf{K}_h\}, \\ \mathcal{W}_h &= \{\mathbf{w} \in \mathbf{C}^0(\overline{\Omega})^d \cap \mathcal{W} : \mathbf{w}|_{\mathbb{K}} \in P_{r+1}(\mathbb{K})^d, \forall \mathbf{K} \in \mathbf{K}_h\}, \\ \mathcal{Q}_h &= \{\omega \in C^0(\overline{\Omega}) \cap \mathcal{Q} : \omega|_{\mathbb{K}} \in P_{r+1}(\mathbb{K}), \forall \mathbf{K} \in \mathbf{K}_h\} \\ \mathcal{X}_{0h} &= \{\mathbf{v} \in \mathcal{X}_h : d(\mathbf{v}, w) = 0, \forall w \in \mathcal{M}_h\}.\end{aligned}$$

As is noted that the $(\mathcal{X}_h, \mathcal{M}_h)$ is Taylor-Hood finite element pair. Therefore, the finite element pair $(\mathcal{X}_h, \mathcal{M}_h)$ satisfies the discrete LBB condition [28, 12]:

Assumption A1: The following discrete LBB condition holds:

$$\exists \beta_0 > 0, \quad \sup_{\mathbf{v}_h \in \mathcal{X}_h, \mathbf{v}_h \neq 0} \frac{d(\mathbf{v}_h, q_h)}{\|\nabla \mathbf{v}_h\|} \geq \beta_0 \|q_h\|, \quad \forall q_h \in \mathcal{M}_h.$$

Moreover, we suppose the finite element spaces satisfy following inverse inequality and finite element approximation properties:

Assumption A2: The following inverse inequality holds

$$\begin{aligned}\|v_h\|_{m,q} &\leq Ch^{l-m+d(\frac{1}{q}-\frac{1}{p})} \|v_h\|_{l,p}, \quad \forall v_h \in \mathcal{X}_h, \mathcal{W}_h \text{ or } \mathcal{Q}_h, \\ 0 \leq l \leq m \leq 1, \quad 1 \leq p \leq q \leq \infty.\end{aligned}$$

Assumption A3: There exists $k \geq 1$, such that for all $1 \leq l \leq k$,

$$\begin{aligned}\inf_{\mathbf{v}_h \in \mathcal{X}_h} [\|\mathbf{v} - \mathbf{v}_h\| + h\|\nabla(\mathbf{v} - \mathbf{v}_h)\|] &\leq Ch^{l+1} \|\mathbf{v}\|_{l+1}, \quad \forall \mathbf{v} \in H^{l+1}(\Omega)^d, \\ \inf_{\mathbf{B}_h \in \mathcal{W}_h} [\|\mathbf{B} - \mathbf{B}_h\| + h\|\nabla(\mathbf{B} - \mathbf{B}_h)\|] &\leq Ch^{l+1} \|\mathbf{B}\|_{l+1}, \quad \forall \mathbf{B} \in H^{l+1}(\Omega)^d, \\ \inf_{\varphi_h \in \mathcal{Q}_h} [\|\varphi - \varphi_h\| + h\|\nabla(\varphi - \varphi_h)\|] &\leq Ch^{l+1} \|\varphi\|_{l+1}, \quad \forall \varphi \in H^{l+1}(\Omega), \\ \inf_{q_h \in \mathcal{M}_h} \|q - q_h\| &\leq Ch^l \|q\|_l, \quad \forall q \in H^l(\Omega).\end{aligned}$$

Let N be a positive integer and $0 = t_0 < t_1 < \dots < t_N = T$ be a uniform partition of $[0, T]$, with $\Delta t = t_i - t_{i-1}$ for $i = 1, \dots, N$. Let us denote by ϕ^n the value $\phi(t_n)$ at the time t_n , and denote $\delta_t \phi := \frac{\phi^n - \phi^{n-1}}{\Delta t}$. Denote $\kappa^n := \kappa(\mathbf{x}, t_n, \phi)$ $\nu^n := \nu(\mathbf{x}, t_n, \phi)$ and $\eta^n := \eta(\mathbf{x}, t_n, \phi)$.

With above preparation, we give the fully discrete scheme of problem (10).

Scheme 3.1: Given $(\phi_h^0, \mathbf{u}_h^0, \mathbf{B}_h^0) \in \mathcal{Q}_h \times \mathcal{X}_h \times \mathcal{W}_h$, find $(\phi_h^n, \mu_h^n, \mathbf{u}_h^n, p_h^n, \mathbf{B}_h^n) \in \mathcal{Q}_h \times \mathcal{Q}_h \times \mathcal{X}_h \times \mathcal{M}_h \times \mathcal{W}_h$, such that

$$(\delta_t \phi_h^n, \psi_h) + \epsilon a_\phi(\phi_h^{n-1}; \mu_h^n, \psi_h) + (\nabla \phi_h^{n-1} \cdot \mathbf{u}_h^n, \psi_h) = 0, \quad (17a)$$

$$\epsilon^{-1}((\phi_h^n)^3 - \phi_h^{n-1}, \theta_h) + \epsilon(\nabla \phi_h^n, \nabla \theta_h) = (\mu_h^n, \theta_h), \quad (17b)$$

$$(\delta_t \mathbf{u}_h^n, \mathbf{v}_h) + a_f(\nu^n(\phi_h^{n-1}); \mathbf{u}_h^n, \mathbf{v}_h) + b_f(\mathbf{u}_h^{n-1}, \mathbf{u}_h^n, \mathbf{v}_h) \quad (17c)$$

$$+ S_c b_{\hat{B}}(\mathbf{B}_h^{n-1}, \mathbf{B}_h^n, \mathbf{v}_h) - d(\mathbf{v}_h, p_h^n) = \lambda(\mu_h^n \nabla \phi_h^{n-1}, \mathbf{v}_h),$$

$$d(\mathbf{u}_h^n, q_h) = 0, \quad (17d)$$

$$(\delta_t \mathbf{B}_h^n, \mathbf{H}_h) + a_B(\eta^n(\phi_h^{n-1}); \mathbf{B}_h^n, \mathbf{H}_h) - c_{\bar{B}}(\mathbf{u}_h^n, \mathbf{B}_h^{n-1}, \mathbf{H}_h) = 0, \quad (17e)$$

for all $(\psi_h, \theta_h, \mathbf{v}_h, q_h, \mathbf{H}_h) \in \mathcal{Q}_h \times \mathcal{Q}_h \times \mathcal{X}_h \times \mathcal{M}_h \times \mathcal{W}_h$.

In addition, the given functions κ, ν, η satisfy following assumption.

Assumption A4: For all $(\mathbf{x}, t, \phi) \in \overline{\Omega} \times (0, T] \times R$, the given functions κ, ν, η satisfy

$$0 < \kappa_1 \leq \kappa(\mathbf{x}, t, \phi) \leq \kappa_2,$$

$$0 < \nu_1 \leq \nu(\mathbf{x}, t, \phi) \leq \nu_2,$$

$$0 < \eta_1 \leq \eta(\mathbf{x}, t, \phi) \leq \eta_2.$$

Denote $\dot{S}_h = \mathcal{Q}_h \cap \mathcal{M}_h$. Let $\hat{\kappa} \in C^{0,1}(\overline{\Omega}; R^+)$ satisfy **Assumption A4**, we define the operator $T_h : \dot{S}_h \rightarrow \dot{S}_h$ by the following weak problem: given $\sigma \in \dot{S}_h$, find $T_h(\sigma) \in \dot{S}_h$ such that

$$a_\phi(\hat{\kappa}; T_h(\sigma), \varrho) = (\sigma, \varrho), \quad \forall \varrho \in \dot{S}_h. \quad (18)$$

As in the work [5], we have similar results.

Lemma 1 Let $\zeta, \vartheta \in \dot{S}_h$ and denote

$$(\zeta, \vartheta)_{-1,h} := a_\phi(\hat{\kappa}; T_h(\zeta), T_h(\vartheta)) = (\zeta, T_h(\vartheta)) = (T_h(\zeta), \vartheta),$$

and define the following negative norm:

$$\|\zeta\|_{-1,h} := \sqrt{(\zeta, \zeta)_{-1,h}} = \sup_{0 \neq \theta \in \dot{S}_h} \frac{(\zeta, \theta)}{\|\nabla \theta\|}.$$

Therefore, $\forall \vartheta \in \mathcal{X}_h$ and $\forall \zeta \in \dot{S}_h$, we have

$$|(\zeta, \vartheta)| \leq \|\zeta\|_{-1,h} \|\nabla \vartheta\|.$$

Then the following Poincaré inequality holds:

$$\|\vartheta\|_{-1,h} \leq C \|\vartheta\|, \quad \forall \vartheta \in \dot{S}_h.$$

We now are ready to state and prove unconditionally energy stability for the fully discrete scheme (17).

Theorem 3 Suppose **Assumptions A1-A4** hold. Then the finite element approximate solution $(\phi_h^n, \mu_h^n, \mathbf{u}_h^n, p_h^n, \mathbf{B}_h^n)$ of problem (17) satisfy the following discrete energy law:

$$\begin{aligned} E(\phi_h^n, \mathbf{u}_h^n, \mathbf{B}_h^n) + \Delta t \sum_{m=1}^n [\kappa_1 \epsilon \|\nabla \mu_h^m\|^2 + \Delta t c_0 \nu_1 \|\nabla \mathbf{u}_h^m\|^2 + S_c \Delta t c_1 \eta_1 \|\nabla \mathbf{B}_h^m\|^2] \\ + \Delta t^2 \sum_{m=1}^n \left[\frac{\epsilon}{2} \|\nabla \delta_t \phi_h^m\|^2 + \frac{1}{2\lambda} \|\delta_t \mathbf{u}_h^m\|^2 + \frac{S}{2} \|\delta_t \mathbf{B}_h^m\|^2 \right. \\ \left. + \frac{1}{4\epsilon} \|\delta_t (\phi_h^m)^2\|^2 + \frac{1}{2\epsilon} \|\phi_h^m \delta_t \phi_h^m\|^2 + \frac{1}{2\epsilon} \|\delta_t \phi_h^m\|^2 \right] \leq E(\phi_h^0, \mathbf{u}_h^0, \mathbf{B}_h^0). \end{aligned}$$

Proof: Taking $\psi_h = \mu_h^n$ in (17a), $\theta_h = \delta_t \phi_h^n$ in (17b), $\mathbf{v}_h = \mathbf{u}_h^n$ in (17c), $q_h = p_h^n$ in (17d), $\mathbf{H}_h = \mathbf{B}_h^n$ in (17e), one finds that

$$\begin{aligned} (\delta_t \phi_h^n, \mu_h^n) + \epsilon a_\phi(\phi_h^{n-1}; \mu_h^n, \mu_h^n) + (\nabla \phi_h^{n-1} \cdot \mathbf{u}_h^n, \mu_h^n) &= 0, \\ \epsilon^{-1}((\phi_h^n)^3 - \phi_h^{n-1}, \delta_t \phi_h^n) + \epsilon(\nabla \phi_h^n, \nabla \delta_t \phi_h^n) &= (\mu_h^n, \delta_t \phi_h^n), \\ (\delta_t \mathbf{u}_h^n, \mathbf{u}_h^n) + a_f(\nu^n(\phi_h^{n-1}); \mathbf{u}_h^n, \mathbf{u}_h^n) + b_f(\mathbf{u}_h^{n-1}, \mathbf{u}_h^n, \mathbf{u}_h^n) \\ + S_c b_{\widehat{B}}(\mathbf{B}_h^{n-1}, \mathbf{B}_h^n, \mathbf{u}_h^n) - d(\mathbf{u}_h^n, p_h^n) &= \lambda(\mu_h^n \nabla \phi_h^{n-1}, \mathbf{u}_h^n), \\ d(\mathbf{u}_h^n, p_h^n) &= 0, \\ (\delta_t \mathbf{B}_h^n, \mathbf{B}_h^n) + a_B(\eta^n(\phi_h^{n-1}); \mathbf{B}_h^n, \mathbf{B}_h^n) - c_{\widehat{B}}(\mathbf{u}_h^n, \mathbf{B}_h^{n-1}, \mathbf{B}_h^n) &= 0. \end{aligned}$$

Using the elementary identity

$$2a(a - b) = a^2 - b^2 + (a - b)^2, \quad \text{for all } a, b \in \mathbf{R}^d.$$

It follows that

$$\begin{aligned} \epsilon(\nabla \phi_h^n, \nabla \delta_t \phi_h^n) &= \frac{\epsilon}{2} [\delta_t \|\nabla \phi_h^n\|^2 + \Delta t \|\nabla \delta_t \phi_h^n\|^2], \\ \epsilon^{-1}((\phi_h^n)^3 - \phi_h^{n-1}, \delta_t \phi_h^n) &= \frac{\epsilon^{-1}}{4} \delta_t \|(\phi_h^n)^2 - 1\|^2 \\ &\quad + \frac{\epsilon^{-1} \Delta t}{4} [\|\delta_t (\phi_h^n)^2\|^2 + 2\|\phi_h^n \delta_t \phi_h^n\|^2 + 2\|\delta_t \phi_h^n\|^2], \\ (\delta_t \mathbf{u}_h^n, \mathbf{u}_h^n) &= \frac{1}{2} [\delta_t \|\mathbf{u}_h^n\|^2 + \Delta t \|\delta_t \mathbf{u}_h^n\|^2], \\ (\delta_t \mathbf{B}_h^n, \mathbf{B}_h^n) &= \frac{1}{2} [\delta_t \|\mathbf{B}_h^n\|^2 + \Delta t \|\delta_t \mathbf{B}_h^n\|^2]. \end{aligned}$$

Then using the operator $\Delta t \sum_{n=1}^m$ to the combined equations, the desired result is derived. The proof is completed. \square

The following Theorem give some bounds for the fully discrete solution. It is very important to derive optimal error estimates of problem (17).

Theorem 4 [5, 18, 33] Suppose **Assumptions A1-A4** hold. Then the fully discrete solution $(\phi_h^n, \mu_h^n, \mathbf{u}_h^n, p_h^n, \mathbf{B}_h^n)$ of (17) satisfy the following bounds:

$$\max_{0 \leq m \leq n} [\|\mathbf{u}_h^n\|^2 + \|\mathbf{B}_h^n\|^2 + \|\nabla \phi_h^n\|^2 + \|(\phi_h^n)^2 - 1\|^2 + \|\phi_h^n\|^4 + \|\phi_h^n\|^2 + \|\phi_h^n\|_{H^1}^2] \leq C, \quad (19)$$

$$\Delta t \sum_{m=1}^n [\|\nabla \mathbf{u}_h^m\|^2 + \|\nabla \mathbf{B}_h^m\|^2 + \|\nabla \mu_h^m\|^2] \leq C, \quad (20)$$

$$\sum_{m=1}^n [\|\nabla(\phi_h^n - \phi_h^{n-1})\|^2 + \|\phi_h^n - \phi_h^{n-1}\|^2 + \|\phi_h^n(\phi_h^n - \phi_h^{n-1})\|^2 + \|(\phi_h^n)^2 - (\phi_h^{n-1})^2\|^2 + \|\phi_h^n - \phi_h^{n-1}\|_{-1,h}^2 + \|\mathbf{u}_h^n - \mathbf{u}_h^{n-1}\|^2 + \|\mathbf{B}_h^n - \mathbf{B}_h^{n-1}\|^2] \leq C, \quad (21)$$

$$\Delta t \sum_{m=1}^n [\|\delta_t \phi_h^n\|_{H^{-1}}^2 + \|\delta_t \phi_h^n\|_{-1,h}^2 + \|\nabla_h \phi_h^n\|^2 + \|\mu_h^n\|^2 + \|\mu_h^n\|_{L^\infty}^{\frac{4(6-d)}{d}} + \|\delta_t \phi_h^n\|^2] \leq C, \quad (22)$$

$$\max_{0 \leq m \leq n} [\|\mu_h^n\|^2 + \|\nabla_h \phi_h^n\|^2 + \|\mu_h^n\|_{L^\infty}^{\frac{4(6-d)}{d}}] \leq C, \quad (23)$$

$$\Delta t^2 \sum_{m=1}^n [\|\delta_t \mathbf{u}_h^n\|^2 + \|\delta_t \mathbf{B}_h^n\|^2] \leq C. \quad (24)$$

4 Error analysis

In this section we drive some optimal error estimates of the fully discrete scheme (17). Therefore, we assume that the solution $(\phi, \mu, \mathbf{u}, p, \mathbf{B})$ has the following regularity.

Assumption A5: The weak solution $(\phi, \mu, \mathbf{u}, p, \mathbf{B})$ of Cahn-Hilliard-Magneto-hydrodynamics (2)-(4) is sufficiently smooth such that

$$\begin{aligned} \partial_{tt} \phi &\in L^\infty(0, T; H^1(\Omega)), & \phi, \partial_t \phi &\in L^\infty(0, T; H^{k+2}(\Omega) \cap W^{1,\infty}(\Omega)), \\ \partial_{tt} \mathbf{u} &\in L^\infty(0, T; H^1(\Omega)^d), & \mathbf{u}, \partial_t \mathbf{u} &\in L^\infty(0, T; H^{k+2}(\Omega)^d \cap W^{1,\infty}(\Omega)^d), \\ \partial_{tt} \mathbf{B} &\in L^\infty(0, T; H^1(\Omega)^d), & \mathbf{B}, \partial_t \mathbf{B} &\in L^\infty(0, T; H^{k+2}(\Omega)^d \cap W^{1,\infty}(\Omega)^d), \\ \mu &\in L^\infty(0, T; H^{k+2}(\Omega)), & p, \partial_t p &\in L^\infty(0, T; H^{k+1}(\Omega)). \end{aligned}$$

We need the following technical lemmas to obtain the rate of convergence of the fully discrete scheme (17).

Lemma 2 [5] Assume $\varpi \in H^1(\Omega)$, and $v \in \dot{S}_h$, Then there exists $C > 0$, independent of h , such that

$$|(\varpi, v)| \leq C \|\nabla \varpi\| \|v\|_{-1,h}. \quad (25)$$

Lemma 3 [5] *Assume that $(\phi, \mu, \mathbf{u}, p, \mathbf{B})$, $(\phi_h^n, \mu_h^n, \mathbf{u}_h^n, p_h^n, \mathbf{B}_h^n)$ are weak solution to problem (10) and (17), respectively. Then for any h , $\Delta t > 0$, the following inequality holds*

$$\|\nabla(\phi^3 - (\phi_h^n)^3)\| \leq C\|\nabla(\phi - \phi_h^n)\|. \quad (26)$$

Let $\hat{\nu}, \hat{\eta}, \hat{\kappa} \in C^{0,1}(\overline{\Omega}; R^+)$ satisfy **Assumption A4**, we introduce the following four projections. Let $(\mathbf{u}, p) \in \mathcal{X} \times \mathcal{M}$, $\mathbf{B} \in \mathcal{W}$, $\theta, \phi \in \mathcal{Q}$, we introduce Stokes projection $(R_h \mathbf{u}, J_h p) \in \mathcal{X}_h \times \mathcal{M}_h$ as the solution of the weak problem as follows

$$a_f(\hat{\nu}; \mathbf{u} - R_h \mathbf{u}, \mathbf{v}) + d(\mathbf{v}, p - J_h p) = 0, \quad \forall \mathbf{v} \in \mathcal{X}_h, \quad (27)$$

$$d(\mathbf{u} - R_h \mathbf{u}, q) = 0, \quad \forall q \in \mathcal{M}_h. \quad (28)$$

We have the following approximation result

$$\|\mathbf{u} - R_h \mathbf{u}\| + h(\|\nabla(\mathbf{u} - R_h \mathbf{u})\| + \|p - J_h p\|) \leq Ch^{k+2}(\|\mathbf{u}\|_{k+2} + \|p\|_{k+1}). \quad (29)$$

Similarly, Maxwell projection $R_{mh} \mathbf{B} \in \mathcal{W}_h$ satisfying

$$a_B(\hat{\eta}; \mathbf{B} - R_{mh} \mathbf{B}, \mathbf{H}) = 0, \quad \forall \mathbf{H} \in \mathcal{W}_h, \quad (30)$$

$$\|\mathbf{B} - R_{mh} \mathbf{B}\| + h\|\nabla(\mathbf{B} - R_{mh} \mathbf{B})\| \leq Ch^{k+2}\|\mathbf{B}\|_{k+2}, \quad (31)$$

Ritz projections $r_h \theta \in \mathcal{Q}_h$ and $\hat{r}_h \theta \in \mathcal{Q}_h$ satisfying

$$a_\phi(\hat{\kappa}; \theta - r_h \theta, \psi) = 0, \quad \forall \psi \in \mathcal{Q}_h, \quad (32)$$

$$\|\theta - r_h \theta\| + h\|\nabla(\theta - r_h \theta)\| \leq Ch^{k+2}\|\theta\|_{k+2}, \quad (33)$$

$$(\nabla(\phi - \hat{r}_h \phi), \nabla \psi) = 0, \quad \forall \psi \in \mathcal{Q}_h, \quad (34)$$

$$\|\phi - \hat{r}_h \phi\| + h\|\nabla(\phi - \hat{r}_h \phi)\| \leq Ch^{k+2}\|\phi\|_{k+2}. \quad (35)$$

For convenience, let us define, for every $n \geq 0$,

$$\begin{aligned} e_\phi^n &:= \phi^n - \phi_h^n, & e_\mu^n &:= \mu^n - \mu_h^n, \\ e_{\mathbf{u}}^n &:= \mathbf{u}^n - \mathbf{u}_h^n, & e_p^n &:= p^n - p_h^n, \\ e_{\mathbf{B}}^n &:= \mathbf{B}^n - \mathbf{B}_h^n. \end{aligned}$$

Which gives

$$\begin{aligned} e_\phi^n &= \eta_\phi^n - \epsilon_\phi^n, & e_\mu^n &= \eta_\mu^n - \epsilon_\mu^n, \\ e_{\mathbf{u}}^n &= \eta_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^n, & e_p^n &= \eta_p^n - \epsilon_p^n, \\ e_{\mathbf{B}}^n &= \eta_{\mathbf{B}}^n - \epsilon_{\mathbf{B}}^n, \end{aligned}$$

with

$$\begin{aligned}
\eta_\phi^n &:= \phi^n - r_h \phi^n \in \mathcal{Q}, & \epsilon_\phi^n &:= \phi_h^n - r_h \phi^n \in \mathcal{Q}_h, \\
\eta_\mu^n &:= \mu^n - \hat{r}_h \mu^n \in \mathcal{Q}, & \epsilon_\mu^n &:= \mu_h^n - \hat{r}_h \mu^n \in \mathcal{Q}_h, \\
\eta_{\mathbf{u}}^n &:= \mathbf{u}^n - R_h \mathbf{u}^n \in \mathcal{X}, & \epsilon_{\mathbf{u}}^n &:= \mathbf{u}_h^n - R_h \mathbf{u}^n \in \mathcal{X}_h, \\
\eta_p^n &:= p^n - J_h p^n \in \mathcal{M}, & \epsilon_p^n &:= p_h^n - J_h p^n \in \mathcal{M}_h, \\
\eta_{\mathbf{B}}^n &:= \mathbf{B}^n - R_{mh} \mathbf{B}^n \in \mathcal{W}, & \epsilon_{\mathbf{B}}^n &:= \mathbf{B}_h^n - R_{mh} \mathbf{B}^n \in \mathcal{W}_h.
\end{aligned}$$

We are now in a position to give and derive the first main theorem of this section for the concentration field, the chemical potential, the velocity field and the magnetic field.

Theorem 5 *Suppose that **Assumption A1-A5** hold with two positive constants $h_0, \Delta t_0$, and suppose the scheme (17) is initialized such that*

$$\|\phi^0 - \phi_h^0\| + \|\mathbf{u}^0 - \mathbf{u}_h^0\| + \|\mathbf{B}^0 - \mathbf{B}_h^0\| \leq C(\Delta t + h^{k+1}). \quad (36)$$

For $h \in (0, h_0]$ and $\Delta t \in (0, \Delta t_0]$, the finite element approximate solution $(\phi_h^n, \mu_h^n, \mathbf{u}_h^n, p_h^n, \mathbf{B}_h^n)$ of (17) satisfy the following error estimates:

$$\begin{aligned}
\max_{1 \leq n \leq N} \|\nabla(\phi(t_n) - \phi_h^n)\|^2 + \Delta t \sum_{n=1}^N \|\nabla(\mu(t_n) - \mu_h^n)\|^2 &\leq C(\Delta t^2 + h^{2k+2}), \\
\max_{1 \leq n \leq N} \|\mathbf{u}(t_n) - \mathbf{u}_h^n\|^2 + \Delta t \sum_{n=1}^N \|\nabla(\mathbf{u}(t_n) - \mathbf{u}_h^n)\|^2 &\leq C(\Delta t^2 + h^{2k+2}), \\
\max_{1 \leq n \leq N} \|\mathbf{B}(t_n) - \mathbf{B}_h^n\|^2 + \Delta t \sum_{n=1}^N \|\nabla(\mathbf{B}(t_n) - \mathbf{B}_h^n)\|^2 &\leq C(\Delta t^2 + h^{2k+2}).
\end{aligned}$$

Proof: Appying (10), (17), (27)-(28), (30), (32) and (34), we can obtain the following error equations:

$$(\delta_t \epsilon_\phi^n, \psi_h) + \epsilon a_\phi(\kappa^n(\phi_h^{n-1}); \epsilon_\mu^n, \psi_h) = (\Lambda_h^n, \psi_h) + (R_\phi^n, \psi_h), \quad (37a)$$

$$\epsilon(\nabla \epsilon_\phi^n, \nabla \theta_h) - (\epsilon_\mu^n, \theta_h) = (\hat{\Lambda}_h^n, \theta_h), \quad (37b)$$

$$(\delta_t \epsilon_{\mathbf{u}}^n, \mathbf{v}_h) + a_f(\nu^n(\phi_h^{n-1}); \epsilon_{\mathbf{u}}^n, \mathbf{v}_h) - d(\mathbf{v}_h, \epsilon_p^n) = (\Phi_h^n, \mathbf{v}_h) + (R_{\mathbf{u}}^n, \mathbf{v}_h), \quad (37c)$$

$$d(\epsilon_{\mathbf{u}}^n, q_h) = 0, \quad (37d)$$

$$S_c(\delta_t \epsilon_{\mathbf{B}}^n, \mathbf{H}_h) + S_c a_B(\eta^n(\phi_h^{n-1}); \epsilon_{\mathbf{B}}^n, \mathbf{H}_h) = (\hat{\Phi}_h^n, \mathbf{H}_h) + S_c(R_{\mathbf{B}}^n, \mathbf{H}_h). \quad (37e)$$

where (R_ϕ^n, ψ_h) , $(R_{\mathbf{u}}^n, \mathbf{v}_h)$, $(R_{\mathbf{B}}^n, \mathbf{H}_h)$, (Λ_h^n, ψ_h) , $(\hat{\Lambda}_h^n, \theta_h)$, (Φ_h^n, \mathbf{v}_h) and $(\hat{\Phi}_h^n, \mathbf{H}_h)$ are denoted by

$$(R_\phi^n, \psi_h) := (\partial_t \phi^n - \delta_t r_h \phi^n, \psi_h), \quad (38)$$

$$(R_{\mathbf{u}}^n, \mathbf{v}_h) := (\partial_t \mathbf{u}^n - \delta_t R_h \mathbf{u}^n, \mathbf{v}_h), \quad (39)$$

$$(R_{\mathbf{B}}^n, \mathbf{H}_h) := (\partial_t \mathbf{B}^n - \delta_t R_{mh} \mathbf{B}^n, \mathbf{H}_h), \quad (40)$$

$$\begin{aligned} (R_h^n, \psi_h) &:= a_\phi(\kappa^n(\phi^n); \mu^n, \psi_h) - a_\phi(\kappa^n(\phi_h^{n-1}); \mu^n, \psi_h) \\ &\quad + (\mathbf{u}^n \cdot \nabla \phi^n, \psi_h) - (\mathbf{u}_h^n \cdot \nabla \phi_h^{n-1}, \psi_h), \end{aligned} \quad (41)$$

$$\begin{aligned} (\tilde{\Lambda}_h^n, \theta_h) &:= (\eta_\mu^n, \theta_h) - \frac{\Delta t}{\epsilon} (\delta_t \phi^n, \theta_h) \\ &\quad - \epsilon^{-1} (\phi^{n-1} - \phi_h^{n-1}, \theta_h) + \epsilon^{-1} ((\phi^n)^3 - (\phi_h^n)^3, \theta_h), \end{aligned} \quad (42)$$

$$\begin{aligned} (\Phi_h^n, \mathbf{v}_h) &:= a_f(\nu^n(\phi^n); \mathbf{u}^n, \mathbf{v}_h) - a_f(\nu^n(\phi_h^{n-1}); \mathbf{u}^n, \mathbf{v}_h) \\ &\quad + b(\mathbf{u}^n, \mathbf{u}^n, \mathbf{v}_h) - b(\mathbf{u}_h^{n-1}, \mathbf{u}_h^n, \mathbf{v}_h) \\ &\quad + S_c c_{\hat{B}}(\mathbf{B}^n, \mathbf{B}^n, \mathbf{v}_h) - S_c c_{\hat{B}}(\mathbf{B}_h^{n-1}, \mathbf{B}_h^n, \mathbf{v}_h) \\ &\quad + \lambda(\mu_h^n \nabla \phi_h^{n-1}, \mathbf{v}_h) - \lambda(\mu^n \nabla \phi^n, \mathbf{v}_h), \end{aligned} \quad (43)$$

$$\begin{aligned} (\hat{\Phi}_h^n, \mathbf{H}_h) &:= S_c a_B(\eta^n(\phi^n); \mathbf{B}^n, \mathbf{H}_h) - S_c a_B(\eta^n(\phi_h^{n-1}); \mathbf{B}^n, \mathbf{H}_h) \\ &\quad + S_c c_{\hat{B}}(\mathbf{u}_h^n, \mathbf{B}_h^{n-1}, \mathbf{H}_h) - S_c c_{\hat{B}}(\mathbf{u}^n, \mathbf{B}^n, \mathbf{H}_h). \end{aligned} \quad (44)$$

Setting $\psi_h = \epsilon_\phi^n \in \mathcal{Q}_h$, $\theta_h = \kappa_1 \epsilon_\mu^n \in \mathcal{Q}_h$ in (37a)-(37b), respectively, we have

$$(\delta_t \epsilon_\phi^n, \epsilon_\phi^n) + \epsilon \kappa_1 (\nabla \epsilon_\mu^n, \nabla \epsilon_\phi^n) \leq (\Lambda_h^n, \epsilon_\phi^n) + (R_\phi^n, \epsilon_\phi^n), \quad (45a)$$

$$-\kappa_1 \epsilon (\nabla \epsilon_\phi^n, \nabla \epsilon_\mu^n) + \kappa_1 (\epsilon_\mu^n, \epsilon_\mu^n) = -\kappa_1 (\hat{\Lambda}_h^n, \epsilon_\mu^n). \quad (45b)$$

Setting $\psi_h = \lambda \epsilon_\mu^n \in \mathcal{Q}_h$, $\theta_h = \lambda \delta_t \epsilon_\phi^n \in \mathcal{Q}_h$, $\mathbf{v}_h = \epsilon_{\mathbf{u}}^n \in \mathcal{X}_h$, $\mathbf{H}_h = \epsilon_{\mathbf{B}}^n \in \mathcal{W}_h$ in (37), respectively, it follows that

$$\lambda (\delta_t \epsilon_\phi^n, \epsilon_\mu^n) + \lambda \epsilon \kappa_1 \|\nabla \epsilon_\mu^n\|^2 \leq \lambda (\Lambda_h^n, \epsilon_\mu^n) + \lambda (R_\phi^n, \epsilon_\mu^n), \quad (46a)$$

$$\epsilon \lambda (\nabla \epsilon_\phi^n, \nabla \delta_t \epsilon_\phi^n) - \lambda (\epsilon_\mu^n, \delta_t \epsilon_\phi^n) = \lambda (\hat{\Lambda}_h^n, \delta_t \epsilon_\phi^n), \quad (46b)$$

$$(\delta_t \epsilon_{\mathbf{u}}^n, \epsilon_{\mathbf{u}}^n) + c_0 \nu_1 \|\nabla \epsilon_{\mathbf{u}}^n\|^2 \leq (\Phi_h^n, \epsilon_{\mathbf{u}}^n) + (R_{\mathbf{u}}^n, \epsilon_{\mathbf{u}}^n), \quad (46c)$$

$$S_c (\delta_t \epsilon_{\mathbf{B}}^n, \epsilon_{\mathbf{B}}^n) + c_1 \eta_1 S_c \|\nabla \epsilon_{\mathbf{B}}^n\|^2 \leq (\hat{\Phi}_h^n, \epsilon_{\mathbf{B}}^n) + S_c (R_{\mathbf{B}}^n, \epsilon_{\mathbf{B}}^n). \quad (46d)$$

Combining (45) and (46), we obtain

$$\begin{aligned}
& \frac{1}{2} (\|\epsilon_\phi^n\|^2 - \|\epsilon_\phi^{n-1}\|^2 + \|\epsilon_\phi^n - \epsilon_\phi^{n-1}\|^2) \\
& + \frac{\epsilon\lambda}{2} (\|\nabla \epsilon_\phi^n\|^2 - \|\nabla \epsilon_\phi^{n-1}\|^2 + \|\nabla \epsilon_\phi^n - \nabla \epsilon_\phi^{n-1}\|^2) \\
& + \frac{1}{2} (\|\epsilon_{\mathbf{u}}^n\|^2 - \|\epsilon_{\mathbf{u}}^{n-1}\|^2 + \|\epsilon_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^{n-1}\|^2) \\
& + \frac{S_c}{2} (\|\epsilon_{\mathbf{B}}^n\|^2 - \|\epsilon_{\mathbf{B}}^{n-1}\|^2 + \|\epsilon_{\mathbf{B}}^n - \epsilon_{\mathbf{B}}^{n-1}\|^2) + \Delta t \lambda \epsilon \kappa_1 \|\nabla \epsilon_\mu^n\|^2 \\
& + \Delta t \kappa_1 \|\epsilon_\mu^n\|^2 + \Delta t c_0 \nu_1 \|\nabla \epsilon_{\mathbf{u}}^n\|^2 + \Delta t c_1 \eta_1 S_c \|\nabla \epsilon_{\mathbf{B}}^n\|^2 \\
& \leq \Delta t \lambda (R_\phi^n, \epsilon_\mu^n) + \Delta t (R_\phi^n, \epsilon_\phi^n) + \Delta t (R_{\mathbf{u}}^n, \epsilon_{\mathbf{u}}^n) \\
& + \Delta t S_c (R_{\mathbf{B}}^n, \epsilon_{\mathbf{B}}^n) + \Delta t \lambda (\Lambda_h^n, \epsilon_\mu^n) + \Delta t (\Lambda_h^n, \epsilon_\phi^n) \\
& + \Delta t \lambda (\hat{\Lambda}_h^n, \delta_t \epsilon_\phi^n) - \Delta t \kappa_1 (\hat{\Lambda}_h^n, \epsilon_\mu^n) \\
& + \Delta t (\Phi_h^n, \epsilon_{\mathbf{u}}^n) + \Delta t (\hat{\Phi}_h^n, \epsilon_{\mathbf{B}}^n) \\
& = \sum_{i=1}^{10} \mathcal{Y}_i.
\end{aligned} \tag{47}$$

We now bound the terms on the RHS of (47). Define time dependent spatial mass average as follows

$$\bar{\epsilon}_\mu^n := |\Omega|^{-1} (\epsilon_\mu^n, 1), \quad 1 \leq n \leq N.$$

For terms $\mathcal{Y}_1 - \mathcal{Y}_4$, using (6), Taylor's theorem, **Assumption 5** and Young's inequality, one finds that

$$\mathcal{Y}_1 = \Delta t \lambda (R_\phi^n, \epsilon_\mu^n - \bar{\epsilon}_\mu^n) \tag{48}$$

$$\begin{aligned}
& \leq \Delta t \lambda \|\partial_t \phi^n - \delta_t r_h \phi^n\| \|\nabla \epsilon_\mu^n\| \\
& \leq C \Delta t \left\{ \Delta t \int_{t-\Delta t}^t \|\partial_{ss}^2 \phi(s)\|_{L^2}^2 ds \right. \\
& \quad \left. + \frac{h^{k+1}}{\sqrt{\Delta t}} \int_{t-\Delta t}^t \|\partial_s \phi(s)\|_{H^{k+1}}^2 ds \right\}^{\frac{1}{2}} \|\nabla \epsilon_\mu^n\| \\
& \leq C \Delta t (\Delta t^2 + h^{2k+2}) + \frac{\kappa_1 \lambda \Delta t}{8} \|\nabla \epsilon_\mu^n\|^2,
\end{aligned}$$

$$\begin{aligned}
\mathcal{Y}_2 & \leq C \Delta t \left\{ \Delta t \int_{t-\Delta t}^t \|\partial_{ss}^2 \phi(s)\|_{L^2}^2 ds \right. \\
& \quad \left. + \frac{h^{k+1}}{\sqrt{\Delta t}} \int_{t-\Delta t}^t \|\partial_s \phi(s)\|_{H^{k+1}}^2 ds \right\}^{\frac{1}{2}} \|\epsilon_\phi^n\| \\
& \leq C \Delta t (\Delta t^2 + h^{2k+2}) + C \Delta t \|\epsilon_\phi^n\|^2.
\end{aligned} \tag{49}$$

Similarly we have

$$\begin{aligned}
\mathcal{I}_3 &\leq C\Delta t \|\nabla \epsilon_{\mathbf{u}}^n\| \|\partial_t \mathbf{u}^n - \delta_t R_h \mathbf{u}^n\| \\
&\leq C\Delta t \left\{ \Delta t \int_{t-\Delta t}^t \|\partial_{ss}^2 \mathbf{u}(s)\|_{L^2}^2 ds \right. \\
&\quad \left. + \frac{h^{k+1}}{\sqrt{\Delta t}} \int_{t-\Delta t}^t \|(\partial_s \mathbf{u}(s), \partial_s p(s))\|_{H^{k+1} \times H^k}^2 ds \right\}^{\frac{1}{2}} \|\nabla \epsilon_{\mathbf{u}}^n\|^2 \\
&\leq C\Delta t (\Delta t^2 + h^{2k+2}) + \frac{c_0 \nu_1 \Delta t}{8} \|\nabla \epsilon_{\mathbf{u}}^n\|^2,
\end{aligned} \tag{50}$$

and

$$\begin{aligned}
\mathcal{I}_4 &\leq C\Delta t \|\nabla \epsilon_{\mathbf{B}}^n\| \|\partial_t \mathbf{B}^n - \delta_t R_{mh} \mathbf{B}^n\| \\
&\leq C\Delta t \left\{ \Delta t \int_{t-\Delta t}^t \|\partial_{ss}^2 \mathbf{B}\|_{L^2}^2 ds \right. \\
&\quad \left. + \frac{h^{k+1}}{\sqrt{\Delta t}} \int_{t-\Delta t}^t \|\partial_t \mathbf{B}\|_{H^{k+1}}^2 ds \right\}^{\frac{1}{2}} \|\nabla \epsilon_{\mathbf{B}}^n\| \\
&\leq C\Delta t (\Delta t^2 + h^{2k+2}) + \frac{\Delta t c_1 \eta_1 S_c}{4} \|\nabla \epsilon_{\mathbf{B}}^n\|^2.
\end{aligned} \tag{51}$$

For nonlinear term \mathcal{I}_5 , adding and subtracting some terms, we can rewrite

$$\begin{aligned}
\mathcal{I}_5 &= \Delta t \lambda \left\{ \epsilon a_\phi(\kappa^n(\phi^n) - \kappa^n(\phi^{n-1}); \mu^n, \epsilon_\mu^n) \right. \\
&\quad + \epsilon a_\phi(\kappa^n(\phi^{n-1}) - \kappa^n(\phi_h^{n-1}); \mu^n, \epsilon_\mu^n) + (\mathbf{u}^n \cdot \nabla(\phi^n - \phi^{n-1}), \epsilon_\mu^n) \\
&\quad + ((\mathbf{u}^n - R_h \mathbf{u}^n) \cdot \nabla \phi^{n-1}, \epsilon_\mu^n) + (R_h \mathbf{u}^n \cdot \nabla(\phi^{n-1} - r_h \phi^{n-1}), \epsilon_\mu^n) \\
&\quad \left. + (R_h \mathbf{u}^n \cdot \nabla \epsilon_\phi^{n-1}, \epsilon_\mu^n) + (\epsilon_\mu^n \cdot \nabla \phi_h^{n-1}, \epsilon_\mu^n) \right\} \\
&= \sum_{i=1}^6 A_i^n + \Delta t \lambda (\epsilon_\mu^n \cdot \nabla \phi_h^{n-1}, \epsilon_\mu^n).
\end{aligned} \tag{52}$$

For terms $A_1^n - A_2^n$, thanks to Hölder inequality, Taylor's theorem and (33), one finds that

$$\begin{aligned}
A_1^n + A_2^n &\leq C\Delta t |\kappa|_{C^{0,1}(\overline{\Omega} \times R; R)} \|\nabla \mu^n\|_{L^\infty} \left\{ \|\Delta t \delta_t \phi^n\| \right. \\
&\quad \left. + \|\phi^{n-1} - \hat{r}_h \phi^{n-1}\| + \|\hat{r}_h \phi^{n-1} - \phi_h^{n-1}\| \right\} \|\nabla \epsilon_\mu^n\| \\
&\leq C\Delta t \left\{ \left(\Delta t \int_{t-\Delta t}^t \|\partial_s \phi(s)\|_{L^2}^2 ds \right)^{\frac{1}{2}} + h^{k+1} \|\partial_t \phi\|_{H^{k+1}} \right. \\
&\quad \left. + \|\epsilon_\phi^{n-1}\| \right\} \|\nabla \epsilon_\mu^n\|,
\end{aligned}$$

where

$$|\kappa|_{C^{0,1}(\overline{\Omega} \times R; R)} := \sup \left\{ \frac{|\kappa(\mathbf{x}, \phi) - \nu(\mathbf{y}, \varphi)|}{|(\mathbf{x}, \phi) - (\mathbf{y}, \varphi)|} : (\mathbf{x}, \phi), (\mathbf{y}, \varphi) \in \overline{\Omega} \times \mathbf{R} \right\}.$$

Using (6), (29) and (33), the terms $A_3^n - A_6^n$ can be bounded by

$$\begin{aligned}
A_3^n &\leq C\Delta t \|\mathbf{u}^n\|_{L^4} \|\Delta t \nabla \delta_t \phi^n\| \|\epsilon_\mu^n - \bar{\epsilon}_\mu^n\|_{L^4} \\
&\leq C\Delta t \left(\Delta t \int_{t-\Delta t}^t \|\partial_s \nabla \phi(s)\|_{L^2}^2 ds \right)^{\frac{1}{2}} \|\nabla \epsilon_\phi^n\|_{L^2}, \\
A_4^n &\leq C\Delta t \|\mathbf{u}^n - R_h \mathbf{u}^n\| \|\nabla \phi^{n-1}\|_{L^4} \|\epsilon_\mu^n - \bar{\epsilon}_\mu^n\|_{L^4} \\
&\leq C\Delta t h^{k+1} \|(\mathbf{u}, p)\|_{H^{k+1} \times H^k} \|\nabla \epsilon_\mu^n\|, \\
A_5^n &\leq C\Delta t \|R_h \mathbf{u}^n\|_{L^4} \|\nabla(\phi^{n-1} - r_h \phi^{n-1})\| \|\epsilon_\mu^n - \bar{\epsilon}_\mu^n\|_{L^4} \\
&\leq C\Delta t h^{k+1} \|\phi\|_{H^{k+2}} \|\nabla \epsilon_\phi^n\|, \\
A_6^n &\leq C\Delta t \|R_h \mathbf{u}^n\|_{L^4} \|\nabla \epsilon_\phi^{n-1}\| \|\epsilon_\mu^n - \bar{\epsilon}_\mu^n\|_{L^4} \leq C\Delta t \|\nabla \epsilon_\phi^{n-1}\| \|\nabla \epsilon_\mu^n\|.
\end{aligned}$$

Combining (52) with above estimates, it follows that

$$\begin{aligned}
\mathcal{I}_5 &\leq C\Delta t (h^{2k+2} + \Delta t^2) + \Delta t \|\epsilon_\phi^{n-1}\|^2 \\
&\quad + \frac{\Delta t \lambda \kappa_1}{8} \|\nabla \epsilon_\mu^n\|^2 + \Delta t \lambda (\epsilon_\mu^n \cdot \nabla \phi_h^{n-1}, \epsilon_\mu^n).
\end{aligned} \tag{53}$$

Similarly, for nonlinear term \mathcal{I}_6 , we rewrite

$$\begin{aligned}
\mathcal{I}_6 &= \Delta t \left\{ \epsilon a_\phi(\kappa^n(\phi^n) - \kappa^n(\phi^{n-1}); \mu^n, \epsilon_\phi^n) \right. \\
&\quad + \epsilon a_\phi(\kappa^n(\phi^{n-1}) - \kappa^n(\phi_h^{n-1}); \mu^n, \epsilon_\phi^n) + (\mathbf{u}^n \cdot \nabla(\phi^n - \phi^{n-1}), \epsilon_\phi^n) \\
&\quad + ((\mathbf{u}^n - R_h \mathbf{u}^n) \cdot \nabla \phi^{n-1}, \epsilon_\phi^n) + (R_h \mathbf{u}^n \cdot \nabla(\phi^{n-1} - r_h \phi^{n-1}), \epsilon_\phi^n) \\
&\quad \left. + (R_h \mathbf{u}^n \cdot \nabla \epsilon_\phi^{n-1}, \epsilon_\phi^n) + (\epsilon_\mu^n \cdot \nabla \phi_h^{n-1}, \epsilon_\phi^n) \right\} \\
&= \sum_{i=7}^{13} A_i^n.
\end{aligned} \tag{54}$$

Applying (6), (29) and (33), we bound the terms $A_7^n - A_{13}^n$ as follows

$$\begin{aligned}
A_7^n + A_8^n &\leq C\Delta t \left\{ \left(\Delta t \int_{t-\Delta t}^t \|\partial_s \nabla \phi(s)\|_{L^2}^2 ds \right)^{\frac{1}{2}} \right. \\
&\quad \left. + h^{k+1} + \|\epsilon_\phi^{n-1}\| \right\} \|\nabla \epsilon_\phi^n\|, \\
A_9^n &\leq C\Delta t \|\mathbf{u}^n\|_{L^\infty} \|\nabla(\phi^n - \phi^{n-1})\| \|\epsilon_\phi^n\| \\
&\leq C\Delta t \left(\Delta t \int_{t-\Delta t}^t \|\partial_s \nabla \phi(s)\|_{L^2}^2 ds \right)^{\frac{1}{2}} \|\epsilon_\phi^n\|, \\
A_{10}^n &\leq C\Delta t \|\mathbf{u}^n - R_h \mathbf{u}^n\| \|\nabla \phi^{n-1}\|_{L^\infty} \|\epsilon_\phi^n\| \\
&\leq C\Delta t h^{k+1} \|(\mathbf{u}, p)\|_{H^{k+1} \times H^k} \|\epsilon_\phi^n\|, \\
A_{11}^n &\leq C\Delta t \|R_h \mathbf{u}^n\|_{L^\infty} \|\nabla(\phi^{n-1} - r_h \phi^{n-1})\| \|\epsilon_\phi^n\| \\
&\leq C\Delta t h^{k+1} \|\phi\|_{H^{k+2}} \|\epsilon_\phi^n\|, \\
A_{12}^n &\leq C\Delta t \|R_h \mathbf{u}^n\|_{L^\infty} \|\nabla \epsilon_\phi^{n-1}\| \|\epsilon_\phi^n\| \leq C\Delta t \|\nabla \epsilon_\phi^{n-1}\| \|\epsilon_\phi^n\|, \\
A_{13}^n &\leq C\Delta t \|\epsilon_\mu^n\|_{L^6} \|\nabla \phi_h^{n-1}\|_{L^3} \|\epsilon_\phi^n\| \leq C\Delta t \|\nabla \epsilon_\mu^n\| \|\epsilon_\phi^n\|.
\end{aligned}$$

Combining (54) with above estimates, one finds that

$$\begin{aligned} \mathcal{R}_6 \leq & C\Delta t(h^{2k+2} + \Delta t^2) + C\Delta t\|\epsilon_\phi^{n-1}\|^2 + C\Delta t\|\nabla\epsilon_\phi^{n-1}\|^2 \\ & + C\Delta t\|\nabla\epsilon_\phi^n\|^2 + \frac{\Delta t c_0 \nu_1}{8}\|\nabla\epsilon_\mathbf{u}^n\|^2 + C\Delta t\|\epsilon_\phi^n\|^2. \end{aligned} \quad (55)$$

For nonlinear term \mathcal{R}_7 , we can rewrite

$$\begin{aligned} \mathcal{R}_7 := & \Delta t \left\{ \lambda(\eta_\mu^n, \delta_t \epsilon_\phi^n) - \frac{\Delta t \lambda}{\epsilon}(\delta_t \phi^n, \delta_t \epsilon_\phi^n) \right. \\ & \left. - \lambda \epsilon^{-1}(\epsilon_\phi^{n-1}, \delta_t \epsilon_\phi^n) + \lambda \epsilon^{-1}((\phi^n)^3 - (\phi_h^n)^3, \delta_t \epsilon_\phi^n) \right\} \\ = & \Delta t \sum_{i=1}^4 \widehat{A}_i^n. \end{aligned} \quad (56)$$

Using (6), (35) and **Lemmas 2-3**, we bound the terms $\widehat{A}_1^n - \widehat{A}_4^n$ as follows

$$\begin{aligned} \widehat{A}_1^n & \leq C\|\nabla\eta_\mu^n\|\|\delta_t \epsilon_\phi^n\|_{-1,h} \leq Ch^{k+1}\|\mu\|_{k+2}\|\delta_t \epsilon_\phi^n\|_{-1,h}, \\ \widehat{A}_2^n & \leq \frac{1}{\epsilon}\|\Delta t \nabla \delta_t \phi^n\|\|\delta_t \epsilon_\phi^n\|_{-1,h} \leq C\left(\Delta t \int_{t-\Delta t}^t \|\partial_s \nabla \phi(s)\|_{L^2}^2 ds\right)^{\frac{1}{2}}\|\delta_t \epsilon_\phi^n\|_{-1,h}, \\ \widehat{A}_3^n & \leq C\|\epsilon_\phi^{n-1}\|\|\delta_t \epsilon_\phi^n\|_{-1,h} \leq C\|\epsilon_\phi^{n-1}\|\|\delta_t \epsilon_\phi^n\|_{-1,h} + C\|\eta_\phi^{n-1}\|\|\delta_t \epsilon_\phi^n\|_{-1,h}, \\ \widehat{A}_4^n & \leq C\|\nabla(\phi^n)^3 - (\phi_h^n)^3\|\|\delta_t \epsilon_\phi^n\|_{-1,h} \leq C\|\nabla(\phi^n - \phi_h^n)\|\|\delta_t \epsilon_\phi^n\|_{-1,h} \\ & \leq C\|\nabla\epsilon_\phi^n\|\|\delta_t \epsilon_\phi^n\|_{-1,h} + C\|\nabla\eta_\phi^n\|\|\delta_t \epsilon_\phi^n\|_{-1,h}. \end{aligned}$$

Combining (56) with above estimates, it follows that

$$\begin{aligned} \mathcal{R}_7 \leq & C\Delta t(h^{2k+2} + \Delta t^2) + C\Delta t\|\epsilon_\phi^{n-1}\|^2 \\ & + C\Delta t\|\nabla\epsilon_\phi^n\|^2 + \alpha\Delta t\|\delta_t \epsilon_\phi^n\|_{-1,h}^2. \end{aligned} \quad (57)$$

Taking $T_h(\delta_t \epsilon_\phi^n)$ in (37a), and using (6), **Lemmas 1-2**, we derive

$$\begin{aligned} \|\delta_t \epsilon_\phi^n\|_{-1,h}^2 = & -\epsilon a_\phi(\kappa^n(\phi_h^{n-1}); \epsilon_\mu^n, T_h(\delta_t \epsilon_\phi^n)) + (R_\phi^n, T_h(\delta_t \epsilon_\phi^n)) + (A_h^n, T_h(\delta_t \epsilon_\phi^n)) \\ & (58) \\ \leq & \epsilon \kappa_2 \|\nabla \epsilon_\mu^n\| \|\delta_t \epsilon_\phi^n\|_{-1,h} + \|\partial_t \phi^n - \delta_t r_h \phi^n\| \|T_h(\delta_t \epsilon_\phi^n)\| \\ & + \|\mathbf{u}^n\|_{L^3} \|\Delta t \nabla \delta_t \phi^n\| \|T_h(\delta_t \epsilon_\phi^n)\|_{L^6} + \|\eta_\mathbf{u}^n\|_{L^3} \|\nabla \phi^{n-1}\| \|T_h(\delta_t \epsilon_\phi^n)\|_{L^6} \\ & + \|R_h \mathbf{u}^n\|_{L^3} \|\nabla \eta_\phi^{n-1}\| \|T_h(\delta_t \epsilon_\phi^n)\|_{L^6} + \|R_h \mathbf{u}^n\|_{L^3} \|\nabla \epsilon_\phi^{n-1}\| \|T_h(\delta_t \epsilon_\phi^n)\|_{L^6} \\ & + \|\epsilon_\mathbf{u}^n\|_{L^3} \|\nabla \phi_h^{n-1}\| \|T_h(\delta_t \epsilon_\phi^n)\|_{L^6} \\ \leq & \frac{1}{2} \|\delta_t \epsilon_\phi^n\|_{-1,h}^2 + C_1 \|\nabla \epsilon_\mu^n\|^2 + C_2 \|\nabla \epsilon_\mathbf{u}^n\|^2 \\ & + C \|\nabla \epsilon_\phi^{n-1}\|^2 + C(h^{2k+2} + \Delta t^2). \end{aligned}$$

Then we obtain

$$\|\delta_t \epsilon_\phi^n\|_{-1,h}^2 \leq 2C_1 \|\nabla \epsilon_\mu^n\|^2 + 2C_2 \|\nabla \epsilon_\mathbf{u}^n\|^2 + C \|\nabla \epsilon_\phi^{n-1}\|^2 + C(h^{2k+2} + \Delta t^2). \quad (59)$$

From (57) and (59), and taking $\alpha = \min\{\frac{\lambda\kappa_1}{8C_1}, \frac{c_0\nu_1}{8C_2}\}$ in (57), one finds that

$$\begin{aligned} \mathcal{R}_7 \leq & C\Delta t(h^{2k+2} + \Delta t^2) + \frac{\Delta t c_0 \nu_1}{8} \|\nabla \epsilon_{\mathbf{u}}^n\|^2 + \frac{\Delta t \lambda \kappa_1}{8} \|\nabla \epsilon_{\mu}^n\|^2 \\ & + C\Delta t \|\nabla \epsilon_{\phi}^{n-1}\|^2 + C\Delta t \|\nabla \epsilon_{\phi}^n\|^2. \end{aligned} \quad (60)$$

Similarly, for nonlinear term \mathcal{R}_8 , we rewrite

$$\begin{aligned} \mathcal{R}_8 := & \Delta t \left\{ (\eta_{\mu}^n, \epsilon_{\mu}^n) + \frac{\Delta t}{\epsilon} (\delta_t \phi^n, \epsilon_{\mu}^n) \right. \\ & \left. + \epsilon^{-1} (e_{\phi}^{n-1}, \epsilon_{\mu}^n) + \epsilon^{-1} ((\phi^n)^3 - (\phi_h^n)^3, \epsilon_{\mu}^n) \right\} \\ = & \sum_{i=5}^8 \hat{A}_i^n. \end{aligned} \quad (61)$$

Using (6), (33), (35) and **Lemma 3**, we bound the terms $\hat{A}_5^n - \hat{A}_8^n$ as follows

$$\begin{aligned} \hat{A}_5^n & \leq C\Delta t \|\eta_{\mu}^n\| \|\epsilon_{\mu}^n\| \leq Ch^{k+1} \|\mu\|_{k+1} \|\epsilon_{\mu}^n\|, \\ \hat{A}_6^n & \leq \Delta t \frac{1}{\epsilon} \|\Delta t \nabla \delta_t \phi^n\| \|\epsilon_{\mu}^n\| \leq C\Delta t \left(\Delta t \int_{t-\Delta t}^t \|\partial_s \nabla \phi(s)\|_{L^2}^2 ds \right)^{\frac{1}{2}} \|\epsilon_{\mu}^n\|, \\ \hat{A}_7^n & \leq C\Delta t \|e_{\phi}^{n-1}\| \|\epsilon_{\mu}^n\| \leq C\Delta t \|\epsilon_{\phi}^{n-1}\| \|\epsilon_{\mu}^n\| + C\Delta t \|\eta_{\phi}^{n-1}\| \|\epsilon_{\mu}^n\| \\ \hat{A}_8^n & \leq C\Delta t \|(\phi^n)^3 - (\phi_h^n)^3\| \|\epsilon_{\mu}^n\| \leq C\Delta t \|(\phi^n - \phi_h^n)\| \|\epsilon_{\mu}^n\| \\ & \leq C\Delta t \|\epsilon_{\phi}^n\| \|\epsilon_{\mu}^n\| + C\Delta t \|\nabla \eta_{\phi}^n\| \|\epsilon_{\mu}^n\|. \end{aligned}$$

Combining (61) with above inequalities, we have

$$\begin{aligned} \mathcal{R}_8 \leq & C\Delta t(h^{2k+2} + \Delta t^2) + \frac{\Delta t \kappa_1}{8} \|\epsilon_{\mu}^n\|^2 \\ & + C\Delta t \|\epsilon_{\phi}^n\|^2 + C\Delta t \|\epsilon_{\phi}^{n-1}\|^2. \end{aligned} \quad (62)$$

For nonlinear term \mathcal{R}_9 , by adding and subtracting some terms, we rewrite

$$\begin{aligned} \mathcal{R}_9 = & \Delta t \left\{ a_f(\nu^n(\phi^n) - \nu^n(\phi^{n-1}); \mathbf{u}^n, \epsilon_{\mathbf{u}}^n) \right. \\ & + a_f(\nu^n(\phi^{n-1}) - \nu^n(\phi_h^{n-1}); \mathbf{u}^n, \epsilon_{\mathbf{u}}^n) + b(\eta_{\mathbf{u}}^n, \mathbf{u}^n, \epsilon_{\mathbf{u}}^n) \\ & + b(R_h \mathbf{u}^n - R_h \mathbf{u}^{n-1}, \mathbf{u}^n, \epsilon_{\mathbf{u}}^n) + b(\epsilon_{\mathbf{u}}^{n-1}, \mathbf{u}^n, \epsilon_{\mathbf{u}}^n) \\ & + b(\mathbf{u}_h^{n-1}, \eta_{\mathbf{u}}^n, \epsilon_{\mathbf{u}}^n) + b(\mathbf{u}_h^{n-1}, \epsilon_{\mathbf{u}}^n, \epsilon_{\mathbf{u}}^n) + S_c c_{\widehat{B}}(\eta_{\mathbf{B}}^n, \mathbf{B}^n, \epsilon_{\mathbf{u}}^n) \\ & + S_c c_{\widehat{B}}(R_{mh} \mathbf{B}^n - R_{mh} \mathbf{B}^{n-1}, \mathbf{B}^n, \epsilon_{\mathbf{u}}^n) + S_c c_{\widehat{B}}(\epsilon_{\mathbf{B}}^{n-1}, \mathbf{B}^n, \epsilon_{\mathbf{u}}^n) \\ & + S_c c_{\widehat{B}}(\mathbf{B}_h^{n-1}, \eta_{\mathbf{B}}^n, \epsilon_{\mathbf{u}}^n) + S_c c_{\widehat{B}}(\mathbf{B}_h^{n-1}, \epsilon_{\mathbf{B}}^n, \epsilon_{\mathbf{u}}^n) \\ & - \lambda(\mu^n \nabla \eta_{\phi}^n, \epsilon_{\mathbf{u}}^n) - \lambda(\mu^n \nabla(r_h \phi^n - r_h \phi^{n-1}), \epsilon_{\mathbf{u}}^n) \\ & - \lambda(\mu^n \nabla \epsilon_{\phi}^{n-1}, \epsilon_{\mathbf{u}}^n) - \lambda(\eta_{\mu}^n \nabla \phi_h^{n-1}, \epsilon_{\mathbf{u}}^n) - \lambda(\epsilon_{\mu}^n \nabla \phi_h^{n-1}, \epsilon_{\mathbf{u}}^n) \\ & = \Delta t \left\{ \sum_{i=1}^{14} \Phi_i^n + b(\mathbf{u}_h^{n-1}, \epsilon_{\mathbf{u}}^n, \epsilon_{\mathbf{u}}^n) + S_c c_{\widehat{B}}(\mathbf{B}_h^{n-1}, \epsilon_{\mathbf{B}}^n, \epsilon_{\mathbf{u}}^n) \right. \\ & \left. - \lambda(\epsilon_{\mu}^n \nabla \phi_h^{n-1}, \epsilon_{\mathbf{u}}^n) \right\}. \end{aligned} \quad (63)$$

For the terms $\Phi_1^n - \Phi_2^n$, using Hölder inequality, (35) and Taylor's theorem, we have

$$\begin{aligned} \Phi_1^n + \Phi_2^n &\leq C|\nu|_{C^{0,1}(\overline{\Omega} \times R; R)} \|\mathbb{D}(\mathbf{u}^n)\|_{L^\infty} \left\{ \|\phi^n - \phi^{n-1}\| \right. \\ &\quad \left. + \|\phi^{n-1} - \hat{r}_h \phi^{n-1}\| + \|\hat{r}_h \phi^{n-1} - \phi_h^{n-1}\| \right\} \|\nabla \epsilon_{\mathbf{u}}^n\| \\ &\leq C \left\{ \left(\Delta t \int_{t-\Delta t}^t \|\partial_s \phi(s)\|_{L^2}^2 ds \right)^{\frac{1}{2}} + h^{k+1} \right. \\ &\quad \left. + \|\epsilon_\phi^{n-1}\| \right\} \|\nabla \epsilon_{\mathbf{u}}^n\|, \end{aligned}$$

Making use of (6) and (29), we estimate the terms $\Phi_3^n - \Phi_6^n$ as follows

$$\begin{aligned} \Phi_3^n &\leq C \|\nabla \eta_{\mathbf{u}}^n\| \|\nabla \mathbf{u}^n\| \|\nabla \epsilon_{\mathbf{u}}^n\| \\ &\leq Ch^{k+1} \|(\mathbf{u}, p)\|_{H^{k+2} \times H^{k+1}} \|\nabla \epsilon_{\mathbf{u}}^n\|, \\ \Phi_4^n &\leq C \|\Delta t R_{mh} \delta_t \mathbf{B}^n\| \|\nabla \mathbf{B}^n\|_{L^3} \|\nabla \epsilon_{\mathbf{u}}^n\| \\ &\leq C \left(\Delta t \int_{t-\Delta t}^t \|\partial_s \mathbf{B}(s)\|_{L^2}^2 ds \right)^{\frac{1}{2}} \|\nabla \epsilon_{\mathbf{u}}^n\|, \\ \Phi_5^n &\leq C \|\epsilon_{\mathbf{u}}^{n-1}\| \|\nabla \mathbf{u}^n\|_{L^3} \|\nabla \epsilon_{\mathbf{u}}^n\| \\ &\leq \|\epsilon_{\mathbf{u}}^{n-1}\| \|\nabla \epsilon_{\mathbf{u}}^n\|, \\ \Phi_6^n &\leq C \|\nabla \mathbf{u}_h^{n-1}\| \|\nabla \eta_{\mathbf{u}}^n\| \|\nabla \epsilon_{\mathbf{u}}^n\| \\ &\leq Ch^{k+1} \|(\mathbf{u}, p)\|_{H^{k+2} \times H^{k+1}} \|\nabla \epsilon_{\mathbf{u}}^n\|. \end{aligned}$$

Thanks to (6) and (31), the terms $\Phi_7^n - \Phi_{10}^n$ can be bounded by

$$\begin{aligned} \Phi_7^n &\leq C \|\eta_{\mathbf{B}}^n\| \|\mathbf{B}^n\|_{L^3} \|\nabla \epsilon_{\mathbf{u}}^n\| \\ &\leq Ch^{k+1} \|\mathbf{B}\|_{H^{k+1}} \|\nabla \epsilon_{\mathbf{u}}^n\|, \\ \Phi_8^n &\leq C \|\Delta t R_{mh} \delta_t \mathbf{B}^n\| \|\operatorname{curl} \mathbf{B}^n\|_{L^3} \|\nabla \epsilon_{\mathbf{u}}^n\| \\ &\leq C \left(\Delta t \int_{t-\Delta t}^t \|\partial_s \mathbf{B}(s)\|_{L^2}^2 ds \right)^{\frac{1}{2}} \|\nabla \epsilon_{\mathbf{u}}^n\|, \\ \Phi_9^n &\leq C \|\epsilon_{\mathbf{B}}^{n-1}\| \|\operatorname{curl} \mathbf{B}^n\|_{L^3} \|\nabla \epsilon_{\mathbf{u}}^n\| \leq C \|\epsilon_{\mathbf{B}}^{n-1}\| \|\nabla \epsilon_{\mathbf{u}}^n\|, \\ \Phi_{10}^n &\leq C \|\eta_{\mathbf{B}}^{n-1}\| \|\operatorname{curl} \mathbf{B}^n\|_{L^3} \|\nabla \epsilon_{\mathbf{u}}^n\| \\ &\leq Ch^{k+1} \|\mathbf{B}\|_{H^{k+1}} \|\nabla \epsilon_{\mathbf{u}}^n\|. \end{aligned}$$

Applying Hölder inequality, (6), (33) and (35), the terms $\Phi_{11}^n - \Phi_{14}^n$ can be estimated as follows

$$\begin{aligned}
\Phi_{11}^n &\leq C \|\mu^n\|_{L^3} \|\nabla \eta_\phi^n\| \|\epsilon_{\mathbf{u}}^n\|_{L^6} \\
&\leq Ch^{k+1} \|\phi\|_{H^{k+2}} \|\nabla \epsilon_{\mathbf{u}}^n\|, \\
\Phi_{12}^n &\leq C \|\mu^n\|_{L^3} \|\Delta t \nabla \delta_t r_h \phi^n\| \|\epsilon_{\mathbf{u}}^n\|_{L^6} \\
&\leq C \left(\Delta t \int_{t-\Delta t}^t \|\partial_s \nabla \phi(s)\|_{L^2}^2 ds \right)^{\frac{1}{2}} \|\nabla \epsilon_{\mathbf{u}}^n\|, \\
\Phi_{13}^n &\leq C \|\mu^n\|_{L^3} \|\nabla \epsilon_\phi^{n-1}\| \|\epsilon_{\mathbf{u}}^n\|_{L^6} \\
&\leq C \|\nabla \epsilon_\phi^{n-1}\| \|\nabla \epsilon_{\mathbf{u}}^n\|, \\
\Phi_{14}^n &\leq C \|\eta_\mu^n\| \|\nabla \phi_h^{n-1}\| \|\epsilon_{\mathbf{u}}^n\|_{L^6} \\
&\leq Ch^{k+1} \|\mu\|_{H^{k+1}} \|\nabla \epsilon_{\mathbf{u}}^n\|.
\end{aligned}$$

Combining (63) with above bounds, one finds that

$$\begin{aligned}
\Upsilon_9 &\leq C \Delta t \left\{ h^{k+1} + \Delta t + \|\epsilon_{\mathbf{u}}^{n-1}\| + \|\epsilon_{\mathbf{B}}^{n-1}\| + \|\epsilon_\phi^{n-1}\| \right\} \|\nabla \epsilon_{\mathbf{u}}^n\| \\
&\quad + \Delta t b(\mathbf{u}_h^{n-1}, \epsilon_{\mathbf{u}}^n, \epsilon_{\mathbf{u}}^n) + \Delta t S_c c_{\tilde{B}}(\mathbf{B}_h^{n-1}, \epsilon_{\mathbf{B}}^n, \epsilon_{\mathbf{u}}^n) - \Delta t \lambda(\epsilon_\mu^n \nabla \phi_h^{n-1}, \epsilon_{\mathbf{u}}^n) \\
&\leq C \left\{ \Delta t (h^{2k+2} + \Delta t^2) + \Delta t \|\epsilon_{\mathbf{u}}^{n-1}\|^2 + \Delta t \|\epsilon_{\mathbf{B}}^{n-1}\|^2 + \Delta t \|\epsilon_\phi^{n-1}\|^2 \right\} + \frac{\Delta t c_0 \nu_1}{8} \|\nabla \epsilon_{\mathbf{u}}^n\|^2 \\
&\quad + \Delta t b(\mathbf{u}_h^{n-1}, \epsilon_{\mathbf{u}}^n, \epsilon_{\mathbf{u}}^n) + \Delta t S_c c_{\tilde{B}}(\mathbf{B}_h^{n-1}, \epsilon_{\mathbf{B}}^n, \epsilon_{\mathbf{u}}^n) - \Delta t \lambda(\epsilon_\mu^n \nabla \phi_h^{n-1}, \epsilon_{\mathbf{u}}^n).
\end{aligned} \tag{64}$$

For nonlinear term Υ_{10} , by adding and subtracting some terms, we can rewrite

$$\begin{aligned}
\Upsilon_{10} &= \Delta t \left\{ S_c a_{\mathbf{B}}(\eta^n(\phi^n) - \eta^n(\phi^{n-1}); \mathbf{B}^n, \epsilon_{\mathbf{B}}^n) \right. \\
&\quad + S_c a_{\mathbf{B}}(\eta^n(\phi^{n-1}) - \eta^n(\phi_h^{n-1}); \mathbf{B}^n, \epsilon_{\mathbf{B}}^n) - S_c c_{\tilde{B}}(\mathbf{u}^n, \eta_{\mathbf{B}}^n, \epsilon_{\mathbf{B}}^n) \\
&\quad - S_c c_{\tilde{B}}(\mathbf{u}^n, R_{mh} \mathbf{B}^n - R_{mh} \mathbf{B}^{n-1}, \epsilon_{\mathbf{B}}^n) - S_c c_{\tilde{B}}(\mathbf{u}^n, \epsilon_{\mathbf{B}}^{n-1}, \epsilon_{\mathbf{B}}^n) \\
&\quad \left. - S_c c_{\tilde{B}}(\eta_{\mathbf{u}}^n, \mathbf{B}_h^{n-1}, \epsilon_{\mathbf{B}}^n) - S_c c_{\tilde{B}}(\epsilon_{\mathbf{u}}^n, \mathbf{B}_h^{n-1}, \epsilon_{\mathbf{B}}^n) \right\} \\
&= \Delta t \left\{ \sum_{i=1}^6 \widehat{\Phi}_i^n - S_c c_{\tilde{B}}(\epsilon_{\mathbf{u}}^n, \mathbf{B}_h^{n-1}, \epsilon_{\mathbf{B}}^n) \right\}.
\end{aligned} \tag{65}$$

For the terms $\widehat{\Phi}_1^n - \widehat{\Phi}_2^n$, thanks to Hölder inequality, (35) and Taylor's theorem, we get

$$\begin{aligned}
\widehat{\Phi}_1^n + \widehat{\Phi}_2^n &\leq C |\eta|_{C^{0,1}(\overline{\Omega} \times R; R)} \|\nabla \mathbf{B}^n\|_{L^\infty} \left\{ \|\Delta t \delta_t \phi^n\| \right. \\
&\quad \left. + \|\phi^{n-1} - \widehat{r}_h \phi^{n-1}\| + \|\widehat{r}_h \phi^{n-1} - \phi_h^{n-1}\| \right\} \|\nabla \epsilon_{\mathbf{B}}^n\| \\
&\leq C \left\{ \left(\Delta t \int_{t-\Delta t}^t \|\partial_s \phi(s)\|_{L^2}^2 ds \right)^{\frac{1}{2}} + h^{k+1} \right. \\
&\quad \left. + \|\epsilon_\phi^{n-1}\| \right\} \|\nabla \epsilon_{\mathbf{B}}^n\|
\end{aligned}$$

By (6), (29) and (31), the terms $\widehat{\Phi}_3^n - \widehat{\Phi}_6^n$ can be bounded by

$$\begin{aligned}
\widehat{\Phi}_3^n &\leq C \|\mathbf{u}^n\|_{L^\infty} \|\eta_{\mathbf{B}}^n\| \|\operatorname{curl} \epsilon_{\mathbf{B}}^n\| \\
&\leq Ch^{k+1} \|\mathbf{B}\|_{H^{k+1}} \|\nabla \epsilon_{\mathbf{B}}^n\|, \\
\widehat{\Phi}_4^n &\leq C \|\mathbf{u}^n\|_{L^\infty} \|\Delta t R_{mh} \delta_t \mathbf{B}^n\| \|\operatorname{curl} \epsilon_{\mathbf{B}}^n\| \\
&\leq C \left(\Delta t \int_{t-\Delta t}^t \|\partial_s \mathbf{B}(s)\|_{L^2}^2 ds \right)^{\frac{1}{2}} \|\nabla \epsilon_{\mathbf{B}}^n\|, \\
\widehat{\Phi}_5^n &\leq C \|\mathbf{u}^n\|_{L^\infty} \|\epsilon_{\mathbf{B}}^{n-1}\| \|\operatorname{curl} \epsilon_{\mathbf{B}}^n\| \\
&\leq C \|\epsilon_{\mathbf{B}}^{n-1}\| \|\nabla \epsilon_{\mathbf{B}}^n\|, \\
\widehat{\Phi}_6^n &\leq C \|\eta_{\mathbf{u}}^n\|_{L^6} \|\mathbf{B}_h^{n-1}\|_{L^3} \|\operatorname{curl} \epsilon_{\mathbf{B}}^n\| \\
&\leq Ch^{k+1} \|(\mathbf{u}, p)\|_{H^{k+2} \times H^{k+1}} \|\mathbf{B}_h^{n-1}\|_{L^3} \|\nabla \epsilon_{\mathbf{B}}^n\|.
\end{aligned}$$

Combining (65) with above inequalities, it follows that

$$\begin{aligned}
\Upsilon_{10} &\leq C \Delta t \left\{ h^{k+1} + \Delta t + \|\epsilon_{\mathbf{B}}^{n-1}\| + \|\epsilon_{\phi}^{n-1}\| \right\} \|\nabla \epsilon_{\mathbf{B}}^n\| \\
&\quad - \Delta t S_c c_{\tilde{B}}(\epsilon_{\mathbf{u}}^n, \mathbf{B}_h^{n-1}, \epsilon_{\mathbf{B}}^n) \\
&\leq C \Delta t (h^{2k+2} + \Delta t^2) + \Delta t \|\epsilon_{\mathbf{B}}^{n-1}\|^2 + \Delta t \|\epsilon_{\phi}^{n-1}\|^2 \\
&\quad + \frac{c_1 \eta_1 S_c \Delta t}{4} \|\nabla \epsilon_{\mathbf{B}}^n\|^2 - \Delta t S_c c_{\tilde{B}}(\epsilon_{\mathbf{u}}^n, \mathbf{B}_h^{n-1}, \epsilon_{\mathbf{B}}^n).
\end{aligned} \tag{66}$$

Combining (47) with (7)-(8), (48)-(51), (53), (55), (60), (62), (64) and (66), we have

$$\begin{aligned}
&\frac{1}{2} (\|\epsilon_{\phi}^n\|^2 - \|\epsilon_{\phi}^{n-1}\|^2 + \|\epsilon_{\phi}^n - \epsilon_{\phi}^{n-1}\|^2) \\
&\quad + \frac{\epsilon \lambda}{2} (\|\nabla \epsilon_{\phi}^n\|^2 - \|\nabla \epsilon_{\phi}^{n-1}\|^2 + \|\nabla \epsilon_{\phi}^n - \nabla \epsilon_{\phi}^{n-1}\|^2) \\
&\quad + \frac{1}{2} (\|\epsilon_{\mathbf{u}}^n\|^2 - \|\epsilon_{\mathbf{u}}^{n-1}\|^2 + \|\epsilon_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^{n-1}\|^2) \\
&\quad + \frac{S_c}{2} (\|\epsilon_{\mathbf{B}}^n\|^2 - \|\epsilon_{\mathbf{B}}^{n-1}\|^2 + \|\epsilon_{\mathbf{B}}^n - \epsilon_{\mathbf{B}}^{n-1}\|^2) + \Delta t \lambda \kappa_1 \|\nabla \epsilon_{\mu}^n\|^2 \\
&\quad + \Delta t \kappa_1 \|\epsilon_{\mu}^n\|^2 + \Delta t c_0 \nu_1 \|\nabla \epsilon_{\mathbf{u}}^n\|^2 + \Delta t c_1 \eta_1 S_c \|\nabla \epsilon_{\mathbf{B}}^n\|^2 \\
&\leq C \Delta t (h^{2k+2} + \Delta t^2) + C \Delta t (\|\epsilon_{\phi}^n\|^2 + \|\nabla \epsilon_{\phi}^n\|^2) \\
&\quad + C \Delta t (\|\epsilon_{\mathbf{u}}^{n-1}\|^2 + \|\epsilon_{\mathbf{B}}^{n-1}\|^2 + \|\epsilon_{\phi}^{n-1}\|^2 + \|\nabla \epsilon_{\phi}^{n-1}\|^2).
\end{aligned} \tag{67}$$

Summing (67) from $n = 1$ to m and using (36), we obtain

$$\begin{aligned}
& \|\epsilon_\phi^m\|^2 + \epsilon\lambda\|\nabla\epsilon_\phi^m\|^2 + \|\epsilon_{\mathbf{u}}^m\|^2 + \|\epsilon_{\mathbf{B}}^m\|^2 + \Delta t\lambda\kappa_1 \sum_{n=1}^m \|\nabla\epsilon_\mu^n\|^2 \\
& + \Delta t\kappa_1 \sum_{n=1}^m \|\epsilon_\mu^n\|^2 + \Delta tc_0\nu_1 \sum_{n=1}^m \|\nabla\epsilon_{\mathbf{u}}^n\|^2 + \Delta tc_1\eta_1 S_c \sum_{n=1}^m \|\nabla\epsilon_{\mathbf{B}}^n\|^2 \\
& \leq C(h^{2k+2} + \Delta t^2) + C_3\Delta t \sum_{n=1}^m \|\epsilon_\phi^n\|^2 + C_4\Delta t \sum_{n=1}^m \|\nabla\epsilon_\phi^n\|^2 \\
& + C_5\Delta t \sum_{n=1}^m \|\epsilon_{\mathbf{u}}^{n-1}\|^2 + C_6\Delta t \sum_{n=1}^m \|\epsilon_{\mathbf{B}}^{n-1}\|^2 \\
& + C_7\Delta t \sum_{n=1}^m \|\epsilon_\phi^{n-1}\|^2 + C_8\Delta t \sum_{n=1}^m \|\nabla\epsilon_\phi^{n-1}\|^2.
\end{aligned} \tag{68}$$

If $0 < \Delta t \leq \Delta t_0 := \frac{1}{2\max\{C_3, C_4\epsilon^{-1}\}} < \frac{1}{\max\{C_3, C_4\epsilon^{-1}\}}$, since $1 \leq \frac{1}{1-\max\{C_3, C_4\epsilon^{-1}\}\Delta t} \leq 2$, it follows from (68) that

$$\begin{aligned}
& \|\epsilon_\phi^m\|^2 + \epsilon\lambda\|\nabla\epsilon_\phi^m\|^2 + \|\epsilon_{\mathbf{u}}^m\|^2 + \|\epsilon_{\mathbf{B}}^m\|^2 + \Delta t\lambda\kappa_1 \sum_{n=1}^m \|\nabla\epsilon_\mu^n\|^2 \\
& + \Delta t\kappa_1 \sum_{n=1}^m \|\epsilon_\mu^n\|^2 + \Delta tc_0\nu_1 \sum_{n=1}^m \|\nabla\epsilon_{\mathbf{u}}^n\|^2 + \Delta tc_1\eta_1 S_c \sum_{n=1}^m \|\nabla\epsilon_{\mathbf{B}}^n\|^2 \\
& \leq C(h^{2k+2} + \Delta t^2) + \frac{(C_3 + C_7)\Delta t}{1 - \max\{C_3, C_4\epsilon^{-1}\}} \sum_{n=1}^m \|\epsilon_\phi^{n-1}\|^2 \\
& + \frac{(C_4 + C_8)\Delta t}{1 - \max\{C_3, C_4\epsilon^{-1}\}} \sum_{n=1}^m \|\nabla\epsilon_\phi^{n-1}\|^2 + \frac{C_5\Delta t}{1 - \max\{C_3, C_4\epsilon^{-1}\}} \sum_{n=1}^m \|\epsilon_{\mathbf{u}}^{n-1}\|^2 \\
& + \frac{C_6\Delta t}{1 - \max\{C_3, C_4\epsilon^{-1}\}} \sum_{n=1}^m \|\epsilon_{\mathbf{B}}^{n-1}\|^2.
\end{aligned} \tag{69}$$

By using the discrete Grönwall inequality, one finds that

$$\begin{aligned}
& \|\epsilon_\phi^m\|^2 + \epsilon\lambda\|\nabla\epsilon_\phi^m\|^2 + \|\epsilon_{\mathbf{u}}^m\|^2 + \|\epsilon_{\mathbf{B}}^m\|^2 + \Delta t\lambda\kappa_1 \sum_{n=1}^m \|\nabla\epsilon_\mu^n\|^2 \\
& + \Delta t\kappa_1 \sum_{n=1}^m \|\epsilon_\mu^n\|^2 + \Delta tc_0\nu_1 \sum_{n=1}^m \|\nabla\epsilon_{\mathbf{u}}^n\|^2 + \Delta tc_1\eta_1 S_c \sum_{n=1}^m \|\nabla\epsilon_{\mathbf{B}}^n\|^2 \\
& \leq C(h^{2k+2} + \Delta t^2).
\end{aligned} \tag{70}$$

The desired result follow from (29), (31), (33), (35) and the triangle inequality. The proof is completed. \square

In order to derive optimal error estimates on pressure, we define the semi-norm for $\mathbf{v} \in H^{-1}(\Omega)^d$:

$$\|\mathbf{v}\|_{*,h} = \sup_{\mathbf{v}_h \in \mathcal{X}_h} \frac{(\mathbf{v}, \mathbf{v}_h)}{\|\nabla \mathbf{v}_h\|}.$$

And we define the following inverse Stokes operator: $S : H^{-1}(\Omega)^d \rightarrow \mathcal{X}$ and discrete inverse Stokes operator: $S_h : H^{-1}(\Omega)^d \rightarrow \mathcal{X}_h$. Let $\hat{\nu} \in C^{0,1}(\overline{\Omega}; R^+)$ satisfy **Assumption A4**, for all $\mathbf{v} \in H^{-1}(\Omega)^d$, $(S(\mathbf{v}), r) \in \mathcal{X} \times \mathcal{M}$ is the weak solution of the following problem:

$$\begin{aligned} a_f(\hat{\nu}; S(\mathbf{v}), \mathbf{v}) + d(\mathbf{v}, r) &= (\mathbf{v}, \mathbf{v}), & \forall \mathbf{v} \in \mathcal{X}, \\ d(S(\mathbf{v}), w) &= 0, & \forall w \in \mathcal{M}. \end{aligned}$$

We have H^2 regularity results [28]:

$$\|S(\mathbf{v})\|_2 + \|\nabla r\| \leq C\|\mathbf{v}\|, \forall \mathbf{v} \in L^2(\Omega)^d.$$

And for all $\mathbf{v} \in H^{-1}(\Omega)^d$, $(S_h(\mathbf{v}), r_h) \in \mathcal{X}_h \times \mathcal{M}_h$ is the weak solution of the following discrete problem:

$$\begin{aligned} a_f(\hat{\nu}; S_h(\mathbf{v}), \mathbf{v}_h) + d(\mathbf{v}, r_h) &= (\mathbf{v}, \mathbf{v}_h), & \forall \mathbf{v}_h \in \mathcal{X}_h, \\ d(S_h(\mathbf{v}), w_h) &= 0, & \forall w_h \in \mathcal{M}_h. \end{aligned}$$

Lemma 4 For $\mathbf{v} \in \mathcal{X}_{0h}$, there exists constant $C > 0$ independent of h such that

$$\|\mathbf{v}\|_{*,h} \leq C\|\nabla S_h(\mathbf{v})\|. \quad (71)$$

Proof: The proof follows the similar lines as in [3]. Here we skip it. \square

Theorem 6 Suppose that assumptions of **Theorem 5** hold. We have the following estimates

$$\begin{aligned} \|\nabla S_h(e_{\mathbf{u}}^n - e_{\mathbf{u}}^{n-1})\| &\leq C\sqrt{\Delta t}(\Delta t + h^{k+1}), \\ \left(\Delta t \sum_{n=1}^N \|\nabla S_h(e_{\mathbf{u}}^n - e_{\mathbf{u}}^{n-1})\|^2\right)^{1/2} &\leq C\Delta t(\Delta t + h^{k+1}). \end{aligned}$$

Proof: Using **Assumption A1** and (37a), we obtain

$$(\delta_t \epsilon_{\mathbf{u}}^n, \mathbf{v}_h) + a_f(\nu^n(\phi_h^{n-1}); \epsilon_{\mathbf{u}}^n, \mathbf{v}_h) - d(\mathbf{v}_h, \epsilon_p^n) = (\Phi_h^n, \mathbf{v}_h) + (R_{\mathbf{u}}^n, \mathbf{v}_h), \quad (72)$$

$$d(\epsilon_{\mathbf{u}}^n, q_h) = 0. \quad (73)$$

Setting $\mathbf{v}_h = S_h(\epsilon_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^{n-1}) \in \mathcal{X}_h$ in (72), noting that $d(S_h(\epsilon_{\mathbf{u}}^n) - S_h(\epsilon_{\mathbf{u}}^{n-1}), q_h) = 0$, one finds that

$$\begin{aligned} \frac{\|\nabla S_h(\epsilon_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^{n-1})\|^2}{\Delta t} &= -a_f(\nu^n(\theta_h^{n-1}); \epsilon_{\mathbf{u}}^n, S_h(\epsilon_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^{n-1})) \\ &\quad + (R_{\mathbf{u}}^n, S_h(\epsilon_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^{n-1})) + (\Phi_h^n, S_h(\epsilon_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^{n-1})). \end{aligned} \quad (74)$$

Applying the same arguments as those in the proof of **Theorem 5**, the RHS of (74) can be estimated as

$$\begin{aligned} -a_f(\nu^n(\theta_h^{n-1}); \epsilon_{\mathbf{u}}^n, S_h(\epsilon_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^{n-1})) &\leq \frac{1}{6\Delta t} \|\nabla S_h(\epsilon_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^{n-1})\|^2 + C\Delta t \|\nabla \epsilon_{\mathbf{u}}^n\|^2, \\ (R_{\mathbf{u}}^n, S_h(\epsilon_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^{n-1})) &\leq \frac{1}{6\Delta t} \|\nabla S_h(\epsilon_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^{n-1})\|^2 + C\Delta t(\Delta t^2 + h^{2k+2}), \\ (\Phi_h^n, S_h(\epsilon_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^{n-1})) &\leq \frac{1}{6\Delta t} \|\nabla S_h(\epsilon_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^{n-1})\|^2 + C\Delta t(\Delta t^2 + h^{2k+2}) \\ &\quad + C\Delta t(\|\nabla \epsilon_{\mu}^n\|^2 + \|\nabla \epsilon_{\mathbf{u}}^n\|^2 + \|\nabla \epsilon_{\phi}^{n-1}\|^2 \\ &\quad + \|\nabla \epsilon_{\mathbf{u}}^{n-1}\|^2 + \|\nabla \epsilon_{\mathbf{B}}^n\|^2 + \|\nabla \epsilon_{\mathbf{B}}^{n-1}\|^2). \end{aligned}$$

Combining (74) with above bounds, it follows that

$$\begin{aligned} \frac{\|\nabla S_h(\epsilon_{\mathbf{u}}^n - \epsilon_{\mathbf{u}}^{n-1})\|^2}{2\Delta t} &\leq C\left\{\Delta t(\Delta t^2 + h^{2k+2}) + C\Delta t(\|\nabla \epsilon_{\mu}^n\|^2 + \|\nabla \epsilon_{\mathbf{u}}^n\|^2 \right. \\ &\quad \left. + \|\nabla \epsilon_{\phi}^{n-1}\|^2 + \|\nabla \epsilon_{\mathbf{u}}^{n-1}\|^2 + \|\nabla \epsilon_{\mathbf{B}}^n\|^2 + \|\nabla \epsilon_{\mathbf{B}}^{n-1}\|^2)\right\}. \end{aligned}$$

Applying **Theorems 5** and the triangle inequality, the desired result is obtained. The proof is finished. \square

Then we give and prove the second main result of this section for the pressure.

Theorem 7 *Suppose that assumptions of **Theorem 5** and $\Delta t \leq Ch$ hold. Then the finite element approximate p_h^n in (17) satisfy following bound:*

$$\left(\Delta t \sum_{n=1}^N \|p(t_n) - p_h^n\|^2\right)^{1/2} \leq C(\Delta t + h^{k+1}).$$

Proof: From **Assumption A1** and (37a), one finds that

$$\begin{aligned}
\beta_0 \|\epsilon_p^n\| &\leq \sup_{\mathbf{v}_h \in \mathcal{X}_h, \mathbf{v}_h \neq \mathbf{0}} \frac{d(\epsilon_p^n, \mathbf{v}_h)}{\|\nabla \mathbf{v}_h\|} \\
&\leq \sup_{\mathbf{v}_h \in \mathcal{X}_h, \mathbf{v}_h \neq \mathbf{0}} \left\{ \frac{(\delta_t \epsilon_{\mathbf{u}}^n, \mathbf{v}_h) + a_f(\nu^n(\phi_h^{n-1}); \epsilon_{\mathbf{u}}^n, \mathbf{v}_h)}{\|\nabla \mathbf{v}_h\|} \right\} \\
&\quad + \sup_{\mathbf{v}_h \in \mathcal{X}_h, \mathbf{v}_h \neq \mathbf{0}} \left\{ \frac{-(\Phi_h^n, \mathbf{v}_h) - (\partial_t \mathbf{u}^n - \delta_t R_h \mathbf{u}^n, \mathbf{v}_h)}{\|\nabla \mathbf{v}_h\|} \right\} \\
&\leq \sup_{\mathbf{v}_h \in \mathcal{X}_h, \mathbf{v}_h \neq \mathbf{0}} \frac{1}{\|\nabla \mathbf{v}_h\|} \left\{ b(\mathbf{u}_h^{n-1}, \epsilon_{\mathbf{u}}^n, \mathbf{v}_h) + S_c c_{\widehat{B}}(\mathbf{B}_h^{n-1}, \epsilon_{\mathbf{B}}^n, \mathbf{v}_h) \right. \\
&\quad \left. - \lambda(\epsilon_{\mu}^n \nabla \phi_h^{n-1}, \mathbf{v}_h) \right\} + C \left\{ \|\delta_t \epsilon_{\mathbf{u}}^n\|_{*,h} + \|\nabla \epsilon_{\mathbf{u}}^n\| \right. \\
&\quad \left. + C(\Delta t + h^{k+1}) + (\|\epsilon_{\mathbf{u}}^{n-1}\| + \|\epsilon_{\mathbf{B}}^{n-1}\| + \|\epsilon_{\phi}^{n-1}\|) \right\} \\
&\leq \sup_{\mathbf{v}_h \in \mathcal{X}_h, \mathbf{v}_h \neq \mathbf{0}} \frac{1}{\|\nabla \mathbf{v}_h\|} \left\{ b(\mathbf{u}_h^{n-1} - \mathbf{u}^{n-1}, \epsilon_{\mathbf{u}}^n, \mathbf{v}_h) + b(\mathbf{u}^{n-1}, \epsilon_{\mathbf{u}}^n, \mathbf{v}_h) \right. \\
&\quad \left. + S_c c_{\widehat{B}}(\mathbf{B}_h^{n-1} - \mathbf{B}^{n-1}, \epsilon_{\mathbf{B}}^n, \mathbf{v}_h) + S_c c_{\widehat{B}}(\mathbf{B}^{n-1}, \epsilon_{\mathbf{B}}^n, \mathbf{v}_h) \right. \\
&\quad \left. - \lambda(\epsilon_{\mu}^n \nabla(\phi_h^{n-1} - \phi^{n-1}), \mathbf{v}_h) - \lambda(\epsilon_{\mu}^n \nabla \phi^{n-1}, \mathbf{v}_h) \right\} + C \left\{ \|\delta_t \epsilon_{\mathbf{u}}^n\|_{*,h} \right. \\
&\quad \left. + \|\nabla \epsilon_{\mathbf{u}}^n\| + C(\Delta t + h^{k+1}) + (\|\epsilon_{\mathbf{u}}^{n-1}\| + \|\epsilon_{\mathbf{B}}^{n-1}\| + \|\epsilon_{\phi}^{n-1}\|) \right\}.
\end{aligned} \tag{75}$$

By **Assumption A2** and **Theorem 5**, it follows that

$$\begin{aligned}
\|\nabla e_{\mathbf{u}}^{n-1}\| &\leq C \min \left\{ h^{-1} \|e_{\mathbf{u}}^{n-1}\|, \|\nabla e_{\mathbf{u}}^{n-1}\| \right\} \\
&\leq C \min \left\{ h^{-1}(\Delta t + h^{k+1}), \Delta t^{-1/2}(\Delta t + h^{k+1}) \right\} \leq C, \\
\|\nabla e_{\mathbf{B}}^{n-1}\| &\leq C \min \left\{ h^{-1} \|e_{\mathbf{B}}^{n-1}\|, \|\nabla e_{\mathbf{B}}^{n-1}\| \right\} \\
&\leq C \min \left\{ h^{-1}(\Delta t + h^{k+1}), \Delta t^{-1/2}(\Delta t + h^{k+1}) \right\} \leq C, \\
\|\nabla e_{\phi}^{n-1}\| &\leq C \min \left\{ h^{-1} \|e_{\phi}^{n-1}\|, \|\nabla e_{\phi}^{n-1}\| \right\} \\
&\leq C \min \left\{ h^{-1}(\Delta t + h^{k+1}), \Delta t^{-1/2}(\Delta t + h^{k+1}) \right\} \leq C.
\end{aligned}$$

Using Hölder inequality and (6), we get

$$\begin{aligned}
b(\mathbf{u}_h^{n-1} - \mathbf{u}^{n-1}, \epsilon_{\mathbf{u}}^n, \mathbf{v}_h) &\leq C \|\nabla e_{\mathbf{u}}^{n-1}\| \|\nabla \epsilon_{\mathbf{u}}^n\| \|\nabla \mathbf{v}_h\| \leq C \|\nabla e_{\mathbf{u}}^{n-1}\| \|\nabla \mathbf{v}_h\|, \\
b(\mathbf{u}^{n-1}, \epsilon_{\mathbf{u}}^n, \mathbf{v}_h) &\leq C \|\nabla \mathbf{u}^{n-1}\| \|\nabla \epsilon_{\mathbf{u}}^n\| \|\nabla \mathbf{v}_h\| \leq C \|\nabla \epsilon_{\mathbf{u}}^n\| \|\nabla \mathbf{v}_h\|, \\
S_c c_{\widehat{B}}(\mathbf{B}_h^{n-1} - \mathbf{B}^{n-1}, \epsilon_{\mathbf{B}}^n, \mathbf{v}_h) &\leq C \|\nabla e_{\mathbf{B}}^{n-1}\| \|\nabla \epsilon_{\mathbf{B}}^n\| \|\nabla \mathbf{v}_h\| \leq C \|\nabla \epsilon_{\mathbf{B}}^n\| \|\nabla \mathbf{v}_h\|, \\
S_c c_{\widehat{B}}(\mathbf{B}^{n-1}, \epsilon_{\mathbf{B}}^n, \mathbf{v}_h) &\leq C \|\nabla \mathbf{B}^{n-1}\| \|\nabla \epsilon_{\mathbf{B}}^n\| \|\nabla \mathbf{v}_h\| \leq C \|\nabla \epsilon_{\mathbf{B}}^n\| \|\nabla \mathbf{v}_h\|, \\
-\lambda(\epsilon_{\mu}^n \nabla(\phi_h^{n-1} - \phi^{n-1}), \mathbf{v}_h) &\leq C \|\epsilon_{\mu}^n - \bar{\epsilon}_{\mu}^n\|_{L^4} \|\nabla e_{\phi}^{n-1}\| \|\nabla \mathbf{v}_h\|_{L^4} \leq C \|\nabla \epsilon_{\mu}^n\| \|\nabla \mathbf{v}_h\|.
\end{aligned}$$

Combining (75) with above inequalities, we derive

$$\begin{aligned} \|\epsilon_p^n\| \leq C \Big\{ & \|\delta_t \epsilon_{\mathbf{u}}^n\|_{*,h} + \|\nabla \epsilon_{\mathbf{u}}^n\| + \|\nabla \epsilon_{\mathbf{B}}^n\| + \|\nabla \epsilon_{\phi}^n\| + (\Delta t + h^{k+1}) \\ & + \|\epsilon_{\mathbf{u}}^{n-1}\| + \|\epsilon_{\mathbf{B}}^{n-1}\| + \|\epsilon_{\phi}^{n-1}\| \Big\} \end{aligned}$$

Due to

$$\begin{aligned} \|\delta_t \epsilon_{\mathbf{u}}^n\|_{*,h} & \leq C \left(\frac{1}{\Delta t} \|\nabla S_h(e_{\mathbf{u}}^n - e_{\mathbf{u}}^{n-1})\| \right) \\ & \leq C \frac{\sqrt{\Delta t}}{\Delta t} (\Delta t + h^{k+1}), \\ \left(\Delta t \sum_{n=1}^N \|\delta_t \epsilon_{\mathbf{u}}^n\|_{*,h}^2 \right)^{1/2} & \leq C (\Delta t + h^{k+1}). \end{aligned}$$

Using **Theorems 5, 6** and the triangle inequality, we obtain the desired result. The proof is finished. \square

5 Numerical results

In this section, some numerical tests are shown to confirm the theoretical convergence rates of the fully discrete scheme (17). The experiments have been finished with applying the finite element libraries from the Fenics Project [16]. The following functions κ, ν, μ are given

$$\kappa(\phi) = e^{\phi}, \quad \nu(\phi) = e^{-\phi}, \quad \eta(\phi) = e^{\phi}.$$

The finite element pair $P_2 - P_2 - \mathbf{P}_2 - \mathbf{P}_2 - P_1$ for the concentration field, the chemical potential, the velocity field, the magnetic field and the pressure is considered.

Note that the fully discrete scheme (17) is a nonlinear problem, thus we solve the problem (17) by a Picard type iteration. Namely, we fix the velocity field \mathbf{u}_h , the magnetic field \mathbf{B}_h and the pressure p_h at a given time step, then compute for the phase field ϕ_h and the chemical potential μ_h . Then we compute the velocity field \mathbf{u}_h , the magnetic field \mathbf{B}_h and the pressure p_h with these updated.

We consider a square domain $\Omega = [0, 1]^2$, and the following functions are given

$$\begin{aligned} \phi(x, y, t) &= 2 + \sin(t) \cos(\pi x) \cos(\pi y), \\ u_1(x, y, t) &= \pi \sin(2\pi y) \sin^2(\pi x) \sin(t), \\ u_2(x, y, t) &= -\pi \sin(2\pi x) \sin^2(\pi y) \sin(t), \\ p(x, y, t) &= \cos(\pi x) \sin(\pi y) \sin(t), \\ B_1(x, y, t) &= \sin(\pi x) \cos(\pi y) \sin(t), \\ B_2(x, y, t) &= -\sin(\pi y) \cos(\pi x) \sin(t), \end{aligned}$$




Fig. 1 Convergence rates: H^1 errors of the concentration field ϕ , the velocity field \mathbf{u} , the chemical potential μ , the magnetic field \mathbf{B} and L^2 the pressure as the space mesh size h .

as the exact solutions, and some source terms are taken such that the exact solutions satisfy (2)-(4).

The parameters are set to $S_c = 1$, $\epsilon = 0.05$, $\lambda = 1$ and $\Delta t = 0.1h^2$, and the uniform triangles meshes are employed. We plot the error estimates of the phase field, the velocity field, the magnetic field and the pressure between the numerical solution and the exact solution at $t = 0.5$ with different space sizes in Figure 1. We observe that the rates of convergence the fully discrete numerical scheme (17) are second order accurate for all variables.

Conclusions

In this paper, we have analyzed a fully discrete scheme for computing Cahn-Hilliard-Magneto-hydrodynamics system. The scheme is based on using conforming finite element method in space and Euler semi-implicit discretization with convex splitting in time. We have prove our scheme is unconditionally

energy stable and obtain optimal error estimates for the concentration field, the chemical potential, the velocity field, the magnetic field and the pressure. Numerical tests are shown to confirm the theoretical rates of the our scheme.

References

1. Anderson D. M., McFadden G. B., and Wheeler A. A.: Diffuse-interface methods in fluid mechanics, *Annu. Rev. Fluid Mech.*, vol. 30, Annual Reviews, Palo Alto, CA, 139-165 (1998).
2. Bermudez, A., Munoz-Sola, R., Vazquez R.: Analysis of two stationary magneto-hydrodynamic systems of equations including Joule heating. *J. Math. Anal. Appl.* 368, 444-468 (2010).
3. Cai Y., Shen J.: Error estimates for a fully discretized scheme to a Cahn-Hilliard phase-field model for two-phase incompressible flows, *Math. Comp.* 313, 2057-2090 (2018).
4. Davidson, P.: *An Introduction to Magnetohydrodynamics*, Cambridge University Press (2001).
5. Diegel A., Feng X., Wise S. M.: Analysis of a mixed finite element method for a Cahn-Hilliard-Darcy-Stokes system, *SIAM J. Numer. Anal.* 53, 127-152 (2015).
6. Elliott C. M., French D.A., Milner F. A.: A second order splitting method for the Cahn-Hilliard equation, *Numer. Math.* 54, 575-590 (1989).
7. Feng X., Prohl A.: Error analysis of a mixed finite element method for the Cahn-Hilliard equation, *Numer. Math.* 99, 47-84 (2004).
8. Feng X.: Fully discrete finite element approximations of the Navier-Stokes-Cahn-Hilliard diffuse interface model for two-phase fluid flows, *SIAM J. Numer. Anal.* 44, 1049-1072 (2006).
9. Feng X., He Y., Liu C.: Analysis of finite element approximations of a phase field model for two-phase fluids, *Math. Comp.* 258, 539-571 (2007).
10. Feng X., Li Y., Xing Y.: Analysis of mixed interior penalty discontinuous Galerkin methods for the Cahn-Hilliard equation and the Hele-Shaw flow, *SIAM J. Numer. Anal.* 54, 825-847 (2016).
11. Gurtin M. E., Polignone D. and Vinals J.: Two-phase binary fluids and immiscible fluids described by an order parameter, *Math. Models Methods Appl. Sci.* 6, 815-831 (1996).
12. Girault, V., Raviart, P. A.: *Finite Element Methods for Navier-Stokes Equations*, Series in Computational Mathematics, Springer-Verlag, New York (1986).
13. Gunzburger, M. D., Meir, A. J., Peterson, J. S.: On the existence and uniqueness and finite element approximation of solutions of the equations of stationary incompressible magnetohydrodynamics. *Math. Comput.* 56, 523-563 (1991).
14. Lowengrub J., Truskinovsky L.: Quasi-incompressible Cahn-Hilliard fluids and topological transitions, *R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci.* 454, 2617-2654 (1998).
15. Gerbeau, J. F., Le Bris, C., Lelièvre, T.: *Mathematical Methods for the Magnetohydrodynamics of Liquid Metals*. Numerical Mathematics and Scientific Computation, Oxford University Press, New York (2006).
16. Logg A., Mardal K-A., Wells G.: *Automated Solution of Differential Equations by the Finite Element Method: The FEniCS Book*, Springer-Verlag, Berlin, 2012.
17. Han D., Wang X., Wu H.: Existence and uniqueness of global weak solutions to a Cahn-Hilliard-Stokes-Darcy system for two phase incompressible flows in karstic geometry, *Journal of Differential Equations* 257, 3887-3933 (2014).
18. He, Y.: Unconditional convergence of the Euler semi-implicit scheme for the three-dimensional incompressible MHD equations. *IMA J. Numer. Anal.* 35, 767-801 (2015).
19. Kim, S., Lee, E. B., Choi, W.: Newton's algorithm for magnetohydrodynamic equations with the initial guess from Stokes-like problem. *J. Comput. Appl. Math.* 309, 1-10 (2017).
20. Lorca, S.A., Boldrini, J.L.: The initial value problem for a generalized Boussinesq model. *Nonlinear Anal.* 36, 457-480 (1999).
21. Liu C., Shen J.: A phase field model for the mixture of two incompressible fluids and its approximation by a Fourier-spectral method, *Phys. D* 179, 211-228 (2003).

22. Layton, W., Tran, W., Trenchea, H.: Stability of partitioned methods for magnetohydrodynamics flows at small magnetic Reynolds number. *Contemp. Math.* 586, 231-238 (2013).
23. Moreau, R.: *Magneto-hydrodynamics*. Kluwer Academic Publishers (1990).
24. Morley N. B., Smolentsev S., Barleon L., Kirillov I.R., Takahashi M.: Liquid magnetohydrodynamics-recent progress and future directions for fusion, *Fusion Eng. Des.* 51-52, 701-713 (2000).
25. Prohl, A.: Convergent finite element discretizations of the nonstationary incompressible magnetohydrodynamic system. *Math. Model. Numer. Anal.* 42, 1065-1087 (2008).
26. Szekely J.: *Fluid Flow Phenomena in Metals Processing*, Academic Press, New York, 1979.
27. Shen J., Yang X.: Decoupled, energy stable schemes for phase-field models of two-phase incompressible flows, *SIAM J. Numer. Anal.* 53, 279-296 (2015).
28. Temam R., *Navier-Stokes equations: Theory and numerical analysis*, 3rd ed., *Studies in Mathematics and its Applications*, vol. 2, North-Holland Publishing Co., Amsterdam, 1984.
29. Tabata, M., Tagami, D.: Error estimates of finite element methods for nonstationary thermal convection problems with temperature-dependent coefficients. *Numer. Math.* 100, 351-372 (2005).
30. Tone, F.: On the long-time H^2 -stability of the implicit Euler scheme for the 2D magnetohydrodynamics equations. *J. Sci. Comput.* 38, 331-348 (2009).
31. Wiedmer, M.: Finite element approximation for equations of magnetohydrodynamics. *Math. Comp.* 69, 83-101 (2000).
32. Yue P., Feng J. J., Liu C., Shen J.: A diffuse-interface method for simulating two-phase flows of complex fluids, *J. Fluid Mech.* 515, 293-317 (2004).
33. Yang J., Mao S., He X., Yang X., He Y.: A diffuse interface model and semi-implicit energy stable finite element method for two-phase magnetohydrodynamic flows, *Comput. Methods Appl. Mech. Engrg.* 356, 435-464 (2019).