Log-Sobolev inequality for the φ_2^4 and φ_3^4 measures

Roland Bauerschmidt^{*} Benoit Dagallier[†]

Abstract

The continuum φ_2^4 and φ_3^4 measures are shown to satisfy a log-Sobolev inequality uniformly in the lattice regularisation under the optimal assumption that their susceptibility is bounded. In particular, this applies to all coupling constants in any finite volume, and uniformly in the volume in the entire high temperature phases of the φ_2^4 and φ_3^4 models.

The proof uses a general criterion for the log-Sobolev inequality in terms of the Polchinski (renormalisation group) equation, a recently proved remarkable correlation inequality for Ising models with general external fields, the Perron–Frobenius theorem, and bounds on the susceptibilities of the φ_2^4 and φ_3^4 measures obtained using skeleton inequalities.

1 The φ_d^4 measure

Let d = 2 or d = 3, and let $\Lambda_{\varepsilon,L} = L\mathbb{T}^d \cap \varepsilon \mathbb{Z}^d$ (and always assume L is a multiple of ε). For $\lambda > 0$ and $\mu \in \mathbb{R}$, the lattice regularised (continuum) φ_d^4 measure is defined as the probability measure

(1.1)
$$\nu_{\lambda,\mu}^{\varepsilon,L}(d\varphi) \propto \exp\left[-\varepsilon^d \sum_{x \in \Lambda_{\varepsilon,L}} \left[\frac{1}{2}\varphi_x(-\Delta^{\varepsilon}\varphi)_x + \frac{\lambda}{4}\varphi_x^4 + \frac{\mu + a^{\varepsilon}(\lambda)}{2}\varphi_x^2\right]\right] d\varphi,$$

where $d\varphi$ denotes the Lebesgue measure on $\mathbb{R}^{\Lambda_{\varepsilon,L}}$, the lattice Laplacian Δ^{ε} is:

(1.2)
$$\forall \varphi \in \mathbb{R}^{\Lambda_{\varepsilon,L}}, \qquad (\Delta^{\varepsilon} \varphi)_x = \varepsilon^{-2} \sum_{y \sim x} \left[\varphi_y - \varphi_x \right],$$

and $a^{\varepsilon}(\lambda)$ is a dimension-dependent divergent counterterm which ensures that the $\varepsilon \to 0$ limit of the measure (on a suitable space of generalised functions on $L\mathbb{T}^d$) exists and is non-Gaussian. The construction of the limiting measures has a long history, and we give some references further below. The division of $\mu + a^{\varepsilon}(\lambda)$ into a finite mass term μ (which plays the role of a temperature of the model) and a divergent counterterm $a^{\varepsilon}(\lambda)$ is only determined up to an additive bounded constant. Explicitly, for an arbitrary fixed $m^2 > 0$, which we will usually take to be $m^2 = 1$, one can take $a^{\varepsilon}(\lambda) = a^{\varepsilon}(\lambda, m^2)$ with

(1.3)
$$a^{\varepsilon}(\lambda, m^2) := -3\lambda \left(-\Delta^{\varepsilon} + m^2 \right)^{-1}(0, 0) + 6\lambda^2 \left\| \left(-\Delta^{\varepsilon} + m^2 \right)^{-1}(0, \cdot) \right\|_{L^3}^3.$$

In this definition and subsequently, the matrix elements of $(-\Delta^{\varepsilon} + m^2)^{-1}$ are normalised with respect to the inner product $(u, v)_{\varepsilon} = \varepsilon^d \sum_{x \in \Lambda_{\varepsilon,L}} u(x)v(x)$ so that $(-\Delta^{\varepsilon} + m^2)^{-1}(x, y)$ converges to its continuum counterpart, and L^p norms are defined by $||f||_{L^p}^p = ||f||_{L^p(\Lambda_{\varepsilon,L})}^p = \varepsilon^d \sum_{x \in \Lambda_{\varepsilon,L}} |f(x)|^p$. This implies the following scaling (with dimension-dependent constants $c_i > 0$):

(1.4)
$$a^{\varepsilon}(\lambda, m^2) = \begin{cases} -c_1 \lambda \log(\varepsilon^{-2}) + O_{m^2,\lambda}(1) & (d=2), \\ -c_1 \lambda \varepsilon^{-1} + c_2 \lambda^2 \log(\varepsilon^{-2}) + O_{m^2,\lambda}(1) & (d=3). \end{cases}$$

^{*}University of Cambridge, Statistical Laboratory, DPMMS. E-mail: rb812@cam.ac.uk.

[†]University of Cambridge, Statistical Laboratory, DPMMS. E-mail: bd444@cam.ac.uk.

Note that when d = 2 the λ^2 counterterm in (1.3) is bounded and part of $O_{m^2,\lambda}(1)$ in (1.4), and could thus be dropped, but we will keep it to unify the presentation of both dimensions. Moreover, as discussed already, any value of $m^2 > 0$ can be used, and for this reason, we often do not explicitly mention m^2 in the notation $\nu_{\lambda,\mu}^{\varepsilon,L}$. Different choices of m^2 only correspond to a different choice of origin for μ . Expectation with respect to (1.1) is denoted by $\langle \cdot \rangle_{\lambda,\mu}^{\varepsilon,L}$.

The Glauber dynamics with invariant measure $\nu_{\lambda,\mu}^{\varepsilon,L}$, also known as the dynamical φ_d^4 model, is the solution to the systems of SDEs

(1.5)
$$d\varphi_t = (\Delta^{\varepsilon}\varphi_t - \lambda\varphi_t^3 - (\mu + a^{\varepsilon}(\lambda))\varphi_t) dt + \sqrt{2}dW_t^{\varepsilon,L}$$

where $dW_t^{\varepsilon,L}$ is space-time white noise on $\mathbb{R}_+ \times \Lambda_{\varepsilon,L}$, i.e., $W^{\varepsilon,L}$ is a Brownian motion on $L^2(\Lambda_{\varepsilon,L})$ (or, in other words, the $W_t^{\varepsilon,L}(x)$ are independent Brownian motions of variance ε^{-d} for $x \in \Lambda_{\varepsilon,L}$). For d = 3, the pathwise existence of the $\varepsilon \to 0$ limit is a main result of Hairer's theory of regularity structures [20–22]; for d = 2 this existence is simpler [12]; for d = 3 see also [10, 18, 36] and [24] for alternative approaches to the pathwise limit. By standard results for SDEs, the solution to (1.5) determines a Markov process with Dirichlet form given by

(1.6)
$$D(F) = \varepsilon^{-d} \sum_{x \in \Lambda_{\varepsilon,L}} \left\langle \left(\frac{\partial F}{\partial \varphi_x}\right)^2 \right\rangle_{\lambda,\mu}^{\varepsilon,L}$$

Our main result is a log-Sobolev inequality for the φ_d^4 measures, stated in Theorem 1.1 below. We formulate it for $\varepsilon > 0$ with uniform ε dependence (and also uniform L dependence when the susceptibility is bounded). All constants that appear in its statement could be made explicit. The log-Sobolev constant of (1.1) is the largest constant $\gamma = \gamma^{\varepsilon,L}(\lambda,\mu)$ such that

(1.7)
$$\forall F : \mathbb{R}^{\Lambda_{\varepsilon,L}} \to \mathbb{R}_+, \quad \operatorname{Ent}(F) \leqslant \frac{2}{\gamma} D(\sqrt{F}),$$

where, for F nonnegative, the relative entropy is defined by

(1.8)
$$\operatorname{Ent}(F) = \langle F \log F \rangle_{\lambda,\mu}^{\varepsilon,L} - \langle F \rangle_{\lambda,\mu}^{\varepsilon,L} \log \langle F \rangle_{\lambda,\mu}^{\varepsilon,L}.$$

The log-Sobolev inequality has many general implications, see the references [1,19] for reviews of these. In particular, it is equivalent to the hypercontractivity of the Markov semigroup associated with the Dirichlet form, and to its exponential relaxation to equilibrium in relative entropy sense for initial conditions of finite relative entropy.

For the continuum φ_d^4 measures, the log-Sobolev inequality was conjectured in [34]. Spectral gap inequalities (quantifying the relaxation in L^2 norm with respect to the invariant measure) and exponential ergodicity (in total variation) for the φ_d^4 measures in finite volume have been proven in [32] for d = 2 and extended to d = 3 in [23]. These results apply directly in the continuum limit $\varepsilon \to 0$. The Dirichlet form in the continuum limit was identified in [35]. A main ingredient of the proof of the above spectral gap inequalities is a qualitative support theorem for the continuum SPDEs and a compactness argument. In particular, the spectral gap obtained this way is not uniform in L and not explicit, and the validity of the log-Sobolev inequality also remained open.

We remark that, for any $\varepsilon > 0$ and $L < \infty$, the regularised φ_d^4 measure is convex at infinity and standard techniques (see, e.g., [25]) show that $\gamma^{\varepsilon,L}(\lambda,\mu) > 0$, but the bound obtained in this way tends to 0 (extremely quickly) as $\varepsilon \to 0$ or $L \to \infty$, as the counterterms (1.3)–(1.4) make the microscopic measure very non-convex as $\varepsilon \to 0$ (for any $\lambda > 0$ and $\mu \in \mathbb{R}$).

Theorem 1.1. Let d = 2 or d = 3 and $\varepsilon > 0$. Let $L \ge 1$ assumed to be a multiple of $\varepsilon > 0$. (i) Let $\lambda > 0$, $\mu \in \mathbb{R}$, and assume that there is a constant $\overline{\chi} \in (0, \infty)$ such that

(1.9)
$$\chi^{\varepsilon,L}(\lambda,\mu) := \varepsilon^d \sum_{x \in \Lambda_{\varepsilon,L}} \langle \varphi_0 \varphi_x \rangle_{\lambda,\mu}^{\varepsilon,L} \leqslant \bar{\chi}.$$

Then the log-Sobolev constant $\gamma^{\varepsilon,L}(\lambda,\mu)$ of (1.1) is bounded below by a positive constant $\bar{\gamma} = \bar{\gamma}(\lambda,\mu,\bar{\chi})$ uniformly in ε and the bound depends only on $(\lambda,\mu,\bar{\chi})$ and not directly on L (or ε):

(1.10)
$$\gamma^{\varepsilon,L}(\lambda,\mu) \ge \bar{\gamma}(\lambda,\mu,\bar{\chi}).$$

(ii) For any $\lambda, \mu > 0$, there are $\mu(d, \lambda), \lambda(d, \mu) > 0$ such that if the counterterms in (1.3)–(1.4) are chosen with $m^2 = \mu$ instead of $m^2 = 1$, and if either $\mu > \mu(d, \lambda)$ or $\lambda \in [0, \lambda(d, \mu)]$, then the log-Sobolev constant satisfies, uniformly in ε and L,

(1.11)
$$C^{-1}\mu \leqslant \gamma^{\varepsilon,L}(\lambda,\mu) \leqslant C\mu.$$

The choice $m^2 = \mu$ instead of $m^2 = 1$ in item (ii) amounts to a different choice of origin for μ , and (1.11) states that this choice ensures that the log-Sobolev constant is comparable to the mass term μ . Equivalently, this statement could be formulated in terms of $m^2 = 1$ when replacing μ in (1.11) on both sides by $\mu + a^{\varepsilon}(\lambda, \mu) - a^{\varepsilon}(\lambda, 1)$, where $a^{\varepsilon}(\lambda, \mu) - a^{\varepsilon}(\lambda, \mu)$ converges as $\varepsilon \to 0$.

Item (i) says that the log-Sobolev constant is bounded below as soon as the susceptibility is bounded above. This condition is optimal, in the sense that the converse implication also holds, i.e., the susceptibility is bounded above if the log-Sobolev constant is bounded below. Indeed, by using the trial function $F(\varphi) = \varepsilon^d L^{-d/2} \sum_{x \in \Lambda_{\varepsilon,L}} \varphi_x$ and observing that $\operatorname{Var}(F) = \chi^{\varepsilon,L}(\lambda,\mu)$ and D(F) = 1, the general fact that the spectral gap is bounded below by the log-Sobolev constant implies that the optimal log-Sobolev constant $\gamma^{\varepsilon,L}(\lambda,\mu)$ of (1.1) satisfies

(1.12)
$$\gamma^{\varepsilon,L}(\lambda,\mu) \leqslant \frac{1}{\chi^{\varepsilon,L}(\lambda,\mu)}.$$

Together with Theorem 1.1 (i) this shows that $\gamma^{\varepsilon,L}(\lambda,\mu)$ is bounded below if and only if $\chi^{\varepsilon,L}(\lambda,\mu)$ is bounded above.

The boundedness of the susceptibility $\chi^{\varepsilon,L}(\lambda,\mu)$ in both ε and L provides a standard definition of the critical temperature $\mu_c(\lambda)$. Thus, in other words, Theorem 1.1 (i) shows that the log-Sobolev constant of the φ_d^4 measure is bounded below throughout the high temperature phase $\mu > \mu_c(\lambda)$. For background, we summarise the following known properties of the susceptibility of φ_d^4 models (d = 2, 3), say with the counterterms (1.3) defined with $m^2 = 1$:

- (i) For any $\lambda > 0$ and $\mu \in \mathbb{R}$, the finite volume susceptibility $\chi^{\varepsilon,L}(\lambda,\mu)$ is bounded in ε for any fixed $L < \infty$. For example, this can be shown using the methods of the recent works [2,17].
- (ii) For any $\lambda > 0$, there is $\mu_0(\lambda) \in \mathbb{R}$ such that for $\mu > \mu_0(\lambda)$, the susceptibility is also bounded both in ε and L. For example, a very simple proof of this was given in [8] whose method of skeleton inequalities we also apply as an ingredient; see Section 4.
- (iii) For any $\lambda > 0$, there is $\mu_1(\lambda) \in \mathbb{R}$ such that for $\mu < \mu_1(\lambda)$, the susceptibility is not bounded in L. For d = 2, see [16], and for d = 3, see [14] and the recent work [11].
- (iv) By the second Griffiths inequality, the susceptibility is decreasing in μ for any fixed $\lambda > 0$, provided the counterterms are defined in terms of m^2 independent of μ , e.g., $m^2 = 1$. Thus with this choice (ii) and (iii) determine a unique $\mu_c(\lambda) \in \mathbb{R}$ that separates the μ for which the susceptibility is bounded or unbounded.

While the above shows that the log-Sobolev constant is positive throughout the high temperature phase, the bound obtained as $\mu \downarrow \mu_c(\lambda)$ is explicit in terms of the susceptibility but far from the expected truth. For large μ , item (ii) of Theorem 1.1 determines the log-Sobolev constant up to multiplicative constants.

For general background on the φ_2^4 measures, see the textbooks [15,30], and for φ_3^4 the discussions of the existing references in the introductions of [8,17]. Aside from the references already mentioned above, recent progress of the SPDE perspective include [26–28].

Our proof of Theorem 1.1 relies on four main ingredients: the criterion for the log-Sobolev inequality in terms of the Polchinski (renormalisation group) equation from [3], the recently proved correlation inequality for the Ising model with general external fields from [13], the Perron– Frobenius theorem, and improvements of the estimates for the φ_d^4 measures from [8]. As a simple application of the first part of the proof, we also show (the also new result) that the log-Sobolev constant of the lattice φ^4 model is bounded below uniformly in the volume up to the critical point (see Section 3). In [4], we apply related methods to the Ising model.

2 Criterion for the log-Sobolev inequality

In this section, we formulate a sufficient criterion for a log-Sobolev inequality for φ^4 measures. The criterion is first stated for φ^4 measures in a general setting (thus applying in particular to any graph and any dimension), in Theorem 2.1, and then specialised to the continuum φ_d^4 measure (1.1) in Theorem 2.6. The criterion could be further extended to more general spin systems in the Griffiths–Simon class [31]. We do not pursue this generalisation explicitly, but in [4], we present a variant of the criterion for the Ising model (with an additional ingredient needed to pass from continuous to discrete spins).

Let Λ be a finite set, let $A = (A_{x,y})_{x,y \in \Lambda} \subset \mathbb{R}$ a symmetric matrix, let g > 0 and $\nu \in \mathbb{R}$, and consider the general φ^4 measure with external field $h \in \mathbb{R}^{\Lambda}$ on Λ :

(2.1)
$$\mu_{A,g,\nu}^{\Lambda,h}(d\varphi) \propto \exp\left[-\frac{1}{2}(\varphi,A\varphi) - V_0(\varphi) + (h,\varphi)\right] d\varphi,$$

with $(u, v) = \sum_{x \in \Lambda} u_x v_x$ for $u, v \in \mathbb{R}^{\Lambda}$ (there is no ε here), and V_0 the potential:

(2.2)
$$\forall \varphi \in \mathbb{R}^{\Lambda}, \qquad V_0(\varphi) = \sum_{x \in \Lambda} \left(\frac{1}{4}g\varphi_x^4 + \frac{1}{2}\nu\varphi_x^2\right).$$

Since the set Λ is fixed in the following, we omit it and write $\mu_{A,g,\nu}^h$ instead of $\mu_{A,g,\nu}^{\Lambda,h}$, and $\langle \cdot \rangle_{A,g,\nu}^h$ for the associated expectation. Also, when h = 0, we simply write $\mu_{A,g,\nu}$ and $\langle \cdot \rangle_{A,g,\nu}$. In this setting, the log-Sobolev constant of (2.1) is the best constant $\gamma = \gamma(A, g, \nu)$ such that:

(2.3)
$$\forall F : \mathbb{R}^{\Lambda} \to \mathbb{R}_{+}, \quad \operatorname{Ent}(F) \leqslant \frac{2}{\gamma} D(\sqrt{F}),$$

with:

(2.4)
$$\operatorname{Ent}(F) = \langle F \log F \rangle_{A,g,\nu} - \langle F \rangle_{A,g,\nu} \log \langle F \rangle_{A,g,\nu}, \qquad D(F) = \sum_{x \in \Lambda} \left\langle \left(\frac{\partial F}{\partial \varphi_x}\right)^2 \right\rangle_{A,g,\nu}.$$

For t > 0, consider the measure $\mu_{A,g,\nu+1/t}$ with additional mass 1/t, and write $S_t = S_{A,g,\nu+1/t}$ for its two-point correlation function:

(2.5)
$$\forall x, y \in \Lambda, \qquad S_t(x, y) := \left\langle \varphi_x \varphi_y \right\rangle_{A, g, \nu + 1/t}.$$

Introduce also the susceptibility $\chi_t = \chi_{A,g,\nu+1/t}$:

(2.6)
$$\chi_t := \max_{x \in \Lambda} \sum_{y \in \Lambda} S_t(x, y)$$

In particular, when Λ is a torus and $A_{x,y} = A_{0,x-y}$ for all $x, y \in \Lambda$, then S_t is translation invariant as well, and χ_t is the usual susceptibility. Define then:

(2.7)
$$\dot{\kappa}_t = \frac{1}{t} - \frac{\chi_t}{t^2}$$

Theorem 2.1. Assume that A is positive definite, and that offdiagonal coefficients of A are nonpositive. The log-Sobolev constant is then bounded by:

(2.8)
$$\frac{1}{\gamma} \leqslant \int_0^\infty \exp\left[-2\int_0^t \dot{\kappa}_s \, ds\right] dt$$

Remark 2.2. The proof only uses that χ_t is an upper bound on the spectral radius of the covariance matrix $(S_t(x, y))_{x,y \in \Lambda}$ and one could therefore instead define χ_t in this way for a slightly more general statement.

The proof of Theorem 2.1 has several ingredients. The starting point is the general criterion for the log-Sobolev inequality derived from the Polchinski (renormalisation group) equation in [3]. Let us recall this general criterion. In [3, Section 2], this criterion is formulated for a general covariance decomposition (C_t) . We will make the particular choice (known as Pauli–Villars regularisation):

(2.9)
$$C_t = (A + 1/t)^{-1} \quad (t > 0), \qquad C_0 = 0,$$

and recall that the matrix A is positive definite. We write \dot{C}_t , \ddot{C}_t for the first and second derivatives of C_t . Let \mathbf{E}_{C_t} denote expectation under the Gaussian measure with covariance C_t , and introduce a renormalised potential V_t by:

(2.10)
$$\forall \varphi \in \mathbb{R}^{\Lambda}, \qquad V_t(\varphi) = -\log \mathbf{E}_{C_t} \Big[e^{-V_0(\varphi + \cdot)} \Big].$$

Equivalently, V_t satisfies the Polchinski (renormalisation group) equation

(2.11)
$$\frac{\partial V_t}{\partial t} = \frac{1}{2} \Delta_{\dot{C}_t} V_t - \frac{1}{2} (\nabla V_t)_{\dot{C}_t}^2,$$

see [3] for notation and discussion of this equation, and [29] for its original use. The Polchinski equation is not used directly in this paper, but it underlies the following proposition.

Proposition 2.3 (Theorem 2.5 in [3]). The log-Sobolev constant $\gamma = \gamma(A, g, \nu)$ for the measure $\mu_{A,g,\nu}$ is bounded in terms of V_t as follows:

(2.12)
$$\frac{1}{\gamma} \leqslant \int_0^\infty \exp\left[-2\int_0^t \dot{\ell}_s \, ds\right] dt,$$

where, for t > 0, $\dot{\ell}_t$ is any real number such that, for all $\varphi \in \mathbb{R}^{\Lambda}$:

(2.13)
$$\dot{C}_t \operatorname{Hess} V_t(\varphi) \dot{C}_t - \frac{1}{2} \ddot{C}_t \ge \dot{\ell}_t \dot{C}_t.$$

Thus we need to show that $\dot{\ell}_t$ may be chosen to be $\dot{\kappa}_t$. To this end, observe that V_t defined in (2.10) can be expressed, by changing variables from ζ to $\zeta - \varphi$, as

(2.14)
$$V_{t}(\varphi) + \text{const} = -\log \int e^{-\frac{1}{2}(\zeta, C_{t}^{-1}\zeta)} e^{-V_{0}(\varphi+\zeta)} d\zeta$$
$$= \frac{1}{2}(\varphi, C_{t}^{-1}\varphi) - \log \int e^{-\frac{1}{2}(\zeta, C_{t}^{-1}\zeta)} e^{-V_{0}(\zeta)} e^{(C_{t}^{-1}\varphi, \zeta)} d\zeta,$$

and hence the Hessian of V_t is given by:

(2.15)
$$\operatorname{Hess} V_t(\varphi) = C_t^{-1} - C_t^{-1} \Sigma_t(\varphi) C_t^{-1},$$

where

(2.16)
$$\Sigma_t(\varphi) = \left(\left\langle \zeta_x; \zeta_y \right\rangle_{A,g,\nu+1/t}^{C_t^{-1}\varphi} \right)_{x,y \in \Lambda},$$

with the expectation $\langle \cdot \rangle$ here acting on the field ζ and the following notation for the covariance:

(2.17)
$$\langle F;G\rangle_{A,g,\nu}^h = \langle FG\rangle_{A,g,\nu}^h - \langle F\rangle_{A,g,\nu}^h \langle G\rangle_{A,g,\nu}^h, \quad F,G:\mathbb{R}^\Lambda \to \mathbb{R}, \quad h \in \mathbb{R}^\Lambda.$$

The second ingredient of the proof of Theorem 2.1 is the remarkable correlation inequality [13, Corollary 1.3] which shows that the truncated two-point function of ferromagnetic Ising models with general external field is maximised pointwise when the external field is 0, recently proved by Ding–Song–Sun. By the Griffiths–Simon construction [31], this inequality extends to φ^4 models as follows (the proof of this extension is exactly as the extension of the other correlation inequalities from Ising models to φ^4 models in [31]).

Proposition 2.4. Let g > 0, $\nu \in \mathbb{R}$, and let A be a matrix with nonpositive off-diagonal coefficients. For any $h \in \mathbb{R}^{\Lambda}$, then

(2.18)
$$\forall x, y \in \Lambda, \qquad 0 \leqslant \left\langle \varphi_x; \varphi_y \right\rangle_{A,g,\nu}^h \leqslant \left\langle \varphi_x; \varphi_y \right\rangle_{A,g,\nu}^0 = \left\langle \varphi_x \varphi_y \right\rangle_{A,g,\nu}^0$$

To use this inequality to bound the Hessian in (2.15) requires a third additional ingredient, as a quadratic form instead of pointwise bound is needed. This ingredient in the proof of Theorem 2.1 is the Perron–Frobenius theorem, which enables us to bound the eigenvalues of the Hessian of V_t in (2.15) through Proposition 2.4. The combination of the last two ingredients culminates in the following proposition from which Theorem 2.1 follows immediately.

Proposition 2.5. For each t > 0, one can take $\dot{\ell}_t = \dot{\kappa}_t$ in (2.13), where $\dot{\kappa}_t$ is defined by (2.7).

Proof of Proposition 2.5. Fix t > 0, take $X \in \mathbb{R}^{\Lambda}$, and using the expression (2.15) for the Hessian notice that

(2.19)
$$X^{T} \operatorname{Hess} V_{t}(\varphi) X \ge X^{T} C_{t}^{-1} X - \|C_{t}^{-1} X\|_{2}^{2} \sup_{\|Y\|_{2}=1} Y^{T} \Sigma_{t}(\varphi) Y.$$

The correlation matrix $\Sigma_t(\varphi)$ has nonnegative entries, as correlations under $\mu_{A,g,\nu}^h$ are nonnegative for any $h \in \mathbb{R}^{\Lambda}$ by the FKG inequality. By the Perron–Frobenius theorem, the spectral radius of the correlation matrix is therefore an eigenvalue, and there is an associated (normalised) eigenvector $Y_0 = Y_0(\varphi)$ with nonnegative entries. The pointwise correlation bound of Proposition 2.4 thus implies:

(2.20)
$$\sup_{\|Y\|_{2}=1} Y^{T} \Sigma_{t}(\varphi) Y = Y_{0}^{T} \Sigma_{t}(\varphi) Y_{0} \leqslant Y_{0}^{T} \Sigma_{t}(0) Y_{0} \leqslant \|\Sigma_{t}(0)\| = \sup_{\|Y\|_{2}=1} Y^{T} \Sigma_{t}(0) Y.$$

The spectral radius $\|\Sigma\|$ of a matrix Σ is a lower bound for any matrix norm. For a matrix Σ with nonnegative entries, this implies:

(2.21)
$$\|\Sigma\| \leqslant \max_{x} \sum_{y} \Sigma_{x,y}.$$

For $\Sigma = \Sigma_t(0)$ given by (2.16), the right-hand side of (2.21) is precisely the susceptibility (2.6). This and (2.19) imply the following bound on the lowest eigenvalue of the Hessian of V_t :

(2.22)
$$X^T \operatorname{Hess}(V_t)(\varphi) X \ge X^T \left(C_t^{-1} - \chi_t C_t^{-2} \right) X.$$

To conclude the proof of Proposition 2.5, notice that, for each t > 0:

(2.23)
$$\dot{C}_t = \frac{1}{t^2} (A + 1/t)^{-2} = \frac{C_t^2}{t^2}, \qquad \ddot{C}_t = -\frac{2}{t^3} A (A + 1/t)^{-3} = -\frac{2}{t} A C_t \dot{C}_t.$$

Injecting these expressions in the criterion (2.13) for the log-Sobolev inequality concludes the proof of Proposition 2.5 (C_t , \dot{C}_t , \ddot{C}_t , and A all commute):

$$\dot{C}_{t} \operatorname{Hess}(V_{t})(\varphi)\dot{C}_{t} - \frac{1}{2}\ddot{C}_{t} \geqslant \left[\frac{C_{t}}{t}\left(C_{t}^{-1} - \chi_{t}C_{t}^{-2}\right)\frac{C_{t}}{t} + \frac{1}{t}AC_{t}\right]\dot{C}_{t}$$

$$= \left[\frac{1}{t}\left(\frac{1}{t} + A\right)C_{t} - \frac{\chi_{t}}{t^{2}}\right]\dot{C}_{t}$$

$$= \left[\frac{1}{t} - \frac{\chi_{t}}{t^{2}}\right]\dot{C}_{t} = \dot{\kappa}_{t}\dot{C}_{t}.$$

$$\Box$$

Proof of Theorem 2.1. As already mentioned, the proof of the theorem is immediate from Propositions 2.3 and 2.5. \Box

Finally, we consider the special choice of Λ , A, g, ν corresponding to the continuum φ_d^4 measure $\nu_{\lambda,\mu}^{\varepsilon,L}$ defined in (1.1), with an arbitrary $m^2 > 0$ to define the counterterms (1.3). To emphasise the value of $m^2 > 0$ used to define the counterterms, we write $\nu_{\lambda,\mu,m^2}^{\varepsilon,L}$ and $\langle \cdot \rangle_{\lambda,\mu,m^2}^{\varepsilon,L}$ for the measure and its expectation. This choice is:

(2.25)
$$\Lambda = \Lambda_{\varepsilon,L}, \quad A = \varepsilon^d \left(-\Delta^{\varepsilon} + m^2 \right), \quad g = \varepsilon^d \lambda, \quad \nu = \varepsilon^d (\mu - m^2 + a^{\varepsilon} (\lambda, m^2)).$$

The factors of ε^d result from the fact that we did not include factors ε^d in (2.2) and the use of the standard inner product $(u, v) = \sum_x u_x v_x$ in this section. In the application to the continuum model, it makes sense to normalise the scale parameter t in continuum scaling also, i.e., to rescale t by ε^d . This means that we consider the measures $\nu_{\lambda,\mu+1/t,m^2}^{\varepsilon,L}$ and the correspondingly normalised susceptibility:

(2.26)
$$\chi_t^{\varepsilon,L}(\lambda,\mu,m^2) := \max_{x \in \Lambda_{\varepsilon,L}} \varepsilon^d \sum_{y \in \Lambda_{\varepsilon,L}} \left\langle \varphi_x \varphi_y \right\rangle_{\lambda,\mu+1/t,m^2}^{\varepsilon,L} = \varepsilon^d \sum_{x \in \Lambda_{\varepsilon,L}} \left\langle \varphi_0 \varphi_x \right\rangle_{\lambda,\mu+1/t,m^2}^{\varepsilon,L},$$

where the last equality is by translation invariance, and we set:

(2.27)
$$\dot{\kappa}_t^{\varepsilon,L} := \frac{1}{t} - \frac{\chi_t^{\varepsilon,L}(\lambda,\mu,m^2)}{t^2}$$

Theorem 2.1 then takes the following form. The log-Sobolev constant in its statement is normalised as in (1.7) with respect to the continuum normalised Dirichlet form (1.6). Note that A in (2.25) is indeed positive definite with nonpositive offdiagonal entries.

Theorem 2.6. Let $\lambda > 0$ and $\mu \in \mathbb{R}$. The log-Sobolev constant $\gamma^{\varepsilon,L}(\lambda,\mu,m^2)$ of (1.1) normalised as in (1.7) and with counterterms defined in terms of mass $m^2 > 0$ satisfies the bound:

(2.28)
$$\frac{1}{\gamma^{\varepsilon,L}(\lambda,\mu,m^2)} \leqslant \int_0^\infty \exp\left[-2\int_0^t \dot{\kappa}_s^{\varepsilon,L} \, ds\right] dt.$$

3 Simple application: Log-Sobolev inequality for lattice φ^4 models

To illustrate the criterion for the log-Sobolev inequality from Theorem 2.1, we give a simple proof that the log-Sobolev constant of the lattice φ^4 spin model in any dimension is bounded uniformly in the volume throughout its entire high temperature phase. For unbounded spin systems including the lattice φ^4 measure, the existence of a spectral gap uniformly in the volume was previously known only away from the critical temperature [5, 25, 33]. Fix a dimension $d \ge 1$. For simplicity of the exposition, we assume that $\Lambda \subset \mathbb{Z}^d$ is a hypercube with periodic boundary conditions, i.e., a discrete torus, although Dirichlet boundary conditions could be considered with essentially no change. For g > 0, $\nu \in \mathbb{R}$, recall that the lattice φ^4 model on Λ is defined by

(3.1)
$$\mu_{g,\nu}^{\Lambda}(d\varphi) = e^{-\frac{1}{2}(\varphi, -\Delta\varphi) - \sum_{x \in \Lambda} (\frac{1}{4}g\varphi^4 + \nu\varphi^2)},$$

where Δ is the lattice Laplacian on Λ , and its susceptibility is defined by

(3.2)
$$\chi^{\Lambda}(g,\nu) = \sum_{x \in \Lambda} \langle \varphi_0 \varphi_x \rangle_{g,\nu}.$$

The critical value of the *d*-dimensional lattice φ^4 model can be defined by

(3.3)
$$\nu_c(g) = \inf\{\nu \in \mathbb{R} : \sup_{\Lambda} \chi^{\Lambda}(g,\nu) < +\infty\},$$

where the supremum is over all d-dimensional discrete tori. In fact, $\nu_c(g) < 0$ for g > 0.

Example 3.1. The log-Sobolev constant $\gamma = \gamma_{q,\nu}^{\Lambda}$ of the measure (3.1) satisfies

(3.4)
$$\frac{1}{\gamma} \leqslant \frac{e^2}{2|\nu|+1} + (2|\nu|+1)^3 e^{2+2(2|\nu|+1)\chi^{\Lambda}(g,\nu)} < \infty.$$

Thus for any g > 0 and $\nu \in \mathbb{R}$ such that $\chi^{\Lambda}(g,\nu)$ is bounded uniformly in Λ , the log-Sobolev constant γ is bounded below uniformly in Λ . Since on the other hand the lower boundedness of the log-Sobolev constant also implies that the susceptibility is bounded (analogously to the discussion below Theorem 1.1), it thus follows that the log-Sobolev constant is bounded below if and only if the susceptibility is bounded.

Proof. Note that the interesting case is $\nu \leq 0$; otherwise the potential is convex and one could use the Bakry–Emery criterion to show the log-Sobolev inequality. In bounding the susceptibility χ_t , let us separately treat the small and large t cases as follows.

In the small t case given by $t \leq 1/(2|\nu|+1)$, the measure $\mu_{g,\nu+1/t}$ is of the form $e^{-U(\varphi)} d\varphi$, where $U = \frac{1}{2}(\varphi, (-\Delta + \nu + 1/t)\varphi)$ + convex is a strictly convex potential, with Hessian bounded below by $(1/t + \nu)$ id > 0 id. Therefore the Brascamp–Lieb inequality [6, Theorem 4.1] implies

(3.5)
$$\chi_t = \operatorname{Var}_{\mu_{g,\nu+1/t}^{\Lambda}} \left((\varphi, \mathbf{1}) \right) \leq \frac{1}{\frac{1}{t} + \nu}, \qquad \mathbf{1} := \frac{1}{|\Lambda|^{1/2}} (1, \dots, 1).$$

Recalling that $\dot{\kappa}_t = 1/t - \chi_t/t^2$ is the quantity appearing in the bound of the log-Sobolev constant in Theorem 2.1, one has:

(3.6)
$$\forall t \leq 1/(2|\nu|+1), \qquad \dot{\kappa}_t \geq \frac{1}{t} \left(1 - \frac{1}{1+t\nu}\right) = \frac{\nu}{1+t\nu}.$$

Consider now the large t case where $t > 1/(2|\nu| + 1)$. Since $\chi^{\Lambda}(g,\nu)$ is decreasing in $\nu \in \mathbb{R}$ by the second Griffiths inequality,

(3.7)
$$\forall t > 0, \qquad \dot{\kappa}_t \ge \frac{1}{t} - \frac{\chi^{\Lambda}(g, \nu + 1/t)}{t^2} \ge \frac{1}{t} - \frac{\chi^{\Lambda}(g, \nu)}{t^2}.$$

Using this bound on $\dot{\kappa}_t$ for $t > 1/(2|\nu|+1)$ and the small t bound (3.6) for $t \leq 1/(2|\nu|+1)$:

$$\forall t > 0, \qquad \kappa_t := \int_0^t \dot{\kappa}_s \, ds \ge \int_0^{t \wedge 1/(2|\nu|+1)} (-2|\nu|) \, dt + \int_{1/(2|\nu|+1)}^{t \vee 1/(2|\nu|+1)} \left(\frac{1}{t} - \frac{\chi^{\Lambda}(g,\nu)}{t^2}\right) dt$$

$$(3.8) \qquad \ge -1 + \mathbf{1}_{t > 1/(2|\nu|+1)} \Big[\log(t/(2|\nu|+1)) - (2|\nu|+1)\chi^{\Lambda}(g,\nu) \Big].$$

This implies:

(3.9)
$$\forall t > 0, \quad e^{-2\kappa_t} \leqslant e^2 \left[\mathbf{1}_{t \leqslant 1/(2|\nu|+1)} + \frac{(2|\nu|+1)^2}{t^2} e^{2(2|\nu|+1)\chi^{\Lambda}(g,\nu)} \mathbf{1}_{t > 1/(2|\nu|+1)} \right].$$

Therefore, by Theorem 2.1, the log-Sobolev constant satisfies

(3.10)
$$\frac{1}{\gamma_{\Lambda}} \leq \frac{e^2}{2|\nu|+1} + (2|\nu|+1)^3 e^{2+2(2|\nu|+1)\chi^{\Lambda}(g,\nu)}$$

as needed.

4 Bound on the susceptibility of the continuum φ_d^4 model

In dimensions $d \in \{2,3\}$, consider the continuum φ_d^4 measure (1.1) with parameters $\lambda > 0$ and $\mu \in \mathbb{R}$. In this section, we will make the dependence on the mass m^2 of the counterterms (1.3) explicit. The measure (1.1) is thus denoted by $\nu_{\lambda,\mu,m^2}^{\varepsilon,L}$, and the associated expectation by $\langle \cdot \rangle_{\lambda,\mu,m^2}^{\varepsilon,L}$.

4.1. Statement of the result. In order to apply Theorem 2.6, which will be done in Section 5, our goal in this section is to estimate, for t > 0, the susceptibility $\chi_t^{\varepsilon,L}(\lambda,\mu,m^2)$ of the φ_d^4 measure with mass $\mu + 1/t$ (and counterterm defined in terms of $m^2 > 0$), defined by

(4.1)
$$\chi_t^{\varepsilon,L}(\lambda,\mu,m^2) := \varepsilon^d \sum_{x \in \Lambda_{\varepsilon,L}} \left\langle \varphi_0 \varphi_x \right\rangle_{\lambda,\mu+1/t,m^2}^{\varepsilon,L}.$$

The parameter t is the scale parameter in the Polchinski (renormalisation group) equation (2.11). Informally, we speak of large scale when t is large, and of small scale when t is small.

For t > 0, further introduce the notation:

(4.2)
$$S_t(x) := \left\langle \varphi_0 \varphi_x \right\rangle_{\lambda, \mu+1/t, m^2}^{\varepsilon, L}, \qquad x \in \Lambda_{\varepsilon, L},$$

(4.3)
$$C_t(x) := \left(-\Delta^{\varepsilon} + m^2 + \frac{1}{t}\right)^{-1}(0, x), \qquad x \in \Lambda_{\varepsilon, L},$$

and note that both functions are positive (e.g., by the FKG inequality). Define also the shorthand:

(4.4)
$$m_t^2 := m^2 + \frac{1}{t},$$

and recall the definition of the norms $\|\cdot\|_{L^p}, \|\cdot\|_{L^{\infty}}$ norms for $p \ge 1$:

(4.5)
$$\|f\|_{L^p} := \left(\varepsilon^d \sum_{x \in \Lambda_{\varepsilon,L}} |f(x)|^p\right)^{1/p}, \qquad \|f\|_{L^\infty} := \max_{x \in \Lambda_{\varepsilon,L}} |f(x)|, \qquad f \in \mathbb{R}^{\Lambda_{\varepsilon,L}}.$$

Finally, introduce for t > 0 the differences η_t, γ_t between counterterms defined at masses m^2 and m_t^2 (recall the definition (1.3) of the counterterm $a^{\varepsilon}(\lambda, m^2)$):

(4.6)
$$\eta_t = C_{\infty}(0) - C_t(0) \ge 0, \qquad \gamma_t = \|C_{\infty}^3\|_{L^1} - \|C_t^3\|_{L^1} \ge 0.$$

In the following c > 0 denotes a constant that is independent from the parameters $\varepsilon, L, \lambda, \mu, m^2$ of the model and may change from line to line. To express dependence on a parameter, say μ , we write $c(\mu)$.

Since $\chi_t^{\check{\varepsilon},L}(\lambda,\mu,m^2) = \|S_t\|_{L^1}$ and $\|C_t\|_{L^1} = m_t^{-2}$ the desired estimates on the susceptibility $\chi_t^{\varepsilon,L}(\lambda,\mu,m^2)$ will be seen to be a direct consequence of the following propositions.

Proposition 4.1 (Small scale, all couplings). Let $d \in \{2,3\}, \lambda > 0, \mu \in \mathbb{R}$, and $m^2 > 0$. Then there is $t_0 = t_0(d, \lambda, \mu, m^2) > 0$ and a polynomial $p_{d,t,\mu,m^2}(\lambda)$ in λ of degree independent of the parameters, with coefficients functions of $\eta_t, \gamma_t, \mu, m^2$ times positive powers of m_t^{-1} (in particular independent of ε , L), such that:

(4.7)
$$\forall t \in (0, t_0], \qquad \|S_t - C_t\|_{L^1} \leq p_{d,t,\mu,m}(\lambda),$$

and $t \mapsto t^{-2} p_{d,t,\mu,m}(\lambda)$ is integrable on $(0, t_0]$.

When $\mu > 0$, one can make the special choice $m^2 = \mu$ as in the statement of Theorem 1.1 (ii). In this case, the results of Proposition 4.1 are valid for large t (i.e., large scale) provided either λ is small enough, or $\mu = m^2$ is large enough.

Proposition 4.2 (All scales, one small coupling). Let $d \in \{2, 3\}$, $\lambda > 0$, $\mu > 0$, and set $m^2 := \mu$. There are then $\lambda(d,\mu), \mu(d,\lambda) > 0$ and a polynomial $\tilde{p}_{d,t,\mu}(\lambda)$ in λ , again of degree independent of the parameters and with coefficients functions of η_t, γ_t, μ times positive powers of m_t^{-1} (but independent of ε , L), such that the following holds: if either $\lambda \in [0, \lambda(d, \mu))]$ or $\mu \ge \mu(d, \lambda)$, then:

(4.8)
$$\forall t > 0, \qquad \|S_t - C_t\|_{L^1} \leq \tilde{p}_{d,t,\mu}(\lambda),$$

the function $t \mapsto t^{-2} \tilde{p}_{d,t,\mu}(\lambda)$ is integrable on $(0,\infty)$, and the integral is independent of λ and μ if either $\lambda \in [0, \lambda(d, \mu)]$ or $\mu \ge \mu(d, \lambda)$. Under the same conditions,

(4.9)
$$\limsup_{t \to \infty} \tilde{p}_{d,t,\mu}(\lambda) \underset{\mu \to \infty}{=} o(\mu^{-1}).$$

4.2. Skeleton inequalities. To prove Propositions 4.1–4.2, the starting point is a bound on correlations presented in Proposition 4.3 below, obtained by Brydges, Fröhlich, and Sokal [8], using the method of skeleton inequalities [9] (which is based on the random walk representation [7]). To state it, let \star denote the convolution:

(4.10)
$$(f \star g)(x) = \varepsilon^d \sum_{y \in \Lambda_{\varepsilon,L}} f(x-y)g(y), \qquad f, g \in \mathbb{R}^{\Lambda_{\varepsilon,L}}.$$

For a function $f \in \mathbb{R}^{\Lambda_{\varepsilon,L}}$, write also:

(4.11)
$$(\mathbf{1}_0^{\varepsilon}f)(x) := \varepsilon^{-d}\mathbf{1}_{x=0}f(0).$$

The following inequalities are exact upper and lower bounds on the two-point function S of the φ_d^4 model, consistent with naive perturbation theory except that the right-hand sides also involve the interacting two-point function (or propagator) S rather than only the noninteracting one C.

Proposition 4.3 (Equations (5.12)–(5.13) in [8]). Consider the φ_d^4 measure (1.1) with parameters $\mu + 1/t \in \mathbb{R}$ (in place of μ), $\lambda > 0$, and $m^2 > 0$. Then, for each $x \in \Lambda_{\varepsilon,L}$:

n .

(4.12)
$$S_t(x) - C_t(x) \ge -3\lambda S_t(0)(C_t \star S_t)(x) + 6\lambda^2 (C_t \star S_t^3 \star S_t)(x) - 54\lambda^3 (C_t \star Q_t \star S_t)(x) - (a^{\varepsilon}(\lambda, m^2) + \mu - m^2)(S_t \star C_t)(x),$$

and:

(4.13)
$$S_t(x) - C_t(x) \leqslant -3\lambda S_t(0)(C_t \star S_t)(x) + 6\lambda^2 (C_t \star S_t^3 \star S_t)(x) - (a^{\varepsilon}(\lambda, m^2) + \mu - m^2)(S_t \star C_t)(x),$$

where:

(4.14)
$$Q_t := S_t \left(S_t^2 \star S_t^2 \right).$$

By writing the counterterms $a^{\varepsilon}(\lambda, m^2)$ from (1.3) in terms of $C_{\infty}(x) = (-\Delta^{\varepsilon} + m^2)^{-1}(0, x)$, the last proposition implies:

$$|S_t(x) - C_t(x)| \leq 3\lambda |S_t(0) - C_{\infty}(0)| (C_t \star S_t)(x) + 6\lambda^2 | (C_t \star (S_t^3 - \mathbf{1}_0^{\varepsilon} || C_{\infty}^3 ||_{L^1}) \star S_t)(x)| + 54\lambda^3 (C_t \star Q_t \star S_t)(x) + |\mu - m^2| (C_t \star S_t)(x).$$

Equation (4.15) is the starting point for the proof of Propositions 4.1–4.2, which takes the rest of Section 4. We use the short-hands:

(4.16)
$$C := C_t, \quad S := S_t, \quad Q := Q_t, \quad E := S - C.$$

When using (4.15) to prove Proposition 4.1–4.2, there are two main differences to the computations done in [8]. The first (minor) difference is that we allow a possibly negative mass $\mu+1/t$, and the counterterms are defined at mass $m^2 > 0$. In contrast, the counterterms in [8] are defined at the same mass $m^2 > 0$ as the one defining the massive Gaussian measure to which the φ^4 measure is compared to. The second (more significant) difference is that we need to pay attention to the *t*-dependence in all bounds: Firstly, the counterterms are defined at mass m^2 , not $m_t^2 = m^2 + 1/t$. Secondly, the resulting bounds on $||S_t - C_t||_{L^1}$ need to be integrable with respect to $t^{-2} dt$ since we are after an estimate on the susceptibility $||S_t||_{L^1}$ in order to estimate the log-Sobolev constant using Theorem 2.6.

Due to the above observations, it is convenient to split the the differences $S(0) - C_{\infty}(0)$ and $S^3 - \mathbf{1}_0^{\varepsilon} \| C_{\infty}^3 \|_{L^1}$ as follows:

(4.17)
$$S(0) - C_{\infty}(0) = S(0) - C(0) + C(0) - C_{\infty}(0) =: S(0) - C(0) - \eta_{t}$$
$$S^{3} - \mathbf{1}_{0}^{\varepsilon} \| C_{\infty}^{3} \|_{L^{1}} = \left(S^{3} - C^{3} \right) + \left(C^{3} - \mathbf{1}_{0}^{\varepsilon} \| C^{3} \|_{L^{1}} \right) - \mathbf{1}_{0}^{\varepsilon} \gamma_{t},$$

where we recall:

(4.18)
$$\eta_t = C_{\infty}(0) - C(0) \ge 0, \qquad \gamma_t = \|C_{\infty}^3\|_{L^1} - \|C^3\|_{L^1} \ge 0.$$

Let us lastly remark that, while proving Proposition 4.1–4.2 ultimately requires bounding $||E||_{L^1}$, due to the term $S(0) - C_{\infty}(0)$ in (4.15), $||E||_{L^1}$ can be bounded through (4.15) only in terms of both $||E||_{L^1}$ and $||E||_{L^{\infty}}$. To prove Propositions 4.1–4.2, we thus first estimate:

$$(4.19) ||E||_{L^1 \cap L^\infty} := ||E||_{L^1} + ||E||_{L^\infty}.$$

The resulting bound on $||E||_{L^1 \cap L^\infty}$ of course implies a bound on $||E||_{L^1}$, but the *t*-dependence of this bound turns out to be neither optimal, nor sufficient for our purposes. However, using that $||E||_{L^\infty}$ is already controlled as input, we improve the bound by rerunning the argument in the L^1 norm only, which then yields the claim of both propositions.

4.3. Estimate in $\|\cdot\|_{L^1\cap L^\infty}$ **norm.** We separately compute the terms related to each power of λ appearing in (4.15). The next lemma gathers useful properties and estimates. The bounds on moments of C are proven in Appendix A.

Lemma 4.4. (i) For t > 0, recall the shorthand $m_t^2 := m^2 + 1/t$ and $C := C_t$. One has:

(4.20)
$$||C||_{L^1} = \frac{1}{m_t^2}, \qquad ||C^2||_{L^1} \leqslant \frac{c}{m_t^2} \mathbf{1}_{d=2} + \frac{c}{m_t} \mathbf{1}_{d=3}.$$

(ii) If $f, g \in \mathbb{R}^{\Lambda_{\varepsilon,L}}$, then:

$$(4.21) ||f \star g||_{L^1} \leqslant ||f||_{L^1} ||g||_{L^1}, ||f \star g||_{L^\infty} \leqslant ||f||_{L^2} ||g||_{L^2}, ||f \star g||_{L^\infty} \leqslant ||f||_{L^1} ||g||_{L^\infty}.$$

In fact $||f \star g||_{L^1} = ||f||_{L^1} ||g||_{L^1}$ when $f, g \ge 0$.

4.3.1. Order 0 term. The only term independent of λ in the bound (4.15) is:

(4.22)
$$T_{\lambda^0} := |\mu - m^2| C \star S = |\mu - m^2| (C \star C + C \star E).$$

Using Lemma 4.4, its $\|\cdot\|_{L^1}$ and $\|\cdot\|_{L^{\infty}}$ norms satisfy:

(4.23)
$$\|T_{\lambda^{0}}\|_{L^{\infty}} \leq |\mu - m^{2}| \Big(\|C\|_{L^{1}} \|E\|_{L^{\infty}} + \|C\|_{L^{2}}^{2} \Big)$$
$$\leq |\mu - m^{2}| \Big(\frac{\|E\|_{L^{\infty}}}{m_{t}^{2}} + \frac{c}{m_{t}^{2}} \mathbf{1}_{d=2} + \frac{c}{m_{t}} \mathbf{1}_{d=3} \Big),$$

and:

(4.24)
$$\|T_{\lambda^0}\|_{L^1} \leq |\mu - m^2| \left(\frac{\|E\|_{L^1}}{m_t^2} + \frac{1}{m_t^4}\right).$$

As a result,

(4.25)
$$||T_{\lambda^0}||_{L^1 \cap L^\infty} \leq |\mu - m^2| \Big[\frac{||E||_{L^1 \cap L^\infty}}{m_t^2} + \frac{1}{m_t^4} + \frac{c}{m_t^2} \mathbf{1}_{d=2} + \frac{c}{m_t} \mathbf{1}_{d=3} \Big].$$

4.3.2. Order 1 term. The first order term is defined by:

(4.26)
$$T_{\lambda} := 3\lambda \big[S(0) - C(0) + C(0) - C_{\infty}(0) \big] S \star C = 3\lambda \big[S(0) - C(0) - \eta_t \big] S \star C,$$

where η_t is the difference (4.18) between counterterms with mass m^2 and m_t^2 . It is computed in Lemma A.3. As for T_{λ^0} , the $\|\cdot\|_{L^{\infty}}$ norm of the first order term T_{λ} reads:

(4.27)
$$\|T_{\lambda}\|_{L^{\infty}} \leq 3\lambda (\|E\|_{L^{\infty}} + \eta_t) \Big(\|C\|_{L^1} \|E\|_{L^{\infty}} + \|C\|_{L^2}^2 \Big)$$
$$\leq 3\lambda \big(\eta_t + \|E\|_{L^{\infty}}\big) \Big(\frac{\|E\|_{L^{\infty}}}{m_t^2} + \frac{c}{m_t^2} \mathbf{1}_{d=2} + \frac{c}{m_t} \mathbf{1}_{d=3} \Big)$$

In contrast, its $\|\cdot\|_{L^1}$ norm is simply:

(4.28)
$$\|T_{\lambda}\|_{L^{1}} \leq 3\lambda \big(\eta_{t} + \|E\|_{L^{\infty}}\big) \|C \star S\|_{L^{1}} \leq \frac{3\lambda \big(\eta_{t} + \|E\|_{L^{\infty}}\big)}{m_{t}^{2}} \Big(\|E\|_{L^{1}} + \frac{1}{m_{t}^{2}}\Big).$$

We conclude on the first order term:

(4.29)
$$\|T_{\lambda}\|_{L^{1}\cap L^{\infty}} \leq c\lambda \Big(\eta_{t} + \|E\|_{L^{1}\cap L^{\infty}}\Big) \Big[\frac{\|E\|_{L^{1}\cap L^{\infty}}}{m_{t}^{2}} + \frac{1}{m_{t}^{4}} + \frac{1}{m_{t}^{2}}\mathbf{1}_{d=2} + \frac{1}{m_{t}}\mathbf{1}_{d=3}\Big].$$

4.3.3. Order 2 terms. The second order contribution T_{λ^2} corresponds to the following terms:

(4.30)
$$T_{\lambda^2} := 6\lambda^2 \Big[C \star (S^3 - C^3) \star S + C \star (\mathbf{1}_0^{\varepsilon} \| C^3 \|_{L^1} - \mathbf{1}_0^{\varepsilon} \| C_\infty^3 \|_{L^1}) \star S + C \star \psi \star S \Big],$$

where ψ is the function:

(4.31)
$$\psi := C^3 - \mathbf{1}_0^{\varepsilon} \| C^3 \|_{L^1}.$$

To compute the term involving $S^3 - C^3$, write:

(4.32)
$$C \star (S^3 - C^3) \star S = C \star E (3C^2 + 3EC + E^2) \star (E + C).$$

The $\|\cdot\|_{L^{\infty}}$ norm of this term then reads:

$$\|C \star (S^{3} - C^{3}) \star S\|_{L^{\infty}} \leq 3\|E\|_{L^{\infty}} \|C \star C^{2} \star C\|_{L^{\infty}} + 3\|E\|_{L^{\infty}}^{2} \left(\|C \star C \star C\|_{L^{\infty}} + \|C \star C^{2}\|_{L^{1}}\right)$$

$$+ 4\|E\|_{L^{\infty}}^{3} \|C\|_{L^{1}}^{2} + \|E\|_{L^{\infty}}^{3} \|E\|_{L^{1}} \|C\|_{L^{1}}.$$

Cauchy-Schwarz inequality and Lemma 4.4 yield an estimate of the terms involving C:

(4.34)
$$\|C \star C \star C\|_{L^{\infty}} \leq \|C\|_{L^{1}} \|C\|_{L^{2}}^{2} = \frac{1}{m_{t}^{2}} \cdot \begin{cases} \frac{c}{m_{t}^{2}} & \text{if } d = 2, \\ \frac{c}{m_{t}} & \text{if } d = 3, \end{cases}$$

(4.35)
$$\|C \star C^2 \star C\|_{L^{\infty}} \leq \|C \star C\|_{L^{\infty}} \|C^2\|_{L^1} \leq \|C^2\|_{L^1}^2 \leq \begin{cases} \frac{c}{m_t^4} & \text{if } d = 2\\ \frac{c}{m_t^2} & \text{if } d = 3 \end{cases}$$

(4.36)
$$\|C^2 \star C\|_{L^1} = \|C^2\|_{L^1} \|C\|_{L^1} \leqslant \frac{1}{m_t^2} \cdot \begin{cases} \frac{c}{m_t^2} & \text{if } d = 2, \\ \frac{c}{m_t} & \text{if } d = 3. \end{cases}$$

It follows that (4.33) becomes:

Similarly, we find for the $\|\cdot\|_{L^1}$ norm:

$$\|C \star (S^{3} - C^{3}) \star S\|_{L^{1}} \leqslant \|C \star S\|_{L^{1}} \|E(3C^{2} + 3EC + E^{2}\|_{L^{1}} \\ \leqslant \frac{1}{m_{t}^{2}} \Big(\frac{1}{m_{t}^{2}} + \|E\|_{L^{1}}\Big) \Big(3\|E\|_{L^{\infty}} \|C^{2}\|_{L^{1}} + 3\|E\|_{L^{\infty}}^{2} \|C\|_{L^{1}} + \|E\|_{L^{\infty}}^{2} \|E\|_{L^{1}}\Big) \\ \leqslant \frac{1}{m_{t}^{2}} \Big(\frac{1}{m_{t}^{2}} + \|E\|_{L^{1}}\Big) \Big[c\|E\|_{L^{\infty}} \Big(\frac{1}{m_{t}^{2}} \mathbf{1}_{d=2} + \frac{1}{m_{t}} \mathbf{1}_{d=3}\Big) + \|E\|_{L^{\infty}}^{2} \Big(\frac{3}{m_{t}^{2}} + \|E\|_{L^{1}}\Big) \Big].$$

$$(4.38)$$

We conclude on the $\|\cdot\|_{L^1\cap L^{\infty}}$ norm of $C \star (S^3 - C^3) \star S$:

$$(4.39) \|C \star (S^3 - C^3) \star S\|_{L^1 \cap L^\infty} \leq c \|E\|_{L^1 \cap L^\infty} \Big[\Big(\frac{1}{m_t^4} + \frac{1}{m_t^6} \Big) \mathbf{1}_{d=2} + \Big(\frac{1}{m_t^2} + \frac{1}{m_t^5} \Big) \mathbf{1}_{d=3} \Big] \\ + c \|E\|_{L^1 \cap L^\infty}^2 \Big(\frac{1}{m_t^6} + \frac{1}{m_t^4} \mathbf{1}_{d=2} + \frac{1}{m_t^3} \mathbf{1}_{d=3} \Big) \\ + \frac{c \|E\|_{L^1 \cap L^\infty}^3}{m_t^4} + \frac{c \|E\|_{L^1 \cap L^\infty}^4}{m_t^2}.$$

Another λ^2 term is $C \star (\mathbf{1}_0^{\varepsilon} || C^3 ||_{L^1} - \mathbf{1}_0^{\varepsilon} || C_\infty^3 ||_{L^1}) \star S$, which simply reads:

(4.40)

$$C \star (\mathbf{1}_{0}^{\varepsilon} \| C^{3} \|_{L^{1}} - \mathbf{1}_{0}^{\varepsilon} \| C_{\infty}^{3} \|_{L^{1}}) \star S = (\| C^{3} \|_{L^{1}} - \| C_{\infty}^{3} \|_{L^{1}}) C \star S$$

$$=: -\gamma_{t} C \star S,$$

with

(4.41)
$$0 \leqslant \gamma_t \leqslant \begin{cases} \frac{1}{tm^2 m_t^2} & \text{if } d = 2, \\ c \log\left(1 + \frac{1}{m^2 t}\right) & \text{if } d = 3, \end{cases}$$

and the bound on γ_t obtained in Lemma A.3. Thus, for the $||E||_{L^{\infty}}$ norm:

(4.42)
$$\begin{aligned} \left\| C \star (\mathbf{1}_{0}^{\varepsilon} \| C^{3} \|_{L^{1}} - \mathbf{1}_{0}^{\varepsilon} \| C_{\infty}^{3} \|_{L^{1}}) \star S \right\|_{L^{\infty}} &\leq \gamma_{t} \left(\| C \|_{L^{1}} \| E \|_{L^{\infty}} + \| C \|_{L^{2}}^{2} \right) \\ &= \frac{\gamma_{t} \| E \|_{L^{\infty}}}{m_{t}^{2}} + \gamma_{t} \| C \|_{L^{2}}^{2} \\ &\leq \frac{\gamma_{t} \| E \|_{L^{\infty}}}{m_{t}^{2}} + \gamma_{t} \cdot \left\{ \frac{c}{m_{t}^{2}} & \text{if } d = 2, \\ \frac{c}{m_{t}} & \text{if } d = 3. \end{cases} \end{aligned}$$

Similarly, the $\|\cdot\|_{L^1}$ norm reads:

(4.43)
$$\left\| C \star (\mathbf{1}_0^{\varepsilon} \| C^3 \|_{L^1} - \mathbf{1}_0^{\varepsilon} \| C_\infty^3 \|_{L^1}) \star S \right\|_{L^1} \leqslant \frac{\gamma_t}{m_t^2} \Big(\| E \|_{L^1} + \frac{1}{m_t^2} \Big),$$

so that:

$$(4.44) \quad \left\| C \star (\mathbf{1}_{0}^{\varepsilon} \| C^{3} \|_{L^{1}} - \mathbf{1}_{0}^{\varepsilon} \| C_{\infty}^{3} \|_{L^{1}}) \star S \right\|_{L^{1} \cap L^{\infty}} \leq \frac{\gamma_{t} \| E \|_{L^{1} \cap L^{\infty}}}{m_{t}^{2}} + \gamma_{t} \Big(\frac{1}{m_{t}^{4}} + \frac{c}{m_{t}^{2}} \mathbf{1}_{d=2} + \frac{c}{m_{t}} \mathbf{1}_{d=3} \Big).$$

The last λ^2 term is $C \star \psi \star S$, with:

(4.45)
$$\psi := (C^3 - \mathbf{1}_0^{\varepsilon} \| C^3 \|_{L^1}).$$

One then has:

(4.46)
$$\left\| C \star \psi \star S \right\|_{L^1 \cap L^\infty} \leq \left\| C \star \psi \right\|_{L^1} \|E\|_{L^1 \cap L^\infty} + \left\| C \star \psi \star C \right\|_{L^1 \cap L^\infty}.$$

In Lemma A.1, the norms of the above quantities are estimated. Indeed, in dimension d = 2, one has $\|\psi\|_{L^1} \leq c/m_t^2$, giving

$$(4.47) \|C \star \psi\|_{L^1} \leqslant \frac{c}{m_t^4}, \|C \star \psi \star C\|_{L^1 \cap L^\infty} \leqslant \|\psi\|_{L^1} \|C \star C\|_{L^1 \cap L^\infty} \leqslant \frac{c}{m_t^4} + \frac{c}{m_t^6}.$$

In dimension d = 3, Lemma A.1 shows that

(4.48)
$$\|C \star \psi\|_{L^{1}} \leqslant \frac{c}{m_{t}^{1/2}} + \frac{c}{m_{t}^{5/2}}, \\\|C \star \psi \star C\|_{L^{1}} \leqslant \|C\|_{L^{1}} \|C \star \psi\|_{L^{1}} \leqslant \frac{c}{m_{t}^{5/2}} + \frac{c}{m_{t}^{9/2}}, \\\|C \star \psi \star C\|_{L^{\infty}} \leqslant \|C\|_{L^{2}} \|C \star \psi\|_{L^{2}} \leqslant \frac{c}{m_{t}}.$$

As a result, we find:

(4.49)
$$\begin{aligned} \left\| C \star \psi \star S \right\|_{L^{1} \cap L^{\infty}} &\leq c \|E\|_{L^{1} \cap L^{\infty}} \left[\frac{1}{m_{t}^{4}} \mathbf{1}_{d=2} + \left(\frac{1}{m_{t}^{1/2}} + \frac{1}{m_{t}^{5/2}} \right) \mathbf{1}_{d=3} \right] \\ &+ c \left(\frac{1}{m_{t}^{4}} + \frac{1}{m_{t}^{6}} \right) \mathbf{1}_{d=2} + c \left(\frac{1}{m_{t}} + \frac{1}{m_{t}^{9/2}} \right) \mathbf{1}_{d=3}. \end{aligned}$$

For future reference, note also the following better bound on $\|C \star \psi \star S\|_{L^1}$:

(4.50)
$$\|C \star \psi \star S\|_{L^1} \leqslant c \Big(\frac{1}{m_t^4} \mathbf{1}_{d=2} + \Big(\frac{1}{m_t^{1/2}} + \frac{1}{m_t^{5/2}}\Big) \mathbf{1}_{d=3}\Big) \Big(\|E\|_{L^1} + \frac{1}{m_t^2}\Big).$$

4.3.4. Order 3 terms. The only remaining term is the λ^3 term:

(4.51)
$$T_{\lambda^3} := 54\lambda^3 C \star Q \star S, \quad Q := S(S^2 \star S^2).$$

Notice first:

$$(4.52) \|T_{\lambda^3}\|_{L^1 \cap L^{\infty}} \leq 54\lambda^3 \Big(\|C \star Q \star C\|_{L^{\infty}} + \|C\|_{L^1} \|Q\|_{L^1} \|E\|_{L^{\infty}} + \|Q\|_{L^1} \|C \star S\|_{L^1} \Big)$$

$$\leq 54\lambda^3 \|Q\|_{L^1} \Big(\|C\|_{L^2}^2 + \frac{\|E\|_{L^1 \cap L^{\infty}}}{m_t^2} + \frac{1}{m_t^4} \Big).$$

It is therefore enough to estimate $\|Q\|_{L^1}$. Writing S = C + E, it reads:

(4.53)
$$\|Q\|_{L^1} \leq \left\| (C+E) \left[\left(C^2 + 2EC + E^2 \right) \star \left(C^2 + 2EC + E^2 \right) \right] \right\|_{L^1}.$$

Bounding E^n by $|E| ||E||_{L^{\infty}}^{n-1}$ for each $n \ge 1$, one finds:

$$\begin{aligned} \|Q\|_{L^{1}} &\leqslant \|C(C^{2} \star C^{2})\|_{L^{1}} + \|E\|_{L^{\infty}} \left(\|C^{2}\|_{L^{1}}^{2} + 4\|C(C \star C^{2})\|_{L^{1}}\right) \\ &+ \|E\|_{L^{\infty}}^{2} \left(6\|C^{2}\|_{L^{1}}\|C\|_{L^{1}} + 4\|C(C \star C)\|_{L^{1}}\right) \\ &+ \|E\|_{L^{\infty}}^{2} \left(2\|E\|_{L^{1}}\|C^{2}\|_{L^{1}} + 8\|E\|_{L^{\infty}}\|C\|_{L^{1}}^{2}\right) \\ &+ 5\|E\|_{L^{\infty}}^{3}\|E\|_{L^{1}}\|C\|_{L^{1}} \\ &+ \|E\|_{L^{\infty}}^{3}\|E\|_{L^{1}}^{2}. \end{aligned}$$

$$(4.54)$$

The first term on the right-hand side of (4.54) is estimated in Lemma A.2 as:

(4.55)
$$\|C(C^2 \star C^2)\|_{L^1} \leqslant \frac{c}{m_t^4} \mathbf{1}_{d=2} + \frac{c}{m_t} \mathbf{1}_{d=3}.$$

The inequality $C(x)C(y) \leq \frac{1}{2}C(x)^2 + \frac{1}{2}C(y)^2$ is enough to also obtain:

(4.56)
$$\|C(C \star C^2)\|_{L^1} \leqslant \|C^2\|_{L^1}^2 \leqslant \frac{c}{m_t^4} \mathbf{1}_{d=2} + \frac{c}{m_t^2} \mathbf{1}_{d=3},$$
$$\|C(C \star C)\|_{L^1} \leqslant \|C^2\|_{L^1} \|C\|_{L^1} \leqslant \frac{c}{m_t^4} \mathbf{1}_{d=2} + \frac{c}{m_t^3} \mathbf{1}_{d=3}.$$

Plugging (4.55)–(4.56) into the bound (4.54) on $||Q||_{L^1}$ yields:

$$||Q||_{L^{1}} \leqslant \frac{c}{m_{t}^{4}} \mathbf{1}_{d=2} + \frac{c}{m_{t}} \mathbf{1}_{d=3} + ||E||_{L^{1} \cap L^{\infty}} \left(\frac{c}{m_{t}^{4}} \mathbf{1}_{d=2} + \frac{c}{m_{t}^{2}} \mathbf{1}_{d=3}\right) + ||E||_{L^{1} \cap L^{\infty}}^{2} \left(\frac{c}{m_{t}^{4}} \mathbf{1}_{d=2} + \frac{c}{m_{t}^{3}} \mathbf{1}_{d=3}\right) + ||E||_{L^{1} \cap L^{\infty}}^{3} \left(\frac{c}{m_{t}^{2}} \mathbf{1}_{d=2} + \frac{c}{m_{t}} \mathbf{1}_{d=3} + \frac{8}{m_{t}^{4}}\right) + \frac{5||E||_{L^{1} \cap L^{\infty}}^{4}}{m_{t}^{2}} + ||E||_{L^{1} \cap L^{\infty}}^{5}.$$

$$(4.57)$$

Injecting the bound (4.57) into the bound (4.52) for T_{λ^3} , one finds that there is a polynomial P_3 of degree 6, with no constant term, such that:

(4.58)
$$\|T_{\lambda^3}\|_{L^1 \cap L^\infty} \leqslant c\lambda^3 \Big[\Big(\frac{1}{m_t^6} + \frac{1}{m_t^8}\Big) \mathbf{1}_{d=2} + \Big(\frac{1}{m_t^2} + \frac{1}{m_t^5}\Big) \mathbf{1}_{d=3} + P_3 \Big(\|E\|_{L^1 \cap L^\infty}\Big) \Big].$$

Moreover, P_3 has coefficients given by sums of negative powers of m_t , the lowest power in absolute value being 1 (if d = 3) or 2 (if d = 2).

4.3.5. Conclusion on $||E||_{L^1 \cap L^\infty}$. Putting together the estimates of T_{λ^i} $(0 \le i \le 3)$ obtained in the previous sections, one finds for $||E||_{L^1 \cap L^\infty}$:

(4.59)
$$||E||_{L^{1}\cap L^{\infty}} \leq c \left(\lambda \eta_{t} + \lambda^{2} \gamma_{t} + |\mu - m^{2}|\right) \left(\frac{1}{m_{t}^{2}} \mathbf{1}_{d=2} + \frac{1}{m_{t}} \mathbf{1}_{d=3} + \frac{1}{m_{t}^{4}}\right) + \frac{c ||E||_{L^{1}\cap L^{\infty}}}{m_{t}^{2}} \left(\lambda \eta_{t} + \lambda^{2} \gamma_{t} + |\mu - m^{2}|\right) + c \sum_{i=1}^{3} \lambda^{i} R_{t}^{(i)} \left(||E||_{L^{1}\cap L^{\infty}}\right),$$

where, for $i \in \{1, 2, 3\}$, $R_t^{(i)}$ is a polynomial with positive coefficients given by sums of negative powers of m_t and with no constant term. The coefficient of the $R_t^{(i)}$ with the lowest power of m_t in absolute value is proportional to $m_t^{-1/2}$ in dimension d = 3, and to m_t^{-2} in dimension d = 2.

The bound (4.59) yields the following estimates of $||E||_{L^1 \cap L^\infty}$.

Lemma 4.5. Let $d \in \{2, 3\}$, $\lambda > 0$, $\mu \in \mathbb{R}$, and $m^2 > 0$.

(i) (Small scale). There is $t_0(d, \lambda, \mu, m^2) > 0$ (independent of ε, L) and a numerical constant c > 0 such that:

(4.60)
$$\forall t \in (0, t_0(d, \lambda, \mu, m^2)], \qquad \|E\|_{L^1 \cap L^\infty} \leqslant c\lambda.$$

(ii) (All scales, one small coupling). If $\mu > 0$ and with the special choice $m^2 = \mu$, there are parameters $\lambda(d,\mu), \mu(d,\lambda) > 0$ (independent of ε, L) with the following property. If either $\lambda \in [0, \lambda(d,\mu)]$ or $\mu \ge \mu(d,\lambda)$, then there is $c(\mu) > 0$ with:

(4.61)
$$\forall t > 0, \qquad \|E\|_{L^1 \cap L^\infty} \leqslant c(\mu)\lambda, \qquad \lim_{\mu \to \infty} c(\mu) = 0.$$

In the proof of the lemma, we will use the following qualitative properties of $t \mapsto E = E_t$, which hold for any fixed $\varepsilon > 0$ and $L < \infty$:

(4.62)
$$t \in (0, \infty) \mapsto ||E||_{L^1 \cap L^\infty} \text{ is continuous;}$$
$$\lim_{t \to 0} ||E||_{L^1 \cap L^\infty} = 0.$$

Indeed, since $\varepsilon > 0$ and $L < \infty$ are fixed, the above continuity follows from the continuity of $t \mapsto S(x)$ and $t \mapsto C(x)$ for each $x \in \Lambda_{\varepsilon,L}$. Similarly, the $t \to 0$ limit follows from $\lim_{t\to 0} C(x) = \lim_{t\to 0} S(x) = 0$ for each $x \in \Lambda_{\varepsilon,L}$. This is clear for $C(x) = (-\Delta^{\varepsilon} + m^2 + 1/t)^{-1}(0,x)$, and for S(x) it follows, for example, from the Brascamp–Lieb inequality which shows $S(x) \leq S(0) \leq (-\Delta^{\varepsilon} + a^{\varepsilon}(\lambda, m^2) + \mu + 1/t)^{-1}(0,0)$ provided $a^{\varepsilon}(\lambda, m^2) + \mu + 1/t > 0$ (similarly to (3.5)).

Proof. Let $d \in \{2,3\}$, and let $f_{\lambda,\mu,m^2,t}$ be the polynomial such that (4.59) corresponds to:

(4.63)
$$\|E\|_{L^1 \cap L^{\infty}} \leqslant \frac{c\lambda\eta_t}{m_t} \mathbf{1}_{d=3} + f_{\lambda,\mu,m^2,t} (\|E\|_{L^1 \cap L^{\infty}}).$$

The reason why the first term is taken out is due to (4.66) below.

Consider first item (i). The difference of the counterterms η_t , γ_t are estimated in Lemma A.3:

(4.64)
$$\eta_t \leqslant c \log\left(1 + \frac{1}{m^2 t}\right) \mathbf{1}_{d=2} + cm\left(\sqrt{1 + \frac{1}{tm^2}} - 1\right) \mathbf{1}_{d=3},$$
$$\gamma_t \leqslant \frac{c}{m^2(m^2 t + 1)} \mathbf{1}_{d=2} + c \log\left(1 + \frac{1}{m^2 t}\right) \mathbf{1}_{d=3}.$$

In particular, they satisfy, for a numerical constant $c_0 > 0$:

(4.65)
$$\sup_{t>0} \frac{c\eta_t}{m_t} \leq \mathbf{1}_{d=2}c(m^2) + c_0 \mathbf{1}_{d=3} \quad \text{with} \quad \lim_{m \to \infty} c(m^2) = 0,$$

and:

(4.66)
$$\limsup_{t \downarrow 0} \frac{c\eta_t}{m_t} \leqslant 0 \mathbf{1}_{d=2} + c_0 \mathbf{1}_{d=3}, \qquad \lim_{t \downarrow 0} \frac{\gamma_t}{m_t} = 0.$$

On the other hand, note the following elementary property:

(4.67)
$$m_t^2 := m^2 + \frac{1}{t} \quad \Rightarrow \quad \lim_{t \downarrow 0} m_t = +\infty.$$

It follows from the above that the (positive) coefficients of the polynomial $f_{\lambda,\mu,m^2,t}$ tend to 0 as $t \downarrow 0$. In particular, there is a largest value $t_0(d, \lambda, \mu, m^2) > 0$ such that:

(4.68)
$$\forall t \in (0, t_0(d, \lambda, \mu, m^2)], \qquad f_{\lambda, \mu, m^2, t}(2c_0\lambda) \leqslant \frac{c_0\lambda}{2},$$

where c_0 is the constant from (4.65) (and we assume $c(m^2) \leq c_0$ in d = 2). Note also that $x \mapsto f_{\lambda,\mu,m^2,t}(x)$ is increasing in $x \geq 0$ (the coefficients of $f_{\lambda,\mu,m^2,t}$ are positive). Since $t \in (0,\infty) \mapsto \|E_t\|_{L^1 \cap L^{\infty}}$ is continuous and $\lim_{t\to\infty} \|E_t\|_{L^1 \cap L^{\infty}} = 0$, both by (4.62), it follows that $\|E\|_{L^1 \cap L^{\infty}} \leq 2c_0\lambda$ on $(0, t_0(d, \lambda, \mu, m^2)]$. Indeed, suppose the contrary, and take $t_* \in (0, t_0(d, \lambda, \mu, m^2)]$ with $\|E_{t_*}\|_{L^1 \cap L^{\infty}} > 2c_0\lambda$. Then (4.68) and the definition (4.63) of $f_{\lambda,\mu,m^2,t}$ yield a contradiction:

(4.69)
$$2c_0\lambda < \|E_{t_*}\|_{L^1 \cap L^\infty} \le c_0\lambda + f_{\lambda,\mu,m^2,t_*}(2c_0\lambda) \le \frac{3c_0\lambda}{2} < 2c_0\lambda.$$

This concludes the proof of item (i).

For item (ii), the reasoning is similar, so we only give a sketch. As $\mu = m^2$, the right-hand side of (4.59) only contains linear combinations of the following terms:

(4.70)
$$c(\lambda\eta_t + \lambda^2\gamma_t) \Big(\frac{1}{m_t^2} \mathbf{1}_{d=2} + \frac{1}{m_t} \mathbf{1}_{d=3} \Big), \qquad c(\lambda\eta_t + \lambda^2\gamma_t) m_t^{-a}, \qquad a \ge 2,$$
$$and \quad c\lambda^n m_t^{-a}, \qquad a > 0, n \in \{1, 2, 3\}.$$

As for item (i), none of these terms depend on ε or L, and $c\lambda^n m_t^{-a}$ is bounded by $c\lambda^n m^{-a} = c\lambda^n \mu^{-a/2}$ for each a, n. To see what happens for large μ or small λ , recall from (4.65) the bound:

(4.71)
$$\sup_{t>0} \frac{c\eta_t}{m_t} \leqslant c(\mu) \mathbf{1}_{d=2} + c_0 \quad \text{with} \quad \lim_{\mu \to \infty} c(\mu) = 0,$$

and from (4.64) the corresponding bound for γ_t :

(4.72)
$$\sup_{t>0} \frac{c\gamma_t}{m_t} \leqslant c'(\mu) \quad \text{with} \quad \lim_{\mu \to \infty} c'(\mu) = 0.$$

It follows that the right-hand side of (4.59) is bounded uniformly in t by $c(\mu)\lambda$ (for a different $c(\mu)$ that nonetheless vanishes when μ is large), provided either λ is smaller than some $\lambda(d, \mu) > 0$ or μ larger than some $\mu(d, \lambda) > 0$. Thus one has an analogue of (4.68) with $t_0 = +\infty$ and repeating the argument of item (i) concludes the proof of item (ii) and of the lemma.

4.4. Estimate of the susceptibility. Fix a dimension $d \in \{2,3\}$, let $\lambda > 0$, $\mu \in \mathbb{R}$, $m^2 > 0$, and t > 0. We now obtain the bound of Proposition 4.1 on the norm $||E||_{L^1}$. The starting point is again the formula (4.15) with the splitting (4.17):

(4.73)
$$|S(x) - C(x)| \leq 3\lambda (\eta_t + ||E||_{L^{\infty}}) (C \star S)(x) + 6\lambda^2 |(C \star (S^3 - \mathbf{1}_0^{\varepsilon} ||C_{\infty}^3||_{L^1}) \star S)(x)| + 54\lambda^3 (C \star Q \star S)(x) + |\mu - m^2| (C \star S)(x).$$

The quantity $||E||_{L^{\infty}}$ is estimated by Lemma 4.5. The $||\cdot||_{L^1}$ norm of E then reads:

(4.74)
$$\|E\|_{L^{1}} \leqslant \frac{3\lambda \left(\eta_{t} + \|E\|_{L^{\infty}} + |\mu - m^{2}|\right)}{m_{t}^{2}} \left(\|E\|_{L^{1}} + \frac{1}{m_{t}^{2}}\right) + \|T_{\lambda^{2}}\|_{L^{1}} + \|T_{\lambda^{3}}\|_{L^{1}}.$$

Lemma 4.6. There are polynomials $P_t^{(i)} = P_{d,t,\mu,m^2,||E||_{L^{\infty}}}^{(i)}$ with $i \in \{1, 2, 3\}$, of degree at most 3 and with no constant term, such that:

$$||E||_{L^{1}} \leqslant \frac{3\lambda \left(\eta_{t} + ||E||_{L^{\infty}} + |\mu - m^{2}|\right)}{m_{t}^{4}} + \frac{c\lambda^{2}}{m_{t}^{4}} \left[\gamma_{t} + \frac{||E||_{L^{\infty}} + ||E||_{L^{\infty}}^{2}}{m_{t}^{2}} + \frac{1}{m_{t}^{2}} \mathbf{1}_{d=2} + \left(\frac{1}{m_{t}} + \frac{1}{m_{t}^{1/2}}\right) \mathbf{1}_{d=3}\right] + \lambda^{3} \left[\frac{c}{m_{t}^{8}} + \left(\frac{c}{m_{t}^{7}} + \frac{c}{m_{t}^{5}}\right) \mathbf{1}_{d=3}\right] + \sum_{i=1}^{3} \lambda^{i} P_{t}^{(i)} \left(||E||_{L^{1}}\right).$$

$$(4.75)$$

The coefficients of the polynomials $P_t^{(i)}$ are nonnegative and of the following form:

(4.76)
$$c \frac{\eta_t + |\mu - m^2|}{m_t^4} \quad or \quad c \frac{\gamma_t}{m_t^2} \quad or \quad c \|E\|_{L^{\infty}}^n m_t^{-a}, \qquad c, a > 0; \ n \in \{0, 1, 2, 3\}.$$

The lowest value of a is 2 in dimension 2, and 1/2 in dimension 3.

Proof. The norms $||T_{\lambda^2}||_{L^1}$ and $||T_{\lambda^3}||_{L^1}$ have been estimated in Sections 4.3.3–4.3.4 respectively. Precisely, $||T_{\lambda^2}||_{L^1}$ is estimated in (4.38)–(4.43)–(4.50) and, bounding E^n by $|E|||E||_{L^{\infty}}^{n-1}$, it reads:

$$\lambda^{-2} \| T_{\lambda^2} \|_{L^1} \leqslant \frac{c\gamma_t}{m_t^4} + \frac{c(\|E\|_{L^{\infty}} + 1)}{m_t^6} \mathbf{1}_{d=2} + c \Big(\frac{\|E\|_{L^{\infty}}}{m_t^5} + \frac{1}{m_t^{5/2}} + \frac{1}{m_t^{9/2}} \Big) \mathbf{1}_{d=3} + \frac{\|E\|_{L^{\infty}}^2}{m_t^6} \\ + c \|E\|_{L^1} \Big[\frac{\gamma_t}{m_t^2} + \frac{\|E\|_{L^{\infty}}^2}{m_t^4} + \frac{(\|E\|_{L^{\infty}} + 1)}{m_t^4} \mathbf{1}_{d=2} + \Big(\frac{1}{m_t^{1/2}} + \frac{1}{m_t^{5/2}} + \frac{\|E\|_{L^{\infty}}}{m_t^3} \Big) \mathbf{1}_{d=3} \Big]$$

$$(4.77) \qquad + \frac{c \|E\|_{L^1}^2}{m_t^2}.$$

On the other hand, for $||T_{\lambda^3}||_{L^1} = 54\lambda^3 ||C \star S||_{L^1} ||Q||_1$, one has:

(4.78)
$$\lambda^{-3} \| T_{\lambda^3} \|_{L^1} \leqslant \frac{54}{m_t^2} \Big(\| E \|_{L^1} + \frac{1}{m_t^2} \Big) \| Q \|_{L^1}.$$

Recalling the bound (4.54) on $||Q||_{L^1}$ concludes the proof of Lemma 4.6:

$$\|Q\|_{L^{1}} \leq \frac{\|E\|_{L^{\infty}}^{3}}{m_{t}^{4}} + \frac{c\left(1 + \|E\|_{L^{\infty}} + \|E\|_{L^{\infty}}^{2}\right)}{m_{t}^{4}} \mathbf{1}_{d=2} + c\left(\frac{1}{m_{t}} + \frac{\|E\|_{L^{\infty}}}{m_{t}^{2}} + \frac{\|E\|_{L^{\infty}}^{2}}{m_{t}^{3}}\right) \mathbf{1}_{d=3}$$

$$(4.79) \qquad + c\|E\|_{L^{1}} \left[\frac{\|E\|_{L^{\infty}}^{3}}{m_{t}^{2}} + \|E\|_{L^{\infty}}^{2} \left(\frac{1}{m_{t}^{2}} \mathbf{1}_{d=2} + \frac{c}{m_{t}} \mathbf{1}_{d=3}\right)\right] + \|E\|_{L^{\infty}}^{3} \|E\|_{L^{1}}^{2}.$$

Lemma 4.6 is sufficient to conclude the proof of Propositions 4.1–4.2.

Proof of Proposition 4.1. Let $t_0(d, \lambda, \mu, m^2) > 0$ be the scale defined in Lemma 4.5, such that, for some numerical constant c:

(4.80)
$$\forall t \in (0, t_0(d, \lambda, \mu, m^2)], \qquad \|E\|_{L^1 \cap L^\infty} \leqslant c\lambda.$$

Henceforth $t \in (0, t_0(d, \lambda, \mu, m^2)]$. Let $G_{d,t,\mu,m^2}(\lambda) + \sum_{i=1}^3 \lambda^i P_{d,t,\mu,m^2,c\lambda}^{(i)}(\|E\|_{L^1})$ denote the right-hand side of the bound in Lemma 4.6 with $\|E\|_{L^{\infty}}$ replaced by $c\lambda$:

(4.81)
$$\|E\|_{L^1} \leqslant G_{d,t,\mu,m^2}(\lambda) + \sum_{i=1}^3 \lambda^i P_{d,t,\mu,m^2,c\lambda}^{(i)} \big(\|E\|_{L^1} \big).$$

Using the already established bound $||E||_{L^1} \leq c\lambda$ on the right-hand side would give a bound on $||E||_{L^1}$ in which each term has a factor $m_t^{-a} \sim t^{a/2}$ as $t \to 0$. However, the smallest powers obtained in this way (for example, a = 1/2 in dimension 3) are not integrable with respect to $t^{-2} dt$ near t = 0. On the other hand, the term in $G_{d,t,\mu,m^2}(\lambda)$ with the most singular t-dependence as $t \to 0$ is $\eta_t m_t^{-4}$ which is integrable with respect to $t^{-2} dt$. (Indeed, recall $m_t^{-4} \sim t^2$, that (4.64) shows that η_t diverges as $\log t^{-1}$ in d = 2 and as $t^{-1/2}$ in d = 3, and that γ_t is bounded in d = 2whereas γ_t diverges as $\log t^{-1}$ in d = 3). In addition:

(4.82)
$$\sup_{t>0} G_{d,t,\mu,m^2}(\lambda) \leqslant c(d,\lambda,\mu,m^2).$$

Therefore, since the $P^{(i)}$ do not have constant terms, by iterating the bound (4.81) a finite number of times, and using the bound $||E||_{L^1} \leq c\lambda$ from (4.80) at the last iteration, it follows that

(4.83)
$$||E||_{L^1} \leqslant p_{d,t,\mu,m^2}(\lambda),$$

with the right-hand side above a polynomial in λ of degree independent of the parameters $\lambda, \mu, m^2, \varepsilon, L$, and coefficients functions of $\eta_t, \gamma_t, |\mu - m^2|$ times powers of m_t^{-1} such that

(4.84)
$$t \mapsto \frac{p_{d,t,\mu,m^2}(\lambda)}{t^2}$$
 is integrable around $t = 0$.

This concludes the proof of Proposition 4.1.

Proof of Proposition 4.2. Fix $\mu = m^2$, and consider either $\lambda \in [0, \lambda(d, \mu)]$, or $\mu \ge \mu(d, \lambda)$, where these quantities are defined in item (ii) of Lemma 4.5. Then:

(4.85)
$$\forall t > 0, \qquad \|E\|_{L^1 \cap L^\infty} \leqslant c(\mu)\lambda, \qquad \lim_{\mu \to \infty} c(\mu) = 0.$$

The argument is then identical to the above proof of Proposition 4.1, replacing $||E||_{L^{\infty}}$ by $c(\mu)\lambda$ for each t > 0 and writing the right-hand side of the bound in Lemma 4.6 as:

(4.86)
$$\|E\|_{L^1} \leq \tilde{G}_{d,t,\mu}(\lambda) + \sum_{i=1}^3 \lambda^i P_{d,t,\mu,\mu,c(\mu)\lambda}^{(i)} \big(\|E\|_{L^1} \big)$$

Then, again by iteration, for each t > 0:

$$(4.87) ||E||_{L^1} \leqslant \tilde{p}_{d,t,\mu}(\lambda)$$

with $t \mapsto t^{-2} \tilde{p}_{d,t,\mu}(\lambda)$ integrable around 0. Using the bound of Lemma 4.6 together with the estimates on the differences of counterterms from Lemma A.3, it also follows that $\tilde{p}_{d,t,\mu}(\lambda)$ has coefficients bounded uniformly on t. It is therefore integrable at ∞ with respect to $t^{-2} dt$, and thus on the whole of $(0,\infty)$. Moreover, since each term appearing in $\tilde{p}_{d,t,\mu}(\lambda)$ vanishes when $\lambda \to 0$ or $\mu \to \infty$, one has, up to decreasing $\lambda(d,\mu)$ or increasing $\mu(d,\lambda)$:

(4.88)
$$\sup_{\lambda>0} \sup_{\mu \geqslant \mu(d,\lambda)} \int_0^\infty \frac{\tilde{p}_{d,t,\mu}(\lambda)}{t^2} dt < \infty, \qquad \sup_{\mu>0} \sup_{\lambda \in [0,\lambda(d,\mu)]} \int_0^\infty \frac{\tilde{p}_{d,t,\mu}(\lambda)}{t^2} dt < \infty.$$

The final claim $\limsup_{t\to\infty} \tilde{p}_{d,t,\mu}(\lambda) = o(\mu^{-1})$ is also clear from the above (recall that the differences η_t, γ_t of the counterterms at time t and ∞ vanish by definition in the limit $t \to \infty$, so only the inverse powers of m_t in $\tilde{p}_{d,t,\mu}$ matter in the limit).

5 Proof of Theorem 1.1

Proof of Theorem 1.1 (i). Fix $m^2 = 1$ (though the value of m^2 is irrelevant). Let $\bar{\chi} \in (0, +\infty)$ and assume that ε and L are such that

(5.1)
$$\chi_{\infty}^{\varepsilon,L} := \varepsilon^d \sum_{x \in \Lambda_{\varepsilon,L}} \left\langle \varphi_0 \varphi_x \right\rangle_{\lambda,\mu}^{\varepsilon,L} \leqslant \bar{\chi}$$

By Theorem 2.6, the log-Sobolev constant $\gamma = \gamma^{\varepsilon,L}(\lambda,\mu)$ of the continuum φ_d^4 measure (1.1) is bounded by

(5.2)
$$\frac{1}{\gamma} \leqslant \int_0^\infty \exp\left[-2\int_0^t \dot{\kappa}_s^{\varepsilon,L} \, ds\right] dt, \qquad \dot{\kappa}_t^{\varepsilon,L} := \frac{1}{t} - \frac{\chi_t^{\varepsilon,L}}{t^2}.$$

To bound γ , we estimate the susceptibility for small scale t and large scale t in a different way.

For t smaller than the $t_0 := t_0(d, \lambda, \mu, m^2 = 1) > 0$ of Proposition 4.1, one has:

(5.3)
$$\chi_t^{\varepsilon,L} \leqslant \frac{1}{1+\frac{1}{t}} + p_{d,t,\mu,m^2=1}(\lambda), \qquad \int_0^{t_0} \frac{p_{d,t,\mu,m^2=1}(\lambda)}{t^2} dt \leqslant c_d(\lambda,\mu,m^2=1),$$

where the $(1+1/t)^{-1}$ term is simply the Gaussian susceptibility $||C_t||_{L^1}$ with mass $m^2 = 1$. This gives:

(5.4)
$$\forall t \in (0, t_0], \quad \dot{\kappa}_t^{\varepsilon, L} \ge \frac{1}{t} - \frac{1}{t^2} \left(\frac{1}{1 + 1/t} + p_{d, t, \mu, m^2 = 1}(\lambda) \right) = \frac{1}{t+1} - \frac{p_{d, t, \mu, m^2 = 1}(\lambda)}{t^2}.$$

and

(5.5)
$$\int_0^t \dot{\kappa}_s^{\varepsilon,L} \, ds \ge \log(t+1) - c_d(\lambda,\mu,m^2=1).$$

On the other hand, the second Griffiths inequality implies that $\chi_t^{\varepsilon,L}$ is decreasing in 1/t, i.e., increasing in t. For large scales $t > t_0$, one thus has $\chi_t^{\varepsilon,L} \leq \chi_{\infty}^{\varepsilon,L} \leq \bar{\chi}$ by (5.1), so that:

(5.6)
$$\forall t > t_0, \qquad \dot{\kappa}_t^{\varepsilon,L} \ge \frac{1}{t} - \frac{\bar{\chi}}{t^2}.$$

This bound implies:

(5.7)
$$\forall t > t_0, \qquad \int_{t_0}^t \dot{\kappa}_s^{\varepsilon,L} \, ds \ge \log t - \log t_0 - \bar{\chi}/t_0 = \log(t/t_0) - C(\lambda, \mu, \bar{\chi}).$$

Combined, (5.5) and (5.7) give for each t > 0:

(5.8)
$$\exp\left[-2\int_0^t \kappa_s^{\varepsilon,L} ds\right] \leqslant \exp\left[-2\int_0^{t_0\wedge t} \kappa_s^{\varepsilon,L} ds\right] \exp\left[-2\int_{t_0\wedge t}^t \kappa_s^{\varepsilon,L} ds\right] \\ \leqslant (t\wedge t_0+1)^{-2} (1\vee t/t_0)^{-2} e^{2c_d(\lambda,\mu,m^2=1)+2C(\lambda,\mu,\bar{\chi})}$$

which is integrable on $(0, \infty)$ with a bound that depends only on $(\lambda, \mu, \overline{\chi})$.

Proof of Theorem 1.1 (ii). Under the assumptions of (ii) $(m^2 > 0, \mu = m^2$ and either $\lambda \in [0, \lambda(d, \mu)]$ or $\mu \ge \mu(d, \lambda)$), we can directly apply Proposition 4.2 for all t > 0. As in the proof of (i), this gives

(5.9)
$$\dot{\kappa}_t^{\varepsilon,L} \ge \frac{m^2}{m^2 t + 1} - \frac{\tilde{p}_{d,t,m^2}(\lambda)}{t^2}$$

and

(5.10)
$$\kappa_t^{\varepsilon,L} \ge \log(m^2 t + 1) - c, \qquad c := \int_0^\infty \frac{\tilde{p}_{d,t,m^2}(\lambda)}{t^2}.$$

Recall from Proposition 4.2 that c is independent of λ, μ under the assumptions of (ii). The log-Sobolev constant $\gamma = \gamma^{\varepsilon,L}(\lambda,\mu)$ thus satisfies:

(5.11)
$$\frac{1}{\gamma} \leqslant \int_0^\infty \exp\left[-2\int_0^t \kappa_s^{\varepsilon,L} \, ds\right] \, dt \leqslant e^c \int_0^\infty \frac{1}{(m^2t+1)^2} \, dt = \frac{e^c}{m^2}.$$

Thus $\gamma \ge e^{-c}m^2 = e^{-c}\mu$. Since $\chi^{\varepsilon,L}(\lambda,\mu) = \|S_{\infty}\|_{L^1}$ and $\|S_{\infty}\|_{L^1} = \mu^{-1} + o(\mu^{-1})$ by the last item of Proposition 4.2, the bound on γ in the other direction was established just below the statement of Theorem 1.1. This concludes the proof of item (ii) of Theorem 1.1.

A Bounds on diagrams

A.1. Moments of Gaussian covariance. In this section, $m^2 > 0$ is fixed and we consider:

(A.1)
$$C_{(m)} := \left(-\Delta^{\varepsilon} + m^2\right)^{-1}$$

with the matrix elements of the inverse normalised with respect to the inner product $(u, v)_{\varepsilon} = \varepsilon^d \sum_{x \in \Lambda_{\varepsilon,L}} u(x)v(x)$. In other words, $C_{(m)}$ is given by:

(A.2)
$$\forall x \in \Lambda_{\varepsilon,L}, \qquad C_{(m)}(x) := C_{(m)}(0,x) = \frac{1}{L^d} \sum_{k \in \Lambda^*} e^{ik \cdot x} \frac{1}{m^2 + \theta(k)}.$$

Above, Λ^* is the set:

(A.3)
$$\Lambda^* := \left\{ k \in \frac{2\pi}{L} \mathbb{Z}^d : \forall i \in \{1, \dots, d\}, \ -\frac{\pi}{\varepsilon} < k_i \leqslant \frac{\pi}{\varepsilon} \right\},$$

and Δ^{ε} is the Laplacian on the torus $\Lambda_{\varepsilon,L}$ defined in (1.2), with eigenvalues θ given by:

(A.4)
$$\theta(k) := \sum_{k=1}^{3} \theta(k_i), \qquad \theta(k_i) := \frac{4}{\varepsilon^2} \sin^2\left(\frac{k_i\varepsilon}{2}\right), \qquad k \in \Lambda^*.$$

For later use, recall that convexity of the sine function on $[0, \pi/2]$ yields, for each $0 \le u \le \pi/2$:

(A.5)
$$\frac{2u}{\pi} \leqslant \sin(u) \leqslant u \quad \Rightarrow \quad \frac{4u^2}{\pi^2} \leqslant \sin^2(\pm u) \leqslant u^2.$$

In particular:

(A.6)
$$\forall k \in \Lambda^*, \qquad \frac{1}{m^2 + \theta(k)} \leqslant \frac{1}{m^2 + \frac{4\|k\|_2^2}{\pi^2}}.$$

Lemma A.1. Let ψ be given by:

(A.7)
$$\psi = C_{(m)}^3 - \mathbf{1}_0^{\varepsilon} \| C_{(m)}^3 \|_{L^1},$$

where $C_{(m)}$ is the Gaussian covariance (A.1), and $\mathbf{1}_0^{\varepsilon} := \varepsilon^{-d} \mathbf{1}_0$. In dimension d = 2, one has:

(A.8)
$$\|\psi\|_{L^1} \leqslant \frac{c}{m^2}$$

When instead d = 3, for each $k \in \Lambda^*$, the Fourier transform $\hat{\psi}(k)$ of ψ satisfies:

(A.9)
$$|\hat{\psi}(k)| \leqslant c \log\left(1 + \frac{2\|k\|_2}{m\pi}\right),$$

and

(A.10)
$$||C_{(m)} \star \psi||_{L^1} \leq \frac{c}{m^{1/2}} + \frac{c}{m^{5/2}}, \qquad ||C_{(m)} \star \psi||_{L^2} \leq \frac{c}{m^{1/2}}.$$

Proof. Let us start with the bound on $\|\psi\|_{L^1}$ in dimension d = 2, which is easiest. Indeed, one has in that case:

(A.11)
$$|\psi| \leq C_{(m)}^3 + \mathbf{1}_0^{\varepsilon} \|C_{(m)}^3\|_{L^1} \Rightarrow \|\psi\|_{L^1} \leq 2 \|C_{(m)}^3\|_{L^1}.$$

One has then:

(A.12)
$$\|C_{(m)}^3\|_{L^1} = \frac{1}{L^{2d}} \sum_{k,p \in \Lambda^*} \frac{1}{m^2 + \theta(p)} \cdot \frac{1}{m^2 + \theta(q)} \cdot \frac{1}{m^2 + \theta(p+q)}.$$

Using convexity to bound the sines in the definition (A.4) of θ and comparing sums to integrals yields:

(A.13)
$$\|C^3\|_{L^1} \leqslant \int_{[-\frac{\pi}{\varepsilon} - 1, \frac{\pi}{\varepsilon}]^4} dp \, dq \, \frac{1}{m^2 + \frac{4\|p\|_2^2}{\pi^2}} \cdot \frac{1}{m^2 + \frac{4\|q\|_2^2}{\pi^2}} \cdot \frac{1}{m^2 + \frac{4\|p+q\|_2^2}{\pi^2}} \\ \leqslant \frac{\pi^4}{16m^2} \int_{\mathbb{R}^4} dp \, dq \, \frac{1}{1 + \|p\|_2^2} \cdot \frac{1}{1 + \|q\|_2^2} \cdot \frac{1}{1 + \|p+q\|_2^2} \leqslant \frac{c}{m^2}$$

where we used that the integral on the last line is finite which follows, e.g., from the Cauchy-Schwarz inequality. This concludes the estimate of $\|\psi\|_{L^1}$ in the d = 2 case.

Consider now d = 3. The Fourier transform $\hat{\psi}(k)$ reads, for each $k \in \Lambda^*$ (defined in (A.3)):

$$\hat{\psi}(k) = \frac{1}{L^6} \sum_{p,q \in \Lambda^*} \frac{1}{m^2 + \theta(p)} \cdot \frac{1}{m^2 + \theta(q)} \cdot \left[\frac{1}{m^2 + \theta(k - p - q)} - \frac{1}{m^2 + \theta(-p - q)} \right]$$
$$= -\int_0^1 d\alpha \, \frac{1}{L^6} \sum_{p,q \in \Lambda^*} \frac{1}{m^2 + \theta(p)} \cdot \frac{1}{m^2 + \theta(q)}$$
$$\cdot \frac{4\pi\varepsilon^{-1} \sum_{i=1}^3 k_i \sin\left(\frac{\pi\varepsilon(\alpha k_i - p_i - q_i)}{2}\right) \cos\left(\frac{\pi\varepsilon(\alpha k_i - p_i - q_i)}{2}\right)}{\left[m^2 + \theta(\alpha k - p - q)\right]^2}.$$
(A.14)

In the following it is convenient to extend the above definition to $k \in \mathbb{R}^3$. Using convexity to bound the sines and comparing sums to integrals, then bounding these integrals by integrals on \mathbb{R}^d and changing variables $p, q \to 2p/m\pi, 2q/m\pi$ as before, we find that $\hat{\psi}(k)$ satisfies:

(A.15)
$$\forall k \in \mathbb{R}^3, \qquad |\hat{\psi}(k)| \leqslant c\hat{f}(2k/m\pi),$$

with:

(A.16)
$$\hat{f}(k) := \|k\|_2 \int_0^1 d\alpha \int_{\mathbb{R}^6} dp \, dq \, \frac{1}{1+\|p\|_2^2} \cdot \frac{1}{1+\|q\|_2^2} \cdot \frac{\|\alpha k - p - q\|_2}{\left[1+\|\alpha k - p - q\|_2^2\right]^2}.$$

Let us estimate \hat{f} . We claim that, for each $\tilde{k} \in \mathbb{R}^3$,

(A.17)
$$\int_{\mathbb{R}^3} \frac{1}{1+\|p\|^2} \frac{\|\tilde{k}-p\|_2}{\left[1+\|\tilde{k}-p\|_2^2\right]^2} \, dp = \int_{\mathbb{R}^3} \frac{1}{1+\|\tilde{k}+p\|^2} \frac{\|p\|_2}{\left[1+\|p\|_2^2\right]^2} \, dp \leqslant \frac{c}{1+\|\tilde{k}\|_2^2}.$$

Indeed, to see this, split the integral between the regions $\{\|p\|_2 \leq \|\tilde{k}\|_2/2\}, \{\|p\|_2 \geq 3\|\tilde{k}\|_2/2\}$, and the complement of these. In the first two regions, $\|\tilde{k} + p\|_2 \geq \|\tilde{k}\|_2/2$. In the complement, bound $\|p\|_2$ by $\|\tilde{k}\|_2$ to find:

$$\int_{\|\tilde{k}\|_{2}/2 \leqslant \|p\|_{2} \leqslant 3\|\tilde{k}\|_{2}/2} \frac{1}{1+\|\tilde{k}+p\|^{2}} \frac{\|p\|_{2}}{\left[1+\|p\|_{2}^{2}\right]^{2}} dp \leqslant \frac{c\|\tilde{k}\|_{2}^{3}}{\left[1+\|\tilde{k}\|_{2}^{2}\right]^{2}} \int_{1/2}^{3/2} \frac{1}{1+\|\tilde{k}\|_{2}^{2}u^{2}} du$$

$$\leqslant \frac{c}{1+\|\tilde{k}\|_{2}^{2}}.$$
(A.18)

Using the above bound with $\tilde{k} = \alpha k - q$ yields:

(A.19)

$$\hat{f}(k) \leq c \|k\|_2 \int_0^1 d\alpha \int_{\mathbb{R}^3} \frac{1}{1 + \|q\|_2^2} \cdot \frac{1}{1 + \|\alpha k - q\|_2^2} dq$$

$$=: \|k\|_2 \int_0^1 d\alpha I(\alpha k).$$

For each $\alpha \in [0, 1]$, $I(\alpha k)$ reads:

$$I(\alpha k) = c \int_{0}^{\infty} \int_{0}^{\infty} ds_{1} ds_{2} \int_{\mathbb{R}^{3}} \exp\left[-s_{1}\left(1+\|q\|_{2}^{2}\right) - s_{2}\left(1+\|\alpha k-q\|_{2}^{2}\right)\right] dq$$

$$\leqslant c \int_{0}^{\infty} \int_{0}^{\infty} ds_{1} ds_{2} e^{-s_{1}-s_{2}} \int_{\mathbb{R}^{3}} \exp\left[-(s_{1}+s_{2})\left\|q-\frac{s_{2}\alpha k}{s_{1}+s_{2}}\right\|_{2}^{2} - \frac{s_{1}s_{2}\|\alpha k\|_{2}^{2}}{s_{1}+s_{2}}\right] dq$$

(A.20)
$$\leqslant c \int_{0}^{\infty} \int_{0}^{\infty} ds_{1} ds_{2} \frac{1}{(s_{1}+s_{2})^{3/2}} e^{-s_{1}-s_{2}} \exp\left[-\frac{s_{1}s_{2}\|\alpha k\|_{2}^{2}}{s_{1}+s_{2}}\right].$$

Changing variable from s_1 to $s = s_1/s_2$ yields:

(A.21)

$$I(\alpha k) \leqslant c \int_{0}^{\infty} \int_{0}^{\infty} \exp\left[-s_{2}\left(\frac{s\|\alpha k\|_{2}^{2}}{s+1}+s+1\right)\right] \frac{1}{\sqrt{s_{2}(s+1)^{3/2}}} \, ds \, ds_{2}$$

$$\leqslant c \int_{0}^{\infty} \frac{ds}{(s+1)\sqrt{(s+1)^{2}+s\|\alpha k\|_{2}^{2}}}$$

$$\leqslant \frac{c}{1+\|\alpha k\|_{2}}$$

using

(A.22)
$$\sqrt{(s+1)^2 + s \|\alpha k\|_2^2} \ge s + 1 + \sqrt{s} \|\alpha k\|_2 \ge \sqrt{s} (1 + \|\alpha k\|_2).$$

Thus:

(A.23)
$$\hat{f}(k) \leq c \log(1 + ||k||_2) \Rightarrow |\hat{\psi}(k)| \leq c \log\left(1 + \frac{2||k||_2}{m\pi}\right).$$

We now prove the bound on $\|C_{(m)} \star \psi\|_{L^1 \cap L^2}$. One has, first for the L^2 norm:

(A.24)
$$\|C_{(m)} \star \psi\|_{L^2} \leqslant c \left(\int_{\mathbb{R}^3} \frac{1}{\left(m^2 + \frac{4\|k\|_2^2}{\pi^2}\right)^2} \log\left(1 + \frac{2\|k\|_2}{m\pi}\right)^2 dk \right)^{1/2} \leqslant \frac{c}{m^{1/2}}.$$

By similar computations as in the bound on $\hat{\psi}$, one can prove $|\partial^{\alpha}\hat{\psi}(k)| \leq c_{\alpha}m^{-|\alpha|}(1+\frac{4||k||_2^2}{m^2\pi^2})^{-|\alpha|/2}$ for any multi-index $\alpha \in \mathbb{N}^3$ and any $k \in \mathbb{R}^3$. In addition, for each $j \in \{1, \ldots, d\}$ and each $F \in \mathbb{R}^{\Lambda_{\varepsilon,L}}$,

$$L^{2}\left[1-\cos\left(\frac{2\pi x_{j}}{L}\right)\right]F(x) = \frac{L^{2}}{2L^{d}}\sum_{k\in\Lambda^{*}}e^{ik\cdot x}\hat{F}(k)\left[2-e^{\frac{2i\pi}{L}\mathbf{e}_{j}\cdot x}-e^{-\frac{2i\pi}{L}\mathbf{e}_{j}\cdot x}\right]$$
$$= -\frac{1}{2L^{d}}\sum_{k\in\Lambda^{*}}e^{ik\cdot x}L^{2}\left[\hat{F}\left(k+\frac{2\pi}{L}\mathbf{e}_{j}\right)+\hat{F}\left(k-\frac{2\pi}{L}\mathbf{e}_{j}\right)-2\hat{F}(k)\right].$$
$$(A.25) \qquad \qquad = -\frac{1}{L^{d}}\sum_{k\in\Lambda^{*}}e^{ik\cdot x}\int_{0}^{1}(1-\alpha)\partial_{j}^{2}\hat{F}\left(k+\frac{2\pi\alpha}{L}\mathbf{e}_{j}\right)d\alpha.$$

where, for the last line above, the Fourier transform \hat{F} is extended from Λ^* to \mathbb{R}^3 as in (A.14) for $\hat{\psi}$. Letting $U(x) := L^2 \left(d - \sum_{j=1}^d \cos(2\pi x_j/L) \right)$ for $x \in \Lambda_{\varepsilon,L}$, noticing that $\|U^{-1}\|_{L^2} \leq c$ and

using the Cauchy-Schwarz inequality, this implies

$$\begin{split} \|C_{(m)} \star \psi\|_{L^{1}} &\leqslant \|U^{-1}\|_{L^{2}} \|U \cdot C_{(m)} \star \psi\|_{L^{2}} \\ &\leqslant \|U^{-1}\|_{L^{2}} \left[\left(\frac{1}{L^{d}} \sum_{k \in \Lambda^{*}} \int_{0}^{1} \Delta \left(\hat{C}_{(m)} \hat{\psi} \right) \left(k + \frac{2\pi\alpha \mathbf{e}_{j}}{L} \right) d\alpha \right)^{2} \right]^{1/2} \\ &\leqslant c \left[\int_{\mathbb{R}^{3}} \left(\frac{1}{(m^{2} + \frac{4\|k\|_{2}^{2}}{\pi^{2}})^{2}} + \frac{1}{(m^{2} + \frac{4\|k\|_{2}^{2}}{\pi^{2}})^{3}} \right) \left(1 + \log \left(1 + \frac{2\|k\|_{2}}{m\pi} \right)^{2} \right) dk \\ &\qquad + \int_{\mathbb{R}^{3}} \left(\frac{1}{(m^{2} + \frac{4\|k\|_{2}^{2}}{\pi^{2}})^{3}} + \frac{1}{(m^{2} + \frac{4\|k\|_{2}^{2}}{\pi^{2}})^{4}} \right) dk \right]^{1/2} \\ &A.26) \qquad \leqslant \frac{c}{m^{1/2}} + \frac{c}{m^{3/2}} + \frac{c}{m^{5/2}} \leqslant \frac{c}{m^{1/2}} + \frac{c}{m^{5/2}}, \end{split}$$

where the Laplace operator Δ on the second line above acts on the Fourier variable.

Lemma A.2. The term $\|C_{(m)}(C^2_{(m)} \star C^2_{(m)})\|_{L^1}$ satisfies:

(A.27)
$$\|C_{(m)}(C_{(m)}^2 \star C_{(m)}^2)\|_{L^1} \leq \frac{c}{m^4} \mathbf{1}_{d=2} + \frac{c}{m} \mathbf{1}_{d=3}.$$

Proof. The proof is similar to that of Lemma A.1, so we only give a sketch. Using the Fourier representation, one has:

(A.28)
$$\|C_{(m)}(C_{(m)}^2 \star C_{(m)}^2)\|_{L^1} = \frac{1}{L^d} \sum_{k \in \Lambda^*} \frac{1}{m^2 + \theta(k)} \Big(\frac{1}{L^d} \sum_{p \in \Lambda^*} \frac{1}{m^2 + \theta(p)} \frac{1}{m^2 + \theta(k-p)} \Big)^2.$$

The idea is again to compare sums to integrals, and then change variables to factor out the mass dependence:

(A.29)
$$\|C_{(m)}(C_{(m)}^2 \star C_{(m)}^2)\|_{L^1} \leq \frac{cm^{3d}}{m^{10}} \int_{\mathbb{R}^d} \frac{dk}{1+\|k\|_2^2} \Big(\int_{\mathbb{R}^d} dp \, \frac{1}{1+\|p\|_2^2} \frac{1}{1+\|k-p\|_2^2}\Big)^2$$

It remains to check that the last integral is finite. Notice however that the integral in the square above is exactly I(k), as defined in (A.19) in the three-dimensional case. It follows from (A.21) that $I(k) \leq C/(1+||k||_2)$ in dimension 3. The same arguments also work in dimension 2, which concludes the proof.

A.2. Differences of the counterterms. In this section (and as in Section 4), C_t is the covariance $C_t = (-\Delta^{\varepsilon} + m^2 + 1/t)^{-1}$ for t > 0.

Lemma A.3. In dimension $d \in \{2,3\}$, the difference of the counterterms at parameters t > 0and $+\infty$ satisfies:

(A.30)

(

$$0 \leqslant C_{\infty}(0) - C_{t}(0) =: \eta_{t} \leqslant c \log \left(1 + \frac{1}{m^{2}t}\right) \mathbf{1}_{d=2} + cm \left(\sqrt{1 + \frac{1}{tm^{2}}} - 1\right) \mathbf{1}_{d=3},$$

$$0 \leqslant \|C_{\infty}^{3}\|_{L^{1}} - \|C_{t}^{3}\|_{L^{1}} := \gamma_{t} \leqslant \frac{c}{m^{2}(m^{2}t+1)} \mathbf{1}_{d=2} + c \log \left(1 + \frac{1}{m^{2}t}\right) \mathbf{1}_{d=3}.$$

Proof. The proof is similar to and uses notations of the proof of Lemma A.1. The key argument consists in passing from discrete sums to integrals, using the relation $f(1) - f(0) = \int_0^1 f'(t) dt$. For $C_{\infty}(0) - C_t(0)$, one has:

(A.31)

$$C_{\infty}(0) - C_{t}(0) = \frac{1}{L^{d}} \sum_{k \in \Lambda^{*}} \left[\frac{1}{m^{2} + \theta(k)} - \frac{1}{m^{2} + \frac{1}{t} + \theta(k)} \right]$$

$$= \frac{1}{t} \int_{0}^{1} d\alpha \frac{1}{L^{d}} \sum_{k \in \Lambda^{*}} \frac{1}{\left(m^{2} + \frac{\alpha}{t} + \theta(k)\right)^{2}}$$

$$\leqslant \frac{c}{t} \int_{0}^{1} d\alpha \int_{\mathbb{R}^{d}} \frac{dk}{\left(m^{2} + \frac{\alpha}{t} + \frac{4||k||_{2}^{2}}{\pi^{2}}\right)^{2}}.$$

Changing variables from k to $p = 2k(\pi\sqrt{m^2 + \alpha/t})^{-1}$ yields:

(A.32)

$$C_{\infty}(0) - C_{t}(0) \leq \int_{0}^{1} \frac{1}{t\left(m^{2} + \frac{\alpha}{t}\right)^{2 - \frac{d}{2}}} d\alpha \times \int_{\mathbb{R}^{d}} \frac{1}{(1 + \|p\|_{2}^{2})^{2}} dp$$

$$\leq c \int_{0}^{1} \frac{1}{t\left(m^{2} + \frac{\alpha}{t}\right)^{2 - \frac{d}{2}}} d\alpha$$

$$= c \log\left(1 + \frac{1}{m^{2}t}\right) \mathbf{1}_{d=2} + cm\left(\sqrt{1 + \frac{1}{tm^{2}}} - 1\right) \mathbf{1}_{d=3}.$$

For the C^3 counterterm γ_t , one has similarly:

$$\|C_{\infty}^{3}\|_{L^{1}} - \|C_{t}^{3}\|_{L^{1}} = \frac{1}{L^{2d}} \sum_{p,q \in \Lambda^{*}} \left[\frac{1}{m^{2} + \theta(p)} \frac{1}{m^{2} + \theta(q)} \frac{1}{m^{2} + \theta(p+q)} - \frac{1}{m^{2} + \frac{1}{t} + \theta(p)} \frac{1}{m^{2} + \frac{1}{t} + \theta(q)} \frac{1}{m^{2} + \frac{1}{t} + \theta(p+q)} \right]$$
(A.33)
$$= \frac{3}{t} \int_{0}^{1} d\alpha \frac{1}{L^{2d}} \sum_{p,q \in \Lambda^{*}} \frac{1}{m^{2} + \frac{\alpha}{t} + \theta(p)} \frac{1}{(m^{2} + \frac{\alpha}{t} + \theta(q))^{2}} \frac{1}{m^{2} + \frac{\alpha}{t} + \theta(p+q)}.$$

Bounding θ by convexity and comparing discrete sums to integrals, then changing variables from p, q to $2p(\pi\sqrt{m^2 + \alpha/t})^{-1}$ and $2q(\pi\sqrt{m^2 + \alpha/t})^{-1}$ yields:

$$\|C_{\infty}^{3}\|_{L^{1}} - \|C_{t}^{3}\|_{L^{1}} \leqslant \frac{3c}{t} \int_{0}^{1} d\alpha \int_{\mathbb{R}^{2d}} \frac{1}{m^{2} + \frac{\alpha}{t} + \frac{4\|p\|_{2}^{2}}{\pi^{2}}} \frac{1}{(m^{2} + \frac{\alpha}{t} + \frac{4\|q\|_{2}^{2}}{\pi^{2}})^{2}} \frac{1}{m^{2} + \frac{\alpha}{t} + \frac{4\|p+q\|_{2}^{2}}{\pi^{2}}} dp dq$$
(A.34)
$$= \int_{0}^{1} d\alpha \frac{3c\pi^{2d}}{t(m^{2} + \frac{\alpha}{t})^{4-d}} \int_{\mathbb{R}^{6}} \frac{1}{1 + \|p\|_{2}^{2}} \frac{1}{(1 + \|q\|_{2}^{2})^{2}} \frac{1}{1 + \|p+q\|_{2}^{2}} dp dq.$$

Since the integral over p, q is bounded, therefore:

(A.35)
$$\|C_{\infty}^{3}\|_{L^{1}} - \|C_{t}^{3}\|_{L^{1}} \leq c \int_{0}^{1} \frac{1}{t\left(m^{2} + \frac{\alpha}{t}\right)^{4-d}} d\alpha$$
$$\leq \frac{c}{m^{2}(m^{2}t+1)} \mathbf{1}_{d=2} + c \log\left(1 + \frac{1}{m^{2}t}\right) \mathbf{1}_{d=3}.$$

Acknowledgements

We thank T. Bodineau and J. Ding for helpful discussions, as well as G. Ferrando for help with the Fourier computations. This work was supported by the European Research Council under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 851682 SPINRG). R.B. also acknowledges the hospitality of the Department of Mathematics at McGill University where part of this work was carried out.

References

- [1] D. Bakry, I. Gentil, and M. Ledoux. Analysis and geometry of Markov diffusion operators, volume 348 of Grundlehren der Mathematischen Wissenschaften. Springer, Cham, 2014.
- [2] N. Barashkov and M. Gubinelli. A variational method for Φ_3^4 . Duke Math. J., 169(17):3339–3415, 2020.

- [3] R. Bauerschmidt and T. Bodineau. Log-Sobolev inequality for the continuum sine-Gordon model. Comm. Pure Appl. Math., 74(10):2064–2113, 2021.
- [4] R. Bauerschmidt and B. Dagallier. Log-Sobolev inequality for near critical Ising models. 2022. Preprint.
- [5] T. Bodineau and B. Helffer. The log-Sobolev inequality for unbounded spin systems. J. Funct. Anal., 166(1):168–178, 1999.
- [6] H.J. Brascamp and E.H. Lieb. On Extensions of the Brunn-Minkowski and Prékopa-Leindler Theorems, Including Inequalities for Log Concave Functions, and with an Application to the Diffusion Equation, pages 441–464. Springer Berlin Heidelberg, 2002.
- [7] D. Brydges, J. Fröhlich, and T. Spencer. The random walk representation of classical spin systems and correlation inequalities. *Commun. Math. Phys.*, 83(1):123–150, 1982.
- [8] D.C. Brydges, J. Fröhlich, and A.D. Sokal. A new proof of the existence and nontriviality of the continuum φ_2^4 and φ_3^4 quantum field theories. *Commun. Math. Phys.*, 91(2):141–186, 1983.
- [9] D.C. Brydges, J. Fröhlich, and A.D. Sokal. The random-walk representation of classical spin systems and correlation inequalities. II. The skeleton inequalities. *Commun. Math. Phys.*, 91(1):117–139, 1983.
- [10] R. Catellier and K. Chouk. Paracontrolled distributions and the 3-dimensional stochastic quantization equation. Ann. Probab., 46(5):2621–2679, 2018.
- [11] A. Chandra, T.S. Gunaratnam, and H. Weber. Phase transitions for ϕ_3^4 . Commun. Math. Phys., 392(2):691–782, 2022.
- [12] G. Da Prato and A. Debussche. Strong solutions to the stochastic quantization equations. Ann. Probab., 31(4):1900–1916, 2003.
- [13] J. Ding, J. Song, and R. Sun. A new correlation inequality for Ising models with external fields. Probab. Theory Related Fields, 2022+.
- [14] J. Fröhlich, B. Simon, and T. Spencer. Infrared bounds, phase transitions and continuous symmetry breaking. *Commun. Math. Phys.*, 50(1):79–95, 1976.
- [15] J. Glimm and A. Jaffe. Quantum physics. Springer-Verlag, second edition, 1987. A functional integral point of view.
- [16] J. Glimm, A. Jaffe, and T. Spencer. Phase transitions in P (φ)₂ quantum fields. Bull. Amer. Math. Soc., 82(5):713–715, 1976.
- [17] M. Gubinelli and M. Hofmanová. A PDE construction of the Euclidean ϕ_3^4 quantum field theory. Commun. Math. Phys., 384(1):1–75, 2021.
- [18] M. Gubinelli, P. Imkeller, and N. Perkowski. Paracontrolled distributions and singular PDEs. Forum Math. Pi, 3:e6, 75, 2015.
- [19] A. Guionnet and B. Zegarlinski. Lectures on logarithmic Sobolev inequalities. In Séminaire de Probabilités, XXXVI, volume 1801 of Lecture Notes in Math., pages 1–134. Springer, Berlin, 2003.
- [20] M. Hairer. A theory of regularity structures. Invent. Math., 198(2):269–504, 2014.
- [21] M. Hairer. Regularity structures and the dynamical Φ_3^4 model. In *Current developments in mathematics 2014*, pages 1–49. Int. Press, Somerville, MA, 2016.

- [22] M. Hairer and K. Matetski. Discretisations of rough stochastic PDEs. Ann. Probab., 46(3):1651–1709, 2018.
- [23] M. Hairer and P. Schönbauer. The support of singular stochastic PDEs. Forum Math. Pi, 2022.
- [24] A. Kupiainen. Renormalization group and stochastic PDEs. Ann. Henri Poincaré, 17(3):497– 535, 2016.
- [25] M. Ledoux. Logarithmic Sobolev inequalities for unbounded spin systems revisited. In Séminaire de Probabilités, XXXV, volume 1755 of Lecture Notes in Math., pages 167–194. Springer, Berlin, 2001.
- [26] A. Moinat and H. Weber. Space-time localisation for the dynamic Φ_3^4 model. Comm. Pure Appl. Math., 73(12):2519–2555, 2020.
- [27] J.-C. Mourrat and H. Weber. The dynamic Φ_3^4 model comes down from infinity. Commun. Math. Phys., 356(3):673–753, 2017.
- [28] J.-C. Mourrat and H. Weber. Global well-posedness of the dynamic Φ^4 model in the plane. Ann. Probab., 45(4):2398–2476, 2017.
- [29] J. Polchinski. Renormalization and effective lagrangians. Nuclear Physics B, 231(2):269 295, 1984.
- [30] B. Simon. The $P(\phi)_2$ Euclidean (quantum) field theory. Princeton University Press, 1974. Princeton Series in Physics.
- [31] B. Simon and R.B. Griffiths. The $(\phi^4)_2$ field theory as a classical Ising model. Commun. Math. Phys., 33:145–164, 1973.
- [32] P. Tsatsoulis and H. Weber. Spectral gap for the stochastic quantization equation on the 2-dimensional torus. Ann. Inst. Henri Poincaré Probab. Stat., 54(3):1204–1249, 2018.
- [33] N. Yoshida. The log-Sobolev inequality for weakly coupled lattice fields. Probab. Theory Related Fields, 115(1):1–40, 1999.
- [34] B. Zegarlinski. The strong decay to equilibrium for the stochastic dynamics of unbounded spin systems on a lattice. *Commun. Math. Phys.*, 175(2):401–432, 1996.
- [35] R. Zhu and X. Zhu. Dirichlet form associated with the Φ_3^4 model. *Electron. J. Probab.*, 23:Paper No. 78, 31, 2018.
- [36] R. Zhu and X. Zhu. Lattice approximation to the dynamical Φ_3^4 model. Ann. Probab., 46(1):397-455, 2018.