

Euclidean preferences in the plane under ℓ_1 , ℓ_2 and ℓ_∞ norms

Bruno Escoffier^{1,2}

Olivier Spanjaard¹

Magdaléna Tydrichová¹

¹Sorbonne Université, CNRS, LIP6, F-75005 Paris, France

²Institut Universitaire de France

Abstract

We present various results about Euclidean preferences in the plane under ℓ_1 , ℓ_2 and ℓ_∞ norms. When there are four candidates, we show that the maximal size (in terms of the number of pairwise distinct preferences) of Euclidean preference profiles in \mathbb{R}^2 under norm ℓ_1 or ℓ_∞ is 19. Whatever the number of candidates, we prove that at most four distinct candidates can be ranked in last position of a two-dimensional Euclidean preference profile under norm ℓ_1 or ℓ_∞ , which generalizes the case of one-dimensional Euclidean preferences (for which it is well known that at most two candidates can be ranked last). We generalize this result to 2^d (resp. $2d$) for ℓ_1 (resp. ℓ_∞) for d -dimensional Euclidean preferences. We also establish that the maximal size of a two-dimensional Euclidean preference profile on m candidates under norm ℓ_1 is in $\Theta(m^4)$, i.e., the same order of magnitude as under norm ℓ_2 . Finally, we provide a new proof that two-dimensional Euclidean preference profiles under norm ℓ_2 for four candidates can be characterized by three voter-maximal two-dimensional Euclidean profiles. This proof is a simpler alternative to that proposed by Kamiya et al (2011).

1 Introduction

The study of domain restrictions is a long standing research topic in modern social choice theory, dating back to the work of Black (1948) on the single-peaked domain. As emphasized by Barberà et al (2020), Arrow already attached importance to studying the role of domain conditions in determining the validity of his impossibility theorem, with two chapters of *Social Choice and Individual Values* (Arrow, 1951) devoted to this topic. For a survey about domain restrictions, the reader may refer to the works of Gaertner (2001; 2002) and Barberà et al (2013). For a computational perspective, one may refer to the survey recently conducted by Elkind et al (2022).

The spatial model of social choice is an important stream of research in this topic, pioneered by the works of Hotelling (1929) and Downs (1957, chapter 8). We focus here on *Euclidean preferences*, where candidates and voters are viewed as points in \mathbb{R}^d , and the preferences of voters are decreasing with their Euclidean distance to the candidates. Note that in this article, by abuse of language, we will use the expression *Euclidean preferences in \mathbb{R}^d under norm ℓ_k* ($k \in \{1, 2, \infty\}$) when measuring distances using norm ℓ_k .

The most widely studied Euclidean preferences are those that are derived by measuring the distances with the ℓ_2 norm (see, e.g., the works of Bennett and Hays, 1960; Bogomolnaia and Laslier, 2004), but the ℓ_1 and ℓ_∞ norms have also been considered in the literature (see, e.g., the work of Peters, 2017). From a more operational point of view, spatial representations are used in particular in voting advice applications (e.g., Wahl-O-Mat in Germany, Smartvote in Switzerland, Vote Compass in the United States, and many others in multiple countries), i.e., online tools that helps the voter choose the candidate closest to her political stances, and actually often provides her a full ranking of candidates according to her answers to a survey on a range of policy statements. The answers are indeed converted into positions on different dimensions, each position reporting on the level of agreement on a particular policy statement. Norm ℓ_1 is typically used when there are many dimensions, while norm ℓ_2 is used when the number of dimensions is lower (Moreno et al, 2022; Isotalo, 2020). For an overview of the topic of voting advice applications, the reader may refer to the survey by Garzia and Marschall (2019).

In this work, we put a special emphasis on the case $d = 2$ and we consider the norms ℓ_2 , ℓ_1 and ℓ_∞ . An original focus of our work is to try to identify the differences between the three norms. Hence, we are interested in the following questions:

- Are there forbidden structures that make a profile not Euclidean, under some of the three norms?

- Given a set of m candidates, what is the maximal size (in terms of the number of pairwise distinct preferences) of profiles that are Euclidean?
- Are there some differences or similarities between the norms in the expressivity of Euclidean preferences? Put another way, are there profiles that are Euclidean with one norm and not with another one?

We first show a structural result on ℓ_1 -Euclidean and ℓ_∞ -Euclidean profiles in \mathbb{R}^d , namely that in an ℓ_∞ -Euclidean profile there are at most $2d$ candidates ranked last by at least one voter, while there are at most 2^d such candidates for an ℓ_1 -Euclidean profile. While this result is not hard to prove, it is interesting in several aspects: first, it provides a strong difference with ℓ_2 , as we can easily build ℓ_2 -Euclidean profiles in the plane where each candidate is ranked last at least once¹. Second, it is an interesting generalization of the case of 1-dimensional Euclidean preferences, where it is well known that at most 2 candidates can be ranked last. Finally, while it is known that ℓ_1 and ℓ_∞ are equivalent when $d \leq 2$ (meaning that, for $d \in \{1, 2\}$, a profile is ℓ_1 -Euclidean if and only if it is ℓ_∞ -Euclidean), an immediate corollary of this structural result is that this equivalence does *not* hold for $d \geq 3$.

We then focus on the case $d = 2$. As it can easily be seen that every profile with 2 or 3 candidates is ℓ_1 -Euclidean (thus ℓ_∞ -Euclidean) and ℓ_2 -Euclidean (Bogomolnaia and Laslier, 2004), we focus in Section 5 on the case of $m = 4$ candidates. We first give an explicit example of a profile which is ℓ_1 -Euclidean but not ℓ_2 -Euclidean. We then focus on the maximal size (in terms of the number of pairwise distinct preferences) of profiles on 4 candidates that are Euclidean. It is known since the work of Bennett and Hays (1960) that the maximal size is 18 for ℓ_2 . We show that this maximal size is exactly 19 for ℓ_1 . Then, we give a new proof that a profile on 4 candidates is ℓ_2 -Euclidean if and only if it is a subprofile of one of three voter-maximal two-dimensional Euclidean profiles (involving 18 voters). Kamiya et al (2011) proved the same result, but they rely on a link they establish with the problem of enumerating chambers of hyperplane arrangements (for an introduction to the topic, see, e.g., the chapter of Stanley et al, 2004), while we use simpler and purely geometrical arguments.

We then focus on the case $m \geq 5$. We focus on the maximal size of profiles which are Euclidean. We show that, despite the strong restriction on the number of candidates ranked last by some voter (at most $2d = 4$), the maximal size of an ℓ_1 -Euclidean profile is $\Theta(m^4)$, i.e., of the same order of magnitude as for ℓ_2 (as shown by Bennett and Hays, 1960).

Organization of the article. We provide a brief overview of the related work in Section 2. We then give in Section 3 some formal definitions, examples, and focus on the relation between ℓ_1 and ℓ_∞ norms for Euclidean profiles. Some geometric properties of representations of ℓ_1 -Euclidean profiles are given in Section 4. We highlight some differences with the ℓ_2 norm, and derive some properties that will be useful for the results of subsequent sections. Then we present in Section 5 our results with $m = 4$ candidates, while Section 6 deals with the general case, i.e., for an arbitrary number of candidates. We conclude the article in Section 7 by providing some research directions.

2 Related work

The one-dimensional case. Chen et al (2015) proved that one-dimensional Euclidean preference profiles cannot be characterized in terms of finitely many forbidden substructures, i.e., one cannot enumerate a finite set of substructures (also called *obstructions*) such that a profile is one-dimensional Euclidean if and only if it contains none of the substructures in the list. It is nevertheless known that one-dimensional Euclidean preference profiles can be recognised in polynomial time in the number of voters and candidates, as first shown by Doignon and Falmagne (1994), and then by Knoblauch (2010) and Elkind and Faliszewski (2014). Very recently, Chen and Grottko (2021) characterized one-dimensional Euclidean preference profiles with a small number of candidates and voters. In particular, they showed that any profile with at most 5 candidates is Euclidean if and only if it is single-peaked and single-crossing (where both single-peaked and single-crossing profiles can be characterized via finitely many finite obstructions). They finally identified the smallest single-peaked and single-crossing profile which is not Euclidean.

¹Place for instance the candidates on a circle, and for each candidate c place a voter v which is the antipode of c with respect to the center of the circle.

The multidimensional case. Bennett and Hays (1960) as well as Hays and Bennett (1961) proposed several methods to estimate the minimum value of d to be able to embed a preference profile in a d -dimensional space, i.e., to associate a point in \mathbb{R}^d to each voter and each candidate so that the voters' preferences are decreasing with the distance to the candidates. In particular, they established that the maximum cardinality of a ℓ_2 -Euclidean profile on m candidates in dimension d is equal to $\sum_{k=m-d}^m |s(m, k)|$, where $s(m, k)$ are the (unsigned) Stirling numbers of the first kind. The same result has been found by Good and Tideman (1977). Later on, Bogomolnaia and Laslier (2004) showed that to guarantee any profile of n preferences on m candidates to be d -Euclidean, it is necessary and sufficient to have d between $\min\{n-1, m-1\}$ and $\min\{n, m-1\}$. Recently, an analogous result was shown by Chen et al (2022) for preference profiles using an ℓ_1 metric. More precisely, Chen et al (2022) showed that each preference profile with m alternatives and n voters is d -Euclidean with respect to the norm ℓ_1 whenever $d \geq \min\{n, m-1\}$. Also, they studied the smallest non-Euclidean profiles in case of $d = 2$. As mentioned earlier, Kamiya et al (2011) studied the question of counting and enumerating voter-maximal ℓ_2 -Euclidean profiles in \mathbb{R}^d , according to the number m of candidates. They provide a formula for the number of voter-maximal profiles if $d = m-2$, and they were able to enumerate them for $m = 4$. Regarding the computational aspects, Peters (2017) proved that the recognition problem (i.e., deciding whether or not a preference profile is ℓ_2 -Euclidean in dimension d) is NP-hard for $d > 1$, and that some Euclidean preference profiles require exponentially many bits in order to specify any Euclidean embedding.

3 Preliminaries

3.1 Euclidean preference profile

We consider a (finite) set V of n voters, and a (finite) set C of m candidates. Each voter gives her preference $>_v$ over the set of candidates as a ranking (total ranking, without tie). The set $\{>_{v_1}, >_{v_2}, \dots, >_{v_n}\}$ of preferences of voters in V on candidates in C is denoted by R , and the couple $\mathcal{P} = (C, R)$ is called a preference profile. We write $c_1 >_v c_2$ if voter v prefers c_1 to c_2 . For conciseness, we will often write the preference $c_i >_v c_j >_v \dots >_v c_k$ for a voter v as (c_i, c_j, \dots, c_k) .

Definition 1. Let $d \geq 1$ be an integer, and $\|\cdot\|_\ell$ a norm on \mathbb{R}^d . The profile $\mathcal{P} = (C, R)$ of preferences of n voters over m candidates is ℓ -Euclidean in \mathbb{R}^d if there exists a mapping $f : V \cup C \rightarrow \mathbb{R}^d$ such that for each $v \in V$ and each $\{c_1, c_2\} \subseteq C$:

$$c_1 >_v c_2 \Rightarrow \|f(v) - f(c_1)\|_\ell < \|f(v) - f(c_2)\|_\ell$$

The mapping is called an ℓ -Euclidean representation of the profile in \mathbb{R}^d . Obviously, such a representation is not necessarily unique. A profile for which there exists a ℓ -Euclidean representation in \mathbb{R}^d is called ℓ -Euclidean in \mathbb{R}^d .

We note that if two voters v and v' have the same preference, then $\mathcal{P} = (C, R)$ is ℓ -Euclidean in \mathbb{R}^d if and only if $(C, R \setminus \{>_{v'}\})$ is ℓ -Euclidean in \mathbb{R}^d . So, without loss of generality, throughout the article we consider preference profiles where any pair of voters have different preferences. We define the *size* (or cardinality) of a profile as the number of votes (or, equivalently, voters).

We note also that, as preferences in \mathcal{P} are strict orders, we could replace the implication in Definition 1 by an equivalence - hence, the last line of the definition becomes:

$$c_1 >_v c_2 \Leftrightarrow \|f(v) - f(c_1)\|_\ell < \|f(v) - f(c_2)\|_\ell.$$

3.2 Boundary hypersurfaces

Consider a profile $\mathcal{P} = (C, R)$, an integer d and a norm $\|\cdot\|_\ell$ on \mathbb{R}^d . Given a set of points $p_1, \dots, p_m \in \mathbb{R}^d$, we now study the question of determining whether there exists a mapping $f : V \cup C \rightarrow \mathbb{R}^d$ such that:

1. for each $i \in \{1, \dots, m\}$, $f(c_i) = p_i$;
2. f is a ℓ -Euclidean representation of \mathcal{P} in \mathbb{R}^d .

To build an ℓ -Euclidean representation in \mathbb{R}^d , it is sufficient (and necessary) to find for each $v \in V$ a value $f(v)$ such that f fulfills the condition in Definition 1. Let us define, for each v , the set $D_f(v)$ of such possible values:

$$D_f(v) = \{f(v) \in \mathbb{R}^d : \forall \{c_1, c_2\} \subseteq C, c_1 >_v c_2 \Rightarrow \|f(v) - f(c_1)\|_\ell < \|f(v) - f(c_2)\|_\ell\}$$




Figure 1: An ℓ_2 -representation of the complete profile on 3 candidates.

With this notation, the profile is ℓ -Euclidean in \mathbb{R}^d if and only if there exists a mapping f such that for each v , $D_f(v)$ is a non-empty set. The natural question is to characterise $D_f(v)$ for each voter v . To this end, we introduce the following notion:

Definition 2. For a pair $\{c_1, c_2\} \subseteq C$ of candidates mapped in positions $f(c_1)$ and $f(c_2)$, the set of points $p \in \mathbb{R}^d$ such that $\|f(c_1) - p\|_\ell = \|f(c_2) - p\|_\ell$ is called the boundary hypersurface of c_1 and c_2 (or just hypersurface in what follows), and is denoted by $H_f(c_1, c_2)$. We denote then by $D_f(c_1, c_2)$ the set of points $p \in \mathbb{R}^d$ such that $\|f(c_1) - p\|_\ell < \|f(c_2) - p\|_\ell$, and by $D_f(c_2, c_1)$ the set of points $p \in \mathbb{R}^d$ such that $\|f(c_1) - p\|_\ell > \|f(c_2) - p\|_\ell$.

It is easy to convince oneself that:

$$D_f(v) = \bigcap_{c_i >_v c_j} D_f(c_i, c_j)$$

Note that if $\ell = \ell_2$, $D_f(v)$ is convex (as an intersection of half spaces bounded by a hyperplane) for each $v \in V$. However, we will see later that $D_f(v)$ is not necessarily convex if $\ell = \ell_1$ or $\ell = \ell_\infty$. For conciseness, and only if no confusion is possible, we will omit the representation function f in the notions introduced in Definition 2. Thus, we will write $H(c_i, c_j)$, resp. $D(c_i, c_j)$ and $D(v)$, instead of $H_f(c_i, c_j)$, resp. $D_f(c_i, c_j)$ and $D_f(v)$.

As $D_f(c_1, c_2)$ depends only on the positions of c_1 and c_2 in \mathbb{R}^d , and hence $D_f(v)$ on the positions of c_1, \dots, c_m , the definition of ℓ -Euclidean profiles in \mathbb{R}^d can be reformulated as follows:

Proposition 1. Let $d \geq 1$ be an integer, and $\|\cdot\|_\ell$ be a norm on \mathbb{R}^d . The profile $\mathcal{P} = (C, R)$ of preferences of n voters over m candidates is ℓ -Euclidean in \mathbb{R}^d if and only if there exists a mapping $f : C \rightarrow \mathbb{R}^d$ such that $D_f(v)$ is non-empty for each $v \in V$.

Given a representation function f and a voter v , we will call the set $D_f(v)$ an *area*, as geometrically, it corresponds to an area of the plane. Thus, $D_f(v)$ is an area of preference ranking $>_f$ with respect to the representation f . By abuse of notation, the terms of area and (its corresponding) preference ranking will be used interchangeably.

Example 1. Consider a preference profile \mathcal{P} with 3 candidates $\{c_1, c_2, c_3\}$ and the 6 possible preferences on these 3 candidates. Figure 1 shows a ℓ_2 -Euclidean representation of \mathcal{P} in \mathbb{R}^2 : the 3 hypersurfaces $H(c_1, c_2)$, $H(c_1, c_3)$ and $H(c_2, c_3)$, as well as the 6 non-empty areas $D(v)$ (with the corresponding preference written in the area).

We now give a profile on 4 candidates which is *not* ℓ_2 -Euclidean in \mathbb{R}^2 . As we will see, the fact that it is not Euclidean is proved by geometric arguments.




Figure 2: The different preference areas obtained from candidates c_1 , c_2 and c_3 forming a triangle. Each preference R_v is a subset of an area, the precise contours of which depends on the position of c_4 in \mathbb{R}^2 .

Example 2. Let us consider the following profile $\mathcal{P} = (C, R)$ with 9 voters and 4 candidates, where R_v is the preference of voter v :

$$\begin{aligned} R_1 &: (c_4, c_3, c_1, c_2) & R_2 &: (c_3, c_4, c_1, c_2) \\ R_3 &: (c_4, c_3, c_2, c_1) & R_4 &: (c_3, c_4, c_2, c_1) \\ R_5 &: (c_2, c_1, c_4, c_3) & R_6 &: (c_2, c_1, c_3, c_4) \\ R_7 &: (c_1, c_2, c_4, c_3) & R_8 &: (c_1, c_2, c_3, c_4) \\ R_9 &: (c_2, c_3, c_1, c_4) \end{aligned}$$

Let us show that this profile is not ℓ_2 -Euclidean in \mathbb{R}^2 (while we will see later that it is ℓ_1 -Euclidean in \mathbb{R}^2). By contradiction, assume that a ℓ_2 -Euclidean representation in \mathbb{R}^2 exists. The points c_1, c_2, c_3 form necessarily a (non-degenerate) triangle, as 5 different rankings over $\{c_1, c_2, c_3\}$ are present in the profile, and at most 4 can be represented if c_1, c_2, c_3 are aligned in \mathbb{R}^2 . Figure 2 illustrates the different preference areas obtained from candidates c_1, c_2, c_3 forming a triangle.

Note that for each $i \in \{1, 2, 3, 4\}$, we obtain R_{2i-1} from R_{2i}^2 by swapping c_3 and c_4 . Thus, $H(c_3, c_4)$ has to go through the area $c_3 > c_1 > c_2$ to separate R_1 and R_2 , through the area $c_3 > c_2 > c_1$ to separate R_3 and R_4 and finally through the areas $c_2 > c_1 > c_3$ and $c_1 > c_2 > c_3$ to separate R_5 and R_6 , and R_7 and R_8 (see Figure 2 for more clarity). This is not possible, as any straight line can cross at most 3 of these 4 areas. Indeed, if a straight line crosses both the area containing $\{R_7, R_8\}$ and the one containing $\{R_5, R_6\}$, then it must intersect $H(c_1, c_2)$ below point A . Similarly, if it crosses both the area containing $\{R_1, R_2\}$ and the one containing $\{R_3, R_4\}$, then it must intersect $H(c_1, c_2)$ above point A . Thus, to cross the 4 areas, it must intersect $H(c_1, c_2)$ twice, a contradiction.

Hence, no ℓ_2 -Euclidean representation of \mathcal{P} exists in \mathbb{R}^2 . \square

3.3 Relation between norms ℓ_1 and ℓ_∞

We consider here the case where $\ell = \ell_1$ or $\ell = \ell_\infty$. Given $\delta \geq 0$, we denote by $\mathcal{S}_\delta^\ell(p)$ the ℓ -sphere of radius δ centered in $p \in \mathbb{R}^d$. Formally:

$$\mathcal{S}_\delta^\ell(p) = \{q \in \mathbb{R}^d : \|p - q\|_\ell = \delta\}$$

With this notation, we characterise $H(c_1, c_2)$ as:

$$H(c_1, c_2) = \bigcup_{\delta \geq 0} (\mathcal{S}_\delta^\ell(f(c_1)) \cap \mathcal{S}_\delta^\ell(f(c_2))) \quad (1)$$

²These pairs correspond to rows in the display of the profile given at the beginning of example




Figure 3: For $d = 2$, the spheres $\mathcal{S}_\delta^{\ell_1}(p)$ and $\mathcal{S}_{\frac{\delta}{\sqrt{2}}}^{\ell_\infty}(p)$ are homothetic via the rotation of 45° .

For $d = 2$, for all $\delta \geq 0$ and $p \in \mathbb{R}^2$, the spheres $\mathcal{S}_\delta^{\ell_1}(p)$ and $\mathcal{S}_{\frac{\delta}{\sqrt{2}}}^{\ell_\infty}(p)$ are homothetic via the rotation of 45° (see Figure 3). Together with the characterisation of $H(c_1, c_2)$ in Equation (1), this yields the following observation, already noted by Lee and Wong (1980).

Observation 1 (Lee and Wong, 1980). *A preference profile is ℓ_1 -Euclidean in \mathbb{R}^2 if and only if it is ℓ_∞ -Euclidean in \mathbb{R}^2 .*

We now show that this equivalence is *not* true for $d \geq 3$. This is actually a corollary of the following proposition, which provides a structural property of ℓ_∞ -Euclidean and ℓ_1 -Euclidean profiles in \mathbb{R}^d .

Proposition 2. *In an ℓ_∞ -Euclidean profile in \mathbb{R}^d , at most $2d$ candidates are ranked last by at least one voter. In an ℓ_1 -Euclidean profile in \mathbb{R}^d , at most 2^d candidates are ranked last by at least one voter. These bounds are tight for all d .*

Proof. Let us first consider an ℓ_∞ -Euclidean profile $\mathcal{P} = (V, C)$, and a corresponding mapping f . We denote by $f_i(x)$ the position of candidate/voter x on the i^{th} coordinate. For $i = 1, \dots, d$, let us denote by $c_{j_i}^m$ and $c_{j_i}^M$ the candidates that have minimal and maximal i^{th} coordinate. There are at most $2d$ of them (some candidates may be extremal on several coordinates). Take a candidate c which is *not* among these extremal candidates, and take a voter v . We show that c cannot be ranked last by v . Let i be such that $\|f(c) - f(v)\|_{\ell_\infty} = |f_i(c) - f_i(v)|$.

If $f_i(c) \geq f_i(v)$, then

$$\|f(c) - f(v)\|_{\ell_\infty} = f_i(c) - f_i(v) \leq f_i(c_{j_i}^M) - f_i(v) \leq \|f(c_{j_i}^M) - f(v)\|_{\ell_\infty}.$$

As the two distances cannot be the same (no tie in the preferences), c is ranked before $c_{j_i}^M$ by v .

If $f_i(c) < f_i(v)$, then

$$\|f(c) - f(v)\|_{\ell_\infty} = f_i(v) - f_i(c) \leq f_i(v) - f_i(c_{j_i}^m) \leq \|f(c_{j_i}^m) - f(v)\|_{\ell_\infty}.$$

Again, c is ranked before $c_{j_i}^m$ by v . In both cases, c is not ranked last.

To show the tightness of the bound, we consider a profile on $2d$ candidates where $f(c_{2i-1})$ is -1 on coordinate i and 0 on all other coordinates, and $f(c_{2i}) = -f(c_{2i-1})$. There are also $2d$ voters, with $f(v_i) = f(c_i)$ for $i = 1, \dots, 2d$. Then it is easy to see that c_{2i-1} is ranked last by v_{2i} , and c_{2i} is ranked last by v_{2i-1} .³

Let us now focus on ℓ_1 -Euclidean profiles. For each vector u in $\{-1, 1\}^d$, let c_u be a candidate which maximizes $u \cdot f(c) = \sum_{i=1}^d u_i \cdot f_i(c)$. As previously, consider a candidate c which is not among these (at most) 2^d extreme candidates, and take a voter v . We show that c cannot be ranked last by v . By definition, $\|f(c) - f(v)\|_{\ell_1} = \sum_{i=1}^d |f_i(c) - f_i(v)|$. Define the vector u as $u_i = 1$ if $f_i(c) \geq f_i(v)$ and $u_i = -1$ otherwise. Then:

$$\begin{aligned} \|f(c) - f(v)\|_{\ell_1} &= \sum_{i=1}^d u_i \cdot (f_i(c) - f_i(v)) = u \cdot f(c) - u \cdot f(v) \\ &\leq u \cdot f(c_u) - u \cdot f(v) \leq \sum_{i=1}^d |f_i(c_u) - f_i(v)| = \|c_u - v\|_{\ell_1}. \end{aligned}$$

³Note that, defined like this, there are some ties in the distances among the candidates that are not ranked last, but these can be easily broken by slightly moving the positions, for instance moving c_{2i-1} (resp. c_{2i}) by $-\epsilon_i$ (resp. $+\epsilon_i$) on the i^{th} coordinate, with $\epsilon_i \neq \epsilon_j$ for $i \neq j$.




Figure 4: A (boundary) hypersurface separating c_1 and c_2 : $\Delta x \neq \Delta y$, $\Delta x, \Delta y > 0$.

As the two distances must be different (no tie in the preferences), c is not ranked last by v .

To show the tightness, let us consider the following profile on 2^d candidates and 2^d voters. For each vector $u \in \{-1, 1\}^d$, we define a candidate c_u with $f(c_u) = u$, and a voter v_u with $f(v_u) = -u$. Then we have $\|f(c_u) - f(v_u)\|_{\ell_1} = 2d$, while if $u \neq u'$ we have $\|f(c_{u'}) - f(v_u)\|_{\ell_1} \leq 2(d-1)$ (as $f_i(c_{u'}) = f_i(v_u)$ on at least one coordinate i). So c_u is ranked last by v_u ⁴. \square

From now on, and throughout the remainder of the article, we fix $d=2$, i.e., we consider a representation of the preferences in the plane. For the sake of brevity, we omit to mention “in \mathbb{R}^2 ” in the following. Given Observation 1, we can use ℓ_1 or ℓ_∞ indifferently. We choose to use ℓ_1 .

4 Properties of hypersurfaces under ℓ_1 in the plane

We give in this section some properties of (boundary) hypersurfaces under ℓ_1 . These properties will be useful to obtain the results on ℓ_1 -Euclidean profiles in Sections 5 and 6. They have also their own interest, giving some geometric insights on the differences between Euclidean profiles under ℓ_1 and ℓ_2 .

Note. For ease of notation, when the position of candidates are fixed, c_i will denote both the candidate and her position in \mathbb{R}^2 (i.e., $f(c_i)$ in the above notation).

4.1 Types of hypersurfaces

We first focus on the description of the hypersurfaces separating two points c_1 and c_2 . In contrast to the ℓ_2 metrics where the hypersurface is always a straight line (if $d=2$), the shape of this hypersurface depends on the relative positions of c_1 and c_2 when using the ℓ_1 metrics, as we will now show. We denote by (x_1, y_1) (resp. (x_2, y_2)) the coordinates of c_1 (resp. c_2), and we use the notations $\Delta x = |x_1 - x_2|$ and $\Delta y = |y_1 - y_2|$.

1. Let us first consider the case $\Delta x \neq \Delta y$, with $\Delta x > 0$ and $\Delta y > 0$. This case is illustrated in Figure 4. Without loss of generality, assume that $\Delta x > \Delta y$ (the case $\Delta x < \Delta y$ can be treated analogously). The positions c_1 and c_2 can be seen as two opposite vertices of a rectangle (see Fig. 4).

- By definition of the ℓ_1 metrics, there are two points M_1, M_2 on the rectangle boundary that belong to $H(c_1, c_2)$: these are the points that are at distance $\frac{\Delta x + \Delta y}{2}$ from both points c_1 and c_2 . Points M_1 and M_2 are symmetric with respect to the rectangle center, and we observe that the segment $[M_1, M_2]$ belongs to $H(c_1, c_2)$ - in fact, we observe that:

$$[M_1, M_2] = \mathcal{S}_{\frac{\Delta x + \Delta y}{2}}^{\ell_1}(c_1) \cap \mathcal{S}_{\frac{\Delta x + \Delta y}{2}}^{\ell_1}(c_2)$$

⁴As previously, the ties between distances among candidates that are not ranked last can be removed by slightly moving the positions.




Figure 5: The hypersurface separating c_1 and c_2 if $\Delta x = \Delta y > 0$.

- The half-line $\{(x_{M_1}, y) : y \geq y_{M_1}\}$ also belongs to $H(c_1, c_2)$, where x_{M_1} and y_{M_1} denote the coordinates of M_1 , as for $y \geq y_{M_1}$, each point (x_{M_1}, y) is at distance $\frac{\Delta x + \Delta y}{2} + y - y_{M_1}$ both from c_1 and c_2 .
- Similarly, the half-line $\{(x_{M_2}, y) | y \leq y_{M_2}\}$ belongs to $H(c_1, c_2)$.

To sum it up, we have identified three parts of $H(c_1, c_2)$: two vertical half-lines connected by a diagonal segment. We can easily prove that for each $z \in \mathbb{R}^2$ that does not belong to one of these parts, we have $\|z - c_1\|_{\ell_1} \neq \|z - c_2\|_{\ell_1}$. More precisely, the points to the left-hand side of the hypersurface are closer to c_1 , while the ones on the right-hand side are closer to c_2 .

2. Let us now consider the case $\Delta x = \Delta y > 0$. This case is illustrated in Figure 5. In this special case, the rectangle is a square where c_1 and c_2 are opposite vertices, and M_1 and M_2 are the two other opposite vertices. The hypersurface $H(c_1, c_2)$ is then composed of the three following parts:

- (a) the quadrant $\{(x, y) \in \mathbb{R}^2 | x \leq x_{M_1}, y \geq y_{M_1}\}$,
- (b) the segment $[M_1, M_2]$,
- (c) the quadrant $\{(x, y) \in \mathbb{R}^2 | x \geq x_{M_1}, y \leq y_{M_1}\}$.

3. Consider now the case $\Delta x = 0$ or $\Delta y = 0$. Clearly, the hypersurface is then the same as for the ℓ_2 metrics (i.e., a straight line at equal ℓ_2 distance from c_1 and c_2).

The following result states that, to recognise an ℓ_1 -Euclidean profile, we can assume without loss of generality that all hypersurfaces are of the first type described above:

Lemma 1. *Let \mathcal{P} be an ℓ_1 -Euclidean profile. There exists a representation of \mathcal{P} in which all hypersurfaces are of type 1, i.e., $\Delta x \neq \Delta y$ and $\Delta x, \Delta y > 0$.*

Proof. In an ℓ_1 -Euclidean representation of a preference profile, as we consider only strict preferences, we have for any candidates c_i, c_j and voter v :

$$|\|f(v) - f(c_i)\|_{\ell_1} - \|f(v) - f(c_j)\|_{\ell_1}| > 0 \quad (2)$$

Then, let us denote by ε_d the minimum difference in absolute value of distances as in (2), over all pairs $\{c_i, c_j\}$ of candidates and voters v . Moreover, let (x_i, y_i) be the position of candidate c_i in the representation, and S_x (resp. S_y, S_{xy}) the set of pairs of candidates $\{c_i, c_j\}$ with $|x_i - x_j| > 0$ (resp. $|y_i - y_j| > 0$, $||x_i - x_j| - |y_i - y_j|| > 0$). We define also:

$$\begin{aligned} \varepsilon_x &= \min_{\{c_i, c_j\} \in S_x} |x_i - x_j|, \\ \varepsilon_y &= \min_{\{c_i, c_j\} \in S_y} |y_i - y_j|, \\ \text{and } \varepsilon_{xy} &= \min_{\{c_i, c_j\} \in S_{xy}} ||x_i - x_j| - |y_i - y_j||. \end{aligned}$$




Figure 6: The different ℓ_1 -hypersurfaces for $x_1 < x_2$.

If there is a pair $\{c_i, c_j\}$ such that $x_i = x_j$ (in other words, $\{c_i, c_j\} \notin S_x$), we can move one of these candidates, say c_i , by adding ε to x_i with $\varepsilon = \frac{1}{2} \min\{\varepsilon_d, \varepsilon_x, \varepsilon_y, \varepsilon_{xy}\}$. We then get $|x_i - x_j| > 0$. We note that after this operation, we have $S_x \leftarrow S_x \cup \{\{c_i, c_j\}\}$ and S_y and S_{xy} are not modified. An analogous reasoning can be done for every pair $\{c_i, c_j\}$ of candidates such that $\{c_i, c_j\} \notin S_y$ (by moving one candidate on the y -axis), resp. $\{c_i, c_j\} \notin S_{xy}$ (by moving one candidate on one axis). This way, by iterating these modifications, we finally get a representation without the degenerated cases $\Delta x = \Delta y$, $\Delta x = 0$, or $\Delta y = 0$. \square

Hence, without loss of generality, we assume that all ℓ_1 -hypersurfaces are of type 1 in the following.

We can go further into the classification of the different hypersurfaces of type 1. First, notice that if $\Delta x < \Delta y$, both half-line parts of the hypersurface are horizontal. In the opposite case, when $\Delta x > \Delta y$, these half-lines are vertical.

Now, let us look at the segment $[M_1, M_2]$ of the hypersurface. In the following, the numbering of the quadrants of the Cartesian coordinate system goes counter-clockwise starting from the upper right quadrant. Without loss of generality, assume that $x_1 < x_2$, where $c_1 = (x_1, y_1)$ and $c_2 = (x_2, y_2)$. If $y_1 < y_2$, the segment $[M_1, M_2]$ is parallel to the II-IV quadrant diagonal, also called the “minus diagonal” (see the upper part of Figure 6). If $y_1 > y_2$, the segment $[M_1, M_2]$ is parallel to the I-III quadrant diagonal, also called the “plus diagonal” (see the lower part of Figure 6).

Now that we have seen the shape of hypersurfaces for ℓ_1 , we illustrate this by giving a representation of the profile on 3 candidates that includes all $3! = 6$ possible strict preferences over the 3 candidates (*complete profile*).

Example 3 (Example 1 continued). *Figure 7 shows that the complete profile on 3 candidates (with 6 preferences) is ℓ_1 -Euclidean, by providing a ℓ_1 -Euclidean representation of the profile.*

More interestingly, we show that the profile on 4 candidates given in Example 2 is ℓ_1 -Euclidean (while it is not ℓ_2 -Euclidean).

Example 4 (Example 2 continued). *Figure 8 shows that profile \mathcal{P} of Example 2 is ℓ_1 -Euclidean in \mathbb{R}^2 , by providing a ℓ_1 -Euclidean representation of the profile.*




Figure 7: A ℓ_1 -Euclidean representation of the complete profile on 3 candidates.




Figure 8: A ℓ_1 -Euclidean representation of the profile \mathcal{P} of Example 2.

4.2 Intersection of boundary hypersurfaces

It will come as no surprise that many geometrical properties holding for ℓ_2 do not hold for ℓ_1 -hypersurfaces. Let us mention some of them that are useful for the rest of this paper.

It is well-known that given two distinct lines (i.e., ℓ_2 -hypersurfaces), the intersections of these lines is either empty (if the lines are parallel) or contains a unique point. In the case of ℓ_1 -hypersurfaces, more situations may arise, as stated in the following proposition (several examples of possible intersections are given in Figure 9 for illustration. The proof, as well as complete figures illustrating the different situations, are given in Appendix A.1).

Proposition 3. *The intersection of two distinct ℓ_1 -hypersurfaces is either empty or contains a unique point, or two distinct points, or an infinite number of points.*

The following result states that, to recognise a ℓ_1 -Euclidean profile, we can assume without loss of generality that the last case of Proposition 3 (corresponding to a degenerate case) never occurs. Thus, in the remainder of the article, we assume w.l.o.g. that hypersurfaces intersect in at most 2 points.

Lemma 2. *Let \mathcal{P} be an ℓ_1 -Euclidean profile. There exists a representation of \mathcal{P} in which any pair of hypersurfaces intersect in at most 2 points.*

Proof. The proof is similar to that of Lemma 1. Let us suppose that for a given ℓ_1 -Euclidean representation f , there are two hypersurfaces $H(c_i, c_j)$ and $H(c_k, c_l)$ that intersect in infinitely many points. We will show




Figure 9: The intersection of two distinct ℓ_1 -hypersurfaces : several examples

that it is always possible to slightly change the position of one of the points $f(c_i)$, $f(c_j)$, $f(c_k)$ and $f(c_l)$ so that the hypersurfaces intersect in at most two points, without modifying the types of hypersurfaces and the set of representation areas.

Suppose that one of the hypersurfaces, say $H(c_i, c_j)$, is vertical (the other case being symmetrical). We move the point $f(c_i)$. To do so, we denote

$$\varepsilon_t = \min_{c_k \in C \setminus \{c_i\}} \left| |x_i - x_k| - |y_i - y_k| \right|$$

and, as in the Lemma 1,

$$\varepsilon_d = \min_{v \in V} \min_{c_i, c_j \in C} \left| \|f(v) - f(c_i)\|_{\ell_1} - \|f(v) - f(c_j)\|_{\ell_1} \right|.$$

As we consider only strict preferences, $\varepsilon_d > 0$. Also, thanks to Lemma 1 that excluded a degeneration $\Delta x = \Delta y$, we have $\varepsilon_t > 0$. Let $\varepsilon = \frac{1}{2} \min\{\varepsilon_d, \varepsilon_t\}$.⁵ We can now move the point $f(c_i)$ by adding ε to x_i . As $\varepsilon < \varepsilon_d$, we do not change the set of preferences corresponding to representation areas. As $\varepsilon < \varepsilon_t$, we do not change the type of any hypersurface involving c_i . Finally, as x_i increased by $\varepsilon > 0$, the value of $|x_i - x_j| + |y_i - y_j|$ changes, and the hypersurface (both the vertical extremities and the middle segment) slightly moves to the right on the x -axis. Therefore, $H(c_i, c_j)$ and $H(c_k, c_l)$ no more intersect in an infinity of points. \square

In Euclidean geometry under norm ℓ_2 , the bisectors of the three sides of a non-degenerate triangle intersect in a unique point, and do not intersect otherwise. In terms of hypersurfaces, given three points c_1 , c_2 and c_3 , the hypersurfaces $H(c_1, c_2)$, $H(c_1, c_3)$ and $H(c_2, c_3)$ intersect in at most one point under ℓ_2 . We have the following analogous result in case of ℓ_1 -hypersurfaces:

Proposition 4. *Given three points c_1 , c_2 and c_3 :*

- *If $H(c_1, c_2)$, $H(c_1, c_3)$ and $H(c_2, c_3)$ are all vertical (or all horizontal), then the intersection of each pair of hypersurfaces is empty. In particular, the intersection of the 3 hypersurfaces is empty.*
- *If two of them are vertical and one is horizontal (or vice-versa), then the intersection of the 3 hypersurfaces is a unique point.*

The proof of this Proposition (see Appendix A.2) uses the following easy lemma, that we will use in some other proofs as well.

Lemma 3. *Given three points c_1 , c_2 and c_3 , we have:*

$$H(c_1, c_2) \cap H(c_1, c_3) \cap H(c_2, c_3) = H(c_i, c_j) \cap H(c_j, c_k)$$

for all i, j, k such that $\{i, j, k\} = \{1, 2, 3\}$.

⁵For completeness, we should also choose ε smaller than ε_x , ε_y and ε_{xy} introduced in Lemma 1, to ensure that we do not create any degeneration excluded by this Lemma while moving the point $f(c_i)$.




Figure 10: The parallelogram determined by c_i and c_j .

Proof. The left-right inclusion is obvious. For the right-left inclusion, without loss of generality, assume that $i=1, j=2$ and $k=3$, and consider $x \in H(c_i, c_j) \cap H(c_i, c_k)$. Then,

$$\|x - c_1\|_{\ell_1} = \|x - c_2\|_{\ell_1} = \|x - c_3\|_{\ell_1}$$

because

$$\begin{aligned} x \in H(c_i, c_j) &\Rightarrow \|x - c_i\|_{\ell_1} = \|x - c_j\|_{\ell_1}, \\ x \in H(c_i, c_k) &\Rightarrow \|x - c_i\|_{\ell_1} = \|x - c_k\|_{\ell_1}. \end{aligned}$$

Hence, $x \in H(c_1, c_2) \cap H(c_1, c_3) \cap H(c_2, c_3)$. \square

The next result is a direct corollary of Proposition 4 (see Appendix A.3 for the proof).

Corollary 1. *Given three points c_i, c_j, c_k , the hypersurfaces $H(c_i, c_j)$ and $H(c_i, c_k)$ intersect in at most one point. In other words, given four points c_i, c_j, c_k and c_l , if two hypersurfaces $H(c_i, c_j)$ and $H(c_k, c_l)$ intersect in two different points, then c_i, c_j, c_k and c_l are all distinct.*

Proposition 4 can be reformulated by giving conditions on the relative positions of c_1, c_2 and c_3 rather than the types of hypersurfaces $H(c_1, c_2), H(c_1, c_3)$ and $H(c_2, c_3)$. For this reformulation, let us first define the following notion of parallelogram associated with the positions of 2 candidates (see Figure 10 for an illustration).

Definition 3. *Let c_i and c_j be two candidates, and $(x_i, y_i), (x_j, y_j)$ their positions in the two-dimensional plane. Let us denote by:*

- $d_i^+ = \{(x, y) | y = x - x_i + y_i\}$ the "+" diagonal going through the point c_i ,
- $d_i^- = \{(x, y) | y = -x + x_i + y_i\}$ the "-" diagonal going through the point c_i ,
- $d_j^+ = \{(x, y) | y = x - x_j + y_j\}$ the "+" diagonal going through the point c_j ,
- $d_j^- = \{(x, y) | y = -x + x_j + y_j\}$ the "-" diagonal going through the point c_j .

Let us call A the intersection point of d_i^+ and d_j^- and B the intersection point of d_i^- and d_j^+ . We call parallelogram determined by c_i and c_j the parallelogram whose vertices are c_i, A, c_j and B , and we denote by $\text{paral}(c_i, c_j)$ the interior of the parallelogram⁶.

We are now able to reformulate Proposition 4 as Proposition 5 below.

Proposition 5. *Given three points $c_1 = (x_1, y_1), c_2 = (x_2, y_2)$ and $c_3 = (x_3, y_3)$:*

⁶Note that as we consider non degenerated profile following Lemma 1, no point (besides c_i and c_j) lies on one of the 4 diagonals - and in particular on the boarder of the parallelogram.

- If c_1, c_2 or c_3 is inside the parallelogram determined by the two other points, then $H(c_1, c_2)$, $H(c_1, c_3)$ and $H(c_2, c_3)$ do not (pairwise) intersect.
- Otherwise, the intersection of the three hypersurfaces is a unique point.

The proof of Proposition 5 can be found in Appendix A.4. This proposition has a direct consequence on the preferences within an ℓ_1 -Euclidean profile. It is given in the following corollary, which will be used in Section 6 to show an upper bound on the number of candidates ranked last by at least one voter in a ℓ_1 -Euclidean profile.

Corollary 2. *Let $\mathcal{P} = (V, C)$ be an ℓ_1 -Euclidean profile, and consider three candidates $c_1 = (x_1, y_1)$, $c_2 = (x_2, y_2)$ and $c_3 = (x_3, y_3)$ in a given ℓ_1 -Euclidean representation of \mathcal{P} . If $c_2 \in \text{paral}(c_1, c_3)$, then there is no voter $v \in V$ for whom both $c_1 >_v c_2$ and $c_3 >_v c_2$. In other words, c_2 is never ranked last among c_1, c_2, c_3 .*

Proof. Assume that c_2 is inside the parallelogram determined by c_1 and c_3 . Proposition 5 implies that $H(c_1, c_2)$, $H(c_1, c_3)$ and $H(c_2, c_3)$ do not (pairwise) intersect. Hence, they are all horizontal, or all vertical (as a vertical hypersurface always intersects a horizontal one). Without loss of generality, assume that all three hypersurfaces are vertical, and that $x_1 < x_2 < x_3$. As each point (x, y) of $H(c_i, c_j)$ satisfies $x_i < x < x_j$, we have $H(c_1, c_2)$ on the left of $H(c_2, c_3)$.

We now show by contradiction that $H(c_1, c_3)$ lies between these two hypersurfaces. Assume the left-to-right order of hypersurfaces is $H(c_1, c_3), H(c_1, c_2), H(c_2, c_3)$. As c_1 lies in the leftmost area, it is necessarily the top-ranked candidate there. The second-ranked candidate in this area must be c_3 , the leftmost hypersurface being $H(c_1, c_3)$. Thus, the ranking of the leftmost area is $c_1 > c_3 > c_2$. By moving from the leftmost to the rightmost area, we obtain consecutively (by crossing the hypersurfaces one by one) the four following rankings: (c_1, c_3, c_2) (the leftmost one), (c_3, c_1, c_2) (after crossing $H(c_1, c_3)$), (c_3, c_2, c_1) (after crossing $H(c_1, c_2)$) and finally (c_2, c_3, c_1) (the rightmost one, after crossing $H(c_2, c_3)$). We get a contradiction: as c_3 lies in the rightmost area (because we have $x_1 < x_2 < x_3$), it must be a top-ranked candidate there.

The case where $H(c_1, c_3)$ is the rightmost hypersurface can be treated similarly. Hence, the only possible order of hypersurfaces is $H(c_1, c_2), H(c_1, c_3), H(c_2, c_3)$, and we see, with similar arguments as previously, that c_2 is never ranked last. \square

Note that Proposition 3 only gives the possible number of intersection points between two hypersurfaces, however, it does not specify the conditions in which each of the cases appears. The following result (see Appendix A.5 for the proof) gives some more precise statement, which will be needed in the next sections in order to compute, based on geometrical arguments, the size of ℓ_1 -Euclidean profiles.

Proposition 6. *Given four points c_1, c_2, c_3 and c_4 , there is at most one pair of hypersurfaces $H(c_i, c_j), H(c_k, c_l)$ (with $\{i, j, k, l\} = \{1, 2, 3, 4\}$) intersecting in two distinct points.*

5 Euclidean profiles on 4 candidates in the plane

As we have seen, all the profiles with 3 candidates are ℓ_2 -Euclidean and ℓ_1 -Euclidean. We focus here on the case with 4 candidates. In Section 5.1, we study the maximum size of ℓ -Euclidean profiles (for $\ell = \ell_1$ and $\ell = \ell_2$). In Section 5.2, we provide a concise characterization of ℓ_2 -Euclidean profiles.

5.1 Maximum size of a Euclidean profile on 4 candidates

Bennett and Hays (1960) gave a recursive formula to compute the maximum cardinality of ℓ_2 -Euclidean profiles in \mathbb{R}^d . For $d=2$ and 4 candidates, their formula gives the following result:

Proposition 7 (Bennett and Hays, 1960). *The max cardinality of a ℓ_2 -Euclidean profile on 4 candidates is 18.*

We examine this question for the norm ℓ_1 , and show that the maximum cardinality is 19 (Theorem 1). The core of the proof is to show that it is at most 19 (Lemma 4): this is done by counting the (maximal) number of areas delimited by hypersurfaces. For this, we use several results of Section 4, as well as Euler's formula for planar graphs. An explicit construction of a ℓ_1 -Euclidean profile with 19 preferences is then given in Lemma 5, which shows that the upper bound of Lemma 4 is tight.




Figure 11: A ℓ_1 -Euclidean representation of a profile \mathcal{P} , and its corresponding graph. The intersection of $H(c_i, c_j)$, $H(c_i, c_k)$, $H(c_j, c_k)$ (resp. $H(c_i, c_j)$ and $H(c_k, c_l)$) yields a vertex $I_{i,j,k}$ (resp. $I_{i,j} \cap I_{k,l}$).

Lemma 4. Any ℓ_1 -Euclidean profile on 4 candidates has at most 19 (pairwise distinct) preferences.

Proof. To prove this proposition, given an ℓ_1 -Euclidean representation of a profile \mathcal{P} , we define a graph whose vertices are all hypersurface intersections, and where there is an edge between two intersections (denoted by I_1 and I_2) if and only if both I_1 and I_2 lie on the same hypersurface, and there is any other intersection on the segment of extremities I_1 and I_2 (see Figure 11 for an illustration).

The corresponding graph is by construction planar. We note that each inner face of the graph corresponds to a bounded area in the representation of the profile, while unbounded areas in the representation of the preference profile are all merged into the outer face of the planar graph.

We can then use Euler's formula in the corresponding graph. It states that the number of faces of a planar graph is $n_f = n_e - n_v + 2$, where n_e is the number of edges and n_v the number of vertices.

Let us denote by n_z the number of areas in the ℓ_1 -Euclidean representation of the profile. Note that each area corresponds to a single preference, so $n \leq n_z$. For 4 candidates, there are 6 hypersurfaces, leading to at most 12 unbounded areas. As mentioned above, these 12 unbounded areas are merged into the outer face of the planar graph. As the bounded areas yield $n_f - 1$ inner faces, we have $n_z \leq n_f - 1 + 12 = n_f + 11$, and therefore (by Euler's formula):

$$n_z \leq n_e - n_v + 13 \quad (3)$$

If k different hypersurfaces intersect in a common point, we call this point a k -intersection. We can assume there are only 2-intersections and 3-intersections: let f be a representation of a given ℓ_1 -Euclidean profile containing a 4-intersection I . As 3 points give only 3 different hypersurfaces, the 4 hypersurfaces intersecting in I involve the four points $f(c_1), f(c_2), f(c_3), f(c_4)$ corresponding to the positions of the four candidates c_1, c_2, c_3, c_4 . By definition, I is equidistant from all candidates - more formally, we have $\|f(c) - I\|_{\ell_1} = \delta > 0$ for each $c \in \{c_1, c_2, c_3, c_4\}$. As in the Lemma 1, we define:

$$\varepsilon_d = \min_{v \in V} \min_{c_i, c_j \in C} | \|f(v) - f(c_i)\|_{\ell_1} - \|f(v) - f(c_j)\|_{\ell_1} |.$$

As we consider only strict preferences, $\varepsilon_d > 0$. We can then add $\varepsilon = \frac{\varepsilon_d}{2}$ to x_1 .⁷ Doing that, I will be no more equidistant from all four points and therefore, there will no more be a 4-intersection in such a modified representation. By iterating the processus, all k -intersections can be excluded for any $k \geq 4$.

As there are 4 candidates, there are at most four 3-intersections:

- $I_{123} = H(c_1, c_2) \cap H(c_1, c_3) \cap H(c_2, c_3)$,
- $I_{124} = H(c_1, c_2) \cap H(c_1, c_4) \cap H(c_2, c_4)$,
- $I_{134} = H(c_1, c_3) \cap H(c_1, c_4) \cap H(c_3, c_4)$,
- $I_{234} = H(c_2, c_3) \cap H(c_2, c_4) \cap H(c_3, c_4)$.

⁷More precisely, ε should be smaller than the minimum of ε_d and $\min\{\varepsilon_x, \varepsilon_y, \varepsilon_{xy}\}$ as defined in Lemma 1, to ensure that we do not create one of the degenerations excluded by this Lemma.

By Lemma 3, we have covered all intersections of type $H(c_i, c_j) \cap H(c_i, c_k)$. That means, all 2-intersections will be of type $H(c_i, c_j) \cap H(c_k, c_l)$ with i, j, k, l pairwise distinct. There are 3 pairs of hypersurfaces of this type:

- $H(c_1, c_2) \cap H(c_3, c_4)$,
- $H(c_1, c_3) \cap H(c_2, c_4)$,
- $H(c_1, c_4) \cap H(c_2, c_3)$.

Each of these three pairs can give us one 2-intersection. In addition, Proposition 6 implies that at most one of these pairs of hypersurfaces can intersect twice. To sum up, we have at most four 2-intersections. Therefore $n_v \leq 8$ (at most four 3-intersections and four 2-intersections).

If $n_v = 8$, there are four 2-intersections and four 3-intersections. Each 2-intersection generates four outgoing half-lines, and each 3-intersection generates six outgoing half-lines. We then get $4 \cdot 4 + 4 \cdot 6 = 40$ outgoing half-lines. However, 12 of them are delimiting outer non-bounded areas, so they are not responsible for any graph edge. Therefore, $40 - 12 = 28$ half-lines are left for forming edges. We observe that each of these half-lines is used in the creation of exactly one edge, and that each edge is a segment corresponding to the common part of exactly two half-lines (as each edge has two extremities which are two different intersections). Thus, we have $n_e = 28/2 = 14$. Finally, using Equation 3:

$$n_z \leq 14 - 8 + 13 = 19.$$

It is easy to check that if $n_v < 8$, then $n_z < 19$: in fact, each 2-intersection (resp. 3-intersection) generate four (resp. six) outgoing half-lines. In both cases at most a half of them are delimiting outer non-bounded areas - which means that at least half of them has another 2-intersection or 3-intersection lying on it. Therefore, each vertex allows to create at least two edges, so in the Euler formula the benefit of deleting a vertex is outweighed by the drawback of deleting two edges. Thus, in any case, $n_z \leq 19$. The size n of the profile therefore satisfies $n \leq n_z \leq 19$. \square

Let us now consider the following profile \mathcal{P}_0^* with 19 voters and 4 candidates (for more conciseness and readability, preferences are in columns, so for instance the first preference is (c_1, c_2, c_3, c_4)).

$$\mathcal{P}_0^* = \begin{pmatrix} c_1 & c_1 & c_1 & c_1 & c_1 & c_1 & c_2 & c_2 & c_2 & c_2 & c_3 & c_3 & c_3 & c_4 & c_4 & c_4 & c_4 & c_4 & c_4 \\ c_2 & c_2 & c_3 & c_3 & c_4 & c_4 & c_1 & c_1 & c_4 & c_4 & c_1 & c_4 & c_4 & c_1 & c_1 & c_2 & c_2 & c_3 & c_3 \\ c_3 & c_4 & c_2 & c_4 & c_2 & c_3 & c_3 & c_4 & c_1 & c_3 & c_4 & c_1 & c_2 & c_2 & c_3 & c_1 & c_3 & c_1 & c_2 \\ c_4 & c_3 & c_4 & c_2 & c_3 & c_2 & c_4 & c_3 & c_3 & c_1 & c_2 & c_2 & c_1 & c_3 & c_2 & c_3 & c_1 & c_2 & c_1 \end{pmatrix}$$

Lemma 5. \mathcal{P}_0^* is ℓ_1 -Euclidean.

Proof. Figure 12 provides a ℓ_1 -Euclidean representation of \mathcal{P}_0^* . Preference p_1 corresponds to (c_1, c_2, c_3, c_4) (the first column in \mathcal{P}_0^*), preference p_7 to (c_2, c_1, c_3, c_4) (the 7th column in \mathcal{P}_0^*) as we cross $H(c_1, c_2)$ to go from p_1 to p_7 , etc. The representation function $f : C \rightarrow \mathbb{R}^2$ leading to Figure 12 corresponds to the following positions: $f(c_1) = (0, 8)$, $f(c_2) = (10, 10)$, $f(c_3) = (4, 1)$ and $f(c_4) = (8, 3)$. These positions are sufficient to plot the hypersurfaces and to convince ourselves that there are 19 non-empty preference areas. For example, let us place a voter v in the area corresponding to preference p_2 , concretely on the coordinates $(5.5, 8)$. We will check that her preference is indeed p_2 . Denoting by P_v the position of voter v (i.e., the point $(5.5, 8)$), we have

$$\begin{aligned} \|P_v - f(c_1)\|_{\ell_1} &= |5.5 - 0| + |8 - 8| = 5.5, \\ \|P_v - f(c_2)\|_{\ell_1} &= |5.5 - 10| + |8 - 10| = 6.5, \\ \|P_v - f(c_3)\|_{\ell_1} &= |5.5 - 4| + |8 - 1| = 8.5, \\ \|P_v - f(c_4)\|_{\ell_1} &= |5.5 - 8| + |8 - 3| = 7.5. \end{aligned}$$

We see that, indeed, the preference of voter v corresponds to $p_2 = (c_1, c_2, c_4, c_3)$. \square

As a direct consequence of Lemmata 4 and 5, we have the following result, which concludes the section.

Theorem 1. *The maximum cardinality of an ℓ_1 -Euclidean profile on 4 candidates is 19.*

We note that \mathcal{P}_0^* is another example of a preference profile on 4 candidates which is ℓ_1 -Euclidean but not ℓ_2 -Euclidean (because there are more than 18 preferences).




Figure 12: An ℓ_1 -Euclidean representation of a profile with 4 candidates and 19 (pairwise) distinct votes.

5.2 Characterization of ℓ_2 -Euclidean profiles

A central question in structured preferences is to determine whether a given profile is structured or not. As we have seen before, with 4 candidates, any profile with more than 18 (resp. 19) preferences is not ℓ_2 -Euclidean (resp. not ℓ_1 -Euclidean). However, there are smaller profiles which are not ℓ_1 - or ℓ_2 -Euclidean (Example 2 gives such a profile for ℓ_2).

In the sequel, we give a concise description of all ℓ_2 -Euclidean profiles on 4 candidates, that moreover enables to easily determine whether a given profile on 4 candidates is ℓ_2 -Euclidean or not. As noted in the introduction, this result has also been proved by Kamiya et al (2011). The interest of the proof presented below is that it is based on simpler and purely geometrical arguments. Furthermore, it might be possible to consider the same type of approach to prove a result of the same nature for $m \geq 5$ candidates *in the plane*, while the result by Kamiya et al only applies for $d = m - 2$ (thus for 4 candidates in the plane). Also, a similar geometric approach might be useful for the ℓ_1 norm, for which we conjecture that the profile \mathcal{P}_0^* is the *unique* maximal ℓ_1 -Euclidean profile.

Back to ℓ_2 -Euclidean profiles on 4 candidates, we show that the number of maximal Euclidean profiles is very small. More precisely, we prove that there are only 3 maximal ℓ_2 -Euclidean profiles \mathcal{P}_1^* , \mathcal{P}_2^* and \mathcal{P}_3^* (up to a permutation of the candidates), each of them of size 18. Thus, a profile is ℓ_2 -Euclidean if and only if it is a subprofile of \mathcal{P}_1^* , \mathcal{P}_2^* or \mathcal{P}_3^* (up to a permutation of the candidates).

We say that two profiles are *isomorphic* if they contain the same set of preferences up to a permutation of the candidates.

Let us consider the three following profiles \mathcal{P}_1^* , \mathcal{P}_2^* and \mathcal{P}_3^* :

$$\mathcal{P}_1^* = \begin{pmatrix} c_1 & c_1 & c_1 & c_2 & c_2 & c_2 & c_2 & c_2 & c_2 & c_3 & c_3 & c_3 & c_4 & c_4 & c_4 & c_4 & c_4 & c_4 \\ c_2 & c_2 & c_4 & c_1 & c_1 & c_3 & c_3 & c_4 & c_4 & c_2 & c_2 & c_4 & c_1 & c_1 & c_2 & c_2 & c_3 & c_3 \\ c_3 & c_4 & c_2 & c_3 & c_4 & c_1 & c_4 & c_1 & c_3 & c_1 & c_4 & c_2 & c_2 & c_3 & c_1 & c_3 & c_1 & c_2 \\ c_4 & c_3 & c_3 & c_4 & c_3 & c_4 & c_1 & c_3 & c_1 & c_4 & c_1 & c_1 & c_3 & c_2 & c_3 & c_1 & c_2 & c_1 \end{pmatrix},$$

$$\mathcal{P}_2^* = \begin{pmatrix} c_1 & c_1 & c_1 & c_1 & c_1 & c_1 & c_2 & c_2 & c_3 & c_3 & c_3 & c_3 & c_3 & c_3 & c_4 & c_4 & c_4 & c_4 \\ c_2 & c_2 & c_3 & c_3 & c_4 & c_4 & c_1 & c_3 & c_1 & c_1 & c_2 & c_2 & c_4 & c_4 & c_1 & c_1 & c_3 & c_3 \\ c_3 & c_4 & c_2 & c_4 & c_2 & c_3 & c_3 & c_1 & c_2 & c_4 & c_1 & c_4 & c_1 & c_2 & c_2 & c_3 & c_1 & c_2 \\ c_4 & c_3 & c_4 & c_2 & c_3 & c_2 & c_4 & c_4 & c_4 & c_2 & c_4 & c_1 & c_2 & c_1 & c_3 & c_2 & c_2 & c_1 \end{pmatrix},$$

$$\mathcal{P}_3^* = \begin{pmatrix} c_1 & c_1 & c_1 & c_1 & c_2 & c_2 & c_2 & c_2 & c_2 & c_2 & c_3 & c_3 & c_3 & c_3 & c_4 & c_4 & c_4 & c_4 \\ c_2 & c_2 & c_3 & c_4 & c_1 & c_1 & c_3 & c_3 & c_4 & c_4 & c_1 & c_2 & c_2 & c_4 & c_1 & c_2 & c_2 & c_3 \\ c_3 & c_4 & c_2 & c_2 & c_3 & c_4 & c_1 & c_4 & c_1 & c_3 & c_2 & c_1 & c_4 & c_2 & c_2 & c_1 & c_3 & c_2 \\ c_4 & c_3 & c_4 & c_3 & c_4 & c_3 & c_4 & c_1 & c_3 & c_1 & c_4 & c_4 & c_1 & c_1 & c_3 & c_3 & c_1 & c_1 \end{pmatrix}.$$

Theorem 2. A profile on 4 candidates is ℓ_2 -Euclidean if and only if it is isomorphic to a subprofile of \mathcal{P}_1^* , \mathcal{P}_2^* or \mathcal{P}_3^* .




Figure 13: The possible relative positions of the 3-intersections in the plane.

Proof. For any maximal profile there is a representation of it such that no pair of hypersurfaces (which are simple lines of the plane in the present case) are parallel. In fact, there will be two parallel lines if there are (at least) three aligned candidates, or if two pairs of candidates are the extremities of two parallel segments. In each of these cases, we can always slightly move one of the candidates (using the same technique as in Lemma 1) so that the two concerned lines are no more parallel and such a modified mapping is still a representation of the given profile.

Assuming that, we have:

- One 2-intersection $H(c_i, c_j) \cap H(c_k, c_l)$ for each pair of hypersurfaces with i, j, k, l pairwise distinct. For 4 candidates, it yields three 2-intersections (because there are three such pairs).
- One 3-intersection $I_{i,j,k} = H(c_i, c_j) \cap H(c_i, c_k) \cap H(c_j, c_k)$ for each triple of hypersurfaces with i, j, k pairwise distinct. For a profile on 4 candidates, it yields four 3-intersections (because there are four such triples).

Let us study the relative positions of the 3-intersections in the plane. There are two possible scenarios (see Figure 13):

1. The 3-intersections are the vertices of a convex quadrilateral (left part of Figure 13). No pair of opposite sides of this quadrilateral can be parallel, otherwise there would be two parallel hypersurfaces, and the profile would not be maximal.
2. Three of the 3-intersections are the vertices of a triangle, and the fourth one is inside this triangle (right part of Figure 13).

We will now take a closer look to each of these cases, and we will construct all maximal profiles corresponding to each of them.

Case 1:

Assume that the 3-intersections are the vertices of a convex quadrilateral (see Figure 14), as described above. There are $\binom{4}{2} = 6$ hypersurfaces, each of them goes through exactly two 3-intersections (because, for an hypersurface $H(c_i, c_j)$, there are two ways to choose c_k with $k \notin \{i, j\}$). We recall that, without loss of generality, we can assume that there is no pair of parallel hypersurfaces. There are then four hypersurfaces that form the sides of a convex quadrilateral. Each of the two pairs of hypersurfaces corresponding to opposite sides of the quadrilateral results in an intersection outside the quadrilateral, which yields two distinct 2-intersections. The remaining two hypersurfaces represent the diagonals of the quadrilateral, and will hence intersect inside it - it results in the third (and last) 2-intersection. Whatever the positions of the four candidates, if the 3-intersections form a convex quadrilateral, the partitioning of the plane will always look like in Figure 14a (where $c_1 = (1, 5)$, $c_2 = (4, 2)$, $c_3 = (6, 8)$ and $c_4 = (9, 3)$): one 2-intersection lies inside the convex quadrilateral, and the two remaining 2-intersections (of hypersurfaces forming opposite sides of the quadrilateral) outside of it.

Note that in Figure 14a some areas are small. For readability reasons, in what follows, we use instead Figure 14b (with a similar arrangement of areas) where the areas are larger but without the explicit positions of candidates.

To enumerate all possible maximal profiles corresponding to this configuration of the 3-intersections, the hypersurfaces (and hence the intersections) need to be labeled so we can list the preferences corresponding




Figure 14: Case 1: The plane is divided into 18 areas, with the 3-intersections forming a convex quadrilateral. The candidates are plotted with empty circles, 3-intersections with blue circles and 2-intersections with red squares.




Figure 15: Labeled representation: $H(c_i, c_j)$ is noted as ij due to lack of space.

to the different areas (see Figure 15). Without loss of generality, we label one of the 3-intersections as $I_{1,2,3}$, and one of the hypersurfaces going through it as $H(c_1, c_2)$. The second 3-intersection involving $H(c_1, c_2)$ is then necessarily $I_{1,2,4}$. The two remaining hypersurfaces going through $I_{1,2,3}$ are $H(c_1, c_3)$ and $H(c_2, c_3)$, that we can arbitrarily label (because it will turn out to be symmetrical). From these labels $I_{1,2,3}$, $I_{1,2,4}$, $H(c_1, c_2)$, $H(c_1, c_3)$ and $H(c_2, c_3)$, we can infer the labels of the two remaining 3-intersections, and so the labels of the remaining hypersurfaces. As mentioned earlier, both ways of labeling $H(c_1, c_3)$ and $H(c_2, c_3)$ are symmetric: it is sufficient to rename c_1 as c_2 and c_2 as c_1 to switch from one complete labeling to the other one (see Figure 15). Hence, the labels of $H(c_1, c_3)$ and $H(c_2, c_3)$ can be fixed without loss of generality, and there is only one possible complete labeling, up to a renaming of the candidates.

Once the hypersurfaces are labeled, we can list the preferences associated with the different areas. Let us focus on the areas A_1 , A_2 , A_3 and A_{13} , as well as on the corresponding preferences p_1 , p_2 , p_3 and p_{13} in Figure 16. To switch from p_1 to p_2 , candidate c_3 is swapped with c_4 (because $H(c_3, c_4)$ is crossed between areas A_1 and A_2), while c_4 is swapped with c_2 to switch from p_2 to p_3 (as $H(c_2, c_4)$ is crossed between areas A_2 and A_3), and finally c_4 is swapped with c_1 to obtain p_{13} . Necessarily, c_4 is ranked either in the first or in the last position in p_1 (resp. p_4), as it is successively swapped with all the remaining candidates. Hence, the area A_1 corresponds to one of the following preferences:

- $p_1 = (c_1, c_2, c_3, c_4)$,
- $p'_1 = (c_4, c_3, c_2, c_1)$.

Once at least one preference is known, we can list all the preferences of the profile. Both profiles $\mathcal{P} = \{p_1, \dots, p_{18}\}$ and $\mathcal{P}' = \{p'_1, \dots, p'_{18}\}$ are listed in Table 1. Profile \mathcal{P} corresponds to \mathcal{P}_1^* in the statement of the theorem, while \mathcal{P}' corresponds to \mathcal{P}_2^* .




Figure 16: Listing the different areas A_1, \dots, A_{18} into which the plane is divided in case 1.

Note that, for each $1 \leq i \leq 18$, p_i is the “opposite” of p'_i . Nevertheless, \mathcal{P}' can not be obtained from \mathcal{P} by renaming the candidates: indeed, while in \mathcal{P} candidates c_1 and c_3 are each ranked first 3 times, and c_2 and c_4 ranked first 6 times, in \mathcal{P}' in contrast, we have c_1 and c_3 that are ranked first 6 times, c_2 ranked first 2 times and c_4 ranked first 4 times.

i	p_i	p'_i	i	p_i	p'_i
1	(c_1, c_2, c_3, c_4)	(c_4, c_3, c_2, c_1)	10	(c_3, c_2, c_1, c_4)	(c_4, c_1, c_2, c_3)
2	(c_1, c_2, c_4, c_3)	(c_3, c_4, c_2, c_1)	11	(c_3, c_2, c_4, c_1)	(c_1, c_4, c_2, c_3)
3	(c_1, c_4, c_2, c_3)	(c_3, c_2, c_4, c_1)	12	(c_3, c_4, c_2, c_1)	(c_1, c_2, c_4, c_3)
4	(c_2, c_1, c_3, c_4)	(c_4, c_3, c_1, c_2)	13	(c_4, c_1, c_2, c_3)	(c_3, c_2, c_1, c_4)
5	(c_2, c_1, c_4, c_3)	(c_3, c_4, c_1, c_2)	14	(c_4, c_1, c_3, c_2)	(c_2, c_3, c_1, c_4)
6	(c_2, c_3, c_1, c_4)	(c_4, c_1, c_3, c_2)	15	(c_4, c_2, c_1, c_3)	(c_3, c_1, c_2, c_4)
7	(c_2, c_3, c_4, c_1)	(c_1, c_4, c_3, c_2)	16	(c_4, c_2, c_3, c_1)	(c_1, c_3, c_2, c_4)
8	(c_2, c_4, c_1, c_3)	(c_3, c_1, c_4, c_2)	17	(c_4, c_3, c_1, c_2)	(c_2, c_1, c_3, c_4)
9	(c_2, c_4, c_3, c_1)	(c_1, c_3, c_4, c_2)	18	(c_4, c_3, c_2, c_1)	(c_1, c_2, c_3, c_4)

Table 1: The two maximal profiles $\mathcal{P} = \{p_1, \dots, p_{18}\}$ and $\mathcal{P}' = \{p'_1, \dots, p'_{18}\}$ obtained in case 1.

Case 2:

To begin, let us denote by T the triangle consisting of areas A_5, A_6, A_7, A_8, A_9 and A_{10} . Using the same method as in the previous case, we note that there are two possible rankings for area A_1 (see Figure 17, and the succession of areas A_1, A_2, A_4 and A_{15}):

- $p_1 = (c_1, c_2, c_3, c_4)$,
- $p'_1 = (c_4, c_3, c_2, c_1)$.

However, if $p'_1 = (c_4, c_3, c_2, c_1)$, candidate c_2 is ranked in last position inside the triangle T : in fact, none of the hypersurfaces crossing the triangle involves c_2 . Let us now discuss the position of c_2 to show that p'_1 is not feasible:

- Denoting by $D(c_i, c_j)$ the set of points that are closer to c_i than to c_j , we have:

$$D(c_2, c_1) = A_1 \cup A_2 \cup A_3 \cup A_4 \cup A_{11} \cup A_{15}.$$

In fact, c_1 is preferred to c_2 in triangle T . Therefore, c_1 must lie on the same side of $H(c_1, c_2)$ as this triangle, and c_2 must then lie on the opposite side of $H(c_1, c_2)$, i.e. on the same side as the area A_1 .

- Analogously, candidate c_2 is necessarily on the same side of $H(c_2, c_3)$ as the area A_3 :

$$D(c_2, c_3) = A_3 \cup A_{11} \cup A_{12} \cup A_{13} \cup A_{14} \cup A_{18}.$$

- Finally, candidate c_2 is necessarily on the same side of $H(c_2, c_4)$ as the area A_4 :

$$D(c_2, c_4) = A_4 \cup A_{14} \cup A_{15} \cup A_{16} \cup A_{17} \cup A_{18}.$$




Figure 17: Listing the different areas A_1, \dots, A_{18} into which the plane is divided in case 2.

As $c_2 \in D(c_2, c_i)$ for each $i \in \{1, 3, 4\}$, and as $D(c_2, c_1) \cap D(c_2, c_3) \cap D(c_2, c_4) = \emptyset$, we cannot have $p'_1 = (4, 3, 2, 1)$.

The case $p_1 = (1, 2, 3, 4)$ is feasible, leading to the profile described in Table 2, which corresponds to profile \mathcal{P}_3^* in the statement of the theorem. \square

i	p_i	i	p_i
1	(c_1, c_2, c_3, c_4)	10	(c_2, c_4, c_3, c_1)
2	(c_1, c_2, c_4, c_3)	11	(c_3, c_1, c_2, c_4)
3	(c_1, c_3, c_2, c_4)	12	(c_3, c_2, c_1, c_4)
4	(c_1, c_4, c_2, c_3)	13	(c_3, c_2, c_4, c_1)
5	(c_2, c_1, c_3, c_4)	14	(c_3, c_4, c_2, c_1)
6	(c_2, c_1, c_4, c_3)	15	(c_4, c_1, c_2, c_3)
7	(c_2, c_3, c_1, c_4)	16	(c_4, c_2, c_1, c_3)
8	(c_2, c_3, c_4, c_1)	17	(c_4, c_2, c_3, c_1)
9	(c_2, c_4, c_1, c_3)	18	(c_4, c_3, c_2, c_1)

Table 2: The maximal profile obtained in case 2.

6 Euclidean profiles on $m \geq 5$ candidates in the plane

Let us now focus on the general case, by giving some results on the relative expressive power of ℓ_2 -Euclidean and ℓ_1 -Euclidean preference profiles. We first note that, as shown in Proposition 2, at most 4 candidates are ranked in last position (by at least one voter), regardless of the number of candidates in the profile. This is in sharp contrast to the ℓ_2 -Euclidean case, in which profiles where each candidate is ranked last at least once can easily be built, as mentioned in the introduction.

This property might indicate that being Euclidean is much more restrictive for ℓ_1 than for ℓ_2 . We show however that if we are interested in the maximum size of a Euclidean profile, then there is no such difference. We show indeed that the maximum size of a ℓ_1 -Euclidean profile on m candidates is $\Theta(m^4)$ (Theorem 3), which is the same asymptotic bound as the one found by Bennett and Hays (1960) for ℓ_2 .

Actually, a precise formula can be easily derived from their result: this maximal size is precisely $\frac{m(3m-10)(m-1)(m+1)}{24} + m(m-1) + 1$. While such a precise formula seems to be tricky to establish for ℓ_1 and is left as an open question, we show that the asymptotical bound is the same:

Theorem 3. *The maximum size of an ℓ_1 -Euclidean profile in \mathbb{R}^2 over m candidates is in $\Theta(m^4)$.*

sketch of proof. We first show that the size of such a profile is in $O(m^4)$. There are $\frac{m(m-1)}{2}$ hypersurfaces. With a non-degenerated profile, each pair of hypersurfaces intersects at most twice, hence, there are at most $2(\frac{m(m-1)}{2})^2$ points of intersections. As in the case of norm ℓ_2 , we have at most 3 hypersurfaces intersecting in one point. If a point is at the intersection of 2 (resp 3) hypersurfaces, it is incident to (i.e., a vertex of) 4 areas (resp. 6 areas). Then, as each area has at least one intersection point in its border, the number of areas

is upper bounded by 6 times the number of intersection points, i.e., in $O(m^4)$.

Let us now show a profile for which this bound is reached. The idea is quite straightforward: as there are only vertical or horizontal hypersurfaces, and as each vertical and each horizontal hypersurface intersect, the positions of candidates c_1 to c_m will be iteratively fixed in such a way that approximately half of hypersurfaces are vertical and half are horizontal. The number of intersections will then be in $\Theta(m^4)$, and the construction ensures that the number of areas is in $\Theta(m^4)$.

The explicit construction of this profile is deferred to Appendix B.1. \square

7 Future work

Because of their novelty, multiple avenues of research regarding ℓ_1 -Euclidean preference profiles can be considered. For instance, we conjecture that there is a unique maximal ℓ_1 -Euclidean preference profile for four candidates (and numerical tests seem to confirm this), but it remains to be proved. A broader research question is to investigate the existence of a general formula giving the maximal size of a ℓ_1 -Euclidean preference profile (as there is for ℓ_2). Regarding the computational aspects, Peters (2017) proved that the problem of recognising ℓ_1 -Euclidean preference profiles in \mathbb{R}^d is in NP, but a more specific complexity class remains to be determined, and efficient recognition procedures are still to be proposed.

Although ℓ_2 -Euclidean preferences have been more studied than ℓ_1 -Euclidean preferences, various works can also be considered following those presented here, among which the identification of the maximal ℓ_2 -Euclidean preference profiles in \mathbb{R}^2 for $m \geq 5$, or a thorough study of ℓ_2 -Euclidean preference profiles in \mathbb{R}^3 .

References

- Arrow KJ (1951) Social choice and individual values. John Wiley & Sons
- Barberà S, Berga D, Moreno B (2013) Some new domain restrictions in social choice, and their consequences. In: Torra V, Narukawa Y, Navarro-Arribas G, Megías D (eds) Modeling Decisions for Artificial Intelligence, Springer Berlin Heidelberg, Berlin, Heidelberg, pp 11–24
- Barberà S, Berga D, Moreno B (2020) Arrow on domain conditions: a fruitful road to travel. Social Choice and Welfare 54(2):237–258
- Bennett JF, Hays WL (1960) Multidimensional unfolding: Determining the dimensionality of ranked preference data. Psychometrika 25(1):27–43
- Black D (1948) On the rationale of group decision-making. The Journal of Political Economy 56(1):23–34
- Bogomolnaia A, Laslier JF (2004) Euclidean preferences. HAL, Working Papers
- Chen J, Grottko S (2021) Small one-dimensional Euclidean preference profiles. Social Choice and Welfare pp 1–28
- Chen J, Pruhs K, Woeginger GJ (2015) The one-dimensional Euclidean domain: Finitely many obstructions are not enough. arXiv:1506.03838
- Chen J, Nöllenburg M, Simola S, Villedieu A, Wallinger M (2022) Multidimensional Manhattan preferences. arXiv:2201.09691
- Doignon JP, Falmagne JC (1994) A polynomial time algorithm for unidimensional unfolding representations. Journal of Algorithms 16(2):218–233
- Downs A (1957) An economic theory of democracy. Harper & Row New York
- Elkind E, Faliszewski P (2014) Recognizing 1-Euclidean preferences: An alternative approach. In: Proceedings of the 7th International Symposium on Algorithmic Game Theory (SAGT), pp 146–157
- Elkind E, Lackner M, Peters D (2022) Preference restrictions in computational social choice: A survey. arXiv:2205.09092

- Gaertner W (2001) Domain conditions in social choice theory. Cambridge University Press
- Gaertner W (2002) Domain restrictions. *Handbook of social choice and welfare* 1:131–170
- Garzia D, Marschall S (2019) Voting advice applications. Oxford University Press
- Good I, Tideman T (1977) Stirling numbers and a geometric structure from voting theory. *Journal of Combinatorial Theory, Series A* 23(1):34–45
- Hays WL, Bennett JF (1961) Multidimensional unfolding: Determining configuration from complete rank order preference data. *Psychometrika* 26(2):221–238
- Hotelling H (1929) Stability in competition. *Economic Journal* 39:41–57
- Isotalo V (2020) Designing voting advice applications: The finnish case. Master’s thesis, URL <http://urn.fi/URN:NBN:fi:aalto-2020112918207>
- Kamiya H, Takemura A, Terao H (2011) Ranking patterns of unfolding models of codimension one. *Advances in Applied Mathematics* 47(2):379–400
- Knoblauch V (2010) Recognizing one-dimensional Euclidean preference profiles. *Journal of Mathematical Economics* 46:1–5
- Lee DT, Wong CK (1980) Voronoi diagrams in $l_1(l_\infty)$ metrics with 2-dimensional storage applications. *SIAM Journal on Computing* 9(1):200–211
- Moreno G, Padilla J, Chueca E (2022) Learning VAA: A new method for matching users to parties in voting advice applications. *Journal of Elections, Public Opinion and Parties* 32(2):339–357
- Peters D (2017) Recognising multidimensional Euclidean preferences. In: *Proceedings of the 31st AAAI Conference on Artificial Intelligence (AAAI)*, pp 642–648
- Stanley RP, et al (2004) An introduction to hyperplane arrangements. *Geometric combinatorics* 13(389-496):24

A Missing proofs of Section 4

A.1 Proof of Proposition 3

Proposition 3. *The intersection of two distinct ℓ_1 -hypersurfaces is either empty or contains a unique point, two distinct points or an infinite number of points.*

Proof. We can assume, without loss of generality, that the hypersurfaces are given by two distinct pairs of points. Let us denote by $c_i = (x_i, y_i)$, $i \in \{1, 2, 3, 4\}$ these points and their coordinates. Still without loss of generality, let $H(c_1, c_2)$ be of type V^- . There are four basic cases to distinguish (see Figs 18–21 for illustrations):

1. $H(c_1, c_2)$ is of type V^- and $H(c_3, c_4)$ of type H^+ (see Figure 18):
In this case, the hypersurfaces intersect in a unique point as the half-lines (resp. the middle segments) of $H(c_1, c_2)$ and $H(c_3, c_4)$ are of opposite orientations.
2. $H(c_1, c_2)$ is of type V^- and $H(c_3, c_4)$ of type H^- (see Figure 19):
As in the previous case, there will be at least one intersection as a horizontal hypersurface and a vertical hypersurface always intersect. Contrary to the above, the middle segments of both hypersurfaces have the same orientation, so they can overlap: in such a case, the intersection contains this overlapping segment, thus an infinite number of points.
3. $H(c_1, c_2)$ is of type V^- and $H(c_3, c_4)$ of type V^+ (see Figure 20):
In this case, the hypersurfaces may not intersect: let us assume that $\max\{x_1, x_2\} < \min\{x_3, x_4\}$. By definition, we have $x \in [\min\{x_i, x_j\}, \max\{x_i, x_j\}]$ for each $(x, y) \in H(c_i, c_j)$. The above inequality then implies that the intersection of $H(c_1, c_2)$ and $H(c_3, c_4)$ is empty (graphically, $H(c_1, c_2)$ will be “on the left” of $H(c_3, c_4)$ - see the first case of Figure 20).
The hypersurfaces may also intersect in a unique point: a middle segment of one of the hypersurfaces can intersect one of the half-lines of the second hypersurface, or its middle segment, as the middle segments are not of the same type (see the second case of Figure 20).
Finally, as the half-lines of both hypersurfaces are of the same type, one of the half-lines of $H(c_1, c_2)$ may (partially) overlap one of the half-lines of $H(c_3, c_4)$ (see the third case of Figure 20). In this case, the intersection will contain an infinity of points.
4. Both hypersurfaces $H(c_1, c_2)$ and $H(c_3, c_4)$ are of type V^- (see Figure 21):
This is the most complex case. For the same reason as above, the hypersurfaces may not intersect. They may also intersect in a unique point if the middle segment of one hypersurface intersects one of the half-lines of the second one (see the first case of the Figure 21). As the types of half-lines and middle segments are both the same for $H(c_1, c_2)$ and $H(c_3, c_4)$, they can also intersect in two distinct points if the middle segment of $H(c_1, c_2)$ intersects one of the half-lines of $H(c_3, c_4)$ and the middle segment of $H(c_3, c_4)$ intersects one of the half-lines of $H(c_1, c_2)$ (see the second case of Figure 21). Finally, the intersection can contain an infinity of points: as the half-lines are of the same type, a half-line of $H(c_1, c_2)$ may (partially) overlap a half-line of $H(c_3, c_4)$. In addition, the middle segments being also of the same type, they can (partially) overlap. See cases 3 and 4 of Figure 21 (the case when both the half-lines and the middle segments overlap is not presented in Figure 21, but it is obviously possible).

□




Figure 18: Intersection of two hypersurfaces: $H(c_1, c_2)$ is of type V^- and $H(c_3, c_4)$ of type H^+ .




Figure 19: Intersection of two hypersurfaces: $H(c_1, c_2)$ is of type V^- and $H(c_3, c_4)$ of type H^- .




Figure 20: Intersection of two hypersurfaces: $H(c_1, c_2)$ is of type V^- and $H(c_3, c_4)$ of type V^+ .

A.2 Proof of Proposition 4

Proposition 4. *Given three points c_1, c_2 and c_3 :*

- *If $H(c_1, c_2)$, $H(c_1, c_3)$ and $H(c_2, c_3)$ are all vertical (or all horizontal), then the intersection of each pair of hypersurfaces is empty. In particular, the intersection of the 3 hypersurfaces is empty.*
- *If two of them are vertical and one is horizontal (or vice-versa), then the intersection of the 3 hypersurfaces is a unique point.*

Proof. Assume first that the three hypersurfaces are vertical. Let (x_1, y_1) , (x_2, y_2) and (x_3, y_3) denote the positions in the plane of c_1, c_2 and c_3 . Without loss of generality, we assume that $x_1 < x_2 < x_3$ and that $H(c_1, c_3)$ is of type V^- .

Given a vertical hypersurface $H(c_i, c_j)$, for each point $(x, y) \in H(c_i, c_j)$ we have $x_i < x < x_j$, because $x_i < x_{M_1} < x_j$ and $x_i < x_{M_2} < x_j$ for the extremities M_1, M_2 of the middle segment of the hypersurface (see Figure 4, page 7). Thus, $H(c_1, c_2)$ and $H(c_2, c_3)$ do not intersect, as we have $x_1 < x_2 < x_3$. Using Lemma 3 (page 11), we conclude that $H(c_1, c_2) \cap H(c_1, c_3) \cap H(c_2, c_3) = \emptyset$.

Let us now assume that two hypersurfaces are vertical (resp. horizontal) and the third one is horizontal (resp. vertical). Without loss of generality, we can assume that $H(c_1, c_2)$ is horizontal and both remaining




Figure 21: Intersection of two hypersurfaces: both hypersurfaces $H(c_1, c_2)$ and $H(c_3, c_4)$ are of type V^- .

hypersurfaces $H(c_1, c_3)$ and $H(c_2, c_3)$ are vertical. Any vertical hypersurface intersects any horizontal hypersurface in a unique point (by assuming w.l.o.g. that the representation is non-degenerate, see Lemma 2). In particular, $|H(c_1, c_2) \cap H(c_1, c_3)| = 1$. Lemma 3 states that $H(c_1, c_2) \cap H(c_1, c_3) \cap H(c_2, c_3) = H(c_i, c_j) \cap H(c_i, c_k)$ for $\{i, j, k\} = \{1, 2, 3\}$. We have therefore the three hypersurfaces intersecting in a unique point. \square

A.3 Proof of Corollary 1

Corollary 1. *Given three points c_i, c_j, c_k , the hypersurfaces $H(c_i, c_j)$ and $H(c_i, c_k)$ intersect in at most one point. In other words, if two hypersurfaces $H(c_i, c_j)$ and $H(c_k, c_l)$ intersect in two different points, then c_i, c_j, c_k and c_l are all distinct.*

Proof. By contradiction, assume that $H(c_i, c_j)$ and $H(c_i, c_k)$ intersect in two distinct points. According to Lemma 3 (page 11), $|H(c_i, c_j) \cap H(c_i, c_k) \cap H(c_j, c_k)| \geq 2$. We get a contradiction with Proposition 4 which states that if the three hypersurfaces intersect, then the point of intersection is unique. \square

A.4 Proof of Proposition 5

Proposition 5 *Given three points $c_1 = (x_1, y_1)$, $c_2 = (x_2, y_2)$ and $c_3 = (x_3, y_3)$:*

- *If one of these 3 points is inside the parallelogram determined by the two other points, then the three hypersurfaces $H(c_1, c_2)$, $H(c_1, c_3)$ and $H(c_2, c_3)$ do not (pairwise) intersect.*
- *Otherwise, the intersection of the three hypersurfaces is a unique point.*

Proof. This proposition is a direct consequence of Proposition 4. To prove the first point, we assume without loss of generality that c_2 lies inside the parallelogram determined by c_1 and c_3 , and we prove that in this case, the three hypersurfaces are all vertical or horizontal - Proposition 4 then implies that they do not (pairwise) intersect.

Up to exchanging the roles of c_1 and c_3 , we can assume, still without loss of generality, that $x_1 < x_3$. There are then 4 cases to distinguish (see Figure 22):

- (a) $H(c_1, c_3)$ is of type V^- (see Figure 22a).

In this case, we have $y_1 < y_3$ (see the classification of hypersurfaces given in Figure 6). Moreover, we have $x_1 < x_2 < x_3$. In the parallelogram given in Figure 22a, c_2 lies then above the diagonal d_1^- (i.e., $y_2 > -x_2 + x_1 + y_1$) and below the diagonal d_1^+ (i.e., $y_2 < x_2 - x_1 + y_1$). Put together, we get

$$-x_2 + x_1 < y_2 - y_1 < x_2 - x_1.$$

In other words, $|y_1 - y_2| < x_2 - x_1$, hence $H(c_1, c_2)$ is vertical. We show similarly that $H(c_2, c_3)$ is vertical, as c_2 lies above the diagonal d_3^+ and below the diagonal d_3^- . All three hypersurfaces being vertical, they do not (pairwise) intersect.

- (b) Let us now suppose that $H(c_1, c_3)$ is of type H^- (see Figure 22b).

We have $y_1 < y_2 < y_3$. As c_2 lies above diagonals d_1^- and d_1^+ , we have $y_2 > -x_2 + x_1 + y_1$ and $y_2 > x_2 - x_1 + y_1$. Put together, we have $y_2 - y_1 > x_1 - x_2$ and $y_2 - y_1 > -(x_1 - x_2)$ - in other words, $y_2 - y_1 > |x_1 - x_2|$. Therefore, the hypersurface $H(c_1, c_2)$ is horizontal. We show similarly that $H(c_2, c_3)$ is horizontal, as c_2 lies below diagonals d_3^+ and d_3^- , so we obtain $y_3 - y_2 > |x_2 - x_3|$.

- (c) We suppose here that $H(c_1, c_3)$ is of type V^+ . We have $y_1 > y_3$ and $x_1 < x_2 < x_3$. Analogously to the previous case, we show that $x_2 - x_1 > |y_2 - y_1|$, so $H(c_1, c_2)$ is vertical, and that $x_3 - x_2 > |y_2 - y_3|$, which implies that $H(c_2, c_3)$ is also vertical.

- (d) Finally, we consider $H(c_1, c_3)$ of type H^+ . We have $y_1 > y_2 > y_3$. As in previous cases, we show that $H(c_1, c_2)$ is horizontal as c_2 lies below diagonals d_1^+ and d_1^- , and $H(c_2, c_3)$ is also horizontal as c_2 lies above diagonals d_3^+ and d_3^- .

To prove the second point of the proposition, we suppose that any point does not lie in the parallelogram determined by the remaining two points, and we will show that in such a case, there is at least one horizontal and one vertical hypersurface. As a vertical hypersurface and a horizontal hypersurface intersect in a unique point, Lemma 3 allows us to conclude that the three hypersurfaces intersect in a unique point.

Suppose first that $H(c_1, c_3)$ is vertical (see Figure 23a). The diagonals d_1^+ , d_1^- , d_3^+ and d_3^- divide the plane into 9 areas A_1, A_2, \dots, A_9 . The point c_2 does not lie in A_1 (resp. A_5, A_9) because c_1 (resp. c_2, c_3) does not lie in the parallelogram determined by the remaining two points. If c_2 lies in area A_2, A_3, A_4 or A_7 , the hypersurface $H(c_1, c_2)$ is horizontal, so it intersects the vertical hypersurface $H(c_1, c_3)$. If c_2 lies in one of the remaining areas A_6 or A_8 , the hypersurface $H(c_2, c_3)$ is horizontal, so it intersects $H(c_1, c_3)$. We note that whether the oblique middle-segment of the hypersurface $H(c_1, c_3)$ is ascending (case (c) of Figure 22) or descending (case (a)) has no impact on this reasoning and it can therefore be used without change for both cases (a) and (c) of Figure 22.

Analogously, we treat the case in which $H(c_1, c_3)$ is horizontal: c_2 cannot lie in areas A_3, A_5 and A_7 as any point does not lie within the parallelogram determined by the remaining two points. If c_2 lies in A_1, A_2, A_6 or A_9 , the hypersurface $H(c_1, c_2)$ is vertical. If it lies in one of the two remaining areas A_4 or A_8 , the hypersurface $H(c_2, c_3)$ is vertical. To sum up, there is always at least one horizontal and one vertical hypersurface.

□




Figure 22: The parallelogram determined by c_1 and c_3 with $x_1 < x_3$: 4 cases to distinguish.

A.5 Proof of Proposition 6

Proposition 6. *Given four points c_1, c_2, c_3 and c_4 , there is at most one pair of hypersurfaces $H(c_i, c_j), H(c_k, c_l)$ (with $\{i, j, k, l\} = \{1, 2, 3, 4\}$) intersecting in two distinct points.*

Proof. For $i \in \{1, 2, 3, 4\}$, we denote by (x_i, y_i) the position of candidate c_i in the plane. Assume there are two pairs of hypersurfaces intersecting in two distinct points. Thanks to Corollary 1, we can assume, w.l.o.g.,




Figure 23: The parallelogram determined by c_1 and c_3 with $x_1 < x_3$, and the possible placements of c_2 .

that the first pair involves the hypersurfaces $H(c_1, c_2)$ and $H(c_3, c_4)$. Moreover, still w.l.o.g, we can assume that they are both of type V^- , and that $H(c_1, c_2)$ is “on the left” of $H(c_3, c_4)$ (as in Figure 24), and that $x_1 < x_2$ and $x_3 < x_4$. According to the classification of hypersurfaces (see Figure 6, page 9), as $H(c_1, c_2)$ and $H(c_3, c_4)$ are of type V^- , we have $y_1 < y_2$ and $y_3 < y_4$. Note that we necessarily have:

$$\{x_1, x_3\} < \{x_2, x_4\} \quad (4)$$

and

$$y_2 > \{y_1, y_4\} > y_3. \quad (5)$$

Equation (4) directly follows from the fact that for each point (x, y) of a hypersurface $H(c_i, c_j)$, we have $x \in [x_i, x_j]$: indeed, if $x_3 > x_2$, we would have $x_1 < x_2 < x_3 < x_4$ and the x-coordinate of each point of $H(c_1, c_2)$ would be smaller than the x-coordinate of each point of $H(c_3, c_4)$. In other words, the hypersurfaces would not intersect. An analogous reasoning can be done to show that $x_1 < x_4$. Equation (5) follows from the fact that the hypersurfaces do not even intersect if these inequalities are not satisfied.

Furthermore, Equation (5) means that when two vertical hypersurfaces $H(c_1, c_2)$ and $H(c_3, c_4)$ intersect twice, if the highest point in $\{c_1, \dots, c_4\}$ belongs to $\{c_1, c_2\}$ (resp. $\{c_3, c_4\}$) then the lowest point belongs to $\{c_3, c_4\}$ (resp. $\{c_1, c_2\}$).




Figure 24: Two intersecting hypersurfaces.

Assume first that the second pair of hypersurfaces that intersect twice are also vertical. As $y_3 = \min\{y_1, y_2, y_3, y_4\}$ and $y_2 = \max\{y_1, y_2, y_3, y_4\}$, by the discussion above $H(c_2, c_3)$ and $H(c_1, c_4)$ cannot intersect twice vertically. So the unique possibility is that $H(c_1, c_3)$ and $H(c_2, c_4)$ intersect twice. This is, however, not possible:

any point in $H(c_1, c_3)$ has x -coordinate in $[x_1, x_3]$, any point in $H(c_2, c_4)$ has x -coordinate in $[x_2, x_4]$, but $[x_1, x_3] \cap [x_2, x_4] = \emptyset$ by Equation (4).

Suppose now that the second pair of hypersurfaces that intersect twice are horizontal. This pair can be either $H(c_1, c_4)$ and $H(c_2, c_3)$, or $H(c_1, c_3)$ and $H(c_2, c_4)$.

- Let us first consider the case where it is $H(c_1, c_4)$ and $H(c_2, c_3)$, which is illustrated in Figure 25. Let us look at the preference p_1 in the upper left part. We have $c_1 > c_2$ and $c_3 > c_4$ (by the positions of $H(c_1, c_2)$ and $H(c_3, c_4)$). As $H(c_2, c_3)$ is horizontal, we have $c_2 > c_3$ (because $y_2 > y_3$ from Equation (5), see also Figure 24). Therefore $p_1 = (c_1, c_2, c_3, c_4)$. As $H(c_1, c_4)$ and $H(c_2, c_3)$ are horizontal, $H(c_2, c_3)$ is necessarily *above* $H(c_1, c_4)$ on the (infinite) left part of the figure, since starting from $p_1 = (c_1, c_2, c_3, c_4)$ and going down we need to cross $H(c_2, c_3)$ before $H(c_1, c_4)$: in fact, going down from the area corresponding to p_1 , we will not cross nor $H(c_1, c_2)$ neither $H(c_3, c_4)$ as they are vertical. We can only cross the remaining hypersurfaces $H(c_1, c_3)$, $H(c_1, c_4)$, $H(c_2, c_3)$ and $H(c_2, c_4)$. However, the hypersurface $H(c_1, c_3)$ (resp. $H(c_1, c_4)$, $H(c_2, c_4)$) cannot be the first hypersurface to be crossed, as c_2 is ranked between c_1 and c_3 (resp. c_2 and c_3 are ranked between c_1 and c_4 , c_3 between c_2 and c_4). Therefore, the first hypersurface to be crossed is necessarily $H(c_2, c_3)$ (c_2 and c_3 are ranked one next to other in p_1 , so they can be swapped).

Similarly, we get $p_2 = (c_4, c_3, c_2, c_1)$ in the lower right part, thus $H(c_2, c_3)$ is *below* $H(c_1, c_4)$ on the (infinite) right part of the figure, since starting from $p_2 = (c_4, c_3, c_2, c_1)$ and going up we need to cross $H(c_2, c_3)$ before $H(c_1, c_4)$, using the same reasoning as in the case of p_1 .




Figure 25: Relative positions of $H(c_1, c_4)$ and $H(c_2, c_3)$.

Hence, $H(c_2, c_3)$ and $H(c_1, c_4)$ cannot intersect twice (otherwise the same hypersurface would be above the other one both on the left part and the right part of the figure).

- Let us finally focus on the case where $H(c_1, c_3)$ and $H(c_2, c_4)$ are horizontal and intersect twice. They are necessarily both of type H^+ or both of type H^- (otherwise, they cannot intersect in two different points). We will show that none of these two cases is possible - in other words, that we cannot have $H(c_1, c_3)$ and $H(c_2, c_4)$ intersecting twice.

1. Firstly, let us assume that $H(c_1, c_3)$ and $H(c_2, c_4)$ are of type H^- :

We recall that $y_2 > \{y_1, y_4\} > y_3$ and $\{x_1, x_3\} < \{x_2, x_4\}$ (see Equations 5 and 4). However, the information on the type of $H(c_1, c_3)$ and $H(c_2, c_4)$ allows us to complete these partial orders on the coordinates of the candidates: using the classification of hypersurfaces (see Figure 6), we must have $x_3 < x_1$ and $x_4 < x_2$ if the hypersurfaces are of type H^- . Put together, we have $x_3 < x_1 < x_4 < x_2$. Moreover, the necessary order on y -coordinates is $y_2 > y_1 > y_4 > y_3$ - if the order would be $y_2 > y_4 > y_1 > y_3$, $H(c_1, c_3)$ and $H(c_2, c_4)$ would not intersect as any point of $H(c_i, c_j)$ has its y -coordinate in $[x_i, x_j]$. Let us denote by $U^{ij} = (U_x^{ij}, U_y^{ij})$ (resp. $L^{ij} = (L_x^{ij}, L_y^{ij})$) the upper extreme point (resp. the lower extreme point) of the middle-segment of $H(c_i, c_j)$. As $H(c_1, c_2)$ and $H(c_3, c_4)$ are of type V^- and intersect twice, the given orders on

both x -coordinates and y -coordinates of candidates implies that $U_x^{12} < U_x^{34}$ and $L_x^{12} < L_x^{34}$ (see Figure 24). When we express the segment extremities positions using the candidates coordinates, these two inequalities rewrite, after simplifying, as follows:

$$\begin{aligned} x_2 + x_1 + y_1 - y_2 &< x_4 + x_3 + y_3 - y_4 \\ x_1 + x_2 + y_2 - y_1 &< x_3 + x_4 + y_4 - y_3 \end{aligned}$$

If we sum the both inequalities, we get:

$$2(x_4 + x_3) > 2(x_1 + x_2),$$

in other words,

$$x_4 - x_1 > x_2 - x_3.$$

But this is in contradiction with the order on x -coordinates which states that $x_3 < x_1 < x_4 < x_2$. Therefore, the hypersurfaces $H(c_1, c_3)$ and $H(c_2, c_4)$ cannot be of type H^- .

2. Let us now assume that $H(c_1, c_3)$ and $H(c_2, c_4)$ are of type H^+ :

This case is illustrated in Figure 26. Note that the “upper” horizontal part of $H(c_1, c_3)$ (starting at the x -position x_3) is below the line (c_1, L^{12}) , as $y_3 < y_1$. Similarly, the lower horizontal part of $H(c_2, c_4)$ (ending at the x -position x_2) is above the line (U^{34}, c_4) as $y_2 > y_4$. If $H(c_1, c_3)$ and $H(c_2, c_4)$ intersect, then $H(c_1, c_3)$ is above $H(c_2, c_4)$ in the central part, see Figure 26.

Then in the (non empty) rectangle delimited by the 4 hypersurfaces (in the center of Figure 26), we have: $c_1 > c_2$, $c_4 > c_3$, $c_2 > c_4$ and $c_3 > c_1$, which yields $c_1 > c_2 > c_4 > c_3 > c_1$, a contradiction.

To conclude, we have proved by contradiction that $H(c_1, c_3)$ and $H(c_2, c_4)$ can neither be both of type H^+ nor both of type H^- . Therefore, they cannot intersect twice. \square




Figure 26: The case where $H(c_1, c_3)$ and $H(c_2, c_4)$ are horizontal and intersect twice.

\square

B Missing proofs of Section 6

B.1 Missing part in the proof of Theorem 3

Explicit construction of a family of profiles with $\Theta(m^4)$ distinct preferences

Proof. We set $c_1 = (0, 0)$ and $c_2 = (1, 2)$. According to the classification of hypersurfaces, $H(c_1, c_2)$ is horizontal (more precisely of type H^-). We then place c_3 in such a way that both $H(c_1, c_3)$ and $H(c_2, c_3)$ are vertical. To do so, we need to fix the values of x_3 and y_3 (coordinates of c_3) such that:

$$\begin{aligned} |x_1 - x_3| &> |y_1 - y_3| \\ \text{and } |x_2 - x_3| &> |y_2 - y_3|. \end{aligned}$$

This can be done by setting, for instance,

$$y_3 = \frac{y_1 + y_2}{2}$$

and

$$x_3 = \max\{x_1, x_2\} + 2|y_1 - y_2|$$

We check that, indeed, for $i \in \{1, 2\}$, we have

$$\begin{aligned} |x_3 - x_i| &= |\max\{x_1, x_2\} - x_i + 2|y_1 - y_2|| \\ &\geq 2|y_1 - y_2| > \frac{1}{2}|y_1 - y_2| \geq \left| \frac{y_1 + y_2}{2} - y_i \right| = |y_3 - y_i| \end{aligned}$$

where the strict inequality follows from the fact that $y_1 \neq y_2$. Geometrically, choosing y_3 between y_1 and y_2 ensures that $|y_3 - y_i|$ is upper bounded by $|y_1 - y_2|$. To guarantee that $|x_i - x_3| > |y_i - y_3|$, it is then sufficient that x_3 is taken large enough - here, the distance from $\max\{x_1, x_2\}$ (and so in particular from both x_1 and x_2) to x_3 is greater than the above mentioned upper bound $|y_1 - y_2|$.

We will now generalize the idea: we want $H(c_{2k}, c_i)$ to be horizontal for all $k \geq 1, i < 2k$, and $H(c_{2k+1}, c_i)$ to be vertical for $k \geq 1, i < 2k + 1$. Let us detail only the case of horizontal hypersurfaces (the case of vertical ones being symmetric).

We set

$$x_{2k} = \frac{\max_{i < 2k} \{x_i\} + \min_{i < 2k} \{x_i\}}{2}$$

and

$$y_{2k} = \max_{i < 2k} \{y_i\} + 2 \left(\max_{i < 2k} \{x_i\} - \min_{i < 2k} \{x_i\} \right).$$

The geometrical intuition remains the same - as we need, for all $i < 2k$, $|x_{2k} - x_i| < |y_{2k} - y_i|$, we choose the value of x_{2k} so that $|x_{2k} - x_i|$ is upper bounded by $\max_{i < 2k} \{x_i\} - \min_{i < 2k} \{x_i\}$, and we chose then y_{2k} in such a way that $|y_{2k} - y_i|$ is greater than this upper bound. Formally, we have:

$$\begin{aligned} |y_{2k} - y_i| &= \left| \max_{i < 2k} \{y_i\} + 2 \left(\max_{i < 2k} \{x_i\} - \min_{i < 2k} \{x_i\} \right) - y_i \right| \\ &= \left| \max_{i < 2k} \{y_i\} - y_i + 2 \left(\max_{i < 2k} \{x_i\} - \min_{i < 2k} \{x_i\} \right) \right| \\ &\geq 2 \left(\max_{i < 2k} \{x_i\} - \min_{i < 2k} \{x_i\} \right) \\ &> \frac{1}{2} \left(\max_{i < 2k} \{x_i\} - \min_{i < 2k} \{x_i\} \right) \\ &\geq \left| \frac{\max_{i < 2k} \{x_i\} + \min_{i < 2k} \{x_i\}}{2} - x_i \right| = |x_{2k} - x_i| \end{aligned}$$

Therefore we have $|y_{2k} - y_i| > |x_{2k} - x_i|$, so $H(c_{2k}, c_i)$ is horizontal.

Analogously, we set

$$y_{2k+1} = \frac{\max_{i < 2k+1} \{y_i\} + \min_{i < 2k+1} \{y_i\}}{2}$$

and

$$x_{2k+1} = \max_{i < 2k+1} \{x_i\} + 2 \left(\max_{i < 2k+1} \{y_i\} - \min_{i < 2k+1} \{y_i\} \right).$$

We prove as above (just by swapping the roles of x and y) that, in this case, the hypersurfaces $H(c_{2k+1}, c_i)$ are all vertical.

For $k \leq m$, we denote by H_k (resp. V_k) the number of horizontal (resp. vertical) hypersurfaces after adding the k -th candidate.

As all horizontal hypersurfaces intersect all vertical hypersurfaces, these intersections already define $(H_k + 1)(V_k + 1)$ different areas (with distinct preferences). Hence, denoting by A_m the number of areas after adding the k -th candidate, we have $A_m \geq (H_m + 1)(V_m + 1)$.

Each time we add a candidate c_k , we obtain $k - 1$ new hypersurfaces $H(c_1, c_k), \dots, H(c_{k-1}, c_k)$, all horizontal if k is even, or all vertical if k is odd. Consequently:

- if k is even, $H_k = H_{k-1} + (k - 1)$ and $V_k = V_{k-1}$;
- if k is odd, $H_k = H_{k-1}$ and $V_k = V_{k-1} + k - 1$.

We can deduce that $H_m \in \Theta(m^2)$ and $V_m \in \Theta(m^2)$, and thus $A_m \in \Omega(m^4)$. □