

Fate of higher-order topological insulator under Coulomb interaction

Jing-Rong Wang^{1,*} and Chang-Jin Zhang^{1,2,†}

¹*High Magnetic Field Laboratory of Anhui Province,
Chinese Academy of Sciences, Hefei 230031, China*

²*Institute of Physical Science and Information Technology, Anhui University, Hefei 230601, China*

In this article, we study the influence of long-range Coulomb interaction on three-dimensional second-order topological insulator (TI) by renormalization group theory. We find that both the analysis method and conclusions in the recent Letter Phys. Rev. Lett. **127**, 176601 (2021) are unreliable. There are two problems in this Letter. Firstly, the characteristic described by the RG flows $m \rightarrow \infty$ and $D \rightarrow 0$ can not be used as the criterion for transition from second-order TI to TI, since this characteristic could be essentially not induced by Coulomb interaction but only results from the trivial power counting contribution of fermion action. Indeed, this characteristic is satisfied even for free second-order TI. Second, the flow of B is not paid attention, which is very important and should be seriously studied. In this article, we analyze carefully the corrections for the flows of the model parameters induced by Coulomb interaction. We find that the sign of m changes but the sign of B holds if the initial Coulomb strength is large enough, while the sign of m holds but the sign of B changes if the initial Coulomb strength takes small values. These results indicate that second-order TI is unstable to trivial band insulator not only under strong Coulomb interaction but also under weak Coulomb interaction. We also study the effects of disorder scattering in second-order TI by renormalization group theory. According to the criterion in Phys. Rev. Lett. **127**, 176601 (2021), weak disorder drives second-order TI to TI. However, we find that second-order TI is robust against weak disorder, since weak disorder does not give qualitative modification for second-order TI. This result is consistent with recent studies based on other methods. Interplay of Coulomb interaction and disorder in second-order TI is also investigated.

I. INTRODUCTION

The studies about topological materials including topological insulators (TIs), topological superconductors, and various topological semimetals, have become one of the most important fields in condensed matter physics [1–11]. Topological materials have critical potential applications, including quantum computation, thermoelectric devices *etc.*, due to their fantastic properties. Dirac, Weyl, and Majornana fermions have been observed in some topological materials [1–11]. These fermion excitations resemble the elementary particles in high-energy physics. Thus, topological materials could provide a platform to simulate the concepts and phenomena in high-energy physics. In some topological materials, there are also unusual fermion excitations, such as semi-Dirac fermions, double-Weyl fermions, triple-Weyl fermions, multi-fold degenerate fermions *etc.*, which have not counterparts in high-energy physics [9–11]. These unusual fermion excitations could result in novel physical behaviors.

Recently, higher-order topological materials attracted a lot of interest [12–30]. For d dimensional TI, the system has $d - 1$ gapless Dirac edge states. For d dimensional second-order TI, the system hosts $d - 2$ gapless edge states. Concretely, three-dimensional (3D) second-order TI hosts 1D hinge states, and 2D second-order TI

has 0D corner states.

Study about correlated interaction effects in topological materials is an important direction and attracted particular interest [31–77]. For example, the theoretical studies showed that long-range Coulomb interaction induces singular fermion velocity renormalization for 2D Dirac fermions, which has been observed experimentally [31–35].

Recently, Zhao *et al.* studied the influence of long-range Coulomb interaction on 3D second-order TIs by renormalization group (RG) theory [73]. They concluded that 3D second-order TIs are always unstable under Coulomb interaction. They showed that there are two types of transitions: second-order TI to TI and second-order TI to trivial band insulator.

However, after careful studies, we find that both of the analysis method and conclusions in Ref. [73] are unreliable. There are two problems in Ref. [73]. First, the characteristic described by the RG flows $m \rightarrow \infty$ and $D \rightarrow 0$ can not be used as the criterion for transition from second-order TI to TI, since this characteristic could be essentially not induced by Coulomb interaction but only results from the trivial power counting contribution of fermion action. Actually, this characteristic is satisfied even for free second-order TI. Second, the flow of B is not paid attention, which is very important and should be seriously studied.

In the recent Comment [74], Lee and Yang have also pointed the problems in Ref. [73] and indicated the conclusions are misleading. The problems in Ref. [73] pointed by Lee and Yang and the ones pointed by us are similar to each other.

*Corresponding author: wangjr@hmf.ac.cn

†Corresponding author: zhangcj@hmf.ac.cn

Whereas, there are also differences between the studies in the Comment [74] and the studies by us. Based on further calculations, they concluded that second-order TI is robust against weak Coulomb interaction. However, we find that the sign of m holds but the sign of the parameter B changes for weak Coulomb interaction. Namely mB changes under the weak Coulomb interaction. It represents that second-order TI is unstable to trivial band insulator under weak Coulomb interaction.

We also study the influence of disorder on second-order TI. According to the criterion in Ref. [73], weak disorder drives second-order TI to TI. However, we find that weak disorder does not give qualitative modification for second-order TI through RG analysis. Namely, second-order TI is robust against weak disorder. Our result is consistent with recent studies about disorder effects in second-order TI based on other methods [24, 25]. The interplay of long-range Coulomb interaction and disorder is also investigated.

The rest of paper is structured as follows. The model for second-order TI with long-range Coulomb interaction is defined in Sec. II. In Sec. III, we present the RG analysis for influence of Coulomb interaction on second-order TI based on numerical and analytical calculations. We compare interaction effects in related systems and give discussions for some related questions in Sec. IV. In Sec. V, we analyze the effects of disorder in second-order TI. The interplay of long-range Coulomb interaction and disorder in second-order TI is studied in Sec. VI. A brief summary is given in Sec. VII. The detailed calculations and derivations are presented in Appendices.

II. MODEL

The free action of fermions is

$$S_\Psi = \int \frac{dk_0}{2\pi} \frac{d^3\mathbf{k}}{(2\pi)^3} \bar{\Psi}(k_0, \mathbf{k}) (ik_0\gamma_0 + \mathcal{H}_f) \Psi(k_0, \mathbf{k}), \quad (1)$$

where Ψ is four-component spinor and $\bar{\Psi} = \Psi^\dagger\gamma_0$. The fermion Hamiltonian density takes the form

$$\begin{aligned} \mathcal{H}_f = & i[v(k_x\gamma_x + k_y\gamma_y) + v_z k_z\gamma_z + D(k_x^2 - k_y^2)\gamma_5] \\ & + m - B_\perp k_\perp^2 - B_z k_z^2. \end{aligned} \quad (2)$$

v , v_z , D , $B_\perp$, B_z are model parameters. If $mB_{\perp,z} > 0$ and $D \neq 0$, it corresponds to second-order TI. If $mB_{\perp,z} > 0$ and $D = 0$, it corresponds to TI. If $mB_{\perp,z} < 0$, it corresponds to trivial band insulator. The matrices γ_0 , γ_x , γ_y , γ_z and γ_5 satisfy the anticommuting relation $\{\gamma_\mu, \gamma_\nu\} = 2\delta_{\mu\nu}$.

The long-range Coulomb interaction between fermions can be written as

$$H_C = \frac{1}{4\pi} \int d^3\mathbf{x} d^3\mathbf{x}' \rho(\mathbf{x}) \frac{e^2}{\epsilon|\mathbf{x} - \mathbf{x}'|} \rho(\mathbf{x}'), \quad (3)$$

where $\rho(\mathbf{x}) = \Psi^\dagger(\mathbf{x})\Psi(\mathbf{x})$ is the fermion density operator, e electric charge, and ϵ dielectric constant. The

Coulomb interaction can be decoupled by introducing a bosonic field ϕ through Hubbard-Stratonovich transformation. Accordingly, the Coulomb interaction between fermions can be described by the action of fermion-boson coupling as

$$S_{\Psi\phi} = ig \int d\tau d^3\mathbf{x} \bar{\Psi}\gamma_0\Psi\phi, \quad (4)$$

where $g = e/\sqrt{\epsilon}$. In energy-momentum space, it takes the form

$$\begin{aligned} S_{\Psi\phi} = & ig \int \frac{dk_{0,1}}{2\pi} \frac{d^3\mathbf{k}_1}{(2\pi)^3} \frac{dk_{0,2}}{2\pi} \frac{d^3\mathbf{k}_2}{(2\pi)^3} \bar{\Psi}(k_{0,1}, \mathbf{k}_1)\gamma_0 \\ & \times \Psi(k_{0,2}, \mathbf{k}_2)\phi(k_{0,1} - k_{0,2}, \mathbf{k}_1 - \mathbf{k}_2). \end{aligned} \quad (5)$$

The free action of boson field ϕ can be written as

$$S_\phi = \int \frac{dk_0}{2\pi} \frac{d^3\mathbf{k}}{(2\pi)^3} \phi(k_0, \mathbf{k}) (k_\perp^2 + \eta k_z^2) \phi(k_0, \mathbf{k}), \quad (6)$$

where η is used to describe the anisotropy of ϕ .

III. RG RESULTS

In this article, we study the influence of long-range Coulomb interaction on second-order TI through RG method [78]. After detailed calculations and derivations shown in Appendices, we obtain the RG equations as following,

$$\frac{dv}{d\ell} = \alpha\mathcal{R}_v v, \quad (7)$$

$$\frac{dv_z}{d\ell} = \alpha\mathcal{R}_{v_z} v_z, \quad (8)$$

$$\frac{dm}{d\ell} = m + \alpha\mathcal{R}_m, \quad (9)$$

$$\frac{dB_\perp}{d\ell} = -B_\perp + \alpha\mathcal{R}_{B_\perp} \quad (10)$$

$$\frac{dB_z}{d\ell} = -B_z + \alpha\mathcal{R}_{B_z} \quad (11)$$

$$\frac{dD}{d\ell} = -D + \alpha\mathcal{R}_D D, \quad (12)$$

$$\frac{d\alpha}{d\ell} = -\alpha^2\mathcal{R}_\alpha, \quad (13)$$

$$\frac{d\zeta}{d\ell} = \alpha\mathcal{R}_\zeta\zeta, \quad (14)$$

where ℓ is the RG running parameter. The strength of Coulomb interaction α and the parameter ζ are defined as

$$\alpha = \frac{g^2}{4\pi^2 v \sqrt{\eta}}, \quad (15)$$

$$\zeta = \frac{v_z}{v\sqrt{\eta}}. \quad (16)$$

The transformations

$$\frac{m}{v\Lambda} \rightarrow m, \quad (17)$$

$$\frac{B_{\perp}\Lambda}{v} \rightarrow B_{\perp}, \quad (18)$$

$$\frac{B_z\Lambda}{v\eta} \rightarrow B_z, \quad (19)$$

$$\frac{D\Lambda}{v} \rightarrow D, \quad (20)$$

have been utilized. $\mathcal{R}_v, \mathcal{R}_{v_z}, \mathcal{R}_m, \mathcal{R}_{B_{\perp}}, \mathcal{R}_{B_z}, \mathcal{R}_D, \mathcal{R}_{\alpha}$, and $\mathcal{R}_{\zeta}$ are functions of $m, B_{\perp}, B_z, D$, and ζ , whose concrete expressions can be found in Appendix F.

For free fermions, the RG equations become

$$\frac{dv^f}{d\ell} = 0, \quad (21)$$

$$\frac{dv_z^f}{d\ell} = 0, \quad (22)$$

$$\frac{dm^f}{d\ell} = m, \quad (23)$$

$$\frac{dB_{\perp}^f}{d\ell} = -B_{\perp}, \quad (24)$$

$$\frac{dB_z^f}{d\ell} = -B_z, \quad (25)$$

$$\frac{dD^f}{d\ell} = -D, \quad (26)$$

where the superscript f corresponds to free fermions. The corresponding solutions are

$$v^f(\ell) = v_0, \quad (27)$$

$$v_z^f(\ell) = v_{z0}, \quad (28)$$

$$m^f(\ell) = m_0 e^{\ell}, \quad (29)$$

$$B_{\perp}^f(\ell) = B_{\perp 0} e^{-\ell}, \quad (30)$$

$$B_z^f(\ell) = B_{z0} e^{-\ell}, \quad (31)$$

$$D^f(\ell) = D_0 e^{-\ell}, \quad (32)$$

where $v_0, v_{z0}, B_{\perp 0}, B_{z0}$, and D_0 are initial values. It is easy to find that $v^f(\ell)$ and $v_z^f(\ell)$ are fixed, and

$$m^f(\ell) \rightarrow \infty, \quad (33)$$

$$B_{\perp}^f(\ell) \rightarrow 0, \quad (34)$$

$$B_z^f(\ell) \rightarrow 0, \quad (35)$$

$$D^f(\ell) \rightarrow 0, \quad (36)$$

in the lowest energy limit $\ell \rightarrow \infty$. These results are directly related to the fact that v and v_z are coefficients of linear terms of momentum components, m is coefficient of zero power of momentum components, and $B_{\perp}, B_z, D$ are coefficients of quadratic terms of momentum components.

The physical meaning of these results shown in Eqs. (27)-(32) and Eqs. (33)-(36) is: Taking the linear terms of Hamiltonian as reference energy, the zero power term becomes larger and larger, and the quadratic terms become smaller and smaller, with lowering of momentum. Namely,

$$m/E_{linear}(\mathbf{k}) \rightarrow \infty, \quad (37)$$




FIG. 1: (a)-(f): Flows of $v(\ell)$, $v_z(\ell)$, $\alpha(\ell)$, $\zeta(\ell)$, $D(\ell)$, and $D(\ell)/D^f(\ell)$ with different initial values of Coulomb strength. Blue, red, green, black, and magenta lines correspond to the initial values $\alpha_0 = 0.1, 0.2, 0.3, 0.4, 0.5$ respectively. $m_0 = 0.1$, $B_{\perp 0} = 1$, $B_{z0} = 1$, $D_0 = 1$, $\zeta_0 = 0.1$ are taken.

$$Bk_{\perp}^2/E_{linear}(\mathbf{k}) \rightarrow 0, \quad (38)$$

$$Bk_z^2/E_{linear}(\mathbf{k}) \rightarrow 0, \quad (39)$$

$$D(k_x^2 - k_y^2)/E_{linear}(\mathbf{k}) \rightarrow 0, \quad (40)$$

where

$$E_{linear}(\mathbf{k}) = \sqrt{v^2 k_{\perp}^2 + v_z^2 k_z^2}, \quad (41)$$

in the limit $k \rightarrow 0$.

In Ref. [73], the characteristic described by the RG flows $m(\ell) \rightarrow \infty$ and $D(\ell) \rightarrow 0$ in the limit $\ell \rightarrow \infty$ are used as the criterion for transition from second-order TI to TI. It is clear that using this characteristic as the criterion for transition from second-order TI to TI is invalid. This characteristic is even satisfied for free second-order TI.

The flows of $v(\ell)$, $v_z(\ell)$, $m(\ell)$, $\alpha(\ell)$, $D(\ell)$, and $D(\ell)/D^f(\ell)$ with different initial values of Coulomb strength are presented in Figs. 1(a)-1(f) respectively. We find that $v(\ell)/v_0$ and $v_z(\ell)/v_{z0}$ flow to positive constants. Thus, Coulomb interaction results in quantitative corrections for v and v_z . According to Figs. 1(c) and 1(d),

$$\alpha(\ell) \rightarrow \alpha^*, \quad (42)$$

$$\zeta(\ell) \rightarrow \zeta^*, \quad (43)$$

where α^* and ζ^* are positive constants. As shown in Fig. 1(e), $D(\ell)$ approaches to zero quickly with lowering of energy scale. From Fig. 1(f), we find that $D(\ell)/D^f(\ell)$ flows to a positive constant value in the low energy regime. It indicates that $D(\ell)$ only acquires quantitative correction in presence of Coulomb interaction, and takes qualitatively same behavior as $D^f(\ell)$. It means that qualitative behaviors of the term $D(k_x^2 - k_y^2)$ are not changed under Coulomb interaction. As the term




FIG. 2: (a)-(f): Flows of $m(\ell)$, $m(\ell)/m^f(\ell)$, $B_\perp(\ell)$, $B_\perp(\ell)/B_\perp^f(\ell)$, $B_z(\ell)$, and $B_z(\ell)/B_z^f(\ell)$ with different initial values of Coulomb strength. Blue, red, green, black, and magenta lines correspond to the initial values $\alpha_0 = 0.1, 0.2, 0.3, 0.4, 0.5$ respectively. $m_0 = 0.1$, $B_{\perp 0} = 1$, $B_{z0} = 1$, $D_0 = 1$, $\zeta_0 = 0.1$ are taken.

$D(k_x^2 - k_y^2)$ is not changed qualitatively by Coulomb interaction, the transition from second-order TI to TI stated in Ref. [73] does not exist.

We presented the flows of $m(\ell)$, $m(\ell)/m^f(\ell)$, $B_\perp(\ell)$, $B_\perp(\ell)/B_\perp^f(\ell)$, $B_z(\ell)$, $B_z(\ell)/B_z^f(\ell)$ with different initial values of Coulomb strength in Figs. 2(a)-2(f) respectively. If the initial Coulomb strength is small, we find that

$$m(\ell)/m^f(\ell) \rightarrow c_m^*, \quad \text{with } c_m^* > 0, \quad (44)$$

where c_m^* is a positive constant. It means that $m(\ell)$ takes the qualitatively same behavior as $m^f(\ell)$. According to Figs. 2(c) and 2(e),

$$B_\perp(\ell) \rightarrow c_{B_\perp}^*, \quad \text{with } c_{B_\perp}^* < 0, \quad (45)$$

$$B_z(\ell) \rightarrow c_{B_z}^*, \quad \text{with } c_{B_z}^* < 0, \quad (46)$$

with lowering of energy scale, where $c_{B_\perp}^*$ and $c_{B_z}^*$ are negative constants. We can find that signs of $B_\perp(\ell)$ and $B_z(\ell)$ change. In the low energy regime, it is easy to obtain

$$B_\perp(\ell)/B_\perp^f(\ell) \sim \frac{c_{B_\perp}^*}{B_{\perp 0}} e^\ell \rightarrow -\infty, \quad (47)$$

$$B_z(\ell)/B_z^f(\ell) \sim \frac{c_{B_z}^*}{B_{z0}} e^\ell \rightarrow -\infty. \quad (48)$$

These results indicate that the behaviors of $B_\perp$ and B_z are obviously modified by Coulomb interaction. These results can be also noticed from Figs. 2(d) and 2(f). The behaviors shown in Eqs. (47) and (48) reveal that the positive quadratic terms $B_\perp k_\perp^2$ and $B_z k_z^2$ become negative linear terms of momentum components induced by Coulomb interaction.

If the initial Coulomb strength is large enough, we notice that

$$m(\ell)/m^f(\ell) \rightarrow c_m^*, \quad \text{with } c_m^* < 0, \quad (49)$$

where c_m^* is a negative constant. The sign of $m(\ell)$ changes in this case. As shown in Figs. 2(c) and 2(e),

$$B_\perp(\ell) \rightarrow c_{B_\perp}^*, \quad \text{with } c_{B_\perp}^* > 0, \quad (50)$$

$$B_z(\ell) \rightarrow c_{B_z}^*, \quad \text{with } c_{B_z}^* > 0, \quad (51)$$

where $c_{B_\perp}^*$ and $c_{B_z}^*$ are positive constants. The signs of $B_\perp(\ell)$ and $B_z(\ell)$ hold. We can further get

$$B_\perp(\ell)/B_\perp^f(\ell) \sim \frac{c_{B_\perp}^*}{B_{\perp 0}} e^\ell \rightarrow \infty, \quad (52)$$

$$B_z(\ell)/B_z^f(\ell) \sim \frac{c_{B_z}^*}{B_{z0}} e^\ell \rightarrow \infty, \quad (53)$$

which can be also viewed from Figs. 2(d) and 2(f). The flows as shown in Eq. (52) and (53) mean that the positive quadratic terms $B_\perp k_\perp^2$ and $B_z k_z^2$ are modified to positive linear terms of momentum components by Coulomb interaction.

In order to better understanding this question, we give analytical calculation for the asymptotic behaviors of $B_\perp(\ell)$ and $B_z(\ell)$. We have find that $\alpha(\ell) \rightarrow \alpha^*$, $m(\ell) \rightarrow \infty$ for weak Coulomb interaction and $m(\ell) \rightarrow -\infty$ for strong enough Coulomb interaction. Accordingly, in the low energy regime, the RG equations for $B_\perp$ and B_z can be approximated by

$$\frac{dB_\perp}{d\ell} \sim -B_\perp - \frac{1}{3}\alpha^* \frac{m}{|m|} \sim -B_\perp - \frac{1}{3}\alpha^* \text{sgn}(m), \quad (54)$$

$$\frac{dB_z}{d\ell} \sim -B_z - \frac{1}{3}\alpha^* \frac{m}{|m|} \sim -B_z - \frac{1}{3}\alpha^* \text{sgn}(m). \quad (55)$$

Accordingly, we can find that

$$B_\perp(\ell) \rightarrow -\frac{1}{3}\alpha^* \text{sgn}(m), \quad (56)$$

$$B_z(\ell) \rightarrow -\frac{1}{3}\alpha^* \text{sgn}(m), \quad (57)$$

in the limit $\ell \rightarrow \infty$. The numerical results of $B_\perp(\ell)/(\alpha(\ell)/3)$ and $B_z(\ell)/(\alpha(\ell)/3)$ are shown in Fig. 3. We can find these numerical results verify the asymptotic behaviors as shown in Eqs. (56) and (57).

Therefore, we can find that the sign of m holds but the signs of $B_\perp$ and B_z change for weak Coulomb interaction, while the sign of m changes but the signs of $B_\perp$ and B_z hold if Coulomb interaction is strong enough. Namely, signs of $mB_\perp$, mB_z usually always change. Thus, second-order TI usually becomes a trivial band insulator not only under strong Coulomb interaction but also under weak Coulomb interaction.

Taking proper initial conditions, m , $B_\perp$, B_z may flow to zero simultaneously. In this case, the system may become a second-order Dirac semimetal (DSM).




FIG. 3: (a) and (b): Flows of $B_{\perp}(\ell)/(\alpha(\ell)/3)$ and $B_z(\ell)/(\alpha(\ell)/3)$ with different initial values of Coulomb strength. Blue, red, green, black, and magenta lines correspond to the initial values $\alpha_0 = 0.1, 0.2, 0.3, 0.4, 0.5$ respectively. $m_0 = 0.1$, $B_{\perp 0} = 1$, $B_{z0} = 1$, $D_0 = 1$, $\zeta_0 = 0.1$ are taken.

In order to avoid possible misunderstanding, it should be strengthened that result $m(\ell) \rightarrow \infty$ does not mean the physical mass becomes to infinity, but represents that the physical mass becomes larger and larger comparing to the linear terms of Hamiltonian with lowering of momentum. The physical mass for free second-order TI is a constant. The physical mass for second-order TI with weak Coulomb interaction only receives quantitative correction and is also a constant with same sign. Thus, we can find that the RG results shown in above will not break down at the scale where $m(\ell)$ is order of one. Actually, the RG results are valid in the limit $\ell \rightarrow \infty$, namely $k \rightarrow 0$.

IV. RESULTS FOR OTHER RELATED SYSTEMS AND SOME DISCUSSIONS

A. Second-order Dirac semimetal

Taking the initial conditions $m_0 = 0$, $B_{\perp 0} = 0$, and $B_{z0} = 0$, we can obtain the RG equations for second-order DSM. The corresponding numerical results are shown in Fig. 4. We notice that α flows to zero slowly, and v and v_z increase slowly with lowering of energy scale. As shown in Fig. 4(f), $D(\ell)/D^f(\ell)$ decreases slowly with lowering of energy scale. These results imply that the terms $vk_{\perp}$, vk_z , $D(k_x^2 - k_y^2)$ receive weak logarithmic-like corrections of momentum components under Coulomb interaction. It indicates that the




FIG. 4: (a)-(f): Flows of $v(\ell)$, $v_z(\ell)$, $\alpha(\ell)$, $\gamma(\ell)$, $D(\ell)$, and $D(\ell)/D^f(\ell)$ for second-order DSM with different initial Coulomb strength. Blue, red, green, black, and magenta lines correspond to the initial values $\alpha_0 = 0.1, 0.2, 0.3, 0.4, 0.5$ respectively. $D_0 = 1$, $\zeta_0 = 0.1$ are taken.

observable quantities such as density of states, specific heat, compressibility acquire logarithmic-like corrections of energy or temperature, and second-order DSM is robust against Coulomb interaction. These results are consistent with recent studies in Ref. [75].

B. 3D TI

Taking the initial value $D_0 = 0$, we get the RG equations for TI. From numerical and analytical calculations, we also find that α approaches to constant value α^* , and

$$B_{\perp}(\ell) \rightarrow -\frac{1}{3}\alpha^* \text{sgn}(m), \quad (58)$$

$$B_z(\ell) \rightarrow -\frac{1}{3}\alpha^* \text{sgn}(m). \quad (59)$$

The flows of $\alpha(\ell)$, $m(\ell)$, $B_{\perp}(\ell)$, $B_{\perp}(\ell)/(\alpha(\ell)/3)$, $B_z(\ell)$, and $B_z(\ell)/(\alpha(\ell)/3)$ for TI are presented in Fig. 5. These results indicate that TI is also usually unstable to trivial band insulator not only under strong Coulomb interaction but also under weak Coulomb interaction.

In the study by Goswami *et al.* [38], the RG equations for the TI with long-range Coulomb interaction are given. However, the RG flow of the parameter B is not analyzed carefully and changing of sign of B is not noticed in Ref. [38]. Through numerical and analytical calculations for the RG equations shown in Ref. [38], we obtain that α flows to a constant value α^* , and

$$B(\ell) \rightarrow \frac{\pi}{3}\alpha^* \text{sgn}(m), \quad (60)$$

in the limit $\ell \rightarrow \infty$ in their notations. It should be noticed that the definition of B in Ref. [38] is different




FIG. 5: (a)-(f): Flows of $\alpha(\ell)$, $m(\ell)$, $B_{\perp}(\ell)$, $B_{\perp}(\ell)/(\alpha(\ell)/3)$, $B_z(\ell)$, and $B_z(\ell)/(\alpha(\ell)/3)$ for TI with different initial values of Coulomb strength. Blue, red, green, black, and magenta lines correspond to the initial values $\alpha_0 = 0.1, 0.2, 0.3, 0.4, 0.5$ respectively. $m_0 = 0.1$, $B_{\perp 0} = 1$, $B_{z0} = 1$, $\zeta_0 = 0.1$ are taken.

from ours with a minus. Namely, $mB < 0$ corresponds to TI and $mB > 0$ stands for normal insulator in Ref. [38]. Whereas, $mB_{\perp,z} > 0$ correspond to TI and $mB_{\perp,z} < 0$ stand for normal insulator in our article. Additionally, the definition of α in our article is different from the one in Ref. [38] with a factor $1/\pi$. Thus, the results we obtain through the RG equations in Ref. [38] are completely consistent with the ones shown in Eqs. (58) and (59).

C. Short-range four-fermion interactions

Roy *et al.* have showed the phase diagrams of TI under short-range four-fermions interactions [76]. Influence of short-range four-fermion interactions on second-order TI is also interesting. For short-range interactions with the form

$$\mathcal{L}_{four\,fermion} = \sum_i \lambda_i (\bar{\Psi} \Gamma_i \Psi)^2, \quad (61)$$

through RG calculation, we find that $B_{\perp,z}(\ell)$ and $D(\ell)$ take the behaviors

$$B_{\perp}(\ell) = B_{\perp 0} e^{-\ell}, \quad (62)$$

$$B_z(\ell) = B_{z0} e^{-\ell}, \quad (63)$$

$$D_z(\ell) = D_0 e^{-\ell}, \quad (64)$$

which are same as the free system, and m will not change sign under weak short-range four-fermion interactions. Thus, second-order TI is robust against weak short-range four-fermion interactions. If the strength of short-range four-fermion interaction is large enough, some quantum

phase transitions could be triggered. The detailed studies about second-order TI under short-range four-fermion interactions are left in further studies.

The important point we should strength here is that weak long-range Coulomb interaction is different from weak short-range four-fermion interactions. Weak long-range Coulomb interaction drives second-order TI to trivial band insulator, whereas second-order TI is robust against weak short-range four-fermion interactions.

D. Comments for the arguments in a recent reference

In a recent reference, Liu *et al.* give some arguments for our results about sign change of $B_{\perp,z}$, and they believe that second-order TI is robust against weak Coulomb interaction [77]. We should indicate that their arguments are unfounded. First, as shown in Sec. III, through numerical and analytical calculations, we have clearly showed that $B_{\perp,z}$ indeed change sign under weak long-range Coulomb interaction. Second, the Coulomb strength $\alpha(\ell)$ decreases from the initial value α_0 gradually and approaches to a small finite value α^* in the lowest energy limit $\ell \rightarrow \infty$. Thus, the RG is controlled and the corresponding results are valid in the lowest energy regime. Their argument that RG becomes invalid in large ℓ (low energy regime) is incorrect. Third, their arguments about the high order terms of momenta are invalid. Liu *et al.* considered that $B_{\perp} k_{\perp}^2$, $B_z k_z^2$ terms come from the expansion of $\sum_i t_i \cos(k_i a)$ as

$$\sum_i t_i \cos(k_i a) = \text{const} - B_{\perp} k_{\perp}^2 - B_z k_z^2 + O(k^4, k^6), \quad (65)$$

and higher order terms beyond k^2 terms exist in general. They stated that even if $B_{\perp,z}$ change sign, then the high order terms k^4 , k^6 can not be ignored. However, we should indicate that their arguments are unfounded. It is easy to find that $\sum_i t_i \cos(k_i a)$ can be expand as

$$\begin{aligned} & \sum_i t_i \cos(k_i a) \\ &= \text{const} - B_{\perp} k_{\perp}^2 - B_z k_z^2 \\ & \quad + B_{\perp(4)} (k_x^4 + k_y^4) + B_{z(4)} k_z^4 \\ & \quad - B_{\perp(6)} (k_x^6 + k_y^6) - B_{z(6)} k_z^6 \\ & \quad \dots + (-1)^n [B_{\perp(2n)} (k_x^{2n} + k_y^{2n}) + B_{z(2n)} k_z^{2n}] \\ & \quad \dots, \end{aligned} \quad (66)$$

where the coefficients $B_{\perp}$, B_z , $B_{\perp(4)}$, $B_{z(4)}$, $B_{\perp(6)}$, $B_{z(6)}$ etc. are all positive. We can find the terms k_i^{2n} with n being odd number are negative, whereas the terms k_i^{2n} with n being even number are positive. These terms could be modified by long-range Coulomb interaction. From the numerical and analytical calculations shown in Sec. III, we have showed that $B_{\perp}$ and B_z change sign under the weak Coulomb interaction. The high order terms k_i^4 , k_i^6

could be also modified by long-range Coulomb interaction. Through tedious calculations, we find that the RG equations for $B_{\perp(4)}$, $B_{z(4)}$, $B_{\perp(6)}$, and $B_{z(6)}$ in the low energy regime can be approximately written as

$$\frac{dB_{\perp(4)}}{d\ell} \sim -3B_{\perp(4)} + \frac{1}{5}\alpha^*\text{sgn}(m), \quad (68)$$

$$\frac{dB_{z(4)}}{d\ell} \sim -3B_{z(4)} + \frac{1}{5}\alpha^*\text{sgn}(m), \quad (69)$$

$$\frac{dB_{\perp(6)}}{d\ell} \sim -5B_{\perp(6)} - \frac{1}{7}\alpha^*\text{sgn}(m), \quad (70)$$

$$\frac{dB_{z(6)}}{d\ell} \sim -5B_{z(6)} - \frac{1}{7}\alpha^*\text{sgn}(m), \quad (71)$$

where the transformations

$$\frac{B_{\perp(4)}\Lambda^3}{v} \rightarrow B_{\perp(4)}, \quad (72)$$

$$\frac{B_{z(4)}\Lambda^3}{v\eta^2} \rightarrow B_{z(4)}, \quad (73)$$

$$\frac{B_{\perp(6)}\Lambda^5}{v} \rightarrow B_{\perp(6)}, \quad (74)$$

$$\frac{B_{z(6)}\Lambda^5}{v\eta^3} \rightarrow B_{z(6)}, \quad (75)$$

have been utilized. Solving Eqs. (68)-(71), we obtain

$$B_{\perp(4)}(\ell) \rightarrow \frac{1}{15}\alpha^*\text{sgn}(m), \quad (76)$$

$$B_{z(4)}(\ell) \rightarrow \frac{1}{15}\alpha^*\text{sgn}(m), \quad (77)$$

$$B_{\perp(6)}(\ell) \rightarrow -\frac{1}{35}\alpha^*\text{sgn}(m), \quad (78)$$

$$B_{z(6)}(\ell) \rightarrow -\frac{1}{35}\alpha^*\text{sgn}(m), \quad (79)$$

in the lowest energy limit $\ell \rightarrow \infty$. We can find that for weak Coulomb interaction $B_{\perp(4)}$ and $B_{z(4)}$ are still positive, whereas $B_{\perp(6)}$ and $B_{z(6)}$ become to negative. Further, we notice that the RG equations for the coefficients of high order terms can be generally expressed by

$$\frac{dB_{\perp(2n)}}{d\ell} \sim -(2n-1)B_{\perp(2n)} + \frac{(-1)^n}{2n+1}\alpha^*\text{sgn}(m), \quad (80)$$

$$\frac{dB_{z(2n)}}{d\ell} \sim -(2n-1)B_{z(2n)} + \frac{(-1)^n}{2n+1}\alpha^*\text{sgn}(m), \quad (81)$$

where the transformations

$$\frac{B_{\perp(2n)}\Lambda^{2n-1}}{v} \rightarrow B_{\perp(2n)}, \quad (82)$$

$$\frac{B_{z(2n)}\Lambda^{2n-1}}{v\eta^n} \rightarrow B_{z(2n)}, \quad (83)$$

have been employed. From Eqs. (80) and (81), we get

$$B_{\perp(2n)}(\ell) \rightarrow (-1)^n \frac{1}{(2n+1)(2n-1)}\alpha^*\text{sgn}(m), \quad (84)$$

$$B_{z(2n)}(\ell) \rightarrow (-1)^n \frac{1}{(2n+1)(2n-1)}\alpha^*\text{sgn}(m), \quad (85)$$

in the limit $\ell \rightarrow \infty$. Thus, under weak Coulomb interaction, $B_{\perp, z(2n)}$ change sign if n is odd number, but hold sign if n is even number. From the above results we know that the original negative terms k_i^{2n} with n being odd number become positive, and the original positive terms k_i^{2n} with n being even number are still positive, under weak Coulomb interaction. Therefore, we find that the system becomes trivial insulator under the influence of weak long-range Coulomb interaction even if the high order terms k_i^4 , k_i^6 etc. are considered.

V. INFLUENCE OF DISORDER SCATTERING

In Ref. [73], Zhao *et al.* also studied the influence of disorder on 3D second-order TI based on RG method. Their studies about disorder effects in second-order TI have several problems. First, they used the invalid criterion for transition from second-order TI to TI to study the influence of disorder on second-order TI. According to their criterion, considering only disorder, weak disorder drives second-order TI to TI. However, recent studies about disorder effects in 3D second-order TI through other methods showed that 3D second-order TI is robust against weak disorder [24, 25].

Second, generation new type of disorder by one type of disorder is not considered in Ref. [73]. If the fermion Hamiltonian of a system satisfies

$$\mathcal{H}_f(\mathbf{k}) + \mathcal{H}_f(-\mathbf{k}) = 0, \quad (86)$$

one type of disorder can exist solely. For examples, in DSM or Weyl semimetal (WSM), the Hamiltonian satisfies Eq. (86), thus one type of disorder can exist solely [45, 79-84]. However, if the Hamiltonian of a system satisfies

$$\mathcal{H}_f(\mathbf{k}) + \mathcal{H}_f(-\mathbf{k}) \neq 0. \quad (87)$$

Some type of disorder can not exist solely and generate other types of disorder. For examples, the generation effect of new disorder results from Eq. (87) was considered in the studies about disorder effects in 2D semi-DSM [85], 3D anisotropic WSM [86-88], 3D double-WSM [56, 89], and Luttinger semimetal [90, 91], in which the Hamiltonian satisfies Eq. (87). We can find that the Hamiltonian of second-order TI as shown in Eq. (2) satisfies Eq. (87). Therefore, the generation of new type of disorder by one type of disorder should be carefully studied.

Third, generation of new type of disorder by two types of disorder was not considered in Ref. [73]. If two types of disorder exist initially, third type of disorder usually is generated. This effect could be quite important. For example, for 2D DSM, if any two types of disorder among random chemical potential, random vector potential, and random mass exist initially, the third one is generated

and the the system becomes compressible diffusive metal [80, 81]. In Ref. [73], they considered random chemical potential and random mass. Other types of disorder could be generated by random chemical potential and random mass. Whereas, this effect was not considered in Ref. [73].

In order to avoid the second and third problems above-mentioned, we find that twelve kinds of disorder should be considered for second-order TI. The corresponding action for fermion-disorder coupling reads

$$S_{\text{dis}} = \int d\tau d^3\mathbf{x} \bar{\Psi} \left[V_C(\mathbf{x})\gamma_0 + V_M(\mathbf{x})\mathbb{1}_{4\times 4} \right. \\ + V_{AC}(\mathbf{x})\gamma_0\gamma_5 + \sum_{j=x,y} V_{SO\perp}(\mathbf{x})\gamma_0\gamma_j \\ + V_{SOz}(\mathbf{x})\gamma_0\gamma_z + V_{PM}(\mathbf{x})i\gamma_5 \\ + V_{MN\perp}(\mathbf{x})(i\gamma_y\gamma_z + i\gamma_z\gamma_x) + V_{MNz}(\mathbf{x})i\gamma_x\gamma_y \\ \left. + V_{AMN\perp}(\mathbf{x}) \sum_{j=x,y} i\gamma_5\gamma_j + V_{AMNz}(\mathbf{x})i\gamma_5\gamma_z \right] \Psi.$$

$$+ V_{CR\perp}(\mathbf{x}) \sum_{j=x,y} i\gamma_j + V_{CRz}(\mathbf{x})i\gamma_z \Big] \Psi. \quad (88)$$

$V_C(\mathbf{x})$ corresponds to random chemical potential. $V_M(\mathbf{x})$ stands for random mass. $V_{AC}(\mathbf{x})$ represents random axial chemical potential. $V_{SO\perp}$ and V_{SOz} correspond to random spin-orbital scattering within xy plane and along z axis. $V_{PM}(\mathbf{x})$ stands for random pseudo mass. $V_{MN\perp}$ and V_{MNz} represent random magnetization within xy plane and along z axis. $V_{AMN\perp}$ and V_{AMNz} are corresponding to random axial magnetization within xy plane and along z axis. $V_{CR\perp}$ and V_{CRz} stand for random current within xy plane and along z axis.

The quenched random field V_j is taken as a Gaussian white noise distribution that satisfies $\langle V_j(\mathbf{x}) \rangle = 0$ and $\langle V_j(\mathbf{x})V_j(\mathbf{x}') \rangle = \Delta_j \delta^3(\mathbf{x} - \mathbf{x}')$. The disorder can be treated by the replica method. The effective action in replica formalism can be written as

$$S_{\text{dis}} = -\frac{\Delta_C}{2} \int d\tau d\tau' d^3\mathbf{x} (\bar{\Psi}_a\gamma_0\Psi_a)_\tau (\bar{\Psi}_b\gamma_0\Psi_b)_{\tau'} - \frac{\Delta_M}{2} \int d\tau d\tau' d^3\mathbf{x} (\bar{\Psi}_a\Psi_a)_\tau (\bar{\Psi}_b\Psi_b)_{\tau'} \\ - \frac{\Delta_{AC}}{2} \int d\tau d\tau' d^3\mathbf{x} (\bar{\Psi}_a\gamma_0\gamma_5\Psi_a)_\tau (\bar{\Psi}_b\gamma_0\gamma_5\Psi_b)_{\tau'} - \frac{\Delta_{SO\perp}}{2} \sum_{j=x,y} \int d\tau d\tau' d^3\mathbf{x} (\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)_\tau (\bar{\Psi}_b\gamma_0\gamma_j\Psi_b)_{\tau'} \\ - \frac{\Delta_{SOz}}{2} \int d\tau d\tau' d^3\mathbf{x} (\bar{\Psi}_a\gamma_0\gamma_z\Psi_a)_\tau (\bar{\Psi}_b\gamma_0\gamma_z\Psi_b)_{\tau'} - \frac{\Delta_{PM}}{2} \int d\tau d\tau' d^3\mathbf{x} (\bar{\Psi}_ai\gamma_5\Psi_a)_\tau (\bar{\Psi}_bi\gamma_5\Psi_b)_{\tau'} \\ - \frac{\Delta_{MN\perp}}{2} \sum_{j=x,y} \int d\tau d\tau' d^3\mathbf{x} [(\bar{\Psi}_ai\gamma_y\gamma_z\Psi_a)_\tau (\bar{\Psi}_bi\gamma_y\gamma_z\Psi_b)_{\tau'} + (\bar{\Psi}_ai\gamma_z\gamma_x\Psi_a)_\tau (\bar{\Psi}_bi\gamma_z\gamma_x\Psi_b)_{\tau'}] \\ - \frac{\Delta_{MNz}}{2} \int d\tau d\tau' d^3\mathbf{x} (\bar{\Psi}_ai\gamma_x\gamma_y\Psi_a)_\tau (\bar{\Psi}_bi\gamma_x\gamma_y\Psi_b)_{\tau'} - \frac{\Delta_{AMN\perp}}{2} \sum_{j=x,y} \int d\tau d\tau' d^3\mathbf{x} (\bar{\Psi}_ai\gamma_5\gamma_j\Psi_a)_\tau \\ \times (\bar{\Psi}_bi\gamma_5\gamma_j\Psi_b)_{\tau'} - \frac{\Delta_{AMNz}}{2} \int d\tau d\tau' d^3\mathbf{x} (\bar{\Psi}_ai\gamma_5\gamma_z\Psi_a)_\tau (\bar{\Psi}_bi\gamma_5\gamma_z\Psi_b)_{\tau'} \\ - \frac{\Delta_{CR\perp}}{2} \sum_{j=x,y} \int d\tau d\tau' d^3\mathbf{x} (\bar{\Psi}_ai\gamma_j\Psi_a)_\tau (\bar{\Psi}_bi\gamma_j\Psi_b)_{\tau'} - \frac{\Delta_{CRz}}{2} \int d\tau d\tau' d^3\mathbf{x} (\bar{\Psi}_ai\gamma_z\Psi_a)_\tau (\bar{\Psi}_bi\gamma_z\Psi_b)_{\tau'}, \quad (89)$$

with $a, b = 1, 2, \dots, n$ being the replica indices. At the end of the calculation, the limit $n \rightarrow 0$ will be taken.

Through the detailed calculations shown in the Appendices, we get the RG equations as following

$$\frac{dv}{d\ell} = -C_0^{\text{dis}}v, \quad (90)$$

$$\frac{dv_z}{d\ell} = -C_0^{\text{dis}}v_z, \quad (91)$$

$$\frac{dm}{d\ell} = m + C_{\text{dis}}^m m, \quad (92)$$

$$\frac{dB_\perp}{d\ell} = -B_\perp, \quad (93)$$

$$\frac{dB_z}{d\ell} = -B_z, \quad (94)$$

$$\frac{dD}{d\ell} = -D, \quad (95)$$

$$\frac{d\Delta_C}{d\ell} = -\Delta_C + 2\Delta_C (\Delta_C + \Delta_M + \Delta_{AC} + 2\Delta_{SO\perp} + \Delta_{SOz} + \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} + 2\Delta_{AMN\perp}$$

$$\begin{aligned}
& +\Delta_{AMNz} + 2\Delta_{CR\perp} + \Delta_{CRz}) (\mathcal{G}_1^\perp + \mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 4 [(\Delta_M \Delta_{SO\perp} + \Delta_{PM} \Delta_{MN\perp} + \Delta_{AMN\perp} \Delta_{CRz} + \Delta_{AMNz} \Delta_{CR\perp}) \mathcal{G}_1^\perp + (\Delta_M \Delta_{SOz} + \Delta_{PM} \Delta_{MNz} \\
& + 2\Delta_{AMN\perp} \Delta_{CR\perp}) \mathcal{G}_1^z + (\Delta_C \Delta_{PM} + 2\Delta_{SO\perp} \Delta_{AMN\perp} + \Delta_{SOz} \Delta_{AMNz}) \mathcal{G}_1^D \\
& + (\Delta_C \Delta_M + 2\Delta_{MN\perp} \Delta_{AMN\perp} + \Delta_{MNz} \Delta_{AMNz}) \mathcal{G}_1^m], \tag{96}
\end{aligned}$$

$$\begin{aligned}
\frac{d\Delta_M}{d\ell} = & -\Delta_M + 2\Delta_M (\Delta_C + \Delta_M - \Delta_{AC} - 2\Delta_{SO\perp} - \Delta_{SOz} - \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} + 2\Delta_{AMN\perp} \\
& + \Delta_{AMNz} - 2\Delta_{CR\perp} - 2\Delta_{CRz}) (-\mathcal{G}_1^\perp - \mathcal{G}_1^z - \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 4 [(\Delta_C \Delta_{SO\perp} + \Delta_{PM} \Delta_{AMN\perp} + \Delta_{MN\perp} \Delta_{CRz} + \Delta_{MNz} \Delta_{CR\perp}) \mathcal{G}_1^\perp + (\Delta_C \Delta_{SOz} + \Delta_{PM} \Delta_{AMNz} \\
& + 2\Delta_{MN\perp} \Delta_{CR\perp}) \mathcal{G}_1^z + (\Delta_M \Delta_{PM} + 2\Delta_{SO\perp} \Delta_{MN\perp} + \Delta_{SOz} \Delta_{MNz}) \mathcal{G}_1^D \\
& + (\Delta_C^2 + \Delta_M^2 + \Delta_{AC}^2 + 2\Delta_{SO\perp}^2 + \Delta_{SOz}^2 + \Delta_{PM}^2 + 2\Delta_{MN\perp}^2 + \Delta_{MNz}^2 + 2\Delta_{AMN\perp}^2 + \Delta_{AMNz}^2 \\
& + 2\Delta_{CR\perp}^2 + \Delta_{CRz}^2) \mathcal{G}_1^m], \tag{97}
\end{aligned}$$

$$\begin{aligned}
\frac{d\Delta_{AC}}{d\ell} = & -\Delta_{AC} + 2\Delta_{AC} (-\Delta_C + \Delta_M - \Delta_{AC} + 2\Delta_{SO\perp} + \Delta_{SOz} + \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} \\
& - 2\Delta_{AMN\perp} - \Delta_{AMNz} - 2\Delta_{CR\perp} - \Delta_{CRz}) (-\mathcal{G}_1^\perp - \mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 4 [(\Delta_{SO\perp} \Delta_{SOz} + \Delta_{MN\perp} \Delta_{MNz} + \Delta_{AMN\perp} \Delta_{AMNz} + \Delta_{CR\perp} \Delta_{CRz}) \mathcal{G}_1^\perp \\
& + 2(\Delta_{SO\perp}^2 + \Delta_{MN\perp}^2 + \Delta_{AMN\perp}^2 + \Delta_{CR\perp}^2) \mathcal{G}_1^z + (\Delta_{AC} \Delta_{PM} + 2\Delta_{SO\perp} \Delta_{CR\perp} + \Delta_{SOz} \Delta_{CRz}) \mathcal{G}_1^D \\
& + (\Delta_M \Delta_{AC} + \Delta_{MNz} \Delta_{CRz} + 2\Delta_{MN\perp} \Delta_{CR\perp}) \mathcal{G}_1^m], \tag{98}
\end{aligned}$$

$$\begin{aligned}
\frac{d\Delta_{SO\perp}}{d\ell} = & -\Delta_{SO\perp} + 2\Delta_{SO\perp} (-\Delta_C + \Delta_M + \Delta_{AC} + \Delta_{SOz} - \Delta_{PM} - \Delta_{MNz} + \Delta_{AMNz} - \Delta_{CRz}) \\
& \times (-\mathcal{G}_1^z - \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 2 [(\Delta_C \Delta_M + \Delta_{AC} \Delta_{SOz} + \Delta_{PM} \Delta_{CRz} + 4\Delta_{MN\perp} \Delta_{AMN\perp} + \Delta_{MNz} \Delta_{AMNz}) \mathcal{G}_1^\perp \\
& + 2(\Delta_{AC} \Delta_{SO\perp} + \Delta_{PM} \Delta_{CR\perp} + \Delta_{MN\perp} \Delta_{AMNz} + \Delta_{MNz} \Delta_{AMN\perp}) \mathcal{G}_1^z \\
& + 2(\Delta_C \Delta_{AMN\perp} + \Delta_M \Delta_{MN\perp} + \Delta_{AC} \Delta_{CR\perp} + \Delta_{SO\perp} \Delta_{PM}) \mathcal{G}_1^D \\
& + 2(\Delta_M \Delta_{SO\perp} + \Delta_{PM} \Delta_{MN\perp} + \Delta_{AMN\perp} \Delta_{CRz} + \Delta_{AMNz} \Delta_{CR\perp}) \mathcal{G}_1^m], \tag{99}
\end{aligned}$$

$$\begin{aligned}
\frac{d\Delta_{SOz}}{d\ell} = & -\Delta_{SOz} + 2\Delta_{SOz} (-\Delta_C + \Delta_M + \Delta_{AC} + 2\Delta_{SO\perp} - \Delta_{SOz} - \Delta_{PM} - 2\Delta_{MN\perp} + \Delta_{MNz} \\
& + 2\Delta_{AMN\perp} - \Delta_{AMNz} - 2\Delta_{CR\perp} + \Delta_{CRz}) (-\mathcal{G}_1^\perp + \mathcal{G}_1^z - \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 4 [(\Delta_{AC} \Delta_{SO\perp} + \Delta_{PM} \Delta_{CR\perp} + \Delta_{MN\perp} \Delta_{AMNz} + \Delta_{MNz} \Delta_{AMN\perp}) \mathcal{G}_1^\perp \\
& + (\Delta_C \Delta_M + 2\Delta_{MN\perp} \Delta_{AMN\perp} + \Delta_{MNz} \Delta_{AMNz}) \mathcal{G}_1^z \\
& + (\Delta_C \Delta_{AMNz} + \Delta_M \Delta_{MNz} + \Delta_{AC} \Delta_{CRz} + \Delta_{SOz} \Delta_{PM}) \mathcal{G}_1^D \\
& + (\Delta_M \Delta_{SOz} + \Delta_{PM} \Delta_{MNz} + 2\Delta_{AMN\perp} \Delta_{CR\perp}) \mathcal{G}_1^m], \tag{100}
\end{aligned}$$

$$\begin{aligned}
\frac{d\Delta_{PM}}{d\ell} = & -\Delta_{PM} + 2\Delta_{PM} (-\Delta_C + \Delta_M + \Delta_{AC} - 2\Delta_{SO\perp} - \Delta_{SOz} - \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} \\
& - 2\Delta_{AMN\perp} - \Delta_{AMNz} + 2\Delta_{CR\perp} + \Delta_{CRz}) (\mathcal{G}_1^\perp + \mathcal{G}_1^z - \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 4 [(\Delta_C \Delta_{MN\perp} + \Delta_M \Delta_{AMN\perp} + \Delta_{SO\perp} \Delta_{CRz} + \Delta_{SOz} \Delta_{CR\perp}) \mathcal{G}_1^\perp \\
& + (\Delta_C \Delta_{MNz} + \Delta_M \Delta_{AMNz} + 2\Delta_{SO\perp} \Delta_{CR\perp}) \mathcal{G}_1^z + (\Delta_C^2 + \Delta_M^2 + \Delta_{AC}^2 + 2\Delta_{SO\perp}^2 + \Delta_{SOz}^2 \\
& + \Delta_{PM}^2 + 2\Delta_{MN\perp}^2 + \Delta_{MNz}^2 + 2\Delta_{AMN\perp}^2 + \Delta_{AMNz}^2 + 2\Delta_{CR\perp}^2 + \Delta_{CRz}^2) \mathcal{G}_1^D \\
& + (\Delta_M \Delta_{PM} + 2\Delta_{SO\perp} \Delta_{MN\perp} + \Delta_{SOz} \Delta_{MNz}) \mathcal{G}_1^m], \tag{101}
\end{aligned}$$

$$\begin{aligned}
\frac{d\Delta_{MN\perp}}{d\ell} = & -\Delta_{MN\perp} + 2\Delta_{MN\perp} (\Delta_C + \Delta_M - \Delta_{AC} + \Delta_{SOz} - \Delta_{PM} - \Delta_{MNz} - \Delta_{AMNz} + \Delta_{CRz}) \\
& \times (\mathcal{G}_1^z - \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 2 [(\Delta_C \Delta_{PM} + \Delta_M \Delta_{CRz} + \Delta_{AC} \Delta_{MNz} + 4\Delta_{SO\perp} \Delta_{AMN\perp} + \Delta_{SOz} \Delta_{AMNz}) \mathcal{G}_1^\perp \\
& + 2(\Delta_M \Delta_{CR\perp} + \Delta_{AC} \Delta_{MN\perp} + \Delta_{SO\perp} \Delta_{AMNz} + \Delta_{SOz} \Delta_{AMN\perp}) \mathcal{G}_1^z \\
& + 2(\Delta_M \Delta_{SO\perp} + \Delta_{PM} \Delta_{MN\perp} + \Delta_{AMN\perp} \Delta_{CRz} + \Delta_{AMNz} \Delta_{CR\perp}) \mathcal{G}_1^D \\
& + 2(\Delta_C \Delta_{AMN\perp} + \Delta_M \Delta_{MN\perp} + \Delta_{AC} \Delta_{CR\perp} + \Delta_{SO\perp} \Delta_{PM}) \mathcal{G}_1^m], \tag{102}
\end{aligned}$$

$$\begin{aligned}
\frac{d\Delta_{MNz}}{d\ell} = & -\Delta_{MNz} + 2\Delta_{MNz}(\Delta_C + \Delta_M - \Delta_{AC} + 2\Delta_{SO\perp} - \Delta_{SOz} - \Delta_{PM} - 2\Delta_{MN\perp} + \Delta_{MNz} \\
& - 2\Delta_{AMN\perp} + \Delta_{AMNz} + 2\Delta_{CR\perp} - \Delta_{CRz})(\mathcal{G}_1^\perp - \mathcal{G}_1^z - \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 4[(\Delta_M\Delta_{CR\perp} + \Delta_{AC}\Delta_{MN\perp} + \Delta_{SO\perp}\Delta_{AMNz} + \Delta_{SOz}\Delta_{AMN\perp})\mathcal{G}_1^\perp \\
& + (\Delta_C\Delta_{PM} + 2\Delta_{SO\perp}\Delta_{AMN\perp} + \Delta_{SOz}\Delta_{AMNz})\mathcal{G}_1^z \\
& + (\Delta_M\Delta_{SOz} + \Delta_{PM}\Delta_{MNz} + 2\Delta_{AMN\perp}\Delta_{CR\perp})\mathcal{G}_1^D \\
& + (\Delta_C\Delta_{AMNz} + \Delta_M\Delta_{MNz} + \Delta_{AC}\Delta_{CRz} + \Delta_{SOz}\Delta_{PM})\mathcal{G}_1^m], \tag{103}
\end{aligned}$$

$$\begin{aligned}
\frac{d\Delta_{AMN\perp}}{d\ell} = & -\Delta_{AMN\perp} + 2\Delta_{AMN\perp}(\Delta_C + \Delta_M + \Delta_{AC} - \Delta_{SOz} + \Delta_{PM} - \Delta_{MNz} - \Delta_{AMNz} - \Delta_{CRz}) \\
& \times (-\mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 2[(\Delta_C\Delta_{CRz} + \Delta_M\Delta_{PM} + \Delta_{AC}\Delta_{AMNz} + 4\Delta_{SO\perp}\Delta_{MN\perp} + \Delta_{SOz}\Delta_{MNz})\mathcal{G}_1^\perp \\
& + 2(\Delta_C\Delta_{CR\perp} + \Delta_{AC}\Delta_{AMN\perp} + \Delta_{SO\perp}\Delta_{MNz} + \Delta_{SOz}\Delta_{MN\perp})\mathcal{G}_1^z \\
& + 2(\Delta_C\Delta_{SO\perp} + \Delta_{PM}\Delta_{AMN\perp} + \Delta_{MN\perp}\Delta_{CRz} + \Delta_{MNz}\Delta_{CR\perp})\mathcal{G}_1^D \\
& + 2(\Delta_C\Delta_{MN\perp} + \Delta_M\Delta_{AMN\perp} + \Delta_{SO\perp}\Delta_{CRz} + \Delta_{SOz}\Delta_{CR\perp})\mathcal{G}_1^m], \tag{104}
\end{aligned}$$

$$\begin{aligned}
\frac{d\Delta_{AMNz}}{d\ell} = & -\Delta_{AMNz} + 2\Delta_{AMNz}(\Delta_C + \Delta_M + \Delta_{AC} - 2\Delta_{SO\perp} + \Delta_{SOz} + \Delta_{PM} - 2\Delta_{MN\perp} + \Delta_{MNz} \\
& - 2\Delta_{AMN\perp} + \Delta_{AMNz} - 2\Delta_{CR\perp} + \Delta_{CRz})(-\mathcal{G}_1^\perp + \mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 4[(\Delta_C\Delta_{CR\perp} + \Delta_{AC}\Delta_{AMN\perp} + \Delta_{SO\perp}\Delta_{MNz} + \Delta_{SOz}\Delta_{MN\perp})\mathcal{G}_1^\perp \\
& + (\Delta_M\Delta_{PM} + 2\Delta_{SO\perp}\Delta_{MN\perp} + \Delta_{SOz}\Delta_{MNz})\mathcal{G}_1^z \\
& + (\Delta_C\Delta_{SOz} + \Delta_{PM}\Delta_{AMNz} + 2\Delta_{MN\perp}\Delta_{CR\perp})\mathcal{G}_1^D \\
& + (\Delta_C\Delta_{MNz} + \Delta_M\Delta_{AMNz} + 2\Delta_{SO\perp}\Delta_{CR\perp})\mathcal{G}_1^m], \tag{105}
\end{aligned}$$

$$\begin{aligned}
\frac{d\Delta_{CR\perp}}{d\ell} = & -\Delta_{CR\perp} + 2\Delta_{CR\perp}(-\Delta_C + \Delta_M - \Delta_{AC} - \Delta_{SOz} + \Delta_{PM} - \Delta_{MNz} + \Delta_{AMNz} + \Delta_{CRz}) \\
& \times (\mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 2[(\Delta_C\Delta_{AMNz} + \Delta_M\Delta_{MNz} + \Delta_{AC}\Delta_{CRz} + \Delta_{SOz}\Delta_{PM})\mathcal{G}_1^\perp \\
& + 2(\Delta_C\Delta_{AMN\perp} + \Delta_M\Delta_{MN\perp} + \Delta_{AC}\Delta_{CR\perp} + \Delta_{SO\perp}\Delta_{PM})\mathcal{G}_1^z \\
& + 2(\Delta_{AC}\Delta_{SO\perp} + \Delta_{PM}\Delta_{CR\perp} + \Delta_{MN\perp}\Delta_{AMNz} + \Delta_{MNz}\Delta_{AMN\perp})\mathcal{G}_1^D \\
& + 2(\Delta_M\Delta_{CR\perp} + \Delta_{AC}\Delta_{MN\perp} + \Delta_{SO\perp}\Delta_{AMNz} + \Delta_{SOz}\Delta_{AMN\perp})\mathcal{G}_1^m], \tag{106}
\end{aligned}$$

$$\begin{aligned}
\frac{d\Delta_{CRz}}{d\ell} = & -\Delta_{CRz} + 2\Delta_{CRz}(-\Delta_C + \Delta_M - \Delta_{AC} - 2\Delta_{SO\perp} + \Delta_{SOz} + \Delta_{PM} - 2\Delta_{MN\perp} + \Delta_{MNz} \\
& + 2\Delta_{AMN\perp} - \Delta_{AMNz} + 2\Delta_{CR\perp} - \Delta_{CRz})(\mathcal{G}_1^\perp - \mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 4[(\Delta_C\Delta_{AMN\perp} + \Delta_M\Delta_{MN\perp} + \Delta_{AC}\Delta_{CR\perp} + \Delta_{SO\perp}\Delta_{PM})\mathcal{G}_1^\perp \\
& + (\Delta_{AC}\Delta_{SOz} + \Delta_{PM}\Delta_{CRz} + 2\Delta_{MN\perp}\Delta_{AMN\perp})\mathcal{G}_1^D \\
& + (\Delta_M\Delta_{CRz} + \Delta_{AC}\Delta_{MNz} + 2\Delta_{SO\perp}\Delta_{AMN\perp})\mathcal{G}_1^m], \tag{107}
\end{aligned}$$

The transformations

$$\Delta_j \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \rightarrow \Delta_j \tag{108}$$

have been used. The expressions of C_0^{dis} , C_m^{dis} , $\mathcal{G}_1^\perp$, $\mathcal{G}_1^z$, $\mathcal{G}_1^D$, and $\mathcal{G}_1^m$ can be found in Appendix F.

It is easy to find that

$$B_\perp(\ell) = B_{\perp,0}e^{-\ell}, \tag{109}$$

$$B_z(\ell) = B_{z,0}e^{-\ell}, \tag{110}$$

$$D(\ell) = D_0e^{-\ell}, \tag{111}$$

which take the same behaviors as the clean second-order TI.

Considering random chemical potential initially with different initial values Δ_{C0} , the flows of different disorder coupling parameters are shown in Figs. 6(a)-6(l). The flows of $m(\ell)$ and $m(\ell)/m^f(\ell)$ are depicted in Figs. 6(m) and 6(n). If Δ_{C0} is small, $\Delta_C(\ell)$ approaches to zero in the lowest energy limit $\ell \rightarrow \infty$, other disorder coupling parameters Δ_M , Δ_{AC} , $\Delta_{SO\perp}$, Δ_{SOz} , Δ_{PM} , $\Delta_{MN\perp}$, Δ_{MNz} , $\Delta_{AMN\perp}$, Δ_{AMNz} , $\Delta_{CR\perp}$, and Δ_{CRz} increase from zero at beginning and also approach to zero in the lowest energy limit. In this case, as shown in Fig. 6(m), $m(\ell) \rightarrow \infty$ in the lowest energy limit.

In Ref. [73], the behaviors $D(\ell) \rightarrow 0$ and $m(\ell) \rightarrow \infty$ in the lowest energy limit are used as the criterion for




FIG. 6: (a)-(m): Flows of parameters $\Delta_C(\ell)$, $\Delta_M(\ell)$, $\Delta_{AC}(\ell)$, $\Delta_{SO\perp}(\ell)$, $\Delta_{SOz}(\ell)$, $\Delta_{PM}(\ell)$, $\Delta_{MN\perp}(\ell)$, $\Delta_{MNz}(\ell)$, $\Delta_{AMN\perp}(\ell)$, $\Delta_{AMNz}(\ell)$, $\Delta_{CR\perp}(\ell)$, $\Delta_{CRz}(\ell)$, $m(\ell)$ and $m(\ell)/m^f(\ell)$. Blue, red, green, black, and magenta lines correspond to the initial values $\Delta_{C0} = 0.05, 0.1, 0.2, 0.5, 1.0$ respectively. Initial values of other disorder parameters are taken to be zero. $m_0 = 0.1$, $B_{\perp 0} = 1$, $B_{z0} = 1$, $D_0 = 1$ are taken.

the transition from second-order TI to TI. According to their criterion, weak disorder drives second-order TI to TI. However, recent studies based on other methods showed that second-order TI is robust against weak disorder [24, 25]. This contradiction is just due to that the criterion used in Ref. [73] is invalid.

As depicted in Fig. 6(m), $m(\ell)/m^f(\ell)$ approaches to a positive constant value in the lowest energy limit if Δ_{C0} is small. This indicates that $m(\ell)$ takes the qualitatively same behavior as the clean second-order TI. Additionally, $B_{\perp}(\ell)$, $B_z(\ell)$, $D(\ell)$ take the same behaviors as clean second-order TI. These behaviors clearly reflect that weak disorder does not induce qualitative modification for second-order TI. Namely, second-order TI is robust against weak disorder.

If Δ_{C0} is large enough, we can find that Δ_C approaches to infinity at a finite running parameter ℓ_c . Δ_M , Δ_{AC} , $\Delta_{SO\perp}$, Δ_{SOz} , Δ_{PM} , $\Delta_{MN\perp}$, Δ_{MNz} , $\Delta_{AMN\perp}$, Δ_{AMNz} , $\Delta_{CR\perp}$, and Δ_{CRz} grow from zero and also flow to infinity at ℓ_c . As shown in Figs. 6(m) and 6(n), we notice that $m(\ell)$ approaches to finite value when $\ell \rightarrow \ell_c$. These behaviors represent that second-order TI is driven to a diffusive metal phase.

Considering random mass initially with different initial values Δ_{M0} , the flows of disorder coupling parameters Δ_M , Δ_{AC} , Δ_{PM} , $\Delta_{AMN\perp}$, and Δ_{AMNz} are displayed in Figs. 7(a)-7(e). Δ_C , $\Delta_{SO\perp}$, Δ_{SOz} , $\Delta_{MN\perp}$, Δ_{MNz} , $\Delta_{CR\perp}$, Δ_{CRz} are not generated, and equal to the initial

value zero. Thus, the flows of these seven disorder coupling parameters are not shown in Fig. 7. The flow of $m(\ell)/m^f(\ell)$ is shown in 7(f). If Δ_{M0} is small enough, Δ_M flows to zero in the lowest energy limit $\ell \rightarrow \infty$. Δ_{AC} , Δ_{PM} , $\Delta_{AMN\perp}$, Δ_{AMNz} increase from zero at the beginning, but become to decrease when ℓ is large enough and finally approach to zero in the lowest energy limit. $m(\ell)/m^f(\ell)$ approaches to a positive constant which is greater than 1. It represents that disorder only induces quantitative increment but does not result in qualitative modification for m . Thus, second-order TI is stable in this case.

If Δ_{M0} is large enough, Δ_M flows to infinity when ℓ approaches to a critical value ℓ_c . Δ_M , Δ_{AC} , Δ_{PM} , $\Delta_{AMN\perp}$, and Δ_{AMNz} are generated from zero and finally also approach to infinity when $\ell \rightarrow \ell_c$. $m(\ell)/m^f(\ell)$ flows to a finite value in the limit $\ell \rightarrow \ell_c$. These behaviors should represent that the system is driven to a diffusive metal state. Whereas, it was shown that the system can not be driven to diffusive metal phase even Δ_{M0} is large in Ref. [73]. This is due to that the generation of new types of disorder by Δ_M is not considered in Ref. [73]

In proper initial conditions, we notice that the second-order TI is driven to trivial band insulator with intermediate disorder strength. For example, considering random axial chemical initially with different values of Δ_{AC0} , the flows of Δ_M , Δ_{AC} , Δ_{PM} , $\Delta_{AMN\perp}$, and Δ_{AMNz} are shown in Figs. 8(a)-8(e). If Δ_{AC0} is




FIG. 7: (a)-(f): Flows of parameters $\Delta_M(\ell)$, $\Delta_{AC}(\ell)$, $\Delta_{PM}(\ell)$, $\Delta_{AMN\perp}(\ell)$, $\Delta_{AMNz}(\ell)$, and $m(\ell)/m^f(\ell)$. Blue, red, green, black, and magenta lines correspond to the initial values $\Delta_{M0} = 0.05, 0.1, 0.2, 0.5, 1.0$ respectively. Initial values of other disorder parameters are taken to be zero. $m_0 = 0.1$, $B_{\perp 0} = 1$, $B_{z0} = 1$, $D_0 = 1$ are taken.




FIG. 8: (a)-(f): Flows of parameters $\Delta_M(\ell)$, $\Delta_{AC}(\ell)$, $\Delta_{PM}(\ell)$, $\Delta_{AMN\perp}(\ell)$, $\Delta_{AMNz}(\ell)$, and $m(\ell)/m^f(\ell)$. Blue, red, green, black, and magenta lines correspond to the initial values $\Delta_{AC0} = 0.05, 0.1, 0.2, 0.5, 1.0$ respectively. Initial values of other disorder parameters are taken to be zero. $m_0 = 0.1$, $B_{\perp 0} = 1$, $B_{z0} = 1$, $D_0 = 1$ are taken.

small, the disorder coupling parameters Δ_M , Δ_{AC} , Δ_{PM} , $\Delta_{AMN\perp}$, and Δ_{AMNz} all flow to zero in the lowest energy limit $\ell \rightarrow \infty$. According to Fig. 8(f), $m(\ell)/m^f(\ell)$ flows to a positive constant value which is smaller than one. It represents that m only decreases quantitatively but not receive qualitative modification. Thus, the system is still in second-order TI phase with small Δ_{AC0} . If Δ_{AC0} is large enough, Δ_M , Δ_{AC} , Δ_{PM} , $\Delta_{AMN\perp}$, and Δ_{AMNz} all approach to infinity at a finite running




FIG. 9: (a)-(f): Flows of parameters $\alpha(\ell)$, $\Delta_C(\ell)$, $\Delta_C(\ell)/\alpha(\ell)$, $D(\ell)$, $m(\ell)/m^f(\ell)$, and $B_{\perp}(\ell)/(\alpha(\ell)/3)$ initially considering both of long-range Coulomb interaction and random chemical potential. Blue, red, green, black, and magenta lines correspond to the initial values $\alpha_0 = 0.1, 0.2, 0.3, 0.4, 0.5$ respectively. $\Delta_{C0} = 0.1$, and initial values of other disorder parameters are taken to be zero. $m_0 = 0.1$, $B_{\perp 0} = 1$, $B_{z0} = 1$, $D_0 = 1$ are taken.

parameter ℓ_c . $m(\ell)/m^f(\ell)$ approaches to finite value at same ℓ_c . In this case, the system is driven to diffusive metal phase. If Δ_{AC0} takes intermediate values, Δ_M , Δ_{AC} , Δ_{PM} , $\Delta_{AMN\perp}$, and Δ_{AMNz} all flow to zero in the lowest energy limit $\ell \rightarrow \infty$. Whereas, $m(\ell)/m^f(\ell)$ flows to a negative constant value. It represents that the system becomes a trivial band insulator.

VI. INTERPLAY OF COULOMB INTERACTION AND DISORDER

In this section, we study the interplay of long-range Coulomb interaction and disorder. The RG equations for the model parameters considering both of Coulomb interaction and disorder can be found in Appendix F. Considering initially both of long-range Coulomb interaction and random chemical potential, we show the flows of $\alpha(\ell)$, $\Delta_C(\ell)$, $\Delta_C(\ell)/\alpha(\ell)$, $D(\ell)$, $m(\ell)/m^f(\ell)$, and $B_{\perp}(\ell)/(\alpha(\ell)/3)$ in Figs. 9 and 10 with different initial conditions. The flow of $B_z(\ell)/(\alpha(\ell)/3)$ is not shown since it take the qualitatively same behavior as $B_{\perp}(\ell)/(\alpha(\ell)/3)$. If Δ_{C0} and α_0 both are small, $\alpha(\ell)$ flows to a constant value α^* , $\Delta_C(\ell)$ approaches to zero, $D(\ell)/D^f(\ell)$ approaches to a positive constant value, $m(\ell)/m^f(\ell)$ flows to a positive constant value, and $B_{\perp}(\ell)/(\alpha(\ell)/3) \rightarrow -1$ in the lowest energy limit. These behaviors represent that second-order TI is driven to trivial band insulator. For a given Δ_{C0} , if α_0 is large enough, we can find that $\alpha(\ell)$ flows to a constant value α^* , $\Delta_C(\ell)$ approaches to zero, $D(\ell)/D^f(\ell)$ approaches to a positive constant value, $m(\ell)/m^f(\ell)$ flows to a nega-




FIG. 10: (a)-(f): Flows of parameters $\alpha(\ell)$, $\Delta_C(\ell)$, $\Delta_C(\ell)/\alpha(\ell)$, $D(\ell)$, $m(\ell)/m^f(\ell)$, and $B_\perp(\ell)/(\alpha(\ell)/3)$ initially considering both of long-range Coulomb interaction and random chemical potential. Blue, red, green, black, and magenta lines correspond to the initial values $\Delta_{C0} = 0.05, 0.1, 0.2, 0.5, 1.0$ respectively. Initial values of other disorder parameters are taken to be zero. $\alpha_0 = 0.2$, $m_0 = 0.1$, $B_{\perp 0} = 1$, $B_{z0} = 1$, $D_0 = 1$ are taken.

tive constant value, and $B_\perp(\ell)/(\alpha(\ell)/3) \rightarrow 1$ in the limit $\ell \rightarrow \infty$. These results indicate that second-order TI becomes to trivial band insulator. For a given α_0 , if Δ_{C0} is large enough, we notice that $\alpha(\ell) \rightarrow \infty$, $\Delta_C(\ell) \rightarrow \infty$, $\Delta_C(\ell)/\alpha(\ell) \rightarrow \infty$, $D(\ell)/D^f(\ell)$, $m(\ell)/m^f(\ell)$, and $B_\perp(\ell)$ flow to a finite values at a finite running parameter ℓ_c . These behaviors represent that second-order TI is driven to diffusive metal phase.

VII. SUMMARY

In this article, we study the influence of Coulomb interaction on second-order TI by RG theory. We show that both the analysis method and conclusions in recent studies in Ref. [73] are unreliable. First, the criterion for transition from second-order TI to TI adopted in Ref. [73] is invalid. Second, the flow of B is important, but is not paid attention in Ref. [73]. Through analyzing the corrections for flows of model parameters induced by Coulomb interaction, we find that second-order TI is unstable to trivial band insulator not only under strong Coulomb interaction but also under weak Coulomb interaction. We also study the effects of disorder in second-order TI. According to the criterion adopted in Ref. [73], weak disorder will drive second-order TI to TI. Whereas, we find that weak disorder does not result in qualitative modification for the Hamiltonian of second-order TI. It indicates that second-order TI is robust against weak disorder. This result is consistent with recent studies about disorder effects in second-order TI through other methods [24, 25]. We also obtain the behaviors of second-order

TI considering both of long-range Coulomb interaction and disorder.

For very strong Coulomb interaction, the particle-hole pairs may be formed, and second-order TI may become an excitonic insulator. In this article, we do not consider this possibility. This possibility could be studied by self-consistent Dyson-Schwinger equations [36, 53, 92, 93].

ACKNOWLEDGEMENTS

We acknowledge the support from the National Natural Science Foundation of China under Grants 11974356, and U1832209. A portion of this work was supported by the High Magnetic Field Laboratory of Anhui Province under Grant AHM-FX-2020-01.

Appendix A: The propagators of fermion and boson

The propagator of fermion is given by

$$G_0(k_0, \mathbf{k}) = \frac{1}{ik_0\gamma_0 + \mathcal{H}_f}, \quad (\text{A1})$$

where

$$\begin{aligned} \mathcal{H}_f = & i[v(k_x\gamma_x + k_y\gamma_y) + v_z\gamma_z + D(k_x^2 - k_y^2)\gamma_5] \\ & + m - B_\perp k_\perp^2 - B_z k_z^2, \end{aligned} \quad (\text{A2})$$

with $k_\perp^2 = k_x^2 + k_y^2$. The propagator of boson field ϕ can be written as

$$D_0(k_0, \mathbf{k}) = \frac{1}{k_\perp^2 + \eta k_z^2}. \quad (\text{A3})$$

Appendix B: The self-energy of fermion




FIG. 11: Self-energy of fermions due to (a) Coulomb interaction and (b) disorder. The solid line represents the fermion propagator, and the wavy line stands for the boson propagator that is equivalent to the Coulomb interaction function. The dashed line denotes disorder scattering.

1. The self-energy of fermion induced by Coulomb interaction

As shown in Fig. 11, the self-energy of fermion induced by Coulomb interaction is given by

$$\Sigma(k_0, \mathbf{k}) = -g^2 \int_{-\infty}^{+\infty} \frac{dq_0}{2\pi} \int' \frac{d^3\mathbf{q}}{(2\pi)^3} \gamma_0 G_0(q_0, \mathbf{q}) \gamma_0$$

$$\times D_0(k_0 - q_0, \mathbf{k} - \mathbf{q}). \quad (\text{B1})$$

$\int'$ represents that a proper momentum shell will be chosen. Substituting Eqs. (A1) and (A3) into Eq. (B1), to the leading order, we obtain

$$\begin{aligned} \Sigma(k_0, \mathbf{k}) = & -\frac{g^2}{2} \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{(m - B_\perp q_\perp^2 - B_z q_z^2)}{E_{\mathbf{q}}(q_\perp^2 + \eta q_z^2)} \\ & -ivk_x \gamma_x g^2 \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{q_x^2}{E_{\mathbf{q}}(q_\perp^2 + \eta q_z^2)^2} \\ & -ivk_y \gamma_y g^2 \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{q_y^2}{E_{\mathbf{q}}(q_\perp^2 + \eta q_z^2)^2} \\ & -ivk_z \gamma_z \eta g^2 \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{q_z^2}{E_{\mathbf{q}}(q_\perp^2 + \eta q_z^2)^2} \\ & + (k_x^2 + k_y^2) \frac{g^2}{2} \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{m - B_\perp q_\perp^2 - B_z q_z^2}{E_{\mathbf{q}}(q_\perp^2 + \eta q_z^2)^2} \\ & + k_z^2 \eta \frac{g^2}{2} \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{m - B_\perp q_\perp^2 - B_z q_z^2}{E_{\mathbf{q}}(q_\perp^2 + \eta q_z^2)^2} \\ & -iDk_x^2 \gamma_5 2g^2 \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{(q_x^2 - q_y^2) q_x^2}{E_{\mathbf{q}}(q_\perp^2 + \eta q_z^2)^3} \\ & -iDk_y^2 \gamma_5 2g^2 \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{(q_x^2 - q_y^2) q_y^2}{E_{\mathbf{q}}(q_\perp^2 + \eta q_z^2)^3} \\ & -k_x^2 2g^2 \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{[m - B_\perp q_\perp^2 - B_z q_z^2] q_x^2}{E_{\mathbf{q}}(q_\perp^2 + \eta q_z^2)^3} \\ & -k_y^2 2g^2 \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{[m - B_\perp q_\perp^2 - B_z q_z^2] q_y^2}{E_{\mathbf{q}}(q_\perp^2 + \eta q_z^2)^3} \\ & -k_z^2 2\eta^2 g^2 \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{[m - B_\perp q_\perp^2 - B_z q_z^2] q_z^2}{E_{\mathbf{q}}(q_\perp^2 + \eta q_z^2)^3}, \end{aligned} \quad (\text{B2})$$

where

$$\begin{aligned} E_{\mathbf{q}} = & \left[v^2 q_\perp^2 + v_z^2 q_z^2 + D^2 (q_x^2 - q_y^2)^2 \right. \\ & \left. + (m - B_\perp q_\perp^2 - B_z q_z^2)^2 \right]^{\frac{1}{2}}. \end{aligned} \quad (\text{B3})$$

Using the transformation

$$\mathbf{q}' = (q_x, q_y, \sqrt{\eta} q_z), \quad (\text{B4})$$

and employing the RG scheme

$$\int \frac{d^3 \mathbf{q}'}{(2\pi)^3} = \frac{1}{8\pi^3} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \int_{b\Lambda}^\Lambda dq' q'^2, \quad (\text{B5})$$

where $b = e^{-\ell}$ with ℓ being RG running parameter and Λ the momentum cutoff, we finally obtain

$$\begin{aligned} \Sigma(k_0, \mathbf{k}) = & -mC_m \ell - iv(k_x \gamma_x + k_y \gamma_y) C_v \ell \\ & -iv_z k_z \gamma_z C_{v_z} \ell - B_\perp (k_x^2 + k_y^2) C_{B_\perp} \ell \\ & -B_z k_z^2 C_{B_z} \ell - iD(k_x^2 - k_y^2) \gamma_5 C_D \ell, \end{aligned} \quad (\text{B6})$$

where

$$\begin{aligned} C_m = & \alpha \frac{v\Lambda}{m} \left[\frac{m}{v\Lambda} (\mathcal{F}_0^\perp + \mathcal{F}_0^z) - \frac{B_\perp \Lambda}{v} \mathcal{F}_0^\perp \right. \\ & \left. - \frac{B_z \Lambda}{v\eta} \mathcal{F}_0^z \right], \end{aligned} \quad (\text{B7})$$

$$C_v = \alpha \mathcal{F}_0^\perp, \quad (\text{B8})$$

$$C_{v_z} = 2\alpha \mathcal{F}_0^z, \quad (\text{B9})$$

$$\begin{aligned} C_{B_\perp} = & \alpha \frac{v}{B_\perp \Lambda} \left[\frac{m}{v\Lambda} (\mathcal{F}_1^\perp + \mathcal{F}_1^z) - \frac{B_\perp \Lambda}{v} \mathcal{F}_1^\perp \right. \\ & \left. - \frac{B_z \Lambda}{v\eta} \mathcal{F}_1^z \right], \end{aligned} \quad (\text{B10})$$

$$\begin{aligned} C_{B_z} = & \alpha \frac{\eta v}{B_z \Lambda} \left\{ \frac{m}{v\Lambda} [\mathcal{F}_0^\perp + \mathcal{F}_0^z - 2(\mathcal{F}_1^\perp + \mathcal{F}_1^z)] - \frac{B_\perp \Lambda}{v} \right. \\ & \left. \times (\mathcal{F}_0^\perp - 2\mathcal{F}_1^\perp) - \frac{B_z \Lambda}{v\eta} (\mathcal{F}_0^z - 2\mathcal{F}_1^z) \right\}, \end{aligned} \quad (\text{B11})$$

$$C_D = \alpha \mathcal{F}_1^D, \quad (\text{B12})$$

with

$$\alpha = \frac{g^2}{4\pi^2 v \sqrt{\eta}}. \quad (\text{B13})$$

$\mathcal{F}_0^\perp, \mathcal{F}_0^z, \mathcal{F}_1^\perp, \mathcal{F}_1^z, \mathcal{F}_1^D$ are given by

$$\mathcal{F}_0^\perp = \frac{1}{4\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \frac{\sin^2(\varphi)}{\Xi}, \quad (\text{B14})$$

$$\mathcal{F}_0^z = \frac{1}{4\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \frac{\cos^2(\varphi)}{\Xi}, \quad (\text{B15})$$

$$\mathcal{F}_1^\perp = \frac{1}{4\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \frac{-\sin^2(\varphi) \cos(2\varphi)}{\Xi}, \quad (\text{B16})$$

$$\mathcal{F}_1^z = \frac{1}{4\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \frac{-\cos^2(\varphi) \cos(2\varphi)}{\Xi}, \quad (\text{B17})$$

$$\mathcal{F}_1^D = \frac{1}{2\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \frac{\sin^4(\varphi) \cos^2(2\theta)}{\Xi}, \quad (\text{B18})$$

where

$$\begin{aligned} \Xi = & \left[\sin^2(\varphi) + \left(\frac{v_z}{v\sqrt{\eta}} \right)^2 \cos^2(\varphi) \right. \\ & + \frac{D^2}{v^2} \Lambda^2 \sin^4(\varphi) \cos^2(2\theta) \\ & \left. + \left(\frac{m}{v\Lambda} - \frac{B_\perp \Lambda}{v} \sin^2(\varphi) - \frac{B_z \Lambda}{v\eta} \cos^2(\varphi) \right)^2 \right]^{\frac{1}{2}} \end{aligned} \quad (\text{B19})$$

We should notice that there is a relation

$$\mathcal{F}_1^\perp + \mathcal{F}_1^z = \mathcal{F}_0^\perp - \mathcal{F}_0^z. \quad (\text{B20})$$

2. The self-energy of fermion induced by disorder scattering

As depicted in Fig. 11, the self-energy of fermions induced by disorder scattering is expressed by

$$\Sigma_{dis}(k_0) = \sum_j \Delta_j \int' \frac{d^3 \mathbf{k}}{(2\pi)^3} \Gamma_j G_0(k_0, \mathbf{k}) \Gamma_j. \quad (\text{B21})$$

Substituting Eq. (A1) into Eq. (B21), we obtain

$$\Sigma_{dis}(k_0) = -ik_0 \gamma_0 C_0^{dis} \ell - m C_m^{dis} \ell, \quad (\text{B22})$$

where

$$C_0^{dis} = (\Delta_C + \Delta_M + \Delta_{AC} + 2\Delta_{SO\perp} + \Delta_{SOz} + \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} + 2\Delta_{AMN\perp} + \Delta_{AMNz} + 2\Delta_{CR\perp} + \Delta_{CRz}) \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} (\mathcal{G}_0^\perp + \mathcal{G}_0^z), \quad (\text{B23})$$

$$C_m^{dis} = -(\Delta_C + \Delta_M - \Delta_{AC} - 2\Delta_{SO\perp} - \Delta_{SOz} - \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} + 2\Delta_{AMN\perp}$$

with

$$\mathcal{G}_0^\perp = \frac{1}{4\pi} \int_0^\pi d\varphi \sin(\varphi) \int_0^{2\pi} d\theta \frac{\sin^2(\varphi)}{\Xi^2}, \quad (\text{B25})$$

$$\mathcal{G}_0^z = \frac{1}{4\pi} \int_0^\pi d\varphi \sin(\varphi) \int_0^{2\pi} d\theta \frac{\cos^2(\varphi)}{\Xi^2}. \quad (\text{B26})$$

Ξ is given by Eq. (B19).




FIG. 12: Self-energy of bosonic field.

Appendix C: The self-energy of boson

As shown in Fig. 12, the self-energy of boson is defined as

$$\Pi(k_0, \mathbf{k}) = g^2 \int \frac{dq_0}{2\pi} \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \text{Tr} [\gamma_0 G_0(q_0, \mathbf{q}) \gamma_0 G_0(k_0 + q_0, \mathbf{k}, \mathbf{q})]. \quad (\text{C1})$$

Substituting Eq. (A1) into Eq. (C1), we arrive

$$\begin{aligned} \Pi(k_0, \mathbf{k}) = & 4g^2 \int \frac{dq_0}{2\pi} \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{1}{(q_0^2 + E_{\mathbf{q}}^2) [(k_0 + q_0)^2 + E_{\mathbf{k}+\mathbf{q}}^2]} \left\{ -q_0(k_0 + q_0) + v^2 q_x(k_x + q_x) + v^2 q_y(k_y + q_y) \right. \\ & \left. + v_z^2 q_z(k_z + q_z) + D^2(q_x^2 - q_y^2) [(k_x + q_x)^2 - (k_y + q_y)^2] + (m - B_i q_i^2) [m - B_i(k_i + q_i)^2] \right\}. \end{aligned} \quad (\text{C2})$$

Taking $k_0 = 0$, and performing further simplification, we get

$$\begin{aligned} \Pi(\mathbf{k}) = & -g^2 \frac{1}{2} \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{1}{E_{\mathbf{q}}^3} F_1(\mathbf{k}, \mathbf{q}) + g^2 \frac{1}{2} \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{1}{E_{\mathbf{q}}^5} F_1^2(\mathbf{k}, \mathbf{q}) + g^2 \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{1}{E_{\mathbf{q}}^3} F_2(\mathbf{k}, \mathbf{q}) \\ & - g^2 \frac{3}{4} \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{1}{E_{\mathbf{q}}^5} F_1(\mathbf{k}, \mathbf{q}) F_2(\mathbf{k}, \mathbf{q}), \end{aligned} \quad (\text{C3})$$

where

$$\begin{aligned} F_1(\mathbf{k}, \mathbf{q}) = & (2v^2 q_x + 4D^2 q_x^3 - 4D^2 q_x q_y^2 - 4m B_\perp q_x + 4B_\perp^2 q_x q_\perp^2 + 4B_\perp B_z q_x q_z^2) k_x \\ & + (2v^2 q_y + 4D^2 q_y^3 - 4D^2 q_y q_x^2 - 4m B_\perp q_y + 4B_\perp^2 q_y q_\perp^2 + 4B_\perp B_z q_y q_z^2) k_y \\ & + (2v_z^2 q_z - 4m B_z q_z + 4B_z^2 q_z^3 + 4B_z B_\perp q_z q_\perp^2) k_z \\ & + (v^2 + 6D^2 q_x^2 - 2D^2 q_y^2 + 4B_\perp^2 q_x^2 - 2m B_\perp + 2B_\perp^2 q_\perp^2 + 2B_\perp B_z q_z^2) k_x^2 \\ & + (v^2 + 6D^2 q_y^2 - 2D^2 q_x^2 + 4B_\perp^2 q_y^2 - 2m B_\perp + 2B_\perp^2 q_\perp^2 + 2B_\perp B_z q_z^2) k_y^2 \\ & + (v_z^2 + 4B_z^2 q_z^2 - 2m B_z + 2B_\perp B_z q_\perp^2 + 2B_z^2 q_z^2) k_z^2 \\ & + (-8D^2 q_x q_y + 8B_\perp^2 q_x q_y) k_x k_y + 8B_\perp B_z q_x q_z k_x k_z + 8B_\perp B_z q_y q_z k_y k_z, \end{aligned} \quad (\text{C4})$$

and

$$\begin{aligned}
F_2(\mathbf{k}, \mathbf{q}) = & (v^2 q_x + 2D^2 q_x^3 - 2D^2 q_x q_y^2 - 2mB_\perp q_x + 2B_\perp^2 q_\perp^2 q_x + 2B_\perp B_z q_z^2 q_x) k_x \\
& + (v^2 q_y + 2D^2 q_y^3 - 2D^2 q_y q_x^2 - 2mB_\perp q_y + 2B_\perp^2 q_\perp^2 q_y + 2B_\perp B_z q_z^2 q_y) k_y \\
& + (v^2 q_z - 2mB_z q_z + 2mB_\perp B_z q_\perp^2 q_z + 2B_z^2 q_z^3) k_z \\
& + (D^2 q_x^2 - D^2 q_y^2 - mB_\perp + B_\perp^2 q_\perp^2 + B_\perp B_z q_z^2) k_x^2 \\
& + (-D^2 q_x^2 + D^2 q_y^2 - mB_\perp + B_\perp^2 q_\perp^2 + B_\perp B_z q_z^2) k_y^2 \\
& + (-mB_z + B_\perp B_z q_\perp^2 + B_z^2 q_z^2) k_z^2.
\end{aligned} \tag{C5}$$

Retaining the quadratic order of k_i , $\Pi(\mathbf{k})$ can be further written as

$$\Pi(\mathbf{k}) \tag{C6}$$

$$\begin{aligned}
= & k_x^2 \frac{1}{2} g^2 \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \left\{ -\frac{1}{E_{\mathbf{q}}^3} (v^2 + 4D^2 q_x^2 + 4B_\perp^2 q_x^2) + \frac{1}{E_{\mathbf{q}}^5} [v^2 + 2D^2 (q_x^2 - q_y^2) - 2mB_\perp + 2B_\perp^2 q_\perp^2 + 2B_\perp B_z q_z^2]^2 q_x^2 \right\} \\
& + k_y^2 \frac{1}{2} g^2 \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \left\{ -\frac{1}{E_{\mathbf{q}}^3} (v^2 + 4D^2 q_y^2 + 4B_\perp^2 q_y^2) + \frac{1}{E_{\mathbf{q}}^5} [v^2 + 2D^2 (q_y^2 - q_x^2) - 2mB_\perp + 2B_\perp^2 q_\perp^2 + 2B_\perp B_z q_z^2]^2 q_y^2 \right\} \\
& + k_z^2 \frac{1}{2} g^2 \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \left[-\frac{1}{E_{\mathbf{q}}^3} (v_z^2 + 4B_z^2 q_z^2) + \frac{1}{E_{\mathbf{q}}^5} (v_z^2 - 2mB_z + 2B_z^2 q_z^2 + 2B_z B_\perp q_\perp^2)^2 q_z^2 \right].
\end{aligned} \tag{C7}$$

Adopting the transformation as shown in Eq. (B4), and utilizing the RG scheme as shown in Eq. (B5), we obtain

$$\Pi(\mathbf{k}) = -C_x k_x^2 \ell - C_y k_y^2 \ell - \eta C_z k_z^2 \ell, \tag{C8}$$

where

$$C_x = \alpha \mathcal{F}_2^x, \tag{C9}$$

$$C_y = \alpha \mathcal{F}_2^y, \tag{C10}$$

$$C_z = \alpha \mathcal{F}_2^z. \tag{C11}$$

$\mathcal{F}_2^x, \mathcal{F}_2^y, \mathcal{F}_2^z$ are expressed by

$$\begin{aligned}
\mathcal{F}_2^x = & \frac{1}{4\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \left\{ \frac{1 + 4 \left(\left(\frac{D\Lambda}{v} \right)^2 + \left(\frac{B_\perp \Lambda}{v} \right)^2 \right) \sin^2(\varphi) \cos^2(\theta)}{\Xi^3} \right. \\
& \left. - \frac{\left[1 + 2 \left(\frac{D\Lambda}{v} \right)^2 \sin^2(\varphi) \cos(2\theta) - 2 \frac{m}{v\Lambda} \frac{B_\perp \Lambda}{v} + 2 \left(\frac{B_\perp \Lambda}{v} \right)^2 \sin^2(\varphi) + 2 \frac{B_\perp \Lambda}{v} \frac{B_z \Lambda}{v\eta} \cos^2(\varphi) \right]^2 \sin^2(\varphi) \cos^2(\theta)}{\Xi^5} \right\} \tag{C12}
\end{aligned}$$

$$\begin{aligned}
\mathcal{F}_2^y = & \frac{1}{4\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \left\{ \frac{1 + 4 \left(\left(\frac{D\Lambda}{v} \right)^2 + \left(\frac{B_\perp \Lambda}{v} \right)^2 \right) \sin^2(\varphi) \sin^2(\theta)}{\Xi^3} \right. \\
& \left. - \frac{\left[1 - 2 \left(\frac{D\Lambda}{v} \right)^2 \sin^2(\varphi) \cos(2\theta) - 2 \frac{m}{v\Lambda} \frac{B_\perp \Lambda}{v} + 2 \left(\frac{B_\perp \Lambda}{v} \right)^2 \sin^2(\varphi) + 2 \frac{B_\perp \Lambda}{v} \frac{B_z \Lambda}{v\eta} \cos^2(\varphi) \right]^2 \sin^2(\varphi) \sin^2(\theta)}{\Xi^5} \right\} \tag{C13}
\end{aligned}$$

$$\begin{aligned}
\mathcal{F}_2^z = & \frac{1}{4\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \left\{ \frac{\left(\frac{v_z}{v\sqrt{\eta}} \right)^2 + 4 \left(\frac{B_z \Lambda}{v\eta} \right)^2 \cos^2(\varphi)}{\Xi^3} \right. \\
& \left. - \frac{\left(\left(\frac{v_z}{v\sqrt{\eta}} \right)^2 - 2 \frac{m}{v\Lambda} \frac{B_z \Lambda}{v\eta} + 2 \left(\frac{B_z \Lambda}{v\eta} \right)^2 \cos^2(\varphi) + 2 \frac{B_z \Lambda}{v\eta} \frac{B_\perp \Lambda}{v} \sin^2(\varphi) \right)^2 \cos^2(\varphi)}{\Xi^5} \right\}, \tag{C14}
\end{aligned}$$

where Ξ is given by Eq. (B19). It easy to verify that $\mathcal{F}_2^x = \mathcal{F}_2^y$. Then we can define

$$\mathcal{F}_2^\perp \equiv \mathcal{F}_2^x = \mathcal{F}_2^y, \tag{C15}$$

$$C_\perp = \alpha \mathcal{F}_2^\perp. \tag{C16}$$

Accordingly, the polarization can be written as

$$\Pi(\mathbf{k}) = -C_{\perp} k_{\perp}^2 \ell - \eta C_z k_z^2 \ell. \quad (\text{C17})$$

Appendix D: The corrections to fermion-boson coupling vertex

As shown in Fig. 13(a), the correction to fermion-boson coupling induced by Coulomb interaction takes the form

$$\begin{aligned} \Gamma^{(1)} = & -ig^3 \int_{-\infty}^{+\infty} \frac{dq_0}{2\pi} \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \gamma_0 G_0(q_0, \mathbf{q}) \\ & \times \gamma_0 G_0(q_0, \mathbf{q}) \gamma_0 D_0(q_0, \mathbf{q}). \end{aligned} \quad (\text{D1})$$

Substituting the expressions of fermion propagator and boson propagator into Eq. (D1), we find

$$\Gamma^{(1)} = 0, \quad (\text{D2})$$

which indicates

$$\delta g^{(1)} = 0. \quad (\text{D3})$$

The correction to fermion-boson coupling induced by disorder scattering as shown in Fig. 13(b) can be written as

$$\begin{aligned} \Gamma^{(2)} = & ig \sum_j \Delta_j \int' \frac{d^3 \mathbf{k}}{(2\pi)^3} \Gamma_j G_0(0, \mathbf{k}) \gamma_0 \\ & \times G_0(0, \mathbf{k}) \Gamma_j \end{aligned} \quad (\text{D4})$$

Substituting Eq. (A1) into Eq. (D4), one can get

$$\Gamma^{(2)} = ig \gamma_0 C_0^{dis} \ell, \quad (\text{D5})$$

where C_0^{dis} is given by Eq. (B23). Thus,

$$\delta g^{(2)} = g C_0^{dis} \ell. \quad (\text{D6})$$

Then we obtain

$$\delta g = \delta g^{(1)} + \delta g^{(2)} = g C_0^{dis} \ell. \quad (\text{D7})$$




FIG. 13: Corrections to fermion-boson coupling due to (a) Coulomb interaction and (b) disorder.




FIG. 14: One-loop Feynman diagrams for the corrections to the fermion-disorder coupling.

Appendix E: Corrections to fermion-disorder coupling vertex

The correction induced by disorder scattering from Fig. 14(a) can be written as

$$W^{(1)} = \sum_i W_i^{(1)}, \quad (\text{E1})$$

where

$$\begin{aligned} W_i^{(1)} = & \Delta_i \sum_j \Delta_j (\bar{\Psi}_a \Gamma_i \Psi) \int' \frac{d^3 \mathbf{k}}{(2\pi)^3} [\bar{\Psi}_b \Gamma_j G_0(0, \mathbf{k}) \Gamma_i \\ & \times G_0(0, \mathbf{k}) \Gamma_j \Psi_b]. \end{aligned} \quad (\text{E2})$$

The correction generated by disorder scattering from Figs. 14(b) and 14(c) takes the form

$$W^{(2)+(3)} = \sum_i \sum_{i \leq j} W_{ij}^{(2)+(3)}, \quad (\text{E3})$$

where

$$\begin{aligned} W_{ij}^{(2)+(3)} = & \Delta_i \Delta_j \int' \frac{d^3 \mathbf{k}}{(2\pi)^3} [\bar{\Psi}_a \Gamma_i G_0(0, \mathbf{k}) \Gamma_j \Psi_a] \\ & \times \{ \bar{\Psi}_b [\Gamma_j G_0(0, \mathbf{k}) \Gamma_i \\ & + \Gamma_i G_0(0, -\mathbf{k}) \Gamma_j] \Psi_b \}. \end{aligned} \quad (\text{E4})$$

Fig. 14(d) induces the correction

$$W^{(4)} = \sum_i W_i^{(4)}, \quad (\text{E5})$$

where

$$\begin{aligned} W_i^{(4)} = & -\Delta_i g^2 (\bar{\Psi}_a \Gamma_i \Psi_a) \int \frac{dq_0}{2\pi} \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} [\bar{\Psi}_b \gamma_0 \\ & \times G_0(q_0, \mathbf{q}) \Gamma_i G_0(q_0, \mathbf{q}) \gamma_0 \Psi_b] D_0(q_0, \mathbf{q}). \end{aligned} \quad (\text{E6})$$

The contribution from Fig. 14(e) is given by

$$\begin{aligned} W_i^{(5)} = & \Delta_i g^2 (\bar{\Psi}_a \Gamma_i \Psi_a) \int \frac{dq_0}{2\pi} \int' \frac{d^3 \mathbf{q}}{(2\pi)^3} \text{Tr} [\Gamma_i G_0(q_0, \mathbf{q}) \\ & \times \gamma_0 G_0(q_0, \mathbf{k} + \mathbf{q})] D_0(\mathbf{k}) (\bar{\Psi}_b \gamma_0 \Psi_b). \end{aligned} \quad (\text{E7})$$

1. Correction induced by disorder scattering from Feynman diagram 14(a)

Substituting the Eq. (A1) into Eq. (E2), we arrive

$$W_i^{(1)} = \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \Delta_i (\bar{\Psi}_a \Gamma_i \Psi_a) \sum_j \Delta_j \times \left\{ \bar{\Psi}_b \Gamma_j \left[-\frac{1}{2} (\gamma_x \Gamma_i \gamma_x + \gamma_y \Gamma_i \gamma_y) \mathcal{G}_1^\perp - \gamma_z \Gamma_i \gamma_z \mathcal{G}_1^z - \gamma_5 \Gamma_i \gamma_5 \mathcal{G}_1^D + \Gamma_i \mathcal{G}_1^m \right] \Gamma_j \Psi_b \right\} \quad (\text{E8})$$

with

$$\mathcal{G}_1^\perp = \frac{1}{4\pi} \int_0^\pi d\varphi \sin(\varphi) \int_0^{2\pi} d\theta \frac{\sin^2(\varphi)}{\Xi^4}, \quad (\text{E9})$$

$$\mathcal{G}_1^z = \frac{1}{4\pi} \int_0^\pi d\varphi \sin(\varphi) \int_0^{2\pi} d\theta \frac{\left(\frac{v_z}{v\sqrt{\eta}}\right)^2 \cos^2(\varphi)}{\Xi^4} \quad (\text{E10})$$

$$\mathcal{G}_1^D = \frac{1}{4\pi} \int_0^\pi d\varphi \sin(\varphi) \int_0^{2\pi} d\theta \times \frac{\frac{D^2 \Lambda^2}{v^2} \sin^4(\varphi) (\cos^2(\theta) - \sin^2(\theta))^2}{\Xi^4}, \quad (\text{E11})$$

$$\mathcal{G}_1^m = \frac{1}{4\pi} \int_0^\pi d\varphi \sin(\varphi) \int_0^{2\pi} d\theta \times \frac{\left(\frac{m}{v\Lambda} - \frac{B_\perp \Lambda}{v} \sin^2(\varphi) - \frac{B_z \Lambda}{v\eta} \cos^2(\varphi)\right)^2}{\Xi^4}, \quad (\text{E12})$$

where Ξ is expressed by Eq. (B19).

$W_i^{(1)}$ can be further written as

$$W_i^{(1)} = \frac{\delta \Delta_i^{(1)}}{2} (\bar{\Psi}_a i \gamma_z \Psi_a) (\bar{\Psi}_b i \gamma_z \Psi_b), \quad (\text{E13})$$

where

$$\delta \Delta_C^{(1)} = 2\Delta_C (\Delta_C + \Delta_M + \Delta_{AC} + 2\Delta_{SO\perp} + \Delta_{SOz} + \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} + 2\Delta_{AMN\perp} + \Delta_{AMNz} + 2\Delta_{CR\perp} + \Delta_{CRz}) \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} (\mathcal{G}_1^\perp + \mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \ell, \quad (\text{E14})$$

$$\delta \Delta_M^{(1)} = 2\Delta_M (\Delta_C + \Delta_M - \Delta_{AC} - 2\Delta_{SO\perp} - \Delta_{SOz} - \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} + 2\Delta_{AMN\perp} + \Delta_{AMNz} - 2\Delta_{CR\perp} - 2\Delta_{CRz}) \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} (-\mathcal{G}_1^\perp - \mathcal{G}_1^z - \mathcal{G}_1^D + \mathcal{G}_1^m) \ell, \quad (\text{E15})$$

$$\delta \Delta_{AC}^{(1)} = 2\Delta_{AC} (-\Delta_C + \Delta_M - \Delta_{AC} + 2\Delta_{SO\perp} + \Delta_{SOz} + \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} - 2\Delta_{AMN\perp} - \Delta_{AMNz} - 2\Delta_{CR\perp} - \Delta_{CRz}) \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} (-\mathcal{G}_1^\perp - \mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \ell, \quad (\text{E16})$$

$$\delta \Delta_{SO\perp}^{(1)} = 2\Delta_{SO\perp} (-\Delta_C + \Delta_M + \Delta_{AC} + \Delta_{SOz} - \Delta_{PM} - \Delta_{MNz} + \Delta_{AMNz} - \Delta_{CRz}) \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \times (-\mathcal{G}_1^z - \mathcal{G}_1^D + \mathcal{G}_1^m) \ell, \quad (\text{E17})$$

$$\delta \Delta_{SOz}^{(1)} = 2\Delta_{SOz} (-\Delta_C + \Delta_M + \Delta_{AC} + 2\Delta_{SO\perp} - \Delta_{SOz} - \Delta_{PM} - 2\Delta_{MN\perp} + \Delta_{MNz} + 2\Delta_{AMN\perp} - \Delta_{AMNz} - 2\Delta_{CR\perp} + \Delta_{CRz}) \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} (-\mathcal{G}_1^\perp + \mathcal{G}_1^z - \mathcal{G}_1^D + \mathcal{G}_1^m) \ell, \quad (\text{E18})$$

$$\delta \Delta_{PM}^{(1)} = 2\Delta_{PM} (-\Delta_C + \Delta_M + \Delta_{AC} - 2\Delta_{SO\perp} - \Delta_{SOz} - \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} - 2\Delta_{AMN\perp} - \Delta_{AMNz} + 2\Delta_{CR\perp} + \Delta_{CRz}) \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} (\mathcal{G}_1^\perp + \mathcal{G}_1^z - \mathcal{G}_1^D + \mathcal{G}_1^m) \ell, \quad (\text{E19})$$

$$\delta \Delta_{MN\perp}^{(1)} = 2\Delta_{MN\perp} (\Delta_C + \Delta_M - \Delta_{AC} + \Delta_{SOz} - \Delta_{PM} - \Delta_{MNz} - \Delta_{AMNz} + \Delta_{CRz}) \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \times (\mathcal{G}_1^z - \mathcal{G}_1^D + \mathcal{G}_1^m) \ell, \quad (\text{E20})$$

$$\delta \Delta_{MNz}^{(1)} = 2\Delta_{MNz} (\Delta_C + \Delta_M - \Delta_{AC} + 2\Delta_{SO\perp} - \Delta_{SOz} - \Delta_{PM} - 2\Delta_{MN\perp} + \Delta_{MNz} - 2\Delta_{AMN\perp} + \Delta_{AMNz} + 2\Delta_{CR\perp} - \Delta_{CRz}) \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} (\mathcal{G}_1^\perp - \mathcal{G}_1^z - \mathcal{G}_1^D + \mathcal{G}_1^m) \ell, \quad (\text{E21})$$

$$\delta \Delta_{AMN\perp}^{(1)} = 2\Delta_{AMN\perp} (\Delta_C + \Delta_M + \Delta_{AC} - \Delta_{SOz} + \Delta_{PM} - \Delta_{MNz} - \Delta_{AMNz} - \Delta_{CRz}) \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \times (-\mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \ell, \quad (\text{E22})$$

$$\begin{aligned}\delta\Delta_{AMNz}^{(1)} &= 2\Delta_{AMNz}(\Delta_C + \Delta_M + \Delta_{AC} - 2\Delta_{SO\perp} + \Delta_{SOz} + \Delta_{PM} - 2\Delta_{MN\perp} + \Delta_{MNz} - 2\Delta_{AMN\perp} + \Delta_{AMNz} \\ &\quad - 2\Delta_{CR\perp} + \Delta_{CRz}) \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} (-\mathcal{G}_1^\perp + \mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \ell, \end{aligned} \quad (E23)$$

$$\begin{aligned}\delta\Delta_{CR\perp}^{(1)} &= 2\Delta_{CR\perp}(-\Delta_C + \Delta_M - \Delta_{AC} - \Delta_{SOz} + \Delta_{PM} - \Delta_{MNz} + \Delta_{AMNz} + \Delta_{CRz}) \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \\ &\quad \times (\mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \ell, \end{aligned} \quad (E24)$$

$$\begin{aligned}\delta\Delta_{CRz}^{(1)} &= 2\Delta_{CRz}(-\Delta_C + \Delta_M - \Delta_{AC} - 2\Delta_{SO\perp} + \Delta_{SOz} + \Delta_{PM} - 2\Delta_{MN\perp} + \Delta_{MNz} + 2\Delta_{AMN\perp} - \Delta_{AMNz} \\ &\quad + 2\Delta_{CR\perp} - \Delta_{CRz}) \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} (\mathcal{G}_1^\perp - \mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \ell. \end{aligned} \quad (E25)$$

It should be noticed that

$$\mathcal{G}_1^\perp + \mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m = \mathcal{G}_0^\perp + \mathcal{G}_0^z. \quad (E26)$$

2. Correction induced by disorder scattering from Feynman diagrams 14(b) and 14(c)

Substituting Eq. (A1) into Eq. (E4), we obtain

$$\begin{aligned}W_{i,j}^{(2)+(3)} &= -\frac{1}{2}\Delta_i\Delta_j(\bar{\Psi}_a\Gamma_i\gamma_x\Gamma_j\Psi_a)[\bar{\Psi}_b(\Gamma_j\gamma_x\Gamma_i - \Gamma_i\gamma_x\Gamma_j)\Psi_b] \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^\perp \\ &\quad -\frac{1}{2}\Delta_i\Delta_j(\bar{\Psi}_a\Gamma_i\gamma_y\Gamma_j\Psi_a)[\bar{\Psi}_b(\Gamma_j\gamma_y\Gamma_i - \Gamma_i\gamma_y\Gamma_j)\Psi_b] \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^\perp \\ &\quad -\Delta_i\Delta_j(\bar{\Psi}_a\Gamma_i\gamma_z\Gamma_j\Psi_a)[\bar{\Psi}_b(\Gamma_j\gamma_z\Gamma_i - \Gamma_i\gamma_z\Gamma_j)\Psi_b] \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^z \\ &\quad -\Delta_i\Delta_j(\bar{\Psi}_a\Gamma_i\gamma_5\Gamma_j\Psi_a)[\bar{\Psi}_b(\Gamma_j\gamma_5\Gamma_i + \Gamma_i\gamma_5\Gamma_j)\Psi_b] \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^D \\ &\quad +\Delta_i\Delta_j(\bar{\Psi}_a\Gamma_i\Gamma_j\Psi_a)[\bar{\Psi}_b(\Gamma_j\Gamma_i + \Gamma_i\Gamma_j)\Psi_b] \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^m. \end{aligned} \quad (E27)$$

Concretely, we find that

$$W_{C,C}^{(2)+(3)} = 2\Delta_C^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^D \ell (\bar{\Psi}_a i\gamma_5 \Psi_a) (\bar{\Psi}_b i\gamma_5 \Psi_b) + 2\Delta_C^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^m \ell (\bar{\Psi}_a \Psi_a) (\bar{\Psi}_b \Psi_b), \quad (E28)$$

$$W_{M,M}^{(2)+(3)} = 2\Delta_M^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^D \ell (\bar{\Psi}_a i\gamma_5 \Psi_a) (\bar{\Psi}_b i\gamma_5 \Psi_b) + 2\Delta_M^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^m \ell (\bar{\Psi}_a \Psi_a) (\bar{\Psi}_b \Psi_b), \quad (E29)$$

$$W_{AC,AC}^{(2)+(3)} = 2\Delta_{AC}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^D \ell (\bar{\Psi}_a i\gamma_5 \Psi_a) (\bar{\Psi}_b i\gamma_5 \Psi_b) + 2\Delta_{AC}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^m \ell (\bar{\Psi}_a \Psi_a) (\bar{\Psi}_b \Psi_b), \quad (E30)$$

$$\begin{aligned}W_{SO\perp,SO\perp}^{(2)+(3)} &= 4\Delta_{SO\perp}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^z (\bar{\Psi}_a \gamma_0 \gamma_5 \Psi_a) (\bar{\Psi}_b \gamma_0 \gamma_5 \Psi_b) + 4\Delta_{SO\perp}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^D (\bar{\Psi}_a i\gamma_5 \Psi_a) (\bar{\Psi}_b i\gamma_5 \Psi_b) \\ &\quad + 4\Delta_{SO\perp}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^m (\bar{\Psi}_a \Psi_a) (\bar{\Psi}_b \Psi_b), \end{aligned} \quad (E31)$$

$$W_{SOz,SOz}^{(2)+(3)} = 2\Delta_{SOz}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^D \ell (\bar{\Psi}_a i\gamma_5 \Psi_a) (\bar{\Psi}_b i\gamma_5 \Psi_b) + 2\Delta_{SOz}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^m \ell (\bar{\Psi}_a \Psi_a) (\bar{\Psi}_b \Psi_b), \quad (E32)$$

$$W_{PM,PM}^{(2)+(3)} = 2\Delta_{PM}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^D \ell (\bar{\Psi}_a i\gamma_5 \Psi_a) (\bar{\Psi}_b i\gamma_5 \Psi_b) + 2\Delta_{PM}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^m \ell (\bar{\Psi}_a \Psi_a) (\bar{\Psi}_b \Psi_b), \quad (E33)$$

$$\begin{aligned}W_{MN\perp,MN\perp}^{(2)+(3)} &= 4\Delta_{MN\perp}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^z (\bar{\Psi}_a \gamma_0 \gamma_5 \Psi_a) (\bar{\Psi}_b \gamma_0 \gamma_5 \Psi_b) + 4\Delta_{MN\perp}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^D (\bar{\Psi}_a i\gamma_5 \Psi_a) (\bar{\Psi}_b i\gamma_5 \Psi_b) \\ &\quad + 4\Delta_{MN\perp}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^m (\bar{\Psi}_a \Psi_a) (\bar{\Psi}_b \Psi_b), \end{aligned} \quad (E34)$$

$$W_{MNz,MNz}^{(2)+(3)} = 2\Delta_{MNz}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^D (\bar{\Psi}_a i\gamma_5 \Psi_a) (\bar{\Psi}_b i\gamma_5 \Psi_b)$$

$$+2\Delta_{MNz}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^m (\bar{\Psi}_a \Psi_a) (\bar{\Psi}_b \Psi_b), \quad (\text{E35})$$

$$\begin{aligned} W_{AMN\perp, AMN\perp}^{(2)+(3)} &= 4\Delta_{AMN\perp}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^z (\bar{\Psi}_a \gamma_0 \gamma_5 \Psi_a) (\bar{\Psi}_b \gamma_0 \gamma_5 \Psi_b) + 4\Delta_{AMN\perp}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^D (\bar{\Psi}_a i\gamma_5 \Psi_a) (\bar{\Psi}_b i\gamma_5 \Psi_b) \\ &\quad + 4\Delta_{AMN\perp}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^m (\bar{\Psi}_a \Psi_a) (\bar{\Psi}_b \Psi_b), \end{aligned} \quad (\text{E36})$$

$$W_{AMNz, AMNz}^{(2)+(3)} = 2\Delta_{AMNz}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^D (\bar{\Psi}_a i\gamma_5 \Psi_a) (\bar{\Psi}_b i\gamma_5 \Psi_b) + 2\Delta_{AMNz}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^m (\bar{\Psi}_a \Psi_a) (\bar{\Psi}_b \Psi_b), \quad (\text{E37})$$

$$\begin{aligned} W_{CR\perp, CR\perp}^{(2)+(3)} &= 4\Delta_{CR\perp}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^z (\bar{\Psi}_a \gamma_0 \gamma_5 \Psi_a) (\bar{\Psi}_b \gamma_0 \gamma_5 \Psi_b) + 4\Delta_{CR\perp}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^D (\bar{\Psi}_a i\gamma_5 \Psi_a) (\bar{\Psi}_b i\gamma_5 \Psi_b) \\ &\quad + 4\Delta_{CR\perp}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^m (\bar{\Psi}_a \Psi_a) (\bar{\Psi}_b \Psi_b), \end{aligned} \quad (\text{E38})$$

$$W_{CRz, CRz}^{(2)+(3)} = 2\Delta_{CRz}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^D (\bar{\Psi}_a i\gamma_5 \Psi_a) (\bar{\Psi}_b i\gamma_5 \Psi_b) + 2\Delta_{CRz}^2 \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^m (\bar{\Psi}_a \Psi_a) (\bar{\Psi}_b \Psi_b), \quad (\text{E39})$$

$$\begin{aligned} W_{C,M}^{(2)+(3)} &= \Delta_C \Delta_M \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^\perp \ell \sum_{j=x,y} (\bar{\Psi}_a \gamma_0 \gamma_j \Psi_a) (\bar{\Psi}_b \gamma_0 \gamma_j \Psi_b) \\ &\quad + 2\Delta_C \Delta_M \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^z \ell (\bar{\Psi}_a \gamma_0 \gamma_z \Psi_a) (\bar{\Psi}_b \gamma_0 \gamma_z \Psi_b) \\ &\quad + 2\Delta_C \Delta_M \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^m \ell (\bar{\Psi}_a \gamma_0 \Psi_a) (\bar{\Psi}_b \gamma_0 \Psi_b), \end{aligned} \quad (\text{E40})$$

$$W_{C,AC}^{(2)+(3)} = 0, \quad (\text{E41})$$

$$\begin{aligned} W_{C,SO\perp}^{(2)+(3)} &= 2\Delta_C \Delta_{SO\perp} \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^\perp \ell (\bar{\Psi}_a \Psi_a) (\bar{\Psi}_b \Psi_b) \\ &\quad + 2\Delta_C \Delta_{SO\perp} \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^D \ell \sum_{j=x,y} (\bar{\Psi}_a i\gamma_5 \gamma_j \Psi_a) (\bar{\Psi}_b i\gamma_5 \gamma_j \Psi_b), \end{aligned} \quad (\text{E42})$$

$$\begin{aligned} W_{C,SOz}^{(2)+(3)} &= 2\Delta_C \Delta_{SOz} \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^z (\bar{\Psi}_a \Psi_a) (\bar{\Psi}_b \Psi_b) \\ &\quad + 2\Delta_C \Delta_{SOz} \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^D (\bar{\Psi}_a i\gamma_5 \gamma_z \Psi_a) (\bar{\Psi}_b i\gamma_5 \gamma_z \Psi_b), \end{aligned} \quad (\text{E43})$$

$$\begin{aligned} W_{C,PM}^{(2)+(3)} &= \Delta_C \Delta_{PM} \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^\perp \ell [(\bar{\Psi}_a i\gamma_y \gamma_z \Psi_a) (\bar{\Psi}_b i\gamma_y \gamma_z \Psi_b) + (\bar{\Psi}_a i\gamma_z \gamma_x \Psi_a) (\bar{\Psi}_b i\gamma_z \gamma_x \Psi_b)] \\ &\quad + 2\Delta_C \Delta_{PM} \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^z \ell (\bar{\Psi}_a i\gamma_x \gamma_y \Psi_a) (\bar{\Psi}_b i\gamma_x \gamma_y \Psi_b) \\ &\quad + 2\Delta_C \Delta_{PM} \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \mathcal{G}_1^D \ell (\bar{\Psi}_a \gamma_0 \Psi_a) (\bar{\Psi}_b \gamma_0 \Psi_b), \end{aligned} \quad (\text{E44})$$

$$\begin{aligned} W_{C,MN\perp}^{(2)+(3)} &= 2\Delta_C \Delta_{MN\perp} \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^\perp (\bar{\Psi}_a i\gamma_5 \Psi_a) (\bar{\Psi}_b i\gamma_5 \Psi_b) \\ &\quad + 2\Delta_C \Delta_{MN\perp} \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^m \sum_{j=x,y} (\bar{\Psi}_a i\gamma_5 \gamma_j \Psi_a) (\bar{\Psi}_b i\gamma_5 \gamma_j \Psi_b), \end{aligned} \quad (\text{E45})$$

$$\begin{aligned} W_{C,MNz}^{(2)+(3)} &= 2\Delta_C \Delta_{MNz} \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^z (\bar{\Psi}_a i\gamma_5 \Psi_a) (\bar{\Psi}_b i\gamma_5 \Psi_b) \\ &\quad + 2\Delta_C \Delta_{MNz} \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^m (\bar{\Psi}_a i\gamma_5 \gamma_z \Psi_a) (\bar{\Psi}_b i\gamma_5 \gamma_z \Psi_b), \end{aligned} \quad (\text{E46})$$

$$\begin{aligned} W_{C,AMN\perp}^{(2)+(3)} &= 2\Delta_C \Delta_{AMN\perp} \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^\perp (\bar{\Psi}_a i\gamma_z \Psi_a) (\bar{\Psi}_b i\gamma_z \Psi_b) \\ &\quad + 2\Delta_C \Delta_{AMN\perp} \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell \mathcal{G}_1^z \sum_{j=x,y} (\bar{\Psi}_a i\gamma_j \Psi_a) (\bar{\Psi}_b i\gamma_j \Psi_b) \end{aligned}$$

$$\begin{aligned}
& +2\Delta_C\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D\sum_{j=x,y}(\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b) \\
& +2\Delta_C\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m[(\bar{\Psi}_ai\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_y\gamma_z\Psi_b)+(\bar{\Psi}_ai\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_bi\gamma_z\gamma_x\Psi_b)], \quad (E47)
\end{aligned}$$

$$\begin{aligned}
W_{C,AMNz}^{(2)+(3)} &= \Delta_C\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp\sum_{j=x,y}(\bar{\Psi}_ai\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_j\Psi_b) \\
& +2\Delta_C\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_a\gamma_0\gamma_z\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_z\Psi_b) \\
& +2\Delta_C\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_ai\gamma_x\gamma_y\Psi_a)(\bar{\Psi}_bi\gamma_x\gamma_y\Psi_b), \quad (E48)
\end{aligned}$$

$$\begin{aligned}
W_{C,CR\perp}^{(2)+(3)} &= 2\Delta_C\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_ai\gamma_5\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_5\gamma_z\Psi_b) \\
& +2\Delta_C\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z\sum_{j=x,y}(\bar{\Psi}_ai\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_5\gamma_j\Psi_b), \quad (E49)
\end{aligned}$$

$$W_{C,CRz}^{(2)+(3)} = \Delta_C\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp\sum_{j=x,y}(\bar{\Psi}_ai\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_5\gamma_j\Psi_b), \quad (E50)$$

$$W_{M,AC}^{(2)+(3)} = 2\Delta_M\Delta_{AC}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_a\gamma_0\gamma_5\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_5\Psi_b), \quad (E51)$$

$$\begin{aligned}
W_{M,SO\perp}^{(2)+(3)} &= 2\Delta_M\Delta_{SO\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a\gamma_0\Psi_a)(\bar{\Psi}_b\gamma_0\Psi_b) \\
& +2\Delta_M\Delta_{SO\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D[(\bar{\Psi}_ai\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_y\gamma_z\Psi_b)+(\bar{\Psi}_ai\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_bi\gamma_z\gamma_x\Psi_b)] \\
& +2\Delta_M\Delta_{SO\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m\sum_{j=x,y}(\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b), \quad (E52)
\end{aligned}$$

$$\begin{aligned}
W_{M,SOz}^{(2)+(3)} &= 2\Delta_M\Delta_{SOz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z(\bar{\Psi}_a\gamma_0\Psi_a)(\bar{\Psi}_b\gamma_0\Psi_b) \\
& +2\Delta_M\Delta_{SOz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_ai\gamma_x\gamma_y\Psi_a)(\bar{\Psi}_bi\gamma_x\gamma_y\Psi_b) \\
& +2\Delta_M\Delta_{SOz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_a\gamma_0\gamma_z\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_z\Psi_b), \quad (E53)
\end{aligned}$$

$$\begin{aligned}
W_{M,PM}^{(2)+(3)} &= \Delta_M\Delta_{PM}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp\sum_{j=x,y}(\bar{\Psi}_ai\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_5\gamma_j\Psi_b) \\
& +2\Delta_M\Delta_{PM}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z(\bar{\Psi}_ai\gamma_5\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_5\gamma_z\Psi_b)+2\Delta_M\Delta_{PM}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_a\Psi_a)(\bar{\Psi}_b\Psi_b) \\
& +2\Delta_M\Delta_{PM}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_ai\gamma_5\Psi_a)(\bar{\Psi}_bi\gamma_5\Psi_b), \quad (E54)
\end{aligned}$$

$$\begin{aligned}
W_{M,MN\perp}^{(2)+(3)} &= 2\Delta_M\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_ai\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_z\Psi_b) \\
& +2\Delta_M\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z\sum_{j=x,y}(\bar{\Psi}_ai\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_j\Psi_b) \\
& +2\Delta_M\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D\sum_{j=x,y}(\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b) \\
& +2\Delta_M\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m[(\bar{\Psi}_ai\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_y\gamma_z\Psi_b)+(\bar{\Psi}_ai\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_bi\gamma_z\gamma_x\Psi_b)], \quad (E55)
\end{aligned}$$

$$W_{M,MNz}^{(2)+(3)} = \Delta_M\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp\sum_{j=x,y}(\bar{\Psi}_ai\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_j\Psi_b)$$

$$\begin{aligned}
& +2\Delta_M\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_a\gamma_0\gamma_z\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_z\Psi_b) \\
& +2\Delta_M\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_ai\gamma_x\gamma_y\Psi_a)(\bar{\Psi}_bi\gamma_x\gamma_y\Psi_b),
\end{aligned} \tag{E56}$$

$$\begin{aligned}
W_{M,AMN\perp}^{(2)+(3)} &= 2\Delta_M\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_ai\gamma_5\Psi_a)(\bar{\Psi}_bi\gamma_5\Psi_b) \\
& +2\Delta_M\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m\sum_{j=x,y}(\bar{\Psi}_ai\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_5\gamma_j\Psi_b),
\end{aligned} \tag{E57}$$

$$\begin{aligned}
W_{M,AMNZ}^{(2)+(3)} &= 2\Delta_M\Delta_{AMNZ}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z(\bar{\Psi}_ai\gamma_5\Psi_a)(\bar{\Psi}_bi\gamma_5\Psi_b) \\
& +2\Delta_M\Delta_{AMNZ}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_ai\gamma_5\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_5\gamma_z\Psi_b),
\end{aligned} \tag{E58}$$

$$\begin{aligned}
W_{M,CRL}^{(2)+(3)} &= 2\Delta_M\Delta_{CRL}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_ai\gamma_x\gamma_y\Psi_a)(\bar{\Psi}_bi\gamma_x\gamma_y\Psi_b) \\
& +2\Delta_M\Delta_{CRL}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z[(\bar{\Psi}_ai\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_y\gamma_z\Psi_b) + (\bar{\Psi}_ai\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_bi\gamma_z\gamma_x\Psi_b)] \\
& +2\Delta_M\Delta_{CRL}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m\sum_{j=x,y}(\bar{\Psi}_ai\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_j\Psi_b),
\end{aligned} \tag{E59}$$

$$\begin{aligned}
W_{M,CRz}^{(2)+(3)} &= \Delta_M\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp[(\bar{\Psi}_ai\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_y\gamma_z\Psi_b) + (\bar{\Psi}_ai\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_bi\gamma_z\gamma_x\Psi_b)] \\
& +2\Delta_M\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_ai\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_z\Psi_b),
\end{aligned} \tag{E60}$$

$$\begin{aligned}
W_{AC,SO\perp}^{(2)+(3)} &= 2\Delta_{AC}\Delta_{SO\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a\gamma_0\gamma_z\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_z\Psi_b) \\
& +2\Delta_{AC}\Delta_{SO\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z\sum_{j=x,y}(\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b) \\
& +2\Delta_{AC}\Delta_{SO\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D\sum_{j=x,y}(\bar{\Psi}_ai\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_j\Psi_b),
\end{aligned} \tag{E61}$$

$$\begin{aligned}
W_{AC,SOz}^{(2)+(3)} &= \Delta_{AC}\Delta_{SOz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp\sum_{j=x,y}(\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b) \\
& +2\Delta_{AC}\Delta_{SOz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_ai\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_z\Psi_b),
\end{aligned} \tag{E62}$$

$$W_{AC,PM}^{(2)+(3)} = 2\Delta_{AC}\Delta_{PM}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_a\gamma_0\gamma_5\Psi_a)(\bar{\Psi}_bi\gamma_0\gamma_5\Psi_b), \tag{E63}$$

$$\begin{aligned}
W_{AC,MN\perp}^{(2)+(3)} &= 2\Delta_{AC}\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_ai\gamma_x\gamma_y\Psi_a)(\bar{\Psi}_bi\gamma_x\gamma_y\Psi_b) \\
& +2\Delta_{AC}\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z[(\bar{\Psi}_ai\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_y\gamma_z\Psi_b) + (\bar{\Psi}_ai\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_bi\gamma_z\gamma_x\Psi_b)] \\
& +2\Delta_{AC}\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m\sum_{j=x,y}(\bar{\Psi}_ai\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_j\Psi_b),
\end{aligned} \tag{E64}$$

$$\begin{aligned}
W_{AC,MNZ}^{(2)+(3)} &= \Delta_{AC}\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp[(\bar{\Psi}_ai\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_y\gamma_z\Psi_b) + (\bar{\Psi}_ai\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_bi\gamma_z\gamma_x\Psi_b)] \\
& +2\Delta_{AC}\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_ai\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_z\Psi_b),
\end{aligned} \tag{E65}$$

$$W_{AC,AMN\perp}^{(2)+(3)} = 2\Delta_{AC}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_ai\gamma_5\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_5\gamma_z\Psi_b)$$

$$+2\Delta_{AC}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z\sum_{j=x,y}(\bar{\Psi}_a i\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_b i\gamma_5\gamma_j\Psi_b), \quad (\text{E66})$$

$$W_{AC,AMNz}^{(2)+(3)} = \Delta_{AC}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp\sum_{j=x,y}(\bar{\Psi}_a i\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_b i\gamma_5\gamma_j\Psi_b), \quad (\text{E67})$$

$$\begin{aligned} W_{AC,CR\perp}^{(2)+(3)} &= 2\Delta_{AC}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a i\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_z\Psi_b) \\ &+ 2\Delta_{AC}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z\sum_{j=x,y}(\bar{\Psi}_a i\gamma_j\Psi_a)(\bar{\Psi}_b i\gamma_j\Psi_b) \\ &+ 2\Delta_{AC}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D\sum_{j=x,y}(\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b) \\ &+ 2\Delta_{AC}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m[(\bar{\Psi}_a i\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_y\gamma_z\Psi_b) + (\bar{\Psi}_a i\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_b i\gamma_z\gamma_x\Psi_b)], \end{aligned} \quad (\text{E68})$$

$$\begin{aligned} W_{AC,CRz}^{(2)+(3)} &= \Delta_{AC}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp\sum_{j=x,y}(\bar{\Psi}_a i\gamma_j\Psi_a)(\bar{\Psi}_b i\gamma_j\Psi_b) \\ &+ 2\Delta_{AC}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_a\gamma_0\gamma_z\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_z\Psi_b) \\ &+ 2\Delta_{AC}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_a i\gamma_x\gamma_y\Psi_a)(\bar{\Psi}_b i\gamma_x\gamma_y\Psi_b), \end{aligned} \quad (\text{E69})$$

$$W_{SO\perp,SOz}^{(2)+(3)} = 2\Delta_{SO\perp}\Delta_{SOz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a\gamma_0\gamma_5\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_5\Psi_b), \quad (\text{E70})$$

$$\begin{aligned} W_{SO\perp,PM}^{(2)+(3)} &= 2\Delta_{SO\perp}\Delta_{PM}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a i\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_z\Psi_b) \\ &+ 2\Delta_{SO\perp}\Delta_{PM}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z\sum_{j=x,y}(\bar{\Psi}_a i\gamma_j\Psi_a)(\bar{\Psi}_b i\gamma_j\Psi_b) \\ &+ 2\Delta_{SO\perp}\Delta_{PM}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D\sum_{j=x,y}(\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b) \\ &+ 2\Delta_{SO\perp}\Delta_{PM}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m[(\bar{\Psi}_a i\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_y\gamma_z\Psi_b) + (\bar{\Psi}_a i\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_b i\gamma_z\gamma_x\Psi_b)], \end{aligned} \quad (\text{E71})$$

$$\begin{aligned} W_{SO\perp,MN\perp}^{(2)+(3)} &= 4\Delta_{SO\perp}\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp\sum_{j=x,y}(\bar{\Psi}_a i\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_b i\gamma_5\gamma_j\Psi_b) \\ &+ 4\Delta_{SO\perp}\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z(\bar{\Psi}_a i\gamma_5\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_5\gamma_z\Psi_b) \\ &+ 4\Delta_{SO\perp}\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_a\Psi_a)(\bar{\Psi}_b\Psi_b) \\ &+ 4\Delta_{SO\perp}\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_a i\gamma_5\Psi_a)(\bar{\Psi}_b i\gamma_5\Psi_b), \end{aligned} \quad (\text{E72})$$

$$\begin{aligned} W_{SO\perp,MNz}^{(2)+(3)} &= 2\Delta_{SO\perp}\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a i\gamma_5\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_5\gamma_z\Psi_b) \\ &+ 2\Delta_{SO\perp}\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z\sum_{j=x,y}(\bar{\Psi}_a i\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_b i\gamma_5\gamma_j\Psi_b), \end{aligned} \quad (\text{E73})$$

$$\begin{aligned} W_{SO\perp,AMN\perp}^{(2)+(3)} &= 4\Delta_{SO\perp}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp[(\bar{\Psi}_a i\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_y\gamma_z\Psi_b) + (\bar{\Psi}_a i\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_b i\gamma_z\gamma_x\Psi_b)] \\ &+ 4\Delta_{SO\perp}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z(\bar{\Psi}_a i\gamma_x\gamma_y\Psi_a)(\bar{\Psi}_b i\gamma_x\gamma_y\Psi_b) \\ &+ 4\Delta_{SO\perp}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_a\gamma_0\Psi_a)(\bar{\Psi}_b\gamma_0\Psi_b) \end{aligned}$$

$$+4\Delta_{SO\perp}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_a i\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_z\Psi_b), \quad (\text{E74})$$

$$\begin{aligned} W_{SO\perp,AMNz}^{(2)+(3)} &= 2\Delta_{SO\perp}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a i\gamma_x\gamma_y\Psi_a)(\bar{\Psi}_b i\gamma_x\gamma_y\Psi_b) \\ &+ 2\Delta_{SO\perp}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z[(\bar{\Psi}_a i\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_y\gamma_z\Psi_b) + (\bar{\Psi}_a i\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_b i\gamma_z\gamma_x\Psi_b)] \\ &+ 2\Delta_{SO\perp}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m\sum_{j=x,y}(\bar{\Psi}_a i\gamma_j\Psi_a)(\bar{\Psi}_b i\gamma_j\Psi_b), \end{aligned} \quad (\text{E75})$$

$$\begin{aligned} W_{SO\perp,CR\perp}^{(2)+(3)} &= 4\Delta_{SO\perp}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z(\bar{\Psi}_a i\gamma_5\Psi_a)(\bar{\Psi}_b i\gamma_5\Psi_b) \\ &+ 4\Delta_{SO\perp}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_a\gamma_0\gamma_5\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_5\Psi_b) \\ &+ 4\Delta_{SO\perp}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_a i\gamma_5\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_5\gamma_z\Psi_b), \end{aligned} \quad (\text{E76})$$

$$\begin{aligned} W_{SO\perp,CRz}^{(2)+(3)} &= 2\Delta_{SO\perp}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a i\gamma_5\Psi_a)(\bar{\Psi}_b i\gamma_5\Psi_b) \\ &+ 2\Delta_{SO\perp}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m\sum_{j=x,y}(\bar{\Psi}_a i\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_b i\gamma_5\gamma_j\Psi_b), \end{aligned} \quad (\text{E77})$$

$$\begin{aligned} W_{SOz,PM}^{(2)+(3)} &= \Delta_{SOz}\Delta_{PM}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\mathcal{G}_1^\perp\ell\sum_{j=x,y}(\bar{\Psi}_a i\gamma_j\Psi_a)(\bar{\Psi}_b i\gamma_j\Psi_b) \\ &+ 2\Delta_{SOz}\Delta_{PM}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\mathcal{G}_1^D\ell(\bar{\Psi}_a\gamma_0\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_0\gamma_z\Psi_b) \\ &+ 2\Delta_{SOz}\Delta_{PM}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\mathcal{G}_1^m\ell(\bar{\Psi}_a i\gamma_x\gamma_y\Psi_a)(\bar{\Psi}_b i\gamma_x\gamma_y\Psi_b), \end{aligned} \quad (\text{E78})$$

$$\begin{aligned} W_{SOz,MN\perp}^{(2)+(3)} &= 2\Delta_{SOz}\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a i\gamma_5\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_5\gamma_z\Psi_b) \\ &+ 2\Delta_{SOz}\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z\sum_{j=x,y}(\bar{\Psi}_a i\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_b i\gamma_5\gamma_j\Psi_b), \end{aligned} \quad (\text{E79})$$

$$\begin{aligned} W_{SOz,MNz}^{(2)+(3)} &= \Delta_{SOz}\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp\sum_{j=x,y}(\bar{\Psi}_a i\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_b i\gamma_5\gamma_j\Psi_b) \\ &+ 2\Delta_{SOz}\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z(\bar{\Psi}_a i\gamma_5\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_5\gamma_z\Psi_b) \\ &+ 2\Delta_{SOz}\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_a\Psi_a)(\bar{\Psi}_b\Psi_b) \\ &+ 2\Delta_{SOz}\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_a i\gamma_5\Psi_a)(\bar{\Psi}_b i\gamma_5\Psi_b), \end{aligned} \quad (\text{E80})$$

$$\begin{aligned} W_{SOz,AMN\perp}^{(2)+(3)} &= 2\Delta_{SOz}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a i\gamma_x\gamma_y\Psi_a)(\bar{\Psi}_b i\gamma_x\gamma_y\Psi_b) \\ &+ 2\Delta_{SOz}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z[(\bar{\Psi}_a i\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_y\gamma_z\Psi_b) + (\bar{\Psi}_a i\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_b i\gamma_z\gamma_x\Psi_b)] \\ &+ 2\Delta_{SOz}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m\sum_{j=x,y}(\bar{\Psi}_a i\gamma_j\Psi_a)(\bar{\Psi}_b i\gamma_j\Psi_b), \end{aligned} \quad (\text{E81})$$

$$\begin{aligned} W_{SOz,AMNz}^{(2)+(3)} &= \Delta_{SOz}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp[(\bar{\Psi}_a i\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_y\gamma_z\Psi_b) + (\bar{\Psi}_a i\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_b i\gamma_z\gamma_x\Psi_b)] \\ &+ 2\Delta_{SOz}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z(\bar{\Psi}_a i\gamma_x\gamma_y\Psi_a)(\bar{\Psi}_b i\gamma_x\gamma_y\Psi_b) \end{aligned}$$

$$+2\Delta_{SOz}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_a\gamma_0\Psi_a)(\bar{\Psi}_b\gamma_0\Psi_b), \quad (\text{E82})$$

$$\begin{aligned} W_{SOz,CR\perp}^{(2)+(3)} &= 2\Delta_{SOz}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_ai\gamma_5\Psi_a)(\bar{\Psi}_bi\gamma_5\Psi_b) \\ &+ 2\Delta_{SOz}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m\sum_{j=x,y}(\bar{\Psi}_ai\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_5\gamma_j\Psi_b), \end{aligned} \quad (\text{E83})$$

$$W_{SOz,CRz}^{(2)+(3)} = 2\Delta_{SOz}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_a\gamma_0\gamma_5\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_5\Psi_b), \quad (\text{E84})$$

$$\begin{aligned} W_{PM,MN\perp}^{(2)+(3)} &= 2\Delta_{PM}\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a\gamma_0\Psi_a)(\bar{\Psi}_b\gamma_0\Psi_b) \\ &+ 2\Delta_{PM}\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D[(\bar{\Psi}_ai\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_y\gamma_z\Psi_b) + (\bar{\Psi}_ai\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_bi\gamma_z\gamma_x\Psi_b)] \\ &+ 2\Delta_{PM}\Delta_{MN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m\sum_{j=x,y}(\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b), \end{aligned} \quad (\text{E85})$$

$$\begin{aligned} W_{PM,MNz}^{(2)+(3)} &= 2\Delta_{PM}\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z(\bar{\Psi}_a\gamma_0\Psi_a)(\bar{\Psi}_b\gamma_0\Psi_b) \\ &+ 2\Delta_{PM}\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_ai\gamma_x\gamma_y\Psi_a)(\bar{\Psi}_bi\gamma_x\gamma_y\Psi_b) \\ &+ 2\Delta_{PM}\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_a\gamma_0\gamma_z\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_z\Psi_b), \end{aligned} \quad (\text{E86})$$

$$\begin{aligned} W_{PM,AMN\perp}^{(2)+(3)} &= 2\Delta_{PM}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a\Psi_a)(\bar{\Psi}_b\Psi_b) \\ &+ 2\Delta_{PM}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D\sum_{j=x,y}(\bar{\Psi}_ai\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_5\gamma_j\Psi_b), \end{aligned} \quad (\text{E87})$$

$$\begin{aligned} W_{PM,AMNz}^{(2)+(3)} &= 2\Delta_{PM}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z(\bar{\Psi}_a\Psi_a)(\bar{\Psi}_b\Psi_b) \\ &+ 2\Delta_{PM}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_ai\gamma_5\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_5\gamma_z\Psi_b), \end{aligned} \quad (\text{E88})$$

$$\begin{aligned} W_{PM,CR\perp}^{(2)+(3)} &= 2\Delta_{PM}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a\gamma_0\gamma_z\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_z\Psi_b) \\ &+ 2\Delta_{PM}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z\sum_{j=x,y}(\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b) \\ &+ 2\Delta_{PM}\Delta_{CR\perp}\sum_{j=x,y}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_ai\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_j\Psi_b), \end{aligned} \quad (\text{E89})$$

$$\begin{aligned} W_{PM,CRz}^{(2)+(3)} &= \Delta_{PM}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp\sum_{j=x,y}(\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b) \\ &+ 2\Delta_{PM}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_ai\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_z\Psi_b), \end{aligned} \quad (\text{E90})$$

$$W_{MN\perp,MNz}^{(2)+(3)} = 2\Delta_{MN\perp}\Delta_{MNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a\gamma_0\gamma_5\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_5\Psi_b), \quad (\text{E91})$$

$$\begin{aligned} W_{MN\perp,AMN\perp}^{(2)+(3)} &= 4\Delta_{MN\perp}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp\sum_{j=x,y}(\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b) \\ &+ 4\Delta_{MN\perp}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z(\bar{\Psi}_a\gamma_0\gamma_z\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_z\Psi_b) \\ &+ 4\Delta_{MN\perp}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_ai\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_z\Psi_b) \end{aligned}$$

$$+4\Delta_{MN\perp}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_a\gamma_0\Psi_a)(\bar{\Psi}_b\gamma_0\Psi_b), \quad (\text{E92})$$

$$\begin{aligned} W_{MN\perp,AMNz}^{(2)+(3)} &= 2\Delta_{MN\perp}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a\gamma_0\gamma_z\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_z\Psi_b) \\ &+ 2\Delta_{MN\perp}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z\sum_{j=x,y}(\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b) \\ &+ 2\Delta_{MN\perp}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D\sum_{j=x,y}(\bar{\Psi}_ai\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_j\Psi_b), \end{aligned} \quad (\text{E93})$$

$$\begin{aligned} W_{MN\perp,CR\perp}^{(2)+(3)} &= 4\Delta_{MN\perp}\Delta_{CR\perp}(\bar{\Psi}_a\Psi_a)(\bar{\Psi}_b\Psi_b)\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z \\ &+ 4\Delta_{MN\perp}\Delta_{CR\perp}(\bar{\Psi}_ai\gamma_5\gamma_z\Psi_a)(\bar{\Psi}_bi\gamma_5\gamma_z\Psi_b)\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D \\ &+ 4\Delta_{MN\perp}\Delta_{CR\perp}(\bar{\Psi}_a\gamma_0\gamma_5\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_5\Psi_b)\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m, \end{aligned} \quad (\text{E94})$$

$$\begin{aligned} W_{MN\perp,CRz}^{(2)+(3)} &= 2\Delta_{MN\perp}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a\Psi_a)(\bar{\Psi}_b\Psi_b) \\ &+ 2\Delta_{MN\perp}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D\sum_{j=x,y}(\bar{\Psi}_ai\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_5\gamma_j\Psi_b), \end{aligned} \quad (\text{E95})$$

$$\begin{aligned} W_{MNz,AMN\perp}^{(2)+(3)} &= 2\Delta_{MNz}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a\gamma_0\gamma_z\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_z\Psi_b) \\ &+ 2\Delta_{MNz}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z\sum_{j=x,y}(\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b) \\ &+ 2\Delta_{MNz}\Delta_{AMN\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D\sum_{j=x,y}(\bar{\Psi}_ai\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_j\Psi_b), \end{aligned} \quad (\text{E96})$$

$$\begin{aligned} W_{MNz,AMNz}^{(2)+(3)} &= \Delta_{MNz}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp\sum_{j=x,y}(\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b) \\ &+ 2\Delta_{MNz}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z(\bar{\Psi}_a\gamma_0\gamma_z\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_z\Psi_b) \\ &+ 2\Delta_{MNz}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_a\gamma_0\Psi_a)(\bar{\Psi}_b\gamma_0\Psi_b), \end{aligned} \quad (\text{E97})$$

$$\begin{aligned} W_{MNz,CR\perp}^{(2)+(3)} &= 2\Delta_{MNz}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a\Psi_a)(\bar{\Psi}_b\Psi_b) \\ &+ 2\Delta_{MNz}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D\sum_{j=x,y}(\bar{\Psi}_ai\gamma_5\gamma_j\Psi_a)(\bar{\Psi}_bi\gamma_5\gamma_j\Psi_b), \end{aligned} \quad (\text{E98})$$

$$W_{MNz,CRz}^{(2)+(3)} = 2\Delta_{MNz}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_a\gamma_0\gamma_5\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_5\Psi_b), \quad (\text{E99})$$

$$W_{AMN\perp,AMNz}^{(2)+(3)} = 2\Delta_{AMN\perp}\Delta_{AMNz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a\gamma_0\gamma_5\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_5\Psi_b), \quad (\text{E100})$$

$$\begin{aligned} W_{AMN\perp,CR\perp}^{(2)+(3)} &= 4\Delta_{AMN\perp}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^z(\bar{\Psi}_a\gamma_0\Psi_a)(\bar{\Psi}_b\gamma_0\Psi_b) \\ &+ 4\Delta_{AMN\perp}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D(\bar{\Psi}_ai\gamma_x\gamma_y\Psi_a)(\bar{\Psi}_bi\gamma_x\gamma_y\Psi_b) \\ &+ 4\Delta_{AMN\perp}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m(\bar{\Psi}_a\gamma_0\gamma_z\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_z\Psi_b), \end{aligned} \quad (\text{E101})$$

$$W_{AMN\perp,CRz}^{(2)+(3)} = 2\Delta_{AMN\perp}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp(\bar{\Psi}_a\gamma_0\Psi_a)(\bar{\Psi}_b\gamma_0\Psi_b)$$

$$\begin{aligned}
& +2\Delta_{AMN\perp}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D [(\bar{\Psi}_a i\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_y\gamma_z\Psi_b) + (\bar{\Psi}_a i\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_b i\gamma_z\gamma_x\Psi_b)] \\
& +2\Delta_{AMN\perp}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m \sum_{j=x,y} (\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b), \tag{E102}
\end{aligned}$$

$$\begin{aligned}
W_{AMNz,CR\perp}^{(2)+(3)} &= 2\Delta_{AMNz}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp (\bar{\Psi}_a\gamma_0\Psi_a)(\bar{\Psi}_b\gamma_0\Psi_b) \\
& +2\Delta_{AMNz}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^D [(\bar{\Psi}_a i\gamma_y\gamma_z\Psi_a)(\bar{\Psi}_b i\gamma_y\gamma_z\Psi_b) + (\bar{\Psi}_a i\gamma_z\gamma_x\Psi_a)(\bar{\Psi}_b i\gamma_z\gamma_x\Psi_b)] \\
& +2\Delta_{AMNz}\Delta_{CR\perp}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^m \sum_{j=x,y} (\bar{\Psi}_a\gamma_0\gamma_j\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_j\Psi_b), \tag{E103}
\end{aligned}$$

$$W_{AMNz,CRz}^{(2)+(3)} = 0 \tag{E104}$$

$$W_{CR\perp,CRz}^{(2)+(3)} = 2\Delta_{CR\perp}\Delta_{CRz}\frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell\mathcal{G}_1^\perp (\bar{\Psi}_a\gamma_0\gamma_5\Psi_a)(\bar{\Psi}_b\gamma_0\gamma_5\Psi_b). \tag{E105}$$

$W^{(2)+(3)}$ can be written as

$$\begin{aligned}
W^{(2)+(3)} &= \sum_i \sum_{i \leq j} W_{i,j}^{(2)+(3)} \\
&= \frac{\delta\Delta_i^{(2)+(3)}}{2} (\bar{\Psi}_a\Gamma_i\Psi_a)(\bar{\Psi}_b\Gamma_i\Psi_b), \tag{E106}
\end{aligned}$$

where

$$\begin{aligned}
\delta\Delta_C^{(2)+(3)} &= 4 [(\Delta_M\Delta_{SO\perp} + \Delta_{PM}\Delta_{MN\perp} + \Delta_{AMN\perp}\Delta_{CRz} + \Delta_{AMNz}\Delta_{CR\perp})\mathcal{G}_1^\perp + (\Delta_M\Delta_{SOz} + \Delta_{PM}\Delta_{MNz} \\
& +2\Delta_{AMN\perp}\Delta_{CR\perp})\mathcal{G}_1^z + (\Delta_C\Delta_{PM} + 2\Delta_{SO\perp}\Delta_{AMN\perp} + \Delta_{SOz}\Delta_{AMNz})\mathcal{G}_1^D \\
& + (\Delta_C\Delta_M + 2\Delta_{MN\perp}\Delta_{AMN\perp} + \Delta_{MNz}\Delta_{AMNz})\mathcal{G}_1^m] \frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell, \tag{E107}
\end{aligned}$$

$$\begin{aligned}
\delta\Delta_M^{(2)+(3)} &= 4 [(\Delta_C\Delta_{SO\perp} + \Delta_{PM}\Delta_{AMN\perp} + \Delta_{MN\perp}\Delta_{CRz} + \Delta_{MNz}\Delta_{CR\perp})\mathcal{G}_1^\perp + (\Delta_C\Delta_{SOz} + \Delta_{PM}\Delta_{AMNz} \\
& +2\Delta_{MN\perp}\Delta_{CR\perp})\mathcal{G}_1^z + (\Delta_M\Delta_{PM} + 2\Delta_{SO\perp}\Delta_{MN\perp} + \Delta_{SOz}\Delta_{MNz})\mathcal{G}_1^D \\
& + (\Delta_C^2 + \Delta_M^2 + \Delta_{AC}^2 + 2\Delta_{SO\perp}^2 + \Delta_{SOz}^2 + \Delta_{PM}^2 + 2\Delta_{MN\perp}^2 + \Delta_{MNz}^2 + 2\Delta_{AMN\perp}^2 + \Delta_{AMNz}^2 \\
& +2\Delta_{CR\perp}^2 + \Delta_{CRz}^2)\mathcal{G}_1^m] \frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell, \tag{E108}
\end{aligned}$$

$$\begin{aligned}
\delta\Delta_{AC}^{(2)+(3)} &= 4 [(\Delta_{SO\perp}\Delta_{SOz} + \Delta_{MN\perp}\Delta_{MNz} + \Delta_{AMN\perp}\Delta_{AMNz} + \Delta_{CR\perp}\Delta_{CRz})\mathcal{G}_1^\perp \\
& +2(\Delta_{SO\perp}^2 + \Delta_{MN\perp}^2 + \Delta_{AMN\perp}^2 + \Delta_{CR\perp}^2)\mathcal{G}_1^z + (\Delta_{AC}\Delta_{PM} + 2\Delta_{SO\perp}\Delta_{CR\perp} + \Delta_{SOz}\Delta_{CRz})\mathcal{G}_1^D \\
& + (\Delta_M\Delta_{AC} + \Delta_{MNz}\Delta_{CRz} + 2\Delta_{MN\perp}\Delta_{CR\perp})\mathcal{G}_1^m] \frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell, \tag{E109}
\end{aligned}$$

$$\begin{aligned}
\delta\Delta_{SO\perp}^{(2)+(3)} &= 2 [(\Delta_C\Delta_M + \Delta_{AC}\Delta_{SOz} + \Delta_{PM}\Delta_{CRz} + 4\Delta_{MN\perp}\Delta_{AMN\perp} + \Delta_{MNz}\Delta_{AMNz})\mathcal{G}_1^\perp \\
& +2(\Delta_{AC}\Delta_{SO\perp} + \Delta_{PM}\Delta_{CR\perp} + \Delta_{MN\perp}\Delta_{AMNz} + \Delta_{MNz}\Delta_{AMN\perp})\mathcal{G}_1^z \\
& +2(\Delta_C\Delta_{AMN\perp} + \Delta_M\Delta_{MN\perp} + \Delta_{AC}\Delta_{CR\perp} + \Delta_{SO\perp}\Delta_{PM})\mathcal{G}_1^D \\
& +2(\Delta_M\Delta_{SO\perp} + \Delta_{PM}\Delta_{MN\perp} + \Delta_{AMN\perp}\Delta_{CRz} + \Delta_{AMNz}\Delta_{CR\perp})\mathcal{G}_1^m] \frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell, \tag{E110}
\end{aligned}$$

$$\begin{aligned}
\delta\Delta_{SOz}^{(2)+(3)} &= 4 [(\Delta_{AC}\Delta_{SO\perp} + \Delta_{PM}\Delta_{CR\perp} + \Delta_{MN\perp}\Delta_{AMNz} + \Delta_{MNz}\Delta_{AMN\perp})\mathcal{G}_1^\perp \\
& + (\Delta_C\Delta_M + 2\Delta_{MN\perp}\Delta_{AMN\perp} + \Delta_{MNz}\Delta_{AMNz})\mathcal{G}_1^z \\
& + (\Delta_C\Delta_{AMNz} + \Delta_M\Delta_{MNz} + \Delta_{AC}\Delta_{CRz} + \Delta_{SOz}\Delta_{PM})\mathcal{G}_1^D \\
& + (\Delta_M\Delta_{SOz} + \Delta_{PM}\Delta_{MNz} + 2\Delta_{AMN\perp}\Delta_{CR\perp})\mathcal{G}_1^m] \frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell, \tag{E111}
\end{aligned}$$

$$\begin{aligned}
\delta\Delta_{PM}^{(2)+(3)} &= 4 [(\Delta_C\Delta_{MN\perp} + \Delta_M\Delta_{AMN\perp} + \Delta_{SO\perp}\Delta_{CRz} + \Delta_{SOz}\Delta_{CR\perp})\mathcal{G}_1^\perp \\
& + (\Delta_C\Delta_{MNz} + \Delta_M\Delta_{AMNz} + 2\Delta_{SO\perp}\Delta_{CR\perp})\mathcal{G}_1^z + (\Delta_C^2 + \Delta_M^2 + \Delta_{AC}^2 + 2\Delta_{SO\perp}^2 + \Delta_{SOz}^2 \\
& + 2\Delta_{PM}^2 + 2\Delta_{MN\perp}^2 + \Delta_{MNz}^2 + 2\Delta_{AMN\perp}^2 + \Delta_{AMNz}^2 + 2\Delta_{CR\perp}^2 + \Delta_{CRz}^2)\mathcal{G}_1^m] \frac{\Lambda}{2\pi^2v^2\sqrt{\eta}}\ell,
\end{aligned}$$

$$\begin{aligned}
& + \Delta_{PM}^2 + 2\Delta_{MN\perp}^2 + \Delta_{MNz}^2 + 2\Delta_{AMN\perp}^2 + \Delta_{AMNz}^2 + 2\Delta_{CR\perp}^2 + \Delta_{CRz}^2) \mathcal{G}_1^D \\
& + (\Delta_M \Delta_{PM} + 2\Delta_{SO\perp} \Delta_{MN\perp} + \Delta_{SOz} \Delta_{MNz}) \mathcal{G}_1^m \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell,
\end{aligned} \tag{E112}$$

$$\begin{aligned}
\delta\Delta_{MN\perp}^{(2)+(3)} &= 2 [(\Delta_C \Delta_{PM} + \Delta_M \Delta_{CRz} + \Delta_{AC} \Delta_{MNz} + 4\Delta_{SO\perp} \Delta_{AMN\perp} + \Delta_{SOz} \Delta_{AMNz}) \mathcal{G}_1^\perp \\
& + 2(\Delta_M \Delta_{CR\perp} + \Delta_{AC} \Delta_{MN\perp} + \Delta_{SO\perp} \Delta_{AMNz} + \Delta_{SOz} \Delta_{AMN\perp}) \mathcal{G}_1^z \\
& + 2(\Delta_M \Delta_{SO\perp} + \Delta_{PM} \Delta_{MN\perp} + \Delta_{AMN\perp} \Delta_{CRz} + \Delta_{AMNz} \Delta_{CR\perp}) \mathcal{G}_1^D \\
& + 2(\Delta_C \Delta_{AMN\perp} + \Delta_M \Delta_{MN\perp} + \Delta_{AC} \Delta_{CR\perp} + \Delta_{SO\perp} \Delta_{PM}) \mathcal{G}_1^m] \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell,
\end{aligned} \tag{E113}$$

$$\begin{aligned}
\delta\Delta_{MNz}^{(2)+(3)} &= 4 [(\Delta_M \Delta_{CR\perp} + \Delta_{AC} \Delta_{MN\perp} + \Delta_{SO\perp} \Delta_{AMNz} + \Delta_{SOz} \Delta_{AMN\perp}) \mathcal{G}_1^\perp \\
& + (\Delta_C \Delta_{PM} + 2\Delta_{SO\perp} \Delta_{AMN\perp} + \Delta_{SOz} \Delta_{AMNz}) \mathcal{G}_1^z \\
& + (\Delta_M \Delta_{SOz} + \Delta_{PM} \Delta_{MNz} + 2\Delta_{AMN\perp} \Delta_{CR\perp}) \mathcal{G}_1^D \\
& + (\Delta_C \Delta_{AMNz} + \Delta_M \Delta_{MNz} + \Delta_{AC} \Delta_{CRz} + \Delta_{SOz} \Delta_{PM}) \mathcal{G}_1^m] \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell,
\end{aligned} \tag{E114}$$

$$\begin{aligned}
\delta\Delta_{AMN\perp}^{(2)+(3)} &= 2 [(\Delta_C \Delta_{CRz} + \Delta_M \Delta_{PM} + \Delta_{AC} \Delta_{AMNz} + 4\Delta_{SO\perp} \Delta_{MN\perp} + \Delta_{SOz} \Delta_{MNz}) \mathcal{G}_1^\perp \\
& + 2(\Delta_C \Delta_{CR\perp} + \Delta_{AC} \Delta_{AMN\perp} + \Delta_{SO\perp} \Delta_{MNz} + \Delta_{SOz} \Delta_{MN\perp}) \mathcal{G}_1^z \\
& + 2(\Delta_C \Delta_{SO\perp} + \Delta_{PM} \Delta_{AMN\perp} + \Delta_{MN\perp} \Delta_{CRz} + \Delta_{MNz} \Delta_{CR\perp}) \mathcal{G}_1^D \\
& + 2(\Delta_C \Delta_{MN\perp} + \Delta_M \Delta_{AMN\perp} + \Delta_{SO\perp} \Delta_{CRz} + \Delta_{SOz} \Delta_{CR\perp}) \mathcal{G}_1^m] \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell,
\end{aligned} \tag{E115}$$

$$\begin{aligned}
\delta\Delta_{AMNz}^{(2)+(3)} &= 4 [(\Delta_C \Delta_{CR\perp} + \Delta_{AC} \Delta_{AMN\perp} + \Delta_{SO\perp} \Delta_{MNz} + \Delta_{SOz} \Delta_{MN\perp}) \mathcal{G}_1^\perp \\
& + (\Delta_M \Delta_{PM} + 2\Delta_{SO\perp} \Delta_{MN\perp} + \Delta_{SOz} \Delta_{MNz}) \mathcal{G}_1^z \\
& + (\Delta_C \Delta_{SOz} + \Delta_{PM} \Delta_{AMNz} + 2\Delta_{MN\perp} \Delta_{CR\perp}) \mathcal{G}_1^D \\
& + (\Delta_C \Delta_{MNz} + \Delta_M \Delta_{AMNz} + 2\Delta_{SO\perp} \Delta_{CR\perp}) \mathcal{G}_1^m] \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell,
\end{aligned} \tag{E116}$$

$$\begin{aligned}
\delta\Delta_{CR\perp}^{(2)+(3)} &= 2 [(\Delta_C \Delta_{AMNz} + \Delta_M \Delta_{MNz} + \Delta_{AC} \Delta_{CRz} + \Delta_{SOz} \Delta_{PM}) \mathcal{G}_1^\perp \\
& + 2(\Delta_C \Delta_{AMN\perp} + \Delta_M \Delta_{MN\perp} + \Delta_{AC} \Delta_{CR\perp} + \Delta_{SO\perp} \Delta_{PM}) \mathcal{G}_1^z \\
& + 2(\Delta_{AC} \Delta_{SO\perp} + \Delta_{PM} \Delta_{CR\perp} + \Delta_{MN\perp} \Delta_{AMNz} + \Delta_{MNz} \Delta_{AMN\perp}) \mathcal{G}_1^D \\
& + 2(\Delta_M \Delta_{CR\perp} + \Delta_{AC} \Delta_{MN\perp} + \Delta_{SO\perp} \Delta_{AMNz} + \Delta_{SOz} \Delta_{AMN\perp}) \mathcal{G}_1^m] \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell,
\end{aligned} \tag{E117}$$

$$\begin{aligned}
\delta\Delta_{CRz}^{(2)+(3)} &= 4 [(\Delta_C \Delta_{AMN\perp} + \Delta_M \Delta_{MN\perp} + \Delta_{AC} \Delta_{CR\perp} + \Delta_{SO\perp} \Delta_{PM}) \mathcal{G}_1^\perp \\
& + (\Delta_{AC} \Delta_{SOz} + \Delta_{PM} \Delta_{CRz} + 2\Delta_{MN\perp} \Delta_{AMN\perp}) \mathcal{G}_1^D \\
& + (\Delta_M \Delta_{CRz} + \Delta_{AC} \Delta_{MNz} + 2\Delta_{SO\perp} \Delta_{AMN\perp}) \mathcal{G}_1^m] \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \ell.
\end{aligned} \tag{E118}$$

3. Correction induced by long-range Coulomb interaction from Feynman diagram 14 (d)

Substituting the expressions of fermion and boson propagators into Eq. (E6), we get

$$\begin{aligned}
W_i^{(4)} &= \Delta_i (\bar{\Psi}_a \Gamma_i \Psi_a) (\bar{\Psi}_b \Gamma_i \Psi_b) \alpha \frac{1}{2} (\mathcal{F}_0^\perp + \mathcal{F}_0^z) \ell \\
& + \Delta_i (\bar{\Psi}_a \Gamma_i \Psi_a) (\bar{\Psi}_b \gamma_0 \gamma_x \Gamma_i \gamma_x \gamma_0 \Psi_b) \alpha \mathcal{F}_3^\perp \ell \\
& + \Delta_i (\bar{\Psi}_a \Gamma_i \Psi_a) (\bar{\Psi}_b \gamma_0 \gamma_y \Gamma_i \gamma_y \gamma_0 \Psi_b) \alpha \mathcal{F}_3^\perp \ell \\
& + \Delta_i (\bar{\Psi}_a \Gamma_i \Psi_a) (\bar{\Psi}_b \gamma_0 \gamma_z \Gamma_i \gamma_z \gamma_0 \Psi_b) \alpha \mathcal{F}_3^z \ell \\
& + \Delta_i (\bar{\Psi}_a \Gamma_i \Psi_a) (\bar{\Psi}_b \gamma_0 \gamma_5 \Gamma_i \gamma_5 \gamma_0 \Psi_b) \alpha \mathcal{F}_3^D \ell \\
& - \Delta_i (\bar{\Psi}_a \Gamma_i \Psi_a) (\bar{\Psi}_b \gamma_0 \Gamma_i \gamma_0 \Psi_b) \alpha \mathcal{F}_3^m \ell, \tag{E119}
\end{aligned}$$

where

$$\mathcal{F}_3^\perp = \frac{1}{16\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \frac{\sin^2(\varphi)}{\Xi^3}, \tag{E120}$$

$$\begin{aligned}
\mathcal{F}_3^z &= \frac{1}{8\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \\
& \times \frac{\left(\frac{v_z}{v\sqrt{\eta}}\right)^2 \cos^2(\varphi)}{\Xi^3},
\end{aligned} \tag{E121}$$

$$\begin{aligned}
\mathcal{F}_3^D &= \frac{1}{8\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \\
& \times \frac{\frac{D^2 \Lambda^2}{v^2} \sin^2(\varphi) \cos^2(2\theta)}{\Xi^3},
\end{aligned} \tag{E122}$$

$$\mathcal{F}_3^m = \frac{1}{8\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \times \frac{\left(\frac{m}{v\Lambda} - \frac{B_\perp \Lambda}{v} \sin^2(\varphi) - \frac{B_z \Lambda}{v\eta} \cos^2(\varphi) \right)^2}{\Xi^3} \quad (\text{E123})$$

Ξ is given by Eq. (B19).

$W_i^{(4)}$ can be also further written as

$$W_i^{(4)} = \frac{\delta\Delta_i^4}{2} (\bar{\Psi}_a \Gamma_i \Psi_a) (\bar{\Psi}_b \Gamma_i \Psi_b), \quad (\text{E124})$$

where

$$\delta\Delta_C^{(4)} = 2\Delta_C \alpha \left[\frac{1}{2} (\mathcal{F}_0^\perp + \mathcal{F}_0^z) - 2\mathcal{F}_3^\perp - \mathcal{F}_3^z - \mathcal{F}_3^D - \mathcal{F}_3^m \right] \ell, \quad (\text{E125})$$

$$\delta\Delta_M^{(4)} = 2\Delta_M \alpha \left[\frac{1}{2} (\mathcal{F}_0^\perp + \mathcal{F}_0^z) + 2\mathcal{F}_3^\perp + \mathcal{F}_3^z + \mathcal{F}_3^D - \mathcal{F}_3^m \right] \ell, \quad (\text{E126})$$

$$\delta\Delta_{AC}^{(4)} = 2\Delta_{AC} \alpha \left[\frac{1}{2} (\mathcal{F}_0^\perp + \mathcal{F}_0^z) - 2\mathcal{F}_3^\perp - \mathcal{F}_3^z + \mathcal{F}_3^D + \mathcal{F}_3^m \right] \ell, \quad (\text{E127})$$

$$\delta\Delta_{SO\perp}^{(4)} = 2\Delta_{SO\perp} \alpha \left[\frac{1}{2} (\mathcal{F}_0^\perp + \mathcal{F}_0^z) - \mathcal{F}_3^z - \mathcal{F}_3^D + \mathcal{F}_3^m \right] \ell, \quad (\text{E128})$$

$$\delta\Delta_{SOz}^{(4)} = 2\Delta_{SOz} \alpha \left[\frac{1}{2} (\mathcal{F}_0^\perp + \mathcal{F}_0^z) - 2\mathcal{F}_3^\perp + \mathcal{F}_3^z - \mathcal{F}_3^D + \mathcal{F}_3^m \right] \ell, \quad (\text{E129})$$

$$\delta\Delta_{PM}^{(4)} = 2\Delta_{PM} \alpha \left[\frac{1}{2} (\mathcal{F}_0^\perp + \mathcal{F}_0^z) + 2\mathcal{F}_3^\perp + \mathcal{F}_3^z - \mathcal{F}_3^D + \mathcal{F}_3^m \right] \ell, \quad (\text{E130})$$

$$\delta\Delta_{MN\perp}^{(4)} = 2\Delta_{MN\perp} \alpha \left[\frac{1}{2} (\mathcal{F}_0^\perp + \mathcal{F}_0^z) - \mathcal{F}_3^z + \mathcal{F}_3^D - \mathcal{F}_3^m \right] \ell, \quad (\text{E131})$$

$$\delta\Delta_{MNz}^{(4)} = 2\Delta_{MNz} \alpha \left[\frac{1}{2} (\mathcal{F}_0^\perp + \mathcal{F}_0^z) - 2\mathcal{F}_3^\perp + \mathcal{F}_3^z + \mathcal{F}_3^D - \mathcal{F}_3^m \right] \ell, \quad (\text{E132})$$

$$\delta\Delta_{AMN\perp}^{(4)} = 2\Delta_{AMN\perp} \alpha \left[\frac{1}{2} (\mathcal{F}_0^\perp + \mathcal{F}_0^z) + \mathcal{F}_3^z - \mathcal{F}_3^D - \mathcal{F}_3^m \right] \ell, \quad (\text{E133})$$

$$\delta\Delta_{AMNz}^{(4)} = 2\Delta_{AMNz} \alpha \left[\frac{1}{2} (\mathcal{F}_0^\perp + \mathcal{F}_0^z) + 2\mathcal{F}_3^\perp - \mathcal{F}_3^z - \mathcal{F}_3^D - \mathcal{F}_3^m \right] \ell, \quad (\text{E134})$$

$$\delta\Delta_{CR\perp}^{(4)} = 2\Delta_{CR\perp} \alpha \left[\frac{1}{2} (\mathcal{F}_0^\perp + \mathcal{F}_0^z) + \mathcal{F}_3^z + \mathcal{F}_3^D + \mathcal{F}_3^m \right] \ell, \quad (\text{E135})$$

$$\delta\Delta_{CRz}^{(4)} = 2\Delta_{CRz} \alpha \left[\frac{1}{2} (\mathcal{F}_0^\perp + \mathcal{F}_0^z) + 2\mathcal{F}_3^\perp - \mathcal{F}_3^z \right]$$

$$+ \mathcal{F}_3^D + \mathcal{F}_3^m \Big] \ell, \quad (\text{E136})$$

It should be notice that

$$2\mathcal{F}_3^\perp + \mathcal{F}_3^z + \mathcal{F}_3^D + \mathcal{F}_3^m = \frac{1}{2} (\mathcal{F}_0^\perp + \mathcal{F}_0^z). \quad (\text{E137})$$

Therefore,

$$\delta\Delta_C^{(4)} = 0. \quad (\text{E138})$$

4. Correction induced by long-range Coulomb interaction from Feynman diagram 14 (e)

Substituting Eqs. (A1) and (A3) into Eq. (E7), one can obtain

$$\begin{aligned} W_C^{(5)} &= \Delta_C (\bar{\Psi}_a \gamma_0 \Psi_a) \Pi(\mathbf{k}) D_0(\mathbf{k}) (\bar{\Psi}_b \gamma_0 \Psi_b) \\ &= \Delta_C (-C_\perp k_\perp^2 \ell - C_z \eta k_z^2 \ell) \frac{1}{k_\perp^2 + \eta k_z^2} \\ &\quad \times (\bar{\Psi}_a \gamma_0 \Psi_a) (\bar{\Psi}_b \gamma_0 \Psi_b). \end{aligned} \quad (\text{E139})$$

It can be approximated as

$$\begin{aligned} W_C^{(5)} &\approx -\Delta_C \frac{C_\perp + C_z}{2} \ell (\bar{\Psi}_a \gamma_0 \Psi_a) (\bar{\Psi}_b \gamma_0 \Psi_b) \\ &= -\Delta_C \alpha \frac{\mathcal{F}_2^\perp + \mathcal{F}_2^z}{2} \ell \\ &\quad \times (\bar{\Psi}_a \gamma_0 \Psi_a) (\bar{\Psi}_b \gamma_0 \Psi_b). \end{aligned} \quad (\text{E140})$$

It is easy to verify that

$$W_M^{(5)} = 0, \quad (\text{E141})$$

$$W_{AC}^{(5)} = 0, \quad (\text{E142})$$

$$W_{SO\perp}^{(5)} = 0, \quad (\text{E143})$$

$$W_{SOz}^{(5)} = 0, \quad (\text{E144})$$

$$W_{PM}^{(5)} = 0, \quad (\text{E145})$$

$$W_{MN\perp}^{(5)} = 0, \quad (\text{E146})$$

$$W_{MNz}^{(5)} = 0, \quad (\text{E147})$$

$$W_{AMN\perp}^{(5)} = 0, \quad (\text{E148})$$

$$W_{AMNz}^{(5)} = 0, \quad (\text{E149})$$

$$W_{CR\perp}^{(5)} = 0, \quad (\text{E150})$$

$$W_{CRz}^{(5)} = 0. \quad (\text{E151})$$

$W_i^{(5)}$ can be written as

$$W_i^{(5)} = \frac{\delta\Delta_i^{(5)}}{2} (\bar{\Psi}_a \Gamma_i \Psi_a) (\bar{\Psi}_b \Gamma_i \Psi_b), \quad (\text{E152})$$

where

$$\delta\Delta_C^{(5)} = -\Delta_C \alpha (\mathcal{F}_2^\perp + \mathcal{F}_2^z) \ell, \quad (\text{E153})$$

$$\delta\Delta_M^{(5)} = 0, \quad (\text{E154})$$

$$\delta\Delta_{AC}^{(5)} = 0, \quad (\text{E155})$$

$$\delta\Delta_{SO\perp}^{(5)} = 0, \quad (\text{E156})$$

$$\delta\Delta_{SOz}^{(5)} = 0, \quad (\text{E157})$$

$$\delta\Delta_{PM}^{(5)} = 0, \quad (\text{E158})$$

$$\delta\Delta_{MN\perp}^{(5)} = 0, \quad (\text{E159})$$

$$\delta\Delta_{MNz}^{(5)} = 0, \quad (\text{E160})$$

$$\delta\Delta_{AMN\perp}^{(5)} = 0, \quad (\text{E161})$$

$$\delta\Delta_{AMNz}^{(5)} = 0, \quad (\text{E162})$$

$$\delta\Delta_{CR\perp}^{(5)} = 0, \quad (\text{E163})$$

$$\delta\Delta_{CRz}^{(5)} = 0. \quad (\text{E164})$$

5. Summary of the corrections

The total contribution is given by

$$\delta\Delta_i = \delta\Delta_i^{(1)} + \delta\Delta_i^{(2)+(3)} + \delta\Delta_i^{(4)} + \delta\Delta_i^{(5)}. \quad (\text{E165})$$

The concrete expressions of Δ_i can be obtained through Eqs. (E14)-(E25), Eqs. (E107)-(E118), (E125)-(E136), (E153)-(E164) into Eq. (E165).

Appendix F: Derivation of RG equations

The free action of fermions is

$$S_\Psi = \int \frac{dk_0}{2\pi} \frac{d^3\mathbf{k}}{(2\pi)^3} \bar{\Psi}(k_0, \mathbf{k}) \left\{ i [k_0\gamma_0 + v(k_x\gamma_x + k_y\gamma_y) + v_z k_z\gamma_z + D(k_x^2 - k_y^2)\gamma_5] + m - B_\perp k_\perp^2 - B_z k_z^2 \right\} \Psi(k_0, \mathbf{k}). \quad (\text{F1})$$

Considering the corrections of fermion self-energies induced by Coulomb interaction and disorder, the action of fermions becomes

$$\begin{aligned} S_\Psi &= \int \frac{dk_0}{2\pi} \frac{d^3\mathbf{k}}{(2\pi)^3} \bar{\Psi}(k_0, \mathbf{k}) \left\{ i [k_0\gamma_0 + v(k_x\gamma_x + k_y\gamma_y) + v_z k_z\gamma_z + D(k_x^2 - k_y^2)\gamma_5] + m - B_\perp k_\perp^2 - B_z k_z^2 - \Sigma(k_0, \mathbf{k}) - \Sigma_{dis}(k_0) \right\} \Psi(k_0, \mathbf{k}) \\ &\approx \int \frac{dk_0}{2\pi} \frac{d^3\mathbf{k}}{(2\pi)^3} \bar{\Psi}(k_0, \mathbf{k}) \left\{ i \left[k_0\gamma_0 e^{C_0^{dis}\ell} + v(k_x\gamma_x + k_y\gamma_y) e^{C_v\ell} + v_z k_z\gamma_z e^{C_{vz}\ell} + D(k_x^2 - k_y^2)\gamma_5 e^{C_D\ell} \right] + m e^{(C_m + C_m^{dis})\ell} - B_\perp k_\perp^2 e^{-C_{B\perp}\ell} - B_z k_z^2 e^{-C_{Bz}\ell} \right\} \Psi(k_0, \mathbf{k}). \quad (\text{F2}) \end{aligned}$$

Employing the transformations

$$k_0 = k'_0 e^{-\ell}, \quad (\text{F3})$$

$$k_x = k'_x e^{-\ell}, \quad (\text{F4})$$

$$k_y = k'_y e^{-\ell}, \quad (\text{F5})$$

$$k_z = k'_z e^{-\ell}, \quad (\text{F6})$$

$$\Psi = \Psi' e^{\frac{1}{2}(5 - C_0^{dis})\ell}, \quad (\text{F7})$$

$$v = v' e^{-(C_v - C_0^{dis})\ell}, \quad (\text{F8})$$

$$v_z = v'_z e^{-(C_{vz} - C_0^{dis})\ell}, \quad (\text{F9})$$

$$m = m' e^{-(1 + C_m + C_m^{dis} - C_0^{dis})\ell}, \quad (\text{F10})$$

$$B_\perp = B'_\perp e^{(1 + C_{B\perp} + C_0^{dis})\ell}, \quad (\text{F11})$$

$$B_z = B'_z e^{(1 + C_{Bz} + C_0^{dis})\ell}, \quad (\text{F12})$$

$$D = D' e^{(1 - C_D + C_0^{dis})\ell}, \quad (\text{F13})$$

the action of fermions can be further written as

$$\begin{aligned} S_{\Psi'} &= \int \frac{dk'_0}{2\pi} \frac{d^3\mathbf{k}'}{(2\pi)^3} \bar{\Psi}'(k'_0, \mathbf{k}') \left\{ i [k'_0\gamma_0 + v' (k'_x\gamma_x + k'_y\gamma_y) + v'_z k'_z\gamma_z e^{C_{vz}\ell} + D' (k'^2_x - k'^2_y)\gamma_5] \right. \\ &\quad \left. + m' - B'_\perp k'^2_\perp - B'_z k'^2_z \right\} \Psi'(k'_0, \mathbf{k}'), \quad (\text{F14}) \end{aligned}$$

which recovers the original form of fermion action.

The free action of boson ϕ is

$$S_\phi = \int \frac{dk_0}{2\pi} \frac{d^3\mathbf{k}}{(2\pi)^3} \phi(k_0, \mathbf{k}) (k_\perp^2 + \eta k_z^2) \phi(k_0, \mathbf{k}). \quad (\text{F15})$$

Including the correction to self-energy of boson, the action of ϕ becomes

$$\begin{aligned} S_\phi &= \int \frac{dk_0}{2\pi} \frac{d^3\mathbf{k}}{(2\pi)^3} \phi(k_0, \mathbf{k}) (k_\perp^2 + \eta k_z^2 - \Pi(0, \mathbf{k})) \\ &\quad \times \phi(k_0, \mathbf{k}) \\ &\approx \int \frac{dk_0}{2\pi} \frac{d^3\mathbf{k}}{(2\pi)^3} \phi(k_0, \mathbf{k}) (k_\perp^2 e^{C_\perp\ell} + \eta k_z^2 e^{C_z\ell}) \\ &\quad \times \phi(k_0, \mathbf{k}). \quad (\text{F16}) \end{aligned}$$

Using the transformations as shown in Eqs. (F3)-(F6), and

$$\phi = \phi' e^{(3 - \frac{C_\perp}{2})\ell}, \quad (\text{F17})$$

$$\eta = \eta' e^{(C_\perp - C_z)\ell}, \quad (\text{F18})$$

the action of boson can be written as

$$\begin{aligned} S_{\phi'} &= \int \frac{dk'_0}{2\pi} \frac{d^3\mathbf{k}'}{(2\pi)^3} \phi'(k'_0, \mathbf{k}') (k'^2_\perp + \eta' k'^2_z) \\ &\quad \times \phi(k'_0, \mathbf{k}'), \quad (\text{F19}) \end{aligned}$$

which has the same form as the original boson action.

The action of fermion-boson coupling is

$$\begin{aligned} S_{\Psi\phi} &= ig \int \frac{dk_{0,1}}{2\pi} \frac{d^3\mathbf{k}_1}{(2\pi)^3} \frac{dk_{0,2}}{2\pi} \frac{d^3\mathbf{k}_2}{(2\pi)^3} \bar{\Psi}(k_{0,1}, \mathbf{k}_1) \\ &\quad \times \gamma_0 \Psi(k_{0,2}, \mathbf{k}_2) \phi(k_{0,1} - k_{0,2}, \mathbf{k}_1 - \mathbf{k}_2). \quad (\text{F20}) \end{aligned}$$

Considering the vertex correction, the action has the from

$$\begin{aligned}
S_{\Psi\phi} &= i(g + \delta g) \int \frac{dk_{0,1}}{2\pi} \frac{d^3\mathbf{k}_1}{(2\pi)^3} \frac{dk_{0,2}}{2\pi} \frac{d^3\mathbf{k}_2}{(2\pi)^3} \bar{\Psi}(k_{0,1}, \mathbf{k}_1) \\
&\quad \times \gamma_0 \Psi(k_{0,2}, \mathbf{k}_2) \phi(k_{0,1} - k_{0,2}, \mathbf{k}_1 - \mathbf{k}_2) \\
&\approx ig e^{C_0^{dis} \ell} \int \frac{dk_{0,1}}{2\pi} \frac{d^3\mathbf{k}_1}{(2\pi)^3} \frac{dk_{0,2}}{2\pi} \frac{d^3\mathbf{k}_2}{(2\pi)^3} \bar{\Psi}(k_{0,1}, \mathbf{k}_1) \\
&\quad \times \gamma_0 \Psi(k_{0,2}, \mathbf{k}_2) \phi(k_{0,1} - k_{0,2}, \mathbf{k}_1 - \mathbf{k}_2). \quad (F21)
\end{aligned}$$

Applying the transformations as shown in Eqs. (F3)-(F7), (F17), and

$$g = g' e^{\frac{C_{\perp}}{2} \ell}, \quad (F22)$$

the action can be written as

$$\begin{aligned}
S_{\Psi'\phi'} &= ig' \int \frac{dk'_{0,1}}{2\pi} \frac{d^3\mathbf{k}'_1}{(2\pi)^3} \frac{dk'_{0,2}}{2\pi} \frac{d^3\mathbf{k}'_2}{(2\pi)^3} \bar{\Psi}'(k'_{0,1}, \mathbf{k}'_1) \\
&\quad \times \gamma_0 \Psi'(k'_{0,2}, \mathbf{k}'_2) \phi(k'_{0,1} - k'_{0,2}, \mathbf{k}'_1 - \mathbf{k}'_2), \quad (F23)
\end{aligned}$$

which recovers the original form of the action of fermion-boson coupling.

The action of fermion-disorder coupling is given by

$$\begin{aligned}
S_{dis} &= - \sum_{j=0}^3 \frac{\Delta_j}{2} \int \frac{dk_{0,1} dk_{0,2} d^3\mathbf{k}_1 d^3\mathbf{k}_2 d^3\mathbf{k}_3}{(2\pi)^{11}} \\
&\quad \times \bar{\Psi}_a(k_{0,1}, \mathbf{k}_1) \Gamma_j \Psi_a(k_{0,1}, \mathbf{k}_2) \bar{\Psi}_b(k_{0,2}, \mathbf{k}_3) \Gamma_j \\
&\quad \times \Psi_b(k_{0,2}, \mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3). \quad (F24)
\end{aligned}$$

Including the corrections to one-loop order, the action of fermion-disorder coupling becomes

$$\begin{aligned}
S_{dis} &= - \sum_{j=0}^3 \frac{(\Delta_j + \delta\Delta_j)}{2} \int \frac{dk_{0,1} dk_{0,2} d^3\mathbf{k}_1 d^3\mathbf{k}_2 d^3\mathbf{k}_3}{(2\pi)^{11}} \\
&\quad \times \bar{\Psi}_a(k_{0,1}, \mathbf{k}_1) \Gamma_j \Psi_a(k_{0,1}, \mathbf{k}_2) \bar{\Psi}_b(k_{0,2}, \mathbf{k}_3) \Gamma_j \\
&\quad \times \Psi_b(k_{0,2}, \mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3). \quad (F25)
\end{aligned}$$

Using the transformations as shown in Eqs. (F3)-(F7), we obtain

$$\begin{aligned}
S_{dis} &= - \sum_{j=0}^3 \frac{(\Delta_j + \delta\Delta_j)}{2} e^{(-1-2C_0^{dis})\ell} \\
&\quad \times \int \frac{dk'_{0,1} dk'_{0,2} d^3\mathbf{k}'_1 d^3\mathbf{k}'_2 d^3\mathbf{k}'_3}{(2\pi)^{11}} \bar{\Psi}'_a(k'_{0,1}, \mathbf{k}'_1) \Gamma_j \\
&\quad \times \Psi'_a(k'_{0,1}, \mathbf{k}'_2) \bar{\Psi}'_b(k'_{0,2}, \mathbf{k}'_3) \Gamma_j \\
&\quad \times \Psi'_b(k'_{0,2}, \mathbf{k}'_1 + \mathbf{k}'_2 + \mathbf{k}'_3) \\
&\approx - \sum_{j=0}^3 \frac{[\Delta_j + \Delta_j (-1 - 2C_0^{dis}) \ell + \delta\Delta_j]}{2} \\
&\quad \times \int \frac{dk'_{0,1} dk'_{0,2} d^3\mathbf{k}'_1 d^3\mathbf{k}'_2 d^3\mathbf{k}'_3}{(2\pi)^{11}} \bar{\Psi}'_a(k'_{0,1}, \mathbf{k}'_1) \Gamma_j \\
&\quad \times \Psi'_a(k'_{0,1}, \mathbf{k}'_2) \bar{\Psi}'_b(k'_{0,2}, \mathbf{k}'_3) \Gamma_j
\end{aligned}$$

$$\times \Psi'_b(k'_{0,2}, \mathbf{k}'_1 + \mathbf{k}'_2 + \mathbf{k}'_3). \quad (F26)$$

Adopting the transformation

$$\Delta'_j = \Delta_j + \Delta_j (-1 - 2C_0^{dis}) \ell + \delta\Delta_j, \quad (F27)$$

the action for fermion-disorder coupling can be written as

$$\begin{aligned}
S_{dis} &= - \sum_{j=0}^3 \frac{\Delta'_j}{2} \int \frac{dk'_{0,1} dk'_{0,2} d^3\mathbf{k}'_1 d^3\mathbf{k}'_2 d^3\mathbf{k}'_3}{(2\pi)^{11}} \bar{\Psi}'_a(k'_{0,1}, \mathbf{k}'_1) \\
&\quad \times \Gamma_j \Psi'_a(k'_{0,1}, \mathbf{k}'_2) \bar{\Psi}'_b(k'_{0,2}, \mathbf{k}'_3) \Gamma_j \\
&\quad \times \Psi'_b(k'_{0,2}, \mathbf{k}'_1 + \mathbf{k}'_2 + \mathbf{k}'_3), \quad (F28)
\end{aligned}$$

which recovers the original form of the action of fermion-disorder coupling.

From Eqs. (F8)-(F13), (F18), (F22), and (F27), we obtain the RG equations

$$\frac{dv}{d\ell} = (C_v - C_0^{dis}) v, \quad (F29)$$

$$\frac{dv_z}{d\ell} = (C_{v_z} - C_0^{dis}) v_z, \quad (F30)$$

$$\frac{dm}{d\ell} = (1 + C_m + C_m^{dis} - C_0^{dis}) m, \quad (F31)$$

$$\frac{dB_{\perp}}{d\ell} = (-1 - C_{B_{\perp}} - C_0^{dis}) B_{\perp}, \quad (F32)$$

$$\frac{dB_z}{d\ell} = (-1 - C_{B_z} - C_0^{dis}) B_z, \quad (F33)$$

$$\frac{dD}{d\ell} = (-1 + C_D - C_0^{dis}) D, \quad (F34)$$

$$\frac{d\eta}{d\ell} = (C_z - C_{\perp}) \eta, \quad (F35)$$

$$\frac{dg}{d\ell} = -\frac{C_{\perp}}{2} g, \quad (F36)$$

$$\frac{d\alpha}{d\ell} = -\alpha \left[C_v - C_0^{dis} + \frac{1}{2} (C_{\perp} + C_z) \right], \quad (F37)$$

$$\frac{d\zeta}{d\ell} = \left[C_{v_z} - C_v - \frac{1}{2} (C_z - C_{\perp}) \right] \zeta, \quad (F38)$$

$$\frac{d\Delta_j}{d\ell} = \Delta_j (-1 - 2C_0^{dis}) + \frac{d\delta\Delta_j}{d\ell}, \quad (F39)$$

where ζ is defined as

$$\zeta = \frac{v_z}{v\sqrt{\eta}}. \quad (F40)$$

Adopting the transformations

$$\frac{m}{v\Lambda} \rightarrow m, \quad (F41)$$

$$\frac{B_{\perp}\Lambda}{v} \rightarrow B_{\perp}, \quad (F42)$$

$$\frac{B_z\Lambda}{v\eta} \rightarrow B_z, \quad (F43)$$

$$\frac{D\Lambda}{v} \rightarrow D, \quad (F44)$$

$$\Delta_j \frac{\Lambda}{2\pi^2 v^2 \sqrt{\eta}} \rightarrow \Delta_j, \quad (\text{F45})$$

the RG equations can be further compactly written as

$$\frac{dv}{d\ell} = (\alpha \mathcal{R}_v - C_0^{dis}) v, \quad (\text{F46})$$

$$\frac{dv_z}{d\ell} = (\alpha \mathcal{R}_{v_z} - C_0^{dis}) v_z, \quad (\text{F47})$$

$$\frac{dm}{d\ell} = m + \alpha \mathcal{R}_m + C_{dis}^m m, \quad (\text{F48})$$

$$\frac{dB_\perp}{d\ell} = -B_\perp + \alpha \mathcal{R}_{B_\perp} \quad (\text{F49})$$

$$\frac{dB_z}{d\ell} = -B_z + \alpha \mathcal{R}_{B_z} \quad (\text{F50})$$

$$\frac{dD}{d\ell} = -D + \alpha \mathcal{R}_D D, \quad (\text{F51})$$

$$\frac{d\eta}{d\ell} = \alpha \mathcal{R}_\eta \eta, \quad (\text{F52})$$

$$\frac{dg}{d\ell} = -\alpha \mathcal{R}_g g, \quad (\text{F53})$$

$$\frac{d\alpha}{d\ell} = -\alpha^2 \mathcal{R}_\alpha + \alpha C_0^{dis}, \quad (\text{F54})$$

$$\frac{d\zeta}{d\ell} = \alpha \mathcal{R}_\gamma \zeta, \quad (\text{F55})$$

$$\begin{aligned} \frac{d\Delta_C}{d\ell} = & -\Delta_C + 2\Delta_C (\Delta_C + \Delta_M + \Delta_{AC} + 2\Delta_{SO\perp} + \Delta_{SOz} + \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} + 2\Delta_{AMN\perp} \\ & + \Delta_{AMNz} + 2\Delta_{CR\perp} + \Delta_{CRz}) (\mathcal{G}_1^\perp + \mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \\ & + 4 [(\Delta_M \Delta_{SO\perp} + \Delta_{PM} \Delta_{MN\perp} + \Delta_{AMN\perp} \Delta_{CRz} + \Delta_{AMNz} \Delta_{CR\perp}) \mathcal{G}_1^\perp + (\Delta_M \Delta_{SOz} + \Delta_{PM} \Delta_{MNz} \\ & + 2\Delta_{AMN\perp} \Delta_{CR\perp}) \mathcal{G}_1^z + (\Delta_C \Delta_{PM} + 2\Delta_{SO\perp} \Delta_{AMN\perp} + \Delta_{SOz} \Delta_{AMNz}) \mathcal{G}_1^D \\ & + (\Delta_C \Delta_M + 2\Delta_{MN\perp} \Delta_{AMN\perp} + \Delta_{MNz} \Delta_{AMNz}) \mathcal{G}_1^m] \\ & - \Delta_C \alpha \left(2\mathcal{F}_0^\perp + \frac{1}{2}\mathcal{F}_2^\perp + \frac{3}{2}\mathcal{F}_2^z \right), \end{aligned} \quad (\text{F56})$$

$$\begin{aligned} \frac{d\Delta_M}{d\ell} = & -\Delta_M + 2\Delta_M (\Delta_C + \Delta_M - \Delta_{AC} - 2\Delta_{SO\perp} - \Delta_{SOz} - \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} + 2\Delta_{AMN\perp} \\ & + \Delta_{AMNz} - 2\Delta_{CR\perp} - 2\Delta_{CRz}) (-\mathcal{G}_1^\perp - \mathcal{G}_1^z - \mathcal{G}_1^D + \mathcal{G}_1^m) \\ & + 4 [(\Delta_C \Delta_{SO\perp} + \Delta_{PM} \Delta_{AMN\perp} + \Delta_{MN\perp} \Delta_{CRz} + \Delta_{MNz} \Delta_{CR\perp}) \mathcal{G}_1^\perp + (\Delta_C \Delta_{SOz} + \Delta_{PM} \Delta_{AMNz} \\ & + 2\Delta_{MN\perp} \Delta_{CR\perp}) \mathcal{G}_1^z + (\Delta_M \Delta_{PM} + 2\Delta_{SO\perp} \Delta_{MN\perp} + \Delta_{SOz} \Delta_{MNz}) \mathcal{G}_1^D \\ & + (\Delta_C^2 + \Delta_M^2 + \Delta_{AC}^2 + 2\Delta_{SO\perp}^2 + \Delta_{SOz}^2 + \Delta_{PM}^2 + 2\Delta_{MN\perp}^2 + \Delta_{MNz}^2 + 2\Delta_{AMN\perp}^2 + \Delta_{AMNz}^2 \\ & + 2\Delta_{CR\perp}^2 + \Delta_{CRz}^2) \mathcal{G}_1^m] \\ & + 2\Delta_M \alpha \left[\frac{1}{2} (-\mathcal{F}_0^\perp + \mathcal{F}_0^z) + \frac{1}{4}\mathcal{F}_2^\perp - \frac{1}{4}\mathcal{F}_2^z + 2\mathcal{F}_3^\perp + \mathcal{F}_3^z + \mathcal{F}_3^D - \mathcal{F}_3^m \right], \end{aligned} \quad (\text{F57})$$

$$\begin{aligned} \frac{d\Delta_{AC}}{d\ell} = & -\Delta_{AC} + 2\Delta_{AC} (-\Delta_C + \Delta_M - \Delta_{AC} + 2\Delta_{SO\perp} + \Delta_{SOz} + \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} \\ & - 2\Delta_{AMN\perp} - \Delta_{AMNz} - 2\Delta_{CR\perp} - \Delta_{CRz}) (-\mathcal{G}_1^\perp - \mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \\ & + 4 [(\Delta_{SO\perp} \Delta_{SOz} + \Delta_{MN\perp} \Delta_{MNz} + \Delta_{AMN\perp} \Delta_{AMNz} + \Delta_{CR\perp} \Delta_{CRz}) \mathcal{G}_1^\perp \\ & + 2 (\Delta_{SO\perp}^2 + \Delta_{MN\perp}^2 + \Delta_{AMN\perp}^2 + \Delta_{CR\perp}^2) \mathcal{G}_1^z + (\Delta_{AC} \Delta_{PM} + 2\Delta_{SO\perp} \Delta_{CR\perp} + \Delta_{SOz} \Delta_{CRz}) \mathcal{G}_1^D \\ & + (\Delta_M \Delta_{AC} + \Delta_{MNz} \Delta_{CRz} + 2\Delta_{MN\perp} \Delta_{CR\perp}) \mathcal{G}_1^m] \\ & + 2\Delta_{AC} \alpha \left[\frac{1}{2} (-\mathcal{F}_0^\perp + \mathcal{F}_0^z) + \frac{1}{4}\mathcal{F}_2^\perp - \frac{1}{4}\mathcal{F}_2^z - 2\mathcal{F}_3^\perp - \mathcal{F}_3^z + \mathcal{F}_3^D + \mathcal{F}_3^m \right], \end{aligned} \quad (\text{F58})$$

$$\begin{aligned} \frac{d\Delta_{SO\perp}}{d\ell} = & -\Delta_{SO\perp} + 2\Delta_{SO\perp} (-\Delta_C + \Delta_M + \Delta_{AC} + \Delta_{SOz} - \Delta_{PM} - \Delta_{MNz} + \Delta_{AMNz} - \Delta_{CRz}) \\ & \times (-\mathcal{G}_1^z - \mathcal{G}_1^D + \mathcal{G}_1^m) \\ & + 2 [(\Delta_C \Delta_M + \Delta_{AC} \Delta_{SOz} + \Delta_{PM} \Delta_{CRz} + 4\Delta_{MN\perp} \Delta_{AMN\perp} + \Delta_{MNz} \Delta_{AMNz}) \mathcal{G}_1^\perp \\ & + 2 (\Delta_{AC} \Delta_{SO\perp} + \Delta_{PM} \Delta_{CR\perp} + \Delta_{MN\perp} \Delta_{AMNz} + \Delta_{MNz} \Delta_{AMN\perp}) \mathcal{G}_1^z \\ & + 2 (\Delta_C \Delta_{AMN\perp} + \Delta_M \Delta_{MN\perp} + \Delta_{AC} \Delta_{CR\perp} + \Delta_{SO\perp} \Delta_{PM}) \mathcal{G}_1^D \\ & + 2 (\Delta_M \Delta_{SO\perp} + \Delta_{PM} \Delta_{MN\perp} + \Delta_{AMN\perp} \Delta_{CRz} + \Delta_{AMNz} \Delta_{CR\perp}) \mathcal{G}_1^m] \end{aligned}$$

$$+2\Delta_{SO\perp}\alpha\left[\frac{1}{2}(-\mathcal{F}_0^\perp+\mathcal{F}_0^z)+\frac{1}{4}\mathcal{F}_2^\perp-\frac{1}{4}\mathcal{F}_2^z-\mathcal{F}_3^z-\mathcal{F}_3^D+\mathcal{F}_3^m\right], \quad (\text{F59})$$

$$\begin{aligned} \frac{d\Delta_{SOz}}{d\ell} = & -\Delta_{SOz}+2\Delta_{SOz}(-\Delta_C+\Delta_M+\Delta_{AC}+2\Delta_{SO\perp}-\Delta_{SOz}-\Delta_{PM}-2\Delta_{MN\perp}+\Delta_{MNz} \\ & +2\Delta_{AMN\perp}-\Delta_{AMNz}-2\Delta_{CR\perp}+\Delta_{CRz})(-\mathcal{G}_1^\perp+\mathcal{G}_1^z-\mathcal{G}_1^D+\mathcal{G}_1^m) \\ & +4[(\Delta_{AC}\Delta_{SO\perp}+\Delta_{PM}\Delta_{CR\perp}+\Delta_{MN\perp}\Delta_{AMNz}+\Delta_{MNz}\Delta_{AMN\perp})\mathcal{G}_1^\perp \\ & +(\Delta_C\Delta_M+2\Delta_{MN\perp}\Delta_{AMN\perp}+\Delta_{MNz}\Delta_{AMNz})\mathcal{G}_1^z \\ & +(\Delta_C\Delta_{AMNz}+\Delta_M\Delta_{MNz}+\Delta_{AC}\Delta_{CRz}+\Delta_{SOz}\Delta_{PM})\mathcal{G}_1^D \\ & +(\Delta_M\Delta_{SOz}+\Delta_{PM}\Delta_{MNz}+2\Delta_{AMN\perp}\Delta_{CR\perp})\mathcal{G}_1^m] \\ & +2\Delta_{SOz}\alpha\left[\frac{1}{2}(-\mathcal{F}_0^\perp+\mathcal{F}_0^z)+\frac{1}{4}\mathcal{F}_2^\perp-\frac{1}{4}\mathcal{F}_2^z-2\mathcal{F}_3^\perp+\mathcal{F}_3^z-\mathcal{F}_3^D+\mathcal{F}_3^m\right], \end{aligned} \quad (\text{F60})$$

$$\begin{aligned} \frac{d\Delta_{PM}}{d\ell} = & -\Delta_{PM}+2\Delta_{PM}(-\Delta_C+\Delta_M+\Delta_{AC}-2\Delta_{SO\perp}-\Delta_{SOz}-\Delta_{PM}+2\Delta_{MN\perp}+\Delta_{MNz} \\ & -2\Delta_{AMN\perp}-\Delta_{AMNz}+2\Delta_{CR\perp}+\Delta_{CRz})(\mathcal{G}_1^\perp+\mathcal{G}_1^z-\mathcal{G}_1^D+\mathcal{G}_1^m) \\ & +4[(\Delta_C\Delta_{MN\perp}+\Delta_M\Delta_{AMN\perp}+\Delta_{SO\perp}\Delta_{CRz}+\Delta_{SOz}\Delta_{CR\perp})\mathcal{G}_1^\perp \\ & +(\Delta_C\Delta_{MNz}+\Delta_M\Delta_{AMNz}+2\Delta_{SO\perp}\Delta_{CR\perp})\mathcal{G}_1^z+(\Delta_C^2+\Delta_M^2+\Delta_{AC}^2+2\Delta_{SO\perp}^2+\Delta_{SOz}^2 \\ & +\Delta_{PM}^2+2\Delta_{MN\perp}^2+\Delta_{MNz}^2+2\Delta_{AMN\perp}^2+\Delta_{AMNz}^2+2\Delta_{CR\perp}^2+\Delta_{CRz}^2)\mathcal{G}_1^D \\ & +(\Delta_M\Delta_{PM}+2\Delta_{SO\perp}\Delta_{MN\perp}+\Delta_{SOz}\Delta_{MNz})\mathcal{G}_1^m] \\ & +2\Delta_{PM}\alpha\left[\frac{1}{2}(-\mathcal{F}_0^\perp+\mathcal{F}_0^z)+\frac{1}{4}\mathcal{F}_2^\perp-\frac{1}{4}\mathcal{F}_2^z+2\mathcal{F}_3^\perp+\mathcal{F}_3^z-\mathcal{F}_3^D+\mathcal{F}_3^m\right], \end{aligned} \quad (\text{F61})$$

$$\begin{aligned} \frac{d\Delta_{MN\perp}}{d\ell} = & -\Delta_{MN\perp}+2\Delta_{MN\perp}(\Delta_C+\Delta_M-\Delta_{AC}+\Delta_{SOz}-\Delta_{PM}-\Delta_{MNz}-\Delta_{AMNz}+\Delta_{CRz}) \\ & \times(\mathcal{G}_1^z-\mathcal{G}_1^D+\mathcal{G}_1^m) \\ & +2[(\Delta_C\Delta_{PM}+\Delta_M\Delta_{CRz}+\Delta_{AC}\Delta_{MNz}+4\Delta_{SO\perp}\Delta_{AMN\perp}+\Delta_{SOz}\Delta_{AMNz})\mathcal{G}_1^\perp \\ & +2(\Delta_M\Delta_{CR\perp}+\Delta_{AC}\Delta_{MN\perp}+\Delta_{SO\perp}\Delta_{AMNz}+\Delta_{SOz}\Delta_{AMN\perp})\mathcal{G}_1^z \\ & +2(\Delta_M\Delta_{SO\perp}+\Delta_{PM}\Delta_{MN\perp}+\Delta_{AMN\perp}\Delta_{CRz}+\Delta_{AMNz}\Delta_{CR\perp})\mathcal{G}_1^D \\ & +2(\Delta_C\Delta_{AMN\perp}+\Delta_M\Delta_{MN\perp}+\Delta_{AC}\Delta_{CR\perp}+\Delta_{SO\perp}\Delta_{PM})\mathcal{G}_1^m] \\ & +2\Delta_{MN\perp}\alpha\left[\frac{1}{2}(-\mathcal{F}_0^\perp+\mathcal{F}_0^z)+\frac{1}{4}\mathcal{F}_2^\perp-\frac{1}{4}\mathcal{F}_2^z-\mathcal{F}_3^z+\mathcal{F}_3^D-\mathcal{F}_3^m\right], \end{aligned} \quad (\text{F62})$$

$$\begin{aligned} \frac{d\Delta_{MNz}}{d\ell} = & -\Delta_{MNz}+2\Delta_{MNz}(\Delta_C+\Delta_M-\Delta_{AC}+2\Delta_{SO\perp}-\Delta_{SOz}-\Delta_{PM}-2\Delta_{MN\perp}+\Delta_{MNz} \\ & -2\Delta_{AMN\perp}+\Delta_{AMNz}+2\Delta_{CR\perp}-\Delta_{CRz})(\mathcal{G}_1^\perp-\mathcal{G}_1^z-\mathcal{G}_1^D+\mathcal{G}_1^m) \\ & +4[(\Delta_M\Delta_{CR\perp}+\Delta_{AC}\Delta_{MN\perp}+\Delta_{SO\perp}\Delta_{AMNz}+\Delta_{SOz}\Delta_{AMN\perp})\mathcal{G}_1^\perp \\ & +(\Delta_C\Delta_{PM}+2\Delta_{SO\perp}\Delta_{AMN\perp}+\Delta_{SOz}\Delta_{AMNz})\mathcal{G}_1^z \\ & +(\Delta_M\Delta_{SOz}+\Delta_{PM}\Delta_{MNz}+2\Delta_{AMN\perp}\Delta_{CR\perp})\mathcal{G}_1^D \\ & +(\Delta_C\Delta_{AMNz}+\Delta_M\Delta_{MNz}+\Delta_{AC}\Delta_{CRz}+\Delta_{SOz}\Delta_{PM})\mathcal{G}_1^m] \\ & +2\Delta_{MNz}\alpha\left[\frac{1}{2}(-\mathcal{F}_0^\perp+\mathcal{F}_0^z)+\frac{1}{4}\mathcal{F}_2^\perp-\frac{1}{4}\mathcal{F}_2^z-2\mathcal{F}_3^\perp+\mathcal{F}_3^z+\mathcal{F}_3^D-\mathcal{F}_3^m\right], \end{aligned} \quad (\text{F63})$$

$$\begin{aligned} \frac{d\Delta_{AMN\perp}}{d\ell} = & -\Delta_{AMN\perp}+2\Delta_{AMN\perp}(\Delta_C+\Delta_M+\Delta_{AC}-\Delta_{SOz}+\Delta_{PM}-\Delta_{MNz}-\Delta_{AMNz}-\Delta_{CRz}) \\ & \times(-\mathcal{G}_1^z+\mathcal{G}_1^D+\mathcal{G}_1^m) \\ & +2[(\Delta_C\Delta_{CRz}+\Delta_M\Delta_{PM}+\Delta_{AC}\Delta_{AMNz}+4\Delta_{SO\perp}\Delta_{MN\perp}+\Delta_{SOz}\Delta_{MNz})\mathcal{G}_1^\perp \\ & +2(\Delta_C\Delta_{CR\perp}+\Delta_{AC}\Delta_{AMN\perp}+\Delta_{SO\perp}\Delta_{MNz}+\Delta_{SOz}\Delta_{MN\perp})\mathcal{G}_1^z \\ & +2(\Delta_C\Delta_{SO\perp}+\Delta_{PM}\Delta_{AMN\perp}+\Delta_{MN\perp}\Delta_{CRz}+\Delta_{MNz}\Delta_{CR\perp})\mathcal{G}_1^D \\ & +2(\Delta_C\Delta_{MN\perp}+\Delta_M\Delta_{AMN\perp}+\Delta_{SO\perp}\Delta_{CRz}+\Delta_{SOz}\Delta_{CR\perp})\mathcal{G}_1^m] \\ & +2\Delta_{AMN\perp}\alpha\left[\frac{1}{2}(-\mathcal{F}_0^\perp+\mathcal{F}_0^z)+\frac{1}{4}\mathcal{F}_2^\perp-\frac{1}{4}\mathcal{F}_2^z+\mathcal{F}_3^z-\mathcal{F}_3^D-\mathcal{F}_3^m\right], \end{aligned} \quad (\text{F64})$$

$$\begin{aligned}
\frac{d\Delta_{AMNz}}{d\ell} = & -\Delta_{AMNz} + 2\Delta_{AMNz}(\Delta_C + \Delta_M + \Delta_{AC} - 2\Delta_{SO\perp} + \Delta_{SOz} + \Delta_{PM} - 2\Delta_{MN\perp} + \Delta_{MNz} \\
& - 2\Delta_{AMN\perp} + \Delta_{AMNz} - 2\Delta_{CR\perp} + \Delta_{CRz})(-\mathcal{G}_1^\perp + \mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 4[(\Delta_C\Delta_{CR\perp} + \Delta_{AC}\Delta_{AMN\perp} + \Delta_{SO\perp}\Delta_{MNz} + \Delta_{SOz}\Delta_{MN\perp})\mathcal{G}_1^\perp \\
& + (\Delta_M\Delta_{PM} + 2\Delta_{SO\perp}\Delta_{MN\perp} + \Delta_{SOz}\Delta_{MNz})\mathcal{G}_1^z \\
& + (\Delta_C\Delta_{SOz} + \Delta_{PM}\Delta_{AMNz} + 2\Delta_{MN\perp}\Delta_{CR\perp})\mathcal{G}_1^D \\
& + (\Delta_C\Delta_{MNz} + \Delta_M\Delta_{AMNz} + 2\Delta_{SO\perp}\Delta_{CR\perp})\mathcal{G}_1^m] \\
& + 2\Delta_{AMNz}\alpha\left[\frac{1}{2}(-\mathcal{F}_0^\perp + \mathcal{F}_0^z) + \frac{1}{4}\mathcal{F}_2^\perp - \frac{1}{4}\mathcal{F}_2^z + 2\mathcal{F}_3^\perp - \mathcal{F}_3^z - \mathcal{F}_3^D - \mathcal{F}_3^m\right], \tag{F65}
\end{aligned}$$

$$\begin{aligned}
\frac{d\Delta_{CR\perp}}{d\ell} = & -\Delta_{CR\perp} + 2\Delta_{CR\perp}(-\Delta_C + \Delta_M - \Delta_{AC} - \Delta_{SOz} + \Delta_{PM} - \Delta_{MNz} + \Delta_{AMNz} + \Delta_{CRz}) \\
& \times (\mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 2[(\Delta_C\Delta_{AMNz} + \Delta_M\Delta_{MNz} + \Delta_{AC}\Delta_{CRz} + \Delta_{SOz}\Delta_{PM})\mathcal{G}_1^\perp \\
& + 2(\Delta_C\Delta_{AMN\perp} + \Delta_M\Delta_{MN\perp} + \Delta_{AC}\Delta_{CR\perp} + \Delta_{SO\perp}\Delta_{PM})\mathcal{G}_1^z \\
& + 2(\Delta_{AC}\Delta_{SO\perp} + \Delta_{PM}\Delta_{CR\perp} + \Delta_{MN\perp}\Delta_{AMNz} + \Delta_{MNz}\Delta_{AMN\perp})\mathcal{G}_1^D \\
& + 2(\Delta_M\Delta_{CR\perp} + \Delta_{AC}\Delta_{MN\perp} + \Delta_{SO\perp}\Delta_{AMNz} + \Delta_{SOz}\Delta_{AMN\perp})\mathcal{G}_1^m] \\
& + 2\Delta_{CR\perp}\alpha\left[\frac{1}{2}(-\mathcal{F}_0^\perp + \mathcal{F}_0^z) + \frac{1}{4}\mathcal{F}_2^\perp - \frac{1}{4}\mathcal{F}_2^z + \mathcal{F}_3^\perp + \mathcal{F}_3^z + \mathcal{F}_3^D + \mathcal{F}_3^m\right], \tag{F66}
\end{aligned}$$

$$\begin{aligned}
\frac{d\Delta_{CRz}}{d\ell} = & -\Delta_{CRz} + 2\Delta_{CRz}(-\Delta_C + \Delta_M - \Delta_{AC} - 2\Delta_{SO\perp} + \Delta_{SOz} + \Delta_{PM} - 2\Delta_{MN\perp} + \Delta_{MNz} \\
& + 2\Delta_{AMN\perp} - \Delta_{AMNz} + 2\Delta_{CR\perp} - \Delta_{CRz})(\mathcal{G}_1^\perp - \mathcal{G}_1^z + \mathcal{G}_1^D + \mathcal{G}_1^m) \\
& + 4[(\Delta_C\Delta_{AMN\perp} + \Delta_M\Delta_{MN\perp} + \Delta_{AC}\Delta_{CR\perp} + \Delta_{SO\perp}\Delta_{PM})\mathcal{G}_1^\perp \\
& + (\Delta_{AC}\Delta_{SOz} + \Delta_{PM}\Delta_{CRz} + 2\Delta_{MN\perp}\Delta_{AMN\perp})\mathcal{G}_1^D \\
& + (\Delta_M\Delta_{CRz} + \Delta_{AC}\Delta_{MNz} + 2\Delta_{SO\perp}\Delta_{AMN\perp})\mathcal{G}_1^m] \\
& + 2\Delta_{CRz}\alpha\left[\frac{1}{2}(-\mathcal{F}_0^\perp + \mathcal{F}_0^z) + \frac{1}{4}\mathcal{F}_2^\perp - \frac{1}{4}\mathcal{F}_2^z + 2\mathcal{F}_3^\perp - \mathcal{F}_3^z + \mathcal{F}_3^D + \mathcal{F}_3^m\right], \tag{F67}
\end{aligned}$$

where

$$\mathcal{R}_v = \mathcal{F}_0^\perp, \tag{F68}$$

$$\mathcal{R}_{vz} = 2\mathcal{F}_0^z, \tag{F69}$$

$$\mathcal{R}_m = m\mathcal{F}_0^z - B_\perp\mathcal{F}_0^\perp - B_z\mathcal{F}_0^z, \tag{F70}$$

$$\begin{aligned}
\mathcal{R}_{B\perp} = & -m(\mathcal{F}_1^\perp + \mathcal{F}_1^z) + B_\perp(\mathcal{F}_1^\perp - \mathcal{F}_0^\perp) \\
& + B_z\mathcal{F}_1^z, \tag{F71}
\end{aligned}$$

$$\begin{aligned}
\mathcal{R}_{Bz} = & -m[\mathcal{F}_0^\perp + \mathcal{F}_0^z - 2(\mathcal{F}_1^\perp + \mathcal{F}_1^z)] \\
& + B_\perp(\mathcal{F}_0^\perp - 2\mathcal{F}_1^\perp) \\
& + B_z(-\mathcal{F}_0^\perp + \mathcal{F}_0^z - 2\mathcal{F}_1^z - \mathcal{F}_2^z + \mathcal{F}_2^\perp), \tag{F72}
\end{aligned}$$

$$\mathcal{R}_D = (\mathcal{F}_1^D - \mathcal{F}_0^\perp), \tag{F73}$$

$$\mathcal{R}_\eta = (\mathcal{F}_2^z - \mathcal{F}_2^\perp), \tag{F74}$$

$$\mathcal{R}_g = \frac{1}{2}\mathcal{F}_2^\perp, \tag{F75}$$

$$\mathcal{R}_\alpha = \mathcal{F}_0^\perp + \frac{1}{2}(\mathcal{F}_2^\perp + \mathcal{F}_2^z), \tag{F76}$$

$$\mathcal{R}_\zeta = 2\mathcal{F}_0^z - \mathcal{F}_0^\perp - \frac{1}{2}(\mathcal{F}_2^z - \mathcal{F}_2^\perp), \tag{F77}$$

$$\begin{aligned}
C_0^{dis} = & (\Delta_C + \Delta_M + \Delta_{AC} + 2\Delta_{SO\perp} + \Delta_{SOz} \\
& + \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} + 2\Delta_{AMN\perp}
\end{aligned}$$

$$\begin{aligned}
& + \Delta_{AMNz} + 2\Delta_{CR\perp} + \Delta_{CRz}) \\
& \times (\mathcal{G}_0^\perp + \mathcal{G}_0^z), \tag{F78}
\end{aligned}$$

$$\begin{aligned}
C_m^{dis} = & -(\Delta_C + \Delta_M - \Delta_{AC} - 2\Delta_{SO\perp} - \Delta_{SOz} \\
& - \Delta_{PM} + 2\Delta_{MN\perp} + \Delta_{MNz} + 2\Delta_{AMN\perp} \\
& + \Delta_{AMNz} - 2\Delta_{CR\perp} - \Delta_{CRz}) \\
& \times \frac{1}{m}[m(\mathcal{G}_0^\perp + \mathcal{G}_0^z) - B_\perp\mathcal{G}_0^\perp - B_z\mathcal{G}_0^z], \tag{F79}
\end{aligned}$$

with

$$\mathcal{F}_0^\perp = \frac{1}{4\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \frac{\sin^2(\varphi)}{\Xi}, \tag{F80}$$

$$\mathcal{F}_0^z = \frac{1}{4\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \frac{\cos^2(\varphi)}{\Xi}, \tag{F81}$$

$$\begin{aligned}
\mathcal{F}_1^\perp = & -\frac{1}{4\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \\
& \times \frac{\sin^2(\varphi) \cos(2\varphi)}{\Xi}, \tag{F82}
\end{aligned}$$

$$\mathcal{F}_1^z = -\frac{1}{4\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta$$

$$\times \frac{\cos^2(\varphi) \cos(2\varphi)}{\Xi}, \quad (\text{F83})$$

$$\times \frac{\sin^4(\varphi) \cos^2(2\theta)}{\Xi}, \quad (\text{F84})$$

$$\mathcal{F}_1^D = \frac{1}{2\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta$$

$$\mathcal{F}_2^x = \frac{1}{4\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \left\{ \frac{1 + 4(D^2 + B_\perp^2) \sin^2(\varphi) \cos^2(\theta)}{\Xi^3} - \frac{[1 + 2D^2 \sin^2(\varphi) \cos(2\theta) - 2mB_\perp + 2B_\perp^2 \sin^2(\varphi) + 2B_\perp B_z \cos^2(\varphi)]^2 \sin^2(\varphi) \cos^2(\theta)}{\Xi^5} \right\}, \quad (\text{F85})$$

$$\mathcal{F}_2^y = \frac{1}{4\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \left\{ \frac{1 + 4(D^2 + B_\perp^2) \sin^2(\varphi) \sin^2(\theta)}{\Xi^3} - \frac{[1 - 2D^2 \sin^2(\varphi) \cos(2\theta) - 2mB_\perp + 2B_\perp^2 \sin^2(\varphi) + 2B_\perp B_z \cos^2(\varphi)]^2 \sin^2(\varphi) \sin^2(\theta)}{\Xi^5} \right\}, \quad (\text{F86})$$

$$\mathcal{F}_2^z = \frac{1}{4\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \left\{ \frac{\zeta^2 + 4B_z^2 \cos^2(\varphi)}{\Xi^3} - \frac{(\zeta^2 - 2mB_z + 2B_z^2 \cos^2(\varphi) + 2B_z B_\perp \sin^2(\varphi))^2 \cos^2(\varphi)}{\Xi^5} \right\}, \quad (\text{F87})$$

$$\mathcal{F}_3^\perp = \frac{1}{16\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \frac{\sin^2(\varphi)}{\Xi^3}, \quad (\text{F88})$$

$$\mathcal{F}_3^z = \frac{1}{8\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \frac{\zeta^2 \cos^2(\varphi)}{\Xi^3}, \quad (\text{F89})$$

$$\mathcal{F}_3^D = \frac{1}{8\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \times \frac{D^2 \sin^4(\varphi) \cos^2(2\theta)}{\Xi^3}, \quad (\text{F90})$$

$$\mathcal{F}_3^m = \frac{1}{8\pi} \int_0^\pi \sin(\varphi) d\varphi \int_0^{2\pi} d\theta \times \frac{(m - B_\perp \sin^2(\varphi) - B_z \cos^2(\varphi))^2}{\Xi^3}, \quad (\text{F91})$$

$$\mathcal{G}_0^\perp = \frac{1}{4\pi} \int_0^\pi d\varphi \sin(\varphi) \int_0^{2\pi} d\theta \frac{\sin^2(\varphi)}{\Xi^2}, \quad (\text{F92})$$

$$\mathcal{G}_0^z = \frac{1}{4\pi} \int_0^\pi d\varphi \sin(\varphi) \int_0^{2\pi} d\theta \frac{\cos^2(\varphi)}{\Xi^2}, \quad (\text{F93})$$

$$\mathcal{G}_1^\perp = \frac{1}{4\pi} \int_0^\pi d\varphi \sin(\varphi) \int_0^{2\pi} d\theta \frac{\sin^2(\varphi)}{\Xi^4}, \quad (\text{F94})$$

$$\mathcal{G}_1^z = \frac{1}{4\pi} \int_0^\pi d\varphi \sin(\varphi) \int_0^{2\pi} d\theta \frac{\zeta^2 \cos^2(\varphi)}{\Xi^4}, \quad (\text{F95})$$

$$\mathcal{G}_1^D = \frac{1}{4\pi} \int_0^\pi d\varphi \sin(\varphi) \int_0^{2\pi} d\theta$$

$$\times \frac{D^2 \sin^4(\varphi) \cos^2(2\theta)}{\Xi^4}, \quad (\text{F96})$$

$$\mathcal{G}_1^m = \frac{1}{4\pi} \int_0^\pi d\varphi \sin(\varphi) \int_0^{2\pi} d\theta \times \frac{(m - B_\perp \sin^2(\varphi) - B_z \cos^2(\varphi))^2}{\Xi^4}, \quad (\text{F97})$$

where

$$\Xi = \left[\sin^2(\varphi) + \zeta^2 \cos^2(\varphi) + D^2 \sin^4(\varphi) \cos^2(2\theta) + (m - B_\perp \sin^2(\varphi) - B_z \cos^2(\varphi))^2 \right]^{\frac{1}{2}}. \quad (\text{F98})$$

Employing the relation

$$\mathcal{F}_1^\perp + \mathcal{F}_1^z = \mathcal{F}_0^\perp - \mathcal{F}_0^z, \quad (\text{F99})$$

we can find that $\mathcal{R}_{B_z}$ can be also written as

$$\mathcal{R}_{B_z} = -m(-\mathcal{F}_0^\perp + 3\mathcal{F}_0^z) + B_\perp(\mathcal{F}_0^\perp - 2\mathcal{F}_1^\perp) + B_z(-\mathcal{F}_1^\perp - 3\mathcal{F}_1^z - \mathcal{F}_2^z + \mathcal{F}_2^\perp). \quad (\text{F100})$$

[1] M. Z. Hasan and C. L. Kane, Colloquium: Topological insulators, Rev. Mod. Phys. **82**, 3045 (2010).

[2] X.-L. Qi and S.-C. Zhang, Topological insulators and superconductors, Rev. Mod. Phys. **83**, 1057 (2011).

- [3] O. Vafek and A. Vishwanath, Dirac fermions in solids: From high- T_c cuprates and graphene to topological insulators and Weyl semimetals, *Annu. Rev. Condens. Matter Phys.* **5**, 83 (2014).
- [4] T. O. Wehling, A. M. Black-Schaffer, and A. V. Balatsky, Dirac materials, *Adv. Phys.* **63**, 1 (2014).
- [5] M. Sato and Y. Ando, Topological superconductors: a review, *Rep. Prog. Phys.* **80**, 076501 (2017).
- [6] B. Yan and C. Felser, Topological materials: Weyl semimetals, *Annu. Rev. Condens. Matter Phys.* **8**, 337 (2017).
- [7] M. Z. Hasan, S.-Y. Xu, I. Belopolski, and S.-M. Huang, Discovery of Weyl fermion semimetals and topological Fermi arc states, *Annu. Rev. Condens. Matter Phys.* **8**, 289 (2017).
- [8] N. P. Armitage, E. J. Mele, and A. Vishwanath, Weyl and Dirac semimetals in three-dimensional solids, *Rev. Mod. Phys.* **90**, 015001 (2018).
- [9] B. Q. Lv, T. Qian, and H. Ding, Experimental perspective on three-dimensional topological semimetals, *Rev. Mod. Phys.* **93**, 025002 (2021).
- [10] M. Z. Hasan, G. Chang, I. Belopolski, G. Bian, S.-Y. Xu, J.-X. Yin, Weyl, Dirac and high-fold chiral fermions in topological quantum matter, *Nat. Rev. Mat.* **6**, 784 (2021).
- [11] B. J. Wieder, B. Bradlyn, J. Cano, Z. Wang, M. G. Vergniorty, L. Elcoro, A. A. Soluyanov, C. Felser, T. Neuper, N. Regnault, and B. A. Bernevig, Topological materials discovery from crystal symmetry, *Nat. Rev. Phys.* **7**, 196 (2022).
- [12] B. Xie, H.-X. Wang, X. Zhang, P. Zhan, J.-H. Jiang, M. Lu, and Y. Chen, Higher-order band topology, *Nat. Rev. Phys.* **3**, 520 (2021).
- [13] W. A. Benalcazar, B. A. Bernevig, and T. L. Hughes, Quantized electric multipole insulators, *Science* **357**, 61 (2017).
- [14] W. A. Benalcazar, B. A. Bernevig, and T. L. Hughes, Electric multipole moments, topological multipole moment pumping, and chiral hinge states in crystalline insulators, *Phys. Rev. B* **96**, 245115 (2017).
- [15] J. Langbehn, Y. Peng, L. Trifunovic, F. v. Oppen, and P. W. Brouwer, Reflection-Symmetric Second-Order Topological Insulators and Superconductors, *Phys. Rev. Lett.* **119**, 246401 (2017).
- [16] Z. Song, Z. Fang, and C. Fang, $(d - 2)$ -Dimensional Edge States of Rotation Symmetry Protected Topological States, *Phys. Rev. Lett.* **119**, 246402 (2017).
- [17] F. Schindler, A. M. Cook, M. G. Vergniorty, Z. Wang, S. S. P. Parkin, B. A. Bernevig, and T. Neupert, Higher-order topological insulators, *Sci. Adv.* **4**, eaat0346 (2018).
- [18] M. Ezawa, Higher-Order Topological Insulators and Semimetals on the Breathing Kagome and Pyrochlore Lattices, *Phys. Rev. Lett.* **120**, 026801 (2018).
- [19] R. Chen, C.-Z. Chen, J.-H. Gao, B. Zhou, and D.-H. Xu, Higher-Order Topological Insulators in Quasicrystals, *Phys. Rev. Lett.* **124**, 036803 (2020).
- [20] C.-A. Li, B. Fu, Z.-A. Hu, J. Li, and S.-Q. Shen, Higher-Order Topological Insulators and Semimetals on the Breathing Kagome and Pyrochlore Lattices, *Phys. Rev. Lett.* **125**, 166801 (2020).
- [21] B. Roy and V. Jurićić, Dislocation as a bulk probe of higher-order topological insulators, *Phys. Rev. Research* **3**, 033107 (2021).
- [22] B. Fu, Z.-A. Hu, and S.-Q. Shen, Bulk-hinge correspondence and three-dimensional quantum anomalous Hall effect in second-order topological insulators, *Phys. Rev. Research* **3**, 033177 (2021).
- [23] Y.-S. Hu, Y.-R. Ding, J. Zhang, Z.-Q. Zhang, and C.-Z. Chen, Disorder and phase diagrams of higher-order topological insulators, *Phys. Rev. B* **104**, 094201 (2021).
- [24] C. Wang and X.R. Wang, Disorder-induced quantum phase transitions in three-dimensional second-order topological insulators, *Phys. Rev. Research* **2**, 033521 (2020).
- [25] C. Wang and X. R. Wang, Robustness of helical hinge states of weak second-order topological insulators, *Phys. Rev. B* **103**, 115118 (2021).
- [26] S. A. A. Ghorashi, T. Li, and T. L. Hughes, Higher-Order Weyl Semimetals, *Phys. Rev. Lett.* **125**, 266804 (2020).
- [27] H.-X. Wang, Z.-K. Lin, B. Jiang, G.-Y. Guo, and J.-H. Jiang, Higher-Order Weyl Semimetals, *Phys. Rev. Lett.* **125**, 146401 (2020).
- [28] A. L. Szabo and B. Roy, Dirty higher-order Dirac semimetal: Quantum criticality and bulk-boundary correspondence, *Phys. Rev. Research* **2**, 043197 (2020).
- [29] Z.-Q. Zhang, B.-L. Wu, C.-Z. Chen, and H. Jiang, Global phase diagram of disordered higher-order Weyl semimetals, *Phys. Rev. B* **104**, 014203 (2021).
- [30] D. González-Cuadra, Higher-order topological quantum paramagnets, *Phys. Rev. B* **105**, L020403 (2022).
- [31] V. N. Kotov, B. Uchoa, V. M. Pereira, F. Guinea, and A. H. Castro Neto, Electron-electron interactions in graphene: Current status and perspectives, *Rev. Mod. Phys.* **84**, 1067 (2012).
- [32] D. C. Elias, R. V. Gorbachev, A. S. Mayorov, S. V. Morozov, A. A. Zhukov, P. Blake, L. A. Ponomarenko, I. V. Grigorieva, K. S. Novoselov, F. Guinea, and A. K. Geim, Dirac cones reshaped by interaction effects in suspended graphene, *Nat. Phys.* **7**, 701 (2011).
- [33] D. A. Siegel, C.-H. Park, C. Hwang, J. Deslippe, A. V. Fedorov, S. G. Louie, and A. Lanzara, Many-body interactions in quasifreestanding graphene, *Proc. Natl. Acad. Sci. U.S.A.* **108**, 11365 (2011).
- [34] G. L. Yu, R. Jalil, B. Belle, A. S. Mayorov, P. Blake, F. Schedin, S. V. Morozov, L. A. Ponomarenko, F. Chiappini, S. Wiedmann, U. Zeitler, M. I. Katsnelson, A. K. Geim, K. S. Novoselov, and D. C. Elias, Interaction phenomena in graphene seen through quantum capacitance, *Proc. Natl. Acad. Sci. U.S.A.* **110**, 3282 (2013).
- [35] L. Miao, Z. F. Wang, W. Ming, M.-Y. Yao, M. Wang, F. Yang, Y. R. Song, F. Zhu, A. V. Fedorov, Z. Sun, C. L. Gao, C. Liu, Q.-X. Xue, C.-X. Liu, F. Liu, D. Qian, and J.-F. Jia, Quasiparticle dynamics in reshaped helical Dirac cone of topological insulators, *Proc. Natl. Acad. Sci. U.S.A.* **110**, 2758 (2013).
- [36] J.-R. Wang and G.-Z. Liu, Absence of dynamical gap generation in suspended graphene, *New J. Phys.* **14**, 043036 (2012).
- [37] J. Hofmann, E. Barnes, and S. Das Sarma, Why does graphene behave as a weakly interacting system?, *Phys. Rev. Lett.* **113**, 105502 (2014).
- [38] P. Goswami and S. Chakravarty, Quantum criticality between topological and band insulators in 3+1 dimensions, *Phys. Rev. Lett.* **107**, 196803 (2011).
- [39] P. Hosur, S. A. Parameswaran, and A. Vishwanath, Charge transport in Weyl semimetals, *Phys. Rev. Lett.* **108**, 046602 (2012).
- [40] H.-K. Tang, J. N. Leaw, J. N. B. Rodrigues, I. F. Herbut, P. Sengupta, F. F. Assaad, and S. Adam, The role

- of electron-electron interactions in two-dimensional Dirac fermions, *Science* **361**, 570 (2018).
- [41] J. N. Leaw, H.-K. Tang, M. Trushin, F. F. Assaad, and S. Adam, Universal Fermi-surface anisotropy renormalization for interacting Dirac fermions with long-range interactions, *Proc. Natl. Acad. Sci. U.S.A.* **116**, 24631 (2019).
 - [42] I. F. Herbut, Interactions and phase transitions on graphene's honeycomb lattice, *Phys. Rev. Lett.* **97**, 146401 (2006).
 - [43] I. F. Herbut, V. Juričić, and B. Roy, Theory of interacting electrons on the honeycomb lattice, *Phys. Rev. B* **79**, 085116 (2009).
 - [44] J. Maciejko and R. Nandkishore, Weyl semimetals with short-range interactions, *Phys. Rev. B* **90**, 035126 (2014).
 - [45] B. Roy and S. Das Sarma, Quantum phases of interacting electrons in three-dimensional dirty Dirac semimetals, *Phys. Rev. B* **94**, 115137 (2016).
 - [46] A. L. Szabó and B. Roy, Emergent chiral symmetry in a three-dimensional interacting Dirac liquid, *J. High Energy Phys.* **01** (2021) 004.
 - [47] E.-G. Moon, C. Xu, Y. B. Kim, and L. Balents, Non-Fermi-liquid and topological states with strong spin-orbit coupling, *Phys. Rev. Lett.* **111**, 206401 (2013).
 - [48] I. F. Herbut and L. Janssen, Topological Mott insulator in three-dimensional systems with quadratic band touching, *Phys. Rev. Lett.* **113**, 106401 (2014).
 - [49] B.-J. Yang, E.-G. Moon, H. Isobe, and N. Nagaosa, Quantum criticality of topological phase transitions in three-dimensional interacting electronic systems, *Nat. Phys.* **10**, 774 (2014).
 - [50] A. A. Abrikosov, Gapless state of bismuth-type semimetals, *J. Low. Temp. Phys.* **8**, 315 (1972).
 - [51] H. Isobe, B.-J. Yang, A. Chubukov, J. Schmalian, and N. Nagaosa, Emergent non-Fermi-liquid at the quantum critical point of a topological phase transition in two dimensions, *Phys. Rev. Lett.* **116**, 076803 (2016).
 - [52] G. Y. Cho and E.-G. Moon, Novel quantum criticality in two dimensional topological phase transitions, *Sci. Rep.* **6**, 19198 (2016).
 - [53] J.-R. Wang, G.-Z. Liu, and C.-J. Zhang, Excitonic pairing and insulating transition in two-dimensional semi-Dirac semimetals, *Phys. Rev. B* **95**, 075129 (2017).
 - [54] H.-H. Lai, Correlation effects in double-Weyl semimetals, *Phys. Rev. B* **91**, 235131 (2015).
 - [55] S.-K. Jian and H. Yao, Correlated double-Weyl semimetals with Coulomb interactions: Possible applications to HgCr_2Se_4 and SrSi_2 , *Phys. Rev. B* **92**, 045121 (2015).
 - [56] J.-R. Wang, G.-Z. Liu, and C.-J. Zhang, Quantum phase transition and unusual critical behavior in multi-Weyl semimetals, *Phys. Rev. B* **96**, 165142 (2017).
 - [57] S.-X. Zhang, S.-K. Jian, and H. Yao, Correlated triple-Weyl semimetals with Coulomb interactions, *Phys. Rev. B* **96**, 241111(R) (2017).
 - [58] J.-R. Wang, G.-Z. Liu, and C.-J. Zhang, Breakdown of Fermi liquid theory in topological multi-Weyl semimetals, *Phys. Rev. B* **98**, 205113 (2018).
 - [59] J.-R. Wang, G.-Z. Liu, and C.-J. Zhang, Topological quantum critical point in a triple-Weyl semimetal: Non-Fermi-liquid behavior and instabilities, *Phys. Rev. B* **99**, 195119 (2019).
 - [60] S. Han, C. Lee, E.-G. Moon, and H. Min, Emergent anisotropic non-Fermi liquid at a topological phase transition in three dimensions, *Phys. Rev. Lett.* **122**, 187601 (2019).
 - [61] S.-X. Zhang, S.-K. Jian, and H. Yao, Quantum criticality preempted by nematicity, *Phys. Rev. B* **103**, 165129 (2021).
 - [62] B. Roy, M. P. Kennett, K. Yang, and V. Juričić, From birefringent electrons to a marginal or non-Fermi liquid of relativistic spin-1/2 fermions: An emergent superuniversality, *Phys. Rev. Lett.* **121**, 157602 (2018).
 - [63] V. N. Kotov, B. Uchoa, and O. P. Sushov, Coulomb interactions and renormalization of semi-Dirac fermions near a topological Lifshitz transition, *Phys. Rev. B* **103**, 045403 (2021).
 - [64] B. Roy, P. Goswami, and V. Juričić, Interacting Weyl fermions: Phases, phase transitions, and global phase diagram, *Phys. Rev. B* **95**, 201102(R) (2017).
 - [65] B. Roy and M. S. Foster, Quantum multicriticality near the Dirac-semimetal to band-insulator critical point in two dimensions: A controlled ascent from one dimension, *Phys. Rev. X* **8**, 011049 (2018).
 - [66] J. Wang, Role of four-fermion interaction and impurity in the states of two-dimensional semi-Dirac materials, *J. Phys.: Condens. Matter* **30**, 125401 (2018).
 - [67] A. L. Szabó, R. Moessner, and B. Roy, Interacting spin-3/2 fermions in a Luttinger semimetal: competing phases and their selection in the global phase diagram, *Phys. Rev. B* **103**, 165139 (2021).
 - [68] I. Boettcher, Interplay of topology and electron-electron interactions in Rarita-Schwinger-Weyl semimetals, *Phys. Rev. Lett.* **124**, 127602 (2020).
 - [69] L. Savary, E.-G. Moon, and L. Balents, New type of quantum criticality in the pyrochlore iridates, *Phys. Rev. X* **4**, 041027 (2014).
 - [70] M. D. Uryszek, E. Christou, A. Jaefari, F. Krüger, and B. Uchoa, Quantum criticality of semi-Dirac fermions in $2 + 1$ dimensions, *Phys. Rev. B* **100**, 155101 (2019).
 - [71] S. Sur and B. Roy, Unifying interacting nodal semimetals: A new route to strong coupling, *Phys. Rev. Lett.* **123**, 207601 (2019).
 - [72] M. D. Uryszek, F. Krüger, and E. Christou, Fermionic criticality of anisotropic nodal point semimetals away from the upper critical dimension: Exact exponents to leading order in $\frac{1}{N_f}$, *Phys. Rev. Research* **2**, 043265 (2020).
 - [73] P.-L. Zhao, X.-B. Qing, H.-Z. Lu, and X.C. Xie, Coulomb Instabilities of a Three-Dimensional Higher-Order Topological Insulator, *Phys. Rev. Lett.* **127**, 176601 (2021).
 - [74] Y.-W. Lee and M.-F. Yang, Comment on “Coulomb Instabilities of a Three-Dimensional Higher-Order Topological Insulator”, *arXiv:2202.01642v1*.
 - [75] Y.-W. Lee and Y.-L. Lee, Correlated second-order Dirac semimetals with Coulomb interactions, *Phys. Rev. B* **105**, 125110 (2022).
 - [76] B. Roy, P. Goswami, and J. D. Sau, Continuous and discontinuous topological quantum phase transitions, *Phys. Rev. B* **94**, 041101(R) (2016).
 - [77] H. Li, H.-Y. Kee, and Y. B. Kim, Green's Function Approach to Interacting Higher-order Topological Insulators, *arXiv:2206.00713*.
 - [78] R. Shankar, Renormalization-group approach to interacting fermions, *Rev. Mod. Phys.* **66**, 129 (1994).
 - [79] A. W. W. Ludwig, M. P. A. Fisher, R. Shankar, and G. Grinstein, Integer quantum Hall transition: An alternative approach and exact results, *Phys. Rev. B* **50**, 7526 (1994).

- [80] F. Evers and A. D. Mirlin, Anderson transitions, *Rev. Mod. Phys.* **80**, 1355 (2008).
- [81] M. S. Foster, Multifractal nature of the surface local density of states in three-dimensional topological insulators with magnetic and nonmagnetic disorder, *Phys. Rev. B* **85**, 085122 (2012).
- [82] B. Roy and S. Das Sarma, Diffusive quantum criticality in three-dimensional disordered Dirac semimetals, *Phys. Rev. B* **90**, 241112(R) (2014).
- [83] S. V. Syzranov, P. M. Ostrovsky, V. Gurarie, and L. Radzihovsky, Critical exponents at the unconventional disorder-driven transition in a Weyl semimetal, *Phys. Rev. B* **93**, 155113 (2016).
- [84] B. Sbierski, K. S. C. Decker, and P. W. Brouwer, Weyl node with random vector potential, *Phys. Rev. B* **94**, 220202(R) (2016).
- [85] D. Carpentier, A. A. Fedorenko, and E. Orignac, Effect of disorder on 2D topological merging transition from a Dirac semi-metal to a normal insulator, *Euro. Phys. Lett.* **102**, 67010 (2013).
- [86] B. Roy, R.-J. Slager, and V. Juričić, Global phase diagram of a Dirty Weyl liquid and emergent superuniversality, *Phys. Rev. X* **8**, 031076 (2018).
- [87] X. Luo, B. Xu, T. Ohtsuki, and R. Shindou, Quantum multicriticality in disordered Weyl semimetals, *Phys. Rev. B* **97**, 045129 (2018).
- [88] J.-R. Wang, W. Li, G. Wang, and C.-J. Zhang, Global phase diagram of Coulomb-interacting anisotropic Weyl semimetal with disorder, *J. Phys.: Condens. Matter* **33**, 125601 (2021).
- [89] S. Bera, J. D. Sau, and B. Roy, Dirty Weyl semimetals: Stability, phase transition, and quantum criticality, *Phys. Rev. B* **93**, 201302(R) (2016).
- [90] R. M. Nandkishore and S. A. Parameswaran, Disorder-driven destruction of a non-Fermi liquid semimetal studied by renormalization group analysis, *Phys. Rev. B* **95**, 205106 (2017).
- [91] I. Mandal and R. M. Nandkishore, Interplay of Coulomb interactions and disorder in three-dimensional quadratic band crossings without time-reversal symmetry and with unequal masses for conduction and valence bands, *Phys. Rev. B* **97**, 125121 (2018).
- [92] M. E. Carrington, C. S. Fischer, L. von Smekal, and M. H. Thoma, Dynamical gap generation in graphene with frequency-dependent renormalization effects, *Phys. Rev. B* **94**, 125102 (2016).
- [93] M. E. Carrington, C. S. Fischer, L. von Smekel, and M. H. Thoma, Role of frequency dependence in dynamical gap generation in graphene, *Phys. Rev. B* **97**, 115411 (2018).