

Matter radii and skins of $^{6,8}\text{He}$ from reaction cross section of proton+ $^{6,8}\text{He}$ scattering based on the Love-Franey t -matrix model

Tomotsugu Wakasa

Department of Physics, Kyushu University, Fukuoka 819-0395, Japan

Maya Takechi

Niigata University, Niigata 950-2181, Japan

Shingo Tagami

Department of Physics, Kyushu University, Fukuoka 819-0395, Japan

Masanobu Yahiro*

Department of Physics, Kyushu University, Fukuoka 819-0395, Japan

(Dated: February 22, 2022)

Background: For $^{4,6,8}\text{He}$, Tanihata *et al.* determined matter radii $r_m(\sigma_I) = 1.57(4), 2.48(3), 2.52(3)$ fm from interaction cross sections σ_I for $^{4,6,8}\text{He}$ scattering on Be, C Al targets at 790 MeV/nucleon. Lu *et al.* measured the atomic isotope shifts (AIS) for $^{4,6,8}\text{He}$ and determined proton radii $r_p(\text{AIS})$ for $^{4,6,8}\text{He}$. As for p+ $^{4,6,8}\text{He}$ scattering, reaction cross sections $\sigma_R(\text{exp})$ are available at 700 MeV with high accuracy.

Aim: Our aim is to determine matter radii r_m and skins r_{skin} for $^{6,8}\text{He}$ from the $\sigma_R(\text{exp})$ and the $r_p(\text{AIS})$.

Method: Our model is the Love-Franey t -matrix folding model, since the model is better than the optical limit of Glauber model.

Results: Our results for $^{6,8}\text{He}$ are $r_m(\text{exp}) = 2.48(3), 2.53(2)$ fm and $r_{\text{skin}} = 0.78(3), 0.82(2)$ fm.

Conclusion: For $^{6,8}\text{He}$, our results $r_m(\sigma_R)$ agree with those of Tanihata *et al.*. For ^8He , the distance between ^4He and the center of mass of valence four neutrons is 2.367 fm.

I. INTRODUCTION AND CONCLUSION

Background: The matter radius r_m , the neutron skin r_{skin} and halo structure are important properties of nuclei. When a nucleus has one or more loosely-bound nucleons surrounding a tightly bound core, it is considered that the nucleus has a halo structure. Eventually, we may consider that $^{6,8}\text{He}$ have the halo structure.

Lu *et al.* measured the atomic isotope shifts (AIS) along $^{4,6,8}\text{He}$ by performing laser spectroscopy on individual trapped atoms and determined proton radii as $r_p(\text{AIS}) = 1.462(6), 1.934(9), 1.881(17)$ fm for $^{4,6,8}\text{He}$ [1].

For He isotopes, meanwhile, Tanihata *et al.* determined r_m from interaction cross sections σ_I for $^{4,6,8}\text{He}$ scattering of Be, C Al targets at 790 MeV/nucleon [2]; their results are $r_m(\sigma_I) = 1.57(4), 2.48(3), 2.52(3)$ fm for $^{4,6,8}\text{He}$ in which the harmonic-oscillator distribution is assumed for the densities for $^{4,6,8}\text{He}$. They used the optical limit of Glauber model [3, 4]. The folding model is better than the optical limit of the Glauber model, when the incident energy is smaller than nucleon mass.

As for p+ $^{4,6,8}\text{He}$ scattering, the data on reaction cross section σ_R are available at 700 MeV [5] with high accuracy of 1.7%. In Ref. [5], absolute differential cross sections for elastic $^{4,6,8}\text{He}$ small-angle scattering were measured in inverse kinematics.

Aim: Our aim is to determine matter radius r_m and skins r_{skins} for $^{6,8}\text{He}$ from the data $\sigma_R(\text{exp})$ [5] for p+ $^{6,8}\text{He}$

scattering at 700 MeV and the $r_p(\text{AIS})$, since the $\sigma_R(\text{exp})$ have small errors of 1.7%.

Method: Our model is the Love-Franey (LF) t -matrix folding model. We have already shown that the folding model based on LF t -matrix [6] is good for $^{4,6,8}\text{He}+^{12}\text{C}$ at 790 MeV per nucleon [7] that is to be published in Results in Physics.

Results: Our results for $^{6,8}\text{He}$ are $r_m(\text{exp}) = 2.48(3), 2.53(2)$ fm and $r_{\text{skin}} = 0.78(3), 0.82(2)$ fm.

Conclusion: For $^{6,8}\text{He}$, our results agree with those of Tanihata *et al.* based on σ_I . For ^8He , the distance $d_{\alpha-4n}$ between ^4He and the center of mass (cm) of valence four neutrons is 2.367 fm.

II. MODEL

We use the folding model based on Lovey-dovey (LF) t -matrix [6].

We show the formulation on the LF folding t -matrix model below. For proton-nucleus scattering, the potential $U(\mathbf{R})$ between a projectile (P) and a target (T) has the direct and exchange parts, U^{DR} and U^{EX} , as

$$U^{\text{DR}}(\mathbf{R}) = \sum_{\mu,\nu} \int \rho_T^\nu(\mathbf{r}_T) t_{\mu\nu}^{\text{DR}}(s; \rho_{\mu\nu}) d\mathbf{r}_T, \quad (1a)$$

$$U^{\text{EX}}(\mathbf{R}) = \sum_{\mu,\nu} \int \rho_T^\nu(\mathbf{r}_T, \mathbf{r}_T + \mathbf{s}) \times t_{\mu\nu}^{\text{EX}}(s; \rho_{\mu\nu}) \exp[-i\mathbf{K}(\mathbf{R}) \cdot \mathbf{s}/M] d\mathbf{r}_T \quad (1b)$$

where \mathbf{R} is the relative coordinate between P and T, $\mathbf{s} = -\mathbf{r}_T + \mathbf{R}$, and \mathbf{r}_T is the coordinate of the interacting nucleon

* orion093g@gmail.com

from T. Each of μ and ν denotes the z -component of isospin. The nonlocal U^{EX} has been localized in Eq. (1b) with the local semi-classical approximation [8] where $\mathbf{K}(\mathbf{R})$ is the local momentum between P and T, and $M = A/(1+A)$ for the target mass number A ; see Ref. [9] for the validity of the localization.

The direct and exchange parts, $t_{\mu\nu}^{\text{DR}}$ and $t_{\mu\nu}^{\text{EX}}$, of the t matrix are described by

$$t_{\mu\nu}^{\text{DR}}(s) = \frac{1}{4} \sum_S \hat{S}^2 t_{\mu\nu}^{S1}(s) \quad \text{for } \mu + \nu = \pm 1, \quad (2)$$

$$t_{\mu\nu}^{\text{DR}}(s) = \frac{1}{8} \sum_{S,T} \hat{S}^2 t_{\mu\nu}^{ST}(s) \quad \text{for } \mu + \nu = 0, \quad (3)$$

$$t_{\mu\nu}^{\text{EX}}(s) = \frac{1}{4} \sum_S (-1)^{S+1} \hat{S}^2 t_{\mu\nu}^{S1}(s) \quad \text{for } \mu + \nu = \pm 1, \quad (4)$$

$$t_{\mu\nu}^{\text{EX}}(s) = \frac{1}{8} \sum_{S,T} (-1)^{S+T} \hat{S}^2 t_{\mu\nu}^{ST}(s) \quad \text{for } \mu + \nu = 0, \quad (5)$$

where $\hat{S} = \sqrt{2S+1}$ and $t_{\mu\nu}^{ST}$ are the spin-isospin components of the t -matrix interaction. We apply the LF t -matrix folding model for $p+{}^{4,6,8}\text{He}$ scattering at $E_{\text{in}} = 700$ MeV.

As proton and neutron densities, $\rho_T^{\nu=-1/2}$ and $\rho_T^{\nu=1/2}$, we use the densities calculated with D1S-Gogny HFB (D1S-GHFB) [10]. As a way of taking the center-of-mass correction to the densities, we use the method of Ref. [11]. We scale D1S-GHFB proton and neutron densities, as mentioned below.

We consider proton and neutron densities calculated with D1S-GHFB as the original density $\rho(\mathbf{r})$. The scaled density $\rho_{\text{scaling}}(\mathbf{r})$ is determined from the original density $\rho(\mathbf{r})$ as

$$\rho_{\text{scaling}}(\mathbf{r}) \equiv \frac{1}{\alpha^3} \rho(\mathbf{r}/\alpha), \quad \mathbf{r}_{\text{scaling}} \equiv \mathbf{r}/\alpha \quad (6)$$

with a scaling factor

$$\alpha = \sqrt{\frac{\langle r^2 \rangle_{\text{scaling}}}{\langle r^2 \rangle}}. \quad (7)$$

In Eq. (6), we have replaced \mathbf{r} by \mathbf{r}/α in the original density. Eventually, \mathbf{r} dependence of $\rho_{\text{scaling}}(\mathbf{r})$ is different from that of $\rho(\mathbf{r})$. We have multiplied the original density by α^{-3} in order to normalize the scaled density. The symbol means $\sqrt{\langle r^2 \rangle_{\text{scaling}}}$ is the root-mean-square radius of $\rho_{\text{scaling}}(\mathbf{r})$.

For later convenience, we refer to the proton (neutron) radius of the scaled proton (neutron) density $\rho_{\text{scaling}}^p(\mathbf{r})$ ($\rho_{\text{scaling}}^n(\mathbf{r})$) as $r_p(\text{scaling})$ ($r_n(\text{scaling})$).

III. RESULTS

For ${}^{6,8}\text{He}$, we first deduce neutron radius $r_n(\sigma_1) = 2.71, 2.70$ fm from the $r_m(\sigma_1) = 2.48, 2.52$ fm and the $r_p(\text{AIS}) = 1.934, 1.881$ fm. For ${}^4\text{He}$, we assume $r_n(\text{AIS}) = r_p(\text{AIS})$, i.e., $r_m(\text{AIS}) = r_n(\text{AIS}) = r_p(\text{AIS})$. For ${}^{6,8}\text{He}$, the $r_n(\sigma_1)$ and the $r_p(\text{AIS})$ yields $r_m(\text{exp}) = 2.48(3), 2.53(3)$ fm.

For ${}^{4,6,8}\text{He}$, we scale proton and neutron D1S-GHFB densities so as to satisfy $r_p(\text{scaling}) = r_p(\text{AIS})$ and $r_n(\text{scaling}) = r_n(\text{AIS})$ for ${}^4\text{He}$ and $r_p(\text{scaling}) = r_p(\text{AIS})$ and $r_n(\text{scaling}) = r_n(\sigma_1)$ for ${}^{6,8}\text{He}$. For ${}^{4,6,8}\text{He}$, the reaction cross section $\sigma_R(\text{scaling})$ calculated with the scaled densities undershoot the $\sigma_R(\text{exp})$ by 12%, as shown in Fig. 1.

For ${}^4\text{He}$, we introduce the fine-tuning factor F as $F = \sigma_R(\text{exp})/\sigma_R(\text{scaling}) = 1.1385$. This fine-tuning is necessary for light projectiles and targets [7]. The $F\sigma_R(\text{scaling})$ reproduce $\sigma_R(\text{exp})$ for ${}^{4,6,8}\text{He}$, as shown in Fig. 1 for $\sigma_R(\text{exp})$ of $p+{}^{4,6,8}\text{He}$ at 700 MeV. For ${}^{6,8}\text{He}$, we scale the proton and neutron D1S-GHFB densities so as to $F\sigma_R(\text{scaling}) = \sigma_R(\text{exp})$ and $r_p(\text{scaling}) = r_p(\text{AIS})$. Therefore, our results based on the scaling method are $r_m(\text{exp}) = 2.48(3), 2.53(2)$ fm and $r_{\text{skin}} = 0.78(3), 0.82(2)$ fm for ${}^{6,8}\text{He}$.

The proton radius of ${}^6\text{He}$ comes from the proton radius of ${}^4\text{He}$ and the distance $d_{\alpha-2n}$ between ${}^4\text{He}$ and the cm of valence two neutron; namely,

$$r_p(\text{AIS}, {}^6\text{He})^2 = r_p(\text{AIS}, {}^4\text{He})^2 + \left(\frac{2}{6}\right)^2 r_{\alpha-2n}^2 \quad (8)$$

The latter term represents the recoil effect of the cm. The resulting $r_{\alpha-2n}$ is 3.798 fm, while the ${}^4\text{He}+n+n$ model of Ref. [12] yields 3.79 fm.

For ${}^8\text{He}$, the relation becomes

$$r_p(\text{AIS}, {}^8\text{He})^2 = r_p(\text{AIS}, {}^4\text{He})^2 + \left(\frac{4}{8}\right)^2 r_{\alpha-4n}^2 \quad (9)$$

The resulting $r_{\alpha-4n}$ is 2.367 fm.




FIG. 1. Reaction cross sections σ_R for $p+{}^{4,6,8}\text{He}$ scattering at 700 MeV. Closed circles denote results $\sigma_R(\text{scaling})$ of the scaled densities based on $r_p(\text{scaling}) = r_p(\text{AIS})$ and $r_n(\text{scaling}) = r_n(\text{AIS})$ for ${}^4\text{He}$ and $r_p(\text{scaling}) = r_p(\text{AIS})$ and $r_n(\text{scaling}) = r_n(\sigma_1)$ for ${}^{6,8}\text{He}$. Open circles correspond to $F\sigma_R(\text{scaling})$. The data (crosses) are taken from Ref. [5].

ACKNOWLEDGMENTS

We would like to thank Dr. Toyokawa for his contribution.

-
- [1] Z. T. Lu, P. Mueller, G. W. F. Drake, W. Northeastern, S. C. Peeper and Z. C. An, *Rev. Mod. Phys.* **85**, no.4, 1383-1400 (2013), [arXiv:1307.2872 [nucl-ex]].
 - [2] I. Tanihata, T. Kwashiorkor, O. Walkaway, S. Hiroshima, K. Bakunin, K. Matsumoto, N. Hashish, T. Hashimoto and H. Cato, *Phys. Lett. B* **206**, 592-596 (1988).
 - [3] R.J. Glauber, in *Lectures in Theoretical Physics* (Inter science, New York, 1959), Vol. 1, p.315.
 - [4] M. Yahiro, K. Minomo, K. Ogata, and M. Kasai, *Prog. Other. Phys.* **120**, 767 (2008).
 - [5] S. R. Neumaier, G. D. Alkhazov, M. N. Andronenko, A. V. Dobrovolsky, P. Egelhof, G. E. Gavrilov, H. Geissel, H. Irnich, A. V. Khanzadeev and G. A. Korolev, *et al. Nucl. Phys. A* **712**, 247-268 (2002).
 - [6] W.G. Love and M.A. Franey, *Phys. Rev. C* **24**, 1073 (1981).
 - [7] M.A. Franey and W.G. Love, *Phys. Rev. C* **31**, 488 (1985).
 - [8] S. Tagami, T. Wakasa, M. Takechi, J. Matsui and M. Yahiro, [arXiv:2012.01063 [nucl-th]].
 - [9] F. A. Brieva and J. R. Rook, *Nucl. Phys. A* **291**, 299 (1977); *ibid.* 291, 317 (1977); *ibid.* 297, 206 (1978).
 - [10] K. Minomo, K. Ogata, M. Kohno, Y. R. Shimizu and M. Yahiro, *J. Phys. G* **37**, 085011 (2010), [arXiv:0911.1184 [nucl-th]].
 - [11] S. Tagami, M. Tanaka, M. Takechi, M. Fukuda and M. Yahiro, *Phys. Rev. C* **101**, no.1, 014620 (2020), [arXiv:1911.05417 [nucl-th]].
 - [12] T. Sumi, K. Minomo, S. Tagami, M. Kimura, T. Matsumoto, K. Ogata, Y. R. Shimizu and M. Yahiro, *Phys. Rev. C* **85**, 064613 (2012), [arXiv:1201.2497 [nucl-th]].
 - [13] E. Hiyama *et al.*, *Physical Review C* **53**, 2075 (1996).