

Exact quasiparticle properties of a heavy polaron in BCS Fermi superfluids

Jia Wang,¹ Xia-Ji Liu,¹ and Hui Hu¹

¹Centre for Quantum Technology Theory, Swinburne University of Technology, Melbourne 3122, Australia

(Dated: February 14, 2022)

We present the Ramsey response and radio-frequency spectroscopy of a heavy impurity immersed in an interacting Fermi superfluid, using exact functional determinant approach. We describe the Fermi superfluid through the conventional Bardeen-Cooper-Schrieffer theory and investigate the role of the pairing gap on quasiparticle properties revealed by the two spectroscopies. The energy cost for pair breaking prevents Anderson's orthogonality catastrophe that occurs in a non-interacting Fermi gas and allows the existence of polaron quasiparticles in the exactly solvable heavy impurity limit. Hence, we rigorously confirm the remarkable features such as dark continuum, molecule-hole continuum and repulsive polaron. For a magnetic impurity scattering at finite temperature, we predict additional resonances related to the sub-gap Yu-Shiba-Rusinov bound state, whose positions can be used to measure the superfluid pairing gap. For a non-magnetic scattering at zero temperature, we surprisingly find undamped repulsive polarons. These exact results might be readily observed in quantum gas experiments with Bose-Fermi mixtures that have a large-mass ratio.

Thanks to the unprecedented controllability recently achieved in ultracold quantum gases, investigations on non-equilibrium quantum dynamics in many-body systems have progressed rapidly [1]. One of such intriguing problems is how a quantum gas medium responds to a suddenly introduced impurity [2, 3]. The quantum gas can be either a degenerate Fermi gas or a Bose-Einstein condensate (BEC). The impurity-medium interaction can essentially be tuned arbitrarily via Feshbach resonance [4], and a variety of impurities, such as Rydberg atoms [5–7] or quantum rotor [8], can be introduced. A unique advantage of these impurity-medium systems is that they present probably the simplest non-trivial many-particle problem, where the medium response can be directly measured (i.e., by Ramsey and radio-frequency spectroscopies) and efficiently calculated even in the non-perturbative strong-coupling regime [3, 9]. Consequently, they can serve as a critical meeting point for theoretical and experimental efforts to understand the complicated quantum dynamics of interacting many-particle systems.

Historically, the first research of impurity-medium systems in 1933 led Landau to introduce a general concept of polarons – quasiparticles formed by dressing the impurity with elementary excitations of the medium [10]. The new platform of ultracold quantum gases has enabled the exploration of polaron quasiparticle properties in a controllable and quantitative manner over the last decade, both experimentally [11–21] and theoretically [22–42]. In particular, a number of salient features of polarons have been predicted by approximate theories and Monte Carlo simulations, including the excited repulsive polaron with finite lifetime [26] and the dark continuum [35] and molecule-hole continuum [2] that separate the attractive and repulsive polaron branches. While the repulsive polaron has been unambiguously observed in experiments [13, 14], the existence of the dark and molecule-hole continua remains elusive due to the uncertainty in theoretical calculations. The purpose of this Letter is to present an

exact calculation of polaron quasiparticle properties in the heavy impurity limit and in the experimentally unexplored regime with a Fermi superfluid medium [43–47].

Our work naturally extends the well-known exactly solvable many-body problem of the Fermi-edge singularity of x-ray absorption spectra in metals [48, 49], which is the first and most important example of non-equilibrium many-body physics [50, 51]. In this impurity-medium problem, the suddenly introduced infinitely heavy impurity can excite particle-hole pairs close to Fermi surfaces without costing finite recoil energy [3, 9]. The multiple particle-hole excitations completely changes the many-particle states in the limit of a large particle number. As a result, the many-particle states with and without impurity become orthogonal, i.e., Anderson's "orthogonality catastrophe" (OC) [52]. In the context of ultracold quantum gases, the Fermi-edge singularity has been quantitatively re-examined via the functional determinant approach (FDA) [53–56], providing insightful understanding of polaron physics [2, 3, 9]. Unfortunately, strictly speaking, due to OC the attractive and repulsive polarons do not exist, as indicated by the vanishing quasiparticle residue [3, 9].

Here, we propose an exactly solvable model of a heavy impurity immersed in a Fermi superfluid medium described by the standard Bardeen-Cooper-Schrieffer (BCS) pairing theory [57–59]. As multiple particle-hole excitations can be efficiently suppressed by the energy cost of pairing breaking, Anderson's OC is avoided and polarons acquire *nonzero* quasiparticle residue. Therefore, we obtain a benchmark theoretical model with well-defined polaron quasiparticles, in which all the speculated characteristics of polarons can be rigorously examined. Our results are also highly experimentally relevant, as a BCS Fermi superfluid (of ⁶Li or ⁴⁰K atoms) has now been routinely realized using Feshbach resonance at the so-called BEC-BCS crossover and a heavy atomic species such as ¹³³Cs can be manipulated at will as impurity.




FIG. 1. A sketch of the occupation and structure of the single-particle dispersion spectrum of a two-component superfluid Fermi gas with a positive chemical potential $\mu > 0$ and the presence of a static impurity (black dot). (a) shows the spectrum E_ν when the impurity is in the non-interacting state (black arrow up) at zero temperature. When the impurity is in the interacting polaron state (black arrow down), the spectrum \tilde{E}_ν are shown in (b) at zero and (c) finite temperature.

In Fig. 1, we outline an injection scheme of interest in this work: the impurity is driven from a non-interacting hyperfine state [Fig. 1 (a)] into an interacting state [Fig. 1 (b)] at time $t = 0$. The dynamical evolution at later time, namely the Ramsey response, is then exactly calculated via an extension of the FDA, from which we extract the spectral function. The existence of pairing gap prevents the OC and preserves well-defined polaron quasi-particle features in the spectral function. In addition to rigorously confirming the remarkable characteristics of polarons, our exact results also reveal two novel unique features related to the Fermi superfluid medium: the resonances related to the sub-gap Yu-Shiba-Rusinov (YSR) state bound to a magnetic impurity [60–64] and repulsive polarons with infinitely long lifetime in the case of a non-magnetic impurity scattering.

Theory. The fundamental Ramsey response is the real-time overlap function between the many-body state with and without the impurity, $S(t) = \langle e^{i\mathcal{H}_i t} e^{-i\mathcal{H}_f t} \rangle \equiv \text{Tr}[e^{i\mathcal{H}_i t} e^{-i\mathcal{H}_f t} \hat{\rho}_0]$, where \mathcal{H}_i (\mathcal{H}_f) is the many-body Hamiltonian in the absence (presence) of the impurity scattering and $\hat{\rho}_0$ is the initial state of the Fermi system (Hereafter, we use the units of $\hbar \equiv 1$). Complementarily, the frequency-resolved spectral function $A(\omega) = \text{Re} \int_0^\infty e^{i\omega t} S(t) dt / \pi$, which determines the radio-frequency (rf) spectroscopy, can be obtained by a Fourier transformation [9, 65]. Since the complexity of the many-body Hamiltonians increases exponentially with the numbers of particles N in the system, an exact calculation of $S(t)$ is usually inaccessible. However, in the case that \mathcal{H}_i and \mathcal{H}_f are both fermionic, bilinear many-body operators, the overlap function can reduce to a determinant in single-particle Hilbert space that grows only linearly to N [53–56]:

$$S(t) = e^{-i\omega_0 t} \det[1 - \hat{n} + e^{i\mathcal{H}_i t} e^{-i\mathcal{H}_f t} \hat{n}], \quad (1)$$

where \hat{n} is the occupation number operator, and \hat{h}_i (\hat{h}_f)

are the single-particle representatives of \mathcal{H}_i (\mathcal{H}_f) up to some constant terms that ω_0 compensates. For example, $\mathcal{H}_f = K_0 + \omega_0 + \int d\mathbf{r} \hat{\phi}^\dagger(\mathbf{r}) \underline{h}_f(\mathbf{r}) \hat{\phi}(\mathbf{r})$, where K_0 is an unimportant constant and

$$\underline{h}_f(\mathbf{r}) = \begin{pmatrix} -\frac{\nabla^2}{2m} + V_\uparrow(\mathbf{r}) - \mu & \Delta \\ \Delta & \frac{\nabla^2}{2m} - V_\downarrow(\mathbf{r}) + \mu \end{pmatrix}, \quad (2)$$

with $V_\sigma(\mathbf{r})$ being the potential between impurity and σ -component fermion. Note that, here we already extend the FDA to the case of a BCS Fermi superfluid, which is characterized by the pairing gap Δ and chemical potential μ to be determined by a given scattering length a between unlike fermions, temperature T and Fermi momentum $k_F = (3\pi^2 N/\mathcal{V})^{1/3}$, where \mathcal{V} is the system volume. It is convenient to use the Nambu spinor operators as $\hat{\phi}^\dagger(\mathbf{r}) = [c_\uparrow^\dagger(\mathbf{r}), c_\downarrow(\mathbf{r})]$, where $c_\sigma^\dagger(\mathbf{r})$ [$c_\sigma(\mathbf{r})$] being the creation (annihilation) operator for a σ -component fermion at position \mathbf{r} . We also have $\omega_0 = \text{Tr} V_\downarrow$, which corresponds to the phase factor in Eq. (1) with \underline{V}_\downarrow being the matrix format of $V_\downarrow(\mathbf{r})$ in a complete orthogonal set of basis. Finally, $\underline{h}_i(\mathbf{r})$ can be obtained by setting $V_\sigma(\mathbf{r})$ equals zero in Eq. (2). In what follows, we briefly describe the computation procedure and present our main physical results, but relegate numerical details and additional discussions to a complementary paper [66].

We consider a finite system confined in a sphere of radius R and take the system size towards infinity, while keeping the density constant, until numerical results are converged [9]. We focus on the s -wave channel and use finite-range potentials $V_\sigma(r)$ whose corresponding energy-dependent scattering length $a_\sigma(E_F) = -\tan \eta_\sigma(k_F)/k_F$, where $\eta_\sigma(k_F)$ is the s -wave scattering length between the impurity and σ -component fermions at the Fermi energy $E_F = k_F^2/(2m)$. We find that numerical results are not sensitive to other short-range details of the potential. Therefore, for simplicity we denote $a_\sigma \equiv a_\sigma(E_F)$ hereafter [67]. Finally, for a given set of parameters $\{k_F, a, a_\uparrow, a_\downarrow, T\}$, we can calculate Eq. (1) by finding the eigenpairs $E_\nu, \phi_\nu \equiv [\phi_{\nu,\uparrow}(r), \phi_{\nu,\downarrow}(r)]$ for $\underline{h}_i(r)$ and $\tilde{E}_\nu, \tilde{\phi}_\nu$ for $\underline{h}_f(r)$. In this presentation, the occupation operator \hat{n} is given by a diagonal matrix with elements $n_{\nu\nu} = [e^{-E_\nu/(k_B T)} + 1]^{-1}$, where k_B is the Boltzmann constant. Figure 1(a) gives a sketch of the occupation and structure of single-particle spectrum E_ν without impurity at $T = 0$, which includes a completely filled Fermi sea (filled circles in the lower branch) and an empty one (empty circles on the top) separated by 2Δ . The presence of impurity scattering leads to a shift of the single-particle levels $E_\nu \rightarrow \tilde{E}_\nu$ as shown in Fig. 1(b), where the small blue dots shows E_ν for comparison. In the case of a magnetic impurity scattering ($a_\uparrow \neq a_\downarrow$), there also exists a sub-gap YSR bound state with energy E_{YSR} [60–63].

Ramsey response. As reported in Fig. 2, our numerical examples here focus on the BCS side of the crossover $k_F a = -2 < 0$, where $\mu \approx 0.85 E_F$ and $\Delta \approx 0.40 E_F$




FIG. 2. Ramsey response $|S(t)|$ for a magnetic impurity scattering with (a) attractive scattering lengths $a_{\uparrow} < 0$ and (b) repulsive scattering lengths $a_{\uparrow} > 0$ are shown for different values of the scattering length a between the two-component fermions and different temperature T ; see legend.

at zero temperature. While our method applies to the whole crossover regime, mean-field description becomes only qualitatively reliable on the BEC side. We also focus on the simplest case, where the impurity only interacts with the spin-up component fermion, i.e., $V_{\downarrow}(r) = 0$. For comparison, we show also the results for $k_F a = 0$ (with $\mu = E_F$ and $\Delta = 0$), where the \uparrow -component of medium reduces to a non-interacting Fermi gas that couples with the impurity, and the \downarrow -component being simply a spectator. These results agree with previous studies for both zero and finite temperature [9].

At $k_F a = 0$ and $T = 0$, the asymptotic behavior of $|S(t)|$ at large t exhibits a power-law decay $|S(t)| \sim t^{-\alpha}$, reflecting Anderson's OC and x-ray infrared singularity $\epsilon^{\alpha-1}$ at the low-energy scale set by the inverse time $\epsilon \sim \hbar/t$ [3, 9, 48]. In contrast, in the presence of a pairing gap $\Delta \neq 0$, $|S(t)| \sim t^0$ at $T = 0$, indicating OC is prevented as the low-energy scale is now cut by Δ [68]. At finite temperature, although both with or without the pairing gap, $|S(t)|$ shows an exponential decay at large t , such behavior appears at a much later time for finite Δ . In particular, for nonzero Δ the results at $T = 0$ and $T = 0.05E_F/k_B$ are almost overlapping at $tE_F \leq 500$, showing that the pairing gap can also protect the response signal against thermal fluctuation if $k_B T \ll \Delta$.

More quantitatively, at nonzero pairing gap we have the following analytic result,

$$S(t \rightarrow \infty) \simeq \begin{cases} D_a e^{-iE_a t}, & a_{\uparrow} < 0 \\ D_a e^{-iE_a t} + D_r e^{-iE_r t}, & a_{\uparrow} > 0 \end{cases} \quad (3)$$

which fits excellently well to our numerical results, with




FIG. 3. The spectral function $A(\omega)$ at $k_F a = -2$ for different temperature (see legend) and different scattering length $a_{\uparrow} < 0$ in (a) and $a_{\uparrow} > 0$ in (b).

D_a , E_a , D_r , and E_r being fitting parameters. At small Δ , the coefficients $|D_a| \propto (\Delta/E_F)^{\alpha_a}$ and $|D_r| \propto (\Delta/E_F)^{\alpha_r}$, making the asymptotic form that agrees with the modification of the analytic expression of $S(t)$ for a non-interacting Fermi gas medium via replacing the low-energy cut-off $1/t \rightarrow \Delta$ (see, i.e., Eqs. (12) and (15) of Ref. [9]). However, our numerical results indicate the power-law exponents α and α_a are close to but not exactly the same as the analytical results given in [9], see Ref. [66] for details. At $T = 0$, E_a is purely real and corresponds to the attractive polaron energy that satisfies $E_a = \sum_{\nu} n_{\nu\nu}(E_{\nu} - \tilde{E}_{\nu})$, indicating that the attractive polaron can be regarded as the renormalization of the Fermi sea due to the impurity level. The repulsive polaron energy E_r is in general complex, where we denote the real and imaginary part as $\text{Re}E_r$ and $\text{Im}E_r$.

rf-spectroscopy. One of the key observations of this Letter is the saturation of $|S(t)|$ at large time, which implies a finite polaron quasiparticle residue $Z = |D_a| \propto \Delta^{\alpha_a}$. To check this, we calculate the frequency response $A(\omega)$ accurately with a Fourier transformation of $S(t)$. We choose a large cut-off $t^* \sim 500/E_F$, evaluate $S(t)$ numerically for $t < t^*$ and use the fitting formula in Eq. (3) for $t \geq t^*$. As shown in Fig. 3 by thick blue solid curves for zero-temperature results, the attractive polaron is characterized by a δ -function peak at E_a (with a small artificial width for visibility), unambiguously confirming the existence of a well-defined quasiparticle. The attractive polaron peak separates with a molecule-hole continuum by a region of anomalously low spectral weight, namely the “dark continuum”. This spectral gap has previously been shown in other polaron systems with approximate calculations, where the anomalously low spectral weight might be an artifact of the adopted approx-




FIG. 4. The spectral function $A(\omega)$ at $k_F a = -2$ as a function of a_\uparrow for different temperature: $k_B T = 0.05, 0.15, 0.2 E_F$ for (a), (b) and (c), respectively. The black dashed (dotted) curves show the attractive (repulsive) polaron energy and the red dashed (dotted) curves show $E_{\text{YSR}}^{(+)}$ ($E_{\text{YSR}}^{(-)}$) at finite temperature.

imations. Only recently, a diagrammatic Monte Carlo study indicates that this dark continuum might be indeed physical [35]. Here, the heavy polaron spectral function is calculated via FDA, and hence can be regarded as an exact proof of the dark continuum. For $a > 0$, a Lorentzian lineshape with a peak at $\text{Re}(E_r)$ corresponds to the repulsive polaron. The finite width determined by $\text{Im}(E_r)$ implies that the repulsive polaron has a finite lifetime.

In Fig. 3, finite-temperature results are indicated by the red thin (purple thinner) curves for $k_B T = 0.1 E_F$ ($0.15 E_F$). Other than the expected thermal broadening, some additional surprising features show up. An enhancement of spectral weight appears sharply at the energy $E_{\text{YSR}}^{(-)} = E_a - (\Delta - E_{\text{YSR}})$ below the attractive polaron. This spectral feature corresponds to the decay process highlighted by the purple arrow in Fig. 1(c), where an additional particle initially excited to the upper Fermi sea by thermal fluctuation is driven to the the YSR state. For the case of $k_F a_\uparrow > 0$, a feature associated with the repulsive polaron appears at $E_{\text{YSR}}^{(+)} = \text{Re}(E_r) - (E_{\text{YSR}} + \Delta)$, as indicated by the green arrow in Fig. 1(c): an additional particle decays from the YSR state to the lower Fermi sea. These features can be better observed in the whole spectrum of a_\uparrow across a resonance, as shown in Fig. 4 for different temperatures. The YSR features are negligible at $k_B T = 0.05 E_F$, and the spectrum in Fig. 4(a) is almost the same as zero-temperature results. This shows the protection against finite temperature provided by the pairing gap. The YSR features become apparent in Fig. 4(b) at $k_B T = 0.15 E_F$ and shows broadening at $k_B T = 0.2 E_F$. We emphasize that this range of temperature is accessible for current experiments. The polaron spectrum can be applied to measure the superfluid gap Δ and E_{YSR} . In particular,

we notice, on the positive side $a_\uparrow > 0$, if E_a , $\text{Re}(E_r)$, $E_{\text{YSR}}^{(-)}$ and $E_{\text{YSR}}^{(+)}$ can all be measured accurately, we have $2\Delta = E_a + \text{Re}(E_r) - E_{\text{YSR}}^{(-)} - E_{\text{YSR}}^{(+)}$ that does not depend on E_{YSR} . Since this formula only relies on the existence of the gap and a mid-gap state, we anticipate it can be used to measure Δ accurately for a Fermi superfluid that can not be quantitatively described by the BCS theory.

Finally, we discuss briefly our observations for the case of a non-magnetic impurity scattering with $a_\uparrow = a_\downarrow$, where the YSR features are absent as expected. Interestingly, we also discover that the repulsive polaron exhibits itself as a δ -function peak in the spectral function at zero temperature. We believe the underlying physics might be due to the gapless density fluctuations in the Fermi superfluid excited by the perfect balance of the two scattering lengths. As a result, the impurity couples to phonon excitations of the superfluid and forms a long-lived repulsive polaron. For more details, we refer to the complementary paper [66].

Experimental realization. Our predictions can be readily confirmed by immersing heavy ^{133}Cs impurities in a BCS Fermi superfluid of ^6Li atoms routinely observed near a broad Feshbach resonance $B_0 \simeq 832$ G. The two interspecies broad resonances located nearby at 843 G and 889 G [69] allow us to independently control the ^{133}Cs - ^6Li scattering lengths $a_{\uparrow,\downarrow}$. Both magnetic and non-magnetic impurity scatterings can therefore be realized by tuning the magnetic field [66].

Conclusions. We have calculated the response functions of driving a heavy impurity in a BCS superfluid from non-interacting to interacting hyperfine states. Due to the existence of a pairing gap in the superfluid, the OC is prevented and genuine polaron quasiparticles exit. The underlying physical reason is apparent: exciting

particle-hole pairs in this system requires energy cost for Cooper-pair breaking, and hence multiple particle-hole excitations are energetic unfavored. We emphasize that our FDA can support this conclusion since it is essentially exact, unlike some approximations such as extended Chevy's ansatz [22, 26] or T -matrix method [2, 36] that allow only a few particle-hole excitations. In this respect, our calculation can be regarded as an exact theoretical model of polarons. Many features of the spectrum structure, such as the existence of a δ -function peak for the attractive polaron and a dark continuum, are rigorously confirmed to be universal. The pairing gap also protects the polaron against thermal fluctuation, preserving clear polaron features in response functions at a finite temperature $k_B T \sim \Delta$. Furthermore, we discover that the polaron spectrum can be applied to measure the background superfluid excitation spectrum, such as the pairing gap Δ . Interestingly, in the case of a magnetic impurity, the polaron spectrum at finite but low temperature has sharp features that can be used to measure the sub-gap YSR bound state. For non-magnetic impurity, we predict the existence of a long-lived repulsive polaron.

We are grateful to Xing-Can Yao for insightful discussions. This research was supported by the Australian Research Council's (ARC) Discovery Program, Grants No. DE180100592 and No. DP190100815 (J.W.), and Grant No. DP180102018 (X.-J.L).

-
- [1] Immanuel Bloch, Jean Dalibard, and Wilhelm Zwerger, "Many-body physics with ultracold gases," *Rev. Mod. Phys.* **80**, 885–964 (2008).
- [2] P. Massignan, M. Zaccanti, and G. M. Bruun, "Polarons, dressed molecules and itinerant ferromagnetism in ultracold Fermi gases," *Rep. Prog. Phys.* **77**, 034401 (2014).
- [3] R. Schmidt, M. Knap, D. A. Ivanov, J.-S. You, M. Cetina, and E. Demler, "Universal many-body response of heavy impurities coupled to a Fermi sea: a review of recent progress," *Rep. Prog. Phys.* **81**, 024401 (2018).
- [4] Cheng Chin, Rudolf Grimm, Paul Julienne, and Eite Tiesinga, "Feshbach resonances in ultracold gases," *Rev. Mod. Phys.* **82**, 1225–1286 (2010).
- [5] Jonathan B. Balewski, Alexander T. Krupp, Anita Gaj, David Peter, Hans Peter B"uchler, Robert Löw, Sebastian Hofferberth, and Tilman Pfau, "Coupling a single electron to a Bose-Einstein condensate," *Nature (London)* **502**, 664–667 (2013).
- [6] Jia Wang, Marko Gacesa, and R. Côté, "Rydberg electrons in a Bose-Einstein condensate," *Phys. Rev. Lett.* **114**, 243003 (2015).
- [7] John Sous, H. R. Sadeghpour, T. C. Killian, Eugene Demler, and Richard Schmidt, "Rydberg impurity in a fermi gas: Quantum statistics and rotational blockade," *Phys. Rev. Research* **2**, 023021 (2020).
- [8] Richard Schmidt and Mikhail Lemeshko, "Rotation of quantum impurities in the presence of a many-body environment," *Phys. Rev. Lett.* **114**, 203001 (2015).
- [9] Michael Knap, Aditya Shashi, Yusuke Nishida, Adilet Imambekov, Dmitry A. Abanin, and Eugene Demler, "Time-dependent impurity in ultracold fermions: Orthogonality catastrophe and beyond," *Phys. Rev. X* **2**, 041020 (2012).
- [10] L. Landau, "Über die bewegung der elektronen im kristallgitter," *Phys. Z. Soviet.* **3**, 664 (1933).
- [11] André Schirotzek, Cheng-Hsun Wu, Ariel Sommer, and Martin W. Zwierlein, "Observation of Fermi polarons in a tunable Fermi liquid of ultracold atoms," *Phys. Rev. Lett.* **102**, 230402 (2009).
- [12] Y. Zhang, W. Ong, I. Arakelyan, and J. E. Thomas, "Polaron-to-polaron transitions in the radio-frequency spectrum of a quasi-two-dimensional fermi gas," *Phys. Rev. Lett.* **108**, 235302 (2012).
- [13] C. Kohstall, M. Zaccanti, M. Jag, A. Trenkwalder, P. Massignan, G. M. Bruun, F. Schreck, and R. Grimm, "Metastability and coherence of repulsive polarons in a strongly interacting Fermi mixture," *Nature (London)* **485**, 615 (2012).
- [14] Marco Koschorreck, Daniel Pertot, Enrico Vogt, Bernd Fröhlich, Michael Feld, and Michael Köhl, "Attractive and repulsive Fermi polarons in two dimensions," *Nature (London)* **485**, 619 (2012).
- [15] M. Cetina, M. Jag, R. S. Lous, I. Fritsche, J. T. M. Walraven, R. Grimm, J. Levinsen, M. M. Parish, R. Schmidt, M. Knap, and E. Demler, "Ultrafast many-body interferometry of impurities coupled to a fermi sea," *Science* **354**, 96 (2016).
- [16] Ming-Guang Hu, Michael J. Van de Graaff, Dhruv Kedar, John P. Corson, Eric A. Cornell, and Deborah S. Jin, "Bose polarons in the strongly interacting regime," *Phys. Rev. Lett.* **117**, 055301 (2016).
- [17] Nils B. Jørgensen, Lars Wacker, Kristoffer T. Skalmstang, Meera M. Parish, Jesper Levinsen, Rasmus S. Christensen, Georg M. Bruun, and Jan J. Arlt, "Observation of attractive and repulsive polarons in a Bose-Einstein condensate," *Phys. Rev. Lett.* **117**, 055302 (2016).
- [18] F. Scazza, G. Valtolina, P. Massignan, A. Recati, A. Amico, A. Burchianti, C. Fort, M. Inguscio, M. Zaccanti, and G. Roati, "Repulsive fermi polarons in a resonant mixture of ultracold ${}^6\text{Li}$ atoms," *Phys. Rev. Lett.* **118**, 083602 (2017).
- [19] Zhenjie Yan, Parth B. Patel, Biswaroop Mukherjee, Richard J. Fletcher, Julian Struck, and Martin W. Zwierlein, "Boiling a unitary fermi liquid," *Phys. Rev. Lett.* **122**, 093401 (2019).
- [20] Z. Z. Yan, Y. Ni, C. Robens, and M.W. Zwierlein, "Bose polarons near quantum criticality," *Science* **368**, 190 (2020).
- [21] Gal Ness, Constantine Shkedrov, Yanay Florshaim, Oriana K. Diessel, Jonas von Milczewski, Richard Schmidt, and Yoav Sagi, "Observation of a smooth polaron-molecule transition in a degenerate Fermi gas," *Phys. Rev. X* **10**, 041019 (2020).
- [22] F. Chevy, "Universal phase diagram of a strongly interacting Fermi gas with unbalanced spin populations," *Phys. Rev. A* **74**, 063628 (2006).
- [23] C. Lobo, A. Recati, S. Giorgini, and S. Stringari, "Normal state of a polarized Fermi gas at unitarity," *Phys. Rev. Lett.* **97**, 200403 (2006).
- [24] R. Combescot, A. Recati, C. Lobo, and F. Chevy, "Normal state of highly polarized Fermi gases: Simple many-body approaches," *Phys. Rev. Lett.* **98**, 180402 (2007).

- [25] M. Punk, P. T. Dumitrescu, and W. Zwerger, “Polaron-to-molecule transition in a strongly imbalanced Fermi gas,” *Phys. Rev. A* **80**, 053605 (2009).
- [26] Xiaoling Cui and Hui Zhai, “Stability of a fully magnetized ferromagnetic state in repulsively interacting ultracold Fermi gases,” *Phys. Rev. A* **81**, 041602 (2010).
- [27] Charles J. M. Mathy, Meera M. Parish, and David A. Huse, “Trimers, molecules, and polarons in mass-imbalanced atomic Fermi gases,” *Phys. Rev. Lett.* **106**, 166404 (2011).
- [28] Richard Schmidt, Tilman Enss, Ville Pietilä, and Eugene Demler, “Fermi polarons in two dimensions,” *Phys. Rev. A* **85**, 021602 (2012).
- [29] Steffen Patrick Rath and Richard Schmidt, “Field-theoretical study of the Bose polaron,” *Phys. Rev. A* **88**, 053632 (2013).
- [30] Aditya Shashi, Fabian Grusdt, Dmitry A. Abanin, and Eugene Demler, “Radio-frequency spectroscopy of polarons in ultracold Bose gases,” *Phys. Rev. A* **89**, 053617 (2014).
- [31] Weiran Li and S. Das Sarma, “Variational study of polarons in Bose-Einstein condensates,” *Phys. Rev. A* **90**, 013618 (2014).
- [32] Peter Kroiss and Lode Pollet, “Diagrammatic monte carlo study of a mass-imbalanced fermi-polaron system,” *Phys. Rev. B* **91**, 144507 (2015).
- [33] Jesper Levinsen, Meera M. Parish, and Georg M. Bruun, “Impurity in a bose-einstein condensate and the efimov effect,” *Phys. Rev. Lett.* **115**, 125302 (2015).
- [34] Hui Hu, An-Bang Wang, Su Yi, and Xia-Ji Liu, “Fermi polaron in a one-dimensional quasiperiodic optical lattice: The simplest many-body localization challenge,” *Phys. Rev. A* **93**, 053601 (2016).
- [35] Olga Goulko, Andrey S. Mishchenko, Nikolay Prokof’ev, and Boris Svistunov, “Dark continuum in the spectral function of the resonant fermi polaron,” *Phys. Rev. A* **94**, 051605 (2016).
- [36] Hui Hu, Brendan C. Mulkerin, Jia Wang, and Xia-Ji Liu, “Attractive fermi polarons at nonzero temperatures with a finite impurity concentration,” *Phys. Rev. A* **98**, 013626 (2018).
- [37] L. A. Peña Ardila, N. B. Jørgensen, T. Pohl, S. Giorgini, G. M. Bruun, and J. J. Arlt, “Analyzing a bose polaron across resonant interactions,” *Phys. Rev. A* **99**, 063607 (2019).
- [38] B. C. Mulkerin, X.-J. Liu, and H. Hu, “Breakdown of the fermi polaron description near fermi degeneracy at unitarity,” *Ann. Phys. (NY)* **407**, 29 (2019).
- [39] Jia Wang, Xia-Ji Liu, and Hui Hu, “Roton-induced bose polaron in the presence of synthetic spin-orbit coupling,” *Phys. Rev. Lett.* **123**, 213401 (2019).
- [40] Felipe Isaule, Ivan Morera, Pietro Massignan, and Bruno Juliá-Díaz, “Renormalization-group study of bose polarons,” *Phys. Rev. A* **104**, 023317 (2021).
- [41] Renato Pessoa, S. A. Vitiello, and L. A. Peña Ardila, “Finite-range effects in the unitary fermi polaron,” *Phys. Rev. A* **104**, 043313 (2021).
- [42] Kushal Seetharam, Yulia Shchadilova, Fabian Grusdt, Mikhail B. Zvonarev, and Eugene Demler, “Dynamical quantum cherenkov transition of fast impurities in quantum liquids,” *Phys. Rev. Lett.* **127**, 185302 (2021).
- [43] Yusuke Nishida, “Polaronic atom-trimer continuity in three-component fermi gases,” *Phys. Rev. Lett.* **114**, 115302 (2015).
- [44] Wei Yi and Xiaoling Cui, “Polarons in ultracold fermi superfluids,” *Phys. Rev. A* **92**, 013620 (2015).
- [45] M. Pierce, X. Leyronas, and F. Chevy, “Few versus many-body physics of an impurity immersed in a superfluid of spin 1/2 attractive fermions,” *Phys. Rev. Lett.* **123**, 080403 (2019).
- [46] Hui Hu, Jia Wang, Jing Zhou, and Xia-Ji Liu, “Crossover polarons in a strongly interacting fermi superfluid,” (2021), arXiv: 2111.01372.
- [47] A. Bigué, F. Chevy, and X. Leyronas, “Mean-field vs rpa calculation of the energy of an impurity immersed in a spin 1/2 superfluid,” (2022), arXiv: 2202.03222.
- [48] P. Nozières and C. T. De Dominicis, “Singularities in the x-ray absorption and emission of metals. iii. one-body theory exact solution,” *Phys. Rev.* **178**, 1097–1107 (1969).
- [49] Gerald D. Mahan, *Many Particle Physics*, 3rd ed. (Kluwer, New York, 2000).
- [50] U. Weiss, *Quantum Dissipative Systems*, Vol. 10 (World Scientific, Singapore, 1999).
- [51] A. Rosch, “Quantum-coherent transport of a heavy particle in a fermionic bath,” *Adv. Phys.* **48**, 295 (1999).
- [52] P. W. Anderson, “Infrared catastrophe in fermi gases with local scattering potentials,” *Phys. Rev. Lett.* **18**, 1049–1051 (1967).
- [53] Leonid S. Levitov and Hyunwoo Lee, “Electron counting statistics and coherent states of electric current,” *J. Math. Phys.* **37**, 4845 (1996).
- [54] I. Klich, *Full Counting Statistics: an Elementary Derivation of Levitov’s Formula* (Kluwer, Dordrecht, 2003).
- [55] K. Schönhammer, “Full counting statistics for noninteracting fermions: Exact results and the Levitov-Lesovik formula,” *Phys. Rev. B* **75**, 205329 (2007).
- [56] Dmitri A Ivanov and Alexander G Abanov, “Fisher-Hartwig expansion for Toeplitz determinants and the spectrum of a single-particle reduced density matrix for one-dimensional free fermions,” *J. Phys. A: Math. Theor.* **46**, 375005 (2013).
- [57] A. J. Leggett, “Cooper pairing in spin-polarized Fermi systems,” *J. Phys. (Paris)* **42**, C7 (1980).
- [58] P. Nozières and S. Schmitt-Rink, “Bose condensation in an attractive fermion gas: From weak to strong coupling superconductivity,” *J. Low Temp. Phys.* **59**, 195 (1985).
- [59] H. Hu, X.-J. Liu, and P. D. Drummond, “Equation of state of a superfluid fermi gas in the bcs-bec crossover,” *Europhys. Lett.* **74**, 574 (2006).
- [60] L. Yu, “Bound state in superconductors with paramagnetic impurities,” *Acta. Phys. Sin.* **21**, 75 (1965).
- [61] H. Shiba, “Classical spin in superconductors,” *Prog. Theor. Phys.* **40**, 435 (1968).
- [62] A. I. Rusinov, “Superconductivity near a paramagnetic impurity,” *JETP Lett. (USSR)* **9**, 85 (1969).
- [63] Eric Vernier, David Pekker, Martin W. Zwierlein, and Eugene Demler, “Bound states of a localized magnetic impurity in a superfluid of paired ultracold fermions,” *Phys. Rev. A* **83**, 033619 (2011).
- [64] Lei Jiang, Leslie O. Baksmaty, Hui Hu, Yan Chen, and Han Pu, “Single impurity in ultracold fermi superfluids,” *Phys. Rev. A* **83**, 061604 (2011).
- [65] J. Goold, T. Fogarty, N. Lo Gullo, M. Paternostro, and Th. Busch, “Orthogonality catastrophe as a consequence of qubit embedding in an ultracold fermi gas,” *Phys. Rev. A* **84**, 063632 (2011).
- [66] Jia Wang, Xia-Ji Liu, and Hui Hu, “Heavy polarons

in ultracold atomic fermi superfluids at the bec-bcs crossover: formalism and applications,” submitted to Phys. Rev. A. (2022).

- [67] One should not confuse a_σ with the scattering length at zero scattering energy, although the differences are negligible except very close to resonances.
- [68] Yanjun Ma, “X-ray edges of superconducting metals,” Phys. Rev. B **32**, 1472–1475 (1985).
- [69] Shih-Kuang Tung, Colin Parker, Jacob Johansen, Cheng Chin, Yujun Wang, and Paul S. Julienne, “Ultracold mixtures of atomic ^6Li and ^{133}Cs with tunable interactions,” Phys. Rev. A **87**, 010702 (2013).