

A characterization of g_2 -minimal normal 3-pseudomanifolds with at most four singularities

Biplab Basak¹, Raju Kumar Gupta and Sourav Sarkar

Department of Mathematics, Indian Institute of Technology Delhi, New Delhi 110016, India².

February 08, 2024

Abstract

Let Δ be a g_2 -minimal normal 3-pseudomanifold. A vertex in Δ whose link is not a sphere is called a singular vertex. When Δ contains at most two singular vertices, its combinatorial characterization is known [9]. In this article, we present a combinatorial characterization of such a Δ when it has three singular vertices, including one $\mathbb{RP}^2$ -singularity, or four singular vertices, including two $\mathbb{RP}^2$ -singularities. In both cases, we prove that Δ is obtained from a one-vertex suspension of a surface, and some boundary complexes of 4-simplices by applying the combinatorial operations of types connected sums, vertex foldings, and edge foldings.

MSC 2020: Primary 57Q05; Secondary 05E45, 51B10, 51E20, 52B70.

Keywords: Normal pseudomanifolds, vertex folding, edge folding, one-vertex suspension.

1 Introduction

The study of pseudomanifolds, and in particular normal pseudomanifolds, has been one of the central topics in combinatorial topology. Structures for such objects from a combinatorial and topological viewpoint in terms of simplicial and cell complexes are being developed by various researchers throughout the world. A basic and fundamental enumerative invariant of any n -dimensional simplicial complex, Δ , is its f -vector $(f_{-1}, f_0, f_1, \dots, f_n)$, where f_i is the number of i -dimensional faces of Δ for $-1 \leq i \leq n$. The empty set is considered to be the only face of dimension -1 . Inequalities in the lower and upper bound conjectures involving f_i have been used as a very interesting and powerful tool to analyze the interplay between the f -vector of a complex and its geometric carrier. In 1970, a breakthrough in this direction was given by Walkup [19]. Inspired by [19], a new combinatorial invariant $g_2(\Delta)$ has been defined as $g_2(\Delta) = f_1 - (d+1)f_0 + \binom{d+2}{2}$ for every d -dimensional simplicial complex Δ . From [19], it is known that for a triangulated 3-manifold Δ , $g_2(\Delta) \geq 0$ with the equality occurring when Δ is a triangulation of a stacked sphere. Barnette [4, 5] achieved the same results in higher dimensions. In 1987, Kalai [13] used the concept of rigidity and framework on graphs to establish the lower bounds of f -vector for normal pseudomanifolds of dimension at least three, where 2-dimensional links were triangulated spheres.

¹*Corresponding author*: Biplab Basak

²*E-mail addresses*: biplab@iitd.ac.in (B. Basak), Raju.Kumar.Gupta@maths.iitd.ac.in (R. K. Gupta), Sourav.Sarkar@maths.iitd.ac.in (S. Sarkar).

Kalai's findings establish a lower bound on the value of g_2 for a normal d -pseudomanifold, where 2-dimensional links are spheres, expressed in terms of the same for a link. Subsequently, Fogelsanger's results in [8, Chapter 8] removed the restriction on 2-dimensional links. The combined implications of Kalai and Fogelsanger's results assert that if $d \geq 0$ and Δ is a normal d -pseudomanifold, then for any face σ of Δ with co-dimension 3 or more, the inequality $g_2(\Delta) \geq g_2(\text{lk}(\sigma, \Delta))$ holds. In [12], the non-negativity of the invariant g_2 has been described in a different way of looking at rigidity, and in [3], a detailed study on the lower bound theorem has been given in terms of Gromov rigidity. For the case of the normal 3-pseudomanifold Δ , the result turned out to be $g_2(\Delta) \geq g_2(\text{lk}(v, \Delta))$ for every vertex v in Δ . If $g_2(\Delta) = g_2(\text{lk}(v, \Delta))$ for some vertex v , then such a complex Δ is said to be g_2 -minimal (or have relatively minimal g_2 [9]). Basak and Swartz [9] proved that if Δ is a g_2 -minimal normal 3-pseudomanifold with at most two singular vertices, then Δ is obtained from a one-vertex suspension of a surface and some boundary complexes of 4-simplices by a sequence of operations of the form vertex foldings and connected sums. In this article, we have worked with the g_2 -minimal normal 3-pseudomanifolds with at most four singularities. More precisely, we proved the following:

Theorem 1.1. *Let Δ be a normal 3-pseudomanifold such that Δ has n singular vertices among which $n - 2$ have $\mathbb{RP}^2$ -singularities, where $3 \leq n \leq 4$. Then Δ is obtained from a one-vertex suspension of a surface, and some boundary complexes of 4-simplices by a sequence of operations of types connected sums, vertex foldings, and edge foldings.*

Several classifications and descriptions of triangulated manifolds and pseudomanifolds have been done based on the f -vector and the value of g_2 . In [2], the d -pseudomanifold with $d + 4$ vertices has been described. The work in [16] established that only a finite number of combinatorial manifolds are possible for a given upper bound of g_2 . In [17, 15], the f -vector was the key factor where homology manifolds and specific types of pseudomanifolds were involved. In [14] and [20], normal d -pseudomanifolds with $g_2 = 1$ and $g_2 = 2$ have been characterized, respectively. The concepts of rigidity and the frameworks described in [13, 18] are utilized in each case. In the recent article [7], normal 3-pseudomanifolds with at most two singular vertices are characterized. Additionally, [6] delves into the characterization of normal 3-pseudomanifolds with $g_2 \leq 4$. For a characterization of homology d -manifolds with $g_2 \leq 3$, one can refer to [8].

2 Preliminaries

A *simplicial complex* K is a finite collection of simplices in $\mathbb{R}^m$ for some $m \in \mathbb{N}$, such that for every simplex $\sigma \in K$, all of its faces are in K , and for any two simplices $\sigma, \tau \in K$, $\sigma \cap \tau$ is a face of both σ and τ . We assume that the empty set $\emptyset$ (considered a simplex of dimension -1) is a member of every simplicial complex. For a simplicial complex Δ , the notation $|\Delta|$ will be used to denote the geometric carrier, i.e., $|\Delta| := \cup_{\sigma \in \Delta} \sigma$. Two simplices, $\sigma = u_0 u_1 \cdots u_k$ and $\tau = v_0 v_1 \cdots v_l$ in $\mathbb{R}^n$ for some $n \in \mathbb{N}$, are considered skew if $u_0, \dots, u_k, v_0, \dots, v_l$ are affinely independent. In that case, $u_0 \cdots u_k v_0 \cdots v_l$ is a $(k + l + 1)$ -simplex and is denoted by $\sigma \star \tau$ or $\sigma\tau$. Two simplicial complexes Δ_1 and Δ_2 in some $\mathbb{R}^m$ are called *skew* if σ and τ are skew for every simplex $\sigma \in \Delta_1$ and $\tau \in \Delta_2$. The join of two skew simplicial complexes Δ_1 and Δ_2 is defined as $\{\sigma \star \tau : \sigma \in \Delta_1, \tau \in \Delta_2\}$, and is denoted by $\Delta_1 \star \Delta_2$. In particular, the join $\sigma \star \Delta_1$, where σ is an i -simplex and Δ_1 is a j -dimensional simplicial complex, is defined as the $(i + j + 1)$ -dimensional simplicial complex $\{\tau : \tau \leq \sigma\} \star \Delta_1$. If a simplex τ is a face of another simplex σ , then we denote it by $\tau \leq \sigma$.

If S is a collection of simplices, then the sub-collection of all the simplices of dimension at most one in S is called the *graph* of S and is denoted by $G(S)$. The link of a face σ in Δ is the collection of faces in Δ that do not intersect σ and whose join with σ lies in Δ and is denoted by $\text{lk}(\sigma, \Delta)$. The star of a face σ in Δ is $\{\gamma : \gamma \leq \sigma \star \alpha \text{ and } \alpha \in \text{lk}(\sigma, \Delta)\}$ and is denoted by $\text{st}(\sigma, \Delta)$. If the underlying simplicial complex is specified, we then simply write $\text{lk}(\sigma)$ and $\text{st}(\sigma)$ for $\text{lk}(\sigma, \Delta)$ and $\text{st}(\sigma, \Delta)$, respectively. A maximal face in Δ (i.e., a maximal simplex in Δ) is called a *facet* of Δ , and if all the facets are of the same dimension, then we say Δ is a pure simplicial complex. For a simplex σ , its boundary complex is defined as $\{\tau : \tau \leq \sigma \text{ and } \tau \neq \sigma\}$.

Several combinatorial tools have been defined to classify simplicial complexes, and taking the suspension of a simplicial complex is one such tool. In [3] and [10], another type of suspension has been defined for the class of normal pseudomanifolds by introducing one extra vertex instead of two. Let Δ be a normal $(d-1)$ -pseudomanifold, and $v \in \Delta$ be a vertex. Then, for a new symbol $u \notin \Delta$, consider the normal d -pseudomanifold $\sum_{v,u} \Delta := (v \star \{\tau \in \Delta : v \not\leq \tau\}) \cup (u \star \Delta)$. The geometric carrier of $\sum_{v,u} \Delta$ is the suspension of $|\Delta|$. The complex $\sum_{v,u} \Delta$ is called the *one-vertex suspension* of Δ with respect to the vertex v . Note that in $\sum_{v,u} \Delta$, $\text{lk } u$ and $\text{lk } v$ are simplicially isomorphic to Δ .

A d -pseudomanifold is a pure d -dimensional simplicial complex Δ such that every face of dimension $d-1$ is contained in exactly two facets, and for any pair of facets σ and τ in Δ , there is a sequence of facets $\sigma_1, \sigma_2, \dots, \sigma_m$ in Δ such that $\sigma = \sigma_1, \tau = \sigma_m$, and $\sigma_i \cap \sigma_{i+1}$ is a $(d-1)$ -simplex in Δ for $1 \leq i \leq m-1$. The d -pseudomanifolds in which the links of all faces of dimensions up to $d-2$ are connected are called *normal*. In particular, a normal 3-pseudomanifold is a strongly connected 3-dimensional pure simplicial complex where the geometric carrier of the link of each vertex is a closed and connected surface. A vertex in a normal pseudomanifold Δ whose link is a triangulated sphere is called a *non-singular vertex* of Δ . If t is a vertex in a normal pseudomanifold Δ whose link is not a triangulated sphere, we call t a *singular vertex* of Δ , and if $|\text{lk}(t, \Delta)| \cong S$, we say t has S -singularity. Two combinatorial tools, vertex folding, and edge folding, are first introduced in [9], and these two are going to be the main key factors in our result.

Definition 2.1. Let σ_1 and σ_2 be two facets of a simplicial complex Δ whose intersection is a single vertex x . Suppose there is a simplicial bijection $\psi : \sigma_1 \rightarrow \sigma_2$ such that $\psi(x) = x$ and for all other vertices $y \leq \sigma_1$, the only path of length two or less from y to $\psi(y)$ is $(y, x, \psi(y))$. Then we can obtain a complex Δ_x^ψ by identifying all the faces $\rho \leq \sigma_1$ with $\psi(\rho) \leq \sigma_2$ and then removing the identified facet. We call Δ_x^ψ a *vertex folding* of Δ at x , and in this case, Δ is called a *vertex unfolding* of Δ_x^ψ .

Definition 2.2. Let σ_1 and σ_2 be two facets of a simplicial complex Δ whose intersection is a single edge xy . Suppose there is a simplicial bijection $\psi : \sigma_1 \rightarrow \sigma_2$ such that $\psi(x) = x$, $\psi(y) = y$, and for all other vertices $u \leq \sigma_1$, all paths of length two or less from u to $\psi(u)$ pass through either x or y . Then we can obtain a complex Δ_{xy}^ψ by identifying all the faces $\rho \leq \sigma_1$ with $\psi(\rho) \leq \sigma_2$ and then removing the identified facet. We call Δ_{xy}^ψ an *edge folding* of Δ at xy , and in this case, Δ is called an *edge unfolding* of Δ_{xy}^ψ .

If Δ is a normal d -pseudomanifold, then Δ_{xy}^ψ is a pseudomanifold that need not be normal. Moreover,

$$g_2(\Delta_{xy}^\psi) = g_2(\Delta) + \binom{d}{2}, \quad \text{and}$$

$$\text{lk}(x, \Delta_{xy}^\psi) = \text{lk}(x, \Delta)_y^\psi.$$

After applying vertex folding and edge folding to a simplicial complex, the vertices of two facets get identified by the admissible map ψ in the new complex. However, to avoid ambiguity in notations in the new complex, we denote the identified vertices with the same notation as in the original complex.

The definition of a simplicial complex implies that sometimes Δ may contain all the proper faces of a simplex σ , but $\sigma \notin \Delta$. In such a case, we say σ is a missing face of Δ . If S is a subset of Δ such that $xy \in S$ but the vertices x and y are not in S , then we say that xy is an open edge of S and denote it by (x, y) . The presence of a missing tetrahedron and an open edge in a normal 3-pseudomanifold and its subsets provides information regarding the construction of the same.

Proposition 2.3 ([9]). *Let Δ be a normal 3-pseudomanifold. Suppose $pqrs$ is a missing facet in Δ such that $\partial(qrs)$ and $\partial(prs)$ separate $\text{lk}(p, \Delta)$ and $\text{lk}(q, \Delta)$, respectively, and a small neighborhood of $|\partial(pqr)|$ in $|\text{lk}(s, \Delta)|$ is a Möbius strip. Then a small neighborhood of $|\partial(pqs)|$ in $|\text{lk}(r, \Delta)|$ is a Möbius strip. Further, there exists a normal 3-pseudomanifold Δ' such that Δ is obtained from Δ' by an edge folding at $rs \in \Delta'$, and $pqrs$ is the image of the removed facet.*

Proposition 2.4 ([9]). *Let Δ be a g_2 -minimal normal 3-pseudomanifold with at most two singular vertices, t and s , such that $g_2(\text{lk}(t, \Delta)) \geq g_2(\text{lk}(s, \Delta))$. Then Δ is obtained from a one-vertex suspension of $\text{lk}(s, \Delta)$ and some boundary complexes of 4-simplices by a sequence of operations of the form vertex foldings and connected sums. In particular, if Δ has exactly one singular vertex, then Δ is obtained from some boundary complexes of 4-simplices by a sequence of operations of the form vertex foldings and connected sums.*

In Proposition 2.4, it is evident that if $\text{lk}(t, \Delta) \cong \text{lk}(s, \Delta) \#_n H$, where H is either Torus or Klein bottle, then Δ is obtained from a one-vertex suspension of $\text{lk}(s, \Delta)$ and some boundary complexes of 4-simplices by exactly n vertex foldings and finitely many connected sums.

Proposition 2.5 ([9]). *Let Δ be a g_2 -minimal normal 3-pseudomanifold. Then there exists a normal 3-pseudomanifold $\bar{\Delta}$ and a vertex $t \in \bar{\Delta}$ such that Δ is obtained from $\bar{\Delta}$ by (possibly zero) facet subdivisions, and $G(\bar{\Delta}) = G(\text{st}(t, \bar{\Delta}))$.*

3 Normal 3-pseudomanifolds with three or four singularities

Let Δ be a normal 3-pseudomanifold and uv be an edge in Δ . Then $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : u \leq \sigma\}$ is a triangulation of a surface with boundary. Let K be a triangulation of a surface with a boundary. An edge e is called an *interior edge* of K if e is contained in exactly two triangles in K , and an edge e is called a *boundary edge* of K if e is contained in exactly one triangle in K . A vertex v is called a *boundary vertex* of K if there is a boundary edge of K containing v . A vertex u is referred to as an *interior vertex* of K if there is no boundary edge of K containing u .

Lemma 3.1. *Let Δ be a normal 3-pseudomanifold such that $G(\Delta) = G(\text{st}(t, \Delta))$, and let v be a vertex in Δ such that $v \neq t$. Then $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$ does not contain any interior vertex.*

Proof. If $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$ has an interior vertex, say x , then $x \in \text{lk}(t, \Delta)$, and vx is an edge in Δ , but $vx \notin \text{lk}(t, \Delta)$. This leads to a contradiction. $\square$

Let v be a singular vertex in Δ . If $|\text{lk}(v, \Delta)| \cong \mathbb{RP}^2$, then $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$ is a Möbius strip, and by Lemma 3.1, $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$ has no interior vertices. Let M be a Möbius strip that does not have any interior vertices. An edge $xy \in M$ will be called a *cut edge* if the edge xy is an interior edge of M , and xy separates M into two portions. Then one portion will be a disc, and the other portion will be another Möbius strip. If xy is an interior edge of M , and xy does not separate M , then we say xy is a *non-cut edge* of the Möbius strip M .

Lemma 3.2. *Let Δ be a normal 3-pseudomanifold such that $G(\Delta) = G(\text{st}(t, \Delta))$ and Δ has exactly three singular vertices, including t . Let v be a singular vertex in Δ such that $v \neq t$ and $|\text{lk}(v, \Delta)| \cong \mathbb{RP}^2$. Then $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$ contains a non-cut edge xy , where x and y are non-singular vertices in Δ . Moreover, the non-cut edge xy creates a missing tetrahedron $xyvt$ in Δ such that a small neighborhood of $|\partial(xyvt)|$ in $|\text{lk}(v, \Delta)|$ is a Möbius strip.*

Proof. From Lemma 3.1, we know that the Möbius strip $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$ does not contain any interior vertex. If the Möbius strip $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$ has a cut edge, then we can cut along the cut edge, and we get another Möbius strip. Continuing this a finite number of times, we have a Möbius strip M where each interior edge will be a non-cut edge. Note that a non-cut edge of M is also a non-cut edge of $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$. Furthermore, any vertex $x \in M$ is on the boundary of M and has a degree of at least 3. Moreover, x is adjacent to two vertices in the boundary of M , and for the other vertex y , if xy is an edge in M , then xy is an interior edge, and hence a non-cut edge of M . Let p be the singular vertex other than t and v in Δ . If possible, let M not contain an interior edge xy , where x and y are non-singular vertices in Δ . Then, all the interior edges of M are incident to p in M . If a non-singular vertex u is not adjacent to p in M , then $d(u) \geq 3$ implies that there is a non-singular vertex x such that ux is an interior edge of M , leading to a contradiction. Thus, all non-singular vertices of M must be adjacent to p in M . Consequently, the Möbius strip M is the same as $\text{st}(p, M)$. This is contradictory, as $\text{st}(p, M)$ is a 2-ball. Therefore, M contains an interior edge xy , where x and y are non-singular vertices in Δ . Since each interior edge of M is a non-cut edge of M , and a non-cut edge of M is also a non-cut edge of $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$, the result is achieved.

It follows from Lemma 3.1 that $x, y \in \text{lk}(tv, \Delta)$. Since xy is an interior edge of $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$, $xy \notin \text{lk}(tv, \Delta)$. Thus, $xyvt \notin \Delta$. On the other hand, $xy \in \text{lk}(v, \Delta)$, $x, y \in \text{lk}(tv, \Delta)$, and $G(\Delta) = G(\text{st}(t, \Delta))$ imply that $\partial(xyvt) \subset \Delta$. Therefore, $xyvt$ is a missing tetrahedron in Δ . Since xy is a non-cut edge of $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$, $\partial(txy)$ does not separate $\text{lk}(v, \Delta)$. Since $|\text{lk}(v, \Delta)| \cong \mathbb{RP}^2$, a small neighborhood of $|\partial(txy)|$ in $|\text{lk}(v, \Delta)|$ is a Möbius strip. $\square$

Lemma 3.3. *Let Δ be a normal 3-pseudomanifold such that $G(\Delta) = G(\text{st}(t, \Delta))$ and Δ has exactly four singular vertices, including t . Let v be a singular vertex in Δ such that $v \neq t$ and $|\text{lk}(v, \Delta)| \cong \mathbb{RP}^2$. Then $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$ contains a non-cut edge xy , where x and y are non-singular vertices in Δ . Moreover, the non-cut edge xy creates a missing tetrahedron $xyvt$ in Δ such that a small neighborhood of $|\partial(xyvt)|$ in $|\text{lk}(v, \Delta)|$ is a Möbius strip.*

Proof. Let p and q be two singular vertices other than v and t in Δ . From Lemma 3.1, we know that the Möbius strip $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$ does not contain any interior vertex. If the Möbius strip $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$ has a cut edge, then we can

cut along the cut edge and get another Möbius strip. Continuing this a finite number of times, we have a Möbius strip M where each interior edge is a non-cut edge. Furthermore, any vertex $x \in M$ is in the boundary of M and has a degree of at least 3. Moreover, x is adjacent to two vertices in the boundary of M , and for the other vertex y , if xy is an edge in M , then xy is an interior edge and hence a non-cut edge of M . If M contains exactly one singular vertex, then we achieve our result by using similar arguments as in Lemma 3.2.




Figure 1

Assume that M contains both singular vertices p and q . If possible, suppose there is no interior edge xy in M , where x and y are non-singular vertices. In such a case, all the interior edges of M are incident to either p or q . If a non-singular vertex u is adjacent to neither p nor q in M , then $d(u) \geq 3$ implies the existence of a non-singular vertex x such that ux is an interior edge of M , leading to a contradiction. Thus, all non-singular vertices of M must be connected to either p or q in M . Consequently, the Möbius strip M is the same as the union of $\text{st}(p, M)$ and $\text{st}(q, M)$. Now we consider three possibilities.

Case 1: Let the edge $pq \notin M$. Then, the stars $\text{st}(p, M)$ and $\text{st}(q, M)$ take the forms depicted in Figure 1, where p_i and q_j are non-singular vertices, $1 \leq i \leq n$, $1 \leq j \leq m$. If we identify $p_i p_{i+1}$ with $q_j q_{j+1}$ or $q_{j+1} q_j$, then the identified edge will be in the interior of M , which is a contradiction. Therefore, $\text{st}(p, M)$ and $\text{st}(q, M)$ are disjoint, which contradicts the fact that the Möbius strip M is the same as the union of $\text{st}(p, M)$ and $\text{st}(q, M)$.

Case 2: Let pq be a boundary edge of M . Suppose $\text{lk}(p, M) = P(p_1, \dots, p_n, x, q)$. Then, no edge of the form qz can be identified with some $p_i p_{i+1}$ or $p_{i+1} p_i$. Let $\text{lk}(q, M) = P(q_1, \dots, q_m, x, p)$. Here, x, p_i , and q_j are non-singular vertices, $1 \leq i \leq n$, $1 \leq j \leq m$. If we identify $p_i p_{i+1}$ with $q_j q_{j+1}$ or $q_{j+1} q_j$, then the identified edge will be an interior edge of M , which is a contradiction. Therefore, the union of $\text{st}(p, M)$ and $\text{st}(q, M)$ is a 2-ball (cf. Figure 2 (a)), contradicting the fact that the Möbius strip M is the same as the union of $\text{st}(p, M)$ and $\text{st}(q, M)$.

Case 3: Let pq be an interior edge of M . Let $\text{lk}(p, M) = P(p_1, p_2, \dots, p_{n_1}, x, q, y, p'_{n_2}, p'_{n_2-1}, \dots, p'_2, p'_1)$. Then, no edge of the form qz can be identified with an edge in the path $P(p_{n_1}, p_{n_1-1}, \dots, p_1, p, p'_1, p'_2, \dots, p'_{n_2})$. Let $\text{lk}(q, M) = P(q_1, q_2, \dots, q_{m_1}, x, p, y, q'_{m_2}, q'_{m_2-1}, \dots, q'_2, q'_1)$. Here, x, y, p_i, p'_j, q_k , and q'_l are non-singular vertices, $1 \leq i \leq n_1$, $1 \leq j \leq n_2$, $1 \leq k \leq m_1$, $1 \leq l \leq m_2$. If we identify any edge from the paths $P(p_{n_1}, p_{n_1-1}, \dots, p_1)$ or $P(p'_1, p'_2, \dots, p'_{n_2})$ with an edge from the paths $P(q_{m_1}, q_{m_1-1}, \dots, q_1)$ or $P(q'_1, q'_2, \dots, q'_{m_2})$, then the identified edge will be an interior edge of M , leading to a contradiction. Consequently, the union of $\text{st}(p, M)$ and $\text{st}(q, M)$ is a 2-ball (cf. Figure 2 (b)), contradicting the fact that the Möbius strip M is the same as the union of $\text{st}(p, M)$ and $\text{st}(q, M)$.




Figure 2

Therefore, M contains an interior edge xy , where x and y are non-singular vertices in Δ . Since every interior edge of M is a non-cut edge, and a non-cut edge of M is likewise a non-cut edge of $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$, we have successfully established the result.

Using similar arguments as those in Lemma 3.2, it can be concluded that $xyvt$ is a missing tetrahedron in Δ . Since xy is a non-cut edge of $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$, the boundary $\partial(txy)$ does not separate $\text{lk}(v, \Delta)$. Since $|\text{lk}(v, \Delta)| \cong \mathbb{RP}^2$, a small neighborhood of $|\partial(txy)|$ in $|\text{lk}(v, \Delta)|$ is a Möbius strip. $\square$

Theorem 3.4. *Let Δ be a g_2 -minimal normal 3-pseudomanifold, and let t be a singular vertex in Δ such that $G(\Delta) = G(\text{st}(t, \Delta))$.*

1. *If Δ has three singular vertices, including one $\mathbb{RP}^2$ -singularity, then Δ is obtained by an edge folding from a g_2 -minimal normal 3-pseudomanifold with exactly two singularities.*
2. *If Δ has four singular vertices, including two $\mathbb{RP}^2$ -singularities, then Δ is obtained by two repeated edge foldings on a g_2 -minimal normal 3-pseudomanifold with exactly two singularities.*

Proof. First, we assume that Δ has three singular vertices, and let the singular vertices of Δ other than t be s and v_1 , where $|\text{lk}(v_1, \Delta)| \cong \mathbb{RP}^2$. Since $|\text{lk}(v_1, \Delta)| \cong \mathbb{RP}^2$, by Lemma 3.2, there is a missing tetrahedron tv_1xy , where x and y are non-singular vertices. Moreover, a small neighborhood of $|\partial(txy)|$ in $|\text{lk}(v_1, \Delta)|$ is a Möbius strip. According to Proposition 2.3, there exists a normal 3-pseudomanifold Δ' such that $\Delta = (\Delta')_{tv_1}^\psi$. Since Δ is g_2 -minimal with respect to t , Δ' is also g_2 -minimal with respect to t . On the other hand, $g_2(\text{lk}(v_1, \Delta')) = g_2(\text{lk}(v_1, \Delta)) - 3 = 0$, and hence v_1 is a non-singular vertex in Δ' . Since $\text{lk}(s, \Delta)$ is isomorphic to $\text{lk}(s, \Delta')$, Δ' is a g_2 -minimal normal pseudomanifold with two singular vertices t and s . Moreover, $g_2(\Delta') = g_2(\text{lk}(t, \Delta'))$, and Δ is obtained from Δ' by an edge folding along the edge tv_1 , where v_1 is a non-singular vertex in Δ' .

Now, let Δ have four singular vertices, including two $\mathbb{RP}^2$ -singularities. Let v_1 and v_2 be the vertices in Δ such that $|\text{lk}(v_1, \Delta)| \cong |\text{lk}(v_2, \Delta)| \cong \mathbb{RP}^2$. Since $|\text{lk}(v_2, \Delta)| \cong \mathbb{RP}^2$, by Lemma 3.3, there is a missing tetrahedron tv_2wz , where w and z are non-singular vertices. Moreover, a small neighborhood of $|\partial(twz)|$ in $|\text{lk}(v_2, \Delta)|$ is a Möbius strip. Therefore, by Proposition 2.3, Δ is formed via an edge folding along the edge tv_2 from a normal 3-pseudomanifold Δ' . In Δ' , v_2 is a non-singular vertex because $g_2(\text{lk}(v_2, \Delta')) = g_2(\text{lk}(v_2, \Delta)) - 3 = 0$. Since the singular vertices other than t and v_2 did not participate in the process of edge folding, their link in Δ' will be isomorphic to the links in Δ . On the other hand, Δ' is also a g_2 -minimal normal pseudomanifold, with $g_2(\Delta') = g_2(\text{lk}(t, \Delta'))$. Thus, Δ' is a g_2 minimal normal pseudomanifold with three singular vertices, and $v_1 \in \Delta'$ is a vertex such that $|\text{lk}(v_1, \Delta')| \cong \mathbb{RP}^2$. By the same argument as in the last paragraph, Δ' is obtained from a g_2 -minimal normal 3-pseudomanifold Δ'' with exactly two singular vertices by an edge folding along the edge tv_1 , where v_1 is a non-singular vertex in Δ'' . Therefore, Δ is obtained from Δ'' by two edge foldings along the edges tv_1 and tv_2 , where v_1 and v_2 are non-singular vertices in Δ'' . $\square$

Let Δ be a g_2 -minimal normal 3-pseudomanifold with three singular vertices, t, s , and v , where $|\text{lk}(v, \Delta)| \cong \mathbb{RP}^2$. Also, let $g_2(\Delta) = g_2(\text{lk}(t, \Delta))$. It follows from Proposition 2.5 that $\Delta = \bar{\Delta} \# \Delta_1 \# \cdots \# \Delta_n$, where each Δ_i is the boundary complex of a 4-simplex, and $G(\bar{\Delta}) = G(\text{st}(t, \bar{\Delta}))$. Note that $\bar{\Delta}$ contains a copy of each of the vertices t, s , and v , with the links having the same Betti numbers as the link of those vertices in Δ . Let b_1 and b_2 be the first Betti numbers of $\text{lk}(t, \Delta)$ and $\text{lk}(s, \Delta)$, respectively. By Theorem 3.4, $\bar{\Delta}$ is obtained from a g_2 -minimal normal 3-pseudomanifold Δ' by an edge folding along an edge of the form tv , where v is a non-singular vertex in Δ' . Further, Δ' is g_2 -minimal with respect to the singular vertex $t \in \Delta'$, and the first Betti number of the link of s in Δ' is unchanged. Notice from the definition that an edge folding along an edge tx reduces the first Betti number of the corresponding vertices by 1 in the resulting complex, and it follows that $b_1 - b_2 \geq 1$. On the other hand, if Δ has four singular vertices, including two $\mathbb{RP}^2$ -singularities, and $G(\Delta) = G(\text{st}(t, \Delta))$, then Δ is obtained from a g_2 -minimal normal pseudomanifold by two repeated applications of edge folding. Therefore, by Proposition 2.5 and Theorem 3.4, we have the following:

Corollary 3.5. *Let Δ be a g_2 -minimal normal 3-pseudomanifold with four singular vertices t, s, v_1 , and v_2 , where $|\text{lk}(v_1, \Delta)| \cong |\text{lk}(v_2, \Delta)| \cong \mathbb{RP}^2$ and Δ is g_2 -minimal with respect to the vertex t . If b_1 and b_2 are the first Betti numbers of $\text{lk}(t, \Delta)$ and $\text{lk}(s, \Delta)$, respectively, then $b_1 - b_2 \geq 2$.*

Let Δ be a g_2 -minimal normal 3-pseudomanifold with three singular vertices, including one $\mathbb{RP}^2$ -singularity, or four singular vertices, including two $\mathbb{RP}^2$ -singularities. Let s and t be the other singular vertices in Δ , where $g_2(\Delta) = g_2(\text{lk}(t, \Delta))$. Then in the first case, $\text{lk}(t, \Delta) \cong \text{lk}(s, \Delta) \#_n \mathbb{T}^2 \# \mathbb{RP}^2$, and for the second case, $\text{lk}(t, \Delta) \cong \text{lk}(s, \Delta) \#_n \mathbb{T}^2 \# \mathbb{RP}^2 \# \mathbb{RP}^2$, where $\mathbb{T}^2$ denotes a triangulated torus. Thus, using Proposition 2.5, Theorem 3.4, and Proposition 2.4, we have the following corollaries:

Corollary 3.6. *Let Δ be a g_2 -minimal normal 3-pseudomanifold with three singular vertices, including one $\mathbb{RP}^2$ -singularity. Let t be the singular vertex in Δ such that $g_2(\Delta) = g_2(\text{lk}(t, \Delta))$. Then, Δ is obtained from a one-vertex suspension of $\text{lk}(s, \Delta)$ and some boundary complexes of 4-simplices by n vertex foldings, one edge folding, and finitely many connected sums, where $\text{lk}(t, \Delta) \cong \text{lk}(s, \Delta) \#_n \mathbb{T}^2 \# \mathbb{RP}^2$ for a singular vertex s .*

Corollary 3.7. *Let Δ be a g_2 -minimal normal 3-pseudomanifold with four singular vertices, including two $\mathbb{RP}^2$ -singularities. Let t be the singular vertex in Δ such that $g_2(\Delta) = g_2(\text{lk}(t, \Delta))$. Then Δ is obtained from a one-vertex suspension of $\text{lk}(s, \Delta)$ and some boundary complexes of 4-simplices by n vertex foldings, two edge foldings, and finitely many connected sums, where $\text{lk}(t, \Delta) \cong \text{lk}(s, \Delta) \#_n \mathbb{T}^2 \# \mathbb{RP}^2 \# \mathbb{RP}^2$ for a singular vertex s .*

Proof of Theorem 1.1. The proof follows from Corollaries 3.6 and 3.7. □

Remark 3.8. The proof of Theorem 1.1 depends on Lemmas 3.2 and 3.3, which give the existence of a non-cut edge xy in $\text{lk}(v, \Delta) \setminus \{\sigma \in \text{lk}(v, \Delta) : t \leq \sigma\}$, where v is a singular vertex and x, y are non-singular vertices in Δ . If Δ has more than four singular vertices, then such a non-cut edge may not exist. Therefore, if Δ has more than four singular vertices, then such a characterization for Δ may not be possible. In fact, we have an example of a g_2 -minimal normal 3-pseudomanifold Δ with five singular vertices, among which one has $\mathbb{T}^2$ -singularity and the remaining four have $\mathbb{RP}^2$ -singularities (see [1]), where Δ cannot be obtained from a normal 3-pseudomanifold Δ' by an edge folding.

Acknowledgement: The authors would like to thank the anonymous referees for their many useful comments and suggestions. The first author is supported by the Science and Engineering Research Board (CRG/2021/000859). The second author is supported by CSIR (India). The third author is supported by the Prime Minister's Research Fellows (PMRF) Scheme.

References

- [1] T. Akhmejanov, Triangulations of normal 3-pseudomanifolds on 9 Vertices, <https://pi.math.cornell.edu/~takhmejanov/pseudoManifolds.html>
- [2] B. Bagchi and B. Datta, A structure theorem for pseudomanifolds, *Discrete Math.* **188** (1998), no. 1-3, 41–60.
- [3] B. Bagchi and B. Datta, Lower bound theorem for normal pseudomanifolds, *Expo. Math.* **26** (2008), 327–351.
- [4] D. Barnette, A proof of the lower bound conjecture for convex polytopes, *Pacific J. Math.* **46** (1973), 349–354.
- [5] D. Barnette, Graph theorems for manifolds, *Israel J. Math.* **16** (1973), 62–72.
- [6] B. Basak and R. K. Gupta: A characterization of normal 3-pseudomanifolds with $g_2 \leq 4$, 2022, 13 pages, arXiv: 2202.06638v1.
- [7] B. Basak, R. K. Gupta and S. Sarkar, A characterization of normal 3-pseudomanifolds with at most two singularities, *Discrete Math.* **346** (12) (2023), Paper No. 113588, 15 pp.
- [8] B. Basak and S. Sarkar: On a construction of some homology d -manifolds, 2023, 22 pages.

- [9] B. Basak and E. Swartz, Three-dimensional normal pseudomanifolds with relatively few edges, *Advances in Mathematics* **365** (2020) 107035, 1–25.
- [10] A. Björner and F. Lutz, H. Simplicial manifolds, bistellar flips and a 16-vertex triangulation of the Poincaré homology 3-sphere. *Experiment. Math.* **9** (2000), 275–289.
- [11] A. Fogelsanger, The generic rigidity of minimal cycles, Ph.D. thesis, Cornell University, 1988.
- [12] M. Gromov, *Partial differential relations*, Springer, Berlin Heidelberg New York, 1986.
- [13] G. Kalai, Rigidity and the lower bound theorem I, *Invent. Math.* **88** (1987), 125–151.
- [14] E. Nevo and E. Novinsky, A characterization of simplicial polytopes with $g_2 = 1$, *J. Combin. Theory Ser. A* **118** (2011), 387–395.
- [15] I. Novik and E. Swartz, Face numbers of pseudomanifolds with isolated singularities, *Math. Scan.*, **110** (2012), 198–212.
- [16] E. Swartz, Topological finiteness for edge-vertex enumeration, *Adv. Math.*, **219** (2008), 1722–1728.
- [17] E. Swartz, Face enumeration: From spheres to manifolds, *J. Eur. Math. Soc.*, **11** (2009), 449–485.
- [18] T. Tay, N. White and W. Whiteley, Skeletal rigidity of simplicia complexes II, *European J. Combin.* **16** (1995), 503–525.
- [19] D. Walkup, The lower bound conjecture for 3- and 4-manifolds, *Acta Math.* **125** (1970), 75–107.
- [20] H. Zheng, A characterization of homology manifolds with $g_2 \leq 2$, *J. Combin. Theory Ser. A* **153** (2018), 31–45.