

A summary of several meteorological properties of the moist-air entropy variables θ_s and $PV(\theta_s)$.

by Pascal Marquet

(retired from Météo-France, CNRM/GMAP, Toulouse, France)

Version 3 of December 5, 2023.

E-mail: pascalmarquet@yahoo.com

1 Introduction - Motivations

The aim of this note is to describe several meteorological properties shown in Marquet (2022) for the moist-air specific entropy, the associated potential temperature (θ_s) defined in Marquet (2011) and the associated potential vorticity $PV(\theta_s)$ defined in Marquet (2014).

The three Tables 1 to 3 provide the numerical values of all thermodynamic constants used in this note.

Table 1: Dry- and moist-air thermodynamic constants

R_d	R_v	c_{pd}	c_{pv}	c_l	c_i	$L_v(T_0)$	$L_s(T_0)$
287.06	461.52	1004.7	1846.1	4218	2106	2501	2835
J/K/kg	J/K/kg	J/K/kg	J/K/kg	J/K/kg	J/K/kg	kJ/kg	kJ/kg

Table 2: Standard and reference values

$T_0 = T_r$	$p_0 = p_r$	$s_{d0}(T_0, p_0)$	$s_{v0}(T_0, p_0)$	$e_r = e_s(T_r)$	$p_{dr} = p_r - e_r$	$s_{dr}(T_r, p_{dr})$	$s_{vr}(T_r, e_r)$	r_r
273.15	1000	6775	10320	6.1064	993.89	6777	12673	3.82
K	hPa	J/K/kg	J/K/kg	hPa	hPa	J/K/kg	J/K/kg	g/kg

Table 3: Non-dimensional derived constants

η	ε	γ	κ	λ	δ	Λ_r
1.608	0.622	0.4594	0.2857	0.8375	0.6078	5.869 ± 0.03

The way the old dry-air potential temperature (θ) was defined in the nineteenth century is recalled in section 2. The definitions of other moist-air potential temperatures (θ , θ'_w , θ_e , θ_v , θ_l , θ_{es} , θ_{il} , θ'_e) are provided in section 3. The definition of the new moist-air entropy potential temperature (θ_s) published in Marquet (2011) is recalled in section 4, before to be compared to the other potential temperatures in section 5.

The moist-air entropy potential vorticity $PV(\theta_s)$ published in Marquet (2014) is recalled in section 6, together with the other previous definitions $PV(\psi)$ with $\psi = \theta$, θ'_w , θ_e , θ_v or θ_{es} . The section 7 shows comparisons between the potential vorticity functions $PV(\theta) \approx PV(\theta_v)$, $PV(\theta_s)$ and $PV(\theta_e)$, with in particular regions of negative values of $PV(\theta_s)$ close to the cold fronts that seems to indicate regions of symmetric instability.

General conclusions are provided in section 8.

2 The old dry-air potential temperature (θ)

von Bezold (1888b) with von Helmholtz (1888) coined the term “potential temperature” for the dry-air quantity:

$$\boxed{\theta(T, p) = T \left(\frac{p_0}{p} \right)^{R_d/c_{pd}} = T \left(\frac{p_0}{p} \right)^\kappa}, \quad (1)$$

where T is the absolute temperature, p the pressure, $p_0 = 1000$ hPa a standard value, R_d the dry-air gas constant, c_{pd} the dry-air specific heat at constant pressure and $\kappa = R_d/c_{pd}$.

The link between the dry-air potential temperature (θ) and the dry-air specific entropy (s_d) was derived by Bauer (1908, 1910), through a relationship which can be written as

$$\boxed{s_d = c_{pd} \ln(\theta)} + [s_{d0}(T_0, p_0) - c_{pd} \ln(T_0)] = c_{pd} \ln\left(\frac{\theta}{\theta_0}\right) + s_{d0}(T_0, p_0) \quad (2)$$

$$\text{and thus } \boxed{s_d = c_{pd} \ln(\theta) + s_0(p_0)}, \quad \text{where } s_0(p_0) = s_{d0}(T_0, p_0) - c_{pd} \ln(T_0). \quad (3)$$

The dry-air specific heat c_{pd} , the reference (standard) temperature $T_0 = \theta_0 = \theta(T_0, p_0)$ and the standard dry-air entropy $s_{d0}(T_0, p_0)$ are all constant terms. Moreover, the quantity $s_0(p_0)$ in (3) only depends on the reference standard pressure $p_0 = 1000$ hPa, because the changes of $s_{d0}(T_0, p_0)$ with T_0 are balanced (and cancelled) by the second term $-c_{pd} \ln(T_0)$.

3 The old moist-air potential temperatures ($\theta'_w, \theta_v, \theta_e, \theta_l, \theta_{es}, \theta_{il}, \theta'_e$)

von Bezold (1888a) coined the definition of the “pseudo-adiabatic” processes, corresponding to the wet-bulb potential temperature (θ'_w) and defined by integration of saturation moist-air differential equations, without analytic expression for

$$\boxed{\theta'_w}. \quad (4)$$

The concept of “virtual potential temperature” was coined by Guldberg and Mohn (1876, 1910), with the more general definition of it given by

$$\boxed{\theta_v = \theta \left[1 + \left(\frac{R_v}{R_d} - 1 \right) q_v - (q_l + q_i) \right]}, \quad (5)$$

where R_v is the water-vapor gas constant and q_v , q_l and q_i are the specific content for water vapor, liquid water and ice species, respectively.

The difference between the wet-bulb (θ'_w) and equivalent (θ_e) potential temperatures was studied by Normand (1921), von Stüve (1927), Robitzsch (1928), Rossby (1932), Bleeker (1939), among others, with the equivalent potential temperature defined by quantities close to the first-order approximation

$$\boxed{\theta_e \approx \theta \exp\left(\frac{L_v q_v}{c_{pd} T}\right)}, \quad (6)$$

where L_v is the latent heat of vaporization.

Hewson (1936, 1938) studied the application of the wet-bulb potential temperature (θ'_w) to air mass analysis, with its meaning and significance studied in Hewson (1946a,b).

The same differential equations derived in von Bezold (1888a) are used in Saunders (1957) to define the adiabatic and pseudo-adiabatic equations.

The three quantities θ'_w , θ_v and θ_e were considered in Lilly (1968)

As a companion of θ_e , the “liquid-water potential temperature” (θ_l) was defined and studied in Betts (1973), with the first-order approximation relationship

$$\theta_l \approx \theta \exp\left(-\frac{L_v q_l}{c_{pd} T}\right). \quad (7)$$

Similar values of θ_l and/or θ_e were derived and studied by Emanuel (1994); Pauluis et al. (2008, 2010); Laliberté et al. (2013); Pauluis and Mrowiec (2013); Pauluis (2016); Mrowiec et al. (2016); Pauluis and Zhang (2017); Fang et al. (2017); Dauhut et al. (2017); Fang et al. (2019).

Empirical formula for the saturation pseudo-adiabats and the “saturation equivalent potential temperature” (θ_{es}) were derived in Betts and Dugan (1973), leading to the first-order approximation relationship

$$\theta_{es} \approx \theta \exp\left(\frac{L_v r_{sw}}{c_{pd} T}\right), \quad (8)$$

where r_{sw} is the saturation water-vapor mixing ratio.

The “ice-liquid water potential temperature” (θ_{il}) defined in Tripoli and Cotton (1981) can be considered as a generalization of the Betts (1973) “liquid-water potential temperature” (θ_l) to the case of ice ($q_i \neq 0$), with the relationship

$$\theta_{il} \approx \theta \exp\left(-\frac{L_v q_l + L_s q_i}{c_{pd} T}\right), \quad (9)$$

where L_s is the latent heat of sublimation.

The Betts (1973) conservative quantity θ_l is nowadays used in most of parameterization of atmospheric turbulence (ex. based on: Deardorff, 1980; Mellor and Yamada, 1982; Cuxart et al., 2000, etc), and to study the subgrid-condensation schemes (ex. those based on: Deardorff and Sommeria, 1977; Bougeault, 1982, etc), with both θ_l and θ_e usually considered to study the cloud-top entrainment instability (ex. those based on: Deardorff, 1980; MacVean and Mason, 1990; Lock, 2004, etc).

The equivalent potential temperature computed in Bolton (1980), Bryan (2008) and Davies-Jones (2009), among others, are different from the “adiabatic” version θ_e approximated by Eq. (6), and rather corresponds to the “pseudo-adiabatic” value θ'_w , which is not analytically defined in Eq. (4). As an example, the (adjusted up to about 0.03 K?) approximation derived in Eqs. (21), (22), (24) and (39) of Bolton (1980) correspond to

$$T_L \approx 55 + \frac{2840}{3.5 \ln(T) - \ln(e/100) - 4.805} \approx 55 + \frac{1}{\frac{1}{T-55} - \frac{\ln(H_l)}{2840}}, \quad (10)$$

$$\theta_{DL} \approx T \left(\frac{p_0}{p-e}\right)^{0.2854} \left(\frac{T}{T_L}\right)^{0.28 r_v}, \quad (11)$$

$$\theta'_e \approx \theta_{DL} \exp\left[\left(\frac{3036}{T_L} - 1.78\right) (1 + 0.448 r_v) r_v\right], \quad (12)$$

where T is the absolute temperature (K), p the total pressure (Pa), e the water-vapor partial pressure (Pa), $H_l = e/e_{sw}$ the relative humidity (Pa/Pa), and $r_v = q_v/(1 - q_t)$ the water-vapor mixing ratio (kg/kg).

This formulation of Bolton (1980) is different from θ_e given by (6), and especially in non-saturated regions where $H_l \neq 1$ and $T_L \neq T$, since:

- $\theta_{DL} \neq \theta = T (p_0/p)^\kappa \approx T (p_0/p)^{0.2857}$, because $T_L \neq T$, $p - e \neq p$ and $0.2854 \neq 0.2857$ in (11);
- $3036/T_L - 1.78 \neq [L_v(T_0) + (c_{pv} - c_l)(T - T_0)] / (c_{pd} T) \approx 3134/T - 2.36$ in (12),
- even if $(1 + 0.448 r_v) r_v \approx [q_v]$ because $r_v = q_v / (1 - q_v) \approx q_v$ and $(1 + 0.448 r_v) \approx 1$.

4 The moist-air entropy potential temperature (θ_s)

The more recent ‘‘moist-air entropy potential temperature’’ (θ_s) published in Marquet (2011) can be considered as an improvement of the quantity θ_s already derived in Hauf and Höller (1987) and Marquet (1993).

The relationship between the moist-air specific entropy (s) and the corresponding potential temperature (θ_s) was assumed in Marquet (2011) to be the same as the one derived by Bauer (1908, 1910) for the dry-air values (s_d and θ), leading to

$$\boxed{s = c_{pd} \ln \left(\frac{\theta_s}{\theta_0} \right) + s_{d0}(T_0, p_0)} = c_{pd} \ln(\theta_s) + [s_{d0}(T_0, p_0) - c_{pd} \ln(T_0)] , \quad (13)$$

$$\text{and } \boxed{s = c_{pd} \ln(\theta_s) + s_0(p_0)} , \quad \text{where } s_0(p_0) = s_{d0}(T_0, p_0) - c_{pd} \ln(T_0) . \quad (14)$$

The specific heat capacity (at constant pressure) c_{pd} , the reference (standard) temperature $T_0 = \theta_0 = \theta(T_0, p_0)$ and the reference dry-air entropy $s_{d0}(T_0, p_0)$ are all constant. Moreover, the quantity $s_0(p_0)$ in (14) only depends on the same reference standard pressure $p_0 = 1000$ hPa as the one introduced to define the dry-air potential temperature θ in (3), because the changes of $s_{d0}(T_0, p_0)$ with T_0 are balanced (and cancelled) by the second term $-c_{pd} \ln(T_0)$.

These properties (13) and (14) show that θ_s becomes a true synonym for the specific moist-air entropy s . This property did not hold in the previous formulations of θ_s and θ^* derived in Hauf and Höller (1987) and Marquet (1993), nor in those of θ_l and θ_e derived in Betts (1973), Emanuel (1994) or Pauluis et al. (2010).

A general version of θ_s , which generalizes those of Marquet (2011, 2017), can be written from (51) as:

$$\boxed{\begin{aligned} \theta_s &= \theta \exp \left[- \frac{L_v (q_l + q_{\text{rain}}) + L_s (q_i + q_{\text{snow}})}{c_{pd} T} \right] \exp(\Lambda_r q_t) \\ &\times \left(\frac{T}{T_r} \right)^{\lambda q_t} \left(\frac{p_r}{p} \right)^{\kappa \delta q_t} \left(\frac{r_r}{r_v} \right)^{\gamma q_t} \frac{(1 + \eta r_v)^{\kappa(1 + \delta q_t)}}{(1 + \eta r_r)^{\kappa \delta q_t}} \\ &\times (H_l)^{\gamma(q_l + q_{\text{rain}})} (H_i)^{\gamma(q_i + q_{\text{snow}})} \left(\frac{T_{\text{rain}}}{T} \right)^{c_l q_{\text{rain}}/c_{pd}} \left(\frac{T_{\text{snow}}}{T} \right)^{c_i q_{\text{snow}}/c_{pd}} \end{aligned}} \quad (15)$$

where $\theta = T (p_0/p)^{R_d/c_{pd}}$ is the dry-air potential temperature. The relative humidities are noted $H_l = e/e_{sw}$ and $H_i = e/e_{si}$. The temperatures of precipitations (rain and snow) are noted T_{rain} and T_{snow} . See the Tables 1 to 3 for the other thermodynamic constants.

Differently, the potential temperature $\theta_{s/\text{HH87}}$ defined in Hauf and Höller (1987, see Eqs. 3.23

to 3.25 page 2893) was linked to the specific entropy through the relationship

$$s = q_d c_{pe}^* \ln\left(\frac{\theta_{s/\text{HH87}}}{T_0}\right) + q_d s^*, \quad (16)$$

$$\text{with } \theta_{s/\text{HH87}} = T \left(\frac{p-e}{p_0}\right)^{-R_d/c_{pe}^*} \exp\left(\frac{L_v r_v - L_s r_i}{c_{pe}^* T}\right) (H_l)^{-R_v r_v/c_{pe}^*} (H_i)^{R_v r_i/c_{pe}^*}. \quad (17)$$

Hauf and Höller (1987, p.2890) explained that, in (16) and (17): “ T_0 and p_0 denote reference values of T and p , and $s_d(T_0, p_0)$, $s_v(T_0, p_0)$, $s_l(T_0)$ and $s_i(T_0)$ are the values of dry-air and water entropies at this reference point.” This means that we must consider that $T_r = T_0$ and $p_r = p_0$ in Hauf and Höller (1987), without justification.

In (16) both $q_d c_{pe}^* = (1 - q_t) c_{pd} + q_t c_l$ and $q_d s^* = [(1 - q_t) s_d(T_0, p_0) + q_t s_l(T_0)]$ depend on q_t . Therefore, the first drawback is that $\theta_{s/\text{HH87}}$ cannot represent the moist-air entropy (s) unless the total-water specific content $q_t = r_t/(1 + r_t)$ is a constant. The words of Hauf and Höller (1987, p.2893) were: “*The entropy temperature $\theta_{s/\text{HH87}}$ is related to entropy (s) by the functional relationship (16), and thus can be considered as a measure of entropy. However, Eq. (16) involved, beside entropy, the concentration of dry air $q_d = 1 - q_t$ and the total water mixing ratio $r_t = q_t/(1 - q_t)$. Therefore, a constant entropy does not generally imply that the entropy temperature $\theta_{s/\text{HH87}}$ is also a constant.*”

The other drawback is the term $(H_l)^{-R_v r_v/c_{pe}^*}$ in (17) that is different from 1 in unsaturated regions, where the relative humidity $H_l = e/e_{sw} < 1$ but $r_v > 0$ is different from 0. The last term $(H_i)^{R_v r_i/c_{pe}^*}$ is most often equal to 1 for both unsaturated ($r_i = 0$) and saturated regions ($H_i = 1$). This issue (see Hauf and Höller, 1987, p.2893) is due to “*the definition of entropy temperature which presumes a system of cloudy air and especially the existence of liquid water, as can be seen from $q_d c_{pe}^* = (1 - q_t) c_{pd} + q_t c_l$ and $q_d s^* = [(1 - q_t) s_d(T_0, p_0) + q_t s_l(T_0)]$,*” which both depend on c_l and $s_l(T_0)$. “*Problems arise if no liquid water is present, either locally or in principle.*”

The potential temperature θ_{M93}^* defined in Marquet (1993, before Eq.42 and Eq.43) partially solved the issues of Hauf and Höller (1987), with θ_{M93}^* linked to the specific entropy through the relationship

$$s = q_d c_{pl}^* \ln\left(\frac{\theta_{\text{M93}}^*}{\theta_0^*}\right) + [(1 - q_t) s_d(T_0, p_0 - e_0) + q_t s_v(T_0, e_0)], \quad (18)$$

with $e_0 = e_{\text{sat}}(T_0)$ and

$$\theta_{\text{M93}}^* = T \left(\frac{p-e}{p_0}\right)^{-R_d/c_{pl}^*} \exp\left(-\frac{L_v r_l + L_s r_i}{c_{pl}^* T}\right) \left(\frac{e}{p_0}\right)^{-R_v r_t/c_{pl}^*} (H_l)^{R_v r_l/c_{pl}^*} (H_i)^{R_v r_i/c_{pl}^*}. \quad (19)$$

Like in (16), both $q_d c_{pl}^* = (1 - q_t) c_{pd} + q_t c_{pv}$ and the bracketed terms (with the reference entropies) depend on q_t in (18), and therefore θ_s^* cannot always represent the moist-air entropy (s) in (18) unless the total-water specific content $q_t = r_t/(1 + r_t)$ is a constant.

However, the first advantage is the term c_l used in Hauf and Höller (1987) that is replaced by c_{pv} in θ_{M93}^* , with the more comfortable idea that there is always some water-vapor content in real parcels of moist air, whereas there was no guaranty of existing liquid water in the formulation of Hauf and Höller (1987).

The other advantage is the symmetric use of the two mixing ratios (r_l, r_i) in the exponential, instead of the use of (r_v, r_i) with two different signs in Hauf and Höller (1987), with also the possibility to neglect the last two terms $(H_l)^{R_v r_l/c_p^*}$ and $(H_i)^{R_v r_i/c_p^*}$ in (19) for both unsaturated (since $r_l = r_i = 0$) and saturated regions (if $H_l = H_i = 1$).

In fact, the formulation (17) of Hauf and Höller (1987) corresponds to the equivalent potential temperature derived in Emanuel (1994) and Pauluis et al. (2010), but with the ice content included. Therefore: $\theta_{s/\text{HH87}} \approx \theta_{e/\text{E94}} \approx \theta_{e/\text{P10}}$, but with no citation to the JAS paper of Hauf and Höller (1987) by K. Emanuel nor by O. Pauluis.

The words of Hauf and Höller (1987, p.2895) were: “*The equivalent potential temperature is a constant in a closed adiabatic system where water vapor is saturated. If in a rising parcel of cloudy air the liquid water remains in the parcel and does not fall out, then θ_e is a constant.*” And then (Hauf and Höller, 1987, p.2896): “*As a result, we see that for a saturated warm cloud the entropy temperature*” ($\theta_{s/\text{HH87}}$) “*is identical with the equivalent potential temperature θ_e* ” (i.e. this equivalence is valid only if q_t is a constant).

Moreover $\theta_{s/\text{HH87}} = \theta_{e/\text{HH87}}$ can be approximated by θ_e of Betts (1973) given by (6), which can be considered as a first-order approximation of the more general formulations of Hauf and Höller (1987), Emanuel (1994) and Pauluis et al. (2010). This is the reason why the “first-order approximation” θ_e of Betts (1973) given by (6) is used in almost all practical applications to study convection and is considered as a “conservative variable” in almost all moist-air turbulence schemes. The words of Hauf and Höller (1987, p.2895) were: “*other definitions of equivalent potential temperature exist (e.g. Betts, 1973), which arise from approximations of the definition of*” $\theta_{s/\text{HH87}}$. “*The present one*” (i.e. $\theta_{s/\text{HH87}} \approx \theta_{e/\text{HH87}}$ for constant q_t) “*is the most general definition, as no further approximations or assumptions have been made.*”

The main result is that the equivalent potential temperatures θ_e derived in Betts (1973), Hauf and Höller (1987), Emanuel (1994) and Pauluis et al. (2010) cannot represent the moist-air entropy (s) unless the total-water specific content $q_t = r_t/(1 + r_t)$ and the mixing ratio $r_t = q_t/(1 - q_t)$ are constant. Only θ_s given by (13) is really synonymous with the specific moist-air entropy, due to the relationship (15) where both c_{pd} and $s_{d0}(T_0, p_0)$ are constant.

The same is true for the liquid-water potential entropy θ_l that can be written, following the same method described in Hauf and Höller (1987), as

$$s = q_d c_{pl}^* \ln\left(\frac{\theta_l}{T_0}\right) + [(1 - q_t) s_d(T_0, p_0) + q_t s_v(T_0, e_0)] , \quad (20)$$

where

$$\theta_l = T \left(\frac{p - e}{p_0}\right)^{-R_d/c_{pl}^*} \exp\left(-\frac{L_v r_l + L_s r_i}{c_{pl}^* T}\right) \left(\frac{e}{e_0}\right)^{-r_t R_v/c_{pl}^*} (H_l)^{R_v r_l/c_{pl}^*} (H_i)^{R_v r_i/c_{pl}^*} . \quad (21)$$

Both $(1 - q_t) c_{pl}^* = (1 - q_t) c_{pd} + q_t c_{pv}$ and the bracketed terms of (20) depend on q_t , and both prevent θ_l to represent the moist-air entropy in all conditions if q_t and r_t are not constant.

The last terms of (20) $(H_l)^{R_v r_l/c_{pl}^*}$ and $(H_i)^{R_v r_i/c_{pl}^*}$ are most often equal to 1 for both unsaturated (because $r_l = r_i = 0$) and saturated regions (if either $H_l = 1$ or $H_i = 1$).

However, the term $(e/e_0)^{-r_t R_v/c_{pl}^*}$ may be significantly different from unity and was missing in the formulation of Betts (1973).

The only difference between θ_l defined by (21) and θ_{M93}^* defined by (19) is the term $(e/e_0)^{-r_t R_v/c_{pl}^*}$ in (21) replaced by $(e/p_0)^{-r_t R_v/c_{pl}^*}$ in (19), with a ratio of these terms $(e_0/p_0)^{-r_t R_v/c_{pl}^*}$ that is constant if r_t is a constant. However, e/e_0 with $e_0 = 6$ hPa is likely more often closer to unity than e/p_0 with $p_0 = 1000$ hPa, and for this reason (21) is likely more relevant than (19).

4.1 Establishment/demonstration of the formulation for θ_s

The formulation (15) can be derived starting from the Appendix B of Marquet (2011), but:

- with a more direct method;
- with all computations shown;
- with the additional hypotheses of T_l and T_i different from T for condensed species (precipitations at the wet-bulb temperature?); and
- with the possibility that $H_l \neq 1$ and $H_i \neq 1$ where $q_l \neq 0$ and $q_i \neq 0$ (unsaturated conditions with condensed species), or the possibility that $H_l > 1$ or $H_i > 1$ (surer-saturation or mixed-phase conditions).

The specific moist-air entropy is first assumed to be a weighted sum of the partial entropies for dry air $s_d(T, p - e)$, water vapor $s_v(T, e)$, liquid water $s_l(T)$, ice $s_i(T)$, rain $s_{\text{rain}}(T_{\text{rain}})$ and snow $s_{\text{snow}}(T_{\text{snow}})$ species, with $q_d + q_v + q_l + q_i + q_{\text{rain}} + q_{\text{snow}} = q_d + q_t = 1$ the specific contents of these species (i.e. the mass concentrations $q_x = m_x/m$ with $m = m_d + m_v + m_l + m_i + m_{\text{rain}} + m_{\text{snow}} = m_d + m_t$ the total mass of a small and homogeneous parcel of the moist-air fluid), leading to

$$s = q_d s_d(T, p - e) + q_v s_v(T, e) + q_l s_l(T) + q_i s_i(T) + q_{\text{rain}} s_l(T_{\text{rain}}) + q_{\text{snow}} s_i(T_{\text{snow}}). \quad (22)$$

The moist-air entropy formulation (22) depends on the partial pressure of dry air ($p - e$) and water vapor (e). This formulation (22) can then be transformed by using $q_v = q_t - q_l - q_i - q_{\text{rain}} - q_{\text{snow}}$ in the first line, and by inserting/removing $s_l(T)$ and $s_i(T)$ in the second line, leading to

$$s = q_d s_d(T, p - e) + q_t s_v(T, e) + q_l [s_l(T) - s_v(T, e)] + q_i [s_i(T) - s_v(T, e)] + q_{\text{rain}} [s_l(T_{\text{rain}}) - s_l(T)] + q_{\text{rain}} [s_l(T) - s_v(T, e)] + q_{\text{snow}} [s_i(T_{\text{snow}}) - s_i(T)] + q_{\text{snow}} [s_i(T) - s_v(T, e)], \quad (23)$$

$$s = q_d s_d(T, p - e) + q_t s_v(T, e) + q_{\text{rain}} [s_l(T_{\text{rain}}) - s_l(T)] - (q_l + q_{\text{rain}}) [s_v(T, e) - s_l(T)] + q_{\text{snow}} [s_i(T_{\text{snow}}) - s_i(T)] - (q_i + q_{\text{snow}}) [s_v(T, e) - s_i(T)]. \quad (24)$$

If the “gas constants” (R_d and R_v) and the “specific heat at constant pressure” (c_{pd} , c_{pv} , c_l , c_i) are assumed to be constant within the range of atmospheric temperatures, the partial entropies can be computed from the relationships

$$s_d(T, p - e) = c_{pd} \ln\left(\frac{T}{T_r}\right) - R_d \ln\left(\frac{p - e}{p_r - e_r}\right) + s_{dr}(T_r, p_r - e_r), \quad (25)$$

$$s_v(T, e) = c_{pv} \ln\left(\frac{T}{T_r}\right) - R_v \ln\left(\frac{e}{e_r}\right) + s_{vr}(T_r, e_r), \quad (26)$$

$$s_l(T) = c_l \ln\left(\frac{T}{T_r}\right) + s_{lr}(T_r), \quad (27)$$

$$s_i(T) = c_i \ln\left(\frac{T}{T_r}\right) + s_{ir}(T_r), \quad (28)$$

where, at this stage, the reference temperature $T_r \neq T_0$ and the reference total pressure $p_r \neq p_0$ are, a priori, different from the standard values $T_0 = 273.15$ K and $p_0 = 1000$ hPa. The reference partial pressure for water vapor is assumed to be the saturation water-vapor pressure $e_r = e_s(T_r)$, which is the only possible value different from 0 and depending on the reference temperature T_r .

The differences in entropy $s_l(T_{\text{rain}}) - s_l(T)$ and $s_i(T_{\text{snow}}) - s_i(T)$ in (23) can be computed from (29) and (30), yielding

$$s_l(T_{\text{rain}}) - s_l(T) = c_l \ln\left(\frac{T_{\text{rain}}}{T}\right) \quad (29)$$

$$\text{and } s_i(T_{\text{snow}}) - s_i(T) = c_i \ln\left(\frac{T_{\text{snow}}}{T}\right), \quad (30)$$

whereas the other differences in entropy can be computed in terms of the latent heats $L_v(T)$ and $L_s(T)$ and the relative humidities $H_l = e/e_{sw}$ and $H_i = e/e_{si}$ according to Hauf and Höller (1987, Eqs. 3.10 to 3.12, p. 2891), yielding

$$s_v(T, e) - s_l(T) = \frac{L_v(T)}{T} - R_v \ln(H_l), \quad (31)$$

$$s_v(T, e) - s_i(T) = \frac{L_s(T)}{T} - R_v \ln(H_i). \quad (32)$$

These differences in water-species entropy (29)-(32) can be inserted into (24), together with $s_d(T, p - e)$ and $s_v(T, e)$ given by (25)-(26), to give

$$\begin{aligned} s = & q_d s_{dr}(T_r, p_r - e_r) + q_t s_{vr}(T_r, e_r) - \frac{(q_l + q_{\text{rain}}) L_v(T) + (q_i + q_{\text{snow}}) L_s(T)}{T} \\ & + (q_d c_{pd} + q_t c_{pv}) \ln\left(\frac{T}{T_r}\right) - q_d R_d \ln\left(\frac{p - e}{p_r - e_r}\right) - q_t R_v \ln\left(\frac{e}{e_r}\right) \\ & + q_{\text{rain}} c_l \ln\left(\frac{T_{\text{rain}}}{T}\right) + q_{\text{snow}} c_i \ln\left(\frac{T_{\text{snow}}}{T}\right) + (q_l + q_{\text{rain}}) R_v \ln(H_l) + (q_i + q_{\text{snow}}) R_v \ln(H_i). \end{aligned} \quad (33)$$

The property $q_d = 1 - q_t$ can be used to rewrite the first two terms of (33) as

$$q_d s_{dr}(T_r, p_r - e_r) + q_t s_{vr}(T_r, e_r) = s_{dr}(T_r, p_r - e_r) + c_{pd} \Lambda_r q_t, \quad (34)$$

where the non-dimensional term $\Lambda_r = [s_{vr}(T_r, e_r) - s_{dr}(T_r, p_r - e_r)]/c_{pd}$ depends on the reference entropies for water vapor and dry air, to be computed for the reference temperature T_r and partial pressures p_r and e_r . Similarly, the term in factor of the logarithm of T can be rewritten as

$$q_d c_{pd} + q_t c_{pv} = c_{pd} + c_{pd} \lambda q_t, \quad \text{where } \lambda = (c_{pv} - c_{pd})/c_{pd}. \quad (35)$$

The dry-air and water species reference entropies can be computed for any set of reference values (T_r, p_r, e_r) according to

$$s_{dr}(T_r, p_r - e_r) = s_{d0}(T_0, p_0) + c_{pd} \ln\left(\frac{T_r}{T_0}\right) - R_d \ln\left(\frac{p_r - e_r}{p_0}\right), \quad (36)$$

$$s_{vr}(T_r, e_r) = s_{v0}(T_0, p_0) + c_{pv} \ln\left(\frac{T_r}{T_0}\right) - R_v \ln\left(\frac{e_r}{p_0}\right), \quad (37)$$

$$s_{lr}(T_r) = s_{l0}(T_0) + c_l \ln\left(\frac{T_r}{T_0}\right), \quad (38)$$

$$s_{ir}(T_r) = s_{i0}(T_0) + c_i \ln\left(\frac{T_r}{T_0}\right). \quad (39)$$

Eqs. (34), (35) and (36) can be inserted into (33), also with the term “ $R_d \ln(p_0/p)$ ” added and removed in the first line, to give¹

$$\begin{aligned}
s = & s_{d0}(T_0, p_0) + c_{pd} \ln\left(\frac{T}{T_0}\right) + R_d \ln\left(\frac{p_0}{p}\right) - \frac{(q_l + q_{\text{rain}}) L_v(T) + (q_i + q_{\text{snow}}) L_s(T)}{T} \\
& + c_{pd} \Lambda_r q_t + c_{pd} \lambda q_t \ln\left(\frac{T}{T_r}\right) - R_d \ln\left(\frac{p_r - e_r}{p}\right) - q_d R_d \ln\left(\frac{p - e}{p_r - e_r}\right) - q_t R_v \ln\left(\frac{e}{e_r}\right) \\
& + q_{\text{rain}} c_l \ln\left(\frac{T_{\text{rain}}}{T}\right) + q_{\text{snow}} c_i \ln\left(\frac{T_{\text{snow}}}{T}\right) + (q_l + q_{\text{rain}}) R_v \ln(H_l) + (q_i + q_{\text{snow}}) R_v \ln(H_i). \quad (40)
\end{aligned}$$

An important property can be deduced from (40): because the first and third lines do not depend on the reference values T_r , p_r and $e_r(T_r)$, the second line cannot depend on, and is thus independent on, these reference value. This means that any changes in T_r , p_r and $e_r(T_r)$ corresponds to changes in all terms that leave the whole second line unchanged.

The next step is to write the equations for the dry air, water vapor and moist air

$$p_d = p - e = \rho_d R_d T = \rho q_d R_d T, \quad (41)$$

$$e = \rho_v R_v T = \rho q_v R_v T, \quad (42)$$

$$p = p_d + e_r = \rho R T = \rho (q_d R_d + q_v R_v) T = \rho q_d R_d (1 + \eta r_v) T, \quad (43)$$

$$\text{and thus with } p = (1 + \eta r_v) (p - e), \quad (44)$$

$$\text{and } e = \rho q_d R_d \eta r_v T = \eta r_v (p - e), \quad (45)$$

where $r_v = q_v/q_d = \rho_v/\rho_d$ and $\eta = R_v/R_d$, leading to

$$p - e = \frac{1}{1 + \eta r_v} p, \quad (46)$$

$$e = \frac{\eta r_v}{1 + \eta r_v} p. \quad (47)$$

The same relationship exist for the “just saturated” (i.e. with $q_{lr} = q_{ir} = 0$ and thus $q_{tr} = q_{vr} = q_{\text{sat}}(T_r, p_r) = q_r$) reference state

$$p_r - e_r = \frac{1}{1 + \eta r_r} p_r, \quad (48)$$

$$e_r = \frac{\eta r_r}{1 + \eta r_r} p_r = \eta r_r (p_r - e_r), \quad (49)$$

$$\text{and thus with } r_r = \frac{e_r}{\eta (p_r - e_r)} \text{ and } q_r = \frac{r_r}{1 + r_r}. \quad (50)$$

The partial pressures relationships (46), (47), (48) and (49), can then be inserted into (40) with

¹Many thanks to Almut Gassman who see a typo in the version 1 of the arXiv note for the term $-q_t R_v \ln(e/e_r)$ at the end of the second line of (40), which was badly written $c_{pd} \gamma q_t \ln(r_r/r_v)$, in that wrongly anticipating the end of the third line of (51) and via a bad LaTeX copy/paste... sorry!

$\eta = R_v/R_d$, $\gamma = R_v/c_{pd}$, $\kappa = R_d/c_{pd}$ and $\delta = R_v/R_d - 1$, leading to:

$$\begin{aligned}
s &= s_{d0}(T_0, p_0) - c_{pd} \ln(T_0) + c_{pd} \ln(T) + c_{pd} \kappa \ln\left(\frac{p_0}{p}\right) \\
&\quad - c_{pd} \left[\frac{(q_l + q_{\text{rain}}) L_v(T) + (q_i + q_{\text{snow}}) L_s(T)}{c_{pd} T} \right] + c_{pd} \Lambda_r q_t \\
&\quad + c_{pd} \lambda q_t \ln\left(\frac{T}{T_r}\right) + c_{pd} \kappa \delta q_t \ln\left(\frac{p_r}{p}\right) + c_{pd} \gamma q_t \ln\left(\frac{r_r}{r_v}\right) \\
&\quad - c_{pd} \kappa \delta q_t \ln(1 + \eta r_r) + c_{pd} \kappa (1 + \delta q_t) \ln(1 + \eta r_v) \\
&\quad + q_{\text{rain}} c_l \ln\left(\frac{T_{\text{rain}}}{T}\right) + q_{\text{snow}} c_i \ln\left(\frac{T_{\text{snow}}}{T}\right) \\
&\quad + c_{pd} (q_l + q_{\text{rain}}) \gamma \ln(H_l) + c_{pd} (q_i + q_{\text{snow}}) \gamma \ln(H_i). \tag{51}
\end{aligned}$$

The relationship (15) for θ_s results from the definition (13) $s = s_{d0}(T_0, p_0) + c_{pd} \ln(\theta_s/T_0) = s_{d0}(T_0, p_0) - c_{pd} \ln(T_0) + c_{pd} \ln(\theta_s)$, namely with the same first two terms $s_{d0}(T_0, p_0)$ and $-c_{pd} \ln(T_0)$ as in (51), and then via an identification of θ_s with all other terms of (51) expressed as $c_{pd} \ln[(\dots)^{(\dots)}]$ or $c_{pd} \ln[\exp(\dots)]$.

The absolute (third-law) standard entropies $s_{d0}(T_0, p_0)$, $s_{v0}(T_0, p_0)$, $s_{l0}(T_0)$ and $s_{i0}(T_0)$ appearing in (36)-(39) are computed at the standard values $T_0 = 273.15$ K and $p_0 = 1000$ hPa.

The same standard values considered in Marquet (2011) were first published in atmospheric science by Hauf and Höller (1987, “HH87”, p.2891, however without references), who explained that: “*the values of the zero-entropies have to be determined experimentally or by quantum statistical considerations,*” leading to

$$\begin{array}{cccccc}
& \text{HH87} & \text{M83} & \text{L00} & \text{MS22} & \text{M22} \\
s_{d0}(T_0, p_0) \approx & 6775 & [---] & [6776] & [6776.2] & [6776.3] \text{ J/K/kg}
\end{array} \tag{52}$$

$$\begin{array}{cccccc}
s_{v0}(T_0, p_0) \approx & 10320 & [---] & [---] & [10319.7] & [10318.1] \text{ J/K/kg}
\end{array} \tag{53}$$

$$\begin{array}{cccccc}
s_{l0}(T_0) \approx & 3517 & [3516.2] & [---] & [3516.7] & [3514.7] \text{ J/K/kg}
\end{array} \tag{54}$$

$$\begin{array}{cccccc}
s_{i0}(T_0) \approx & 2296 & [---] & [---] & [---] & [2291.9] \text{ J/K/kg}
\end{array} \tag{55}$$

The other statistical-physics (standard and absolute) values indicated into brackets in (52)-(55) are from

- Millero (1983, “M83”) for liquid-water only (s_{l0}) and from the value 70.00 J/K/mol at 298.15 K of Lewis and Randall (1961);
- Lemmon et al. (2000, “L00”) for dry air only (s_{d0}) and for dry-air composed of $\text{N}_2 + \text{O}_2 + \text{Ar}$ (without CO_2);
- Marquet and Stevens (2022, “MS22”) without the ice value (s_{i0});
- from recent full computations made in Marquet (2022, “M22”, to appear on arXiv) based on the NIST-JANAF4 equations of Chase (1998) for H_2O (vapour+liquid+ice) and for dry-air composed of $\text{N}_2 + \text{O}_2 + \text{Ar} + 400$ ppm of CO_2 .

The numerical values listed in (52)-(55) show that the uncertainties on dry-air and water-vapour standard entropies s_{d0} and s_{v0} are small: about 1 and 2 J/K/kg, respectively.

It is worthwhile to remember that the standard entropies computed at 298.15 K and 1000 hPa (or 1013.25 hPa) for N_2 , O_2 , Ar , CO_2 and H_2O were already fully available since Kelley (1932), and

then in the Thermodynamic Tables of Rossini et al. (1952), Lewis and Randall (1961), Wagman et al. (1965) and Robie et al. (1978), thus long before the computations made in atmospheric science by Betts (1973), Iribarne and Godson (1973, 1981), Emanuel (1994), Pauluis et al. (2010) and others, with (very) little changes in comparison to the more modern computations available in the Thermodynamic Tables and book of Gokcen and Reddy (1996), Chase (1998) and Atkins and de Paula (2014).

Figure 1 summarizes the way all absolute entropies can be computed with either the calorimetric or statistical-physics methods, leading to the same values at 273.15 K. The third law must be applied to the “more stable crystalline states” at 0 K (and not to the gases as wrongly stated in the Appendix of Pauluis et al., 2010, where criticisms of the third law of thermodynamics were unfounded). It is also needed to take into account the special residual entropy of 189 J/K/kg for ice-Ih, according to Pauling (1935) and Nagle (1966).

Similarly, IAPWS and TEOS10 computations of the entropies of ocean and moist-air (Wagner and Pruss, 2002; Feistel et al., 2008; IAPWS, 2010; Feistel et al., 2010; Feistel, 2012, 2019; Feistel and Hellmuth, 2020) may have use the values of Millero (1983) for sea water and Lemmon et al. (2000) for dry air, which are moreover available (but commented and unused) in the TEOS10 FORTRAN software. Differently, IAPWS and TEOS10 computations are made by canceling some standard entropies of dry air and water species at 273.15 K, in that being in contradiction with the third law of thermodynamics and with statistical physics principles, like in Iribarne and Godson (1973, 1981), Emanuel (1994), Pauluis et al. (2010) and others.

It was similarly assumed by Emanuel (1994), Pauluis et al. (2010) and many others that $s_d(T_0, p_0)$ and either $s_v(T_0, p_0)$ for θ_l or $s_l(T_0)$ for θ_e can be freely set to zero, in clear disagreement with the third law of thermodynamics and differently from Hauf and Höller (1987), Millero (1983), Lemmon et al. (2000) and Marquet (2011, 2017), where the third law is fully taken into account for the solid states of all dry-air and water species (N_2 , O_2 , Ar, CO_2 , H_2O).

Moreover, in fact, there is no need to set to zero these standard values $s_d(T_0, p_0)$, $s_v(T_0, p_0)$ or $s_l(T_0)$ to defined θ_e and θ_l from the formulation $\theta_{s/HH87}$ of Hauf and Höller (1987), and the same should have been for Emanuel (1994), Pauluis et al. (2010) and others.

The only (however big) price to pay for leaving terms depending on the variable water content q_t outside the logarithm in (16) and (18), including the terms depending on the non-zero entropies $s_d(T_0, p_0)$, $s_v(T_0, p_0)$ and $s_l(T_0)$, is that none of θ_e , θ_l , $\theta_{s/HH87}$ and $\theta_{s/M93}$ can represent the moist-air entropy in the real world, where q_t is neither a constant due to evaporation, precipitations, entrainment, detrainment and turbulent-mixing processes.

Anyhow, since the entropy is a state function, there is only one definition of it, and only θ_s can represent the moist-air entropy in all conditions. In order to understand more precisely this property, let us imagine two parcels of the moist-air fluid labeled by A and B , with the thermodynamic conditions $(T_A, p_A, q_{vA}, q_{lA}, q_{iA}, \dots)$ and $(T_B, p_B, q_{vB}, q_{lB}, q_{iB}, \dots)$, respectively. The total water contents $q_{tA} = q_{vA} + q_{iA} + \dots$ and $q_{tB} = q_{vB} + q_{iB} + \dots$ are thus different. In that case, according to (16), (18) or (20), the difference in specific entropy $\Delta s = s_B - s_A$ between the two points A and B depends on differences like $[s_l(T_0) - s_d(T_0, p_0)] \Delta q_t$ for (16), where $\Delta q_t = q_{tB} - q_{tA}$.




Figure 1: Entropies for dry-air (N_2 , O_2 , Ar, CO_2 and water H_2O) species plotted against the absolute temperature and computed at 1000 hPa. The calorimetric method $\left(\int_0^T c_p(T') d\ln(T') + \sum_j L(T_j)/T_j\right)$ corresponds to the coloured solid lines. The third-law hypothesis is applied at 0 K with zero entropies for all the solid phases, but with the residual entropy of 189 J/K/kg for ice-Ih. The vertical jumps correspond to phase changes at T_j with the phase-change enthalpies $L(T_j)$ between solids phases (for N_2 and O_2), then from solid to liquid phases, then from liquid to vapour phases. The statistical-physics values (black dashed lines) are computed from $S = k \ln(W)$ and $F = -k T \ln(Z)$ for the vapour phases according to the method described in Chase (1998) for translational, rotational, vibrational and electronic partition functions (Z).

Therefore, since the specific entropy “ s ” is a state function, then the difference between two points $\Delta s = s_B - s_A$ must have a unique (non-arbitrary) value independent on any thermodynamic path leading from A to B , and then it is not possible to modify at will and arbitrarily the difference $[s_l(T_0) - s_d(T_0, p_0)]$, because it is in factor of the variable term Δq_t and any arbitrarily change in $s_l(T_0)$ or $s_d(T_0, p_0)$ would generate arbitrary changes in Δs associated with Δq_t . This would be in contradiction with the second law of thermodynamics and the need to know precisely if the (moist-air) entropy decreases, increases or remain constant. It is therefore needed to rely on the third law of thermodynamics (applied at 0 K for the more stable crystalline states) and the absolute definitions of the entropies for dry-air and water species (N_2 , O_2 , Ar , CO_2 , H_2O).

Another way to avoid arbitrarily canceling the reference entropies is to keep the terms depending on q_t and the reference entropies in (16), (18) or (20) as they are, in order to define θ_l and θ_e without trying to make them synonymous with the specific entropy s .

This approach was chosen in Marquet and Stevens (2022), where θ_l , θ_s and θ_e are defined (for ice-free conditions $q_i = q_{snow} = 0$) in a similar way without canceling any of the reference partial entropies. The new method is the one previously described in Stevens and Siebesma (2020, from a course with the same name given at MPI) for deriving both θ_l and θ_e , but generalizes in Marquet and Stevens (2022) to the definition of θ_s .

Stevens and Siebesma (2020, after Eq. 2.30) reinterpret entropy equations like (16) in the following way: “*Given a completely specified reference state it provides an expression for the entropy. This was the sense in which it was derived. Alternatively, one can use this equation to ask what would the reference state temperature need to be, for the system in the reference state configuration (as specified through the pressure, amount and distribution of water mass) to have the same entropy as in the given state.*”

Accordingly, the three equations

$$s(T, p, q_v, q_l) = c_{pe} \ln \left(\frac{\theta_e}{T_r} \right) + [(1 - q_t) s_d(T_r, p_0) + q_t s_l(T_r)] , \quad (56)$$

$$s(T, p, q_v, q_l) = c_{pl} \ln \left(\frac{\theta_\ell}{T_r} \right) + [(1 - q_t) s_d(T_r, p_0 - e_0) + q_t s_v(T_r, e_0)] , \quad (57)$$

$$\text{and } s(T, p, q_v, q_l) = c_{pd} \ln \left(\frac{\theta_s}{T_r} \right) + [s_d(T_r, p_0)] = c_{pd} \ln \left(\frac{\theta_s}{T_0} \right) + [s_d(T_0, p_0)] \quad (58)$$

provide the new definitions:

θ_e , θ_ℓ and θ_s are the reference temperature T_r for which the actual entropy $s(T, p, q_v, q_l)$ on the left-hand-sides of (56)-(58) is equal to the reference entropies into brackets on the right-hand-sides of (56)-(58).

These definitions avoid canceling any of the reference partial entropies $s_d(T_r, p_0)$, $s_d(T_r, p_0 - e_0)$, $s_v(T_r, e_0)$ or $s_l(T_r)$ for arbitrary values of T_r , as done in Iribarne and Godson (1973, 1981), Emanuel (1994), Pauluis et al. (2010) and others.

The peculiar feature valid for θ_s only is that the second part of (58) is also valid for T_r replaced by the standard value $T_0 = 273.15$ K. This is simply due to the dry-air equation (36) and the corresponding property $s_d(T_0, p_0) - c_{pd} \ln(T_0) = s_d(T_r, p_0) - c_{pd} \ln(T_r)$, whatever T_r may be. This is precisely why θ_s given by (15) is independent on any value of T_r .

The method of Stevens and Siebesma (2020) corresponding to (56)-(58) is used in Marquet and Stevens (2022) to define and compute the three potential temperatures, leading to

$$\theta_e = T \left(\frac{p_0}{p} \right)^{\kappa_e} \exp\left(\frac{q_v L_v}{c_{pe} T} \right) \Omega_e, \text{ where } \Omega_e = \left(\frac{R}{R_e} \right)^{\kappa_e} (H_l)^{-q_v R_v / c_{pe}}, \quad (59)$$

$$\theta_\ell = T \left(\frac{p_0}{p} \right)^{\kappa_\ell} \exp\left(-\frac{q_l L_v}{c_{pl} T} \right) \Omega_\ell, \text{ where } \Omega_\ell = \left(\frac{R}{R_\ell} \right)^{\kappa_\ell} \left(\frac{q_t}{q_v} \right)^{q_t R_v / c_{pl}}, \quad (60)$$

$$\theta_s = \theta \exp\left(-\frac{q_l L_v}{c_{pd} T} + \Lambda_r q_t \right) \Omega_s, \text{ where } \Omega_s = \left(\frac{R}{R_e} \right)^{\kappa_d} \left(\frac{p - e}{p_0 - e_0} \right)^{q_t \kappa_d} \left(\frac{e_0}{e} \right)^{q_t \gamma} \left(\frac{T}{T_0} \right)^{q_t \lambda} \quad (61)$$

and with $\theta = T (p_0/p)^{\kappa_d}$ the usual dry-air potential temperature.

The moist-air exponent terms appearing in (59)-(61) are $R = R_d(1 - q_t) + R_v q_v$ (i.e; the moist-air gas constant), $R_e = R_d(1 - q_t)$, $c_{pe} = c_{pd}(1 - q_t) + c_l q_t$, $\kappa_e = R_e/c_{pe}$, $R_\ell = R_d(1 - q_t) + R_v q_t$, $c_{pl} = c_{pd}(1 - q_t) + c_{pv} q_t$ and $\kappa_\ell = R_\ell/c_{pl}$.

The hope is that the Omega terms (Ω_e , Ω_ℓ , Ω_s) in (59)-(61) may remain close to unity in all atmospheric conditions, with therefore the “first-order approximations” of θ_e , θ_ℓ and θ_s given by the other terms depending on T , (p_0/p) and the exponentials. In particular, we can recognize for θ_s in (61) the first line of (15) with $q_i = q_{snow} = 0$, this first line being interpreted as the “first-order approximation of θ_s ” hereafter.

Indeed, the vertical profiles of the FIRE-I (RF03B) observed radial flight studied in Marquet (2011) plotted between 0 and 1.5 km are shown in the Fig. (8) below, together with the vertical profiles plotted between 0 and 4 km in the Fig 1 of Marquet and Stevens (2022), which are the average for 757 drop-sounds observed during the EUREC⁴A-Circle. It is possible to check with these vertical profiles that the dashed or dotted lines (first-order approximations) remain close to the solid lines (exact formulations), up to about 0.6 K for θ_e and θ_s and better than 0.1 K for θ_ℓ . Moreover, these differences are by far smaller than the differences between θ_ℓ , θ_s and θ_e , this comforting the status of “first-order approximations”.

A rigorous method is described in Marquet (2015, 2019a) to derive not only the first-order approximation $(\theta_s)_1$, but also a second-order approximation $(\theta_s)_2$

The relationship already suggested in Marquet (2011) is indeed a first-order approximation: it corresponds to the first line of (15)

$$\theta_s \approx \left[(\theta_s)_1 = \theta \exp \left[-\frac{L_v (q_l + q_{rain}) + L_s (q_i + q_{snow})}{c_{pd} T} \right] \exp(\Lambda_r q_t) \approx \theta_{il} \exp(\Lambda_r q_t) \right], \quad (62)$$

where

$$\theta_{il} = \theta \exp \left[-\frac{L_v (q_l + q_{rain}) + L_s (q_i + q_{snow})}{c_{pd} T} \right] \quad (63)$$

is close to the ice-liquid formulation of Tripoli and Cotton (1981), except the constant values T_0 , $L_v(T_0)$ and $L_s(T_0)$ are replaced by the variable values T , $L_v(T)$ and $L_s(T)$ that depend on T in the exponential of (63), and with the rain and snow contents added from (9). The second-order relationship derived in Marquet (2015, 2019a) corresponds to:

$$\theta_s \approx \left[(\theta_s)_2 = (\theta_s)_1 \exp \left[-\gamma q_t \ln \left(\frac{r_v}{r_*} \right) - \gamma (q_l + q_{\text{rain}} + q_i + q_{\text{snow}}) \right] \right], \quad (64)$$

where $r_* \approx 12.4$ g/kg is a tuning parameter.

The reference entropies are computed with the standard temperature $T_r = T_0 = 273.15$ K and pressures $p_r = p_0 = 1000$ hPa and $e_r = e_{sw}(T_r) \approx 6.11$ hPa, with Eq. (36) and (37) leading to $s_{dr}(T_r, p_r - e_r) \approx 6777$ J/K/kg, $s_{vr}(T_r, e_r) \approx 12673$ J/K/kg, $r_r = \varepsilon e_r / (p_r - e_r) \approx 3.8214$ g kg⁻¹ and $q_r = r_r / (1 + r_r) \approx 3.8068$ g kg⁻¹

These values of the dry air and water vapor reference entropies lead to

$$\Lambda_r = \frac{s_{vr}(T_r, e_r) - s_{dr}(T_r, p_r - e_r)}{c_{pd}} \approx 5.869 \pm 0.003. \quad (65)$$

The accuracy of Λ_r can be computed from $\Delta s_{vr} \approx 2$ J/K/kg and $\Delta s_{dr} \approx 1$ J/K/kg, leading to $\Delta \Lambda_r \approx 3/1000 = 0.003$.

Furthermore, for ice-free conditions ($q_i = 0$), θ_{s1} can be expressed in terms of the Betts (1973) ‘‘conservative variables’’ (q_t , θ_l and θ_e), according to

$$\theta_{s1} \approx \theta_l \exp(\Lambda q_t) \approx \theta_e \exp \left[- \left(\frac{L_v}{c_{pd} T} - \Lambda_r \right) q_t \right], \quad (66)$$

$$\text{and thus } \theta_{s1} \approx \theta_l \exp(6 q_t) \approx \theta_e \exp(-3 q_t) \quad (67)$$

because $L_v / (c_{pd} T) \approx 9$ and $\Lambda_r \approx 6$.

This result shows that θ_s is different from both θ_l and θ_e if $q_t \neq 0$, and is in a two-third position in between these two variables, a property indeed valid in Fig. (8):

- (i) for the FIRE-I (RF03B) observed radial flight (Marquet, 2011);
- (ii) for drop-sounds observed during the EUREC⁴A - Circle (Marquet and Stevens, 2022);
- (iii) for (2.5 km) DYAMOND simulation outputs (Marquet and Stevens, 2022); and
- (iv) for all studied undergone since December 2008 (the date of the discovery of θ_s).

An important question is now to explain and justify the choices $T_r = T_0 = 273.15$ K and $p_r = p_0 = 1000$ hPa for the reference values equal to the standard values for temperature and pressure. Indeed, if θ_s given by (15) is independent on any values of T_r and p_r , both $(\theta_s)_1$ and $(\theta_s)_2$ given by (62) and (64) do depend on Λ_r given by (65), and thus on T_r and p_r .

Fig. 2 shows that $\Lambda(T_r, p_r)$ varies relatively little with the reference pressure p_r . Also, why not choose the standard value $p_r = p_0 = 1000$ hPa? And among all the possible values of T_r , the zero Celcius seems relevant since it represents a kind of average value according to the variations of T . Also, why choose another value than 273.15 K?

Moreover, it is the product of Λ_r with q_t that influences θ_s , and therefore with a maximum impact in the lower layers and temperatures and pressures close to $p_0 = 1000$ hPa and $T_0 = 273.15$ K.

In any case, if the aim is to go beyond the first order to the second order, the impact of humidity must be taken into account. Indeed, let us first define (without approximation) the more general variable Λ_s by the formula $\theta_s = \theta_{il} \exp(\Lambda_s q_t)$, where θ_{il} , q_t and θ_s given by (15) are three known




Figure 2: An unpublished plot of $\Lambda(T_r, p_r)$ for T_r from 220 K to 315 K and p_r from 1000 hPa to 200 hPa.

quantities, and thus with Λ_s given by the reciprocal relationship valid for the liquid case (with $q_i = 0$ and $\theta_{il} = \theta_l$):

$$\Lambda_s(T, p, q_v, q_l) = \frac{1}{q_t} \ln\left(\frac{\theta_s}{\theta_l}\right). \quad (68)$$

Fig. 3 (top) shows that, as a first guess, the term Λ_s varies from about 5.7 to 7.6 for 16 vertical profiles of cumulus and stratocumulus (FIRE-I RF02/03/04/08/10, EPIC, DYCOMS-II, ARM-Cu, ASTEX, ATEX, GATE, BOMEX, SCMS-RF12). A detailed study indicates that Λ_s is:

- smaller for the moister (solid-red) cumulus (q_t and q_v from 9 to 23 g/kg);
- larger for the moderate (dashed-blue) stratocumulus (q_t and q_v from 1 to 15 g/kg); and
- even larger for the dryer (solid-black) ASTEX case (q_t and q_v from 0.3 to 11 g/kg).





Figure 3: **Top:** an unpublished plot of $\Lambda(T_r, p_r)$ for T_r from 220 K to 315 K and p_r from 1000 hPa to 200 hPa. **Bottom:** a plot from Marquet (2019a) showing Λ_s a function of $\ln(r_v)$.

As a consequence, we are encouraged to seek for a value of Λ_s which would decrease with q_t or q_v . To do so, according to the second-order formulations derived in Marquet (2015) and Marquet (2019a) and leading to (64), the dependence of Λ_s with moisture variables could be of the form

$$\Lambda_s = \frac{1}{q_t} \ln\left(\frac{\theta_s}{\theta_l}\right) \approx \Lambda_r - \gamma \ln(r_v) - \gamma \frac{q_l}{q_t} + Cste, \quad (69)$$

where $\Lambda_r = 5.87$ and $\gamma \approx 0.46$. Fig. 3 (bottom) clearly shows that, for the same 16 vertical profiles of cumulus and stratocumulus as in the top figure, the term Λ_s indeed varies like $-\gamma \ln(r_v) = -0.46 \ln(r_v)$. This linear law appears to be valid for a large range of r_v (from 0.2 to 24 g kg⁻¹). This corresponds to a linear decrease of

$$\Lambda_s + \gamma \frac{q_l}{q_t} = \frac{1}{q_t} \left[\ln\left(\frac{\theta_s}{\theta_l}\right) + \gamma q_l \right] \approx \Lambda_r - \gamma \ln\left(\frac{r_v}{r_\star}\right) \quad (70)$$

with $\ln(r_v)$, where the constant in (69) is written without loss of generality as $-\gamma \ln(r_\star)$ in (70) in terms of a tuning parameter r_\star to be determined.

It is then possible to plot in Fig. 4 (top) the vertical profiles of this tuning parameter r_\star for the same 16 vertical profiles of cumulus and stratocumulus as in the Figs. 3, with r_\star obtained from (70) and written as

$$r_\star \approx r_v \exp\left\{ \frac{1}{\gamma q_t} \left[\ln\left(\frac{\theta_s}{\theta_l}\right) + \gamma q_l \right] - \frac{\Lambda_r}{\gamma} \right\}. \quad (71)$$

A first-guess value for r_\star derived in Marquet (2015) and Marquet (2019a) is $r_r \times \exp(1) \approx 3.82 \times 2.718 \approx 10.4$ g/kg.

According to the second-order formulation derived in Marquet (2019a), it is then possible to find the tuning mixing ratio r_\star for which hold true, where r_\star will play the role of positioning the dashed-dotted thick black line of slope $-\gamma \approx -0.46$ in between the cumulus and stratocumulus profiles. This corresponds to a linear fitting of r_v against $\exp[(\Lambda_r - \Lambda_s)/\gamma + \gamma q_l]$, r_\star being the average slope of the scattered data points. It is shown in Fig. 4 (top) that the value $r_\star \approx 12.4$ g kg⁻¹ corresponds to a relevant fitting of all cumulus and stratocumulus vertical profiles for a range of r_v from 0.2 up to 24 g kg⁻¹.

It is finally shown in Fig. 4 (bottom) that Λ_s can indeed be approximated by $\Lambda_\star(r_v, r_\star)$ given by (69), with a clear improved accuracy in comparison with the constant value $\Lambda_r \approx 5.87$ for a range of r_v between 0.2 and 24 g kg⁻¹. Curves of $\Lambda_\star(r_v, r_\star)$ with $r_\star = 10.4$ and 12.4 g kg⁻¹ (solid black lines) both simulate with a good accuracy the non-linear variation of Λ_s with r_v , and both simulate the rapid increase of Λ_s for $r_v < 2$ g kg⁻¹.





Figure 4: Top: Bottom:




Figure 5: **Top: Bottom:**

All definitions of $\theta_{s/\text{HH87}}$ or θ_l by Hauf and Höller (1987), θ_e by Betts (1973) recalled in (16), (6)

Previously, the relationship (16) for $\theta_{s/\text{HH87}}$ is usually approximated by the first-order approximation θ_e of Betts (1973) given by (6). Similarly, θ_l defined by (21) is usually approximated by the first-order approximation of Betts (1973) given by (7).

By doing this, the first-order approximation neglects the impact of the ratio e/e_0 for θ_l defined by (21), and also the impact of the choice of T_0 and e_0 in the additional terms left outside the logarithms in (16) and (21), which depends on the reference values $s_d(T_0, p_0)$, $s_v(T_0, e_0)$ and $s_l(T_0)$.

The moist-air entropy variables θ_s , θ_{s1} and θ_{s2} have been studied in several papers since 2011:

- for computing the moist-air Brunt-Väisälä frequency (Marquet and Geleyn, 2013);
- for computing and studying the associated Potential Vorticity $PV(\theta_s)$ (Marquet, 2014);
- in a synthetic chapter about moist-air thermodynamics (Marquet and Geleyn, 2015);
- for computing improved adjustment of marine bulk formulas (Marquet and Belamari, 2017);
- for computing the work- and stream-functions for the hurricane Dumilé (Marquet, 2017);
- to answers O. Pauluis's criticisms with Hector-the-Convecton (Marquet and Dauhut, 2018);
- to explain in French and English the third-law of thermodynamics (Marquet, 2019b,c);
- to improve a θ_s -based E.I.S. in the IFS-ECMWF model (Marquet and Bechtold, 2020);
- to compare the H2O pathways with lines of constant θ_e or θ_s (Marquet and Bailey, 2021);
- to provide a common method to derive θ_l , θ_e and θ_s (Marquet and Stevens, 2022);
- to use θ_s and q_t in a 1D-Var assimilation scheme (Marquet et al., 2022).

5 Comparison of dry and moist-air potential temperature

All previous dry- and moist-air potential temperatures (θ'_w , θ_v , θ_l , θ_{il} , θ , θ_s , θ_e , θ_{es} and θ'_e) have been computed for a realistic updraft based on (a slightly modified version of) the vertical profile of the Hurricane “Season” described in Table 5 of Jordan (1958), with precipitating and entrainment processes and with liquid water, ice and mixed phases. Figs 6 shows that:

- a clear-air region extends from the surface up to about 820 hPa;
- a mixed-phase exists between about 500 hPa and 300 hPa;
- θ_l and θ_{il} (in orange) remain close to θ (dashed black), with differences (in the cloud region between 850 and 200 hPa) of the same order of magnitude than the difference between θ and θ_v (dashed grey);
- the four versions of θ_e (in blue) are not so different from each others (they all increases with height with discrepancies of less than 1 to 2 K);
- the vertical changes in θ'_w (in violet) and θ_e (in blue) are very different (θ'_w is almost conservative whereas the four θ_e increase with height);
- the first- and second-order approximations θ_{s1} and θ_{s2} (dashed and dotted-dashed red) remain very close to the exact version θ_s (solid red), up to less than 0.6 K for θ_{s1} and 0.05 K for θ_{s2} ;
- the entropy potential temperature θ_s (in red) is indeed in a two-third position in between θ or θ_l (dashed black or orange) and θ_e (in blue), and is significantly different from all the other potential temperatures;
- the saturated-equivalent temperature θ_{es} (in black) is very different from the four θ_e (in blue) in the unsaturated low-levels region below the cloud (θ_{es} is much larger by more than 50 K);
- the pseudo-adiabatic equivalent potential temperature θ'_e (in green) computed from Eqs. (10)-(12), which looks like θ'_w but with enhanced variations close to those of the (about 6 times) enhanced (dashed violet) vertical profile of $299.5 + 6 \times (\theta'_w - 299.5)$.





Figure 6: Vertical profiles for an updraft based on the Hurricane “Season” (Jordan, 1958). Left: water vapor (q_v), liquid water ($10 \times q_l$) and ice ($10 \times q_i$) specific contents (in g/kg). Right: potential temperatures (in K).

Entropy in Arpege+Arome+Meso-NH : from 1987 ... to 2022



After more than 30 years of work, and looking ahead to 2022:

→ The more general formulation for θ_s (Marquet, JAS, 2017):

either of these 2 formulations used hereafter

$$\theta_s = \theta \exp\left(-\frac{L_v q_l + L_s q_i}{c_{pd} T}\right) \exp\{\Lambda_r q_t\} \times \left(\frac{T}{T_r}\right)^{\lambda q_t} \left(\frac{p}{p_r}\right)^{-\kappa \delta q_t} \left(\frac{r_r}{r_v}\right)^{\gamma q_t} \frac{(1+\eta r_v)^{\kappa(1+\delta q_t)}}{(1+\eta r_r)^{\kappa \delta q_t}} \times (H_l)^{\gamma q_l} (H_i)^{\gamma q_i} \left(\frac{T_l}{T}\right)^{c_l q_l/c_{pd}} \left(\frac{T_i}{T}\right)^{c_i q_i/c_{pd}}$$

→ With a relevant first-order formulation (Marquet, 2011, 2015, 2019) made by the first line only:

$$\theta_{s1} = \theta \exp\left(-\frac{L_v q_l + L_s q_i}{c_{pd} T}\right) \exp\{\Lambda_r q_t\} \approx \theta_{il} \exp\{\Lambda_r q_t\}$$

→ Impact of the Third Law:

$$\Lambda_r \approx 5.869 \pm 0.003$$


→ And with even more relevant simple formulations (if needed):

$$\theta_s \approx \theta_{il} \times \exp\{\Lambda_r q_t\} \approx \theta \times (1 + \Lambda_r q_v)$$

!! not used hereafter (just for pedagogy)

Figure 7: Two slides from Marquet (2022), where the main properties of the moist-air specific entropy $s = c_{pd} \ln(\theta_s) + s_0$, the associated potential temperature θ_s and the potential vorticity $PV(\theta_s)$ are summarized.

Observations : FIRE-I air flights / Marquet (2011)



→ θ_s was indeed in a 2/3 position
 between θ_l et θ_e
 → Only moist-air entropy (θ_s) is
 well-mixed ← (constant) from
 the surface up to 1050 m includ-
 ing in the entrainment region!

$$\theta_l \approx \theta_l \times (1 + 0 q_t)$$

$$\theta_s \approx \theta_l \times (1 + 6 q_t)$$

$$\theta_e \approx \theta_l \times (1 + 9 q_t)$$

Observations & Simulation : Marquet and Stevens (JAS 2022)

Average for 757 drop-sounds during the EUREC4-A campaign

Stevens et al. (2021) ; George et al. (2021)




Figure 1: Mean profiles of the liquid water, θ_l , entropy, θ_s , equivalent θ_e potential temperatures from 757 dropsondes launched along the EUREC⁴A-Circle

Observations

$$\theta_l \approx \theta_l \times (1 + 0 q_t)$$

$$\theta_s \approx \theta_l \times (1 + 6 q_t)$$

$$\theta_e \approx \theta_l \times (1 + 9 q_t)$$




Figure 3: Profiles of θ_e , θ_l and θ_s for a composite sounding over the dry (dotted) and moist (solid) tropics.

derived from global storm-resolving (2.5 km) simulations from the DYAMOND project (Stevens et al., 2019) using the ICON model (Hohenegger et al., 2020).


Simulation

→ This implies that θ_s
is in a 2/3 position
 between θ_l et θ_e

P. Marquet (January 2022)

Figure 8: From Marquet (2022). **Top:** vertical profiles of potential temperatures from the Radial-Flight 03B of FIRE-I (Marquet, 2011). **Bottom:** vertical profiles of potential temperatures from dropsondes along the EUREC⁴A-Circle and from tropical simulated composite soundings (Marquet and Stevens, 2022)

Observations : Marquet (La Météorologie, 2019a,b)
Marquet and Stevens (JAS, 2022)



Meso-NH L.E.S. Shallow + Deedp / Didier RICARD (2018 / unpublished)






Figure 9: From Marquet (2022). **Top:** half of the ASTEX Lagrangien vertical profiles for θ_l , q_t and θ_s computed from the radio-sounding dataset for T , q_v , q_l (Marquet, 2019b,c; Marquet and Stevens, 2022). The special property $\partial\theta_s/\partial p \approx 0$ at the transition between Stratocumulus and Cumulus regimes has been used to build a new Estimated-Inersion-Strength in the IFS model at ECWF (Marquet and Bechtold, 2020). **Bottom:** diagnostic outputs from the Meso-NH models (Ricard, 2018, unpublished), where the moist-air entropy (and θ_s) entering the plumes of convective cells are more similar to mid-tropospheric values at 6 to 8 km (differently from colder PBL values for θ_l and warmer PBL values for θ_e).

Analyses from Arpege NWP model : Marquet (ENM course 2019)




Simulations : Marquet and Bailey (WGNE-Blue-Book, 2021)



- \rightarrow The H₂O pathways emitted from 40S-30S almost follow iso-300K θ_s lines
- \rightarrow with up to “8 degrees & 8 K” of bias with the iso-300K θ_e lines
- \rightarrow a validation that θ_s and the “true isentropes” have a physical impact?

Figure 10: From Marquet (2022). **Top:** the surface of equal values of θ , θ_s and θ_e do not coincide in annual and zonal average, with a 2/3 position of θ_s in between θ and θ_e (unpublished result; from a 2019 course at the ENM / French School of Meteorology). Only θ_s represents the moist-air entropy and generates the true moist-air isentropes. One of the consequence is an impact on the definition of moist-air baroclinic waves in mid-latitudes (different slopes of “isentropic” surfaces). **Bottom:** the H₂O pathways originating from a certain band of latitude seems to preferentially follow the moist-air isentropes labeled with θ_s (Marquet and Bailey, 2021).



Isobaric charts of Potential Temperatures (colour scales + dashed contours) θ'_w and θ_s at 850 hPa + MSLP (solid contours)


ARPEGE 17 Oct. 2012 r00 (12h)

Almost the same patterns ... but with different colour scales

Since this difference depends on the level, this must correspond to more different cross-sections?

$$\theta_s \approx \theta \times (1 + 6 q_v)$$


Marquet (2012) ... Marquet QJRMS (2014)



P. Marquet (January 2022)

Figure 11: From Marquet (2022). **Top:** It is possible to plot air-mass charts (here at 850 hPa) by replacing the pseudo-adiabatic variable θ'_w (or θ_e) by the moist-air entropy variable θ_s (Blot, 2013). It is just necessary to change the color scale. **Bottom:** Vertical cross-sections published in Marquet (2014) showing how a front can be described with the variables θ , θ'_w , θ_s and θ_e . The moist-air entropy variable θ_s allows a better view of the air masses in the lower layers in relation to the frontal surfaces, with a boundary layer that is better mixed (homogeneous) in θ_s , and with the convective regions associated with some sort of "aspiration of moist-air isentropes".

M2 internship of Etienne BLOT (2013)



M2 internship of Etienne BLOT (2013)






Figure 12: From Marquet (2022) and Blot (2013). **Top:** the structure of a split cold front can be represented by the moist-air entropy variable θ_s , as relevant as with the θ'_w variable. **Bottom:** the same is true for a squall-line line, which is clearly visible in terms of θ_s , with the convective regions associated with some sort of “aspiration of moist-air isentropes”.

M2 internship of Etienne BLOT (2013)



→ Eye-walls (solid black lines)
 + Spiral bands (dashed black lines)
 plotted from a cross-section for RH
 (not shown)

→ Almost the « same visions » for θ'_w & θ_e
 (thanks to different colour scales)





Etienne BLOT (2013) ... Marquet (2019)

 Cross-sections for the Hurricane Dumilé

 ALADIN-Réunion 3 Jan. 2013 r00 (12h)

M2 internship of Etienne BLOT (2013)



- A severe test for the entropy (θ_s), because the more intense/moist atmospheric phenomena!
- Eye-walls (solid black) and spiral bands (dashed black) lines are clearly visible with the entropy (θ_s)
- ... like an "aspiration" of the (true) isentropes (θ_s)
- No more "warm region" close to the surface with θ_s
- (better) well-mixed (constant) θ_s in the boundary layer

Etienne BLOT (2013) ... Marquet (2019)

 Cross-sections for the Hurricane Dumilé

 ALADIN-Réunion 3 Jan. 2013 r00 (12h)

$$\theta_s \approx \theta \times (1 + 6 q_v)$$

Figure 13: From Marquet (2022) and Blot (2013). **Top:** the structure of a hurricane (here Dumilé, simulated by the Aladin 8 km model) can be equally represented by the pseudo-adiabatic variables θ'_w and θ_e (thanks to different color scales). **Bottom:** the moist-air entropy variable θ_s is as relevant as θ'_w and θ_s variable to see the core, the eye-walls and the spiral bands, with more homogeneous boundary layers and with the convective regions associated with some sort of "aspiration of mid-troposphere moist-air isentropes" labeled with θ_s .

Recent results about Arpege and Arome outputs?






Figure 14: From Marquet (2022). **Top:** The surface fronts for the storm "Barra" are clearly located with the MSLP simulated with the Arome 1.3 km model. **Bottom:** The cross-sections at 46 N allow the plot of cold (solid blue) and warm (solid red) fronts up to 700 to 400 hPa.


ANASYG
2021 Dec. 07
r00 / 12 UTC



(with the "epygram" tool / A. Mary)
12h Forecast from 2021 Dec. 07 00UTC

Cross-section 46 N
(-11.5 W / 15.5 E)

- « air masses » more visible and homogeneous with θ_s than θ'_w !
- only low-level cold fronts?
- two separate low- & mid-level warm front?




$$\theta_s \approx \theta \times (1 + 6 q_t)$$

- The same vision for ARPEGE (9 km) and AROME-NH (1.3 km) NWP models


P. Marquet (January 2022)

ANASYG
2021 Dec. 07
r00 / 12 UTC



(with the "epygram" tool / A. Mary)
12h Forecast from 2021 Dec. 07 00UTC

Cross-section 45 N
(-11.5 W / 15.5 E)



$$\theta_s \approx \theta \times (1 + 6 q_t)$$

- The same at 45 N

P. Marquet (January 2022)

Figure 15: From Marquet (2022). **Top:** The cold (solid blue) and warm (solid red) fronts seems to be better associated with “air masses” labeled with the moist-air entropy variable θ_s for the cross section at 46 N (no more marked minimum at 850 or 800 hPa as for θ'_w). This result is valid for the Arome 1.3 km model and the Arpege 8 km models as well. **Bottom:** The same is true for the cross section at 45 N.




Figure 16: From Marquet (2022). Historical examples justifying the plot of fronts with limited vertical extension on the previous cross-sections.

6 The dry- and moist-air potential vorticity formulations $PV(\theta)$

The original formulation of “potential vorticity” by Ertel (1942a)

$$PV(\psi) = \frac{1}{\rho} \boldsymbol{\zeta}_a \cdot \nabla(\psi) \quad (72)$$

was suitable for any “hydrodynamical invariant” (i.e. “conservative variable”) ψ , with ρ the density, $\boldsymbol{\zeta}_a$ the 3D absolute vorticity vector and ∇ the 3D gradient operator (see the English translations of all Ertel’s papers in Schubert et al., 2004).

Ertel suggested different kinds of such “conservative variable” ψ :

- the “polytropic temperature under polytropic conditions” (for which the soleonidal term $N(p, \rho, \psi)$ cancels out), and thus the (dry-air) “potential temperature” $\psi = \theta$, or the “total water content” $\psi = q_t = q_v + q_l + q_i$ under conditions of no precipitation and without moisture input (Ertel, 1942a);
- the (dry-air) potential temperature $\psi = \theta$, i.e. a special case of the polytropic temperature $\psi = \Theta_x$ with $\Theta_x = T(\rho_0/\rho)^{(1/x)}$ and with $x = c_v/(c_p - c_v)$ (Ertel, 1942b);
- the (dry-air) polytropic temperature or the (dry-air) entropy $\psi = s_d$ (Ertel, 1942c), due to the Bauer (1908, 1910) relationship $s_d = c_{pd} \ln(\theta) + cste$, and thus with $PV(s_d) = PV[c_{pd} \ln(\theta)] = (c_{pd}/\theta) PV(\theta)$ indeed proportional to $PV(\theta)$.

After Ertel’s papers, different dry- and moist-air “(more or less) conservative” quantities have been studied:

- $PV(\theta)$ in Hoskins and Bretherton (1972); Hoskins (1974); Hoskins et al. (1985), Hoskins and Berrisford (1988); Hoskins (1991, 1997); Hoskins et al. (2003), (...) Hoskins (2015);
- $PV(\theta'_w)$ in Bennetts and Hoskins (1979);
- $PV(\theta_e)$ in Rotunno and Klemp (1985), Cao and Cho (1995), Persson (1995), Gao et al. (2002), Deng and Gao (2009);
- $PV(\theta_{es})$ in Emanuel et al. (1987);
- $PV(\theta_s)$ in Rivas Soriano and García Díez (1997) / for θ_s defined in Hauf and Höller (1987);
- $PV(\theta_v)$ in Schubert et al. (2001) and Schubert (2004);
- $PV(q_v)$, $PV(\theta)$ and $PV(\theta_e)$ in Gao and Zhou (2008);
- $PV(\theta_s)$, $PV(\theta_v)$ and $PV(q_t)$ in Marquet (2014) / for θ_s defined in Marquet (2011).

7 Comparison of dry and moist-air Potential Vorticity




Figure 17: From Marquet (2022). **Top:** It is recalled in this figure of R.A. Houze that regions with negative values of $PV(\theta_e)$ are associated with symmetric instability and slantwise convection. The first-order formulations for $PV(\theta_s)$ and $PV(\theta_e)$ show that the impact of the moisture part $PV(q_v)$ in $PV(\theta_s)$ is about 2/3 of the moisture part in $PV(\theta_e)$. **Bottom** (from Marquet, 2014): Average low-level values seems too positive for $PV(\theta) \approx PV(\theta_v)$, too negative for $PV(\theta_e)$ and well-balanced for $PV(\theta_s)$.

Old results: moist-air entropy Potential Vorticity?

- $PV(\theta_s)$, $PV(\theta_v)$ and $PV(q_t)$ in Marquet (2014) / for θ_s defined in Marquet (2011).





Figure 18: From Marquet (2022) and Marquet (2014). **Top:** the low-level (850 hPa) cold fronts seems to be associated with the negative values of $PV(\theta_s)$, whereas $PV(\theta)$ is positive everywhere and $PV(\theta_e)$ negative almost everywhere. **Bottom:** the negative values of $PV(\theta)$ seems clearly correspond to (are located with) the cold fronts in these cross-sections (same fronts as in Fig. 11), and possibly to symmetric instability and slantwise convection. On the one hand there is no negative values for $PV(\theta)$, whereas on the other hand there are negative values of $PV(\theta_e)$ almost everywhere in the low-level boundary layer.

Recent results: moist-air entropy Potential Vorticity?





Figure 19: From Marquet (2022). **Top:** The potential vorticity $PV(\theta)$ and $PV(\theta_s)$ for the storm "Barra" simulated with the global Arpege NWP (8 km) model (cross-sections at 46 N). Some negative values of the moist-air entropy variable $PV(\theta_s)$ appears close to the cold front (they already exist and are the same between the warm fronts), with almost unchanged upper-levels dynamic-meteorology signals above 500 hPa in the dry regions (tropospheric intrusion of high stratospheric values). **Bottom:** The same is true for the Arome 1.3 km (NH) model.

Recent results: moist-air entropy Potential Vorticity?




Figure 20: From Marquet (2022). Same results as in Fig 19 but at 45 N.

8 Conclusions

This note is a mere synthesis of the most important results I have discovered since 2009, with the aim to mainly show Figures instead of formulas (which can be found in my papers published since 2011 and available on arXiv).

However, I am now I am in retirement (since May, 2022) from the CNRM and Météo-France, and I am preparing a (huge) book showing in detail why it is needed, and how it is possible to defined and compute the third-law (absolute) definitions of not only the moist-air entropy for the atmosphere, but also the sea-salt entropy for the ocean.

In this book I will recall (and show) almost all the bibliography I have studied since 1988 about the atmosphere and ocean thermodynamics (even though many of them, but not all of them, have been recalled in my papers published between 2011 and 2022).

References

- Atkins, P., and J. de Paula, 2014: *Physical chemistry. Thermodynamics, structure, and change*, 1008 Pp. W. H. Freeman and Company, New-York.
- Bauer, L. A., 1908: The Relation between “Potential Temperature” and “Entropy”. *Phys. Rev. (Series I)*, **26 (2)**, 177–183, doi:10.1103/PhysRevSeriesI.26.177.
- Bauer, L. A., 1910: *The relation between “potential temperature” and “entropy”*, Art. XXII. 495–500 Pp. The Mechanics of the Earth Atmosphere. Collection of translations by Cleveland Abbe. Smithsonian Miscellaneous Collections.
- Bennetts, D. A., and B. J. Hoskins, 1979: Conditional symmetric instability - a possible explanation for frontal rainbands. *Quart. J. Roy. Meteorol. Soc.*, **105 (446)**, 945–962, doi:10.1002/qj.49710544615.
- Betts, A. K., 1973: Non-precipitating cumulus convection and its parameterization. *Quart. J. Roy. Meteorol. Soc.*, **99 (419)**, 178–196, doi:10.1002/qj.49709941915.
- Betts, A. K., and F. J. Dugan, 1973: Empirical formula for saturation pseudoadiabats and saturation equivalent potential temperature. *J. Appl. Meteor.*, **12 (4)**, 731–732, doi:10.1175/1520-0450(1973)012<0731:EFFSPA>2.0.CO;2.
- Bleeker, W., 1939: On the conservatism of the equivalent potential and the wet-bulb potential temperatures. *Quart. J. Roy. Meteor. Soc.*, **65 (282)**, 542–550, doi:10.1002/qj.49706528212.
- Blot, E., 2013: Etude de l’entropie humide dans un contexte d’analyse et de prévision du temps. Météo-France, Ecole Nationale de la Météorologie. Toulouse, Rapport de stage d’approfondissement EIENM3.
- Bolton, D., 1980: The computation of equivalent potential temperature. *Mon. Weather. Rev.*, **108 (7)**, 1046–1053, doi:10.1175/1520-0493(1980)108<1046:TCOEPT>2.0.CO;2.
- Bougeault, P., 1982: Cloud-ensemble relations based on the gamma probability distribution for the higher-order models of the planetary boundary layer. *J. Atmos. Sci.*, **39 (12)**, 2691–2700, doi:10.1175/1520-0469(1982)039<2691:CERBOT>2.0.CO;2.
- Bryan, G. H., 2008: On the computation of pseudoadiabatic entropy and equivalent potential temperature. *Mon. Weather. Rev.*, **136 (12)**, 5239–5245.
- Cao, Z., and H.-R. Cho, 1995: Generation of moist potential vorticity in extratropical cyclones. *J. Atmos. Sci.*, **52 (18)**, 3263–3281, doi:10.1175/1520-0469(1995)052<3263:GOMPVI>2.0.CO;2.
- Chase, M. W., 1998: *Journal of physical and chemical reference data. Monograph No 9. NIST-JANAF thermochemical tables. Fourth Edition*, 1951 Pp. American Chemical Society and American Institute of Physics.
- Cuxart, J., P. Bougeault, and J.-L. Redelsperger, 2000: A turbulence scheme allowing for mesoscale and large-eddy simulations. *Quart. J. Roy. Meteorol. Soc.*, **126 (562)**, 1–30, doi:10.1002/qj.49712656202.

- Dauhut, T., J.-P. Chaboureaud, P. Mascart, and O. Pauluis, 2017: The atmospheric overturning induced by Hector the convective. *J. Atmos. Sci.*, **74** (10), 3271–3284, doi:10.1175/JAS-D-17-0035.1.
- Davies-Jones, R., 2009: On formulas for equivalent potential temperature. *Monthly Weather Review*, **137** (9), 3137 – 3148, doi:10.1175/2009MWR2774.1.
- Deardorff, J. W., 1980: Cloud top entrainment instability. *J. Atmos. Sci.*, **37** (1), 131–147, doi:10.1175/1520-0469(1980)037<0131:CTEI>2.0.CO;2.
- Deardorff, J. W., and G. Sommeria, 1977: Subgrid-scale condensation in models of non-precipitating clouds. *J. Atmos. Sci.*, **34** (2), 344–355, doi:/10.1175/1520-0469(1977)034<0344:SSCIMO>2.0.CO;2.
- Deng, G., and S.-t. Gao, 2009: The theory of moist potential vorticity and its application in the diagnosis of typhoon rainfall and intensity. *J. Trop. Meteorol.*, **15** (2), 204–209.
- Emanuel, K. A., 1994: *Atmospheric Convection*, 580 Pp. Oxford University Press.
- Emanuel, K. A., M. Fantini, and A. J. Thorpe, 1987: Baroclinic instability in an environment of small stability to slantwise moist convection. Part I: Two-dimensional models. *J. Atmos. Sci.*, **44** (12), 1559–1573, doi:10.1175/1520-0469(1987)044<1559:BIIAEO>2.0.CO;2.
- Ertel, H., 1942a: (a) Ein neuer hydrodynamischer Wirbelsatz [A new hydrodynamical vorticity equation]. *Meteorologische Zeitschrift*, **59** (9), 277–281, doi:10.1007/BF01475602.
- Ertel, H., 1942b: (b) Ein neuer hydrodynamischer Erhaltungssatz [A new hydrodynamical conservation principle]. *Die Naturwiss (Berlin)*, **36** (30), 543–544.
- Ertel, H., 1942c: (c) Über hydrodynamische Wirbelsätze [On hydrodynamical vorticity equations]. *Physik. Z. (Leipzig)*, **43**, 526–529.
- Fang, J., O. Pauluis, and F. Zhang, 2017: Isentropic analysis on the intensification of Hurricane Edouard (2014). *J. Atmos. Sci.*, **74** (12), 4177–4197, doi:10.1175/JAS-D-17-0092.1.
- Fang, J., O. Pauluis, and F. Zhang, 2019: The thermodynamic cycles and associated energetics of Hurricane Edouard (2014) during its intensification. *J. Atmos. Sci.*, **76** (6), 1769–1784, doi:10.1175/JAS-D-18-0221.1.
- Feistel, R., 2012: A new international oceanographic standard for seawater, ice, fluid water and humid air. *Int. J. Thermophys.*, **33** (8), 1335–1351, doi:10.1007/s10765-010-0901-y.
- Feistel, R., 2019: Distinguishing between Clausius, Boltzmann and Pauling entropies of frozen non-equilibrium states. *Entropy*, **21** (799), 1–24, doi:10.3390/e21080799.
- Feistel, R., and O. Hellmuth, 2020: Zur Rolle des Wassers in der Energiebilanz des Klimasystems (On the role of water in the energy balance of the climate system). English translation available on https://www.researchgate.net/publication/346785071_On_the_Role_of_Water_in_the_Energy_Balance_of_the_Climate_System). *Sitzungsberichte der Leibniz-Sozietät der Wissenschaften zu Berlin*, **144**, 51–139, doi:10.13140/RG.2.2.19649.17766, URL <https://leibnizsozietat.de/wp-content/uploads/2020/12/Feistel-R-und-Hellmuth-O.pdf>.

- Feistel, R., D. G. Wright, H.-J. Kretzschmar, E. Hagen, S. Herrmann, and R. Span, 2010: Thermodynamic properties of sea air. *Ocean Sci.*, **6** (1), 91–141, doi:10.5194/os-6-91-2010, URL <http://www.ocean-sci.net/6/91/2010/>.
- Feistel, R., D. G. Wright, K. Miyagawa, A. H. Harvey, J. Hraby, D. R. Jackett, T. J. McDougall, and W. Wagner, 2008: Mutually consistent thermodynamic potentials for fluid water, ice and seawater: a new standard for oceanography. *Ocean Sci.*, **4** (4), 275–291, doi:10.5194/os-4-275-2008, URL <http://www.ocean-sci.net/4/275/2008/>.
- Gao, S.-T., T. Lei, and Z. Yu-shu, 2002: Moist potential vorticity anomaly with heat and mass forcings in torrential rain system. *Chin. Phys. Lett.*, **19** (6), 878–880.
- Gao, S.-T., and F.-F. Zhou, 2008: Water vapour potential vorticity and its applications in tropical cyclones. *Chin. Phys. Lett.*, **25** (10), 3830–3833, doi:10.1088/0256-307X/25/10/085.
- Gokcen, N., and R. Reddy, 1996: *Thermodynamics (Chapter VII on the Third Law)*, 400 Pp. Springer Science+Business Media, New York.
- Guldberg, C. M., and H. Mohn, 1876: *Études sur les mouvements de l’atmosphère. Première partie. Programme de l’Université pour le 2e semestre 1876 (Studies on the movements of the atmosphere. First part. University program for the 2nd semester of 1876)*, 44 Pp. Imprimerie de Brøgger, Institutions scientifiques et littéraires de la Norvège. Christiania (Oslo).
- Guldberg, C. M., and H. Mohn, 1910: Studies on the movements of the atmosphere. Part I. (Christiania/Oslo), 1876, revised 1883). In: *The Mechanics of the Earth Atmosphere, a collection of translations by Cleveland Abbe, 1910. Smithsonian Miscellaneous Collections. [Translated from: Études sur les mouvements de l’atmosphère., 44 pp. (1876)]*, **Art. XI**, 122–174.
- Hauf, T., and H. Höller, 1987: Entropy and potential temperature. *J. Atmos. Sci.*, **44** (20), 2887–2901, doi:10.1175/1520-0469(1987)044<2887:EAPT>2.0.CO;2.
- Hewson, E. W., 1936: The application of wet-bulb potential temperature to air mass analysis. I. *Q. J. R. Meteorol. Soc.*, **62** (266), 387–420, doi:10.1002/qj.49706226604.
- Hewson, E. W., 1938: The application of wet-bulb potential temperature to air mass analysis. IV. *Q. J. R. Meteorol. Soc.*, **64** (276), 407–422, doi:10.1002/qj.49706427606.
- Hewson, E. W., 1946a: The wet-bulb thermometer-its meaning and significance. Part I. *Weather*, **1** (7), 194–197, doi:10.1002/j.1477-8696.1946.tb00084.x.
- Hewson, E. W., 1946b: The wet-bulb thermometer-its meaning and significance. Part II. *Weather*, **1** (8), 226–230, doi:10.1002/j.1477-8696.1946.tb00098.x.
- Hoskins, B., 1997: A potential vorticity view of synoptic development. *Meteorological Applications*, **4** (4), 325–334, doi:10.1017/S1350482797000716.
- Hoskins, B., and P. Berrisford, 1988: A potential vorticity perspective of the storm of 15-16 October 1987. *Weather*, **43** (3), 122–129, doi:10.1002/j.1477-8696.1988.tb03890.x.
- Hoskins, B., M. Pedder, and D. W. Jones, 2003: The omega equation and potential vorticity. *Quart. J. Roy. Meteorol. Soc.*, **129** (595), 3277–3303, doi:10.1256/qj.02.135.

- Hoskins, B. J., 1974: The role of potential vorticity in symmetric stability and instability. *Quart. J. Roy. Meteorol. Soc.*, **100** (425), 480–482, doi:10.1002/qj.49710042520.
- Hoskins, B. J., 1991: Towards a PV- θ view of the general circulation. *Tellus*, **43A** (4), 27–35, doi:10.3402/tellusa.v43i4.11936.
- Hoskins, B. J., 2015: Potential vorticity and the PV perspective. *Adv. Atmos. Sci.*, **32**, 2–9, doi:10.1007/s00376-014-0007-8.
- Hoskins, B. J., and F. P. Bretherton, 1972: Atmospheric frontogenesis models: Mathematical formulation and solution. *J. Atmos. Sci.*, **29** (1), 11–37, doi:10.1175/1520-0469(1972)029<0011:AFMMFA>2.0.CO;2.
- Hoskins, B. J., M. E. McIntyre, and A. W. Robertson, 1985: On the use and significance of isentropic potential vorticity maps. *Q. J. R. Meteorol. Soc.*, **111** (470), 877–946, doi:https://doi.org/10.1002/qj.49711147002.
- IAPWS, 2010: *Guideline on an equation of state for humid air in contact with seawater and ice, consistent with the IAPWS formulation 2008 for the thermodynamic properties of seawater*. International Association for the Properties of Water and Steam, URL <http://www.iapws.org/relguide/SeaAir.pdf>, Authorized by the IAPWS at its meeting in Niagara Falls, Canada, July 2010. Pp.19.
- Iribarne, J. V., and W. L. Godson, 1973: *Atmospheric Thermodynamics*, 222 Pp. D. Reidel Publishing Company, Dordrecht, Holland, doi:https://doi.org/10.1007/978-94-010-2642-0.
- Iribarne, J. V., and W. L. Godson, 1981: *Atmospheric Thermodynamics*, 260 Pp. D. Reidel Publishing Company, Dordrecht, Holland.
- Jordan, C. L., 1958: Mean soundings for the west indies area. *J. Atmos. Sci.*, **15** (1), 91–97, doi:10.1175/1520-0469(1958)015<0091:MSFTWI>2.0.CO;2.
- Kelley, K. K., 1932: *Contributions to the Data on Theoretical Metallurgy: I. The entropies of inorganic substances. Bulletin 350*, 63 Pp. United States, Government Printing Office, Washington.
- Laliberté, F., T. Shaw, and O. M. Pauluis, 2013: A theory for the lower-tropospheric structure of the moist isentropic circulation. *J. Atmos. Sci.*, **70** (3), 843–854, doi:10.1175/JAS-D-12-066.1.
- Lemmon, E. W., R. T. Jacobsen, S. G. Penoncello, and D. G. Friend, 2000: Thermodynamic properties of air and mixtures of Nitrogen, Argon, and Oxygen from 60 to 2000 K at pressures to 2000 MPa. *Journal of Physical and Chemical Reference Data.*, **29** (3), 331–385.
- Lewis, G. N., and M. Randall, 1961: *Thermodynamics*, 723 Pp. McGraw-Hill Book Company. New York.
- Lilly, D. K., 1968: Models of cloud-topped mixed layers under a strong inversion. *Quart. J. Roy. Meteorol. Soc.*, **94** (401), 292–309, doi:10.1002/qj.49709440106.
- Lock, A. P., 2004: The sensitivity of a GCM’s marine stratocumulus to cloud-top entrainment. *Quart. J. Roy. Meteorol. Soc.*, **130** (604), 3323–3338, doi:https://doi.org/10.1256/qj.03.114.

- MacVean, M. K., and P. J. Mason, 1990: Cloud-top entrainment instability through small-scale mixing and its parameterization in numerical models. *J. Atmos. Sci.*, **47** (8), 1012–1030, doi:10.1175/1520-0469(1990)047<1012:CTEITS>2.0.CO;2.
- Marquet, P., 1993: Exergy in meteorology: definition and properties of moist available enthalpy. *Quart. J. Roy. Meteorol. Soc.*, **119** (511), 567–590, doi:10.1002/qj.49711951112, URL <https://arxiv.org/abs/1807.05830>.
- Marquet, P., 2011: Definition of a moist entropy potential temperature: application to FIRE-I data flights. *Quart. J. Roy. Meteorol. Soc.*, **137** (656), 768–791, doi:10.1002/qj.787, URL <http://arxiv.org/abs/1401.1097>.
- Marquet, P., 2014: On the definition of a moist-air potential vorticity. *Quart. J. Roy. Meteorol. Soc.*, **140** (680), 917–929, doi:10.1002/qj.2182, URL <http://arxiv.org/abs/1401.2006>.
- Marquet, P., 2015: An improved approximation for the moist-air entropy potential temperature θ_s . *Research Activities in Atmospheric and Oceanic Modelling. WRCP-WGNE Blue-Book*, **4**, 14–15, URL http://bluebook.meteoinfo.ru/uploads/2015/individual-articles/04_Marquet_Pascal_Entropie.pdf.
- Marquet, P., 2017: A Third-Law Isentropic Analysis of a Simulated Hurricane. *J. Atmos. Sci.*, **74** (10), 3451–3471, doi:10.1175/JAS-D-17-0126.1, URL <https://arxiv.org/abs/1704.06098>.
- Marquet, P., 2019a: The first and second order approximations of the third-law moist-air entropy potential temperature. *Submitted (and rejected) to the Quart. J. Roy. Meteorol. Soc., then to the Mon. Wea. Rev., and then to the J. Atmos. Sci.*, -, 1–14, URL <https://arxiv.org/abs/1901.08108>.
- Marquet, P., 2019b: Le troisième principe de la thermodynamique ou une définition absolue de l'entropie - Partie 1 : Origines et applications en thermodynamique. *La Météorologie*, **107** (Série 8), 45–52, doi:10.4267/2042/70558, URL <https://arxiv.org/abs/1904.11696>.
- Marquet, P., 2019c: Le troisième principe de la thermodynamique ou une définition absolue de l'entropie - Partie 2 : Définitions et applications en météorologie et en climat. *La Météorologie*, **107** (Série 8), 53–63, doi:10.4267/2042/70559, URL <https://arxiv.org/abs/1904.11699>.
- Marquet, P., 2022: Entropy in Arpege/Arome NWP models : computations / post-treatments / use by forecasters? *Workshop for Forecasting and Modelling, 24th of January, 2022 (Météo-France / CNRM-GMAP), Toulouse,*, -, 38 slides, URL https://www.researchgate.net/publication/358395622_Entropy_in_ArpegeArome_NWP_models_computations_posttreatments_use_by_forecasters_Workshop_for_Forecasting_and_Modelling_24th_of_January_2022_Meteo-France_CNRM-GMAP_Toulouse_France.
- Marquet, P., and A. Bailey, 2021: Comparisons of H₂O pathways with moist isentropes. *Research activities in Earth system modelling. Working Group on Numerical Experimentation. Report No51. WCRP Report No13/2021. Ed. E. Astakhova, WMO, Geneva*, **51**, 1–2, URL http://bluebook.meteoinfo.ru/uploads/2021/docs/04_MARQUET_Pascal_IsentropesH20Path.pdf.
- Marquet, P., and P. Bechtold, 2020: A new estimated inversion strength (EIS) based on the moist-air entropy. *Research activities in Earth system modelling. Working Group on Numerical Experimentation. Report No50. WCRP Report No12/2020. Ed. E. Astakhova, WMO,*

- Geneva*, **50**, 4–07, URL http://bluebook.meteoinfo.ru/uploads/2020/docs/04_Marquet.Pascal.NewEntropyEIS.pdf.
- Marquet, P., and S. Belamari, 2017: On new bulk formulas based on moist-air entropy. *Research Activities in Atmospheric and Oceanic Modelling. WRCP-WGNE Blue-Book*, **4**, 9–10, URL http://bluebook.meteoinfo.ru/uploads/2017/docs/04_MARQUET.Pascal.moist_entropy_bulk.pdf.
- Marquet, P., and T. Dauhut, 2018: Reply to “Comments on ‘A third-law isentropic analysis of a simulated hurricane’”. *J. Atmos. Sci.*, **75** (10), 3735–3747, doi:10.1175/JAS-D-18-0126.1, URL <https://arxiv.org/abs/1805.00834>.
- Marquet, P., and J.-F. Geleyn, 2013: On a general definition of the squared Brunt-Väisälä frequency associated with the specific moist entropy potential temperature. *Quart. J. Roy. Meteorol. Soc.*, **139** (670), 85–100, doi:10.1002/qj.1957, URL <http://arxiv.org/abs/1401.2379>.
- Marquet, P., and J.-F. Geleyn, 2015: Formulations of moist thermodynamics for atmospheric modelling. *Parameterization of Atmospheric Convection. Vol II: Current Issues and New Theories*, R. S. Plant, and J.-I. Yano, Eds., *World Scientific*, Imperial College Press, 221–274, doi:10.1142/9781783266913_0026, URL <http://arxiv.org/abs/1510.03239>.
- Marquet, P., J.-F. Mahfouf, A. L. Barbu, and B. Ménétrier, 2022: Data assimilation using conservative thermodynamic variables. *Atmospheric Measurement Techniques (In revision)*, **XX** (nn), ?–?, doi:10.5194/amt-2021-361, URL <https://arxiv.org/abs/2111.05883>.
- Marquet, P., and B. Stevens, 2022: On moist potential temperatures and their ability to characterize differences in the properties of air parcels. *Accepted in the J. Atmos. Sci.*, **XX** (nn), ?–?, URL <https://arxiv.org/abs/2104.01376>.
- Mellor, G. L., and T. Yamada, 1982: Development of a turbulence closure model for geophysical fluid problems. *Reviews of Geophysics*, **20** (4), 851–875, doi:<https://doi.org/10.1029/RG020i004p00851>.
- Millero, F. J., 1983: The thermodynamics of seawater, Part II. Thermochemical properties. *Ocean Science and Engineering.*, **8** (1), 1–40.
- Mrowiec, A. A., O. M. Pauluis, and F. Zhang, 2016: Isentropic analysis of a simulated hurricane. *J. Atmos. Sci.*, **73** (5), 1857–1870, doi:10.1175/JAS-D-15-0063.1.
- Nagle, J. F., 1966: Lattice statistics of hydrogen bonded crystals. i. the residual entropy of ice. *J. Math. Phys.*, **7** (8), 1484–1491.
- Normand, C. W. B., 1921: Wet bulb temperatures and the thermodynamics of the air. *Memoirs of the India Meteorological Department. Meteorological Office*, **23**, 1–21.
- Pauling, L., 1935: The structure and entropy of ice and of other crystals with some randomness of atomic arrangement. *J. Am. Chem. Soc.*, **57** (12), 2680–2684.
- Pauluis, O., A. Czaja, and R. Korty, 2010: The global atmospheric circulation in moist isentropic coordinates. *J. Climate*, **23** (11), 3077–3093, doi:10.1175/2009JCLI2789.1.
- Pauluis, O. M., 2016: The mean air flow as lagrangian dynamics approximation and its application to moist convection. *J. Atmos. Sci.*, **73** (11), 4407–4425, doi:10.1175/JAS-D-15-0284.1.

- Pauluis, O. M., A. Czaja, and R. Korty, 2008: The global atmospheric circulation on moist isentropes. *Science*, **321** (5892), 1075–1078, doi:10.1126/science.1159649.
- Pauluis, O. M., and A. A. Mrowiec, 2013: Isentropic analysis of convective motions. *J. Atmos. Sci.*, **70** (11), 3673–3688, doi:10.1175/JAS-D-12-0205.1.
- Pauluis, O. M., and F. Zhang, 2017: Reconstruction of thermodynamic cycles in a high-resolution of a hurricane. *J. Atmos. Sci.*, **74** (10), 3367–3381, doi:10.1175/JAS-D-16-0353.1.
- Persson, P. O. G., 1995: Simulations of the potential vorticity structure and budget of fronts 87 IOP8. *Q. J. R. Meteorol. Soc.*, **121** (525), 1041–1081, doi:https://doi.org/10.1002/qj.49712152506.
- Rivas Soriano, L. J., and E. L. García Díez, 1997: Effect of ice on the generation of a generalized potential vorticity. *J. Atmos. Sci.*, **54** (10), 1385–1387, doi:10.1175/1520-0469(1997)054<1385:EOIOTG>2.0.CO;2.
- Robie, R. A., B. S. Hemingway, and J. R. Fisher, 1978: *Thermodynamic properties of minerals and related substances at 298.15 K and 1 Bar (10⁵ Pascals) pressure and at higher temperatures*. U.S. Geological Survey Bulletin 1452, 456 Pp. United States Government Printing Office. Washington : 1978.
- Robitzsch, M., 1928: Äquivalenttemperatur und Äquivalentthermometer [Equivalent temperature and equivalent thermometer]. *Meteorologische Zeitschrift (45th year)*. *Zeitschrift der Österreichischen Gesellschaft für Meteorologie (Journal of the Austrian Meteorological Society)*, **63**, 313–315.
- Rossby, C.-G., 1932: Thermodynamics applied to air mass analysis. *Papers in Physical Oceanography and Meteorology*, **I** (3), 57 pp., URL <https://darchive.mblwhoilibrary.org/handle/1912/1139>, Cambridge, Massachusetts.
- Rossini, F. D., D. D. Wagman, W. H. Evans, S. Levine, and I. Jaffe, 1952: *Selected values of chemical thermodynamic properties*, 1268 Pp. U. S. Department of Commerce. National Bureau of Standards.
- Rotunno, R., and J. Klemp, 1985: On the rotation and propagation of simulated supercell thunderstorms. *J. Atmos. Sci.*, **42** (3), 271–292, doi:10.1175/1520-0469(1985)042<0271:OTRAPO>2.0.CO;2.
- Saunders, P. M., 1957: The thermodynamics of saturated air: A contribution to the classical theory. *Quart. J. Roy. Meteorol. Soc.*, **83** (357), 342–350, doi:10.1002/qj.49708335707.
- Schubert, W., 2004: A generalization of Ertel’s potential vorticity to a cloudy, precipitating atmosphere. *Meteorologische Zeitschrift*, **13** (6), 465–471, doi:10.1127/0941-2948/2004/0013-0465.
- Schubert, W., E. Ruprecht, R. Hertenstein, R. N. Ferreira, R. Taft, C. Rozoff, P. Ciesielski, and H.-C. Kuo, 2004: English translations of twenty-one of Ertel’s papers on geophysical fluid dynamics. *Meteorologische Zeitschrift*, **13** (6), 527–576, doi:10.1127/0941-2948/2004/0013-0527.
- Schubert, W. H., S. A. Hausman, M. Garcia, K. V. Ooyama, and H.-C. Kuo, 2001: Potential vorticity in a moist atmosphere. *J. Atmos. Sci.*, **58** (21), 3148–3157, doi:10.1175/1520-0469(2001)058<3148:PVIAMA>2.0.CO;2.

- Stevens, B., and P. Siebesma, 2020: Clouds as fluids. *Clouds and Climate*, A. Siebesma, S. Bony, C. Jakob, and B. Stevens, Eds., *Climate Science's Greatest Challenge*, Cambridge Univ. Press., UK, p.35–73 in 1–409, URL <https://doi.org/10.1017/9781107447738>.
- Tripoli, G. J., and W. R. Cotton, 1981: The use of ice-liquid water potential temperature as a thermodynamic variable in deep atmospheric models. *Mon. Wea. Rev.*, **5 (14)**, 1094–1102, doi:10.1175/1520-0493(1981)109<1094:TUOLLW>2.0.CO;2.
- von Bezold, W., 1888a: *On the thermo-dynamics of the atmosphere (first communication)*. In: *The Mechanics of the Earth Atmosphere, a collection of translations by Cleveland Abbe, 1891. Smithsonian Miscellaneous Collections. [Translated from the Proceedings of the Royal Prussian Academy of Science at Berlin (Sitzungsberichte der König. Preuss. Akademie der Wissenschaften zu Berlin), 485–522 April 26 (1888)]*, **Art. XV**, 212–242.
- von Bezold, W., 1888b: *On the thermo-dynamics of the atmosphere (second communication)*. In: *The Mechanics of the Earth Atmosphere, a collection of translations by Cleveland Abbe, 1891. Smithsonian Miscellaneous Collections. [Translated from the Proceedings of the Royal Prussian Academy of Science at Berlin (Sitzungsberichte der König. Preuss. Akademie der Wissenschaften zu Berlin), Vol. XLVI 1189–1206 November 15 (1888)]*, **Art. XVI**, 243–256.
- von Helmholtz, H., 1888: On atmospheric movements (first paper). In: *The Mechanics of the Earth Atmosphere, a collection of translations by Cleveland Abbe, 1891. Smithsonian Miscellaneous Collections. [Translated from the Proceedings of the Royal Prussian Academy of Science at Berlin (Sitzungsberichte der König. Preuss. Akademie der Wissenschaften zu Berlin), 647–663 (1888)]*, **Art. V**, 78–93.
- von Stüve, G., 1927: Potentielle und pseudopotentielle Temperatur [Potential and pseudopotential temperature]. *Beitr. Phys. Atm.*, **13**, 218–233.
- Wagman, D. D., W. H. Evans, I. Halow, B. Parker, S. M. Bailey, and R. H. Schumm, 1965: *Selected values of chemical thermodynamic properties. Part 1. Technical Note 270-1*, 124 Pp. Institute for Basic Standards. National Bureau of Standards, Washington, D.C., USA.
- Wagner, W., and A. Pruss, 2002: The IAPWS formulation 1995 for the thermodynamic properties of ordinary water substance for general and scientific use. *Journal of Physical and Chemical Reference Data*, **31 (2)**, 387–535, doi:<http://dx.doi.org/10.1063/1.1461829>, URL <http://scitation.aip.org/content/aip/journal/jpcrd/31/2/10.1063/1.1461829>.