

Causal Effect Identification in Cluster DAGs

Tara V. Anand^{*1}, Adèle H. Ribeiro^{*2}, Jin Tian³, Elias Bareinboim²

¹Department of Biomedical Informatics, Columbia University

²Department of Computer Science, Columbia University

³Department of Computer Science, Iowa State University

tara.v.anand@columbia.edu, adele@cs.columbia.edu, jtian@iastate.edu, eb@cs.columbia.edu

Abstract

Reasoning about the effect of interventions and counterfactuals is a fundamental task found throughout the data sciences. A collection of principles, algorithms, and tools has been developed for performing such tasks in the last decades (Pearl 2000). One of the pervasive requirements found throughout this literature is the articulation of assumptions, which commonly appear in the form of causal diagrams. Despite the power of this approach, there are significant settings where the knowledge necessary to specify a causal diagram over all variables is not available, particularly in complex, high-dimensional domains. In this paper, we introduce a new graphical modeling tool called *cluster DAGs* (for short, C-DAGs) that allows for the partial specification of relationships among variables based on limited prior knowledge, alleviating the stringent requirement of specifying a full causal diagram. A C-DAG specifies relationships between clusters of variables, while the relationships between the variables within a cluster are left unspecified, and can be seen as a graphical representation of an equivalence class of causal diagrams that share the relationships among the clusters. We develop the foundations and machinery for valid inferences over C-DAGs about the clusters of variables at each layer of Pearl’s Causal Hierarchy (Pearl and Mackenzie 2018; Bareinboim et al. 2020) - \mathcal{L}_1 (probabilistic), \mathcal{L}_2 (interventional), and \mathcal{L}_3 (counterfactual). In particular, we prove the soundness and completeness of d-separation for probabilistic inference in C-DAGs. Further, we demonstrate the validity of Pearl’s do-calculus rules over C-DAGs and show that the standard ID identification algorithm is sound and complete to systematically compute causal effects from observational data given a C-DAG. Finally, we show that C-DAGs are valid for performing counterfactual inferences about clusters of variables.

1 Introduction

One of the central tasks found in data-driven disciplines is to infer the effect of a treatment X on an outcome Y , which is formally written as the interventional distribution $P(Y|do(X = x))$, from observational (non-experimental) data collected from the phenomenon under investigation. These relations are considered essential in the construction of explanations and for making decisions about interventions that have never been implemented before (Pearl 2000;

Spirtes, Glymour, and Scheines 2000; Bareinboim and Pearl 2016; Peters, Janzing, and Schölkopf 2017).

Standard tools necessary for identifying the aforementioned do-distribution, such as d-separation, do-calculus (Pearl 1995), and the ID-algorithm (Tian and Pearl 2002a; Shpitser and Pearl 2006; Huang and Valtorta 2006; Lee, Correa, and Bareinboim 2019) take as input a combination of an observational distribution and a qualitative description of the underlying causal system, often articulated in the form of a causal diagram (Pearl 2000). However, specifying a causal diagram requires knowledge about the causal relationships among all pairs of observed variables, which is not always available in many real-world applications. This is especially true and acute in complex, high-dimensional settings, which curtails the applicability of causal inference theory and tools.

In the context of medicine, for example, electronic health records include data on lab tests, drugs, demographic information, and other clinical attributes, but medical knowledge is not yet advanced enough to lead to the construction of causal diagrams over all of these variables, limiting the use of the graphical approach to inferring causality (Kleinberg and Hripacsak 2011). In many cases, however, contextual or temporal information about variables is available, which may partially inform how these variables are situated in a causal diagram relative to other key variables. For instance, a data scientist may know that covariates occur temporally before a drug is prescribed or an outcome occurs. They may even suspect that some pre-treatment variables are causes of the treatment and the outcome variables. However, they may be uncertain about the relationships among each pair of covariates, or it may be burdensome to explicitly define them. Given that a misspecified causal diagram may lead to wrong causal conclusions, this issue raises the question of whether a coarser representation of the causal diagram, where no commitment is made to the relationship between certain variables, would still be sufficient to determine the causal effect of interest.

In this paper, our goal is to develop a framework for causal inferences in partially understood domains such as the medical domain discussed above. We will focus on formalizing the problem of causal effect identification considering that the data scientist does not have prior knowledge to fully specify a causal diagram over all pairs of variables. First, we formally define and characterize a novel class of graphs

^{*}These authors contributed equally.

called *cluster DAGs* (or C-DAG, for short), which will allow for encoding of partially understood causal relationships between variables in different abstracted clusters, representing a group of variables among which causal relationships are not understood or specified. Then, we develop the foundations and machinery for valid probabilistic and causal inferences, akin to Pearl’s d-separation and do-calculus for when such a coarser graphical representation of the system is provided based on the limited prior knowledge available. In particular, we follow Pearl’s Causal Hierarchy (Pearl and Mackenzie 2018; Bareinboim et al. 2020) and develop the machinery for inferences in C-DAGs at all three inferential layers – \mathcal{L}_1 (associational), \mathcal{L}_2 (interventional), and \mathcal{L}_3 (counterfactual). The results are fundamental first steps in terms of semantics and graphical conditions to perform probabilistic, interventional, and counterfactual inferences over clusters of variables.

Specifically, we outline our technical contributions below.

1. We introduce a new graphical modelling tool called *cluster DAGs* (or C-DAGs) over macro-variables representing clusters of variables where the relationships among the variables inside the clusters are left unspecified (*Definition 1*). Semantically, a C-DAG represents an equivalence class of all underlying graphs over the original variables that share the relationships among the clusters.
2. We show that a C-DAG is a (probabilistic) Bayesian Network (BN) over macro-variables and Pearl’s d-separation is sound and complete for extracting conditional independencies over macro-variables if the underlying graph over the original variables is a BN (*Theorems 1 and 2*).
3. We show that a C-DAG is a Causal Bayesian Network (CBN) over macro-variables and Pearl’s do-calculus is sound and complete for causal inferences about macro-variables in C-DAGs if the underlying graph over the original variables is a CBN (*Theorems 3, 4, and 5*). The results can be used to show that the ID-algorithm is sound and complete to systematically infer causal effects from the observational distribution and partial domain knowledge encoded as a C-DAG (*Theorem 6*).
4. We show that, assuming the underlying graph G is induced by an SCM \mathcal{M} , then there exists an SCM \mathcal{M}_C over macro-variables \mathbf{C} such that its induced causal diagram is G_C and it is equivalent to \mathcal{M} on statements about the macro-variables (*Theorem 7*). Therefore, the CTFID algorithm (Correa, Lee, and Bareinboim 2021) for the identification of nested counterfactuals from an arbitrary combination of observational and experimental distributions can be extended to the C-DAGs.

1.1 Related work

Since a group of variables may constitute a semantically meaningful entity, causal models over abstracted clusters of variables have attracted increasing attention for the development of more interpretable tools (Schölkopf et al. 2021; Shen, Choi, and Darwiche 2018). (Parviainen and Kaski 2016) studied the problem of, given a DAG, under what assumptions a DAG over macro-variables can represent the same conditional independence relations between

the macro-variables. Recent developments on causal abstraction have focused on the distinct problem of investigating mappings of a cluster of (micro-)variables to a single (macro-)variable, while preserving some causal properties (Chalupka, Perona, and Eberhardt 2015; Chalupka, Eberhardt, and Perona 2016; Rubenstein et al. 2017; Beckers and Halpern 2019). The result is a new structural causal model defined on a higher level of abstraction, but with causal properties similar to those in the low-level model.¹ Other related works include chain graphs (Lauritzen and Richardson 2002) and ancestral causal graphs (Zhang 2008) developed to represent collections of causal diagrams equivalent under certain properties. By contrast, our work proposes a new graphical representation of a class of compatible causal diagrams, representing limited causal knowledge when the full structural causal model is unknown.

Causal discovery algorithms can be an alternative for when prior knowledge is insufficient to fully delineate a causal diagram (Pearl 2000; Spirtes, Glymour, and Scheines 2000; Peters, Janzing, and Schölkopf 2017). However, in general, it is impossible to fully recover the causal diagram based solely on observational data, without making strong assumptions about the underlying causal model, including causal sufficiency (all variables have been measured), the form of the functions (e.g., linearity, additive noise), and the distributions of the error terms (e.g. Gaussian, non-Gaussian, etc) (Glymour, Zhang, and Spirtes 2019). Then, there are cases where a meaningful causal diagram cannot be learned and prior knowledge is necessary for its construction. Our work focuses on establishing a language and corresponding machinery to encode partial knowledge and infer causal effects over clusters, alleviating some challenges in causal modeling in high-dimensional settings.

2 Preliminaries

Notation. A single variable is denoted by a (non-boldface) uppercase letter X and its realized value by a small letter x . A boldfaced uppercase letter \mathbf{X} denotes a set (or a cluster) of variables. We use kinship relations, defined along the full edges in the graph, ignoring bidirected edges. We denote by $Pa(\mathbf{X})_G$, $An(\mathbf{X})_G$, and $De(\mathbf{X})_G$, the sets of parents, ancestors, and descendants in G , respectively. A vertex V is said to be *active* on a path relative to \mathbf{Z} if 1) V is a collider and V or any of its descendants are in \mathbf{Z} or 2) V is a non-collider and is not in \mathbf{Z} . A path p is said to be *active* given (or conditioned on) \mathbf{Z} if every vertex on p is active relative to \mathbf{Z} . Otherwise, p is said to be *inactive*. Given a graph G , \mathbf{X} and \mathbf{Y} are d-separated by \mathbf{Z} if every path between \mathbf{X} and \mathbf{Y} is inactive given \mathbf{Z} . We denote this d-separation by $(\mathbf{X} \perp\!\!\!\perp \mathbf{Y} \mid \mathbf{Z})_G$. The mutilated graph $G_{\overline{\mathbf{XZ}}}$ is the result of removing from G edges with an arrowhead into \mathbf{X} (e.g., $A \rightarrow \mathbf{X}$, $A \leftrightarrow \mathbf{X}$), and edges with a tail from \mathbf{Z} (e.g., $A \leftarrow \mathbf{Z}$).

Structural Causal Models (SCMs) Formally, an SCM \mathcal{M} is a 4-tuple $\langle \mathbf{U}, \mathbf{V}, \mathcal{F}, P(\mathbf{U}) \rangle$, where \mathbf{U} is a set of exogenous (latent) variables and \mathbf{V} is a set of endogenous (mea-

¹In (Beckers and Halpern 2019)’s notation, we investigate the case of a *constructive τ -abstraction* where the mapping τ only groups the low-level variables into high-level (cluster) variables.

	Level (Symbol)	Typical Activity	Typical Model	Typical Question
\mathcal{L}_1	Associational $P(y x)$	Seeing	BN	What is? How would seeing X change my belief in Y ?
\mathcal{L}_2	Interventional $P(y do(x), c)$	Doing	CBN	What if? What if I do X ?
\mathcal{L}_3	Counterfactual $P(y_x x', y')$	Imagining	SCM	Why? What if I had acted differently?

Table 1: The Ladder of Causation / Pearl’s Causal Hierarchy

sured) variables. \mathcal{F} is a collection of functions $\{f_i\}_{i=1}^{|\mathbf{V}|}$ such that each endogenous variable $V_i \in \mathbf{V}$ is a function $f_i \in \mathcal{F}$ of $\mathbf{U}_i \cup Pa(V_i)$, where $\mathbf{U}_i \subseteq \mathbf{U}$ and $Pa(V_i) \subseteq \mathbf{V} \setminus V_i$. The uncertainty is encoded through a probability distribution over the exogenous variables, $P(\mathbf{U})$. Each SCM \mathcal{M} induces a directed acyclic graph (DAG) with bidirected edges – or an acyclic directed mixed graph (ADMG) – $G(\mathbf{V}, \mathbf{E})$, known as a *causal diagram*, that encodes the structural relations among $\mathbf{V} \cup \mathbf{U}$, where every $V_i \in \mathbf{V}$ is a vertex, there is a directed edge $(V_j \rightarrow V_i)$ for every $V_i \in \mathbf{V}$ and $V_j \in Pa(V_i)$, and there is a dashed bidirected edge $(V_j \leftrightarrow V_i)$ for every pair $V_i, V_j \in \mathbf{V}$ such that $\mathbf{U}_i \cap \mathbf{U}_j \neq \emptyset$ (V_i and V_j have a common exogenous parent). Performing an intervention $\mathbf{X} = \mathbf{x}$ is represented through the do-operator, $do(\mathbf{X} = \mathbf{x})$, which represents the operation of fixing a set \mathbf{X} to a constant \mathbf{x} , and induces a submodel $\mathcal{M}_{\mathbf{x}}$, which is \mathcal{M} with f_X replaced to x for every $X \in \mathbf{X}$. The post-interventional distribution induced by $\mathcal{M}_{\mathbf{x}}$ is denoted by $P(\mathbf{v} \setminus \mathbf{x} | do(\mathbf{x}))$. For any subset $\mathbf{Y} \subseteq \mathbf{V}$, the *potential response* $\mathbf{Y}_{\mathbf{x}}(\mathbf{u})$ is defined as the solution of \mathbf{Y} in the submodel $\mathcal{M}_{\mathbf{x}}$ given $\mathbf{U} = \mathbf{u}$. $P(\mathbf{U})$ then induces a *counterfactual variable* $\mathbf{Y}_{\mathbf{x}}$.

Pearl’s Causal Hierarchy (PCH) / The Ladder of Causation (Pearl and Mackenzie 2018; Bareinboim et al. 2020) is a formal framework that divides inferential tasks into three different layers, namely, 1) associational, 2) interventional, and 3) counterfactual (see Table 1). An important result formalized under the rubric of the Causal Hierarchy Theorem (CHT) (Bareinboim et al. 2020, Thm. 1) states that inferences at any layer of the PCH almost never can be obtained by using solely information from lower layers.

3 C-DAGs: Definition and Properties

Standard causal inference tools typically require assumptions articulated through causal diagrams. We investigate the situations where the knowledge necessary to specify the underlying causal diagram $G(\mathbf{V}, \mathbf{E})$ over the individual variables in \mathbf{V} may not be available. In particular, we assume that variables are grouped into a set of clusters of variables $\mathbf{C}_1, \dots, \mathbf{C}_k$ that form a partition of \mathbf{V} (note that a variable may be grouped in a cluster by itself) such that we do not have knowledge about the relationships amongst the variables inside the clusters \mathbf{C}_i but we have some knowledge about the relationships between variables in different groups. We are interested in performing probabilistic and causal inferences about these clusters of variables; one may consider each cluster as defining a *macro-variable* and our aim is to reason about these macro-variables.




Figure 1: (a): a possible ADMG over lisinopril (X), stroke (Y), age (A), blood pressure (B), comorbidities (C), medication history (D), and sleep quality (S). (b): a C-DAG of (a) with $\mathbf{Z} = \{A, B, C, D\}$. (c): a C-DAG of (a) with $\mathbf{W} = \{S, B\}$, $\mathbf{Z} = \{A, C\}$. (d): an invalid C-DAG of (a), as the partition $\{X, Y, \mathbf{W}, \mathbf{Z}\}$, with $\mathbf{W} = \{S, B\}$, $\mathbf{Z} = \{A, C, D\}$ is inadmissible due to the cycle among $(X, \mathbf{W}, \mathbf{Z})$.

Formally, we address the following technical problem:

Problem Statement: Consider a causal diagram G^* over \mathbf{V} and a set of clusters of variables $\mathbf{C} = \{\mathbf{C}_1, \dots, \mathbf{C}_k\}$ forming a partition of \mathbf{V} . We aim to perform probabilistic, interventional, or counterfactual inferences about the macro-variables. Can we construct a causal diagram $G_{\mathbf{C}}^*$ over the macro-variables in \mathbf{C} such that inferences by applying standard tools (d-separation, do-calculus, ID algorithm) on $G_{\mathbf{C}}^*$ are valid in the sense they lead to the same conclusions as inferred on G^* ?

To this end, we propose a graphical object called *cluster DAGs* (or C-DAGs) to capture our partial knowledge about the underlying causal diagram over individual variables.

Definition 1 (Cluster DAG or C-DAG). Given an ADMG $G(\mathbf{V}, \mathbf{E})$ (a DAG with bidirected edges) and a partition $\mathbf{C} = \{\mathbf{C}_1, \dots, \mathbf{C}_k\}$ of \mathbf{V} , construct a graph $G_{\mathbf{C}}(\mathbf{C}, \mathbf{E}_{\mathbf{C}})$ over \mathbf{C} with a set of edges $\mathbf{E}_{\mathbf{C}}$ defined as follows:

1. An edge $\mathbf{C}_i \rightarrow \mathbf{C}_j$ is in $\mathbf{E}_{\mathbf{C}}$ if exists some $V_i \in \mathbf{C}_i$ and $V_j \in \mathbf{C}_j$ such that $V_i \in Pa(V_j)$ in G ;
2. A dashed bidirected edge $\mathbf{C}_i \leftrightarrow \mathbf{C}_j$ is in $\mathbf{E}_{\mathbf{C}}$ if exists some $V_i \in \mathbf{C}_i$ and $V_j \in \mathbf{C}_j$ such that $V_i \leftrightarrow V_j$ in G .

If $G_{\mathbf{C}}(\mathbf{C}, \mathbf{E}_{\mathbf{C}})$ contains no cycles, then we say that \mathbf{C} is an *admissible partition* of \mathbf{V} . We then call $G_{\mathbf{C}}$ a *cluster DAG*, or *C-DAG*, compatible with G .

Throughout the paper, we will use the same symbols (e.g. \mathbf{C}_i) to represent both a cluster node in a C-DAG $G_{\mathbf{C}}$ and the set of variables contained in the cluster.

Remark 1. The definition of C-DAGs does not allow for cycles in order to utilize standard graphical modeling tools that work only in DAGs. An inadmissible partition of \mathbf{V} means that the partial knowledge available for constructing $G_{\mathbf{C}}$ is not enough for drawing conclusions using the tools developed in this paper.

Remark 2. Although a C-DAG is defined in terms of an underlying graph G , in practice, one will construct a C-DAG when complete knowledge of the graph G is unavailable. As an example of this construction, consider the model of the effect of lisinopril (X) on the outcome of having a stroke (Y) in Fig. 1(a). If not all the relationships specified in Fig. 1(a) are known, a data scientist cannot construct a full causal diagram, but may still have enough knowledge to create a C-DAG. For instance, they may have partial knowledge that the




Figure 2: G_{C_1} is the C-DAG for diagrams (a) and (b) and G_{C_2} is the C-DAG for diagrams (c) and (d), where $\mathbf{Z} = \{Z_1, Z_2, Z_3\}$. $P(y|do(x))$ is identifiable in G_{C_1} by backdoor adjustment over \mathbf{Z} and is not identifiable in G_{C_2} .

covariates occur temporally before lisinopril is prescribed, or that a stroke occurs and the suspicion that some of the pre-treatment variables are causes of X and Y . Specifically, they can create the cluster $\mathbf{Z} = \{A, B, C, D\}$ with all the covariates, and then construct a C-DAG with edges $\mathbf{Z} \rightarrow X$ and $\mathbf{Z} \rightarrow Y$. Further, the data scientist may also suspect that some of the variables in \mathbf{Z} are confounded with X and others with Y , an uncertainty that is encoded in the C-DAG through the bidirected edges $\mathbf{Z} \leftrightarrow X$ and $\mathbf{Z} \leftrightarrow Y$. With the additional knowledge that sleep quality (S) acts as a mediator between the treatment and outcome, the C-DAG in Fig. 1(b) can be constructed. Note that this C-DAG is consistent with the true underlying causal diagram in Fig. 1(a), but was constructed without knowing this diagram and using much less knowledge than what is encoded in it. Alternatively, if clusters $\mathbf{W} = \{S, B\}$ and $\mathbf{Z} = \{A, C\}$ are created, then the C-DAG shown in Fig. 1(c) would be constructed. Note that both (a) and (b) are considered valid C-DAGs because no cycles are created. Finally, if a clustering with $\mathbf{W} = \{S, B\}$ and $\mathbf{Z} = \{A, C, D\}$ is created, this would lead to the C-DAG shown in Fig. 1(d), which is invalid. The reason is that a cycle $X \rightarrow \mathbf{W} \rightarrow \mathbf{Z} \rightarrow X$ is created due to the connections $X \rightarrow S$, $B \rightarrow C$, and $D \rightarrow X$ in the diagram (a).

Remark 3. It is important to note that a C-DAG G_C as defined in Def. 1 is merely a graph over clusters of nodes C_1, \dots, C_k , and does not have a priori the semantics and properties of a BN or CBN over macro-variables C_i . It's not clear, for example, whether the cluster nodes C_i satisfy the Markov properties w.r.t. the graph G_C . Rather, a C-DAG can be seen as a graphical representation of an equivalence class (EC, for short) of graphs that share the relationships among the clusters while allowing for any possible relationships among the variables within each cluster. For instance, in Fig. 2, the diagrams (a) and (b) can be represented by C-DAG G_{C_1} (top) and can, therefore, be thought of as being members of an EC represented by G_{C_1} . The same can be concluded for diagrams (c) and (d), both represented by C-DAG G_{C_2} . The graphical representation of this ECs are shown in Fig. 3, where on the left we have the space of all possible ADMGs, and on the right the space of C-DAGs.




Figure 3: Identifying $P(y|do(x))$ in a C-DAG means identifying such an effect for the entire equivalence class. In G_{C_1} , the effect is identifiable (blue) because it is identifiable in $G_{(a)}$, $G_{(b)}$, and all the other causal diagrams represented. In G_{C_2} , the same effect is non-identifiable (red), as the encoded partial knowledge is compatible with some causal diagrams of C-DAGs where the effect is not identifiable (e.g., $G_{(d)}$).

Given the semantics of a C-DAG as an equivalence class of ADMGs, what valid inferences can one perform about the cluster variables given a C-DAG G_C ? What properties of C-DAGs are shared by all compatible ADMGs?

In principle, definite conclusions from C-DAGs can only be drawn about properties shared among all EC's members. Going back to Fig. 3, we identify an effect in a C-DAG (e.g., G_{C_1} in Fig. 2) whenever this effect is identifiable in all members of the EC; e.g., causal diagrams (a), (b), and all other diagrams compatible with G_{C_1} . Note that in this particular EC, all dots are marked with blue, which means that the effect is identifiable in each one of them. On the other hand, if there exists one diagram in the EC where the effect is not identifiable, this effect will not be identifiable in the corresponding C-DAG. For instance, the effect is not identifiable in the C-DAG G_{C_2} due to diagram (d) in Fig. 2.

Once the semantics of C-DAGs is well-understood, we turn our attention to computational issues. One naive approach to causal inference with cluster variables, e.g. identifying $Q = P(C_i|do(C_k))$, goes as follows – first enumerate all causal diagrams compatible with G_C ; then, evaluate the identifiability of Q in each diagram; finally, output $P(C_i|do(C_k))$ if all the diagrams entail the same answer, otherwise output “non-identifiable”. However, in practice, this approach is intractable in high-dimensional settings – given a cluster C_i of size m , the number of possible causal diagrams over the variables in C_i is super-exponential in m . Can valid inferences be performed about cluster variables using C-DAGs directly, without going through exhaustive enumeration? What properties of C-DAGs are shared by all the compatible causal diagrams? The next sections present theoretical results to address these questions.

Finally, note that not all properties of C-DAGs are shared across all compatible diagrams. To illustrate, consider the case of *backdoor paths*, i.e., paths between X and Y with an arrowhead into X , in Fig. 2. The path $X \leftrightarrow \mathbf{Z} \rightarrow Y$

in G_{C_2} is active when not conditioning on \mathbf{Z} . However, the corresponding backdoor paths in diagram (c) are all inactive. Therefore, a d-connection in a C-DAG does not necessarily correspond to a d-connection in all diagrams in the EC.

4 C-DAGs for \mathcal{L}_1 -Inferences

In this section, we study probabilistic inference with C-DAGs - \mathcal{L}_1 inferences. We assume the underlying graph G over \mathbf{V} is a Bayesian Network (BN) *with no causal interpretation*.² We aim to perform probabilistic inferences about macro-variables with G_C that are valid in G regardless of the unknown relationships within each cluster.

First, we extend d-separation (Pearl 1988), a fundamental tool in probabilistic reasoning in BNs, to C-DAGs. As noticed earlier, a d-connecting path in a C-DAG does not necessarily imply that the corresponding paths in a compatible ADMG G are connecting. Such paths can be either active or inactive. However, d-separated paths in a C-DAG correspond to only d-separated paths in all compatible ADMGs.³ These observations together, lead to the following definition where the symbol $*$ represents either an arrow head or tail:

Definition 2 (d-Separation in C-DAGs). A path p in a C-DAG G_C is said to be d-separated (or blocked) by a set of clusters $\mathbf{Z} \subset \mathbf{C}$ if and only if p contains a triplet

1. $C_i * \text{---} C_m \rightarrow C_j$ such that the non-collider cluster C_m is in \mathbf{Z} , or
2. $C_i * \rightarrow C_m \leftarrow * C_j$ such that the collider cluster C_m and its descendants are not in \mathbf{Z} .

A set of clusters \mathbf{Z} is said to d-separate two sets of clusters $\mathbf{X}, \mathbf{Y} \subset \mathbf{C}$, denoted by $(\mathbf{X} \perp\!\!\!\perp \mathbf{Y} \mid \mathbf{Z})_{G_C}$, if and only if \mathbf{Z} blocks every path from a cluster in \mathbf{X} to a cluster in \mathbf{Y} .

We show in the following proposition that the d-separation rules are sound and complete in C-DAGs in the following sense: whenever a d-separation holds in a C-DAG, it holds for all ADMGs compatible with it; on the other hand, if a d-separation does not hold in a C-DAG, then there exists at least one ADMG compatible with it for which the same d-separation statement does not hold.

Theorem 1. (Soundness and completeness of d-separation in C-DAGs). In a C-DAG G_C , let $\mathbf{X}, \mathbf{Z}, \mathbf{Y} \subset \mathbf{C}$. If \mathbf{X} and \mathbf{Y} are d-separated by \mathbf{Z} in G_C , then, in any ADMG G compatible with G_C , \mathbf{X} and \mathbf{Y} are d-separated by \mathbf{Z} in G , i.e.,

$$(\mathbf{X} \perp\!\!\!\perp \mathbf{Y} \mid \mathbf{Z})_{G_C} \implies (\mathbf{X} \perp\!\!\!\perp \mathbf{Y} \mid \mathbf{Z})_G. \quad (1)$$

If \mathbf{X} and \mathbf{Y} are not d-separated by \mathbf{Z} in G_C , then, there exists an ADMG G compatible with G_C where \mathbf{X} and \mathbf{Y} are not d-separated by \mathbf{Z} in G .

Theorem 1 implies that G_C does not imply any conditional independence that is not implied by the underlying G .

ADMGs are commonly used to represent BNs with latent variables (which may imply Verma constraints on $P(v)$)

²For a more detailed discussion on the tension between layers \mathcal{L}_1 and \mathcal{L}_2 , please refer to (Bareinboim et al. 2020, Sec. 1.4.1).

³In Appendix A.1, we investigate in detail how path analysis is extended to C-DAGs. We note that d-separation in ADMGs has also been called m-separation (Richardson 2003).

not captured by independence relationships (Tian and Pearl 2002b)) where the observational distribution $P(\mathbf{v})$ factorizes according to the ADMG G as follows

$$P(\mathbf{v}) = \sum_{\mathbf{u}} P(\mathbf{u}) \prod_{k: V_k \in \mathbf{V}} P(v_k | pa_{V_k}, \mathbf{u}_k), \quad (2)$$

where Pa_{V_k} are the endogenous parents of V_k in G and $\mathbf{U}_k \subseteq \mathbf{U}$ are the latent parents of V_k . We show next that the observational distribution $P(\mathbf{v})$ factorizes according to the graphical structure of the C-DAG G_C as well.

Theorem 2. (C-DAG as BN). Let G_C be a C-DAG compatible with an ADMG G . If the observational distribution $P(\mathbf{v})$ factorizes according to G by Eq. (2), then the observational distribution $P(\mathbf{v}) = P(\mathbf{c})$ factorizes according to G_C , i.e.,

$$P(\mathbf{c}) = \sum_{\mathbf{u}} P(\mathbf{u}) \prod_{k: \mathbf{C}_k \in \mathbf{C}} P(\mathbf{c}_k | pa_{\mathbf{C}_k}, \mathbf{u}'_k), \quad (3)$$

where $Pa_{\mathbf{C}_k}$ are the parents of the cluster \mathbf{C}_k , and $\mathbf{U}'_k \subseteq \mathbf{U}$ such that, for any i, j , $\mathbf{U}'_i \cap \mathbf{U}'_j \neq \emptyset$ if and only if there is a bidirected edge $(\mathbf{C}_i \leftrightarrow \mathbf{C}_j)$ between \mathbf{C}_i and \mathbf{C}_j in G_C .

Thm. 2 implies that if the underlying ADMG G represents a BN with latent variables over \mathbf{V} , then the C-DAG G_C represents a BN over micro-variables \mathbf{C} .

5 C-DAGs for \mathcal{L}_2 -Inferences

We study now interventional (\mathcal{L}_2) reasoning with C-DAGs. We assume the underlying graph G over \mathbf{V} is a CBN. Our goal is to perform causal reasoning about macro-variables with G_C that are guaranteed to be valid in each G of the underlying EC. We focus on extending to C-DAGs Pearl's celebrated *do-calculus* (Pearl 1995) and the ID-algorithm (Tian 2002; Shpitser and Pearl 2006; Huang and Valtorta 2006).

Do-Calculus in C-DAGs

Do-calculus is a fundamental tool in causal inference from causal diagrams and has been used extensively for solving a variety of identification tasks. We show that if the underlying ADMG G is a CBN on which do-calculus rules hold, then do-calculus rules are valid in the corresponding C-DAG G_C . We first present a key lemma for proving the soundness of do-calculus in C-DAGs that the mutilation operations in a C-DAG to create $G_{C_{\mathbf{X}}}$ and $G_{C_{\mathbf{X}}}$ carry over to all compatible underlying ADMGs. This result is shown in the following:

Lemma 1. If a C-DAG G_C is compatible with an ADMG G , then, for $\mathbf{X}, \mathbf{Z} \subset \mathbf{C}$, the mutilated C-DAG $G_{C_{\mathbf{X}\mathbf{Z}}}$ is compatible with the mutilated ADMG $G_{\mathbf{X}\mathbf{Z}}$.

The soundness of do-calculus in C-DAGs as stated next follows from Theorem 1 and Lemma 1.

Theorem 3. (Do-calculus in causal C-DAGs). Let G_C be a C-DAG compatible with an ADMG G . If G is a CBN encoding interventional distributions $P(\cdot | do(\cdot))$, then for any disjoint subsets of clusters $\mathbf{X}, \mathbf{Y}, \mathbf{Z}, \mathbf{W} \subset \mathbf{C}$, the following three rules hold:

$$\begin{aligned} \text{Rule 1: } P(\mathbf{y} | do(\mathbf{x}), \mathbf{z}, \mathbf{w}) &= P(\mathbf{y} | do(\mathbf{x}), \mathbf{w}) \\ &\text{if } (\mathbf{Y} \perp\!\!\!\perp \mathbf{Z} | \mathbf{X}, \mathbf{W})_{G_{C_{\mathbf{X}\mathbf{Z}}}} \end{aligned}$$

Rule 2: $P(y|do(x), do(z), w) = P(y|do(x), z, w)$
if $(Y \perp\!\!\!\perp Z|X, W)_{G_{C_{\overline{XZ}}}}$

Rule 3: $P(y|do(x), do(z), w) = P(y|do(x), w)$
if $(Y \perp\!\!\!\perp Z|X, W)_{G_{C_{\overline{XZ}(W)}}$

where $G_{C_{\overline{XZ}}}$ is obtained from G_C by removing the edges into X and out of Z , and $Z(W)$ is the set of Z -clusters that are non-ancestors of any W -cluster in $G_{C_{\overline{XZ}}}$.

We also show next that the do-calculus rules in C-DAGs are complete in the following sense:

Theorem 4. (Completeness of do-calculus). *If in a C-DAG G_C a do-calculus rule does not apply, then there is a CBN G compatible with G_C for which it also does not apply.*

Truncated Factorization

An ADMG G represents a CBN if the interventional distributions factorizes according to the graphical structure, known as the truncated factorization, i.e., for any $X \subseteq V$

$$P(v \setminus x|do(x)) = \sum_{\mathbf{u}} P(\mathbf{u}) \prod_{k: V_k \in V \setminus X} P(v_k|pa_{v_k}, \mathbf{u}_k), \quad (4)$$

where pa_{V_k} are the endogenous parents of V_k in G and $U_k \subseteq U$ are the latent parents of V_k .

We show that the truncated factorization holds in C-DAGs as if the underlying ADMG is a CBN, in the following sense.

Theorem 5. (C-DAG as CBN) *Let G_C be a C-DAG compatible with an ADMG G . If G satisfies the truncated factorization (4) with respect to the interventional distributions, then, for any $X \subseteq C$, the interventional distribution $P(c \setminus x|do(x))$ factorizes according to G_C , i.e.,*

$$P(c \setminus x|do(x)) = \sum_{\mathbf{u}} P(\mathbf{u}) \prod_{k: C_k \in C \setminus X} P(c_k|pa_{C_k}, \mathbf{u}'_k), \quad (5)$$

where pa_{C_k} are the parents of the cluster C_k , and $U'_k \subseteq U$ such that, for any i, j , $U'_i \cap U'_j \neq \emptyset$ if and only if there is a bidirected edge $(C_i \leftrightarrow C_j)$ between C_i and C_j in G_C .

Theorem 5 essentially shows that a C-DAG G_C can be treated as a CBN over the macro-variables C if the underlying ADMG is a CBN.

ID-Algorithm

Equipped with d-separation, do-calculus, and the truncated factorization in C-DAGs, causal inference algorithms developed for a variety of tasks that rely on a known causal diagram can be extended to C-DAGs (Bareinboim and Pearl 2016). In this paper, we consider the problem of identifying causal effects from observational data using C-DAGs.

There exists a complete algorithm to determine whether $P(y|do(x))$ is identifiable from a causal diagram G and the observational distribution $P(V)$ (Tian 2002; Shpitser and Pearl 2006; Huang and Valtorta 2006). This algorithm, or ID-algorithm for short, is based on the truncated factorization, therefore, Theorem 5 allows us to prove that the ID-algorithm is sound and complete to systematically infer causal effects from the observational distribution $P(V)$ and partial domain knowledge encoded as a C-DAG G_C .

Theorem 6. (Soundness and Completeness of ID-algorithm). *The ID-algorithm is sound and complete when applied to a C-DAG G_C for identifying causal effects of the form $P(y|do(x))$ from the observational distribution $P(V)$, where X and Y are sets of clusters in G_C .*

The ID algorithm returns a formula for identifiable $P(y|do(x))$ that is valid in all causal diagrams compatible with the C-DAG G_C . The completeness result ensures that if the ID-algorithm fails to identify $P(y|do(x))$ from G_C , then there exists a causal diagram G compatible with G_C where the effect $P(y|do(x))$ is not identifiable. Appendix A.3 contains an experimental study evaluating the ability of C-DAGs to accurately assess the identifiability of effects while requiring less domain knowledge for their construction.

Examples of Causal Identifiability in C-DAGs

We show examples of identification in C-DAGs in practice. Besides illustrating identification of causal effects in the coarser graphical representation of a C-DAG, these examples demonstrate that clustering variables may lead to diagrams where effects are not identifiable. Therefore, care should be taken when clustering variables, to ensure not so much information is lost in a resulting C-DAG, such that identifiability is maintained when possible.

Identification in Fig. 1. In diagram (a) the effect of X on Y is identifiable through backdoor adjustment (Pearl 2000, pp. 79-80) over the set of variables $\{B, D\}$. In the C-DAG in Fig. 1(b), with cluster $Z = \{A, B, C, D\}$, the effect of X on Y is identifiable through front-door adjustment (Pearl 2000, p. 83) over S , given by $P(y|do(x)) = \sum_s P(s|x) \sum_{x'} P(y|x', s) P(x')$. Because this front-door adjustment holds for the C-DAG in Fig. 1(b) with which diagram (a) is compatible, this front-door adjustment identification formula is equivalent to the adjustment in the case of diagram (a) and gives the correct causal effect in any other compatible causal diagram. In the C-DAG in (c), the loss of separations from the creation of clusters $Z = \{A, B, C, D\}$ and $W = \{B, S\}$ render the effect no longer identifiable, indicating that there exists another graph compatible with (c) for which the effect cannot be identified.




Figure 4: (a): causal diagram G where the effect $P(y_1, y_2|do(x_1, x_2))$ is identifiable. (b): C-DAG G_{C_1} with clustering $X = \{X_1, X_2\}$, $Y = \{Y_1, Y_2\}$, and $Z = \{Z_1, Z_2\}$. (c): C-DAG G_{C_2} with clustering $X = \{X_1, X_2\}$ and $Y = \{Y_1, Y_2\}$. (d): C-DAG G_{C_3} with clustering $Y = \{Y_1, Y_2\}$ and $Z = \{Z_1, Z_2\}$. The effect $P(y|do(x))$ is not identifiable in G_{C_1} , but is identifiable in G_{C_2} and $P(y|do(x_1, x_2))$ is identifiable in G_{C_3} .

Identification in Fig. 4. In causal diagram (a), the ef-

fect of $\{X_1, X_2\}$ on $\{Y_1, Y_2\}$ is identifiable by backdoor adjustment over $\{Z_1, Z_2\}$ as follows: $P(y_1, y_2 | do(x_1, x_2)) = \sum_{z_1, z_2} P(y_1, y_2 | x_1, x_2, z_1, z_2) P(z_1, z_2)$. Note, however, that the backdoor path cannot be blocked in the C-DAG G_1 (b) with clusters $\mathbf{X} = \{X_1, X_2\}$, $\mathbf{Y} = \{Y_1, Y_2\}$, and $\mathbf{Z} = \{Z_1, Z_2\}$. In this case, the effect $P(y | do(\mathbf{x}))$ is not identifiable. If the covariates Z_1 and Z_2 are not clustered together as shown in the C-DAG G_{C_2} (c), the backdoor paths relative to \mathbf{X} and \mathbf{Y} can still be blocked despite the unobserved confounders between Z_1 and \mathbf{X} and between Z_2 and \mathbf{Y} . So the effect $P(y | do(\mathbf{x}))$ is identifiable by backdoor adjustment over $\{Z_1, Z_2\}$ as follows: $P(y | do(\mathbf{x})) = \sum_{z_1, z_2} P(y | \mathbf{x}, z_1, z_2) P(z_1, z_2)$. If the treatments X_1 and X_2 are not clustered together as shown in the C-DAG G_{C_3} (d), then the joint effect of X_1 and X_2 on the cluster \mathbf{Y} is identifiable and given by the following expression: $P(y | do(x_1, x_2)) = \sum_{z, x'_1} P(y | x'_1, x_2, z) P(x'_1, z)$.




Figure 5: (a), (b), and (c) are causal diagrams compatible with the C-DAG G_C in (d) where $\mathbf{X} = \{X_1, X_2\}$ and $\mathbf{Y} = \{Y_1, Y_2\}$. The causal effect $P(y_1, y_2 | do(x_1, x_2))$ is identifiable in (a) but not in (b) or (c). Consequently, the effect $P(y | do(\mathbf{x}))$ is not identifiable from the C-DAG G_C .

Identification in Fig. 5. In the causal diagram (a), the effect of the joint intervention to $\{X_1, X_2\}$ on both outcomes $\{Y_1, Y_2\}$ is identifiable as follows: $P(y_1, y_2 | do(x_1, x_2)) = P(y_1 | x_1, x_2) \sum_{x'_1} P(y_2 | x'_1, x_2, y_1) P(x'_1)$. By clustering the two treatments as \mathbf{X} and the two outcomes as \mathbf{Y} , we lose the information that X_2 is not a confounded effect of X_1 and that Y_1 and Y_2 are not confounded. If this is the case, as in causal diagrams G_2 (b) and G_3 (c), the effect would not be identifiable. Note that the C-DAG (d), representing causal diagrams (a), (b), and (c), is the bow graph, where the effect $P(y | do(\mathbf{x}))$ is also not identifiable.

6 C-DAGs for \mathcal{L}_3 -Inferences

Now, we study counterfactual (\mathcal{L}_3) inferences in C-DAGs. We assume that the underlying graph G over \mathbf{V} is induced by an SCM \mathcal{M} , and our goal is to perform counterfactual reasoning about macro-variables with G_C that are always valid in \mathcal{M} (while both G and the SCM \mathcal{M} are unknown).

We show that for any SCM over \mathbf{V} with causal diagram G , there is an equivalent SCM over macro-variables \mathbf{C} that induces C-DAG G_C and makes the same predictions about counterfactual distributions over the macro-variables.

Theorem 7. *Let G_C be a C-DAG compatible with an ADMG G . Assume G is induced by an SCM \mathcal{M} , then there exists an SCM \mathcal{M}_C over macro-variables \mathbf{C} such that its induced causal diagram is G_C and, for any set of counterfactual variables $\mathbf{Y}_x, \dots, \mathbf{Z}_w$ where $\mathbf{Y}, \mathbf{X}, \dots, \mathbf{Z}, \mathbf{W} \subseteq \mathbf{C}$,*

$$P_{\mathcal{M}}(\mathbf{y}_x, \dots, \mathbf{z}_w) = P_{\mathcal{M}_C}(\mathbf{y}_x, \dots, \mathbf{z}_w).$$

Following this result, algorithms developed for a variety of counterfactual inference tasks that rely on a known causal diagram, such as the CTFID algorithm (Correa, Lee, and Bareinboim 2021), can be used in the context of C-DAGs.

For example, consider the C-DAG G_{C_1} in Fig. 2, where X is a drug, Y is a disease, and \mathbf{Z} is a cluster of factors potentially affecting X and Y . Suppose that a patient who took the drug ($X = 1$) would like to know what his chances of being cured ($Y = 1$) would have been had he not taken the drug ($X = 0$). This quantity is defined by $P(Y_{X=0} = 1 | X = 1)$. The CTFID algorithm applied to G_{C_1} will conclude that $P(Y_{X=0} = 1 | X = 1)$ is identifiable and given by $\sum_z P(Y = 1 | X = 0, Z = z) P(Z = z | X = 1)$. This formula is correct in all ADMGs compatible with G_{C_1} , regardless of the relationships within cluster \mathbf{Z} .

After all, we note that inferences in the lower layers assume less knowledge than the higher layers. On the one hand, some results about the lower layers are implied by the higher layers. For instance, if G is induced by an SCM, then G represents a CBN and a BN, and Thm. 7 implies that if the underlying G is induced by an SCM, then G_C represents a CBN and a BN. If G is a CBN, then it is necessarily a BN, therefore Thm. 5 implies that if the underlying G represents a CBN, then G_C represents a BN. On the other hand, if one does not want to commit to the SCM generative process, but can only ascertain that the truncated factorization holds (e.g., for \mathcal{L}_2), it's still possible to leverage the machinery developed without any loss of inferential power or making unnecessary assumptions about the upper layers.

7 Conclusions

Causal diagrams provide an intuitive language for specifying the necessary assumptions for causal inferences. Despite all their power and successes, the substantive knowledge required to construct a causal diagram – i.e., the causal and confounded relationships among all pairs of variables – is unattainable in some critical settings found across society, including in the health and social sciences. This paper introduces a new class of graphical models that allow for a more relaxed encoding of knowledge. In practice, when a researcher does not fully know the relationships among certain variables, under some mild assumptions delineated by Def. 1, these variables can be clustered together. (A causal diagram is an extreme case of a C-DAG where each cluster has exactly one variable.) We prove fundamental results to allow causal inferences within C-DAG's equivalence class, which translate to statements about all diagrams compatible with the encoded constraints. We develop the formal machinery for probabilistic, interventional, and counterfactual reasoning in C-DAGs following Pearl's hierarchy assuming the (unknown) underlying model over individual variables are BN (\mathcal{L}_1), CBN (\mathcal{L}_2), and SCM (\mathcal{L}_3), respectively. These results are critical for enabling C-DAGs use in ways comparable to causal diagrams. We hope these new tools will allow researchers to represent complex systems in a simplified way, allowing for more relaxed causal inferences when substantive knowledge is largely unavailable and coarse.

Acknowledgements

This work was done in part while Jin Tian was visiting the Simons Institute for the Theory of Computing. Jin Tian was partially supported by NSF grant IIS-2231797. This research was supported by the NSF, ONR, AFOSR, DoE, Amazon, JP Morgan, The Alfred P. Sloan Foundation and the United States NLM T15LM007079.

References

- Bareinboim, E.; Correa, J. D.; Ibeling, D.; and Icard, T. 2020. On Pearl's Hierarchy and the Foundations of Causal Inference. Technical Report R-60, Causal Artificial Intelligence Lab, Columbia University.
- Bareinboim, E.; and Pearl, J. 2016. Causal inference and the data-fusion problem. *Proceedings of the National Academy of Sciences*, 113(27): 7345–7352.
- Beckers, S.; and Halpern, J. Y. 2019. Abstracting Causal Models. In *Proceedings of the Thirty-Third AAAI Conference on Artificial Intelligence*, AAAI'19.
- Chalupka, K.; Eberhardt, F.; and Perona, P. 2016. Multi-Level Cause-Effect Systems. In Gretton, A.; and Robert, C. C., eds., *Proceedings of the 19th International Conference on Artificial Intelligence and Statistics*, volume 51 of *Proceedings of Machine Learning Research*, 361–369. Cadiz, Spain: PMLR.
- Chalupka, K.; Perona, P.; and Eberhardt, F. 2015. Visual Causal Feature Learning. In *Proceedings of the Thirty-First Conference on Uncertainty in Artificial Intelligence*, UAI'15, 181–190. Arlington, Virginia, USA: AUAI Press.
- Correa, J.; Lee, S.; and Bareinboim, E. 2021. Nested Counterfactual Identification from Arbitrary Surrogate Experiments. In Ranzato, M.; Beygelzimer, A.; Dauphin, Y.; Liang, P. S.; and Vaughan, J. W., eds., *Advances in Neural Information Processing Systems*, volume 34, 6856–6867. Curran Associates, Inc.
- Csardi, G.; and Nepusz, T. 2006. The igraph software package for complex network research. *InterJournal*, Complex Systems: 1695.
- Glymour, C.; Zhang, K.; and Spirtes, P. 2019. Review of Causal Discovery Methods Based on Graphical Models. *Frontiers in Genetics*, 10.
- Hauser, A.; and Bühlmann, P. 2012. Characterization and greedy learning of interventional Markov equivalence classes of directed acyclic graphs. *Journal of Machine Learning Research*, 13: 2409–2464.
- Huang, Y.; and Valtorta, M. 2006. Pearl's Calculus of Intervention Is Complete. In T.S. Richardson, R. D. a., ed., *Proceedings of the Twenty-Second Conference on Uncertainty in Artificial Intelligence*, 217–224. AUAI Press.
- Kleinberg, S.; and Hripcsak, G. 2011. A review of causal inference for biomedical informatics. *Journal of Biomedical Informatics*, 44(6): 1102–1112.
- Lauritzen, S. L.; and Richardson, T. S. 2002. Chain graph models and their causal interpretations. *Royal Statistical Society*, 64(Part 2): 1–28.
- Lee, S.; Correa, J. D.; and Bareinboim, E. 2019. General Identifiability with Arbitrary Surrogate Experiments. In *Proceedings of the Thirty-Fifth Conference Annual Conference on Uncertainty in Artificial Intelligence*. AUAI Press.
- Parviainen, P.; and Kaski, S. 2016. Bayesian Networks for Variable Groups. In Antonucci, A.; Corani, G.; and Campos, C. P., eds., *Proceedings of the Eighth International Conference on Probabilistic Graphical Models*, volume 52 of *Proceedings of Machine Learning Research*, 380–391. PMLR.
- Pearl, J. 1988. *Probabilistic Reasoning in Intelligent Systems*. San Mateo, CA: Morgan Kaufmann.
- Pearl, J. 1995. Causal diagrams for empirical research. *Biometrika*, 82(4): 669–688.
- Pearl, J. 2000. *Causality: Models, Reasoning, and Inference*. NY, USA: Cambridge University Press, 2nd edition.
- Pearl, J.; and Mackenzie, D. 2018. *The Book of Why*. New York: Basic Books.
- Peters, J.; Janzing, D.; and Schölkopf, B. 2017. *Elements of Causal Inference: Foundations and Learning Algorithms*. Adaptive Computation and Machine Learning. Cambridge, MA: MIT Press.
- R Core Team. 2021. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Richardson, T. 2003. Markov Properties for Acyclic Directed Mixed Graphs. *Scandinavian Journal of Statistics*, 30(1): 145–157.
- Rubenstein, P. K.; Weichwald, S.; Bongers, S.; Mooij, J. M.; Janzing, D.; Grosse-Wentrup, M.; and Schölkopf, B. 2017. Causal Consistency of Structural Equation Models. In Eli-dan, G.; Kersting, K.; and Ihler, A. T., eds., *Proceedings of the Thirty-Third Conference on Uncertainty in Artificial Intelligence, UAI 2017, Sydney, Australia, August 11-15, 2017*. AUAI Press.
- Schölkopf, B.; Locatello, F.; Bauer, S.; Ke, N. R.; Kalchbrenner, N.; Goyal, A.; and Bengio, Y. 2021. Toward Causal Representation Learning. *Proceedings of the IEEE*, 109(5): 612–634.
- Shen, Y.; Choi, A.; and Darwiche, A. 2018. Conditional PSDDs: Modeling and Learning With Modular Knowledge. *Proceedings of the AAAI Conference on Artificial Intelligence*, 32(1).
- Shpitser, I.; and Pearl, J. 2006. Identification of Joint Interventional Distributions in Recursive semi-Markovian Causal Models. In *Proceedings of the Twenty-First AAAI Conference on Artificial Intelligence*, 1219–1226.
- Spirtes, P.; Glymour, C. N.; and Scheines, R. 2000. *Causation, Prediction, and Search*. Cambridge, MA: MIT Press, 2nd edition.
- Textor, J.; van der Zander, B.; Gilthorpe, M. S.; Liśkiewicz, M.; and Ellison, G. T. 2016. Robust causal inference using directed acyclic graphs: the R package 'dagitty'. *International Journal of Epidemiology*, 45(6): 1887–1894.
- Tian, J. 2002. *Studies in Causal Reasoning and Learning*. Ph.D. thesis, Computer Science Department, University of California, Los Angeles, CA.

Tian, J.; and Pearl, J. 2002a. A General Identification Condition for Causal Effects. In *Proceedings of the Eighteenth National Conference on Artificial Intelligence (AAAI 2002)*, 567–573. Menlo Park, CA: AAAI Press/MIT Press.

Tian, J.; and Pearl, J. 2002b. On the Testable Implications of Causal Models with Hidden Variables. *Proceedings of the Eighteenth Conference on Uncertainty in Artificial Intelligence (UAI-02)*, 519–527.

Tikka, S.; and Karvanen, J. 2017. Identifying Causal Effects with the R Package causaleffect. *Journal of Statistical Software*, 76(12): 1–30.

Zhang, J. 2008. On the completeness of orientation rules for causal discovery in the presence of latent confounders and selection bias. *Artificial Intelligence*, 172(16): 1873–1896.

A Appendix: Supplemental material for “Causal Effect Identification in Cluster DAGs”

This appendix consists of four parts. In Section A.1, we describe how path analysis is extended to C-DAGs. In Section A.2 we provide a more detailed explanation on how a C-DAG can be characterized as a Causal Bayesian Network (CBN) over clustered variables, representing possibly multiple underlying CBNs over the original variables. In Section A.3, we empirically illustrate the validity of C-DAGs as an abstraction of causal diagrams. Lastly, in Section A.4, we present the proofs of all results in the paper, namely Propositions 1 and 2, Lemmas 2, 3, and 1, and Theorems 1 to 6, along with some auxiliary results necessary for those proofs.

A.1 Path Analysis in C-DAGs

Since a C-DAG represents a class of compatible causal diagrams, we must make sure that the status of a path in a C-DAG reflects the status of all corresponding paths in all compatible causal diagrams.

Let $\mathbf{X}, \mathbf{Y}, \mathbf{Z}$ be clusters in the C-DAG G_C . Let the symbol $*$ represent either an arrow head or tail. Let p be a path in G_C between \mathbf{X} and \mathbf{Y} that goes only through \mathbf{Z} .

We say that p is *inactive* in G_C if in all causal diagrams G compatible with G_C , all corresponding paths are inactive according to the d-separation rules. Note that if p is in the form $\mathbf{X} \ast \mathbf{Z} \rightarrow \mathbf{Y}$ (i.e., \mathbf{Z} is a non-collider), then the corresponding paths in G are those between \mathbf{X} and \mathbf{Y} going only through variables in \mathbf{Z} and containing the edge $Z \rightarrow Y$ for some $Z \in \mathbf{Z}$ and some $Y \in \mathbf{Y}$. Alternatively, if p is in the form $\mathbf{X} \ast \mathbf{Z} \leftarrow \mathbf{Y}$ (i.e., \mathbf{Z} is a collider), then the corresponding paths in G are those between \mathbf{X} and \mathbf{Y} going only through variables in \mathbf{Z} and containing the edge $X \ast Z$ for some $X \in \mathbf{X}$ and some $Z \in \mathbf{Z}$ and the edge $Z' \leftarrow Y$ for some $Z' \in \mathbf{Z}$ and some $Y \in \mathbf{Y}$.

We say p is *either active or inactive* if in some compatible causal diagram G all the paths between \mathbf{X} and \mathbf{Y} going through only variables in \mathbf{Z} are inactive, while in some compatible causal diagram G there exists at least one active path between \mathbf{X} and \mathbf{Y} going through only variables in \mathbf{Z} .

Next, we investigate the status of paths in a C-DAG under two possible scenarios for a cluster \mathbf{Z} : 1) as a non-collider, or 2) as a collider or a descendant of a collider.

Cluster as a Non-Collider (Chains & Forks)

Consider a C-DAG where \mathbf{Z} is a non-collider between \mathbf{X} and \mathbf{Y} as shown in Fig. 6(a) and (d). The causal diagrams (b) and (c) in Fig. 6, over $\mathbf{V} = \{X, Z_1, Z_2, Z_3, Y\}$, are compatible with the C-DAG in Fig. 6(a), where $\mathbf{X} = \{X\}$, $\mathbf{Z} = \{Z_1, Z_2, Z_3\}$, and $\mathbf{Y} = \{Y\}$. Note that in diagram (b) the path between X and Y is active when \mathbf{Z} is not conditioned on. However, in diagram (c), the path between X and Y is inactive, even though the abstracted cluster \mathbf{Z} is a non-collider. Similarly, in Fig. 6, the causal diagrams (e) and (f) over $\mathbf{V} = \{X, Z_1, Z_2, Z_3, Y\}$ are compatible with the C-DAG G_C in (d), where $\mathbf{X} = \{X\}$, $\mathbf{Z} = \{Z_1, Z_2, Z_3\}$, and $\mathbf{Y} = \{Y\}$. In diagram (e), the path is active while in diagram (f) the path is inactive when \mathbf{Z} is not conditioned on.

From these examples, it is clear that when a cluster \mathbf{Z} acts as a non-collider between two other clusters \mathbf{X} and \mathbf{Y} , the path between \mathbf{X} and \mathbf{Y} through \mathbf{Z} may be either active or inactive.

Remark 1. In a C-DAG, the path $\mathbf{X} \ast \mathbf{Z} \rightarrow \mathbf{Y}$ may be either active or inactive when \mathbf{Z} is not conditioned on.

Conditioning on a cluster will be considered as equivalent to conditioning on all the variables in it. In all diagrams (b), (c), (e) and (f), the path between X and Y is inactive when conditioning on $\{Z_1, Z_2, Z_3\}$ and we show that this property will hold regardless of the connections within a cluster as stated in the following lemma.

Lemma 2. In a C-DAG, the path $\mathbf{X} \ast \mathbf{Z} \rightarrow \mathbf{Y}$ is inactive when non-collider \mathbf{Z} is conditioned on.

Proof. Let G be any causal diagram compatible with the C-DAG G_C : $\mathbf{X} \ast \mathbf{Z} \rightarrow \mathbf{Y}$. We show \mathbf{X} is blocked from \mathbf{Y} by \mathbf{Z} in G . We show the contrapositive.

Assume there is an active path between \mathbf{X} and \mathbf{Y} not blocked by \mathbf{Z} in G . Then there is path p between a variable $X \in \mathbf{X}$ and a variable $Y \in \mathbf{Y}$ such that all the non-terminal nodes on p are in \mathbf{Z} (otherwise X will be adjacent to Y in G_C) and are colliders (otherwise p will be blocked by \mathbf{Z}). Let $V \in \mathbf{Z}$ be the collider adjacent

to Y . For V to be a collider there must either be an edge from Y into V or a bidirected edge between Y and V . In both cases, the edge is into V . This contradicts the hypothesis of \mathbf{Z} being connected to \mathbf{Y} by the edge $\mathbf{Z} \rightarrow \mathbf{Y}$, which does not allow a variable in \mathbf{Y} to be either an ancestor of a variable in \mathbf{Z} or connected to a variable in \mathbf{Z} by a bidirected edge. \square

Cluster as a Collider or a Descendant of a Collider

Consider a C-DAG where \mathbf{Z} is a collider between \mathbf{X} and \mathbf{Y} as shown in Fig. 6(g), or a descendant of a collider \mathbf{W} between \mathbf{X} and \mathbf{Y} as shown in Fig. 6(j). The causal diagrams (h) and (i) in Fig. 6 over $\mathbf{V} = \{X, Z_1, Z_2, Z_3, Y\}$ are compatible with the C-DAG in Fig. 6(g), where $\mathbf{X} = \{X\}$, $\mathbf{Z} = \{Z_1, Z_2, Z_3\}$, and $\mathbf{Y} = \{Y\}$. The path between X and Y is active when conditioning on $\{Z_1, Z_2, Z_3\}$ in diagram (h), but it is inactive in diagram (i). Similarly, the causal diagrams (k) and (l) in Fig. 6 over $\mathbf{V} = \{X, W_1, W_2, Z_1, Z_2, Y\}$ are compatible with the C-DAG in Fig. 6(j), where $\mathbf{X} = \{X\}$, $\mathbf{Z} = \{Z_1, Z_2\}$, $\mathbf{W} = \{W_1, W_2\}$, and $\mathbf{Y} = \{Y\}$. In diagram (k), the path from X and Y is active when conditioning on $\{Z_1, Z_2\}$, while in diagram (l) the path is inactive when conditioning on $\{Z_1, Z_2\}$. These examples illustrate that when a cluster \mathbf{Z} acts as a collider or a descendant of a collider between two other clusters \mathbf{X} and \mathbf{Y} , the path between \mathbf{X} and \mathbf{Y} through \mathbf{Z} may be either active or inactive when conditioning on \mathbf{Z} .

Remark 2. In a C-DAG, the path $\mathbf{X} \ast \mathbf{Z} \leftarrow \mathbf{Y}$ may be either active or inactive when \mathbf{Z} or some descendant of \mathbf{Z} is conditioned on.

Note that the path between X and Y is inactive in all diagrams (h), (i), (k), (l). We show this property holds in general




Figure 6: Causal diagrams (b), (c) are compatible with the C-DAG (a). Causal diagrams (e), (f) are compatible with the C-DAG (d). In C-DAGs (a), (d), Z is a non-collider between X and Y . The path is active in (b), (e), but inactive in (c) and (f). Causal diagrams (h), (i) are compatible with the C-DAG (g), where Z is a collider between X and Y . Causal diagrams (k), (l) are compatible with the C-DAG (j), where W is a collider between X and Y , and Z is a descendant of W . When Z is conditioned on, the path is active in (h), (k), but inactive in (i), (l).

regardless of the connections within a cluster as stated in the following lemma.

Lemma 3. *In a C-DAG, the path $X \rightarrow Z \leftarrow Y$ is inactive when none of the descendants of Z (nor Z) are conditioned on.*

Proof. Let G be any causal diagram compatible with the C-DAG $X \rightarrow Z \leftarrow Y$. We show X is blocked from Y in G when conditioning neither on Z nor on a descendant of Z . Consider any path p in G between a variable $X \in \mathbf{X}$ and a variable $Y \in \mathbf{Y}$ going through some variables in \mathbf{Z} . Then there exist $X' \in \mathbf{X}$ and $V \in \mathbf{Z}$, both in p , such that $X' \rightarrow V$. Further, there exist $V' \in \mathbf{Z}$ and $Y' \in \mathbf{Y}$, both in p , such that $Y' \rightarrow V'$. Then p is of the form $X \rightarrow \dots \rightarrow X' \rightarrow V \leftarrow \dots \leftarrow V' \leftarrow Y'$ and must contain a collider between V and V' that is in \mathbf{Z} . Therefore, p must be inactive when none of the descendants of such a collider (nor the collider) is conditioned on. \square

A.2 Characterization of C-DAGs as Causal Bayesian Networks

In Section 3, Remark 3, we noted that it does not follow directly from Def. 1 that a C-DAG G_C has the semantics and properties of a causal diagram or Causal Bayesian Network (CBN) over the cluster nodes \mathbf{C}_i , for $i = 1, \dots, k$. Here we clarify how the results proved in this work allow us to characterize a C-DAG as a CBN over clustered variables, representing possibly multiple underlying CBNs over the original variables.

The semantics of a C-DAG is defined in terms of the class of all compatible causal diagrams and corresponding SCMs over \mathbf{V} . In this way, a C-DAG can be seen as an equivalence class of causal diagrams sharing the relationships among the clusters while allowing for any possible relationships among the variables within each cluster. For instance, in Fig. 7, the DAGs G_1, G_2 induce different col-




Figure 7: With cluster $\mathbf{Z} = \{Z_1, Z_2, Z_3\}$, G_{C_1} is the C-DAG for G_1 and G_2 ; G_{C_2} is the C-DAG for G_3 and G_4 ; and G_{C_3} is the C-DAG for G_5 and G_6 .

lections of interventional distributions over the set of variables \mathbf{V} (i.e., $\mathbf{P}_*^1(\mathbf{v}) \neq \mathbf{P}_*^2(\mathbf{v})$). However, when considering the cluster $\mathbf{Z} = \{Z_1, Z_2, Z_3\}$, both G_1 and G_2 are represented by the C-DAG G_{C_1} , which induces the same collection of interventional distributions over the clusters \mathbf{C} (i.e., $\mathbf{P}_*^1(\mathbf{c}) = \mathbf{P}_*^2(\mathbf{c}) = \mathbf{P}_{C_1}^1(\mathbf{c})$). In this sense, G_1 and G_2 can be thought of as being members of the equivalence class represented by G_{C_1} . The same can be concluded for G_3, G_4 , represented by G_{C_2} , and for G_5, G_6 , represented by G_{C_3} .

In Section 4, Theorem 1, we showed that a d-separation in a C-DAG G_C implies the same d-separation in all compatible causal diagrams. Given this pleasant result, we can characterize a C-DAG as a graphical encoder of conditional

independencies over the set of clusters \mathbf{C} , but this is still insufficient to give the causal interpretation necessary for the application of causal inferences tools. In fact, the same set of conditional independencies can be encoded by different DAGs, where vertices are not necessarily arranged following a causal order. This means that different DAGs can induce the same observational distribution, while differing with respect to the induced collection of interventional distributions. For example, in Fig. 7, for DAGs G_i , with $i = 1, 3, 5$, it holds that $P^i(\mathbf{v}) = P(\mathbf{v})$, even though $P^i(\mathbf{v}|do(\mathbf{x})) \neq P^j(\mathbf{v}|do(\mathbf{x}))$ for any $i \neq j$ and some $\mathbf{X} \subseteq \mathbf{V}$. Now, at the cluster level, for all DAGs G_i , with $i = 1, \dots, 6$, $P^i(\mathbf{c}) = P(\mathbf{c})$, while only DAGs represented by the same C-DAG induce the same collection of interventional distributions over \mathbf{C} .

To formalize a C-DAG $G_{\mathbf{C}}$ as a CBN over the cluster variables $\mathbf{C}_i \in \mathbf{C}$, it is necessary to show that any distribution $P_{\mathbf{x}}(\mathbf{v})$ resulting from the intervention $do(\mathbf{X} = \mathbf{x})$, for any $\mathbf{X} \subseteq \mathbf{C}$, including the observational distribution $P(\mathbf{v})$, where $\mathbf{X} = \emptyset$, factorizes according to the graphical structure of $G_{\mathbf{C}}$. This result, proved in Theorem 5, allows us to characterize a single C-DAG $G_{\mathbf{C}}$ as an encoder of the space of all interventional distributions \mathbf{P}_* at the cluster level, i.e., over the clusters of variables $\mathbf{C}_i \in \mathbf{C}$. More formally, a C-DAG $G_{\mathbf{C}}$, compatible with a causal diagram G , is a CBN for $\mathbf{P}_*^{\mathcal{M}}(\mathbf{c})$, the collection of interventional distributions induced by \mathcal{M} , over the clusters \mathbf{C} .

This characterization is crucial for the correct abstraction of the possible underlying causal models and validity of causal tools applied to causal C-DAGs. Identifying an effect $P(y|do(x))$ in a causal C-DAG means identifying such an effect for the entire class of causal diagrams represented. In $G_{\mathbf{C}_2}$, the identification formula $P(y|do(x)) = P(y|x)$ is valid for G_3 and G_4 , as well as all other diagrams in the class. This effect is different in the causal diagrams represented by $G_{\mathbf{C}_3}$ (e.g., G_5 and G_6) where $P(y|do(x)) = P(y)$. On the other hand, no expression can be derived for $P(y|do(x))$ from $G_{\mathbf{C}_1}$ since the encoded partial knowledge is compatible with some causal diagram in which the effect is not identifiable such as G_1 .

A.3 Simulations

We evaluate C-DAGs in their ability to determine an equivalent causal effect from an observational dataset to the effect determined from any compatible causal diagram, by their respective identification formulas. Given a causal diagram, representing true and complete causal structural knowledge, an identifiable effect should be computable by an identification formula from a compatible C-DAG, representing partial structural knowledge, in which the effect is also identifiable. This means that given the limited knowledge of a C-DAG, an effect can be determined that would be equivalent to whatever the underlying (unknown) causal diagram is.

In Fig. 8, $P(y|do(x))$ is identifiable by backdoor adjustment in $G_{\mathbf{C}_1}$ (i.e., $P(y|do(x)) = \sum_{\mathbf{z}} P(y|x, \mathbf{z})P(\mathbf{z})$), but it is not identifiable in $G_{\mathbf{C}_2}$. Diagrams (a)-(b) are compatible with $G_{\mathbf{C}_1}$ and diagrams (c)-(d) are compatible with $G_{\mathbf{C}_2}$. Those are examples of compatible causal diagrams in which two variables are clustered to create \mathbf{Z} .




Figure 8: $G_{\mathbf{C}_1}$ is the C-DAG for diagrams (a) and (b). $G_{\mathbf{C}_2}$ is the C-DAG for diagrams (c) and (d). $P(y|do(x))$ is identifiable by backdoor adjustment over the cluster $\mathbf{Z} = \{Z_1, Z_2\}$ in $G_{\mathbf{C}_1}$ and, therefore, in *all* compatible causal diagrams (e.g., diagrams (a)-(b)). However, $P(y|do(x))$ is not identifiable in $G_{\mathbf{C}_2}$. This means that there is *some* compatible causal diagram for which $P(y|do(x))$ is not identifiable (e.g., diagram (d)).

For our simulations, we consider more complex causal diagrams involving 10 variables clustered to create \mathbf{Z} .

For C-DAG $G_{\mathbf{C}_1}$, we randomly generated 100 causal diagrams that are compatible with this C-DAG. Each generated compatible causal diagram includes randomly generated relationships among the 10 variables in cluster \mathbf{Z} , and to \mathbf{X} and \mathbf{Y} , while adhering to the constraints imposed by the C-DAG.

Identification expressions were determined by the ID-algorithm given the C-DAG as well as the compatible causal diagrams. According to each causal diagram, we generated 100 data sets consisting of N observations. For each data set, we computed effect sizes using the identification formula derived from the causal diagram used to generate the data set and the one derived from the C-DAG. Then, we obtained the average difference between these two calculated effect sizes over the samples for all causal diagrams. The simulation was repeated with increasing values of $N = \{5000, 10000, 50000, 100000\}$. The simulations were implemented in R v4.1.2 (R Core Team 2021), using the packages *dagitty* v0.3.1 (Textor et al. 2016), *igraph* v1.2.8.9014 (Csardi and Nepusz 2006), *pcalg* v2.7.3 (Hauser and Bühlmann 2012), and *causaleffect* v1.3.13 (Tikka and Karvanen 2017). The results are shown in Fig. 9-A).

These empirical results illustrate a negligible average difference between the effect sizes computed by the identification formula for a causal diagram, and by the identification formula for the compatible C-DAG. Moreover, the standard error greatly decreases as the number of observations in the dataset increases. This illustrates that C-DAGs can be used to determine equivalent, accurate effect sizes, while necessitating significantly less domain knowledge for their construction.

For the C-DAG $G_{\mathbf{C}_2}$ in Fig. 8, we evaluate the non-identifiability of the causal effect in a subset of the causal




Figure 9: (A) Average and standard error of the difference between the effects computed using the identification formula returned by the ID algorithm when the C-DAG G_{C_1} in Fig. 8 is given and when a random compatible causal diagram is given. The average is over 100 causal diagrams with 100 data sets consisting of N observations, with $N = \{5000, 10000, 50000, 100000\}$. (B) Percentage of causal diagrams compatible with the C-DAG G_{C_2} in Fig. 8 in which the causal effect was identifiable (ID) and non-identifiable (Non-ID), in a sample of 100 randomly generated causal diagrams.

diagrams compatible with the C-DAG. Since the interventional distribution $P(y|do(x))$ is not identifiable given G_{C_2} , it's expected that there is at least one causal diagram compatible with G_{C_2} for which the effect is not identifiable. For these experiments, we generated 100 random causal diagrams compatible with the C-DAG G_{C_2} and used the ID-algorithm to determine if the effect was identifiable or not in the true causal diagram. This simulation was repeated 100 times. As shown in Fig. 9-B), the simulation shows that the effect was not identifiable in $86.13\% \pm 3.41$ (mean \pm standard error) of the random causal diagrams generated and compatible with C-DAG G_{C_2} .

A.4 Proofs

We start by showing in Proposition 1 that any adjacency between variables is preserved in a compatible C-DAG.

Proposition 1. (Preservation of adjacencies) *Let $G_C(\mathbf{C}, \mathbf{E}_C)$ be a C-DAG compatible with an ADMG $G(\mathbf{V}, \mathbf{E})$. Consider distinct clusters $\mathbf{C}_i, \mathbf{C}_j \in \mathbf{C}$. If $V_i, V_j \in \mathbf{V}$ are adjacent in G and belong to $\mathbf{C}_i, \mathbf{C}_j$ respectively, then \mathbf{C}_i and \mathbf{C}_j are adjacent in G_C . Further, if \mathbf{C}_i and \mathbf{C}_j are adjacent in G_C , then there exists $V_i \in \mathbf{C}_i$ and $V_j \in \mathbf{C}_j$ such that V_i and V_j are adjacent in G .*

Proof. Assume that two variables $V_i, V_j \in \mathbf{V}$ are connected by an edge in G . If $V_i \in \mathbf{C}_i$ and $V_j \in \mathbf{C}_j$, where \mathbf{C}_i and \mathbf{C}_j are two distinct clusters in G_C , then by the definition of cluster DAGs (Definition 1), if $V_i \rightarrow V_j$, then $\mathbf{C}_i \rightarrow \mathbf{C}_j$. Further, if $V_i \leftrightarrow V_j$, then $\mathbf{C}_i \leftrightarrow \mathbf{C}_j$. So, \mathbf{C}_i and \mathbf{C}_j are also connected by an edge in G_C .

Now, assume that \mathbf{C}_i and \mathbf{C}_j are connected by an edge in G_C . If $\mathbf{C}_i \rightarrow \mathbf{C}_j$, then, by Definition 1, there exists $V_i \in \mathbf{C}_i$ and $V_j \in \mathbf{C}_j$ such that $V_i \in Pa(V_j)$ in G . Therefore, V_i and V_j are connected by the edge $V_i \rightarrow V_j$ in G . Similarly, if $\mathbf{C}_i \leftrightarrow \mathbf{C}_j$, then, by Definition 1, there exists $V_i \in \mathbf{C}_i$ and $V_j \in \mathbf{C}_j$ such that $V_i \leftrightarrow V_j$, that is, V_i and V_j are connected by an edge in G . \square

We show in Proposition 2 that connections via directed paths are also preserved. This implies that both order and ancestral relationships are preserved in C-DAGs. To prove, we need first the following lemma:

Lemma 4. (Preservation of paths) *Let $G_C(\mathbf{C}, \mathbf{E}_C)$ be a C-DAG compatible with an ADMG $G(\mathbf{V}, \mathbf{E})$ and $\mathbf{C}_i, \mathbf{C}_j \in \mathbf{C}$ be two distinct clusters. If two variables $V_i \in \mathbf{C}_i$ and $V_j \in \mathbf{C}_j$ are connected in G by a path (of any sorts), then the clusters \mathbf{C}_i and \mathbf{C}_j are connected by a path in G_C .*

Proof. First by Proposition 1, if V_i and V_j are adjacent, then \mathbf{C}_i and \mathbf{C}_j are connected in G_C by an edge. Now, assume that V_i and V_j are not adjacent and let p be a path connecting V_i and V_j in G . Let p' be any subpath of p that is split into two different clusters of the form $V_1 ** \dots ** V_k ** V_{k+1} ** \dots ** V_n$, where $V_1, \dots, V_k \in \mathbf{C}_k$, $V_{k+1}, \dots, V_n \in \mathbf{C}_{k+1}$, and $*$ represents either an edge head or tail. By Definition 1, \mathbf{C}_k and \mathbf{C}_{k+1} will be connected by the same type of edge between V_k and V_{k+1} in G_C . By using the same argument for every subpath of p that is split into two different clusters, we conclude that all clusters containing variables in p are connected by a path and, thus, \mathbf{C}_i and \mathbf{C}_j are connected by a path in G_C . \square

Proposition 2. (Preservation of directed paths) Let $G_C(\mathbf{C}, \mathbf{E}_C)$ be a C-DAG compatible with an ADMG $G(\mathbf{V}, \mathbf{E})$ and $\mathbf{C}_i, \mathbf{C}_j \in \mathbf{C}$ be two distinct clusters. If two variables $V_i \in \mathbf{C}_i$ and $V_j \in \mathbf{C}_j$ are connected in G by a directed path from V_i to V_j , then the clusters \mathbf{C}_i and \mathbf{C}_j are connected by a directed path from \mathbf{C}_i to \mathbf{C}_j in G_C .

Proof. The proof is analogous to that of Lemma 4, noticing that the edges forming the path connecting \mathbf{C}_i to \mathbf{C}_j in G_C are always pointing towards \mathbf{C}_j . If V_i and V_j are adjacent, then $V_i \in Pa(V_j)$ and, by definition of C-DAGs, $\mathbf{C}_i \rightarrow \mathbf{C}_j$. Now, assume that V_i and V_j are connected, but not adjacent, and let p be a directed path from V_i to V_j . Let p' be any subpath of p that is split into two different clusters of the form $V_1 \rightarrow \dots \rightarrow V_k \rightarrow V_{k+1} \rightarrow \dots \rightarrow V_n$, where $V_1, \dots, V_k \in \mathbf{C}_k$, $V_{k+1}, \dots, V_n \in \mathbf{C}_{k+1}$. Since $V_k \rightarrow V_{k+1}$ in G , we have, by Definition 1, that $\mathbf{C}_k \rightarrow \mathbf{C}_{k+1}$ in G_C . By using the same argument for every subpath of p that is split into two different clusters, we conclude that all clusters containing variables in p are connected by a directed path pointing away from V_i and towards V_j , thus forming a directed path from \mathbf{C}_i to \mathbf{C}_j in G_C . \square

Next, we show the proofs for the main results in this work, including soundness and completeness of d-separation, do-calculus, and ID-algorithm in C-DAGs.

Theorem 1. (Soundness and completeness of d-separation in C-DAGs). In a C-DAG G_C , let $\mathbf{X}, \mathbf{Z}, \mathbf{Y} \subset \mathbf{C}$. If \mathbf{X} and \mathbf{Y} are d-separated by \mathbf{Z} in G_C , then, in any ADMG G compatible with G_C , \mathbf{X} and \mathbf{Y} are d-separated by \mathbf{Z} in G , i.e.,

$$(\mathbf{X} \perp\!\!\!\perp \mathbf{Y} \mid \mathbf{Z})_{G_C} \implies (\mathbf{X} \perp\!\!\!\perp \mathbf{Y} \mid \mathbf{Z})_G. \quad (1)$$

If \mathbf{X} and \mathbf{Y} are not d-separated by \mathbf{Z} in G_C , then, there exists an ADMG G compatible with G_C where \mathbf{X} and \mathbf{Y} are not d-separated by \mathbf{Z} in G .

Proof. We first prove the soundness of d-separation in C-DAGs by showing that if \mathbf{X} and \mathbf{Y} are d-separated by \mathbf{Z} in G_C , then, in any ADMG G compatible with G_C , \mathbf{X} and \mathbf{Y} are d-separated by \mathbf{Z} in G .

We show the contrapositive. Assume that \mathbf{X} and \mathbf{Y} are d-separated by \mathbf{Z} in G_C , but in a compatible ADMG G there exists a path p' between a variable $X \in \mathbf{X}$ and $Y \in \mathbf{Y}$ that is active when the set of variables contained in clusters in \mathbf{Z} are conditioned on. By Proposition 1 and Lemma 4, no connection is destroyed through clustering, so p' is contained in a path p of G_C between clusters \mathbf{X} and \mathbf{Y} . Since \mathbf{X} and \mathbf{Y} are d-separated by \mathbf{Z} in G_C , p is blocked and clusters \mathbf{X} and \mathbf{Y} are not adjacent. Therefore, by Definition 2, there is at least one triplet of clusters in p that is blocked. Let this triplet be $\langle \mathbf{C}_i, \mathbf{C}_m, \mathbf{C}_j \rangle$, where \mathbf{C}_m is distinct from \mathbf{X} and \mathbf{Y} . Consider the subpath p'_{ij} of p' contained in the triplet $\langle \mathbf{C}_i, \mathbf{C}_m, \mathbf{C}_j \rangle$ in p . Since p' is active by assumption, every subpath of p' is active, including p'_{ij} . The triplet $\langle \mathbf{C}_i, \mathbf{C}_m, \mathbf{C}_j \rangle$ is blocked either if \mathbf{C}_m is a mediator or common cause such that $\mathbf{C}_m \in \mathbf{Z}$ or if \mathbf{C}_m is a collider such that $\mathbf{C}_m \notin \mathbf{Z}$ and no descendant of \mathbf{C}_m is in \mathbf{Z} . In the former case, Lemma 2 shows that any path between \mathbf{C}_i and \mathbf{C}_j in a compatible ADMG G is inactive when \mathbf{C}_m is conditioned on. In the latter case, Lemma 3 shows that any path

between \mathbf{C}_i and \mathbf{C}_j in a compatible ADMG G is inactive when \mathbf{C}_m is not conditioned on. So, the subpath p'_{ij} of p' is inactive, and, thus, p' is also active.

We now prove the completeness of d-separation for C-DAGs by showing that if \mathbf{X} and \mathbf{Y} are not d-separated by \mathbf{Z} in G_C , then there exists an ADMG G compatible with G_C where \mathbf{X} and \mathbf{Y} are not d-separated by \mathbf{Z} in G .

If \mathbf{X} and \mathbf{Y} are adjacent, then, by Definition 1, in any ADMG G compatible with G_C , there always exists a pair of variables $X \in \mathbf{X}$ and $Y \in \mathbf{Y}$ that are adjacent. Now, if \mathbf{X} and \mathbf{Y} are not adjacent and are not d-separated by \mathbf{Z} in G_C , then by Definition 2, there exists a path p between \mathbf{X} and \mathbf{Y} in G_C such that every triplet $\langle \mathbf{C}_i, \mathbf{C}_m, \mathbf{C}_j \rangle$ in it is active. This means that for all mediators and common causes \mathbf{C}_m contained in p , \mathbf{C}_m must not be in \mathbf{Z} , and for all colliders \mathbf{C}_n contained in p , either $\mathbf{C}_n \in \mathbf{Z}$ or some descendant of \mathbf{C}_n is in \mathbf{Z} . By remarks 1, 2, 3, and 4, these triplets may be either active or inactive. Therefore, there exists some ADMG G compatible with G_C in which a path p' between \mathbf{X} and \mathbf{Y} in G goes through all (and only) clusters in p and all of its subpaths contained in a triplet of clusters $\langle \mathbf{C}_i, \mathbf{C}_m, \mathbf{C}_j \rangle$ in p are active. In this case, p' is active and \mathbf{X} and \mathbf{Y} are not d-separated by \mathbf{Z} in G . \square

Theorem 2. (C-DAG as BN). Let G_C be a C-DAG compatible with an ADMG G . If the observational distribution $P(\mathbf{v})$ factorizes according to G by Eq. (2), then the observational distribution $P(\mathbf{v}) = P(\mathbf{c})$ factorizes according to G_C , i.e.,

$$P(\mathbf{c}) = \sum_{\mathbf{u}} P(\mathbf{u}) \prod_{k: \mathbf{C}_k \in \mathbf{C}} P(\mathbf{c}_k | pa_{\mathbf{C}_k}, \mathbf{u}'_k), \quad (3)$$

where $Pa_{\mathbf{C}_k}$ are the parents of the cluster \mathbf{C}_k , and $\mathbf{U}'_k \subseteq \mathbf{U}$ such that, for any i, j , $\mathbf{U}'_i \cap \mathbf{U}'_j \neq \emptyset$ if and only if there is a bidirected edge $(\mathbf{C}_i \leftrightarrow \mathbf{C}_j)$ between \mathbf{C}_i and \mathbf{C}_j in G_C .

Proof. Consider the following topological order over all clusters in \mathbf{C} : $\mathbf{C}_1 \prec \mathbf{C}_2 \prec \dots \prec \mathbf{C}_K$. By Proposition 2, the ancestral relationships are preserved relative to any compatible ADMG. Therefore, such a topological order is consistent with a topological order of the compatible graph G , regardless of the internal order of the variables in each cluster.

Given the topological order over clusters, we construct a consistent topological order for the compatible ADMG G over \mathbf{V} as follows: $(V_{11} \prec \dots \prec V_{1n_1}) \prec (V_{21} \prec \dots \prec V_{2n_2}) \prec \dots \prec (V_{K1} \prec \dots \prec V_{Kn_K})$.

By assumption, the observational distribution $P(\mathbf{v}) = P(\mathbf{c})$ factorizes according to G by the following:

$$P(\mathbf{c}) = \sum_{\mathbf{u}} P(\mathbf{u}) \prod_{k: \mathbf{C}_k \in \mathbf{C}} \prod_{i: V_{ki} \in \mathbf{C}_k} P(v_{ki} | pa(v_{ki}), \mathbf{u}_{ki}). \quad (6)$$

where $Pa(V_{ki})$ are the endogenous parents of V_{ki} and \mathbf{U}_{ki} are the exogenous parents of V_{ki} , including those that are shared with some other variable $V_{k'j}$ (represented by $V_{k'j} \leftrightarrow V_{ki}$ in G).

Consider $V_{ki} \in \mathbf{C}_k$. The set $Pa(V_{ki})$ of endogenous parents of V_{ki} consists of the union of the set of the endogenous

parents of V_{ki} within the cluster \mathbf{C}_k , denoted as $Pa_{ki}^{(k)}$, and the sets of the endogenous parents of V_{ki} in all of the clusters that come before \mathbf{C}_k in the topological order, denoted as $Pa_{ki}^{(1)}, \dots, Pa_{ki}^{(k-1)}$. Define $\mathbf{U}_{ki}^{(k')} = \mathbf{U}_{ki} \cap (\bigcup_j \mathbf{U}_{k'j})$ for $k' \neq k$. Also let any exogenous parent of V_{ki} that is not shared with other variables outside of \mathbf{C}_k be in $\mathbf{U}_{ki}^{(k)}$. Note that $\mathbf{U}_{ki} = \bigcup_{k'} \mathbf{U}_{ki}^{(k')}$. In words, \mathbf{U}_{ki} is the union of $\mathbf{U}_{ki}^{(1)}, \dots, \mathbf{U}_{ki}^{(K)}$, which are the sets of exogenous parents of V_{ki} that are also exogenous parents of some variables in clusters $\mathbf{C}_1, \dots, \mathbf{C}_K$, respectively. Note that each variable $U \in \mathbf{U}_{ki}^{(k')}$ indicates a connection $V_{k'j} \leftrightarrow V_{ki}$ in G and, therefore, a connection $\mathbf{C}_{k'} \leftrightarrow \mathbf{C}_k$ in $G_{\mathbf{C}}$. Also note that $\mathbf{U}_{ki}^{(k)}$ is the set of exogenous parents of V_{ki} that may also be exogenous parents of some other variable V_{kj} in \mathbf{C}_k . The connection $V_{kj} \leftrightarrow V_{ki}$ in G is part of the internal structure of the cluster \mathbf{C}_k , which is abstracted away in $G_{\mathbf{C}}$.

Using this notation, we can rewrite the conditional distribution for V_{ki} as follows:

$$P(v_{ki} | pa(v_{ki}), \mathbf{u}_{ki}) = P(v_{ki} | pa_{ki}^{(1)}, \dots, pa_{ki}^{(k-1)}, pa_{ki}^{(k)}, \mathbf{u}_{ki}^{(1)}, \dots, \mathbf{u}_{ki}^{(K)})$$

Note that all variables $V_{k1}, \dots, V_{k,i-1} \in \mathbf{C}_k$ precede V_{ki} in the topological order and, therefore, are non-descendants of V_{ki} . Then, we have

$$(V_{ki} \perp\!\!\!\perp V_{k1}, \dots, V_{k,i-1} | Pa_{ki}^{(1)}, \dots, Pa_{ki}^{(k-1)}, Pa_{ki}^{(k)}, \mathbf{U}_{ki}^{(1)}, \dots, \mathbf{U}_{ki}^{(K)}). \quad (7)$$

Now, consider the set of endogenous parents of all other variables in the cluster, excluding those that are in \mathbf{C}_k or are also endogenous parents of V_{ki} :

$$Pa^*(\{V_{k1}, \dots, V_{kn_k}\}) = Pa(\{V_{k1}, \dots, V_{kn_k}\}) \setminus (Pa(V_{ki}) \cup \mathbf{C}_k) = (Pa_{k1}^{(1)} \cup \dots \cup Pa_{kn_k}^{(1)} \cup \dots \cup Pa_{k1}^{(k-1)} \cup \dots \cup Pa_{kn_k}^{(k-1)}) \setminus (Pa(V_{ki}) \cup \mathbf{C}_k)$$

Note that all variables in $Pa^*(\{V_{k1}, \dots, V_{kn_k}\})$ are non-descendants of V_{ki} . Then, we have:

$$(V_{ki} \perp\!\!\!\perp Pa^*(\{V_{k1}, \dots, V_{kn_k}\}) | Pa_{ki}^{(1)}, \dots, Pa_{ki}^{(k-1)}, Pa_{ki}^{(k)}, \mathbf{U}_{ki}^{(1)}, \dots, \mathbf{U}_{ki}^{(K)}). \quad (8)$$

Further, consider the set of variables that are in the same cluster as some parents of $\{V_{k1}, \dots, V_{kn_k}\}$, excluding the parents themselves, denoted as $\mathbf{C}^*(\{Pa_{k1}, \dots, Pa_{kn_k}\})$. Denote by $\mathbf{C}_{kj}^{*(k')}$, for $k' = 1, \dots, k-1$, the variables in cluster k' that are together with a parent of V_{kj} , but are not parents of any variable $V_{kj'} \in \mathbf{C}_k$, for $j' = 1, \dots, n_k$.

$$\mathbf{C}^*(\{Pa_{k1}, \dots, Pa_{kn_k}\}) = \mathbf{C}_{k1}^{*(1)} \cup \dots \cup \mathbf{C}_{kn_k}^{*(1)} \cup \dots \cup \mathbf{C}_{k1}^{*(k-1)} \cup \dots \cup \mathbf{C}_{kn_k}^{*(k-1)}.$$

Since all these variables are also non-descendants of V_{ki} , we have:

$$(V_{ki} \perp\!\!\!\perp \mathbf{C}^*(\{Pa_{k1}, \dots, Pa_{kn_k}\}) | Pa_{ki}^{(1)}, \dots, Pa_{ki}^{(k-1)}, Pa_{ki}^{(k)}, \mathbf{U}_{ki}^{(1)}, \dots, \mathbf{U}_{ki}^{(K)}). \quad (9)$$

Lastly, consider the set of exogenous parents of V_{k1}, \dots, V_{kn_k} that are not exogenous parents of V_{ki} :

$$\mathbf{U}_{\{k1, \dots, kn_k\}}^* = (\mathbf{U}_{k1}^{(1)} \cup \dots \cup \mathbf{U}_{kn_k}^{(1)} \cup \dots \cup \mathbf{U}_{k1}^{(K)} \cup \dots \cup \mathbf{U}_{kn_k}^{(K)}) \setminus (\mathbf{U}_{ki}^{(1)}, \dots, \mathbf{U}_{ki}^{(K)}).$$

We also have

$$(V_{ki} \perp\!\!\!\perp \mathbf{U}_{\{k1, \dots, kn_k\}}^* | Pa_{ki}^{(1)}, \dots, Pa_{ki}^{(k-1)}, Pa_{ki}^{(k)}, \mathbf{U}_{ki}^{(1)}, \dots, \mathbf{U}_{ki}^{(K)}). \quad (10)$$

Given the conditional independence relations in (7), (8), (9), and (10), we can rewrite the conditional distribution of V_{ki} to also condition on such non-descendants of V_{ki} as follows:

$$P(v_{ki} | pa(v_{ki}), \mathbf{u}_{ki}) = P(v_{ki} | v_{k1}, \dots, v_{k,i-1}, pa_{k1}^{(1)}, \dots, pa_{kn_k}^{(1)}, \dots, pa_{k1}^{(k-1)}, \dots, pa_{kn_k}^{(k-1)}, \mathbf{c}_{k1}^{*(1)}, \dots, \mathbf{c}_{kn_k}^{*(1)}, \dots, \mathbf{c}_{k1}^{*(k-1)}, \dots, \mathbf{c}_{kn_k}^{*(k-1)}, \mathbf{u}_{k1}^{(1)}, \dots, \mathbf{u}_{kn_k}^{(1)}, \dots, \mathbf{u}_{k1}^{(K)}, \dots, \mathbf{u}_{kn_k}^{(K)}).$$

Note that the parents of V_{ki} that are in the cluster \mathbf{C}_k are contained in $\{V_{k1}, \dots, V_{k,i-1}\}$. Also note that $(Pa(\{V_{k1}, \dots, V_{kn_k}\}) \setminus \mathbf{C}_k) \cup \mathbf{C}^*(\{Pa_{k1}, \dots, Pa_{kn_k}\}) = \bigcup Pa(\mathbf{C}_k)_{G_{\mathbf{C}}} \setminus \mathbf{C}_k$. Denote as $\mathbf{U}'_k = \bigcup_i \mathbf{U}_{ki} = \bigcup_{j,k'} \mathbf{U}_{kj}^{(k')}$, which is the set of the exogenous parents of all variables $V_{kj} \in \mathbf{C}_k$.

Given that, we can rewrite the conditional distribution of each $V_{ki} \in \mathbf{C}_i$ as follows:

$$P(v_{ki} | pa(v_{ki}), \mathbf{u}_{ki}) = P(v_{ki} | v_{k1}, \dots, v_{k,i-1}, pa(\mathbf{c}_k), \mathbf{u}'_k). \quad (11)$$

By considering the factors for every variable $V_{ki} \in \mathbf{C}_k$, for $i = 1 \dots n_k$, we can write the conditional distribution of the set of variables in the cluster \mathbf{C}_k as follows:

$$P(\mathbf{c}_k | pa(\mathbf{c}_k), \mathbf{u}'_k) = \prod_{i: V_{ki} \in \mathbf{C}_k} P(v_{ki} | v_{k1}, \dots, v_{k,i-1}, pa(\mathbf{c}_k), \mathbf{u}'_k). \quad (12)$$

Note that, for any two clusters $\mathbf{C}_i, \mathbf{C}_j \in \mathbf{C}$, $\mathbf{U}'_i \cap \mathbf{U}'_j \neq \emptyset$ if, and only if, there exist a variable $V_{il} \in \mathbf{C}_i$ and a variable $V_{jl'}$ in \mathbf{C}_j with a common exogenous parent. This follows from the construction of the sets \mathbf{U}'_i and \mathbf{U}'_j . Note that \mathbf{U}'_i includes all and only the exogenous parents of V_{il} , for $l = 1, \dots, n_i$, that are also exogenous parents of any other variable in the other clusters, including $V_{jl'}$. Similarly, \mathbf{U}'_j includes all and only the exogenous parents of $V_{jl'}$, for $l' = 1, \dots, n_j$, that are also exogenous parents of any other variable in the other clusters, including V_{il} . Since, by Definition 1, a bidirected edge between \mathbf{C}_i and \mathbf{C}_j exists if $\exists V_{il} \in \mathbf{C}_i$ and $V_{jl'}$ in \mathbf{C}_j such that $V_{il} \leftrightarrow V_{jl'}$ (i.e., V_{il} and $V_{jl'}$ have a common exogenous parent), we have $\mathbf{U}'_i \cap \mathbf{U}'_j \neq \emptyset$ if and only if $\mathbf{C}_i \leftrightarrow \mathbf{C}_j$ in $G_{\mathbf{C}}$.

Therefore we can rewrite the factorization over the variables $V_{ki} \in \mathbf{V}$ in G , shown in Equation 6, as a factorization over the clusters $\mathbf{C}_k \in \mathbf{C}$ in G_C as follows:

$$P(\mathbf{c}) = \sum_{\mathbf{u}} P(\mathbf{u}) \prod_{k: \mathbf{C}_k \in \mathbf{C}} P(\mathbf{c}_k | pa(\mathbf{c}_k), \mathbf{u}'_k). \quad (13)$$

□

Lemma 1. *If a C-DAG G_C is compatible with an ADMG G , then, for $\mathbf{X}, \mathbf{Z} \subset \mathbf{C}$, the mutilated C-DAG $G_{C_{\overline{\mathbf{XZ}}}}$ is compatible with the mutilated ADMG $G_{\overline{\mathbf{XZ}}}$.*

Proof. Note that we abuse the notation by letting $\mathbf{X}, \mathbf{Z} \subset \mathbf{C}$ denote sets of clusters in G_C and also letting $\mathbf{X}, \mathbf{Z} \subset \mathbf{V}$ denote, in G , the set of variables contained in the clusters in $\mathbf{X}, \mathbf{Z} \subset \mathbf{C}$.

We prove that the mutilated graph $G_{\overline{\mathbf{X}}}$ is compatible with $G_{C_{\overline{\mathbf{X}}}}$ by showing that the C-DAG $G_{C_{\overline{\mathbf{X}}}}$ constructed by removing from G_C the edges into \mathbf{X} is the same as the one constructed from $G_{\overline{\mathbf{X}}}$ using Definition 1.

An edge in G is cut in the construction of $G_{\overline{\mathbf{X}}}$ if and only if it is into some variable $X \in \mathbf{X}_i \in \mathbf{X}$. Note that, if every such an edge is between two variables that are in the same cluster \mathbf{X}_i , then the C-DAG constructed from $G_{\overline{\mathbf{X}}}$ will be the same as the one constructed from G . In other words, $G_{C_{\overline{\mathbf{X}}}} = G_C$. Further, the C-DAG G_C constructed from G contains no edge into any cluster $\mathbf{X}_i \in \mathbf{X}$. Therefore, no edge will be cut in the construction of $G_{C_{\overline{\mathbf{X}}}}$ from G_C , leading to the same conclusion that $G_{C_{\overline{\mathbf{X}}}} = G_C$.

Now, consider any edge of the type $V \rightarrow X$ in G , where $X \in \mathbf{X}_i \in \mathbf{X}$ and $V \in \mathbf{C}_i \in \mathbf{C}$ such that $\mathbf{C}_i \neq \mathbf{X}_i$ (note that \mathbf{C}_i can be a cluster in \mathbf{X}). Every of such an edge is cut in the construction of $G_{\overline{\mathbf{X}}}$ from G . Therefore, the C-DAG constructed from $G_{\overline{\mathbf{X}}}$ has no edge into any cluster $\mathbf{X}_i \in \mathbf{X}$. Further, by Definition 1, the existence of $V \rightarrow X$ in G guarantees the existence of the edge $\mathbf{C}_i \rightarrow \mathbf{X}_i$ in G_C and every such an edge in G_C will be cut in the construction of $G_{C_{\overline{\mathbf{X}}}}$ from G_C . Note that all other edges that are cut from G to construct $G_{\overline{\mathbf{X}}}$ are those between two variables that are in the same cluster $\mathbf{X}_i \in \mathbf{X}$. As previously discussed, such cuts do not change the C-DAG constructed from $G_{\overline{\mathbf{X}}}$. Therefore, the C-DAG constructed from $G_{\overline{\mathbf{X}}}$ is exactly the G_C after cutting the edges into every cluster $\mathbf{X}_i \in \mathbf{X}$.

Similar to the proof that for the compatibility between $G_{\overline{\mathbf{X}}}$ and $G_{C_{\overline{\mathbf{X}}}}$, we prove that the mutilated graph $G_{\overline{\mathbf{Z}}}$ is compatible with $G_{C_{\overline{\mathbf{Z}}}}$ by showing that the C-DAG $G_{C_{\overline{\mathbf{Z}}}}$ constructed by removing from G_C the edges out of \mathbf{Z} is the same as the one constructed from $G_{\overline{\mathbf{Z}}}$ using Definition 1.

An edge in G is cut in the construction of $G_{\overline{\mathbf{Z}}}$ if and only if it is out of some variable $Z \in \mathbf{Z}_i \in \mathbf{Z}$. If every such an edge is between two variables that are in the same cluster \mathbf{Z}_i , then the C-DAG constructed from $G_{\overline{\mathbf{Z}}}$ will be the same as the one constructed from G . In other words, $G_{C_{\overline{\mathbf{Z}}}} = G_C$. Further, the C-DAG G_C constructed from G contains no edge out of any cluster $\mathbf{Z}_i \in \mathbf{Z}$. Therefore, no edge will be cut in the construction of $G_{C_{\overline{\mathbf{Z}}}}$ from G_C , leading to the same conclusion that $G_{C_{\overline{\mathbf{Z}}}} = G_C$.

Now, consider any edge of the type $Z \rightarrow V$ in G , where $Z \in \mathbf{Z}_i \in \mathbf{Z}$ and $V \in \mathbf{C}_i \in \mathbf{C}$ such that $\mathbf{C}_i \neq \mathbf{Z}_i$ (note

that \mathbf{C}_i can be a cluster in \mathbf{Z}). Every of such an edge is cut in the construction of $G_{\overline{\mathbf{Z}}}$ from G . Therefore, the C-DAG constructed from $G_{\overline{\mathbf{Z}}}$ has no edge out of any cluster $\mathbf{Z}_i \in \mathbf{Z}$. Further, by Definition 1, the existence of $Z \rightarrow V$ in G guarantees the existence of the edge $\mathbf{Z}_i \rightarrow \mathbf{C}_i$ in G_C and every such an edge in G_C will be cut in the construction of $G_{C_{\overline{\mathbf{Z}}}}$ from G_C . Note that all other edges that are cut from G to construct $G_{\overline{\mathbf{Z}}}$ are those between two variables that are in the same cluster $\mathbf{Z}_i \in \mathbf{Z}$. As previously discussed, such cuts do not change the C-DAG constructed from $G_{\overline{\mathbf{Z}}}$. Therefore, the C-DAG constructed from $G_{\overline{\mathbf{Z}}}$ is exactly the G_C after cutting the edges into every cluster $\mathbf{Z}_i \in \mathbf{Z}$.

Since the processes of cutting edges for constructing $G_{\overline{\mathbf{X}}}$ and $G_{\overline{\mathbf{Z}}}$ are independent of each other, we conclude that $G_{C_{\overline{\mathbf{XZ}}}}$ is compatible with $G_{\overline{\mathbf{XZ}}}$, i.e., the C-DAG $G_{C_{\overline{\mathbf{XZ}}}}$ constructed by cutting from G_C the edges into \mathbf{X} and out of \mathbf{Z} is the same as the one constructed from $G_{\overline{\mathbf{XZ}}}$ using Def. 1. □

Theorem 3. (Do-calculus in causal C-DAGs). *Let G_C be a C-DAG compatible with an ADMG G . If G is a CBN encoding interventional distributions $P(\cdot | do(\cdot))$, then for any disjoint subsets of clusters $\mathbf{X}, \mathbf{Y}, \mathbf{Z}, \mathbf{W} \subseteq \mathbf{C}$, the following three rules hold:*

$$\textbf{Rule 1: } P(\mathbf{y} | do(\mathbf{x}), \mathbf{z}, \mathbf{w}) = P(\mathbf{y} | do(\mathbf{x}), \mathbf{w}) \\ \text{if } (\mathbf{Y} \perp\!\!\!\perp \mathbf{Z} | \mathbf{X}, \mathbf{W})_{G_{C_{\overline{\mathbf{X}}}}}$$

$$\textbf{Rule 2: } P(\mathbf{y} | do(\mathbf{x}), do(\mathbf{z}), \mathbf{w}) = P(\mathbf{y} | do(\mathbf{x}), \mathbf{z}, \mathbf{w}) \\ \text{if } (\mathbf{Y} \perp\!\!\!\perp \mathbf{Z} | \mathbf{X}, \mathbf{W})_{G_{C_{\overline{\mathbf{XZ}}}}}$$

$$\textbf{Rule 3: } P(\mathbf{y} | do(\mathbf{x}), do(\mathbf{z}), \mathbf{w}) = P(\mathbf{y} | do(\mathbf{x}), \mathbf{w}) \\ \text{if } (\mathbf{Y} \perp\!\!\!\perp \mathbf{Z} | \mathbf{X}, \mathbf{W})_{G_{C_{\overline{\mathbf{XZ}(\mathbf{W})}}}}$$

where $G_{C_{\overline{\mathbf{XZ}}}}$ is obtained from G_C by removing the edges into \mathbf{X} and out of \mathbf{Z} , and $\mathbf{Z}(\mathbf{W})$ is the set of \mathbf{Z} -clusters that are non-ancestors of any \mathbf{W} -cluster in $G_{C_{\overline{\mathbf{X}}}}$.

Proof. Note that we abuse the notation by letting $\mathbf{W}, \mathbf{X}, \mathbf{Y}, \mathbf{Z} \subset \mathbf{V}$ also denote the set of variables contained in the clusters in $\mathbf{W}, \mathbf{X}, \mathbf{Y}, \mathbf{Z} \subset \mathbf{C}$, respectively.

We first show that Rule 1 is sound. By Lemma 1, $G_{\overline{\mathbf{X}}}$ is compatible with $G_{C_{\overline{\mathbf{X}}}}$. Then by Theorem 1, if $(\mathbf{Y} \perp\!\!\!\perp \mathbf{Z} | \mathbf{X}, \mathbf{W})_{G_{C_{\overline{\mathbf{X}}}}}$ then $(\mathbf{Y} \perp\!\!\!\perp \mathbf{Z} | \mathbf{X}, \mathbf{W})_{G_{\overline{\mathbf{X}}}}$. Therefore Rule 1 holds by the Rule 1 of do-calculus in causal diagrams.

We now show that Rule 2 is sound. By Lemma 1, $G_{C_{\overline{\mathbf{XZ}}}}$ is compatible with $G_{\overline{\mathbf{XZ}}}$. Then by Theorem 1, if $(\mathbf{Y} \perp\!\!\!\perp \mathbf{Z} | \mathbf{X}, \mathbf{W})_{G_{C_{\overline{\mathbf{XZ}}}}}$ then $(\mathbf{Y} \perp\!\!\!\perp \mathbf{Z} | \mathbf{X}, \mathbf{W})_{G_{\overline{\mathbf{XZ}}}}$. Therefore Rule 2 holds by the Rule 2 of do-calculus in causal diagrams.

Lastly, we show that Rule 3 is sound. By Lemma 1, $G_{\overline{\mathbf{X}}}$ is compatible with $G_{C_{\overline{\mathbf{X}}}}$. Let $\mathbf{Z}_i \in \mathbf{Z}$ be a cluster in G_C . If $\exists \mathbf{W} \in \mathbf{W}$ and $\mathbf{Z} \in \mathbf{Z}_i$ such that $\mathbf{Z} \in An(\mathbf{W})$ in $G_{\overline{\mathbf{X}}}$, then, by Proposition 2, there exists in $G_{C_{\overline{\mathbf{X}}}}$ a directed path from \mathbf{Z}_i to some cluster $\mathbf{W}_i \in \mathbf{W}$ and, thus, \mathbf{Z}_i is an ancestor of \mathbf{W} in $G_{C_{\overline{\mathbf{X}}}}$. In this case, any existing edge $\mathbf{C}_i \rightarrow \mathbf{Z}_i$, where $\mathbf{C}_i \in \mathbf{C}$, in $G_{C_{\overline{\mathbf{X}}}}$ also exists in the graph $G_{C_{\overline{\mathbf{XZ}(\mathbf{W})}}}$, which is obtained by cutting from $G_{C_{\overline{\mathbf{X}}}}$ the edges into the \mathbf{Z} -clusters

that are not ancestors of any \mathbf{W} -clusters. On the other hand, an edge $\mathbf{C}_i \rightarrow \mathbf{Z}_i$ in $G_{\mathbf{C}_{\overline{\mathbf{X}}}}$ also exists in a C-DAG $G'_{\mathbf{C}_{\overline{\mathbf{XZ}}(\mathbf{W})}}$ compatible with $G_{\overline{\mathbf{XZ}}(\mathbf{W})}$ if $\exists W \in \mathbf{W}$, $Z \in \mathbf{Z}_i$, and $V \in \mathbf{C}_i$ such that $Z \in \text{An}(W)$ and $V \in \text{Pa}(Z)$ in $G'_{\overline{\mathbf{X}}}$. Therefore, $G_{\mathbf{C}_{\overline{\mathbf{XZ}}(\mathbf{W})}}$ has the same or more edges than $G'_{\mathbf{C}_{\overline{\mathbf{XZ}}(\mathbf{W})}}$ and, thus, a separation in $G_{\mathbf{C}_{\overline{\mathbf{XZ}}(\mathbf{W})}}$ implies the same separation in $G'_{\mathbf{C}_{\overline{\mathbf{XZ}}(\mathbf{W})}}$ which, by Theorem 1, implies the same separation in any compatible graph $G_{\overline{\mathbf{XZ}}(\mathbf{W})}$. In particular, if $(\mathbf{Y} \perp\!\!\!\perp \mathbf{Z}|\mathbf{X}, \mathbf{W})_{G_{\mathbf{C}_{\overline{\mathbf{XZ}}(\mathbf{W})}}}$, then $(\mathbf{Y} \perp\!\!\!\perp \mathbf{Z}|\mathbf{X}, \mathbf{W})_{G_{\overline{\mathbf{XZ}}(\mathbf{W})}}$. Therefore Rule 3 holds by the Rule 3 of do-calculus in causal diagrams. \square

Theorem 4. (Completeness of do-calculus). *If in a C-DAG $G_{\mathbf{C}}$ a do-calculus rule does not apply, then there is a CBN G compatible with $G_{\mathbf{C}}$ for which it also does not apply.*

Proof. Let $G_{\mathbf{C}}$ be a C-DAG over a set of clusters \mathbf{C} and G be a compatible causal diagram over a set of variables \mathbf{V} . Also, let $\mathbf{W}, \mathbf{X}, \mathbf{Y}$, and \mathbf{Z} be disjoint subsets of clusters in \mathbf{C} . Note that we abuse the notation by letting $\mathbf{W}, \mathbf{X}, \mathbf{Y}, \mathbf{Z} \subset \mathbf{V}$ also denote the set of variables contained in the clusters in $\mathbf{W}, \mathbf{X}, \mathbf{Y}, \mathbf{Z} \subset \mathbf{C}$, respectively.

We first show that if $(\mathbf{Y} \not\perp\!\!\!\perp \mathbf{Z}|\mathbf{X}, \mathbf{W})_{G_{\mathbf{C}_{\overline{\mathbf{X}}}}}$ then $(\mathbf{Y} \not\perp\!\!\!\perp \mathbf{Z}|\mathbf{X}, \mathbf{W})_{G_{\overline{\mathbf{X}}}}$ for at least one causal diagram G that is compatible with the C-DAG $G_{\mathbf{C}}$. Lemma 1 ensures that the C-DAG $G_{\mathbf{C}_{\overline{\mathbf{X}}}}$ is compatible with $G_{\overline{\mathbf{X}}}$. This, along with the completeness part of Theorem 1, implies that if $(\mathbf{Y} \not\perp\!\!\!\perp \mathbf{Z}|\mathbf{X}, \mathbf{W})_{G_{\mathbf{C}_{\overline{\mathbf{X}}}}}$, then there exists a causal diagram G compatible with $G_{\mathbf{C}}$ for which $(\mathbf{Y} \not\perp\!\!\!\perp \mathbf{Z}|\mathbf{X}, \mathbf{W})_{G_{\overline{\mathbf{X}}}}$ and, therefore, Rule 1 does not hold by the Rule 1 of do-calculus in causal diagrams.

We now show that if $(\mathbf{Y} \not\perp\!\!\!\perp \mathbf{Z}|\mathbf{X}, \mathbf{W})_{G_{\mathbf{C}_{\overline{\mathbf{XZ}}}}}$ then $(\mathbf{Y} \not\perp\!\!\!\perp \mathbf{Z}|\mathbf{X}, \mathbf{W})_{G_{\overline{\mathbf{XZ}}}}$ for at least one causal diagram G that is compatible with the C-DAG $G_{\mathbf{C}}$. Lemma 1 ensures that the C-DAG $G_{\mathbf{C}_{\overline{\mathbf{XZ}}}}$ is compatible with $G_{\overline{\mathbf{XZ}}}$. This, along with the completeness part of Theorem 1, implies that if $(\mathbf{Y} \not\perp\!\!\!\perp \mathbf{Z}|\mathbf{X}, \mathbf{W})_{G_{\mathbf{C}_{\overline{\mathbf{XZ}}}}}$, then there exists a causal diagram G compatible with $G_{\mathbf{C}}$ for which $(\mathbf{Y} \not\perp\!\!\!\perp \mathbf{Z}|\mathbf{X}, \mathbf{W})_{G_{\overline{\mathbf{XZ}}}}$ and, therefore, Rule 2 does not hold by the Rule 2 of do-calculus in causal diagrams.

Lastly, we show that if $(\mathbf{Y} \not\perp\!\!\!\perp \mathbf{Z}|\mathbf{X}, \mathbf{W})_{G_{\mathbf{C}_{\overline{\mathbf{XZ}}(\mathbf{W})}}}$ then $(\mathbf{Y} \not\perp\!\!\!\perp \mathbf{Z}|\mathbf{X}, \mathbf{W})_{G_{\overline{\mathbf{XZ}}(\mathbf{W})}}$ for at least one causal diagram G that is compatible with the C-DAG $G_{\mathbf{C}}$. Consider the causal diagram G over \mathbf{V} compatible with $G_{\mathbf{C}}$ where, for every $\mathbf{C}_i, \mathbf{C}_j \in \mathbf{C}$, any two variables $V_i, V_j \in \mathbf{V}$ such that $V_i \in \mathbf{C}_i$ and $V_j \in \mathbf{C}_j$, there is a connection $V_i \rightarrow V_j$ if $\mathbf{C}_i \rightarrow \mathbf{C}_j$ and $V_i \leftrightarrow V_j$ if $\mathbf{C}_i \leftrightarrow \mathbf{C}_j$. In this case, if for some cluster $\mathbf{Z}_i \in \mathbf{Z}$ there exist $Z \in \mathbf{Z}_i$ and $W \in \mathbf{W}$, such that $Z \in \text{An}(W)$, then, $\forall Z' \in \mathbf{Z}_i$, we have $Z' \in \text{An}(W)$. As previously discussed in the proof of the soundness of the Rule 3 (Theorem 3), in this case $G_{\mathbf{C}_{\overline{\mathbf{XZ}}(\mathbf{W})}}$ is compatible with $G_{\overline{\mathbf{XZ}}(\mathbf{W})}$. Further, in this case, for every path p in $G_{\mathbf{C}}$, there exists a path p' in G of the same form of p

and going through only one variable in each cluster in p . Therefore, every d-connection in $G_{\mathbf{C}}$ corresponds to a d-connection in G . In particular, if $(\mathbf{Y} \not\perp\!\!\!\perp \mathbf{Z}|\mathbf{X}, \mathbf{W})_{G_{\mathbf{C}_{\overline{\mathbf{XZ}}(\mathbf{W})}}}$, then $(\mathbf{Y} \not\perp\!\!\!\perp \mathbf{Z}|\mathbf{X}, \mathbf{W})_{G_{\overline{\mathbf{XZ}}(\mathbf{W})}}$ and Rule 3 does not hold in G by the Rule 3 of do-calculus in causal diagrams. \square

Theorem 5. (C-DAG as CBN) *Let $G_{\mathbf{C}}$ be a C-DAG compatible with an ADMG G . If G satisfies the truncated factorization (4) with respect to the interventional distributions, then, for any $\mathbf{X} \subseteq \mathbf{C}$, the interventional distribution $P(\mathbf{c} \setminus \mathbf{x} | \text{do}(\mathbf{x}))$ factorizes according to $G_{\mathbf{C}}$, i.e.,*

$$P(\mathbf{c} \setminus \mathbf{x} | \text{do}(\mathbf{x})) = \sum_{\mathbf{u}} P(\mathbf{u}) \prod_{k: \mathbf{C}_k \in \mathbf{C} \setminus \mathbf{X}} P(\mathbf{c}_k | \text{pa}_{\mathbf{C}_k}, \mathbf{u}'_k), \quad (5)$$

where $\text{Pa}_{\mathbf{C}_k}$ are the parents of the cluster \mathbf{C}_k , and $\mathbf{U}'_k \subseteq \mathbf{U}$ such that, for any i, j , $\mathbf{U}'_i \cap \mathbf{U}'_j \neq \emptyset$ if and only if there is a bidirected edge $(\mathbf{C}_i \leftrightarrow \mathbf{C}_j)$ between \mathbf{C}_i and \mathbf{C}_j in $G_{\mathbf{C}}$.

Proof. By assumption, the post-interventional distribution $P(\mathbf{v} \setminus \mathbf{x} | \text{do}(\mathbf{x})) = P(\mathbf{c} \setminus \mathbf{x} | \text{do}(\mathbf{x}))$ factorizes according to G by the following truncated factorization:

$$P(\mathbf{c} \setminus \mathbf{x} | \text{do}(\mathbf{x})) = \sum_{\mathbf{u}} P(\mathbf{u}) \prod_{k: \mathbf{C}_k \in \mathbf{C} \setminus \mathbf{X}} \prod_{i: V_{ki} \in \mathbf{C}_k} P(v_{ki} | \text{pa}(v_{ki}), \mathbf{u}_{ki}). \quad (14)$$

where $\text{Pa}(V_{ki})$ are the endogenous parents of V_{ki} and \mathbf{U}_{ki} are the exogenous parents of V_{ki} , including those that are shared with some other variable $V_{k'j}$ (represented by $V_{k'j} \leftrightarrow V_{ki}$ in G).

We can use the same procedure in the proof of Theorem 2 to show that, for any cluster $\mathbf{C}_k \in \mathbf{C}$, the conditional distribution $P(\mathbf{c}_k | \text{pa}(\mathbf{c}_k), \mathbf{u}'_k)$ factorizes as a product of conditional distributions over the variables in \mathbf{C}_k as shown in Equation 12. Further, from the discussion regarding the construction of the sets $\mathbf{U}'_i \cap \mathbf{U}'_j$ in the proof of Theorem 2, we also conclude that $\mathbf{U}'_i \cap \mathbf{U}'_j \neq \emptyset$ iff $\mathbf{C}_i \leftrightarrow \mathbf{C}_j$ in $G_{\mathbf{C}}$.

Therefore we can rewrite the truncated factorization over the variables $V_{ki} \in \mathbf{V}$ in G , shown in Equation 14, as a truncated factorization over the clusters $\mathbf{C}_k \in \mathbf{C}$ in $G_{\mathbf{C}}$ as follows:

$$P(\mathbf{c} \setminus \mathbf{x} | \text{do}(\mathbf{x})) = \sum_{\mathbf{u}} P(\mathbf{u}) \prod_{k: \mathbf{C}_k \in \mathbf{C} \setminus \mathbf{X}} P(\mathbf{c}_k | \text{pa}(\mathbf{c}_k), \mathbf{u}'_k). \quad (15)$$

\square

Lemma 5. (Preservation of ancestral relationships) *Let $G_{\mathbf{C}}(\mathbf{C}, \mathbf{E}_{\mathbf{C}})$ be a C-DAG compatible with causal diagram $G(\mathbf{V}, \mathbf{E})$. For any $\mathbf{C}_i \in \mathbf{C}$, $\forall V \in \mathbf{C}_i$, $(\text{An}(V)_G \cup \{V\}) \subseteq (\text{An}(\mathbf{C}_i)_{G_{\mathbf{C}}} \cup \mathbf{C}_i)$ and $(\text{De}(V)_G \cup \{V\}) \subseteq (\text{De}(\mathbf{C}_i)_{G_{\mathbf{C}}} \cup \mathbf{C}_i)$.*

Proof. Suppose that exists $V_j \in \mathbf{C}_j$ and $V \in \mathbf{C}_i$ such that V_j is an ancestor of V . If $\mathbf{C}_j = \mathbf{C}_i$, then $V_j \in \mathbf{C}_i$. Otherwise, by Proposition 2, the directed path p from V_j to V in

G is preserved and represented by a directed path from \mathbf{C}_j to \mathbf{C}_i in $G_{\mathbf{C}}$. Therefore $C_j \in An(\mathbf{C}_i)_{G_{\mathbf{C}}} \cup \mathbf{C}_i$. Then $V_j \in An(\mathbf{C}_i)_{G_{\mathbf{C}}} \cup \mathbf{C}_i$. This holds for any $V_j \in An(V)_G \cup \{V\}$, hence we obtain $(An(V)_G \cup \{V\}) \subseteq (An(\mathbf{C}_i)_{G_{\mathbf{C}}} \cup \mathbf{C}_i)$.

Analogously, suppose that there exists $V_j \in \mathbf{C}_j$ and $V \in \mathbf{C}_i$ such that V_j is a descendant of V . If $\mathbf{C}_j = \mathbf{C}_i$, then $V_j \in \mathbf{C}_i$. Otherwise, by Proposition 2, the directed path p from V to V_j in G is preserved and represented by a directed path from \mathbf{C}_i to \mathbf{C}_j in $G_{\mathbf{C}}$. Therefore, $C_j \in De(\mathbf{C}_i)_{G_{\mathbf{C}}} \cup \mathbf{C}_i$. Then $V_j \in De(\mathbf{C}_i)_{G_{\mathbf{C}}} \cup \mathbf{C}_i$ and we obtain $(De(V)_G \cup \{V\}) \subseteq (De(\mathbf{C}_i)_{G_{\mathbf{C}}} \cup \mathbf{C}_i)$. \square

Lemma 6. (Reduction to an ancestral set) *Given a C-DAG $G_{\mathbf{C}}$, let $\mathbf{A} \subseteq \mathbf{C}$. If \mathbf{A} is an ancestral set in $G_{\mathbf{C}}$ then, $Q[\mathbf{A}] = \sum_{\mathbf{C} \setminus \mathbf{A}} Q[\mathbf{C}]$ in any compatible causal diagram G , where \mathbf{A}, \mathbf{C} represent the sets of variables contained in the clusters in \mathbf{A}, \mathbf{C} , respectively.*

Proof. For any C-DAG $G_{\mathbf{C}}$, let $\mathbf{A} \subseteq \mathbf{C}$. We first show that if \mathbf{A} is an ancestral set in $G_{\mathbf{C}}$ then, for any compatible causal diagram G , the set of all variables contained in \mathbf{A} is ancestral in G .

Again we will use the same notation \mathbf{A} to also indicate the set of variables in \mathbf{V} in G contained in the cluster $\mathbf{A} \subset \mathbf{C}$ in $G_{\mathbf{C}}$.

Assume that \mathbf{A} is ancestral in $G_{\mathbf{C}}$. This means \mathbf{A} contains all $An(\mathbf{A})_{G_{\mathbf{C}}}$. Now, consider any graph G compatible with the C-DAG, $G_{\mathbf{C}}$. By Lemma 5, $An(\mathbf{A})_G \subseteq An(\mathbf{A})_{G_{\mathbf{C}}}$, so if \mathbf{A} is an ancestral set in $G_{\mathbf{C}}$, then \mathbf{A} is ancestral in G . This means that the distribution $Q[\mathbf{A}]$ induced by any compatible graph G can be calculated directly from $Q[\mathbf{C}]$ by marginalizing the clusters in $\mathbf{C} \setminus \mathbf{A}$. \square

Lemma 7. (Preservation of c-components) *Given a C-DAG $G_{\mathbf{C}}$, for any causal diagram G compatible with $G_{\mathbf{C}}$, if any two variables $V_i, V_j \in \mathbf{V}$ are in the same c-component in G , then the clusters $\mathbf{C}_i, \mathbf{C}_j \in \mathbf{C}$ containing V_i and V_j , respectively, are in the same c-component in $G_{\mathbf{C}}$.*

Proof. We show that if two variables $V_i, V_j \in \mathbf{V}$ belong to a c-component in G , then they will belong to the same c-component in $G_{\mathbf{C}}$. If V_i and V_j are in the same cluster in $G_{\mathbf{C}}$, then they are always together in a c-component of $G_{\mathbf{C}}$. If $V_i \in \mathbf{C}_i$ and $V_j \in \mathbf{C}_j$, where $\mathbf{C}_i, \mathbf{C}_j \in \mathbf{C}$ are distinct clusters, then, since V_i and V_j are in the same c-component in G , there exists a path of bidirected edges of the form $V_i \leftrightarrow \dots \leftrightarrow V'_i \leftrightarrow V'_j \leftrightarrow \dots \leftrightarrow V_j$, where $V'_i \in \mathbf{C}_i$ and $V'_j \in \mathbf{C}_j$. Then, by Definition 1, $\mathbf{C}_i \leftrightarrow \mathbf{C}_j$ in $G_{\mathbf{C}}$. This follows for every pair of variables V_i, V_j that are in the same c-component. Therefore, any path of bidirected edges between variables in G is represented by a path of bidirected edges between clusters in $G_{\mathbf{C}}$. Consequently, any c-component in G implies a c-component in $G_{\mathbf{C}}$. \square

Theorem 6. (Soundness and Completeness of ID-algorithm). *The ID-algorithm is sound and complete when applied to a C-DAG $G_{\mathbf{C}}$ for identifying causal effects of the*

form $P(\mathbf{y}|do(\mathbf{x}))$ from the observational distribution $P(\mathbf{V})$, where \mathbf{X} and \mathbf{Y} are sets of clusters in $G_{\mathbf{C}}$.

Proof. (Soundness): For any set of clusters $\mathbf{S} \subseteq \mathbf{C}$, let the quantity $Q[\mathbf{S}]$ denote the post-intervention distribution of \mathbf{S} under the intervention to all other clusters in \mathbf{C} , i.e.: $Q[\mathbf{S}] = P(\mathbf{s}|do(\mathbf{c} \setminus \mathbf{s}))$. By the truncated factorization in C-DAGs shown in Theorem 5, we have:

$$Q[\mathbf{S}] = P(\mathbf{s}|do(\mathbf{c} \setminus \mathbf{s})) = \sum_{\mathbf{u}} P(\mathbf{u}) \prod_{k: \mathbf{C}_k \in \mathbf{S}} P(\mathbf{c}_k | pa(\mathbf{c}_k), \mathbf{u}_k).$$

Let $\mathbf{X}, \mathbf{Y} \subseteq \mathbf{C}$ be distinct sets of clusters in a C-DAG $G_{\mathbf{C}}$. Also, let $\mathbf{D} = An(\mathbf{Y})$ in $G_{\mathbf{C}}[\mathbf{C} \setminus \mathbf{X}]$, where $G_{\mathbf{C}}[\mathbf{C} \setminus \mathbf{X}]$ is the subgraph of $G_{\mathbf{C}}$ containing only the clusters in \mathbf{C} not in \mathbf{X} . Since \mathbf{D} contains all ancestors of \mathbf{Y} in $G_{\mathbf{C}}[\mathbf{C} \setminus \mathbf{X}]$ and $Q[\mathbf{C} \setminus \mathbf{X}] = P(\mathbf{c} \setminus \mathbf{x}|do(\mathbf{x}))$, we have:

$$P(\mathbf{y}|do(\mathbf{x})) = \sum_{\mathbf{c} \setminus (\mathbf{x} \cup \mathbf{y})} Q[\mathbf{C} \setminus \mathbf{X}] = \sum_{\mathbf{d} \setminus \mathbf{y}} Q[\mathbf{D}]. \quad (16)$$

Note that, by Lemma 5, \mathbf{D} contains all ancestors of \mathbf{Y} in $G[\mathbf{V} \setminus \mathbf{X}]$, for any causal diagram G compatible with $G_{\mathbf{C}}$. Therefore, by Lemma 6, the reduction shown in Equation (16) is valid in any G compatible with $G_{\mathbf{C}}$.

Now, let $\mathbf{D}_1, \dots, \mathbf{D}_l$ be the c-components of the subgraph $G_{\mathbf{C}}[\mathbf{D}]$. Also, let $\mathbf{S}_1, \dots, \mathbf{S}_k$ be the c-components of $G_{\mathbf{C}}$.

Consider the quantity $Q[\mathbf{S}_j] = P(\mathbf{s}_j|do(\mathbf{c} \setminus \mathbf{s}_j))$. Given a topological order $\mathbf{C}_1 \prec \dots \prec \mathbf{C}_n$ over all clusters in $G_{\mathbf{C}}$, $Q[\mathbf{S}_j]$ is computable from $P(\mathbf{c})$ as follows (Tian 2002; Tian and Pearl 2002a):

$$Q[\mathbf{S}_j] = \prod_{\{i | \mathbf{C}_i \in \mathbf{S}_j\}} P(\mathbf{C}_i | \mathbf{C}_1, \dots, \mathbf{C}_{i-1}). \quad (17)$$

By Lemma 7, any c-component of a causal diagram G compatible with $G_{\mathbf{C}}$ is contained in some c-component \mathbf{S}_j of $G_{\mathbf{C}}$. Further, Theorem 5 shows that a factorization of any intervention distribution over the clusters in \mathbf{C} according to a C-DAG $G_{\mathbf{C}}$ is valid over the variables in \mathbf{V} in any compatible causal diagram G . Thus, in any G compatible with $G_{\mathbf{C}}$, the quantity $Q[\mathbf{S}_j]$ shown in Equation (17) corresponds to the distribution $P(\mathbf{s}_j|do(\mathbf{v} \setminus \mathbf{s}_j))$, where \mathbf{S}_j represents the set of variables contained in all clusters in \mathbf{S}_j .

By considering the Q-decomposition of $Q[\mathbf{D}]$ (i.e., $Q[\mathbf{D}] = \prod_i Q[\mathbf{D}_i]$), the ID-algorithm decomposes the problem of identifying $P(\mathbf{y}|do(\mathbf{x}))$ into the smaller sub-problems of identifying the quantities $Q[\mathbf{D}_i]$, for $i = 1, \dots, l$, from some $Q[\mathbf{S}_j]$, where $\mathbf{D}_i \subseteq \mathbf{S}_j$:

$$P(\mathbf{y}|do(\mathbf{x})) = \sum_{\mathbf{d} \setminus \mathbf{y}} \prod_i Q[\mathbf{D}_i].$$

The ID-algorithm repeatedly applies the reduction to an ancestral set and the factorization into c-components until it fails or outputs an expression for $Q[\mathbf{D}_i]$ in terms of $Q[\mathbf{S}_j]$. As previously indicated, such manipulations of the probability distribution over clusters correspond to valid manipulations in the probability distribution over variables. Therefore, if all $Q[\mathbf{D}_i]$'s are identifiable in $G_{\mathbf{C}}$, then, in any G compatible with $G_{\mathbf{C}}$, all $Q[\mathbf{D}_i]$'s are also identifiable and thus, $P(\mathbf{y}|do(\mathbf{x}))$ is identifiable.

(Completeness): Let G_C be a C-DAG and $\mathbf{X}, \mathbf{Y} \subseteq \mathbf{C}$ be two sets of clusters. We will show that if the ID-algorithm fails to identify $P(\mathbf{y}|do(\mathbf{x}))$ in G_C , then there exists a causal diagram G compatible with G_C where the effect $P(\mathbf{y}|do(\mathbf{x}))$ is not identifiable and, therefore, the ID-algorithm would fail to identify the effect $P(\mathbf{y}|do(\mathbf{x}))$ in G .

Assume that the ID-algorithm fails in G_C . Then, for some $\mathbf{X}' \subseteq \mathbf{X}$ and $\mathbf{Y}' \subseteq \mathbf{Y}$, there exists a hedge $\langle \mathcal{F}_C, \mathcal{F}'_C \rangle$ for $P(\mathbf{y}'|do(\mathbf{x}'))$ in G_C . Note that $\mathcal{F}_C, \mathcal{F}'_C$ are \mathbf{R} -rooted C-forests, where $\mathbf{R} \subseteq An(\mathbf{Y})_{G_{C\bar{\mathbf{X}}}}$, such that $\mathcal{F}'_C \subseteq \mathcal{F}_C$, $\mathcal{F}'_C \cap \mathbf{X} = \emptyset$, and $\mathcal{F}_C \cap \mathbf{X} \neq \emptyset$. We will show that we can construct a causal diagram G , compatible with G_C , with a hedge $\langle \mathcal{F}, \mathcal{F}' \rangle$ for $P(\mathbf{y}'|do(\mathbf{x}'))$.

Consider the causal diagram G over \mathbf{V} where, for every $\mathbf{C}_i, \mathbf{C}_j \in \mathbf{C}$, any two variables $V_i, V_j \in \mathbf{V}$ such that $V_i \in \mathbf{C}_i$ and $V_j \in \mathbf{C}_j$, there is a connection $V_i \rightarrow V_j$ if $\mathbf{C}_i \rightarrow \mathbf{C}_j$ and $V_i \leftrightarrow V_j$ if $\mathbf{C}_i \leftrightarrow \mathbf{C}_j$. Note that, by definition of cluster causal diagrams, such edges do not violate the compatibility of G with G_C . Now, for each $\mathbf{C}_i \in \mathbf{C}$, let $V_{i1} \prec \dots \prec V_{iN_i}$ be a topological order over the set of variables that constitute \mathbf{C}_i . Let the internal structure of \mathbf{C}_i be a chain with an edge $V_{ik} \rightarrow V_{i(k+1)}$ and a bidirected edge $V_{ik} \leftrightarrow V_{i(k+1)}$, for $k = 1, \dots, N_i$. Note that such an internal structure is consistent with the topological order and does not violate the compatibility with G_C .

Construct \mathcal{F}' and \mathcal{F} , edge subgraphs of G , as follows: consider the subgraphs over the variables that are contained in the clusters in \mathcal{F}_C and \mathcal{F}'_C , respectively; remove all outgoing edges from any variable in \mathbf{R} ; then remove a set of directed edges so that every observable variable has at most one child, while preserving \mathbf{R} as the maximal root set. This construction process ensures that the resulting edge subgraphs \mathcal{F}' and \mathcal{F} satisfy the properties of \mathbf{R} -rooted C-forests in G_C . Note that, since the variables within the clusters form a chain from a cluster to another, the ancestral relations in \mathcal{F}_C and \mathcal{F}'_C are preserved so that \mathcal{F}' and \mathcal{F} have \mathbf{R} as a root set. Also, since no bidirected edge is removed, both \mathcal{F}' and \mathcal{F} have only one c-component. Lastly, since \mathcal{F} and \mathcal{F}' are constructed over the variables in \mathcal{F}_C and \mathcal{F}'_C , respectively, we have $\mathcal{F}' \subseteq \mathcal{F}$, $\mathcal{F}' \cap \mathbf{X} = \emptyset$, and $\mathcal{F} \cap \mathbf{X} \neq \emptyset$. Therefore, for some $\mathbf{X}' \subseteq \mathbf{X}$ and $\mathbf{Y}' \subseteq \mathbf{Y}$, there exist \mathbf{R} -rooted C-forests $\mathcal{F}, \mathcal{F}'$ that form a hedge for $P(\mathbf{y}'|do(\mathbf{x}'))$ in G , which, in turn, implies that the ID-algorithm will fail to identify $P(\mathbf{y}|do(\mathbf{x}))$ in G . \square

Theorem 7. Let G_C be a C-DAG compatible with an ADMG G . Assume G is induced by an SCM \mathcal{M} , then there exists an SCM \mathcal{M}_C over macro-variables \mathbf{C} such that its induced causal diagram is G_C and, for any set of counterfactual variables $\mathbf{Y}_x, \dots, \mathbf{Z}_w$ where $\mathbf{Y}, \mathbf{X}, \dots, \mathbf{Z}, \mathbf{W} \subseteq \mathbf{C}$, $P_{\mathcal{M}}(\mathbf{y}_x, \dots, \mathbf{z}_w) = P_{\mathcal{M}_C}(\mathbf{y}_x, \dots, \mathbf{z}_w)$.

Proof. Given a partition $\mathbf{C} = \{\mathbf{C}_1, \dots, \mathbf{C}_K\}$ of \mathbf{V} , denote V_{ij} the j th variable in cluster $\mathbf{C}_i \in \mathbf{C}$. We assume that G over \mathbf{V} is induced by an SCM $\mathcal{M} = \langle \mathbf{V}, \mathbf{U}, \mathcal{F}, P(\mathbf{U}) \rangle$, where each variable $V_{ij} \in \mathbf{C}_i$ is given by a function $f_{ij} \in \mathcal{F}$ as follows:

$$V_{ij} \leftarrow f_{ij}(\mathbf{Pa}_{ij}, \mathbf{U}_{ij}),$$

where $\mathbf{Pa}_{ij} \subseteq \mathbf{V}$ is the set of endogenous variables representing parent nodes of V_{ij} in G and $\mathbf{U}_{ij} \subseteq \mathbf{U}$ is the set of exogenous variable representing the latent parents of V_{ij} .

Now, let $\tilde{\mathbf{C}} = \{\tilde{\mathbf{C}}_1, \dots, \tilde{\mathbf{C}}_K\}$ be a set of “macro-variables” where, for $i = 1 \dots K$, letting N_i be the number of variables in cluster \mathbf{C}_i , $\tilde{\mathbf{C}}_i = (V_{i1}, \dots, V_{iN_i})$ is a N_i -dimensional random vector. We construct a function for $\tilde{\mathbf{C}}_i$ as follows:

$$\tilde{\mathbf{C}}_i = \begin{pmatrix} V_{i1} \\ \vdots \\ V_{iN_i} \end{pmatrix} = \begin{pmatrix} f'_{i1}(\mathbf{Pa}'_{i1}, \mathbf{U}'_{i1}) \\ \vdots \\ f'_{iN_i}(\mathbf{Pa}'_{iN_i}, \mathbf{U}'_{iN_i}) \end{pmatrix}, \quad (18)$$

where, for $j = 1, \dots, N_i$, $\mathbf{Pa}'_{ij} = Pa(An(V_{ij}) \cap \mathbf{C}_i) \setminus \mathbf{C}_i$, i.e., the set of endogenous variables that are not in \mathbf{C}_i but are parents of nodes in G that are both ancestors of V_{ij} and in \mathbf{C}_i ; and $\mathbf{U}'_{ij} = \{\mathbf{U}_{ik} : V_{ik} \in An(V_{ij}) \cap \mathbf{C}_i\}$, i.e., the set of exogenous variables representing latent parents of nodes that are both ancestors of V_{ij} and in \mathbf{C}_i . f'_{ij} results from $f_{ij}(\mathbf{Pa}_{ij}, \mathbf{U}_{ij})$ in which each endogenous variable V_{ik} in \mathbf{Pa}_{ij} that is in \mathbf{C}_i is recursively replaced by its own function f_{ik} until none of the arguments are in \mathbf{C}_i (this is feasible due to the acyclicity of SCMs and can be done following the topological order over the variables in \mathbf{C}_i induced by G). Therefore, f'_{ij} is just an expanded form of f_{ij} in terms of endogenous variables that are not in \mathbf{C}_i , and the assignment $V_{ij} \leftarrow f'_{ij}(\mathbf{Pa}'_{ij}, \mathbf{U}'_{ij})$ does not change the original assignment of V_{ij} given a fixed \mathbf{U} value.

Now, define $\mathbf{Pa}'_{\tilde{\mathbf{C}}_i} = \mathbf{Pa}'_{i1} \cup \dots \cup \mathbf{Pa}'_{iN_i}$ and $\mathbf{U}'_{\tilde{\mathbf{C}}_i} = \mathbf{U}'_{i1} \cup \dots \cup \mathbf{U}'_{iN_i}$. Take the smallest subset \mathbf{C}' of \mathbf{C} such that $\mathbf{Pa}'_{\tilde{\mathbf{C}}_i} \subseteq \mathbf{C}'$ and denote $\mathbf{Pa}_{\tilde{\mathbf{C}}_i}$ as the set of macro-variables in $\tilde{\mathbf{C}}$ corresponding to the clusters in \mathbf{C}' . We rewrite the function for $\tilde{\mathbf{C}}_i$ in Eq. (18) as

$$\tilde{\mathbf{C}}_i \leftarrow f_i(\mathbf{Pa}_{\tilde{\mathbf{C}}_i}, \mathbf{U}_{\tilde{\mathbf{C}}_i}) \equiv \begin{pmatrix} f'_{i1}(\mathbf{Pa}'_{i1}, \mathbf{U}'_{i1}) \\ \vdots \\ f'_{iN_i}(\mathbf{Pa}'_{iN_i}, \mathbf{U}'_{iN_i}) \end{pmatrix} \quad (19)$$

We construct an SCM $\mathcal{M}_{\tilde{\mathbf{C}}} = \langle \tilde{\mathbf{C}}, \mathbf{U}, \mathcal{F}_{\tilde{\mathbf{C}}}, P(\mathbf{U}) \rangle$, where $\mathcal{F}_{\tilde{\mathbf{C}}}$ consists of the set of functions f_i specified in Eq. (19), for $i = 1, \dots, K$.

This SCM $\mathcal{M}_{\tilde{\mathbf{C}}}$ induces a causal diagram $G_{\tilde{\mathbf{C}}}$ over macro-variables $\tilde{\mathbf{C}}$ where $\tilde{\mathbf{C}}_i \leftarrow \tilde{\mathbf{C}}_j$ iff $\tilde{\mathbf{C}}_j \in \mathbf{Pa}_{\tilde{\mathbf{C}}_i}$; and $\tilde{\mathbf{C}}_i \leftrightarrow \tilde{\mathbf{C}}_j$ iff $\mathbf{U}_{\tilde{\mathbf{C}}_i} \cap \mathbf{U}_{\tilde{\mathbf{C}}_j} \neq \emptyset$. We have that

1. $\tilde{\mathbf{C}}_j \in \mathbf{Pa}_{\tilde{\mathbf{C}}_i}$ iff $\mathbf{C}_j \cap \mathbf{Pa}'_{\mathbf{C}_i} \neq \emptyset$ iff $\exists k, \mathbf{C}_j \cap \mathbf{Pa}'_{ik} \neq \emptyset$ iff a node in $\tilde{\mathbf{C}}_j$ is a parent of some node in \mathbf{C}_i since every node in \mathbf{Pa}'_{ik} is a parent of some node in \mathbf{C}_i .
2. $\mathbf{U}_{\tilde{\mathbf{C}}_i} \cap \mathbf{U}_{\tilde{\mathbf{C}}_j} \neq \emptyset$ iff $\exists k, k', \mathbf{U}'_{ik} \cap \mathbf{U}'_{jk'} \neq \emptyset$ iff there exists a latent node in $\mathbf{U}'_{ik} \cap \mathbf{U}'_{jk'}$ that is a shared parent of a node in \mathbf{C}_i and a node in \mathbf{C}_j since every latent node in \mathbf{U}'_{ik} is a parent of some node in \mathbf{C}_i and every latent node in $\mathbf{U}'_{jk'}$ is a parent of some node in \mathbf{C}_j .

Therefore, the causal diagram $G_{\tilde{\mathbf{C}}}$ is exactly the C-DAG G_C of G given by Definition 1.

By the construction of the set of functions f_i in Eq. (19), we have $\tilde{\mathbf{C}}_i(\mathbf{u}) = (V_{i1}(\mathbf{u}), \dots, V_{iN_i}(\mathbf{u})) = \mathbf{C}_i(\mathbf{u})$ for any fixed $\mathbf{U} = \mathbf{u}$ value. Performing an intervention $do(\tilde{\mathbf{C}}_i = \mathbf{c}_i)$ in $\mathcal{M}_{\tilde{\mathbf{C}}}$ for any i replaces the function f_i in Eq. (19) with $\tilde{\mathbf{C}}_i \leftarrow \mathbf{c}_i$ which is the same operation as performing an intervention $do(\mathbf{C}_i = \mathbf{c}_i)$ in \mathcal{M} . Therefore, for any $\mathbf{Y}, \mathbf{X} \subseteq \mathbf{C}$ and corresponding macro-variables $\tilde{\mathbf{Y}}, \tilde{\mathbf{X}}$, we have $\mathbf{Y}_{\mathbf{x}}(\mathbf{u}) = \tilde{\mathbf{Y}}_{\tilde{\mathbf{x}}}(\mathbf{u})$. We obtain that for any set of counterfactual variables $\mathbf{Y}_{\mathbf{x}} \dots \mathbf{Z}_{\mathbf{w}}$ where $\mathbf{Y}, \mathbf{X}, \dots, \mathbf{Z}, \mathbf{W} \subseteq \mathbf{C}$,

$$\begin{aligned}
P_{\mathcal{M}}(\mathbf{y}_{\mathbf{x}} \dots \mathbf{z}_{\mathbf{w}}) &= \sum_{\mathbf{u} | \mathbf{Y}_{\mathbf{x}}(\mathbf{u})=\mathbf{y}, \dots, \mathbf{Z}_{\mathbf{w}}(\mathbf{u})=\mathbf{z}} P(\mathbf{u}) \\
&= \sum_{\mathbf{u} | \tilde{\mathbf{Y}}_{\tilde{\mathbf{x}}}(\mathbf{u})=\tilde{\mathbf{y}}, \dots, \tilde{\mathbf{Z}}_{\tilde{\mathbf{w}}}(\mathbf{u})=\tilde{\mathbf{z}}} P(\mathbf{u}) \\
&= P_{\mathcal{M}_{\tilde{\mathbf{C}}}}(\tilde{\mathbf{y}}_{\tilde{\mathbf{x}}} \dots \tilde{\mathbf{z}}_{\tilde{\mathbf{w}}}). \tag{20}
\end{aligned}$$

□