EXTENSION OF IRREDUCIBILITY RESULTS ON GENERALISED LAGUERRE POLYNOMIALS $L_n^{(-1-n-s)}(x)$

SARANYA G. NAIR AND T. N. SHOREY

ABSTRACT. We consider the irreducibility of Generalised Laguerre Polynomials for negative integral values given by $L_n^{(-1-n-s)}(x) = \sum_{j=0}^n \binom{n-j+s}{n-j} \frac{x^j}{j!}$. For different values of s, this family gives polynomials which are of great interest. It was proved earlier that for $0 \le s \le 60$, these polynomials are irreducible over \mathbb{Q} . In this paper we improve this result up to $s \le 88$.

1. Introduction

For a positive integer n and real number α , the Generalised Laguerre Polynomial (GLP) is defined as

(1)
$$L_n^{(\alpha)}(x) = \sum_{j=0}^n \frac{(n+\alpha)(n-1+\alpha)\cdots(j+1+\alpha)}{j!(n-j)!} (-x)^j.$$

These polynomials were discovered around 1880 and they have been extensively studied in various branches of mathematics and mathematical physics. The algebraic properties of GLP were first studied by Schur [8],[9] where he established the irreducibility of $L_n^{(\alpha)}(x)$ for $\alpha \in \{0, 1, -n - 1\}$, gave a formula for the discriminant $\Delta_n^{(\alpha)}$ of $\mathcal{L}_n^{(\alpha)}(x) = n! L_n^{(\alpha)}(x)$ by

$$\Delta_n^{(\alpha)} = \prod_{j=1}^n j^j (\alpha + j)^{j-1}$$

and calculated their assosciated Galois groups. For an account of results obtained on GLP, we refer to [3],[7].

Let $f(x) \in \mathbb{Q}[x]$ with deg f = n. By irreducibility of a polynomial, we shall always mean its irreducibility over \mathbb{Q} . We observe that if a polynomial of degree n has a factor of degree k < n, then it has a factor of degree n - k. Therefore given a polynomial of degree n, we always consider factors of degree k where $1 \le k \le \frac{n}{2}$. If the argument α of (1) is a negative integer, we see that the constant term of $L_n^{(\alpha)}(x)$ vanishes if and only if $n \ge |\alpha| = -\alpha$ and then $L_n^{(\alpha)}(x)$ is reducible. Therefore we assume that

²⁰¹⁰ Mathematics Subject Classification: Primary 11A41, 11B25, 11N05, 11N13, 11C08, 11Z05. Keywords: Irreducibility, Laguerre Polynomials, Primes, Newton Polygons.

Nair and Shorey

 $\alpha \leq -n-1$. We write $\alpha = -n-s-1$ where s is a non-negative integer. We have

(2)
$$L_n^{(-n-s-1)}(x) = \sum_{j=0}^n (-1)^n \frac{(n+s-j)!}{(n-j)!s!} \frac{x^j}{j!}.$$

Borrowing the notation from [11], we consider the following polynomial

$$g(x) := g(x, n, s) = (-1)^n L_n^{(-n-s-1)} = \sum_{j=0}^n \binom{n+s-j}{n-j} \frac{x^j}{j!} = \sum_{j=0}^n b_j \frac{x^j}{j!}$$

where $b_j = \binom{n+s-j}{n-j}$ for $0 \le j \le n$. Thus $b_n = 1, b_0 = \binom{n+s}{s} = \frac{(n+1)\cdots(n+s)}{s!}$. We observe that g(x) is irreducible if and only if $L_n^{(-n-s-1)}(x)$ is irreducible. The aim of this paper is to discuss the irreducibility of g(x). We consider more general polynomial

$$G(x) := G(x, n, s) = \sum_{j=0}^{n} a_j b_j \frac{x^j}{j!}$$

such that $a_j \in \mathbb{Z}$ for $0 \leq j \leq n$ with $|a_0| = |a_n| = 1$. If $a_j = 1$ for $0 \leq j \leq n$, we have G(x) = g(x). We write

$$g_1(x) := n! g(x) = n! \sum_{j=0}^n \binom{n+s-j}{n-j} \frac{x^j}{j!}$$

and

$$G_1(x) = n!G(x)$$

so that g_1 and G_1 are monic polynomials with integer coefficients of degree n. The irreducibility of $g_1(x)$ and $G_1(x)$ implies the irreducibility of g(x) and G(x) respectively. We begin with the following result by Sinha and Shorey [11] on G(x).

Lemma 1.1. Let $s \le 92$. Then G(x) = G(x, n, s) has no factor of degree $k \ge 2$ except when $(n, k, s) \in \{(4, 2, 7), (4, 2, 23), (9, 2, 19), (9, 2, 47), (16, 2, 14), (16, 2, 34), (16, 2, 89), (9, 3, 47), (16, 3, 19), (10, 5, 4)\}.$

We re-state Lemma 1.1 as follows:

Let $s \leq 92$. Assume that G(x) has a factor of degree greater than or equal to 2. Then $(n, k, s) \in \{(4, 2, 7), (4, 2, 23), (9, 2, 19), (9, 2, 47), (16, 2, 14), (16, 2, 34), (16, 2, 89), (9, 3, 47), (16, 3, 19), (10, 5, 4)\}$. We check that g(x) is irreducible when $n \in \{4, 9, 10, 16\}$. Therefore by Lemma 1.1 with G(x) = g(x) we have

Lemma 1.2. Let $n \ge 3$ and $s \le 92$. Then $g_1(x)$ is either irreducible or linear factor times an irreducible polynomial.

The irreducibility of $g_1(x)$ was proved by Schur [8] for s = 0, by Hajir [2] for s = 1, by Sell [10] for s = 2 and by Hajir [3] for $3 \le s \le 8$. We shall prove

Theorem 1. $g_1(x)$ is irreducible for $9 \le s \le 88$.

Nair and Shorey [6] and Jindal, Laishram and Sarma [5] already proved the irreducibility of $g_1(x)$ in the range of $9 \le s \le 22$ and $23 \le s \le 60$ respectively. But our proof of Theorem 1 is new. The proofs of [6] and [5] depend on the method of Hajir in [3] whereas our proof depends on Lemma 1.2 which is a direct consequence of Lemma 1.1. We could not cover the cases of $88 \le s \le 92$ due to computational limitations.

At this point, we pause with a digression concerning notation. In order to dispell possible confusion in the reader concerning notation, it is worth pointing out that the notation $L_n^{\langle s \rangle}(x)$ (resp., $\mathcal{L}_n^{\langle s \rangle}(x)$) was used by the authors in [3], [5] and [7] to denote the polynomial g(x, n, s) (resp., $g_1(x, n, s)$).

2. Preliminaries

From now onwards we shall assume that $s \ge 9$. For a real number α , we write $[\alpha]$ to be the largest integer not exceeding α .

Let
$$f(x) = \sum_{j=0}^{m} d_j x^j \in \mathbb{Z}[x]$$
 with $d_0 d_m \neq 0$ and let p be a prime. For an integer x ,

let $\nu(x) = \nu_p(x)$ be the highest power of p dividing x and we write $\nu(0) = \infty$. Let S be the following set of points in the extended plane

$$S = \{(0, \nu(d_m)), (1, \nu(d_{m-1})), (2, \nu(d_{m-2})), \dots, (m, \nu(d_0))\}.$$

Consider the lower edges along the convex hull of these points. The left most endpoint is $(0, \nu(d_m))$ and the right most endpoint is $(m, \nu(d_0))$. The endpoints of each edge belong to S and the slopes of the edges increase strictly from left to right. The polygonal path formed by these edges is called the Newton polygon of f(x) with respect to the prime p and we denote it by $NP_p(f)$. The endpoints of the edges of $NP_p(f)$ are called the vertices of $NP_p(f)$. We begin with a very useful result, due to Filaseta [1], giving a criterion on the factorisation of a polynomial in terms of the maximum slope of the edges of its Newton polygon.

Lemma 2.1. Let l, k, m be integers with $m \ge 2k > 2l \ge 0$. Suppose $h(x) = \sum_{j=0}^{m} b_j x^j \in \mathbb{Z}[x]$ and p be a prime such that $p \nmid b_m$ and $p \mid b_j$ for $0 \le j \le m - l - 1$ and the right most edge of $NP_p(h)$ has slope $< \frac{1}{k}$. Then for any integers a_0, a_1, \ldots, a_m with $p \nmid a_0 a_m$, the polynomial $f(x) = \sum_{j=0}^{m} a_j b_j x^j$ cannot have a factor with degree in [l+1,k].

The next result is Lemma 4.2 of [12] with a = 0.

Lemma 2.2. Let a_0, a_1, \dots, a_n denote arbitrary integers and

$$h(x) = \sum_{j=0}^{n} a_j \frac{x^j}{j!}.$$

Nair and Shorey

Assume that h(x) has a factor of degree $k \ge 1$. Suppose that there exists a prime p > k such that p divides $n(n-1)\cdots(n-k+1)$. Then p divides a_0a_n .

For a positive integer l and a prime p, let $\nu_p(l)$ be the maximal power of p dividing l.

Lemma 2.3. Let p be a prime. For any integer $l \ge 1$, write l in base p as $l = l_t p^t + l_{t-1} p^{t-1} + \cdots + l_1 p + l_0$ where $0 \le l_i \le p-1$ for $0 \le i \le t$ and $l_t > 0$. Then

$$\nu_p(l!) = \frac{l - \sigma_p(l)}{p - 1}$$

where $\sigma_p(l) = l_t + l_{t-1} + \dots + l_1 + l_0$.

This is due to Legendre. For a proof, see [4, Ch.17, p 263]. As a consequence we have

(3)
$$\nu_p\left(\binom{m}{t}\right) = \frac{\sigma_p(t) + \sigma_p(m-t) - \sigma_p(m)}{p-1}$$

Lemma 2.4. Assume that $g_1(x)$ is a linear factor times an irreducible polynomial. Let p be a prime dividing n and $s < p^2$. Then

$$d + \left[\frac{s}{p}\right] \ge p$$

where $d \equiv \frac{n}{p} \pmod{p}$ for $1 \le d < p$.

The assertion of Lemma 2.4 was proved in [5, Corollary 3.2] under the assumption of p dividing n_1 where

(4)
$$n = n_0 \cdot n_1$$
 with $gcd(n_0, n_1) = 1$

and

(5)
$$n_1 = \prod_{\substack{p \mid \gcd(n, \binom{n+s}{s})}} p^{\operatorname{ord}_p(n)}$$

Therefore n_0 is the largest divisor of n which is coprime to $\binom{n+s}{s}$. Thus the assumption p dividing n_1 in [5, Corollary 3.2] is replaced by p dividing n in Lemma 2.4 when $g_1(x)$ is linear factor times an irreducible polynomial.

Proof. We apply Lemma 2.2 with $h(x) = g_1(x)$ and k = 1 to conclude that

(6)
$$p|\frac{(n+1)\cdots(n+s)}{s!}$$

If $\nu_p(n) > s$, then $\nu_p\left(\frac{(n+1)\cdots(n+s)}{s!}\right) \le \nu_p(\frac{s!}{s!}) \le 0$ contradicting (6). Therefore $\nu_p(n) \le s < p^2$ and hence $1 \le \nu_p(n) < 2$. We write,

n = pD where gcd(D, p) = 1

and $s = ps_1 + s_0$ where $1 \le s_1 < p, 0 \le s_0 < p$. Then $n + s = p(D + s_1) + s_0$ which implies that $\sigma_p(n + s) = \sigma_p(D + s_1) + s_0$. Now we argue as in [5, Lemma 3.1] for deriving from (3) and (6) that

$$1 \leq \nu_p \left(\binom{n+s}{s} \right) = \frac{\sigma_p(n) + \sigma_p(s) - \sigma_p(n+s)}{p-1}$$
$$= \frac{\sigma_p(D) + s_1 + s_0 + -\sigma_p(D+s_1) - s_0}{p-1}$$
$$= \frac{\sigma_p(D) + s_1 - \sigma_p(D+s_1)}{p-1}$$
$$= \nu_p \left(\binom{D+s_1}{s_1} \right)$$
$$= \nu_p \left(\binom{(D+1)\cdots(D+s_1)}{s_1!} \right)$$
$$= \nu_p((D+1)\cdots(D+s_1))$$
$$= \nu_p(D+j) \text{ for exactly one } j \text{ with } 1 \leq j \leq s_1$$

since $s_1 < p$. Hence $D + \left[\frac{s}{p}\right] = D + s_1 \ge D + j \equiv 0 \pmod{p}$. Since gcd(D, p) = 1, we observe from (3) that $D = \frac{n}{p} \equiv d \pmod{p}$ where $1 \le d < p$ by looking at p- adic representation of n. Hence $d + \left[\frac{s}{p}\right] \ge p$ where $d \equiv \frac{n}{p} \pmod{p}$ for $1 \le d < p$. \Box

Next, we prove

Lemma 2.5. Assume that $g_1(x)$ is linear factor times an irreducible polynomial. Then for $n \leq 127$ and $s \leq 103$, $g_1(x)$ is irreducible.

This result is proved in [6, Lemma 2.10] without the assumption that $g_1(x)$ is linear factor times an irreducible polynomial. But Lemma 2.5 suffices for our purpose. We give here a proof of this particular case, as it involves considerably less computations.

Proof. Let $n \leq 127$ and $s \leq 103$. Since $g_1(x)$ is linear factor times an irreducible polynomial, we see that $n_0 = 1$. Assume that $n \geq s+2$. Then $\max(\frac{n+s}{2}, n-1) = n-1$ and we derive from [3, Lemma 4.1] that $g_1(x)$ is irreducible if n is prime. Now we check that $g_1(x)$ is irreducible for all pairs (n, s) with n composite and $n \geq s+2$. Next we assume that n < s+2. Then $\max(\frac{n+s}{2}, n-1) = \frac{n+s}{2}$. We determine all pairs (n, s) such that n < s+2 and there exists a prime p satisfying $\frac{n+s}{2} . Then$ $<math>g_1(x)$ is irreducible for all these pairs (n, s) by [3, Lemma 4.1]. For the remaining pairs (n, s) with n < s+2, we check that $g_1(x)$ is irreducible. \Box

We close this section by stating the following result which is Lemma 3.6 from [5] with i = 1.

Lemma 2.6. Let p|n(s+1) and $\nu_p\left(\binom{n+s}{s}\right) = u$. Then $g_1(x)$ cannot have a factor of degree 1 if any of the following condition holds:

Nair and Shorey

(i)
$$u = 0$$

(ii) $u > 0, p > 2$ and $\max\left\{\frac{u+1}{p}, \frac{\nu_p(n+s-z_0)-\nu_p(n)}{z_0+1}\right\} < 1$, where $z_0 \equiv n+s \pmod{p}$
with $1 \le z_0 < p$.

3. Proof of Theorem 1

For c > 1 and $s \ge c^2$, we consider the following set given by

$$H_{s,c} = \{n \in \mathbb{N}, n > 127 \text{ and for } p | n, p^{\nu_p(n)} \le s \text{ and if } p > sc^{-1} \text{ then } d + \left[\frac{s}{p}\right] \ge p\}$$

where $1 \leq d < p$ and $d \equiv \frac{n}{p} \pmod{p}$. Since $p \geq sc^{-1} \geq \sqrt{s}$, we derive from Lemmas 2.4 and 2.5 that it suffices to prove the irreducibility of $g_1(x) = g_1(x, n, s)$ with $n \in H_{s,c}$. We partition $H_{s,c}$ as $H_{s,c,1}$ and $H_{s,c,2}$ given by

$$H_{s,c,1} = \{n \in H_{s,c} \text{ such that } P(n) \le [sc^{-1}]\}$$

and

$$H_{s,c,2} = \{n \in H_{s,c} \text{ such that } P(n) > [sc^{-1}]\}.$$

Let $9 \le s \le 88$. By taking $c \in \{3, 3.42, 5.5, 7.7\}$ we compute $H_{s,c,1}$ and $H_{s,c,2}$ and hence $H_{s,c}$ for $9 \le s \le 88$. We give some details regarding the computations of $H_{s,c}$ below. For example, for s = 80, and c = 7.7, the cardinality of $H_{s,c}$, $|H_{s,c}| = 1538$ and for s = 85, and c = 7.7, the cardinality of $H_{s,c}$, $|H_{s,c}| = 2466$. The following table gives the c values which we have chosen for each s to compute the set $H_{s,c}$.

s	С
$9 \le s \le 11$	3
$12 \le s \le 35$	3.42
$36 \le s \le 60$	5.5
$61 \le s \le 88$	7.7

For each $n \in H_{s,c}$, we apply Lemma 2.6 to derive that $g_1(x) = g(x, n, s)$ is irreducible except for $(n, s) \in T$ where

$$\begin{split} T = & \{(272, 17), (144, 21), (144, 23), (144, 25), (144, 26), (312, 26), (600, 26), (216, 29), (216, 31), \\ & (720, 31), (240, 35), (1440, 35), (288, 40), (288, 41), (216, 42), (216, 44), (216, 47), (288, 47), \\ & (288, 48), (216, 49), (144, 51), (288, 51), (144, 53), (216, 53), (288, 53), (4320, 55), (216, 59), \\ & (216, 63), (288, 63), (432, 63), (672, 63), (180, 71), (192, 71), (216, 71), (216, 79), (576, 79), \\ & (144, 80), (192, 80), (216, 80), (320, 80), (432, 80), (576, 80), (720, 80), (4320, 80) \} \end{split}$$

We use IrreducibilityQ command in Mathematica to check that $g_1(x)$ is irreducible for all these values of $(n, s) \in T$.

Remark 3.1. It is not necessary to compute $H_{s,c}$ for all values of s. For a fixed c, if $[sc^{-1}] = [(s+1)c^{-1}]$ and $P(s+1) \leq [sc^{-1}]$, then $H_{s,c} = H_{s+1,c}$. The assertion follows from the definitions of $H_{s,c,1}$ and $H_{s,c,2}$. Therefore $H_{s,c} = H_{s+1,c}$ for $s \in C$.

 $\mathbf{6}$

Extension of Irreducibility results on Generalised Laguerre Polynomials $L_n^{(-1-n-s)}(x)$

7

 $\{19, 27, 29, 34\}$ with c = 3.42, $s \in \{39, 41, 47, 49, 53, 55, 59\}$ with c = 5.5 and $s \in \{62, 69, 71, 74, 79, 83, 87\}$ with c = 7.7.

Remark 3.2. We can take c = 5.5 or c = 7.7 according as s at least 36 or $s \ge 60$ respectively. But we cannot take c more than $3.42\frac{s}{s-1}$ without sharpening [6, Corollary 6]. Consequently we get sets $H_{s,c}$ of smaller size when $s \ge 36$ and this reduces the computations.

Acknowledgements. The computations for calculating $H_{s,c}$ have been carried out by using SAGE, and all other computations have been done in MATHEMATICA. We thank Ankita Jindal for helping us with the computations. The second author is supported by INSA during the period of this work.

References

- M. Filaseta, The irreducibility of all but finitely many Bessel polynomials, Acta Math. 174 (1993), 383-397.
- [2] F. Hajir, Some A_n -extensions obtained from generalized Laguerre polynomials Journal of Number Theory 50 (1995), 206-212.
- [3] F. Hajir, Algebraic properties of a family of generalized Laguerre polynomials, Canad. J. Math. 61 (2009), 583-603.
- [4] H. Hasse, *Number theory*, Classics in Mathematics, Springer-Verlag, Berlin 2002.
- [5] A. Jindal, S. Laishram and R. Sarma, Irreducibility and Galois groups of Generalised Laguerre Polynomials $L_n^{(-1-n-r)}(x)$, J. Number Theory **183** (2018), 388-406.
- [6] S. G. Nair and T. N Shorey, Irreducibility of Laguerre Polynomial $L_n^{(-1-n-r)}(x)$, Indag. Math. **26** (2015), 615-625.
- [7] S. G. Nair and T.N. Shorey, Generalised Laguerre Polynomials with applications, Math. Student 86, (2017), 87-101.
- [8] I. Schur, Gleichungen ohne Affekt In: Gesammelte Abhandlungen, Band III, Springer-Verlag, Berlin, 1973, 191-197.
- [9] I. Schur, Affektlose Gleichungen in der Theorie der Laguerreschen und Hermiteschen Polynome, In: Gesammelte Abhandlungen, Band III, Springer-Verlag, Berlin-New York, 1973, 227-233.
- [10] E.A Sell, On a certain family of generalized Laguerre polynomials, J. Number Theory 107 (2004), 266-281.
- [11] S. B. Sinha and T.N. Shorey, Extension of Laguerre Polynomials with negative arguments, Indag. Math, to appear.
- [12] T. N. Shorey and R. Tijdeman, Generalizations of some irreducibility results by Schur, Acta Arith. 145 (2010), 341-371.

DEPARTMENT OF MATHEMATICS, BITS PILANI, K K BIRLA GOA CAMPUS, GOA- 403726

Email address: saranyan@goa.bits-pilani.ac.in

NATIONAL INSTITUTE OF ADVANCED STUDIES, IISC CAMPUS, BANGALORE, 560012

Email address: shorey@math.iitb.ac.in