

## ON SYMBOL CORRESPONDENCES FOR QUARK SYSTEMS

P.A.S. ALCÂNTARA AND P. DE M. RIOS

ABSTRACT. We present the characterization of symbol correspondences for mechanical systems that are symmetric under  $SU(3)$ , which we refer to as quark systems. The quantum systems are the unitary irreducible representations of  $SU(3)$ , denoted by  $Q(p, q)$ ,  $p, q \in \mathbb{N}_0$ , with their operator algebras. We study the cases when the classical phase space is a coadjoint orbit: either the complex projective plane  $\mathbb{C}P^2$  or the flag manifold that is the total space of fiber bundle  $\mathbb{C}P^1 \hookrightarrow \mathcal{E} \rightarrow \mathbb{C}P^2$ . In the first case, we refer to pure-quark systems and the characterization of their correspondences is given in terms of characteristic numbers, similarly to the case of spin systems, cf. [18]. In the second case, we refer to generic quark systems and the characterization of their correspondences is given in terms of characteristic matrices, which introduces various novel features. Furthermore, we present the  $SU(3)$  decomposition of the product of quantum operators and their corresponding twisted products of classical functions, for both pure and generic quark systems. In preparation for asymptotic analysis of these twisted products, we also present the  $SU(3)$  decomposition of the pointwise product of classical functions.

## CONTENTS

1. Introduction	2
2. On the representations of $SU(3)$	5
2.1. Irreducible unitary representations	7
2.2. Clebsch-Gordan series and space of operators	10
2.3. Decomposition of the operator product	13
2.4. Mixed Casimir operators and symmetric CG coefficients	15
2.5. Wigner symbols and the operator product	23
2.6. Coadjoint orbits	27
3. Pure-quark systems	30
3.1. Classical pure-quark system	30
3.2. Quantum pure-quark systems	31
3.3. Symbol correspondences for pure-quark systems	35
3.4. Twisted product for pure-quark system	45
4. Generic quark systems	49
4.1. Classical generic quark system	49
4.2. Quantum generic quark system	52
4.3. Symbol correspondences for generic quark systems	53
4.4. Twisted products for generic quark systems	60

2010 *Mathematics Subject Classification.* 17B08, 20C35, 22E46, 22E70, 43A85, 53D99, 81Q99, 81S10, 81S30 .

*Key words and phrases.* Dequantization, Quantization, Symmetric mechanical systems, Symbol correspondences, Quark systems.

The authors thank CAPES (finance code 001) for support.

5. Concluding remarks	64
6. Appendix	65
References	66

## 1. INTRODUCTION

Inspired by the treatment of symbol correspondences between quantum and classical mechanical systems which are symmetric under  $SU(2)$ , the so-called spin systems, cf. [18], we proceed to expand our knowledge about the correspondence principle for systems that are symmetric under a compact Lie group by studying symbol correspondences for mechanical systems with  $SU(3)$  symmetry<sup>1</sup>. Since  $SU(3)$  is the symmetry group of the strong force, we call such systems *quark systems*.

The first remarkable difference between the  $SU(2)$  and the  $SU(3)$  cases is that in the former case there is just one classical system, namely the Poisson algebra of smooth functions on  $\mathbb{C}P^1 \simeq S^2$ , whereas in the latter case there are two types of symplectic phase space: the complex projective plane  $\mathbb{C}P^2$  and the flag manifold that is a fiber bundle  $\mathcal{E}$  over  $\mathbb{C}P^2$  with fibers  $\mathbb{C}P^1$ , denoted  $\mathbb{C}P^1 \hookrightarrow \mathcal{E} \rightarrow \mathbb{C}P^2$ . The quantum systems of interest here are the Hilbert spaces  $\mathcal{H}_{p,q}$  with an irreducible representation  $Q(p, q)$  of  $SU(3)$ , for  $p, q \in \mathbb{N}_0$ , together with their operator algebras.

Then the main question posed here can be addressed in the following way: when/how is it possible to injectively map the space of operators on  $\mathcal{H}_{p,q}$  to the space of smooth functions on  $\mathbb{C}P^2$ , or  $\mathcal{E}$ , in an  $SU(3)$ -equivariant way which also ensures that quantum observables give rise to classical observables?

To answer the question above we define *symbol correspondences*<sup>2</sup> in the spirit of what is already done in literature, especially in [18], as linear injective maps

$$W : \mathcal{B}(\mathcal{H}_{p,q}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathcal{O}) , \quad A \mapsto W_A ,$$

where  $\mathcal{B}(\mathcal{H}_{p,q})$  is the space of operators and  $\mathcal{O}$  is either  $\mathbb{C}P^2$  or  $\mathcal{E}$ , satisfying a few extra properties: (i) any such map  $W$  is  $SU(3)$ -equivariant, (ii) the image of any Hermitian operator is a real function and (iii) the normalization condition

$$(1.1) \quad \int_{\mathcal{O}} W_A(x) dx = \frac{1}{\dim Q(p, q)} \operatorname{tr}(A)$$

applies to every operator  $A \in \mathcal{B}(\mathcal{H}_{p,q})$ , with respect to a normalized left invariant integral on  $\mathcal{O}$ . Condition (ii) encodes that  $W$  maps observables to observables and condition (iii) means it preserves expected values.

It turns out that for  $\mathcal{O} = \mathbb{C}P^2$  we can only define symbol correspondences for irreducible representations of type  $Q(p, 0)$  or  $Q(0, q)$ . We refer to the classical and quantum systems associated to  $\mathbb{C}P^2$  and Hilbert spaces  $\mathcal{H}_{p,0}$  or  $\mathcal{H}_{0,q}$  as *pure-quark systems*, since the pertinent irreducible representations of  $SU(3)$  emerge from systems of  $p$  quarks only, or  $q$  antiquarks only.<sup>3</sup> Characterization of symbol correspondences for pure-quark systems is very similar to what is known for spin systems.

<sup>1</sup>Initial efforts in this direction can be found in [15, 16, 19].

<sup>2</sup>It may be fair to call them *symplectic symbol correspondences* because we are working only with symplectic manifolds. For  $SU(3)$ , one could also try to work with a Poisson manifold irregularly foliated by symplectic leaves, trying to define *Poissonian symbol correspondences*.

<sup>3</sup>One can also interpret such pure-quark systems as solution spaces for the three dimensional isotropic oscillator, cf. Remark 3.9.

In particular, the correspondences for  $\mathcal{H}_{p,0}$ , or  $\mathcal{H}_{0,p}$ , are unequivocally determined by an ordered set of nonzero real numbers

$$(1.2) \quad c_n \in \mathbb{R}^* , \quad 1 \leq n \leq p ,$$

called *characteristic numbers*, so that the moduli space of symbol correspondences for pure-quark systems is  $(\mathbb{R}^*)^p$ , cf. Theorem 3.13 and Corollary 3.14.

However, when  $\mathcal{O}$  is the flag manifold  $\mathcal{E}$ , we can define symbol correspondences for any  $\mathcal{H}_{p,q}$ , i.e. any irreducible representation  $\rho$  of class  $Q(p,q)$  of  $SU(3)$ , so we refer to the classical and quantum systems associated to  $\mathcal{E}$  and  $\mathcal{H}_{p,q}$  as *generic quark systems*<sup>4</sup>. Here we see some novel features and the characterization of symbol correspondences for generic quark systems is now presented via full-rank complex matrices, called *characteristic matrices*, cf. Theorem 4.11, so that the moduli space of symbol correspondences for generic quark systems is a product of non compact Stiefel manifolds, cf. Corollary 4.12.

Just as it happens for spin systems, for both pure-quark and generic quark systems, Theorems 3.15 and 4.13 show that a symbol correspondence  $W$  can be realized in terms of expectations over a Hermitian operator with unitary trace  $K$ , called the *operator kernel*, via

$$(1.3) \quad W_A(gx_0) = \text{tr}(AK^g) ,$$

for any  $A \in \mathcal{B}(\mathcal{H}_{p,q})$  and  $g \in SU(3)$ , where  $x_0 \in \mathcal{O}$  is a suitable point related to a choice of basis for  $\mathcal{H}_{p,q}$ , with  $gx_0$  denoting the (co)adjoint action of  $g$  on  $x_0 \in \mathcal{O}$  and  $K^g$  denoting the action of  $g$  on  $K \in \mathcal{B}(\mathcal{H}_{p,q})$ ,

$$SU(3) \ni g : K \mapsto \rho(g)K\rho(g)^{-1} = K^g ,$$

where  $\rho$  is the irreducible  $SU(3)$ -representation on the respective quantum system.

Thus, one can also interpret symbol correspondences for quark systems as expectation values over pseudo-states (Hermitian operators with unit trace which are not necessarily positive). When the operator kernel is also a positive operator, i.e. when  $K$  is precisely a state, the correspondence maps positive(-definite) operators to (strictly-)positive functions and is called a *mapping-positive correspondence*.

On the other hand, if the correspondence  $W$  induces an isometry between  $\mathcal{B}(\mathcal{H}_{p,q})$  and the image set of  $W$ , for appropriately normalized invariant inner products in  $\mathcal{B}(\mathcal{H}_{p,q})$  and  $C_c^\infty(\mathcal{O})$ , then  $W$  is called a *Stratonovich-Weyl correspondence*. As for spin systems, Theorem 3.26 shows that no Stratonovich-Weyl correspondence is mapping-positive, for the case of pure-quark systems.

Adaptations of a proof from [18] show that the projection on the lowest weight vector of  $\mathcal{H}_{p,0}$  is an operator kernel as well as the projection on highest weight vector of  $\mathcal{H}_{0,p}$ , so the mapping-positive symbol correspondences they define are called *Berezin correspondences*, cf. Propositions 3.28 and 3.29 and Definitions 3.31 and 4.26. Berezin correspondences also present examples of a natural relation between correspondences for dual quark systems (a concealed feature for spin systems since these are self dual systems). Accordingly, in [18] the authors defined *alternate correspondences*, whilst here we shall use the term *antipodal correspondences*, cf. Definitions 3.34 and 4.28, Propositions 3.36 and 4.29, Remarks 3.33 and 3.35.

---

<sup>4</sup>We emphasize that quantum pure-quark systems are special cases of quantum generic quark systems.

In this paper, we also start the study of *twisted products of symbols*, that is, non-commutative products of functions in some finite dimensional subspaces of  $C_c^\infty(\mathcal{O})$ , which are induced by the operator product on  $\mathcal{B}(\mathcal{H}_{p,q})$  via symbol correspondences.

In order to do so, first we develop the  $SU(3)$ -invariant decomposition of the operator product on  $\mathcal{B}(\mathcal{H}_{p,q})$ , cf. Lemma 2.14, Proposition 2.27 and Corollary 2.29, defining the Wigner product symbol that is a product of Wigner coupling and recoupling symbols, cf. Definitions 2.25, 2.26 and 2.28. From this, in Theorems 3.42, 3.43, 4.36 and 4.37 we are able to present some explicit expressions for twisted products. And in Propositions 3.48 and 4.42 we show that antipodal correspondences induce a “reverse symbolic dynamics” via twisted product, cf. Remark 3.50.

This paper is organized as follows.

In Section 2 we first present some general facts concerning the Lie group  $SU(3)$  and proceed in subsection 2.1 with the characterization of its irreducible unitary representations using the Gelfand-Tsetlin technology for defining a basis of  $\mathcal{H}_{p,q}$ . Then, in subsection 2.2 we present the Clebsch-Gordan series of  $SU(3)$ , followed by the  $SU(3)$ -invariant decomposition of the operator product, in subsection 2.3. Using mixed Casimir operators, we present a way to distinguish subrepresentations with multiplicities in the  $CG$  series, in subsection 2.4, from which we define various symmetric Wigner symbols which are used to rewrite the operator product, in subsection 2.5. Finally, in subsection 2.6 we present the description of  $\mathbb{C}P^2$  and  $\mathcal{E}$  as (co)adjoint orbits of  $SU(3)$ .

In Sections 3 and 4 we work out the description of symbol correspondences for pure-quark systems and generic quark systems, respectively. The first section of each chapter presents the construction of harmonic functions on the respective classical phase space, from which we identify the pertinent irreducible representations of  $SU(3)$ , and proceeds with the  $SU(3)$ -invariant decomposition of their pointwise product. The last two sections of each chapter are devoted to symbol correspondences and their twisted products.

Section 5 presents a brief discussion of some results obtained in this paper, with indication of some topics for future investigations.

*Acknowledgements:* We thank Eldar Straume and Luiz San Martin for some interesting comments.

2. ON THE REPRESENTATIONS OF  $SU(3)$ 

Let  $SU(3)$  denote the special unitary subgroup of  $GL_3(\mathbb{C})$ , satisfying  $\det g = 1$  and  $gg^\dagger = g^\dagger g = e$ , for all  $g \in SU(3)$ . As a manifold,  $SU(3)$  can be seen as a fiber bundle over  $S^5$  whose fibers are  $S^3 \simeq SU(2)$  (see discussion in section 2.6), hence it is a compact Lie group of real dimension 8. The Lie algebra of  $SU(3)$ , denoted by  $\mathfrak{su}(3)$ , can be generated by  $i\lambda_k$ , for  $k = 1, \dots, 8$ , where

$$(2.1) \quad \begin{aligned} \lambda_1 &= \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda_2 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\ \lambda_4 &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad \lambda_5 = \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}, \quad \lambda_6 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \\ \lambda_7 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad \lambda_8 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix} \end{aligned}$$

are Hermitian matrices, known as *Gell-Mann matrices*, satisfying

$$(2.2) \quad \text{tr}(\lambda_a \lambda_b) = 2\delta_{ab},$$

and

$$(2.3) \quad [\lambda_a, \lambda_b] = 2i \sum_{c=1}^8 f^{abc} \lambda_c$$

with  $f^{abc}$  totally antisymmetric and completely determined by Table 1 – see e.g. [12]. It follows that  $SU(3)$  is not only compact, but also a simple Lie group.

$abc$	123	147	156	246	257	345	367	458	678
$f^{abc}$	1	1/2	-1/2	1/2	1/2	1/2	-1/2	$\sqrt{3}/2$	$\sqrt{3}/2$

TABLE 1. Structure constants for Gell-Mann matrices.

In order to describe irreducible representations of  $SU(3)$ , it is useful to take the complexification of  $\mathfrak{su}(3)$ . Thus, we will work with complex linear combinations of the so-called *F-spin* operators:

$$(2.4) \quad F_k = \frac{1}{2} \lambda_k.$$

As expected, due to their names, using the *F-spin* operators we get a Lie bracket structure for  $SU(3)$  more similar to the canonical one for  $SU(2)$ . By defining

$$(2.5) \quad \begin{aligned} T_\pm &= F_1 \pm iF_2, \quad T_3 = F_3, \\ V_\pm &= F_4 \pm iF_5, \quad U_\pm = F_6 \pm iF_7, \\ Y &= \frac{2}{\sqrt{3}} F_8, \end{aligned}$$

one can easily verify that

$$(2.6) \quad T_\pm^\dagger = T_\mp, \quad U_\pm^\dagger = U_\mp, \quad V_\pm^\dagger = V_\mp, \quad T_3^\dagger = T_3, \quad Y^\dagger = Y$$

and

$$(2.7) \quad [Y, T_3] = 0 .$$

For  $Y$ , we also have

$$(2.8) \quad [Y, T_\pm] = 0 \quad , \quad [Y, U_\pm] = \pm U_\pm \quad , \quad [Y, V_\pm] = \pm V_\pm .$$

For the other operators, the following holds:

$$(2.9) \quad [T_3, T_\pm] = \pm T_\pm \quad , \quad [T_3, U_\pm] = \mp \frac{1}{2} U_\pm \quad , \quad [T_3, V_\pm] = \pm \frac{1}{2} V_\pm \quad , \quad [T_+, T_-] = 2T_3 ,$$

$$(2.10) \quad [U_+, U_-] = \frac{3}{2} Y - T_3 \quad , \quad [V_+, V_-] = \frac{3}{2} Y + T_3 .$$

$$(2.11) \quad [T_+, U_+] = V_+ \quad , \quad [T_+, U_-] = 0 \quad , \quad [T_+, V_+] = 0 \quad , \quad [T_+, V_-] = -U_-$$

$$(2.12) \quad [U_+, V_+] = 0 \quad , \quad [U_+, V_-] = T_- .$$

From (2.7) and (2.10), we are compelled to define

$$(2.13) \quad U_3 = \frac{3}{4} Y - \frac{1}{2} T_3 \quad , \quad V_3 = \frac{3}{4} Y + \frac{1}{2} T_3 ,$$

so we get<sup>5</sup>

$$(2.14) \quad [U_3, U_\pm] = \pm U_\pm \quad , \quad [U_3, V_\pm] = \pm \frac{1}{2} V_\pm \\ [V_3, V_\pm] = \pm V_\pm \quad , \quad [V_3, U_\pm] = \pm \frac{1}{2} U_\pm ,$$

$$(2.15) \quad [T_3, U_3] = [U_3, V_3] = [V_3, T_3] = 0 .$$

The remaining commutation relations can be computed using the Hermitian conjugate of the above expressions. We thus conclude that  $SU(3)$  is of rank 2 and a straightforward calculation gives us

$$(2.16) \quad \text{tr}(T_3 T_3) = \frac{1}{2} \quad , \quad \text{tr}(Y Y) = \frac{2}{3} \quad , \quad \text{tr}(T_3 Y) = 0 ,$$

so the set  $\{iT_3, iY\}$  forms an orthogonal, but not normal basis of the Cartan subalgebra of  $\mathfrak{su}(3)$ .


Equations (2.14)-(2.15) together with equations (2.8)-(2.9), show that the root system of  $SU(3)$  is composed by three root systems of  $SU(2)$ , with the same length, framing a regular hexagon as in Figure 1.

The roots  $\alpha_1$ ,  $\alpha_2$  and  $\alpha_3$  are associated to the ladder operators  $T_+$ ,  $U_+$  and  $V_+$ , respectively. We choose the fundamental Weyl chamber as the blue hatched one, enclosed by the dashed lines, so that  $\{\alpha_1, \alpha_2, \alpha_3\}$  is the set of positive roots and  $\{\alpha_1, \alpha_2\}$  is the set of simple roots. Let  $\omega_1$  and  $\omega_2$  be the fundamental weights satisfying

$$(2.17) \quad 2 \frac{\langle \omega_j | \alpha_k \rangle}{\|\alpha_k\|^2} = \delta_{j,k} \quad , \quad j, k \in \{1, 2\} ,$$

---

<sup>5</sup>Among the infinitely many  $SU(2)$  subgroups of  $SU(3)$ , the ones associated to  $\{T_3, T_\pm\}$ ,  $\{U_3, U_\pm\}$  and  $\{V_3, V_\pm\}$  are singled out as the three *standard*  $SU(2)$  subgroups of  $SU(3)$ .

FIGURE 1. Root diagram of  $\mathfrak{su}(3)$ .

where  $\langle | \rangle$  is the canonical Euclidean inner product on the root space. Writing the fundamental weights as linear combination of the simple roots  $\{\alpha_1, \alpha_2\}$ , and using that

$$(2.18) \quad \langle \alpha_1 | \alpha_2 \rangle = -\|\alpha_1\| \|\alpha_2\| / 2 \quad , \quad \|\alpha_1\| = \|\alpha_2\| \quad ,$$

the orthogonality relations  $\langle \omega_1 | \alpha_2 \rangle = \langle \omega_2 | \alpha_1 \rangle = 0$  imply

$$\omega_1 = a(2\alpha_1 + \alpha_2) \quad , \quad \omega_2 = b(\alpha_1 + 2\alpha_2) \quad .$$

The remaining relations given by (2.17) imply  $a = b = 1/3$ , so we finally get

$$(2.19) \quad w_1 = \frac{1}{3}(2\alpha_1 + \alpha_2) \quad , \quad w_2 = \frac{1}{3}(\alpha_1 + 2\alpha_2) \quad .$$

**2.1. Irreducible unitary representations.** Thus, we label the classes of irreducible unitary representations of  $SU(3)$  by two nonnegative integers  $p$  and  $q$ , denoting each class by  $Q(p, q)$ , where  $(p, q) \equiv p\omega_1 + q\omega_2$  is the highest weight of the representation. In addition, we shall often refer to an unitary irreducible representation class  $Q(p, q)$  in a less specific way simply as an unitary irreducible representation  $Q(p, q)$ , or just as a representation  $Q(p, q)$ . Accordingly, for a representation  $Q(p, q)$ ,  $p$  and  $q$  are the maximal integers such that  $(T_-)^p$  and  $(U_-)^q$  can be applied to the highest weight vector before vanishing, and an orthonormal basis of weight vectors for the representation  $Q(p, q)$ , where<sup>6</sup>

$$(2.20) \quad \dim Q(p, q) = \frac{(p+1)(q+1)(p+q+2)}{2} \quad ,$$

is obtained via linear combinations of the action of  $(T_-)^a (U_-)^b (T_-)^c$  on the highest weight vector.

**Remark 2.1.** *The weights of  $Q(p, q)$  can be placed on a plane diagram that is very useful for seeing the action of the step operators. In the examples below, the highest weights are highlighted as square dots and the multiplicities of a weight are represented by rings around the weight. Also, the axes  $t$  and  $u$  express the eigenvalues for the operators  $T_3$  and  $U_3$ , respectively.*

Now, let  $\rho$  be an unitary irreducible representation of class  $Q(p, q)$  acting on a complex Hilbert space  $(\mathcal{H}, \langle | \rangle)$  of dimension  $\dim Q(p, q)$ . Then, we use the fact that

<sup>6</sup>Cf. Weyl dimensionality formula [13].




FIGURE 2. Examples of weight diagram for irreducible representations of  $SU(3)$ .

the weights of  $Q(p, q)$  can be labeled by  $(\nu_1, \nu_2, \nu_3)$  and  $I$  satisfying the Gelfand-Tsetlin pattern<sup>7</sup>

$$(2.21) \quad \begin{aligned} 0 \leq r_- \leq q \leq r_+ \leq p + q, \quad r_- \leq \nu_1 \leq r_+, \\ \nu_2 = (r_+ + r_-) - \nu_1, \quad \nu_3 = p + 2q - (r_+ + r_-), \\ I = \frac{1}{2}(r_+ - r_-), \end{aligned}$$

where  $r_+$  and  $r_-$  are integers. The quantities

$$(2.22) \quad t = (\nu_1 - \nu_2)/2, \quad u = (\nu_2 - \nu_3)/2, \quad v = (\nu_1 - \nu_3)/2$$

are the eigenvalues<sup>8</sup> for the operators  $T_3$ ,  $U_3$  and  $V_3$ , respectively, and the index  $I$ , which is the spin number of the subrepresentation of  $SU(2)$  related to  $\{T_3, T_\pm\}$ , is usually referred to as the *isospin*, in the physics literature. In this way,  $T_\pm$  changes  $(\nu_1, \nu_2, \nu_3) \mapsto (\nu_1 \pm 1, \nu_2 \mp 1, \nu_3)$ ,  $U_\pm$  changes  $(\nu_1, \nu_2, \nu_3) \mapsto (\nu_1, \nu_2 \pm 1, \nu_3 \mp 1)$  and  $V_\pm$  changes  $(\nu_1, \nu_2, \nu_3) \mapsto (\nu_1 \pm 1, \nu_2, \nu_3 \mp 1)$ .

In particular, the highest weight is given by

$$(2.23) \quad \nu_1 = p + q$$

and

$$(2.24) \quad r_+ = p + q, \quad r_- = q,$$

so that

$$(2.25) \quad \nu_2 = q, \quad \nu_3 = 0, \quad I = p/2.$$

<sup>7</sup>The same method is used in [3]. For a general description of Gelfand-Tsetlin pattern, see [17, 27], for instance.

<sup>8</sup>Henceforth the weights from which we are taking the eigenvalues will be specified or will be clear from the presence or absence of subscript and superscript in  $t$ ,  $u$  and  $v$ .

**Definition 2.2** (cf. e.g. [3]). A Gelfand-Tsetlin basis  $\{e((p, q); \nu, I)\}$ , or simply a GT basis, of a  $SU(3)$ -representation of class  $Q(p, q)$  is an orthonormal basis indexed by the isospin  $I$  (the spin number of  $\{T_3, T_\pm\}$ ) and

$$(2.26) \quad \nu = (\nu_1, \nu_2, \nu_3) ,$$

as specified above, cf. (2.21)-(2.22), constructed by fixing a highest weight vector

$$e((p, q); (p + q, q, 0), p/2)$$

and choosing the other basis vectors such that

$$(2.27) \quad T_- e((p, q); (\nu_1, \nu_2, \nu_3), I) = \sqrt{(I+t)(I-t+1)} e((p, q); (\nu_1 - 1, \nu_2 + 1, \nu_3), I) ,$$

$$(2.28)$$

$$U_- e((p, q); (\nu_1, \nu_2, \nu_3), I) = \sqrt{\frac{(I-t)(p+q-I+t-\nu_1+1)(I-t+\nu_1-q)(I-t+\nu_1+1)}{2I(2I+1)}} \times e((p, q); (\nu_1, \nu_2 - 1, \nu_3 + 1), I - 1/2) + \sqrt{\frac{(I+t+1)(p+q-\nu_1+I+t+2)(q-\nu_1+I+t+1)(\nu_1-I-t)}{(2I+2)(2I+1)}} \times e((p, q); (\nu_1, \nu_2 - 1, \nu_3 + 1), I + 1/2) .$$

Considering the anti-isomorphism  $\mathcal{H}^* \leftrightarrow \mathcal{H}$  via inner product, for the dual representation  $\check{\rho} \leftrightarrow \rho$ , we get that  $\check{T}_3 \leftrightarrow -T_3$ ,  $\check{U}_3 \leftrightarrow -U_3$ ,  $\check{V}_3 \leftrightarrow -V_3$ ,  $\check{T}_\pm \leftrightarrow -T_\mp$ ,  $\check{U}_\pm \leftrightarrow -U_\mp$  and  $\check{V}_\pm \leftrightarrow -V_\mp$ . Thus, the states of  $\check{\rho}$  are related to the states of  $\rho$  by

$$(2.29) \quad \check{I} = I ; \quad \check{t} = -t , \quad \check{u} = -u ,$$

which implies

$$(2.30) \quad \check{\nu}_1 = p + q - \nu_1 \quad , \quad \check{\nu}_2 = p + q - \nu_2 \quad , \quad \check{\nu}_3 = p + q - \nu_3 .$$

Therefore, the highest weight of the representation  $\check{\rho}$  on  $\mathcal{H}^*$  is  $-\mu$ , where  $\mu$  is the lowest weight of  $\rho$ . Since the lowest weight is the image of the highest weight by the longest element of the Weyl group, we find that  $(q, p) = q\omega_1 + p\omega_2$  is the highest weight of  $\check{\rho}$ , that is,  $\check{\rho}$  is of class  $Q(q, p)$ , so that

$$(2.31) \quad Q(p, q)^* = Q(q, p) .$$

**Notation 1.** In the light of this dualization symmetry, we introduce the simplifying notation<sup>9</sup>:

$$(2.32) \quad \mathbf{p} = (p, q) \in \mathbb{N}_0 \times \mathbb{N}_0 \leftrightarrow \check{\mathbf{p}} = (q, p) , \quad \text{with } |\mathbf{p}| = |\check{\mathbf{p}}| = p + q .$$

Then, to better state the relations in (2.30), we define

$$(2.33)$$

$$\Delta_{\nu, \mu}^{|\mathbf{p}|} \equiv \Delta_{\nu, \mu}^{p+q} := \begin{cases} 1 & , \quad \text{if } \nu + \mu = (p + q, p + q, p + q) = (|\mathbf{p}|, |\mathbf{p}|, |\mathbf{p}|) \\ 0 & , \quad \text{otherwise} \end{cases} .$$

<sup>9</sup>We use the convention which identifies the set of natural numbers as  $\mathbb{N} = \{1, 2, 3, \dots\}$  and denote  $\mathbb{N}_0 = \{0, 1, 2, 3, \dots\}$ .

**Definition 2.3.** Given a Gelfand-Tsetlin basis  $\{e(\mathbf{p}; \boldsymbol{\nu}, I)\}$  of a  $SU(3)$ -representation of class  $Q(\mathbf{p}) \equiv Q(p, q)$ , the induced Gelfand-Tsetlin basis of the dual representation of class  $Q(p, q)^* = Q(q, p) \equiv Q(\check{\mathbf{p}})$  is the basis comprised by the vectors

$$(2.34) \quad \check{e}(\check{\mathbf{p}}; \check{\boldsymbol{\nu}}, I) = (-1)^{2(t_\nu + u_\nu)} e^*(\mathbf{p}; \boldsymbol{\nu}, I) ,$$

where  $e^*(\mathbf{p}; \boldsymbol{\nu}, I) \in Q(\check{\mathbf{p}})$  is the Hermitian dual of  $e(\mathbf{p}; \boldsymbol{\nu}, I)$ , that is, the dual of  $e(\mathbf{p}; \boldsymbol{\nu}, I)$  via Hermitian inner product on  $Q(\mathbf{p})$ , and where  $t_\nu$  and  $u_\nu$  stand as in (2.22), with  $\boldsymbol{\nu}$  and  $\check{\boldsymbol{\nu}}$  satisfying the duality relations (2.30), that is, using (2.33),

$$(2.35) \quad \text{duality: } \boldsymbol{\nu} \leftrightarrow \check{\boldsymbol{\nu}} \iff \Delta_{\boldsymbol{\nu}, \check{\boldsymbol{\nu}}}^{|\mathbf{p}|} = 1 .$$

**Remark 2.4.** A GT basis  $\{e(\mathbf{p}; \boldsymbol{\nu}, I)\}$  and the induced GT basis  $\{\check{e}(\check{\mathbf{p}}; \check{\boldsymbol{\nu}}, I)\}$  are related by an involution. Considering the natural isomorphism between a finite dimensional vector space and its double dual, the dual GT basis induced by  $\{\check{e}(\check{\mathbf{p}}; \check{\boldsymbol{\nu}}, I)\}$  is precisely  $\{e(\mathbf{p}; \boldsymbol{\nu}, I)\}$ , that is,

$$(2.36) \quad e(\mathbf{p}; \boldsymbol{\nu}, I) = (-1)^{2(t+u)} \check{e}^*(\check{\mathbf{p}}; \check{\boldsymbol{\nu}}, I) .$$

This contrasts with standard convention for irreducible representations of  $SU(2)$ , since for a  $SU(2)$ -representation with spin number  $j$ , there is a phase  $(-1)^{2j}$  between a standard basis and the basis of the double dual space induced by the basis of the dual space, c.f. [18].

**Definition 2.5.** The Wigner  $D$ -functions (in the GT basis) of an irreducible unitary  $SU(3)$ -representation  $\rho$  of class  $Q(\mathbf{p}) \equiv Q(p, q)$  are the functions

$$(2.37) \quad D_{\boldsymbol{\nu}I, \boldsymbol{\mu}J}^{\mathbf{p}}(g) = \langle e(\mathbf{p}; \boldsymbol{\nu}, I) | \rho(g) e(\mathbf{p}; \boldsymbol{\mu}, J) \rangle .$$

Using the conjugate symmetry of the inner product and the relation  $\langle v|w \rangle = \langle w^*|v^* \rangle$  between inner products of  $\mathcal{H}$  and  $\mathcal{H}^*$ , we get

$$(2.38) \quad \begin{aligned} \overline{D_{\boldsymbol{\nu}I, \boldsymbol{\mu}J}^{\mathbf{p}}(g)} &= \langle \rho(g) e(\mathbf{p}; \boldsymbol{\mu}, J) | e(\mathbf{p}; \boldsymbol{\nu}, I) \rangle \\ &= \langle e^*(\mathbf{p}; \boldsymbol{\nu}, I) | \check{\rho}(g) e^*(\mathbf{p}; \boldsymbol{\mu}, J) \rangle \\ &= (-1)^{2(t_\nu + u_\nu + t_\mu + u_\mu)} \langle \check{e}(\check{\mathbf{p}}; \check{\boldsymbol{\nu}}, I) | \check{\rho}(g) \check{e}(\check{\mathbf{p}}; \check{\boldsymbol{\mu}}, J) \rangle \\ &= (-1)^{2(t_\nu + u_\nu + t_\mu + u_\mu)} D_{\check{\boldsymbol{\nu}}I, \check{\boldsymbol{\mu}}J}^{\check{\mathbf{p}}}(g) \end{aligned}$$

for  $\Delta_{\boldsymbol{\nu}, \check{\boldsymbol{\nu}}}^{|\mathbf{p}|} = \Delta_{\boldsymbol{\mu}, \check{\boldsymbol{\mu}}}^{|\mathbf{p}|} = 1$ .

**2.2. Clebsch-Gordan series and space of operators.** An irreducible unitary representation  $\rho$  of class  $Q(\mathbf{p})$  on  $\mathcal{H}$  extends to an unitary representation (with respect to the trace inner product) on  $\mathcal{B}(\mathcal{H})$  via adjoint action: for  $A \in \mathcal{B}(\mathcal{H})$  and  $g \in SU(3)$ , the action of  $g$  on  $A$  is

$$(2.39) \quad A^g = \rho(g) A \rho(g)^{-1} .$$

Now, we recall that  $\mathcal{B}(\mathcal{H})$  is naturally isomorphic to  $\mathcal{H} \otimes \mathcal{H}^*$  in a manner that (2.39) matches the representation  $\rho \otimes \check{\rho}$  of class  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}})$  on  $\mathcal{H} \otimes \mathcal{H}^*$ .

The decomposition of a tensor product of irreducible unitary  $SU(3)$ -representations into a direct sum of irreducible unitary  $SU(3)$ -representations is known as the *Clebsch-Gordan series* of  $SU(3)$ .

**Theorem 2.6** (see e.g. [9]). *The Clebsch-Gordan series of  $SU(3)$  are given by*

$$(2.40) \quad Q(p_1, q_1) \otimes Q(p_2, q_2) = \bigoplus_{n=0}^{\min(p_1, q_2)} \bigoplus_{m=0}^{\min(p_2, q_1)} Q(p_1 - n, p_2 - m; q_1 - m, q_2 - n),$$

where

$$(2.41) \quad \begin{aligned} Q(r_1, r_2; s_1, s_2) = & Q(r_1 + r_2, s_1 + s_2) \oplus \left( \bigoplus_{k=1}^{\min(r_1, r_2)} Q(r_1 + r_2 - 2k, s_1 + s_2 + k) \right) \\ & \oplus \left( \bigoplus_{k=1}^{\min(s_1, s_2)} Q(r_1 + r_2 + k, s_1 + s_2 - 2k) \right). \end{aligned}$$

**Corollary 2.7.** *For  $p_1 = q_2 = p$  and  $q_1 = p_2 = q$ , the Clebsch-Gordan series assumes the form*

$$(2.42) \quad \begin{aligned} Q(p, q) \otimes Q(q, p) = & \bigoplus_{n=0}^p \bigoplus_{m=0}^q \left\{ Q(p + q - n - m, p + q - n - m) \right. \\ & \oplus \left[ \bigoplus_{k=1}^{\min(p-n, q-m)} \left( Q(p + q - n - m - 2k, p + q - n - m + k) \right. \right. \\ & \left. \left. \oplus Q(p + q - n - m + k, p + q - n - m - 2k) \right) \right] \left. \right\}. \end{aligned}$$

Note that an irreducible unitary representation of class  $Q(\mathbf{a})$  may appear more than once in the Clebsch-Gordan series of  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2)$  and of  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}})$ , in general.

**Notation 2.** *We shall denote the multiplicity of  $Q(\mathbf{a}) = Q(a, b)$  in the Clebsch-Gordan series of  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2)$  by*

$$(2.43) \quad \mathbf{m}(\mathbf{p}_1, \mathbf{p}_2; \mathbf{a}).$$

*To distinguish multiple appearances of the same class of representation in a Clebsch-Gordan series, we will write*

$$(2.44) \quad (\mathbf{a}; \sigma) = (a, b; \sigma), \quad Q(\mathbf{a}; \sigma) \equiv Q(a, b; \sigma),$$

where the index  $\sigma$  counts the multiplicity starting from 1 to  $\mathbf{m}(\mathbf{p}_1, \mathbf{p}_2; \mathbf{a})$ .

We thus provide two basis for  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2)$ .

**Definition 2.8.** *An uncoupled GT basis of the tensor product representation  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2)$  is a basis comprised by the tensor product of GT basis  $\{\mathbf{e}(\mathbf{p}_1; \boldsymbol{\nu}_1, I_1)\}$  of  $Q(\mathbf{p}_1)$  and  $\{\mathbf{e}(\mathbf{p}_2; \boldsymbol{\nu}_2, I_2)\}$  of  $Q(\mathbf{p}_2)$ .*

**Definition 2.9.** *A coupled GT basis of the tensor product representation  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2)$  is the union of GT basis  $\{\mathbf{e}(\mathbf{a}; \sigma; \boldsymbol{\nu}, I)\}$  of each  $Q(\mathbf{a}; \sigma) \equiv Q(a, b; \sigma)$  in the Clebsch-Gordan series of  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2)$ .*

**Remark 2.10.** *For the sake of good reading, we will not write the labels of the representations of the tensor product on the elements of a coupled basis, but it will be clear from the context. Also, unless specified otherwise, from now on we shall*

always refer to the uncoupled and coupled basis of the tensor product as meaning their respective GT basis, and likewise for the Clebsch-Gordan coefficients.

Both these basis are orthonormal, so they are related by an unitary transformation.

**Definition 2.11.** *The Clebsch-Gordan coefficients (in the GT basis) are the entries of the transformation that relate a coupled and an uncoupled GT basis of  $Q(p_1, q_1) \otimes Q(p_2, q_2)$ :*

$$(2.45) \quad C_{\nu_1 I_1, \nu_2 I_2, \nu I}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a}; \sigma)} = \langle e((\mathbf{a}; \sigma); \nu, I) | e(\mathbf{p}_1; \nu_1, I_1) \otimes e(\mathbf{p}_2; \nu_2, I_2) \rangle ,$$

where  $\langle \cdot | \cdot \rangle$  is the  $SU(3)$ -invariant inner product induced by the ones on each representation of the tensor product.

We are able to fix a relative phase between a coupled and an uncoupled basis so that all Clebsch-Gordan are real. Usually, one chooses some set of Clebsch-Gordan coefficients to be positive and the remaining coefficients are completely determined via the action of the step operators on the basis vectors – cf. e.g. [24]. In the next section, we shall return to this problem. What is important now is that we take Clebsch-Gordan coefficients as real numbers, so that we have

$$(2.46) \quad e(\mathbf{p}_1; \nu_1, I_1) \otimes e(\mathbf{p}_2; \nu_2, I_2) = \sum_{\substack{(\mathbf{a}; \sigma) \\ \nu, I}} C_{\nu_1 I_1, \nu_2 I_2, \nu I}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a}; \sigma)} e((\mathbf{a}; \sigma); \nu, I) ,$$

$$(2.47) \quad e((\mathbf{a}; \sigma); \nu, I) = \sum_{\substack{\nu_1, I_1 \\ \nu_2, I_2}} C_{\nu_1 I_1, \nu_2 I_2, \nu I}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a}; \sigma)} e(\mathbf{p}_1; \nu_1, I_1) \otimes e(\mathbf{p}_2; \nu_2, I_2) ,$$

$$(2.48) \quad \sum_{\substack{(\mathbf{a}; \sigma) \\ \nu, I}} C_{\nu_1 I_1, \nu_2 I_2, \nu I}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a}; \sigma)} C_{\nu'_1 I'_1, \nu'_2 I'_2, \nu I}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a}; \sigma)} = \delta_{\nu_1, \nu'_1} \delta_{\nu_2, \nu'_2} \delta_{I_1, I'_1} \delta_{I_2, I'_2} ,$$

$$\sum_{\substack{\nu_1, I_1 \\ \nu_2, I_2}} C_{\nu_1 I_1, \nu_2 I_2, \nu I}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a}; \sigma_1)} C_{\nu_1 I_1, \nu_2 I_2, \nu' I'}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{b}; \sigma_2)} = \delta_{\mathbf{a}, \mathbf{b}} \delta_{\sigma_1, \sigma_2} \delta_{\nu, \nu'} \delta_{I, I'} .$$

**Remark 2.12.** *We also point out that, from the way the GT basis for  $SU(3)$  representations were constructed using  $SU(2)$  subrepresentations, the  $SU(3)$  Clebsch-Gordan coefficients in the GT basis are related to the  $SU(2)$  Clebsch-Gordan coefficients by*

$$(2.49) \quad C_{\nu_1 I_1, \nu_2 I_2, \nu I}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a}; \sigma)} = C_{t_{\nu_1}, t_{\nu_2}, t_{\nu}}^{I_1, I_2, I} C_{u_{\nu_1} I_1, u_{\nu_2} I_2, u_{\nu} I}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a}; \sigma)} ,$$

where the second coefficient on the r.h.s. is called isoscalar factor, and this provides explicit equations for the  $SU(3)$  Clebsch-Gordan coefficients in the GT basis in terms of explicit equations for the  $SU(2)$  Clebsch-Gordan coefficients, as found in [18] and [25], for instance.

From decompositions (2.46)-(2.47), we obtain some sufficient conditions for the Clebsch-Gordan coefficients to be zero. Since  $e(\mathbf{p}_1; \nu_1, I_1)$  and  $e(\mathbf{p}_2; \nu_2, I_2)$  are basis vectors of  $SU(2)$ -representations with spin numbers  $I_1$  and  $I_2$ , their tensor product is a vector of the tensor product of the  $SU(2)$ -representations they belong to, that is, the Clebsch-Gordan coefficients are zero if  $I_1, I_2$  and  $I$  do not satisfy the

triangle inequality. Also, using superscripts to identify the  $SU(3)$ -representations, the operators  $T_3$  and  $U_3$  in  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2)$  have the form

$$(2.50) \quad \begin{aligned} \bigoplus_{(\mathbf{a};\sigma)} T_3^{(\mathbf{a};\sigma)} &= T_3^{\mathbf{p}_1} \otimes \mathbf{1} + \mathbf{1} \otimes T_3^{\mathbf{p}_2} , \\ \bigoplus_{(\mathbf{a};\sigma)} U_3^{(\mathbf{a};\sigma)} &= U_3^{\mathbf{p}_1} \otimes \mathbf{1} + \mathbf{1} \otimes U_3^{\mathbf{p}_2} , \end{aligned}$$

where  $\mathbf{1}$  is the identity operator. Thus, the Clebsch-Gordan coefficients are zero if  $t \neq t_1 + t_2$  or  $u \neq u_1 + u_2$ , for  $t, t_1, t_2, u, u_1$  and  $u_2$  being the eigenvalues of  $T_3$  and  $U_3$  related to the weights  $\boldsymbol{\nu}, \boldsymbol{\nu}_1$  and  $\boldsymbol{\nu}_2$ .

To summarize, let  $\delta(x, y, z)$  be equal to 1 if  $x, y$  and  $z$  satisfy the triangle inequality, or 0 otherwise, and let

$$(2.51) \quad \nabla_{\boldsymbol{\nu}, \boldsymbol{\mu}} := \delta_{t_{\boldsymbol{\nu}}, t_{\boldsymbol{\mu}}} \delta_{u_{\boldsymbol{\nu}}, u_{\boldsymbol{\mu}}} ,$$

where  $\delta_{m,n}$  is the Kronecker delta. Then,

$$(2.52) \quad C_{\boldsymbol{\nu}_1 I_1, \boldsymbol{\nu}_2 I_2, \boldsymbol{\nu} I}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a};\sigma)} \neq 0 \quad \Longrightarrow \quad \begin{cases} \nabla_{\boldsymbol{\nu}_1 + \boldsymbol{\nu}_2, \boldsymbol{\nu}} = 1 \\ \delta(I_1, I_2, I) = 1 \end{cases} .$$

**2.3. Decomposition of the operator product.** That established, we proceed to establish the decomposition of the pointwise product of Wigner  $D$ -functions and the decomposition of the operator product.

**Lemma 2.13.** *The pointwise product of Wigner  $D$ -functions of  $SU(3)$  can be decomposed into a sum of the form*

$$(2.53) \quad D_{\boldsymbol{\nu}_1 I_1, \boldsymbol{\nu}'_1 I'_1}^{\mathbf{p}_1} D_{\boldsymbol{\nu}_2 I_2, \boldsymbol{\nu}'_2 I'_2}^{\mathbf{p}_2} = \sum_{(\mathbf{a};\sigma)} \sum_{\substack{\boldsymbol{\nu}, I \\ \boldsymbol{\nu}', I'}} C_{\boldsymbol{\nu}_1 I_1, \boldsymbol{\nu}_2 I_2, \boldsymbol{\nu} I}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a};\sigma)} C_{\boldsymbol{\nu}'_1 I'_1, \boldsymbol{\nu}'_2 I'_2, \boldsymbol{\nu}' I'}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a};\sigma)} D_{\boldsymbol{\nu} I, \boldsymbol{\nu}' I'}^{\mathbf{a}} ,$$

where the summations are restricted to  $\nabla_{\boldsymbol{\nu}_1 + \boldsymbol{\nu}_2, \boldsymbol{\nu}} = \nabla_{\boldsymbol{\nu}'_1 + \boldsymbol{\nu}'_2, \boldsymbol{\nu}'} = 1$ ,  $\delta(I_1, I_2, I) = \delta(I'_1, I'_2, I') = 1$  and  $(\mathbf{a}; \sigma)$  such that  $Q(\mathbf{a}; \sigma)$  is in the Clebsch-Gordan series of  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2)$ .

*Proof.* The proof is analogous to the  $SU(2)$  case. Let  $g \in SU(3)$ . From (2.46) and (2.52), we have

$$(2.54) \quad \begin{aligned} &\sum_{\substack{\boldsymbol{\nu}_1, I_1 \\ \boldsymbol{\nu}_2, I_2}} D_{\boldsymbol{\nu}_1 I_1, \boldsymbol{\nu}'_1 I'_1}^{\mathbf{p}_1}(g) D_{\boldsymbol{\nu}_2 I_2, \boldsymbol{\nu}'_2 I'_2}^{\mathbf{p}_2}(g) e(\mathbf{p}_1; \boldsymbol{\nu}_1, I_1) \otimes e(\mathbf{p}_2; \boldsymbol{\nu}_2, I_2) = \\ &\sum_{\substack{(\mathbf{a};\sigma) \\ \boldsymbol{\nu}', I'}} C_{\boldsymbol{\nu}'_1 I'_1, \boldsymbol{\nu}'_2 I'_2, \boldsymbol{\nu}' I'}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a};\sigma)} \sum_{\boldsymbol{\nu}, I} D_{\boldsymbol{\nu} I, \boldsymbol{\nu}' I'}^{\mathbf{a}}(g) e((\mathbf{a}; \sigma), \boldsymbol{\nu}, I) , \end{aligned}$$

where the sum over  $(\mathbf{a}; \sigma)$ ,  $\boldsymbol{\nu}'$  and  $I'$  satisfies the statement. Now, we use (2.47) and (2.52) to obtain

$$(2.55) \quad \begin{aligned} &\sum_{\substack{\boldsymbol{\nu}_1, I_1 \\ \boldsymbol{\nu}_2, I_2}} D_{\boldsymbol{\nu}_1 I_1, \boldsymbol{\nu}'_1 I'_1}^{\mathbf{p}_1}(g) D_{\boldsymbol{\nu}_2 I_2, \boldsymbol{\nu}'_2 I'_2}^{\mathbf{p}_2}(g) e(\mathbf{p}_1; \boldsymbol{\nu}_1, I_1) \otimes e(\mathbf{p}_2; \boldsymbol{\nu}_2, I_2) = \\ &\sum_{\substack{\boldsymbol{\nu}_1, I_1 \\ \boldsymbol{\nu}_2, I_2}} \sum_{(\mathbf{a};\sigma)} \sum_{\substack{\boldsymbol{\nu}, I \\ \boldsymbol{\nu}', I'}} C_{\boldsymbol{\nu}'_1 I'_1, \boldsymbol{\nu}'_2 I'_2, \boldsymbol{\nu}' I'}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a};\sigma)} C_{\boldsymbol{\nu}_1 I_1, \boldsymbol{\nu}_2 I_2, \boldsymbol{\nu} I}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a};\sigma)} D_{\boldsymbol{\nu} I, \boldsymbol{\nu}' I'}^{\mathbf{a}}(g) \\ &\quad \times e(\mathbf{p}_1; \boldsymbol{\nu}_1, I_1) \otimes e(\mathbf{p}_2; \boldsymbol{\nu}_2, I_2) , \end{aligned}$$

where  $\nu$ ,  $\nu_1$ ,  $\nu_2$ ,  $I$ ,  $I_1$  and  $I_2$  are related as in the statement. The decomposition in a basis is unique, so this finishes the proof.  $\square$

Again, let  $\mathcal{H}$  be a Hilbert space with an irreducible  $SU(3)$ -representation of class  $Q(\mathbf{p})$ . Also, let  $\{e(\mathbf{p}; \nu, I)\}$  be a GT basis of such space and  $\{\check{e}(\check{\mathbf{p}}; \check{\nu}, \check{I})\}$  be the induced GT basis of its dual space. The trivial representation within  $\mathcal{B}(\mathcal{H})$  is spanned by the normalized operator

$$(2.56) \quad \begin{aligned} \frac{1}{\sqrt{\dim Q(\mathbf{p})}} \mathbb{1} &= \frac{1}{\sqrt{\dim Q(\mathbf{p})}} \sum_{\nu, I} e(\mathbf{p}; \nu, I) \otimes e^*(\mathbf{p}; \nu, I) \\ &= \frac{1}{\sqrt{\dim Q(\mathbf{p})}} \sum_{\nu, I} (-1)^{2(t+u)} e(\mathbf{p}; \nu, I) \otimes \check{e}(\check{\mathbf{p}}; \check{\nu}, \check{I}) . \end{aligned}$$

For operators  $A, R \in \mathcal{B}(\mathcal{H})$ ,

$$(2.57) \quad A = e(\mathbf{p}; \nu, I) \otimes \check{e}(\check{\mathbf{p}}; \nu', I') \quad , \quad R = e(\mathbf{p}; \mu', J') \otimes \check{e}(\check{\mathbf{p}}; \mu, J) ,$$

their product is given by

$$(2.58) \quad AR = \delta_{I', J'} \Delta_{\nu', \mu'}^{|\mathbf{p}|} (-1)^{2(t_{\nu'} + u_{\nu'})} e(\mathbf{p}; \nu, I) \otimes \check{e}(\check{\mathbf{p}}; \mu, J) ,$$

where  $t_{\nu'} = (\nu'_1 - \nu'_2)/2$  and  $u_{\nu'} = (\nu'_2 - \nu'_3)/2$  for  $\nu' = (\nu'_1, \nu'_2, \nu'_3)$ , cf. Definition 2.3.

**Lemma 2.14.** *Let  $(\mathbf{a}_1; \sigma_1)$  and  $(\mathbf{a}_2; \sigma_2)$  label  $SU(3)$ -representations in the Clebsch-Gordan series of  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}})$ . The operator product of elements of a coupled basis of  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}})$  can be decomposed into*

$$(2.59) \quad e((\mathbf{a}_1; \sigma_1); \nu_1, I_1) e((\mathbf{a}_2; \sigma_2); \nu_2, I_2) = \sum_{\substack{(\mathbf{a}; \sigma) \\ \nu, I}} \mathcal{M}[\mathbf{p}]_{\nu_1 I_1, \nu_2 I_2, \nu I}^{(\mathbf{a}_1; \sigma_1), (\mathbf{a}_2; \sigma_2), (\mathbf{a}; \sigma)} e((\mathbf{a}; \sigma); \nu, I) ,$$

with summation over  $(\mathbf{a}; \sigma)$  restricted to  $Q(\mathbf{a}; \sigma)$  in the Clebsch-Gordan series of  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}})$ , and

$$(2.60) \quad \begin{aligned} \mathcal{M}[\mathbf{p}]_{\nu_1 I_1, \nu_2 I_2, \nu I}^{(\mathbf{a}_1; \sigma_1), (\mathbf{a}_2; \sigma_2), (\mathbf{a}; \sigma)} &= \sum_{\substack{\mu_1, \mu_2, \mu_3 \\ J_1, J_2, J_3}} (-1)^{2(t_{\mu_2} + u_{\mu_2})} C_{\mu_1 J_1, \mu_2 J_2, \nu_1 I_1}^{\mathbf{p}, \check{\mathbf{p}}, (\mathbf{a}_1; \sigma_1)} \\ &\times C_{\mu_2 J_2, \mu_3 J_3, \nu_2 I_2}^{\mathbf{p}, \check{\mathbf{p}}, (\mathbf{a}_2; \sigma_2)} C_{\mu_1 J_1, \mu_3 J_3, \nu I}^{\mathbf{p}, \check{\mathbf{p}}, (\mathbf{a}; \sigma)} . \end{aligned}$$

In particular,  $\mathcal{M}[\mathbf{p}]_{\nu_1 I_1, \nu_2 I_2, \nu I}^{(\mathbf{a}_1; \sigma_1), (\mathbf{a}_2; \sigma_2), (\mathbf{a}; \sigma)} \neq 0$  implies that  $\nabla_{\nu_1 + \nu_2, \nu} = 1$  and  $I \leq I_1 + I_2 + |\mathbf{p}|$ .

*Proof.* Using (2.47), we write

$$(2.61) \quad e((\mathbf{a}_1; \sigma_1); \nu_1, I_1) = \sum_{\substack{\mu_1, J_1 \\ \mu_2, J_2}} C_{\mu_1 J_1, \mu_2 J_2, \nu_1 I_1}^{\mathbf{p}, \check{\mathbf{p}}, (\mathbf{a}_1; \sigma_1)} e(\mathbf{p}; \mu_1, J_1) \otimes \check{e}(\check{\mathbf{p}}; \mu_2, J_2)$$

and

$$(2.62) \quad e((\mathbf{a}_2; \sigma_2); \nu_2, I_2) = \sum_{\substack{\mu_3, J_3 \\ \mu_4, J_4}} C_{\mu_4 J_4, \mu_3 J_3, \nu_2 I_2}^{\mathbf{p}, \check{\mathbf{p}}, (\mathbf{a}_2; \sigma_2)} e(\mathbf{p}; \mu_4, J_4) \otimes \check{e}(\check{\mathbf{p}}; \mu_3, J_3) .$$

From (2.58), we have

$$(2.63) \quad e((\mathbf{a}_1; \sigma_1); \boldsymbol{\nu}_1, I_1) e((\mathbf{a}_2; \sigma_2); \boldsymbol{\nu}_2, I_2) = \sum_{\substack{\mu_1, \mu_2, \mu_3 \\ J_1, J_2, J_3}} (-1)^{2(t_{\mu_2} + u_{\mu_2})} C_{\mu_1 J_1, \mu_2 J_2, \nu_1 I_1}^{\mathbf{p}, \check{\mathbf{p}}, (\mathbf{a}_1; \sigma_1)} C_{\mu_2 J_2, \mu_3 J_3, \nu_2 I_2}^{\mathbf{p}, \check{\mathbf{p}}, (\mathbf{a}_2; \sigma_2)} \times e(\mathbf{p}; \boldsymbol{\mu}_1, J_1) \otimes \check{e}(\check{\mathbf{p}}; \boldsymbol{\mu}_3, J_3) .$$

Now, from (2.46) and (2.52),

$$(2.64) \quad e((\mathbf{a}_1; \sigma_1); \boldsymbol{\nu}_1, I_1) e((\mathbf{a}_2; \sigma_2); \boldsymbol{\nu}_2, I_2) = \sum_{\substack{(\mathbf{a}; \sigma) \\ \boldsymbol{\nu}, I}} \sum_{\substack{\mu_1, \mu_2, \mu_3 \\ J_1, J_2, J_3}} (-1)^{2(t_{\mu_2} + u_{\mu_2})} C_{\mu_1 J_1, \mu_2 J_2, \nu_1 I_1}^{\mathbf{p}, \check{\mathbf{p}}, (\mathbf{a}_1; \sigma_1)} C_{\mu_2 J_2, \mu_3 J_3, \nu_2 I_2}^{\mathbf{p}, \check{\mathbf{p}}, (\mathbf{a}_2; \sigma_2)} \times C_{\mu_1 J_1, \mu_3 J_3, \nu I}^{\mathbf{p}, \check{\mathbf{p}}, (\mathbf{a}; \sigma)} e((\mathbf{a}; \sigma); \boldsymbol{\nu}, I) ,$$

where  $\nabla_{\mu_1 + \mu_2, \nu_1} = \nabla_{\mu_2 + \mu_3, \nu_2} = \nabla_{\mu_1 + \mu_3, \nu} = 1$ ,  $J_1 \leq I_1 + J_2$ ,  $J_3 \leq I_2 + J_2$  and  $I \leq J_1 + J_3$ . It follows that  $\nabla_{\nu_1 + \nu_2, \nu} = 1$  and, considering that  $J_2 \leq |\mathbf{p}|/2$ , which can be inferred from (2.21), we have  $I \leq I_1 + I_2 + |\mathbf{p}|$ .  $\square$

**2.4. Mixed Casimir operators and symmetric CG coefficients.** We have avoided until now the problem of specifying a decomposition for degenerate representations in general Clebsch-Gordan series  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2)$ . If  $Q(\mathbf{a})$  is such that  $\mathbf{m}(\mathbf{p}_1, \mathbf{p}_2; \mathbf{a}) > 1$  (cf. Notation 2), there is no canonically unique way to decompose

$$(2.65) \quad \bigoplus_{\sigma=1}^{\mathbf{m}(\mathbf{p}_1, \mathbf{p}_2; \mathbf{a})} Q(\mathbf{a}; \sigma) \subset Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2)$$

into irreducible representations of class  $Q(\mathbf{a})$ . To fix a unique convention for such a decomposition, we shall use the method of mixed Casimir operators, based on [8, 7, 21] and described below. The envisaged decomposition provides more symmetric Clebsch-Gordan coefficients.

Let  $\mathcal{H}$  be a Hilbert space carrying an irreducible representation of class  $Q(p, q)$  and let  $A_{jk}$ , for  $j, k \in \{1, 2, 3\}$ , be the generators satisfying:

$$(2.66) \quad A_{12} = T_+ , \quad A_{23} = U_+ , \quad T_3 = \frac{1}{2}(A_{11} - A_{22}) , \quad U_3 = \frac{1}{2}(A_{22} - A_{33}) ,$$

$$(2.67) \quad A_{jk}^\dagger = A_{kj} , \quad A_{11} + A_{22} + A_{33} = 0 ,$$

$$(2.68) \quad [A_{jk}, A_{lm}] = \delta_{l,k} A_{jm} - \delta_{j,m} A_{lk} .$$

Then,  $Q(q, p)$  is generated by the operators

$$(2.69) \quad \check{A}_{jk} = -A_{kj} .$$

For  $Q(p, q)$ , the *quadratic and cubic Casimir operators* are

$$(2.70) \quad C_2 := \frac{1}{2} \sum_{j,k=1}^3 A_{jk} A_{kj} , \quad C_3 := \sum_{j,k,l=1}^3 A_{jk} A_{kl} A_{lj} ,$$

so that

$$(2.71) \quad \begin{aligned} C_2 &= \frac{1}{3}[(p+q)(p+q+3) - pq]\mathbb{1} , \\ C_3 - 3C_2 &= \frac{1}{9}(p-q)(p+2q+3)(2p+q+3)\mathbb{1} , \end{aligned}$$

cf. [23].

Now, for  $x \in \{1, 2, 3\}$ , consider the Hilbert space  $\mathcal{H}_x$  carrying the representation  $Q(\mathbf{p}_x) = Q(p_x, q_x)$  and the triple tensor product  $\mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \mathcal{H}_3$ . Also, let  $A_{jk}^{(x)}$ ,  $C_2^{(x)}$  and  $C_3^{(x)}$  be operators relative to  $Q(\mathbf{p}_x)$  as defined above. For simplicity, given an operator  $A^{(x)}$  on  $\mathcal{H}_x$ , we just write  $A^{(x)}$  to denote its tensor product with the identity operator. Thus,

$$(2.72) \quad A_{jk}^{(x)} A_{lm}^{(y)} = A_{lm}^{(y)} A_{jk}^{(x)}$$

for  $x \neq y$ . The *mixed Casimir operators* are defined by

$$(2.73) \quad C_2^{xy} := \sum_{j,k} A_{jk}^{(x)} A_{kj}^{(y)} = \sum_{j,k} A_{jk}^{(y)} A_{kj}^{(x)} =: C_2^{yx} ,$$

$$(2.74) \quad C_3^{xyz} := \sum_{j,k,l} A_{jk}^{(x)} A_{kl}^{(y)} A_{lj}^{(z)} ,$$

for indices not all equal. If  $x, y, z$  are all distinct, equation (2.72) implies

$$(2.75) \quad \sum_{j,k,l} A_{jk}^{(x)} A_{kl}^{(y)} A_{lj}^{(z)} = \sum_{j,k,l} A_{kl}^{(y)} A_{lj}^{(z)} A_{jk}^{(x)} = \sum_{j,k,l} A_{lj}^{(z)} A_{jk}^{(x)} A_{kl}^{(y)} ,$$

so

$$(2.76) \quad C_3^{xyz} = C_3^{yzx} = C_3^{zxy} ,$$

and

$$(2.77) \quad C_3^{xyy} = \sum_{j,k,l} A_{jk}^{(x)} A_{kl}^{(y)} A_{lj}^{(y)} = \sum_{j,k,l} A_{kl}^{(y)} A_{lj}^{(y)} A_{jk}^{(x)} = C_3^{yyx} ,$$

but

$$(2.78) \quad \begin{aligned} C_3^{xyx} &= \sum_{j,k,l} A_{jk}^{(x)} A_{kl}^{(y)} A_{lj}^{(x)} = \sum_{j,k,l} A_{kl}^{(y)} A_{jk}^{(x)} A_{lj}^{(x)} \\ &= \sum_{j,k,l} A_{kl}^{(y)} \left( [A_{jk}^{(x)}, A_{lj}^{(x)}] + A_{lj}^{(x)} A_{jk}^{(x)} \right) \\ &= \sum_{j,k,l} A_{kl}^{(y)} \left( \delta_{l,k} A_{jj}^{(x)} - A_{lk}^{(x)} + A_{lj}^{(x)} A_{jk}^{(x)} \right) \\ &= C_3^{yxx} - C_2^{yx} . \end{aligned}$$

Such mixed Casimir operators arise, for example, from the Casimir operators on  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2) \otimes Q(\mathbf{p}_3)$ :

$$(2.79) \quad \frac{1}{2} \sum_{j,k} \left( A_{jk}^{(1)} + A_{jk}^{(2)} + A_{jk}^{(3)} \right) \left( A_{kj}^{(1)} + A_{kj}^{(2)} + A_{kj}^{(3)} \right) = C_2^{(1)} + C_2^{(2)} + C_2^{12} + C_2^{23} + C_2^{31} ,$$

$$\begin{aligned}
(2.80) \quad & \sum_{j,k,l} \left( A_{jk}^{(1)} + A_{jk}^{(2)} + A_{jk}^{(3)} \right) \left( A_{kl}^{(1)} + A_{kl}^{(2)} + A_{kl}^{(3)} \right) \left( A_{lj}^{(1)} + A_{lj}^{(2)} + A_{lj}^{(3)} \right) \\
& = C_3^{(1)} + C_3^{(2)} + C_3^{(3)} + C_3^{112} + C_3^{113} + C_3^{121} + C_3^{122} + C_3^{123} + C_3^{131} + C_3^{132} \\
& \quad + C_3^{133} + C_3^{211} + C_3^{212} + C_3^{213} + C_3^{221} + C_3^{223} + C_3^{231} + C_3^{232} + C_3^{233} \\
& \quad + C_3^{311} + C_3^{312} + C_3^{313} + C_3^{321} + C_3^{322} + C_3^{323} + C_3^{331} + C_3^{332} .
\end{aligned}$$

From the mixed cubic Casimir operators, we set

$$(2.81) \quad \mathbf{S}_{xy} := \frac{1}{2}(C_3^{xyy} - C_3^{yxx})$$

and

$$(2.82) \quad \mathbf{S}_{xyz} := \frac{1}{3}(\mathbf{S}_{xy} + \mathbf{S}_{yz} + \mathbf{S}_{zx}) .$$

**Lemma 2.15.** *The operators  $\mathbf{S}_{xy}$  and  $\mathbf{S}_{xyz}$  are anti-symmetric under odd permutation of the indices, that is,*

$$(2.83) \quad \mathbf{S}_{xy} = -\mathbf{S}_{yx} ,$$

$$(2.84) \quad \mathbf{S}_{xyz} = -\mathbf{S}_{yxz} = -\mathbf{S}_{xzy} .$$

*Proof.* We have that

$$\mathbf{S}_{xy} = \frac{1}{2}(C_3^{xyy} - C_3^{yxx}) = -\frac{1}{2}(C_3^{yxx} - C_3^{xyy}) = -\mathbf{S}_{yx} ,$$

so

$$\mathbf{S}_{xyz} = \frac{1}{3}(\mathbf{S}_{xy} + \mathbf{S}_{yz} + \mathbf{S}_{zx}) = -\frac{1}{3}(\mathbf{S}_{yx} + \mathbf{S}_{zy} + \mathbf{S}_{xz}) .$$

Thus,  $\mathbf{S}_{xyz} = -\mathbf{S}_{yxz} = -\mathbf{S}_{xzy}$  .  $\square$

For representations  $Q(\check{\mathbf{p}}_x)$  generated by the operators

$$(2.85) \quad \check{A}_{jk}^{(x)} = -A_{kj}^{(x)} ,$$

cf. (2.69), we have

$$(2.86) \quad \check{C}_2^{xy} = \sum_{j,k} \check{A}_{jk}^{(x)} \check{A}_{kj}^{(y)} = \sum_{j,k} A_{kj}^{(x)} A_{jk}^{(y)} = C_2^{xy} ,$$

$$(2.87) \quad \check{C}_3^{xyz} = \sum_{j,k,l} \check{A}_{jk}^{(x)} \check{A}_{kl}^{(y)} \check{A}_{lj}^{(z)} = -\sum_{j,k,l} A_{kj}^{(x)} A_{lk}^{(y)} A_{jl}^{(z)} .$$

In particular,

$$\begin{aligned}
(2.88) \quad \check{C}_3^{xyy} & = -\sum_{j,k,l} A_{kj}^{(x)} A_{lk}^{(y)} A_{jl}^{(y)} \\
& = -\sum_{j,k,l} A_{kj}^{(x)} \left( [A_{lk}^{(y)}, A_{jl}^{(y)}] + A_{jl}^{(y)} A_{lk}^{(y)} \right) \\
& = -\sum_{j,k,l} A_{kj}^{(x)} (\delta_{j,k} A_{ll}^{(y)} - A_{jk}^{(y)} + A_{jl}^{(y)} A_{lk}^{(y)}) \\
& = C_2^{xy} - C_3^{xyy}
\end{aligned}$$

Therefore, we have:

**Lemma 2.16.** *The operators  $\mathbf{S}_{xy}$  and  $\mathbf{S}_{xyz}$  are anti-symmetric under dualization, that is,*

$$(2.89) \quad \check{\mathbf{S}}_{xy} = -\mathbf{S}_{xy} ,$$

$$(2.90) \quad \check{\mathbf{S}}_{xyz} = -\mathbf{S}_{xyz} .$$

*Proof.* From (2.73), (2.81) and (2.88), we have that

$$\check{\mathbf{S}}_{xy} = \frac{1}{2}(\check{C}_3^{xyy} - \check{C}_3^{yxx}) = -\frac{1}{2}(C_3^{xyy} - C_3^{yxx}) = -\mathbf{S}_{xy} .$$

From (2.89), we have

$$\check{\mathbf{S}}_{xyz} = \frac{1}{3}(\check{\mathbf{S}}_{xy} + \check{\mathbf{S}}_{yz} + \check{\mathbf{S}}_{zx}) = -\frac{1}{3}(\mathbf{S}_{xy} + \mathbf{S}_{yz} + \mathbf{S}_{zx}) = -\mathbf{S}_{xyz} .$$

□

Now, let  $\mathcal{H}^0 \equiv \mathcal{H}_{123}^0$  be a maximal subspace of  $\mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \mathcal{H}_3$  where  $SU(3)$  acts trivially. That means

$$(2.91) \quad A_{jk}^{(1)} + A_{jk}^{(2)} + A_{jk}^{(3)} = 0$$

on  $\mathcal{H}^0$ , for all  $j, k \in \{1, 2, 3\}$ .

**Lemma 2.17.**  *$\mathcal{H}_{123}^0$  is not null if and only if there is a representation of class  $Q(\check{\mathbf{p}}_3)$  in the Clebsch-Gordan series of  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2)$ .*

*Proof.* Considering the Clebsch-Gordan series

$$(2.92) \quad Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2) = \bigoplus_{(\mathbf{a}; \sigma)} Q(\mathbf{a}; \sigma) ,$$

we have

$$(2.93) \quad Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2) \otimes Q(\mathbf{p}_3) = \bigoplus_{(\mathbf{a}; \sigma)} Q(\mathbf{a}; \sigma) \otimes Q(\mathbf{p}_3) .$$

Note that  $k \geq 1$  in (2.41), so there exists a factor of class  $Q(0, 0)$  in the CG series of  $Q(a, b) \otimes Q(p_3, q_3)$  if and only if there is  $0 \leq n \leq \min(a, q_3)$  and  $0 \leq m \leq \min(p_3, b)$  satisfying

$$(2.94) \quad a - n + p_3 - m = b - m + q_3 - n = 0 .$$

The only possible solution is  $n = a = q_3$  and  $m = p_3 = b$ . Thus,  $\mathcal{H}^0$  is not null if and only if there is a representation of class  $Q(\mathbf{a}) = Q(\check{\mathbf{p}}_3)$  in the l.h.s. of (2.92). □

**Lemma 2.18.** *On  $\mathcal{H}^0$ , the following holds:*

$$(2.95) \quad \mathbf{S}_{123} = \mathbf{S}_{12} - \frac{1}{3}C_3^{(1)} + \frac{1}{3}C_3^{(2)} .$$

*Proof.* From (2.91),

$$(2.96) \quad \begin{aligned} C_3^{233} &= \sum_{j,k,l} A_{jk}^{(2)} \left( A_{kl}^{(1)} + A_{kl}^{(2)} \right) \left( A_{lj}^{(1)} + A_{lj}^{(2)} \right) \\ &= C_3^{211} + C_3^{212} + C_3^{221} + C_3^{(2)} . \end{aligned}$$

For  $x = 1$  and  $y = 2$ , equation (2.77) reads

$$(2.97) \quad C_3^{221} = C_3^{122}$$

and likewise, for  $x = 2$  and  $y = 1$ ,

$$(2.98) \quad C_3^{211} = C_3^{112} .$$

Also,

$$(2.99) \quad C_3^{212} = C_3^{122} - C_2^{12} ,$$

cf. (2.78), and

$$(2.100) \quad C_3^{233} = C_3^{211} + 2C_3^{122} + C_3^{(2)} - C_2^{12} .$$

For  $C_3^{322}$  we have

$$(2.101) \quad \begin{aligned} C_3^{322} &= - \sum_{j,k,l} (A_{jk}^{(1)} + A_{jk}^{(2)}) A_{kl}^{(2)} A_{lj}^{(2)} \\ &= -C_3^{122} - C_3^{(2)} . \end{aligned}$$

Then

$$(2.102) \quad \mathbf{S}_{23} = \frac{1}{2} (C_3^{211} + 3C_3^{122} + 2C_3^{(2)} - C_2^{12}) .$$

By formal identification of the formulas, we get in a straightforward way

$$(2.103) \quad \mathbf{S}_{31} = -\mathbf{S}_{13} = -\frac{1}{2} (C_3^{122} + 3C_3^{211} + 2C_3^{(1)} - C_2^{21}) ,$$

so

$$(2.104) \quad \mathbf{S}_{23} + \mathbf{S}_{31} = C_3^{122} - C_3^{211} - C_3^{(1)} + C_3^{(2)} = 2\mathbf{S}_{12} - C_3^{(1)} + C_3^{(2)} ,$$

where we used  $C_2^{12} = C_2^{21}$ . Therefore,

$$(2.105) \quad \mathbf{S}_{123} = \frac{1}{3} (\mathbf{S}_{12} + \mathbf{S}_{23} + \mathbf{S}_{31}) = \mathbf{S}_{12} - \frac{1}{3} C_3^{(1)} + \frac{1}{3} C_3^{(2)} ,$$

on  $\mathcal{H}^0$ . □

**Notation 3.** In view of the previous lemma, for  $\mathcal{H}^0 \neq 0$ , we shall denote

$$(2.106) \quad \mathbf{S}_{123}^0 := \mathbf{S}_{123}|_{\mathcal{H}^0} .$$

**Lemma 2.19.** The operators  $\mathbf{S}_{12}$  and  $\mathbf{S}_{123}^0$  are Hermitian and  $SU(3)$ -invariant.

*Proof.* By straightforward calculation, we have

$$(2.107) \quad (C_3^{xyy})^\dagger = \sum_{j,k,l} \left( A_{jk}^{(x)} A_{kl}^{(y)} A_{lj}^{(y)} \right)^\dagger = \sum_{j,k,l} A_{jl}^{(y)} A_{lk}^{(y)} A_{kj}^{(x)} = C_3^{yyx} ,$$

so  $(C_3^{122})^\dagger = C_3^{122}$  and  $(C_3^{211})^\dagger = C_3^{211}$ , which implies

$$(2.108) \quad (\mathbf{S}_{12})^\dagger = \frac{1}{2} (C_3^{122} - C_3^{211})^\dagger = \frac{1}{2} (C_3^{122} - C_3^{211}) = \mathbf{S}_{12} .$$

For the  $SU(3)$ -invariance, we will need the following equalities:

$$(2.109) \quad A_{cd}^{(x)} A_{jk}^{(x)} = \left[ A_{cd}^{(x)}, A_{jk}^{(x)} \right] + A_{jk}^{(x)} A_{cd}^{(x)} = \delta_{j,d} A_{ck}^{(x)} - \delta_{c,k} A_{jd}^{(x)} + A_{jk}^{(x)} A_{cd}^{(x)} ,$$

$$\begin{aligned}
(2.110) \quad A_{cd}^{(x)} A_{kl}^{(x)} A_{lj}^{(x)} &= \left( \left[ A_{cd}^{(x)}, A_{kl}^{(x)} \right] + A_{kl}^{(x)} A_{cd}^{(x)} \right) A_{lj}^{(x)} \\
&= \left( \delta_{k,d} A_{cl}^{(x)} - \delta_{c,l} A_{kd}^{(x)} + A_{kl}^{(x)} A_{cd}^{(x)} \right) A_{lj}^{(x)} \\
&= \left( \delta_{k,d} A_{cl}^{(x)} - \delta_{c,l} A_{kd}^{(x)} \right) A_{lj}^{(x)} \\
&\quad + A_{kl}^{(x)} \left( \left[ A_{cd}^{(x)}, A_{lj}^{(x)} \right] + A_{lj}^{(x)} A_{cd}^{(x)} \right) \\
&= \left( \delta_{k,d} A_{cl}^{(x)} - \delta_{c,l} A_{kd}^{(x)} \right) A_{lj}^{(x)} \\
&\quad + A_{kl}^{(x)} \left( \delta_{l,d} A_{cj}^{(x)} - \delta_{c,j} A_{ld}^{(x)} \right) + A_{kl}^{(x)} A_{lj}^{(x)} A_{cd}^{(x)} .
\end{aligned}$$

With this, we obtain

$$(2.111) \quad A_{cd}^{(x)} C_3^{xyy} = \sum_{k,l} A_{ck}^{(x)} A_{kl}^{(y)} A_{ld}^{(y)} - \sum_{j,l} A_{jd}^{(x)} A_{cl}^{(y)} A_{lj}^{(y)} + C_3^{xyy} A_{cd}^{(x)} ,$$

$$(2.112) \quad A_{cd}^{(x)} C_3^{yxx} = \sum_{j,l} A_{jd}^{(y)} A_{cl}^{(x)} A_{lj}^{(x)} - \sum_{k,l} A_{ck}^{(y)} A_{kl}^{(x)} A_{ld}^{(x)} + C_3^{yxx} A_{cd}^{(x)} ,$$

Thus,

$$(2.113) \quad \left[ A_{cd}^{(1)}, C_3^{122} - C_3^{211} \right] = \left[ A_{cd}^{(2)}, C_3^{211} - C_3^{122} \right] \implies \left[ A_{cd}^{(1)} + A_{cd}^{(2)}, \mathbf{S}_{12} \right] = 0 .$$

Hence,  $\mathbf{S}_{12}$  is Hermitian and  $SU(3)$ -invariant. The result for  $\mathbf{S}_{123}^0$  follows immediately from Lemma 2.18.  $\square$

In this way, we decompose degenerate representations in the Clebsch-Gordan series of  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2)$  via diagonalization of the operator

$$(2.114) \quad \mathbf{S} := \mathbf{S}_{12} - \frac{1}{3} C_3^{(1)} + \frac{1}{3} C_3^{(2)} ,$$

satisfying

$$(2.115) \quad \mathbf{S}|_{\mathcal{H}_{123}^0} = \mathbf{S}_{123}^0 ,$$

cf. (2.106). Because  $\mathbf{S}$  is built from Casimir operators, the eigenvalues  $s_{123}$  of  $\mathbf{S}_{123}^0$  depend only on the representations comprising  $\mathcal{H}_{123}^0 \subset \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \mathcal{H}_3$ .

We then have the following:

**Theorem 2.20** ([8]). *The eigenvalues of  $\mathbf{S}_{123}^0$  are distinct, for distinct representations in  $\mathcal{H}_{123}^0$ .*

Thus, for

$$(2.116) \quad \bigoplus_{\sigma=1}^{\mathbf{m}(\mathbf{p}_1, \mathbf{p}_2; \mathbf{a})} Q(\mathbf{a}; \sigma) \subset Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2) ,$$

with  $Q(\check{\mathbf{a}}) = Q(\mathbf{p}_3)$ , cf. Lemma 2.17, we can define a function

$$(2.117) \quad s_{\mathbf{p}_1, \mathbf{p}_2; \mathbf{a}} : \{1, \dots, \mathbf{m}(\mathbf{p}_1, \mathbf{p}_2; \mathbf{a})\} \rightarrow \mathbb{R} , \quad \sigma \mapsto s_{\mathbf{p}_1, \mathbf{p}_2; \mathbf{a}}(\sigma) ,$$

which indexes the eigenvalues of  $\mathbf{S}_{123}^0$  in this case. For simplicity, we shall denote

$$(2.118) \quad s_{\mathbf{p}_1, \mathbf{p}_2; \mathbf{a}} \equiv s_{\mathbf{a}}$$

and, henceforth, we shall adopt the convention that  $s_{\mathbf{a}}$  is an increasing function of the multiplicity counting index, that is,

$$(2.119) \quad \{1, \dots, \mathbf{m}(\mathbf{p}_1, \mathbf{p}_2; \mathbf{a})\} \ni \sigma \mapsto s_{\mathbf{a}}(\sigma) \in \mathbb{R}, \quad s_{\mathbf{a}}(\sigma) < s_{\mathbf{a}}(\sigma + 1).$$

In other words, the eigenvalues of  $\mathbf{S}_{123}^0$  are indexed by increasing order.

**Remark 2.21.** In [8], the authors make use of polynomial basis for  $SU(3)$  representations as constructed in [7] to compute the matrix entries of an operator equivalent to  $\mathbf{S}$  in order to prove Theorem 2.20. In the Appendix, we illustrate the computation for the degenerate representation of class  $Q(1, 1)$  within  $Q(1, 1) \otimes Q(1, 1)$ .

**Notation 4.** For  $\bigoplus_{\sigma} Q(\mathbf{a}; \sigma)$  in the CG series of  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2)$ , the following involution in the set of multiplicity indices will be relevant:

$$(2.120) \quad \{1, \dots, \mathbf{m}(\mathbf{p}_1, \mathbf{p}_2; \mathbf{a})\} \ni \sigma \mapsto \check{\sigma} = \mathbf{m}(\mathbf{p}_1, \mathbf{p}_2; \mathbf{a}) - \sigma + 1.$$

**Remark 2.22.** The involution (2.120) is at par with the involution  $s \mapsto -s$  which is consequence of (2.83)-(2.84) and (2.89)-(2.90), taking into account the convention (2.119).

From (2.56) and Lemmas 2.15-2.19, considering the convention in (2.119), we can choose coupled basis  $\{e_{12}(\check{\mathbf{p}}_3; \sigma; \boldsymbol{\nu}, I)\}$ ,  $\{e_{21}(\check{\mathbf{p}}_3; \sigma; \boldsymbol{\nu}, I)\}$ ,  $\{e_{13}(\check{\mathbf{p}}_2; \sigma; \boldsymbol{\nu}, I)\}$  and  $\{\check{e}_{12}(\mathbf{p}_3; \sigma; \boldsymbol{\nu}, I)\}$  for  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2)$ ,  $Q(\mathbf{p}_2) \otimes Q(\mathbf{p}_1)$ ,  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_3)$  and  $Q(\check{\mathbf{p}}_1) \otimes Q(\check{\mathbf{p}}_2)$ , respectively, such that

$$(2.121) \quad 1 \leq \sigma \leq \mathbf{m}(\mathbf{p}_1, \mathbf{p}_2; \check{\mathbf{p}}_3) = \mathbf{m}(\mathbf{p}_2, \mathbf{p}_1; \check{\mathbf{p}}_3) = \mathbf{m}(\mathbf{p}_1, \mathbf{p}_3; \check{\mathbf{p}}_2) = \mathbf{m}(\check{\mathbf{p}}_1, \check{\mathbf{p}}_2; \mathbf{p}_3)$$

and  $\mathcal{H}^0$  is spanned by

$$(2.122) \quad \begin{aligned} & \sum_{\boldsymbol{\nu}, I} \frac{(-1)^{2(t+u)}}{\sqrt{\dim Q(\mathbf{p}_3)}} e_{12}(\check{\mathbf{p}}_3; \sigma; \boldsymbol{\nu}, I) \otimes e(\mathbf{p}_3; \check{\boldsymbol{\nu}}, I) \\ &= (-1)^{|\mathbf{p}_1|+|\mathbf{p}_2|+|\mathbf{p}_3|} \sum_{\boldsymbol{\nu}, I} \frac{(-1)^{2(t+u)}}{\sqrt{\dim Q(\mathbf{p}_3)}} e_{21}(\check{\mathbf{p}}_3; \check{\sigma}; \boldsymbol{\nu}, I) \otimes e(\mathbf{p}_3; \check{\boldsymbol{\nu}}, I) \\ &= (-1)^{|\mathbf{p}_1|+|\mathbf{p}_2|+|\mathbf{p}_3|} \sum_{\boldsymbol{\nu}, I} \frac{(-1)^{2(t+u)}}{\sqrt{\dim Q(\mathbf{p}_2)}} e_{13}(\check{\mathbf{p}}_2; \check{\sigma}; \boldsymbol{\nu}, I) \otimes e(\mathbf{p}_2; \check{\boldsymbol{\nu}}, I) \\ &= (-1)^{|\mathbf{p}_1|+|\mathbf{p}_2|+|\mathbf{p}_3|} \sum_{\boldsymbol{\nu}, I} \frac{(-1)^{2(t+u)}}{\sqrt{\dim Q(\check{\mathbf{p}}_3)}} \check{e}_{12}(\mathbf{p}_3; \check{\sigma}; \boldsymbol{\nu}, I) \otimes \check{e}(\check{\mathbf{p}}_3; \check{\boldsymbol{\nu}}, I), \end{aligned}$$

where we have made use of (2.120). As consequence, now the Clebsch-Gordan coefficients satisfy a bigger set of symmetry relations, as follows.

**Theorem 2.23.** For representations of class  $Q(\mathbf{p}_1)$  and  $Q(\mathbf{p}_2)$ , the Clebsch-Gordan coefficients for the Clebsch-Gordan series  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2) = \bigoplus Q(\mathbf{a}; \sigma)$  satisfy

$$(2.123) \quad \begin{aligned} C_{\boldsymbol{\nu}_1 I_1, \boldsymbol{\nu}_2 I_2, \boldsymbol{\nu} I}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a}; \sigma)} &= (-1)^{|\mathbf{p}_1|+|\mathbf{p}_2|+|\mathbf{a}|} C_{\boldsymbol{\nu}_2 I_2, \boldsymbol{\nu}_1 I_1, \boldsymbol{\nu} I}^{\mathbf{p}_2, \mathbf{p}_1, (\mathbf{a}; \check{\sigma})} \\ &= (-1)^{|\mathbf{p}_1|-2(t_{\nu_1}+u_{\nu_1})} \sqrt{\frac{\dim Q(\mathbf{a})}{\dim Q(\mathbf{p}_2)}} C_{\boldsymbol{\nu}_1 I_1, \check{\boldsymbol{\nu}} I, \check{\boldsymbol{\nu}}_2 I_2}^{\mathbf{p}_1, \check{\mathbf{a}}, (\check{\mathbf{p}}_2; \check{\sigma})} \\ &= (-1)^{|\mathbf{p}_1|+|\mathbf{p}_2|+|\mathbf{a}|} C_{\check{\boldsymbol{\nu}}_1 I_1, \check{\boldsymbol{\nu}}_2 I_2, \check{\boldsymbol{\nu}} I}^{\check{\mathbf{p}}_1, \check{\mathbf{p}}_2, (\check{\mathbf{a}}; \check{\sigma})}. \end{aligned}$$

*Proof.* By writing (2.122) with  $\mathbf{a} = \check{\mathbf{p}}_3$  and taking the inner product with suitable uncoupled basis, the result follows straightforwardly from the symmetries of  $\mathbf{S}$  satisfying Lemma 2.18, cf. (2.83)-(2.84) and (2.89)-(2.90), using (2.119) and (2.120).  $\square$

We highlight that each generator  $A_{jk}$  can be realized as a real matrix on a GT basis, so  $\mathbf{S}$  can be seen as a symmetric real matrix acting on

$$(2.124) \quad \text{Span}_{\mathbb{R}}\{e(\mathbf{p}_1; \boldsymbol{\nu}, I) \otimes e(\mathbf{p}_2; \boldsymbol{\mu}, J)\} ,$$

which means that the elements of any basis of the previous paragraph can be constructed as real linear combinations of the respective uncoupled basis. With this convention, all Clebsch-Gordan coefficients are still real, and (2.48) and (2.52) still hold.

In particular, the Hermitian conjugate  $\dagger$  of operators in  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}})$  satisfies

$$(2.125) \quad e^\dagger((\mathbf{a}; \sigma); \boldsymbol{\nu}, I) = (-1)^{2(t+u)} e((\check{\mathbf{a}}; \check{\sigma}); \check{\boldsymbol{\nu}}, I)$$

and the phases are chosen such that

$$(2.126) \quad e((0, 0); (0, 0, 0), 0) = \frac{(-1)^{|\mathbf{p}|}}{\sqrt{\dim Q(\mathbf{p})}} \mathbb{1} ,$$

this being the only element with non vanishing trace.

Besides Hermitian conjugate, the adjoint

$$(2.127) \quad * : Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}}) \rightarrow Q(\check{\mathbf{p}}) \otimes Q(\mathbf{p}) : A \mapsto A^* ,$$

is given in the uncoupled basis by

$$(2.128) \quad \begin{aligned} * : e(\mathbf{p}; \boldsymbol{\nu}, I) \otimes \check{e}(\check{\mathbf{p}}; \boldsymbol{\mu}, J) &= (-1)^{2(t_\mu + u_\mu)} e(\mathbf{p}; \boldsymbol{\nu}, I) \otimes e^*(\mathbf{p}; \check{\boldsymbol{\mu}}, J) \\ &\mapsto (-1)^{2(t_\mu + u_\mu)} e^*(\mathbf{p}; \check{\boldsymbol{\mu}}, J) \otimes e(\mathbf{p}; \boldsymbol{\nu}, I) \\ &= \check{e}(\check{\mathbf{p}}; \boldsymbol{\mu}, J) \otimes e(\mathbf{p}; \boldsymbol{\nu}, I) , \end{aligned}$$

cf. (2.34). Thus, for the coupled basis of  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}})$ , cf. (2.47), the adjoint is given by

$$(2.129) \quad \begin{aligned} * : e((\mathbf{a}; \sigma); \boldsymbol{\nu}, I) &= \sum_{\substack{\boldsymbol{\nu}_1, I_1 \\ \boldsymbol{\nu}_2, I_2}} C_{\boldsymbol{\nu}_1 I_1, \boldsymbol{\nu}_2 I_2, \boldsymbol{\nu} I}^{\mathbf{p}, \check{\mathbf{p}}, (\mathbf{a}; \sigma)} e(\mathbf{p}; \boldsymbol{\nu}_1, I_1) \otimes \check{e}(\check{\mathbf{p}}; \boldsymbol{\nu}_2, I_2) \\ &\mapsto \sum_{\substack{\boldsymbol{\nu}_1, I_1 \\ \boldsymbol{\nu}_2, I_2}} C_{\boldsymbol{\nu}_1 I_1, \boldsymbol{\nu}_2 I_2, \boldsymbol{\nu} I}^{\mathbf{p}, \check{\mathbf{p}}, (\mathbf{a}; \sigma)} \check{e}(\check{\mathbf{p}}; \boldsymbol{\nu}_2, I_2) \otimes e(\mathbf{p}; \boldsymbol{\nu}_1, I_1) \\ &= (-1)^{|\mathbf{a}|} \sum_{\substack{\boldsymbol{\nu}_1, I_1 \\ \boldsymbol{\nu}_2, I_2}} C_{\boldsymbol{\nu}_2 I_2, \boldsymbol{\nu}_1 I_1, \boldsymbol{\nu} I}^{\check{\mathbf{p}}, \mathbf{p}, (\mathbf{a}; \check{\sigma})} \check{e}(\check{\mathbf{p}}; \boldsymbol{\nu}_2, I_2) \otimes e(\mathbf{p}; \boldsymbol{\nu}_1, I_1) . \end{aligned}$$

In the light of Remark 2.4, from the above calculation, we identify

$$(2.130) \quad \check{e}((\mathbf{a}; \check{\sigma}); \boldsymbol{\nu}, I) := \sum_{\substack{\boldsymbol{\nu}_1, I_1 \\ \boldsymbol{\nu}_2, I_2}} C_{\boldsymbol{\nu}_2 I_2, \boldsymbol{\nu}_1 I_1, \boldsymbol{\nu} I}^{\check{\mathbf{p}}, \mathbf{p}, (\mathbf{a}; \check{\sigma})} \check{e}(\check{\mathbf{p}}; \boldsymbol{\nu}_2, I_2) \otimes e(\mathbf{p}; \boldsymbol{\nu}_1, I_1) ,$$

so that

$$(2.131) \quad e^*((\mathbf{a}; \sigma); \boldsymbol{\nu}, I) = (-1)^{|\mathbf{a}|} \check{e}((\mathbf{a}; \check{\sigma}); \boldsymbol{\nu}, I) .$$

**Definition 2.24.** Given a coupled basis  $\{e((\mathbf{a}; \sigma); \boldsymbol{\nu}, I)\}$  of  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}})$ , the induced coupled basis of  $Q(\check{\mathbf{p}}) \otimes Q(\mathbf{p})$  is the basis  $\{\check{e}((\mathbf{a}; \check{\sigma}); \boldsymbol{\nu}, I)\}$  satisfying (2.130)-(2.131).

**2.5. Wigner symbols and the operator product.** For  $SU(2)$ , the Clebsch-Gordan coefficients can be substituted by other coefficients with better symmetry properties, the so-called *Wigner 3jm-symbols* (cf. e.g. [18, 25, 26]). References [10, 6] address such symmetry problem for general compact groups, and generalized Wigner 3jm-symbols are defined. Here, we follow the conventions in [22].

**Definition 2.25.** *The Wigner coupling symbol is the coefficient denoted by the round brackets below:*

$$(2.132) \quad \left( \begin{array}{ccc} \mathbf{p}_1 & \mathbf{p}_2 & (\mathbf{a}; \sigma) \\ \nu_1, I_1 & \nu_2, I_2 & \nu, I \end{array} \right) = \frac{(-1)^{|\mathbf{a}|+2(t_\nu+u_\nu)}}{\sqrt{\dim Q(\check{\mathbf{a}})}} C_{\nu_1 I_1, \nu_2 I_2, \check{\nu} I}^{\mathbf{p}_1, \mathbf{p}_2, (\check{\mathbf{a}}; \sigma)}.$$

Thus, from Theorem 2.23, we have

$$(2.133) \quad \begin{aligned} \left( \begin{array}{ccc} \mathbf{p}_1 & \mathbf{p}_2 & (\mathbf{a}; \sigma) \\ \nu_1, I_1 & \nu_2, I_2 & \nu, I \end{array} \right) &= (-1)^{|\mathbf{p}_1|+|\mathbf{p}_2|+|\mathbf{a}|} \left( \begin{array}{ccc} \mathbf{p}_2 & \mathbf{p}_1 & (\mathbf{a}; \check{\sigma}) \\ \nu_2, I_2 & \nu_1, I_1 & \nu, I \end{array} \right) \\ &= (-1)^{|\mathbf{p}_1|+|\mathbf{p}_2|+|\mathbf{a}|} \left( \begin{array}{ccc} \mathbf{p}_1 & \mathbf{a} & (\mathbf{p}_2; \check{\sigma}) \\ \nu_1, I_1 & \nu, I & \nu_2, I_2 \end{array} \right) \\ &= (-1)^{|\mathbf{p}_1|+|\mathbf{p}_2|+|\mathbf{a}|} \left( \begin{array}{ccc} \check{\mathbf{p}}_1 & \check{\mathbf{p}}_2 & (\check{\mathbf{a}}; \check{\sigma}) \\ \check{\nu}_1, I_1 & \check{\nu}_2, I_2 & \check{\nu}, I \end{array} \right). \end{aligned}$$

Another useful property of Wigner coupling symbols arises when we study the recoupling problem of a triple tensor product  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2) \otimes Q(\mathbf{p}_3)$ , whose decomposition can be realized in any order. Consider, for example, the Clebsch-Gordan series

$$(2.134) \quad Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2) = \bigoplus_{(\mathbf{a}_{12}; \sigma_{12})} Q(\mathbf{a}_{12}; \sigma_{12}), \quad Q(\mathbf{a}_{12}) \otimes Q(\mathbf{p}_3) = \bigoplus_{(\mathbf{a}; \sigma)} Q(\mathbf{a}; \sigma).$$

With this, we get a basis  $\{e((\mathbf{a}_{12}; \sigma_{12}), (\mathbf{a}; \sigma); \boldsymbol{\mu}, J)\}$  for  $Q(\mathbf{p}_1) \otimes Q(\mathbf{p}_2) \otimes Q(\mathbf{p}_3)$  satisfying

$$(2.135) \quad \begin{aligned} e((\mathbf{a}_{12}; \sigma_{12}), (\mathbf{a}; \sigma); \boldsymbol{\mu}, J) &= \sum_{\substack{\mu_{12}, J_{12} \\ \nu_3, I_3}} \sum_{\substack{\nu_1, I_1 \\ \nu_2, I_2}} C_{\mu_{12} J_{12}, \nu_3 I_3, \boldsymbol{\mu} J}^{\mathbf{a}_{12}, \mathbf{p}_3, (\mathbf{a}; \sigma)} C_{\nu_1 I_1, \nu_2 I_2, \mu_{12} J_{12}}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a}_{12}; \sigma_{12})} \\ &\quad \times e(\mathbf{p}_1; \nu_1, I_1) \otimes e(\mathbf{p}_2; \nu_2, I_2) \otimes e(\mathbf{p}_3; \nu_3, I_3) \\ &= \sqrt{\dim Q(\mathbf{a}) \dim Q(\mathbf{a}_{12})} \sum_{\substack{\mu_{12}, J_{12} \\ \nu_3, I_3}} \sum_{\substack{\nu_1, I_1 \\ \nu_2, I_2}} (-1)^{|\mathbf{a}|+|\mathbf{a}_{12}|+2(t_\mu+u_\mu+t_{\mu_{12}}+u_{\mu_{12}})} \\ &\quad \times \left( \begin{array}{ccc} \mathbf{a}_{12} & \mathbf{p}_3 & (\check{\mathbf{a}}; \sigma) \\ \mu_{12}, J_{12} & \nu_3, I_3 & \boldsymbol{\mu}, J \end{array} \right) \left( \begin{array}{ccc} \mathbf{p}_1 & \mathbf{p}_2 & (\check{\mathbf{a}}_{12}; \sigma_{12}) \\ \nu_1, I_1 & \nu_2, I_2 & \mu_{12}, J_{12} \end{array} \right) \\ &\quad \times e(\mathbf{p}_1; \nu_1, I_1) \otimes e(\mathbf{p}_2; \nu_2, I_2) \otimes e(\mathbf{p}_3; \nu_3, I_3). \end{aligned}$$

On the other hand, if we use

$$(2.136) \quad Q(\mathbf{p}_2) \otimes Q(\mathbf{p}_3) = \bigoplus_{(\mathbf{a}_{23}; \sigma_{23})} Q(\mathbf{a}_{23}; \sigma_{23}), \quad Q(\mathbf{a}_{23}) \otimes Q(\mathbf{p}_1) = \bigoplus_{(\mathbf{a}'; \sigma')} Q(\mathbf{a}'; \sigma'),$$

we get a basis  $\{e((\mathbf{a}_{23}; \sigma_{23}), (\mathbf{a}'; \sigma'); \boldsymbol{\mu}', J')\}$  satisfying  
(2.137)

$$\begin{aligned} e((\mathbf{a}_{23}; \sigma_{23}), (\mathbf{a}'; \sigma'); \boldsymbol{\mu}', J') &= \sum_{\substack{\boldsymbol{\mu}_{23}, J_{23} \\ \boldsymbol{\nu}_1, I_1}} \sum_{\substack{\boldsymbol{\nu}_2, I_2 \\ \boldsymbol{\nu}_3, I_3}} C_{\boldsymbol{\mu}_{23}, J_{23}, \boldsymbol{\nu}_1, I_1, \boldsymbol{\mu}', J'}^{\mathbf{a}_{23}, \mathbf{p}_1, (\mathbf{a}'; \sigma')} C_{\boldsymbol{\nu}_2, I_2, \boldsymbol{\nu}_3, I_3, \boldsymbol{\mu}_{23}, J_{23}}^{\mathbf{p}_2, \mathbf{p}_3, (\mathbf{a}_{23}; \sigma_{23})} \\ &\quad \times e(\mathbf{p}_1; \boldsymbol{\nu}_1, I_1) \otimes e(\mathbf{p}_2; \boldsymbol{\nu}_2, I_2) \otimes e(\mathbf{p}_3; \boldsymbol{\nu}_3, I_3) \\ &= \sqrt{\dim Q(\mathbf{a}') \dim Q(\mathbf{a}_{23})} \sum_{\substack{\boldsymbol{\mu}_{23}, J_{23} \\ \boldsymbol{\nu}_1, I_1}} \sum_{\substack{\boldsymbol{\nu}_2, I_2 \\ \boldsymbol{\nu}_3, I_3}} (-1)^{|\mathbf{a}'| + |\mathbf{a}_{23}| + 2(t_{\boldsymbol{\mu}'} + u_{\boldsymbol{\mu}'} + t_{\boldsymbol{\mu}_{23}} + u_{\boldsymbol{\mu}_{23}})} \\ &\quad \times \begin{pmatrix} \mathbf{a}_{23} & \mathbf{p}_1 & (\check{\mathbf{a}}'; \sigma') \\ \boldsymbol{\mu}_{23}, J_{23} & \boldsymbol{\nu}_1, I_1 & \check{\boldsymbol{\mu}}', J' \end{pmatrix} \begin{pmatrix} \mathbf{p}_2 & \mathbf{p}_3 & (\check{\mathbf{a}}_{23}; \sigma_{23}) \\ \boldsymbol{\nu}_2, I_2 & \boldsymbol{\nu}_3, I_3 & \check{\boldsymbol{\mu}}_{23}, J_{23} \end{pmatrix} \\ &\quad \times e(\mathbf{p}_1; \boldsymbol{\nu}_1, I_1) \otimes e(\mathbf{p}_2; \boldsymbol{\nu}_2, I_2) \otimes e(\mathbf{p}_3; \boldsymbol{\nu}_3, I_3). \end{aligned}$$

Of course, there is an unitary transformation relating these basis and  
(2.138)

$$\langle e((\mathbf{a}_{12}; \sigma_{12}), (\mathbf{a}; \sigma); \boldsymbol{\mu}, J) | e((\mathbf{a}_{23}; \sigma_{23}), (\mathbf{a}'; \sigma'); \boldsymbol{\mu}', J') \rangle \neq 0 \implies \begin{cases} \mathbf{a} = \mathbf{a}' \\ (\boldsymbol{\mu}, J) = (\boldsymbol{\mu}', J') \end{cases}$$

Also, the coefficients of the form

$$(2.139) \quad \langle e((\mathbf{a}_{12}; \sigma_{12}), (\mathbf{a}; \sigma); \boldsymbol{\mu}, J) | e((\mathbf{a}_{23}; \sigma_{23}), (\mathbf{a}; \sigma'); \boldsymbol{\mu}, J) \rangle$$

do not depend on the weight since the vectors in the inner product can be generated from the highest weight vectors of their respective representations by applying the ladder operators and we can write a highest weight vector of one basis as a linear combination of highest weight vectors of the other basis for equivalent representations.

**Definition 2.26.** *The Wigner recoupling symbol is the coefficient denoted by the curly brackets below:*

$$\begin{aligned} (2.140) \quad &\left\{ \begin{pmatrix} \mathbf{p}_1 & \mathbf{p}_2 & (\check{\mathbf{a}}_{12}; \sigma_{12}) \\ \boldsymbol{\mu}_{12}, J_{12} & \boldsymbol{\nu}_3, I_3 & \boldsymbol{\mu}, J \end{pmatrix} \right\} = \frac{(-1)^{|\mathbf{a}_{23}| + |\mathbf{p}_2| + |\mathbf{p}_3|}}{\sqrt{\dim Q(\mathbf{a}_{12}) \dim Q(\mathbf{a}_{23})}} \sum_{\substack{\boldsymbol{\nu}_1, I_1 \\ \boldsymbol{\nu}_2, I_2 \\ \boldsymbol{\nu}_3, I_3}} \sum_{\substack{\boldsymbol{\mu}_{12}, J_{12} \\ \boldsymbol{\mu}_{23}, J_{23}}} C_{\boldsymbol{\mu}_{12}, J_{12}, \boldsymbol{\nu}_3, I_3, \boldsymbol{\mu}, J}^{\mathbf{a}_{12}, \mathbf{p}_3, (\mathbf{a}; \sigma)} \\ &\quad \times C_{\boldsymbol{\nu}_1, I_1, \boldsymbol{\nu}_2, I_2, \boldsymbol{\mu}_{12}, J_{12}}^{\mathbf{p}_1, \mathbf{p}_2, (\mathbf{a}_{12}; \sigma_{12})} C_{\boldsymbol{\mu}_{23}, J_{23}, \boldsymbol{\nu}_1, I_1, \boldsymbol{\mu}, J}^{\mathbf{a}_{23}, \mathbf{p}_1, (\mathbf{a}; \sigma')} C_{\boldsymbol{\nu}_2, I_2, \boldsymbol{\nu}_3, I_3, \boldsymbol{\mu}_{23}, J_{23}}^{\mathbf{p}_2, \mathbf{p}_3, (\mathbf{a}_{23}; \sigma_{23})} \\ &= \sum_{\substack{\boldsymbol{\nu}_1, I_1 \\ \boldsymbol{\nu}_2, I_2 \\ \boldsymbol{\nu}_3, I_3}} \sum_{\substack{\boldsymbol{\mu}_{12}, J_{12} \\ \boldsymbol{\mu}_{23}, J_{23} \\ \boldsymbol{\mu}, J}} (-1)^{|\mathbf{p}_2| + |\mathbf{p}_3| + |\mathbf{a}_{12}| + 2(t_{\boldsymbol{\mu}_{12}} + u_{\boldsymbol{\mu}_{12}} + t_{\boldsymbol{\mu}_{23}} + u_{\boldsymbol{\mu}_{23}})} \\ &\quad \times \begin{pmatrix} \mathbf{a}_{12} & \mathbf{p}_3 & (\check{\mathbf{a}}; \sigma) \\ \boldsymbol{\mu}_{12}, J_{12} & \boldsymbol{\nu}_3, I_3 & \check{\boldsymbol{\mu}}, J \end{pmatrix} \begin{pmatrix} \mathbf{p}_1 & \mathbf{p}_2 & (\check{\mathbf{a}}_{12}; \sigma_{12}) \\ \boldsymbol{\nu}_1, I_1 & \boldsymbol{\nu}_2, I_2 & \check{\boldsymbol{\mu}}_{12}, J_{12} \end{pmatrix} \\ &\quad \times \begin{pmatrix} \mathbf{a}_{23} & \mathbf{p}_1 & (\check{\mathbf{a}}; \sigma') \\ \boldsymbol{\mu}_{23}, J_{23} & \boldsymbol{\nu}_1, I_1 & \check{\boldsymbol{\mu}}, J \end{pmatrix} \begin{pmatrix} \mathbf{p}_2 & \mathbf{p}_3 & (\check{\mathbf{a}}_{23}; \sigma_{23}) \\ \boldsymbol{\nu}_2, I_2 & \boldsymbol{\nu}_3, I_3 & \check{\boldsymbol{\mu}}_{23}, J_{23} \end{pmatrix}. \end{aligned}$$

With this definition, we have the following equation:

$$(2.141) \quad e((\mathbf{a}_{23}; \sigma_{23}), (\mathbf{a}; \check{\sigma}'); \boldsymbol{\mu}, J) = \sum_{(\mathbf{a}_{12}, \sigma_{12})} (-1)^{|\mathbf{a}_{23}|+|\mathbf{p}_2|+|\mathbf{p}_3|} \sqrt{\dim Q(\mathbf{a}_{12}) \dim Q(\mathbf{a}_{23})} \\ \times \left\{ \begin{array}{ccc} \mathbf{p}_1 & \mathbf{p}_2 & (\check{\mathbf{a}}_{12}; \sigma_{12}) \\ \mathbf{p}_3 & (\mathbf{a}; \sigma, \sigma') & (\mathbf{a}_{23}; \sigma_{23}) \end{array} \right\} e((\mathbf{a}_{12}; \sigma_{12}), (\mathbf{a}; \sigma); \boldsymbol{\mu}, J) .$$

Then, by (2.135) and (2.137), we have *Wigner's identity*<sup>10</sup>:

$$(2.142) \quad \sum_{\boldsymbol{\mu}_{23}, J_{23}} (-1)^{2(t_{\nu_1}+u_{\nu_1})} \left( \begin{array}{ccc} \mathbf{a}_{23} & \mathbf{p}_1 & (\check{\mathbf{a}}; \check{\sigma}') \\ \boldsymbol{\mu}_{23}, J_{23} & \boldsymbol{\nu}_1, I_1 & \check{\boldsymbol{\mu}}, J \end{array} \right) \left( \begin{array}{ccc} \mathbf{p}_2 & \mathbf{p}_3 & (\check{\mathbf{a}}_{23}; \sigma_{23}) \\ \boldsymbol{\nu}_2, I_2 & \boldsymbol{\nu}_3, I_3 & \check{\boldsymbol{\mu}}_{23}, J_{23} \end{array} \right) \\ = \sum_{\boldsymbol{\mu}_{12}, J_{12}} \sum_{(\mathbf{a}_{12}; \sigma_{12})} (-1)^{|\mathbf{a}_{12}|+|\mathbf{p}_2|+|\mathbf{p}_3|+2(t_{\nu_3}+u_{\nu_3})} \dim Q(\mathbf{a}_{12}) \\ \times \left\{ \begin{array}{ccc} \mathbf{p}_1 & \mathbf{p}_2 & (\check{\mathbf{a}}_{12}; \sigma_{12}) \\ \mathbf{p}_3 & (\mathbf{a}; \sigma, \sigma') & (\mathbf{a}_{23}; \sigma_{23}) \end{array} \right\} \left( \begin{array}{ccc} \mathbf{a}_{12} & \mathbf{p}_3 & (\check{\mathbf{a}}; \sigma) \\ \boldsymbol{\mu}_{12}, J_{12} & \boldsymbol{\nu}_3, I_3 & \check{\boldsymbol{\mu}}, J \end{array} \right) \\ \times \left( \begin{array}{ccc} \mathbf{p}_1 & \mathbf{p}_2 & (\check{\mathbf{a}}_{12}; \sigma_{12}) \\ \boldsymbol{\nu}_1, I_1 & \boldsymbol{\nu}_2, I_2 & \check{\boldsymbol{\mu}}_{12}, J_{12} \end{array} \right) .$$

And using (2.133), we obtain

$$(2.143) \quad \left\{ \begin{array}{ccc} \mathbf{p}_1 & \mathbf{p}_2 & (\check{\mathbf{a}}_{12}; \sigma_{12}) \\ \mathbf{p}_3 & (\mathbf{a}; \sigma, \sigma') & (\mathbf{a}_{23}; \sigma_{23}) \end{array} \right\} = \left\{ \begin{array}{ccc} \check{\mathbf{p}}_2 & \check{\mathbf{p}}_1 & (\mathbf{a}_{12}; \sigma_{12}) \\ \mathbf{a} & (\mathbf{p}_3; \sigma, \sigma_{23}) & (\mathbf{a}_{23}; \sigma') \end{array} \right\} \\ = \left\{ \begin{array}{ccc} \check{\mathbf{p}}_1 & \mathbf{a}_{12} & (\check{\mathbf{p}}_2; \sigma_{12}) \\ \mathbf{p}_3 & (\mathbf{a}_{23}; \sigma_{23}, \sigma') & (\mathbf{a}; \sigma) \end{array} \right\} \\ = \left\{ \begin{array}{ccc} \mathbf{p}_3 & \check{\mathbf{a}} & (\mathbf{a}_{12}; \sigma') \\ \mathbf{p}_1 & (\check{\mathbf{p}}_2; \sigma_{12}, \sigma_{23}) & (\check{\mathbf{a}}_{23}; \sigma) \end{array} \right\} \\ = \left\{ \begin{array}{ccc} \check{\mathbf{p}}_1 & \check{\mathbf{p}}_2 & (\mathbf{a}_{12}; \check{\sigma}_{12}) \\ \check{\mathbf{p}}_3 & (\check{\mathbf{a}}; \check{\sigma}, \check{\sigma}') & (\check{\mathbf{a}}_{23}; \check{\sigma}_{23}) \end{array} \right\} .$$

Thus, using the Wigner coupling and recoupling symbols, we get a more symmetric way of writing the coefficients  $\mathcal{M}[\mathbf{p}]_{\boldsymbol{\nu}_1 I_1, \boldsymbol{\nu}_2 I_2, \boldsymbol{\nu} I}^{(\mathbf{a}_1; \sigma_1), (\mathbf{a}_2; \sigma_2), (\mathbf{a}; \sigma)}$ , as follows.

**Proposition 2.27.** *The coefficients  $\mathcal{M}[\mathbf{p}]_{\boldsymbol{\nu}_1 I_1, \boldsymbol{\nu}_2 I_2, \boldsymbol{\nu} I}^{(\mathbf{a}_1; \sigma_1), (\mathbf{a}_2; \sigma_2), (\mathbf{a}; \sigma)}$  in (2.59)-(2.60) are given by*

$$(2.144) \quad \mathcal{M}[\mathbf{p}]_{\boldsymbol{\nu}_1 I_1, \boldsymbol{\nu}_2 I_2, \boldsymbol{\nu} I}^{(\mathbf{a}_1; \sigma_1), (\mathbf{a}_2; \sigma_2), (\mathbf{a}; \sigma)} = \sqrt{\dim Q(\mathbf{a}_1) \dim Q(\mathbf{a}_2) \dim Q(\mathbf{a})} \sum_{\sigma'} (-1)^{|\mathbf{p}|+2(t_{\nu}+u_{\nu})} \\ \times \left\{ \begin{array}{ccc} \mathbf{a}_1 & \mathbf{a}_2 & (\check{\mathbf{a}}; \sigma') \\ \mathbf{p} & (\mathbf{p}; \sigma, \sigma_1) & (\mathbf{p}; \sigma_2) \end{array} \right\} \left( \begin{array}{ccc} \mathbf{a}_1 & \mathbf{a}_2 & (\check{\mathbf{a}}; \sigma') \\ \boldsymbol{\nu}_1, I_1 & \boldsymbol{\nu}_2, I_2 & \check{\boldsymbol{\nu}}, I \end{array} \right) .$$

<sup>10</sup>An equivalent formula is deduced for  $SU(2)$  in [26] and for a general compact group in [6].

*Proof.* From (2.142), we have

$$\begin{aligned}
& \sum_{\mu_3, J_3} (-1)^{2(t_{\nu_1} + u_{\nu_1})} \begin{pmatrix} \mathbf{p} & \mathbf{a}_1 & (\check{\mathbf{p}}; \check{\sigma}_1) \\ \mu_3, J_3 & \nu_1, I_1 & \check{\mu}_2, J_2 \end{pmatrix} \begin{pmatrix} \mathbf{a}_2 & \mathbf{p} & (\check{\mathbf{p}}; \sigma_2) \\ \nu_2, I_2 & \mu_1, J_1 & \check{\mu}_3, J_3 \end{pmatrix} \\
&= \sum_{\nu', I'} \sum_{\substack{\mathbf{a}' \\ \sigma''}} (-1)^{|\mathbf{a}'| + |\mathbf{a}_2| + |\mathbf{p}| + 2(t_{\mu_1} + u_{\mu_1})} \dim Q(\mathbf{a}') \\
(2.145) \quad & \times \begin{Bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & (\check{\mathbf{a}}'; \sigma') \\ \mathbf{p} & (\mathbf{p}; \sigma'', \sigma_1) & (\mathbf{p}; \sigma_2) \end{Bmatrix} \begin{pmatrix} \mathbf{a}' & \mathbf{p} & (\check{\mathbf{p}}; \sigma'') \\ \nu', I' & \mu_1, J_1 & \check{\mu}_2, J_2 \end{pmatrix} \\
& \times \begin{pmatrix} \mathbf{a}_1 & \mathbf{a}_2 & (\check{\mathbf{a}}'; \sigma') \\ \nu_1, I_1 & \nu_2, I_2 & \check{\nu}', I' \end{pmatrix}.
\end{aligned}$$

Multiplying both sides by

$$(2.146) \quad (-1)^{2(t_{\nu} + u_{\nu} + t_{\mu_2} + u_{\mu_2})} \begin{pmatrix} \mathbf{p} & \check{\mathbf{p}} & (\mathbf{a}; \sigma) \\ \mu_1, J_1 & \check{\mu}_2, J_2 & \nu, I \end{pmatrix}$$

and performing a summation over  $\mu_1, J_1, \mu_2$  and  $J_2$ , by (2.133) and (2.48), we obtain

$$\begin{aligned}
(2.147) \quad & \sum_{\substack{\mu_1, J_1 \\ \mu_2, J_2 \\ \mu_3, J_3}} (-1)^{2(t_{\nu} + u_{\nu} + t_{\mu_3} + u_{\mu_3})} \begin{pmatrix} \mathbf{p} & \check{\mathbf{p}} & (\mathbf{a}; \sigma) \\ \mu_1 & \check{\mu}_2 & \nu, I \end{pmatrix} \begin{pmatrix} \mathbf{p} & \mathbf{a}_1 & (\check{\mathbf{p}}; \check{\sigma}_1) \\ \mu_3 & \nu_1, I_1 & \check{\mu}_2 \end{pmatrix} \\
& \times \begin{pmatrix} \mathbf{a}_2 & \mathbf{p} & (\check{\mathbf{p}}; \sigma_2) \\ \nu_2, I_2 & \mu_1 & \check{\mu}_3 \end{pmatrix} \\
& = \sum_{\sigma'} (-1)^{|\mathbf{a}| + |\mathbf{a}_2| + |\mathbf{p}|} \begin{Bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & (\check{\mathbf{a}}; \sigma') \\ \mathbf{p} & (\mathbf{p}; \sigma, \sigma_1) & (\mathbf{p}; \sigma_2) \end{Bmatrix} \begin{pmatrix} \mathbf{a}_1 & \mathbf{a}_2 & (\check{\mathbf{a}}; \sigma') \\ \nu_1, I_1 & \nu_2, I_2 & \check{\nu}, I \end{pmatrix}.
\end{aligned}$$

Again by (2.133), we get what we want.  $\square$

As a corollary, using (2.52), we get an improvement in Lemma 2.14 with the restriction  $\delta(I_1, I_2, I) = 1$  to summation (2.59). We will state this together with a more symmetric way to write the product of operators by means of the next definition:

**Definition 2.28.** *The Wigner product symbol is the coefficient denoted by the square brackets below:*

$$\begin{aligned}
(2.148) \quad & \left[ \begin{pmatrix} \mathbf{a}_1; \sigma_1 & \mathbf{a}_2; \sigma_2 & \mathbf{a}; \sigma \\ \nu_1, I_1 & \nu_2, I_2 & \nu, I \end{pmatrix} [\mathbf{p}] \right] = \sqrt{\dim Q(\mathbf{a}_1) \dim Q(\mathbf{a}_2) \dim Q(\mathbf{a})} \\
& \times \sum_{\sigma'} \begin{Bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & (\check{\mathbf{a}}; \sigma') \\ \mathbf{p} & (\mathbf{p}; \sigma, \sigma_1) & (\mathbf{p}; \sigma_2) \end{Bmatrix} \begin{pmatrix} \mathbf{a}_1 & \mathbf{a}_2 & (\check{\mathbf{a}}; \sigma') \\ \nu_1, I_1 & \nu_2, I_2 & \nu, I \end{pmatrix}.
\end{aligned}$$

From (2.52), we have

$$(2.149) \quad \left[ \begin{pmatrix} \mathbf{a}_1; \sigma_1 & \mathbf{a}_2; \sigma_2 & \mathbf{a}; \sigma \\ \nu_1, I_1 & \nu_2, I_2 & \nu, I \end{pmatrix} [\mathbf{p}] \right] \neq 0 \quad \Longrightarrow \quad \begin{cases} \nabla_{\nu_1 + \nu_2, \check{\nu}} = 1 \\ \delta(I_1, I_2, I) = 1 \end{cases}.$$

Also, symmetries (2.133) and (2.143) imply

$$\begin{aligned}
(2.150) \quad \begin{bmatrix} (\mathbf{a}_1; \sigma_1) & (\mathbf{a}_2; \sigma_2) & (\mathbf{a}_3; \sigma_3) \\ \boldsymbol{\nu}_1, I_1 & \boldsymbol{\nu}_2, I_2 & \boldsymbol{\nu}_3, I_3 \end{bmatrix} [\mathbf{p}] &= \begin{bmatrix} (\mathbf{a}_3; \sigma_3) & (\mathbf{a}_1; \sigma_1) & (\mathbf{a}_2; \sigma_2) \\ \boldsymbol{\nu}_3, I_3 & \boldsymbol{\nu}_1, I_1 & \boldsymbol{\nu}_2, I_2 \end{bmatrix} [\mathbf{p}] \\
&= \begin{bmatrix} (\mathbf{a}_2; \sigma_2) & (\mathbf{a}_3; \sigma_3) & (\mathbf{a}_1; \sigma_1) \\ \boldsymbol{\nu}_2, I_2 & \boldsymbol{\nu}_3, I_3 & \boldsymbol{\nu}_1, I_1 \end{bmatrix} [\mathbf{p}] \\
&= \begin{bmatrix} (\check{\mathbf{a}}_2; \sigma_2) & (\check{\mathbf{a}}_1; \sigma_1) & (\check{\mathbf{a}}_3; \sigma_3) \\ \check{\boldsymbol{\nu}}_2, I_2 & \check{\boldsymbol{\nu}}_1, I_1 & \check{\boldsymbol{\nu}}_3, I_3 \end{bmatrix} [\mathbf{p}] \\
&= (-1)^{\sum_{k=1}^3 |\mathbf{a}_k|} \begin{bmatrix} (\check{\mathbf{a}}_1; \check{\sigma}_1) & (\check{\mathbf{a}}_2; \check{\sigma}_2) & (\check{\mathbf{a}}_3; \check{\sigma}_3) \\ \check{\boldsymbol{\nu}}_1, I_1 & \check{\boldsymbol{\nu}}_2, I_2 & \check{\boldsymbol{\nu}}_3, I_3 \end{bmatrix} [\check{\mathbf{p}}] \\
&= (-1)^{\sum_{k=1}^3 |\mathbf{a}_k|} \begin{bmatrix} (\mathbf{a}_2; \check{\sigma}_2) & (\mathbf{a}_1; \check{\sigma}_1) & (\mathbf{a}_3; \check{\sigma}_3) \\ \boldsymbol{\nu}_2, I_2 & \boldsymbol{\nu}_1, I_1 & \boldsymbol{\nu}_3, I_3 \end{bmatrix} [\check{\mathbf{p}}] .
\end{aligned}$$

**Corollary 2.29.** *The operator product of elements of a coupled basis of  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}})$  can be decomposed into*

$$\begin{aligned}
(2.151) \quad e((\mathbf{a}_1; \sigma_1); \boldsymbol{\nu}_1, I_1) e((\mathbf{a}_2; \sigma_2); \boldsymbol{\nu}_2, I_2) &= \\
\sum_{\substack{(\mathbf{a}; \sigma) \\ \boldsymbol{\nu}, I}} (-1)^{|\mathbf{p}|+2(t_\nu+u_\nu)} \begin{bmatrix} (\mathbf{a}_1; \sigma_1) & (\mathbf{a}_2; \sigma_2) & (\mathbf{a}; \sigma) \\ \boldsymbol{\nu}_1, I_1 & \boldsymbol{\nu}_2, I_2 & \check{\boldsymbol{\nu}}, I \end{bmatrix} [\mathbf{p}] e((\mathbf{a}; \sigma); \boldsymbol{\nu}, I)
\end{aligned}$$

where summation over  $(\mathbf{a}; \sigma)$  is restricted to  $Q(\mathbf{a}; \sigma)$  in the Clebsch-Gordan series of  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}})$ , and summations over  $\boldsymbol{\nu}$  and  $I$  can be simplified using (2.149).

In particular, if we calculate the product of any  $e((\mathbf{a}; \sigma); \boldsymbol{\nu}, I)$  by the identity operator, recalling (2.126), we obtain

$$\begin{aligned}
(2.152) \quad \begin{bmatrix} (0, 0) & (\mathbf{a}; \sigma) & (\mathbf{a}'; \sigma') \\ (0, 0, 0), 0 & \boldsymbol{\nu}, I & \boldsymbol{\mu}, J \end{bmatrix} [\mathbf{p}] &= \begin{bmatrix} (\mathbf{a}; \sigma) & (0, 0) & (\mathbf{a}'; \sigma') \\ \boldsymbol{\nu}, I & (0, 0, 0), 0 & \boldsymbol{\mu}, J \end{bmatrix} [\mathbf{p}] \\
&= \frac{(-1)^{2(t_\nu+u_\nu)}}{\sqrt{\dim Q(\mathbf{p})}} \delta_{\boldsymbol{\nu}, \check{\boldsymbol{\mu}}} \delta_{I, J} \delta_{\mathbf{a}, \mathbf{a}'} \delta_{\sigma, \sigma'} .
\end{aligned}$$

**2.6. Coadjoint orbits.** Being a simple compact Lie group,  $SU(3)$  has equivalent adjoint and coadjoint actions. That is, the coadjoint and adjoint orbits are isomorphic<sup>11</sup>. We identify the root diagram of  $\mathfrak{su}(3)$  with the Cartan subalgebra generated by  $iT_3$  and  $iU_3$  by making  $\alpha_1 \equiv 2iT_3$  and  $\alpha_2 \equiv 2iU_3$ . Then, we obtain

$$(2.153) \quad w_1 \equiv \frac{i}{2} \lambda_3 + \frac{i}{2\sqrt{3}} \lambda_8 = \frac{i}{3} \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} , \quad w_2 = \frac{i}{\sqrt{3}} \lambda_8 = \frac{i}{3} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix} .$$

It is well known that each orbit intersects the closed positive Weyl chamber

$$(2.154) \quad \overline{\mathcal{C}} = \{x w_2 + y w_1 : x, y \geq 0\}$$

in precisely one single point (see e.g. [5]). Let  $\mathcal{O}_{x,y}$  be the orbit passing through

$$(2.155) \quad \xi_{x,y} = x w_2 + y w_1 \in \overline{\mathcal{C}} \setminus \{\mathbf{0}\} .$$

<sup>11</sup>A general discussion of coadjoint orbits of semisimple Lie groups can be found in [4].

From the commutation relations, it is clear that, if  $x, y > 0$ , then the isotropy subgroup of  $\xi_{x,0}$  is

$$(2.156) \quad H := \left\{ \begin{pmatrix} U & 0 \\ 0 & \det(U)^{-1} \end{pmatrix}, U \in U(2) \right\} \simeq S(U(2) \times U(1)) \simeq U(2),$$

whereas the isotropy subgroup of  $\xi_{0,y}$  is

$$(2.157) \quad H' := g_0 H g_0^{-1} = g_0 H g_0 = \left\{ \begin{pmatrix} \det(U)^{-1} & 0 \\ 0 & U \end{pmatrix}, U \in U(2) \right\} \\ \simeq S(U(1) \times U(2)) \simeq U(2),$$

for

$$(2.158) \quad g_0 = \begin{pmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{pmatrix} \in SU(3).$$

On the other hand, the isotropy subgroup of  $\xi_{x,y}$  is the maximal torus<sup>12</sup>

$$(2.159) \quad T := \left\{ \begin{pmatrix} e^{i\theta_1} & 0 & 0 \\ 0 & e^{i\theta_2} & 0 \\ 0 & 0 & e^{i\theta_3} \end{pmatrix} : \theta_1 + \theta_2 + \theta_3 = 0 \right\} \\ \simeq S(U(1) \times U(1) \times U(1)) \simeq U(1) \times U(1).$$

Therefore, we have two types of non trivial (co)adjoint orbits:

$$\mathcal{O}_{x,0} \simeq SU(3)/H \simeq SU(3)/H' \simeq \mathcal{O}_{0,y}$$

and

$$\mathcal{O}_{x,y} \simeq SU(3)/T \simeq \mathcal{O}_{y,x},$$

for  $x, y > 0$ .

For a better realization of such orbits, we recall the complex projective space  $\mathbb{C}P^2$ : the quotient of  $\mathbb{C}^3 \setminus \{\mathbf{0}\}$  by the equivalence relation

$$(2.160) \quad z \sim z' \iff z = a z', \quad a \in \mathbb{C}^*.$$

To construct  $\mathbb{C}P^2$  using this equivalence relation<sup>13</sup>, we can look only to the unitary vectors of  $\mathbb{C}^3$ , reducing our analysis to the  $SU(3)$ -homogeneous space  $S^5 = \{z \in \mathbb{C}^3 : \|z\| = 1\}$ . Since the point  $(0, 0, 1) \in S^5$  has

$$(2.161) \quad \left\{ \begin{pmatrix} U & 0 \\ 0 & 1 \end{pmatrix} : U \in SU(2) \right\} \subset SU(3)$$

as isotropy subgroup, we have<sup>14</sup>  $S^5 \simeq SU(3)/SU(2)$ . Also note that,  $\forall z \in S^5$ ,  $e^{i\theta} z \sim z$ . So  $\mathbb{C}P^2 \simeq S^5/S^1$  and the isotropy subgroup of the equivalence class  $[0 : 0 : 1] \in \mathbb{C}P^2$  is  $H$ , i.e.,  $SU(3)/H \simeq \mathbb{C}P^2$ . By similar argument we get  $U(2)/(U(1) \times U(1)) \simeq \mathbb{C}P^1$ , so  $SU(3)/T \simeq \mathcal{E}$ , where  $\mathcal{E}$  is the total space of a fiber bundle  $\pi : \mathcal{E} \rightarrow \mathbb{C}P^2$  with fiber  $\mathbb{C}P^1$ , which we denote by  $\mathcal{E}(\mathbb{C}P^2, \mathbb{C}P^1, \pi)$ , that is,

$$(2.162) \quad \mathcal{E} = \mathcal{E}(\mathbb{C}P^2, \mathbb{C}P^1, \pi) := \mathbb{C}P^1 \hookrightarrow \mathcal{E} \xrightarrow{\pi} \mathbb{C}P^2.$$

<sup>12</sup>It is a matter of simple calculation to verify that  $T' = g_0 T g_0 = T$ .

<sup>13</sup>This construction is presented in [2].

<sup>14</sup> $SU(2) \simeq S^3$  hence  $SU(3)$  is a 3-sphere bundle over  $S^5$ .

Thus,

$$(2.163) \quad \mathcal{O}_{x,y} \simeq \begin{cases} \mathbb{C}P^2, & \text{if } x = 0 \text{ or } y = 0 \\ \mathcal{E}, & \text{if } x, y > 0 \end{cases} .$$

The orbits  $\mathcal{O}_{x,y}$  and  $\mathcal{O}_{y,x}$  are related by the involution  $\iota = -id$  on  $\mathfrak{su}(3)$ . Indeed,  $\iota \circ Ad_g = Ad_g \circ \iota$  trivially holds for every  $g \in SU(3)$  and

$$(2.164) \quad \iota(x w_2 + y w_1) = -x w_2 - y w_1 = Ad_{g_0}(y w_2 + x w_1) ,$$

so  $\iota(\mathcal{O}_{x,y}) = \mathcal{O}_{y,x}$ . Thus,  $\iota$  is an involution on  $\mathcal{O}_{x,x}$ .

Let  $\mathbf{x}_0 = [0 : 0 : 1] \in \mathbb{C}P^2$ , whose isotropy subgroup is  $H$ , so that the isotropy subgroup of  $g_0 \mathbf{x}_0 = [1 : 0 : 0]$  is  $H'$ , and let  $\mathbf{z}_0 \in \pi^{-1}(\mathbf{x}_0) \subset \mathcal{E}$  be a point with  $T$  as isotropy subgroup. Then consider the equivariant diffeomorphisms

$$(2.165) \quad \begin{aligned} \psi_{x,0} : \mathcal{O}_{x,0} &\rightarrow \mathbb{C}P^2 : Ad_g \xi_{x,0} \mapsto g \mathbf{x}_0 , \\ \psi_{0,y} : \mathcal{O}_{0,y} &\rightarrow \mathbb{C}P^2 : Ad_g \xi_{0,y} \mapsto g g_0 \mathbf{x}_0 , \\ \psi_{x,y} : \mathcal{O}_{x,y} &\rightarrow \mathcal{E} : Ad_g \xi_{x,y} \mapsto g \mathbf{z}_0 , \end{aligned}$$

for  $x, y > 0$  still holding. Therefore,

$$(2.166) \quad \psi_{x,0} \circ \iota \circ \psi_{0,x}^{-1} : \mathbb{C}P^2 \rightarrow \mathbb{C}P^2$$

is the identity map, and

$$(2.167) \quad \alpha := \psi_{x,y} \circ \iota \circ \psi_{y,x}^{-1} : \mathcal{E} \rightarrow \mathcal{E}$$

is an  $SU(3)$ -equivariant involution. Of course, there is an equivalent involution on each  $\mathcal{O}_{x,y} \simeq \mathcal{E}$ , namely

$$(2.168) \quad \alpha_{x,y} := \psi_{x,y}^{-1} \circ \psi_{y,x} \circ \iota : \mathcal{O}_{x,y} \rightarrow \mathcal{O}_{x,y} ,$$

which reduces to  $\iota$  for  $x = y$ .

Every (co)adjoint orbit of  $SU(3)$  is a symplectic manifold, that is, every  $\mathcal{O}_{x,y}$  carries a  $SU(3)$ -invariant symplectic form, the so-called Kirillov-Kostant-Souriau form<sup>15</sup>, cf. eg. [14, Ch. I, sec. 2]). Furthermore, the  $SU(3)$ -invariant symplectic form on  $\mathcal{O}_{x,y}$  induces a normalized left invariant integral on the orbit  $\mathcal{O}_{x,y}$  such that any other left invariant integral differs from it by a scalar factor. Then, we can fix this factor for  $\mathcal{O}_{x,y}$  so that the lift  $\tilde{f} \in C(SU(3))$  of any  $f \in C(\mathcal{O}_{x,y})$  satisfies

$$(2.169) \quad \int_{SU(3)} \tilde{f}(g) dg = \int_{\mathcal{O}_{x,y}} f(x) dx$$

for the Haar integral on  $SU(3)$  (see [11, (2.49) Theorem]). With no danger of confusion, we may denote the  $SU(3)$ -invariant inner product in  $L^2(\mathcal{O}_{x,y})$  with respect to such integral simply by  $\langle | \rangle$ , that is, for any  $f_1, f_2 \in L^2(\mathcal{O}_{x,y})$ ,

$$(2.170) \quad \langle f_1 | f_2 \rangle = \int_{\mathcal{O}_{x,y}} \overline{f_1}(x) f_2(x) dx .$$

Throughout this text, we consider  $\mathbb{C}P^2$  and  $\mathcal{E}$  equipped with the aforementioned symplectic forms and normalized left invariant integrals.

In what follows, we shall identify as a classical quark system, the Poisson algebra of smooth functions on  $\mathcal{O}$ . When  $\mathcal{O} = \mathbb{C}P^2$ , we shall refer to a *classical pure-quark system*. When  $\mathcal{O} = \mathcal{E}$ , we shall refer to a *generic classical quark system*.

<sup>15</sup>Actually first identified by Sophus Lie.

### 3. PURE-QUARK SYSTEMS

We start to properly attack our main problem by focusing on the simpler possible phase space for a classical quark system:  $\mathbb{C}P^2$ . First, we describe the set of harmonic functions on  $\mathbb{C}P^2$ , which imposes a restriction on the classes of irreducible representations of  $SU(3)$  with possible correspondences to smooth functions on  $\mathbb{C}P^2$ . Then, we proceed to describe the relevant  $SU(3)$ -representations for this case, as quantum quark systems. To finish, we work out the characterization of all correspondences from such quantum quark systems to the classical quark system of interest and describe the induced twisted products of symbols. The construction and characterization of symbol correspondences in this section is very close to what is done for spin systems in [18]. Accordingly, proofs of some propositions are identical to the  $SU(2)$  case. The quantum and classical systems in correspondence, in this chapter, are both called “pure-quark system” and this name is explained right after Definition 3.7.

#### 3.1. Classical pure-quark system.

**Definition 3.1.** *The classical pure-quark system consists of  $\mathbb{C}P^2$  equipped with its  $SU(3)$ -invariant symplectic form, together with its Poisson algebra on  $C_c^\infty(\mathbb{C}P^2)$ .*

Since  $\mathbb{C}P^2 \simeq SU(3)/H$ , where  $H \simeq U(2)$ , cf. (2.156), we look for representations  $Q(p, q)$  with weights satisfying  $t = u = I = 0$  (cf. (2.21)-(2.22)) to determine the harmonic functions on  $\mathbb{C}P^2$ .

**Proposition 3.2.** *The representations of  $SU(3)$  with non null vectors fixed by  $H \simeq U(2)$  are the representations  $Q(n, n)$ . The space fixed by  $H$  is spanned by  $e((n, n); (n, n, n), 0)$ .*

*Proof.* From  $t = u = 0$ , we get  $\nu_1 = \nu_2 = \nu_3 = \nu$ . From the constraints (2.21), we get  $r_+ = r_-$ , which implies  $r_+ = q = \nu = r_-$  that, in turn, implies  $\nu = p$ . Thus, putting  $n = p = q$ , we finish the proof.  $\square$

**Definition 3.3.** *The  $\mathbb{C}P^2$  harmonics are the functions  $X_{\nu, I}^n : \mathbb{C}P^2 \rightarrow \mathbb{C}$ , such that*

$$(3.1) \quad X_{\nu, I}^n(g\mathbf{x}_0) = (n+1)^{3/2} \overline{D_{\nu I, (n, n, n)0}^{(n, n)}}(g) ,$$

for  $\mathbf{x}_0 = [0 : 0 : 1] \in \mathbb{C}P^2$ ,  $g \in SU(3)$  and  $D_{\nu I, (n, n, n)0}^{(n, n)}$  a Wigner  $D$ -function as in Definition 2.5.

The factor  $(n+1)^{3/2}$  in the definition of  $\mathbb{C}P^2$  harmonics is the square root of the dimension of the representation  $Q(n, n)$  and is used to ensure normalization according to Schur’s Orthogonality Relations, so that we have

$$(3.2) \quad \langle X_{\nu, I}^n | X_{\mu, J}^m \rangle = \delta_{n, m} \delta_{\nu, \mu} \delta_{I, J}$$

with respect to the inner product described in section 2.6, cf. (2.169)-(2.170).

We note that

$$(3.3) \quad X_{(0,0,0),0}^0 \equiv 1$$

and, cf. (2.38),

$$(3.4) \quad \overline{X_{\nu, I}^n} = (-1)^{2(t+u)} X_{\tilde{\nu}, \tilde{I}}^n , \text{ for } \Delta_{\nu, \tilde{\nu}}^{2n} = 1 .$$

**Remark 3.4.** Fixed  $x > 0$ , the diffeomorphism  $\psi_{x,0}$  can be used to carry  $\mathbb{C}P^2$  harmonics to  $\mathcal{O}_{x,0}$  by means of the composition  $X_{\nu,I}^n \circ \psi_{x,0}$ , cf. (2.165). Equivalently,  $X_{\nu,I}^n \circ \psi_{0,x}$  are the  $\mathbb{C}P^2$  harmonics carried to the orbit  $\mathcal{O}_{0,x}$ . Consequently, we have a set of harmonic functions on  $\mathcal{O}_{x,0}$  related to a set of harmonic functions on  $\mathcal{O}_{0,x}$  by the map  $\iota$ , cf. (2.166).

Next, we derive the decomposition of the pointwise product of two  $\mathbb{C}P^2$  harmonics from Lemma 2.13.

**Theorem 3.5.** *The decomposition of the pointwise product of  $\mathbb{C}P^2$  harmonics is given by*

$$(3.5) \quad X_{\nu_1, I_1}^{n_1} X_{\nu_2, I_2}^{n_2} = \sum_{\substack{(n,n;\sigma) \\ \nu, I}} \left( \frac{(n_1+1)(n_2+1)}{n+1} \right)^{3/2} C_{\nu_1 I_1, \nu_2 I_2, \nu I}^{(n_1, n_1), (n_2, n_2), (n, n; \sigma)} \\ \times C_{\mathbf{n}_1 0, \mathbf{n}_2 0, \mathbf{n} 0}^{(n_1, n_1), (n_2, n_2), (n, n; \sigma)} X_{\nu, I}^n,$$

where  $\mathbf{n}_j = (n_j, n_j, n_j)$  and  $\mathbf{n} = (n, n, n)$ , and summation is restricted to  $\nabla_{\nu_1 + \nu_2, \nu} = 1$ ,  $\delta(I_1, I_2, I) = 1$  and  $Q(n, n; \sigma)$  in the Clebsch-Gordan series of  $Q(n_1, n_1) \otimes Q(n_2, n_2)$ ; in particular,  $|n_1 - n_2| \leq n \leq n_1 + n_2$ .

*Proof.* With a little abuse of notation, we write

$$(3.6) \quad X_{\nu_j, I_j}^{n_j} = (n_j + 1)^{3/2} \overline{D_{\nu_j I_j, \mathbf{n}_j 0}^{(n_j, n_j)}},$$

and apply Lemma 2.13 to get

$$(3.7) \quad X_{\nu_1, I_1}^{n_1} X_{\nu_2, I_2}^{n_2} = \sum_{(\mathbf{a}; \sigma)} \sum_{\substack{\nu, I \\ \mu, 0}} ((n_1+1)(n_2+1))^{3/2} C_{\nu_1 I_1, \nu_2 I_2, \nu I}^{(n_1, n_1), (n_2, n_2), (\mathbf{a}; \sigma)} \\ \times C_{\mathbf{n}_1 0, \mathbf{n}_2 0, \mu 0}^{(n_1, n_1), (n_2, n_2), (\mathbf{a}; \sigma)} \overline{D_{\nu I, \mu 0}^{\mathbf{a}}},$$

where  $\nabla_{\nu_1 + \nu_2, \nu} = \nabla_{\mathbf{n}_1 + \mathbf{n}_2, \mu} = 1$  and  $\delta(I_1, I_2, I) = 1$ , so  $\mu = (\mu, \mu, \mu)$ . But  $e(\mathbf{a}; (\mu, \mu, \mu), 0)$  only exists if  $\mathbf{a} = (\mu, \mu)$ . Thus, we set  $\mathbf{a} = (n, n)$  and  $\mu = \mathbf{n} = (n, n, n)$ . The restriction over  $n$  follows from Theorem 2.6.  $\square$

**Remark 3.6.** *The fact that  $\mathbb{C}P^2$  is a symplectic manifold plays no part in the decomposition of the pointwise product of  $\mathbb{C}P^2$  harmonics. Accordingly, the next step in the study of the classical pure-quark system amounts to decomposing the Poisson bracket of  $\mathbb{C}P^2$  harmonics. However, this problem is considerably harder and is deferred to a later study.*

**3.2. Quantum pure-quark systems.** From Proposition 3.2, we look for representations  $Q(p, q)$  such that the tensor product  $Q(p, q) \otimes Q(q, p)$  splits into representations of the form  $Q(n, n)$ , without multiplicities. From Corollary 2.7, we have that  $Q(p, 0)$  and  $Q(0, p)$  are the only ones that satisfy these requirements. These are special cases of quantum quark systems, for which we make the following definition:

**Definition 3.7.** Let<sup>16</sup>  $\mathbf{p} \in (\mathbb{N} \times \{0\}) \cup (\{0\} \times \mathbb{N})$  with  $|\mathbf{p}| = p$ . A quantum pure-quark system is a complex Hilbert space  $\mathcal{H}_{\mathbf{p}} \simeq \mathbb{C}^d$ , where

$$d = \frac{(p+1)(p+2)}{2},$$

<sup>16</sup>We are ignoring the trivial representation  $Q(0, 0)$ .

with an irreducible unitary  $SU(3)$ -representation of class  $Q(\mathbf{p})$  together with its operator algebra  $\mathcal{B}(\mathcal{H}_{\mathbf{p}})$ .

In the context of quark systems, such spaces arise in description of systems of  $p$  identical quarks only (or  $p$  identical antiquarks only), as we shall see below. We call them pure-quark systems because the number of antiquarks (or quarks) is zero for such systems.

An explicit way to construct such representations is by the so-called tensor method. Consider the basic representation  $\rho_1$  of class  $Q(1, 0)$  on  $\mathcal{H}_{1,0} \simeq \mathbb{C}^3$ , and let the canonical basis  $\{e_1, e_2, e_3\}$  matches a GT basis, where each vector correspond to the respective weight on the diagram of Figure 3.




FIGURE 3. Each  $e_k$  corresponds to the  $k$ -th weight.

Then, the tensor product space  $\mathcal{H} = \mathcal{H}_{1,0} \otimes \dots \otimes \mathcal{H}_{1,0}$  with  $p$  copies of  $\mathcal{H}_{1,0}$  carries a  $SU(3)$ -representation  $\rho$  given by

$$(3.8) \quad \rho(g) w_1 \otimes \dots \otimes w_p = (\rho_1(g)w_1) \otimes \dots \otimes (\rho_1(g)w_p) .$$

Now, let  $\mathcal{H}_{p,0} = \text{Sym}^p(\mathcal{H}_{1,0}) \subset \mathcal{H}$  be the subspace of totally symmetric tensors, that is,

$$(3.9) \quad \sum_{i_1, \dots, i_p=1}^3 c_{i_1, \dots, i_p} e_{i_1} \otimes \dots \otimes e_{i_p} \in \mathcal{H}_p$$

if and only if  $c_{i_{f(1)}, \dots, i_{f(p)}} = c_{i_1, \dots, i_p}$  for every permutation  $f \in S_p$ . It is immediate that  $\mathcal{H}_{p,0}$  is an invariant subspace.

We can get a basis for  $\mathcal{H}_{p,0}$  by means of symmetrization. For  $e_{i_1} \otimes \dots \otimes e_{i_p} \in \mathcal{H}$ , let  $j$ ,  $k$  and  $l$  be the numbers of occurrence of index 1, 2 and 3, respectively, and take

$$(3.10) \quad e_{j,k,l} = \binom{p}{j, k, l}^{-1/2} \sum_{f \in S_p} e_{i_{f(1)}} \otimes \dots \otimes e_{i_{f(p)}} \in \mathcal{H}_{p,0} ,$$

where

$$(3.11) \quad \binom{p}{j, k, l} = \frac{p!}{j! k! l!}$$

is the respective coefficient of the trinomial expansion of order  $p$ . The set  $\{e_{j,k,l} : j+k+l = p\}$  is an orthonormal basis of  $\mathcal{H}_{p,0}$  considering the inner product induced

by  $\mathcal{H}_{1,0}$  on  $\mathcal{H}$ . Starting with the element  $e_{p,0,0}$ , we can obtain the basis  $\{e_{j,k,l} : j+k+l=p\}$  by recursively applying the ladder operators  $T_-$  and  $U_-$  and normalizing the result. As can be seen from the diagram above,  $e_{j,k,l} = \mu_{j,k,l}(U_-)^l(T_-)^{k+l}e_{p,0,0}$ , where  $\mu_{j,k,l} > 0$ . Since  $\dim \mathcal{H}_{p,0} = (p+1)(p+2)/2$  and  $e_{p,0,0}$  is a highest weight vector<sup>17</sup> with eigenvalues  $p/2$  for  $T_3$  and 0 for  $U_3$ , we conclude that the  $SU(3)$ -representation on  $\mathcal{H}_{p,0}$  is an irreducible representation of class  $Q(p,0)$ . In particular, the map

$$(3.12) \quad \mathcal{H}_1 \rightarrow \mathcal{H}_{p,0} : w = (z_1, z_2, z_3) \mapsto w \otimes \dots \otimes w = \sum_{j+k+l=1} \sqrt{\binom{p}{j, k, l}} z_1^j z_2^k z_3^l e_{j,k,l}$$

is equivariant.

An equivalent procedure starting with  $\mathcal{H}_{0,1} = \mathcal{H}_{1,0}^*$  gives us the space  $\mathcal{H}_{0,p} = \mathcal{H}_{p,0}^*$  with a representation of class  $Q(0,p)$  and the equivariant map

$$(3.13) \quad \mathcal{H}_{0,1} \rightarrow \mathcal{H}_{0,p} : w^* = (z_1, z_2, z_3) \mapsto w^* \otimes \dots \otimes w^* = \sum_{j+k+l=1} \sqrt{\binom{p}{j, k, l}} z_1^j z_2^k z_3^l \check{e}_{j,k,l},$$

where  $\{\check{e}_{j,k,l} : j+k+l=p\}$  is the basis induced by<sup>18</sup>  $\{\check{e}_1 = -e_3^*, \check{e}_2 = e_2^*, \check{e}_3 = -e_1^*\}$  in the same way that  $\{e_{j,k,l} : j+k+l=p\}$  is induced by  $\{e_1, e_2, e_3\}$ , cf. Definition 2.3.

In Physics, the space of colors (resp. flavors) of a quark is precisely the representation  $Q(1,0)$ , with  $e_1 \equiv \text{red}$  (resp. *up quark*),  $e_2 \equiv \text{blue}$  (resp. *down quark*) and  $e_3 \equiv \text{green}$  (resp. *strange quark*). Thus,  $Q(p,0)$  is the totally symmetric part of a system of  $p$  quarks. Analogously,  $Q(0,q)$  is the totally symmetric part of a system of  $q$  antiquarks since the representation  $Q(0,1)$  describes an antiquark,  $\check{e}_1 \equiv \text{antigreen}$  (*strange antiquark*),  $\check{e}_2 \equiv \text{antiblue}$  (*down antiquark*) and  $\check{e}_3 \equiv \text{antired}$  (*up antiquark*).

**Remark 3.8.** *Because the operator space of  $Q(p,0)$ , or  $Q(0,p)$ , have the maximal isotropy subgroup  $H \simeq U(2)$ , these representations are also called symmetric representations, in the literature. So, pure-quark systems could also be referred to as symmetric quark systems.*

**Remark 3.9.** *There is another interpretation of the representation  $Q(p,0)$  as a quantum system that matches our classical phase space  $\mathbb{C}P^2$  in an enlightening way. A three-dimensional isotropic harmonic oscillator is a quantum system which is governed by a Hamiltonian of the form*

$$(3.14) \quad H = \sum_{i=1}^3 \left( \frac{1}{2m} P_i^2 + \frac{1}{2} m \omega^2 X_i^2 \right),$$

where  $P_i$  and  $X_i$  are the component operators of momentum and position, for some positive parameters  $m$  and  $\omega$ . It has degenerate energy levels<sup>19</sup>

$$(3.15) \quad E = \left( n_1 + n_2 + n_3 + \frac{3}{2} \right) \omega$$

<sup>17</sup>Also,  $e_{0,0,p}$  is a lowest weight vector.

<sup>18</sup>Again,  $\check{e}_{p,0,0}$  is a highest weight vector and  $\check{e}_{0,0,p}$  is a lowest weight vector.

<sup>19</sup>We set  $\hbar = 1$  during all this work.

with  $SU(3)$ -symmetry given by representations  $Q(p, 0)$  for  $p = n_1 + n_2 + n_3$  [20]. Thus, a quantum pure-quark system  $\mathcal{H}_{p,0}$  can be interpreted as the solution of a quantum isotropic harmonic oscillator of energy  $E_p = p + 3/2$ , by setting  $\omega = 1$ .

For the classical three-dimensional isotropic harmonic oscillator of same parameters  $m$  and  $\omega$ , the phase space is  $T^*\mathbb{R}^3 \simeq \mathbb{R}^3 \times \mathbb{R}^3 = \mathbb{R}^6$  and the Hamiltonian is

$$(3.16) \quad h = \sum_{i=1}^3 \left( \frac{p_i^2}{2m} + \frac{1}{2}m\omega x_i^2 \right) ,$$

where  $p_i$  and  $x_i$  are the components of momentum and position, in complete analogy to (3.14). By appropriately rescaling, a region of given fixed energy is identified with  $S^5 \subset \mathbb{R}^6$ . The solution passing through a point of  $S^5$  is the orbit of the point under an  $SO(2)$ -action, where  $SO(2)$  acts via rotations on each  $\mathbb{R}^2$  determined by the pairs  $(x_i, p_i)$ . But the action of  $SO(2)$  on  $\mathbb{R}^2$  is equivalent to the action of  $U(1)$  on  $\mathbb{C}$ , so the set of solutions of a classical isotropic three-dimensional harmonic oscillator is identified with  $S^5/S^1 = \mathbb{C}P^2$ .

If  $Q(\mathbf{p}) = Q(p, 0)$ , the Gelfand-Tsetlin pattern (2.21) is reduced to

$$(3.17) \quad \begin{aligned} 0 &\leq r \leq p , \\ 0 &\leq \nu_1 \leq r , \nu_2 = r - \nu_1 , \nu_3 = p - r , \\ I &= \frac{r}{2} . \end{aligned}$$

A similar simplification can be applied to  $Q(\mathbf{p}) = Q(0, p)$ :

$$(3.18) \quad \begin{aligned} 0 &\leq r \leq p , \\ r &\leq \nu_1 \leq p , \nu_2 = p + r - \nu_1 , \nu_3 = p - r , \\ I &= \frac{p - r}{2} . \end{aligned}$$

In both cases, the isospin  $I$  is determined by  $\nu_3$ , so we simplify the notation for  $\mathbf{p} \in (\mathbb{N} \times \{0\}) \cup (\{0\} \times \mathbb{N})$  as

$$(3.19) \quad \mathbf{e}(\mathbf{p}; \boldsymbol{\nu}) := \mathbf{e}(\mathbf{p}; \boldsymbol{\nu}, I) , \check{\mathbf{e}}(\check{\mathbf{p}}; \boldsymbol{\nu}) := \check{\mathbf{e}}(\check{\mathbf{p}}; \boldsymbol{\nu}, I) .$$

To clear even more the notation, we will denote the elements of a coupled basis of  $\mathcal{B}(\mathcal{H}_{\mathbf{p}})$  that lies in the  $Q(n, n)$ -invariant subspace by  $\mathbf{e}(n; \boldsymbol{\nu}, I)$ . Consequently, the notation for the Clebsch-Gordan coefficients can be simplified to

$$(3.20) \quad C_{\boldsymbol{\nu}_1, \boldsymbol{\nu}_2, \boldsymbol{\nu} I}^{\mathbf{p}, \check{\mathbf{p}}, n} := C_{\boldsymbol{\nu}_1 I_1, \boldsymbol{\nu}_2 I_2, \boldsymbol{\nu} I}^{\mathbf{p}, \check{\mathbf{p}}, n} := C_{\boldsymbol{\nu}_1 I_1, \boldsymbol{\nu}_2 I_2, \boldsymbol{\nu} I}^{\mathbf{p}, \check{\mathbf{p}}, (n, n)} .$$

Applying the same simplification to the Wigner product symbol, expressions (2.149) and (2.150) now read as

$$(3.21) \quad \begin{bmatrix} n_1 & n_2 & n_3 \\ \boldsymbol{\nu}_1, I_1 & \boldsymbol{\nu}_2, I_2 & \boldsymbol{\nu}_3, I_3 \end{bmatrix} [\mathbf{p}] \neq 0 \quad \Longrightarrow \quad \begin{cases} \nabla_{\boldsymbol{\nu}_1 + \boldsymbol{\nu}_2, \check{\boldsymbol{\nu}}_3} = 1 \\ \delta(I_1, I_2, I_3) = 1 \end{cases} ,$$

$$\begin{aligned}
(3.22) \quad \begin{bmatrix} n_1 & n_2 & n_3 \\ \nu_1, I_1 & \nu_2, I_2 & \nu_3, I_3 \end{bmatrix} [\mathbf{p}] &= \begin{bmatrix} n_3 & n_1 & n_2 \\ \nu_3, I_3 & \nu_1, I_1 & \nu_2, I_2 \end{bmatrix} [\mathbf{p}] \\
&= \begin{bmatrix} n_2 & n_3 & n_1 \\ \nu_2, I_2 & \nu_3, I_3 & \nu_1, I_1 \end{bmatrix} [\mathbf{p}] \\
&= \begin{bmatrix} n_2 & n_1 & n_3 \\ \check{\nu}_2, I_2 & \check{\nu}_1, I_1 & \check{\nu}_3, I_3 \end{bmatrix} [\mathbf{p}] \\
&= \begin{bmatrix} n_1 & n_2 & n_3 \\ \check{\nu}_1, I_1 & \check{\nu}_2, I_2 & \check{\nu}_3, I_3 \end{bmatrix} [\check{\mathbf{p}}] \\
&= \begin{bmatrix} n_2 & n_1 & n_3 \\ \nu_2, I_2 & \nu_1, I_1 & \nu_3, I_3 \end{bmatrix} [\check{\mathbf{p}}] .
\end{aligned}$$

Therefore, Corollary 2.29 takes the special form:

**Theorem 3.10.** *For a quantum pure-quark system  $\mathcal{H}_{\mathbf{p}}$ ,  $|\mathbf{p}| = p$ , the product of elements of a coupled basis of the space of operators  $\mathcal{B}(\mathcal{H}_{\mathbf{p}})$  decomposes in the form (3.23)*

$$e(n_1; \nu_1, I_1) e(n_2; \nu_2, I_2) = \sum_{n=0}^p \sum_{\nu, I} (-1)^{p+2(t_\nu+u_\nu)} \begin{bmatrix} n_1 & n_2 & n_3 \\ \nu_1, I_1 & \nu_2, I_2 & \check{\nu}, I \end{bmatrix} [\mathbf{p}] e(n; \nu, I)$$

for  $0 \leq n_1, n_2 \leq p$ , where summations over  $\nu$  and  $I$  can be restricted to  $\nabla_{\nu_1+\nu_2, \nu} = 1$  and  $\delta(I_1, I_2, I) = 1$  due to (3.21).

We also identify the operator algebra  $\mathcal{B}(\mathcal{H}_{\mathbf{p}})$  with the matrix algebra  $M_{\mathbb{C}}(d)$  by means of an uncoupled basis of  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}})$ . So let  $\nu$  and  $\check{\nu}$  be such that  $\Delta_{\nu, \check{\nu}}^{|\mathbf{p}|} = 1$ , the operator  $e(\mathbf{p}, \nu) \otimes \check{e}(\check{\mathbf{p}}, \check{\nu})$  is a diagonal matrix and its decomposition in the coupled basis can be written as

$$(3.24) \quad e(\mathbf{p}; \nu) \otimes \check{e}(\check{\mathbf{p}}; \check{\nu}) = \sum_{n=0}^p \sum_{I=0}^n C_{\nu, \check{\nu}, n}^{\mathbf{p}, \check{\mathbf{p}}, n} e(n; (n, n, n), I) .$$

That is, any diagonal matrix is a linear combination of  $\{e(n, (n, n, n), I)\}$ . Since the cardinality of this set is  $(p+1)(p+2)/2$ , it is the set of diagonal matrices of a coupled basis. Clebsch-Gordan coefficients being all real implies that such matrices are also real.

**3.3. Symbol correspondences for pure-quark systems.** Let  $\mathbf{p} \in (\mathbb{N} \times \{0\}) \cup (\{0\} \times \mathbb{N})$  with  $|\mathbf{p}| = p$ .

**Definition 3.11.** *A symbol correspondence for a pure-quark system  $(\mathcal{H}_{\mathbf{p}}, Q(\mathbf{p}))$ , also referred to simply as a symbol correspondence or just as a correspondence, is an injective linear map  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_c^\infty(\mathbb{C}P^2) : A \mapsto W_A$  that satisfies, for any  $A \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$ ,*

- i) *Equivariance:*  $\forall g \in SU(3), W_{A^g} = (W_A)^g$  ;
- ii) *Reality:*  $W_{A^\dagger} = \overline{W_A}$  ;
- iii) *Normalization:*  $\int_{\mathbb{C}P^2} W_A(\mathbf{x}) d\mathbf{x} = \frac{2}{(p+1)(p+2)} \text{tr}(A)$  .

**Remark 3.12.** *If one replace  $\mathbb{C}P^2$  on the definition of a symbol correspondence for a pure quark system by the orbit  $\mathcal{O}_{x,0}$  or  $\mathcal{O}_{0,x}$ , for  $x > 0$ , the definition remains essentially the same. Given any symbol correspondence  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_c^\infty(\mathbb{C}P^2)$ , the map  $W' : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_c^\infty(\mathcal{O}_{x,0})$ ,  $W'_A = W_A \circ \psi_{x,0}$ , satisfies the desired properties*

and  $W_A = W'_A \circ \psi_{x,0}^{-1}$ . Using  $\psi_{0,x}$ , we get symbol correspondences with codomain  $C_{\mathbb{C}}^{\infty}(\mathcal{O}_{0,x})$ . Conveniently, one can define symbol correspondences as maps  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathcal{O}_{\mathbf{p}})$ .

A symbol correspondence is an injective equivariant linear map, thus Schur's Lemma leads us to a simple characterization of these correspondences for a pure-quark system:

**Theorem 3.13.** *A linear map  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$  is a symbol correspondence if and only if it maps*

$$(3.25) \quad W : \sqrt{\frac{(p+1)(p+2)}{2}} e(n; \boldsymbol{\nu}, I) \mapsto c_n X_{\boldsymbol{\nu}, I}^n$$

for  $(c_1, \dots, c_p) \in (\mathbb{R}^*)^p$  and  $c_0 = (-1)^p$ .

*Proof.* By the Schur's Lemma,  $W$  is equivariant and injective if and only if it provides a mapping of the form (3.25) with  $c_n \neq 0$  for every  $n \in \{0, \dots, p\}$ . Since  $e^\dagger(n, \boldsymbol{\nu}, I) = (-1)^{2(t+u)} e(n, \check{\boldsymbol{\nu}}, I)$  and  $\overline{X_{\boldsymbol{\nu}, I}^n} = (-1)^{2(t+u)} X_{\check{\boldsymbol{\nu}}, I}^n$ , reality holds if and only if the constants  $c_n$  are all real numbers. To finish, we have

$$(3.26) \quad (-1)^p \sqrt{\frac{(p+1)(p+2)}{2}} e(0; (0, 0, 0), 0) = \mathbf{1}$$

and  $X_{\mathbf{0},0}^0 = 1$ , then the normalization condition is satisfied if and only if  $c_0 = (-1)^p$ .  $\square$

**Corollary 3.14.** *The moduli space of correspondences for a pure-quark system  $\mathcal{H}_{\mathbf{p}}$  is  $(\mathbb{R}^*)^p$ .*

But there is another way to construct symbol correspondences. Again, let  $\mathbf{x}_0 = [0 : 0 : 1] \in \mathbb{C}P^2$ . Given an operator  $K \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$  fixed by  $H$ , consider the operator-valued function  $\mathbb{C}P^2 \rightarrow \mathcal{B}(\mathcal{H}_{\mathbf{p}}) : \mathbf{x} \mapsto K(\mathbf{x}) = K^g$ , where  $g \in SU(3)$  is such that  $\mathbf{x} = g\mathbf{x}_0$ .

**Theorem 3.15.** *A map  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2) : A \mapsto W_A$  is a symbol correspondence satisfying (3.25) if and only if*

$$(3.27) \quad W_A(\mathbf{x}) = \text{tr}(AK(\mathbf{x})) ,$$

that is,

$$(3.28) \quad W_A(g\mathbf{x}_0) = \text{tr}(AK^g) ,$$

for  $K \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$  of the form

$$(3.29) \quad K = \frac{2}{(p+1)(p+2)} \mathbf{1} + \sum_{n=1}^p c_n \sqrt{\frac{2(n+1)^3}{(p+1)(p+2)}} e(n; (n, n, n), 0) .$$

In particular,  $K$  is a diagonal matrix with real entries and unitary trace.

*Proof.* Suppose  $W$  is a symbol correspondence given by (3.25). The map  $A \mapsto W_A(\mathbf{x}_0)$  is a linear functional, then there exists  $K \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$  such that (3.28) holds for the identity of  $SU(3)$ . From equivariance, we have  $W_A(g^{-1}\mathbf{x}_0) = (W_A)^g(\mathbf{x}_0) = W_{A^g}(\mathbf{x}_0) = \text{tr}(A^g K) = \text{tr}(AK^{g^{-1}})$  for very  $g \in SU(3)$ . Since  $\mathbf{x}_0$  is fixed by  $H$ ,

we have  $\text{tr}(AK^g) = \text{tr}(AK)$  for every  $g \in H$  and every operator  $A$ . Thus, we can write, cf. Proposition 3.2,

$$(3.30) \quad K = \sum_{n=0}^p k_n \mathbf{e}(n; (n, n, n), 0) ,$$

that is,  $K$  is a diagonal matrix. Taking  $A = \mathbf{e}(n; \boldsymbol{\nu}, I) = (-1)^{2(t+u)} \mathbf{e}^\dagger(n; \check{\boldsymbol{\nu}}, I)$ , we have

$$(3.31) \quad W_A(g\mathbf{x}_0) = \text{tr}(AK^g) = k_n (-1)^{2(t+u)} D_{\check{\boldsymbol{\nu}}I, (n, n, n)_0}^{(n, n)}(g) = \overline{k_n D_{\boldsymbol{\nu}I, (n, n, n)_0}^{(n, n)}} ,$$

cf (2.38). Then,

$$(3.32) \quad W : \mathbf{e}(n, \boldsymbol{\nu}, I) \mapsto \frac{k_n}{(n+1)^{3/2}} X_{\boldsymbol{\nu}, I}^n .$$

It follows from Theorem 3.13 that

$$(3.33) \quad k_n = c_n \sqrt{\frac{2(n+1)^3}{(p+1)(p+2)}} .$$

On the other hand, for  $K$  given by (3.29), equations (3.30)-(3.32) imply that (3.28) defines a symbol correspondence given by (3.25).  $\square$

These results are completely analogous to the case of spin systems, cf. [18], so we come with the next definition.

**Definition 3.16.** *An operator kernel  $K \in \mathcal{B}(\mathcal{H}_p)$  is an operator that induces a symbol correspondence via (3.28). That is,  $K$  is given by (3.29) with non-zero real numbers ( $c_n$ ) called characteristic numbers of both the operator kernel and the symbol correspondence.*

If  $K \in \mathcal{B}(\mathcal{H}_p)$  is an operator kernel, it is diagonal with real entries, thus it is a linear combination of projections of the form

$$(3.34) \quad K = \sum_{\boldsymbol{\nu}} a_{\boldsymbol{\nu}} \Pi_{\boldsymbol{\nu}} ,$$

for real coefficients  $a_{\boldsymbol{\nu}}$ . We can separate the summation into different values of isospin:

$$(3.35) \quad K = \sum_{j=0}^p K_j , \quad K_j = \sum_{\boldsymbol{\nu} \in j/2} a_{\boldsymbol{\nu}} \Pi_{\boldsymbol{\nu}} ,$$

where  $\boldsymbol{\nu} \in j/2$  means the weight  $\boldsymbol{\nu}$  has isospin  $j/2$ , cf. (3.17)-(3.18).

**Proposition 3.17.** *If  $K \in \mathcal{B}(\mathcal{H}_p)$  is an operator kernel, then*

$$(3.36) \quad K = \sum_{j=0}^p a_j \sum_{\boldsymbol{\nu} \in j/2} \Pi_{\boldsymbol{\nu}} ,$$

where the coefficients  $a_j$  are real numbers satisfying

$$(3.37) \quad \sum_{j=0}^p a_j (j+1) = 1 .$$

*Proof.* Every operator kernel is fixed by  $H$  of (2.156), so  $K$  must be fixed also by the  $SU(2)$  of (2.161). Decomposing  $K$  as in (3.35), we have that a component  $K_j$  is an operator on a representation  $j/2$  of  $SU(2)$ , therefore it must commute with  $T_3$  and  $T_{\pm}$ , which implies each  $K_j$  is a multiple of the identity operator on the subspace spanned by the states with same isospin  $j/2$ , that is,

$$(3.38) \quad K_j = a_j \sum_{\nu \in j/2} \Pi_{\nu} ,$$

where  $a_j$  is real. To finish,  $\text{tr}(K) = 1$  implies (3.37).  $\square$

We can also use operator kernels to construct symbol correspondences in an implicit way, as follows.

**Proposition 3.18.** *Let  $K$  be an operator kernel with characteristic numbers  $(c_n)$ . The equation*

$$(3.39) \quad A = \frac{(p+1)(p+2)}{2} \int_{\mathbb{C}P^2} \widetilde{W}_A(\mathbf{x}) K(\mathbf{x}) d\mathbf{x} .$$

defines a symbol correspondence  $\widetilde{W}$  with characteristic numbers  $(\tilde{c}_n)$  given by  $\tilde{c}_n = 1/c_n$ .

*Proof.* We have

$$(3.40) \quad \begin{aligned} \int_{\mathbb{C}P^2} X_{\nu,I}^n(\mathbf{x}) K(\mathbf{x}) d\mathbf{x} &= \int_{SU(3)} X_{\nu,I}^n(g\mathbf{x}_0) K^g dg \\ &= \sum_{n', \mu, J} k_{n'} \int_{SU(3)} X_{\nu,I}^n(g\mathbf{x}_0) D_{\mu, J, n'_0}^{(n', n')}(g\mathbf{x}_0) dg \mathbf{e}(n'; \mu, J) \\ &= \sum_{n', \mu, J} \frac{k_{n'}}{(n'+1)^{3/2}} \langle X_{\mu, J}^{n'} | X_{\nu, I}^n \rangle \mathbf{e}(n; \mu, J) \\ &= \frac{k_n}{(n+1)^{3/2}} \mathbf{e}(n; \nu, I) , \end{aligned}$$

where

$$(3.41) \quad k_n = c_n \sqrt{\frac{2(n+1)^3}{(p+1)(p+2)}} .$$

Thus,

$$(3.42) \quad \frac{(p+1)(p+2)}{2} \int_{\mathbb{C}P^2} \frac{1}{c_n} X_{\nu, I}^n(\mathbf{x}) K(\mathbf{x}) d\mathbf{x} = \sqrt{\frac{(p+1)(p+2)}{2}} \mathbf{e}(n; \nu, I) .$$

By Theorem 3.13,  $\widetilde{W}$  is a symbol correspondence with characteristic numbers  $(\tilde{c}_n)$  satisfying  $\tilde{c}_n = 1/c_n$ .  $\square$

Actually, there is a duality relation between the symbol correspondences defined by the same operator kernel via (3.27) and via (3.39) considering the normalized inner product

$$(3.43) \quad \langle A|R \rangle_{\mathbf{p}} = \frac{2}{(p+1)(p+2)} \langle A|R \rangle = \frac{2}{(p+1)(p+2)} \text{tr}(A^\dagger R)$$

for all  $A, R \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$ .

**Definition 3.19.** Given a symbol correspondence  $W : \mathcal{B}(\mathcal{H}_p) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$ , its dual correspondence is the symbol correspondence  $\widetilde{W} : \mathcal{B}(\mathcal{H}_p) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$  that satisfies

$$(3.44) \quad \langle A|R \rangle_p = \left\langle \widetilde{W}_A \middle| W_R \right\rangle = \left\langle W_A \middle| \widetilde{W}_R \right\rangle .$$

The operators kernels of  $W$  and  $\widetilde{W}$  are also said to be dual to each other.

**Proposition 3.20.** Let  $K$  be an operator kernel. The symbol correspondences defined by  $K$  via (3.27) and via (3.39) are dual to each other, that is, for any symbol correspondence with characteristic numbers  $(c_n)$ , its dual correspondence has characteristic numbers  $(1/c_n)$ .

*Proof.* Given any two operators  $A$  and  $R$ , if we write  $A^\dagger$  as in (3.39) and use the reality property, we get

$$(3.45) \quad \frac{2}{(p+1)(p+2)} \operatorname{tr}(A^\dagger R) = \int_{\mathbb{C}P^2} \overline{\widetilde{W}_A(\mathbf{x})} \operatorname{tr}(RK(\mathbf{x})) d\mathbf{x} = \int_{\mathbb{C}P^2} \overline{\widetilde{W}_A(\mathbf{x})} W_R(\mathbf{x}) d\mathbf{x} ,$$

that is,  $\langle A|R \rangle_p = \left\langle \widetilde{W}_A \middle| W_R \right\rangle$ . Writing  $R$  as in (3.39), we get  $\langle A|R \rangle_p = \left\langle W_A \middle| \widetilde{W}_R \right\rangle$ .  $\square$

In particular, a symbol correspondence  $W$  is an isometry if and only if it is self-dual.

**Definition 3.21.** A symbol correspondence  $W : \mathcal{B}(\mathcal{H}_p) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$  is a Stratonovich-Weyl correspondence if it is an isometry, that is,

$$(3.46) \quad \langle A|R \rangle_p = \langle W_A|W_R \rangle$$

for all  $A, R \in \mathcal{B}(\mathcal{H}_p)$ .

**Proposition 3.22.** A symbol correspondence is a Stratonovich-Weyl correspondence if and only if its characteristic numbers  $(c_n)$  satisfy  $|c_n| = 1$ .

*Proof.* The proof follows from Proposition 3.20.  $\square$

Although isometry is a very nice property for a symbol correspondence, there is another property for symbol correspondences that is very reasonable, from a physical perspective. Recall that a Hermitian operator with only nonnegative eigenvalues is called *positive*, or *positive-definite* if all of its eigenvalues are positive, and a real function that takes only nonnegative values is called *positive*, or *strictly-positive* if it only takes positive values.

**Definition 3.23.** A symbol correspondence for a pure-quark system is mapping-positive if it maps positive(-definite) operators to (strictly-)positive functions. A symbol correspondence for a pure-quark system which is dual to a mapping-positive correspondence is a positive-dual correspondence.

In Theorem 3.15, we characterize all symbol correspondences as expected values with respect to  $K^g$ , where  $K$  is an operator kernel, that is, an  $H$ -invariant operator satisfying equation (3.29) with  $c_n \in \mathbb{R}^*$ . From (3.34)-(3.37),  $K$  can be identified with an  $H$ -invariant *pseudo-state*, since, from Physics, we have the following:

**Definition 3.24.** An operator on a complex Hilbert space is a state if it is a positive operator with unitary trace.

On the other hand, from (3.34)-(3.37), because the real numbers  $a_r$  for a general operator kernel  $K$  can be negative, generally  $K$  has unitary trace but is not a positive operator and we can see it as providing pseudo-probabilities, since an operator kernel which is actually a state should provide actual probabilities. In fact, we have:

**Proposition 3.25.** *A symbol correspondence  $W : \mathcal{B}(\mathcal{H}_p) \rightarrow C_c^\infty(\mathbb{C}P^2)$  with operator kernel  $K$  is mapping-positive if and only if  $K$  is a state, that is,  $K$  given by (3.29), with  $c_n \in \mathbb{R}^*$ , is in the convex hull of  $\{\Pi_\nu\}$ , that is,  $K \in \text{Conv}\{\Pi_\nu\}$ .*

*Proof.* We assume that  $K$  is an operator kernel, thus  $K$  is given by (3.29) with  $c_n \in \mathbb{R}^*$ . Now, let  $K$  be decomposed as in (3.36). Suppose  $K \in \text{Conv}\{\Pi_\nu\}$ , so that the coefficients  $a_j$  in the summation are all nonnegative. An operator  $A \in \mathcal{B}(\mathcal{H}_p)$  is positive if and only if  $A = R^\dagger R$  for some  $R \in \mathcal{B}(\mathcal{H}_p)$ , and  $A$  is positive-definite if and only if  $R$  is an automorphism. Thus, for any  $g \in SU(3)$  and  $\tilde{R} = R\rho(g)$ ,

$$(3.47) \quad \begin{aligned} W_A(g\mathbf{x}_0) &= \sum_{j=0}^p a_j \sum_{\nu \in j/2} \text{tr}(R^\dagger R \rho(g) \Pi_\nu \rho(g)^\dagger) = \sum_{j=0}^p a_j \sum_{\nu \in j/2} \text{tr}(R \rho(g) \Pi_\nu \rho(g)^\dagger R^\dagger) \\ &= \sum_{j=0}^p a_j \sum_{\nu \in j/2} \text{tr}(\tilde{R} \Pi_\nu \tilde{R}^\dagger) = \sum_{j=0}^p a_j \sum_{\nu \in j/2} \left\| \tilde{R}(e(\mathbf{p}; \nu)) \right\|^2 \\ &\geq 0, \end{aligned}$$

where the inequality is strict if  $\tilde{R}$  is an automorphism, which is true if  $R$  is an automorphism, that is, if  $A$  is positive-definite.

Now, suppose  $K \notin \text{Conv}\{\Pi_\nu\}$ . Then, there is a negative coefficient  $a_j$  so that any projection  $\Pi_\nu$  with  $\nu \in j/2$  is a positive operator satisfying  $W_{\Pi_\nu}(\mathbf{x}_0) = \text{tr}(K \Pi_\nu) < 0$ .  $\square$

For a pure-quark system  $\mathcal{H}_p$ , let  $\mathcal{S}_\leq^p$ ,  $\mathcal{S}_<^p$  and  $\mathcal{S}_>^p$  be the sets of Stratonovich-Weyl, mapping-positive and positive-dual correspondences, respectively.

**Theorem 3.26.** *The sets  $\mathcal{S}_\leq^p$ ,  $\mathcal{S}_<^p$  and  $\mathcal{S}_>^p$  are mutually disjoint.*

*Proof.* If  $W \in \mathcal{S}_<^p$ , then Proposition 3.25 implies that its operator kernel  $K$  is given by

$$(3.48) \quad K = \sum_{\nu} (-1)^{2(t+u)} a_\nu e(\mathbf{p}; \nu) \otimes \check{e}(\check{\mathbf{p}}; \check{\nu}),$$

where  $a_\nu$  are coefficients of a convex combination, that is, they are non negative and sum up to 1. From (3.29), its characteristic numbers are

$$(3.49) \quad c_n = \sqrt{\frac{(p+1)(p+2)}{2(n+1)^3}} \sum_{\nu} (-1)^{2(t+u)} a_\nu C_{\nu, \check{\nu}, (n, n, n)0}^{\mathbf{p}, \check{\mathbf{p}}, n}.$$

Since CG coefficients are coefficients of an unitary transformation, their absolute values are bounded above by 1, so

$$(3.50) \quad |c_n| \leq \sqrt{\frac{(p+1)(p+2)}{2(n+1)^3}} \sum_{\nu} a_\nu = \sqrt{\frac{(p+1)(p+2)}{2(n+1)^3}}.$$

Thus,

$$(3.51) \quad |c_p| < 1$$

and characteristic numbers  $(\tilde{c}_n)$  of the correspondence  $\widetilde{W} \in \mathcal{S}_>^p$  dual to  $W \in \mathcal{S}_<^p$  satisfy

$$(3.52) \quad |\tilde{c}_p| = \frac{1}{|c_p|} > 1 ,$$

cf. Proposition 3.20. From Proposition 3.22 and inequalities (3.51)-(3.52), we conclude the statement.  $\square$

**Remark 3.27.** *Theorem 3.26 for pure-quark systems mirrors the analogous result for spin systems, cf. Theorem 3.3 in [1].*

To verify the existence of a mapping-positive correspondence, we consider the defining representation  $\rho_1$  of  $SU(3)$  on  $\mathbb{C}^3$ , and invoke the canonical projection

$$\tilde{\pi} : S^5 \rightarrow \mathbb{C}P^2 ,$$

inside  $\mathbb{C}^3 \rightarrow \mathbb{C}P^2$ , and

$$(3.53) \quad \Phi_p : \mathbb{C}^3 \rightarrow \mathbb{C}^{(p+1)(p+2)/2} \\ (z_1, z_2, z_3) \mapsto (z_1^p, \dots, \sqrt{\binom{p}{j, k, l}} z_1^j z_2^k z_3^l, \dots, z_3^p) ,$$

where  $\mathbb{C}^{(p+1)(p+2)/2} \simeq \mathcal{H}_{p,0}$  is endowed with a  $SU(3)$ -representation  $\rho_p$  of class  $Q(p,0)$ , cf. (3.12).

**Proposition 3.28.** *The map  $B : \mathcal{B}(\mathcal{H}_{p,0}) \rightarrow C_c^\infty(\mathbb{C}P^2) : A \mapsto B_A$ , with*

$$B_A(\mathbf{x}) = \langle \Phi_p(\mathbf{n}) | A \Phi_p(\mathbf{n}) \rangle$$

for  $\mathbf{x} \in \mathbb{C}P^2$  and  $\mathbf{n} \in S^5$  related by  $\tilde{\pi}(\mathbf{n}) = \mathbf{x}$ , is a mapping-positive symbol correspondence whose operator kernel is the projection  $\Pi_{(0,0,p)}$  onto the lowest weight vector and whose characteristic numbers are

$$(3.54) \quad b_n = (-1)^p \sqrt{\frac{(p+1)(p+2)}{2(n+1)^3}} C_{(0,0,p),(p,p,0),(n,n,n),0}^{(p,0), (0,p), n} .$$

*Proof.* First of all, we need to verify that  $B_A$  is a well defined function for every  $A \in \mathcal{B}(\mathcal{H}_{p,0})$ . For  $\mathbf{n}, \mathbf{n}' \in S^5$ , we have  $\tilde{\pi}(\mathbf{n}) = \tilde{\pi}(\mathbf{n}')$  if and only if  $\mathbf{n}' = e^{i\theta} \mathbf{n}$ , but  $\Phi_p(e^{i\theta} \mathbf{n}) = e^{ip\theta} \Phi_p(\mathbf{n})$ , so

$$(3.55) \quad \langle \Phi_p(e^{i\theta} \mathbf{n}) | A \Phi_p(e^{i\theta} \mathbf{n}) \rangle = \langle e^{ip\theta} \Phi_p(\mathbf{n}) | e^{ip\theta} A \Phi_p(\mathbf{n}) \rangle = \langle \Phi_p(\mathbf{n}) | A \Phi_p(\mathbf{n}) \rangle .$$

This proves that  $B_A$  is well defined. It is also smooth, since  $\tilde{\pi}$  is a surjective submersion and  $B_A \circ \tilde{\pi}$  is a composition of smooth functions of  $S^5$ .

The linearity of  $B$  is trivial. The equivariance follows straightforwardly from the equivariance of  $\Phi_p$ . For any  $g \in SU(3)$ ,

$$(3.56) \quad B_{A^g}(\mathbf{x}) = \langle \Phi_p(\mathbf{n}) | A^g \Phi_p(\mathbf{n}) \rangle = \langle \Phi_p(\mathbf{n}) | \rho_p(g) A \rho_p(g)^{-1} \Phi_p(\mathbf{n}) \rangle \\ = \langle \rho_p(g)^{-1} \Phi_p(\mathbf{n}) | A \rho_p(g)^{-1} \Phi_p(\mathbf{n}) \rangle = \langle \Phi_p(\rho_1(g)^{-1} \mathbf{n}) | A \Phi_p(\rho_1(g)^{-1} \mathbf{n}) \rangle \\ = B_A(g^{-1} \mathbf{x}) = (B_A)^g(\mathbf{x}) ,$$

cf. (2.39). Equivariance implies that  $\ker B$  is an invariant subspace, and we use that to prove  $B$  is injective by means of contradiction. Suppose  $B$  is not injective, then  $\ker B$  contains an irreducible representation of the form  $Q(n, n)$ , so the diagonal matrix  $e(n, \mathbf{n}, I)$  lies in  $\ker B$ , that is, there exists a non-zero diagonal matrix  $D = \text{diag}(d_{p,0,0}, \dots, d_{j,k,l}, \dots, d_{0,0,p}) \in \ker B$ . Thus, we have

$$(3.57) \quad B_D(\mathbf{x}) = \langle \Phi_p(\mathbf{n}) | D \Phi_p(\mathbf{n}) \rangle = \sum_{j+k+l=1} \binom{p}{j, k, l} d_{j,k,l} |z_1|^{2j} |z_2|^{2k} |z_3|^{2l} = 0$$

for every  $\mathbf{n} = (z_1, z_2, z_3) \in S^5$ . It follows that the polynomials with real coefficients

$$\sum_{j+k+l=1} \binom{p}{j, k, l} \text{Re}(d_{j,k,l}) x_1^{2j} x_2^{2k} x_3^{2l}, \quad \sum_{j+k+l=1} \binom{p}{j, k, l} \text{Im}(d_{j,k,l}) x_1^{2j} x_2^{2k} x_3^{2l}$$

are identically zero, so  $D$  must be a zero matrix, and this is the desired contradiction. Therefore,  $B$  is injective.

We know that

$$(3.58) \quad \begin{aligned} \Pi_{(0,0,p)} &= e((p, 0); (0, 0, p)) \otimes e^*((p, 0); (0, 0, p)) \\ &= (-1)^p e((p, 0); (0, 0, p)) \otimes \check{e}((0, p); (p, p, 0)) \\ &= (-1)^p \sum_{n=0}^p C_{(0,0,p), (p,p,0), (n,n,n), 0}^{(p,0), (0,p), n} e(n; (n, n, n), 0) \\ &= \frac{2}{(p+1)(p+2)} \mathbf{1} + (-1)^p \sum_{n=1}^p C_{(0,0,p), (p,p,0), (n,n,n), 0}^{(p,0), (0,p), n} e(n; (n, n, n), 0), \end{aligned}$$

where the last equality follows from  $\text{tr}(\Pi_{(0,0,p)}) = 1$ . Since the Clebsch-Gordan coefficients are real and  $B$  is an injective map, it is sufficient to show  $B_A(g\mathbf{x}_0) = \text{tr}(A\Pi_{(0,0,p)}^g)$ . We have that  $\mathbf{n}_0 = (0, 0, 1) \in S^5$  satisfies  $\pi(\mathbf{n}_0) = \mathbf{x}_0$  and  $\Phi_p(\mathbf{n}_0) = (0, 0, \dots, 1)$ , so

$$(3.59) \quad B_A(\mathbf{x}_0) = \langle \Phi_p(\mathbf{n}_0) | A \Phi_p(\mathbf{n}_0) \rangle = \text{tr}(A\Pi_{(0,0,p)})$$

and

$$(3.60) \quad B_A(g\mathbf{x}_0) = (B_A)^{g^{-1}}(\mathbf{x}_0) = B_{A^{g^{-1}}}(\mathbf{x}_0) = \text{tr}(A^{g^{-1}}\Pi_{(0,0,p)}) = \text{tr}(A\Pi_{(0,0,p)}^g).$$

This proves that  $B$  is a symbol correspondence sequence with operator kernel  $\Pi_{(0,0,p)}$ . Proposition 3.25 implies that it is a mapping-positive correspondence.  $\square$

One could expect that the projection  $\Pi_{(p,0,0)} \in \mathcal{B}(\mathcal{H}_{p,0})$  onto the highest weight vector is also an operator kernel. This is not the case since  $\Pi_{(p,0,0)}$  is not  $H$ -invariant, cf. Proposition 3.17 and Proposition 3.30. However, this is just a matter of convention, a choice of  $U(2)$  subgroup by which we impose invariance. Recalling (3.10), we have  $\rho_p(g_0)e_{j,k,l} = (-1)^p e_{l,k,j}$  for  $g_0$  as in (2.158), so that  $\Pi_{(p,0,0)} = \Pi_{(0,0,p)}^{g_0}$  and  $\Pi_{(p,0,0)}$  is fixed by  $H' = g_0 H g_0$  as given in (2.157). If we set  $\mathbf{x}'_0 = g_0 \mathbf{x}_0 = [1 : 0 : 0]$ , then we can construct a symbol correspondence  $A \mapsto B'_A$  using  $\Pi_{(p,0,0)}$ :

$$(3.61) \quad B'_A(g\mathbf{x}'_0) = \text{tr}(A\Pi_{(p,0,0)}^g).$$

But

$$(3.62) \quad B'_A(g\mathbf{x}'_0) = \text{tr}\left(A\Pi_{(p,0,0)}^g\right) = \text{tr}\left(A\Pi_{(0,0,p)}^{gg_0}\right) = B_A(gg_0\mathbf{x}_0) = B_A(g\mathbf{x}'_0) \quad ,$$

that is,  $B'$  and  $B$  are the same map.

Notwithstanding, a minor adaptation of the argument of the previous proposition proves that  $\Pi_{(p,p,0)} \in \mathcal{B}(\mathcal{H}_{0,p})$ , the projection onto the highest weight vector, is also an operator kernel. Now, consider  $\check{\rho}_1$  and  $\check{\rho}_p$  the dual representations of  $\rho_1$  and  $\rho_p$ , respectively, so that the maps

$$(3.63) \quad \omega : \mathbb{C}^3 \rightarrow \mathbb{C}^3 : (z_1, z_2, z_3) \mapsto (-\bar{z}_3, \bar{z}_2, -\bar{z}_1)$$

and  $\Phi_p$  are both equivariant in the following sense:

$$(3.64) \quad \begin{aligned} \check{\rho}_p(g) \circ \omega &= \omega \circ \rho_p(g) \quad , \\ \check{\rho}_p(g) \circ \Phi_p &= \Phi_p \circ \check{\rho}_p(g) \quad , \end{aligned}$$

cf. (3.13).

**Proposition 3.29.** *The map  $B^- : \mathcal{B}(\mathcal{H}_{0,p}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2) : A \mapsto B_A^-$ , with*

$$B_A^-(\mathbf{x}) = \langle \Phi_p \circ \omega(\mathbf{n}) | A \Phi_p \circ \omega(\mathbf{n}) \rangle$$

for  $\mathbf{x} \in \mathbb{C}P^2$  and  $\mathbf{n} \in S^5 \subset \mathbb{C}^3$  related by  $\pi(\mathbf{n}) = \mathbf{x}$ , is a mapping-positive symbol correspondence whose operator kernel is the projection  $\Pi_{(p,p,0)}$  onto the highest weight vector and whose characteristic numbers are

$$(3.65) \quad b_{n-} = (-1)^p \sqrt{\frac{(p+1)(p+2)}{2(n+1)^3}} C_{(p,p,0),(0,0,p),(n,n,n),0}^{(0,p),(p,0),n} \quad .$$

*Proof.* The proof goes just as the proof of the previous proposition (linearity is, again, obvious), we just highlight that the following holds:  $\Phi_p \circ \omega(e^{i\theta}\mathbf{n}) = e^{-ip\theta}\Phi_p \circ \omega(\mathbf{n})$ ,

$$(3.66) \quad \begin{aligned} \Pi_{(p,p,0)} &= \check{e}((0,p);(p,p,0)) \otimes \check{e}^*((0,p);(p,p,0)) \\ &= (-1)^p \check{e}((0,p);(p,p,0)) \otimes e((p,0);(0,0,p)) \\ &= (-1)^p \sum_{n=0}^p C_{(p,p,0),(0,0,p),(n,n,n),0}^{(0,p),(p,0),n} e(n;(n,n,n),0) \\ &= \frac{2}{(p+1)(p+2)} \mathbb{1} + (-1)^p \sum_{n=1}^p C_{(p,p,0),(0,0,p),(n,n,n),0}^{(0,p),(p,0),n} e(n;(n,n,n),0) \end{aligned}$$

and  $\mathbf{n}_0 = (0,0,1) \in S^5$  satisfies  $\Phi_p \circ \omega(\mathbf{n}_0) = (1, \dots, 0, 0)$ .  $\square$

**Proposition 3.30.** *The operator  $\Pi_{(0,0,p)} \in \mathcal{B}(\mathcal{H}_{p,0})$  is the unique operator kernel among the projections onto weight vectors. The same is true for  $\Pi_{(p,p,0)} \in \mathcal{B}(\mathcal{H}_{0,p})$ .*

*Proof.* From equations (3.34) and (3.36), if  $K = \Pi_{\nu} \in \mathcal{B}(\mathcal{H}_{p,0})$  is an operator kernel, then (3.37) implies that  $j = 0$ , that is,  $\nu$  has null isospin. Using (3.17), we get that  $\nu = (0,0,p)$ . The proof for  $\mathcal{H}_{0,p}$  is completely analogous, but we use (3.18) to conclude  $\nu = (p,p,0)$ .  $\square$

**Definition 3.31.** *For any  $p \in \mathbb{N}$ , the symbol correspondences  $B : \mathcal{B}(\mathcal{H}_{p,0}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$ , with operator kernel  $\Pi_{(0,0,p)}$ , and  $B^- : \mathcal{B}(\mathcal{H}_{0,p}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$ , with operator kernel  $\Pi_{(p,p,0)}$ , are called Berezin correspondences.*

**Proposition 3.32.** *The Berezin correspondences for  $\mathcal{H}_{p,0}$  and  $\mathcal{H}_{0,p}$  have the same characteristic numbers.*

*Proof.* It follows from Theorem 2.23.  $\square$

**Remark 3.33.** *Our proof of Proposition 3.28 is, in essence, the same given in [18] for spin systems, where the symmetry group is  $SU(2)$  and the orbit is  $\mathbb{C}P^1 \simeq S^2$ . However, in that book, an involution on the space of symbol correspondences for spin systems is defined in this context. Let  $\{u_1, u_2\}$  be a spin standard basis and  $\{\check{u}_1 = -u_2^*, \check{u}_2 = u_1^*\}$  be the dual spin standard basis of  $\mathbb{C}^2$ , so that  $\omega_s(z_1, z_2) = (-\bar{z}_2, \bar{z}_1)$  is the dualization via inner product. Then  $\omega_s$  commutes with the action of  $SU(2)$  and induces an involution on  $\mathbb{C}P^1$  that coincides with the antipodal map  $\alpha_s$ , that is, for the Hopf map  $\pi_s : S^3 \rightarrow S^2$ , we have  $\pi_s \circ \omega_s = \alpha_s \circ \pi_s$ . If  $W^s$  is a symbol correspondence for a spin- $j$  system, we have that  $A \mapsto W_A^s \circ \alpha_s$  is also a symbol correspondence, known as the alternate correspondence of  $W^s$ . This is used to prove that both projections onto lowest and highest weight vectors of the same representation are correspondences. The name ‘‘alternate’’ is due to the fact that symbol correspondences for a spin- $j$  system are characterized by a set of non-zero real numbers  $\{c_l : l = 1, \dots, 2j\}$ , where each  $l$  stands for a representation of  $SU(2)$  in the decomposition of the operator algebra into irreducible representations, just as we have for pure quark systems, and to alternate a correspondence is equivalent to alternate the signs of such numbers, that is, to perform the change  $c_l \mapsto (-1)^l c_l$ .*

*One may try to reproduce the same argument for pure-quark systems, but this is not possible since the dualization  $\omega(z_1, z_2, z_3) = (-\bar{z}_3, \bar{z}_2, -\bar{z}_1)$  on  $\mathbb{C}^3$  does not commute with the action of  $SU(3)$ . In fact, for the defining representation  $\rho_1$  of  $SU(3)$  and its dual  $\check{\rho}_1$ , we have that  $\rho_1(g) \circ \omega = \omega \circ \check{\rho}_1(g)$  and  $\check{\rho}_1(g) \circ \omega = \omega \circ \rho_1(g)$  for every  $g \in SU(3)$ . Thus, in general, a symbol correspondence  $W$  for a pure-quark system does not satisfy  $W_{A^g} \circ \tilde{\alpha} = (W_A \circ \tilde{\alpha})^g$ , where  $\tilde{\alpha}([z_1 : z_2 : z_3]) = [-\bar{z}_3 : \bar{z}_2 : -\bar{z}_1]$  is the involution on  $\mathbb{C}P^2$  such that  $\tilde{\alpha} \circ \pi = \pi \circ \omega$ . Take, for instance,  $\mathbf{z}_0 = (0, 0, 1)$ , where  $\pi(\mathbf{z}_0) = [0 : 0 : 1] = \mathbf{x}_0$ , and*

$$(3.67) \quad g = \begin{pmatrix} 1/\sqrt{2} & -1/\sqrt{2} & 0 \\ 1/\sqrt{2} & 1/\sqrt{2} & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

so

$$(3.68) \quad \begin{aligned} g \tilde{\alpha}(\mathbf{x}_0) &= g \tilde{\alpha} \circ \pi(\mathbf{z}_0) = g \pi \circ \omega(\mathbf{z}_0) \\ &= \pi(\rho_1(g) \omega(\mathbf{z}_0)) = \pi \circ \omega(\check{\rho}_1(g) \mathbf{z}_0) \\ &= \pi \circ \omega(0, -1/\sqrt{2}, 1/\sqrt{2}) = \tilde{\alpha} \circ \pi(0, -1/\sqrt{2}, 1/\sqrt{2}) \\ &\neq \tilde{\alpha}(g \mathbf{x}_0). \end{aligned}$$

*At this point one may think to define alternate correspondences for pure-quark systems by applying  $c_n \mapsto (-1)^n c_n$  to the characteristic numbers. However, this is not the correct notion of alternation of correspondences for pure-quark systems as we show below.*

We now establish a solid link between Berezin correspondences for  $\mathcal{H}_{\mathbf{p}}$  and  $\mathcal{H}_{\check{\mathbf{p}}}$ .

**Definition 3.34.** *For a symbol correspondence  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$ , its antipodal correspondence is the symbol correspondence  $\check{W} : \mathcal{B}(\mathcal{H}_{\check{\mathbf{p}}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$  given*

by

$$(3.69) \quad \widetilde{W}_{A^*} = W_A ,$$

cf. (2.127)-(2.129).

If one defines symbol correspondences from  $\mathcal{B}(\mathcal{H}_{\mathbf{p}})$  to  $C_{\mathbb{C}}^{\infty}(\mathcal{O}_{\mathbf{p}})$  as pointed out in Remark 3.12, two antipodal correspondences  $W$  and  $\widetilde{W}$  are related by

$$\widetilde{W}_{A^*} = W_A \circ \iota ,$$

thus the name antipodal.

**Remark 3.35.** *It turns out that Definition 3.34 is completely analogous to alternation, for spin systems. The classes of irreducible representations of  $SU(2)$  are self-dual and their symbol correspondences satisfy  $W_{A^*}^s = W_A^s \circ \alpha_s$ , maintaining the same notation of Remark 3.33. But, now, for pure-quark systems the irreducible representations are not self-dual ( $\mathcal{H}_{\mathbf{p}} \neq \mathcal{H}_{\bar{\mathbf{p}}}$ ) and the alternation of characteristic numbers does not occur anymore.*

**Proposition 3.36.** *Two symbol correspondences  $W_1 : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$  and  $W_2 : \mathcal{B}(\mathcal{H}_{\bar{\mathbf{p}}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$  are antipodal to each other if and only if their characteristic numbers are equal.*

*Proof.* The result follows from (2.129) and Theorem 3.13.  $\square$

**Corollary 3.37.** *The Berezin correspondences for  $\mathcal{H}_{\mathbf{p}}$  and  $\mathcal{H}_{\bar{\mathbf{p}}}$  are antipodal to each other.*

**Corollary 3.38.** *A symbol correspondence for a pure-quark system is Stratonovich-Weyl correspondence if and only if its antipodal correspondence is a Stratonovich-Weyl correspondence.*

**3.4. Twisted product for pure-quark system.** Again, let  $\mathbf{p} \in (\mathbb{N} \times \{0\}) \cup (\{0\} \times \mathbb{N})$  with  $|\mathbf{p}| = p$ .

It is obvious from Theorem 3.13 that the images of all symbol correspondences for  $\mathcal{H}_{\mathbf{p}}$  and  $\mathcal{H}_{\bar{\mathbf{p}}}$  are the same space, namely, the space spanned by the  $\mathbb{C}P^2$  harmonics  $X_{\nu,I}^n$  with  $0 \leq n \leq p$ . Such space shall be denoted by  $\mathcal{X}_p$ , that is,

$$\mathcal{X}_p = \text{Span}_{\mathbb{C}}\{X_{\nu,I}^n\}_{0 \leq n \leq p} .$$

Now, we translate the operator algebra  $\mathcal{B}(\mathcal{H}_{\mathbf{p}})$  to  $\mathcal{X}_p$  using a symbol correspondence.

**Definition 3.39.** *Given a symbol correspondence  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$ , the twisted product of symbols induced by  $W$  is the binary operation  $\star$  on  $\mathcal{X}_p$  given by*

$$(3.70) \quad W_A \star W_R = W_{AR} ,$$

for any  $A, R \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$ . The algebra  $(\mathcal{X}_p, \star)$  is called a twisted  $\mathbf{p}$ -algebra.

**Proposition 3.40.** *Any twisted  $\mathbf{p}$ -algebra  $(\mathcal{X}_p, \star)$  is*

- i)  *$SU(3)$ -equivariant:  $(f_1 \star f_2)^g = f_1^g \star f_2^g$ ;*
- ii) *Associative:  $(f_1 \star f_2) \star f_3 = f_1 \star (f_2 \star f_3)$ ;*
- iii) *Unital:  $1 \star f = f \star 1 = f$ ;*
- iv) *A  $\ast$ -algebra:  $\overline{f_1 \star f_2} = \overline{f_2} \star \overline{f_1}$ ;*

where  $f_1, f_2, f_3, f \in \mathcal{X}_p$ ,  $g \in SU(3)$  and  $1 \in \mathcal{X}_p$  is the constant function equal to 1 on  $\mathbb{C}P^2$ , cf. (3.3).

*Proof.* The operator space  $\mathcal{B}(\mathcal{H}_{\mathbf{p}})$  is a  $SU(3)$ -equivariant unital associative  $*$ -algebra with respect to Hermitian conjugate, where  $\mathbb{1}$  is the identity. Since any symbol correspondence  $W$  for  $\mathcal{H}_{\mathbf{p}}$  is a  $SU(3)$ -equivariant linear isomorphism between  $\mathcal{B}(\mathcal{H}_{\mathbf{p}})$  and  $\mathcal{X}_{\mathbf{p}}$  satisfying reality and  $W_{\mathbb{1}} = 1$ , the statement is true.  $\square$

**Proposition 3.41.** *Fixed  $\mathbf{p} \in (\mathbb{N} \times \{0\}) \cup (\{0\} \times \mathbb{N})$ , any two twisted  $\mathbf{p}$ -algebras are naturally isomorphic, and any twisted  $\mathbf{p}$ -algebra is naturally anti-isomorphic to any  $\check{\mathbf{p}}$ -algebra.*

*Proof.* Let  $W_1, W_2 : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$  be symbol correspondences. Then  $W_1 \circ W_2^{-1} : \mathcal{X}_{\mathbf{p}} \rightarrow \mathcal{X}_{\mathbf{p}}$  is an isomorphism because each  $W_j$  is an isomorphism onto  $\mathcal{X}_{\mathbf{p}}$ . If, now, we suppose  $W_2 : \mathcal{B}(\mathcal{H}_{\check{\mathbf{p}}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$ , then  $W_1 \circ * \circ W_2^{-1} : \mathcal{X}_{\mathbf{p}} \rightarrow \mathcal{X}_{\check{\mathbf{p}}}$  is an anti-isomorphism since the adjoint map  $*$  is an anti-isomorphism and, again, each  $W_j$  is an isomorphism onto  $\mathcal{X}_{\mathbf{p}}$ .  $\square$

Twisted products of  $\mathbb{C}P^2$  harmonics can be easily computed and determine the twisted product for all functions in  $\mathcal{X}_{\mathbf{p}}$  by bilinearity of the product.

**Theorem 3.42.** *If  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$  is a symbol correspondence with characteristic numbers  $(c_n)$ , then the induced twister product is given by*

$$(3.71) \quad X_{\nu_1, I_1}^{n_1} \star X_{\nu_2, I_2}^{n_2} = \sqrt{\frac{(p+1)(p+2)}{2}} \sum_{n=0}^p \sum_{\nu, I} (-1)^{p+2(t_{\nu}+u_{\nu})} \begin{bmatrix} n_1 & n_2 & n \\ \nu_1, I_1 & \nu_2, I_2 & \check{\nu}, I \end{bmatrix} [p] \\ \times \frac{c_n}{c_{n_1} c_{n_2}} X_{\nu, I}^n$$

for  $0 \leq n_1, n_2 \leq p$ , where summations over  $\nu$  and  $I$  can be restricted to  $\nabla_{\nu_1+\nu_2, \nu} = 1$  and  $\delta(I_1, I_2, I) = 1$  due to (3.21).

*Proof.* The result follows from Theorems 3.10 and 3.13.  $\square$

Any twisted product on  $\mathcal{X}_{\mathbf{p}}$  admits an integral formulation, which supposedly allows one to compute it without decomposition of functions in the basis of  $\mathbb{C}P^2$  harmonics.

**Theorem 3.43.** *If  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathbb{C}P^2)$  is a symbol correspondence with operator kernel  $K$  and characteristic numbers  $(c_n)$ , then the induced twister product is given by*

$$(3.72) \quad f_1 \star f_2(\mathbf{x}) = \int_{\mathbb{C}P^2 \times \mathbb{C}P^2} f_1(\mathbf{x}_1) f_2(\mathbf{x}_2) \mathbb{L}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}) d\mathbf{x}_1 d\mathbf{x}_2$$

for any  $f_1, f_2 \in \mathcal{X}_{\mathbf{p}}$ , where

$$(3.73) \quad \mathbb{L}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) = \left( \frac{(p+1)(p+2)}{2} \right)^2 \operatorname{tr} \left( \tilde{K}(\mathbf{x}_1) \tilde{K}(\mathbf{x}_2) K(\mathbf{x}_3) \right) \\ = (-1)^p \sqrt{\frac{(p+1)(p+2)}{2}} \sum_{n_j=0}^p \sum_{\nu_j, I_j} \begin{bmatrix} n_1 & n_2 & n_3 \\ \nu_1, I_1 & \nu_2, I_2 & \nu_3, I_3 \end{bmatrix} [p] \frac{c_{n_3}}{c_{n_1} c_{n_2}} \\ \times \overline{X_{\nu_1, I_1}^{n_1}}(\mathbf{x}_1) \overline{X_{\nu_2, I_2}^{n_2}}(\mathbf{x}_2) \overline{X_{\nu_3, I_3}^{n_3}}(\mathbf{x}_3)$$

for  $\tilde{K}$  being the operator kernel dual to  $K$ , that is, with characteristic numbers  $(1/c^n)$ .

*Proof.* Let  $A_1, A_2 \in \mathcal{B}(\mathcal{H}_p)$  so that  $f_1 = W_{A_1}$  and  $f_2 = W_{A_2}$ . Then, by Proposition 3.20,

$$(3.74) \quad \begin{aligned} & \left( \frac{(p+1)(p+2)}{2} \right)^2 \int_{\mathbb{C}P^2 \times \mathbb{C}P^2} f_1(\mathbf{x}_1) f_2(\mathbf{x}_2) \operatorname{tr} \left( \tilde{K}(\mathbf{x}_1) \tilde{K}(\mathbf{x}_2) K(\mathbf{x}) \right) d\mathbf{x}_1 d\mathbf{x}_2 \\ &= \operatorname{tr} \left( \left( \frac{(p+1)(p+2)}{2} \right)^2 \int_{\mathbb{C}P^2 \times \mathbb{C}P^2} f_1(\mathbf{x}_1) f_2(\mathbf{x}_2) \tilde{K}(\mathbf{x}_1) \tilde{K}(\mathbf{x}_2) d\mathbf{x}_1 d\mathbf{x}_2 K(\mathbf{x}) \right) \\ &= \operatorname{tr}(A_1 A_2 K(\mathbf{x})) = W_{A_1 A_2}(\mathbf{x}) = f_1 \star f_2(\mathbf{x}) . \end{aligned}$$

It is worth to highlight that  $\tilde{K}(\mathbf{x}_1)$ ,  $\tilde{K}(\mathbf{x}_2)$  and  $K(\mathbf{x}_3)$  can be expanded in the coupled basis using Wigner  $D$ -functions so that  $\mathbb{L}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3)$  is a linear combinations of  $\overline{X_{\nu_1, I_1}^{n_1}}(\mathbf{x}_1) \overline{X_{\nu_2, I_2}^{n_2}}(\mathbf{x}_2) \overline{X_{\nu_3, I_3}^{n_3}}(\mathbf{x}_3)$ . That said, Theorem 3.42 implies the second equality in (3.73).  $\square$

**Definition 3.44.** *The integral trikernel  $\mathbb{L} \in C_c^\infty(\mathbb{C}P^2 \times \mathbb{C}P^2 \times \mathbb{C}P^2)$  of a twisted product induced by a symbol correspondence  $W : \mathcal{B}(\mathcal{H}_p) \rightarrow C_c^\infty(\mathbb{C}P^2)$  is the function given by (3.73) so that the twisted product is given by (3.72).*

Obviously, the integral in (3.73) is well defined for any pair of smooth functions on  $\mathbb{C}P^2$ , so it leads to a product on  $C_c^\infty(\mathbb{C}P^2)$ .

**Proposition 3.45.** *Let  $\mathbb{L}$  be the integral trikernel of a twisted product  $\star$  induced by a symbol correspondence  $W : \mathcal{B}(\mathcal{H}_p) \rightarrow C_c^\infty(\mathbb{C}P^2)$ . Then the binary operation  $\bullet$  on  $C_c^\infty(\mathbb{C}P^2)$ ,*

$$(3.75) \quad f_1 \bullet f_2(\mathbf{x}) = \int_{\mathbb{C}P^2 \times \mathbb{C}P^2} f_1(\mathbf{x}_1) f_2(\mathbf{x}_2) \mathbb{L}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}) d\mathbf{x}_1 d\mathbf{x}_2$$

for any  $f_1, f_2 \in C_c^\infty(\mathbb{C}P^2)$ , defines a  $SU(3)$ -equivariant associative  $\ast$ -algebra with respect to complex conjugation. In particular, if  $f_1, f_2 \in \mathcal{X}_p$ , we have  $f_1 \bullet f_2 = f_1 \star f_2$ . But, if either  $f_1$  or  $f_2$  is orthogonal to  $\mathcal{X}_p$ , we have  $f_1 \bullet f_2 = 0$ .

*Proof.* Linearity of integral implies the product is bilinear, hence it defines an algebra. By definition, it is clear that  $f_1 \bullet f_2 = f_1 \star f_2$  if  $f_1, f_2 \in \mathcal{X}_p$ . Now, suppose  $f_j$  is orthogonal to  $\mathcal{X}_p$ . Thus, it is orthogonal to every  $X_{\nu, I}^n$  with  $n \leq p$ , which implies the integral over  $\mathbf{x}_j$  in (3.75) results in 0, so  $f_1 \bullet f_2 = 0$ . Since any  $f \in C_c^\infty(\mathbb{C}P^2)$  can be decomposed into  $f = f_{\parallel} + f_{\perp}$ , where  $f_{\parallel} \in \mathcal{X}_p$  and  $f_{\perp}$  is orthogonal to  $\mathcal{X}_p$ , the  $SU(3)$ -equivariant, associative and  $\ast$ -algebra properties of  $\star$  extends to  $\bullet$ .  $\square$

For a product  $\bullet$  as in the previous proposition, the constant function 1 is not the identity anymore, now it gives an orthogonal projection  $C_c^\infty(\mathbb{C}P^2) \rightarrow \mathcal{X}_p : f \mapsto 1 \bullet f = f \bullet 1$ .

**Notation 5.** *Before stating general properties of integral trikernels, we establish a convention to denote the reproducing kernel on  $\mathcal{X}_p$ :*

$$(3.76) \quad \mathcal{R}_p(\mathbf{x}_1, \mathbf{x}_2) = \sum_{n=0}^p \sum_{\nu, I} \overline{X_{\nu, I}^n}(\mathbf{x}_1) X_{\nu, I}^n(\mathbf{x}_2) = \mathcal{R}_p(\mathbf{x}_2, \mathbf{x}_1) ,$$

satisfying

$$(3.77) \quad \int_{\mathbb{C}P^2} f(\mathbf{x}_1) \mathcal{R}_p(\mathbf{x}_1, \mathbf{x}_2) d\mathbf{x}_1 = f(\mathbf{x}_2) , \quad \forall f \in \mathcal{X}_p .$$

**Proposition 3.46.** *Let  $\mathbb{L}$  be an integral trikernell of a twisted product  $\star$  on  $\mathcal{X}_p$ . Then, for every  $g \in SU(3)$  and every  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \mathbf{x}_4 \in \mathbb{C}P^2$ ,*

- i)  $\mathbb{L}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) = \mathbb{L}(g\mathbf{x}_1, g\mathbf{x}_2, g\mathbf{x}_3)$ ;
- ii)  $\int_{\mathbb{C}P^2} \mathbb{L}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}) \mathbb{L}(\mathbf{x}, \mathbf{x}_3, \mathbf{x}_4) d\mathbf{x} = \int_{\mathbb{C}P^2} \mathbb{L}(\mathbf{x}_1, \mathbf{x}, \mathbf{x}_4) \mathbb{L}(\mathbf{x}_2, \mathbf{x}_3, \mathbf{x}) d\mathbf{x}$ ;
- iii)  $\int_{\mathbb{C}P^2} \mathbb{L}(\mathbf{x}, \mathbf{x}_1, \mathbf{x}_2) d\mathbf{x} = \int_{\mathbb{C}P^2} \mathbb{L}(\mathbf{x}_1, \mathbf{x}, \mathbf{x}_2) d\mathbf{x} = \mathcal{R}_p(\mathbf{x}_1, \mathbf{x}_2)$ ;
- iv)  $\overline{\mathbb{L}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3)} = \mathbb{L}(\mathbf{x}_2, \mathbf{x}_1, \mathbf{x}_3)$ .

*Proof.* Let  $f_1, f_2 \in \mathcal{X}_p$ . Writing the equality  $(f_1)^g \star (f_2)^g = (f_1 \star f_2)^g$  in the integral form, we get that  $SU(3)$ -equivariance of  $\star$  is equivalent to property (i). In the same vein, we conclude that each property of this statement is equivalent to the property of Proposition 3.40 with same number.  $\square$

**Remark 3.47.** *Although the integral formulation of a twisted product on  $\mathcal{X}_p$  is supposed to circumvent the necessity of decomposing symbols (elements of  $\mathcal{X}_p$ ) in the basis of  $\mathbb{C}P^2$  harmonics, the formula (3.73) for an integral trikernell uses these harmonics explicitly. In [18], new formulas for integral trikernells of spin systems were obtained using  $SU(2)$ -invariant 2-points and 3-points functions on  $\mathbb{C}P^1$ , but a similar exercise in the case of pure-quark systems is much harder and is deferred for later.*

We finish with a relation between twisted algebras induced by antipodal correspondences.

**Proposition 3.48.** *The twisted products  $\star$  and  $\check{\star}$  induced by a symbol correspondence and its antipodal correspondence satisfy*

$$(3.78) \quad f_1 \star f_2 = f_2 \check{\star} f_1 .$$

*Proof.* For  $f_1 = W_{F_1} = \check{W}_{F_1^*}$  and  $f_2 = W_{F_2} = \check{W}_{F_2^*}$ ,

$$(3.79) \quad f_1 \star f_2 = W_{F_1 F_2} = \check{W}_{(F_1 F_2)^*} = \check{W}_{F_2^* F_1^*} = f_2 \check{\star} f_1 .$$

$\square$

**Corollary 3.49.** *For  $\star$  and  $\check{\star}$  as in the previous proposition, their integral trikernells  $\mathbb{L}$  and  $\check{\mathbb{L}}$  satisfy*

$$(3.80) \quad \mathbb{L}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) = \check{\mathbb{L}}(\mathbf{x}_2, \mathbf{x}_1, \mathbf{x}_3) .$$

**Remark 3.50.** *We already mentioned that the notion of antipodal correspondences for quark systems is analogous to alternation for spin systems considering the appropriate characterization. In addition to the previous discussion, we present a related phenomenon encoded in Proposition 3.48 which also happens for spin systems. The commutator  $[\cdot, \cdot]_{\star}$  of a twisted product  $\star$  satisfies*

$$(3.81) \quad [f_1, f_2]_{\star} = [f_2, f_1]_{\check{\star}} ,$$

where  $\check{\star}$  is the twisted product induced by the antipodal correspondence. In this way,  $\check{\star}$  can be seen as defining the reverse symbolic dynamics of the one defined by  $\star$ . In fact, recalling Heisenberg's equation for an operator  $F$  subject to a Hamiltonian  $H$ ,

$$(3.82) \quad \frac{dF}{dt} = [F, H] + \frac{\partial F}{\partial t} ,$$

if  $F$  has no explicit temporal dependence, then under a symbol correspondence  $W$  its symbol  $f$  satisfies

$$(3.83) \quad \frac{df}{dt} = [f, h]_{\star} ,$$

where  $h = W_H$ . It follows that if we set  $H^*$  as the Hamiltonian of the dual space, the symbolic dynamics of  $F^*$  under  $\widetilde{W}$  is given by

$$(3.84) \quad \frac{df}{dt} = [f, h]_{\star}^{\vee} = -[f, h]_{\star} .$$

#### 4. GENERIC QUARK SYSTEMS

In this chapter, we begin a study of correspondences for generic quark systems, that is, representations of generic class  $Q(p, q)$  and generic coadjoint orbit  $\mathcal{E}(\mathbb{C}P^2, \mathbb{C}P^1, \pi)$ . Although we basically reproduce what we have done for  $Q(p, 0)$  (or  $Q(0, q)$ ) and  $\mathbb{C}P^2$ , some new phenomena appear when compared to the previous case.

##### 4.1. Classical generic quark system.

**Definition 4.1.** *The generic classical quark system is the symplectic total space  $\mathcal{E}$  of the fiber bundle  $\mathcal{E}(\mathbb{C}P^2, \mathbb{C}P^1, \pi)$ , with base  $\mathbb{C}P^2$ , fiber  $\mathbb{C}P^1$  and projection  $\pi$ ,*

$$(4.1) \quad \mathbb{C}P^1 \hookrightarrow \mathcal{E} \xrightarrow{\pi} \mathbb{C}P^2 ,$$

together with its Poisson algebra on  $C_c^\infty(\mathcal{E})$ .

We have  $\mathcal{E} \simeq SU(3)/T$ , where  $T$  is the maximal torus (2.159) of  $SU(3)$ . So we look for representations with weights satisfying  $t = u = 0$ , cf. (2.21)-(2.22).

**Proposition 4.2.** *The representations of  $SU(3)$  with non null vectors fixed by  $T$  are of the form  $Q(a, b)$  for  $a \equiv b \pmod{3}$ . For*

$$k = |a - b|/3 ,$$

the space fixed by  $T$  is spanned by the set

$$(4.2) \quad \{ \mathbf{e}((a, b); \boldsymbol{\nu}_{(a,b)}, I_\gamma) : \gamma = 1, \dots, \min\{a, b\} + 1 \} ,$$

where

$$(4.3) \quad \boldsymbol{\nu}_{(a,b)} = \begin{cases} (a + 2k, a + 2k, a + 2k) & , \quad \text{if } \min\{a, b\} = a \\ (b + k, b + k, b + k) & , \quad \text{if } \min\{a, b\} = b \end{cases}$$

and

$$(4.4) \quad I_\gamma = \gamma - 1 + k .$$

*Proof.* From (2.21)-(2.22), we get that  $\nu_1 = \nu_2 = \nu_3 = \nu \in \mathbb{N}_0$ , with

$$(4.5) \quad 2\nu = r_+ + r_-$$

and

$$(4.6) \quad 3\nu = a + 2b ,$$

for

$$r_- \leq b \leq r_+ \leq a + b , \quad r_- \leq \nu \leq a + b .$$

From (4.6),  $a + 2b \equiv 0 \pmod{3}$ , which implies  $a \equiv b \pmod{3}$ .

For representations of class  $Q(a, a + 3k)$ , with  $a, k \in \mathbb{N}_0$ , we have that

$$(4.7) \quad \nu = a + 2k$$

and the GT states fixed by  $T$  are given by  $r_+$  and  $r_-$  satisfying

$$(4.8) \quad \begin{cases} r_+ + r_- = 2a + 4k \\ 0 \leq r_- \leq a + 3k \leq r_+ \leq 2a + 3k \end{cases} .$$

The system has  $a + 1$  solutions:

$$(4.9) \quad \begin{cases} r_+ = 2a + 3k, & r_- = k \\ \vdots \\ r_+ = a + 3k, & r_- = a + k \end{cases}$$

so that

$$(4.10) \quad \{e((a, a + 3k); (a + 2k, a + 2k, a + 2k), I) : I = k, \dots, a + k\}$$

spans the subspace fixed by  $T$ .

For a representation of class  $Q(b + 3k, b)$ , since it is dual to  $Q(b, b + 3k)$ , the subspace fixed by  $T$  is spanned by the dualization of the set  $\{e((b, b + 3k), (b + 2k, b + 2k, b + 2k), I) : I = k, \dots, b + k\}$ , which implies that

$$\nu = b + k$$

and is given by the solutions of

$$(4.11) \quad \begin{cases} r_+ + r_- = 2b + 2k \\ 0 \leq r_- \leq a \leq r_+ \leq 2b + 3k \end{cases} .$$

The same reasoning of the previous case leads us to

$$(4.12) \quad \begin{cases} r_+ = 2b + 2k, & r_- = 0 \\ \vdots \\ r_+ = b + 2k, & r_- = b \end{cases} .$$

That is,

$$(4.13) \quad \{e((b + 3k, b); (b + k, b + k, b + k), I) : I = k, \dots, b + k\}$$

spans the subspace fixed by  $T$ .

To finish, we order the  $I$ -multiplicities by crescent  $I$  in both cases (4.10)-(4.13) by setting  $I_\gamma = \gamma + k - 1$ , where  $1 \leq \gamma \leq \min\{a, b\} + 1$ .  $\square$

**Definition 4.3.** *The  $\mathcal{E}$  harmonics are the functions*

$$(4.14) \quad Z_{\nu, I}^{(a, b, \gamma)}(gz_0) = \sqrt{\frac{(a + 1)(b + 1)(a + b + 2)}{2}} \overline{D_{\nu I, \nu_{(a, b)} I_\gamma}^{(a, b)}}(g)$$

on  $\mathcal{E}$ , for  $\mathbf{z}_0 \in \pi^{-1}([0 : 0 : 1]) \subset \mathcal{E}$  with  $T$  as isotropy subgroup,  $g \in SU(3)$ ,  $a \equiv b \pmod{3}$ ,  $(\nu_{(a, b)}, I_\gamma)$  as in Proposition 4.2 and  $D_{\nu I, \nu_{(a, b)} I_\gamma}^{(a, b)}$  a Wigner  $D$ -function as in Definition 2.5.

Just as in definition of  $\mathbb{C}P^2$  harmonics, the factor  $\sqrt{(a+1)(b+1)(a+b+2)/2}$  is the square root of the dimension of the representation  $Q(a, b)$  and is used to ensure normalization according to Schur's Orthogonality Relations, so that

$$(4.15) \quad \left\langle Z_{\nu, I}^{(a, b, \gamma)} \middle| Z_{\mu, J}^{(c, d, \zeta)} \right\rangle = \delta_{a, c} \delta_{b, d} \delta_{\gamma, \zeta} \delta_{\nu, \mu} \delta_{I, J} .$$

**Remark 4.4.** Analogously to Remark 3.4, for any  $x, y > 0$ , we can take the generic harmonics as functions on  $\mathcal{O}_{x, y}$  via the compositions  $Z_{\nu, I}^{(a, b, \gamma)} \circ \psi_{x, y}$  so that the harmonic functions on  $\mathcal{O}_{x, y}$  are related to the ones on  $\mathcal{O}_{y, x}$  by  $\alpha_{x, y} \circ \iota$ , cf. (2.168). Besides that, the involution  $\alpha_{x, y}$  generates another, but somewhat equivalent, set of harmonic functions on  $\mathcal{O}_{x, y}$ , just as  $\alpha$  does to  $\mathcal{E}$ :

$$(4.16) \quad \tilde{Z}_{\nu, I}^{(a, b, \gamma)}(g\mathbf{z}_0) = Z_{\nu, I}^{(a, b, \gamma)}(gg_0\mathbf{z}_0) ,$$

cf. (2.167).

As expected, we have

$$(4.17) \quad Z_{(0,0,0),0}^{(0,0)} \equiv 1$$

and, cf. (2.38),

$$(4.18) \quad \overline{Z_{\nu, I}^{(a, b, \gamma)}} = (-1)^{2(t+u)} Z_{\tilde{\nu}, \tilde{I}}^{(b, a, \gamma)} , \text{ for } \Delta_{\nu, \tilde{\nu}}^{a+b} = 1 .$$

**Theorem 4.5.** The decomposition of pointwise product of  $\mathcal{E}$  harmonics is given by

$$(4.19) \quad Z_{\nu_1, I_1}^{(\mathbf{a}_1, \gamma_1)} Z_{\nu_2, I_2}^{(\mathbf{a}_2, \gamma_2)} = \sum_{\substack{(\mathbf{a}; \sigma) \\ \nu, I, \gamma}} \sqrt{\frac{\dim Q(\mathbf{a}_1) \dim Q(\mathbf{a}_2)}{\dim Q(\mathbf{a})}} C_{\nu_1 I_1, \nu_2 I_2, \nu I}^{\mathbf{a}_1, \mathbf{a}_2, (\mathbf{a}; \sigma)} \\ \times C_{\nu_{\mathbf{a}_1} I_{\gamma_1}, \nu_{\mathbf{a}_2} I_{\gamma_2}, \nu_{\mathbf{a}} I_{\gamma}}^{\mathbf{a}_1, \mathbf{a}_2, (\mathbf{a}; \sigma)} Z_{\nu, I}^{(\mathbf{a}, \gamma)} ,$$

for  $(\nu_{\mathbf{a}_j}, I_{\gamma_j})$  and  $(\nu_{\mathbf{a}}, I_{\gamma})$  as in Proposition 4.2, and summation restricted to  $\nabla_{\nu_1 + \nu_2, \nu} = 1$ ,  $\delta(I_1, I_2, I) = \delta(I_{\gamma_1}, I_{\gamma_2}, I_{\gamma}) = 1$  and  $Q(\mathbf{a}; \sigma)$  in the Clebsch-Gordan series of  $Q(\mathbf{a}_1) \otimes Q(\mathbf{a}_2)$ .

*Proof.* With a little abuse of notation, again,

$$(4.20) \quad Z_{\nu_j, I_j}^{(\mathbf{a}_j, \gamma_j)} = \sqrt{\dim Q(\mathbf{a}_j)} \overline{D_{\nu_j I_j, \nu_{\mathbf{a}_j} I_{\gamma_j}}^{\mathbf{a}_j}}$$

and Lemma 2.13 give us

$$(4.21) \quad Z_{\nu_1, I_1}^{(\mathbf{a}_1, \gamma_1)} Z_{\nu_2, I_2}^{(\mathbf{a}_2, \gamma_2)} = \sum_{(\mathbf{a}; \sigma)} \sum_{\substack{\nu, I \\ \mu, J}} \sqrt{\dim Q(\mathbf{a}_1) \dim Q(\mathbf{a}_2)} C_{\nu_1 I_1, \nu_2 I_2, \nu I}^{\mathbf{a}_1, \mathbf{a}_2, (\mathbf{a}; \sigma)} \\ \times C_{\nu_{\mathbf{a}_1} I_{\gamma_1}, \nu_{\mathbf{a}_2} I_{\gamma_2}, \mu J}^{\mathbf{a}_1, \mathbf{a}_2, (\mathbf{a}; \sigma)} \overline{D_{\nu I, \mu J}^{\mathbf{a}}} ,$$

where  $\nabla_{\nu_1 + \nu_2, \nu} = \nabla_{\nu_{\mathbf{a}_1} + \nu_{\mathbf{a}_2}, \mu} = 1$  and  $\delta(I_1, I_2, I) = \delta(I_{\gamma_1}, I_{\gamma_2}, J) = 1$ , so  $\mu = (\mu, \mu, \mu)$ . But  $\mathbf{e}(\mathbf{a}; (\mu, \mu, \mu), J)$  only exists if  $\mathbf{a}$  and  $(\mu, J)$  are as in Proposition 4.2. Thus, we set  $\mu = \nu_{\mathbf{a}}$  and  $J = I_{\gamma}$ .  $\square$

**Remark 4.6.** As in the decomposition of the pointwise product of  $\mathbb{C}P^2$  harmonics, the decomposition of the pointwise product of  $\mathcal{E}$  harmonics follows directly as a special case of Lemma 2.13 and does not “see” the symplectic structure on  $\mathcal{E}$ . Thus, as in the pure-quark case, the next step is to decompose the Poisson bracket of  $\mathcal{E}$  harmonics, but this is a much harder problem that is deferred to a later study.

**4.2. Quantum generic quark system.** Now, we want representations  $Q(p, q)$  such that  $Q(p, q) \otimes Q(q, p)$  splits only into representations of the form  $Q(a, b)$ ,  $a \equiv b \pmod{3}$ , with multiplicity less than or equal to  $\min\{a, b\} + 1$ . From Corollary 2.7, if we suppose, without loss of generality, that  $\min\{a, b\} = a$ , then the occurrences of  $Q(a, b) \oplus Q(b, a)$  are given by the solutions of

$$(4.22) \quad \begin{cases} a = p + q - n - m - 2k \\ 0 \leq n \leq p - k \\ 0 \leq m \leq q - k \end{cases},$$

where  $b = a + 3k$ . Of course, we can also assume without loss of generality that  $p \geq q$ . If  $a + k \leq q$ , then we have  $a + 1$  solutions:

$$(4.23) \quad \begin{cases} n = p - k - a, & m = q - k \\ n = p - k - a + 1, & m = q - k - 1 \\ \vdots \\ n = p - k, & m = q - a - k \end{cases}.$$

Otherwise, we need to eliminate some lines of the above solutions, which means  $Q(a, b) \oplus Q(b, a)$  have multiplicity less than  $\min\{a, b\} + 1$ . Then, we are able to define:

**Definition 4.7.** Let<sup>20</sup>  $(p, q) \in (\mathbb{N} \times \mathbb{N}_0) \cup (\mathbb{N}_0 \times \mathbb{N})$ . A quantum generic quark system is a complex Hilbert space  $\mathcal{H}_{p,q} \simeq \mathbb{C}^d$ , where

$$d = \frac{(p+1)(q+1)(p+q+2)}{2},$$

with an irreducible unitary  $SU(3)$ -representation of class  $Q(p, q)$  together with its operator algebra  $\mathcal{B}(\mathcal{H}_{p,q})$ .

If  $p \geq q$ , the pair  $(p, q)$  and the system  $\mathcal{H}_{p,q}$  are called material. If  $p > q$ , they are called baryonic and if  $p = q$  they are called mesonic. Alternatively, if  $p < q$ , they are called antibaryonic.

**Remark 4.8.** The name material refers to a system composed of a larger (or equal) number of quarks than antiquarks. From Theorem 2.6,

$$(4.24) \quad Q(p, 0) \otimes Q(0, q) = \bigoplus_{n=0}^{\min\{p,q\}} Q(p-n, q-n),$$

so a generic representation  $Q(p, q)$  is the invariant space of  $Q(p, 0) \otimes Q(0, q)$  where the product of the highest weight vectors lives in. That means material quark systems can be constructed from systems with number of quarks greater than or equal to number of antiquarks.

The names baryonic and mesonic make reference to systems with positive and null baryon number, respectively. We recall that a system of  $p$  quarks and  $q$  antiquarks has baryon number

$$(4.25) \quad B = \frac{1}{3}(p - q).$$

If  $B > 0$ , the system is classified as a baryon; if  $B < 0$  it is an antibaryon; if  $B = 0$ , we have a meson.

<sup>20</sup>Again, we are ignoring the trivial representation  $Q(0, 0)$ .

In particular, quantum generic quark systems encompass quantum pure-quark systems as special cases. But now, all forms of (2.149) and (2.150) are relevant to us, and we cannot further simplify Corollary 2.29 as we did for pure-quark systems.

**4.3. Symbol correspondences for generic quark systems.** Let  $\mathbf{p} \in (\mathbb{N} \times \mathbb{N}_0) \cup (\mathbb{N}_0 \times \mathbb{N})$ .

The following is completely analogous to Definition 3.11.

**Definition 4.9.** A symbol correspondence for a generic quark system  $(\mathcal{H}_{\mathbf{p}}, Q(\mathbf{p}))$ , also referred to simply as a symbol correspondence or just as a correspondence, is an injective linear map  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathcal{E}) : P \mapsto W_P$  that satisfies, for any  $A \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$ ,

- i) *Equivariance:*  $\forall g \in SU(3), W_{A^g} = (W_A)^g$ ;
- ii) *Reality:*  $W_{A^\dagger} = \overline{W_A}$ ;
- iii) *Normalization:*  $\int_{\mathcal{E}} W_A(\mathbf{z}) d\mathbf{z} = \frac{1}{\dim Q(\mathbf{p})} \text{tr}(A)$ .

**Remark 4.10.** In the spirit of Remark 3.12, one can replace  $\mathcal{E}$  by  $\mathcal{O}_{x,y}$  for any  $x, y > 0$  using the diffeomorphism  $\psi_{x,y}$ .

**Notation 6.** Recalling the notations

$$\mathbf{a} = (a, b), \quad \mathbf{p} = (p, q) \iff \check{\mathbf{p}} = (q, p),$$

from now on, we shall use the notation

$$(4.26) \quad m(\mathbf{a}) = m(a, b) = \min\{a, b\} + 1.$$

and simplify the notation  $\mathbf{m}(\mathbf{p}, \check{\mathbf{p}}; \mathbf{a})$  for the multiplicity of  $Q(\mathbf{a}) = Q(a, b)$  in the Clebsch-Gordan series of  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}}) = Q(p, q) \otimes Q(q, p)$  by setting

$$(4.27) \quad \mathbf{m}(\mathbf{p}; \mathbf{a}) := \mathbf{m}(\mathbf{p}, \check{\mathbf{p}}; \mathbf{a}).$$

Finally, we set

$$(4.28) \quad \mathcal{B}(\mathbf{p}; \mathbf{a}) = \bigoplus_{\sigma=1}^{\mathbf{m}(\mathbf{p}; \mathbf{a})} Q(\mathbf{a}; \sigma) \subset \mathcal{B}(\mathcal{H}_{\mathbf{p}}).$$

**Theorem 4.11.** A linear map  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathcal{E}) : A \mapsto W_A$  is a symbol correspondence if and only if, for each  $Q(\mathbf{a}; \sigma)$  in  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}})$ , it maps

$$(4.29) \quad W : \sqrt{\dim Q(\mathbf{p})} e((\mathbf{a}; \sigma); \boldsymbol{\nu}, I) \mapsto \sum_{\gamma=1}^{m(\mathbf{a})} c_{\gamma}^{\sigma}(\mathbf{a}) Z_{\boldsymbol{\nu}, I}^{(\mathbf{a}; \gamma)} =: Z\mathbf{C}(\mathbf{a})_{\boldsymbol{\nu}, I}^{\sigma},$$

where  $c_{\gamma}^{\sigma}(\mathbf{a})$  is the  $\gamma \times \sigma$  entry of a complex matrix of order  $m(\mathbf{a}) \times \mathbf{m}(\mathbf{p}; \mathbf{a})$  denoted by  $\mathbf{C}(\mathbf{a})$ , that is,

$$(4.30) \quad \mathbf{C}(\mathbf{a}) = [c_{\gamma}^{\sigma}(\mathbf{a})],$$

with  $\mathbf{C}(\mathbf{a})$  being of full rank and satisfying  $\overline{\mathbf{C}(\mathbf{a})} = \mathbf{C}(\check{\mathbf{a}})$  and  $\mathbf{C}(0, 0) = (-1)^{|\mathbf{p}|}$ .

*Proof.* Since  $W$  is injective and equivariant, the image of  $Q(\mathbf{a}; \sigma)$  is a representation isomorphic to  $Q(\mathbf{a})$ . Let  $\{f((\mathbf{a}; \sigma); \boldsymbol{\nu}, I)\}$  be a GT basis of the image of  $Q(\mathbf{a}; \sigma)$ , so that

$$(4.31) \quad W : \sqrt{\dim Q(\mathbf{p})} e((\mathbf{a}; \sigma); \boldsymbol{\nu}, I) \mapsto \alpha_{(\mathbf{a}; \sigma)} f((\mathbf{a}; \sigma); \boldsymbol{\nu}, I), \quad \alpha_{(\mathbf{a}; \sigma)} \neq 0.$$

Because the multiplicity of  $Q(\mathbf{a})$  in  $C_{\mathbb{C}}^{\infty}(\mathcal{E})$  is  $m(\mathbf{a})$ , cf. (4.2), we must have

$$(4.32) \quad f((\mathbf{a}; \sigma); \boldsymbol{\nu}, I) = \sum_{\gamma=1}^{m(\mathbf{a})} \beta_{\gamma}^{(\mathbf{a}; \sigma)} Z_{\boldsymbol{\nu}, I}^{(\mathbf{a}, \gamma)},$$

where  $Z_{\boldsymbol{\nu}, I}^{(\mathbf{a}, \gamma)}$  are the  $\mathcal{E}$  harmonics, cf. Definition 4.3, which implies in particular that

$$(4.33) \quad \sum_{\gamma=1}^{m(\mathbf{a})} \left| \beta_{\gamma}^{(\mathbf{a}; \sigma)} \right|^2 = \|f((\mathbf{a}; \sigma); \boldsymbol{\nu}, I)\|^2 = 1.$$

Let

$$(4.34) \quad c_{\gamma}^{\sigma}(\mathbf{a}) = \alpha_{(\mathbf{a}; \sigma)} \beta_{\gamma}^{(\mathbf{a}; \sigma)}.$$

The injection hypothesis implies that the union of basis

$$\bigcup_{\sigma=1}^{\mathbf{m}(\mathbf{p}; \mathbf{a})} \{f((\mathbf{a}; \sigma); \boldsymbol{\nu}, I)\}$$

is a linearly independent set, hence  $\{(\beta_1^{(\mathbf{a}; \sigma)}, \dots, \beta_{m(\mathbf{a})}^{(\mathbf{a}; \sigma)}) : \sigma = 1, \dots, \mathbf{m}(\mathbf{p}; \mathbf{a})\}$  is a linearly independent set in  $\mathbb{C}^{m(\mathbf{a})}$ , cf. (4.32). This means that  $\{(c_1^{\sigma}(\mathbf{a}), \dots, c_{m(\mathbf{a})}^{\sigma}(\mathbf{a})) : \sigma = 1, \dots, \mathbf{m}(\mathbf{p}; \mathbf{a})\}$  is a linearly independent set too, cf. (4.34), so the complex matrix  $\mathbf{C}(\mathbf{a})$  whose  $\gamma \times \sigma$  entry is  $c_{\gamma}^{\sigma}(\mathbf{a})$  is of full rank.

We have that  $e^{\dagger}((\mathbf{a}; \sigma); \boldsymbol{\nu}, I) = (-1)^{2(t+u)} e((\check{\mathbf{a}}; \sigma); \check{\boldsymbol{\nu}}, I)$ , cf. (2.125), and also that  $\overline{Z_{\boldsymbol{\nu}, I}^{(\mathbf{a}, \gamma)}} = (-1)^{2(t+u)} Z_{\check{\boldsymbol{\nu}}, I}^{(\check{\mathbf{a}}, \gamma)}$ , cf. (4.18), so the reality condition implies

$$(4.35) \quad \overline{c_{\gamma}^{\sigma}(\mathbf{a})} = c_{\gamma}^{\sigma}(\check{\mathbf{a}}),$$

or in a concise form, the matrices  $\mathbf{C}(\mathbf{a})$  satisfy  $\overline{\mathbf{C}(\mathbf{a})} = \mathbf{C}(\check{\mathbf{a}})$ .

The normalization property implies

$$(4.36) \quad W : (-1)^{|\mathbf{p}|} \sqrt{\dim Q(p, q)} e((0, 0); (0, 0, 0), 0) \mapsto Z_{(0, 0, 0), 0}^{(0, 0)},$$

cf. (2.126) and (4.17), hence  $\mathbf{C}(0, 0) = (-1)^{|\mathbf{p}|}$ .

It is more straightforward to prove the converse, that is, to check that a map with these properties is a symbol correspondence, so we leave this to the reader.  $\square$

**Corollary 4.12.** *The moduli space  $\mathbf{S}_{\mathbf{p}}$  of correspondences for a generic quark system  $\mathcal{H}_{\mathbf{p}}$  can be described as*

$$(4.37) \quad \mathbf{S}_{\mathbf{p}} = \left( \prod_{a=0}^{|\mathbf{p}|} V_{\mathbf{m}(\mathbf{p}; a, a)}(\mathbb{R}^{a+1}) \right) \times \left( \prod_{a < b} V_{\mathbf{m}(\mathbf{p}; a, b)}(\mathbb{C}^{a+1}) \right),$$

where  $V_k(\mathbb{K}^n) = GL_n(\mathbb{K})/GL_{n, k}(\mathbb{K})$ , for  $GL_{n, k}(\mathbb{K}) \subset GL_n(\mathbb{K})$  a maximal subgroup that fixes a  $k$ -dimensional subspace, is a non compact Stiefel manifold.

The matrices  $\mathbf{C}(\mathbf{a})$  are matrix representations of the maps  $W|_{\mathcal{B}(\mathbf{p}; \mathbf{a})}$  with respect to a coupled basis of  $\mathcal{B}(\mathcal{H}_{\mathbf{p}})$  and the  $\mathcal{E}$  harmonics. They are analogous to characteristic numbers of symbol correspondences for pure-quark system: in the latter case, the domain and codomain of a symbol correspondence are multiplicity free and have only representations  $Q(n, n)$ , so it provides a  $1 \times 1$  real matrix indexed by  $n$ . The moduli space in that case is a product of  $V_1(\mathbb{R}) = \mathbb{R}^*$ .

We now prove a theorem analogous to Theorem 3.15.

As usual, let  $\mathbf{z}_0 \in \pi^{-1}([0 : 0 : 1]) \subset \mathcal{E}$  be a point with  $T$  as isotropy subgroup. Now, for an operator  $K \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$  fixed by  $T$ , we have  $\mathcal{E} \rightarrow \mathcal{B}(\mathcal{H}_{\mathbf{p}}) : \mathbf{z} \mapsto K(\mathbf{z}) = K^g$ , where  $g \in SU(3)$  is such that  $\mathbf{z} = g\mathbf{z}_0$ .

**Theorem 4.13.** *A map  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathcal{E}) : A \mapsto W_A$  is a symbol correspondence satisfying (4.29) if and only if*

$$(4.38) \quad W_A(\mathbf{z}) = \text{tr}(AK(\mathbf{z})) ,$$

that is,

$$(4.39) \quad W_A(g\mathbf{z}_0) = \text{tr}(AK^g) ,$$

for  $K \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$  of the form

$$(4.40) \quad K = \frac{1}{\dim Q(\mathbf{p})} \mathbb{1} + \sum_{(\mathbf{a}; \sigma)} \sum_{\gamma=1}^{m(\mathbf{a})} \frac{c_{\gamma}^{\sigma}(\mathbf{a})}{c_{\gamma}^{\sigma}(\mathbf{a})} \sqrt{\frac{\dim Q(\mathbf{a})}{\dim Q(\mathbf{p})}} e((\mathbf{a}; \sigma); \nu_{\mathbf{a}}, I_{\gamma}) ,$$

with  $c_{\gamma}^{\sigma}(\mathbf{a}) = [\mathfrak{C}(\mathbf{a})]_{\gamma}^{\sigma}$  as in Theorem 4.11, where the summation is over all  $(\mathbf{a}; \sigma)$  in the CG series of  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}})$ . In particular,  $K$  is Hermitian with unitary trace.

*Proof.* Assuming  $W$  is a symbol correspondence, we can reproduce the proof of Theorem 3.15 to conclude  $W_A(g\mathbf{z}_0) = \text{tr}(AK^g)$ , where  $K$  is a linear combination of the vectors fixed by  $T$ , so

$$(4.41) \quad K = \sum_{\substack{(\mathbf{a}; \sigma) \\ \gamma}} k_{\gamma}^{(\mathbf{a}; \sigma)} e((\mathbf{a}; \sigma); \nu_{\mathbf{a}}, I_{\gamma}) ,$$

cf. Proposition 4.2.

For  $A = e((\mathbf{a}; \sigma); \nu, I) = (-1)^{2(t+u)} e^{\dagger}((\check{\mathbf{a}}; \sigma); \check{\nu}, I)$  we get

$$(4.42) \quad W_A(g\mathbf{z}_0) = \text{tr}(AK^g) = \sum_{\gamma=1}^{m(\mathbf{a})} k_{\gamma}^{(\check{\mathbf{a}}; \sigma)} (-1)^{2(t+u)} D_{\check{\nu}I, \nu_{\check{\mathbf{a}}}I_{\gamma}}^{\check{\mathbf{a}}} (g) = \sum_{\gamma=1}^{m(\mathbf{a})} k_{\gamma}^{(\check{\mathbf{a}}; \sigma)} \overline{D_{\nu I, \nu_{\mathbf{a}}}^{\mathbf{a}}}$$

cf (2.38). Then,

$$(4.43) \quad W_A = \sum_{\gamma=1}^{m(\mathbf{a})} \frac{k_{\gamma}^{(\check{\mathbf{a}}; \sigma)}}{\sqrt{\dim Q(\mathbf{a})}} Z_{\nu, I}^{(\mathbf{a}, \gamma)} .$$

It follows from Theorem 4.11 that

$$(4.44) \quad k_{\gamma}^{(\check{\mathbf{a}}; \sigma)} = c_{\gamma}^{\sigma}(\mathbf{a}) \sqrt{\frac{\dim Q(\mathbf{a})}{\dim Q(\mathbf{p})}} \iff k_{\gamma}^{(\mathbf{a}; \sigma)} = c_{\gamma}^{\sigma}(\check{\mathbf{a}}) \sqrt{\frac{\dim Q(\check{\mathbf{a}})}{\dim Q(\mathbf{p})}} .$$

Using (4.35) and  $\dim Q(\mathbf{a}) = \dim Q(\check{\mathbf{a}})$ , we obtain (4.40).

The Hermitian property of  $K$  follows from (4.35) and (2.125) plus the fact that, if  $Q(\mathbf{a}; \sigma)$  is in the CG series of  $Q(\mathbf{p}) \otimes Q(\check{\mathbf{p}})$ , then so is  $Q(\check{\mathbf{a}}; \sigma)$ . Unitary trace for  $K$  is immediate from the fact that every  $e((\mathbf{a}; \sigma); \nu_{\mathbf{a}}, I_{\gamma})$  is traceless (orthogonal to  $\mathbb{1}$ ).

The converse is, again, analogous to Theorem 3.15, and it is rather straightforward to verify that, for  $K$  given by (4.40), equations (4.41)-(4.43) imply that (4.39) defines a symbol correspondence given by (4.29).  $\square$

**Definition 4.14.** Any  $K \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$  that induces a symbol correspondence via (4.39) is an operator kernel. Thus,  $K$  is given by (4.40), where the numbers  $(c_{\gamma}^{\sigma}(\mathbf{a}))$  are called characteristic parameters and the matrices  $\mathbf{C}(\mathbf{a})$  with  $c_{\gamma}^{\sigma}(\mathbf{a})$  in the  $\gamma \times \sigma$  entry, cf. (4.30), are the characteristic matrices of both the operator kernel and the symbol correspondence.

**Remark 4.15.** It is worth to explicitly point out that an operator kernel  $K$  of a symbol correspondence for a pure quark system  $\mathcal{H}_{p,0}$  (or, equivalently,  $\mathcal{H}_{0,p}$ ) is also an operator kernel of a symbol correspondence for  $\mathcal{H}_{p,0}$  taken as a generic quark system. If  $K$  has characteristic numbers  $(c_n)$ , it has characteristic parameters  $(c_{\gamma}^{\perp}(n, n))$  given by  $c_{\gamma}^{\perp}(n, n) = c_n \delta_{\gamma,1}$ .

Now, in the case of generic quark systems, we cannot unambiguously define symbol correspondences in a implicit way as we did in Proposition 3.18 because characteristic matrices may have more than one left inverse.

**Proposition 4.16.** Let  $K \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$  be an operator kernel with characteristic matrices  $\mathbf{C}(\mathbf{a})$ . A symbol correspondence  $\widetilde{W} : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C^{\infty}(\mathcal{E})$  satisfies

$$(4.45) \quad A = \dim Q(\mathbf{p}) \int_{\mathcal{E}} \widetilde{W}_A(\mathbf{z}) K(\mathbf{z}) d\mathbf{z}$$

if and only if it has characteristic matrices  $\widetilde{\mathbf{C}}(\mathbf{a})$  such that  $(\widetilde{\mathbf{C}}(\mathbf{a}))^{\dagger} \mathbf{C}(\mathbf{a}) = \mathbf{1}$ .

*Proof.* By straightforward calculation, we get

$$(4.46) \quad \begin{aligned} \int_{\mathcal{E}} Z_{\nu, I}^{(\mathbf{a}, \gamma)}(\mathbf{z}) K(\mathbf{z}) d\mathbf{z} &= \int_{SU(3)} Z_{\nu, I}^{(\mathbf{a}, \gamma)}(g\mathbf{z}_0) K^g dg \\ &= \sum_{\substack{(\mathbf{a}'; \sigma') \\ \gamma', J}} k_{\gamma'}^{(\mathbf{a}'; \sigma')} \int_{SU(3)} Z_{\nu, I}^{(\mathbf{a}, \gamma)}(g\mathbf{z}_0) D_{\mu J, \nu \mathbf{a}' I, \gamma'}^{\mathbf{a}'}(g) dg e((\mathbf{a}'; \sigma'); \mu, J) \\ &= \sum_{\substack{(\mathbf{a}'; \sigma') \\ \gamma', J}} \frac{k_{\gamma'}^{(\mathbf{a}'; \sigma')}}{\sqrt{\dim Q(\mathbf{a}')}} \langle Z_{\mu, J}^{(\mathbf{a}', \gamma')} | Z_{\nu, I}^{(\mathbf{a}, \gamma)} \rangle e((\mathbf{a}'; \sigma'); \mu, J) \\ &= \sum_{\sigma'=1}^{\mathfrak{m}(\mathbf{p}; \mathbf{a})} \frac{k_{\gamma}^{(\mathbf{a}; \sigma')}}{\sqrt{\dim Q(\mathbf{a})}} e((\mathbf{a}; \sigma'); \nu, I), \end{aligned}$$

where

$$(4.47) \quad k_{\gamma'}^{(\mathbf{a}'; \sigma')} = \frac{\overline{c_{\gamma'}^{\sigma'}(\mathbf{a}')}}{\sqrt{\dim Q(\mathbf{p})}} \sqrt{\frac{\dim Q(\mathbf{a}')}{\dim Q(\mathbf{p})}},$$

cf. (4.44). So, for  $c_{\gamma}^{\sigma}(\mathbf{a})$  and  $\widetilde{c}_{\gamma}^{\sigma}(\mathbf{a})$  being the characteristic parameters associated to  $\mathbf{C}(\mathbf{a})$  and  $\widetilde{\mathbf{C}}(\mathbf{a})$ , respectively, we have, cf. (4.29),

$$(4.48) \quad \dim Q(\mathbf{p}) \int_{\mathcal{E}} Z \widetilde{\mathbf{C}}(\mathbf{a})_{\nu, I}^{\sigma}(\mathbf{z}) K(\mathbf{z}) d\mathbf{z} = \sqrt{\dim Q(\mathbf{p})} \sum_{\gamma, \sigma'} \widetilde{c}_{\gamma}^{\sigma}(\mathbf{a}) \overline{c_{\gamma}^{\sigma'}(\mathbf{a})} e((\mathbf{a}; \sigma'); \nu, I).$$

Hence, (4.45) holds for  $A = e((\mathbf{a}; \sigma); \boldsymbol{\nu}, I)$  if and only if

$$(4.49) \quad \sum_{\gamma=1}^{m(\mathbf{a})} \tilde{c}_{\gamma}^{\sigma}(\mathbf{a}) \overline{c_{\gamma}^{\sigma'}(\mathbf{a})} = \delta_{\sigma, \sigma'} ,$$

which means  $(\mathbf{C}(\mathbf{a}))^{\dagger} \tilde{\mathbf{C}}(\mathbf{a}) = \mathbf{1}$ , or equivalently  $(\tilde{\mathbf{C}}(\mathbf{a}))^{\dagger} \mathbf{C}(\mathbf{a}) = \mathbf{1}$ .  $\square$

Now, let

$$(4.50) \quad \langle A|R \rangle_{\mathbf{p}} = \frac{1}{\dim Q(\mathbf{p})} \langle A|R \rangle = \frac{1}{\dim Q(\mathbf{p})} \text{tr}(A^{\dagger} R)$$

be the normalized inner product in  $\mathcal{B}(\mathcal{H}_{\mathbf{p}})$  and  $\|\cdot\|_{\mathbf{p}}$  be the induced norm.

**Definition 4.17.** Two symbol correspondences  $W, \tilde{W} : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C^{\infty}(\mathcal{E})$  satisfying

$$(4.51) \quad \langle A|R \rangle_{\mathbf{p}} = \langle \tilde{W}_A | W_R \rangle = \langle W_A | \tilde{W}_R \rangle$$

for every  $A, R \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$  are said to be dual correspondences. The operators kernels of  $W$  and  $\tilde{W}$  are also said to be dual to each other.

**Proposition 4.18.** Two symbol correspondences  $W, \tilde{W} : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C^{\infty}(\mathcal{E})$  with characteristic matrices  $\mathbf{C}(\mathbf{a})$  and  $\tilde{\mathbf{C}}(\mathbf{a})$  are dual to each other if and only if

$$(\tilde{\mathbf{C}}(\mathbf{a}))^{\dagger} \mathbf{C}(\mathbf{a}) = \mathbf{1} .$$

*Proof.* The proof follows analogous to the proof of Proposition 3.20 by writing the operators using (4.45) and symbols using (4.39).  $\square$

**Remark 4.19.** Duality is no longer  $1 \leftrightarrow 1$ . Consider, for instance, the correspondences  $W, \tilde{W} : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathcal{E})$  defined respectively by the characteristic parameters  $c_{\gamma}^{\sigma}(\mathbf{a}) = \delta_{\gamma, \sigma}$  and  $\tilde{c}_{\gamma}^{\sigma}(\mathbf{a}) = \delta_{\gamma, \sigma} + \delta_{\gamma, m(\mathbf{a})} \delta_{\sigma, 1}$ . Then, both  $\tilde{W}$  and  $W$  itself are dual to  $W$ .

The correspondence  $W$  of the previous remark is obviously an isometry. In addition to such special cases of correspondences which are isometric, now we also have correspondences given by a direct sum of conformal maps.

**Definition 4.20.** A symbol correspondence  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathcal{E})$  is a Stratonovich-Weyl correspondence if it is an isometry, that is,

$$(4.52) \quad \langle A|R \rangle_{\mathbf{p}} = \langle W_A | W_R \rangle$$

for all  $A, R \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$ . If  $W$  preserves angles for each  $\mathcal{B}(\mathbf{p}; \mathbf{a})$ , that is,

$$(4.53) \quad \frac{\langle A|R \rangle_{\mathbf{p}}}{\|A\|_{\mathbf{p}} \|R\|_{\mathbf{p}}} = \frac{\langle W_A | W_R \rangle}{\|W_A\| \|W_R\|}$$

for all non null  $A, R \in \mathcal{B}(\mathbf{p}; \mathbf{a})$  and every  $\mathcal{B}(\mathbf{p}; \mathbf{a}) \subset \mathcal{B}(\mathcal{H}_{\mathbf{p}})$ , cf. (4.28), then  $W$  shall be called a semi-conformal correspondence.

**Proposition 4.21.** A symbol correspondence  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathcal{E})$  is a Stratonovich-Weyl correspondence if and only if its characteristic matrices are unitary matrices, that is, they satisfy  $(\mathbf{C}(\mathbf{a}))^{\dagger} \mathbf{C}(\mathbf{a}) = \mathbf{1}$ . Furthermore,  $W$  is a semi-conformal correspondence if and only if its characteristic matrices are conformal matrices, that is,  $(\mathbf{C}(\mathbf{a}))^{\dagger} \mathbf{C}(\mathbf{a}) = \alpha(\mathbf{a}) \mathbf{1}$  for  $\alpha(\mathbf{a}) > 0$ , where  $\alpha(\mathbf{a}) = \alpha(\tilde{\mathbf{a}})$  and  $\alpha(0, 0) = 1$ .

*Proof.* Proposition 4.18 implies that  $W$  is its own dual if and only if  $(\mathfrak{C}(\mathbf{a}))^\dagger \mathfrak{C}(\mathbf{a}) = \mathbb{1}$  holds, which proves the first part of the statement. For the second part, we use that a linear map is conformal if and only if it is a positive real multiple of an unitary linear map, thus  $W$  is a semi-conformal correspondence if and only if there is  $\alpha(\mathbf{a}) > 0$  for each  $\mathcal{B}(\mathbf{p}; \mathbf{a})$  such that  $\alpha(\mathbf{a})^{-1/2} W|_{\mathcal{B}(\mathbf{p}; \mathbf{a})}$  is an unitary map, and this is true if and only if the characteristic matrices of  $W$  satisfy  $(\mathfrak{C}(\mathbf{a}))^\dagger \mathfrak{C}(\mathbf{a}) = \alpha(\mathbf{a}) \mathbb{1}$ . The equations  $\alpha(0, 0) = 1$  and  $\alpha(\mathbf{a}) = \alpha(\tilde{\mathbf{a}})$  follows from  $\overline{\mathfrak{C}(\mathbf{a})} = \mathfrak{C}(\tilde{\mathbf{a}})$  and  $\mathfrak{C}(0, 0) = (-1)^{|\mathbf{p}|}$ .  $\square$

**Remark 4.22.** *A symbol correspondence  $W$  is an actual conformal map if and only if  $W = \sqrt{\alpha} W'$  for  $\alpha > 0$  and some Stratonovich-Weyl correspondence  $W'$ . Since  $W_{\mathbb{1}} = W'_{\mathbb{1}}$ , we must have  $\alpha = 1$ , so the only actual conformal correspondences are the isometric ones.*

*We also point out that, for pure-quark systems (likewise for spin systems), every symbol correspondence is a semi-conformal correspondence, with  $\alpha(\mathbf{a}) = \alpha(n, n) = c_n^2$ .*

Propositions 4.16–4.21 illustrate how characteristic matrices encode all the information about symbol correspondences for generic quark systems in the same vein of characteristic numbers for pure-quark system. The existence of multiple dual representations can be explained by the existence of invariant subspaces  $Q(\mathbf{a})$  with higher degeneracy within  $C_c^\infty(\mathcal{E})$  than within  $\mathcal{B}(\mathcal{H}_{\mathbf{p}})$ , or equivalently by the existence of multiple left inverses of characteristic matrices.

In general, there is no natural way to choose a dual correspondence among all possibilities. For semi-conformal symbol correspondences, however, we make the following definition:

**Definition 4.23.** *Let  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_c^\infty(\mathcal{E})$  be a semi-conformal correspondence with characteristic matrices  $\mathfrak{C}(\mathbf{a})$  satisfying  $(\mathfrak{C}(\mathbf{a}))^\dagger \mathfrak{C}(\mathbf{a}) = \alpha(\mathbf{a}) \mathbb{1}$ . Its canonical dual correspondence is the symbol correspondence  $\tilde{W} : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C^\infty(\mathcal{E})$  with characteristic matrices*

$$(4.54) \quad \tilde{\mathfrak{C}}(\mathbf{a}) = \frac{1}{\alpha(\mathbf{a})} \mathfrak{C}(\mathbf{a}) .$$

Thus, Stratonovich-Weyl correspondences for generic quark systems are their own canonical dual correspondences.

Just as for pure-quark systems, positive operator kernels provide special symbol correspondences for generic quark systems.

**Definition 4.24.** *A symbol correspondence  $W$  for a generic quark system is mapping-positive if it maps positive(-definite) operators to (strictly-)positive functions. If  $\tilde{W}$  is dual to a mapping-positive correspondence, then  $\tilde{W}$  is a positive-dual correspondence.*

**Proposition 4.25.** *A symbol correspondence  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_c^\infty(\mathcal{E})$  with operator kernel  $K$  is mapping-positive if and only if  $K$  is a state, that is,  $K$  is a positive operator.*

*Proof.* Suppose  $K$  is a positive operator, so  $K = R^\dagger R$  for some  $R \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$ , and let  $A = M^\dagger M \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$  be a positive operator. Then, for any  $g \in SU(3)$  and

$$\widetilde{M} = M\rho(g), \quad (4.55)$$

$$W_A(g\mathbf{z}_0) = \text{tr}(M^\dagger M\rho(g)K\rho(g)^\dagger) = \text{tr}(M\rho(g)K\rho(g)^\dagger M^\dagger) = \text{tr}(\widetilde{M}R^\dagger R\widetilde{M}^\dagger) \geq 0,$$

where we used that  $\widetilde{M}R^\dagger R\widetilde{M}^\dagger$  is a positive operator. Since  $K$  is non null,  $R$  is also non null, so there exist  $w_0 \in \mathcal{H}_{\mathbf{p}}$  such that  $\|R(w_0)\|^2 > 0$ . If  $A$  is positive-definite,  $\widetilde{M}^\dagger$  is an automorphism and we can set  $w = (\widetilde{M}^\dagger)^{-1}(w_0)$  so that  $\|w\| > 0$  and

$$\begin{aligned} W_A(\mathbf{z}) &= \text{tr}(\widetilde{M}R^\dagger R\widetilde{M}^\dagger) \geq \frac{\langle w | \widetilde{M}R^\dagger R\widetilde{M}^\dagger(w) \rangle}{\|w\|^2} = \frac{\langle R\widetilde{M}^\dagger(w) | R\widetilde{M}^\dagger(w) \rangle}{\|w\|^2} \\ (4.56) \quad &= \frac{\|R\widetilde{M}^\dagger(w)\|^2}{\|w\|^2} = \frac{\|R(w_0)\|^2}{\|w\|^2} \\ &> 0. \end{aligned}$$

Now, suppose  $K$  is not positive. Then,  $K$  has a negative eigenvalue. Let  $\Pi$  be the projection onto an eigenspace of  $K$  associated to a negative eigenvalue. We have that  $\text{tr}(\Pi K) < 0$ .  $\square$

In the light of Remark 4.15,  $\Pi_{(0,0,p)} \in \mathcal{B}(\mathcal{H}_{p,0})$  and  $\Pi_{(p,p,0)} \in \mathcal{B}(\mathcal{H}_{0,p})$  are operator kernels of mapping-positive correspondences for generic quark systems.

**Definition 4.26.** For any  $p \in \mathbb{N}$ , the mapping-positive correspondences  $\mathcal{B}(\mathcal{H}_{p,0}) \rightarrow C_c^\infty(\mathcal{E})$ , with operator kernel  $\Pi_{(0,0,p)}$ , and  $\mathcal{B}(\mathcal{H}_{0,p}) \rightarrow C_c^\infty(\mathcal{E})$ , with operator kernel  $\Pi_{(p,p,0)}$ , are Berezin correspondences for generic quark systems.

**Remark 4.27.** The Berezin correspondences of Definition 4.26 are rather trivial examples, in the sense that the symbols on  $\mathbb{C}P^1 \rightarrow \mathcal{E} \rightarrow \mathbb{C}P^2$  are constant extensions of functions on  $\mathbb{C}P^2$ . Thus, it would be interesting if we could exhibit some nontrivial examples of mapping-positive correspondences for generic quark systems.

In fact, we would like to emphasize that we still don't know whether a result similar to Proposition 3.26 holds for generic quark systems.

Also, for pure-quark systems every correspondence is semi-conformal, cf. Remark 4.22, and for generic quark system every Stratonovich-Weyl correspondence is semi-conformal, cf. Proposition 4.21. We still don't know if there is any general relation between mapping-positive (or positive-dual) correspondences and semi-conformal correspondences, for generic quark systems.

To finish this section, we define antipodal correspondences for generic quark systems and show some of their general properties.

**Definition 4.28.** For a symbol correspondence  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_c^\infty(\mathcal{E})$ , its antipodal correspondence is the symbol correspondence  $\widetilde{W} : \mathcal{B}(\mathcal{H}_{\bar{\mathbf{p}}}) \rightarrow C^\infty(\mathcal{E})$  given by

$$(4.57) \quad \widetilde{W}_{A^*} = W_A,$$

cf. (2.127)-(2.129).

**Proposition 4.29.** The symbol correspondences  $\widetilde{W} : \mathcal{B}(\mathcal{H}_{\bar{\mathbf{p}}}) \rightarrow C_c^\infty(\mathcal{E})$  with characteristic parameters  $(\check{c}_\gamma^\sigma(\mathbf{a}))$  is antipodal to the symbol correspondence  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_c^\infty(\mathcal{E})$  with characteristic parameters  $(c_\gamma^\sigma(\mathbf{a}))$  if and only if

$$(4.58) \quad c_\gamma^\sigma(\mathbf{a}) = (-1)^{|\mathbf{a}|} \check{c}_\gamma^{\check{\sigma}}(\mathbf{a}).$$

*Proof.* The result follows from (2.131) and Theorem 4.11.  $\square$

Recalling (2.120), we have from (4.58) that the characteristic matrices of a symbol correspondence and of its antipodal differ just by a constant  $(-1)^{|\mathbf{a}|}$  factor and the reverse ordering of columns. We then have the following:

**Corollary 4.30.** *For generic quark systems, a symbol correspondence and its antipodal have the same image in  $C_{\mathbb{C}}^{\infty}(\mathcal{E})$ .*

**Proposition 4.31.** *If  $W, \widetilde{W} : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathcal{E})$  are symbol correspondences dual to each other, then their respective antipodal correspondences are also dual to each other.*

*Proof.* Let  $(c_{\gamma}^{\sigma}(\mathbf{a}))$  and  $(\widetilde{c}_{\gamma}^{\sigma}(\mathbf{a}))$  be the characteristic parameters of  $W$  and  $\widetilde{W}$ , respectively. The result follows from

$$(4.59) \quad \sum_{\gamma=1}^{m(\mathbf{a})} \left( (-1)^{|\mathbf{a}|} \overline{\widetilde{c}_{\gamma}^{\sigma}(\mathbf{a})} \right) \left( (-1)^{|\mathbf{a}|} c_{\gamma}^{\sigma'}(\mathbf{a}) \right) = \sum_{\gamma=1}^{m(\mathbf{a})} \overline{\widetilde{c}_{\gamma}^{\sigma}(\mathbf{a})} c_{\gamma}^{\sigma'}(\mathbf{a}) = \delta_{\sigma, \sigma'} .$$

$\square$

**Corollary 4.32.** *A symbol correspondence for a generic quark system is a semi-conformal (resp. Stratonovich-Weyl) correspondence if and only if its antipodal correspondence is also a semi-conformal (resp. Stratonovich-Weyl) correspondence.*

**4.4. Twisted products for generic quark systems.** Again, let  $\mathbf{p} \in (\mathbb{N} \times \mathbb{N}_0) \cup (\mathbb{N}_0 \times \mathbb{N})$ .

Now, we may have symbol correspondences  $W_1, W_2 : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathcal{E})$  with different image sets, that is, such that  $W_1(\mathcal{B}(\mathcal{H}_{\mathbf{p}})) \neq W_2(\mathcal{B}(\mathcal{H}_{\mathbf{p}}))$ . To see this, consider  $W_1 \neq W_2 : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C^{\infty}(\mathcal{E})$  determined by characteristic matrices  $\mathfrak{C}[1](\mathbf{a})$  and  $\mathfrak{C}[2](\mathbf{a})$  with respective characteristic parameters  $c[1]_{\gamma}^{\sigma}(\mathbf{a})$  and  $c[2]_{\gamma}^{\sigma}(\mathbf{a})$ . Since  $\mathfrak{m}(\mathbf{p}; |\mathbf{p}|, |\mathbf{p}|) = 1$ , for  $Q(\mathbf{a}) = Q(|\mathbf{p}|, |\mathbf{p}|)$ , we can drop the index  $\sigma$  for the characteristic parameters  $c[1]_{\gamma}(|\mathbf{p}|, |\mathbf{p}|)$  and  $c[2]_{\gamma}(|\mathbf{p}|, |\mathbf{p}|)$ . Then, for  $\mathfrak{C}[1](|\mathbf{p}|, |\mathbf{p}|)$  and  $\mathfrak{C}[2](|\mathbf{p}|, |\mathbf{p}|)$  such that

$$\mathfrak{C}[1](|\mathbf{p}|, |\mathbf{p}|) \cdot \mathfrak{C}[2](|\mathbf{p}|, |\mathbf{p}|) = \sum_{\gamma=1}^{|\mathbf{p}|+1} c[1]_{\gamma}(|\mathbf{p}|, |\mathbf{p}|) c[2]_{\gamma}(|\mathbf{p}|, |\mathbf{p}|) = 0 ,$$

we have that

$$W_1(Q(|\mathbf{p}|, |\mathbf{p}|)) = \text{span} \{ Z \mathfrak{C}[1](|\mathbf{p}|, |\mathbf{p}|)_{\nu, I} \}$$

is orthogonal<sup>21</sup> to

$$W_2(Q(|\mathbf{p}|, |\mathbf{p}|)) = \text{span} \{ Z \mathfrak{C}[2](|\mathbf{p}|, |\mathbf{p}|)_{\nu, I} \} ,$$

hence  $W_1(\mathcal{B}(\mathcal{H}_{\mathbf{p}})) \neq W_2(\mathcal{B}(\mathcal{H}_{\mathbf{p}}))$ .

In view of the this fact, for any symbol correspondence  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C^{\infty}(\mathcal{E})$ , we shall denote by  $\mathcal{S}_{\mathbf{p}}(W)$  the image of  $W$ , that is,

$$(4.60) \quad \mathcal{S}_{\mathbf{p}}(W) = W(\mathcal{B}(\mathcal{H}_{\mathbf{p}})) \subset C_{\mathbb{C}}^{\infty}(\mathcal{E}) .$$

<sup>21</sup>We recall that the constant function 1 on  $\mathcal{E}$  is in the image of  $Q(0, 0)$ , thus  $1 \notin W(Q(|\mathbf{p}|, |\mathbf{p}|))$ , for any  $\mathbf{p} \in (\mathbb{N} \times \mathbb{N}_0) \cup (\mathbb{N}_0 \times \mathbb{N})$  and for any correspondence  $W$ .

**Definition 4.33.** Given a symbol correspondence  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathcal{E})$ , the twisted product of symbols induced by  $W$  is the binary operation  $\star$  on  $\mathcal{S}_{\mathbf{p}}(W)$  given by

$$(4.61) \quad W_A \star W_R = W_{AR}$$

for any  $A, R \in \mathcal{B}(\mathcal{H}_{\mathbf{p}})$ . The algebra  $(\mathcal{S}_{\mathbf{p}}(W), \star)$  is called a twisted  $\mathbf{p}$ -algebra.

The proofs of the two next propositions are exactly as the proofs of Propositions 3.40 and 3.41, respectively.

**Proposition 4.34.** Any twisted  $\mathbf{p}$ -algebra  $(\mathcal{S}_{\mathbf{p}}(W), \star)$  is

- i)  $SU(3)$ -equivariant:  $(f_1 \star f_2)^g = f_1^g \star f_2^g$ ;
- ii) Associative:  $(f_1 \star f_2) \star f_3 = f_1 \star (f_2 \star f_3)$ ;
- iii) Unital:  $1 \star f = f \star 1 = f$ ;
- iv)  $A$   $\ast$ -algebra:  $\overline{f_1 \star f_2} = \overline{f_2} \star \overline{f_1}$ ;

where  $f_1, f_2, f_3, f \in \mathcal{S}_{\mathbf{p}}(W)$ ,  $g \in SU(3)$  and  $1 \in \mathcal{S}_{\mathbf{p}}(W)$  is the constant function equal to 1 on  $\mathcal{E}$ .

**Proposition 4.35.** Fixed  $\mathbf{p} \in (\mathbb{N} \times \mathbb{N}_0) \cup (\mathbb{N}_0 \times \mathbb{N})$ , any two twisted  $\mathbf{p}$ -algebras are naturally isomorphic, and any  $\mathbf{p}$ -algebra is naturally anti-isomorphic to any  $\check{\mathbf{p}}$ -algebra.

*Proof.* Although we may have correspondences  $W_1, W_2 : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathcal{E})$  with different images, we still have that each  $W_j$  is an isomorphism onto its image, so  $W_1 \circ W_2^{-1} : \mathcal{S}_{\mathbf{p}}(W_2) \rightarrow \mathcal{S}_{\mathbf{p}}(W_1)$  is an isomorphism. If  $W_2 : \mathcal{B}(\mathcal{H}_{\check{\mathbf{p}}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathcal{E})$ , then  $W_1 \circ \ast \circ W_2^{-1} : \mathcal{S}_{\check{\mathbf{p}}}(W_2) \rightarrow \mathcal{S}_{\mathbf{p}}(W_1)$  is an anti-isomorphism because the adjoint map  $\ast$  is an anti-isomorphism and, again, each  $W_j$  is an isomorphism onto its image.  $\square$

We cannot decompose twisted products for generic quark systems into the harmonic basis as we did for pure-quark systems because, in general,  $\mathcal{S}_{\mathbf{p}}(W)$  is not spanned by the generic harmonics  $Z_{\nu, I}^{(\mathbf{a}, \gamma)}$ , but by the linear combinations expressed in (4.29). Thus, a procedure equivalent to the one executed in Theorem 3.42 leads to the following:

**Theorem 4.36.** If  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C^{\infty}(\mathcal{E})$  is a symbol correspondence with characteristic matrices  $\mathfrak{C}(\mathbf{a})$ , then the induced twisted product is given by

$$(4.62) \quad \begin{aligned} & Z\mathfrak{C}(\mathbf{a}_1)_{\nu_1, I_1}^{\sigma_1} \star Z\mathfrak{C}(\mathbf{a}_2)_{\nu_2, I_2}^{\sigma_2} \\ &= \sqrt{\dim Q(\mathbf{p})} \sum_{\substack{(\mathbf{a}; \sigma) \\ \nu, I}} (-1)^{|\mathbf{p}|+2(t_{\nu}+u_{\nu})} \begin{bmatrix} (\mathbf{a}_1; \sigma_1) & (\mathbf{a}_2; \sigma_2) & (\mathbf{a}; \sigma) \\ \nu_1, I_1 & \nu_2, I_2 & \check{\nu}, I \end{bmatrix} [\mathbf{p}] Z\mathfrak{C}(\mathbf{a})_{\nu, I}^{\sigma}, \end{aligned}$$

where  $Z\mathfrak{C}(\mathbf{a})_{\nu, I}^{\sigma}$  is given by (4.29) and where summations over  $\nu$  and  $I$  can be simplified using (2.149).

*Proof.* The proof follows from Corollary 2.29 and Theorem 4.11.  $\square$

In that light, integral formulation of twisted products may be more useful for the generic case.

**Theorem 4.37.** *If  $W : \mathcal{B}(\mathcal{H}_p) \rightarrow C^\infty(\mathcal{E})$  is a symbol correspondence with operator kernel  $K$  and characteristic matrices  $\mathfrak{C}(\mathbf{a})$ , then the induced twister product is given by*

$$(4.63) \quad f_1 \star f_2(\mathbf{z}) = \int_{\mathcal{E} \times \mathcal{E}} f_1(\mathbf{z}_1) f_2(\mathbf{z}_2) \mathbb{L}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}) d\mathbf{z}_1 d\mathbf{z}_2$$

for any  $f_1, f_2 \in \mathcal{S}_p(W)$ , where

$$(4.64) \quad \begin{aligned} \mathbb{L}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3) &= (\dim Q(\mathbf{p}))^2 \operatorname{tr} \left( \tilde{K}(\mathbf{z}_1) \tilde{K}(\mathbf{z}_2) K(\mathbf{z}_3) \right) \\ &= (-1)^{|\mathbf{p}|} \sqrt{\dim Q(\mathbf{p})} \sum_{\substack{(\mathbf{a}_j; \sigma_j) \\ \nu_j, I_j}} \begin{bmatrix} (\mathbf{a}_1; \sigma_1) & (\mathbf{a}_2; \sigma_2) & (\mathbf{a}_3; \sigma_3) \\ \nu_1, I_1 & \nu_2, I_2 & \nu_3, I_3 \end{bmatrix} [\mathbf{p}] \\ &\quad \times \overline{Z\tilde{\mathfrak{C}}(\mathbf{a}_1)_{\nu_1, I_1}^{\sigma_1}(\mathbf{z}_1)} \overline{Z\tilde{\mathfrak{C}}(\mathbf{a}_2)_{\nu_2, I_2}^{\sigma_2}(\mathbf{z}_2)} \overline{Z\tilde{\mathfrak{C}}(\mathbf{a}_3)_{\nu_3, I_3}^{\sigma_3}(\mathbf{z}_3)} \end{aligned}$$

for  $\tilde{\mathfrak{C}}(\mathbf{a})$  being the characteristic matrices of an operator kernel  $\tilde{K}$  dual to  $K$ .

*Proof.* The proof follows analogously to Theorem 3.43, but now the second equality comes from Theorem 4.36.

We emphasize that, although the expression (4.64) for integral trikernels depends explicitly on the choice of dual representation, the twisted product given by (4.63) does not have such dependence. By definition,

$$(4.65) \quad \begin{aligned} \int_{\mathcal{E}} \overline{Z\tilde{\mathfrak{C}}(\mathbf{a})_{\nu, I}^{\sigma}(\mathbf{z})} Z\mathfrak{C}(\mathbf{a}')_{\nu', I'}^{\sigma'}(\mathbf{z}) d\mathbf{z} &= \left\langle Z\tilde{\mathfrak{C}}(\mathbf{a})_{\nu, I}^{\sigma} \middle| Z\mathfrak{C}(\mathbf{a}')_{\nu', I'}^{\sigma'} \right\rangle \\ &= \dim Q(\mathbf{p}) \langle \mathbf{e}((\mathbf{a}; \sigma); \nu, I) | \mathbf{e}((\mathbf{a}'; \sigma'); \nu', I') \rangle_{\mathbf{p}} \\ &= \langle \mathbf{e}((\mathbf{a}; \sigma); \nu, I) | \mathbf{e}((\mathbf{a}'; \sigma'); \nu', I') \rangle \end{aligned}$$

no matter which dual correspondence is used.  $\square$

**Definition 4.38.** *An integral trikernel  $\mathbb{L} \in C^\infty(\mathcal{E} \times \mathcal{E} \times \mathcal{E})$  of a twisted product induced by a symbol correspondence  $W : \mathcal{B}(\mathcal{H}_p) \rightarrow C^\infty(\mathcal{E})$  is a function of the form (4.64) so that the twisted product is given by (4.63). If  $W$  is a semi-conformal correspondence, the integral trikernel constructed using its canonical dual correspondence is the canonical integral trikernel.*

One may use (4.63) to expand a twisted product on  $\mathcal{S}_p(W)$  induced by some symbol correspondence  $W$  to a product  $\bullet$  on all  $C_{\mathbb{C}}^\infty(\mathcal{E})$  in the same way we did for pure-quark systems in Proposition 3.45, but integral trikernels are not unique, so such expansions are not unique too. In addition, the product  $\bullet$  in general fails to vanish for functions orthogonal to  $\mathcal{S}_p(W)$  since we may find a symbol correspondence  $\tilde{W}$  dual to  $W$  with  $\mathcal{S}_p(\tilde{W}) \neq \mathcal{S}_p(W)$  as exemplified in Remark 4.19. However, for semi-conformal correspondences and canonical integral trikernels, we have the following:

**Proposition 4.39.** *Let  $\mathbb{L}$  be the canonical integral trikernel of a twisted product  $\star$  induced by a semi-conformal correspondence  $W : \mathcal{B}(\mathcal{H}_p) \rightarrow C_{\mathbb{C}}^\infty(\mathcal{E})$  with characteristic matrices satisfying  $(\mathfrak{C}(\mathbf{a}))^\dagger \mathfrak{C}(\mathbf{a}) = \alpha(\mathbf{a}) \mathbb{1}$ . The binary operation  $\bullet$  on  $C_{\mathbb{C}}^\infty(\mathcal{E})$  given by*

$$(4.66) \quad f_1 \bullet f_2(\mathbf{z}) = \int_{\mathcal{E} \times \mathcal{E}} f_1(\mathbf{z}_1) f_2(\mathbf{z}_2) \mathbb{L}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}) d\mathbf{z}_1 d\mathbf{z}_2$$

for any  $f_1, f_2 \in C_{\mathbb{C}}^{\infty}(\mathcal{E})$ , defines a  $SU(3)$ -equivariant associative  $*$ -algebra with respect to complex conjugation. In particular, if  $f_1, f_2 \in \mathcal{S}_{\mathbf{p}}(W)$ , we have  $f_1 \bullet f_2 = f_1 \star f_2$ . But, if either  $f_1$  or  $f_2$  is orthogonal to  $\mathcal{S}_{\mathbf{p}}(W)$ , then  $f_1 \bullet f_2 = 0$ .

*Proof.* The proof follows from the same arguments applied to Proposition 3.45, but now it is needed to point out that

$$(4.67) \quad \begin{aligned} \mathbb{L}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3) &= (-1)^{|\mathbf{p}|} \sqrt{\dim Q(\mathbf{p})} \sum_{\substack{(\mathbf{a}_j; \sigma_j) \\ \nu_j, I_j}} \left[ \begin{array}{ccc} (\mathbf{a}_1; \sigma_1) & (\mathbf{a}_2; \sigma_2) & (\mathbf{a}_3; \sigma_3) \\ \nu_1, I_1 & \nu_2, I_2 & \nu_3, I_3 \end{array} \right] [\mathbf{p}] \\ &\times \frac{1}{\alpha(\mathbf{a})^2} \overline{Z\mathfrak{C}(\mathbf{a}_1)_{\nu_1, I_1}^{\sigma_1}(\mathbf{z}_1)} \overline{Z\mathfrak{C}(\mathbf{a}_2)_{\nu_2, I_2}^{\sigma_2}(\mathbf{z}_2)} \overline{Z\mathfrak{C}(\check{\mathbf{a}}_3)_{\nu_3, I_3}^{\sigma_3}(\mathbf{z}_3)}. \end{aligned}$$

Thus, if  $f_j \in C_{\mathbb{C}}^{\infty}(\mathcal{E})$  is orthogonal to  $\mathcal{S}_{\mathbf{p}}(W)$ , it is orthogonal to every  $Z\mathfrak{C}(\mathbf{a})_{\nu, I}^{\sigma}$  and this implies that the integral over  $\mathbf{z}_j$  in (4.66) vanishes.  $\square$

Analogously to pure-quark systems, a product  $\bullet$  satisfying the previous proposition is such that  $C_{\mathbb{C}}^{\infty}(\mathcal{E}) \rightarrow \mathcal{S}_{\mathbf{p}}(W) : f \mapsto 1 \bullet f = f \bullet 1$  is an orthogonal projection.

**Proposition 4.40.** *Let  $\mathbb{L}$  be an integral trikernel of a twisted product  $\star$  induced by  $W : \mathcal{B}(\mathcal{H}_{\mathbf{p}}) \rightarrow C_{\mathbb{C}}^{\infty}(\mathcal{E})$  as in (4.64). Then, for every  $g \in SU(3)$  and every  $\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3, \mathbf{z}_4 \in \mathcal{E}$ ,*

$$(4.68) \quad \begin{aligned} &i) \mathbb{L}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3) = \mathbb{L}(g\mathbf{z}_1, g\mathbf{z}_2, g\mathbf{z}_3) ; \\ &ii) \int_{\mathcal{E}} \mathbb{L}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}) \mathbb{L}(\mathbf{z}, \mathbf{z}_3, \mathbf{z}_4) d\mathbf{z} = \int_{\mathcal{E}} \mathbb{L}(\mathbf{z}_1, \mathbf{z}, \mathbf{z}_4) \mathbb{L}(\mathbf{z}_2, \mathbf{z}_3, \mathbf{z}) d\mathbf{z} ; \\ &iii) \int_{\mathcal{E}} \mathbb{L}(\mathbf{z}, \mathbf{z}_1, \mathbf{z}_2) d\mathbf{z} = \int_{\mathcal{E}} \mathbb{L}(\mathbf{z}_1, \mathbf{z}, \mathbf{z}_2) d\mathbf{z} = \mathcal{R}_{\mathbf{p}}^W(\mathbf{z}_1, \mathbf{z}_2) , \text{ where} \\ &\mathcal{R}_{\mathbf{p}}^W(\mathbf{z}_1, \mathbf{z}_2) = \sum_{\substack{(\mathbf{a}; \sigma) \\ \nu, I}} \overline{Z\mathfrak{C}(\mathbf{a})_{\nu, I}^{\sigma}(\mathbf{z}_1)} Z\mathfrak{C}(\mathbf{a})_{\nu, I}^{\sigma}(\mathbf{z}_2) , \end{aligned}$$

satisfies

$$(4.69) \quad \int_{\mathcal{E}} f(\mathbf{z}_1) \mathcal{R}_{\mathbf{p}}^W(\mathbf{z}_1, \mathbf{z}_2) d\mathbf{z}_1 = f(\mathbf{z}_2)$$

for every  $f \in \mathcal{S}_{\mathbf{p}}(W)$ ;

$$iv) \overline{\mathbb{L}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3)} = \mathbb{L}(\mathbf{z}_2, \mathbf{z}_1, \mathbf{z}_3).$$

*Proof.* Adapting the proof of Proposition 3.46, we get that each property of this statement is equivalent to the property of Proposition 4.34 with same number. It is just worth to highlight that the expression for  $\mathcal{R}_{\mathbf{p}}^W$  comes from (4.64), orthonormality of  $\mathcal{E}$  harmonics and (2.152).  $\square$

**Remark 4.41.** *In general,  $\mathcal{R}_{\mathbf{p}}^W(\mathbf{z}_1, \mathbf{z}_2) \neq \mathcal{R}_{\mathbf{p}}^W(\mathbf{z}_2, \mathbf{z}_1)$ . But if  $W$  is a semi-conformal correspondence and  $\mathbb{L}$  is its canonical integral trikernel, then (iii) is satisfied with*

$$(4.70) \quad \mathcal{R}_{\mathbf{p}}^W(\mathbf{z}_1, \mathbf{z}_2) = \sum_{\substack{(\mathbf{a}; \sigma) \\ \nu, I}} \frac{1}{\alpha(\mathbf{a})} \overline{Z\mathfrak{C}(\mathbf{a})_{\nu, I}^{\sigma}(\mathbf{z}_1)} Z\mathfrak{C}(\mathbf{a})_{\nu, I}^{\sigma}(\mathbf{z}_2) = \mathcal{R}_{\mathbf{p}}^W(\mathbf{z}_2, \mathbf{z}_1) ,$$

so that  $\mathcal{R}_{\mathbf{p}}^W$  is the reproducing kernel on  $\mathcal{S}_{\mathbf{p}}(W)$ .

The phenomenon described in Remark 3.50 also occurs to generic quark systems.

**Proposition 4.42.** *The twisted products  $\star$  and  $\check{\star}$  induced by a symbol correspondence and its antipodal correspondence satisfy*

$$(4.71) \quad f_1 \star f_2 = f_2 \check{\star} f_1 .$$

*Proof.* The proof is analogous to Proposition 3.48.  $\square$

**Corollary 4.43.** *For  $\star$  and  $\check{\star}$  as in the previous proposition, we can choose integral trikernels  $\mathbb{L}$  and  $\check{\mathbb{L}}$  satisfying*

$$(4.72) \quad \mathbb{L}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3) = \check{\mathbb{L}}(\mathbf{z}_2, \mathbf{z}_1, \mathbf{z}_3) .$$

## 5. CONCLUDING REMARKS

The main problem studied in this paper, the characterization of symbol correspondences for quark systems, is often settled on more general facts, thus some of the features presented here are in common to the case of spin systems. For example, the realization of symbol correspondences as expectation values over an *operator kernel*, which is a special “pseudo-state”, that is, an Hermitian operator with unitary trace. Then, the more restricted cases of mapping-positives correspondences are the ones which are generated as expectation values over an operator kernel that is an “actual state”, thus being also a positive operator.

In particular, symbol correspondences for pure-quark systems show little formal distinction to what is known for spin systems, being also determined by a set of non zero real numbers, the *characteristic numbers*. Nonetheless, a remarkable difference occurs due to the breakdown of the self dual property for representations  $Q(p, 0)$  and  $Q(0, p)$  in the case of quantum pure-quark systems: antipodal correspondences are defined for pure-quark systems dual to each other and have the same characteristic numbers<sup>22</sup>.

However, correspondences for generic quark system present some new features originated from the degeneracy of representations within both the quantum operator space  $\mathcal{B}(\mathcal{H}_{p,q})$  and the classical function space  $C_{\mathbb{C}}^{\infty}(\mathcal{E})$ . Then, the characterization of correspondences for generic quark systems, in the same vein of what is done to pure-quark systems, are given not in terms of characteristic numbers, but in terms of *characteristic matrices*. As consequence, there are multiple correspondences linked by a dual relation and, in addition to isometric (*Stratonovich-Weyl*) correspondences, we have the more general definition of *semi-conformal correspondences* as special cases of symbol correspondences, alongside the special cases of mapping-positive and positive-dual correspondences.

Future work shall be dedicated to the problem of asymptotic behavior of symbol correspondences and verifying the conditions under which the Poisson algebras of classical systems emerge as an asymptotic limit of operator algebras of quantum systems. In this respect, the first problem at hand is to decompose the Poisson bracket of  $\mathbb{C}P^2$ -harmonics, or  $\mathcal{E}$ -harmonics, similarly to what has been done in the case of spin systems and spherical harmonics.

Due to the similarity with spin systems, we hope that the asymptotic analysis of pure-quark systems shall be more feasible. On the other hand, generic quark systems seem to be quite more subtle for asymptotic analysis, since the relevant quantum systems are labeled by two indices,  $(p, q) \in (\mathbb{N} \times \mathbb{N}_0) \cup (\mathbb{N}_0 \times \mathbb{N})$ , so the construction of sequences of symbol correspondences, as done for spin systems, may involve some arbitrary choices (in principle, we would have to deal with bi-sequences of correspondences and study the asymptotic limit  $d \rightarrow \infty$ , where  $d = d(p, q)$  is

<sup>22</sup>The antipodal relation stems from the action of the longest element of the Weyl group, which opens the question of other possible relations associated to the action of other elements of the Weyl group.

the dimension of  $Q(p, q)$ ). Furthermore, different correspondences for the same generic quark systems may have different images, so it might be the case that we could generate sequences (or bi-sequences) of symbol correspondences whose images never reach some harmonic function  $f \in C^\infty(\mathcal{E})$ . Nonetheless, we hope that the factorization obtained in Proposition 2.27 may be useful in a similar approach to the one performed for spin systems in [18].

Another direction to be explored is the study of symbol correspondences from quantum quark systems to  $SU(3)$ -invariant Poisson manifolds, particularly  $S^7 \subset \mathbb{R}^8 \simeq \mathfrak{su}(3)$ . The 7-sphere can be split as  $S^7 \simeq M \cup N$ , where  $M$  is the disjoint union of two copies of  $\mathbb{C}P^2$  and  $N$  is an uncountable disjoint union of  $\mathcal{E}$  copies. In other words,  $\mathcal{E}$  and  $\mathbb{C}P^2$  are isomorphic to the symplectic leaves of  $S^7$ , with  $\mathcal{E}$  isomorphic to the regular leaves of this irregular foliation of  $S^7$ . In this respect, the first problem at hand is to understand how to “glue” the harmonic functions of  $\mathbb{C}P^2$  and  $\mathcal{E}$  in order to obtain  $SU(3)$ -equivariant smooth functions on  $S^7$ .

## 6. APPENDIX

In this appendix, we illustrate the general approach developed in [8] to prove Theorem 2.20, in a simple example. We know from Corollary 2.7 that

$$Q(1, 1) \otimes Q(1, 1) = Q(2, 2) \oplus Q(0, 3) \oplus Q(3, 0) \oplus Q(1, 1) \oplus Q(1, 1) \oplus Q(0, 0) .$$

Using [8, (1.2)], we find that

$$(6.1) \quad \psi_0 = PB_{21}x_1^1x_2^{2*}, \quad \psi_1 = PB_{12}x_1^2x_2^{1*}$$

are highest weight states of  $Q(1, 1) \oplus Q(1, 1)$ , where

$$(6.2) \quad B_{jk} = \sum_{l=1}^3 x_l^j x_l^{k*}$$

and the operator  $P$  is a projection onto the space of polynomials satisfying

$$(6.3) \quad \sum_{j=1}^3 \partial_j^k \partial_j^{k*} \psi = 0$$

for  $k = 1, 2$ . Then, we can write

$$(6.4) \quad \psi_j = \phi_j + \psi'_j ,$$

where  $\phi_j$  is  $\psi_j$  without  $P$  and  $\psi'_j$  contain terms with  $B_{11}$  and  $B_{22}$ .

Using the polynomial basis, we have

$$(6.5) \quad \mathbf{S} = \lambda \mathbf{1} + \mathbf{S}'$$

for some  $\lambda \in \mathbb{R}$  and

$$(6.6) \quad \mathbf{S}' = \frac{1}{2} \text{tr} \left( 3\overline{N}_1 N_2 - 3N_1 \overline{N}_2 - N_1 \overline{N}_1 N_2 + N_2 \overline{N}_2 N_1 + N_1 \overline{N}_1 \overline{N}_2 \right. \\ \left. - N_2 \overline{N}_2 \overline{N}_1 - \overline{N}_1 N_1 N_2 + \overline{N}_2 N_2 N_1 + \overline{N}_1 N_1 \overline{N}_2 - \overline{N}_2 N_2 \overline{N}_1 \right) ,$$

cf. [8, (2.13)-(2.14)], where

$$(6.7) \quad N_j = \begin{pmatrix} x_1^j \partial_1^j & x_1^j \partial_2^j & x_1^j \partial_3^j \\ x_2^j \partial_1^j & x_2^j \partial_2^j & x_2^j \partial_3^j \\ x_3^j \partial_1^j & x_3^j \partial_2^j & x_3^j \partial_3^j \end{pmatrix}, \quad \overline{N}_j = \begin{pmatrix} x_1^{j*} \partial_1^{j*} & x_2^{j*} \partial_1^{j*} & x_3^{j*} \partial_1^{j*} \\ x_1^{j*} \partial_2^{j*} & x_2^{j*} \partial_2^{j*} & x_3^{j*} \partial_2^{j*} \\ x_1^{j*} \partial_3^{j*} & x_2^{j*} \partial_3^{j*} & x_3^{j*} \partial_3^{j*} \end{pmatrix},$$

cf. [7, (2.9)].

From now on, we assume  $\mathbf{S}'$  acts only on  $\text{span}\{\psi_1, \psi_2\}$ . To compute the entries  $[\mathbf{S}']_{j,k}$ ,

$$(6.8) \quad \mathbf{S}'\psi_k = \sum_{j=0}^1 [\mathbf{S}']_{j,k} \psi_j,$$

we ignore terms with  $B_{11}$  and  $B_{22}$ . Thus substituting  $\psi_k$  for  $\phi_k$  in (6.8) and using only the terms

$$(6.9) \quad \text{tr}(\overline{N}_1 N_2) = B_{21} \sum_{j=1}^3 \partial_j^{1*} \partial_j^2, \quad \text{tr}(N_1 \overline{N}_2) = B_{12} \sum_{j=1}^3 \partial_j^1 \partial_j^{2*},$$

$$(6.10) \quad \begin{aligned} \text{tr}(N_1 \overline{N}_1 N_2) &= x_1^1 B_{21} \sum_{j=1}^3 \partial_j^1 \partial_j^{1*} \partial_1^2 + x_2^1 B_{21} \sum_{j=1}^3 \partial_j^1 \partial_j^{1*} \partial_2^2 \\ &\quad + x_3^1 B_{21} \sum_{j=1}^3 \partial_j^1 \partial_j^{1*} \partial_3^2, \end{aligned}$$

$$(6.11) \quad \begin{aligned} \text{tr}(N_2 \overline{N}_2 N_1) &= x_1^2 B_{12} \sum_{j=1}^3 \partial_j^2 \partial_j^{2*} \partial_1^1 + x_2^2 B_{12} \sum_{j=1}^3 \partial_j^2 \partial_j^{2*} \partial_2^1 \\ &\quad + x_3^2 B_{12} \sum_{j=1}^3 \partial_j^2 \partial_j^{2*} \partial_3^1, \end{aligned}$$

$$(6.12) \quad \begin{aligned} \text{tr}(N_1 \overline{N}_1 \overline{N}_2) &= x_1^{1*} B_{12} \sum_{j=1}^3 \partial_j^1 \partial_j^{1*} \partial_1^{2*} + x_2^{1*} B_{12} \sum_{j=1}^3 \partial_j^1 \partial_j^{1*} \partial_2^{2*} \\ &\quad + x_3^{1*} B_{12} \sum_{j=1}^3 \partial_j^1 \partial_j^{1*} \partial_3^{2*}, \end{aligned}$$

$$(6.13) \quad \begin{aligned} \text{tr}(N_2 \overline{N}_2 \overline{N}_1) &= x_1^{2*} B_{21} \sum_{j=1}^3 \partial_j^2 \partial_j^{2*} \partial_1^{1*} + x_2^{2*} B_{21} \sum_{j=1}^3 \partial_j^2 \partial_j^{2*} \partial_2^{1*} \\ &\quad + x_3^{2*} B_{21} \sum_{j=1}^3 \partial_j^2 \partial_j^{2*} \partial_3^{1*}, \end{aligned}$$

we get

$$(6.14) \quad \mathbf{S}' = \frac{1}{2} \begin{pmatrix} 7 & -2 \\ 2 & -7 \end{pmatrix}.$$

It is easy to see that  $\mathbf{S}'$  has a cyclic vector, so its eigenvalues are distinct (more precisely, in this case the eigenvalues of  $\mathbf{S}'$  are  $\pm 3\sqrt{5}/2$ ).

#### REFERENCES

- [1] P.A.S. Alcantara and P. de M. Rios. Asymptotic localization of symbol correspondences for spin systems and sequential quantizations of  $S^2$ . Preprint available at arXiv:2004.03929. 2021.

- [2] G. Alexanian et al. Fuzzy  $CP^2$ . In: *Journal of Geometry and Physics* 42.1-2 (2002), pp. 28–53.
- [3] G. E. Baird and L. C. Biedenharn. On the representations of the semisimple Lie groups. II. In: *Journal of Mathematical Physics* 4.12 (1963), pp. 1449–1466.
- [4] J. Bernatska and P. Holod. *Geometry and topology of coadjoint orbits of semisimple Lie groups*. Tech. rep. 2012.
- [5] R. Bott et al. “The geometry and representation theory of compact Lie groups”. In: *Representation Theory of Lie Groups*. London Mathematical Society Lecture Note Series. Cambridge University Press, 1980, pp. 65–90.
- [6] P. H. Butler. Coupling coefficients and tensor operators for chains of groups. In: *Philosophical Transaction of the Royal Society of London* 277.1272 (1975), pp. 545–585.
- [7] C. K. Chew and R. T. Sharp.  $SU(3)$  isoscalar factors. In: *Nuclear Physics B* 2 (1967), pp. 697–712.
- [8] C. K. Chew and R. T. Sharp. On the degeneracy problem in  $SU(3)$ . In: *Canadian Journal of Physics* 44 (1966), pp. 2789–2795.
- [9] S. Coleman. The Clebsch-Gordan series for  $SU(3)$ . In: *Journal of Mathematical Physics* 5.9 (1964), pp. 1343–1344.
- [10] J.-R. Derome and W. T. Sharp. Racah Algebra for an Arbitrary Group. In: *Journal of Mathematical Physics* 6.10 (1965), pp. 1584–1590.
- [11] G. B. Folland. *A Course in Abstract Harmonic Analysis*. Boca Raton, FL: CRC Press, Taylor & Francis, 2016. ISBN: 1498727131.
- [12] W. Greiner and B. Müller. *Quantum Mechanics*. Springer Berlin Heidelberg, 1994.
- [13] J. E. Humphreys. *Introduction to Lie Algebras and Representation Theory*. Springer New York, 1973. 196 pp.
- [14] A. A. Kirillov. *Lectures on the Orbit Method*. Providence, R.I: American Mathematical Society, 2004. ISBN: 9780821835302.
- [15] A. B Klimov and H. de Guise. General approach to  $SU(n)$  quasi-distribution functions. In: *Journal of Physics A* 43.40 (2010), p. 402001.
- [16] A. B. Klimov, J. L. Romero, and H. de Guise. Generalized  $SU(2)$  covariant Wigner functions and some of their applications. In: *Journal of Physics A* 50.32 (2017), p. 323001.
- [17] J. D. Louck. Recent progress toward a theory of tensor operators in the unitary groups. In: *American Journal of Physics* 38.1 (1970), pp. 3–42.
- [18] P. de M. Rios and E. Straume. *Symbol Correspondences for Spin Systems*. Springer, 2014. 212 pp. ISBN: 3319358111.
- [19] Alex Clésio Nunes Martins, Andrei B Klimov, and Hubert de Guise. Correspondence rules for Wigner functions over  $SU(3)/U(2)$ . In: *Journal of Physics A: Mathematical and Theoretical* 52.28 (2019), p. 285202.
- [20] R. Murgan and A. Zender. Energy eigenvalues of the three-dimensional quantum harmonic oscillator from  $SU(3)$  cubic Casimir operator. In: 40.1 (2018), p. 015405.
- [21] Z Pluhar, Yu F Smirnov, and V N Tolstoy. Clebsch-Gordan coefficients of  $SU(3)$  with simple symmetry properties. In: *Journal of Mathematical Physics A* 19.1 (1986), pp. 21–28.

- [22] Z Pluhar, L J Weigert, and P Holan. Symmetry properties of  $s$ -classified  $SU(3)$   $3j$ -,  $6j$ - and  $9j$ -symbols. In: *Journal of Mathematical Physics A* 19.1 (1986), pp. 29–34.
- [23] R. T. Sharp and H. von Baeyer. Polynomial bases and isoscalar factors for  $SU(3)$ . In: *Journal of Mathematical Physics* 7 (1966), p. 1105.
- [24] J. J. de Swart. The octet model and its Clebsch-Gordan coefficients. In: *Reviews of Modern Physics* 4 (1963), pp. 916–939.
- [25] D. A. Varshalovich, A. N. Moskalev, and V. K. Khersonskii. *Quantum Theory of Angular Momentum*. WSPC, 1988. 526 pp. ISBN: 9971509962.
- [26] E. Wigner. *Group Theory and its Application to the Quantum Mechanics of Atomic Spectra*. New York: Academic Press, 1959. ISBN: 9780127505503.
- [27] D. Zhelobenko. *Compact Lie Groups and Their Representations*. Providence: American Mathematical Society, 1973. ISBN: 9781470444556.

INSTITUTO DE CIÊNCIAS MATEMÁTICAS E DE COMPUTAÇÃO, UNIVERSIDADE DE SÃO PAULO.  
SÃO CARLOS, SP, BRAZIL.

*Email address:* `pedro.antonio.alcantara@usp.br`

*Email address:* `prios@icmc.usp.br`