# The Complete Picard Vessiot Closure

Andy R. Magid

ABSTRACT. Let F be a differential field with field of constants C, which we assume to be algebraically closed and of characteristic 0. The complete Picard– Vessiot closure  $F_{\infty}$  of F is a differential field extension of F with the same constants C as F, which has no Picard–Vessiot extensions, and is minimal over F with these properties. There is a correspondence between subfields of  $F_{\infty}$ and subgroups of its differential automorphism group, which arises because  $F_{\infty}$ comes from F via repeated Picard–Vessiot extensions. This correspondence persists for certain normal subfields of  $F_{\infty}$ , fields which can be characterized independently of their embedding in  $F_{\infty}$ .

### 1. Introduction

Let F be a field of characteristic zero. Polynomials over F have solutions in finite Galois extensions of F, and taking the compositum of all these produces a minimal field extension  $\overline{F} \supseteq F$  in which all polynomials over F have a full set of solutions. Moreover  $\overline{F}$  is a (generally infinite) Galois extension of F, and there is a Galois correspondence bijection between its lattice of subextensions and the lattice of closed subgroups of the profinite group of automorphisms of  $\overline{F}$  over F. Of course  $\overline{F}$  is the algebraic closure of F, and is itself algebraically closed: every polynomial over  $\overline{F}$  has a full set of solutions in  $\overline{F}$ .

The situation with differential fields is analogous but more complicated.

Let F be a differential field of characteristic zero and with algebraically closed field of constants C.

Linear differential equations over F have solutions in Picard–Vessiot extension fields of F, and taking a compositum of all of these produces a minimal differential field extension  $F_1 \supseteq F$  in which every linear differential equation over F has a full set of solutions. It may happen that not every linear differential equation over  $F_1$  has a solution in  $F_1$ , but by repeating the previous procedure we obtain an extension field  $F_2 \supseteq F_1$  in which the equations with  $F_1$  coefficients have solutions. Continuing in this fashion produces a tower  $F = F_0 \subseteq F_1 \dots F_i \subseteq F_{i+1} \dots$  whose union  $F_{\infty}$  has the property that every linear differential equation over  $F_{\infty}$  has a full set of solutions in  $F_{\infty}$ . We will see that  $F_{\infty}$  is minimal over F with respect to this property. (We require that none of the extensions of F introduce a new constant.)

The field  $F_{\infty}$  is a natural object of study when considering differential equations starting from F and their solutions. Among the things we might be concerned

<sup>2020</sup> Mathematics Subject Classification. 12H05.

with are the elements of  $F_{\infty}$ , its automorphisms, and the lattice structure of its differential subfields.

We will characterize the finitely generated differential extensions of F which embedd in  $F_{\infty}$ : these are the differential fields which can be embedded in a finite tower of Picard–Vessiot extensions over F (a finite tower in which each field is a Picard–Vessiot extension of its predecessor).

The group of differential automorphisms  $G(F_{\infty}/F)$  of  $F_{\infty}$  over F is the inverse limit of the groups  $G(F_i/F)$ . Further, the restriction from  $F_i$  to  $F_{i-1}$  induces a surjection  $G(F_i/F) \rightarrow G(F_{i-1}/F)$  whose kernel is  $G(F_i/F_{i-1})$ . This latter is a (pro)affine (pro)algebraic group, although neither  $G(F_i/F)$  nor  $G(F_{\infty}/F)$  need be.

If H is any subgroup of  $G(F_{\infty}/F)$  then the fixed field  $F_{\infty}^{H}$  is a differential subfield of  $F_{\infty}$  which contains F. We show that this observation has a full converse: every differential subfield K of  $F_{\infty}$  containing F is the fixed field of some subgroup of  $G(F_{\infty}/F)$ , and hence of  $G(F_{\infty}/K)$ . (This correspondence property – all subfields are fixed fields of subgroups – is called "semi–Galois" in the literature.)

The field  $F_{\infty}$  should be of special interest in the case F = C, and we include an example in this case.

We fix throughout the notation of this introduction. If  $E \supset F$  is a differential field extension, we use G(E/F) to denote the group of differential automorphisms of E over F. If S is a subset of E, we let  $F\langle S \rangle$  denote the smallest differential subfield of E that contains both F and S. If K is another differential subfield of E, then  $F\langle K \rangle$  is the field theoretic compositum FK. For unreferenced definitions and notation regarding the Picard–Vessiot theory, we refer to [2] and [6]. For general reference for Picard–Vessiot closure, we cite [3] and [6].

### 2. Universal Property

It is important to note that, despite the notation, the symbol  $F_{\infty}$  includes choices: Picard–Vessiot closures of  $F_i$  are not unique, although all are isomorphic [3, Theorem, p.6]. Then, using the automorphism lifting property [3, Theorem, p. 7], it is clear that any two choices of  $F_{\infty}$  are also isomorphic. In this section, we will show that  $F_{\infty}$  has an intrinsic characterization.

In the iterative construction of  $F_{i+1}$  as the Picard–Vessiot closure of  $F_i$  we have that the constants of  $F_{i+1}$  are those of  $F_i$ , that every linear homogeneous differential equation over  $F_i$  has a full set of solutions in  $F_{i+1}$ , and that  $F_{i+1}$  is generated as a differential field over  $F_i$  by solutions of such equations. We use these to characterize  $F_{\infty}$ :

THEOREM 1. The extension  $F_{\infty} \supseteq F$  satisfies

- (1) The constants of  $F_{\infty}$  are those of F.
- (2) Every every linear homogeneous differential equation over  $F_{\infty}$  has a full set of solutions in  $F_{\infty}$ .
- (3) If  $F_{\infty} \supseteq E \supseteq F$  is an intermediate differential subfield such that every linear homogeneous differential equation over E has a full set of solutions in E then  $E = F_{\infty}$ .

Moreover, any differential field  $K \supseteq F$  with the above properties is differentiably F isomorphic to  $F_{\infty}$ .

PROOF. Since  $F_{\infty} = \bigcup F_i$  and no new constants are introduced in any  $F_i$ , the constants of  $F_{\infty}$  are those of F. If  $Y^{(n)} + a_{n-1}Y^{(n-1)} + \cdots + a_1Y^{(1)} + a_0Y = 0$  is a differential equation over  $F_{\infty}$  then for some j we have  $a_i \in F_j$ ,  $0 \le i \le n-1$ . Then there is a full set of solutions in  $F_{j+1}$ , and hence in  $F_{\infty}$ . Now suppose  $F_{\infty} \supseteq E \supseteq F$  is an intermediate field such that every linear homogeneous differential equation over E has a full set of solutions in E. Then  $F_0 = F$  is contained in E, and if we have  $F_i$  contained in E then there is a full set of solutions in E for every linear homogeneous differential equation by such solutions, it follows that  $F_{i+1}$  is contained in E as well. Thus by induction  $F_{\infty}$  is contained in E, so they coincide.

Now suppose we have a differential field  $K \supseteq F$  with the three properties. We trivially have an F differential embedding  $f_0: F_0 \to K$ . Suppose we have an F embedding  $f_i: F_i \to K$ . Then we produce an F embedding  $f_{i+1}: F_{i+1} \to K$ extending  $f_i$ . We consider the set S of pairs (f, E) where  $F_{i+1} \supseteq E \supseteq F_i$  is an intermediate differential field and f extends  $f_i$ ; S is ordered in the obvious way and the union of any chain in  $\mathcal{S}$  is an upper bound for it in  $\mathcal{S}$ . Zorn's lemma then produces a maximal pair  $(f_{\mu}, E_{\mu})$ . If  $E_{\mu} \neq F_{i+1}$ , then there is Picard-Vessiot extension of  $F_i$  not contained in  $E_{\mu}$ , say for the equation  $Y^{(n)} + a_{n-1}Y^{(n-1)} +$  $\cdots + a_1 Y^{(1)} + a_0 Y = 0$ . Denote its left hand side by L. This means that the Picard–Vessiot extension  $E_L$  of  $E_{\mu}$  for L = 0 contained in  $F_{i+1}$  is proper. Let V denote a full set of solutions of L = 0 in  $E_L$  so that  $E_L = E \langle V \rangle$  Let  $f_{\mu}(L)$ denote  $Y^{(n)} + f_{\mu}(a_{n-1})Y^{(n-1)} + \cdots + f_{\mu}(a_1)Y^{(1)} + f_{\mu}(a_0)Y$ . By the properties of K,  $f_{\mu}(L) = 0$  has a full set of solutions W. Then there is an F embedding  $E_L = E_\mu \langle V \rangle \to f_\mu(E_\mu) \langle W \rangle$  which extends  $f_\mu$ . This is contrary to the maximality of  $f_{\mu}$ , and hence  $E_{\mu} = F_{i+1}$  and we can take  $f_{\mu}$  for  $f_{i+1}$ . By induction, we obtain consistent  $f_i$ 's for all *i*, and use them to define  $f_\infty : F_\infty \to K$ . Since  $f_\infty(F_\infty)$ has the property that every linear homogeneous equation over it has a full set of solutions in it, and K is the only subfield of K with this property, we have that  $f_{\infty}(F_{\infty}) = K$  and  $f_{\infty}$  is the desired isomorphism. 

We note that only role F plays in the three conditions of Theorem 1 is in the third, where the minimality condition is phrased over F. If E is a differential subfield of  $F_{\infty}$  containing F, then  $F_{\infty}$  also satisfies the minimality condition over E. Consequently,  $F_{\infty}$  is also  $E_{\infty}$ , a fact we now record:

COROLLARY 1. Let E be a differential subfield of  $F_{\infty}$  with  $F \subseteq E$ . Then  $F_{\infty} = E_{\infty}$ . In particular, all the fields  $E_i$  can be regarded as subfields of  $F_{\infty}$ .

The following lemma is a special case of Corollary 1 which we will need to use below.

LEMMA 1. Let E be a differential subfield of  $F_{\infty}$  with  $F \subseteq E$ , and let L be a monic linear differential operator over E. Then there is a differential subfield  $E_L \subseteq F_{\infty}$  with  $E \subseteq E_L$  such that  $E_L \supseteq E$  is a Picard–Vessiot extension for L.

For the proof, we note that we have  $E_1 \subseteq F_{\infty}$  by the Corollary and  $E_L \subseteq E_1$ .

### 3. Intermediate fields and the semi Galois correspondence

DEFINITION 1. A differential subfield E containing F of  $F_{\infty}$  will be called an intermediate field (of  $F_{\infty}$ ). An intermediate field is called normal (in  $F_{\infty}$ ) if it is preserved set-wise by every differential automorphism of  $F_{\infty}$  over F.

Let *E* be an intermediate field. By Corollary 1 we have that  $E_{\infty} = F_{\infty}$ . It follows that  $G(F_{\infty}/E) = G(E_{\infty}/E)$  and since *E* is the fixed field of the latter, it is also the fixed field of the former. Moreover, since differential automorphisms of *E* extend to  $E_{\infty}$  [3, Theorem, p. 7], differential automorphisms of *E* extend to  $F_{\infty}$  ("automorphism lifting property").

Now suppose  $E \subset M \subset K$  are intermediate fields, with K normal. By the normality of K, restriction defines a group homomorphism  $G(F_{\infty}/E) \to G(K/E)$ , which is surjective by the automorphism lifting property. Since E is the fixed field of  $G(F_{\infty}/E)$ , it follows that E is also the fixed field of G(K/E), and of course that M is the fixed field of the subgroup G(K/M) of G(K/E). In other words, the association  $M \mapsto G(K/M)$  is an injection from the set of differential subfields of K containing E to the set of subgroups of G(K/E), with right inverse  $H \mapsto K^H$ .

An injection of the sort just observed is half of the Galois correspondence, and has been termed a "semi–Galois correspondence" in the literature.

Once we have an internal description of when a differential field extension  $K \supseteq E$  can be embedded in  $F_{\infty}$  so that  $E \supseteq F$  and K is normal, we can state the semi–Galois correspondence as a theorem for such extensions. This we will do in Section 5 below.

## 4. Example

At this point we pause to consider an example. This example is based on the one considered in [3, pp 12-13]. We let F = C(t) be the field of rational functions over C with derivation t' = 1. Inside  $F_{\infty}$  we choose elements y and  $\{z_a \mid a \in C\}$  with  $y' = t^{-1}$  and  $z'_a = ((y+a)t)^{-1}$ . (In fact  $y \in F_1$  and  $z_a \in F_2$ ). Then  $F\langle y \rangle = F(y)$ , and  $F\langle z_a \rangle = F(y, z_a)$ ; we denote the latter by  $E_a$ . Note that  $F \subset F(y)$  and  $F(y) \subset E_a$  are Picard–Vessiot extensions, and  $F \subset E_a$  is an iterated Picard–Vessiot extension: a defining tower is  $F \subset F(y) \subset E_a$ . Let  $\mathcal{E} = \{E_a \mid a \in C\}$ . (The compositium  $E = F(y, \{z_a \mid a \in C\})$  of  $\mathcal{E}$  in  $F_{\infty}$  is, by Lemma 2 below, a LIPV extension of F.)

The differential equations of which y and  $z_a$  are solutions are, respectively, tY'' + Y' = 0 and (y + a)tY'' - (y + a + 1)Y' = 0. The coefficient t of the former is in the base field F (and a solution of the differential equation Y'' = 0), and the coefficient y + a + 1 of the latter is a solution of tY'' + Y' = 0, while the leading coefficient is a solution of tY''' + Y'' = 0. Thus the field E is generated by solutions of differential equations, whose coefficients are solutions of differential equations.

Let  $\sigma \in G(F_{\infty}/F)$ . For some notational convenience below, we will write the action on the right. Since both y and  $y^{\sigma}$  have derivative  $t^{-1}$ , we have  $y^{\sigma} = y + b(\sigma)$  for some  $b(\sigma) \in C$ . If also  $\tau \in G(F_{\infty}/F)$  then comparing  $y^{\sigma\tau}$  and  $(y^{\sigma})^{\tau}$  we see that  $b(\sigma\tau) = b(\sigma) + b(\tau)$ . Also  $(z_a^{\sigma})' = (z_a')^{\sigma} = ((y+a)t)^{-1})^{\sigma} = z'_{a+b(\sigma)}$ , from which it follows that  $z_a^{\sigma} = z_{a+b(\sigma)} + c(\sigma, a)$  for some  $c(\sigma, a) \in C$ . Comparing  $z_a^{\sigma\tau}$  and  $(z_a^{\sigma})^{\tau}$  we find that  $c(\sigma\tau, a) = c(\sigma, a + b(\tau)) + c(\tau, a)$ .

These calculations show that every  $\sigma$  in  $G(F_{\infty}/F)$  carries E to itself and hence that E is a normal LIPV extension of F (see definition 3 below). (In fact, they can be used to show that E is the smallest normal extension of F which contains  $E_0$ .)

The extensions  $F(y) \subset E_a$  are Picard–Vessiot as previously noted; this makes the compositum extension  $F(y) \subset E$  a locally Picard–Vessiot extension [4, p. 153] (or infinite Picard–Vessiot extension [3, p. 2]). The differential Galois group  $G(E_a/F(y))$  is the additive algebraic group  $\mathbb{G}_a$  (more precisely, its C points  $\mathbb{G}_a(C)$  are), and the differential Galois group G(E/F(y)) is the proalagebraic group  $\prod_{c \in C} \mathbb{G}_a$ . (One can see the latter by regarding the product as the set  $\operatorname{Map}(C, \mathbb{G}_a)$  of (all) functions from C to C and map G(E/F(y)) to it by  $\sigma \mapsto (a \mapsto c(\sigma, a))$ . If  $\sigma \in G(E/F(y))$  then  $\sigma(y) = y$  so  $b(\sigma) = 0$  and thus  $G(E/F(y)) \to \operatorname{Map}(C, \mathbb{G}_a)$  is a homomorphism.)

It follows that the extension  $F \subset E$  goes in steps  $F \subset F(y)$  and  $F(y) \subset E$  with each step (locally) Picard–Vessiot, but of course the extension  $F \subset E$  is not.

The group G(E/F) is the semidirect product  $\operatorname{Map}(C, \mathbb{G}_a) \rtimes \mathbb{G}_a$  where  $G_a = C$ acts on  $\operatorname{Map}(C, \mathbb{G}_a)$  by  $a \cdot f(x) = f(x+a)$ : we have  $G(E/F) \to \operatorname{Map}(C, \mathbb{G}_a) \rtimes \mathbb{G}_a$ by  $\sigma \mapsto (c(\sigma, \cdot), b(\sigma))$ .

As abstract groups the semidirect product  $\operatorname{Map}(C, \mathbb{G}_a) \rtimes \mathbb{G}_a$  is the regular wreath product  $\mathbb{G}_a \wr_r \mathbb{G}_a$ .

The semidirect product also has an algebraic (pro)variety structure: it is the product of the proaffine variety  $\prod_{c \in C} \mathbb{G}_a$  and the variety  $\mathbb{G}_a$ . This has as coordinate ring the polynomial ring  $C[\{X_c | c \in C\}, Y]$  where (reverting to semidirect product and the functional notation for the first factor)  $X_c((f, a)) = f(c)$  and Y((f, a)) = a. We use the "dot" notation for the action on functions on a group coming from translation actions on the group:  $\phi \cdot a(b) = \phi(ab)$  and  $b \cdot \phi(a) = \phi(ab)$  [1]. Then  $X_c \cdot (f, a)((g, b)) = X_c((f, a)(g, b)) = f(c) + g(a + c)$  so that

$$X_c \cdot (f, a) = f(c) + X_{c+a}.$$

Also,  $Y \cdot (f, a)((g, b)) = Y((f, a)(g, b)) = a + b$ , so that

 $Y \cdot (f, a) = a + Y.$ 

The displayed formulae show that left translation on  $\operatorname{Map}(C, \mathbb{G}_a) \rtimes \mathbb{G}_a$  by the fixed element (f, a) is a morphism of proaffine varieties.

Since  $(g, b) \cdot X_c((f, a)) = X_c((f, a)(g, b)) = f(c) + g(a + c)$ , however, we can see that right translation is not necessarily a morphism. Consider the case where  $C = \mathbb{C}$  is the complex numbers and  $g = \exp$  the exponential function. Then

$$((\exp, 0) \cdot X_0 - x_0)((f, a)) = \exp(a).$$

It is clear that  $(f, a) \mapsto \exp(a)$  is not in the polynomial ring  $C[\{X_c | c \in C\}, Y]$ .

#### 5. Characterization of intermediate fields

We we now return to an internal characterization of intermediate fields, and normal intermediate fields.

We begin with finitely generated extensions:

DEFINITION 2. A differential extension E of F is an iterated Picard–Vessiot (*IPV*) extension if there is a chain of differential subfields  $F = E_0 \subseteq E_1 \cdots \subseteq E_n = E$  such that for each  $i \in E_{i+1}$  is a Picard–Vessiot extension of  $E_i$ . We call the fields  $E_0, E_1, \ldots, E_n$  a defining tower for E.

The notation for defining towers of IPV extensions overlaps that for the tower of Picard–Vessiot closures. It will be clear from context, or noted, which one is intended.

Note that a Picard–Vessiot extension of an IPV extension of F is also an IPV extension of F. Since Picard–Vessiot extensions of F are finitely generated as fields, iterated Picard–Vessiot extensions of F are finitely generated over F.

#### ANDY R. MAGID

THEOREM 2. Let E be a differential subfield of  $F_{\infty}$  finitely generated over F. Then E is contained in an iterated Picard-Vessiot extension of F. Conversely, if  $E \supseteq F$  is a subfield of an iterated Picard-Vessiot extension then there is a differential embedding of E over F into  $F_{\infty}$ .

**PROOF.** Suppose X is a finite subset of  $F_{\infty}$ . Let k be the least i such that  $X \subset F_i$ . Let E be the differential field generated over F by X (note that  $E \subseteq F_i$ ). We prove that E is contained in an IPV by induction on k. If k = 0 then  $X \subset F$  and the assertion is trivial. So suppose k > 0 and that the assertion holds by induction for values less than k. Assume  $X \subset F_k$ . Since X is finite, there is a Picard–Vessiot extension  $M \supseteq F_{k-1}$  such that  $X \subset M$ . Each element of M is a ratio of elements which satisfy differential equations over  $F_{i-1}$ . Let  $X_0$  be a finite set of elements of  $F_i$  whose ratios contain X and such that each element  $x \in X_0$  satisfies a monic linear homogeneous differential operator  $L_x$  with coefficients in  $F_{i-1}$ . Let Y be the union of the sets of coefficients of the  $L_x$ . Note that Y is finite. Since  $Y \subset F_{i-1}$ , by induction the differential field N generated over F by Y is IPV. By Lemma 1, for each  $x \in X_0$  there is a Picard–Vessiot extension  $E_x$  of N for  $L_x$  contained in  $F_{\infty}$ . Note that  $x \in E_x$ . Let  $E_{X_0}$  be the compositum of the  $E_x$ 's. By [4, Cor. 10] p. 156],  $E_{X_0} \supseteq N$  is Picard–Vessiot (so  $E_{X_0} \supseteq F$  is IPV), and  $X_0 \subset E_{X_0}$  so also  $X \subset E_{X_0}$  and hence  $E \subseteq E_{X_0}$ . This proves the inductive step and shows that finitely generated extensions of F in  $F_{\infty}$  are contained in IPV subextensions.

Conversely, suppose that  $E \supseteq F$  is an IPV extension with defining tower  $F = E_0 \supseteq E_1 \supseteq \cdots \supseteq E_n$ . We prove by induction on n that E embeds over F into  $F_{\infty}$ . For n = 0, this is trivial. So suppose n > 0 and that the result holds for all smaller values. Then  $E_{n-1}$  embeds in  $F_{\infty}$  over F, and we may assume that in fact this embedding is inclusion. Suppose  $E_n \supseteq E_{n-1}$  is Picard–Vessiot for an operator L over  $E_{n-1}$ . By Lemma 1 there is a Picard–Vessiot extension M of  $E_{n-1}$  for L contained in  $F_{\infty}$ , and by uniqueness of Picard–Vessiot extensions [2, Theorem 3.5 p. 25]  $E_n$  and M are isomorphic over  $E_{n-1}$ ; this isomorphism is the desired embedding, and the result follows by induction. And if IPV extensions of F embed in  $F_{\infty}$ , so do their subfields.

Now we expand the definition to locally iterated Picard–Vessiot extensions:

DEFINITION 3. A differential extension E of F is a locally iterated Picard– Vessiot (LIPV) extension if every finite subset of E belongs to an iterated Picard– Vessiot subextension of F contained in E.

The analog of Theorem 2 applies for LIPV extensions. We will need to use the following lemma about composita of IPV extensions.

LEMMA 2. Let  $K \supseteq F$  be a differential extension with the same contants as F, and let  $\mathcal{I}$  be a set of locally iterated Picard–Vessiot subextensions of F in K. Then the compositum E of the fields in  $\mathcal{I}$  is locally iterated Picard–Vessiot.

PROOF. Since by definition a LIPV extension of F is the compositum of its IPV subextensions, it suffices to prove the lemma in the case where the elements of  $\mathcal{I}$  are IPV extensions. Moreover, if the lemma is true in the case that  $\mathcal{I}$  is finite, then E will be the union of its IPV subextensions, and hence LIPV. Hence it suffices to prove the lemma when  $\mathcal{I}$  consists of two subfields M and N with defining towers  $F = M_0 \subseteq \cdots \subseteq M_m = M$  and  $F = N_0 \subseteq \cdots \subseteq N_n = N$ . We prove that the compositum  $M_m N_n$  is IPV by induction on n. In the case  $n = 0, M_m F = M_m$  is

IPV over F by assumption. So suppose it is true for n-1, so that  $M_m N_{n-1}$  is IPV over F. Then since  $N_{n-1} \subseteq N_n$  is Picard–Vessiot, so is  $M_m N_{n-1} \subseteq M_m N_n$  (here is where we use that K has no new contants). Thus  $F \subseteq M_m N_n = MN$  is IPV as desired.

THEOREM 3. Let E be a differential subfield of  $F_{\infty}$ . Then E is contained in a locally iterated Picard-Vessiot extension of F. Conversely, if  $E \supseteq F$  is a subfield of an locally iterated Picard-Vessiot extension then there is a differential embedding of E over F into  $F_{\infty}$ .

PROOF. Let E be a differential subfield of  $F_{\infty}$ . Every finite subset of E belongs to an IPV extension of F by Theorem 2. Let  $\mathcal{I}$  be the set of these subfields; by Lemma 2 their compositum is LIPV.

Now suppose that E is a LIPV extension of F. We construct an embedding of E into  $F_{\infty}$  over F. Considering pairs of subfields of E and embeddings we can use Zorn's lemma to find a differential subfield  $E^0$  of E containing F and an embedding  $f^0: E^0 \to F_{\infty}$  over F which can't be further extended. Suppose  $E^0 \neq E$ , and  $a \in E, a \notin E^0$ . There is an IPV extension K of F in E containing a, and then there is a set in the defining tower of K with  $K_{i-1} \subseteq E^0$  but  $K_i \notin E^0$ . Consider the field  $E^1 = E^0 K_i$ : this is a Picard–Vessiot extension of  $E^0$  with respect to some operator L over  $E^0$ . Let  $f^0(L)$  denote the corresponding operator over  $f^0(E^0)$  obtained by applying  $f^0$  to the coefficients of L. By Lemma 1 we know that there is a Picard–Vessiot extension M of  $f^0(E^0)$  for  $f^0(L)$  contained in  $F_{\infty}$ , and then by [4, Proposition 18 p.161] that  $f^0$  extends to an isomorphism  $f^1$  from  $E^1$  to M. The field  $E^1$  and the map  $f^1: E^1 \to F_{\infty}$  contradict the maximality of  $E^0, f^0$ , and so we conclude that  $E^0 = E$ . Since LIPV extensions embedd over F into  $F_{\infty}$ , so do their subfields.

The extension argument used in the proof of Theorem 3 actually proves a bit more. Rather than simply show the existence of an embedding into  $F_{\infty}$  over F of the LIPV extension E, it shows that any embedding  $f^o: E^o \to F_{\infty}$  of a subextension  $E^o$  of E can be extended to all of E (we use a Zorn's Lemma argument to make  $E^0$  a maximal subfield containing  $E^o$  on which the original embedding  $f^o$  extends to an embedding  $f^0$  and then argue as in the theorem that  $E^0 = E$ ).

Now we turn to the definition of normality:

DEFINITION 4. A locally iterated Picard-Vessiot extension E of F is normal if whenever  $K \supseteq F$  is an extension with no new constants then all embeddings of Ein K over F have the same image.

Note that Picard–Vessiot extensions always have the normality property [2, Proposition 3.3 p. 24]. An interated Picard–Vessiot extension which is not Picard–Vessiot will not be normal.

We will need the following result about extensions of isomorphisms, which is also of independent interest:

PROPOSITION 1. Let  $K^1$  and  $K^2$  be locally iterated Picard-Vessiot extensions of F inside  $F_{\infty}$ , and suppose  $\tau : K^1 \to K^2$  is an F differential isomorphism. Then there is an F differential automorphism  $\sigma$  of  $F_{\infty}$  which restricts to  $\tau$  on  $K^1$ .

PROOF. Suppose  $\sigma: F_{\infty} \to F_{\infty}$  is an F differential embedding. Assume  $\sigma$  is not onto. Let  $E = F_{\infty}$ , let  $E^0 = \sigma(F_{\infty})$ , and let  $f^0 = \sigma^{-1}: E^0 \to F_{\infty}$ . The

### ANDY R. MAGID

extension argument in Theorem 3 (in the notation of that lemma) gives a field  $E^1$ ,  $F_{\infty} \supseteq E^1 \supset E^0$  and an embedding  $f^1 : E^1 \to F_{\infty}$ . If  $y \in E^1$ ,  $y \notin E^0$ , and  $x = \sigma(f^1(y))$ , then  $y - x \neq 0$  but  $f^1(y - x) = 0$ . This contradiction shows that any such  $\sigma$  is onto. Thus to prove the Proposition, it suffices to show that the embedding  $\tau : K^1 \to F_{\infty}$  extends to an embedding  $\sigma : F_{\infty} \to F_{\infty}$ . Since  $F_{\infty}$  is LIPV over F, this again follows by the extension argument of Theorem 3.

THEOREM 4. Let E be a locally iterated Picard-Vessiot extension of F contained in  $F_{\infty}$ . Then the following conditions are equivalent:

- (1) Every differential automorphism of  $F_{\infty}$  over F carries E to itself.
- (2) For any no new constants extension K of F, all differential embeddings  $E \to K$  over F have the same image.

PROOF. Suppose that  $\sigma(E) = E$  for all  $\sigma \in G(F_{\infty}/F)$ , and let  $\tau_i : E \to K$ , i = 1, 2, be embeddings over F into a no new constants extension. By Lemma 2 the compositum  $\tau_1(E)\tau_2(E)$  is a LIPV extension of F, and thus by Theorem 3 embedds over F into  $F_{\infty}$ . We compose the  $\tau_i$  with this embedding, so we may assume that  $K = F_{\infty}$ . Now consider  $\tau_1(E) \subseteq F_{\infty}$  and the embedding  $\tau = \tau_2 \tau_1^{-1} : \tau_1(E) \to F_{\infty}$ . By Proposition (1), there is  $\sigma \in G(F_{\infty}/F)$  which extends  $\tau$ . Since  $\sigma(\tau_1(E)) = \tau_1(E)$ by assumption, it follows that  $\tau_1(E) = \tau_2(E)$ , and so (1) implies (2).

Now suppose that for any no new constants extension K of F, all differential embeddings  $E \to K$  over F have the same image, and let  $\sigma \in G(F_{\infty}/F)$ . Let  $\tau_i : E \to F_{\infty}$  be defined as follows: let  $\tau_1$  be the restriction of  $\sigma$  to E and let  $\tau_2$  be the inclusion. Since by assumption  $\tau_1(E) = \tau_2(E)$ , we have  $\sigma(E) = E$  and so (1) implies (2).

Theorem 4 shows that the above definition of normal, Definition 4, coincides with that for intermediate fields, Definition 1. We have already observed the semi– Galois correspondence for the latter, which means we now have proven, and state as a theorem, the semi–Galois correspondence for the former:

THEOREM 5 (Fundamental Theorem for Normal LIPV Extensions). Let K be a normal locally iterated Picard-Vessiot extension of F. Let  $\mathcal{E}$  denote the set of differential subfields of K containing F ("set of intermediate fields") and let  $\mathcal{H}$ denote the set of closed subgroups of G(K/F). Then correspondences

$$\begin{array}{ccc} \mathcal{E} \to \mathcal{H} & and & \mathcal{H} \to \mathcal{E} \\ & by \\ E \mapsto G(K/E) & and & H \mapsto K^H \end{array}$$

are bijections.

### References

- Hochschild, G., and Mostow, G.D. Representations and representative functions of Lie groups, Ann. of Math. 66 (1957) 495–542.
- [2] Magid, A. Lectures on Differential Galois Theory, University Lecture Series 7, American Mathematical Society, Providence RI, 1997 (second printing with corrections).
- [3] Magid, A. The Picard-Vessiot Antiderivative Closure, J. of Algebra 244 (2001), 1–18.
- Magid, A. The Picard-Vessiot closure in differential Galois theory, Banach Center Publications 58, Polish Academy of Sciences, Warsaw, 2002 151–164
- [5] Magid, A. Left algebraic groups, J. of Algebra 35 (1975), 253–272

[6] van der Put, M. and Singer, M. Differential Galois Theory, Springer-Verlag, New York, 2003
DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN OK 73019