

**TOTAL STIEFEL WHITNEY CLASSES FOR REAL
REPRESENTATIONS OF GL_n OVER \mathbb{F}_q, \mathbb{R} AND \mathbb{C}**

JYOTIRMOY GANGULY, ROHIT JOSHI

ABSTRACT. We compute the total Stiefel Whitney class for a real representation π of $GL_n(\mathbb{F}_q)$, where q is odd in terms of character values of π on order 2 diagonal elements. We also compute the total Stiefel Whitney classes of real representations of $GL_n(\mathbb{R})$ and $GL_n(\mathbb{C})$.

CONTENTS

1. Introduction	1
2. Notation and Preliminaries	4
2.1. Irreducible Representations of C_m^n	4
2.2. Cohomology of $GL_n(\mathbb{F}_q)$, for $q \equiv 1 \pmod{4}$	5
3. SWC of S_n Invariant Representations of C_2^n	5
4. proof of main theorem	7
5. Detection Results	9
6. Calculations of Some SWC	11
6.1. Calculations Using the Main Theorem	11
6.2. Calculations of SWC Using Detection Results	14
7. Some Applications	16
7.1. Lifting Criteria for Real Representations of $GL_n(\mathbb{F}_q)$	16
7.2. Non-Detection by An-isotropic Torus	16
8. The Case of $GL_n(\mathbb{C})$ and $GL_n(\mathbb{R})$	17
9. Examples	17
9.1. Principal Series Representation	18
9.2. Cuspidal Representations	20
9.3. Steinberg Representation	20
10. Data Availability Statement	22
11. Conflict of Interest Statement	22
References	22

1. INTRODUCTION

Let π be a real representation of a finite group G . One can define Stiefel Whitney classes (SWC) $w_i(\pi)$ for $0 \leq i \leq \deg \pi$ as members of the cohomology groups $H^i(G) = H^i(G, \mathbb{Z}/2\mathbb{Z})$. The total Stiefel Whitney class of π in the cohomology ring $H^*(G)$ is defined as

$$w(\pi) = w_0(\pi) + w_1(\pi) + \cdots + w_d(\pi),$$

where $d = \deg \pi$. (see [Ben91, Section 2.6, page no. 50] for details).

In this article, we compute the total SWC for a real representation π of $GL_n(\mathbb{F}_q)$, where q is odd, in terms of the character values on diagonal elements of order 2. The computations for the particular case of $n = 2$ can be found in [GJ22]. We also compute the total SWC of a real representation of $GL_n(\mathbb{C})$ and $GL_n(\mathbb{R})$.

The computation of second SWC for representations of S_n and related groups can be found in [GS20]. Explicit expressions for the first and second SWC of the representations of $O_n(\mathbb{R})$ were found in [GJ21] in terms of highest weights.

We will now describe our results in some more detail. Let \mathfrak{X} denote the set of binary strings of length n . For $a = (a_1, a_2, \dots, a_n) \in \mathfrak{X}$, define $\nu(a) = \sum_{i=1}^n a_i \in \mathbb{Z}$. Put $\mathfrak{X}_i = \{a \in \mathfrak{X} \mid \nu(a) = i\}$. Let C_m denote the additive cyclic group of order m . We fix a generator x for C_m .

Consider the linear character χ^j of C_m given by $\chi^j(x) = \zeta_m^j$, where ζ_m is the primitive m th root of unity. For m even, write ‘sgn’ for $\chi^{m/2}$. For a binary string $a \in \mathfrak{X}$ and m even, consider the representation sgn_a of C_m^n , defined as $\text{sgn}_a = \boxtimes_{j=1}^n (\text{sgn})^{a_j}$ (external tensor product), where

$$(\text{sgn})^\epsilon = \begin{cases} \mathbb{1}, & \text{if } \epsilon = 0, \\ \text{sgn}, & \text{if } \epsilon = 1. \end{cases}$$

Here we use some terminologies from supercharacter theory. For more details we refer the reader to [DI08]. Consider the action of S_n on C_2^n by permuting the copies of C_2 . If we identify C_2^n with set of binary strings \mathfrak{X} , then the superclasses are \mathfrak{X}_i which are orbits of S_n action.

We will identify C_2^n with the subgroup

$$\text{diag}(\pm 1, \dots, \pm 1) \subset \text{GL}_n(F),$$

where $F = \mathbb{F}_q, \mathbb{R}$ or \mathbb{C} depending upon the context. We will write

$$h_i = \text{diag}(\underbrace{-1, \dots, -1}_{i \text{ times}}, 1, \dots, 1) \in \text{GL}_n(F).$$

Then h_i represents the superclass \mathfrak{X}_i . Consider the representations

$$\sigma_i = \bigoplus_{l \in \mathfrak{X}_i} \text{sgn}_l \tag{1}$$

of C_m^n for $1 \leq i \leq n$. The supercharacters are the direct sum of the representations in a single orbit for the action of S_n on characters of C_2^n . Then clearly the supercharacters are σ_i .

For $l \in \mathfrak{X}$, consider the linear character $\chi(l)$ of C_m^n given by $\chi(l) = \bigotimes_{k=1}^n \chi^{l_k}$. For $1 \leq i \leq n$, take $e_i \in \mathfrak{X}$ such that $e_{ij} = 1$ if $i = j$ and $e_{ij} = 0$ otherwise. If $4 \mid m$ then

$$H^*(C_m^n) = \frac{\mathbb{Z}/2\mathbb{Z}[s_1, \dots, s_n, t_1, \dots, t_n]}{(s_1^2, \dots, s_n^2)},$$

where $s_i = w_1(\text{sgn}_{e_i})$ and $t_i = w_2(\chi(e_i)_{\mathbb{R}})$. On the other hand

$$H^*(C_2^n) = \mathbb{Z}/2\mathbb{Z}[v_1, \dots, v_n],$$

where $v_i = w_1(\text{sgn}_{e_i})$. Let $v = (v_1, \dots, v_n)$ and $t = (t_1, \dots, t_n)$. For $l \in \mathfrak{X}$ we write $l \cdot v$ (resp. $l \cdot t$) for $\sum_{i=1}^n l_i v_i$ (resp. $\sum_{i=1}^n l_i t_i$).

For a real representation π of $\text{GL}_n(F)$ where $F = \mathbb{F}_q$ or \mathbb{R} or \mathbb{C} , consider the term

$$c_k = \frac{1}{2^n} \sum_{i=0}^n \chi_{\sigma_i}(h_k) \chi_{\pi}(h_i),$$

where $M = (\chi_{\sigma_i}(h_k))$ is the matrix which is exactly the supercharacter table. Thus we explicitly compute the table by proving that $\chi_{\sigma_i}(h_k)$ is the coefficient of y^i in the expression $(1-y)^k(1+y)^{n-k}$ (see Proposition 2).

For a real representation π of $\mathrm{GL}_n(\mathbb{F}_q)$ we define

$$\delta = \begin{cases} 0, & \text{if } \det \pi = 1, \\ 1, & \text{otherwise.} \end{cases}$$

For a subgroup H of G we denote by $w^H(\pi)$ (resp. $w_i^H(\pi)$) the total Stiefel Whitney class $w(\pi|_H) = \mathrm{res}_H^G(w(\pi))$ (resp. $w_i(\pi|_H)$). Let D denote the diagonal torus of $\mathrm{GL}_n(\mathbb{F}_q)$. We have $\mathrm{res} : H^*(\mathrm{GL}_n(\mathbb{F}_q)) \hookrightarrow H^*(D)$ for $q \equiv 1 \pmod{4}$ and $\mathrm{res} : H^*(\mathrm{GL}_n(\mathbb{F}_q)) \hookrightarrow H^*(C_2^n)$ for $q \equiv 3 \pmod{4}$ are injections (see [AM13, Theorem 4.4, page 227]). Our main theorem, proved in Section 4, says that

Theorem 1 (Main Theorem). *Let π be a real representation of $\mathrm{GL}_n(\mathbb{F}_q)$ where q is odd.*

(1) *If $q \equiv 1 \pmod{4}$ then*

$$w^D(\pi) = (1 + \delta b_1) \prod_{i=1}^n \left(\prod_{p \in \mathfrak{X}_i} (1 + p \cdot t) \right)^{c_i/2},$$

$$\text{where } p \cdot t = \sum_{i=1}^n p_i t_i \text{ and } b_1 = \sum_{i=1}^n s_i.$$

(2) *If $q \equiv 3 \pmod{4}$ then*

$$w^{C_2^n}(\pi) = \prod_{i=1}^n \left(\prod_{l \in \mathfrak{X}_i} (1 + l \cdot v) \right)^{c_i},$$

$$\text{where } l \cdot v = \sum_{i=1}^n l_i v_i.$$

We use the same approach to compute the total Stiefel Whitney class of real representations of $\mathrm{GL}_n(\mathbb{C})$ and $\mathrm{GL}_n(\mathbb{R})$. We have $\mathrm{res} : H^*(\mathrm{GL}_n(k)) \hookrightarrow H^*(C_2^n)$ is injective when $k = \mathbb{R}$ or \mathbb{C} .

Theorem 2. *Let π be a real representation of $\mathrm{GL}_n(\mathbb{C})$ or $\mathrm{GL}_n(\mathbb{R})$. Then the total Stiefel Whitney class of π is given by*

$$w^{C_2^n}(\pi) = \prod_{i=1}^n \left(\prod_{l \in \mathfrak{X}_i} (1 + l \cdot v) \right)^{c_i}. \quad (2)$$

In [Nak60, Corollary 6.7] Nakaoka proved that the cohomology of the symmetric group stabilizes. In particular, the result says that if $i < n/2$, then the map $\mathrm{res} : H^i(S_n) \rightarrow H^i(S_{2i})$ is an isomorphism. We prove a similar result for $\mathrm{GL}_n(\mathbb{F}_q)$.

Theorem 3. *The map $\mathrm{res} : H^i(\mathrm{GL}_m(\mathbb{F}_q)) \rightarrow H^i(\mathrm{GL}_{n-1}(\mathbb{F}_q))$, where $m \geq n$ and $q \equiv 1 \pmod{4}$, is injective for $i < 2n - 1$.*

We have a similar result for the case when $q \equiv 3 \pmod{4}$ (see Theorem 8). We also provide criteria for spinorality of real representations of $\mathrm{GL}_n(\mathbb{F}_q)$ where q is odd (see Theorem 14). This recovers the main result in [JS21, Theorem 1]

Theorem 3 allows further simplification to calculate the second and fourth Stiefel Whitney classes. Let π be a real representation of $\mathrm{GL}_n(\mathbb{F}_q)$, where q is odd. Then we have

$$w_2^D(\pi) = \frac{m_\pi}{2} \left(\sum_{i=1}^n t_i \right), \quad \text{if } q \equiv 1 \pmod{4},$$

$$w_2^{C_2^n}(\pi) = \binom{m_\pi}{2} \left(\sum_{i=1}^n v_i^2 \right), \quad \text{if } q \equiv 3 \pmod{4},$$

where $m_\pi = \frac{\dim \pi - \chi_\pi(h_1)}{2}$. Moreover, if $q \equiv 1 \pmod{4}$, then

$$w_4^D(\pi) = \binom{m_\pi/2}{2} \sum_{i=1}^n t_i^2 + \frac{\dim \pi - \chi_\pi(h_2)}{8} \sum_{1 \leq i < j \leq n} t_i t_j.$$

As an application of the result we obtain the following property for the real principal series representations (see Section 9.1).

Theorem 4. *Let π be a real principal series representation for $\mathrm{GL}_n(\mathbb{F}_q)$ where $q \equiv 1 \pmod{4}$. Then $w_4(\pi) = 0$ for $n \geq 5$.*

We also compute the second SWC for q odd and the fourth SWC for $q \equiv 1 \pmod{4}$.

We now describe the layout of the paper. In Section 2 recall some basic facts on cohomology of the groups $\mathrm{GL}_n(\mathbb{F}_q)$ and C_m^n . In Section 3 we compute the SWC for S_n invariant real representations of C_2^n . We prove the main theorem in Section 4. The theorems 3 and 8 are proved in Section 5. We compute the second and the fourth SWC in Section 6 using the previous results. Section 7 shows some applications of the theory. Section 8 deals with the cases of $\mathrm{GL}_n(\mathbb{C})$ and $\mathrm{GL}_n(\mathbb{R})$. We work out some examples in Section 9.

Acknowledgements: The authors would like to thank Steven Spallone for helpful conversations. The first author of this paper was supported by a post-doctoral fellowship from Chennai Mathematical Institute (CMI). The second author of this paper was supported by a post-doctoral fellowship of National Board of Higher Mathematics India (NBHM).

2. NOTATION AND PRELIMINARIES

A complex representation (π, V) of a finite group G is called orthogonal if it preserves a non degenerate symmetric bilinear form. We call π real if the representation space V is real. For a complex representation π one can consider its realisation $(\pi_{\mathbb{R}}, V_{\mathbb{R}})$ by forgetting the complex structure of V and regarding it as a real vector space.

2.1. Irreducible Representations of C_m^n . Take the set of ordered pairs

$$\mathcal{P} = \{ \tau = (\tau_1, \dots, \tau_n) \mid \tau_j \in C_m, 1 \leq j \leq n \}.$$

Consider the subset \mathcal{P}' of \mathcal{P} given by

$$\mathcal{P}' = \{ \tau = (\tau_1, \dots, \tau_n) \mid \tau_j \in \{0, m/2\}, 1 \leq j \leq n \}.$$

Consider the linear character $\chi_\tau = (\chi^{\tau_1} \otimes \chi^{\tau_2} \otimes \dots \otimes \chi^{\tau_n})$ of C_m^n (See Section 2)). We have $\chi_{-\tau} = \chi_\tau^\vee$. Note that χ_τ is real if $\tau \in \mathcal{P}'$. Define an equivalence relation on \mathcal{P} as

$$\tau \sim (-\tau_1, -\tau_2, \dots, -\tau_n).$$

Put $\widehat{\mathcal{P}} = (\mathcal{P} - \mathcal{P}') / \sim$. Therefore a real irreducible representations of C_m^n is either of the form $(\chi_\tau)_{\mathbb{R}}$ for $\tau \in \widehat{\mathcal{P}}$ or of the form χ_τ where $\tau \in \mathcal{P}'$.

2.2. Cohomology of $GL_n(\mathbb{F}_q)$, for $q \equiv 1 \pmod{4}$. The cohomology ring of $GL_n(\mathbb{F}_q)$, where $q \equiv 1 \pmod{4}$ is given by

$$H^*(GL_n(\mathbb{F}_q), \mathbb{Z}/2\mathbb{Z}) = \mathbb{Z}/2\mathbb{Z}[a_2, a_4, \dots, a_{2n}] \otimes E[b_1, b_3, \dots, b_{2n-1}], \quad (3)$$

where

$$a_{2l} = \sum_{i_1 < i_2 < \dots < i_l} t_{i_1} \cdots t_{i_l}, \quad b_{2l-1} = \sum_{i_1 < i_2 < \dots < i_l, 1 \leq j \leq l} t_{i_1} \cdots \hat{t}_{i_j} \cdots t_{i_l} s_{i_j}, \quad 1 \leq l \leq n,$$

and $E[b_1, \dots, b_{2n-1}]$ denotes the alternating algebra. For details we refer the reader to [FP06, Theorem 8.1, page 288].

We know that the symmetric group S_n is the Weyl group of $GL_n(\mathbb{F}_q)$ with respect to the diagonal torus D . It acts on D by permuting the diagonal entries.

Proposition 1. *Let π be a representation of $GL_n(\mathbb{F}_q)$. Then $\pi|_D$ is S_n invariant.*

Proof. We leave the proof to the reader. \square

This enables us to work with S_n invariant representations of D .

Definition 1. *Let G' be a subgroup of a finite group G . We say that G' detects the (mod 2) cohomology of G , provided that the restriction map*

$$\text{res} : H^*(G, \mathbb{Z}/2\mathbb{Z}) \rightarrow H^*(G', \mathbb{Z}/2\mathbb{Z}) \quad (4)$$

is injective.

The action of S_n on D induces an action of S_n on $H^*(D)$. We write $H^*(D)^{S_n}$ to denote the subalgebra fixed under the action.

Theorem 5 (Corollary of Theorem 3, page 571 Quillen). *The diagonal subgroup D detects the cohomology of $GL_n(\mathbb{F}_q)$. In fact, the homomorphism*

$$H^*(GL_n(\mathbb{F}_q)) \rightarrow H^*(D)$$

induced from the restriction of $GL_n(\mathbb{F}_q)$ to D is injective. Moreover, the image lies inside $H^(D)^{S_n}$.*

3. SWC OF S_n INVARIANT REPRESENTATIONS OF C_2^n

An S_n invariant representation of C_2^n will be of the form

$$\pi = \bigoplus_{i=0}^n c_i \sigma_i, \quad (5)$$

where σ_i is as in (1).

Proposition 2. *The character value $\chi_{\sigma_i}(h_k)$ is equal to the coefficient of y^i in the expression $(1-y)^k(1+y)^{n-k}$.*

Proof. Fix two integers i, k such that $0 \leq i, k \leq n$. Let R_l denote the set of elements $a \in \mathfrak{X}_i$ such that the first k coordinates of a contains l number of 1's. Then for $a \in R_l$ we have $\chi_{\text{sgn}_a}(h_k) = (-1)^l$. Put $\tau_l = \bigoplus_{a \in R_l} \text{sgn}_a$. Then

$$\chi_{\tau_l}(h_k) = \binom{k}{l} \binom{n-k}{i-l} (-1)^l.$$

Observe that $\bigoplus_{l=0}^k \tau_l = \sigma_i$. Thus

$$\chi_{\sigma_i}(h_k) = \sum_{l=0}^k \binom{k}{l} \cdot \binom{n-k}{i-l} (-1)^l. \quad (6)$$

Note that this gives the coefficient of y^i in $(1-y)^k(1+y)^{n-k}$. □

From (5) we have

$$\chi_\pi(h_i) = \sum_{j=0}^n c_j \cdot \chi_{\sigma_j}(h_i) \quad (7)$$

Define matrix $M = (m_{ij})$ where $0 \leq i, j \leq n$ such that

$$m_{ij} = \chi_{\sigma_j}(h_i). \quad (8)$$

Take $a = {}^t(\chi_\pi(h_0), \dots, \chi_\pi(h_n))$ and $m = {}^t(c_0, \dots, c_n)$, where tA denotes the transpose of the matrix A . Observe that

$$a = M \cdot m. \quad (9)$$

Theorem 6. *We have*

$$M^2 = 2^n I_{n+1},$$

where I_{n+1} denotes the identity matrix of rank $n+1$.

Proof. Let

$$R_i = (m_{i0}, \dots, m_{in}),$$

be the i -th row of M . Note that from Proposition 2 the element m_{il} , $0 \leq l \leq n$, is equal to the coefficient of y^l in $(1-y)^i(1+y)^{n-i}$.

We calculate the x -th row in the matrix M^2 .

$$\begin{aligned} \sum_{l=0}^n (M^2)_{xl} y^l &= \sum_{l=0}^n \left(\sum_{i=0}^n m_{xi} m_{il} y^l \right) \\ &= \sum_{i=0}^n m_{xi} \left(\sum_{l=0}^n m_{il} y^l \right) \\ &= \sum_{i=0}^n \chi_{\sigma_i}(h_x) (1-y)^i (1+y)^{n-i} \\ &= \sum_{i=0}^n \left(\sum_{l=0}^x \binom{x}{l} \cdot \binom{n-x}{i-l} (-1)^l \right) (1-y)^i (1+y)^{n-i} \\ &= \sum_{l=0}^x \binom{x}{l} (-1)^l \left(\sum_{i=0}^n \binom{n-x}{i-l} (1-y)^i (1+y)^{n-i} \right) \\ &= \sum_{l=0}^x \binom{x}{l} (-1)^l \left(\sum_{t=0}^{n-x} \binom{n-x}{t} (1-y)^{t+l} (1+y)^{n-t-l} \right) \\ &= \sum_{l=0}^x \binom{x}{l} (-1)^l (1-y)^l \left(\sum_{t=0}^{n-x} \binom{n-x}{t} (1-y)^t (1+y)^{n-t-l-x+x} \right) \\ &= \sum_{l=0}^x \binom{x}{l} (-1)^l (1-y)^l (1+y)^{x-l} \left(\sum_{t=0}^{n-x} \binom{n-x}{t} (1-y)^t (1+y)^{n-x-t} \right) \\ &= \sum_{l=0}^x \binom{x}{l} (y-1)^l (y+1)^{x-l} (1-y+1+y)^{n-x} \\ &= (2y)^x 2^{n-x} \\ &= 2^n y^x. \end{aligned}$$

□

Since M is invertible, we have

$$m = M^{-1} \cdot a. \quad (10)$$

Proposition 3. *Let π be the representation of C_2^n as in equation (5). Then we have*

$$c_k = \frac{1}{2^n} \sum_{i=0}^n \chi_{\sigma_i}(h_k) \chi_{\pi}(h_i).$$

Proof. The proof follows from Equations (8) (10) and Theorem 6. \square

Theorem 7. *Let π be an S_n invariant representation of C_2^n of the form $\pi = \bigoplus_{i=1}^n c_i \sigma_i$. Then we have*

$$w(\pi) = \prod_{i=1}^n \left(\prod_{l \in \mathfrak{X}_i} (1 + l \cdot v) \right)^{c_i} \quad (11)$$

where $c_i = \frac{1}{2^n} \sum_{k=0}^n \chi_{\sigma_k}(h_i) \chi_{\pi}(h_k)$.

Proof. We have

$$w(\sigma_i) = \prod_{l \in \mathfrak{X}_i} (1 + l \cdot v)$$

and

$$w(\pi) = \prod_{i=1}^n (w(\sigma_i))^{c_i}.$$

Now from Proposition 3 we have

$$c_i = \frac{1}{2^n} \sum_{k=0}^n \chi_{\sigma_k}(h_i) \chi_{\pi}(h_k).$$

Hence the result follows. \square

4. PROOF OF MAIN THEOREM

For a subset $I = \{i_1, i_2, \dots, i_l\}$ of $\{1, 2, \dots, n\}$ and for a multiset $J = \{j_1, j_2, \dots, j_k\}$ where $j_i \in \{1, 2, \dots, n\}$ we define

$$s^I = s_{i_1} \cdots s_{i_l} \text{ and } t^J = t_{j_1} t_{j_2} \cdots t_{j_k}.$$

Observe that $H^*(C_{4m}^n) = \frac{\mathbb{Z}/2\mathbb{Z}[s_1, \dots, s_n, t_1, \dots, t_n]}{(s_1^2, \dots, s_n^2)}$ is a $\mathbb{Z}/2\mathbb{Z}$ vector space with basis $s^I t^J$, where I (resp. J) varies over all subsets (resp. multi-sets) over $\{1, 2, \dots, n\}$.

Proposition 4. *Let $q \equiv 1 \pmod{4}$. If π is a real representation of $GL_n(\mathbb{F}_q)$ and if $w(\pi) = \sum_{I, J} a_{I, J} s^I t^J$, then elements of the form s^I have coefficient 0 i.e. $a_{I, \emptyset} = 0$ if $|I| \geq 2$.*

Proof. Consider the projection map

$$\text{pr} : \frac{\mathbb{Z}/2\mathbb{Z}[s_1, \dots, s_n, t_1, \dots, t_n]}{(s_1^2, \dots, s_n^2)} \rightarrow \frac{\mathbb{Z}/2\mathbb{Z}[s_1, \dots, s_n]}{(s_1^2, \dots, s_n^2)}$$

defined by $\text{pr}(s_i) = s_i$ and $\text{pr}(t_i) = 0$ for all i . This gives $\text{pr}(a_{2i}) = 0$ for all i and $\text{pr}(b_{2i-1}) = 0$ for $i > 1$. Then from Equation (3) we have

$$\text{pr}(H^*(GL_n(\mathbb{F}_q))) = \mathbb{Z}/2\mathbb{Z}[b_1]. \quad (12)$$

Now if s^I appears as a term in $w(\pi)$, where $|I| \geq 2$, then it survives in $\text{pr}(w(\pi))$. But this contradicts Equation (12). \square

Proof of Main Theorem 1. Let \widehat{P} be as defined in Section 2. Let π be a real representation of $\text{GL}_n(\mathbb{F}_q)$. Recall that for a subgroup H of $\text{GL}_n(\mathbb{F}_q)$ we write $w_i^H(\pi)$ and $w^H(\pi)$ to denote $w_i(\pi|_H)$ and $w(\pi|_H)$ respectively.

Since $\pi|_D$ is an S_n invariant representation of D we write

$$\pi|_D = \mathbb{1}^{\oplus m_0} \bigoplus_{i=1}^n \sigma_i^{\oplus m_i} \bigoplus_{\tau \in \widehat{P}} ((\chi_\tau)_\mathbb{R})^{\oplus m_\tau}. \quad (13)$$

Therefore we may write

$$\begin{aligned} w^D(\pi) &= w(\mathbb{1}^{\oplus m_0} \bigoplus_{i=1}^n \sigma_i^{\oplus m_i}) \cdot w\left(\bigoplus_{\tau \in \widehat{P}} ((\chi_\tau)_\mathbb{R})^{\oplus m_\tau}\right) \\ &= (1 + Q(s_1, \dots, s_n))(1 + P(t_1, \dots, t_n)) \quad (\text{where } P, Q \text{ do not have constant term}), \\ &= (1 + \delta b_1)(1 + P(t_1, \dots, t_n)) \quad (\text{follows from Proposition 4}), \end{aligned}$$

This calculation shows that $w_1^D(\pi) = \delta b_1$. From this we conclude that

$$\delta = \begin{cases} 0, & \text{if } \det \pi = 1, \\ 1, & \text{otherwise.} \end{cases}$$

Consider the representation $\psi = (\chi^1)_\mathbb{R}$ of C_{q-1} . Let $C_2 < C_{q-1}$ be the subgroup of order 2. Then the restriction map $\text{res} : H^*(C_{q-1}) \rightarrow H^*(C_2)$ maps $t = w_2(\psi)$ to $w_2^{C_2}(\psi) = w_2(\text{sgn} \oplus \text{sgn}) = v^2$. Thus the restriction map $\text{res}_n : H^*(C_{q-1}^n) \rightarrow H^*(C_2^n)$ maps t_i to v_i^2 and s_i to 0.

$$w^{C_2^n}(\pi) = 1 + P(v_1^2, \dots, v_n^2) \quad (14)$$

$$= (1 + P(v_1, \dots, v_n))^2. \quad (15)$$

On the other hand $\pi|_{C_2^n}$ is an S_n invariant representation, therefore by Theorem 7 we have

$$w^{C_2^n}(\pi) = \prod_{i=1}^n \left(\prod_{l \in \mathfrak{X}_i} (1 + \sum_j l_j v_j) \right)^{c_i},$$

where $c_i = \frac{1}{2^n} \sum_{k=0}^n \chi_{\sigma_k}(h_i) \chi_\pi(h_k)$. Now since $H^*(C_2^n) = \mathbb{Z}/2\mathbb{Z}[v_1, \dots, v_n]$ is a

Unique Factorization domain, we have $1 + P(v_1, \dots, v_n) = \prod_{i=1}^n \left(\prod_{l \in \mathfrak{X}_i} (1 + \sum_j l_j v_j) \right)^{c_i/2}$.

Thus we obtain

$$w^D(\pi) = (1 + \delta b_1) \prod_{i=1}^n \left(\prod_{l \in \mathfrak{X}_i} (1 + l \cdot t) \right)^{c_i/2},$$

where $l \cdot t = \sum_j l_j t_j$.

In the case $q \equiv 3 \pmod{4}$ we have

$$\text{res}_{C_2^n}^D : H^*(D) \rightarrow H^*(C_2^n)$$

is an isomorphism as C_2^n is the 2 Sylow subgroup of D . (see [GJ22, Propositions 5 and 6]). Now we apply Theorem 7 to obtain

$$w^{C_2^n}(\pi) = \prod_{i=1}^n \left(\prod_{l \in \mathfrak{X}_i} (1 + \sum_j l_j v_j) \right)^{c_i},$$

where again $c_i = \frac{1}{2^n} \sum_k \chi_{\sigma_k}(h_i) \chi_{\pi}(h_k)$. This completes the proof of the main theorem. \square

Remark 1. Since c_i is the multiplicity of σ_i in $\pi|_{C_2^n}$, it is an integer, where σ_i is a representation of C_2^n . Moreover, if $q \equiv 1 \pmod{4}$, then $\sigma_j|_{C_2^n} = \binom{n}{j} \mathbb{1}$, where σ_j is a representation of D . Also $(\chi)_{\mathbb{R}}|_{C_2^n} = 2\chi|_{C_2^n}$. Therefore the integers c_i are even for $i \geq 1$.

Remark 2. For a real representation of $GL_n(\mathbb{F}_q)$, where $q \equiv 1 \pmod{4}$, one can write the expression for $w(\pi)$ in terms of the generators of $H^*(GL_n(\mathbb{F}_q))$, namely a_{2l}, b_{2l-1} for $1 \leq l \leq n$. This follows from the fact that the terms a_{2l} are elementary symmetric functions in t_i for $1 \leq i \leq n$. For example if $n = 3$, then

$$\begin{aligned} w(\pi) &= \left(1 + \delta \cdot \sum_{i=1}^3 s_i \right) ((1+t_1)(1+t_2)(1+t_3))^{m_1} \\ &\quad ((1+t_1+t_2)(1+t_1+t_3)(1+t_2+t_3))^{m_2} \\ &\quad (1+t_1+t_2+t_3)^{m_3}. \end{aligned}$$

Therefore one has

$$\begin{aligned} w(\pi) &= (1 + \delta \cdot b_1)(1 + a_2 + a_4 + a_6)^{m_1} \\ &\quad (1 + a_2^2 + 2a_2 + a_4 + a_2a_4 - a_6)^{m_2} \\ &\quad (1 + a_2)^{m_3}. \end{aligned}$$

5. DETECTION RESULTS

We write

$$E[x_1, \dots, x_n] = \frac{\mathbb{Z}/2\mathbb{Z}[x_1, \dots, x_n]}{(x_1^2, \dots, x_n^2)}.$$

Let D_n denote the diagonal torus of $GL_n(\mathbb{F}_q)$, where $q \equiv 1 \pmod{4}$. Take $m \geq n$. We have

$$H^*(D_m) = \frac{\mathbb{Z}/2\mathbb{Z}[t_1, \dots, t_m, s_1, \dots, s_m]}{(s_1^2, \dots, s_m^2)},$$

and

$$H^*(D_{n-1}) = \frac{\mathbb{Z}/2\mathbb{Z}[t'_1, \dots, t'_{n-1}, s'_1, \dots, s'_{n-1}]}{(s'_1{}^2, \dots, s'_{n-1}{}^2)}.$$

Consider the inclusion map $j_n : GL_{n-1}(\mathbb{F}_q) \rightarrow GL_m(\mathbb{F}_q)$ given by

$$j_n(A) = \begin{pmatrix} A & 0 \\ 0 & I_{m-n+1} \end{pmatrix},$$

where $A \in GL_{n-1}(\mathbb{F}_q)$. It will induce a map on the diagonal subgroups namely $j_n : D_{n-1} \rightarrow D_m$ (we use the same notation for the induced map). We have

$$j_n^*(t_l) = \begin{cases} t'_l, & \text{if } l < n \\ 0, & \text{if } l \geq n \end{cases}$$

Similarly

$$j_n^*(s_l) = \begin{cases} s'_l, & \text{if } l < n \\ 0, & \text{if } l \geq n \end{cases}$$

It will induce a map

$$j_n^* : \mathbb{Z}/2\mathbb{Z}[a_2, \dots, a_{2m}] \otimes E[b_1, b_3, \dots, b_{2m-1}] \rightarrow \mathbb{Z}/2\mathbb{Z}[a'_2, \dots, a'_{2n-2}] \otimes E[b'_1, \dots, b'_{2n-3}],$$

where

$$j_n^*(a_{2l}) = a'_{2l} = \sum_{1 \leq i_1 < i_2 < \dots < i_l \leq n-1} t'_{i_1} \cdots t'_{i_l},$$

for $l < n$ and

$$j_n^*(b_{2k-1}) = b'_{2k-1} = \sum_{1 \leq j \leq k} \left(\sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n-1} t'_{i_1} \cdots t'_{i_j} \cdots t'_{i_k} s'_{i_j} \right),$$

for $k < n$. Therefore $j_n^*(a_{2l}) = 0$ when $l \geq n$ and $j_n^*(b_{2k-1}) = 0$ for $k \geq n$.

Proof of Theorem 3. Put $R_m = \mathbb{Z}/2\mathbb{Z}[a_2, \dots, a_{2m}]$, $E_m = E[b_1, b_3, \dots, b_{2m-1}]$, $R'_{n-1} = \mathbb{Z}/2\mathbb{Z}[a'_2, \dots, a'_{2n-2}]$ and $E'_{n-1} = E[b'_1, b'_3, \dots, b'_{2n-3}]$. Since the elements a_2, \dots, a_{2n-2} are algebraically independent, we have

$$\ker j_n^* |_{R_m} = \langle a_{2n}, a_{2n+2}, \dots, a_{2m} \rangle.$$

Take $R_m = \bigoplus_i A_i$ and $R'_{n-1} = \bigoplus_i A'_i$, where A_i and A'_i are the i -th degree components. Then comparing degrees we have $A_i \cap \ker j_n^* = 0$, for $i < 2n$. So the map $j_n^* : A_i \rightarrow A'_i$ is injective for $i < 2n - 1$.

Note that $E[b_1, b_3, \dots, b_{2m-1}]$ is a $\mathbb{Z}/2\mathbb{Z}$ vector space with basis elements

$$b_{2i_1-1} \cdots b_{2i_k-1},$$

where $1 \leq i_1 < i_2 < \dots < i_k \leq m$ and $1 \leq k \leq m$. In each basis element a particular b_j can have exponent 1 or 0. Therefore $\dim E[b_1, b_3, \dots, b_{2m-1}] = 2^m$. Similarly $\dim E[b'_1, b'_3, \dots, b'_{2n-3}] = 2^{n-1}$. Since j_n^* is surjective we have $\dim \ker j_n^* |_{E_m} = 2^m - 2^{n-1}$.

Consider the ideal $\mathfrak{J} = \langle b_{2n-1}, b_{2n+1}, \dots, b_{2m-1} \rangle$ of E_m . As a $\mathbb{Z}/2\mathbb{Z}$ vector space $\dim \mathfrak{J} = 2^m - 2^{n-1}$. Also for $x \in \mathfrak{J}$ one has $j_n^*(x) = 0$. Therefore we obtain

$$\ker j_n^* |_{E_m} = \langle b_{2n-1}, b_{2n+1}, \dots, b_{2m-1} \rangle.$$

Let $E_m = \bigoplus_i B_i$ and $E'_{n-1} = \bigoplus_j B'_j$. Comparing the degrees we get $\ker j_n^* \cap B_i = 0$ for $i < 2n - 1$. So the map $j_n^* : B_i \rightarrow B'_i$ is injective for $i < 2n - 1$.

The i th graded part of $R_m \otimes E_m$ is of the form $\bigoplus_{k+l=i} A_k \otimes B_l$. Similarly, the i th graded part of $R'_{(n-1)} \otimes E'_{(n-1)}$ is of the form $\bigoplus_{k+l=i} A'_k \otimes B'_l$. To show the injectivity of the map j_n^* it is enough to show that the map $j_n^* : A_k \otimes B_{i-k} \rightarrow A'_k \otimes B'_{i-k}$ is injective for every $0 \leq k \leq i < 2n - 1$. It follows from the fact that the maps $j_n^* : A_k \rightarrow A'_k$ and $j_n^* : B_k \rightarrow B'_k$ are injective for $0 \leq i < 2n - 1$. \square

Theorem 8. *The map $\text{res} : H^i(\text{GL}_m(\mathbb{F}_q)) \rightarrow H^i(\text{GL}_{n-1}(\mathbb{F}_q))$, where $m \geq n$ and $q \equiv 3 \pmod{4}$ is injective for $i < n$.*

Proof. From Theorem 5 we obtain that the map $\text{res}_m : H^i(\text{GL}_m(\mathbb{F}_q)) \rightarrow H^i(D_m)^{S_m}$ is injective. Let $\alpha : H^i(\text{GL}_m(\mathbb{F}_q)) \rightarrow H^i(\text{GL}_{n-1}(\mathbb{F}_q))$ denote the restriction map.

We have $H^*(D_k)^{S_k} = \mathbb{Z}/2\mathbb{Z}[v_1, \dots, v_k]^{S_k}$. Consider the map

$$\beta_n^* : \mathbb{Z}/2\mathbb{Z}[v_1, \dots, v_m]^{S_m} \rightarrow \mathbb{Z}/2\mathbb{Z}[v'_1, \dots, v'_{n-1}]^{S_{n-1}},$$

given by

$$\beta_n^*(v_i) = \begin{cases} v'_i, & \text{if } i < n, \\ 0, & \text{if } i \geq n. \end{cases}$$

Note that β_n^* induces a map

$$\beta_{ni}^* : H^i(D_m)^{S_m} \rightarrow H^i(D_{n-1})^{S_{n-1}}.$$

Consider the following commutative diagram.

$$\begin{array}{ccc} H^i(GL_m(\mathbb{F}_q)) & \xrightarrow{\alpha} & H^i(GL_{n-1}(\mathbb{F}_q)) \\ \downarrow \text{res}_m & & \downarrow \text{res}_{n-1} \\ H^i(D_m)^{S_m} & \xrightarrow{\beta_{ni}^*} & H^i(D_{n-1})^{S_{n-1}} \end{array}$$

Since the maps res_m and res_{n-1} are injective, it is enough to prove that the map β_{ni}^* is injective. Let f_i^k denote the elementary symmetric polynomial of degree i in k variables v_1, \dots, v_k , where $k \geq i$. Note that $\beta_{ni}^*(f_i^k) \neq 0$ for $1 \leq i \leq n-1 \leq k$. We know that the set of elementary symmetric functions form a basis for the algebra $\mathbb{Z}/2\mathbb{Z}[v_1, \dots, v_m]^{S_m}$. An element in $\mathbb{Z}/2\mathbb{Z}[v_1, \dots, v_m]^{S_m}$ of degree $i \leq n-1$ is of the form $P(f_1^m, \dots, f_i^m)$, where P is a polynomial (see [Lan02, Theorem 6.1, page 191]). We have $\beta_{ni}^*(P(f_1^m, \dots, f_i^m)) = P(f_1^{n-1}, \dots, f_i^{n-1})$. Therefore $\beta_{ni}^*(P(f_1^m, \dots, f_i^m)) \neq 0$ as the elements $f_1^{n-1}, \dots, f_i^{n-1}$ are algebraically independent. So β_{ni}^* is injective for $i \leq n-1$. \square

6. CALCULATIONS OF SOME SWC

6.1. Calculations Using the Main Theorem. Here we calculate small SWCs using the main result.

6.1.1. *For $\mathbf{q} \equiv \mathbf{1} \pmod{4}$.* From the main theorem we have for $q \equiv 1 \pmod{4}$,

$$w(\pi) = (1 + \delta b_1) \prod_{i=1}^n \left(\prod_{p \in \mathfrak{X}_i} (1 + p \cdot t) \right)^{c_i/2},$$

Here we calculate $w_2(\pi)$. It is clear from the above expression that

$$\begin{aligned} w_2(\pi) &= \sum_{j=1}^n (c_j/2) \left(\sum_{1 \leq i_1 < i_2 < \dots < i_j \leq n} (t_{i_1} + t_{i_2} + \dots + t_{i_j}) \right) \\ &= \frac{1}{2} \sum_{j=1}^n c_j \binom{n-1}{j-1} \left(\sum_{i=1}^n t_i \right) \\ &= \frac{1}{2} \sum_{j=1}^n c_j \binom{n-1}{j-1} a_2. \end{aligned}$$

Let $m = {}^t[c_0, c_1, \dots, c_n]$ and $J = [0, \binom{n-1}{0}, \binom{n-1}{1}, \dots, \binom{n-1}{n-1}]$. So we have

$$Jm = \sum_{j=1}^n \binom{n-1}{j-1} c_j. \quad (16)$$

Using Equation (9) and Theorem 6 we have $m = (1/2^n)Ma$. As in the proof of Theorem 1, let $R_i = (m_{i0}, m_{i1}, \dots, m_{in})$ be the i -th row of M which satisfies $\sum_{j=0}^n m_{ij} y^j = (1-y)^i (1+y)^{n-i}$. We calculate JM .

Proposition 5. *We have $Jm = \frac{\chi_\pi(h_0) - \chi_\pi(h_1)}{2}$.*

Proof. We identify a $1 \times (n+1)$ vector (m_{i0}, \dots, m_{in}) with the polynomial $\sum_{j=0}^n m_{ij} y^j = (1-y)^i (1+y)^{n-i}$. We compute

$$\begin{aligned} JM &= \sum_{i=1}^n \binom{n-1}{i-1} (1-y)^i (1+y)^{n-i} \quad \text{Put } i-1 = \alpha \\ &= \sum_{\alpha=0}^{n-1} \binom{n-1}{\alpha} (1-y)^{\alpha+1} (1+y)^{n-1-\alpha} \\ &= (1-y) \sum_{\alpha=0}^{n-1} \binom{n-1}{\alpha} (1-y)^\alpha (1+y)^{n-1-\alpha} \\ &= (1-y)(1-y+1+y)^{n-1} \\ &= 2^{n-1}(1-y) \\ &= [2^{n-1}, -2^{n-1}, 0, \dots, 0]. \end{aligned}$$

Thus we have

$$\begin{aligned} Jm &= (1/2^n) JM a \\ &= (1/2^n) [2^{n-1}, -2^{n-1}, 0, \dots, 0]^t [\chi_\pi(h_0), \chi_\pi(h_1), \dots, \chi_\pi(h_n)] \\ &= (1/2)(\chi_\pi(h_0) - \chi_\pi(h_1)). \end{aligned}$$

□

So for $q \equiv 1 \pmod{4}$ we have

$$w_2(\pi) = \frac{\chi_\pi(h_0) - \chi_\pi(h_1)}{4} a_2.$$

6.1.2. For $\mathbf{q} \equiv \mathbf{3} \pmod{4}$.

Theorem 9. *Let π be a real representation of $\text{GL}_n(\mathbb{F}_q)$, where $q \equiv 3 \pmod{4}$. Then*

$$w_1(\pi) = \frac{\chi_\pi(h_0) - \chi_\pi(h_1)}{2} \left(\sum_{i=1}^n v_i \right).$$

Proof. We have

$$w(\pi) = \prod_{i=1}^n \left(\prod_{b \in \mathfrak{X}_i} (1 + v \cdot b) \right)^{c_i}.$$

We write $v \cdot b$ for $\sum_{i=1}^n v_i b_i$. In particular,

$$w_1(\pi) = \sum_{i=1}^n c_i \binom{n-1}{i-1} \left(\sum_{i=1}^n v_i \right) = Jm \left(\sum_{i=1}^n v_i \right).$$

Now the proof follows from Proposition 5. □

Now we calculate $w_2(\pi)$. For $b \in \mathfrak{X}_i$ let $E_b = (1 + v \cdot b)^{c_i}$ and $P_i = \prod_{b \in \mathfrak{X}_i} E_b$. Note that

$$(1 + v \cdot b)^{c_i} = \left(1 + c_i \left(\sum_{s=1}^i v_{j_s} \right) + \binom{c_i}{2} \left(\sum_{s=1}^i v_{j_s}^2 \right) + 2 \binom{c_i}{2} \left(\sum_{k < l} v_{j_k} v_{j_l} \right) + T_3 \right),$$

where $\{j_1, \dots, j_i\}$ are such that $b_{j_s} = 1$ and T_3 denotes the terms with degree greater than or equal to 3. Let N_i denote the number of elements in \mathfrak{X}_i with 1 in a fixed position. We have $N_i = \binom{n-1}{i-1}$. Put $N'_i = \binom{n-2}{i-2}$ which is the number of

elements in \mathfrak{X}_i with 1 in two fixed positions. Also consider $N_i'' = \binom{n-2}{i-1}$ to be the number of elements in \mathfrak{X}_i with 1 in one fixed position and 0 in other fixed position.

The coefficient of $\sum_{t=1}^n v_t^2$ in the symmetric polynomial P_i is

$$N_i \binom{c_i}{2} + c_i^2 \binom{N_i}{2}.$$

The coefficient of $\sum_{1 \leq l < k \leq n} v_k v_l$ in P_i is

$$2 \binom{n-2}{i-2} \binom{c_i}{2} + c_i^2 N_i' (N_i' - 1) + c_i^2 \cdot 2N_i' N_i'' + c_i^2 N_i''^2 \quad (17)$$

We ignore the term $2 \binom{n-2}{i-2} \binom{c_i}{2}$ since it is even.

We calculate

$$\begin{aligned} c_i^2 N_i' (N_i' - 1) + c_i^2 \cdot 2N_i' N_i'' + c_i^2 N_i''^2 &= c_i^2 (N_i'^2 + 2N_i' N_i'' + N_i''^2 - N_i') \\ &= c_i^2 ((N_i' + N_i'')^2 - N_i') \\ &= c_i^2 (N_i^2 - N_i'), \quad (\text{using identity } N_i' + N_i'' = N_i) \\ &= c_i (N_i - N_i') \pmod{2} \\ &= c_i N_i'' \pmod{2}. \end{aligned}$$

Since $w(\pi) = \prod_i P_i$, we have

$$\begin{aligned} w_2(\pi) &= \left(\sum_{1 \leq i < j \leq n} N_i N_j c_i c_j + \sum_{i=1}^n N_i \binom{c_i}{2} + \sum_{i=1}^n c_i^2 \binom{N_i}{2} \right) \left(\sum_{t=1}^n v_t^2 \right) \\ &\quad + \left(\sum_{i=1}^n c_i N_i'' + 2 \sum_{1 \leq i < j \leq n} N_i N_j c_i c_j \right) \cdot \left(\sum_{k < l} v_k v_l \right). \end{aligned} \quad (18)$$

The Vandermonde identity says

$$\binom{r_1 + \cdots + r_t}{s} = \sum_{k_1 + \cdots + k_t = s} \binom{r_1}{k_1} \cdots \binom{r_t}{k_t}.$$

Using this identity we calculate

$$\begin{aligned} \binom{\sum_{i=1}^n c_i N_i}{2} &= \sum_{i=1}^n \binom{c_i N_i}{2} + \sum_{1 \leq l < k \leq n} c_l c_k N_l N_k \\ &= \sum_{i=1}^n \left(\sum_{k_1 + \cdots + k_{N_i} = 2} \binom{c_i}{k_1} \cdots \binom{c_i}{k_{N_i}} \right) + \sum_{1 \leq l < k \leq n} c_l c_k N_l N_k \\ &= \sum_{i=1}^n N_i \binom{c_i}{2} + \sum_{i=1}^n c_i^2 \binom{N_i}{2} + \sum_{1 \leq l < k \leq n} c_l c_k N_l N_k. \end{aligned}$$

Thus we have

$$w_2(\pi) = \binom{\sum_{i=1}^n c_i N_i}{2} \left(\sum_{l=1}^n v_l^2 \right) + \left(\sum_{i=1}^n c_i N_i'' \right) \left(\sum_{1 \leq l < k \leq n} v_l v_k \right). \quad (19)$$

Put $N = [N_0'', N_1'', N_2'', \dots, N_n'']$. So we have $Nm = \sum_{j=1}^n c_j N_j''$. We calculate NM .

Proposition 6. *We have $\sum_i c_i N_i'' = Nm = \frac{\chi_\pi(h_0) - \chi_\pi(h_2)}{4}$.*

Proof. The proof is similar to Proposition 5. \square

Theorem 10. *Let π be a real representation of $\mathrm{GL}_n(\mathbb{F}_q)$, where $q \equiv 3 \pmod{4}$. Then*

$$w_2(\pi) = \binom{m_\pi}{2} \sum_i v_i^2.$$

Proof. Observe that $\sum_{i=1}^n c_i N_i = \frac{\chi_\pi(h_0) - \chi_\pi(h_1)}{2} = m_\pi$ by equation (16) and Proposition 5. From Equation (19) and Proposition 6 we have

$$w_2(\pi) = \binom{m_\pi}{2} \sum_l v_l^2 + \frac{\chi_\pi(h_0) - \chi_\pi(h_2)}{4} \cdot \sum_{1 \leq l < k \leq n} v_l v_k.$$

Now from [GJ22][Lemma 4.4] it follows that the coefficient of $\sum_{1 \leq l < k \leq n} v_l v_k$ is even. Hence we get the result. \square

6.2. Calculations of SWC Using Detection Results. Let $i_k : \mathrm{GL}_k(\mathbb{F}_q) \hookrightarrow \mathrm{GL}_n(\mathbb{F}_q)$, where $k < n$, denote the injection of $\mathrm{GL}_k(\mathbb{F}_q)$ inside $\mathrm{GL}_n(\mathbb{F}_q)$ at the top left corner with rest diagonal entries 1. When $q \equiv 1 \pmod{4}$ we have $i_1^*(a_2) = t_1$ and when $q \equiv 3 \pmod{4}$ we have $i_2^*(\sum_{i=1}^n v_i^2) = v_1^2 + v_2^2$.

For a real representation π of $\mathrm{GL}_1(\mathbb{F}_q)$, Theorem 1 gives

$$w(\pi) = \begin{cases} (1 + \delta s_1)(1 + t_1)^{c_1/2}, & \text{if } q \equiv 1 \pmod{4}, \\ (1 + v_1)^{c_1}, & \text{if } q \equiv 3 \pmod{4}, \end{cases}$$

For a real representation π of $\mathrm{GL}_2(\mathbb{F}_q)$, Theorem 1 says

$$w(\pi) = \begin{cases} (1 + \delta(s_1 + s_2))((1 + t_1)(1 + t_2))^{c'_1/2} (1 + t_1 + t_2)^{c'_2/2}, & \text{if } q \equiv 1 \pmod{4}, \\ (1 + v_1)^{c'_1} (1 + v_2)^{c'_1} (1 + v_1 + v_2)^{c'_2}, & \text{if } q \equiv 3 \pmod{4}. \end{cases}$$

For the case of $\mathrm{GL}_1(\mathbb{F}_q)$ we have

$$\begin{pmatrix} c_0 \\ c_1 \end{pmatrix} = (1/2) \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \cdot \begin{pmatrix} \chi_\pi(h_0) \\ \chi_\pi(h_1) \end{pmatrix},$$

while for the case of $\mathrm{GL}_2(\mathbb{F}_q)$ we have

$$\begin{pmatrix} c'_0 \\ c'_1 \\ c'_2 \end{pmatrix} = (1/4) \begin{pmatrix} 1 & 2 & 1 \\ 1 & 0 & -1 \\ 1 & -2 & 1 \end{pmatrix} \cdot \begin{pmatrix} \chi_\pi(h_0) \\ \chi_\pi(h_1) \\ \chi_\pi(h_2) \end{pmatrix}.$$

For $q \equiv 3 \pmod{4}$ we compute $w_1(\pi)$ in terms of character values.

Theorem 11. *Let π be a real representation of $\mathrm{GL}_n(\mathbb{F}_q)$, where $q \equiv 3 \pmod{4}$. Then we have*

$$w_1(\pi) = m_\pi \left(\sum_{i=1}^n v_i \right).$$

Proof. Since $i = 1$ in Theorem 8, we have $n = 2$. We calculate for $\mathrm{GL}_1(\mathbb{F}_q)$. So for the representation π we have

$$w_1(\pi) = c_1 v_1.$$

Since $c_1 = m_\pi$, we have the result. \square

Next we compute the second SWC.

Theorem 12. *Let π be a real representation of $GL_n(\mathbb{F}_q)$, where q is odd. Then we have*

$$w_2(\pi) = \begin{cases} \frac{m_\pi}{2} a_2, & \text{if } q \equiv 1 \pmod{4} \\ \binom{m_\pi}{2} \left(\sum_{i=1}^n v_i^2 \right), & \text{if } q \equiv 3 \pmod{4}. \end{cases}$$

Proof. For a real representation π of $GL_1(\mathbb{F}_q)$, where $q \equiv 1 \pmod{4}$, we have

$$w_2(\pi) = \frac{c_1}{2} t_1.$$

We have $c_1 = (1/2)(\chi_\pi(h_0) - \chi_\pi(h_1)) = m_\pi$. Thus by Theorem 3 we have

$$w_2(\pi) = \frac{m_\pi}{2} a_2.$$

For a real representation π of $GL_2(\mathbb{F}_q)$ and $q \equiv 3 \pmod{4}$ we have

$$w_2(\pi) = \binom{c'_1 + c'_2}{2} (v_1^2 + v_2^2) + c_1'^2 v_1 v_2.$$

We know from [GJ22, Lemma 4.4]) that c'_1 is even. Moreover we have $c'_1 + c'_2 = (1/2)(\chi_\pi(h_0) - \chi_\pi(h_1)) = m_\pi$. Hence the Theorem follows from Theorem 8. \square

Using the detection result we compute the fourth SWC in terms of character values when $q \equiv 1 \pmod{4}$.

Theorem 13. *Let π be a real representation of $GL_n(\mathbb{F}_q)$, where $q \equiv 1 \pmod{4}$. Then*

$$w_4(\pi) = \binom{m_\pi/2}{2} \sum_{i=1}^n t_i^2 + \frac{\dim \pi - \chi_\pi(h_2)}{8} \sum_{1 \leq i < j \leq n} t_i t_j.$$

Proof. From Theorem 3 we know that the map $\text{res} : H^4(GL_n(\mathbb{F}_q)) \rightarrow H^4(GL_2(\mathbb{F}_q))$ is injective. We have

$$w^{\text{GL}_2(\mathbb{F}_q)}(\pi) = (1 + \delta(s_1 + s_2)) ((1 + t_1)(1 + t_2))^{c'_1/2} (1 + t_1 + t_2)^{c'_2/2}.$$

Then we calculate

$$\begin{aligned} \text{res}(w_4(\pi)) &= \left(\binom{c'_1/2}{2} + \binom{c'_2/2}{2} + \frac{c'_1 c'_2}{4} \right) (t_1^2 + t_2^2) + \frac{c'_1}{2} t_1 t_2 \\ &= \left(\frac{c'_1}{2} + \frac{c'_2}{2} \right) \binom{c'_1 + c'_2}{2} (t_1^2 + t_2^2) + \frac{c'_1}{2} t_1 t_2 \\ &= \binom{m_\pi/2}{2} (t_1^2 + t_2^2) + \frac{\dim \pi - \chi_\pi(h_2)}{8} t_1 t_2. \end{aligned}$$

So we have

$$w_4(\pi) = \binom{m_\pi/2}{2} \sum_{i=1}^n t_i^2 + \frac{\dim \pi - \chi_\pi(h_2)}{8} \sum_{1 \leq i < j \leq n} t_i t_j.$$

\square

7. SOME APPLICATIONS

7.1. Lifting Criteria for Real Representations of $\mathrm{GL}_n(\mathbb{F}_q)$. A real representation π of a finite group G is called spinorial provided it lifts to the Pin group, which is a double cover of the orthogonal group. One knows that π is spinorial iff $w_2(\pi) + w_1^2(\pi) = 0$ (see [GKT89, page 328]).

The main result in [JS21, Theorem 1] provides a criterion for spinoriality of a real representation of $\mathrm{GL}_n(\mathbb{F}_q)$ in terms of character values except for some particular pairs (n, q) . We recover the result here for q odd.

Theorem 14. *Let π be a real representation of $\mathrm{GL}_n(\mathbb{F}_q)$, where q is odd. Then*

- (1) *for $q \equiv 1 \pmod{4}$, we have π is spinorial iff $m_\pi \equiv 0 \pmod{4}$,*
- (2) *for $q \equiv 3 \pmod{4}$, we have π is spinorial iff $m_\pi \equiv 0$ or $3 \pmod{4}$.*

Proof. We know that π is spinorial iff $w_2(\pi) + w_1^2(\pi) = 0$. From Theorems 3 and 8 we deduce that π is spinorial iff $\mathrm{res}(w_2(\pi) + w_1^2(\pi)) = 0$, where $\mathrm{res} : H^*(\mathrm{GL}_n(\mathbb{F}_q), \mathbb{Z}/2\mathbb{Z}) \rightarrow H^*(\mathrm{GL}_k(\mathbb{F}_q), \mathbb{Z}/2\mathbb{Z})$ denotes the restriction map, where $k = 1$ for $q \equiv 1 \pmod{4}$ and $k = 2$ for $q \equiv 3 \pmod{4}$.

From Theorems 11 and 12, it follows that for a real representation π of $\mathrm{GL}_n(\mathbb{F}_q)$, we have

$$\mathrm{res}(w_2(\pi) + w_1^2(\pi)) = \begin{cases} \frac{m_\pi}{2} t_1, & \text{if } q \equiv 1 \pmod{4}, \\ \frac{m_\pi(m_\pi + 1)}{2} (v_1^2 + v_2^2), & \text{if } q \equiv 3 \pmod{4}. \end{cases}$$

□

7.2. Non-Detection by An-isotropic Torus. Let $a \in \mathbb{F}_{q^n}^\times$ and $G = \mathrm{GL}_n(\mathbb{F}_q)$, where q is odd. We fix a basis of \mathbb{F}_{q^n} over \mathbb{F}_q . Then the map $x \rightarrow a \cdot x$ gives a linear map $M_a \in G$. Then $M = \{M_a, a \in \mathbb{F}_{q^n}^\times\}$ is a cyclic group of order $q^n - 1$ and is called an anisotropic torus.

Theorem 15. *The anisotropic torus M does not detect the mod-2 cohomology of G .*

Proof. Consider the representation $\pi = (\chi)_\mathbb{R}$, where $\chi = \chi^j \circ \det$ and j is odd. One has

$$\begin{aligned} \chi_\pi(h_1) &= 2\chi_j(\det(h_1)) \\ &= 2\chi_j(-1) \\ &= 2(-1)^j. \end{aligned}$$

Since j is odd we have $\chi_\pi(h_1) = -2$. This gives

$$m_\pi = \frac{1}{2}(2 + 2) = 2.$$

Then from Theorem 12 one obtains $w_2(\pi) \neq 0$.

Let \hat{g} be a generator of M . Now we have

$$w_2(\pi|_M) = \kappa(c_1(\chi^j \circ \det|_M)).$$

We have

$$\chi(\hat{g}) = \chi^j \circ \det(\hat{g}) = \zeta_{q-1}^j = \zeta_{q^2-1}^{(q+1)j}.$$

Take $u = c_1(\dot{\chi}) \in H^2(M, \mathbb{Z})$ where $\dot{\chi} \in \mathrm{Hom}(M, \mathbb{C}^\times)$ such that $\dot{\chi}(\hat{g}) = \zeta_{q^2-1}$. As a result we get

$$\begin{aligned} \kappa(c_1(\chi^j \circ \det|_M)) &= \kappa((q+1)ju) \\ &= (q+1)j\kappa(u) \\ &= 0. \end{aligned}$$

This shows that $w_2(\pi|_M) = \text{res}_M^G(w_2(\pi)) = 0$ whereas $w_2(\pi) \neq 0$. \square

8. THE CASE OF $GL_n(\mathbb{C})$ AND $GL_n(\mathbb{R})$

For a Lie group G we write $H^i(G)$ to denote the (singular) cohomology $H^i(BG)$ of a classifying space BG of G . Consider the subgroup

$$\Gamma = \text{diag}(\pm 1, \pm 1, \dots, \pm 1) \cong C_2^n$$

of $GL_n(k)$, where $k = \mathbb{R}$ or \mathbb{C} . Take $T = (S^1)^n$ to be the diagonal torus of the unitary group U_n . Now we establish some useful detection results.

Proposition 7. *The mod 2 cohomology of $GL_n(\mathbb{C})$ is detected by the group Γ .*

Proof. From [T⁺87, Theorem 2.1] we obtain that the maps $i_1^* : H^*(GL_n(\mathbb{C})) \rightarrow H^*(U_n)$ and $i_2^* : H^*(U_n) \rightarrow H^*(T)$ are injective. As a result we conclude that the mod 2 cohomology of $GL_n(\mathbb{C})$ is detected by the diagonal torus T . First we prove that the mod 2 cohomology of S^1 is detected by C_2 . Note that the circle group S^1 is isomorphic to $SO_2(\mathbb{R})$. We know that $H^*(SO_2(\mathbb{R})) = \mathbb{Z}/2\mathbb{Z}[w_2]$, where w_2 denotes the second Stiefel Whitney class of standard representation of $SO_2(\mathbb{R})$. The standard representation of $SO_2(\mathbb{R})$ can be thought of as the representation $\pi = \rho \oplus \rho^\vee$ of S^1 where $\rho(z) = z$ for $z \in S^1$. So $w_2 = w_2(\pi)$. One can consider $C_2 = \{\pm 1\}$ as a subgroup of S^1 to obtain a map $\text{res} : H^*(SO_2(\mathbb{R})) \rightarrow H^*(C_2, \mathbb{Z}/2\mathbb{Z})$. We have $H^*(C_2, \mathbb{Z}/2\mathbb{Z}) = \mathbb{Z}/2\mathbb{Z}[v]$, where $\deg v = 1$. Note that $\pi|_{C_2} = \text{sgn} \oplus \text{sgn}$. Thus we have

$$\text{res } w_2(\pi) = w_2(\text{sgn} \oplus \text{sgn}) = w_1^2(\text{sgn}) \neq 0.$$

So $\text{res}(w_2) = v^2$. This implies that the map ‘res’ is injective. Now the result follows from Kunnetth formula. \square

Proposition 8. *The mod 2 cohomology of $GL_n(\mathbb{R})$ is detected by Γ .*

Proof. From [Mit01, Theorem 7.7] we know that $BGL_n(\mathbb{R})$ is homotopy equivalent to $BO_n(\mathbb{R})$. Thus the restriction map $\text{res}_O : H^*(GL_n(\mathbb{R})) \rightarrow H^*(O_n(\mathbb{R}))$ is an isomorphism. From [T⁺87, Theorem 2.2] we obtain that $\text{res}_\Gamma : H^*(O_n(\mathbb{R})) \rightarrow H^*(\Gamma)$ is an injection. Thus the $\mathbb{Z}/2\mathbb{Z}$ cohomology of $GL_n(\mathbb{R})$ is detected by the subgroup $\Gamma \cong C_2^n$. \square

Proof of Theorem 2. Given a real representation π of $GL_n(\mathbb{R})$ (or $GL_n(\mathbb{C})$), we calculate the total Stiefel Whitney class of $\psi = \pi|_\Gamma$. Moreover ψ is a S_n invariant representation as it is a restriction of representation of $GL_n(\mathbb{R})$ (or $GL_n(\mathbb{C})$). Now the Theorem 2 follows from Theorem 7. \square

Remark 3. Let π be a representation of $O_n(\mathbb{R})$. Then the proof of Proposition 8 gives that the total SWC of π is same as the expression in Equation (2).

9. EXAMPLES

We have

$$w_2(\pi) = \begin{cases} \frac{m_\pi}{2} a_2, & \text{if } q \equiv 1 \pmod{4}, \\ \binom{m_\pi}{2} (\sum v_i^2), & \text{if } q \equiv 3 \pmod{4}, \end{cases}$$

where $m_\pi = (1/2)(\dim \pi - \chi_\pi(h_1))$. Also

$$w_4(\pi) = \binom{m_\pi/2}{2} \sum_i t_i^2 + \frac{n_\pi}{4} \sum_{i < j} t_i t_j,$$

where $n_\pi = \frac{\dim \pi - \chi_\pi(h_2)}{2}$.

9.1. Principal Series Representation. Let $\chi_1, \chi_2, \dots, \chi_n$ are linear characters of \mathbb{F}_q^\times and let B be the Borel subgroup of $\mathrm{GL}_n(\mathbb{F}_q)$. Put

$$\pi = \mathrm{Ind}_B^G(\chi_1 \boxtimes \chi_2 \boxtimes \cdots \boxtimes \chi_n).$$

Such representations are called principal series representations of G . Such a representation π is orthogonal iff

$$\{\chi_1, \dots, \chi_n\} = \{\chi_1^{-1}, \dots, \chi_n^{-1}\},$$

as multisets. We introduce the notation

$$\begin{aligned} [n]_q! &= \prod_{i=1}^n \frac{q^i - 1}{q - 1} \\ &= (q + 1)(q^2 + q + 1) \cdots (q^{n-1} + q^{n-2} + \cdots + q + 1). \end{aligned}$$

From [Car85, Proposition 7.5.3, page no. 233] we obtain the following result.

$$\chi_\pi(h_k) = [k]_q! [n - k]_q! \sum_{1 \leq i_1 < i_2 < \cdots < i_k \leq n} \chi_{i_1}(-1) \chi_{i_2}(-1) \cdots \chi_{i_k}(-1). \quad (20)$$

Put

$$T_i = (1 + q + q^2 + \cdots + q^i).$$

So we calculate

$$\begin{aligned} m_\pi &= \frac{1}{2}(\dim \pi - \chi_\pi(h_1)) \\ &= \frac{1}{2} \left([n]_q! - [n - 1]_q! \left(\sum_{i=1}^n \chi_i(-1) \right) \right) \\ &= \frac{1}{2} [n - 1]_q! \left(T_{n-1} - \sum_{i=1}^n \chi_i(-1) \right). \end{aligned}$$

Similarly one has

$$\begin{aligned} n_\pi &= \frac{1}{2}(\dim \pi - \chi_\pi(h_2)) \\ &= \frac{1}{2} [n - 2]_q! \left(T_{n-2} T_{n-1} - (1 + q) \sum_{i < j} \chi_i(-1) \chi_j(-1) \right). \end{aligned}$$

Theorem 16. *Let π be a real principal series representation for $\mathrm{GL}_n(\mathbb{F}_q)$ where q is odd. Then $w_2(\pi) = 0$ for $n \geq 3$.*

Proof. From [JS21, Theorem 5] we know that if $n \geq 3$, then all the real principal series representations of $\mathrm{GL}_n(\mathbb{F}_q)$ are spinorial. Now π is spinorial iff $w_2(\pi) = w_1^2(\pi)$ (see [GS20, Proposition 6.1]). If $q \equiv 1 \pmod{4}$, then $w_1^2(\pi) = 0$. Therefore $w_2(\pi) = 0$.

Now consider the case for $q \equiv 3 \pmod{4}$. We have

$$m_\pi = \frac{1}{2} [n - 1]_q! \left(T_{n-1} - \sum_{i=1}^n \chi_i(-1) \right).$$

Also

$$T_{n-1} \equiv \sum_{i=1}^n \chi_i(-1) \equiv n \pmod{2}.$$

This gives $T_{n-1} - \sum_{i=1}^n \chi_i(-1) \equiv 0 \pmod{2}$. We have $[m]_q! \equiv 0 \pmod{4}$ for $m \geq 2$.

For $n \geq 3$, we have $v_2(m_\pi) \geq 2$. Therefore $\binom{m_\pi}{2} \equiv 0 \pmod{2}$. Therefore $w_2(\pi) = 0$ for $n \geq 3$. \square

Proof of Theorem 4. We have

$$n_\pi = \frac{1}{2}[n-2]_q! \left(T_{n-2}T_{n-1} - (1+q) \sum_{i<j} \chi_i(-1)\chi_j(-1) \right).$$

Observe that $T_{n-2}T_{n-1} - (1+q) \sum_{i<j} \chi_i(-1)\chi_j(-1)$ is always even. Also $v_2([n-2]_q!) \geq 3$ for $n \geq 6$. Therefore $n_\pi/4$ remains even for $n \geq 6$.

We have

$$m_\pi = \frac{1}{2}[n-1]_q! \left(T_{n-1} - \sum_{i=1}^n \chi_i(-1) \right).$$

Also

$$T_{n-1} \equiv \sum_{i=1}^n \chi_i(-1) \equiv n \pmod{2}.$$

This gives $T_{n-1} - \sum_{i=1}^n \chi_i(-1) \equiv 0 \pmod{2}$. For $n \geq 6$, the term $m_\pi/2$ remains even. So we have $\binom{m_\pi/2}{2} \equiv \frac{m_\pi}{4} \pmod{2}$. Now $v_2(m_\pi) \geq 4$ for $n \geq 6$. Therefore $w_4(\pi) = 0$ for $n \geq 6$.

For $n = 5$ the two possible real Principal series representations are $\pi_1 = \{\mathbb{1}, \chi_i, \chi_i^{-1}, \chi_j, \chi_j^{-1}\}$ and $\pi_2 = \{\text{sgn}, \chi_i, \chi_i^{-1}, \chi_j, \chi_j^{-1}\}$.

We calculate

$$\begin{aligned} m_{\pi_1} &= \frac{1}{2} ([5]_q! - [4]_q!(1 + 2(-1)^i + 2(-1)^j)) \\ &= \frac{1}{2}(q+1)^2(q^2+1)(q^2+q+1)(q(q+1)(q^2+1) - 2(-1)^i - 2(-1)^j). \end{aligned}$$

This shows $v_2(m_{\pi_1}) \geq 4$. Similarly one obtains $v_2(m_{\pi_2}) \geq 4$.

We have

$$\begin{aligned} n_{\pi_1} &= \frac{1}{2} ([5]_q! - [2]_q![3]_q!(2 + 2(-1)^i + 2(-1)^j + 4(-1)^{i+j})) \\ &= \frac{1}{2}(q+1)^2(q^2+q+1)((q^2+1)(q^4+q^3+q^2+q+1) - (2 + 2(-1)^i + 2(-1)^j + 4(-1)^{i+j})) \\ &= \frac{1}{2}(q+1)^2(q^2+q+1)(\{q^6+q^5+2(q^4+q^3+q^2)\} + \{q-1\} - \{2(-1)^i + 2(-1)^j + 4(-1)^{i+j}\}). \end{aligned}$$

Note that the all the terms in the curly brackets and $(q+1)^2$ are divisible by 4. Therefore $v_2(n_{\pi_1}) \geq 3$. Similar calculation shows that $v_2(n_{\pi_2}) \geq 3$. Now the result follows from Theorem 13. \square

Here we list down the value of $w_4(\pi)$, where π is an orthogonal principal series representation of $GL_n(\mathbb{F}_q)$ for $q \equiv 1 \pmod{4}$.

(1) Take $n = 3$ and $\pi = \{\mathbb{1}, \chi_j, \chi_j^{-1}\}$ or $\{\text{sgn}, \chi_j, \chi_j^{-1}\}$ Then

$$w_4(\pi) = \sum_{i=1}^3 t_i^2 + \sum_{1 \leq i < j \leq 3} t_i t_j$$

if $q \equiv 1, 9 \pmod{16}$ and j is odd or $q \equiv 5, 13 \pmod{16}$ and j is even.
Otherwise $w_4(\pi) = 0$.

(2) Take $n = 4$ and $\pi = \{\mathbb{1}, \text{sgn}, \chi_i, \chi_i^{-1}\}$. Then

$$w_4(\pi) = \begin{cases} \sum_{i=1}^4 t_i^2, & \text{if } i \text{ is odd,} \\ 0, & \text{if } i \text{ is even.} \end{cases}$$

(3) Take $n = 4$ and $\pi = \{\chi_j, \chi_j^{-1}, \chi_k, \chi_k^{-1}\}$.

$$w_4(\pi) = \begin{cases} \sum_{i=1}^4 t_i^2, & \text{if } j \text{ and } k \text{ have different parity,} \\ 0, & \text{otherwise.} \end{cases}$$

9.2. Cuspidal Representations. Let θ be a regular character of full anisotropic torus T of $\text{GL}_n(\mathbb{F}_q)$. Then there is a cuspidal representation π_θ associated to it. The representation π_θ is real iff there exists a Weyl group element w such that $w(\theta) = \theta^{-1}$.

Let π be a real Cuspidal representation of $\text{GL}_n(\mathbb{F}_q)$.

9.2.1. Computation of $w_2(\pi)$. We know that $\chi_\pi(h_1) = 0$ and $\dim \pi = \psi_{n-1}(q)$ where

$$\psi_m(q) = \prod_{i=1}^m (q^i - 1).$$

Thus $m_\pi = \psi_{n-1}(q)/2$.

Theorem 17. *Let π be a real Cuspidal representation of $\text{GL}_n(\mathbb{F}_q)$ such that $n \geq 3$. Then $w_2(\pi) = 0$.*

Proof. For $n \geq 3$, the term $\frac{(q-1)(q^2-1)}{2}$ divides m_π , Thus $m_\pi \equiv 0 \pmod{4}$, and therefore $w_2(\pi) = 0$. \square

For $n = 2$ and $q \equiv 1 \pmod{4}$, we have $m_\pi = (q-1)/2$ and $w_2 = \frac{q-1}{4}a_2$.

$$w_2(\pi) = \begin{cases} 0, & \text{if } q \equiv 1 \pmod{8}, \\ a_2, & \text{if } q \equiv 5 \pmod{8}. \end{cases}$$

For $n = 2$ and $q \equiv 3 \pmod{4}$, we have $m_\pi = (q-1)/2$ and $w_2 = \binom{m_\pi}{2} \cdot (v_1^2 + v_2^2)$. Therefore

$$w_2(\pi) = \begin{cases} 0, & \text{if } q \equiv 3 \pmod{8}, \\ v_1^2 + v_2^2, & \text{if } q \equiv 7 \pmod{8}. \end{cases}$$

9.2.2. Computation of $w_4(\pi)$. We summarize the computations in the following table.

9.3. Steinberg Representation. Let π be the Steinberg representation of $\text{GL}_n(\mathbb{F}_q)$.

TABLE 1. Computation of $w_4(\pi)$ where π is the Cuspidal representation of $GL_n(\mathbb{F}_q)$

q mod 16	$\theta(-1)$	$\binom{m_\pi/2}{2}$ (mod 2)	$n_\pi/4$ (mod 2)
1	1	0	0
1	-1	0	0
5	1	0	0
5	-1	0	1
9	1	1	0
9	-1	1	0
13	1	1	0
13	-1	1	1

9.3.1. **Computation of $w_2(\pi)$.** We have $\chi_\pi(h_1) = p^k$ where p^k is the highest power of p dividing $|Z_G(h_1)|$ (see [Car85, Theorem 6.4.7, page 195]). We have $Z_G(h_1) \cong GL_1(\mathbb{F}_q) \times GL_{n-1}(\mathbb{F}_q)$. Thus $\chi_\pi(h_1) = q^{(1/2)(n-1)(n-2)}$. Therefore

$$\begin{aligned} m_\pi &= \frac{q^{(1/2)(n(n-1))} - q^{(1/2)(n-1)(n-2)}}{2} \\ &= (1/2) \cdot q^{(1/2)(n-1)(n-2)}(q^{n-1} - 1) \\ &= \frac{q-1}{2} \cdot q^{(1/2)(n-1)(n-2)} \cdot (1 + q + q^2 + \cdots + q^{n-2}). \end{aligned}$$

For $q \equiv 1 \pmod{4}$ we have

$$w_2(\pi) = \begin{cases} a_2, & \text{if } q \equiv 5 \pmod{8} \text{ and for } n \text{ even,} \\ 0, & \text{otherwise.} \end{cases}$$

For $q \equiv 3 \pmod{4}$ we have

$$w_2(\pi) = \begin{cases} \sum_{i=1}^n v_i^2, & \text{if } q \equiv 3 \pmod{8} \text{ and for } n \equiv 0 \pmod{4}, \\ \sum_{i=1}^n v_i^2, & \text{if } q \equiv 7 \pmod{8} \text{ and for } n \equiv 2 \pmod{4}, \\ 0, & \text{otherwise.} \end{cases}$$

9.3.2. **Computation of $w_4(\pi)$.** We summarize the computations in the following table.

TABLE 2. Computation of $w_4(\pi)$ where π is the Steinberg representation of $\mathrm{GL}_n(\mathbb{F}_q)$

$q \pmod 8$	$n \pmod 4$	$\binom{m_\pi/2}{2} \pmod 2$	$n_\pi/4 \pmod 2$
1	0, 1, 2, 3	0	0
5	0	1	0
5	1	0	1
5	2	0	0
5	3	1	1
9	1, 3	0	0
9	0, 2	1	0
13	0	0	0
13	1	0	1
13	2	1	0
13	3	1	1

10. DATA AVAILABILITY STATEMENT

Our manuscript has no associated data.

11. CONFLICT OF INTEREST STATEMENT

All authors declare that they have no conflicts of interest.

REFERENCES

- [AM13] Alejandro Adem and R James Milgram. *Cohomology of finite groups*, volume 309. Springer Science & Business Media, 2013.
- [Ben91] David J Benson. *Representations and cohomology*, volume 2. Cambridge university press, 1991.
- [Car85] Roger William Carter. Finite groups of lie type: Conjugacy classes and complex characters. *Pure Appl. Math.*, 44, 1985.
- [DI08] Persi Diaconis and I Isaacs. Supercharacters and superclasses for algebra groups. *Transactions of the American Mathematical Society*, 360(5):2359–2392, 2008.
- [FP06] Zbigniew Fiedorowicz and Stewart Priddy. *Homology of classical groups over finite fields and their associated infinite loop spaces*, volume 674. Springer, 2006.
- [GJ21] Jyotirmoy Ganguly and Rohit Joshi. Spinorial representations of orthogonal groups. *J. Lie Theory*, 31(1):265–286, 2021.
- [GJ22] Jyotirmoy Ganguly and Rohit Joshi. Stiefel whitney classes for real representations of $\mathrm{gl}_2(\mathbb{f}_q)$. *International Journal of Mathematics*, page 2250010, 2022.
- [GKT89] J Gunawardena, B Kahn, and C Thomas. Stiefel-whitney classes of real representations of finite groups. *Journal of Algebra*, 126(2):327–347, 1989.
- [GS20] Jyotirmoy Ganguly and Steven Spallone. Spinorial representations of symmetric groups. *Journal of Algebra*, 544:29–46, 2020.

- [JS21] Rohit Joshi and Steven Spallone. Spinoriality of orthogonal representations of $\mathfrak{gl}(n, \mathbb{q})$. *Pacific Journal of Mathematics*, 311(2):369–383, 2021.
- [Lan02] Serge Lang. *Graduate Texts in Mathematics: Algebra*. Springer, 2002.
- [Mit01] Stephen A Mitchell. Notes on principal bundles and classifying spaces. *Lecture Notes. University of Washington*, 2001.
- [Nak60] Minoru Nakaoka. Decomposition theorem for homology groups of symmetric groups. *Annals of Mathematics*, pages 16–42, 1960.
- [T+87] Hiroshi Toda et al. Cohomology of classifying spaces. In *Homotopy theory and related topics*, pages 75–108. Mathematical Society of Japan, 1987.

CHENNAI MATHEMATICAL INSTITUTE, H1, SIPCOT IT PARK, SIRUSERI, KELAMBAKKAM 603103,
TAMIL NADU, INDIA

Email address: `jjyotirmoy.math@gmail.com`

IISER PUNE, DR. HOMI BHABHA ROAD, PUNE-411008, MAHARASHTRA, INDIA

Email address: `rohitsj@students.iiserpune.ac.in`