

MULTIPLICATIVE MAPS ON MATRIX ALGEBRAS

XIAOMEI YANG AND FUHAI ZHU

Abstract: In this paper, we use elementary method to give a classification of the multiplicative maps on matrix algebra $M_n(\mathbb{F})$ over a field \mathbb{F} of characteristic 0. All the multiplicative maps are classified into three classes: the trivial ones, the degenerate ones and the non-degenerate ones.

Keywords: Multiplicative map, ring homomorphism.

Mathematics Subject Classification: 15A33, 15A60

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULT

During the past one hundred years, a great deal of mathematical effort has been devoted to the study of preserver problems in matrix theory. In [4], [8] and [11], the authors deal with linear preserver problems, i.e., linear maps on matrix algebras and operator algebras preserving determinant, spectrum, commutativity, respectively. Chi-Kwong Li and Nam-Kiu Tsing presented a brief description of the subject in [9].

Naturally, some similar preserver problems are raised and investigated for multiplicative maps (i.e., maps Φ satisfying $\Phi(AB) = \Phi(A)\Phi(B)$) instead of linear maps. In [5], Hochwald described the form of those multiplicative maps of a matrix algebra which preserve the spectrum. As a natural generalization, he also raised the question of spectrum-preserving multiplicative maps for infinite-dimensional vector spaces with added condition of surjectivity, see [12]. In [2], An and Hou considered rank preserving multiplicative maps on matrix algebra. In [3], the authors established some general techniques that are used to prove results for multiplicative maps on semigroups of square matrices and obtained some basic results to study multiplicative preserver problems. In recent years, there have been extensive study and application of multiplicative preserver problems on operator algebras over infinite-dimensional spaces. In [6], Hou gave a structure theorem of abstract multiplicative maps Φ from $\mathfrak{B}(X)$ into $\mathfrak{B}(Y)$ which sends some rank-1 operator to an operator of rank not greater than 1. Afterwards, An and Hou obtained some characterizations of rank-preserving multiplicative maps on $\mathfrak{B}(X)$ in [1]. In [10], Molnár described the form of those continuous multiplicative maps on $\mathfrak{B}(H)$ (H is a separable complex Hilbert space of dimension not less than 3) which preserve the rank or the corank.

It is very strange that people always try to deal with multiplicative maps with some additional conditions, say, preserving spectrum, ranks, etc., rather than to classify all multiplicative maps first. In [7], the authors investigated non-degenerate multiplicative maps

(see section 4 for the definition) of $n \times n$ matrices over a principal ideal domain R . In [13], the author considered non-degenerate multiplicative maps of $n \times n$ matrices over a division ring. The purpose of this paper is to deal with the multiplicative maps for matrix algebras over any field of characteristic zero with an elementary method, which may be applicable to positive characteristic case. All the multiplicative maps are separated into three classes: we devote Section 2 to study the trivial ones, which maps $\text{SL}(n, \mathbb{F})$ to the identity. We prove that such multiplicative map is reduced to a multiplicative map $\bar{\Phi} : \mathbb{F} \rightarrow M_k(\mathbb{F})$ with $\bar{\Phi}(1) = I_k$. In particular, we show that any of multiplicative map $\Phi : M_n(\mathbb{F}) \rightarrow M_k(\mathbb{F}) (k < n)$ is actually a trivial multiplicative map; in Section 3, we classify the degenerate ones, which maps every singular matrix to the zero matrix. We present such $\Phi(A)$ is the form of $\lambda(\det(\Phi_1(A)))\Phi_2(A)$, where $\lambda : \mathbb{F}^* \rightarrow \mathbb{F}^*$ is a multiplicative map and Φ_1, Φ_2 are compositions of the following multiplicative maps: (a). $\Phi_R(A) = R^{-1}AR$ for some $n \times n$ invertible matrix R ; (b). $\Phi^*(A) = (A^*)^t$, where A^* is the cofactor matrix of A and A^t is the transpose of A ; (c). $\Phi_\varphi(A) = (\varphi(a_{ij}))$ for any $A = (a_{ij}) \in \text{GL}(n)$, where $\varphi : \mathbb{F} \rightarrow \mathbb{F}$ is a ring homomorphism; the non-degenerate multiplicative maps are considered in Section 4, in which we conclude that $\Phi(A) = R^{-1}(\varphi(a_{ij}))R$ or $\Phi(A) = R^{-1}(\varphi(a_{ij})^*)^t R$, for some ring homomorphism $\varphi : \mathbb{F} \rightarrow \mathbb{F}$ and some invertible matrix R . With such classification in hand, it is easy to find all multiplicative maps which preserve spectrums, ranks, etc.

2. TRIVIAL MULTIPLICATIVE MAPS

Denote by $M_n(\mathbb{F})$ the matrix algebra over a field \mathbb{F} of characteristic zero. A map $\Phi : M_n(\mathbb{F}) \rightarrow M_k(\mathbb{F})$ is called a *multiplicative map* if $\Phi(AB) = \Phi(A)\Phi(B)$ for any $A, B \in M_n(\mathbb{F})$. In this section, we study a special case of multiplicative maps. First, we need the following easy property of multiplicative maps.

Lemma 2.1. *Let $\Phi : M_n(\mathbb{F}) \rightarrow M_k(\mathbb{F})$ be a multiplicative map. Then for any matrix $A \in M_n(\mathbb{F})$, up to a conjugation, we have*

$$\Phi(A) = \begin{pmatrix} \tilde{A} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & I_s \end{pmatrix},$$

where $s = \text{rank}(\Phi(0))$, $l = \text{rank}(\Phi(I_n)) - \text{rank}(\Phi(0))$ and $\tilde{A} \in M_l(\mathbb{F})$.

Proof. Since $\Phi(0), \Phi(I_n)$ are idempotent matrices. We may assume that $\Phi(0), \Phi(I_n)$ are diagonal matrices with the diagonal elements being 0 or 1. Using $\Phi(0)\Phi(I_n) = \Phi(I_n)\Phi(0) = \Phi(0)$, up to a conjugation, we have

$$\Phi(0) = \begin{pmatrix} 0 & 0 \\ 0 & I_s \end{pmatrix}, \quad \Phi(I_n) = \begin{pmatrix} I_l & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & I_s \end{pmatrix},$$

where $s = \text{rank}(\Phi(0))$, $l = \text{rank}(\Phi(I_n)) - \text{rank}(\Phi(0))$.

For any $A \in M_n(\mathbb{F})$, from

$$\Phi(0)\Phi(A) = \Phi(A)\Phi(0) = \Phi(0),$$

$$\Phi(I_n)\Phi(A) = \Phi(A)\Phi(I_n) = \Phi(A),$$

it follows that $\Phi(A) = \begin{pmatrix} \tilde{A} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & I_s \end{pmatrix}$, where \tilde{A} is a $l \times l$ matrix. □

From the proof of the above Lemma, one can define a multiplicative map $\overline{\Phi} : M_n(\mathbb{F}) \rightarrow M_l(\mathbb{F})$ by $\overline{\Phi}(A) = \tilde{A}$. It is easy to see that Φ is determined by $\overline{\Phi}$ and $\text{rank}(\Phi(0))$ uniquely. In the remainder of this paper, we only consider those maps such that $\Phi(0) = 0$ and $\Phi(I_n) = I_k$. Then it is trivial that $\Phi(A^{-1}) = \Phi(A)^{-1}$ and $\Phi^{-1}(I_k)$ is a normal subgroup of $\text{GL}(n, \mathbb{F})$. A classical result says that if N is a normal subgroup of $\text{GL}(n, \mathbb{F})$, then either N is contained in $Z(\text{GL}(n, \mathbb{F}))$, that is the center of $\text{GL}(n, \mathbb{F})$, or N contains $\text{SL}(n, \mathbb{F})$. It is natural to introduce the following definition.

Definition 2.2. A multiplicative map $\Phi : M_n(\mathbb{F}) \rightarrow M_k(\mathbb{F})$ is called trivial if $\text{SL}(n, \mathbb{F}) \subseteq \Phi^{-1}(I_k)$.

Trivial multiplicative maps have the following property.

Lemma 2.3. Let $\Phi : M_n(\mathbb{F}) \rightarrow M_k(\mathbb{F})$ be a trivial multiplicative map. Then $\Phi(A) = 0$ if $\text{rank}(A) < n$.

Proof. Since $\text{rank}(A) = r < n$, there exist $P, Q, S, T \in \text{SL}(n, \mathbb{F})$ such that $A = PIQ = SJT$, where $I = \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}$ and J is a Jordan canonical form with diagonal entries being 0 and $\text{rank}(J) = r$. Thus $\Phi(A) = \Phi(I) = \Phi(J)$. Since $I = I^m$ and $J^m = 0$ for $m > r$, it is easy to see that $\Phi(A) = \Phi(I) = \Phi(I^m) = \Phi(J^m) = 0$. □

Thanks to the above lemma, any trivial multiplicative map Φ is reduced to a multiplicative map $\overline{\Phi} : \mathbb{F} \rightarrow M_k(\mathbb{F})$ with $\overline{\Phi}(1) = I_k$. The following proposition shows that many multiplicative maps are trivial and it is crucial for rest of the paper.

Proposition 2.4. Let $\Phi : M_n(\mathbb{F}) \rightarrow M_k(\mathbb{F}) (k < n)$ be a multiplicative map. Then Φ is a trivial multiplicative map.

Proof. Let N (resp. \overline{N}) denote the set of $n \times n$ (resp. $k \times k$) upper triangular matrices with all the diagonal entries being 1. Consider the lower central sequence: $N^0 = N$, $N^1 = [N, N] = \{xyx^{-1}y^{-1} | x, y \in N\}$, \dots , $N^i = [N, N^{i-1}]$. It is easy to calculate that $N^n = \{I_n\}$ and $(\overline{N})^k = \{I_k\}$. Furthermore, since $(\Phi(N))^n = \Phi(N^n) = \Phi(I_n) = I_k$, it follows that, up to a conjugation, $\Phi(N) \subseteq \overline{N}$. Then we have $\Phi(N)^k \subseteq (\overline{N})^k = \{I_k\}$, which implies that $N^k \subseteq \Phi^{-1}(I_k)$. Therefore $\Phi^{-1}(I_k) \supseteq \text{SL}(n, \mathbb{F})$ since it is not contained in the center of $\text{GL}(n, \mathbb{F})$. □

Thus, any of multiplicative map $\Phi : M_n(\mathbb{F}) \rightarrow M_k(\mathbb{F}) (k < n)$ is the form of $\Phi(A) = \overline{\Phi}(\det(A))$ for some multiplicative map $\overline{\Phi} : \mathbb{F} \rightarrow M_k(\mathbb{F})$.

3. DEGENERATE MULTIPLICATIVE MAPS

In this section, we just need to consider non-trivial multiplicative maps and hence we have $\Phi(0) = 0$ and $\Phi(I_n) = I_n$. Following Jodeit and Lam we call a nontrivial multiplicative map $\Phi : M_n(\mathbb{F}) \rightarrow M_n(\mathbb{F})$ *degenerate*, if $\Phi(A) = 0$ for any degenerate A , and is called *non-degenerate* otherwise. It is trivial that degenerate multiplicative maps are in 1-1 correspondence with multiplicative maps $\Phi : \text{GL}(n, \mathbb{F}) \rightarrow \text{GL}(n, \mathbb{F})$.

Theorem 3.1. *Let $\Phi : \text{GL}(n, \mathbb{F}) \rightarrow \text{GL}(n, \mathbb{F})$ be a non-trivial multiplicative map. Then $\Phi(A) = \lambda(\det(\Phi_1(A)))\Phi_2(A)$, where $\lambda : \mathbb{F}^* \rightarrow \mathbb{F}^*$ is a multiplicative map and Φ_1, Φ_2 are compositions of the following multiplicative maps.*

- (a). $\Phi_R(A) = R^{-1}AR$ for some $n \times n$ invertible matrix R ;
- (b). $\Phi^*(A) = (A^*)^t$, where A^* is the cofactor matrix of A and A^t is the transpose of A ;
- (c). $\Phi_\varphi(A) = (\varphi(a_{ij}))$ for any $A = (a_{ij}) \in \text{GL}(n)$, where $\varphi : \mathbb{F} \rightarrow \mathbb{F}$ is a ring homomorphism.

Proof. The proof of the theorem consists of several steps. For $j = 1, \dots, n$, let $D_j(k) = \text{diag}(1, \dots, 1, k, 1, \dots, 1)$, where the j -th entry is k .

Claim (1): Up to a composition with Φ_R and a multiplication by $\det(A)$ if necessary, we have $\Phi(D_i(-1)) = D_i(-1)$.

From the fact that $\Phi(D_1(-1))^2 = \Phi(D_1(-1)^2) = \Phi(I_n) = I_n$, composing with some R when necessary, we have $\Phi(D_1(-1)) = \begin{pmatrix} -I_m & 0 \\ 0 & I_{n-m} \end{pmatrix}$. Since $D_1(-1)$ commutes with $A = \begin{pmatrix} 1 & 0 \\ 0 & A_1 \end{pmatrix}$, where $A_1 \in \text{GL}(n-1, \mathbb{F})$, it follows that $\Phi(A)$ must be a quasi diagonal matrix. We may therefore assume that

$$\Phi(A) = \begin{pmatrix} B_1 & 0 \\ 0 & B_2 \end{pmatrix}, \text{ where } B_1 \in \text{GL}(m, \mathbb{F}), B_2 \in \text{GL}(n-m, \mathbb{F}).$$

It shows that the maps $\Phi_1 : \text{GL}(n-1, \mathbb{F}) \rightarrow \text{GL}(m, \mathbb{F})$ defined by $\Phi_1(A_1) = B_1$ and $\Phi_2 : \text{GL}(n-1, \mathbb{F}) \rightarrow \text{GL}(n-m, \mathbb{F})$ defined by $\Phi_2(A_1) = B_2$ are both multiplicative maps. Then we conclude that $m = 1$ or $m = n-1$ by Proposition 2.4, otherwise both Φ_1 and Φ_2 are trivials, so is Φ . If $m = n-1$, we may assume that $m = 1$. Since $\Phi(D_i(-1))$ have the same eigenvalues as $\Phi(D_1(-1))$ and $\Phi(D_i(-1))$ commute with $\Phi(D_j(-1))$ for $i, j = 1, \dots, n$, we have $\Phi(D_i(-1)) = D_i(-1)$ up to a conjugation.

Claim (2): Up to a composition with Φ_T , we have $\Phi(S_{ij}) = S_{ij}$, where T is a diagonal matrix and S_{ij} is the identity matrix with rows i and j switched.

We consider the image of $S_{i,i+1}$ under Φ . Note that $S_{i,i+1}$ (resp. $\Phi(S_{i,i+1})$) commutes with $D_k(-1)$ (resp. $\Phi(D_k(-1))$), for $k \neq i, i+1$, and $D_i(-1)S_{i,i+1}D_{i+1}(-1) = S_{i,i+1}$. Then we

see that $\Phi(S_{i,i+1}) = \text{diag}(a_1, \dots, a_{i-1}, B_i, a_{i+2}, \dots, a_n)$, for some $a_i \in \mathbb{F}^*$ and $B_i = \begin{pmatrix} 0 & b_i \\ \frac{1}{b_i} & 0 \end{pmatrix}$ for some nonzero $b_i \in \mathbb{F}$. Furthermore, $S_{i,i+1}$ is similar to $D_1(-1)$, so is $\Phi(S_{i,i+1})$. Thus we have $a_i = 1$. Take $T = \text{diag}(1, b_1, b_1 b_2, \dots, b_1 b_2 \dots b_{n-1})$, one have $T\Phi(S_{i,i+1})T^{-1} = S_{i,i+1}$. Replacing Φ by $T\Phi T^{-1}$, one has $\Phi(S_{ij}) = S_{ij}$.

Claim (3): Up to a composition with some Φ_R and with Φ^* when necessary, we have $\Phi(A) = (\varphi(a_{ij}))$ for any $A = (a_{ij}) \in \text{SL}(n, \mathbb{F})$, where φ is a nontrivial ring homomorphism of \mathbb{F} .

For invertible A , it is easy to see that A is unipotent if and only if A is conjugate to A^n for any $n \in \mathbb{N}$. Thus $\Phi(A)$ is also unipotent.

For $j = 1, \dots, n$, let $P_{j,j+1}(k) = \text{diag}(1, \dots, 1, \begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}, 1, \dots, 1)$, where the $j, j+1$ entry is k . Let $A = P_{12}(k)$. Then A commutes with $D_i(-1)$, $i > 2$. Thus one can easily get $\Phi(A) = \text{diag}(B, 1, \dots, 1)$, where $B = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is a 2×2 unipotent matrix. By $D_1(-1)AD_1(-1) = A^{-1}$, one can get that $a = d = 1$ and $bc = 0$. Replacing $\Phi(A)$ by $\Phi((A^{-1})^t) = \Phi \circ (\det(A)^{-1}\Phi_2(A))$ when necessary, one can assume that $c = 0$. Note that the map $A \rightarrow (A^{-1})^t$ preserves $D_i(-1)$ and S_{ij} . Denote b by $\varphi(k)$. Then one can easily check that

$$\varphi(k+l) = \varphi(k) + \varphi(l).$$

Similarly, by $S_{12}AS_{12} = A^t$, one has that $\Phi(A^t) = \Phi(A)^t$. Since $\Phi(S_{ij}) = S_{ij}$, we have that $\Phi(P_{ij}(k)) = P_{ij}(\varphi(k))$.

Notice that $P_{13}(kl) = P_{12}(k)^{-1}P_{23}(l)^{-1}P_{12}(k)P_{23}(l)$, we have

$$P_{13}(\varphi(kl)) = P_{12}(\varphi(k))^{-1}P_{23}(\varphi(l))^{-1}P_{12}(\varphi(k))P_{23}(\varphi(l)) = P_{13}(\varphi(k)\varphi(l)).$$

Hence

$$\varphi(kl) = \varphi(k)\varphi(l),$$

i.e., φ is a nontrivial ring homomorphism of \mathbb{F} . Consequently $\varphi(1) = 1$. Since $\text{SL}(n, \mathbb{F})$ are generated by $P_{ij}(k)$, one has that $\Phi(A) = (\varphi(a_{ij}))$, for any $A = (a_{ij}) \in \text{SL}(n, \mathbb{F})$.

Now we are ready to complete the proof. For any $A \in \text{GL}(n, \mathbb{F})$, $A = D_1(k)A_0$, where $k = \det(A)$ and $A_0 \in \text{SL}(n, \mathbb{F})$. It is easy to see that $\Phi(D_1(k)) = s\text{diag}(t, 1, \dots, 1)$ for some $s, t \in \mathbb{F}^*$. Similarly, $\Phi(D_2(k)) = s\text{diag}(1, t, 1, \dots, 1)$. Noticing that $D_1(k)D_2(1/k) \in \text{SL}(n, \mathbb{F})$, then we have

$$\Phi(D_1(k)D_2(1/k)) = \Phi(\text{diag}(k, 1/k, 1, \dots, 1)).$$

Therefore, $\text{diag}(t, 1/t, 1, \dots) = \text{diag}(\varphi(k), 1/\varphi(k), 1, \dots, 1)$, i.e., $t = \varphi(k)$. Denoting $s = \lambda(\det(A))$, where λ is a multiplicative map on \mathbb{F}^* , we have $\Phi(A) = \lambda(\det(A))(\varphi(a_{ij}))$, for any $A = (a_{ij}) \in \text{GL}(n, \mathbb{F})$. \square

4. NON-DEGENERATE MULTIPLICATIVE MAPS

In this section, we study non-degenerate multiplicative maps Φ on $M_n(\mathbb{F})$, i.e., there exists some degenerate matrix A such that $\Phi(A) \neq 0$. The main result is the following theorem, which is well-known in some sense (see [7], for example). One can easily see that it is a consequence of the proof of Theorem 3.1.

Theorem 4.1. *Let $\Phi : M_n(\mathbb{F}) \rightarrow M_n(\mathbb{F})$ be a non-degenerate multiplicative map. Then, for any $A = (a_{ij}) \in M_n(\mathbb{F})$,*

$$\Phi(A) = R^{-1}(\varphi(a_{ij}))R \quad \text{or} \quad \Phi(A) = R^{-1}(\varphi(a_{ij})^*)'R,$$

for some ring homomorphism $\varphi : \mathbb{F} \rightarrow \mathbb{F}$ and some invertible matrix R .

Proof. If $\Phi(A) \neq 0$ for some (hence all) matrix of rank 1, letting $F_{ij} = \Phi(E_{ij})$, one has $F_{ij}F_{kl} = \delta_{jk}F_{il}$. It is a standard result that there exists a matrix R such that $RF_{ij}R^{-1} = E_{ij}$. Replacing Φ by $R\Phi R^{-1}$, we may assume that $\Phi(E_{ij}) = E_{ij}$. For any i, j and for any $b \in \mathbb{F}$, it is easy to see that $\Phi(bE_{ij}) = b'E_{ij}$, for some $b' \in \mathbb{F}$. Then we have a map $\varphi : \mathbb{F} \rightarrow \mathbb{F}$ such that $\varphi(b) = b'$. The map φ is independent of i, j since $E_{ki}E_{ij}E_{jl} = E_{kl}$. For any $B = (b_{ij}) \in M_n(\mathbb{F})$, since $E_{ii}\Phi(B)E_{jj} = \Phi(E_{ii}BE_{jj}) = \Phi(b_{ij}E_{ij}) = \varphi(b_{ij})E_{ij}$, we have $\Phi(B) = (\varphi(b_{ij}))$ such that $\varphi(b_{ij}) = b'_{ij}$, ($i, j = 1, 2, \dots, n$). It's an easy exercise to show that $\varphi(1) = 1$, $\varphi(kl) = \varphi(k)\varphi(l)$ and $\varphi(k+l) = \varphi(k) + \varphi(l)$ for any $k, l \in \mathbb{F}$. Thus φ is a nontrivial ring homomorphism of \mathbb{F} . For any $B = (b_{ij}) \in M(n, \mathbb{F})$, we have $E_{ii}\Phi(B)E_{jj} = \Phi(E_{ii}BE_{jj}) = \Phi(b_{ij}E_{ij}) = \varphi(b_{ij})E_{ij}$. Therefore $\Phi(B) = (\varphi(b_{ij}))$.

Now assume that $\Phi(E_{ij}) = 0$. It is easy to see that if $\Phi(A) = 0$ then $\Phi(B) = 0$ provided $\text{rank}(A) = \text{rank}(B)$. Assume that r is the least integer such that $\Phi(A) \neq 0$ if $\text{rank}(A) = r$. Let E_r be the set of diagonal idempotent matrices. Without loss of generality, we may assume that, for any $A \in E_r$, $\Phi(A)$ is also a diagonal idempotent matrix. For any $A, B \in E_r$, $A \neq B$, then $\text{rank}(AB) < r$ and $\Phi(AB) = \Phi(A)\Phi(B) = 0$. Therefore the image $\Phi(E_r)$ has at most n elements. If $r < n-1$, then E_r has $C_n^r > n$ elements, and hence there exist $A, B \in E_r$ such that $\Phi(A) = \Phi(B)$, which is absurd since $\Phi(AB) = 0$. Thus one can easily see that $r = n-1$ and for any A with $\text{rank}(A) = n-1$, $\text{rank}(\Phi(A)) = 1$. Set $F_j = I_n - E_{jj}$. Without loss of generality, we may assume that $\Phi(F_j) = E_{jj}$. Then with the same method as in the proof of Theorem 3.1, we can easily get the result. \square

Remark 4.2. *In the cases of $\mathbb{F} = \mathbb{Q}$ or \mathbb{R} , it is easy to see that the only ring homomorphism is the identity map. For $\mathbb{F} = \mathbb{Q}$, it is trivial. For $\mathbb{F} = \mathbb{R}$, take any $x, y \in \mathbb{R}$, assume that $x - y > 0$, then there exists an element $z \in \mathbb{R}$ such that $z^2 = x - y$. From $\varphi(x - y) = \varphi(x) - \varphi(y) = \varphi(z)^2$, we have $\varphi(x) > \varphi(y)$. It follows that $\varphi(x) = x$ on \mathbb{R} .*

For complex number field \mathbb{C} , there exists ring homomorphisms which are not surjective.

Acknowledgements: This work was supported by the National Natural Science Foundation of China (Grant No. 11571182) and National Science Foundation for Young Scientists of China (Grant No. 11301287 and Grant No. 11401425).

REFERENCES

- [1] G. M. An and J. C. Hou, Rank-preserving multiplicative maps on $B(X)$, *Linear Algebra Appl.* **342** (2002), 59–78.
- [2] G. M. An and J. C. Hou, Multiplication-Preserving Maps On Matrix Algebras (in Chinese), *Act. Math. Sci.* **28A**(6) (2008), 1194–1205.
- [3] W. S. Cheung, S. Fallet and C. K. Li, Multiplicative preservers on semigroups of matrices, *Linear Algebra Appl.* **335** (2002), 173–186.
- [4] G. Dolinar and P. Šemrl, Determinant preserving maps on matrix algebras, *Linear Algebra Appl.* **348** (2002), 189–192.
- [5] S. H. Hochwald, Multiplicative maps on matrices that preserve the spectrum, *Linear Algebra Appl.* **212/213** (1994), 339–351.
- [6] J. C. Hou, Multiplicative maps on $\mathfrak{B}(X)$, *Science. in. China. Ser. A*, Volume **41** (1998), no. 4, 337–345.
- [7] M. Jodeit and T. Y. Lam, Multiplicative maps of matrix semi-groups, *Arch. Math. Ser. A*, Volume **22** (1968), no. 4, 10–16.
- [8] A. A. Jafarian and A. R. Sourour, Spectrum-preserving linear maps, *J. Funct. Anal.* **66** (1986), 255–261.
- [9] C. K. Li and N. K. Tsing, Linear preserver problems: a brief introduction and some special techniques, *Linear Algebra Appl.* **162–164** (1992), 217–235.
- [10] L. Molnár III, Some multiplicative preservers on $B(H)$, *Linear Algebra Appl.* **301** (1999), 1–13.
- [11] M. Omladi, On operators preserving commutativity, *J. Funct. Anal.* **66** (1986), 105–122.
- [12] M. Omladi and P. Šemrl, Spectrum-preserving additive maps, *Linear Algebra Appl.* **153** (1991), 67–72.
- [13] P. Šemrl, Endomorphisms of matrix semigroups over division rings, *Israel. J. Math.* **163** (2008), 125–138.

(Yang) SCHOOL OF MATHEMATICS, SICHUAN NORMAL UNIVERSITY, CHENGDU, SICHUAN 610068,
P. R. CHINA

Email address: xiaomeiyang@mail.nankai.edu.cn

(Zhu) DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING, P. R. CHINA

Email address: zhufuhai@nju.edu.cn