

The Endomorphism Rings of Supersingular Elliptic Curves over \mathbb{F}_p and the Binary Quadratic Forms

Guanju Xiao^a, Zijian Zhou^a, Yingpu Deng^{b,c}, Longjiang Qu^{a,*}

^a*College of Liberal Arts and Sciences, National University of Defense Technology
Changsha 410073, China*

^b*Key Laboratory of Mathematics Mechanization, NCMIS, Academy of Mathematics and Systems
Science, Chinese Academy of Sciences, Beijing 100190, China*

^c*School of Mathematical Sciences, University of Chinese Academy of Sciences
Beijing 100049, China*

Abstract

It is well known that there is a one-to-one correspondence between the supersingular j -invariants up to the action of $\text{Gal}(\mathbb{F}_{p^2}/\mathbb{F}_p)$ and the isomorphism classes of maximal orders in $B_{p,\infty}$ by Deuring's theorem. Interestingly, we establish a one-to-one correspondence between the \mathbb{F}_p -isomorphism classes of supersingular elliptic curves and the primitive reduced binary quadratic forms with discriminant $-p$ or $-16p$. Due to this correspondence and the fact that the isogenies between elliptic curves over \mathbb{F}_p could be represented by quadratic forms, we show that operations of these isogenies on elliptic curves over \mathbb{F}_p are compatible with compositions of corresponding quadratic forms. Based on these results, we could reduce the security of the CSIDH cryptosystem to the computation of this correspondence explicitly.

Keywords: Supersingular Elliptic Curve, Endomorphism Ring, Binary Quadratic Form

1. Introduction

In 1941, Deuring [7] proved that there is a one-to-one correspondence between isomorphism classes of maximal orders in $B_{p,\infty}$ and isomorphism classes of supersingular elliptic curves up to the action of $\text{Gal}(\mathbb{F}_{p^2}/\mathbb{F}_p)$. For a supersingular elliptic curve E defined over \mathbb{F}_p , its endomorphism ring $\text{End}(E)$ contains a root of $x^2 + p = 0$, this means that the order $\mathbb{Z}[\sqrt{-p}]$ can be embedded into $\text{End}(E)$. Let $q \equiv 3 \pmod{8}$ be a prime and $\left(\frac{p}{q}\right) = -1$. Ibukiyama[8] gave two kinds of maximal orders $\mathcal{O}(q, r)$ and $\mathcal{O}'(q, r')$ where r (resp. r') satisfies $r^2 + p \equiv 0 \pmod{q}$ (resp. $r'^2 + p \equiv 0 \pmod{4q}$), and proved that they are isomorphic to the endomorphism rings of some supersingular elliptic curves which are defined over \mathbb{F}_p .

*Corresponding author

Email addresses: gjxiao@amss.ac.cn (Guanju Xiao), zhouzijian122006@163.com (Zijian Zhou), dengyp@amss.ac.cn (Yingpu Deng), ljqu_happy@hotmail.com (Longjiang Qu)

In this paper, we study the structure of the maximal orders $\mathcal{O}(q, r)$'s and $\mathcal{O}'(q, r')$'s. We establish a correspondence between the maximal orders $\mathcal{O}'(q, r')$'s and the reduced binary quadratic forms with discriminant $-p$. We prove this result by ignoring the subring $\mathbb{Z}[\frac{1+\sqrt{-p}}{2}]$ of $\mathcal{O}'(q, r')$, and the rest of the endomorphism ring $\mathcal{O}'(q, r')$ can be represented by a quadratic form, which is equivalent to the form $(q, r', \frac{r'^2+p}{4q})$ or $(q, -r', \frac{r'^2+p}{4q})$. On the contrary, if we have a reduced form (a, b, c) with discriminant $-p$, then we can get a unique supersingular j -invariant by computing the greatest common divisors in \mathbb{F}_p of different Hilbert class polynomials. Similar results also hold for $\mathcal{O}(q, r)$'s, while the corresponding reduced forms are of discriminant $-16p$ and these forms are not in the principal genus.

We also study the connections of this correspondence and the isogenies between supersingular elliptic curves defined over \mathbb{F}_p . Let $\phi : E \rightarrow E'$ be an isogeny defined over \mathbb{F}_p , where $\deg(\phi)$ is an odd prime. As we know, the isogeny ϕ can be represented by an ideal in $\mathbb{Z}[\sqrt{-p}]$ or $\mathbb{Z}[\frac{1+\sqrt{-p}}{2}]$. Equivalently, we can denote it by a reduced form g . If the curve E corresponds to the form f , then we prove that the curve E' corresponds to the form fg^2 . This shows an explicit relation between isogenies of supersingular elliptic curves defined over \mathbb{F}_p and the endomorphism rings of them.

Note that this correspondence may has a great application in isogeny-based cryptography. Isogeny-based cryptography is a relatively new kind of elliptic-curve cryptography, whose security relies on the problem of finding an explicit isogeny between two given isogenous elliptic curves over a finite field. One of the isogeny-based cryptographic protocols is CSIDH proposed by Castryck et al. [3] in 2018. CSIDH uses the action of an ideal class group on the set of supersingular elliptic curves defined over \mathbb{F}_p . In this paper, we reduced the security of the CSIDH cryptosystem to computing the correspondence explicitly.

The remainder of this paper is organized as follows. In Section 2, we review some preliminaries on supersingular elliptic curves defined over \mathbb{F}_p and binary quadratic forms. We prove the correspondence between supersingular elliptic curves over \mathbb{F}_p and binary quadratic forms in Section 3. In Section 4, we prove that the composition of forms corresponds to the action of isogenies on elliptic curves. In Section 5, we discuss the applications of these results in CSIDH. Finally, we make a conclusion in Section 6.

2. Preliminaries

2.1. Elliptic curves and isogenies

We present some basic facts about elliptic curves over finite fields, and the readers can refer to [11] for more details.

Let \mathbb{F}_q be a finite field with characteristic $p > 3$. An elliptic curve E defined over \mathbb{F}_q can be written as a Weierstrass model $E : Y^2 = X^3 + aX + b$ where $a, b \in \mathbb{F}_q$ and $4a^3 + 27b^2 \neq 0$. The j -invariant of E is $j(E) = 1728 \cdot 4a^3 / (4a^3 + 27b^2)$. Different elliptic curves with the same j -invariant are isomorphic over the algebraic closed field. The

chord-and-tangent addition law makes of $E(\mathbb{F}_q) = \{(x, y) \in \mathbb{F}_q^2 : y^2 = x^3 + ax + b\} \cup \{\infty\}$ an abelian group, where ∞ is the point at infinity. For any integer $n \geq 2$ with $p \nmid n$, let $E[n]$ be the group of n -torsion points on E . We have $E[n] \cong \mathbb{Z}/n\mathbb{Z} \oplus \mathbb{Z}/n\mathbb{Z}$.

Let E_1 and E_2 be elliptic curves defined over \mathbb{F}_q . An isogeny from E_1 to E_2 is a morphism $\phi : E_1 \rightarrow E_2$ satisfying $\phi(\infty) = \infty$. If $\phi(E_1) = \{\infty\}$, we say $\phi = 0$. If $\phi \neq 0$, then ϕ is a surjective group homomorphism with finite kernel. The isogeny ϕ is called a \mathbb{F}_q -isogeny if it is defined over \mathbb{F}_q . Every \mathbb{F}_q -isogeny can be represented as $\phi = (r_1(X), r_2(X) \cdot Y)$. If $r_1(X) = p_1(X)/q_1(X)$ with $p_1, q_1 \in \mathbb{F}_q[X]$ and $\gcd(p_1, q_1) = 1$, then the degree of ϕ is $\max(\deg p_1, \deg q_1)$. The isogeny ϕ is said to be separable if $r_1'(X) \neq 0$. Note that all isogenies of prime degree $\ell \neq p$ are separable. If $\phi : E_1 \rightarrow E_2$ is an isogeny of degree m , then there exists a unique isogeny $\hat{\phi} : E_2 \rightarrow E_1$ satisfying $\hat{\phi} \circ \phi = [m]_{E_1}$ and $\phi \circ \hat{\phi} = [m]_{E_2}$, where $[m]$ is the multiplication-by- m map with degree m^2 . We call $\hat{\phi}$ the dual of ϕ .

An endomorphism of E is an isogeny from E to itself. The Frobenius map $\pi : (x, y) \mapsto (x^q, y^q)$ is an inseparable endomorphism. The characteristic polynomial of π is $x^2 - tx + q$, where t is the trace of π . It is well known that E is supersingular (resp. ordinary) if $p \mid t$ (resp. $p \nmid t$). Moreover, the j -invariant of every supersingular elliptic curve over $\overline{\mathbb{F}_p}$ is proved to be in \mathbb{F}_{p^2} and it is called a supersingular j -invariant.

The set $\text{End}(E)$ of all endomorphisms of E form a ring under the usual addition and composition as multiplication. The set $\text{End}_{\mathbb{F}_q}(E)$ is a subring of $\text{End}(E)$ which contains all the endomorphisms defined over \mathbb{F}_q .

2.2. Imaginary quadratic orders and binary quadratic forms

We recall some basic facts about imaginary quadratic orders and binary quadratic forms. The general references are [2, 5].

Let K be an imaginary quadratic field and O an order of K . The conductor of O is $f = [O_K : O]$, and the discriminant of O is $f^2 D_K$.

Let $I(O)$ be the group of proper fractional O -ideals. The principal proper fractional O -ideals give a subgroup $P(O) \subset I(O)$. The ideal class group of O is $C(O) = I(O)/P(O)$ and the class number of O is $h(O) = \#C(O)$. Let $I(f)$ be the group of fractional O_K -ideals prime to f . Let $P_{\mathbb{Z}}(f)$ be the subgroup of $I(f)$ generated by principal ideals of the form αO_K , where $\alpha \in O_K$ satisfies $\alpha \equiv a \pmod{f O_K}$ for $a \in (\mathbb{Z}/f\mathbb{Z})^\times$. The group $C(O)$ is canonically isomorphic to the ring class group $I(f)/P_{\mathbb{Z}}(f)$. Furthermore, we have the following proposition (see [5, Theorem 7.24]).

Proposition 2.1. *Let O be the order of conductor f in an imaginary quadratic field K . Then*

$$h(O) = \frac{h(O_K)f}{[O_K^\times : O^\times]} \prod_{p|f} \left(1 - \left(\frac{D_K}{p}\right) \frac{1}{p}\right).$$

Furthermore, $h(O)$ is always an integer multiple of $h(O_K)$.

We often denote the class number of O by $h(D)$ if the discriminant of O is D . By Proposition 2.1, we can relate the class numbers $h(m^2 D)$ and $h(D)$ as follows.

Corollary 2.1. *Let $D \equiv 0, 1 \pmod{4}$ be a negative integer and m a positive integer. Then*

$$h(m^2D) = \frac{h(D)m}{[O^\times : O'^\times]} \prod_{p|m} \left(1 - \left(\frac{D}{p} \right) \frac{1}{p} \right),$$

where O and O' are the orders of discriminant D and m^2D respectively.

We denote a quadratic form $f(X, Y) = aX^2 + bXY + cY^2$ by (a, b, c) with $a, b, c \in \mathbb{Z}$. Its discriminant is $D = b^2 - 4ac$. Note that all the forms are positive definite in this paper, which means $D < 0$ and $a > 0$. A form (a, b, c) is called primitive if $\gcd(a, b, c) = 1$. Two forms $f(X, Y)$ and $g(X, Y)$ are equivalent if there are integers p, q, r and s such that

$$f(X, Y) = g(pX + qY, rX + sY) \quad \text{and} \quad ps - qr = 1.$$

In this case, we denote $f \sim g$. If a form (a, b, c) satisfies

- (1) $-a < b \leq a$,
- (2) $a \leq c$,
- (3) if $a = c$ then $b \geq 0$,

then it is reduced. Every primitive positive definite form is equivalent to a unique reduced form (see [5, Theorem 2.8]).

For every $D = b^2 - 4ac$, there is an imaginary quadratic order O such that the discriminant of O is D . Every ideal in O has an integral basis, and we use $[\alpha, \beta] = \mathbb{Z}\alpha + \mathbb{Z}\beta, \alpha, \beta \in O$ to represent an ideal in O . We can relate the ideal class group $C(O)$ and the form class group $C(D)$ defined in [5, §3] as follows:

Proposition 2.2. *Let O be the order of discriminant D in an imaginary quadratic field K . Then:*

- (1) *If $f(X, Y) = aX^2 + bXY + cY^2$ is a primitive positive definite quadratic form of discriminant D , then $[a, (-b + \sqrt{D})/2]$ is a proper ideal of O .*
- (2) *The map sending $f(X, Y)$ to $[a, (-b + \sqrt{D})/2]$ induces an isomorphism $\iota : C(D) \rightarrow C(O)$.*
- (3) *A positive integer m is represented by a form $f(X, Y)$ if and only if m is the norm of some ideal \mathfrak{a} in the corresponding ideal class in $C(O)$.*

Remark 2.1. *If $\mathfrak{a} = [\alpha, \beta]$ is a proper O -ideal with $\text{Im}(\beta/\alpha) > 0$, then*

$$f(X, Y) = \frac{N(\alpha X - \beta Y)}{N(\mathfrak{a})}$$

is a positive definite form of discriminant D . On the level of classes, this map is the inverse to the map ι in Proposition 2.2.

By Proposition 2.2, one can use the primitive reduced quadratic forms to represent the ideal classes in an imaginary quadratic order, and the composition of forms is equivalent to the multiplication of ideals.

2.3. The Isogeny Graphs of Supersingular Elliptic Curves over \mathbb{F}_p

We discuss the isogeny graphs of supersingular elliptic curves over \mathbb{F}_p , and the reader can refer to [6] for more details.

Let E be a supersingular elliptic curve over \mathbb{F}_p with $p > 3$. We know that $\text{End}_{\mathbb{F}_p}(E)$ is an order in $K = \mathbb{Q}(\sqrt{-p})$. Furthermore, we have

$$\mathbb{Z}[\sqrt{-p}] \subseteq \text{End}_{\mathbb{F}_p}(E) \subseteq \mathcal{O}_K,$$

where \mathcal{O}_K is the integer ring of K .

- If $p \equiv 1 \pmod{4}$, we always have $\text{End}_{\mathbb{F}_p}(E) = \mathbb{Z}[\sqrt{-p}]$ for a supersingular elliptic curve E over \mathbb{F}_p .
- If $p \equiv 3 \pmod{4}$, then $\mathbb{Z}[\sqrt{-p}]$ has conductor 2 in \mathcal{O}_K and $\text{End}_{\mathbb{F}_p}(E)$ must be one of these two orders.

We say that the curve E is on the surface (resp. floor) if $\text{End}_{\mathbb{F}_p}(E) = \mathcal{O}_K$ (resp. $\mathbb{Z}[\sqrt{-p}]$). Note that surface and floor coincide for $p \equiv 1 \pmod{4}$.

Let ϕ be an ℓ -isogeny between supersingular elliptic curves E and E' over \mathbb{F}_p . If $\text{End}_{\mathbb{F}_p}(E) \cong \text{End}_{\mathbb{F}_p}(E')$, then ϕ is called horizontal. If E is on the floor and E' is on the surface (resp. E on the surface and E' on the floor), ϕ is called ℓ -isogeny up (resp. down).

Proposition 2.3. (see [6, Theorem 2.7]) *Let $p > 3$ be a prime.*

(1) $p \equiv 1 \pmod{4}$: *There are $h(-4p)$ \mathbb{F}_p -isomorphism classes of supersingular elliptic curves over \mathbb{F}_p , all having the same endomorphism ring $\mathbb{Z}[\sqrt{-p}]$. From every one there is one outgoing \mathbb{F}_p -rational horizontal 2-isogeny as well as two horizontal ℓ -isogenies for every prime $\ell > 2$ with $\left(\frac{-p}{\ell}\right) = 1$.*

(2) $p \equiv 3 \pmod{4}$: *There are two levels in the supersingular isogeny graph. From each vertex there are two horizontal ℓ -isogenies for every prime $\ell > 2$ with $\left(\frac{-p}{\ell}\right) = 1$.*

- *If $p \equiv 7 \pmod{8}$, on each level $h(-p)$ vertices are situated. Surface and floor are connected 1:1 with 2-isogenies and on the surface we also have two horizontal 2-isogenies from each vertex.*
- *If $p \equiv 3 \pmod{8}$, there are $h(-p)$ vertices on the surface and $3h(-p)$ on the floor. Surface and floor are connected 1:3 with 2-isogenies, and there are no horizontal 2-isogenies.*

2.4. Endomorphism rings of supersingular elliptic curves

For a supersingular elliptic curve E over \mathbb{F}_{p^2} , the endomorphism ring $\text{End}(E)$ is isomorphic to a maximal order of $B_{p,\infty}$, where $B_{p,\infty}$ is a quaternion algebra defined over \mathbb{Q} and ramified at p and ∞ .

An order \mathcal{O} of $B_{p,\infty}$ is a subring of $B_{p,\infty}$ which is also a lattice, and it is called a maximal order if it is not properly contained in any other order. Two orders \mathcal{O}_1 and \mathcal{O}_2

are equivalent if and only if there exists an element $\alpha \in B_{p,\infty}^\times$ such that $\mathcal{O}_1 = \alpha^{-1}\mathcal{O}_2\alpha$. Let \mathcal{O} be a maximal order of $B_{p,\infty}$ and I a left ideal of \mathcal{O} . Define the left order $\mathcal{O}_L(I)$ and the right order $\mathcal{O}_R(I)$ of I by

$$\mathcal{O}_L(I) = \{x \in B_{p,\infty} : xI \subseteq I\}, \quad \mathcal{O}_R(I) = \{x \in B_{p,\infty} : Ix \subseteq I\}.$$

Moreover, $\mathcal{O}_L(I) = \mathcal{O}$, and $\mathcal{O}_R(I)$ is also a maximal order, in which case we say that I connects \mathcal{O} and \mathcal{O}' . The reduced norm of I can be defined as

$$\text{Nrd}(I) = \text{gcd}(\{\text{Nrd}(\alpha) | \alpha \in I\}).$$

Deuring [7] gave an equivalence of categories between the supersingular j -invariants and the maximal orders in the quaternion algebra $B_{p,\infty}$. Furthermore, if E is a supersingular elliptic curve with $\text{End}(E) = \mathcal{O}$, then there is a one-to-one correspondence between isogenies $\phi : E \rightarrow E'$ and left \mathcal{O} -ideals I . More details on the correspondence can be found in Chapter 42 of [13].

If the supersingular elliptic curve E is defined over \mathbb{F}_p , then $\text{End}(E)$ contains an element α with minimal polynomial $x^2 + p$ (see [6]). Equivalently, the endomorphism ring $\text{End}(E)$ contains a subring $\mathbb{Z}[\sqrt{-p}]$. Ibukiyama [8] has given an explicit description of all maximal orders \mathcal{O} in $B_{p,\infty}$ containing a root of $x^2 + p = 0$.

Choose a prime integer q such that

$$q \equiv 3 \pmod{8}, \quad \left(\frac{p}{q}\right) = -1. \quad (1)$$

Then $B_{p,\infty}$ can be written as $B_{p,\infty} = \mathbb{Q} + \mathbb{Q}\alpha + \mathbb{Q}\beta + \mathbb{Q}\alpha\beta$ where $\alpha^2 = -p$, $\beta^2 = -q$ and $\alpha\beta = -\beta\alpha$. Let r be an integer such that $r^2 + p \equiv 0 \pmod{q}$. Put

$$\mathcal{O}(q, r) = \mathbb{Z} + \mathbb{Z}\frac{1+\beta}{2} + \mathbb{Z}\frac{\alpha(1+\beta)}{2} + \mathbb{Z}\frac{(r+\alpha)\beta}{q}.$$

When $p \equiv 3 \pmod{4}$, we further choose an integer r' such that $r'^2 + p \equiv 0 \pmod{4q}$ and put

$$\mathcal{O}'(q, r') = \mathbb{Z} + \mathbb{Z}\frac{1+\alpha}{2} + \mathbb{Z}\beta + \mathbb{Z}\frac{(r'+\alpha)\beta}{2q}.$$

Both $\mathcal{O}(q, r)$ and $\mathcal{O}'(q, r')$ are independent of the choices of r and r' up to isomorphism respectively. Ibukiyama's results [8] show that both $\mathcal{O}(q, r)$ and $\mathcal{O}'(q, r')$ are maximal orders of $B_{p,\infty}$ and the endomorphism ring $\text{End}(E)$ is isomorphic to $\mathcal{O}(q, r)$ or $\mathcal{O}'(q, r')$ with suitable choice of q if E is a supersingular elliptic curve over \mathbb{F}_p .

Let $j \in \mathbb{F}_p$ be a supersingular j -invariant and $E(j)$ be the corresponding supersingular elliptic curve. If $j \neq 0, 1728$, then $\text{End}(E(j)) \cong \mathcal{O}'(q, r')$ if $\frac{1+\pi}{2} \in \text{End}(E(j))$ and $\text{End}(E(j)) \cong \mathcal{O}(q, r)$ if $\frac{1+\pi}{2} \notin \text{End}(E(j))$ for some q satisfying (1). Especially, we have $\text{End}(E(0)) \cong \mathcal{O}(3, 1)$. By Lemma 1.8 and Proposition 2.1 of [8], the following proposition holds.

Proposition 2.4. *Suppose $q_1 \neq q_2$ are primes satisfying (1). Let $K = \mathbb{Q}(\sqrt{-p})$. Suppose q_1 and q_2 have prime decompositions $q_1 O_K = \mathfrak{q}_1 \bar{\mathfrak{q}}_1$ and $q_2 O_K = \mathfrak{q}_2 \bar{\mathfrak{q}}_2$.*

- (1) $\mathcal{O}(q_1, r_1) \cong \mathcal{O}'(q_2, r_2)$ if and only if $|\mathcal{O}(q_1, r_1)^\times| = |\mathcal{O}'(q_2, r_2)^\times| = 4$. Then $\mathcal{O}(q_1, r_1) \not\cong \mathcal{O}'(q_2, r_2)$ if one of them is isomorphic to $\text{End}(E)$ for $j(E) \neq 1728$.
- (2) $\mathcal{O}'(q_1, r_1) \cong \mathcal{O}'(q_2, r_2) \Leftrightarrow$ the equation $x^2 + py^2 = 4q_1 q_2$ is solvable over $\mathbb{Z} \Leftrightarrow$ either $\mathfrak{q}_1 \mathfrak{q}_2 \in P(2)$ or $\mathfrak{q}_1 \bar{\mathfrak{q}}_2 \in P(2)$.
- (3) $\mathcal{O}(q_1, r_1) \cong \mathcal{O}(q_2, r_2) \Leftrightarrow$ the equation $x^2 + 4py^2 = q_1 q_2$ is solvable over $\mathbb{Z} \Leftrightarrow$ either $\mathfrak{q}_1 \mathfrak{q}_2 \in P_{\mathbb{Z}}(2)$ or $\mathfrak{q}_1 \bar{\mathfrak{q}}_2 \in P_{\mathbb{Z}}(2)$.

Remark 2.2. *For $p \equiv 3 \pmod{4}$ and $p > 3$, if E is a supersingular elliptic curve defined over \mathbb{F}_p , then $|\text{End}(E)^\times| = 4$ if and only if $j(E) = 1728$.*

For $p \equiv 3 \pmod{4}$, $\mathfrak{q}_1 \mathfrak{q}_2 \in P(2)$ (resp. $\mathfrak{q}_1 \bar{\mathfrak{q}}_2 \in P(2)$) if and only if $\bar{\mathfrak{q}}_1 = \mathfrak{q}_2$ (resp. $\mathfrak{q}_1 = \mathfrak{q}_2$) in the class group $C(-p)$ which means q_1 and q_2 can be represented by the same reduced forms with discriminant $-p$. Note that q_i can be represented by a form f if and only if it can be represented by f^{-1} . Moreover, $\mathcal{O}(q_1, r_1) \cong \mathcal{O}(q_2, r_2)$ if and only if q_1 and q_2 can be represented by the same reduce forms with discriminant $-4p$.

For $p \equiv 1 \pmod{4}$, $\mathcal{O}(q_1, r_1) \cong \mathcal{O}(q_2, r_2)$ if and only if q_1 and q_2 can be represented by the same reduce forms with discriminant $-16p$.

3. Endomorphism rings and binary quadratic forms

By Ibukiyama's results [8], there is a correspondence between the supersingular j -invariants over \mathbb{F}_p and the maximal orders $\mathcal{O}(q, r)$'s and $\mathcal{O}'(q, r')$'s.

In this section, we first show that the maximal orders $\mathcal{O}'(q, r')$'s can be reduced to quadratic forms with discriminant $-p$. We also prove that the maximal orders $\mathcal{O}(q, r)$'s can be reduced to quadratic forms with discriminant $-16p$, and these forms are not in the principal genus. Based on the results above, we establish a correspondence between the \mathbb{F}_p -isomorphism classes of supersingular elliptic curves and the reduced quadratic forms with discriminant $-p$ or $-16p$.

Based on Theorems 1.1 and 1.2 in [14], we have the following lemma about the number of \mathbb{F}_p -roots of Hilbert class polynomials. Note that Hilbert class polynomials are class polynomials in [5, §13].

Lemma 1. *Let $p \equiv 3 \pmod{4}$ be a prime. For a positive integer $1 \leq a < p/4$, if $\left(\frac{a}{p}\right) = 1$ and the equation $x^2 \equiv -p \pmod{4a}$ is solvable, then the Hilbert class polynomial $H_{-4a}(X)$ has \mathbb{F}_p -roots. Moreover, if b is a solution of $x^2 \equiv -p \pmod{4a}$ with $-a < b \leq a$ and $c = \frac{b^2+p}{4a}$ is an integer, then the Hilbert class polynomials $H_{-4a}(X)$ and $H_{-4c}(X)$ have only one common root in \mathbb{F}_p .*

Proof. If $a = 1$ and $p \equiv 3 \pmod{4}$, then the equation $x^2 \equiv -p \pmod{4}$ is solvable and the Hilbert class polynomial $H_{-4}(X)$ has a root 1728 in \mathbb{F}_p . In this case, $c = \frac{p+1}{4}$ and

we can show $H_{-4}(X)$ and $H_{-(p+1)}(X)$ have only one common root in \mathbb{F}_p by Theorem 1.5 in [10]. In the following, we assume $1 < a < p/4$.

If $2 \mid a$, then $p \equiv 7 \pmod{8}$ and $\left(\frac{2}{p}\right) = 1$. For any odd prime $\ell \mid a$, $x^2 \equiv -p \pmod{4a}$ is solvable if and only if $\left(\frac{-p}{\ell}\right) = \left(\frac{\ell}{p}\right) = 1$.

Denote $D = -4a > -p$. Then the prime p does not divide the discriminant of $H_{-4a}(X)$ and [14, Theorem 1.1] holds. Let F be the genus field of $\mathbb{Q}(\sqrt{D})$ and E the maximal real field of F . We have that $x^2 \equiv -p \pmod{4a}$, $4 < 4a < p$ is solvable if and only if p splits completely in E . It follows that the Hilbert class polynomial $H_{-4a}(X)$ has roots in \mathbb{F}_p .

If $-a < b \leq a$ is a solution of $x^2 \equiv -p \pmod{4a}$ and $c = \frac{b^2+p}{4a}$, then the equation $x^2 \equiv -p \pmod{4c}$ is solvable. Moreover, we have $p > 4c$ since $4 < 4a < p$, so the Hilbert class polynomial $H_{-4c}(X)$ has roots in \mathbb{F}_p . We have $p \mid (4a \cdot 4c - x^2)$ if and only if $x = \pm 2b$. By Theorem 1.5 in [10], we have $p \parallel J(-4a, -4c)$. It follows that $H_{-4a}(X)$ and $H_{-4c}(X)$ have only one common root in \mathbb{F}_p . □

Remark 3.1. Let (a, b, c) be a reduced form with discriminant $-p$. We have that $H_{-4a}(X)$ and $H_{-4c}(X)$ have only one common \mathbb{F}_p -root j . Moreover, the number $4a = |D_1|$ is the smallest positive integer such that j is a \mathbb{F}_p -root of $H_{D_1}(X)$, and the number $4c = |D_2|$ is the smallest positive integer such that $H_{D_1}(X)$ and $H_{D_2}(X)$ have a common \mathbb{F}_p -root j .

Theorem 1. Let $p \equiv 3 \pmod{4}$ be a prime with $p > 5$. If E is a supersingular elliptic curve over \mathbb{F}_p with $\text{End}(E) \cong \mathcal{O}'(q, r')$, then there exist two reduced forms with discriminant $-p$ which can represent q and one form is the inverse of the other. On the contrary, given a reduced form with discriminant $-p$ which can represent a prime q satisfying (1), there exists a unique supersingular j -invariant $j \in \mathbb{F}_p$ such that $\text{End}(E(j)) \cong \mathcal{O}'(q, r')$. Moreover, the inverse form corresponds to the same supersingular j -invariant.

Proof. Let E be a supersingular elliptic curve defined over \mathbb{F}_p with $\text{End}(E) \cong \mathcal{O}'(q, r')$. If the imaginary quadratic order $O = \mathbb{Z}[\frac{D+\sqrt{D}}{2}]$ can be embedded into $\mathcal{O}'(q, r')$, then there exists a $\gamma = w + x\frac{1+\alpha}{2} + y\beta + z\frac{(r'+\alpha)\beta}{2q} \in \mathcal{O}'(q, r')$ such that:

$$\text{Trd}(\gamma) = 2w + x = D$$

and

$$\text{Nrd}(\gamma) = \left(w + \frac{x}{2}\right)^2 + \frac{p}{4}x^2 + q\left(y + \frac{zr'}{2q}\right)^2 + p\frac{z^2}{4q} = \frac{D^2 - D}{4}.$$

These equations are equivalent to

$$2w + x = D$$

and

$$px^2 + q\left(2y + \frac{zr'}{q}\right)^2 + \frac{pz^2}{q} = -D.$$

We put $x = 0$ in the above equation, so we have the following equation

$$4qy^2 + 4r'yz + \frac{r'^2 + p}{q}z^2 = -D.$$

We need to discuss the choice of r' . As we know, $\pm r' + 2kq, k \in \mathbb{Z}$ are the solutions of the equation $x^2 + p \equiv 0 \pmod{4q}$, so we get two forms $f = (q, r', \frac{r'^2+p}{4q})$ and $f^{-1} = (q, -r', \frac{r'^2+p}{4q})$ from $\mathcal{O}'(q, r')$. Suppose f (resp. f^{-1}) is equivalent to a reduced form (a, b, c) (resp. $(a, -b, c)$). Then the form (a, b, c) (resp. $(a, -b, c)$) can represent the prime q .

On the contrary, let (a, b, c) be a reduced quadratic form with discriminant $-p$ which can represent a prime q satisfying (1). Since $b^2 - 4ac = -p$ and $1 \leq a < \sqrt{p/3} < p/4$, we have that $H_{-4a}(X) \pmod{p}$ and $H_{-4c}(X) \pmod{p}$ have a unique common root $j \in \mathbb{F}_p$ by Lemma 1. By Ibukiyama's theorems, the endomorphism ring of $E(j)$ is isomorphic to the maximal order $\mathcal{O}(q_j, r_j)$ or $\mathcal{O}'(q_j, r'_j)$ for some q_j satisfying (1).

If $a = 1$, then $j = 1728$ and we can assume $\text{End}(E(1728)) \cong \mathcal{O}'(q_j, r'_j)$. In the following, we assume $1 < a < c < p/4$.

Suppose $\text{End}(E(j)) \cong \mathcal{O}(q_j, r_j)$. Since j is the common root of $H_{-4a}(X) \pmod{p}$ and $H_{-4c}(X) \pmod{p}$, we have that the form $(q_j, 4r_j, \frac{4r_j^2+4p}{q_j})$ can represent $4a$ and $4c$ by the proof of Theorem 1 in [9]. Moreover, the number $4a = |D_1|$ is the smallest positive integer such that j is a \mathbb{F}_p -root of $H_{D_1}(X)$, and the number $4c = |D_2|$ is the smallest positive integer such that $H_{D_1}(X) \pmod{p}$ and $H_{D_2}(X) \pmod{p}$ have a common \mathbb{F}_p -root j . The first (resp. second) successive minimum of $(q_j, 4r_j, \frac{4r_j^2+4p}{q_j})$ is $\sqrt{4a}$ (resp. $\sqrt{4c}$). By Proposition 5.7.3 and Theorem 5.7.6 in [1], the form $(q_j, 4r_j, \frac{4r_j^2+4p}{q_j})$ is equivalent to the form $(4a, 4b, 4c)$ or $(4a, -4b, 4c)$, which is a contradiction. We must have $\text{End}(E(j)) \cong \mathcal{O}'(q_j, r'_j)$.

It is easy to show that the form $(q_j, r'_j, \frac{r'^2+p}{4q_j})$ is equivalent to the form (a, b, c) or $(a, -b, c)$. It follows that q can be represented by $(q_j, r'_j, \frac{r'^2+p}{4q_j})$, and the endomorphism ring of $E(j)$ is isomorphic to $\mathcal{O}'(q, r')$. □

For the maximal orders $\mathcal{O}(q, r)$'s, similar result also holds, while the corresponding forms are of discriminant $-16p$.

We review some facts about genera of positive definite quadratic forms. Let $p > 3$ be a prime. The primitive forms with discriminant $-16p$ have two assigned characters $\chi(a) = \left(\frac{a}{p}\right)$ and $\delta(a) = (-1)^{(a-1)/2}$. Furthermore, the primitive forms with discriminant $-16p$ can be divided into two genera (see [5, Theorem 3.15]). The genus which contains the principal form $(1, 0, 4p)$ is called the principal genus, and we call the other non-principal genus.

Let (a, b, c) be a reduced form with discriminant $-16p$ in the non-principal genus. If the prime q can be represented by (a, b, c) , then $1 = \left(\frac{-p}{q}\right) = \left(\frac{-1}{q}\right) \left(\frac{p}{q}\right)$. If $p \equiv 1$

(mod 4), then $\left(\frac{p}{q}\right) = \left(\frac{q}{p}\right) = \left(\frac{-1}{q}\right) = -1$. If $p \equiv 3 \pmod{4}$, then $1 = \left(\frac{-p}{q}\right) = \left(\frac{q}{p}\right)$ and $\left(\frac{-1}{q}\right) = -1$. In this case, we always have $\left(\frac{-q}{p}\right) = -1$.

If a (resp. c) is odd, then $a \equiv 3 \pmod{4}$ (resp. $c \equiv 3 \pmod{4}$). If a is even, we claim $4 \mid a$. If $a \equiv 2 \pmod{4}$, then $8 \mid b^2 (= 4ac - 16p)$. It follows $4 \mid b$ and $2 \mid c$. This is a contradiction. Similarly, if c is even, then $4 \mid c$.

Similar to Lemma 1, we have the following lemma.

Lemma 2. *Let p be a prime and $-a$ the discriminant of an imaginary quadratic order satisfying $3 \leq a < p$. If $\left(\frac{-a}{p}\right) = -1$ and the equation $x^2 \equiv -16p \pmod{4a}$ is solvable, then the Hilbert class polynomial $H_{-4a}(X)$ has \mathbb{F}_p -roots. Moreover, let b be a solution of $x^2 \equiv -16p \pmod{4a}$ with $-a < b \leq a$. If the form (a, b, c) ($c = \frac{b^2 + 16p}{4a}$) is primitive, then the Hilbert class polynomials $H_{-a}(X)$ and $H_{-c}(X)$ have only one common root in \mathbb{F}_p .*

Theorem 2. *Let E be a supersingular elliptic curve over \mathbb{F}_p with $p > 3$. If $\text{End}(E) \cong \mathcal{O}(q, r)$, then there exist two reduced forms with discriminant $-16p$ in the non-principal genus which can represent the prime q and one form is the inverse of the other. On the contrary, given a reduced form with discriminant $-16p$ in the non-principal genus, there is a unique supersingular j -invariant in \mathbb{F}_p with $\text{End}(E(j)) \cong \mathcal{O}(q, r)$. Moreover, the inverse form corresponds to the same supersingular j -invariant.*

Proof. The proof is similar to that of Theorem 1. Suppose the imaginary quadratic order $O = \mathbb{Z}[\frac{D + \sqrt{D}}{2}]$ can be embedded in $\mathcal{O}(q, r)$. Then there exists a $\gamma = w + x\frac{1+\beta}{2} + y\frac{\alpha(1+\beta)}{2} + z\frac{(r+\alpha)\beta}{q} \in \mathcal{O}(q, r)$ such that:

$$\text{Trd}(\gamma) = D \quad \text{and} \quad \text{Nrd}(\gamma) = \frac{D^2 - D}{4}.$$

These equations are equivalent to

$$2w + x = D$$

and

$$py^2 + q\left(x + \frac{2zr}{q}\right)^2 + pq\left(y + \frac{2z}{q}\right)^2 = -D.$$

We put $y = 0$ in the above equation, so we have the following equation

$$qx^2 + 4rxz + \frac{4r^2 + 4p}{q}z^2 = -D.$$

Similarly, we get two forms $f = (q, 4r, \frac{4r^2 + 4p}{q})$ and $g = f^{-1} = (q, -4r, \frac{4r^2 + 4p}{q})$ with discriminant $-16p$, and we have $f(x, z) \equiv g(x, z) \equiv 0, 3 \pmod{4}$ for any $x, z \in \mathbb{Z}$. If the form f (resp. g) is equivalent to a reduced form (a, b, c) (resp. $(a, -b, c)$), then the form (a, b, c) (resp. $(a, -b, c)$) can represent the prime q . Moreover, $\left(\frac{-1}{q}\right) = -1$, so the reduced form (a, b, c) (resp. $(a, -b, c)$) is in the non-principal genus.

On the contrary, if (a, b, c) is a primitive reduced form in the non-principal genus with discriminant $b^2 - 4ac = -16p$, then $\left(\frac{-a}{p}\right) = -1$ and $a, c \equiv 0, 3 \pmod{4}$. By Lemma 2, the Hilbert class polynomials $H_{-a}(X)$ and $H_{-c}(X) \pmod{p}$ have one common root $j \in \mathbb{F}_p$.

If $p \equiv 3 \pmod{4}$ and $a = 4$, then $j = 1728$. Without loss of generality, we can assume the endomorphism ring of $E(j)$ is isomorphic to $\mathcal{O}(q_j, r_j)$. If $p \equiv 2 \pmod{3}$ and $a = 3$, then $j = 0$ and $\text{End}(E(j)) \cong \mathcal{O}(3, 1)$. If $p = 5$, then there is only one supersingular j -invariant 0 in \mathbb{F}_p . In the following, we assume $p > 5$ and $a > 4$, so $a < c < p$.

We claim that j is not a root of the Hilbert class polynomial $H_{-p}(X) \pmod{p}$. We can assume $4 \nmid a$ and $a < p$. If there exist integers x and $k \geq 0$ such that $4pk = ap - x^2$, then $x^2 = p(a - 4k)$. It follows that $p \mid x$ and $p \mid (a - 4k)$. This is a contradiction since $0 < a - 4k \leq a < p$. Let $d_1 = -a$ and $d_2 = -p$. We have $v_p(F(\frac{d_1 d_2 - x^2}{4})) = 0$ for any $|x| < \sqrt{d_1 d_2}$ by Theorem 1.1 in [10]. This shows that the Hilbert class polynomials $H_{-a}(X)$ and $H_{-p}(X)$ have not common root in \mathbb{F}_p , and j is not a root of $H_{-p}(X) \pmod{p}$. It follows that the order $\mathbb{Z}[\frac{1+\sqrt{-p}}{2}]$ is not embedded into the endomorphism ring of $E(j)$, so we can assume $\text{End}(E(j)) \cong \mathcal{O}(q_j, r_j)$ for some q_j satisfying (1).

In fact, the number $a = |D_1|$ is the smallest positive integer such that j is a \mathbb{F}_p -root of $H_{D_1}(X)$, and the number $c = |D_2|$ is the smallest positive integer such that $H_{D_1}(X) \pmod{p}$ and $H_{D_2}(X) \pmod{p}$ have a common \mathbb{F}_p -root j . Since $a < c < p$, the quadratic form $(q_j, 4r_j, \frac{4r_j^2 + 4p}{q_j})$ can represent a and c . It follows that the form $(q_j, 4r_j, \frac{4r_j^2 + 4p}{q_j})$ can be reduced to (a, b, c) or $(a, -b, c)$. Since the prime q can be represented by (a, b, c) , we have $\mathcal{O}(q_j, r_j) \cong \mathcal{O}(q, r)$. Obviously, the inverse form $(a, -b, c)$ corresponds to the same j -invariant. □

If $p \equiv 3 \pmod{4}$, then the reduced form $(1, 1, (p+1)/4)$ is the unique one with order ≤ 2 in the class group $C(-p)$. Moreover, the reduced form $(4, 0, p)$ is the unique one in the non-principal genus with order ≤ 2 in the class group $C(-16p)$. In fact, the forms $(1, 1, (p+1)/4)$ and $(4, 0, p)$ correspond the same supersingular j -invariant 1728. If $p \equiv 1 \pmod{4}$, then there exists no form in the non-principal genus with order ≤ 2 in the class group $C(-16p)$.

For a prime $p > 5$, let S_p be the set of all supersingular j -invariants in \mathbb{F}_p . If $p \equiv 1 \pmod{4}$, the number of reduced forms with discriminant $-16p$ in non-principal genus is $h(-16p)/2 = h(-4p)$ and $\#S_p = h(-4p)/2$. If $p \equiv 3 \pmod{4}$, $\#S_p = (h(-4p) + 1)/2 + (h(-p) + 1)/2 - 1 = h(-4p)/2 + h(-p)/2$. So we have the following formula:

$$\#S_p = \begin{cases} \frac{1}{2}h(-4p) & \text{if } p \equiv 1 \pmod{4}, \\ h(-p) & \text{if } p \equiv 7 \pmod{8}, \\ 2h(-p) & \text{if } p \equiv 3 \pmod{8}, \end{cases}$$

where $h(D)$ is the class number of the imaginary quadratic order with discriminant D . This formula coincides with the result in [6].

Theorems 1 and 2 establish a correspondence between reduced quadratic forms and supersingular j -invariants. Notice that this correspondence is not 1-to-1. In the following of this section, we consider the quadratic twist curves and the isomorphism class of supersingular elliptic curves over \mathbb{F}_p .

Let $p > 3$ be a prime. Choose a non-square element $\mu \in \mathbb{F}_p$. For an elliptic curve $E : Y^2 = X^3 + aX + b$, we define its quadratic twist curve \tilde{E} by the following way:

$$\begin{aligned} (1) \quad \tilde{E} : Y^2 &= X^3 + a\mu^2 X + b\mu^3 & \text{if } j(E) \neq 1728; \\ (2) \quad \tilde{E} : Y^2 &= X^3 + a\mu X & \text{if } j(E) = 1728. \end{aligned}$$

This definition is a simple case in Proposition 5.4 of [11, X.5]. The quadratic twist curves have the same j -invariant, while they are not isomorphic in \mathbb{F}_p .

Let $\text{Iso}(\mathbb{F}_p, \frac{1+\pi}{2})$ (resp. $\text{Iso}(\mathbb{F}_p, \pi)$) denote the set of the \mathbb{F}_p -isomorphism classes of supersingular elliptic curves over \mathbb{F}_p with endomorphism ring $\mathcal{O}'(q, r')$ (resp. $\mathcal{O}(q, r)$). We have the following theorem.

Theorem 3. *Let $p > 3$ be a prime. Every class in $\text{Iso}(\mathbb{F}_p, \frac{1+\pi}{2})$ is 1-to-1 corresponding to a reduced quadratic form with discriminant $-p$. Every class in $\text{Iso}(\mathbb{F}_p, \pi)$ is 1-to-1 corresponding to a reduced form in the non-principal genus with discriminant $-16p$.*

Proof. For $j(E) \neq 1728$, the \mathbb{F}_p -isomorphism class containing E corresponds to the form (a, b, c) , and the \mathbb{F}_p -isomorphism class containing \tilde{E} corresponds to $(a, -b, c)$.

As we know, the root of the Hilbert class polynomial $H_{-4}(X) = X - 1728$ is a supersingular j -invariant if and only if $p \equiv 3 \pmod{4}$. In this case, we use $E(1728) : Y^2 = X^3 - X$ to represent an isomorphism class in \mathbb{F}_p . Its quadratic twist curve can be written as $\tilde{E}(1728) : Y^2 = X^3 + 4X$. Furthermore, the curve $E(1728)$ is in the surface of the isogeny graph, so the curve $E(1728)$ corresponds to the form $(1, 1, (p+1)/4)$. The curve $\tilde{E}(1728)$ is on the floor, and it corresponds to the form $(4, 0, p)$. □

Example 1. If $p = 83$, then there are 12 \mathbb{F}_{83} -isomorphism classes of supersingular elliptic curves. The curves and the corresponding forms are as following.

$$\begin{array}{llll} E_0 : y^2 = x^3 - x & f_0 = (1, 1, 21) & \tilde{E}_0 : y^2 = x^3 + x & g_0 = (4, 0, 83) \\ E_1 : y^2 = x^3 + 13x^2 - x & f_1 = (3, 1, 7) & \tilde{E}_1 : y^2 = x^3 - 13x^2 - x & f_1^{-1} = (3, -1, 7) \\ E_2 : y^2 = x^3 + 11x^2 + x & g_2 = (11, -6, 31) & \tilde{E}_2 : y^2 = x^3 - 11x^2 + x & g_2^{-1} = (11, 6, 31) \\ E_3 : y^2 = x^3 + 12x^2 + x & g_3 = (7, 4, 48) & \tilde{E}_3 : y^2 = x^3 - 12x^2 + x & g_3^{-1} = (7, -4, 48) \\ E_4 : y^2 = x^3 + 6x^2 + x & g_4 = (16, -12, 23) & \tilde{E}_4 : y^2 = x^3 - 6x^2 + x & g_4^{-1} = (16, 12, 23) \\ E_5 : y^2 = x^3 - 13x^2 + x & g_5 = (3, -2, 111) & \tilde{E}_5 : y^2 = x^3 + 13x^2 + x & g_5^{-1} = (3, 2, 111) \end{array}$$

The supersingular elliptic curves are from Remark 5 in [3]. Note that the forms f_i and f_i^{-1} are of discriminant -83 , and the forms g_i and g_i^{-1} are of discriminant -83×16 . In fact, the corresponding form is not unique except for E_0 and \tilde{E}_0 . We just choose one correspondence in this example.

We have established a correspondence between the set of \mathbb{F}_p -isomorphism classes of supersingular elliptic curves over \mathbb{F}_p and set of reduced quadratic forms. In fact, this correspondence is not unique. Moreover, the reduced forms with discriminant $-p$ form a group by composition. This inspires us to research the isogenies between supersingular elliptic curves.

4. Isogenies and ideals

As we have seen, the correspondence which we have established in Section 3 is only about sets. Now, consider the isogenies between supersingular elliptic curves defined over \mathbb{F}_p . Note that these isogenies correspond to ideals in imaginary quadratic orders, hence they could be represented by quadratic forms.

In this section, we present the connection between the quadratic forms corresponding to isogenies over \mathbb{F}_p and the quadratic forms which correspond to supersingular elliptic curves defined over \mathbb{F}_p .

To study the connections between $\mathcal{O}(q, r)$ and $\mathcal{O}'(q, r')$, we recall the connections between quadratic forms with different discriminants.

4.1. Quadratic forms with different discriminants

We now review some basic facts about quadratic forms with different discriminants. If

$$S = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$$

is a 2×2 matrix with integer coefficients and determinant $\alpha\delta - \beta\gamma = s > 0$, then the change of variables takes a form $f = (a, b, c)$ of discriminant D to a form

$$f' = (a', b', c') = (a\alpha^2 + b\alpha\gamma + c\gamma^2, b(\alpha\delta + \beta\gamma) + 2(a\alpha\beta + c\gamma\delta), a\beta^2 + b\beta\delta + c\delta^2)$$

of discriminant Ds^2 . In matrix notation this is

$$\begin{pmatrix} a' & b'/2 \\ b'/2 & c' \end{pmatrix} = \begin{pmatrix} \alpha & \gamma \\ \beta & \delta \end{pmatrix} \begin{pmatrix} a & b/2 \\ b/2 & c \end{pmatrix} \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

which we will write as $f' = S^T f S$ for brevity. We shall call such a matrix S a transformation of determinant s and shall say that f' is derived from f by the transformation of determinant s .

Any primitive form of discriminant Ds^2 can be derived from a primitive form of discriminant D by the application of a transformation S of determinant s . We define two transformations S_1 and S_2 of discriminant s to be $SL_2(\mathbb{Z})$ -right-equivalent if there exists a matrix $M \in SL_2(\mathbb{Z})$ such that $S_1 M = S_2$. Moreover, we have the following proposition [2, Proposition 7.2].

Proposition 4.1. *Let s be a prime integer. The $SL_2(\mathbb{Z})$ -right-equivalent transformations have as equivalence class representatives the $s + 1$ transformations*

$$S_s = \begin{pmatrix} 1 & 0 \\ 0 & s \end{pmatrix} \quad \text{and} \quad S_i = \begin{pmatrix} s & i \\ 0 & 1 \end{pmatrix},$$

for $0 \leq i \leq s-1$.

Remark 4.1. Let q be a prime with $q^2 \nmid Ds^2$. If q can be represented by f , then it must be represented by some of $S_i^T f S_i$ with $0 \leq i \leq s$. On the contrary, if a prime q can be represented by any of $S_i^T f S_i$, then q must be represented by f .

Proposition 4.2. Let $p \equiv 3 \pmod{4}$ be a prime. If $q_1 \neq q_2$ are primes satisfying (1), then $\mathcal{O}(q_1, r_1) \cong \mathcal{O}(q_2, r_2)$ if and only if q_1 and q_2 can be represented by the same primitive reduced quadratic form with discriminant $-16p$.

Proof. By Proposition 2.4, $\mathcal{O}(q_1, r_1) \cong \mathcal{O}(q_2, r_2)$ if and only if q_1 and q_2 can be represented by the same primitive reduced quadratic form with discriminant $-4p$.

Assume that the prime q_i satisfying (1) can be represented by a primitive form (a, b, c) with discriminant $-4p$. We have $2 \mid b$. For $s = 2$, the following three forms with discriminant $-16p$ are derived from (a, b, c) :

$$(a, 2b, 4c) \quad (4a, 2b, c) \quad (4a, 4a + 2b, a + b + c).$$

For $p \equiv 3 \pmod{4}$, we have $ac = \frac{b^2 + 4p}{4} = \left(\frac{b}{2}\right)^2 + p$.

- If $b/2$ is even, then $ac \equiv 3 \pmod{4}$. It follows that only $(a, 2b, 4c)$ and $(4a, 2b, c)$ are primitive. Moreover, only one of $(a, 2b, 4c)$ and $(4a, 2b, c)$ can represent q_i .
- If $b/2$ is odd, then $4 \mid ac$. We can assume $4 \mid a$ and $2 \nmid c$ because the form (a, b, c) is primitive. It follows that only $(4a, 2b, c)$ and $(4a, 4a + 2b, a + b + c)$ are primitive. Moreover, only one of $(4a, 2b, c)$ and $(4a, 4a + 2b, a + b + c)$ can represent q_i .

Moreover, q_1 and q_2 can be represented by the same reduced quadratic form with discriminant $-16p$. On the contrary, if q_1 and q_2 can be represented by the same reduced quadratic form with discriminant $-16p$, then $\mathcal{O}(q_1, r_1) \cong \mathcal{O}(q_2, r_2)$ obviously. \square

In the following of this subsection, we discuss the reduced quadratic forms with discriminant $-16p$ derived from (a, b, c) with discriminant $-p$.

Proposition 4.3. If $p \equiv 3 \pmod{8}$, then every reduced form with discriminant $-p$ divides six reduced forms with discriminant $-16p$ and three of them are in the non-principal genus. If $p \equiv 7 \pmod{8}$, then every reduced form with discriminant $-p$ divides two reduced forms with discriminant $-16p$ and one of them is in the non-principal genus.

Proof. Let (a, b, c) be a reduced form with discriminant $-p$. The following three forms with discriminant $-4p$ are derived from (a, b, c) .

$$(a, 2b, 4c) \quad (4a, 2b, c) \quad (4a, 4a + 2b, a + b + c)$$

(1) If $p \equiv 3 \pmod{8}$, then ac is odd. It follows that the above three forms are primitive. Since $h(-4p) = 3h(-p)$, these three forms are not equivalent. The following nine forms with discriminant $-16p$ are derived from (a, b, c) .

$$\begin{array}{lll} (4a, 4b, 4c) & (4a, 4b, 4c) & (4a, 8a + 4b, 4a + 4b + 4c) \\ (a, 4b, 16c) & (16a, 4b, c) & (16a, 8a + 4b, a + b + c) \\ (4a, 4a + 4b, a + 2b + 4c) & (16a, 16a + 4b, 4a + 2b + c) & (16a, 24a + 4b, 9a + 3b + c) \end{array}$$

It is obvious that the three forms in the first row are not primitive, and the other six forms are not equivalent since $h(-16p) = 2h(-4p)$. Moreover, only one form in every column is in the non-principal genus.

(2) If $p \equiv 7 \pmod{8}$, then ac is even. Only one of the three forms is primitive. Without loss of generality, we assume the form $(a, 2b, 4c)$ is primitive. In this case, one of $(a, 4b, 16c)$ and $(4a, 4a + 4b, a + 2b + 4c)$ is in the non-principal genus. \square

4.2. Isogenies and quadratic forms

Denote the imaginary quadratic field $K = \mathbb{Q}(\sqrt{-p})$. Let O be an order in K and O_K be the maximal order.

Let E_1, E_2 be supersingular elliptic curves defined over \mathbb{F}_p . Assume that the isogeny $\phi : E_1 \rightarrow E_2$ is also defined over \mathbb{F}_p . Without loss of generality, we can restrict to the case when $\deg(\phi) = \ell$ is a prime, since every isogeny can be written as compositions of isogenies of prime degree. We divide the remainder of this subsection into three cases according to the endomorphism rings of E_1 and E_2 .

Case 1: $\text{End}_{\mathbb{F}_p}(E_1) \cong \text{End}_{\mathbb{F}_p}(E_2) \cong \mathbb{Z}[\frac{1+\sqrt{-p}}{2}]$.

In this case, we must have $p \equiv 3 \pmod{4}$. For simplicity, we write $\mathcal{O}'(q, r)$ with $r^2 + p \equiv 0 \pmod{4q}$. We distinct maximal orders $\mathcal{O}'(q, r)$ and $\mathcal{O}'(q, -r)$, even though they are isomorphic as quaternion orders. If $\text{End}(E_1) \cong \mathcal{O}'(q_1, r_1)$, without loss of generality, we can assume that E_1 corresponds to the form $(q_1, r_1, \frac{r_1^2+p}{4q_1})$.

Theorem 4. *For a prime $p \equiv 3 \pmod{4}$, let E_1 be a supersingular elliptic curve with $\text{End}(E_1) \cong \mathcal{O}'(q_1, r_1)$. If $\phi : E_1 \rightarrow E_2$ is an isogeny defined over \mathbb{F}_p of degree ℓ with $\text{End}_{\mathbb{F}_p}(E_2) \cong O_K$, then there is a prime q_2 satisfying (1) which can be represented by the form $(q_1, r_1, \frac{r_1^2+p}{4q_1})(\ell, b, \frac{b^2+p}{4\ell})^2$ such that $\text{End}(E_2) \cong \mathcal{O}'(q_2, r_2)$.*

Proof. For $p \equiv 3 \pmod{4}$, if $\text{End}(E_1) \cong \mathcal{O}'(q_1, r_1)$, then we can assume E_1 corresponds to the quadratic form $(q_1, r_1, \frac{r_1^2+p}{4q_1})$. We also assume $q_1 O_K = \mathfrak{q}_1 \bar{\mathfrak{q}}_1$ with $\bar{\mathfrak{q}}_1 = [q_1, \frac{-r_1+\sqrt{-p}}{2}]$.

If $\phi : E_1 \rightarrow E_2$ is an isogeny defined over \mathbb{F}_p and $\text{End}_{\mathbb{F}_p}(E_1) \cong \text{End}_{\mathbb{F}_p}(E_2) \cong O_K$, then we can assume $\text{Ker}(\phi) = \{P \in E_1[\ell] \mid \frac{-b+\sqrt{-p}}{2}(P) = \infty\}$ with $b^2 \equiv -p \pmod{4\ell}$. Note that $\mathfrak{l} = [\ell, (-b + \sqrt{-p})/2]$ is an ideal in O_K , and it corresponds to the form $(\ell, b, \frac{b^2+p}{4\ell})$ by the isomorphism in Proposition 2.2. It follows that $\mathcal{O}'(q_1, r_1)\mathfrak{l}$ is the ideal kernel of ϕ . Moreover, $\mathfrak{l}^{-1}\mathcal{O}'(q_1, r_1)\mathfrak{l}$ is the right order of $\mathcal{O}'(q_1, r_1)\mathfrak{l}$ with $\mathfrak{l}^{-1} = \bar{\mathfrak{l}}/\ell$.

By Proposition 4.2 in [8], we have $\text{End}(E_2) \cong \mathcal{O}'(q_2, r_2)$ for any q_2 satisfying (1) which is the norm of a prime ideal $\mathfrak{q}_2 \in \bar{\mathfrak{q}}_1 \ell^2 P(O_K)$. The ideal $\bar{\mathfrak{q}}_1 = [q_1, \frac{-r_1 + \sqrt{-p}}{2}]$ corresponds to the form $(q_1, r_1, \frac{r_1^2 + p}{4q_1})$, so E_2 corresponds to the form $(q_2, r_2, \frac{r_2^2 + p}{4q_1}) \sim (q_1, r_1, \frac{r_1^2 + p}{4q_1})(\ell, b, \frac{b^2 + p}{4\ell})^2$ and $\text{End}(E_2) \cong \mathcal{O}'(q, r_2)$. \square

Remark 4.2. For $(\frac{-p}{\ell}) = 1$, there are two ℓ -isogeny $\phi : E_1 \rightarrow E_2$ and $\phi' : E_1 \rightarrow E'_2$ over \mathbb{F}_p with $\text{End}_{\mathbb{F}_p}(E_2) \cong \text{End}_{\mathbb{F}_p}(E'_2) \cong O_K$. The isogeny ϕ' corresponds to the form $(\ell, -b, \frac{b^2 + p}{4\ell})$, and E'_2 corresponds to the form $(q_1, r_1, \frac{r_1^2 + p}{4q_1})(\ell, -b, \frac{b^2 + p}{4\ell})^2$.

Case 2: $\text{End}_{\mathbb{F}_p}(E_1) \cong \text{End}_{\mathbb{F}_p}(E_2) \cong \mathbb{Z}[\sqrt{-p}]$.

If $\text{End}(E_1) \cong \mathcal{O}(q_1, r_1)$ and $\text{End}(E_2) \cong \mathcal{O}(q_2, r_2)$ and $\deg(\phi) = \ell$ is an odd prime, we have the following proposition.

Proposition 4.4. Let \mathfrak{l} be an integral ideal of $O = \mathbb{Z}[2\sqrt{-p}]$ where $N(\mathfrak{l}) = \ell$ is an odd prime. Put $\mathfrak{q}_1 = \mathbb{Z}q_1 + \mathbb{Z}(2r_1 + 2\sqrt{-p})$. We have an isomorphism: $\Gamma^{-1}\mathcal{O}(q_1, r_1)\mathfrak{l} \cong \mathcal{O}(q_2, r_2)$ for some q_2 satisfying (1) which is the norm of a prime ideal $\mathfrak{q}_2 \in \bar{\mathfrak{q}}_1 \ell^2 P_{\mathbb{Z}}(2)$ (resp. $\mathfrak{q}_2 \in \bar{\mathfrak{q}}_1 \ell^2 P_{\mathbb{Z}}(4)$) if $p \equiv 1 \pmod{4}$ (resp. $p \equiv 3 \pmod{4}$).

Proof. We have $\Gamma^{-1}\mathcal{O}(q_1, r_1)\mathfrak{l} \subset (O/2) + \mathfrak{q}_1 \bar{\ell}^2 \beta_1 / 4q_1 \ell$. We can take an isomorphism $\Gamma^{-1}\mathcal{O}(q_1, r_1)\mathfrak{l} \cong \mathcal{O}(q_2, r_2)$ for some q_2 satisfying (1) such that the image of O is O . Let β_2 be the image of β_{q_2} of this isomorphism. So we have $\beta_2 \in \mathfrak{q}_1 \bar{\ell}^2 \beta_1 / 4q_1 \ell$. Put $\kappa = \ell q_1 \beta_2 \beta_1^{-1}$. Then $N(2\kappa) = 4\ell^2 q_1 q_2 \in \mathbb{Z}$ and $\text{tr}(2\kappa) \in \mathbb{Z}$, since $2\kappa \in \mathfrak{q}_1 \bar{\ell}^2 / 2 \subset O/2$. It follows that $2\kappa \in O_K$.

If $p \equiv 1 \pmod{4}$, we can write $2\kappa = a + b\sqrt{-p}$ with $a, b \in \mathbb{Z}$, and we have $N(2\kappa) = a^2 + pb^2$. Since $4 \mid N(2\kappa)$ and $p \equiv 1 \pmod{4}$, we must have $2 \mid a$ and $2 \mid b$. This means $\kappa \in O_K$.

If $p \equiv 3 \pmod{4}$, then $2\kappa \in O/2 \cap O_K = \mathbb{Z} + \mathbb{Z}\sqrt{-p}$. Write $2\kappa = a + b\frac{1+\sqrt{-p}}{2}$ with $2 \mid b$, so $N(2\kappa) = a^2 + ab + \frac{1+p}{4}b^2 \equiv a^2 + ab \pmod{4}$. We have $2 \mid a$ since $4 \mid N(2\kappa)$. It follows that $\kappa \in O_K$.

Moreover, we have

$$(1 + \beta_2) / 2 \in \Gamma^{-1}\mathcal{O}(q_1, r_1)\mathfrak{l}, \quad \text{so} \quad (\ell q_1 + \kappa \beta_1) / 2q_1 \in \ell \Gamma^{-1}\mathcal{O}(q_1, r_1)\mathfrak{l} \subset \mathcal{O}(q_1, r_1).$$

If $p \equiv 1 \pmod{4}$, putting $\kappa = c + d\sqrt{-p}$, we have

$$(\ell q_1 + \kappa \beta_1) / 2q_1 = (\ell q_1 - c + dr) / 2q_1 + (c - dr)(1 + \beta_1) / 2q_1 + d(r + \sqrt{-p})\beta_1 / 2q_1.$$

So we have $2 \mid d$ and $(\kappa) \in P_{\mathbb{Z}}(2)$. Furthermore, since $N(\kappa) = \ell^2 q_1 q_2$ and $\mathfrak{q}_1 \bar{\ell}^2 \mid (\kappa)$, there is an integral ideal \mathfrak{q}_2 such that $(\kappa) = \mathfrak{q}_1 \mathfrak{q}_2 \bar{\ell}^2$ and $N(\mathfrak{q}_2) = q_2$.

If $p \equiv 3 \pmod{4}$, putting $\kappa = c + d\frac{1+\sqrt{-p}}{2}$, we have

$$(\ell q_1 + \kappa \beta_1) / 2q_1 = (2\ell q_1 - 2c - d + dr) / 4q_1 + (2c + d - dr)(1 + \beta_1) / 4q_1 + d(r + \sqrt{-p})\beta_1 / 4q_1.$$

So we have $4 \mid d$ and $(\kappa) \in P_{\mathbb{Z}}(4)$. Similarly, there is an integral ideal \mathfrak{q}_2 such that $(\kappa) = \mathfrak{q}_1 \mathfrak{q}_2 \bar{\ell}^2$ and $N(\mathfrak{q}_2) = q_2$. \square

For $p \equiv 1 \pmod{4}$, if $\phi : E_1 \rightarrow E_2$ is a 2-isogeny over \mathbb{F}_p with $\text{End}_{\mathbb{F}_p}(E_1) \cong \text{End}_{\mathbb{F}_p}(E_2) \cong \mathbb{Z}[\sqrt{-p}]$, then the isogeny ϕ corresponds to the ideal $[2, \sqrt{-p}-1]$. Because the ideals $[2, \sqrt{-p}-1]$ and $[\frac{p+1}{2}, \sqrt{-p}-1]$ are in the same ideal class in $C(\mathbb{Z}[\sqrt{-p}])$, there exists an isogeny $\psi : E_1 \rightarrow E_2$ defined over \mathbb{F}_p with kernel $\text{Ker}(\psi) = \{P \in E_1[\frac{p+1}{2}] | [\sqrt{-p}-1]P = \infty\}$. Note that $\text{Ker}(\psi) = \{P \in E_1[\frac{p+1}{2}] | [2\sqrt{-p}-2]P = \infty\}$ since $\frac{1+p}{2}$ is odd. Moreover, the kernel ideal of ψ is $[\frac{p+1}{2}, 2\sqrt{-p}-2]$, which corresponds to the form $(\frac{p+1}{2}, 4, 8) \sim (8, -4, \frac{p+1}{2})$.

Proposition 4.5. *Let $p \equiv 1 \pmod{4}$ and $O = \mathbb{Z}[2\sqrt{-p}]$. The ideal $\mathfrak{l} = [8, 2+2\sqrt{-p}]$ is proper in O . Put $\mathfrak{q}_1 = \mathbb{Z}q_1 + \mathbb{Z}(2r_1 + 2\sqrt{-p})$. We have an isomorphism: $\mathfrak{l}^{-1}\mathcal{O}(q_1, r_1)\mathfrak{l} \cong \mathcal{O}(q_2, r_2)$ for some q_2 satisfying (1) which is the norm of a prime ideal $\mathfrak{q}_2 \in \bar{\mathfrak{q}}_1 \mathfrak{l}^2 P_{\mathbb{Z}}(2)$.*

Proof. Similar to the proof of Proposition 4.4. □

If ℓ is an odd prime, then $\mathfrak{l} = [\ell, -b + 2\sqrt{-p}]$ with $b^2 \equiv -4p \pmod{\ell}$. If $\ell = 2$, then $\mathfrak{l} = [8, 2 + 2\sqrt{-p}]$. In this case, $\bar{\mathfrak{l}} = [8, -2 + 2\sqrt{-p}]$ and $\mathfrak{l}^2 = \bar{\mathfrak{l}}^2 = [4, 2\sqrt{-p}]$.

Theorem 5. *Let E_1 be a supersingular elliptic curve with $\text{End}(E_1) \cong \mathcal{O}(q_1, r_1)$. If $\phi : E_1 \rightarrow E_2$ is an isogeny defined over \mathbb{F}_p of degree ℓ with $\text{End}_{\mathbb{F}_p}(E_2) \cong \mathbb{Z}[\sqrt{-p}]$, then there is a prime q_2 satisfying (1) which can be represented by the form $(q_1, r_1, \frac{4r_1^2+4p}{q_1})(\ell, 2b, \frac{b^2+p}{\ell})^2$ such that $\text{End}(E_2) \cong \mathcal{O}(q_2, r_2)$.*

Proof. Similar to the proof of Theorem 4. □

Case 3: $\text{End}_{\mathbb{F}_p}(E_1) \not\cong \text{End}_{\mathbb{F}_p}(E_2)$.

Considering the structure of the isogeny graphs of supersingular elliptic curves over \mathbb{F}_p , we just need to study the 2-isogenies defined over \mathbb{F}_p in this case.

Let $p \equiv 3 \pmod{4}$ be a prime and q a prime satisfying (1). As we know, $r^2 + p \equiv 0 \pmod{4q}$ implies $r^2 + p \equiv 0 \pmod{q}$. On the contrary, if the number r satisfies the equation $x^2 + p \equiv 0 \pmod{q}$, then r or $r+q$ satisfies the equation $x^2 + p \equiv 0 \pmod{4q}$. It follows that $\mathcal{O}(q, r)$ and $\mathcal{O}'(q, r)$ are maximal orders in $B_{p, \infty}$ for the same r and q . The following theorem tells us the relations between $\mathcal{O}(q, r)$ and $\mathcal{O}'(q, r)$.

Theorem 6. *Let $p \equiv 3 \pmod{4}$ be a prime and E_2 a supersingular elliptic curve over \mathbb{F}_p with $j(E_2) \neq 1728$. If the endomorphism ring of E_2 is isomorphic to $\mathcal{O}(q, r)$, then there exists a 2-isogeny $\phi : E_1 \rightarrow E_2$ such that the endomorphism ring of E_1 is $\mathcal{O}'(q, r)$.*

Proof. If $j(E_2) \neq 1728$, for the same prime q , we consider the order

$$\tilde{\mathcal{O}}(q, r) = \mathbb{Z} + \mathbb{Z}\beta + \mathbb{Z}\frac{1 + \alpha + \beta + \alpha\beta}{2} + \mathbb{Z}\frac{(r + \alpha)\beta}{q}.$$

It is easy to show $\tilde{\mathcal{O}}(q, r) \subseteq \mathcal{O}'(q, r) \cap \mathcal{O}(q, r)$, and the discriminant of $\tilde{\mathcal{O}}(q, r)$ is $4p^2$. Since $j(E_2) \neq 1728$, we have $\mathcal{O}'(q, r') \not\cong \mathcal{O}(q, r)$ by Proposition 2.4. It follows that

$[\mathcal{O}'(q, r') : \tilde{\mathcal{O}}(q)] = [\mathcal{O}(q, r) : \tilde{\mathcal{O}}(q)] = 2$ and $\tilde{\mathcal{O}}(q, r) = \mathcal{O}'(q, r) \cap \mathcal{O}(q, r)$. Moreover, there exists a 2-isogeny $\phi : E_1 \rightarrow E_2$ with $\text{End}(E_1) \cong \mathcal{O}'(q, r)$. □


Remark 4.3. *If $j(E_2) = 1728$, we can assume $E_2 : y^2 = x^3 + 4x$. There exists a 2-isogeny $\phi : E_1 \rightarrow E_2$ with $E_1 : y^2 = x^3 - x$ and $\phi(x, y) = (\frac{x^2-1}{x}, \frac{x^2y+y}{x^2})$. Moreover, we have $j(E_1) = 1728$ and $\text{End}(E_1) \cong \text{End}(E_2)$.*

In fact, the curve E_1 is on the surface and E_2 is on the floor. If $p \equiv 7 \pmod{8}$, there is only one down 2-isogeny from E_1 which is $\phi : E_1 \rightarrow E_2$. If E_1 corresponds to the form $(q, r, \frac{r^2+p}{4q})$ with q satisfying (1), then E_2 corresponds to the form $(q, 4r, \frac{4r^2+4p}{q})$.


If $p \equiv 3 \pmod{8}$, there are three down 2-isogenies from E_1 . Denote them by $E_{2,i}$ with $i \in \{1, 2, 3\}$. If E_1 corresponds to the form $(q, r, \frac{r^2+p}{4q})$, by Proposition 4.3, there are three forms with discriminant $-16p$ that can be derived from $(q, r, \frac{r^2+p}{4q})$. Moreover, every form corresponds to $E_{2,i}$. On the contrary, suppose that the curve $E_{2,i}$ corresponds to the form $(q_i, 4r_i, \frac{4r_i^2+4p}{q_i})$ with q_i satisfying (1). For every i , the form $(q_i, 4r_i, \frac{4r_i^2+4p}{q_i})$ can be derived from $(q_i, r_i, \frac{r_i^2+p}{4q_i})$. It follows that the three forms $(q_i, r_i, \frac{r_i^2+p}{4q_i})$, $i \in \{1, 2, 3\}$ are equivalent and the curve E_1 corresponds to every one of these three forms.

We use the curves in Example 1 to illustrate results in this section.

Example 2. The notations are as Example 1. The followings are 2-isogeny graphs of supersingular elliptic curves over \mathbb{F}_{83} .




Considering the corresponding forms, we have the following graphs.




Note that the forms g_0, g_4 and g_4^{-1} are derived from f_0 , and the other two graphs are similar. These examples are coincident with Theorem 6. In order to illustrate Theorems


4 and 5, we compute the 3-isogeny graphs of supersingular elliptic curves over \mathbb{F}_{83} .



Considering the corresponding forms, we have the following graphs.



Every arrow means composition with the form $(3, 1, 7)^2$.



Every arrow means composition with the form $(3, 2, 111)^2$.

5. Possibility in CSIDH

In this section, we recall the encipherment of CSIDH cryptosystem, and apply our results to analyze the security of CSIDH.

Let $k \in \mathbb{Z}_+$ and consider a prime p of the form $p = 4\ell_1\ell_2 \dots \ell_k - 1$, where the ℓ_i 's are distinct odd prime numbers. This implies $p \equiv 3 \pmod{8}$. CSIDH chooses the curve in the set $\text{Iso}(\mathbb{F}_p, \pi)$, which contains all the \mathbb{F}_p -isomorphism classes of supersingular elliptic curves over \mathbb{F}_p with endomorphism ring $\mathcal{O}(q, r)$.

The starting curve of CSIDH is $E_0 : y^2 = x^3 + x$, which corresponds to the form $(4, 0, p)$. Alice and Bob secretly sample $[\mathbf{a}]$ resp. $[\mathbf{b}]$ from the class group of $\mathbb{Z}[\sqrt{-p}]$, compute $E_A = [\mathbf{a}]E_0$ resp. $E_B = [\mathbf{b}]E_0$, and publish the result. The shared secret is

then $E_{AB} = [\mathbf{a}][\mathbf{b}]E_0$, which Alice computes as $[\mathbf{a}](([\mathbf{b}]E_0)$ and while Bob computes as $[\mathbf{b}](([\mathbf{a}]E_0)$.

In fact, if the curve E_A (resp. E_B) corresponds to the form (a_1, b_1, c_1) (resp. (a_2, b_2, c_2)), then the shared secret curve E_{AB} corresponds to the form $(4, 0, p)(a_1, b_1, c_1)(a_2, b_2, c_2)$.

Given a supersingular elliptic curve E_A over \mathbb{F}_p , if we know its endomorphism ring $\mathcal{O}(q_1, r_1)$, then it corresponds to the form $(q_1, 4r_1, \frac{4r_1^2+4p}{q_1})$ or $(q_1, -4r_1, \frac{4r_1^2+4p}{q_1})$ by Theorem 2. Similarly, the curve E_B corresponds to the form $(q_2, 4r_2, \frac{4r_2^2+4p}{q_2})$ or $(q_2, -4r_2, \frac{4r_2^2+4p}{q_2})$. In general, we compute 4 forms and one of them corresponds to the curve E_{AB} .

On the other hand, given a reduced form (a, b, c) with discriminant $-16p$ in the non-principal genus, the j -invariant is the unique common root of the Hilbert class polynomials $H_{-a}(X)$ and $H_{-c}(X)$. Sutherland [12] presented a space-efficient algorithm to compute $H_D(X)$ modulo a positive integer p , based on an explicit form of the Chinese Remainder Theorem. Under the Generalized Riemann Hypothesis, the algorithm uses $O(|D|^{1/2+\epsilon} \log p)$ space and has an expected running time of $O(|D|^{1+\epsilon})$. Moreover, in order to compute the j -invariant, we also need to compute the greatest common divisor of $H_{-a}(X) \bmod p$ and $H_{-c}(X) \bmod p$.

In fact, in order to get the corresponding form of a supersingular elliptic curve E defined over \mathbb{F}_p , we just need to know two imaginary quadratic orders O_1 and O_2 with discriminant d_1 and d_2 respectively which can be embedded in the endomorphism ring $\text{End}(E)$, where $|d_i| < p$ and $O_1 \cap O_2 = \mathbb{Z}$.

As we discussed above, if one can compute the correspondence between supersingular elliptic curves over \mathbb{F}_p and binary quadratic forms explicitly, then it is possible to get the shared secret of CSIDH. Compared with the method in [4], we do not need to solve the discrete logarithm problem in the class group of $\mathbb{Z}[\sqrt{-4p}]$, which is in general quite difficult.

6. Conclusion

For a supersingular elliptic curve defined over \mathbb{F}_p , Ibukiyama's results show that its endomorphism ring is isomorphic to one of the maximal orders $\mathcal{O}(q, r)$'s and $\mathcal{O}'(q, r')$'s. We show that a quadratic form in the maximal order plays a crucial role, and the operations of isogenies on elliptic curves defined over \mathbb{F}_p is compatible with the composition of quadratic forms. The advantage of this correspondence is that the composition of quadratic forms is commutative, while the multiplication in maximal orders is not. Computing this correspondence explicitly and applying these results to the isogeny-based cryptography are our future work.

References

- [1] Johannes Buchmann and Ulrich Vollmer, *Binary quadratic forms. an algorithmic approach*, Algorithms and Computation in Mathematics, vol. 20, Springer, Berlin, 2007.

- [2] Duncan A. Buell, *Binary quadratic forms, classical theory and modern computations*, Springer-Verlag, New York, 1989.
- [3] Wouter Castryck, Tanja Lange, Chloe Martindale, Lorenz Panny, and Joost Renes, *CSIDH: an efficient post-quantum commutative group action*, Advances in cryptology—ASIACRYPT 2018. Part III, Lecture Notes in Comput. Sci., vol. 11274, Springer, Cham, 2018, pp. 395–427.
- [4] Wouter Castryck, Lorenz Panny, and Frederik Vercauteren, *Rational isogenies from irrational endomorphisms*, Advances in cryptology—EUROCRYPT 2020. Part II, Lecture Notes in Comput. Sci., vol. 12106, Springer, Cham, [2020] ©2020, pp. 523–548.
- [5] David A. Cox, *Primes of the form $x^2 + ny^2$. fermat, class field theory, and complex multiplication*, second ed., Pure and Applied Mathematics (Hoboken), John Wiley & Sons, Inc., Hoboken, NJ, 2013.
- [6] Christina Delfs and Steven D. Galbraith, *Computing isogenies between supersingular elliptic curves over \mathbb{F}_p* , Des. Codes Cryptogr. **78** (2016), no. 2, 425–440.
- [7] Max Deuring, *Die Typen der Multiplikatorenringe elliptischer Funktionenkörper*, Abh. Math. Sem. Hansischen Univ. **14** (1941), 197–272.
- [8] Tomoyoshi Ibukiyama, *On maximal orders of division quaternion algebras over the rational number field with certain optimal embeddings*, Nagoya Math. J. **88** (1982), 181–195.
- [9] Masanobu Kaneko, *Supersingular j -invariants as singular moduli mod p* , Osaka J. Math. **26** (1989), no. 4, 849–855.
- [10] Kristin Lauter and Bianca Viray, *On singular moduli for arbitrary discriminants*, Int. Math. Res. Not. IMRN (2015), no. 19, 9206–9250.
- [11] Joseph H. Silverman, *The arithmetic of elliptic curves*, second ed., Graduate Texts in Mathematics, vol. 106, Springer, Dordrecht, 2009.
- [12] Andrew V. Sutherland, *Computing Hilbert class polynomials with the Chinese remainder theorem*, Math. Comp. **80** (2011), no. 273, 501–538.
- [13] John Voight, *Quaternion algebras*, Graduate Texts in Mathematics, vol. 288, Springer, Cham, [2021] ©2021.
- [14] Guanju Xiao, Lixia Luo, and Yingpu Deng, *Supersingular j -invariants and the class number of $\mathbb{Q}(\sqrt{-p})$* , International Journal of Number Theory. Doi:10.1142/S1793042122500555.