

# Analytic expressions for some Mellin transforms with their application to prime counting function and interpolation formulas for the zeta function

Omprakash Atale\*

February 26, 2022

## Abstract

The aim of our present work here is to present few results in the theory of Mellin transforms using the method that S. Ramanujan used in proving his Master Theorem. Further applications of our results for some number-theoretic functions such as the prime counting function and the zeta function are established.

## 1 Introduction

**1.1.** This is a technical paper which is an application based extension of Ramanujan's Master Theorem which is a powerful tool for evaluating Mellin type integrals [1129, pg. [5]]. It states that if  $f$  has expansion of the form

$$f(x) = \sum_{n=0}^{\infty} (-1)^n \frac{\phi(n)}{n!} x^n \quad (1.1)$$

where  $\phi(n)$  has a natural and continuous extension such that  $\phi(0) \neq 0$ , then for  $s > 0$ , we have

$$\int_0^{\infty} x^{s-1} \left( \sum_{n=0}^{\infty} (-1)^n \frac{\phi(n)}{n!} x^n \right) dx = \Gamma(s) \phi(-s). \quad (1.2)$$

where  $s$  is any positive integer. Eqn. (1.2) was communicated by Ramanujan in his *Quarterly Reports* [[1], p.298][2] and was used by him in computing the values

---

\*Khandesh College Education Society's Moolji Jaitha College Jalgaon-425001, Maharashtra, India. E-mail: atale.om@outlook.com

of certain definite integrals [3]. We kindly request readers to make themselves familiar with the derivation of Ramanujan's Master Theorem from [[1], p.298][2] whose method of proof is frequently used throughout the paper. Now, for the purpose of application, consider the following binomial expansion for  $a, v > 0$

$$(1 + ax)^{-v} = \sum_{n=0}^{\infty} a^n \frac{\Gamma(v+n)}{\Gamma(v)} \frac{(-x)^n}{n!}. \quad (1.6)$$

Employing Eqn. (1.2) yields

$$\int_0^{\infty} x^{n-1} (1 + ax)^{-v} dx = \frac{\Gamma(n) \Gamma(v-n)}{a^n \Gamma(v)}. \quad (1.7)$$

Further applications and examples of Eqn. (1.2) can be found in [2][4][8].

## 2 On certain Mellin transforms and their analytic expressions

**2.1.** In this section, few theorems are established that are motivated by Ramanujan's method of deriving Eqn. (1.2). Furthermore, certain applications of respective theorems are studied and applied in calculating the Mellin transform of certain infinite series. Throughout this section, it is assumed that  $\phi(n)$  has a natural and continuous extension such that  $\phi(0) \neq 0$ .

**Theorem 2.1.** *If  $0 < \Re(s) < 1$ , then*

(i)

$$\int_0^{\infty} x^{s-1} \sum_{n=0}^{\infty} \phi(2n+1) \frac{(-1)^n}{(2n+1)!} x^{2n+1} dx = \phi(-s) \Gamma(s) \sin \frac{\pi s}{2}, \quad (2.1)$$

(ii)

$$\int_0^{\infty} x^{s-1} \sum_{n=0}^{\infty} \phi(2n) \frac{(-1)^n}{(2n)!} x^{2n} dx = \phi(-s) \Gamma(s) \cos \frac{\pi s}{2}, \quad (2.2)$$

*Proof.* Consider the following Mellin transform of  $\sin(ax)$  [[7], pg. 332]

$$\int_0^{\infty} x^{s-1} \sin(ax) dx = a^{-s} \Gamma(s) \sin \left( \frac{\pi s}{2} \right). \quad (2.3)$$

Substituting  $a = r^k$  with  $r > 0$  in the above equation and expand  $\sin(ax)$  in its

Maclaurin series to get

$$\int_0^{\infty} x^{s-1} \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} (r^k x)^{2n+1} dx = \Gamma(s) r^{-sk} \sin\left(\frac{\pi s}{2}\right).$$

Multiply both sides by  $\frac{f^{(k)}(a)h^k}{k!}$  where  $f$  shall be specified later and sum on  $k$ ,  $0 \leq k < \infty$  to get

$$\begin{aligned} & \sum_{k=0}^{\infty} \frac{f^{(k)}(a)h^k}{k!} \int_0^{\infty} x^{s-1} \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} (r^k x)^{2n+1} dx \\ &= \sum_{k=0}^{\infty} \frac{f^{(k)}(a)(hr^{-s})^k}{k!} \Gamma(s) \sin\left(\frac{\pi s}{2}\right), \\ & \int_0^{\infty} x^{s-1} \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{f^{(k)}(a)(hr^{2n+1})^k (-1)^n}{k! (2n+1)!} x^{2n+1} dx \\ &= \sum_{k=0}^{\infty} \frac{f^{(k)}(a)(hr^{-s})^k}{k!} \Gamma(s) \sin\left(\frac{\pi s}{2}\right). \end{aligned}$$

Now, let

$$\phi(-s) = f(hr^{-s} + a) = \sum_{k=0}^{\infty} \frac{f^{(k)}(a)(hr^{-s})^k}{k!}.$$

Therefore, after further simplification we get

$$\int_0^{\infty} x^{s-1} \sum_{n=0}^{\infty} \phi(2n+1) \frac{(-1)^n}{(2n+1)!} x^{2n+1} dx = \phi(-s) \Gamma(s) \sin\left(\frac{\pi s}{2}\right).$$

Proof of (ii) can be obtained by a similar method using [[6], pg. 332]

$$\int_0^{\infty} x^{s-1} \cos(ax) dx = \Gamma(s) a^{-s} \cos\left(\frac{\pi s}{2}\right). \quad (2.4)$$

■

**Corollary 2.1.1.** For  $0 < \Re(s) < 1$  and  $s \neq \frac{1}{2}, \frac{1}{4}$  we have

(i)

$$\int_0^{\infty} x^{s-1} \left( \zeta(4)x - \zeta(8) \frac{1}{3!}x^3 + \zeta(12) \frac{1}{5!}x^5 + \dots \right) dx = \zeta(2-2s) \Gamma(s) \sin\left(\frac{\pi s}{2}\right), \quad (2.5)$$

(ii)

$$\int_0^{\infty} x^{s-1} \left( \zeta(2) - \zeta(6) \frac{1}{2!}x^2 + \zeta(10) \frac{1}{4!}x^4 + \dots \right) dx = \zeta(2-2s) \Gamma(s) \cos\left(\frac{\pi s}{2}\right), \quad (2.6)$$

where  $\zeta(s)$  is the zeta function.

*Proof.* Let

$$\phi(n) = \zeta(2n+2)$$

which yields

$$\phi(2n+1) = \zeta(4n+4)$$

and

$$\phi(2n) = \zeta(4n+2).$$

Now, use Eqn. (2.1) and (2.2) to get the desired result. ■

**Corollary 2.1.2.** *We have*

(i)

$$\int_0^{\infty} \zeta(4) - \zeta(8) \frac{1}{3!}x^2 + \zeta(12) \frac{1}{5!}x^4 + \dots dx = \frac{\pi^3}{12}, \quad (2.8)$$

(ii)

$$\int_0^{\infty} \frac{\log x}{x} \left( \zeta(2) - \zeta(6) \frac{1}{2!}x^2 + \zeta(10) \frac{1}{4!}x^4 + \dots \right) dx = \frac{\pi^4}{24}, \quad (2.9)$$

where  $\zeta(s)$  is the zeta function.

*Proof.* We have mentioned earlier that  $0 < \Re(s) < 1$ , but there is a particular case where we can apply Theorem 2.1 at  $s = 0$ . This can be done as follows. Using reflection formula for the gamma function of the right hand side of Eqn. (2.5) and taking the limit on  $s$  the both sides to zero, we get

$$\int_0^{\infty} \zeta(4) - \zeta(8) \frac{1}{3!}x^2 + \zeta(12) \frac{1}{5!}x^4 + \dots dx = \frac{\pi^2}{6} \lim_{s \rightarrow 0} \frac{\pi \sin\left(\frac{\pi s}{2}\right)}{\Gamma(1-s) \sin(\pi s)}.$$

Using L'Hospital's rule, we get

$$\int_0^\infty \zeta(4) - \zeta(8) \frac{1}{3!} x^2 + \zeta(12) \frac{1}{5!} x^4 + \dots dx = \frac{\pi^3}{12}.$$

Similarly, after calculating the value of integral (2.6), Eqn. (2.9) readily follows. A detailed proof can be found in Appendix A.1.  $\blacksquare$

**Corollary 2.1.3.** *for  $|t| < |a|$ ,  $0 < \Re(s) < 1$  and  $c > 0$  We have*

$$\begin{aligned} & \frac{1}{2} \int_0^\infty x^{s-1} [\zeta(c, a+x) - \zeta(c, a-x)] dx \\ &= \frac{(-1)^{-s} \Gamma(s) \Gamma(c-s) \zeta(c-s, a)}{\Gamma(c)} \sin\left(\frac{\pi s}{2}\right). \end{aligned} \quad (2.10)$$

*Proof.* From [[9], pg. 412] we have

$$\sum_{k=0}^\infty \frac{(c)_{2k+1}}{(2k+1)!} \zeta(c+2k+1, a) t^{2k+1} = \frac{1}{2} [\zeta(c, a-t) - \zeta(c, a+t)] \quad (2.11)$$

where

$$(c)_{2k+1} = \frac{\Gamma(c+2k+1)}{\Gamma(c)}.$$

Letting  $t = x$  and applying Eqn. (2.1) yields the desire result.  $\blacksquare$

**Theorem 2.2.** *We have*

$$\int_0^\infty \frac{\log x}{x} \sum_{n=0}^\infty c_n x^{2n} d\pi(x) = \sum_{n=0}^\infty c_n A_n \quad (2.12)$$

where

$$c_n = \zeta(4n+2) \frac{(-1)^n}{(2n)!}, \quad (2.13)$$

$$A_n = \sum_{k=1}^\infty \frac{\mu(k)}{(1-2n)k-1} + f(1-2n), \quad (2.14)$$

and  $\pi(x)$  is the prime counting function.

*Proof.* Compress the sum in the integrand of Eqn. (2.8) and instead of integrating from all values from 0 to  $\infty$ , integrate only on primes [[7], pg. 118, Eqn. (9.2)],

that is

$$\int_0^\infty \frac{\log x}{x} \sum_{n=0}^\infty \zeta(4n+2) \frac{(-1)^n}{(2n)!} x^{2n} d\pi(x)$$

where  $\pi(x)$  is the prime counting function. Therefore, we get

$$\int_0^\infty \frac{\log x}{x} \sum_{n=0}^\infty \zeta(4n+2) \frac{(-1)^n}{(2n)!} x^{2n} d\pi(x) = \sum_p \frac{\log p}{p} \sum_{n=0}^\infty \zeta(4n+2) \frac{(-1)^n}{(2n)!} p^{2n}.$$

Now, let

$$c_n = \zeta(4n+2) \frac{(-1)^n}{(2n)!}$$

then substitute the value of  $c_n$  and inverting the order of summation, we get

$$\int_0^\infty \frac{\log x}{x} \sum_{n=0}^\infty c_n x^{2n} d\pi(x) = \sum_{n=0}^\infty c_n \sum_p \frac{\log p}{p} p^{2n} = \sum_{n=0}^\infty c_n \sum_p \frac{\log p}{p^{1-2n}}. \quad (2.15)$$

Now, using Eqn. (6.1) from [7], pg. 116], for  $\Re(s) > 1$ , we have

$$\sum_p \frac{\log p}{p^s} = \sum_{k=1}^\infty \frac{\mu(k)}{sk-1} + f(s)$$

where  $\mu(k)$  is the Mobius function,  $f(k)$  is analytic and is given by

$$f(s) = - \sum_{k=1}^\infty \mu(k) \left\{ \frac{\zeta'(ks)}{\zeta(ks)} + \frac{1}{ks-1} \right\}.$$

Therefore, we get

$$\sum_p \frac{\log p}{p^{1-2n}} = \sum_{k=1}^\infty \frac{\mu(k)}{(1-2n)k-1} + f(1-2n)$$

and

$$f(1-2n) = - \sum_{k=1}^\infty \mu(k) \left\{ \frac{\zeta'(k(1-2n))}{\zeta(k(1-2n))} + \frac{1}{k(1-2n)-1} \right\}.$$

Substituting the above values in Eqn. (2.15) yields the following result

$$\int_0^\infty \frac{\log x}{x} \sum_{n=0}^\infty c_n x^{2n} d\pi(x) = \sum_{n=0}^\infty c_n A_n$$

where

$$A_n = \sum_{k=1}^{\infty} \frac{\mu(k)}{(1-2n)k-1} + f(1-2n).$$

■

**Theorem 2.3.** *If  $p, k, s > 0$ , then*

(i)

$$\int_0^{\infty} x^{s-1} \sum_{n=0}^{\infty} \phi(2n+1) \frac{(-1)^n}{(2n+1)!p^{2n+1}} x^{(2n+1)k} dx = \phi\left(\frac{-s}{k}\right) {}_p\Gamma_k(s) \sin\left(\frac{\pi s}{2k}\right), \quad (2.16)$$

(ii)

$$\int_0^{\infty} x^{s-1} \sum_{n=0}^{\infty} \phi(2n+1) \frac{(-1)^n}{(2n+1)!k^{2n+1}} x^{(2n+1)k} dx = \phi\left(\frac{-s}{k}\right) \Gamma_k(s) \sin\left(\frac{\pi s}{2k}\right), \quad (2.17)$$

(iii)

$$\int_0^{\infty} x^{s-1} \sum_{n=0}^{\infty} \phi(2n) \frac{(-1)^n}{(2n)!p^{2n}} x^{(2n)k} dx = \phi\left(\frac{-s}{k}\right) {}_p\Gamma_k(s) \cos\left(\frac{\pi s}{2k}\right), \quad (2.19)$$

(iv)

$$\int_0^{\infty} x^{s-1} \sum_{n=0}^{\infty} \phi(2n) \frac{(-1)^n}{(2n)!k^{2n}} x^{(2n)k} dx = \phi\left(\frac{-s}{k}\right) \Gamma_k(s) \cos\left(\frac{\pi s}{2k}\right), \quad (2.20)$$

where  ${}_p\Gamma_k(s)$  is the  $p$ - $k$  gamma function [10] and  $\Gamma_k(s)$  is the  $k$  gamma function [11]<sup>1</sup> defined as follows

$${}_p\Gamma_k(s) = \left(\frac{p}{k}\right)^{\frac{s}{k}} \Gamma_k(s) = \frac{p^{\left(\frac{s}{k}\right)}}{k} \Gamma\left(\frac{s}{k}\right).$$

*Proof.* Replace  $x$  with  $x^k/p$  in Eqn. (2.16) to get

$$\int_0^{\infty} \frac{x^{sk-k}}{p^{s-1}} \sum_{n=0}^{\infty} \phi(2n+1) \frac{(-1)^n}{(2n+1)!p^{2n+1}} x^{(2n+1)k} \frac{kx^{k-1}}{p} dx = \phi(-s) \Gamma(s) \sin\left(\frac{\pi s}{2}\right),$$

---

<sup>1</sup> ${}_p\Gamma_k(x) \Rightarrow {}_k\Gamma_k(x) = \Gamma_k(x)$  as  $p = k$  and  ${}_p\Gamma_k(x) \Rightarrow {}_1\Gamma_1(x) = \Gamma(x)$  as  $p, k \rightarrow 1$ .

$$\int_0^\infty \frac{x^{sk-1}}{p^s} \sum_{n=0}^\infty \phi(2n+1) \frac{(-1)^n}{(2n+1)!p^{2n+1}} x^{(2n+1)k} dx = \phi(-s) \Gamma(s) \sin\left(\frac{\pi s}{2}\right).$$

Now, replacing  $s$  with  $s/k$  yields

$$\int_0^\infty x^{s-1} \sum_{n=0}^\infty \phi(2n+1) \frac{(-1)^n}{(2n+1)!p^{2n+1}} x^{(2n+1)k} dx = \phi\left(\frac{-s}{k}\right) \frac{p^{\frac{s}{k}}}{k} \Gamma\left(\frac{s}{k}\right) \sin\left(\frac{\pi s}{2k}\right).$$

By further simplification, the desired result readily follows. (iii) can be derived in a similar manner. (ii) and (iv) are special cases of (i) and (iii) when  $p = k$  respectively.  $\blacksquare$

### 3 Appendix

**A.1.** Take the derivatives both the sides of Eqn. (2.6) with respect to  $s$  and then multiply and divide right hand side of the equation with  $\Gamma(s)$  to get

$$\begin{aligned} \int_0^\infty x^{s-1} \log x \left( \zeta(2) - \zeta(6) \frac{1}{2!} x^2 + \zeta(10) \frac{1}{4!} x^4 \dots \right) dx &= \zeta'(2-2s) \Gamma(s) \cos\left(\frac{\pi s}{2}\right) \\ &\quad - \frac{1}{\Gamma(s)} \zeta(2-4s) \psi(s) \cos\left(\frac{\pi s}{2}\right) + \frac{\pi}{2} \zeta(2-2s) \Gamma(s) \sin\left(\frac{\pi s}{2}\right). \end{aligned}$$

Using the reflection formula for gamma function, we get

$$\begin{aligned} \int_0^\infty x^{s-1} \log x \left( \zeta(2) - \zeta(6) \frac{1}{2!} x^2 + \zeta(10) \frac{1}{4!} x^4 \dots \right) dx &= \zeta'(2-2s) \frac{\pi \cos\left(\frac{\pi s}{2}\right)}{\Gamma(1-s) \sin \pi s} \\ &\quad + \frac{1}{\pi} \zeta(2-2s) \Gamma(1-s) \sin(\pi s) \psi(s) \cos\left(\frac{\pi s}{2}\right) + \frac{\pi}{2} \zeta(2-2s) \frac{\pi \sin\left(\frac{\pi s}{2}\right)}{\Gamma(1-s) \sin \pi s}. \end{aligned}$$

Now, taking limit both the sides of  $s$  from  $s \rightarrow 0$ , and applying L' Hospital's rule, we get

$$\begin{aligned} &\int_0^\infty \frac{\log x}{x} \left( \zeta(2) - \zeta(6) \frac{1}{2!} x^2 + \zeta(10) \frac{1}{4!} x^4 \dots \right) dx \\ &= \lim_{s \rightarrow 0} \zeta'(2-2s) \frac{\pi \cos\left(\frac{\pi s}{2}\right)}{\Gamma(1-s) \sin \pi s} + \lim_{s \rightarrow 0} \frac{\pi}{2} \zeta(2-2s) \frac{\pi \sin\left(\frac{\pi s}{2}\right)}{\Gamma(1-s) \sin \pi s}, \\ &= \zeta'(2) \lim_{s \rightarrow 0} \frac{\pi \cos\left(\frac{\pi s}{2}\right)}{\Gamma(1-s) \sin \pi s} + \frac{\pi^2}{4} \zeta(2) \lim_{s \rightarrow 0} \frac{\cos\left(\frac{\pi s}{2}\right)}{\cos \pi s} \end{aligned}$$



$$= 0 + \frac{\pi^2}{2} \zeta(2) = \frac{\pi^4}{24}.$$

Therefore,

$$\int_0^{\infty} \frac{\log x}{x} \left( \zeta(2) - \zeta(6) \frac{1}{2!} x^2 + \zeta(10) \frac{1}{4!} x^4 \dots \right) dx = \frac{\pi^4}{24}.$$

## References

- [1] B. Berndt, Ramanujan's Notebooks, Part I, *Springer-Verlag, New York*, (1985).
- [2] Berndt, B.C. The quarterly reports of S. Ramanujan. *Amer. Math. Monthly* 90(8), 505–516 (1983).
- [3] G. H. Hardy. Ramanujan. Twelve Lectures on subjects suggested by his life and work. *Chelsea Publishing Company, New York, N.Y.*, 3rd edition, (1978).
- [4] Qureshi, M. I., Quraishi, K. A. and Pal, R. Some Applications of Celebrated Master Theorem of Ramanujan. *British Journal of Mathematics and Computer Science*, 4(20), 2862-287, (2014).
- [5] I.S. Gradshteyn and I.M. Ryzhik, eds., Table of Integrals, Series, and Products, 5th Edition., *Academic Press, San Diego*, (1994).
- [6] George E. Andrews and Bruce C. Berndt. Ramanujan's Lost Notebook - Part IV. *Springer Publishing Company*, (2013)
- [7] B. Berndt, Ramanujan's Notebooks, Part IV, *Springer-Verlag, New York*, (1994).
- [8] Amdeberhan, T., Espinosa, O., Gonzalez, I. et al. Ramanujan's Master Theorem. *Ramanujan J* 29, 103–120 (2012).
- [9] H.M. Srivastava and Junesang Choi, Zeta and q-Zeta Functions and Associated Series and Integrals, 1st Edition, *Elsevier*, (2012).
- [10] Kuldeep Singh Gehlot, Two Parameter Gamma Function and it's Properties, [arXiv:1701.01052v1](https://arxiv.org/abs/1701.01052)(math.CA) (2017).
- [11] Diaz, R. and Pariguan, E. On hypergeometric functions and Pochhammer k-symbol. *Divulgaciones Mathematicas*, Vol. 15 No. 2, 179-192, (2007).