arXiv:2203.02494v1 [math.AT] 4 Mar 2022

DIFFERENT TYPES OF TOPOLOGICAL COMPLEXITY ON HIGHER HOMOTOPIC DISTANCE

MELİH İS AND İSMET KARACA

ABSTRACT. We first study the higher version of the relative topological complexity by using the homotopic distance. We also introduced the generalized version of the relative topological complexity of a topological pair on both the Schwarz genus and the homotopic distance. With these concepts, we give some inequalities including the topological complexity and the Lusternik-Schnirelmann category, the most important parts of the study of robot motion planning in topology. Finally, by defining the parametrised topological complexity via the homotopic distance, we present some estimates on the higher setting of this concept.

1. INTRODUCTION

Studies on determining the topological complexity TC first start with M. Farber [11]. As Farber [13] states, one of the basic methods followed in topological complexity studies is to study the different versions of the number TC and obtain a relationship between these versions and TC. This generally leads to some natural bounds for TC. For instance, if X is a path-connected topological space and Y is a subset of $X \times X$, then the relative topological complexity $TC_X(Y)$ satisfies the following inequality [13]:

$$\operatorname{TC}_X(Y) \leq \operatorname{TC}(X)$$

Along with the definition of higher topological complexity TC_n by Y. Rudyak [24], introducing the improved version of relative topological complexity also presents a natural lower bound for TC_n . In Section 3, we first focus on these higher settings of the relative topological complexity $TC_{n,X}(Y)$ via relative homotopic distance, where Y is a subset of X^n .

Let (A, B) be a pair of topological spaces with the condition $B \subseteq A$. Then the relative topological complexity of a pair TC(A, B) is defined by R. Short [26] on the notion of Schwarz genus and admits the following fact:

$$TC(A, B) = TC_A(A \times B).$$

In other words, if one rewrites TC(A, B) as $TC_{n,A}(B)$ (for n = 2), then one can define the relative higher topological complexity of a pair $TC_n(A, B)$ by using the relative homotopic distance such as the relative higher topological complexity of a space. Before such a definition on the homotopic distance, in Section 4, we first introduce $TC_n(A, B)$ in terms of the Schwarz genus.

Date: March 7, 2022.

²⁰¹⁰ Mathematics Subject Classification. 55M30, 14F35, 55R10, 14M15, 57N99.

Key words and phrases. topological complexity number, parametrised topological complexity number, lusternik-schnirelmann category, schwarz genus, higher homotopic distance.

In recent years, topological robotics studies have focused on computing the new fiber homotopy equivalent invariant parametrised topological complexity [6, 7, 14, 27]. For the Farber's topological complexity TC, one considers the motion planning problem as follows [11].

$$\pi: PX \to X \times X, \ \pi(\gamma) = (\gamma(0), \gamma(1)),$$

is a fibration, where PX contains all continuous paths on the path-connected topological space X. PX has a compact-open topology. TC(X) is the Schwarz genus of π and TC(X) = n means that the motion planning algorithm $s_j : A_j \subset X \times X \rightarrow PX$ must be continuous with the condition $\pi \circ s_j = 1_{A_j}$ for each $j \in \{1, \dots, n\}$.

In the parametrised version, there are some basic changes, but the main idea is familiar. This is to take the endpoints of the space and form a continuous motion between these endpoints. One has some extra conditions. The first is that the initial and final states are located in the same fiber of the fibration $q: E \to B$. The other one is to restrict the motion planning algorithm to the same fiber. We shall explain this idea mathematically. Let $q: E \to B$ be a fibration with a nonempty, path-connected fibre $X = q^{-1}(b)$ for any $b \in B$. Let E_B^I denote the subspace of E that consists of all paths α in E such that $q \circ \alpha$ is the constant map. $E \times_B E$ is a subset of $E \times E$ that includes the point (e_1, e_2) for which $q(e_1)$ equals $q(e_2)$, i.e., the pair (e_1, e_2) is located in the same fiber. Then we consider the fibration $\pi': E_B^I \to E \times_B E$ with $\pi'(\alpha) = (\alpha(0), \alpha(1))$. Note that the path-connected fiber of this fibration is ΩX . The parametrised topological complexity $\mathrm{TC}[q: E \to B]$ is the Schwarz genus of π' [7].

In the sense of robot motion planning problems, the parametrised topological complexity gives an extra meaning to the base space B of the fibration $q: E \to B$. The external conditions in the system can be parameterized by means of the space B. With the algebraic topology (especially homotopy) tools, the parametrised topological complexity determines the degree of navigational complexity as a positive number for the system when the initial and final states have the same external conditions. We first rewrite the definition of the parametrised topological complexity with respect to the homotopic distance and then we update this definition for n > 1 to exhibit the parametrised higher topological complexity TC_n[$q: E \to B$] in Section 5.

Topological complexity number varies such as relative, symmetric, monoidal, parametric etc., and accordingly, describing the higher version of all these numbers is a requirement for understanding the general concept of robot motion planning algorithms. For instance, see [15] and [4] for the simplicial complexity. Davis investigates the symmetric complexity of a circle [9] and the geodesic complexity of Klein bottles with the dimension n [10], respectively. In this study, we investigate the general setting of certain topological complexities. First, we recall some definitions and facts about TC and the related invariants such as cat (Lusternik-Schnirelmann category, see for a detail information [8]), Schwarz genus, and the homotopic distance. Then we give the homotopic distance definitions of relative higher topological complexity of a space, relative higher topological complexity of a pair, and the parametrised higher topological complexity of a fibration in the respective sections. We also mention on some equalities and inequalities including TC_n , cat, and TC of a fibration with considering particular homotopy facts. In addition, we obtain lower and upper bounds for Hopf fibration and examine the Stiefel and Grasmann manifolds in the sense of parametrised topological complexity.

2. Preliminaries

This section is dedicated to providing brief information about the different types of topological complexities and their related invariants. Note that we frequently use these facts in the following sections.

2.1. Schwarz Genus and Homotopic Distance.

Definition 2.1. [25] Let $q: E \to B$ be a fibration. Assume that B has an open cover $\{B_1, B_2, \dots, B_r\}$ for the minimum possible positive integer r such that there is a continuous map $s_j: B_j \to E$ satisfying that $q \circ s_j = 1_{B_j}$ for each $j \in \{1, 2, \dots, r\}$. Then the Schwarz genus of q is r.

The Schwarz genus of q is generally denoted by genus(q) or secat(q).

Definition 2.2. [5, 21] Let $f_j : A \to B$ be a continuous map for each $j \in \{1, 2, \dots, m\}$. Assume that A has an open cover $\{A_1, A_2, \dots, A_r\}$ for the minimum possible positive integer r such that the condition

$$f_1^i \big|_{A_i} \simeq f_2^i \big|_{A_i} \simeq \cdots \simeq f_m^i \big|_{A_i}$$

holds for all $i \in \{1, \dots, r\}$. Then the higher homotopic distance of degree m (or the m-th homotopic distance) is r.

The higher homotopic distance is denoted by $D(f_1, f_2, \dots, f_m)$ and is simply called the higher homotopic distance. For m = 2, the notion is particularly called the homotopic distance [21].

Theorem 2.3. [21] Let f_1 , $f_2 : A \to B$ be two maps and $\pi : PB \to B \times B$. Assume that $p : P \to A$ is the pullback of π by $(f_1, f_2) : A \to B \times B$, where

$$P = \{(a, \beta) \in A \times PB : \beta(0) = f_1(a) \text{ and } \beta(1) = f_2(a)\},\$$

i.e., the following diagram is commutative:

$$P \xrightarrow{p_{T_2}} PB$$

$$\downarrow_p \qquad \qquad \downarrow_{\pi}$$

$$A \xrightarrow{(f_1, f_2)} B \times B$$

Then $D(f_1, f_2) = genus(p)$.

We have some useful notes on the (higher) homotopic distance for the forthcoming sections as follows [5, 21]:

Proposition 2.4. [5, 21] Each of the following satisfies:

a) If $f_j : A \to B$ is homotopic to $g_j : A \to B$ for each $j \in \{1, \dots, n\}$, then $D(f_1, \dots, f_n) = D(g_1, \dots, g_n)$.

b) Let 1 < k < l and $f_1, \dots, f_k, \dots, f_l : A \to B$ be maps. Then we have that $D(f_1, \dots, f_k) \leq D(f_1, \dots, f_l).$

c) Let $f_j : A \to B$ and $p_j : B \to C$ be maps with $p_{j-1} \simeq p_j$ for each $j \in \{2, \dots, n\}$. Then $D(p_1 \circ f_1, \dots, p_n \circ f_n) \leq D(f_1, \dots, f_n)$.

d) Let $f_j : A \to B$ and $p_j : C \to A$ be maps with $p_{j-1} \simeq p_j$ for each $j \in \{2, \dots, n\}$. Then $D(f_1 \circ p_1, \dots, f_n \circ p_n) \leq D(f_1, \dots, f_n)$.

e) The higher homotopic distance is a homotopy invariant.

f) Let $g, g': C \to A$ and $f_1, f_2: A \to B$ be maps. If $f_1 \circ g' \simeq f_2 \circ g'$, then $D(f_1 \circ g, f_2 \circ g) \leq D(g, g')$.

2.2. **Topological Complexity and LS-Category.** Topological complexity, the main part of the studies of topological robotics, can be expressed in two ways. For simplicity, we use SG and HD as abbreviations for any definition in the sense of Schwarz genus and homotopic distance, respectively.

Definition 2.5. [11, 21] Let X be a path-connected topological space.

- (SG) Let $\pi : PX \to X \times X$, $\pi(\alpha) = (\alpha(0), \alpha(1))$, be a path fibration. Then the topological complexity of X is the genus (π) .
- (HD) Let $p_i : X^2 \to X$ be a projection map for each $i \in \{1, 2\}$. Then the topological complexity of X is $D(p_1, p_2)$.

TC of a contractible space is 1 and the converse is also true, that is, if TC(X) = 1, then X is a contractible space [11].

Proposition 2.6. [11] Let X be a paracompact and locally contractible space with dimX = n. Then $TC(X) \le 2n + 1$.

An important result of Proposition 2.6 is that 2cat(X) - 1 is an upper bound for TC(X) when X is paracompact. cat is also a natural lower bound TC. Thus, we conclude that $cat(X) \leq TC \leq 2cat(X) - 1$.

Theorem 2.7. [16] The topological complexity of the complex Stiefel manifold $V_r(\mathbb{C}^k)$, denoted by $TC(V_r(\mathbb{R}^k))$, is less than or equal to 2r(k-r) + 1.

The topological complexity is denoted by TC and generalized for n > 1 as follows.

Definition 2.8. [5, 21, 24] Let X be a path-connected space.

- (SG) Let $e_n : P_n(X) \to X^n$, $e_n(\gamma) = (\gamma_1(1), \gamma_2(1), \dots, \gamma_n(1))$, be a fibration, where $P_n(X)$ is the space of all multipaths in X such that the initial point of all paths in each multipath is the same. Then the n-th topological complexity (simply called the higher topological complexity) of X is the genus(e_n).
- (HD) Let $p_i : X^n \to X$ be a projection map for each $i \in \{1, 2, \dots, n\}$. Then the higher topological complexity of X is $D(p_1, p_2, \dots, p_n)$.

The higher topological complexity is denoted by TC_n and it is useful for n > 1because $\mathrm{TC}_1(X) = 1$ for any space X. One of the important observations on TC_n is the equality $\mathrm{TC}_2(X) = \mathrm{TC}(X)$. Another is the inequality $\mathrm{TC}_n(X) \leq \mathrm{TC}_{n+1}(X)$.

The Lusternik Schnirelmann category (simply called LS-category or denoted by cat) is another important homotopy invariant that inspired TC. It is also a natural bound for the topological complexity number.

Definition 2.9. [8,21] Let X be a path-connected topological space

- (SG) Let $\eta: P_0X \to X$, $\eta(\gamma) = (\gamma(1))$, be a Serre Fibration, where $P_0(X)$ is the special case of $P_n(X)$, i.e., the space of all paths in X for which the first point of these paths is a constant point x' in X. Then the LS-category of X is genus (η) .
- (HD) Let x' be any point of X. For the continuous maps $i_1: X \to X \times X$, $i_1(x) = (x, x')$ and $i_2: X \to X \times X$, $i_2(x) = (x', x)$, the LS-category of X is $D(i_1, i_2)$.

Proposition 2.10. [18] Let A and B be paracompact and path-connected spaces. Then $cat(A \times B) \leq cat(A) + cat(B)$.

Assume that $q: E \to B$ is a fibration with the fiber X. Then we have that $TC(E) \leq TC(X) + cat(B \times B) + 1$ [22]. Another important observation is that the numbers TC and cat coincide when the topological space is a connected Lie group [12].

Theorem 2.11. [3] The LS-category of the complex Grassmannian $G_r(\mathbb{C}^k)$, denoted by $cat(G_r(\mathbb{C}^k))$, equals rk.

Theorem 2.12. [19] The LS-category of the quaternionic Grassmannian $G_r(\mathbb{H}^k)$, denoted by $cat(G_r(\mathbb{H}^k))$, equals k(r-k).

It is possible to compute TC or TC_n of a fibration as well as TC of a pathconnected topological space [17, 20, 23].

Definition 2.13. [17,20,23] Let $q: E \to B$ be a surjective fibration.

- (SG) For the fibration $\pi'': PE \to E \times B$ defined as $\pi''(\gamma) = (\gamma(0), q \circ \gamma(1)),$ the topological complexity of q, denoted by TC(q), is genus (π'') .
- (HD) Let $\pi_1 : E \times B \to E$ be the first projection map, and $\pi_2 : E \times B \to B$ the second projection map. Then the topological complexity of q is $D(\pi_1, \pi_2)$.

Proposition 2.14. [23] Let $q_1 : E_1 \to B_1$ and $q_2 : E_2 \to B_2$ be two fibrations. Then

 $\max\{TC(q_1), TC(q_2)\} \le TC(q_1 \times q_2) \le TC(q_1) + TC(q_2) - 1.$

Proposition 2.15. [23] Let $q: E \to B$ be a fibration. Then $cat(B) \leq TC(q) \leq \min\{TC(B), cat(E \times B)\}.$ The general case of TC(q) is given in [17] with the following definition:

Definition 2.16. [17] Let $q: (q_1, q_2, \dots, q_n): E \to B^n$ be a surjective fibration.

- (SG) For a fibration $e_n^q : P_n(E) \to Y^n$, the *n*-th (higher) topological complexity of q is genus(e_n^q).
- (HD) For each projection map $p_i : E^n \to E$ with $i \in \{1, 2, \dots, n\}$, the *n*-th (higher) topological complexity of q is $D(q \circ p_1, q \circ p_2, \dots, q \circ p_n)$.

Theorem 2.17. [17] If $q_1 \simeq q_2 : E \to B^n$ are homotopic fibrations, then $TC_n(q_1) = TC_n(q_2).$

Theorem 2.18. [2] Let X be a path-connected space. Then $cat(X^{n-1}) \leq TC_n(X) \leq cat(X^n).$

2.3. **Different Types of Topological Complexity.** First, we recall the notions the relative (subspace) homotopic distance and the relative topological complexity. Later, we remind the definition of the parametrised topological complexity.

Definition 2.19. [20] Let $f, g: A \to B$ be two continuous maps with the subset $Y \subset A$. Then the relative homotopic distance (or subspace distance) on Y is the homotopic distance of two maps $f|_Y$ and $g|_Y$, informal saying,

$$D_A(Y; f, g) = D(f|_Y, g|_Y).$$

Definition 2.20. [13, 20] Let X be a path-connected topological space and $Y \subset X \times X$.

- (SG) Let $\pi^Y : P_Y X \to Y, \pi^Y(\alpha) = (\alpha(0), \alpha(1))$, be a path fibration, where $P_Y X$ is a subset of PX and contains all the paths in X with the property that $(\alpha(0), \alpha(1))$ lies in Y. Then the relative topological complexity of X with respect to the subspace Y is genus (π^Y) .
- (HD) Let $p_i : X^2 \to X$ be the projection map for each $i \in \{1, 2\}$ and $i_Y : Y \to X \times X$ be the inclusion map. Then the relative topological complexity of X with respect to the subspace Y is $D_{X \times X}(Y; p_1, p_2)$.

The relative topological complexity of X with respect to the subspace Y is denoted by $TC_Y(X)$.

Definition 2.21. [26] **(SG)** Let A be a path-connected topological space and $B \subset A$. Let $\pi^{A \times B} : P_{A \times B} \to A \times B$, $\pi^{A \times B}(\gamma) = (\gamma(0), \gamma(1))$, be a fibration, where $P_{A \times B}$ is the space of all paths in A with the property that $\gamma(0) \in A$ and $\gamma(1) \in B$. Then the relative topological complexity of a pair (A, B) is genus $(\pi^{A \times B})$.

The relative topological complexity of a pair (A, B) is denoted by TC(A, B). We mention a similar result to the ordinary TC number:

Proposition 2.22. [26] Let (A, B) be the pair. Then $TC(A, B) = 1 \Leftrightarrow A$ is contractible. **Definition 2.23.** [7] **(SG)** Let $q: E \to B$ be a fibration with nonempty, pathconnected topological space $X = q^{-1}(b)$ for any $b \in B$. Let

$$E_B^I = \{\beta \in PE : q \circ \beta \text{ is constant}\} \subset PE$$

and

$$E \times_B E = \{(a, b) \in E \times E : q(a) = q(b)\} \subset E^2$$

Then for the fibration $\pi_B : E_B^I \to E \times_B E$ defined by $\pi_B(\beta) = \beta(0), \beta(1)$, the parametrised topological complexity of q, denoted by $\text{TC}[q : E \to B]$, is genus(π_B).

Proposition 2.24. [7] Let $q_1 : E_1 \to B_1$ and $q_2 : E_2 \to B_2$ be two fibrations with fibers X_1 and X_2 , respectively, for metrisable spaces E_1 , E_2 , B_1 , and B_2 . For the fibration $q = q_1 \times q_2 : E_1 \times E_2 \to B_1 \times B_2$ with the fiber $X_1 \times X_2$, we have

 $TC[q: E_1 \times E_2 \to B_1 \times B_2] \le TC[q_1: E_1 \to B_1] + TC[q_2: E_2 \to B_2].$

3. Relative Higher Topological Complexity of a Space

Definition 3.1. (HD) Let X be a path-connected topological space and $Y \subseteq X^n$ be a subspace. Then the relative higher topological complexity is defined as

$$TC_{n,X}(Y) = D_{X^n}(Y; p_1, \cdots, p_n),$$

where $p_i: X^n \to X$ is a projection onto the *i*-th factor for each $i = 1, \dots, n$.

In particular, if Y is choosen as X^n , then we find $TC_{n,X}(X^n) = TC_n(X)$. Indeed,

 $TC_{n,X}(X^n) = D_{X^n}(X^n; p_1, \cdots, p_n) = D(p_1, \cdots, p_n) = TC_n(X).$

Proposition 3.2. Let Y be a subspace of X^n and $p_i : X^n \to X$ be the corresponding projection map for each $i = 1, \dots, n$. Then $TC_{n,X}(Y) = 1$ if and only if the projections $(p_i)|_Y : Y \to X$ with $i = 1, \dots, n$ are homotopic to each other.

Proof. If Y is a subspace of X^n , then we get

$$\operatorname{TC}_{n,X}(Y) = 1 \Leftrightarrow \operatorname{D}_{X^n}(Y; p_1, \cdots, p_n) = 1 \Leftrightarrow (p_1)|_Y \simeq \cdots \simeq (p_n)|_Y.$$

Proposition 3.3. Let Y be a subspace of X^n . Then $TC_{n,X}(Y) \leq TC_n(X)$.

Proof. If Y is a subspace of X^n , then we have an inclusion map $i_Y : Y \to X^n$. Therefore, by (d) part of Proposition 2.4, we observe that

$$TC_{n,X}(Y) = D_{X^n}(Y; p_1, \cdots, p_n)$$

= $D(p_1 \circ i_Y, \cdots, p_n \circ i_Y)$
 $\leq D(p_1, \cdots, p_n)$
= $TC_n(X).$

Proposition 3.4. Let $Y \subset Z \subset X^n$. Then $TC_{n,X}(Y) \leq TC_{n,X}(Z)$.

Proof. Let $i_Y : Y \to X^n$, $i_Z : Z \to X^n$, and $i : Y \to Z$ be three inclusion maps. Then i_Y can be rewritten as the composition of i and i_Z i.e., $i_Y = i_Z \circ i$. From Proposition 2.4 (d), we have

$$\begin{aligned} \mathrm{TC}_{n,X}(Y) &= \mathrm{D}_{X^n}(Y; p_1, \cdots, p_n) = \mathrm{D}(p_1 \circ i_Y, \cdots, p_n \circ i_Y) \\ &= \mathrm{D}(p_1 \circ i_Z \circ i, \cdots, p_n \circ i_Z \circ i) \leq \mathrm{D}(p_1 \circ i_Z, \cdots, p_n \circ i_Z) \\ &= \mathrm{D}_{X^n}(Z; p_1, \cdots, p_n) \\ &= \mathrm{TC}_{n,X}(Z). \end{aligned}$$

Proposition 3.5. If Y is a retract of X, then $TC_{n,X}(Y^n) \ge TC_n(Y)$.

Proof. Let $r: X \to Y$ be a retraction map. Let $p_j: X^n \to X$ and $q_j: Y^n \to Y$ be the projection maps for each $j = 1, \dots, n$. By (c) part of Proposition 2.4, we find

$$TC_{n,X}(Y^n) = D_{X^n}(Y^n; p_1, \cdots, p_n) = D(p_1 \circ i_{Y^n}, \cdots, p_n \circ i_{Y^n})$$

$$\geq D(r \circ p_1 \circ i_{Y^n}, \cdots, r \circ p_n \circ i_{Y^n}) = D(q_1, \cdots, q_n)$$

$$= TC_n(Y)$$

with considering the fact that $q_j = r \circ p_j \circ i_{Y^n}$ for each j.

Corollary 3.6. If Y is a retract of X then, $TC_n(Y) \leq TC_n(X)$.

Proof. Since $Y \subset X$ is a retract, $Y^n \subset X^n$. Then Proposition 3.4 says that $\operatorname{TC}_{n,X}(Y^n) \leq \operatorname{TC}_{n,X}(X^n) = \operatorname{TC}_n(X)$. Using Proposition 3.5, we conclude that $\operatorname{TC}_n(Y) \leq \operatorname{TC}_n(X)$.

Proposition 3.7. Let $\{Y_1, \dots, Y_m\}$ be an open covering of X^n . Then $TC_{n,X}(Y_1) + \dots + TC_{n,X}(Y_m) \leq m \cdot TC_n(X).$

Proof. Let Y_1, \dots, Y_m be an open subsets of X^n with the fact $X^n = Y_1 \cup \dots \cup Y_m$. Then we have that

$$TC_{n,X}(Y_{1}) + \dots + TC_{n,X}(Y_{m})$$

$$= D_{X^{n}}(Y_{1}; p_{1}, \dots, p_{n}) + \dots + D_{X^{n}}(Y_{m}; p_{1}, \dots, p_{n})$$

$$= D(p_{1} \circ i_{Y_{1}}, \dots, p_{n} \circ i_{Y_{1}}) + \dots + D(p_{1} \circ i_{Y_{m}}, \dots, p_{n} \circ i_{Y_{m}})$$

$$\leq D(p_{1}, \dots, p_{n}) + \dots + D(p_{1}, \dots, p_{n})$$

$$= m \cdot TC_{n}(X)$$

from (d) part of Proposition 2.4.

Theorem 3.8. Let $Y \subseteq X^2$. Then $TC_{2,X}(Y) \leq cat_{X \times X}(Y)$.

Proof. Let $p_1, p_2 : X \times X \to X$ be projections and $* : Y \to X \times X$ a constant map. The fact $p_1 \circ * \simeq p_2 \circ *$ implies $D(p_1 \circ i_Y, p_2 \circ i_Y) \leq D(i_Y, *)$ from (f) of Proposition 2.4. Thus, we conclude that $TC_{2,X}(Y) \leq cat_{X \times X}(Y)$.

It is clear that $TC_{2,X}(Y) = TC_X(Y)$. The previous theorem also confirms the fact that $TC_X(Y) \leq \operatorname{cat}_{X \times X}(Y)$. Theorem 3.8 can be generalized as follows with considering Theorem 2.18:

Corollary 3.9. Let $Y \subseteq X^n$. Then $TC_{n,X}(Y) \leq cat_{X^n}(Y)$.

Theorem 3.10. Let $Y, Z \subset X^n$ such that Y and Z have the same homotopy type. Then $TC_{n,X}(Y) = TC_{n,X}(Z)$.

Proof. Let $\beta : Z \to Y$ be the homotopy equivalence map. Assume that $i \in \{1, \dots, n\}$. Consider the following commutative diagram for the projection $p_i : X^n \to X$ with the inclusions $i_Y : Y \to X^n$ and $i_Z : Z \to X^n$:



Hence, by Theorem 2.17, we get

$$D(p_1 \circ i_Y, \cdots, p_n \circ i_Y) = D(p_1 \circ i_Z, \cdots, p_n \circ i_Z)$$

which concludes that $TC_{n,X}(Y) = TC_{n,X}(Z)$.

4. Relative Higher Topological Complexity of a Pair

Definition 4.1. (SG) Let A be a path-connected space and $B \subseteq A$. Set

$$P'_{A \times B} = \{ \alpha \in A^{J_n} : \alpha(0) \in A, \ \alpha(1) = (\alpha_1(1), \cdots, \alpha_n(1)) \in B \} \subseteq A^{J_n}.$$

Then for a fibration $e'_{n}: P'_{A \times B} \to B^{n}$ defined with $e'_{n}(\alpha) = (\alpha_{1}(1), \cdots, \alpha_{n}(1))$, the relative higher topological complexity of the pair (A, B) is defined as

$$TC_n(A, B) = genus(e'_n).$$

Note that $e'_n : P'_{A \times B} \to B^n$ is indeed a fibration because the restriction of a fibration $e_n : A^{J_n} \to A^n$, $e_n(\alpha) = (\alpha_1(1), \cdots, \alpha_n(1))$, to a subset $B^n \subset A^n$ is e'_n .

Proposition 4.2. a) $TC_1(A, B) = 1$. This means that the notation $TC_n(A, B)$ is significative for n > 1.

b) $TC_n(A,B) = genus(d'_n)$, where $d'_n : B \to B^n$ is a diagonal map since e'_n is a fibrational substitute of d'_n .

c) For
$$n = 2$$
, $TC_2(A, B) = TC(A, B)$.

Proof. **a)** Let n = 1. Then for a fibration $e'_1 : P'_{A \times B} \to B$ with $e'_1(\alpha) = \alpha_1(1)$, we construct a map $s : B \to P'_{A \times B}$ such that s takes any point y of B to the constant path ϵ_y at this point. Therefore, we get

$$e_{1}^{'} \circ s(y) = e_{1}^{'}(\epsilon_{y}) = y = 1_{B}(y).$$

Thus, $genus(e'_1)$ equals 1.

b) Take a homotopy equivalence $h: B \to P'_{A \times B}$ defined as $h(y) = \epsilon_y$, where ϵ_y is a constant path at y. Then we get

$$e'_{n} \circ h(y) = e'_{n}(\epsilon_{y}) = (y, \cdots, y) = d'_{n}(y).$$

c) Let n = 2. Define $e_2^{''}: P_{A \times B} \to A \times B$ as $e_2^{''}(\alpha) = (\alpha(0), \alpha(1))$. Then $e_2^{''}$ is a fibrational substitute of the diagonal map $d_2': B \to B^2$ because $h': B \to P_{A \times B}$, $h(y) = \epsilon_y$, is a homotopy equivalence and the condition $e_2^{''} \circ h' = d_2'$ holds with considering that $B \subseteq A$. Thus, we find $\operatorname{TC}_2(A, B) = \operatorname{genus}(e_2^{''}) = \operatorname{TC}(A, B)$. \Box

One of the well-known results of Schwarz [25] leads to us having an important relationship between the relative higher topological complexity and the Lusternik-Schnirelmann category:

Corollary 4.3. For a path-connected space A with its subset B, we have $TC_n(A,B) \leq cat(B^n)$. In addition, if A is contractible, then we conclude that $TC_n(A,B) = cat(B^n)$.

By Corollary 4.3, we immediately have that if B is contractible, then we get

 $TC_n(A, B) \le \operatorname{cat}(B^n) = 1,$

i.e., $TC_n(A, B) = 1$.

Example 4.4. For any point x_0 in a path-connected space A, we obtain that $TC_n(A, \{x_0\}) = 1$.

It is possible to improve Corollary 4.3 with using Proposition 2.10:

Corollary 4.5. Let A be path-connected and B be a path-connected and paracompact subset of A. Then we have $TC_n(A, B) \leq n \cdot cat(B)$.

Proof. By Corollary 4.3, we obtain $TC_n(A, B) \leq cat(B^n)$. Since B is pathconnected and paracompact, we observe that $cat(B^n) \leq n \cdot cat(B)$.

Besides Schwarz genus, the relative higher topological complexity $TC_n(A, B)$ of a pair (A, B) can also be defined by higher homotopic distance:

Definition 4.6. (HD) Let A be a path-connected space and $B \subseteq A$. Then

 $TC_n(A, B) = D_{A^n}(A \times B \times B \times \dots \times B; p_1, p_2, \dots, p_n)$

for n > 1, where each p_i is a projection from $A \times A \times \cdots \times A$ to A onto the *i*-th factor for $i = 1, \cdots, n$.

We assume that $TC_1(A, B)$ always equals 1. If B = A, then we conclude that $TC_n(A, A) = TC_n(A)$. We observe that $TC_n(A, B) \leq TC_{n+1}(A, B)$ as well as TC_n of a space or a fibration.

Proposition 4.7. Let $B_1 \subset B_2 \subset A$. Then $TC_n(A, B_1) \leq TC_n(A, B_2)$.

Proof. When we consider three inclusion maps $i_{B_1} : B_1 \to A$, $i_{B_2} : B_2 \to A$, $i : B_1 \to B_2$ such that $i_{B_1} = i_{B_2} \circ i$, the remaining part of the proof goes similar to the proof of Proposition 3.4.

Proposition 4.8. If A is path-connected with a subset $B \subset A$, then we have $TC_n(A, B) \leq TC_n(A)$.


Proof. Let $i: A \times B \times B \times \cdots \times B \to A^n$ be an inclusion map. Then Proposition 2.4 (d) gives us that

$$TC_n(A, B) = D_{A^n}(A \times B \times B \times \dots \times B; p_1, p_2, \dots, p_n)$$

= $D(p_1 \circ i, p_2 \circ i, \dots, p_n \circ i)$
 $\leq D(p_1, p_2, \dots, p_n)$
= $TC_n(A).$

5. Parametrised (Higher) Topological Complexity Using Homotopic Distance

The task in this section is to mention the homotopic distance definition of the parametrised topological complexity. Let $q: E \to B$ be a fibration and $X \neq \emptyset$ is a path-connected fiber for the fibration q. Let E_B^I be a set consisting of all continuous paths γ in E such that $q \circ \gamma$ is a constant path. Moreover, $E \times_B E$ is a subset of $E \times E$ and it contains all points (e, e') with the condition q(e) equals q(e'). Recall that the map $\pi: E_B^I \to E \times_B E$ with $\pi(\gamma) = (\gamma(0), \gamma(1))$ is a fibration with fibre ΩX . Then consider the following diagram (see also Theorem 2.3):



Here $p_i^B : E \times_B E \to E$ is a projection map onto the *i*th factor for i = 1, 2. Similarly, π_i is another projection map for each i = 1, 2. Also, P is given by the set $\{(e, e', \gamma) : \gamma(0) = p_1(e, e') = e, \gamma(1) = p_2(e, e') = e'\}$, and it is clearly a subset of $E \times_B E \times E_B^I$. This yields that $D(p_1^B, p_2^B) = genus(\pi_1)$. Since $(p_1^B, p_2^B) = 1_{E \times_B E}$, we observe that $genus(\pi_1) = genus(\pi)$. Combining this result with the Definition 2.23, we have a new statement of the parametrised topological complexity on homotopic distance:

Definition 5.1. (HD) The parametrised topological complexity is defined as

$$TC[q: E \to B] = D(p_1^B, p_2^B)$$

for the projection map $p_i^B : E \times_B E \to E$ with each i = 1, 2.

For the improved version of Definition 5.1, we note that the parametrised higher topological complexity is defined as

$$\operatorname{TC}_n[q: E \to B] = D(p_1^B, \cdots, p_n^B)$$

for n > 1 and the map $p_i^B : E \times_B E \cdots \times_B E \to E$ with each $i = 1, \cdots, n$.

If n = 1, then $\mathrm{TC}_1[q: E \to B]$ is always 1. The second observation states that the inequality $\mathrm{TC}_n[q: E \to B] \leq \mathrm{TC}_{n+1}[q: E \to B]$ holds. Furthermore, one can easily observe the equality $\mathrm{TC}_2[q: E \to B] = \mathrm{TC}[q: E \to B]$ by using the higher homotopic distance.

Proposition 5.2. Let $q : E \to B$ be a fibration. Let $q|_{B'} = q' : E' \to B'$ be another fibration with $B' \subset B$ and $E' = q^{-1}(B')$. Then

$$TC[q': E' \to B'] \le TC[q: E \to B].$$

Remark 5.3. Proposition 5.2 is first expressed in [7]. We now give explicit proof by using the homotopic distance.

Proof. By Definition 5.1, $\operatorname{TC}[q: E \to B]$ and $\operatorname{TC}[q': E' \to B']$ are respectively equal to $\operatorname{D}(p_1^B, p_2^B)$ and $\operatorname{D}(p_1^{B'}, p_2^{B'})$ for the projection maps $p_i^B: E \times_B E \to E$ and $p_i^{B'}: E' \times_{B'} E' \to E'$ for each i = 1, 2. The projection map $p_i^{B'}$ can be thought as $p_i^B \circ j$, where $j: E' \times_{B'} E' \to E \times_B E$ is an inclusion. Therefore, we have

$$D(p_1^{B'}, p_2^{B'}) = D(p_1^{B} \circ j, p_2^{B} \circ j)$$

via (d) of Proposition 2.4. Finally, the inequality $D(p_1^B \circ j, p_2^B \circ j) \leq D(p_1^B, p_2^B)$ concludes that $TC[q' : E' \to B'] \leq TC[q : E \to B]$.

It is possible that the parametrised topological complexity can be defined by the relative topological complexity. Indeed,

$$TC[q: E \to B] = D(p_1^B, p_2^B)$$

= $D(p_1 \circ i_{E \times BE}, p_2 \circ i_{E \times BE})$
= $D_{E^2}(E \times BE; p_1, p_2)$
= $TC_E(E \times BE).$

This fact is also improved with the following equality:

$$\operatorname{TC}_n[q: E \to B] = \operatorname{TC}_{n,E}(E \times_B E \cdots \times_B E).$$

Proposition 5.4. $TC_n[q: E \to B] \leq TC_n(E)$.

Proof. Let $j: E \times_B E \cdots \times_B E \to E \times E \cdots \times E$ be the inclusion map. Then, by Proposition 2.4 (d), we have

$$TC_n[q: E \to B] = D(p_1^B, \cdots, p_n^B)$$

= $D(p_1 \circ j, \cdots, p_n \circ j)$
 $\leq D(p_1, \cdots, p_n) = TC_n(E).$

Corollary 5.5. $TC[q: E \rightarrow B] \leq TC(E)$.

Theorem 5.6. $TC_n(q) \leq TC_n[q: E \rightarrow B].$

Proof. Let $j : E \times_B E \cdots \times_B E \to E \times E \cdots \times E$ be an inclusion map and $p_i : E^n \to E$ be the projection map for each $i = 1, \dots, n$. Then we find

$$TC_n(q) = D(q \circ p_1, \cdots, q \circ p_n)$$

$$\leq D(p_1, \cdots, p_n)$$

$$\leq D(p_1 \circ j, \cdots, p_n \circ j)$$

$$= D(p_1^B, \cdots, p_n^B) = TC_n[q : E \to B]$$

by using (c) and (d) parts of Proposition 2.4, respectively.

Corollary 5.7. $TC(q) \leq TC[q: E \rightarrow B]$.

Corollary 5.8. a) Let $q_1 : E_1 \to B_1$ be a fibration with a path-connected fiber X_1 and $q_2 : E_2 \to B_2$ be another fibration with a path-connected fiber X_2 . If E_1 , B_1 , E_2 and B_2 are metrisable, then

 $\max\{TC(q_1), TC(q_2)\} \le TC[q_1 : E_1 \to B_1] + TC[q_2 : E_2 \to B_2].$ **b)** $cat(B) \le TC[q : E \to B].$

Proof. a) Let $q_1 \times q_2 : E_1 \times E_2 \to B_1 \times B_2$ be a fibration with a path-connected fiber $X_1 \times X_2$. Then by Proposition 2.14, Corollary 5.7, and Proposition 2.24, respectively, we have

$$\max\{TC(q_1), TC(q_2)\} \leq TC(q_1 \times q_2)$$

$$\leq TC[q_1 \times q_2 : E_1 \times E_2 \to B_1 \times B_2]$$

$$\leq TC[q_1 : E_1 \times B_1] + TC[q_2 : E_2 \to B_2].$$

b) This is obvious from Proposition 2.15 and Corollary 5.7.

Example 5.9. Hopf Fibration: Consider the Hopf fibration $q: S^3 \to S^2$ with the path-connected fiber S^1 . We shall show that $2 \leq \operatorname{TC}_n[q: S^3 \to S^2] \leq n$ for $n \geq 2$. Since $\operatorname{TC}_n(S^3) = n$ [24], by Corollary 5.5, $\operatorname{TC}_n[q: S^3 \to S^2] \leq n$. On the other hand, $\operatorname{cat}(S^2) = 2$ yields that $\operatorname{TC}_n[q: S^3 \to S^2] \geq 2$ via (b) part of Corollary 5.8.

Example 5.10. Stiefel and Grassmann Manifolds: Consider the complex case, i.e.,

$$U(r) \longrightarrow V_r(\mathbb{C}^k) \xrightarrow{q_1} G_r(\mathbb{C}^k).$$

Since $\operatorname{cat}(G_r(\mathbb{C}^k)) = rk$ by Theorem 2.11, Corollary 5.8 (b) states that

$$rk \leq \mathrm{TC}[q_1 : V_r(\mathbb{C}^k) \to G_r(\mathbb{C}^k)]$$

On the other hand, we have that $\operatorname{TC}(V_r(\mathbb{C}^k)) \leq 2r(k-r) + 1$ from Theorem 2.7. By Corollary 5.5, we get $\operatorname{TC}[q_1 : V_r(\mathbb{C}^k) \to G_r(\mathbb{C}^k)] \leq 2r(k-r) + 1$. Finally, we conclude that

$$rk \leq \mathrm{TC}[q_1: V_r(\mathbb{C}^k) \to G_r(\mathbb{C}^k)] \leq 2r(k-r) + 1$$

for n = 2 in the sense of TC_n .

If we assume that the quaternionic case, namely that,

$$Sp(r) \longrightarrow V_r(\mathbb{H}^k) \xrightarrow{q_2} G_r(\mathbb{H}^k),$$

then the inequality

$$k(r-k) \leq \mathrm{TC}[q_2: V_r(\mathbb{H}^k) \to G_r(\mathbb{H}^k)]$$

holds from the fact that $k(r-k) = \operatorname{cat}(G_r(\mathbb{H}^k))$ by Theorem 2.12. Now consider the real case:

$$O(n) \longrightarrow V_r(\mathbb{R}^k) \xrightarrow{q_3} G_r(\mathbb{R}^k).$$

First, assume that r = 2 and $k = 2^p + 1$ with any integer p > 0. Then, by Theorem 2.8 of [1], we get $\operatorname{TC}[q_3 : V_2(\mathbb{R}^{2^p+1}) \to G_2(\mathbb{R}^{2^p+1})] \ge 2^{p+1} - 2$. For the upper bound, we shall use Corollary 5.5. Since $\dim(V_2(\mathbb{R}^{2^p+1})) = 2^{p+1} - 1$, Proposition 2.6 says that $\operatorname{TC}(V_2(\mathbb{R}^{2^p+1})) \le 2^{p+2} - 1$. As a consequence,

$$2^{p+1} - 2 \le \mathrm{TC}[q_3 : V_2(\mathbb{R}^{2^p+1}) \to G_2(\mathbb{R}^{2^p+1})] \le 2^{p+2} - 1.$$

Now, assume that r = 2 and $k = 2^p + 2$ with any integer p. Similar to the previous case, by using Theorem 2.11 in [1], we have the following inequalities:

$$2^{p+1} - 1 \le \mathrm{TC}[q_3 : V_2(\mathbb{R}^{2^p+2}) \to G_2(\mathbb{R}^{2^p+2})] \le 2^{p+2} + 3$$


with considering the fact that $\dim(V_2(\mathbb{R}^{2^p+2})) = 2^{p+1} + 1$.

Theorem 5.11. If $q_1 : E_1 \to B_1$ and $q_2 : E_2 \to B_1$ are fiber homotopy equivalent with the same nonempty path-connected fiber X_1 for both two fibrations q_1 and q_2 , then

$$TC_n[q_1: E_1 \to B_1] = TC_n[q_2: E_2 \to B_1],$$

that is, the parametrised higher topological complexity is a fiber homotopy equivalent invariant.

Proof. Let $q_1 : E_1 \to B_1$ and $q_2 : E_2 \to B_1$ are fiber homotopy equivalent with the fiber X_1 . Then we have two maps $h : E_1 \to E_2$ and $k : E_2 \to E_1$ satisfying two conditions $h \circ k \simeq 1_{E_2}$ and $k \circ h \simeq 1_{E_1}$. For simplicity, we rewrite E'_1 and E'_2 as $E_1 \times_{B_1} E_1 \times_{B_1} \cdots \times_{B_1} E_1$ and $E_2 \times_{B_1} E_2 \times_{B_1} \cdots \times_{B_1} E_2$, respectively. Consider the homotopy equivalence map $\beta : E'_2 \to E'_1$. Assume that $p_i^B : E'_1 \to E_1$ and $q_i^B : E'_2 \to E_2$ are projections for each $i \in \{1, \dots, n\}$. Then the following commutative diagram



states that $D(p_1^{B_1}, \dots, p_n^{B_1}) = D(q_1^{B_1}, \dots, q_n^{B_1})$. As a consequence, the equality $TC_n[q_1: E_1 \to B_1] = TC_n[q_2: E_2 \to B_1]$ holds.

6. CONCLUSION

The relative topological complexity is one of the first examples of different types of TC. With this approach, a lower bound for TC is obtained. A similar approach to TC_n is done by the relative higher topological complexity on the notion homotopic distance in this study. Interestingly, it also gives a new way to introduce

the parametrised topological complexity, which has been one of the most popular numbers among several TC versions in the last few years due to the different constructions of its motion planning algorithm. The higher homotopic distance again leads to define the parametrised higher topological complexity. These higher settings of TC always provide a different and strong perspective on the subject of robot motion planning problems in daily life.

Acknowledgment. The first author is granted as a fellowship by the Scientific and Technological Research Council of Turkey TUBITAK-2211-A. In addition, this work was partially supported by the Research Fund of Ege University (Project Number: FDK-2020-21123).

References

- F. Akhtarifar and M.A. Asaidi, A note on topological complexity of real Grassmann manifolds, Journal of Mathematical Extension, 16, (2020).
- [2] I. Basabe, J. Gonzalez, Y. Rudyak and B. Tamaki, Higher topological complexity and its symmetrization, Algebraic & Geometric Topology, 14(4), 2103-2124 (2014).
- [3] I. Berstein, On the Lusternik-Schnirelmann category of grassmannians, Mathematical Proceedings of the Cambridge Philosophical Society, 79(1), 129-134 (1976).
- [4] A. Borat, Higher dimensional simplicial complexity, New York Journal of Mathematics, 26, 1130-1144 (2020).
- [5] A. Borat and T. Vergili, Higher homotopic distance, Topological Methods in Nonlinear Analysis, 57(2), 525-534 (2021).
- [6] D.C. Cohen, M. Farber and S. Weinberger, Parametrized topological complexity of collisionfree motion planning in the plane, arXiv:2010.09809v1 (2020).
- [7] D.C. Cohen, M. Farber, and S. Weinberger, Topology of parametrised motion planning algorithms, SIAM Journal of Applied Algebra and Geometry, 5(2), 229-249 (2021).
- [8] O. Cornea, G. Lupton, J. Oprea, and D. Tanre, Lusternik-Schnirelmann Category, Mathematical Surveys and Monographs, 103, AMS, Providence, RI, (2003).
- [9] D. Davis, The symmetric topological complexity of the circle, New York Journal of Mathematics, 23, 593-602 (2017).
- [10] D. Davis, and D. Recio-Mitter, The geodesic complexity of n-dimensional Klein bottles, New York Journal of Mathematics, 27, 296-318 (2021).
- [11] M. Farber, Topological complexity of motion planning, Discrete and Computational Geometry, 29, 211-221 (2003).
- [12] M. Farber, Instabilities of robot motion, Topology and Its Applications, 140(2-3), 245-266 (2004).
- [13] M. Farber, Invitation to Topological Robotics, Zurich Lectures in Advanced Mathematics, EMS (2008).
- [14] J.M. Garcia-Calcines, Formal aspects of parametrized topological complexity and its pointed version, Journal of Topology and Analysis (accepted), Doi number: 10.1142/S1793525321500631 (2022).
- [15] J. Gonzalez, Simplicial complexity: piecewise linear motion planning in robotics, New York Journal of Mathematics, 24, 279-292 (2018).
- [16] M. Grant, Topological complexity, fibrations and symmetry, Topology and its Applications, 159(1), 88-97 (2012).
- [17] M. Is and I. Karaca, Higher topological complexity for fibrations, FILOMAT (accepted), arXiv:2107.04465 [math.AT] (2022).
- [18] I.M. James, On category, in the sense of Lusternik–Schnirelmann, Topology, 17, 331-348 (1978).
- [19] E. Macias-Virgos, J. Pereira-Saez and D. Tanre, Morse theory and the Lusternik–Schnirelmann category of quaternionic grassmannians, Proceedings of the Edinburgh Mathematical Society, 60(2), 441-449 (2017).
- [20] E. Macias-Virgos, D. Mosquera-Lois and J. Pereira-Saez, Homotopic distance and generalized motion planning, arXiv:2105.13006 [math.AT] (2021).

- [21] E. Macias-Virgos and D. Mosquera-Lois, Homotopic distance between maps, Mathematical Proceedings of the Cambridge Philosophical Society, 172(1), 73-93 (2022).
- [22] B. Naskar and S. Sarkar, On LS-category and topological complexity of some fiber bundles and Dold manifolds, Topology and its Applications, 284, 107367 (2020).
- [23] P. Pavesic, Topological complexity of a map, Homology, Homotopy and Applications, 21, 107-130 (2019).
- [24] Y. Rudyak, On higher analogs of topological complexity, Topology and Its Applications, 157(5), 916-920 (2010). Erratum: Topology and Its Applications 157, 1118 (2010).
- [25] A.S. Schwarz, The genus of a fiber space, American Mathematical Society Translation, 55, 49-140 (1966).
- [26] R. Short, Relative topological complexity of a pair, Topology and its Applications, 248, 7-23 (2018).
- [27] K. Tanaka, Parametrized topological complexity of poset-stratified spaces, Journal of Applied and Computational Topology, 150, 1-26 (2021).

Melih IS, Ege University, Faculty of Sciences, Department of Mathematics, Izmir, Turkey

Email address: melih.is@ege.edu.tr

ISMET KARACA, EGE UNIVERSITY, FACULTY OF SCIENCE, DEPARTMENT OF MATHEMATICS, IZMIR, TURKEY

Email address: ismet.karaca@ege.edu.tr