

Thermal hadron resonances in chiral and $U(1)_A$ restoration

A. Gómez Nicola,^{1,*} J. Ruiz de Elvira,^{1,†} and A. Vioque-Rodríguez^{1,‡}

¹*Universidad Complutense de Madrid, Facultad de Ciencias Físicas,
Departamento de Física Teórica and IPARCOS. Plaza de las Ciencias 1, 28040 Madrid, Spain*

We review recent work on thermal resonances and their connection with chiral symmetry and $U(1)_A$ restoration within the QCD phase diagram. In particular, the $f_0(500)$ and $K_0^*(700)$ states generated from $\pi\pi$ and πK scattering within Unitarized Chiral Perturbation Theory (ChPT) at finite temperature allow one to describe scalar susceptibilities, which combined with Ward Identities yield interesting conclusions regarding the interplay between chiral and $U(1)_A$ restoration, key to understand the nature of the transition.

I. INTRODUCTION

The analysis and understanding of the QCD Phase Diagram is one of the major research lines in hadronic physics where considerable progress has been reached within recent years, mostly from lattice field theory and theoretical developments based on effective theories [1–7]. The coincidence of deconfinement and chiral symmetry restoration in the (T, μ_B) plane of temperature and baryon chemical potential allows one to use genuine chiral-restoring observables as signals of the transition, as we will see here. In fact, although observables related to deconfinement, such as the trace anomaly, or others indicating chiral symmetry breaking, such as the light quark condensate $\langle \bar{q}q \rangle_l$, receive contributions from many massive hadronic states and require the implementation of Hadron Resonance Gas (HRG) based approaches [8–12], we will focus here on those for which an effective theory approach based only on light hadronic degrees of freedom provides the dominant physical description. That is the case of observables constructed out of the following quark bilinears in the light channels:

$$\begin{aligned} I = 0 & : \sigma_l = \bar{\psi}_l \psi_l (S), \quad \eta_l = i\bar{\psi}_l \gamma_5 \psi_l (P) \\ I = 1 & : \pi^a = i\bar{\psi}_l \gamma_5 \tau^a \psi_l (P), \quad \delta^a = \bar{\psi}_l \tau^a \psi_l (S) \\ I = 1/2 & : K^b = i\bar{\psi} \gamma_5 \lambda^b \psi (P), \quad \kappa^b = i\bar{\psi} \lambda^b \psi (S) \end{aligned} \quad (1)$$

with $a = 1, 2, 3$, $b = 4, 5, 6, 7$. We denote by ψ_l the light quark (u, d) field doublet in flavour space; the isospin as well as the pseudoscalar (P) or scalar (S) character of each operator are indicated, and τ^a and λ^a denote respectively Pauli and Gell-Mann matrices.

The lowest lying states in the hadron spectrum with the quantum numbers of the above quark operators are the $f_0(500)$ (σ_l), mostly dominated by its light quark component, the light component of the η (η_l), the pion (π^a), the $a_0(980)$ (δ^a), the kaon (K^b) and the $K_0^*(700)$ (κ^b).

The interest to study the above hadronic states in connection with the QCD transition is that for exact restoration of the chiral symmetry $SU(N_f)_V \times SU(N_f)_A$, with N_f the number of light flavors, the pairs $\pi^a - \sigma_l$, $\delta^a - \eta_l$, $K^b - \kappa^b$ can be connected by the chiral group. In other words, they become chiral partners, so that any observable calculated from their correlators, such as susceptibilities ($p = 0$ correlator in Fourier space) or spatial screening masses ($p^0 = 0$, $|\vec{p}| \rightarrow 0^+$) would degenerate for two given chiral partners in the regime of exact chiral restoration. That would be the case e.g. for $N_f = 2$ massless quarks at the QCD transition temperature T_c [13]. We recall that in the physical case of $N_f = 2 + 1$ massive flavours the transition for $\mu_B = 0$ is most likely a crossover at $T_c \simeq 155$ MeV [1–5]. In addition, the scalar susceptibility, i.e. the σ_l $p = 0$ correlator, develops a peak around T_c which is actually one of the more reliable sources for establishing the chiral restoration crossover in the lattice.

As a bonus, the previous operators provide also information about the possibility of the restoration of the $U(1)_A$ symmetry, which was predicted long time ago as an asymptotic mechanism rather than a sharp transition [14–17]. The relevant issue here, which is still under study in the community, is whether such $U(1)_A$ restoration takes place close enough to the chiral transition. Should that be the case, it would affect many interesting aspects of the phase diagram such as the order and universality class of the transition [13, 18]. There are also phenomenological consequences such

*Electronic address: gomez@ucm.es

†Electronic address: jacobore@ucm.es

‡Electronic address: avioque@ucm.es




FIG. 1: $I = 0$ (left) and $I = 1/2$ (right) scalar susceptibilities within the saturation approach, from [34] and [37] respectively. In the left panel, the lattice points are from [1], only points up to $T_c \simeq 155$ MeV are included in the fit and the uncertainty corresponds to the 95% confidence level. In the right panel the uncertainty bands correspond to variations within the LEC uncertainty range.

as the reduction of the η' mass, since its main contribution comes from the $U(1)_A$ axial anomaly [19]. As a matter of fact, Bose-Einstein correlations can be fitted experimentally with a reduced in-medium η' mass [20]. Such decreasing trend is also measured in the lattice [21].

From the viewpoint of the bilinear operators considered above, now $\pi^a - \delta^a$, $\sigma - \eta_l$, $K^b - \kappa^b$ can be connected by a $U(1)_A$ rotation and therefore degeneration of observables for those pairs would be the relevant signal to study the strength of $U(1)_A$ breaking at the chiral transition. Note that the $K - \kappa$ degenerate both for chiral and $U(1)_A$ restoration.

The results obtained by lattice collaborations regarding degeneration of chiral and $U(1)_A$ partners show some tension among them. On the one hand, $N_f = 2$ analyses close to the light chiral limit show compatibility between chiral and $U(1)_A$ restoration and a small gap between them for massive quarks [22–24]. On the other hand, including strangeness, i.e. for $N_f = 2 + 1$ simulations, and physical masses, $U(1)_A$ symmetry is still significantly broken at the chiral transition [25]. As we will see below, our recent analysis based on Unitarized Effective Theories and Ward Identities helps to reconcile those results and emphasizes the role of thermal resonances.

II. THE ROLE OF THERMAL RESONANCES

The spectral properties of resonances develop a dependence with temperature and chemical potentials when those resonances are generated and decay inside the thermal bath or in-medium environment. In some particular cases, such dependence, which can be inferred from hadron effective theories, turns out to be crucial to explain some of the aspects of the QCD phase diagram discussed in the introduction.

A remarkable example is the $f_0(500)$ (formerly known as σ) resonance. The nature and even the existence of that state has been the object of a myriad of studies in the past and is nowadays firmly established as a broad pole in the second Riemann sheet of the $I = J = 0$ $\pi\pi$ scattering amplitude [26–28], which is successfully described as dynamically generated within unitarized effective theories [29, 30]. Its thermal dependence has been obtained from a Chiral Perturbation Theory (ChPT) calculation of the unitarized finite-temperature $\pi\pi$ scattering amplitude [31, 32] and the connection with chiral symmetry restoration has been established by obtaining the scalar susceptibility through the following saturated approximation [33, 34]

$$\frac{\chi_S(T)}{\chi_S(0)} \simeq \frac{M_S^2(0)}{M_S^2(T)}, \quad (2)$$

where M_S^2 is the real part of the resonance pole $s_p = (M - i\Gamma/2)^2$ of the $\pi\pi$ unitarized scattering amplitude, which plays the role of the scalar $f_0(500)$ $p = 0$ self-energy [34]. In Figure 1 (left) we show the result of a fit to lattice data of the saturated susceptibility within the unitarized ChPT approach, as given in [34]. The fit parameter, denoted A is just a normalization constant, which takes the value $A_{ChPT} = 0.15$ when $\chi_S(0)$ is chosen as the scalar susceptibility calculated in ChPT [35, 36] with the low-energy constants (LEC) used in [34]. These results show that the expected peak of the scalar susceptibility around the transition is well reproduced just with the thermal $f_0(500)$ and the fit

parameter lies within the ChPT value. Actually, even without fitting, the LEC uncertainty covers already the lattice points below T_c . In addition, when comparing this unitarized saturated approach with a HRG calculation there is a significant improvement in the former over the latter, which actually fits numerically the lattice points but does not have a peak profile [34].

A similar approach has been followed for the $I = 1/2$ scalar susceptibility, i.e., that corresponding to the κ channel [37]. In that case the relevant process is πK scattering at finite temperature and the saturated approach is carried out using (2), now for the $K_0^*(700)$ thermal pole. The result is also given in Figure 1 (right) for one of the unitarization methods given in [37] (other methods give similar results). A very interesting conclusion for this channel is that a susceptibility peak also appears, but now significantly above T_c . As we will discuss in section III, the presence of that peak and the behaviour of the susceptibility around is directly linked to the interplay between chiral and $U(1)_A$ restoration, providing direct quantitative information about the role of strangeness and quark masses in that context. Actually, note that as the kaon mass is reduced towards the pion one, the susceptibility peak becomes more pronounced and the peak temperature decreases, approaching T_c . This is nothing but a natural consequence of approaching the regime of degenerated $SU(3)$ flavor symmetry as long as the quark masses are concerned. In such regime, the scalar/pseudoscalar nonet members tend to degenerate. In particular, in that limit the poles of the scalar $K_0^*(700)$ and the octet component of the $f_0(500)$ become degenerate [38] and so they do the π and K pseudoscalar states [39]. Thus, the susceptibility peak in the κ channel inherits naturally the properties of the σ channel one as the kaon mass drops. Another interesting limit displayed also in Figure 1 is the light chiral limit, for which, as we will see below, we expect that the κ -channel peak flattens for temperatures above that peak, due to a more rapid degeneration between the κ and K states.

III. WARD IDENTITIES FOR SUSCEPTIBILITIES AND CONDENSATES

A series of recent works has shown that Ward Identities (WI) derived formally from the QCD generating functional can be used to relate susceptibilities and quark condensates in different channels, which is indeed quite useful regarding the issues commented above for the QCD phase diagram [33, 37, 40–42]. In particular, the following identities turn out to be quite relevant for our present discussion:

$$\chi_P^{ls}(T) = -2 \frac{m_l}{m_s} \chi_{5, disc}(T) \quad (3)$$

$$= -\frac{2}{m_l m_s} \chi_{top}(T) = \frac{1}{2m_l m_s} [m_l \langle \bar{q}q \rangle_l(T) + m_l^2 \chi_P^l], \quad (4)$$

$$\chi_P^K(T) = -\frac{\langle \bar{q}q \rangle_l(T) + 2\langle \bar{s}s \rangle(T)}{m_l + m_s}, \quad (5)$$

$$\chi_S^\kappa(T) = \frac{\langle \bar{q}q \rangle_l(T) - 2\langle \bar{s}s \rangle(T)}{m_s - m_l}. \quad (6)$$

Here, $m_l = m_u = m_d$, χ_P^{ls} is the pseudoscalar crossed $\eta_l \eta_s$ susceptibility with $\eta_s = i\bar{s}\gamma_5 s$, $\chi_{5, disc} = \frac{1}{4} [\chi_P^\pi - \chi_P^l]$ with χ_P^π , χ_P^l , χ_P^K , χ_S^κ the $\pi\pi$, $\eta_l \eta_l$, KK , $\kappa\kappa$ susceptibilities, respectively, and $\chi_{top} = (-1/36)\chi_P^{AA}$ is the topological susceptibility and χ_P^{AA} is the correlator of the anomaly operator $A(x) = \frac{3g^2}{32\pi^2} G_{\mu\nu}^a \tilde{G}_a^{\mu\nu}$ with $G_{\mu\nu}^a$ the gluon field tensor.

The interest of the identity (3) lies in the fact that, as discussed above, the η_l field can be transformed into δ by a chiral $SU(2)_A$ rotation, under which the η_s field remains invariant. Therefore, should the chiral symmetry be exact, χ_P^{ls} would be degenerated with the $p = 0$ $\delta\eta$ correlator, but the latter is odd under parity transformations and therefore vanishes from parity conservation. Thus, if we combine such parity argument with the WI given in eq.(3), one arrives to the conclusion, at least for these channels, that exact chiral symmetry restoration implies exact $U(1)_A$ restoration since $\chi_{5, disc}$ requires *both* symmetries to be restored so that π and η_l degenerate.

The above conclusion based on WI is totally consistent with the result obtained in lattice simulations for $N_f = 2$, which are compatible with $U(1)_A$ restoration close to T_c as the chiral limit is approached. Confirmation of the above results has also been obtained within a recent $U(3)$ ChPT calculation [42]. In that work, degeneration of $U(1)_A$ partners takes place above the chiral ones for physical meson masses, consistently with lattice analyses. As the light chiral limit is approached by reducing the pion mass, one obtains the result showed in Figure 2 for the degeneration temperatures of different partners. Namely, we denote by T_c and T_0 the chiral restoration temperatures corresponding to $\pi - \sigma$ degeneration and the vanishing of $\langle \bar{q}q \rangle_l$, T_{c2} corresponds to the $U(1)_A$ $\pi - \delta$ degeneration, T_{c3} stands for the chiral and $U(1)_A$ restoration of the $\pi - \eta_l$ partners ($\chi_{5, disc}$ vanishing) while $K - \kappa$ channels degenerate at T_{c4} . It becomes pretty clear that all those degeneration temperatures tend to coincide as the light chiral limit is approached, confirming our previous remarks based on WI. On the other hand, a remarkable result, which will help to understand




FIG. 2: Upper left panel: Evolution of different degeneration temperatures with the pion mass, from [42]. Upper right panel: Topological susceptibility at finite temperature from [44]. The lattice points in the right panel correspond to [45, 46] while the bands reflect the LEC uncertainty. Lower panel: Reconstructed kaon and kappa susceptibilities, from the analysis in [37] with condensate lattice points taken from [3].

the role of strangeness, is that $K - \kappa$ degeneration almost coincides with the $\pi - \delta$ one even for the physical pion mass and is therefore linked to $U(1)_A$ degeneration. The latter conclusion has also been confirmed by a PNJL-model analysis of meson screening and pole masses, combined with lattice data [43].

As for the identity (4), it establishes the topological susceptibility as an alternative measure of joint chiral and $U(1)_A$ restoration, providing a neat separation into one contribution proportional to the light quark condensate plus another one where the η_l susceptibility enters. The first contribution is responsible for chiral restoration and dominates for low temperatures and up to the chiral transition, while the second one accounts mostly for the residual $U(1)_A$ breaking above the chiral transition. That behaviour has indeed been reproduced also within $U(3)$ ChPT in [44]. The advantage of that formalism is that it incorporates naturally the effect of η' loops, which turn out to be of the same order as those of π, K, η ones. Actually, the dominant contribution to the topological susceptibility, both at zero and finite temperature, comes from the lightest states, i.e., the pions, as it can be seen already from the leading order ChPT contribution (temperature independent)

$$\chi_{top}^{U(3),LO} = -\frac{1}{2} \langle \bar{q}q \rangle_l^0 \frac{M_0^2 \bar{m}}{M_0^2 + 6B_0 \bar{m}} \quad (7)$$

where $\langle \bar{q}q \rangle_l^0$ is the quark condensate in the chiral limit, M_0 stands for the anomalous part of the η' mass and $\bar{m}^{-1} = \sum_{i=u,d,s} m_i^{-1}$ which shows clearly the dominance of light states. Also for that reason the finite-temperature analysis in this formalism captures reasonably well results from the lattice, even for temperatures well above the ChPT applicability range, as it is shown in Figure 2.

Finally, identities (5)-(6) open up the possibility of considering the $K - \kappa$ sector for the study of the strangeness effect in the interplay between chiral and $U(1)_A$ restoration and explain also the behaviour found for the susceptibility in the kappa channel from the thermal resonance method discussed in section II. Thus, those identities predict a constantly decreasing χ_P^K since both light and strange condensates decrease, but the minus relative sign between those two

condensates in (6) implies that the initially increasing behaviour of χ_S^κ with temperature dominated by the decreasing of (negative) $\langle \bar{q}q \rangle_l$ is taken over by the strange condensate at some point above the chiral transition, from which χ_S^κ would start decreasing tending to degenerate with χ_P^K . Therefore, this argument explains the peak obtained for χ_S^κ in section II and, together with our previous WI and effective theory arguments, suggests that such peak and its behaviour above it is an indication of how $U(1)_A$ is restored above the chiral transition, quantifying the role of strangeness through $\langle \bar{s}s \rangle$. The previous argument is further supported by the lattice, as Figure 2 shows, where the kaon and kappa susceptibilities are reconstructed using identities (5)-(6) and the quark condensates obtained in the lattice collaboration [3]. In particular, from our previous arguments, as the light chiral limit is approached we expect a more rapid growth just below the peak, from chiral restoration, and a flattening just above it from the faster $U(1)_A$ $K - \kappa$ degeneration.

IV. CONCLUSIONS

Light thermal resonances $f_0(500)$ and $K_0^*(700)$ are key to understand chiral and $U(1)_A$ restoration. Combining effective theory approaches with Ward Identities support $U(1)_A$ restoration for fully restored chiral symmetry and allows one to understand the role of quark masses and strangeness in the physical regime of massive quarks and $N_f = 2 + 1$ flavors.

Acknowledgments

Work partially supported by research contract PID2019-106080GB-C21 (“Ministerio de Ciencia e Innovación”), the European Union Horizon 2020 research and innovation program under grant agreement No 824093 and the Swiss National Science Foundation, project No. PZ00P2_174228. A. V-R acknowledges support from a fellowship of the UCM predoctoral program.

-
- [1] Y. Aoki, S. Borsanyi, S. Durr, Z. Fodor, S.D. Katz, S. Krieg, K.K. Szabo, The QCD transition temperature: Results with physical masses in the continuum limit II. JHEP **0906** (2009) 088.
 - [2] S. Borsanyi, Z. Fodor, C. Hoelbling, S.D. Katz, S. Krieg, C. Ratti, K.K. Szabo, Is there still any T_c mystery in lattice QCD? Results with physical masses in the continuum limit III, JHEP **1009** (2010) 073.
 - [3] A. Bazavov, T. Bhattacharya, M. Cheng, C. DeTar, H. T. Ding, S. Gottlieb, R. Gupta and P. Hegde *et al.* [HotQCD Collaboration], The chiral and deconfinement aspects of the QCD transition, Phys. Rev. D **85** (2012) 054503.
 - [4] C. Ratti, Lattice QCD and heavy ion collisions: a review of recent progress, Rept. Prog. Phys. **81** (2018) 084301.
 - [5] A. Bazavov *et al.* [USQCD Collaboration], Hot-dense Lattice QCD: USQCD whitepaper 2018, Eur. Phys. J. A **55** (2019) 194.
 - [6] A. Gómez Nicola, Light quarks at finite temperature: chiral restoration and the fate of the $U(1)_A$ symmetry, Eur. Phys. J. ST **230** (2021) 1645.
 - [7] A. Gómez Nicola, Aspects on Effective Theories and the QCD Transition, Symmetry **12** (2020) 945.
 - [8] F. Karsch, K. Redlich, A. Tawfik, Hadron resonance mass spectrum and lattice QCD thermodynamics, Eur. Phys. J. C **29** (2003) 549.
 - [9] P. Huovinen and P. Petreczky, QCD Equation of State and Hadron Resonance Gas, Nucl. Phys. A **837** (2010) 26.
 - [10] J. Jankowski, D. Blaschke and M. Spalinski, Chiral condensate in hadronic matter, Phys. Rev. D **87** (2013) 105018.
 - [11] A. Andronic, P. Braun-Munzinger, J. Stachel and M. Winn, Interacting hadron resonance gas meets lattice QCD, Phys. Lett. B **718** (2012), 80.
 - [12] P. Huovinen and P. Petreczky, Hadron resonance gas with repulsive interactions and fluctuations of conserved charges, Phys. Lett. B **777** (2018) 125.
 - [13] R. D. Pisarski and F. Wilczek, Remarks on the Chiral Phase Transition in Chromodynamics, Phys. Rev. D **29** (1984) 338.
 - [14] D. J. Gross, R. D. Pisarski and L. G. Yaffe, QCD and Instantons at Finite Temperature, Rev. Mod. Phys. **53** (1981) 43.
 - [15] E. V. Shuryak, Which chiral symmetry is restored in hot QCD?, Comments Nucl. Part. Phys. **21** (1994) 235. [arXiv:hep-ph/9310253 [hep-ph]].
 - [16] T. D. Cohen, The High temperature phase of QCD and $U(1)_A$ symmetry, Phys. Rev. D **54** (1996), R1867-R1870.
 - [17] S. H. Lee and T. Hatsuda, $U_A(1)$ symmetry restoration in QCD with N_f flavors, Phys. Rev. D **54** (1996) R1871.
 - [18] A. Pelissetto and E. Vicari, Relevance of the axial anomaly at the finite-temperature chiral transition in QCD, Phys. Rev. D **88**, 105018 (2013).
 - [19] J. I. Kapusta, D. Kharzeev and L. D. McLerran, The Return of the prodigal Goldstone boson, Phys. Rev. D **53** (1996) 5028.

- [20] T. Csorgo, R. Vertesi and J. Sziklai, Indirect observation of an in-medium η' mass reduction in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions, Phys. Rev. Lett. **105** (2010) 18230.
- [21] A. Y. Kotov, M. P. Lombardo and A. M. Trunin, Fate of the η' in the quark gluon plasma, Phys. Lett. B **794** (2019) 83.
- [22] G. Cossu, S. Aoki, H. Fukaya, S. Hashimoto, T. Kaneko, H. Matsufuru and J. I. Noaki, Finite temperature study of the axial U(1) symmetry on the lattice with overlap fermion formulation, Phys. Rev. D **87** (2013) 114514 [erratum: Phys. Rev. D **88** (2013) 019901].
- [23] A. Tomiya, G. Cossu, S. Aoki, H. Fukaya, S. Hashimoto, T. Kaneko and J. Noaki, Evidence of effective axial U(1) symmetry restoration at high temperature QCD, Phys. Rev. D **96** (2017) 034509.
- [24] B. B. Brandt, A. Francis, H. B. Meyer, O. Philipsen, D. Robaina and H. Wittig, On the strength of the $U_A(1)$ anomaly at the chiral phase transition in $N_f = 2$ QCD, JHEP **12** (2016) 158.
- [25] M. I. Buchoff, M. Cheng, N. H. Christ, H. T. Ding, C. Jung, F. Karsch, Z. Lin, R. D. Mawhinney, S. Mukherjee and P. Petreczky, *et al.*, QCD chiral transition, $U(1)_A$ symmetry and the dirac spectrum using domain wall fermions, Phys. Rev. D **89** (2014) 054514.
- [26] S. M. Berman, Unification of photoproduction and electroproduction, Phys. Rev. **135** (1964) B1249-B1254.
- [27] R. Garcia-Martin, R. Kaminski, J. R. Pelaez and J. Ruiz de Elvira, Phys. Rev. Lett. **107** (2011) 072001.
- [28] P. A. Zyla *et al.* [Particle Data Group], Review of Particle Physics, PTEP **2020** (2020) 083C01.
- [29] J. R. Pelaez, From controversy to precision on the sigma meson: a review on the status of the non-ordinary $f_0(500)$ resonance, Phys. Rept. **658** (2016) 1.
- [30] J. R. Peláez, A. Rodas and J. R. de Elvira, Precision dispersive approaches versus unitarized chiral perturbation theory for the lightest scalar resonances $\sigma/f_0(500)$ and $\kappa/K_0^*(700)$, Eur. Phys. J. ST **230** (2021) 1539.
- [31] A. Gomez Nicola, F. J. Llanes-Estrada and J. R. Pelaez, Finite temperature pion scattering to one loop in chiral perturbation theory, Phys. Lett. B **550** (2002) 55.
- [32] A. Dobado, A. Gomez Nicola, F. J. Llanes-Estrada and J. R. Pelaez, Thermal rho and sigma mesons from chiral symmetry and unitarity, Phys. Rev. C **66** (2002) 055201.
- [33] A. Gomez Nicola, J. Ruiz de Elvira and R. Torres Andres, Chiral Symmetry Restoration and Scalar-Pseudoscalar partners in QCD, Phys. Rev. D **88** (2013) 076007.
- [34] S. Ferreres-Solé, A. Gómez Nicola and A. Vioque-Rodríguez, Role of the thermal $f_0(500)$ in chiral symmetry restoration, Phys. Rev. D **99** (2019) 036018.
- [35] A. Gomez Nicola, J. R. Pelaez and J. Ruiz de Elvira, Non-factorization of four-quark condensates at low energies within Chiral Perturbation Theory, Phys. Rev. D **82** (2010) 074012.
- [36] A. Gomez Nicola, J. R. Pelaez and J. Ruiz de Elvira, Scalar susceptibilities and four-quark condensates in the meson gas within Chiral Perturbation Theory, Phys. Rev. D **87** (2013), 016001.
- [37] A. Gómez Nicola, J. Ruiz de Elvira, A. Vioque-Rodríguez and D. Álvarez-Herrero, The role of strangeness in chiral and $U(1)_A$ restoration, Eur. Phys. J. C **81** (2021) 637.
- [38] J. A. Oller, The Mixing angle of the lightest scalar nonet, Nucl. Phys. A **727** (2003), 353.
- [39] J. Ruiz de Elvira, U. G. Meißner, A. Rusetsky and G. Schierholz, Feynman–Hellmann theorem for resonances and the quest for QCD exotica, Eur. Phys. J. C **77** (2017) 659.
- [40] A. Gómez Nicola and J. Ruiz de Elvira, Pseudoscalar susceptibilities and quark condensates: chiral restoration and lattice screening masses, JHEP **1603** (2016) 186 (2016).
- [41] A. Gómez Nicola and J. Ruiz de Elvira, Patterns and partners for chiral symmetry restoration, Phys. Rev. D **97** (2018) 074016.
- [42] A. Gómez Nicola and J. Ruiz De Elvira, Chiral and $U(1)_A$ restoration for the scalar and pseudoscalar meson nonets, Phys. Rev. D **98** (2018) 014020.
- [43] M. Ishii, H. Kouno and M. Yahiro, Model prediction for temperature dependence of meson pole masses from lattice QCD results on meson screening masses, Phys. Rev. D **95** (2017) 114022.
- [44] A. Gómez Nicola, J. Ruiz De Elvira and A. Vioque-Rodríguez, The QCD topological charge and its thermal dependence: the role of the η' , JHEP **11** (2019) 086.
- [45] C. Bonati, M. D'Elia, M. Mariti, G. Martinelli, M. Mesiti, F. Negro, F. Sanfilippo and G. Villadoro, Axion phenomenology and θ -dependence from $N_f = 2 + 1$ lattice QCD, JHEP **03** (2016) 155.
- [46] S. Borsanyi, Z. Fodor, J. Guenther, K. H. Kampert, S. D. Katz, T. Kawanai, T. G. Kovacs, S. W. Mages, A. Pasztor and F. Pittler, *et al.* Calculation of the axion mass based on high-temperature lattice quantum chromodynamics, Nature **539** (2016) 69-71.