

# New satellites of figure-eight orbit computed with high precision

I. Hristov<sup>1,a</sup>, R. Hristova<sup>1</sup>, I. Puzynin<sup>2</sup>, T. Puzynina<sup>2</sup>, Z. Sharipov<sup>2,b</sup>, Z. Tukhliev<sup>2</sup>

<sup>1</sup> *Faculty of Mathematics and Informatics, Sofia University, Sofia, Bulgaria*

<sup>2</sup> *Meshcheryakov Laboratory of Information Technologies, JINR, Dubna, Russia*

*E-mails:* <sup>a</sup> [ivanh@fmi.uni-sofia.bg](mailto:ivanh@fmi.uni-sofia.bg)    <sup>b</sup> [zarif@jinr.ru](mailto:zarif@jinr.ru)

**Keywords:** Three-body problem, Satellites of figure-eight orbit, Stable choreographies, Modified Newton's method with high precision

**Abstract** In this paper we use a Modified Newton's method based on the Continuous analog of Newton's method and high precision arithmetic for a search of new satellites of the famous figure-eight orbit. By making a purposeful search for such satellites, we found over 300 new satellites, including 7 new stable choreographies.

## 1 Introduction

Numerical algorithms used in finding new periodic orbits are of big importance for further investigation and understanding of the three-body problem. Major progress in this area was made in recent years. In 2013 Shuvakov and Dmitrashinovich found 13 new topological families applying a clever numerical algorithm based on the gradient descent method in the standard double precision arithmetic [1,2]. Since the three-body problem has a sensitive dependence on the initial conditions, working with double precision limits the number of solutions that can be found, especially nonstable solutions. However, a method using double precision arithmetic could be very useful, since it is not time consuming and catches many of the very important stable solutions. In 2017 Li and Liao applied Newton's method to search for new periodic orbits for the planar three-body problem. They used the high order multiple precision Taylor series method to form the linear system at each step of Newton's method and found more than 600 new families of periodic orbits at once [3]. This result shows the great potential of high precision computations for the three-body problem.

The absence of details for the used numerical algorithm in [3] motivated us to present this algorithm in our recent work [4]. In [4] we made a detailed description of Newton's method and also its modification, based on the Continuous analog of

Newton's method [5] for computing periodic orbits of the planar three-body problem. In particular, we gave all needed formulas, based on the rules of the automatic differentiation for the Taylor series method. We tested our programs by making a general search for relatively short periods and a relatively coarse searching grid. As a result we found 33 new topological families, that are not included in the database in [3]. The solutions were computed with 150 correct digits. We concluded that the usage of a relaxed return proximity condition for the initial approximations and the usage of the modified Newton's method instead of the classic Newton's method is a very perspective approach to find more periodic solutions for a given search grid.

In this paper we use the modified Newton's method for a purposeful search of new satellites (topological powers) of the famous figure-eight orbit [6]. There are two known figure-eight solutions: Moore's figure eight orbit [6, 7], which is a stable choreography, and Simo's figure-eight orbit which is also stable, but is not a choreography [8]. A choreography is such a periodic solution, for which all bodies move along one and the same trajectory with a time delay of  $T/3$ , where  $T$  is the period of the solution. Satellites of figure-eight are of special interest as it is expected that some of them are stable periodic orbits and (or) choreographies. Our work is motivated by the works [9, 10], where many new satellites of figure-eight are found, including new choreographies, and particularly one stable choreography, which is until now only the second stable choreography found after the Moore's figure eight orbit. In this work we first make a general search and then concentrate on a small window for initial conditions. As a result we manage to find 427 satellites of the figure-eight orbit (398 new ones), including 7 new stable choreographies.

## 2 Mathematical model

A planar motion of the three bodies is considered. The bodies are with equal masses and are treated as mass points. The normalized differential equations describing the motion of the bodies are:

$$\ddot{r}_i = \sum_{j=1, j \neq i}^3 \frac{(r_j - r_i)}{\|r_i - r_j\|^3}, i = 1, 2, 3. \quad (1)$$

The vectors  $r_i, \dot{r}_i$  have two components:  $r_i = (x_i, y_i), \dot{r}_i = (\dot{x}_i, \dot{y}_i)$ . The system (1) can be written as a first order one this way:

$$\dot{x}_i = vx_i, \dot{y}_i = vy_i, \dot{vx}_i = \sum_{j=1, j \neq i}^3 \frac{(x_j - x_i)}{\|r_i - r_j\|^3}, \dot{vy}_i = \sum_{j=1, j \neq i}^3 \frac{(y_j - y_i)}{\|r_i - r_j\|^3}, i = 1, 2, 3 \quad (2)$$

We numerically solve the problem in this first order form. Hence we have a vector of 12 unknown functions  $u(t) = (x_1, y_1, vx_1, vy_1, x_2, y_2, vx_2, vy_2, x_3, y_3, vx_3, vy_3)^\top$ . We search for periodic planar collisionless orbits as in [1–3]: with zero angular momentum and symmetric initial configuration with parallel velocities:

$$\begin{aligned}
(x_1(0), y_1(0)) &= (-1, 0), \quad (x_2(0), y_2(0)) = (1, 0), \quad (x_3(0), y_3(0)) = (0, 0) \\
(vx_1(0), vy_1(0)) &= (vx_2(0), vy_2(0)) = (v_x, v_y) \\
(vx_3(0), vy_3(0)) &= -2(vx_1(0), vy_1(0)) = (-2v_x, -2v_y)
\end{aligned} \tag{3}$$

Here the velocities  $v_x \in [0, 1]$ ,  $v_y \in [0, 1]$  are parameters. The periods of the orbits are denoted with  $T$ . So, our goal is to find triplets  $(v_x, v_y, T)$  for which the periodicity condition  $u(T) = u(0)$  is fulfilled.

### 3 Description of the numerical procedure

The numerical procedure has two stages. During the first stage we search for candidates for correction with the modified Newton's method, i.e. we compute initial approximations. During the second stage we apply the modified Newton's method, which can converge or diverge. Convergence means that a periodic orbit is found.

A 2D grid with some step is introduced in some rectangular window for parameters  $v_x, v_y$  for the first stage. For every grid point the system (2) with initial conditions (3) is solved numerically up to a pre-fixed time  $T_0$ . Let

$$P(t) = \sqrt{\sum_{i=1}^3 \|r_i(t) - r_i(0)\|^2 + \sum_{i=1}^3 \|\dot{r}_i(t) - \dot{r}_i(0)\|^2}, \quad 0 \leq t \leq T_0$$


be the proximity function. As candidates for the correction method, we take those triplets  $(v_x, v_y, T)$  for which the return proximity  $P(T)$  is less than 0.7:

$$P(T) = \min_{1 < t \leq T_0} P(t) < 0.7$$

and also we want the return proximity be a local minimum on the grid for  $v_x, v_y$ . During the second stage the modified Newton's method is applied for correction of the initial approximations obtained during the first stage. To do this, a linear algebraic system with a  $12 \times 3$  matrix for the corrections  $\Delta v_x, \Delta v_y, \Delta T$  has to be solved [11]. To form the matrix, a system of 36 ODE's (the original differential equations plus the differential equations for the partial derivatives with respect to the parameters  $v_x$  and  $v_y$ ) has to be solved first. The multiple precision Taylor series method, realized by using the rules of automatic differentiation is used during both stages [12, 13]. A variable step-size strategy from [14] is used. The linear algebraic system is solved in linear least square sense using  $QR$  decomposition based on Householder reflections [15]. All the details of the Newton's and the modified Newton's method for the planar three-body problem can be seen in our recent work [4].

In this work we make a purposeful search for figure-eight satellites. We first make a general search in the window  $[0, 0.8] \times [0, 0.8]$  for  $v_x, v_y$  with a grid step  $1/2048$  and  $T_0 = 70$ . After that we add to our found satellites the satellites found in [9, 10, 16] (we use the tables in [9, 10] and the table for sequence V (figure-eight) at [17]). A cluster of the initial velocities' points  $(v_x, v_y)$  is seen, which we

enclose with the rectangle (window)  $[0.1, 0.33] \times [0.49, 0.545]$  (see Fig.1). Then we concentrate on this rectangle with a grid step  $1/4096$  and  $T_0 = 300$ . For both stages of the numerical procedure we use 154-th order of Taylor series and precision of 134 decimal digits. The computations are performed in "Nestum" cluster, Sofia, Bulgaria [18] and "Govorun" supercomputer, Dubna, Russia [19].



**Fig. 1** Initial velocities: 45 black points - previously found satellites, 50 white points - new found satellites after a general search up to  $T_0 = 70$  with a grid-step  $1/2048$

## 4 Results

By adding the solutions found in  $[0.1, 0.33] \times [0.49, 0.545]$  to those found from the general search up to  $T_0 = 70$ , we found all in all 427 figure-eight satellites (including 26 choreographies). 398 of the found satellites are new (not included in [9,10,16]). A topological method from [20] is applied to classify the periodic orbits into families.



Each family corresponds to a different conjugacy class of the free group on two letters (a,b). We use "the free group word reading algorithm" from [2] to obtain the free group elements. Satellites of figure-eight correspond to free group elements  $(abAB)^k$  for some natural power  $k$ . For the found solutions the power  $k$  varies from 4 up to 62, including all values in between.


For each found solution we computed the free group element and the four numbers  $(v_x, v_y, T, T^*)$  with 150 correct digits, where  $T^*$  is the scale-invariant period. The scale-invariant period is defined as  $T^* = T|E|^{\frac{3}{2}}$ , where  $E$  is the energy of our initial configuration:  $E = -2.5 + 3(v_x^2 + v_y^2)$ . Equal  $T^*$  for two different initial conditions means two different representations of the same solution. For computing the numbers  $(v_x, v_y, T, T^*)$  with 150 correct digits we made two additional high precision computations with the classic Newton's method. The first computation was with 220-th order method and 192 decimal digits of precision and the second computation for verification - with 264-th order method and 231 digits of precision. All 427 solutions computed with 150 correct digits and animations in the real  $x - y$  plane for them can be seen in [21].

We made additional numerical tests for the stability of the presented solutions. These tests (although not rigour) show that 87 solutions are most possibly stable and the other solutions are most possibly unstable. The results for stability of each solution can be seen in a table in [21], where the solutions are ordered by  $T^*$ . It is seen from this table that stable solutions tend to be close to each other in the table. The distribution of the initial condition points in the rectangle  $[0.1, 0.33] \times [0.49, 0.545]$  can be seen in Fig.2 (the black points are the stable solutions, the white points - the unstable ones).

An important result is the finding of 7 new stable choreographies which are topological powers of the figure-eight. Two of the choreographies are presented by two different initial conditions (all in all 9 solutions). Until now there were only two stable choreographies - the famous Moore's figure-eight and the Shuvakov's choreography [9] (solutions 24 and 25 in the table in [21]). The numbers  $(v_x, v_y, T, T^*)$  with 17 correct digits and the power  $k$  can be seen in Table 1. The trajectories in the real  $x - y$  plane of these choreographies can be seen in Figures 3 - 9.

$N$	$v_x$	$v_y$	$T$	$T^*$	$k$
1	0.29294799125543721e0	0.52925901352413728e0	0.2058493691047143e3	0.34179371490016304e3	37
2	0.29791817208757154e0	0.53275520940835742e0	0.21032180603055653e3	0.34179379656326958e3	37
3	0.30889168505067089e0	0.52866005593876979e0	0.2119149795956615e3	0.34179379656326958e3	37
4	0.17791680921792612e0	0.52719348724657283e0	0.17814022614781591e3	0.35085277996057519e3	38
5	0.2529928848686255e0	0.51092018996687722e0	0.18632778573173244e3	0.35085277996057519e3	38
6	0.27224186760209537e0	0.524520188831711169e0	0.23223664454023032e3	0.40645290803177171e3	44
7	0.29653206621705968e0	0.52476506719403896e0	0.24274490570242023e3	0.40645421334926255e3	44
8	0.21036388572856618e0	0.51854057011033809e0	0.27480620837192349e3	0.53574606807894121e3	58
9	0.26684473313570582e0	0.52168674682029369e0	0.30063538172570104e3	0.53576756803804106e3	58


**Table 1** Data with 17 correct digits for the 7 stable choreographies




**Fig. 2** Initial velocities for all found solutions (including previously found) in the window  $[0.1, 0.33] \times [0.49, 0.545]$ , black points are the stable solutions, white points - unstable

## 5 Conclusions


A modified Newton's method, based on the Continuous analog of Newton's method and high precision is successfully used in finding more than 300 new satellites of the famous figure-eight orbit, including 7 new stable choreographies. The results show the great potential of the used numerical algorithm in searching for new periodic orbits of the three-body problem.




**Fig. 3** Stable choreography (solution 1)




**Fig. 4** Stable choreography (solution 2)




**Fig. 5** Stable choreography (solution 4)




**Fig. 6** Stable choreography (solution 6)



**Fig. 7** Stable choreography (solution 7)



**Fig. 8** Stable choreography (solution 8)



**Fig. 9** Stable choreography (solution 9)

**Acknowledgements** We thank for the opportunity to use the computational resources of the Nestum cluster, Sofia, Bulgaria and the "Govorun" supercomputer at the Meshcheryakov Laboratory of Information Technologies of JINR, Dubna, Russia. We would also like to thank Veljko Dmitrashinovich from Institute of Physics, Belgrade University, Serbia for a valuable e-mail discussion and advice, and his encouragement to continue our numerical search for new periodic orbits. The work is supported by a grant of the Plenipotentiary Representative of the Republic of Bulgaria at JINR, Dubna, Russia.

## References

1. Shuvakov, Milovan, and Veljko Dmitrashinovich. "Three classes of Newtonian three-body planar periodic orbits." *Physical review letters* 110.11 (2013): 114301.
2. Shuvakov, Milovan, and Veljko Dmitrashinovich. "A guide to hunting periodic three-body orbits." *American Journal of Physics* 82.6 (2014): 609-619.
3. Li, XiaoMing, and ShiJun Liao. "More than six hundred new families of Newtonian periodic planar collisionless three-body orbits." *Science China Physics, Mechanics and Astronomy* 60.12 (2017): 1-7.
4. Hristov, I., Hristova, R., Puzynin, I., et al. "Newton's method for computing periodic orbits of the planar three-body problem." *arXiv preprint arXiv:2111.10839* (2021).

5. Puzynin, I. V., et al. "The generalized continuous analog of Newton's method for the numerical study of some nonlinear quantum-field models." *Physics of Particles and Nuclei* 30.1 (1999): 87-110.
6. Moore, Christopher. "Braids in classical gravity." *Phys. Rev. Lett* 70.24 (1993): 3675-3679.
7. A. Chenciner and R. Montgomery, "A remarkable periodic solution of the three-body problem in the case of equal masses," *Ann. Math.* 152, 881–901 (2000).
8. C. Simo, "Dynamical properties of the figure eight solution of the threebody problem", *Celestial Mechanics*, edited by Alain Chenciner, Richard Cushman, Clark Robinson, and Zhihong Jeff Xia (AMS, Providence, RI, 2002).
9. Shuvakov, Milovan. "Numerical search for periodic solutions in the vicinity of the figure-eight orbit: slaloming around singularities on the shape sphere." *Celestial Mechanics and Dynamical Astronomy* 119.3 (2014): 369-377.
10. Shuvakov, Milovan, and Mitsuru Shibayama. "Three topologically nontrivial choreographic motions of three bodies." *Celestial Mechanics and Dynamical Astronomy* 124.2 (2016): 155-162.
11. Abad, Alberto, Roberto Barrio, and Angeles Dena. "Computing periodic orbits with arbitrary precision." *Physical review E* 84.1 (2011): 016701.
12. Barrio, Roberto, et al. "Breaking the limits: the Taylor series method." *Applied mathematics and computation* 217.20 (2011): 7940-7954.
13. Barrio, Roberto. "Sensitivity analysis of ODEs/DAEs using the Taylor series method." *SIAM Journal on Scientific Computing* 27.6 (2006): 1929-1947.
14. Jorba, Angel, and Maorong Zou. "A software package for the numerical integration of ODEs by means of high-order Taylor methods." *Experimental Mathematics* 14.1 (2005): 99-117.
15. Demmel, James W. *Applied numerical linear algebra*. Society for Industrial and Applied Mathematics, 1997.
16. Dmitrashinovich, V., Hudomal, A., Shibayama, M., Sugita, A. (2018). "Linear stability of periodic three-body orbits with zero angular momentum and topological dependence of Kepler's third law: a numerical test." *Journal of Physics A: Mathematical and Theoretical*, 51(31), 315101.
17. <http://three-body.ipb.ac.rs/sequences.php>
18. <http://hpc-lab.sofiatech.bg/>
19. <http://hlit.jinr.ru/>
20. Montgomery, Richard. "The N-body problem, the braid group, and action-minimizing periodic solutions." *Nonlinearity* 11.2 (1998): 363.
21. <http://db2.fmi.uni-sofia.bg/3body/>