

Hundreds of new satellites of figure-eight orbit computed with high precision

I. Hristov^{1,a}, R. Hristova¹, I. Puzynin², T. Puzynina², Z. Sharipov^{2,b}, Z. Tukhliev²

¹ *Faculty of Mathematics and Informatics, Sofia University, Sofia, Bulgaria*

² *Meshcheryakov Laboratory of Information Technologies, Joint Institute for Nuclear Research, Dubna, Russia*

E-mails: ^a ivanh@fmi.uni-sofia.bg ^b zarif@jinr.ru

Keywords: Three-body problem, Satellites of figure-eight orbit, Modified Newton's method with high precision, Linear stability, Linearly stable choreographies

Abstract Satellites (topological powers) of the famous figure-eight orbit are special periodic solutions of the planar three-body problem. In this paper we use a modified Newton's method based on the Continuous analog of Newton's method and high precision arithmetic for a purposeful numerical search of new satellites of the figure-eight orbit. Over 700 new satellites are found, including 7 new linearly stable choreographies. The linear stability is checked by a high precision computing of the eigenvalues of the monodromy matrices. The initial conditions of all found solutions are given with 150 correct decimal digits.

1 Introduction

Numerical algorithms used in finding new periodic orbits are of big importance for further investigation and understanding of the three-body problem. Major progress in this area was made in recent years. In 2013 Shuvakov and Dmitrashinovich found 13 new topological families of periodic solutions for the planar three-body problem, applying a clever numerical algorithm based on the gradient descent method in the standard double precision arithmetic [1, 2]. Since the three-body problem has a sensitive dependence on the initial conditions, working with double precision limits the number of solutions that can be found, especially nonstable solutions. However, a method using double precision arithmetic could be very useful, since it is not time consuming and catches many of the very important stable solutions. In 2017 Li and Liao applied the Newton's method to search for new periodic orbits of the planar three-body problem. They used the high order multiple precision Taylor series method to form the linear system at each step of Newton's method and found more

than 600 new topological families of periodic orbits [3]. This result shows the great potential of high precision computations for the three-body problem.

The absence of details for the used numerical algorithm in [3] motivated us to present this algorithm in our recent work [4]. In [4] we made a detailed description of the Newton's method and also its modification, based on the Continuous analog of Newton's method [5] for computing periodic orbits of the planar three-body problem. In particular, we gave all needed formulas, based on the rules of the automatic differentiation for the Taylor series method. We tested our programs by making a general search for relatively short periods and a relatively coarse search grid. As a result we found 33 new topological families, that are not included in the database in [3]. The solutions were computed with 150 correct digits. We concluded that the usage of a relaxed return proximity condition for the initial approximations and the usage of the modified Newton's method instead of the classic Newton's method is a very perspective approach to find more periodic solutions for a given search grid.

In this paper we use the presented in [4] modified Newton's method for a purposeful search of new satellites (topological powers) of the famous figure-eight orbit [6]. There are two known figure-eight solutions: Moore's figure eight orbit [6, 7], which is a stable choreography, and Simo's figure-eight orbit which is of hyperbolic-elliptic type and is not a choreography [8]. A choreography is such a periodic solution, for which all bodies move along one and the same trajectory with a time delay of $T/3$, where T is the period of the solution. Satellites of figure-eight are of special interest as it is expected that some of them are stable periodic orbits and (or) choreographies. Our work is motivated by the works [9, 10], where many new satellites of figure-eight with zero angular momentum are found, including new choreographies, and particularly one stable choreography, which is until now only the second stable choreography found after the Moore's figure eight orbit. Many three-body choreographies (345) are also found in [8], but they are with nonzero angular momentum and undetermined topological type.

In this work we first make a general search and then concentrate on a small domain for initial conditions. As a result we manage to find 821 satellites of the figure-eight orbit (778 new ones), including 7 new linearly stable choreographies.

2 Mathematical model

A planar motion of the three bodies is considered. The bodies are with equal masses and are treated as mass points. The normalized differential equations describing the motion of the bodies are:

$$\ddot{r}_i = \sum_{j=1, j \neq i}^3 \frac{(r_j - r_i)}{\|r_i - r_j\|^3}, \quad i = 1, 2, 3. \quad (1)$$

The vectors r_i , \dot{r}_i have two components: $r_i = (x_i, y_i)$, $\dot{r}_i = (\dot{x}_i, \dot{y}_i)$. The system (1) can be written as a first order one this way:

$$\dot{x}_i = vx_i, \quad \dot{y}_i = vy_i, \quad \dot{x}_i = \sum_{j=1, j \neq i}^3 \frac{(x_j - x_i)}{\|r_i - r_j\|^3}, \quad \dot{y}_i = \sum_{j=1, j \neq i}^3 \frac{(y_j - y_i)}{\|r_i - r_j\|^3}, \quad i = 1, 2, 3 \quad (2)$$

We solve numerically the problem in this first order form. Hence we have a vector of 12 unknown functions $X(t) = (x_1, y_1, vx_1, vy_1, x_2, y_2, vx_2, vy_2, x_3, y_3, vx_3, vy_3)^\top$. We search for periodic planar collisionless orbits as in [1–3]: with zero angular momentum and symmetric initial configuration with parallel velocities:

$$\begin{aligned} (x_1(0), y_1(0)) &= (-1, 0), \quad (x_2(0), y_2(0)) = (1, 0), \quad (x_3(0), y_3(0)) = (0, 0) \\ (vx_1(0), vy_1(0)) &= (vx_2(0), vy_2(0)) = (v_x, v_y) \\ (vx_3(0), vy_3(0)) &= -2(vx_1(0), vy_1(0)) = (-2v_x, -2v_y) \end{aligned} \quad (3)$$

Here the velocities $v_x \in [0, 1], v_y \in [0, 1]$ are parameters. The periods of the orbits are denoted with T . So, our goal is to find triplets (v_x, v_y, T) for which the periodicity condition $X(T) = X(0)$ is fulfilled.

3 Description of the numerical searching procedure

The numerical searching procedure for new periodic orbits has three stages. During the first stage we search for candidates for correction with the modified Newton's method, i.e. we compute initial approximations. During the second stage we apply the modified Newton's method, which has a larger domain of convergence than the classic Newton's method. Convergence during the second stage means that a periodic orbit is found. During the third stage we apply the classic Newton's method with a higher precision in order to specify the solutions with more correct digits (150 correct digits in this work).

A square 2D grid with some step is introduced in some 2D domain for parameters v_x, v_y for the first stage. For every grid point the system (2) with initial conditions (3) is solved numerically up to a pre-fixed time T_0 . Let

$$P(t) = \sqrt{\sum_{i=1}^3 \|r_i(t) - r_i(0)\|^2 + \sum_{i=1}^3 \|\dot{r}_i(t) - \dot{r}_i(0)\|^2}, \quad 0 \leq t \leq T_0$$

be the proximity function. As candidates for the correction method, we take those triplets (v_x, v_y, T) for which the return proximity $P(T)$ is less than 0.1:

$$P(T) = \min_{1 < t \leq T_0} P(t) < 0.1$$

We also set the constraint that the return proximity has a local minimum on the grid for v_x, v_y . The modified and the classic Newton's method are applied for correction during the second and the third stage. To do this, a linear algebraic system with a 12×3 matrix for the corrections $\Delta v_x, \Delta v_y, \Delta T$ has to be solved [11]. To form the

matrix, a system of 36 ODEs (the original differential equations plus the differential equations for the partial derivatives with respect to the parameters v_x and v_y) has to be solved first. The linear algebraic system is solved in linear least square sense using *QR* decomposition based on Householder reflections [12]. The multiple precision Taylor series method [13, 14] with a variable step-size strategy from [15] is used during all three stages to solve the corresponding systems of ODEs. During the first and the second stage Taylor series method with order 154 and 134 decimal digits of precision is used. During the third stage we make two computations. The first computation is with 220-th order Taylor series method and 192 digits of precision and the second computation is for verification - with 264-th order method and 231 digits of precision. All the details of the Newton's and the modified Newton's method for the planar three-body problem can be seen in our recent work [4].

A purposeful search for figure-eight satellites is made in this work. We first make a general search in the rectangle $[0, 0.8] \times [0, 0.8]$ for v_x, v_y with a grid step $1/2048$ and $T_0 = 70$. After that we add to our found satellites the satellites found in [9, 10, 16] (we use the tables in [9, 10] and the table for sequence V (figure-eight) at [17]). A cluster of the initial velocities' points (v_x, v_y) is seen, which we enclose with the rectangle $[0.1, 0.33] \times [0.49, 0.545]$ (see Fig. 1). All the points outside of this rectangle are surrounded by circles with radius 0.02. Then we concentrate on this rectangle and the union of circles with a grid step $1/4096$ and $T_0 = 300$. After that we left only those points outside the rectangle $[0.1, 0.33] \times [0.49, 0.545]$ for which we found in their vicinities (circles) some new solutions. Then we again consider the union of vicinities of the left points together with the newly found solutions, i.e. we make a second iteration for searching. All in all we make a search for the rectangle $[0.1, 0.33] \times [0.49, 0.545]$ plus the domain with the curved boundary shown in Fig. 1.

4 Linear stability computations

The linear stability of a given periodic orbit is determined by the eigenvalues λ of the 12×12 monodromy matrix $M(X_{po}; T)$ [18]:

$$M_{ij}[X_{po}; T] = \frac{\partial X_i(T)}{\partial X_j(0)}, \quad M(0) = I$$

The eigenvalues of M come in pairs or quadruplets $(\lambda, \lambda^{-1}, \lambda^*, \lambda^{*-1})$. They are of four types:

- 1) Elliptically stable - $\lambda = \exp(\pm 2\pi i \nu)$, where $\nu > 0$ (real) is the stability angle. In this case the eigenvalues are on the unit circle. Angle ν describes the stable revolution of adjacent trajectories around a periodic orbit.
- 2) Marginally stable - $\lambda = \pm 1$.
- 3) Hyperbolic - $\lambda = \pm \exp(\pm \mu)$, where $\mu > 0$ (real) is the Lyapunov exponent.
- 4) Loxodromic - $\lambda = \exp(\pm \mu \pm i \nu)$, μ, ν (real)




Fig. 1 Initial velocities: 45 black points - previously found satellites, 50 white points - new found satellites after a general search up to $T_0 = 70$ with a grid-step $1/2048$

Eight of the eigenvalues of M are equal to 1 [18]. The other four determine the linear stability. Here we are interested in elliptically stable orbits, i.e. the four eigenvalues to be $\lambda_j = \exp(\pm 2\pi i v_j)$, $j = 1, 2$. The elements of the monodromy matrix M_{ij} are computed in the same way as the partial derivatives in [4] - with the multiple precision Taylor series method using the rules of automatic differentiation. For computing the eigenvalues we use a Multiprecision Computing Toolbox [19] for MATLAB [20]. First the elements of M are obtained with 120 correct digits and then two computations with 70 and 120 digits of precision are made with the toolbox. The four eigenvalues under consideration are verified by a check for matching the first 30 dig-

its of them and the corresponding condition numbers obtained by the two computations (with 70 and 120 digits of precision). The obtained values for Moore's figure-8 are $v_1 = 0.298092529004750122423759217553$, $v_2 = 0.00842272470813137798255957006197$ and for hyperbolic-elliptic Simo's figure-8 are $\lambda = 1.09587441937857035881716297118$, $v = 0.297572835261762177616886515603$. First digits exactly match with given in [8].

5 Results

We found totally 821 figure-eight satellites (including Moore's and Simo's figure-eight orbits and old satellites, and counting different initial conditions as different solutions). 65 of the found solutions are choreographies. 778 of them are new (not included in [9, 10, 16]).

A topological method from [21] is applied to classify the periodic orbits into families. Each family corresponds to a different conjugacy class of the free group on two letters (a,b). We use "the free group word reading algorithm" from [2] to obtain the free group elements. Satellites of figure-eight correspond to free group elements $(abAB)^k$ for some natural power k . For the found solutions the power k varies from 4 up to 62, including all values in between. For each found solution we computed the free group element and the four numbers (v_x, v_y, T, T^*) with 150 correct digits, where T^* is the scale-invariant period. The scale-invariant period is defined as $T^* = T|E|^{\frac{3}{2}}$, where E is the energy of our initial configuration: $E = -2.5 + 3(v_x^2 + v_y^2)$. Equal T^* for two different initial conditions means two different representations of the same solution. All 821 solutions computed with 150 correct digits, plots of the trajectories and animations in the real $x - y$ plane for them can be seen in [22].

We started the investigation for the stability of the presented solutions with simple tests. The simple tests consist of integration of system (2) for the found initial conditions with Taylor series method with order 70 and 57 decimal digits of precision. The solutions that do not escape from the periodic trajectory for a very long time (2000 periods) become candidates for stable solutions. We obtain 115 candidates. Let us mention that if the greatest magnitude of eigenvalues is slightly greater than one, these tests may give false result. Then by computing the eigenvalues of the monodromy matrix we obtain 82 linearly stable orbits (76 new ones). The obtained linearly stable solutions are only among the 115 candidates, which passed the first simple test. The linear stability results for the earlier found satellites of figure eight from [16] are confirmed, i.e. we obtain the same results for linear stability for old solutions as in [16].

The results for linear stability can be seen in a table in [22]. All stability angles $v_{1,2}$ with 30 correct digits for the linearly stable solutions are given in another table in [22]. The distribution of the initial condition points can be seen in Fig. 2 (the black points are the linearly stable solutions, the white points – the unstable ones (more precisely, not confirmed to be linearly stable)). As it is seen from this figure most of the initial condition points for the linearly stable solutions are found in the rectangle $[0.1, 0.33] \times [0.49, 0.545]$ and also these points are clustered.

An important result is the finding of 7 new linearly stable choreographies which are topological powers of the figure-eight. The second and the third of the choreographies are presented by two different initial conditions (all in all we have 9 solutions). Until now there were known only two stable choreographies - the famous Moore's figure-eight and the Shuvakov's choreography [9] (solutions 28 and 29 in the table in [22]). The numbers (v_x, v_y, T, T^*) with 17 correct digits and the power k can be seen in Table 1. The stability angles are given in Table 2. The trajectories in the real $x - y$ plane of these choreographies can be seen in Figures 3 – 9.

N	v_x	v_y	T	T^*	k
1	0.40169296201102497e0	0.53156253772635003e0	0.2706819009639754e3	0.34179371490016304e3	37
2	0.29791817208757154e0	0.53275520940835742e0	0.21032180603055653e3	0.34179379656326958e3	37
3	0.30889168505067089e0	0.52866005593876979e0	0.2119149795956615e3	0.34179379656326958e3	37
4	0.17791680921792612e0	0.52719348724657283e0	0.17814022614781591e3	0.35085277996057519e3	38
5	0.2529928848686255e0	0.51092018996687722e0	0.18632778573173244e3	0.35085277996057519e3	38
6	0.27224186760209537e0	0.52452018883171169e0	0.23223664454023032e3	0.40645290803177171e3	44
7	0.29653206621705968e0	0.52476506719403896e0	0.24274490570242023e3	0.40645421334926255e3	44
8	0.21036388572856618e0	0.51854057011033809e0	0.27480620837192349e3	0.53574606807894121e3	58
9	0.26684473313570582e0	0.52168674682029369e0	0.30063538172570104e3	0.53576756803804106e3	58

Table 1 Data with 17 correct digits for the 7 new linearly stable choreographies

N	v_1	v_2
1	0.486970420626426013258486704454	1.57783479465699085471836250548e-10
2	0.0327953388952792073716032366859	3.22317484874566580526492303709e-06
3	0.0327953388952792073716032366859	3.22317484874566580526492303709e-06
4	0.267131864119191530384832354753	0.127489871368591999957771257333
5	0.267131864119191530384832354753	0.127489871368591999957771257333
6	0.187691189591235263346007262414	2.61504980535709161857184424983e-12
7	0.0937385311958221201020392196211	0.000838678707197693244105921997421
8	0.427373702369188544983378448257	0.00236826906538802133452909107108
9	0.376315052261497474047661511684	4.95737788455128082115378851449e-10

Table 2 Stability angles v_j , $\lambda_j = \exp(\pm 2\pi i v_j)$, $j=1,2$ for the 7 new linearly stable choreographies

The extensive computations for the numerical search are performed in "Nestum" cluster, Sofia, Bulgaria [23] and "Govorun" supercomputer, Dubna, Russia [24]. The GMP library (GNU multiple precision library) [25] for multiple precision floating point arithmetic is used.

6 Conclusions

A modified Newton's method, based on the Continuous analog of Newton's method and high precision is successfully used in finding more than 700 new satellites of the famous figure-eight orbit, including 7 new linearly stable choreographies. The results show the great potential of the used numerical algorithm in searching for new periodic orbits of the three-body problem.

Acknowledgements We thank for the opportunity to use the computational resources of the "Nes-tum" cluster, Sofia, Bulgaria and the "Govorun" supercomputer at the Meshcheryakov Laboratory of Information Technologies of JINR, Dubna, Russia. We would also like to thank Veljko Dmi-trashinovich from Institute of Physics, Belgrade University, Serbia for a valuable e-mail discussion and advice, and his encouragement to continue our numerical search for new periodic orbits. The work is supported by a grant of the Plenipotentiary Representative of the Republic of Bulgaria at JINR, Dubna, Russia.




Fig. 2 Initial velocities for all found solutions (including previously found), black points are the linearly stable solutions, white points - unstable (more precisely, not confirmed to be linearly stable)




Fig. 3 Linearly stable choreography (solution 1)




Fig. 4 Linearly stable choreography (solution 2)




Fig. 5 Linearly stable choreography (solution 4)




Fig. 6 Linearly stable choreography (solution 6)




Fig. 7 Linearly stable choreography (solution 7)




Fig. 8 Linearly stable choreography (solution 8)




Fig. 9 Linearly stable choreography (solution 9)

References

1. Shuvakov, Milovan, and Veljko Dmitrashinovich. "Three classes of Newtonian three-body planar periodic orbits." *Physical review letters* 110.11 (2013): 114301.
2. Shuvakov, Milovan, and Veljko Dmitrashinovich. "A guide to hunting periodic three-body orbits." *American Journal of Physics* 82.6 (2014): 609–619.
3. Li, XiaoMing, and ShiJun Liao. "More than six hundred new families of Newtonian periodic planar collisionless three-body orbits." *Science China Physics, Mechanics and Astronomy* 60.12 (2017): 1–7.
4. Hristov, I., Hristova, R., Puzynin, I., et al. "Newton's method for computing periodic orbits of the planar three-body problem." *arXiv preprint arXiv:2111.10839* (2021).
5. Puzynin, I. V., et al. "The generalized continuous analog of Newton's method for the numerical study of some nonlinear quantum-field models." *Physics of Particles and Nuclei* 30.1 (1999): 87–110.
6. Moore, Christopher. "Braids in classical gravity." *Phys. Rev. Lett* 70.24 (1993): 3675–3679.
7. A. Chenciner and R. Montgomery, "A remarkable periodic solution of the three-body problem in the case of equal masses," *Ann. Math.* 152, 881–901 (2000).

8. C. Simo, "Dynamical properties of the figure eight solution of the threebody problem", *Celestial Mechanics*, edited by Alain Chenciner, Richard Cushman, Clark Robinson, and Zhihong Jeff Xia (AMS, Providence, RI, 2002).
9. Shuvakov, Milovan. "Numerical search for periodic solutions in the vicinity of the figure-eight orbit: slaloming around singularities on the shape sphere." *Celestial Mechanics and Dynamical Astronomy* 119.3 (2014): 369–377.
10. Shuvakov, Milovan, and Mitsuru Shibayama. "Three topologically nontrivial choreographic motions of three bodies." *Celestial Mechanics and Dynamical Astronomy* 124.2 (2016): 155–162.
11. Abad, Alberto, Roberto Barrio, and Angeles Dena. "Computing periodic orbits with arbitrary precision." *Physical review E* 84.1 (2011): 016701.
12. Demmel, James W. *Applied numerical linear algebra*. Society for Industrial and Applied Mathematics, 1997.
13. Barrio, Roberto, et al. "Breaking the limits: the Taylor series method." *Applied mathematics and computation* 217.20 (2011): 7940–7954.
14. Barrio, Roberto. "Sensitivity analysis of ODEs/DAEs using the Taylor series method." *SIAM Journal on Scientific Computing* 27.6 (2006): 1929–1947.
15. Jorba, Angel, and Maorong Zou. "A software package for the numerical integration of ODEs by means of high-order Taylor methods." *Experimental Mathematics* 14.1 (2005): 99–117.
16. Dmitrashinovich, V., Hudomal, A., Shibayama, M., Sugita, A. (2018). "Linear stability of periodic three-body orbits with zero angular momentum and topological dependence of Kepler's third law: a numerical test." *Journal of Physics A: Mathematical and Theoretical*, 51(31), 315101.
17. <http://three-body.ipb.ac.rs/sequences.php>
18. Roberts, Gareth E. "Linear stability analysis of the figure-eight orbit in the three-body problem." *Ergodic Theory and Dynamical Systems* 27.6 (2007): 1947–1963.
19. Advanpix LLC., Multiprecision Computing Toolbox for MATLAB, <http://www.advanpix.com/>, Version 4.8.8.14711, 2022-08-02
20. MATLAB version 9.12.0.1975300 (R2022a) Update 3, The Mathworks, Inc., Natick, Massachusetts, 2021
21. Montgomery, Richard. "The N-body problem, the braid group, and action-minimizing periodic solutions." *Nonlinearity* 11.2 (1998): 363.
22. <http://db2.fmi.uni-sofia.bg/3body821/>
23. <http://hpc-lab.sofiatech.bg/>
24. <http://hlit.jinr.ru/>
25. <https://gmplib.org/>