Explicit expressions for Sapovalov elements in Type A.

Ian M. Musson* Department of Mathematical Sciences University of Wisconsin-Milwaukee email: musson@uwm.edu

March 11, 2022

Abstract

We give explicit expressions for Šapovalov elements in Type A Lie algebras and superalgebras. Explicit expressions were already given in [Mus17] Section 9, using non-commutative determinants, and in fact our first main results, Theorems 2.3 and 2.6 can be viewed as complete expansions of these determinants. But we give new proofs, which seem easier because they avoid induction and cofactor expansion. We also describe Šapovalov elements for $\mathfrak{gl}(m, n)$ with respect to an arbitrary Borel subalgebra in Theorem 3.7, and interpret Šapovalov elements in Type A as determinants of Hessenberg matrices in Theorems 4.5 and 4.9.The exact form of the explicit expressions depends on an ordering on the set of positive roots, and Hessenberg matrices are useful in changing the ordering. Having explicit expressions for Šapovalov elements allows us to give easy proofs of several results in representation theory, see Section 6 and Subsection 4.6.

1 Definition and significance of Sapovalov elements.

2 The Type A Case.

3 Šapovalov elements for arbitrary Borel subalgebras.

4 Šapovalov elements as determinants of Hessenberg matrices.

5 Powers of Sapovalov elements.

6 Survival of Šapovalov elements in factor modules.

1 Definition and significance of Sapovalov elements.

Let \mathfrak{g} be a simple Lie algebra or a contragredient Lie superalgebra with set of simple and positive roots Π and Δ^+ respectively. In the Type A case we consider the cases $\mathfrak{gl}(m)$ and $\mathfrak{gl}(m,n)$. Fix a positive root η and a positive integer m. If η is isotropic, assume m = 1, and if η is odd non-isotropic, assume that m is odd. Let $\overline{\mathbf{P}}(m\eta)$ be the set of partitions of $m\eta$, see [Mus12] Remark 8.4.3, and also (8.4.4) for the notation $e_{-\pi}$ below. Then let $\pi^0 \in \overline{\mathbf{P}}(m\eta)$ be the unique partition of $m\eta$ such that

 $^{^{*}\}mbox{Research}$ partly supported by NSA Grant H98230-12-1-0249, and Simons Foundation grant 318264.

 $\pi^0(\alpha) = 0$ if $\alpha \in \Delta^+ \setminus \Pi$. We say that $\theta = \theta_{\eta,m} \in U(\mathfrak{b}^-)^{-m\eta}$ is a *Šapovalov element* for the pair (η, m) if it has the form

$$\theta = \sum_{\pi \in \overline{\mathbf{P}}(m\eta)} e_{-\pi} H_{\pi}, \tag{1.1}$$

where

$$H_{\pi} \in U(\mathfrak{h}), \ H_{\pi^0} = 1, \tag{1.2}$$

and

$$e_{\alpha}\theta \in U(\mathfrak{g})(h_{\eta} + \rho(h_{\eta}) - m(\eta, \eta)/2) + U(\mathfrak{g})\mathfrak{n}^{+}, \text{ for all } \alpha \in \Delta^{+}.$$
(1.3)

For a semisimple Lie algebra, the existence of such elements was shown by Sapovalov, $[\check{S}ap72]$ Lemma 1. Consider the hyperplane in \mathfrak{h}^* given by

$$\mathcal{H}_{\eta,m} = \{\lambda \in \mathfrak{h}^* | (\lambda + \rho, \eta) = m(\eta, \eta)/2\}.$$
(1.4)

From (1.3), the Sapovalov element $\theta_{\eta,m}$ has the important property that if $\lambda \in \mathcal{H}_{\eta,m}$ then $\theta_{\eta,m}v_{\lambda}$ is a highest weight vector of weight $\lambda - m\eta$ in $M(\lambda)$. The normalization condition $H_{\pi^0} = 1$ guarantees that $\theta_{\eta,m}v_{\lambda}$ is never zero. Hence $xv_{\lambda-m\eta} \longrightarrow x\theta_{\eta,m}v_{\lambda}$ defines a non-zero map between Verma modules $M(\lambda - m\eta) \longrightarrow M(\lambda)$. In the semisimple case, all maps between Vermas are composites of these maps, [Hum08] Theorem 5.1.

Sapovalov elements have also appeared in a number of situations in representation theory, usually in Type A. Though not given this name, they appear in the work of Carter and Lusztig [CL74]. Indeed determinants similar to those in our Theorem 4.5 were introduced in [CL74] Equation (5), and our Theorem 5.2 may be viewed as a version of [CL74] Theorem 2.7. Carter and Lusztig use their result to study tensor powers of the defining representation of GL(V), and homomorphisms between Weyl modules in positive characteristic, see also [CP80] and [Fra88]. Later Carter [Car87] used Šapovalov elements to construct raising and lowering operators for $\mathfrak{sl}(n, \mathbb{C})$, see also [Bru98], [Car95]. In [Car87], these operators are used to construct orthogonal bases for non-integral Verma modules, and all finite dimensional modules for $\mathfrak{sl}(n, \mathbb{C})$.

From the results in Section 5 it is only really necessary to consider the case m = 1, so to simplify notation we set $\mathcal{H}_{\eta} = \mathcal{H}_{\eta,1}$, and denote a Šapovalov element for the pair $(\eta, 1)$ by θ_{η} . In addition in the Type A case, it is only necessary to consider the highest root (resp. highest odd root) of $\mathfrak{gl}(m)$ (resp. $\mathfrak{gl}(m, n)$), since any positive root of the same type is the highest root of some general linear subalgebra in the Lie algebra (resp. superalgebra) case. These two observations greatly simplify our computations.

This paper replaces Sections 9 and 10 of the unpublished manuscript [Mus17]. Earlier sections will be replaced by [Mus22a] and [Mus22b]. I thank Jon Brundan for suggesting that Šapovalov elements could be used to prove Theorem 6.5.

2 The Type A Case.

The main results of this Section are Theorems 2.3 and 2.6. Let $\alpha_i = \epsilon_i - \epsilon_{i+1}$ be the simple roots of $\mathfrak{gl}(m)$ for $1 \leq i \leq m-1$. For the superalgebra case we will also need the simple roots $\gamma_i = \delta_i - \delta_{i+1}$ of $0 \oplus \mathfrak{gl}(n)$ for $1 \leq i \leq n-1$, and $\beta = \epsilon_m - \delta_1$. It may help to remark that

$$e_{\alpha_i} = e_{i,i+1}, \ e_{\beta} = e_{m,m+1}, \ e_{\gamma_j} = e_{m+j,m+j+1},$$

$$h_{\alpha_i} = e_{i,i} - e_{i+1,i+1}, \ h_{\beta} = e_{m,m} + e_{m+1,m+1}, \ h_{\gamma_j} = e_{m+j+1,m+j+1} - e_{m+j,m+j}.$$
(2.1)

For any simple root σ , let $e_{-\sigma}$ be the transpose of e_{σ} . As remarked above, we can assume that $\eta = \epsilon_1 - \epsilon_m$ in the $\mathfrak{gl}(m)$ case, and $\eta = \epsilon_1 - \delta_n$ in the case of $\mathfrak{gl}(m, n)$. The general case follows from this by relabelling the indices. Let $(,)_{m,n}$ be the bilinear form on \mathfrak{h}^* defined by

$$(\epsilon_i, \epsilon_j)_{m,n} = -(\delta_i, \delta_j)_{m,n} = \delta_{i,j} \tag{2.2}$$

for all relevant indices i, j.

Remark 2.1. (a) For a non-isotropic root α , set $\alpha^{\vee} = 2\alpha/(\alpha, \alpha)$. From (2.2), it follows that $(\alpha, \alpha) = 2$, and $\alpha^{\vee} = \alpha$, (resp. $(\gamma, \gamma) = -2$ and $\gamma^{\vee} = -\gamma$) for a simple root α of $\mathfrak{gl}(m) \oplus 0$ (resp. $0 \oplus \mathfrak{gl}(n)$). In the latter case, it is not possible to choose elements $e = e_{\gamma}, f = e_{-\gamma}$ and $h = h_{\gamma} = [e_{\gamma}, e_{-\gamma}]$, such that the elements of e, f, h satisfy the usual defining relations for sl(2), see for example [Mus12] (A.4.2). With the above choices for γ we have $[e_{\gamma}, e_{-\gamma}] = -h_{\gamma}$.

For $\alpha \in \mathfrak{h}^*$, let $h_\alpha \in \mathfrak{h}$ be the unique element such that $(\alpha, \beta) = \beta(h_\alpha)$ for all $\beta \in \mathfrak{h}^*$.

For $k \in [m-1]$, let $\sigma_k = \epsilon_1 - \epsilon_{k+1}$, and

$$h_k = h_{\sigma_k} + (\rho, \sigma_k) - 1.$$
 (2.3)

Lemma 2.2. (a) If α is a simple root of $\mathfrak{g} = \mathfrak{gl}(m, n)$ then $2(\rho, \alpha) = (\alpha, \alpha)$.

(b) For $k \in [m-1]$, $h_k - h_{k-1} = h_{\alpha_k} + 1$ if k > 1, and $h_1 = h_{\alpha_1}$.

Proof. (a) is well known, see for example [Mus12] Corollary 8.5.4. Then (b) follows from (2.3).

Now consider a strictly descending sequence of integers

$$I = \{i_0, i_1, \dots i_s, i_{s+1}\}.$$
(2.4)

If I is a singleton set, then set $f_I = 1$. Otherwise define $f_I \in U(\mathfrak{n}^-)$ by

$$f_I = e_{i_0, 1_1} e_{i_1, i_2} e_{i_2, i_3} \dots e_{i_s, i_{s+1}}.$$
(2.5)

2.1 The case of $\mathfrak{gl}(m)$.

Let

$$\mathbb{I} = \{ I \subseteq [m] | 1, m \in I \},$$
(2.6)

and for $I \in \mathbb{I}$, define

$$r(I) = \{s - 1 | s \in \overline{I}\}$$
(2.7)

where I be the complement of I in I. Recall h_i from (2.3) and define

$$H_J = \prod_{i \in r(J)} h_i.$$
(2.8)

Theorem 2.3. Let $\eta = \epsilon_1 - \epsilon_m$, and suppose v is a highest weight vector in the \mathfrak{g} -Verma module $M(\lambda)$ of weight λ , and set

$$\Theta_{\eta} = \sum_{J \subseteq \mathbb{I}} f_J H_J. \tag{2.9}$$

Then $e_{\alpha_k}\Theta_{\eta}v = 0$ for $k \in [m-2]$. If $(\lambda + \rho, \eta) = 1$, Then $\Theta_{\eta}v$ is a \mathfrak{g} -highest weight vector, and so Θ_{η} is a Sapovalov element for the pair $(\eta, 1)$.

Proof. Clearly subsets of \mathbb{I} correspond to partitions of η , and (1.1) holds. Furthermore the partition π^0 corresponds to $[m] \in \mathbb{I}$ and $e_{\pi^0} = f_{[m]}$. Since $H_{[m]} = 1$, (1.2) holds. We show that $\sum_{J \subseteq \mathbb{I}} f_J H_J$ satisfies Equation (1.3) in the definition of a Šapovalov element. Given a simple root $\alpha_k = \epsilon_k - \epsilon_{k+1}$. To avoid double subscripts we sometimes set $\alpha = \alpha_k$. Thus $e_\alpha = e_{k,k+1}$. The goal is to show that if v is a highest weight vector with weight $\lambda \in \mathcal{H}_\eta$, then $e_\alpha \sum_{J \subseteq \mathbb{I}} f_J H_J v = 0$. Consider the set

$$\mathcal{S}_k = \{ I \subseteq \mathbb{I} | k, k+1 \in I \}.$$

Suppose first $I \in S_k$. In this case $e_{-\alpha}$ is a factor of f_I . There is a unique g such that $k + 1 = i_g$ and $k = i_{g+1}$. With I as in (2.4), $i_0 = m$ and $i_{s+1} = 1$, we set

$$I_1 = \{m, i_1, \dots, i_{g-1}, k+1\}$$
 and $I_2 = \{k, i_{g+2}, \dots, i_s, 1\}.$ (2.10)

Then I is a disjoint union $I = I_1 \cup I_2$, and we have

$$e_{\alpha}f_{I}v = f_{I_{1}}h_{\alpha}f_{I_{2}}v = \begin{cases} f_{I_{1}}h_{\alpha}v & \text{if } I_{2} = \{1\} \text{ is a singleton} \\ f_{I_{1}}f_{I_{2}}(h_{\alpha}+1)v & \text{otherwise.} \end{cases}$$
(2.11)

Define

$$I^+ = I \setminus \{k\}, \text{ and } I^- = I \setminus \{k+1\}.$$
 (2.12)

If k = m - 1, then $I^- \notin \mathbb{I}$ and if k = 1, then $I^+ \notin \mathbb{I}$. We set $f_{I^-} = 0$ or $f_{I^+} = 0$ in these cases. It is easy to check the following result.

Lemma 2.4. We have

(a) $e_{\alpha}f_{I^+}v = f_{I_1}f_{I_2}v$ and $e_{\alpha}f_{I^-}v = -f_{I_1}f_{I_2}v$.

(b)

$$r(I^+) = r(I) \cup \{k-1\} \text{ and } r(I^-) = r(I) \cup \{k\}$$

(c) If $J \subseteq \mathbb{I}$ and $e_{\alpha}f_J v \in f_{I_1}f_{I_2}U(\mathfrak{h})v$, then

$$J = I, I^+ \text{ or } I^-.$$

If I^+ or I^- is not a subset of \mathbb{I} , they should be left out in the above equation.

From (b) in the Lemma and the definition of H_J , we have

$$H_{I^+} = h_{k-1}H_I$$
 and $H_{I^-} = h_kH_I$. (2.13)

Now we apply e_{α} to the sum in (2.9) corresponding to I, I^+ and I^- . Suppose that $k \in [m-2]$. For any highest weight vector v, we have using (2.13), (2.11) and Lemma 2.4,

$$e_{\alpha}(f_{I}H_{I} + f_{I^{+}}H_{I^{+}} + f_{I^{-}}H_{I^{-}})v = \begin{cases} f_{I_{1}}f_{I_{2}}(h_{\alpha_{1}} - h_{1})H_{I}v & \text{if } i = 1\\ f_{I_{1}}f_{I_{2}}(h_{\alpha_{k}} + 1 + h_{k-1} - h_{k})H_{I}v & \text{if } i > 1. \end{cases}$$

$$(2.14)$$

and this is zero by Lemma 2.2, provided that $k \neq m-1$. Now if J does not contain k or k+1, then $e_{\alpha}f_{J}H_{J}v = 0$. Otherwise there exists a set $I \in S_{k}$ such that $J = I, I^{+}$ or I^{-} . Thus we have using (2.14),

$$e_{\alpha} \sum_{J \subseteq \mathbb{I}} f_J H_J v = e_{\alpha} \sum_{I \in \mathcal{S}_k} (f_I H_I + f_{I^+} H_I^+ + f_{I^-} H_I^-) v = 0.$$

This proves the first statement in the Theorem. Now suppose k = m-1, so $f_{I^-} = 0$, and $\lambda \in \mathcal{H}_{\eta}$. For $I \in \mathcal{S}_{m-1}$, we have $e_{\alpha}f_{I}v = h_{\alpha}f_{J}v = f_{J}(h_{\alpha}+1)v$ where $J = I^+ = I \setminus \{m\}$. Set $I^- = I \setminus \{m-1\}$. Then $e_{\alpha}f_{I^-}v = -f_{J}v$. Now by Equation (2.13), $H_{I^+} = h_{m-2}H_I$. Thus using the facts that σ_{m-2} is a sum of m-2 simple roots, and $\alpha_{m-1} + \sigma_{m-2} = \eta$, we have

$$e_{\alpha}(f_{I}H_{I} + f_{I+}H_{I+})v = f_{J}H_{I}(h_{\alpha_{m-1}} + 1 + h_{m-2})v$$

$$= f_{J}H_{I}(h_{\alpha_{m-1}} + h_{\sigma_{m-2}} + m - 2)v$$

$$= f_{J}H_{I}(h_{\eta} + m - 2)v. \qquad (2.15)$$

Now since $\lambda \in \mathcal{H}_{\eta}$ and $(\rho, \eta) = m - 1$, it follows that $(\lambda, \eta) = 2 - m$. Thus the expression in (2.15) is zero.

Remark 2.5. There is a simple method that can be used to determine the linear factors h_k of H_I . Consider the case that I = [m] is as large as possible. Then $H_I = 1$. If $\alpha = \alpha_k$, for $k \in [m-2]$ and k > 1, consider the condition, compare (2.14),

$$0 = e_{\alpha}(f_{I}H_{I} + f_{I^{+}}H_{I^{+}} + f_{I^{-}}H_{I^{-}})v$$

= $f_{I_{1}}f_{I_{2}}(h_{\alpha} + 1 + h_{k-1} - h_{k})v.$ (2.16)

This gives a recurrence relation on the coefficients h_k . Furthermore when k = 1, there are only two terms in the above equation, and this gives h_1 . This remark can be applied to other situations, in particular we used it to develop the proof of Theorem 3.7, see Lemma 3.5. The recurence in (2.16) is obtained by working from left to right along the Dynkin diagram. We could have instead worked from right to left, and obtained an different expression $\sum_{J\subseteq I} f_J H'_J$ for the Šapovalov element θ_η . (Recall that by [Mus17] Theorem 2.1, Šapovalov elements are only unique modulo a left ideal.) However for $\lambda \in \mathcal{H}_\eta$, $\sum_{J\subseteq I} f_J H_J(\lambda) = \sum_{J\subseteq I} f_J H'_J(\lambda)$. Thus both expressions give rise to the same highest weight vector in $M(\lambda)$. In the super case, Theorem 2.6 we work from both ends of the diagram towards the unique grey node. In all the proofs the assumption $\lambda \in \mathcal{H}_\eta$ is used only in the last step.

2.2 The case of $\mathfrak{gl}(m, n)$.

For $i \in [n-1]$, set $\tau_i = \delta_i - \delta_n$. Let $\mathbb{I} = \{I \subseteq [m+n] | 1, m+n \in I\}$. We consider I a strictly descending sequence of integers as in Equation (2.4), and define the corresponding element $f_I \in U(\mathfrak{n}^-)$ as in (2.5). We use the set \mathbb{I} to index an expression for the Šapovalov element $\theta_\eta \in U(\mathfrak{b}^-)$ analogous to that given in Theorem 2.3. For $k \in [m+n-1]$, consider the set

$$\mathcal{S}_k = \{ I \subseteq \mathbb{I} | k, k+1 \in I \}.$$

Define as before,

$$r(I) = \{s - 1 | s \in \overline{I}\}$$
(2.17)

where \overline{I} be the complement of I in \mathbb{I} . Also let I^{\pm} be as in (2.12) If k = m + n - 1, then $I^- \notin \mathbb{I}$ and if k = 1, then $I^+ \notin \mathbb{I}$. As before, set $f_{I^-} = 0$ or $f_{I^+} = 0$ in these cases. Next set

$$h_{i} = \begin{cases} h_{\sigma_{i}} + (\rho, \sigma_{i}) - 1 & \text{if } i \in [m-1] \\ h_{\tau_{i+1-m}} + (\rho, \tau_{i+1-m}) & \text{if } m \le i \le m+n-2. \end{cases}$$
(2.18)

Then we have an analog of Lemma 2.2 (b),

$$h_{m+j-1} - h_{m+j} = h_{\gamma_j} - 1 \text{ for } j \in [n-2].$$
 (2.19)

If we set $h_{m+n-1} = 0$, then (2.19) holds also for j = n - 1. Now define

$$H_J = \prod_{i \in r(J)} h_i.$$

Theorem 2.6. Suppose v is a highest weight vector in the g-Verma module $M(\lambda)$ of weight λ , and set

$$\Theta_{\eta} = \sum_{J \subseteq \mathbb{I}} f_J H_J.$$

Then $\Theta_{\eta}v$ is a \mathfrak{g}_0 -highest weight vector. If $(\lambda + \rho, \eta) = 0$, then $\Theta_{\eta}v$ is a \mathfrak{g} -highest weight vector, and so Θ_{η} is a Šapovalov element for the pair $(\eta, 1)$.

Proof. This follows the same general pattern as the proof of Theorem 2.3, but there are several differences resulting from Remark 2.1, so we give some details. Suppose $I \in \mathcal{S}_k$. First if $\alpha = \alpha_k$ for $k \in [m-1]$ the proof of (2.14) shows that

$$e_{\alpha}(f_{I}H_{I} + f_{I^{+}}H_{I^{+}} + f_{I^{-}}H_{I^{-}})v = 0.$$

Next suppose that k = m + j for $j \in [n - 1]$, and set $\alpha = \gamma_j$, see (2.1). Instead of (2.10), we set

$$I_1 = \{m + n - 1, i_1, \dots i_{g-1}, k+1\} \text{ and } I_2 = \{k, i_{g+2}, \dots i_s, 1\}$$
(2.20)

Note that $e_{-\alpha}$ is a factor of f_I . Thus from Remark 2.1, we have instead of (2.11),

$$e_{\alpha}f_{I}v = -f_{I_{1}}h_{\alpha}f_{I_{2}}v = f_{I_{1}}f_{I_{2}}(1-h_{\alpha})v.$$
(2.21)

By (2.13), we have $H_{I^+} = h_{m+j-1}H_I$ and $H_{I^-} = h_{m+j}H_I$. Note that Lemma 2.4 still holds in this situation. Now in place of (2.14) we have using (2.21),

$$e_{\alpha}(f_{I}H_{I} + f_{I^{+}}H_{I^{+}} + f_{I^{-}}H_{I^{-}})v = f_{I_{1}}f_{I_{2}}(1 - h_{\gamma_{j}} + h_{m+j-1} - h_{m+j})H_{I}v \quad (2.22)$$

and this is zero by (2.19). This proves the first statement of the Theorem.

Finally suppose that k = m so that α is the odd simple root and $e_{\alpha} = e_{m,m+1}$. Fix $I \in S_m$. Then f_{I^+} (resp. f_{I^-}) contains the odd root vector $e_{m+1,i_{g+2}}$ (resp. $e_{i_{g-1},m}$) as a factor. Hence

$$e_{\alpha}f_{I^{+}}H_{I^{+}}v = f_{I_{1}}f_{I_{2}}h_{m-1}H_{I}v$$
 and $e_{\alpha}f_{I^{-}}H_{I^{-}}v = f_{I_{1}}f_{I_{2}}h_{m}H_{I}v$.

On the other hand

$$e_{\alpha}f_{I}H_{I}v = f_{I_{1}}h_{\alpha}f_{I_{2}}H_{I}v = f_{I_{1}}f_{I_{2}}(1+h_{\alpha})H_{I}v.$$

Therefore by (2.18), we have for $\lambda \in \mathcal{H}_{\eta}$,

$$e_{\alpha}(f_{I}H_{I} + f_{I^{+}}H_{I^{+}} + f_{I^{-}}H_{I^{-}})v = f_{I_{1}}f_{I_{2}}(1 + h_{\alpha} + h_{m-1} + h_{m})H_{I}v$$

$$= f_{I_{1}}f_{I_{2}}(h_{\alpha} + h_{\sigma_{m-1}} + h_{\tau_{1}} + (\rho, \sigma_{m-1} + \tau_{1}))H_{I}v.$$

$$= f_{I_{1}}f_{I_{2}}(h_{\eta} + (\rho, \eta))H_{I}v.$$

$$= 0.$$

3 Sapovalov elements for arbitrary Borel subalgebras.

Let $\mathfrak{b}^{\text{dist}}$ denote the distinguished Borel subalgebra of $\mathfrak{gl}(m, n)$. We denote the set of Borel subalgebras with the same even part as $\mathfrak{b}^{\text{dist}}$ by \mathcal{B} . For $\mathfrak{b} \in \mathcal{B}$ and η a positive root of both $\mathfrak{b}^{\text{dist}}$ and \mathfrak{b} we describe the Šapovalov elements with respect to \mathfrak{b} . In [Mus12] 3.3, the set \mathcal{B} is described in terms of shuffles. Here the notation is slightly different. We write a permutation σ of the set $||m+n|| := \{1, 2, ..., m, 1', 2', ..., n'\}$ in one-line notation as

$$\underline{\sigma} = (\sigma(1), \sigma(2), \dots, \sigma(m), \sigma(1'), \sigma(2'), \dots, \sigma(n')).$$

The last *n* entries of ||m + n|| are called *primed*, the others *unprimed*. The shuffle condition is that both $1, 2, \ldots, m$ and $1', 2', \ldots, n'$ are subsequences of $\underline{\sigma}$. A useful way to think about this is by using the Dynkin-Kac diagram of \mathfrak{b} , see [Mus12] section 3.4. The simple roots $\alpha_1, \ldots, \alpha_{m+n-1}$ are used to label the nodes from left to right. Then augment the diagram by adding the entries of $\underline{\sigma}$, in order above the diagram between the nodes, and at both ends. Without loss we assume that $\eta = \epsilon_1 - \delta_n$ is the longest odd positive root of $\mathfrak{g} = \mathfrak{gl}(m, n)$, since any root which is positive in both $\mathfrak{b}^{\text{dist}}$ and \mathfrak{b} is the longest root in a possibly smaller Type A superalgebra. This assumption implies that $\sigma(1) = 1$ and $\sigma(n') = n'$. From the diagram we can immediately give an expression any simple root. Each simple root α_i lies between a pair of neighbors in $\underline{\sigma}$, and there are 4 posibilities. The rule is

if the neighbors of
$$\alpha_i$$
 are $(a, a + 1), (a, b'), (a', b), (b', b + 1'),$
then the root α_i is respectively $\epsilon_a - \epsilon_{a+1}, \epsilon_a - \delta_b, \delta_a - \epsilon_b, \delta_b - \delta_{b+1}.$

Example 3.1. We give an example of an augmented Dynkin-Kac diagram when m = 4, n = 3.

$$1 \qquad 1' \qquad 2 \qquad 3 \qquad 4 \qquad 2' \qquad 3' \\ \otimes \qquad \otimes \qquad \odot \qquad \odot \qquad \odot \qquad \odot \qquad \odot \\ \alpha_1 \qquad \alpha_2 \qquad \alpha_3 \qquad \alpha_4 \qquad \alpha_5 \qquad \alpha_6$$

In this example

$$\alpha_1 = \epsilon_1 - \delta_1, \ \alpha_2 = \delta_1 - \epsilon_2, \ \alpha_3 = \epsilon_2 - \epsilon_3, \ \alpha_4 = \epsilon_3 - \epsilon_4, \ \alpha_5 = \epsilon_4 - \delta_2, \ \alpha_6 = \delta_2 - \delta_3.$$

In a similar way we can read off any positive root from the diagram, and it follows that $\sum_{k=1}^{m+n-1} \alpha_k = \eta$. Under the isomorphism $\mathfrak{h}^* \longrightarrow \mathfrak{h}$, suppose α_k maps to h_k . We deduce that m+n-1

$$\sum_{k=1}^{+n-1} h_k = h_\eta.$$
(3.1)

Let $\mathbb{I} = \{I \subseteq ||m+n|| | 1, n' \in I\}$. The shuffle σ induces a decreasing order on the set ||m+n||. Explicitly

$$\sigma(n') = n' > \sigma(n-1') > \ldots > \sigma(2) > \sigma(1) = 1.$$

and for $I \in \mathbb{I}$, $J \subseteq I$, we use this order to form the products f_J by analogy with Equation (2.5).

Now fix a simple root $\alpha = \alpha_k$ and let i(k) (resp. j(k)) be the number of grey

nodes to the left (resp. right) of α in the diagram. If a and b are the left and right neighbors of α respectively, define

$$S_{\alpha} = \{ I \in \mathbb{I} | a, b \in I \},\$$

and note that a and b can be primed or unprimed numbers. For $I \in \mathbb{I}$ we can write

$$I = \{n', \dots c, b, a, *, \dots, 1\}$$
(3.2)

as a disjoint union, $I_1 \cup I_2$, where

$$I_1 = \{n', \dots, b\}$$
 and $I_2 = \{a, \dots, 1\}.$ (3.3)

The bilnear form $(,)_{m,n}$ on \mathfrak{h}^* defined in (2.2) induces an isomorphism $\mathfrak{h}^* \longrightarrow \mathfrak{h}$ with $\alpha \mapsto h_{\alpha}$. On the other hand there is a bilinear form on \mathfrak{h} induced by the supertrace Str, that is $(h, h') \mapsto \operatorname{Str}(hh')$. These are related by

$$(\alpha,\beta) = \operatorname{Str}(h_{\alpha}h_{\beta}). \tag{3.4}$$

We need an explicit expression for $h_k = h_{\alpha_k}$, in terms of the neighbors of $\alpha_k = \alpha$. For α is even, we already know the answer,

$$h_k = (-1)^{i(k)} (e_{aa} - e_{bb}). aga{3.5}$$

Indeed α is a root of $\mathfrak{gl}(m) \oplus 0$ if i(k) is even, and a root of $0 \oplus \mathfrak{gl}(n)$ if i(k) is odd. Thus (3.5) follows from Remark 2.1.

Lemma 3.2. If α is odd, then

(a)

$$h_k = (-1)^{i(k)} (e_{aa} + e_{bb}). aga{3.6}$$

(b)

$$[e_{\alpha}, e_{-\alpha}] = (-1)^{i(k)} h_k. \tag{3.7}$$

Proof. (a) It seems that the easiest way to proceed is to take (3.6) as the definition, and then show that (3.4) holds. Consider the Cartan matrix (α_k, α_j) . Suppose first that $k \neq 1, m + n - 1$. The row indexed by α_k has consecutive entries -1, 0, 1 (resp. 1, 0, -1) if i(k) is even (resp. odd) with 0 on the main diagonal. All other entries in this row are zero. If k = 1 (resp. k = m + n - 1), the first (resp. last) entry in the above sequences must be deleted, since 0 is on the main diagonal. We show that the same holds for the Gram matrix of the bilinear form on \mathfrak{h} induced by the supertrace form, $(h, h') \mapsto \operatorname{Str}(hh')$. For example if i(k) is even, then a is unprimed. If α_{k-1} is even, then $h_{k-1} = e_{**} - e_{aa}$ where * is the entry in I to the right of a, see (3.2). Then $\operatorname{Str}(h_{k-1}h_k) = -1$. If α_{k-1} is odd, then $h_{k-1} = (-1)^{i(k-1)}(e_{**} + e_{aa})$ and i(k-1) is odd. Again we obtain $\operatorname{Str}(h_{k-1}h_k) = -1$. Similarly $\operatorname{Str}(h_k h_{k+1}) = 1$. We leave the cases where i(k) is odd, k = 1 or k = m + n - 1 to the reader.

(b) This is an easy computation.

Note also that

$$(e_{aa} \pm e_{bb})f_{I_2} = f_{I_2}(e_{aa} \pm e_{bb} + 1)$$
 if I_2 is not a singleton. (3.8)

Lemma 3.3. Suppose that v is a highest weight vector, and $I \in S_{\alpha}$. Then

(a) If α is even we have

$$e_{\alpha}f_{I}v = f_{I_{1}}[e_{\alpha}, e_{-\alpha}]f_{I_{2}}v = \begin{cases} (-1)^{i(k)}f_{I_{1}}h_{k}v & \text{if } I_{2} = \{1\} \text{ is a singleton} \\ f_{I_{1}}f_{I_{2}}(1+(-1)^{i(k)}h_{k})v & \text{otherwise.} \end{cases}$$
(3.9)

(b) If α is odd we have

$$e_{\alpha}f_{I}v = (-1)^{j(k)}f_{I_{1}}[e_{\alpha}, e_{-\alpha}]f_{I_{2}}v = \begin{cases} f_{I_{1}}h_{k}v & \text{if } I_{2} = \{1\} \text{ is a singleton} \\ f_{I_{1}}f_{I_{2}}(h_{k} + (-1)^{i(k)})v & \text{otherwise.} \end{cases}$$
(3.10)

Proof. (a) The first equality follows since e_{α} commutes with f_{I_1} and then the second follows from (3.5) and (3.8).

(b) If α is odd then e_{α} anticommutes with j(k) factors in f_{I_1} and commutes with the others. The condition that $\sigma(1) = 1$ and $\sigma(n') = n'$ implies that the total number of odd simple roots is odd, so if α is odd, then i(k) + j(k) is even. Hence the second equality holds by Lemma 3.2 (b) and (3.8).

Next for $I \in S_{\alpha}$, set

$$I^+ = I \setminus \{a\}, \text{ and } I^- = I \setminus \{b\}.$$

$$(3.11)$$

If $I^{\pm} \notin \mathbb{I}$, then as before, we set $f_{I^{\pm}} = 0$. Then we have an analog of Lemma 2.4.

Lemma 3.4. We have

- (a) If α is even, then $e_{\alpha}f_{I^+}v = f_{I_1}f_{I_2}v$ and $e_{\alpha}f_{I^-}v = -f_{I_1}f_{I_2}v$.
- (b) If α is odd, then

$$e_{\alpha}f_{I^{-}}v = -(-1)^{i(k+1)}f_{I_{1}}f_{I_{2}}v, \qquad (3.12)$$

and

$$e_{\alpha}f_{I^{+}}v = (-1)^{j(k)}f_{I_{1}}f_{I_{2}}v = (-1)^{i(k)}f_{I_{1}}f_{I_{2}}v.$$
(3.13)

Furthermore Lemma 2.4 (c) holds.

Proof. (a) If α is even, then "even rules" apply to the commutator of e_{α} and any root vector which is a factor of $f_{I^{\pm}}$. Thus we get the same result as Lemma 2.4 (a). (b) We prove (3.12). Define

$$I_1^- = \{n', \dots, c\} \text{ and } I_2^- = \{a, \dots, 1\} = I_2$$
 (3.14)

Then $f_{I^-} = f_{I_1^-} e_{ca} f_{I_2^-}$, so

$$e_{\alpha}f_{I^{-}}v = (-1)^{j(k+1)}f_{I_{1}^{-}}e_{ab}e_{ca}f_{I_{2}^{-}}v$$

=
$$\begin{cases} (-1)^{j(k+1)}f_{I_{1}}f_{I_{2}}v & \text{if } \alpha_{k+1} \text{ is even} \\ (-1)^{j(k+1)+1}f_{I_{1}}f_{I_{2}}v & \text{if } \alpha_{k+1} \text{ is odd.} \end{cases}$$

The first equality holds since there are exactly j(k+1) factors in $f_{I_1^-}$ that are odd root vectors. For the second, note that if α_{k+1} is even (resp. odd), then e_{ca} is an odd (resp. even) root vector, and hence $[e_{ab}, e_{ca}] = e_{cb}$ (resp. $[e_{ab}, e_{ca}] = -e_{cb}$). Now if α_{k+1} is even, then mod 2, $j(k+1) \equiv i(k+1) + 1$, while if α_{k+1} is odd, then $j(k+1) + 1 \equiv i(k+1) + 1$. This gives (3.12).

The proof of (3.13) is similar to (3.12) but easier. Note that, if

$$I_1^+ = \{n', \dots, b\} = I_1 \text{ and } I_2^+ = \{*, \dots, 1\},$$
 (3.15)

we have $f_{I+}v = f_{I_1^+}e_{b*}f_{I_2^+}v$. The last statement in the Lemma is easy to prove. Recall h_k defined by (3.5), and set $s_k = h_1 + \ldots + h_k = s_{k-1} + h_k$. Next let

 $\ell(k) = |i \in \{[k-1] | \alpha_i \text{ is a root of } 0 \oplus \mathfrak{gl}(n)\}| - |i \in \{[k-1] | \alpha_i \text{ is a root of } \mathfrak{gl}(m) \oplus 0\}|,$

and define $\bar{i}(k) = (1 - (-1)^{i(k)})/2$. Thus $\bar{i}(k) \in \{0, 1\}$, and $\bar{i}(k) \equiv i(k) \mod 2$. Set $d_k = \ell(k) + \bar{i}(k)$. Thus $d_1 = 0$.

Lemma 3.5. If $k \in [m + n - 2]$, we have the following recurrence,

$$d_{k+1} = d_k - (-1)^{i(k+1)}$$

Proof. We have to show that

$$\ell(k+1) + \overline{i}(k+1) = \ell(k) + \overline{i}(k) - (-1)^{i(k+1)}.$$
(3.16)

If α_k is even, then i(k+1) = i(k). Thus (3.16) becomes

$$\ell(k+1) = \ell(k) - (-1)^{i(k)}.$$
(3.17)

Now if i(k) is even (resp. odd), then α_k is a root of $\mathfrak{gl}(m) \oplus 0$ (resp. $0 \oplus \mathfrak{gl}(n)$) so $\ell(k+1) = \ell(k) - 1$ (resp. $\ell(k+1) = \ell(k) + 1$). This gives (3.17).

If α_k is odd, then $\ell(k+1) = \ell(k)$, and i(k+1) = i(k) + 1, so we have to show

$$\overline{i}(k+1) = \overline{i}(k) + (-1)^{i(k)}.$$

This follows easily by considering the cases, i(k) even and odd separately.

Lemma 3.6. $d_{m+n-1} = -(\rho, \eta).$

Proof. Let r (resp. s) be the total number of roots in the Dynkin-Kac diagram belonging to $0 \oplus \mathfrak{gl}(n)$ (resp. $\mathfrak{gl}(m) \oplus 0$). Thus $(\rho, \eta) = -r + s$ by Remark 2.1 and Lemma 2.2 (a). If $\alpha = \alpha_{m+n-1}$ is even, then it is a root of $0 \oplus \mathfrak{gl}(n)$. Thus $\ell(m+n-1) = r-1-s$. Also i(m+n-1) is odd, so $d_{m+n-1} = \ell(m+n-1)+1 = r-s$. If α is odd, then i(m+n-1) is even, so $d_{m+n-1} = \ell(m+n-1) = r-s$. This gives the result.

If (a, b) are the neighbors of α_k , set

$$h_{\hat{a}} = (-1)^{i(k)} (s_{k-1} - d_{k-1}) \text{ and } h_{\hat{b}} = (-1)^{i(k+1)} (s_k - d_k).$$
 (3.18)

Note that b is also the left neighbor of α_{k+1} , and Equation (3.18) is consistent with this fact. Also (3.18) is very convenient for the proof of the main result because of (3.19) below. However we need another ingredient to define the coefficients H_I . If e is the right neighbor of α_i , then set $\mathbf{r}(e) = j$. Then define

$$t_e = (-1)^{i(\mathbf{r}(e)+1)} (s_{\mathbf{r}(e)} - d_{\mathbf{r}(e)}),$$

and $H_I = \prod_{e \in \overline{I}} t_e$, where \overline{I} is the complement of I in ||m+n||. For example $\mathbf{r}(b) = k$, so by (3.18) $t_b = h_{\hat{b}}$. If (a, b) are as above, it follows from (3.11) that

$$H_{I^+} = h_{\hat{a}} H_I \text{ and } H_{I^-} = h_{\hat{b}} H_I.$$
 (3.19)

Theorem 3.7. Set

$$\Theta_{\eta} = \sum_{J \subseteq \mathbb{I}} f_J H_J.$$

Then Θ_{η} is a Sapovalov element θ_{η} for the pair $(\eta, 1)$ for the Borel subalgebra \mathfrak{b} .

Proof. Assume first that $k \neq 1, m + n - 1$. If α is even then, i(k+1) = i(k). Hence by (3.9) and Lemma 3.4 (a), then (3.18) and Lemma 3.5,

$$e_{\alpha}(f_{I}H_{I} + f_{I^{+}}H_{I^{+}} + f_{I^{-}}H_{I^{-}})v = f_{I_{1}}f_{I_{2}}(1 + (-1)^{i(k)}h_{k} + h_{\hat{a}} - h_{\hat{b}})H_{I}v$$

$$= (-1)^{i(k)}f_{I_{1}}f_{I_{2}}(h_{k} + s_{k-1} - s_{k})H_{I}v$$

$$+ f_{I_{1}}f_{I_{2}}(1 - (-1)^{i(k)}d_{k-1} + (-1)^{i(k)}d_{k})H_{I}v$$

$$= 0.$$
(3.20)

If k = 1, then I_2 is a singleton and $d_1 = i(2) = 0$, so by (3.9), (3.13) and (3.18),

$$e_{\alpha}(f_{I}H_{I} + f_{I} - H_{I} -)v = f_{I_{1}}(h_{1} - h_{\hat{b}})H_{I}v$$

$$= f_{I_{1}}(h_{1} - s_{1} + d_{1})H_{I}v$$

$$= 0.$$

Now assume α is odd and $k \neq 1, m + n - 1$. Then, by Lemma 3.3 (b), (3.13) and (3.12), (3.19), then (3.18) and Lemma 3.5,

$$e_{\alpha}(f_{I}H_{I} + f_{I^{+}}H_{I^{+}} + f_{I^{-}}H_{I^{-}})v = f_{I_{1}}f_{I_{2}}(h_{k} + (-1)^{i(k)} + (-1)^{i(k)}h_{\hat{a}} - (-1)^{i(k+1)}h_{\hat{b}})H_{I}v$$

$$= f_{I_{1}}f_{I_{2}}((h_{k} + s_{k-1} - s_{k})H_{I}v$$

$$+ f_{I_{1}}f_{I_{2}}((-1)^{i(k)} - d_{k-1} + d_{k})H_{I}v$$

$$= 0.$$
(3.21)

If k = 1, then by (3.10), (3.12) and (3.18)

$$e_{\alpha}(f_{I}H_{I} + f_{I^{-}}H_{I^{-}})v = f_{I_{1}}(h_{1} + -(-1)^{i(k+1)}h_{\hat{b}})H_{I}v$$

= $f_{I_{1}}((h_{1} - (s_{1} - d_{1}))H_{I}v$
= $0.$

Now suppose that k = m + n - 1, and that v has weight $\lambda \in \mathcal{H}_{\eta}$. If α is even, then i(k) is odd. In the sequence of equalities below the first follows from (3.9) and (3.19), the second from (3.18). Then the third follows from (3.1) and Lemma 3.5, the fourth from Lemma 3.6 and the final equality holds since $\lambda \in \mathcal{H}_{\eta}$.

$$e_{\alpha}(f_{I}H_{I} + f_{I^{+}}H_{I^{+}})v = f_{I_{2}}(1 - h_{k} + h_{\hat{a}})H_{I}v$$

$$= f_{I_{2}}(1 - h_{k} - s_{k-1} + d_{k-1})H_{I}v$$

$$= -f_{I_{2}}(h_{\eta} - d_{m+n-1})H_{I}v$$

$$= -f_{I_{2}}(h_{\eta} + (\rho, \eta))H_{I}v = 0.$$

Suppose α is odd. Then i(k) is even. For the first two below we use (3.9), (3.13), (3.19) and (3.18). The remainder of the proof is similar to the case where α is even.

$$e_{\alpha}(f_{I}H_{I} + f_{I+}H_{I+})v = f_{I_{2}}((-1)^{i(k)} + h_{k} + (-1)^{i(k)}h_{\hat{a}})H_{I}.$$

$$= f_{I_{2}}(1 + h_{k} + s_{m+n-2} - d_{m+n-2})H_{I}$$

$$= f_{I_{2}}(h_{\eta} - d_{m+n-1})H_{I}$$

$$= f_{I_{2}}(h_{\eta} + (\rho, \eta))H_{I}v = 0.$$

4 Šapovalov elements as determinants of Hessenberg matrices.

4.1 Hessenberg Matrices.

An $n \times n$ matrix $B = (b_{ij})$ is (upper) Hessenberg of order n if $b_{ij} = 0$ unless $i \leq j + 1$. The only assumption on the Hessenberg matrix B is that the entries on the subdiagonal commute with all other entries. We define a noncommutative determinant of the $n \times n$ matrix $B = (b_{ij})$, working from left to right, by

$$\overrightarrow{\det}(B) = \sum_{w \in S_n} sign(w) b_{w(1),1} \dots b_{w(n),n},$$
(4.1)

Cofactor expansions of $\overrightarrow{\det}(B)$ are valid as long as the overall order of the terms is unchanged.

Lemma 4.1. Suppose that B is Hessenberg of order n.

(a) For a fixed $q \in [n-1]$, let $T = -b_{q+1q}$. Then

$$\overrightarrow{\det}B = \operatorname{T}\overrightarrow{\det}B'' + \overrightarrow{\det}B', \qquad (4.2)$$

where B' and B" are obtained from B by setting T = 0, and by deleting the row and column containing T respectively.

- (b) The matrix B' is block upper triangular, with two diagonal blocks which are upper Hessenberg of order q and n q.
- (c) The matrix B'' is upper Hessenberg of order n-1. Also any term in the expression (4.1) for $\overrightarrow{\det}(B)$ which contains a factor of the form b_{iq} or b_{q+1j} cannot occur in $\overrightarrow{\det}B''$.

Proof. Part (a) follows by separating the products in (4.1) that contain T from those that do not. Note that T commutes with all entries in B, and that the order of all other factors of the products is unchanged. The rest is easy.

We consider determinants of Hessenberg matrices with entries on or above the diagonal from \mathfrak{n}^- and subdiagonal entries which commute with all other matrix entries. Explicit expressions in $U(\mathfrak{b}^-)$ for Šapovalov elements can be obtained as complete expansions of suitable determinants of this kind. There are significant differences in the complete expansions depending on the ordering of entries in the matrices. We consider two orderings on the set of positive roots, and explain the relationships between the determinants without reference to Šapovalov elements. First consider the matrix with entries in $U(\mathfrak{gl}(m))$, or any $U(\mathfrak{gl}(r,s))$ with r + s = m.

$$D = \begin{bmatrix} e_{m,m-1} & e_{m,m-2} & \dots & e_{m,2} & e_{m,1} \\ -a_{m-2} & e_{m-1,m-2} & \dots & e_{m-1,2} & e_{m-1,1} \\ 0 & -a_{m-3} & \dots & e_{m-2,2} & e_{m-2,1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & -a_1 & e_{2,1} \end{bmatrix}.$$
(4.3)

Note that in (4.3) the entries from the same row from \mathfrak{n}^- have the form $e_{i,*}$ for some fixed *i*. Another possibility is to require that all entries from the same row have the form $e_{*,i,}$. Thus consider the matrix, where $c_i = a_i + 1$,

$$E = \begin{bmatrix} e_{2,1} & e_{3,1} & \dots & e_{m-1,1} & e_{m,1} \\ -c_1 & e_{3,2} & \dots & e_{m-1,2} & e_{m,2} \\ 0 & -c_2 & \dots & e_{m-1,3} & e_{m,3} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & -c_{m-2} & e_{m,m-1} \end{bmatrix}.$$
(4.4)

The next result is used in the proof of Theorem 4.15.

Proposition 4.2. We have

$$\overrightarrow{\det} D = \overrightarrow{\det} E$$

Proof. This is proved using cofactor expansion and induction. Rather than giving full details, an example might be more helpful here. It is easy to see how the example generalizes. In the general case, we need a commutation relation for cofactors of E.

Lemma 4.3. Let E^1 , E^2 be the cofactors of entry $e_{m,m-1}$ and $-c_{m-2}$ in E respectively. Then $[\overrightarrow{\det} E^1, e_{m,m-1}] = -\overrightarrow{\det} E^2$.

Proof. Note that E^1 is obtained from E^2 by replacing the entries $e_{m,i}$ with $e_{m-1,i}$ for $i \in [m-2]$. Since i < m-1, it is impossible for both $e_{m-1,i}$ and $e_{m,m-1}$ to be odd. Thus $[e_{m-1,i}, e_{m,m-1}] = -e_{m,i}$, the result follows.

Example 4.4. Let

$$E = \begin{bmatrix} e_{2,1} & e_{3,1} & e_{4,1} \\ -c_1 & e_{3,2} & e_{4,2} \\ 0 & -c_2 & e_{4,3} \end{bmatrix}.$$

By cofactor expansion along the last row,

$$\overrightarrow{\det} E = \overrightarrow{\det} E^1 e_{4,3} + c_2 \overrightarrow{\det} E^2$$

where

$$E^{1} = \begin{bmatrix} e_{2,1} & e_{3,1} \\ -c_{1} & e_{3,2} \end{bmatrix}$$
 and $E^{2} = \begin{bmatrix} e_{2,1} & e_{4,1} \\ -c_{1} & e_{4,2} \end{bmatrix}$.

Next set

$$D^{1} = \begin{bmatrix} e_{3,2} & e_{3,1} \\ -a_{1} & e_{2,1} \end{bmatrix} \text{ and } D^{2} = \begin{bmatrix} e_{4,2} & e_{4,1} \\ -a_{1} & e_{2,1} \end{bmatrix}$$

Now by Lemma 4.3,

$$[\overrightarrow{\det} E^1, e_{4,3}] = -\overrightarrow{\det} E^2,$$

 \mathbf{SO}

$$\overrightarrow{\det} E = e_{4,3} \overrightarrow{\det} E^1 + (c_2 - 1) \overrightarrow{\det} E^2$$
$$= e_{4,3} \overrightarrow{\det} D^1 + a_2 \overrightarrow{\det} D^2$$
(4.5)

using induction for the second equality. This is the cofactor expansion of $\overrightarrow{\det} D$ down the first column.

4.2 Application to Sapovalov elements in Type A.

Expansions of Šapovalov elements in Type A were already given in [Mus17] Section 9, using determinants of a certain Hessenberg matrices. However it seems unlikely that this method will generalize, because the determinant of Hessenberg matrix of order m has 2^{m-1} terms, and outside of Type A, the number of partitions of a positive root is rarely a power of 2. Nevertheless the use of Hessenberg matrices gives more insight in the Type A case. In particular they can be used to give expressions for Šapovalov elements using different orderings on the set of positive roots.

Consider the Lie algebra $\mathfrak{gl}(m)$ with simple roots $\alpha_i = \epsilon_i - \epsilon_{i+1}$ for $i \in [m-1]$. Let $\eta' = \epsilon_1 - \epsilon_m$ and $\mu \in \mathcal{H}_{\eta'}$. We show that $\theta_{\eta'}$ can be expressed as a determinant of a certain Hessenberg matrix. Recall the elements

$$h_i = h_{\sigma_i} + (\rho, \sigma_i) - 1$$

from (2.3), for $i \in [m-1]$, Then let $\mathcal{D}^m(\mu)$ be the matrix D from (4.3) with

$$a_i = h_i(\mu) = (\mu + \rho, \sigma_i) - 1.$$
 (4.6)

Theorem 4.5. The Šapovalov element for η' satisfies

$$\theta_{\eta'}(\mu) = \overrightarrow{\det} \mathcal{D}^m(\mu)$$

for all $\mu \in \mathcal{H}_{\eta'}$.

We give two proofs of Theorem 4.5. A comparison of the two approaches is necessary to prove Proposition 5.2. Define H_J from (2.8). In the first we show that the complete expansion of the determinant $\det \mathcal{D}^m(\mu)$ is the evaluation of the element $\Theta_{\eta'}$ from Theorem 2.3 at $\mu \in \mathcal{H}_{\eta'}$. In other words we show

$$\overrightarrow{\det} \mathcal{D}^m(\mu) = \sum_{J \subseteq \mathbb{I}} f_J H_J(\mu).$$
(4.7)

In the second we use induction on m, cofactor expansion and the fundamental Lemma 4.8 below. This Lemma which is [Mus12] Lemma 9.4.3, is the basis for the proof of the existence of Šapovalov elements in the general case. Essentially the same proof is given in [Hum08] Section 4.13.

Theorem 4.5: First Proof. Consider the matrix $\mathcal{D}^m(\mu)$ from Theorem 4.5. We obtain the complete expansion of $\det \mathcal{D}^m(\mu)$. Let \mathbb{I} be as in (2.6). Each term in the complete expansion is obtained by choosing a non-zero product of elements from each column, with each row occurring exactly once. The product of the chosen elements lying above the subdiagonal has the form f_I for some $I \in \mathbb{I}$. The proof of (4.7) is completed by the following Lemma.

Lemma 4.6. The product of subdiagonal terms accompanying f_I is

$$\prod_{i \in r(I)} a_i = \prod_{i \in r(I)} h_i(\mu) = H_I(\mu).$$
(4.8)

Proof. It is easy to see that the rows in which our chosen elements belong to \mathfrak{n}^- are indexed by the set $\{m+1-j|j \in I \setminus \{1\}\}$. The remaining rows must contain an element the subdiagonal of $\mathcal{D}^m(\mu)$. Define $r(I) = \{s-1|s \in \overline{I}\}$ as in Equation (2.7). The subdiagonal element in row k of $\mathcal{D}^m(\mu)$ is $-a_{m-k}$. Since m - (m+1-j) = j - 1, the product of subdiagonal terms accompanying f_I must be

$$\pm \prod_{i \in r(I)} a_i = \pm \prod_{i \in r(I)} h_i(\mu).$$
(4.9)

Next we make a remark about the (non-commutative) determinant of a Hessenberg matrix B of order n. For a general $n \times n$ matrix the complete expansion of the determinant is indexed by permutations w from the symmetric group of degree n. In Equation (4.1) the corresponding term is zero unless $w(i) \ge i - 1$ for $2 \le i \le n$. Now it is easy to see the following.

Lemma 4.7. Assume that in the expression for det (B) given in (4.1), the term indexed by w is nonzero. Let $C = |\{i|w(i) = i - 1\}|$. Then $sign(w) = |(-1)^{|C|}$.

Proof. Each $i \in C$ corresponds to an inversion (i, i - 1) in w. There are no other inversions.

This is fortunate for us because the set C corresponds to the elements on the subdiagonal of B and these all have the form $-a_j$ for some j. This means that (4.9) can be replaced with (4.8) as in the statement of Lemma 4.6.

Theorem 4.5: Second Proof. The Sapovalov element θ_{η} can be constructed inductively using the next Lemma.

Lemma 4.8. Suppose that α is a simple root and set $\mu = s_{\alpha} \cdot \lambda$, $\eta' = s_{\alpha}\eta$. Assume that

- (a) $p = (\mu + \rho, \alpha^{\vee}) \in \mathbb{N} \setminus \{0\}$ and $q = (\eta, \alpha^{\vee}) \in \mathbb{N} \setminus \{0\}$
- (b) $\mu \in \mathcal{H}_{\eta'}$, and consequently $\lambda \in \mathcal{H}_{\eta}$.

Then the evaluation of the Sapovalov elements $\theta_{\eta'}, \theta_{\eta}$ at μ and λ satisfy

$$e_{-\alpha}^{p+q}\theta_{\eta'}(\mu) = \theta_{\eta}(\lambda)e_{-\alpha}^{p}.$$
(4.10)

Proof. See the sources cited above.

Consider the Lie algebra $\mathfrak{gl}(m+1)$ with simple roots $\alpha_i = \epsilon_i - \epsilon_{i+1}$ for $i \in [m]$, and set $\sigma_i = \epsilon_1 - \epsilon_{i+1}$. Let $\alpha = \alpha_m$, $\eta' = \epsilon_1 - \epsilon_m$ and $\eta = \epsilon_1 - \epsilon_{m+1} = s_\alpha \eta'$. Suppose that $\mu \in \mathcal{H}_{\eta'}$. Then $\lambda = s_\alpha \cdot \mu \in \mathcal{H}_{\eta}$. Thus since $(\lambda + \rho, \alpha^{\vee}) = -p$, it follows that

$$1 = (\lambda + \rho, \eta) = (\lambda + \rho, \sigma_{m-1} + \alpha) = -p + (\lambda + \rho, \sigma_{m-1}),$$

which implies that $p = (\lambda + \rho, \sigma_{m-1}) - 1 := a_{m-1}$. If $i \in [m-2]$, then $(\sigma_i, \alpha) = 0$, it follows that the a_i as defined in (4.6) satisfy

$$a_i = (\lambda + \rho, \sigma_i) - 1, \tag{4.11}$$

which means that $\mathcal{D}^{m}(\mu)$ fits into the lower right corner of $\mathcal{D}^{m+1}(\lambda)$. Then $\mathcal{D}^{m+1}(\lambda)$ is obtained from $\mathcal{D}^{m}(\mu)$ by first adding a column on the left, whose only non-zero entry is $-a_{m-1}$ in the first row, and then adding a row so that $e_{m+1,i}$ is directly above $e_{m,i}$ for $i \in [m-1]$, and with $e_{m+1,m}$ as first entry.

Next introduce two determinants $\mathcal{D}_1, \mathcal{D}_2$ by

Observe that $e_{m+1,m}$ commutes with all entries in $\mathcal{D}^m(\mu)$ except for those in the second row, and that

$$e_{m+1,m}^{p+1}e_{m,i} = (e_{m+1,m}e_{m,i} + pe_{m+1,i})e_{m+1,m}^p.$$
(4.13)

This implies that

$$e_{m+1,m}^{p+1} \overrightarrow{\det} \mathcal{D}^m(\mu) = (\overrightarrow{\det} \mathcal{D}_1 + \overrightarrow{\det} \mathcal{D}_2) e_{m+1,m}^p.$$
(4.14)

In (4.14), \mathcal{D}_2 arises from the second term in the factor $(e_{m+1,m}e_{m,i} + pe_{m+1,i})$ from Equation (4.13), and similarly \mathcal{D}_1 corresponds to the first term. But det $\mathcal{D}_1 + \det \mathcal{D}_2$ is just the cofactor expansion of det $\mathcal{D}^{m+1}(\lambda)$ down the first column. Now comparing (4.10) and (4.14), and using the fact that $e_{-\alpha}$ is not a zero divisor in $U(\mathfrak{g})$ it follows that

$$\theta_{\eta}(\lambda) = \overrightarrow{\det} \mathcal{D}^{m+1}(\lambda).$$

We record how Sapovalov elements can be expressed using non-commutative determinants in the case of $\mathfrak{g} = \mathfrak{gl}(m, n)$. First define

$$B_i = \begin{cases} (\lambda + \rho, \sigma_i) - 1 & \text{for } i \in [m-1], \\ (\lambda + \rho, \tau_{i+1-m}) & \text{for } m \le i \le m+n-2. \end{cases}$$
(4.15)

With h_i as in (2.18), we have

$$h_i(\lambda) = B_i \text{ for all } i \in [m+n-2].$$

$$(4.16)$$

,

Next consider the determinant

$$\mathbb{A}(\lambda) = \begin{bmatrix} e_{m+n,m+n-1} & e_{m+n,m+n-2} & \dots & \dots & e_{m+n,2} & e_{m+n,1} \\ -B_{m+n-2} & e_{m+n-1,m+n-2} & \dots & \dots & e_{m+n-1,2} & e_{m+n-1,1} \\ 0 & -B_{m+n-3} & \dots & \dots & e_{m+n-2,2} & e_{m+n-2,1} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \dots & \dots \\ 0 & 0 & -B_m & e_{m+1,m} & \dots & \dots & e_{m+1,1} \\ 0 & 0 & & -B_{m-1} & e_{m,m-1} & \dots & e_{m,1} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \dots & \dots \\ 0 & 0 & \dots & \dots & -B_1 & e_{2,1} \\ \end{bmatrix}$$

Recall the expression for the Šapovalov element θ_{η} from Theorem 2.6.

Theorem 4.9. We have

$$\overrightarrow{\det} \mathbb{A}(\lambda) = \sum_{J \subseteq \mathbb{I}} f_J H_J(\lambda).$$

Thus det $\mathbb{A}(\lambda)$ is the evaluation of the Šapovalov element θ_{η} at $\lambda \in \mathcal{H}_{\eta}$.

Proof. This follows from (4.16), Theorem 2.6 and the argument of Theorem 4.5 showing that f_J is correctly paired with H_J in the above sum.

4.3 Sapovalov elements for arbitrary odd roots.

Assume γ is an odd root. Although as we remarked in Section 1, to compute Šapovalov element θ_{γ} we can assume γ is the highest odd root of $\mathfrak{gl}(m, n)$, we later apply our results when this is not the case. Thus it will be conveneint to have and explicit expressions for a Šapovalov elements for an arbitrary odd root.

Let t = m + 1 - r. We need the Šapovalov element for $\gamma = \epsilon_r - \delta_s$ in $U(\mathfrak{gl}(m, n))$. Let \mathfrak{g}' be the subalgebra $\mathfrak{g} = \mathfrak{gl}(m, n)$ with rows and columns indexed by the set $\{r, \ldots, m+s\}$. Note that \mathfrak{g}' and \mathfrak{g} both share the same odd simple root vector $e_{\beta} = e_{m,m+1}$. Also $\mathfrak{g}' \cong \mathfrak{gl}(t, s)$, and γ is the longest odd positive root for the subalgebra \mathfrak{g}' . The Šapovalov element θ_{γ} is easily found using the matrix

$$\mathbb{A}^{r,s}(\lambda) = \begin{bmatrix} e_{m+s,m+s-1} & e_{m+s,m+s-2} & \dots & e_{m+s,r+1} & e_{m+s,r} \\ -A_{m+s-2} & e_{m+s-1,m+s-2} & \dots & e_{m+s-1,r+1} & e_{m+s-1,r} \\ \vdots & \vdots & \ddots & \dots & \ddots \\ \vdots & \vdots & \ddots & \dots & \ddots \\ 0 & 0 & \ddots & -A_{r+1} & e_{r+2,r+1} & e_{r+2,r} \\ 0 & 0 & \dots & \dots & -A_r & e_{r+1,r} \end{bmatrix},$$

$$(4.18)$$

Note that all entries in the above matrix belong to the subalgebra \mathfrak{g}' of \mathfrak{g} . Set

$$\sigma_i = \epsilon_r - \epsilon_{r+i}, \text{ and } \omega_i = \delta_{i+1-t} - \delta_s.$$
 (4.19)

Then we define A_i by

$$A_{i+r-1} = \begin{cases} (\lambda + \rho, \sigma_i) - 1 & \text{for } i \in [t-1].\\ (\lambda + \rho, \omega_i) & \text{for } t \le i \le t+s-2. \end{cases}$$
(4.20)

We have $(\rho, \alpha) = 1$ or -1, respectively if α is a simple root of $\mathfrak{gl}(t) \oplus 0$ or $0 \oplus \mathfrak{gl}(s)$ and $(\rho, \beta) = 0$ for the unique simple odd root, see Remark 2.1. These are the only properties of ρ we need, so there is no need to introduce the analog of ρ for $\mathfrak{gl}(t, s)$. Let

$$\mathbb{I} = \{ I \subseteq \{ r, r+1, \dots, m+s \} | r, m+s \in I \}.$$
(4.21)

Then for $J \in \mathbb{I}$, define $f_J, r(J)$ as usual. Define $H_{i+r-1} \in U(\mathfrak{h})$ by

$$H_{i+r-1} = \begin{cases} h_{\sigma_i} + \rho(h_{\sigma_i}) - 1 & \text{for } i \in [t-1].\\ h_{\omega_i} + \rho(h_{\omega_i}) & \text{for } t \le i \le t+s-2. \end{cases}$$
(4.22)

Then let $H_J = \prod_{i \in r(J)} H_i$, and note that by (4.20) and (4.22), we have by

$$H_{i+r-1}(\lambda) = A_{i+r-1} \text{ for } i \in [t+s-2].$$
(4.23)

By Theorem 4.9, $\overrightarrow{\det} \mathbb{A}^{r,s}(\lambda)$ is the evaluation of the Šapovalov element $\theta_{\gamma} \xrightarrow{\text{at }} \lambda \in \mathcal{H}_{\gamma}$. We use the same procedure as before to obtain the complete expansion of $\overrightarrow{\det} \mathbb{A}^{r,s}(\lambda)$ and thus the Šapovalov element $\theta_{\gamma} \in U(\mathfrak{gl}(t,s)) \subseteq U(\mathfrak{gl}(m,n))$.

Theorem 4.10. The Šapovalov element θ_{γ} for the root $\gamma = \epsilon_r - \delta_s$ in $U(\mathfrak{gl}(m,n))$ is given by

$$\theta_{\gamma} = \sum_{J \subseteq \mathbb{I}} f_J H_J. \tag{4.24}$$

Proof. This follows from the proof of Theorem 2.6.

Corollary 4.11. For $\lambda \in \mathcal{H}_{\gamma}$, $\theta_{\gamma}(\lambda) = \overrightarrow{\det} \mathbb{A}^{r,s}(\lambda)$.

Proof. By Equation 4.23,

$$\sum_{J \subseteq \mathbb{I}} f_J H_J(\lambda) = \sum_{J \subseteq \mathbb{I}} f_J \prod_{i \in r(J)} A_i$$

and the argument in Theorem 4.5 shows that in the complete expansion of det $\mathbb{A}^{r,s}(\lambda)$, the coefficient of f_J is $\prod_{i \in r(J)} A_i$.

4.4 More Expansions for Sapovalov elements.

We consider two new orders on root vectors, in one the unique odd root vector appears last in each product f_J . In the other they appear first. Applications of each are given in Subsection 4.6.

4.4.1 Putting odd root vectors last.

With I as in (4.21), we give another expansion of the Sapovalov element θ_{γ} using a different order on positive order roots. With this new order for each $I \subseteq I$, the unique odd root vector appears last. Because of this requirement, the definition of f_I is not as easily described using an ordering on I. Thus consider I as an unordered set

$$I = \{m + s, i_1, i_2, \dots, i_g, j_1, \dots, j_h\}$$
(4.25)

where $m + s > i_1 > i_2 \dots > i_g \ge m + 1$ and $r = j_1 < \dots < j_h \le m$. Then we define

$$f_I = e_{m+s,i_1} e_{i_1,i_2} \dots e_{i_{g-1},i_g} e_{j_2,j_1} \dots e_{j_h,j_{h-1}} e_{i_g,j_h}.$$
(4.26)

Note that the odd root vector e_{i_g,j_h} is the last entry in f_I . The definition of H_I is also changed. Instead of (4.22), we define $H_{i+r-1} \in U(\mathfrak{h})$ by

$$H_{i+r-1} = \begin{cases} h_{\sigma_i} + \rho(h_{\sigma_i}) & \text{for } i \in [t-1] \\ h_{\omega_i} + \rho(h_{\omega_i}) & \text{for } t \le i \le t+s-2. \end{cases}$$
(4.27)

Recall the definition of A_i from (4.20) and set $C_i = A_i + 1$ for all *i*. Then

$$H_{i+r-1}(\lambda) = \begin{cases} C_{i+r-1} & \text{for } i \in [t-1] \\ A_{i+r-1} & \text{for } t \le i \le t+s-2. \end{cases}$$
(4.28)

Define r(I) as usual, then set $H_I = \prod_{i \in r(I)} H_i$. Then we have

Theorem 4.12. With the above notation, the Šapovalov element θ_{γ} for the root $\gamma = \epsilon_r - \delta_s$ in $U(\mathfrak{gl}(m, n))$ is given by

$$\theta_{\gamma} = \sum_{I \subseteq \mathbb{I}} f_I H_I. \tag{4.29}$$

Proof. Recall the matrix $\mathbb{A}^{r,s}(\lambda)$ defined in (4.18). The entries in $\mathbb{A}^{r,s}(\lambda)$ belong to the subalgebra $\mathfrak{g}' \cong \mathfrak{gl}(t,s)$. Note that the entries in the last t columns and first s rows of $\mathbb{A}^{r,s}(\lambda)$ are precisely the odd negative root vectors of \mathfrak{g}' . We use the following cofactor procedure. Begin with a cofactor expansion of $\mathbb{A}^{r,s}(\lambda)$ down the first column. Then use a cofactor expansion of the resulting cofactors down the first column. Repeat this a total of s-1 times. Note that every time we use cofactor expansion on a minor, the two resulting cofactors have one fewer row containing odd elements. Thus at the end of the procedure, all the remaining minors contain only odd root vectors in the first row, and all root vectors in this row are odd. Next we use Proposition 4.2 to replace each minor by a matrix containing only odd vectors in the last column. Then the result follows easily after some bookkeeping. Let $\mathbb{K} = \{K \subseteq \{m+1, \dots, m+s\} | m+s \in K\}$. For $K \in \mathbb{K}$, put the entries of K is descending order and define f_K as in (2.5). If \overline{K} is the complement of K in K, set $s(K) = \{p-1 | p \in \overline{K}\}$. The smallest element of K is m+j for some $j \in [s]$, and we set j(K) = j in this situation. Next we describe the minors that can arise at the end of the procedure. First let

$$\mathbb{F}(\lambda) = \begin{bmatrix} -A_{m-1} & e_{m,m-1} & e_{m,m-2} & \dots & e_{m,r+1} & e_{m,r} \\ 0 & -A_{m-2} & e_{m-1,m-2} & \dots & \dots & e_{m-1,r} \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ \vdots & \vdots & \dots & -A_{r+1} & e_{r+2,r+1} & e_{r+2,r} \\ 0 & 0 & 0 & 0 & -A_r & e_{r+1,r} \end{bmatrix}, \quad (4.30)$$

and note that $\mathbb{F}(\lambda)$ fits into the bottom right corner of $\mathbb{A}^{r,s}(\lambda)$. Now let $\mathbb{F}^{(j)}(\lambda)$ be the matrix obtained from $\mathbb{F}(\lambda)$ obtained by adjoining, as the first row the vector

$$(e_{m+j,m}, e_{m+j,m-1}, \dots, e_{m+j,r+1}, e_{m+j,r}).$$
 (4.31)

Note that all entries in (4.31) are odd, while all other entries in $\mathbb{F}^{(j)}(\lambda)$ not on the subdiagonal belong to $\mathfrak{gl}(m) \oplus 0$. Then the result of the cofactor procedure is the following.

Lemma 4.13. If $\lambda \in \mathcal{H}_{\gamma}$, then

$$\theta_{\gamma}(\lambda) = \sum_{K \in \mathbb{K}} f_K \prod_{k \in s(K)} A_k \overrightarrow{\det}(\mathbb{F}^{(j(K))}(\lambda))$$

Proof. In each step of the cofactor procedure we choose either an element from $\mathfrak{n}^$ or a subdiagonal element of $\mathbb{A}^{r,s}(\lambda)$. In column *i*, the entries in \mathfrak{n}^- all have the form $e_{*,m+s-i}$ and the subdiagonal element is $-A_{m+s-i-1}$. We choose an entry from $\mathfrak{n}^$ from this column iff $m + s - i \in K$. Otherwise we choose $-A_{m+s-i-1}$, which agrees with $m + s - i - 1 \in s(K)$. Now in the complete expansion of $\mathbb{A}^{r,s}(\lambda)$, f_K is the initial part of various products f_I with $I \in \mathbb{I}$. If j = j(K), then the last factor in f_K has the form $e_{*,m+j}$. Then to get a valid product the next term should be $e_{m+j,*}$ which means that the minor associated to K at the end of the cofactor procedure is $\mathbb{F}^{(j(K))}(\lambda)$.

Now let $\mathbb{G}^{(j)}(\lambda)$ be the matrix

$$\mathbb{G}^{(j)}(\lambda) = \begin{bmatrix} e_{r+1,r} & e_{r+2,r} & e_{r+3,r} & \dots & e_{m,r} & e_{m+j,r} \\ -C_r & e_{r+2,r+1} & e_{r+2,r} & \dots & e_{m,r+1} & e_{m+j,r+1} \\ 0 & -C_{r+1} & e_{r+3,r+2} & \dots & \dots & e_{m+j,r+2} \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ \vdots & \vdots & \dots & -C_{m-2} & e_{m,m-1} & e_{m+j,m-1} \\ 0 & 0 & 0 & 0 & -C_{m-1} & e_{m+j,m} \end{bmatrix} .$$
(4.32)

Then by Proposition 4.2, $\overrightarrow{\det}(\mathbb{F}^{(j)}(\lambda)) = \overrightarrow{\det}(\mathbb{G}^{(j)}(\lambda))$. Hence

$$\theta_{\gamma}(\lambda) = \sum_{K \in \mathbb{K}} f_K \prod_{k \in s(K)} A_k \overrightarrow{\det}(\mathbb{G}^{(j(K))}(\lambda))$$

Let $\mathbb{J}(j) = \{J \subseteq \{r, r+1, \ldots, m, m+j\} | r, m+j \in J\}$. It is clear that in the complete expansion of the determinant $\det(\mathbb{G}^{(j)}(\lambda))$, we encounter only terms f_J , for some $J \in \mathbb{J}(j)$ ordered as in the last part of (4.26), and that all such J occur. To complete the proof we use a slightly different cofactor procedure than before. We start with a cofactor expansion along the last row, which we call row 1, and continue upwards, numbering rows from bottom to top. In row i the entries from \mathfrak{n}^- have the form $e_{*,m+1-i}$ and the subdiagonal entry is $-C_{m-i}$. Now $m+1-i \in J$ iff we choose an element from \mathfrak{n}^- from this row. Otherwise we choose $-C_{m-i}$, which agrees with $m-i \in r(J)$. Now define t(J) using the complement of J in $\mathbb{J}(j)$. Then we have

$$\overrightarrow{\det}(\mathbb{G}^{(j)}(\lambda)) = \sum_{J \in \mathbb{J}(j)} f_J \prod_{i \in t(J)} C_i$$

Thus

$$\theta_{\gamma}(\lambda) = \sum_{K \in \mathbb{K}} f_K \sum_{J \in \mathbb{J}(j(K))} f_J \prod_{i \in s(K)} A_i \prod_{i \in t(J)} C_i$$

It is clear that for any $K \in \mathbb{K}$ and $J \in \mathbb{J}(j)$ with j = j(K), we have $f_K f_J \in U(\mathfrak{n}^-)^{-\gamma}$ and the terms are correctly ordered so $f_K f_J = f_I$ where $I = K \cup J \in \mathbb{I}$. We have $r(I) = s(K) \cup t(J)$. Now to complete the proof we need to show that

$$\prod_{i \in r(I)} H_i(\lambda) = \prod_{i \in s(K)} A_i \prod_{i \in t(J)} C_i$$

This follows from (4.28).

4.4.2 Putting odd root vectors first.

If I is the set in (4.25) define

$$f_I = e_{i_g, j_h} e_{j_2, j_1} \dots e_{j_h, j_{h-1}} e_{m+s, i_1} e_{i_1, i_2} \dots e_{i_{g-1}, i_g}.$$
(4.33)

Next instead of (4.27), we define $H_{i+r-1} \in U(\mathfrak{h})$ by

$$H_{i+r-1} = \begin{cases} h_{\sigma_i} + \rho(h_{\sigma_i}) - 1 & \text{for } i \in [t-1] \\ h_{\omega_i} + \rho(h_{\omega_i}) + 1 & \text{for } t \le i \le t+s-2. \end{cases}$$
(4.34)

Note that in place of (4.28), we now have

$$H_{i+r-1}(\lambda) = \begin{cases} A_{i+r-1} & \text{for } i \in [t-1] \\ C_{i+r-1} & \text{for } t \le i \le t+s-2. \end{cases}$$
(4.35)

Theorem 4.14. With the above notation, the Šapovalov element θ_{γ} for the root $\gamma = \epsilon_r - \delta_s$ in $U(\mathfrak{gl}(m, n))$ is given by

$$\theta_{\gamma} = \sum_{I \subseteq \mathbb{I}} f_I H_I. \tag{4.36}$$

Proof. This is similar to the proof of Theorem 4.12, so we give fewer details. We use a cofactor expansion of $\mathbb{A}^{r,s}(\lambda)$ along the last t-1 rows working from the bottom up. The subdiagonal entries that emerge in these steps are the A_{i+r-1} for $i \in [t-1]$. At the end of these steps, the unexpanded cofactors have last column consisting of odd root vectors and no other odd root vectors. Then we use Proposition 4.2 to replace each of these cofactors with a matrix containing only odd vectors in the first row, and subdiagonal entries C_{i+r-1} for $t \leq i \leq t+s-2$. The proof concludes as before.

4.5 Partial Expansions of Šapovalov elements.

We obtain two expansions of the Šapovalov element for the root $\gamma = \epsilon_r - \delta_s$ in (4.29) under some additional assumptions. These expansions correspond to partial expansions of the determinant $\det \mathbb{A}^{r,s}(\lambda)$, obtained by separating out the terms containing certain subdiagonal entries, whence the title of this subsection. The main results are Theorems 4.15 and 4.16. Motivation for these results is given in Subsection 4.6. The additional assumptions are

Case 1 We assume that $(\lambda + \rho, \gamma) = (\lambda + \rho, \gamma') = 0$, where $\gamma' = s_{\alpha}\gamma = \gamma - \alpha$ for some posiitive root α . This means that $(\lambda + \rho, \alpha^{\vee}) = 0$ and by [Mus17] Theorem 5.9 we have, in this situation $\theta_{\gamma}v_{\lambda} = \theta_{\alpha,1}\theta_{\gamma'}v_{\lambda}$. Theorem 4.15 below (the main result in Case 1) could be considered as a refinement of this equation when λ is replaced by $\tilde{\lambda} = \lambda + T\xi$.

Case 2 We suppose α_1, α_2 (resp. γ_1, γ_2) are distinct positive even (resp. odd) roots such that

$$\alpha_1 + \gamma_1 + \alpha_2 = \gamma_2. \tag{4.37}$$

It is convenient to use some overlap of notation between Cases 1 and 2. We disregard the notation for α_i, γ_i introduced at the start of Section 2. Then for Case 1 we assume that

$$\gamma = \gamma_2 = \epsilon_r - \delta_s, \alpha = \alpha_1 = \epsilon_r - \epsilon_\ell \text{ and } \gamma' = s_\alpha \gamma = \epsilon_\ell - \delta_s.$$
 (4.38)

In Case 2, without loss of generality we may assume that α_1 and $\gamma = \gamma_2$ are as in Case 1 and we set

$$\alpha_2 = \delta_k - \delta_s, \quad \gamma_1 = \epsilon_\ell - \delta_k. \tag{4.39}$$

4.5.1 Case 1.

We assume that Case 1 holds. Thus (4.38) holds. Before we state the main result there is another issue to deal with. We need to use a different ordering on positive roots. To do this we introduce the matrix

$$\mathbb{B}^{r,s}(\lambda) = \begin{bmatrix} e_{r+1,r} & e_{r+2,r} & \dots & \dots & e_{m+s-1,r} & e_{m+s,r} \\ -C_r & e_{r+2,r+1} & \dots & \dots & e_{m+s-1,r+1} & e_{m+s,r+1} \\ \vdots & \vdots & \ddots & \dots & \dots & \dots \\ \vdots & \vdots & \ddots & \dots & \dots & \dots \\ 0 & 0 & \dots & -C_{m+s-1} & e_{m+s-1,m+s-2} & e_{m+s,m+s-2} \\ 0 & 0 & \dots & \dots & -C_{m+s-2} & e_{m+s,m+s-1} \end{bmatrix},$$

$$(4.40)$$

By Proposition 4.2 and (4.18), we have

$$\overrightarrow{\det} \mathbb{A}^{r,s}(\lambda) = \overrightarrow{\det} \mathbb{B}^{r,s}(\lambda).$$
(4.41)

At this point we bring $\alpha = \epsilon_r - \epsilon_\ell = \sigma_{\ell-r}$ and $T = H_{\ell-1}$ into the story by noting that $(\ell - r) + (r - 1) = \ell - 1$, so by Equations (4.20) and (4.28), we have

$$\mathbf{T}(\lambda) = C_{\ell-1} = (\lambda + \rho, \alpha_1). \tag{4.42}$$

With \mathbb{I} as in (4.21), we define

$$\mathbb{I}_{\ell} = \{ I \subseteq \mathbb{I} | \ell - 1 \in r(I) \}, \ \mathbb{I}^{\ell} = \{ I \subseteq \mathbb{I} | \ell - 1 \notin r(I) \}.$$

Note that $I \subseteq \mathbb{I}^{\ell}$ if and only if $\ell \in I$. Also $T = H_{\ell-1}$ is a factor of H_J iff $J \subseteq \mathbb{I}_{\ell}$. For $J \subseteq \mathbb{I}_{\ell}$, define $H'_J = H_J/T$.

Theorem 4.15. If θ_{γ} is as in Theorem 4.10, then for $\lambda \in \mathcal{H}_{\gamma}$, we have

$$\theta_{\gamma} v_{\lambda} = (\theta_{\alpha,1} \theta_{\gamma'} + \sum_{J \subseteq \mathbb{I}_{\ell}} f_J H'_J \mathtt{T}) v_{\lambda}.$$
(4.43)

Proof. We have

$$\theta_{\gamma} = \sum_{J \subseteq \mathbb{I}^{\ell}} f_J H_J + \sum_{J \subseteq \mathbb{I}_{\ell}} f_J H'_J \mathsf{T}.$$
(4.44)

It is useful to think about this proof in terms of non-commutative determinants. Thus we set $B = \mathbb{B}^{r,s}(\lambda)$ as in (4.40), and recall that $T(\lambda) = C_{\ell-1}$ by (4.42). By Corollary 4.11 and (4.41) we have $\theta_{\gamma}(\lambda) = \overrightarrow{\det B}$. From (4.2),

$$\overrightarrow{\det}B = \overrightarrow{\det}B' + \mathsf{T}(\lambda)\overrightarrow{\det}B'', \qquad (4.45)$$

where B' and B'' are obtained from B by setting $T(\lambda) = 0$, and by deleting the row and column containing $T(\lambda)$ respectively. Thus B'' is the cofactor of entry $T(\lambda)$ in B. Also any term in the expression (4.1) for $\overrightarrow{\det}(B)$ which contains a factor of the form $e_{*,\ell}$ or $e_{\ell,*}$, cannot occur in $\overrightarrow{\det}B''$. This means that $\overrightarrow{\det}B''$ belong to $U(\overline{\mathfrak{g}})$ where $\overline{\mathfrak{g}}$ the subalgebra isomorphic to $\mathfrak{gl}(m-1,n)$ and with rows and columns indexed by the set $\mathbb{I}(\ell) = \{r, \ldots, \widehat{\ell}, \ldots, m+s\}$. As mentioned above $J \subseteq \mathbb{I}_{\ell}$ if and only if $\ell \notin J$, so it follows that the second terms on the right in (4.44) and (4.45) coincide. Note that when $T(\lambda) = 0$, $\theta_{\gamma} v_{\lambda} = \theta_{\alpha,1} \theta_{\gamma'} v_{\lambda}$, as already mentioned, so the proof is complete.

4.5.2 Case 2.

The proof of our main result depends on Theorem 4.12. Hence we define f_I, A_i, H_i as in (4.26), (4.20) and (4.27) respectively. The relation between these elements is given by (4.28). Recall $\alpha_1 = \epsilon_r - \epsilon_\ell = \sigma_{\ell-r}$ and $\alpha_2 = \delta_k - \delta_s$ and we have

$$A_{\ell-1} = (\lambda + \rho, \alpha_1) - 1 \text{ and } A_{m+k-1} = (\lambda + \rho, \alpha_2).$$
(4.46)

It may help to observe that $\omega_i = \delta_k - \delta_s = \alpha_2$, when i = k + t - 1, and thus i + r - 1 = m + k - 1.

We treat

$$T = H_{\ell-1} \text{ and } S = -H_{m+k-1}$$
 (4.47)

as indeterminates which can be evaluated on any $\lambda \in \mathcal{H}_{\gamma}$. By Equation (4.28), we have

$$\mathsf{T}(\lambda) = C_{\ell-1} = (\lambda + \rho, \alpha_1^{\vee}) \text{ and } \mathsf{S}(\lambda) = -A_{m+k-1} = (\lambda + \rho, \alpha_2^{\vee}).$$
(4.48)

For each subset J in (4.29) we are interested in when S or T, from (4.47), or both are factors of H_J . Note that

T is a factor of
$$H_J$$
 iff $\ell - 1 \in r(J)$, iff $J \in \mathbb{I}(\alpha_2)$ or $\mathbb{I}(\alpha_1, \alpha_2)$
S is a factor of H_J iff $m + k - 1 \in r(J)$ iff $J \in \mathbb{I}(\alpha_1)$ or $\mathbb{I}(\alpha_1, \alpha_2)$. (4.49)

where

 $\mathbb{I}(\emptyset) = \{J \subseteq \mathbb{I}|\ell - 1, m + k - 1 \notin r(J)\}, \ \mathbb{I}(\alpha_1) = \{J \subseteq \mathbb{I}|\ell - 1 \notin r(J), m + k - 1 \in r(J)\}, \ (4.50)$ $\mathbb{I}(\alpha_2) = \{J \subseteq \mathbb{I}|m + k - 1 \notin r(J), \ell - 1 \in r(J)\}, \ \mathbb{I}(\alpha_1, \alpha_2) = \{J \subseteq \mathbb{I}|\ell - 1, m + k - 1 \in r(J)\}.$

The notation is inspired by Equation (4.48). Alternatively

$$\mathbb{I}(\emptyset) = \{J \subseteq \mathbb{I}|\ell, m+k \in J\}, \ \mathbb{I}(\alpha_1) = \{J \subseteq \mathbb{I}|\ell \in J, m+k \notin J\},$$
(4.51)
$$\mathbb{I}(\alpha_2) = \{J \subseteq \mathbb{I}|m+k \in J, \ell \notin J\}. \ \mathbb{I}(\alpha_1, \alpha_2) = \{J \subseteq \mathbb{I}|\ell, m+k \notin J\}.$$

Observe that \mathbb{I} is the disjoint union of the above 4 sets. Next for J a subset of $\mathbb{I}(\alpha_1), \mathbb{I}(\alpha_2), \mathbb{I}(\alpha_1, \alpha_2)$ elements $H_J^{(1)}, H_J^{(2)}, H_J^{(3)}$ respectively by

$$H_J^{(1)} = H_J/S, \ H_J^{(2)} = H_J/T, \ H_J^{(3)} = H_J/ST.$$
 (4.52)

By (4.49) these are all elements of $U(\mathfrak{h})$.

Theorem 4.16. The Šapovalov element θ_{γ} for the root $\gamma = \epsilon_r - \delta_s$ in satisfies

$$\theta_{\gamma} = \theta_{\alpha_1} \theta_{\alpha_2} \theta_{\gamma_1} - \sum_{J \in \mathbb{I}(\alpha_1)} f_J H_J^{(1)} S + \sum_{J \in \mathbb{I}(\alpha_2)} f_J H_J^{(2)} T - \sum_{J \in \mathbb{I}(\alpha_1, \alpha_2)} f_J H_J^{(3)} ST. \quad (4.53)$$

Proof. We break the sum in Theorem 4.29 into 4 pieces corresponding to the disjoint union of the sets in (4.51). Then we have to show that $\theta_{\alpha_1}\theta_{\alpha_2}\theta_{\gamma_1} = \sum_{J \in \mathbb{I}(\emptyset)} f_J H_J$. Recall that $T(\lambda) = C_{\ell-1}$ and $S(\lambda) = -A_{m+k-1}$. We first consider what happens when these are zero. Since $C_{\ell-1} = 0$, $\mathbb{B}^{r,s}(\lambda)$ is block upper triangular, and this gives $\overrightarrow{\det} \mathbb{B}^{r,s}(\lambda) = \theta_{\alpha_1} \overrightarrow{\det} \mathbb{B}^{\ell,s}(\lambda) = \theta_{\alpha_1} \overrightarrow{\det} \mathbb{A}^{\ell,s}(\lambda)$. Then $A_{m+k-1} = 0$ implies that $\mathbb{A}^{\ell,s}(\lambda)$ is block upper triangular, and we deduce that $\theta_{\gamma} = \theta_{\alpha_1}\theta_{\alpha_2}\theta_{\gamma_1}$. Now $J \in \mathbb{I}(\emptyset)$ means that $\ell - 1, m + k - 1 \notin r(J)$, and thus neither of S or T is a factor of H_J . So the sum $\sum_{J \in \mathbb{I}(\emptyset)} f_J H_J$ is independent of $T(\lambda)$ and $S(\lambda)$. It follows that $\theta_{\alpha_1}\theta_{\alpha_2}\theta_{\gamma_1} = \sum_{J \in \mathbb{I}(\emptyset)} f_J H_J$ as required.

Remark 4.17. From the proof of Theorem 4.15, that the term $\sum_{J \in \mathbb{I}_{\ell}} f_J H'_J$ in is the Šapovalov element for the longest root $\bar{\gamma}$ of the subalgebra $\bar{\mathfrak{g}}$ of \mathfrak{g} . To see this observe that with $\det B''$ as in the proof, we have $\theta_{\bar{\gamma}}(\lambda) = \det B''$. We can make a similar remark about the last three sums on the right of (4.53). To do this we need the index sets, cf. (4.51),

$$\mathbb{I}_{[\alpha_1]} = \{r, r+1, \dots, \widehat{\ell}, \dots, m+s\}, \\
\mathbb{I}_{[\alpha_2]} = \{r, r+1, \dots, \widehat{m+k}, \dots, m+s\} \\
\mathbb{I}_{[\alpha_1, \alpha_2]} = \{r, r+1, \dots, \widehat{\ell}, \dots, \widehat{m+k}, m+s\},.$$
(4.54)

The set $\mathbb{I}_{[\alpha_1]}$ indexes the rows and columns of the subalgebra $\overline{\mathfrak{g}} \cong \mathfrak{gl}(m-1,n)$ as in Case 1. Define subalgebras $\underline{\mathfrak{g}} \cong \mathfrak{gl}(m,n-1)$ and $\underline{\overline{\mathfrak{g}}} \cong \mathfrak{gl}(m-1,n-1)$, similarly, using the sets $\mathbb{I}_{[\alpha_2]}$ and $\mathbb{I}_{[\alpha_1,\alpha_2]}$ respectively to index the rows and columns. Then in Equation (4.53) from Theorem 4.16, the coefficients of S, T and ST are the Šapovalov elements for the longest odd root of the subalgebras $\underline{\mathfrak{g}}, \overline{\mathfrak{g}}$ and $\underline{\overline{\mathfrak{g}}}$ respectively. More details on this approach were given in [Mus17]. Since we do not need this extra information about Theorems 4.15 or 4.16, we give no further details here. **Example 4.18.** Let $\mathfrak{g} = \mathfrak{gl}(2,2)$ and using the notation of (2.1), set $\alpha = \alpha_1$ and $\gamma = \gamma_1$. We also need

$$e_{-\alpha-\beta} = e_{31}, \quad e_{-\beta-\gamma} = e_{42}, \quad e_{-\alpha-\beta-\gamma} = e_{41}$$

We find the Sapovalov elements for the roots $\alpha + \beta, \beta + \gamma$ and $\alpha + \beta + \gamma$ using Theorem 2.6.

$$\theta_{\alpha+\beta} = e_{-\beta}e_{-\alpha} + e_{-\alpha-\beta}h_{\alpha}$$

$$= e_{-\alpha}e_{-\beta} + e_{-\alpha-\beta}(h_{\alpha}+1).$$

$$(4.55)$$

$$\theta_{\beta+\gamma} = e_{-\gamma}e_{-\beta} + e_{-\beta-\gamma}(h_{\gamma} - 1)$$

$$= e_{-\beta}e_{-\gamma} + e_{-\beta-\gamma}h_{\gamma}.$$
(4.56)

We give four expressions for $\theta_{\alpha+\beta+\gamma}$. The first using Theorem 2.6, the second using Theorem 4.12 or Theorem 4.16, the third using Theorem 4.14 and the fourth by expanding (4.40).

$$\theta_{\alpha+\beta+\gamma} = e_{-\gamma}e_{-\beta}e_{-\alpha} + e_{-\gamma}e_{-\alpha-\beta}h_{\alpha} + e_{-\beta-\gamma}e_{-\alpha}(h_{\gamma}-1) + e_{-\alpha-\beta-\gamma}h_{\alpha}(h_{\gamma}-1)$$

$$= e_{-\gamma}e_{-\alpha}e_{-\beta} + e_{-\gamma}e_{-\alpha-\beta}(h_{\alpha}+1) + e_{-\alpha}e_{-\beta-\gamma}(h_{\gamma}-1) + e_{-\alpha-\beta-\gamma}(h_{\alpha}+1)(h_{\gamma}-1)$$

$$= e_{-\beta}e_{-\gamma}e_{-\alpha} + e_{-\alpha-\beta}e_{-\gamma}h_{\alpha} + e_{-\beta-\gamma}e_{-\alpha}h_{\gamma} + e_{-\alpha-\beta-\gamma}h_{\alpha}h_{\gamma}$$

$$= e_{-\alpha}e_{-\beta}e_{-\gamma} + e_{-\alpha-\beta}e_{-\gamma}(h_{\alpha}+1) + e_{-\alpha}e_{-\beta-\gamma}h_{\gamma} + e_{-\alpha-\beta-\gamma}(h_{\alpha}+1)h_{\gamma}.$$

$$(4.57)$$

Using the relations in $U(\mathfrak{g})$, the right sides of the above expressions are easily seen to be equal.

4.6 Motivation for the Expansions.

Let X be an orthogonal set of positive isotropic roots. Suppose $(\lambda + \rho, \gamma) = 0$ for all $\gamma \in X$. In [Mus17], see also [Mus22b] we constructed some highest weight modules $M^X(\lambda)$ with highest weight λ , and character $e^{\lambda}p_X$, where p_X is a partition function that counts partitions not involving roots in X. The definition of these modules is as follows. Let T be an indeterminant and set $A = \mathbf{k}[T], B = \mathbf{k}(T)$. Now choose $\xi \in \mathfrak{h}^*$ subject to certain conditions which are spelled out in the two cases below, but otherwise generic, and set $\tilde{\lambda} = \lambda + T\xi$, and consider the Verma module $M(\tilde{\lambda})$ over B with highest weight $\tilde{\lambda}$. Then set

$$M^{X}(\widetilde{\lambda})_{B} = M(\widetilde{\lambda})_{B} / \sum_{\gamma \in X} U(\mathfrak{g})_{B} \theta_{\gamma} v_{\widetilde{\lambda}}.$$
(4.58)

Next let $M^X(\lambda)_A$ be the $U(\mathfrak{g})_A$ -submodule of $M^X(\lambda)_B$ generated by the highest weight vector and define

$$M^X(\lambda) = M^X(\widetilde{\lambda})_A / T M^X(\widetilde{\lambda})_A.$$
(4.59)

In [Mus17] Section 11, we evaluate the Šapovalov determinant det F_{η}^{X} and give a Jantzen sum formula for these modules. A difficulty which does not arise in the

case of Verma modules [Mus12] Theorem 10.2.5, is that there is no natural A-basis for the weight space $M^X(\tilde{\lambda})_A^{\tilde{\lambda}-\eta}$ indexed by partitions. To surmount this difficulty we use related determinant det G_η^X of a matix G_η^X with rows and columns indexed by elements $e_{-\pi}v_{\tilde{\lambda}}$, with π a partition of η as in the classical case. The leading term of det G_η^X is easy to compute. In Theorem 11.1 of [Mus17] we determine the relationship between the leading terms of the two determinants det G_η^X and det F_η^X , and evaluate the latter. The comparison of leading terms relies on Equations (4.60) and (4.61) below, which are used repeatedly to improve the A-basis used to calculate det G_η^X . Below we refer to Cases 1 and 2 from the previous Subsection.

Case 1: Let $T = \mathsf{T}$ be as in (4.42). Suppose $(\lambda + \rho, \gamma) = (\lambda + \rho, \gamma') = 0$, where $\gamma' = s_{\alpha}\gamma = \gamma - \alpha$ for some positive root α . Then $(\lambda + \rho, \alpha^{\vee}) = 0$. Choose $\xi \in \mathfrak{h}^*$ so that $(\xi, \gamma) = (\xi, \gamma') = 0 = (\xi, \alpha^{\vee})$ and with ξ otherwise generic. Then set $\lambda = \lambda + T\xi$. Using the notation of (4.43), consider the elements of $M(\lambda)$ given by

$$p = \theta_{\alpha,1} \theta_{\gamma'} v_{\widetilde{\lambda}} \text{ and } q = \sum_{J \subseteq \mathbb{I}_{\ell}} f_J H'_J v_{\widetilde{\lambda}},$$

we have the relation $\theta_{\gamma} v_{\tilde{\lambda}} = p + qT$ in $M(\tilde{\lambda})_A$. Thus if $\gamma \in X$, we have in $M^X(\tilde{\lambda})_A$.

$$p + qT = 0. (4.60)$$

This is a key step in the proof of Theorem 11.4 of [Mus17].

Case 2: Recall by (4.48) that $T(\lambda) = (\lambda + \rho, \alpha_1^{\vee})$ and $S(\lambda) = (\lambda + \rho, \alpha_2^{\vee})$. Set S = T = T + 1. Thus if T = 0, we have $(\lambda + \rho, \alpha_1^{\vee}) = (\lambda + \rho, \alpha_2^{\vee}) = 1$ which is the assumption of Theorem 11.8 of [Mus17]. Assume that $\lambda \in \mathcal{H}_{\gamma_1} \cap \mathcal{H}_{\gamma_2}$. Now choose $\xi \in \mathfrak{h}^*$ so that $(\xi, \gamma_1) = (\xi, \gamma_2) = 0$, and $(\xi, \alpha_1^{\vee}) = (\xi, \alpha_2^{\vee}) = 1$ and with ξ otherwise generic. Then set $\lambda = \lambda + T\xi$, and consider the Verma module $M(\lambda)$ over B with highest weight λ . Let M be the factor module of $M(\lambda)_A$ defined by setting $\theta_{\gamma_1}v_{\lambda} = \theta_{\gamma_2}v_{\lambda} = 0$. Then $(\lambda + \rho, \alpha_1^{\vee}) = (\lambda + \rho, \alpha_2^{\vee}) = T + 1$. Hence in the notation of Theorem 4.16 we have S = T = T + 1. Thus since T + 1 is not a zero divisor in B, if $\theta_{\gamma_1}v_{\lambda} = \theta_{\gamma_2}v_{\lambda} = 0$, then (4.53) yields the following relation in M

$$0 = \left[\sum_{J \in \mathbb{I}(\alpha_1)} f_J H_J^{(1)} - \sum_{J \in \mathbb{I}(\alpha_2)} f_J H_J^{(2)} + \sum_{J \in \mathbb{I}(\alpha_1, \alpha_2)} f_J H_J^{(3)}(T+1)\right] v_{\tilde{\lambda}}$$
(4.61)

This gives a generalization of Equation (B.20) from [Mus17] and is a key step in the proof of Theorem 11.8 of [Mus17].

Finally Theorem 4.14 can be used to give a new proof of a result of Serganova, [Ser96] Theorem 5.5, see Theorem 6.5. For the proof it is important that root vectors are ordered so that the unique odd root vector in each f_J is the first factor. For the proof of Theorem 6.4 the unique odd root vector needs to be the last factor in each f_J .

5 Powers of Šapovalov elements.

Powers of Šapovalov elements are dealt with in detail in Section 5.1 of [Mus17]. The method, which works for all non-exceptional algebras, depends on Lemma 4.8, which is the basis of the original constuction of Šapovalov elements from [Šap72]. We briefly review the details. First write the root η in the form $\eta = w\beta$ with β a simple root and $w \in W$. Next by induction on the length of the w, we construct $\theta_{\eta} \in U(\mathfrak{n}^{-})^{-\eta}$ such that $\theta_{\eta}v_{\lambda}$ is a highest weight vector for $M(\lambda)$ whenever $\lambda \in \mathcal{H}_{\eta}$, and the coefficient of $e_{-\pi^{0}}$ in θ_{η} equals 1, see (1.2). Then we fix a (non-unique) lifting of θ_{η} to an element $\Theta_{\eta} \in U(\mathfrak{b}^{-})^{-\eta} = U(\mathfrak{n}^{-})^{-\eta} \otimes U(\mathfrak{h})$ such that $\Theta_{\eta}v_{\lambda} = \Theta_{\eta}(\lambda)v_{\lambda} = \theta_{\eta}v_{\lambda}$, for all $\lambda \in \mathcal{H}_{\eta}$.

For an isotropic root η , if $\lambda \in \mathcal{H}_{\eta}$, then also $\lambda - \eta \in \mathcal{H}_{\eta}$, and by induction on the length of the w we have

Proposition 5.1. Let η be an odd positive root of $\mathfrak{gl}(m,n)$, and define Θ_{η} as in Theorem 2.6. If $\lambda \in \mathcal{H}_{\gamma}$, then $\Theta_{\eta}^2 v_{\lambda} = 0$.

Proof. Combine Theorem 5.1 from [Mus17] and Theorem 2.6.

For a non-isotropic root η , the situation is more delicate, because it is no longer true that $\lambda \in \mathcal{H}_{\eta,p}$ implies $\lambda - \eta \in \mathcal{H}_{\eta,p}$, and thus we need to evaluate $\Theta_{\eta,p}$ at a point that is not in $\mathcal{H}_{\eta,p}$. To resolve this issue, a uniform inductive construction of Θ_{η} is given in [Mus17] and [Mus22b]. This depends on a specific choice of β and a shortest length expression for w. Then Θ_{η}^{p} is a Šapovalov elements for the pair (η, p) , by [Mus17] Theorem 5.8.

Needless to say, this method of proof is completely different from the direct approach used in most of this paper. However the two approaches are brought together by the two proofs of Theorem 4.5, the second of which uses the same choice of β as in [Mus17]. From this we can deduce the following

Proposition 5.2. Let η be a positive root of $\mathfrak{gl}(m)$, and define Θ_{η} as in Theorem 2.3. Then, for $p \geq 1$, Θ_{η}^{p} is the Šapovalov element for the pair (η, p) . In other words if $\lambda \in \mathcal{H}_{\eta,p}$, we have

$$\Theta^p_\eta(\lambda) = \theta_{\eta,p}(\lambda).$$

Proof. This follows from the above discussion and Theorem 2.3.

The above result may be viewed as a version of [CL74] Theorem 2.7.

6 Survival of Šapovalov elements in factor modules.

Let v_{λ} be a highest weight vector in a Verma module $M(\lambda)$ with highest weight λ , and suppose γ is an odd root with $(\lambda + \rho, \gamma) = 0$. We are interested in the condition that the image of $\theta_{\gamma}v_{\lambda}$ is non-zero in various factor modules of $M(\lambda)$. Throughout this section we assume that $\mathfrak{g} = \mathfrak{gl}(m, n)$.

6.1 Independence of Šapovalov elements.

Given $\lambda \in \mathfrak{h}^*$ set

$$B(\lambda) = \{\gamma \in \Delta_1^+ | (\lambda + \rho, \gamma) = 0\}$$

If γ, γ' are positive non-orthogonal isotropic roots, then $\gamma' = s_{\alpha} \gamma$ for some even root α . Assume that

$$(\gamma, \alpha^{\vee}) = 1 \text{ and } (\lambda + \rho, \alpha^{\vee}) = 0$$
 (6.1)

In this situation the next result relates Sapovalov elements for γ, γ' and α .

Theorem 6.1. Let v_{λ} be a highest weight vector in $M(\lambda)$ and set $\gamma' = s_{\alpha}\gamma$. If $\gamma' \in B(\lambda)$ we have

$$\theta_{\gamma} v_{\lambda} = \theta_{\alpha,1} \theta_{\gamma'} v_{\lambda}. \tag{6.2}$$

Proof. This is a special case of [Mus17] Theorem 5.9 (a). Note that by (6.1) and the hypothesis, $\gamma \in B(\lambda)$.

Define a "Bruhat order" \leq on $B(\lambda)$ as follows. First write $\gamma' \triangleleft \gamma$ if $\gamma - \gamma'$ is a positive even root α , $(\lambda + \rho, \alpha^{\vee}) =$ and $s_{\alpha}\gamma = \gamma'$. The relation \leq is the transitive extension of \triangleleft . Thus $\gamma' \leq \gamma$ if there is a sequence of elements $\gamma = \gamma_0, \gamma_1, \ldots, \gamma_n = \gamma'$ such that $\gamma_i \triangleleft \gamma_{i-1}$ for $i \in [n]$. Then we say that γ is λ -minimal if $\gamma' \leq \gamma$ with $\gamma' \in B(\lambda)$ implies that $\gamma' = \gamma$. For $\gamma \in B(\lambda)$ set $B(\lambda)^{-\gamma} = B(\lambda) \setminus \{\gamma\}$. We say γ is independent at λ if

$$\theta_{\gamma} v_{\lambda} \notin \sum_{\gamma' \in B(\lambda)^{-\gamma}} U(\mathfrak{g}) \theta_{\gamma'} v_{\lambda}.$$

Proposition 6.2. If $\gamma' \triangleleft \gamma$ with $\gamma' \in B(\lambda)$, then $\theta_{\gamma} v_{\lambda} \in U(\mathfrak{g}) \theta_{\gamma'} v_{\lambda}$.

Proof. By hypothesis (6.1) holds. Thus the result follows from Theorem 6.1.

By the Proposition, if we are interested in the independence of the Sapovalov elements θ_{γ} for distinct isotropic roots, it suffices to study only λ -minimal roots γ .

We order the positive roots of \mathfrak{g} as in Theorem 4.12, thus for each $\pi \in \overline{\mathbf{P}}(\gamma)$ the odd root vector is the last factor of $e_{-\pi}$, that is we have $e_{-\pi} \in U(\mathfrak{n}_0^-)\mathfrak{n}_1^-$.

Lemma 6.3. If If γ is λ -minimal, then $e_{-\gamma}v_{\lambda}$ occurs with non-zero coefficient in $\theta_{\gamma}v_{\lambda}$.

Proof. Assume $\gamma = \epsilon_r - \delta_s$. Then if $\alpha = \epsilon_r - \epsilon_i$ with r < i, or $\alpha = \delta_j - \delta_s$ with j < s we have $(\lambda + \rho, \alpha^{\vee}) \neq 0$, since γ is λ -minimal. Thus by Equation (4.27) and Theorem 4.12, the coefficient of $e_{-\gamma}v_{\lambda}$ in $\theta_{\gamma}v_{\lambda}$ is non-zero.

Theorem 6.4. The isotropic root γ is independent at λ if and only if γ is λ -minimal.

Proof. Set $B = B(\lambda)^{-\gamma}$. If γ is not λ -minimal then γ is not independent at λ by Proposition 6.2. Suppose that γ is λ -minimal and

$$\theta_{\gamma} v_{\lambda} \in \sum_{\gamma' \in B} U(\mathfrak{g}) \theta_{\gamma'} v_{\lambda} = \sum_{\gamma' \in B} U(\mathfrak{n}^{-}) \theta_{\gamma'} v_{\lambda}.$$

Hence by Lemma 6.3, there are $a_{\pi} \in \mathbf{k}$ such that

$$e_{-\gamma}v_{\lambda} = \sum_{\pi \in \overline{\mathbf{P}}(\gamma): e_{-\pi} \neq e_{-\gamma}} a_{\pi}e_{-\pi}v_{\lambda}.$$

This contradicts the fact that the set $\{e_{-\pi}v_{\lambda}|\pi\in\overline{\mathbf{P}}(\gamma)\}$ is a basis for $M(\lambda)^{\lambda-\gamma}$. \Box

6.2 Survival of Šapovalov elements in Kac modules.

We have $\mathfrak{g}_1 = \mathfrak{g}_1^+ \oplus \mathfrak{g}_1^-$, where \mathfrak{g}_1^+ (resp. \mathfrak{g}_1^-) is the set of block upper (resp. lower) triangular matrices. Let \mathfrak{h} be the Cartan subalgebra of \mathfrak{g} consisting of diagonal matrices, and set $\mathfrak{p} = \mathfrak{g}_0 \oplus \mathfrak{g}_1^+$. Next let

$$P^+ = \{ \lambda \in \mathfrak{h}^* | (\lambda, \alpha^{\vee}) \in \mathbb{Z}, (\lambda, \alpha^{\vee}) \ge 0 \text{ for all } \alpha \in \Delta_0^+ \}$$

For $\lambda \in P^+$, let $L^0(\lambda)$ be the (finite dimensional) simple \mathfrak{g}_0 -module with highest weight λ . Then $L^0(\lambda)$ is naturally a \mathfrak{p} -module and we define the *Kac module* $K(\lambda)$ by

$$K(\lambda) = U(\mathfrak{g}) \otimes_{U(\mathfrak{g})} L^0(\lambda).$$

By a corollary to the PBW Theorem, see [Mus12] Corollary 6.1.5, we have as a vector space

$$K(\lambda) = \Lambda(\mathfrak{g}_1^-) \otimes L^0(\lambda).$$

The next result is well-known. Indeed two methods of proof are given in Theorem 4.37 of [Bru03]. The second of these is based on Theorem 5.5 in [Ser96]. We give a short proof using Theorem 4.14. Assume that the roots are ordered as in the Theorem, that is with the odd root vector first.

Theorem 6.5. Let $\gamma = \epsilon_r - \delta_s$, and suppose λ and $\lambda - \gamma$ belong to P^+ and $(\lambda + \rho, \gamma) = 0$. Let the highest weight vector v_{λ} be the highest weight vector in $K(\lambda)$ of weight λ . Then

(a) There is a non-zero map of \mathfrak{g} -modules $K(\lambda - \gamma) \longrightarrow K(\lambda)$, sending $xv_{\lambda-\gamma}$ to $x\theta_{\gamma}v_{\lambda}$.

(b)

$$[K(\lambda) : L(\lambda - \gamma)] \neq 0.$$

Proof. Let θ_{γ} be the in the Šapovalov element for the root γ . Because of the way the roots are ordered, applying the right side of (4.36) to the highest weight vector $v_{\lambda} \in K(\lambda)$ yields

$$\theta_{\gamma}v_{\lambda} \in \mathfrak{g}_1^- \otimes L^0(\lambda) \subseteq \Lambda \mathfrak{g}_1^- \otimes L^0(\lambda).$$

Therefore to prove (a) it suffices to show that the coefficient of $e_{-\gamma}v_{\lambda} = e_{-\gamma} \otimes v_{\lambda}$ in $\theta_{\gamma}v_{\lambda}$ is nonzero. Now the term $e_{-\gamma}$ arises in the sum (4.36) when $I \in \mathbb{I}$ is as small as possible, that is $I = \{m + s, r\}$. In this case we have, by (4.19) and (4.34)

$$H_{I} = \prod_{k=1}^{m-r} (h_{\epsilon_{r}-\epsilon_{r+k}-1} + \rho(h_{\epsilon_{r}-\epsilon_{r+k}-1}) - 1) \prod_{j=1}^{s-1} (h_{\delta_{j}-\delta_{s}} + \rho(h_{\delta_{j}-\delta_{s}}) + 1).$$

Thus the coefficient of $e_{-\gamma}v_{\lambda}$ is

$$H_{I}(\lambda) = \prod_{k=1}^{m-r} ((\lambda + \rho, \epsilon_{r} - \epsilon_{r+k}) - 1) \prod_{j=1}^{s-1} ((\lambda + \rho, \delta_{j} - \delta_{s}) + 1).$$

Since $\lambda \in P^+$, the only factors in this product that could be zero arise when k = 1 or j = s - 1. Let $\sigma = \epsilon_r - \epsilon_{r+1}$ and $\tau = \delta_{s-1} - \delta_s$. Then the only factors that could be zero are

$$(\lambda+\rho,\sigma)-1=(\lambda,\sigma) \text{ and } (\lambda+\rho,\tau)+1=(\lambda,\tau).$$

However $(\gamma, \sigma^{\vee}) = (\gamma, \tau^{\vee}) = 1$. So if this happens, then $(\lambda - \gamma, \sigma^{\vee}) = -1$ or $(\lambda - \gamma, \tau^{\vee}) = -1$, and $\lambda - \gamma \notin P^+$. The claim in (b) follows immediately from(a).

References

- [Bru98] J. Brundan, Lowering operators for GL(n) and quantum GL(n), Group representations: cohomology, group actions and topology (Seattle, WA, 1996), Proc. Sympos. Pure Math., vol. 63, Amer. Math. Soc., Providence, RI, 1998, pp. 95–114. MR1603139 (99c:20058)
- [Bru03] _____, Kazhdan-Lusztig polynomials and character formulae for the Lie superalgebra gl(m|n), J. Amer. Math. Soc. 16 (2003), no. 1, 185–231 (electronic). MR1937204 (2003k:17007)
- [Car95] K. J. Carlin, Local systems of Shapovalov elements, Comm. Algebra 23 (1995), no. 8, 3039–3049, DOI 10.1080/00927879508825386. MR1332164 (96d:17009)
- [Car87] R. W. Carter, Raising and lowering operators for sl_n, with applications to orthogonal bases of sl_n-modules, The Arcata Conference on Representations of Finite Groups (Arcata, Calif., 1986), Proc. Sympos. Pure Math., vol. 47, Amer. Math. Soc., Providence, RI, 1987, pp. 351–366. MR933425 (89h:17008)
- [CL74] R. W. Carter and G. Lusztig, On the modular representations of the general linear and symmetric groups, Math. Z. 136 (1974), 193–242. MR0354887 (50 #7364)
- [CP80] R. W. Carter and M. T. J. Payne, On homomorphisms between Weyl modules and Specht modules, Math. Proc. Cambridge Philos. Soc. 87 (1980), no. 3, 419–425, DOI 10.1017/S0305004100056851. MR556922
- [Fra88] J. Franklin, Homomorphisms between Verma modules in characteristic p, J. Algebra 112 (1988), no. 1, 58–85, DOI 10.1016/0021-8693(88)90132-9. MR921964 (89c:17015)
- [Hum08] J. E. Humphreys, Representations of semisimple Lie algebras in the BGG category O, Graduate Studies in Mathematics, vol. 94, American Mathematical Society, Providence, RI, 2008. MR2428237
- [Mus12] I. M. Musson, Lie Superalgebras and Enveloping Algebras, Graduate Studies in Mathematics, vol. 131, American Mathematical Society, Providence, RI, 2012.
- [Mus17] _____, Šapovalov elements and the Jantzen filtration for contragredient Lie superalgebras., arXiv:math/1710.10528. (2017).

- [Mus22a] _____, Coefficients of Šapovalov elements for contragredient Lie superalgebras, in preparation, 2022.
- [Mus22b] _____, The construction of modules with prescribed characters for contragredient Lie superalgebras., in preparation, 2022.
 - [Šap72] N. N. Šapovalov, A certain bilinear form on the universal enveloping algebra of a complex semisimple Lie algebra, Funkcional. Anal. i Priložen. 6 (1972), no. 4, 65–70 (Russian). MR0320103 (47 #8644)
 - [Ser96] V. Serganova, Kazhdan-Lusztig polynomials and character formula for the Lie superalgebra gl(m|n), Selecta Math. (N.S.) 2 (1996), no. 4, 607–651. MR1443186 (98f:17007)