ALMOST CYCLIC REGULAR ELEMENTS IN IRREDUCIBLE REPRESENTATIONS OF SIMPLE ALGEBRAIC GROUPS

D.M. TESTERMAN AND A.E. ZALESSKI

Abstract Let G be a simple linear algebraic group defined over an algebraically closed field of characteristic $p \geq 0$ and let ϕ be a p-restricted irreducible representation of G. Let T be a maximal torus of G and $s \in T$. We say that s is strongly regular if $\alpha(s) \neq \beta(s)$ for all distinct T-roots α and β of G. Our main result states that if all but one of the eigenvalues of $\phi(s)$ are of multiplicity 1 then, with a few specified exceptions, s is strongly regular. This can be viewed as an extension of our earlier result saying that under the same hypotheses, s must be regular and all non-zero weights of ϕ are of multiplicity 1.

1. Introduction

Fix an algebraically closed field F of characteristic $p \geq 0$, let G be a simple linear algebraic group defined over F (hereafter referred to as an algebraic group). Let $\phi: G \to GL(V)$ be a rational irreducible representation (so V is a finite-dimensional FG-module). Recall that $g \in G$ is said to be regular if $\dim C_G(g) = \operatorname{rank}(G)$; so in particular, for a semisimple element $s \in G$, s is regular if and only if $C_G(s)^{\circ}$ is a maximal torus of G. A starting point of our project is the following observation:

(*) Let $s \in G$ be semisimple. If $\phi(s)$ is a regular element in GL(V), then s is regular in G.

Note that the assumption on $\phi(s)$ in (*) is equivalent to saying that all eigenvalue multiplicities of $\phi(s)$ are equal to 1.

Let T be a maximal torus of G. As T meets every conjugacy class of semisimple elements of G, for our considerations, we can assume that $s \in T$. Recall that the roots of G with respect to T are the non-trivial irreducible constituents of the restriction $\mathrm{Ad}|_T$, where Ad is the adjoint representation of G. Then s is regular if and only if $\alpha(s) \neq 1$, for every root α of the root system of G, see [10, Ch. III, §1, Corollary 1.7]. We generalize this in the following

Definition 1.1. Let $s \in T$. We say that s is strongly regular if $\alpha(s) \neq \beta(s)$ whenever $\alpha \neq \beta$ are T-roots of G.

A strongly regular element is regular as $\alpha(s) = 1$ implies $(-\alpha)(s) = 1$; however, there are many elements in T that are regular but not strongly regular. Our first main result enforces (*) as follows:

²⁰²⁰ Mathematics Subject Classification. 20G05.

Key words and phrases. simple algebraic groups, representations, eigenvalue multiplicities, regular elements

The first author acknowledges the support of the Swiss National Science Foundation, research grant number 200020_175571. In addition, this work was initiated while the second author visited the EPFL; he gratefully acknowledges this institute's support.

Theorem 1.2. Let G be a simply connected simple algebraic group defined over F. Let $s \in G$ be a semisimple element and let ϕ be an irreducible representation of G with prestricted highest weight $\omega \neq 0$. Suppose that all of the eigenvalues of $\phi(s)$ are of multiplicity 1. Then one of the following holds:

```
(1) the element s is strongly regular;

(2) G = A_n, n \ge 1, with p \ne 2 if n = 1, and \omega \in \{\omega_1, \ldots, \omega_n\};

(3) G = B_n, n \ge 3, and \omega \in \{\omega_1, \omega_n\};

(4) G = C_n, n \ge 2, p = 2, and \omega \in \{\omega_1, \omega_n\};

(5) G = C_n, n \ge 2, p \ne 2, and either

(a) \omega = \omega_1, or

(b) n \in \{2, 3\} and \omega = \omega_n, or

(c) n \ge 3, p = 3, and \omega \in \{\omega_{n-1}, \omega_n\};

(6) G = D_n, n \ge 4, and \omega \in \{\omega_1, \omega_{n-1}, \omega_n\};

(7) G = E_6 and \omega \in \{\omega_1, \omega_6\};

(8) G = E_7 and \omega = \omega_7;

(9) G = F_4, p = 3, and \omega = \omega_4;

(10) G = G_2 and either \omega = \omega_1, or p = 3 and \omega = \omega_2.
```

(We refer the reader to the end of this section for an explanation of the notation used in the statement. For the reader's convenience the highest weights given in items (2)–(10) are collected in Table 4, at this end of this manuscript.)

The above result, as well as Theorems 1.4, 1.5, 1.7 below, can be viewed as an identification result, in the following sense: the existence of an element whose spectrum satisfies a specified criteria on a given representation either determines the representation or provides structural information about the element. We expect that strongly regular elements can play a certain role in the general theory of algebraic groups and finite groups of Lie type. We mention a paper by Seitz [9, Lemma 2], where he proves that $\alpha|_{T(q)} \neq \beta|_{T(q)}$ for the maximal tori T(q) of finite quasisimple groups of Lie type G(q), with q > 5. This allows him to obtain a certain upper bound for the homogeneous components of T(q) on irreducible G(q)-modules. It was mentioned in [8, p.179] that an analogous bound holds for the s-eigenspace dimensions, $s \in G$ semisimple, provided s is strongly regular.

In fact, we are able to obtain nearly the same conclusion as in Theorem 1.2 under a significantly weaker assumption, specifically, by allowing one of the eigenvalue multiplicities of $\phi(s)$ to be arbitrary (excluding the trivial case of s being central in G). A step in this direction is made in our paper [12, Theorem 2], where we prove the following:

Theorem 1.3. Let G be a simply connected simple algebraic group defined over F and let $s \in G$ be a non-regular non-central semisimple element. Let V be a non-trivial irreducible FG-module with associated representation ϕ . Suppose that at most one eigenvalue of $\phi(s)$ is of multiplicity greater than 1. Then one of the following holds:

```
(1) G is of Lie type A_n, B_n (p \neq 2), C_n or D_n and \dim V = n + 1, 2n + 1, 2n, 2n, respectively;
```

- (2) G is of Lie type B_n , $n \geq 3$, p = 2 and dim V = 2n;
- (3) G is of type A_3 and dim V = 6;
- (4) G is of type C_2 , $p \neq 2$, dim V = 5.

In view of Theorem 1.3, in our consideration of semisimple elements having at most one eigenvalue mutiplicity greater than 1 on some irreducible representation, we may now restrict our attention to regular elements. We have the following result.

Theorem 1.4. Let G be a simply connected simple algebraic group defined over F, $s \in G$ a regular semisimple element, and let ϕ be a non-trivial irreducible representation with prestricted highest weight ω . Suppose that at most one eigenvalue multiplicity of $\phi(s)$ is greater than 1. Then one of the following holds:

- (i) s is strongly regular;
- (ii) G and ω are as in one of the cases (2) (10) of the conclusion of Theorem 1.2;
- (iii) $G = A_n \text{ and } \omega \in \{2\omega_1, 2\omega_2\}, \text{ or } (n, p) = (2, 3) \text{ and } \omega = \omega_1 + \omega_2;$
- (iv) $G = B_n$, $n \ge 3$, p = 2 and $\omega = \omega_2$;
- (v) $G = C_n, n \ge 3, (n, p) \ne (3, 3)$ and $\omega = \omega_2, \text{ or } n = 4, p \ne 2, 3 \text{ and } \omega = \omega_4;$
- (vi) $G = F_4$, $p \neq 3$ and $\omega = \omega_4$, or p = 2 and $\omega = \omega_1$.

The following result shows that the exceptions in Theorems 1.2 and 1.4 are genuine.

Theorem 1.5. Let G be as above and let ϕ be a non-trivial irreducible representation of G.

- (1) Suppose that ϕ is as in items (2) (10) of Theorem 1.2. Then there exists a regular, not strongly regular, semisimple element $s \in G$ such that all eigenvalue multiplicities of $\phi(s)$ are equal to 1.
- (2) Suppose that ϕ is as in items (ii) (vi) of Theorem 1.4. Then there exists a regular, not strongly regular, semisimple element $s \in G$ such that at most one eigenvalue multiplicity of $\phi(s)$ is greater than 1.

There are many examples of irreducible representations ϕ and regular elements $s \in T$ such that $\phi(s)$ has exactly one eigenvalue of multiplicity greater than 1. However, these examples are within a rather narrow class of representations, as the following result shows.

Theorem 1.6. [12, Theorem 2] Let G be a simply connected simple algebraic group defined over F and ϕ a non-trivial irreducible representation of G. Then the following statements are equivalent:

- (i) There exists a non-central semisimple element $s \in G$ such that at most one eigenvalue multiplicity of $\phi(s)$ is equal to 1.
- (2) All non-zero weights of ϕ are of multiplicity 1.

The irreducible representations ϕ whose non-zero weights are of multiplicity 1 have been determined in [11] and those with p-restricted highest weight are reproduced in Tables 2 and 3; see the end of the manuscript.

One can ask whether the assumption that ω be p-restricted can be dropped in Theorems 1.2 and 1.4. Here, we have the following result.

Theorem 1.7. Let G, s be as in Theorem 1.2, where $\operatorname{char}(F) = p > 0$. Let ϕ be an irreducible representation of G with highest weight $\omega = \sum p^i \lambda_i$, where all λ_i are p-restricted weights and at least two of them are non-zero. Suppose that at most one of the eigenvalue multiplicities of $\phi(s)$ is greater than 1. Then either s is strongly regular or for each i with $\lambda_i \neq 0$, we have $(G, \lambda_i) = (G, \omega)$ for (G, ω) as in items (2) - (10) of Theorem 1.2. Moreover,

(i) if $(G, p) = (C_n, 2)$, then for all i, $(\lambda_i, \lambda_{i+1}) \neq (\omega_n, \omega_1)$;

```
(ii) if (G, p) = (G_2, 2), then for all i, (\lambda_i, \lambda_{i+1}) \neq (\omega_1, \omega_1);
(iii) if (G, p) = (G_2, 3), then for all i, (\lambda_i, \lambda_{i+1}) \neq (\omega_2, \omega_1).
```

Notation We fix throughout an algebraically closed field F of characteristic $p \ge 0$. By abuse of notation, when we write $p > p_0$ for some prime p_0 , we allow the case $\operatorname{char}(F) = 0$. Throughout the paper G is a simple simply connected linear algebraic group defined over F. (As mentioned before, we will suppress the adjective "linear".) All G-modules considered are rational finite-dimensional FG-modules.

We fix a maximal torus T in G, which in turn defines the roots of G as well as the weights of G-modules and representations. Recall that the T-weights of a G-module V are the irreducible constituents of the restriction of V to T. When T is fixed (which we assume throughout the paper), we omit the reference to T and write "weights" in place of "T-weights". The set of weights of a G-module V is denoted by $\Omega(V)$. Recall that the roots of G are the non-zero weights of the adjoint representation, equivalently of the G-module $\mathrm{Lie}(G)$. We denote the set of roots by Φ or $\Phi(G)$. The \mathbb{Z} -span of Φ is denoted by R or R(G). The weights in R are called $\mathrm{radical}$. We fix a base of the root system and the simple roots of $\Phi(G)$ are denoted by $\alpha_1, \ldots, \alpha_n$ and ordered as in [1]; we denote the associated set of positive roots by Φ^+ and the $\mathbb{Z}_{>0}$ -span of the simple roots by R^+ .

The Weyl group of G is denoted by W (or W(G)); in fact $W(G) = N_G(T)/T$ so the conjugation action of $N_G(T)$ on T yields an action of W(G) on T and on the T-weights. The W-orbit of $\mu \in \Omega$ is denoted by $W\mu$. Minuscule weights are defined and tabulated in [2, Ch. VIII, §7.3].

The set $\Omega = \operatorname{Hom}(T, F^{\times})$ (the rational homomorphisms of T to F^{\times}) is called the weight lattice, which is a free \mathbb{Z} -module of finite rank called the rank of G. For a G-module $V, t \in T$ and $\mu \in \Omega(V), \mu(t)$ is an eigenvalue of t on V; then $(-\mu)(t) = t^{-1}$ and $(\mu + \nu)(t) = \mu(t)\nu(t)$ for $\nu \in \Omega(V)$. As every semisimple element of G is conjugate to an element of T, in most results which follow, the assumption that s is semisimple is replaced by the assumption that $s \in T$.

We let $\{\omega_1, \ldots, \omega_n\}$ be the set of fundamental dominant weights with respect to the choice of simple roots. These form a \mathbb{Z} -basis of Ω , so every $\nu \in \Omega$ can be expressed in the form $\nu = \sum a_i \omega_i$ $(a_i \in \mathbb{Z})$; the set of ν with $a_i \geq 0$ is denoted by Ω^+ . If p > 0, we set Ω_p^+ to be the set of weights $\nu = \sum a_i \omega_i$ with $0 \leq a_i < p$. If $\operatorname{char}(F) = 0$ we let $\Omega_p^+ = \Omega^+$. The weights of Ω^+ , respectively Ω_p^+ are called dominant, respectively p-restricted weights (so that if $\operatorname{char}(F) = 0$, all dominant weights are by definition p-restricted weights). An irreducible G-module is called p-restricted if its highest weight is p-restricted.

There is a standard partial ordering of elements of Ω ; specifically, for $\mu, \mu' \in \Omega$ we write $\mu \prec \mu'$ and $\mu' \succ \mu$ if and only if $\mu' - \mu \in R^+$ and $\mu \neq \mu'$. We write $\mu \preceq \mu'$ or $\mu' \succeq \mu$ if and only if $\mu' - \mu \in R^+$. For $\mu, \nu \in \Omega^+$, with $\mu \preceq \nu$, we say μ is subdominant to ν . Every irreducible G-module V has a weight ω such that $\mu \prec \omega$ for every $\mu \in \Omega(V)$ with $\mu \neq \omega$. This is called the *highest weight of* V. There is a bijection between Ω^+ and the set of isomorphism classes of irreducible G-modules, so for $\omega \in \Omega^+$ we denote by V_ω the irreducible G-module with highest weight ω . The zero weight space of V is sometimes denoted by V^T . In general, for $S \subseteq G$ we set $V^S = \{v \in V : sv = v \text{ for all } s \in S\}$. The

maximal root of $\Phi(G)$ is denoted by ω_a ; this is the highest weight of the adjoint module Lie(G) and of a composition factor V_a of Lie(G).

Because of natural isomorphisms between certain indecomposable root systems, we will assume throughout that the rank of G is as in Table 1 below. For brevity we write $G = A_n$

G	rank(G)
A_n	$n \ge 1$
B_n	$n \ge 3$
C_n	$n \ge 2$
D_n	$n \ge 4$

Table 1. rank(G)

to say that G is a simple simply connected algebraic group of type A_n , and similarly for the other types. For classical groups G the module with highest weight ω_1 is called the natural module, with one exception, namely the natural module for $G = B_n$ when p = 2 is a reducible (2n + 1)-dimensional module with two composition factors V_{ω_1} and V_0 .

In our proofs, we regularly use so-called "Bourbaki weights", which are elements of a \mathbb{Z} -lattice containing Ω with basis $\varepsilon_1, \varepsilon_2, ...$; the explicit expressions of the fundamental weights and the simple roots of G in terms of the ε_i are given in [1, Planches 1 – XIII].

If $h:G\to G$ is a surjective algebraic group homomorphism and ϕ is a representation of G then the h-twist ϕ^h of ϕ is defined as the mapping $g\to\phi(h(g))$ for $g\in G$. Of fundamental importance is the Frobenius mapping $Fr:G\to G$ arising from the mapping $x\to x^p$ ($x\in F$) when p>0. If V is a G-module then, for integers $k\geq 0$, the modules V^{Fr^k} are called Frobenius twists of V; if V is irreducible with highest weight ω then the highest weight of V^{Fr^k} is $p^k\omega$. Every $\omega=\sum a_i\omega_i\in\Omega^+$ has a unique p-adic expansion $\omega=\lambda_0+p\lambda_1+\cdots+p^k\lambda_k$ for some k, where $\lambda_0,\ldots,\lambda_k\in\Omega_p^+$. The Steinberg tensor product theorem then implies that $V_\omega\cong V_{\lambda_0}\otimes V_{\lambda_1}^{Fr}\otimes\cdots\otimes V_{\lambda_k}^{Fr^k}$.

Finally, we recall that a matrix $M \in M_{n \times n}(F)$ is called *cyclic* if it is diagonalisable and all eigenvalue multiplicities equal 1; we will say M is *almost cyclic*, if it is diagonalizable and at most one eigenvalue multiplicity is greater than 1. This leads us to introduce the following terminology to be used throughout.

Definition 1.8. Let $s \in T$ and let V be a G-module with associated representation $\phi : G \to \operatorname{GL}(V)$. We say that s is cyclic on V, respectively almost cyclic on V if $\phi(s)$ is cyclic, respectively almost cyclic.

2. Preliminaries

We start by recalling a characterization of regular semisimple elements, as mentioned in the Introduction. We will use this frequently without direct reference.

Lemma 2.1. [10, Ch. III, §1, Corollary 1.7] Let G be a semisimple algebraic group, $T \leq G$ a maximal torus and $s \in T$. Then the following conditions are equivalent:

- (1) s is regular;
- (2) if gs = sg for $g \in G$, then g is semisimple;

(3) for all roots $\alpha \in \Phi(G)$, $\alpha(s) \neq 1$.

The following lemma concerns assertion (*) in the Introduction, rephrased in terms of Definition 1.8.

Lemma 2.2. Let V, M be non-trivial G-modules and $s \in T$.

- (1) If s is cyclic on V, then s is regular.
- (2) If s is almost cyclic on $V \otimes M$, then s is regular.

Proof. (1) Let ρ be the representation afforded by V. If s is cyclic on V then $C_{GL(V)}(\rho(s))$ consists of semisimple elements. So the claim follows from Lemma 2.1(2).

(2) By [12, Lemma 1], s is cyclic on V, so the result follows from (1).

Definition 2.3. Let V be a G-module, $\mu, \nu \in \Omega(V)$, and $s \in T$. We say that s separates the weights μ, ν of V if $\mu(s) \neq \nu(s)$. If this holds for every pair of weights $\mu, \nu \in \Omega(V)$, we say that s separates the weights of V.

If s separates the weights of V then the eigenvalue multiplicities of s acting on V are simply the weight multiplicities of V.

Lemma 2.4. Let V, V_1, V_2 be non-trivial G-modules. Let $s \in T \setminus Z(G)$ and assume that s is almost cyclic on V.

- (1) Suppose that $V = V_1 \otimes V_2$. Then s is cyclic on V_1 and on V_2 , all weights of V_1 and V_2 are of multiplicity 1, and s is regular.
- (2) Suppose that $\Omega(V_1) + \Omega(V_2) = \Omega(V)$. Then s separates the weights of V_i , for i = 1, 2.

Proof. This follows from [12, Lemma 1] and [12, Lemma 3].

The set of weights of a G-module V determines the action of a given semisimple element on the module V. We will rely on the following important result [7, Theorem 1] describing precisely, in many cases, the set $\Omega(V)$. We require an additional notation, defined in loc.cit.; we write e(G) for the maximum of the squares of the ratios of the lengths of the roots in $\Phi(G)$. So e(G)=1 for G of types $A_n,D_n,E_n,\ e(G)=2$ for G of type B_n,C_n,F_4 and e(G)=3 for G of type G_2 .

Theorem 2.5. [7, Theorem 1] Assume that p = 0 or p > e(G). Let λ be a p-restricted weight. Then $\Omega(V_{\lambda}) = \{w(\mu) \mid \mu \in \Omega^+, \mu \leq \lambda, w \in W\}$.

As an immediate Corollary of Theorem 2.5, we have:

Lemma 2.6. Assume that p = 0 or p > e(G). Let $\lambda, \mu \in \Omega^+$ with λ p-restricted, and V_{λ} , respectively, V_{μ} the associated irreducible G-modules. Then the following hold.

- (1) If $\mu \prec \lambda$ then $\Omega(V_{\mu}) \subseteq \Omega(V_{\lambda})$.
- (2) If $\lambda + \mu$ is p-restricted then $\Omega(V_{\lambda+\mu}) = \Omega(V_{\lambda} \otimes V_{\mu}) = \Omega(V_{\lambda}) + \Omega(V_{\mu})$.
- (3) If λ is radical then some root is a weight of V_{λ} ; otherwise $\Omega(V_{\lambda})$ contains some minuscule weight.

We begin our considerations of strongly regular elements and their action in certain representations by considering the action on V_a , the irreducible G-module with highest weight the highest root.

Lemma 2.7. Assume that p = 2 if $G = A_1$, $p \neq 3$ if $G = A_2$ or G_2 , and $p \neq 2$ if $G = B_n$, C_n or F_4 . Let $s \in T \setminus Z(G)$. Then s is strongly regular if and only if s is almost cyclic on

Proof. If $G = A_1$ and p = 2, then all non-identity semisimple elements of G act cycically on V_a (which is just a twist of the natural representation of G) and all non-identity semisimple elements are strongly regular. So here the statement is clear.

Now turn to the other cases and suppose that s is strongly regular, so that $\alpha(s) \neq \beta(s)$ for roots $\alpha, \beta \in \Phi(G)$, with $\alpha \neq \beta$. Then, with one exception, the hypothesis of (1), together with Theorem 2.5, implies that $\Omega(V_a) = \Phi(G) \cup \{0\}$ and so s is almost cyclic on V_a . The exception occurs when $G = G_2$ and p = 2. But here as well, we have $\Omega(V_a) = \Phi(G) \cup \{0\}$; see for example [5].

Now suppose that s is almost cyclic on V_a (and rank(G) > 1). Then, in all cases considered here, the zero weight has multiplicity at least 2 on V_a . Thus $\alpha(s) \neq \beta(s)$ for all roots $\alpha, \beta \in \Phi(G)$, with $\alpha \neq \beta$. That is, s is strongly regular.

Remark 2.8. The exceptions in Lemma 2.7 are genuine. Indeed, if $p \neq 2$ and $G = SL_2(F)$, then elements of T which are not strongly regular are of the form $s = \operatorname{diag}(c, c^{-1})$ with $c^4 = 1$, since $\alpha(s) = c^2$. We choose c such that $c^2 = -1$ and then s is non-central, regular, and almost cyclic on V_a . If $G = C_n, n \geq 2$, with p = 2, then $\omega_a = 2\omega_1$. Then V_a is the Frobenius twist of V_{ω_1} , so s is almost cyclic on V_a if and only if s is almost cyclic V_{ω_1} . However, it is easy to construct non-regular elements $s \in T$ that are almost cyclic on V_{ω_1} . For $G = G_2$, p = 3, see Proposition 4.9(4), and for $G = F_4$, p = 2, see Remark 4.4(1). For $G = A_2$ with p = 3 take $s = \operatorname{diag}(a, -a, -a^{-2})$ for $a \in F^{\times}$, so $\alpha_1(s) = (-\alpha_1)(s)$. Then the Jordan normal form of s on V_a is diag $(1, -1, -1, a^3, -a^3, a^{-3}, -a^{-3})$. This matrix is almost cyclic if $a^4 \neq 1$. Finally, for $G = B_n, n \geq 3$, and p = 2, see Lemma 3.15 and Remark 3.16.

Lemma 2.9 and Remark 2.10 will not be required in what follows, but give some additional information about the relationship between a semisimple element being regular and its action on V_a .

Lemma 2.9. Let $s \in T$. Let ω be the highest short root and $V = V_{\omega}$.

- (1) If s is regular, then $V^s = V^T$ and $V^s_a = V^T_a$. (2) Suppose that $V^s_a = V^T_a$. If p > e(G) or if $G = G_2$ and p = 2, then s is regular.

Proof. By Lemma 2.1, s is regular if and only if $\alpha(s) \neq 1$ for all $\alpha \in \Phi(G)$. For (1), we use the fact that all non-zero weights of V_a and V are roots. (2) Using Theorem 2.5, one observes that for p > e(G), $\Omega(V_a) = \Phi(G) \cup \{0\}$. As mentioned in the proof of Lemma 2.7, this also is true for $G = G_2$, when p = 2. So $V_a^T = V_a^s$ implies $\alpha(s) \neq 1$ for $\alpha \in \Phi(G)$, and hence s is regular.

Remark 2.10. If p = e(G), there exists a non-regular element $s \in T$, such that $V_a^s = V_a^T$. See Proposition 4.9(5), for $G = G_2$ with p = 3 and Remark 4.4(2) for $G = F_4$ with p = 2. Now let $G = C_n$ with p = 2. Then $V_a = V_{2\omega_1}$ and hence $\Omega(V_a) = \{\pm 2\varepsilon_1, \ldots, \pm 2\varepsilon_n\}$. So $V_a^T = \{0\}$, and $V_a^s = V_a^T$ if and only if $V_a^s = \{0\}$, equivalently, if and only if 1 is not an eigenvalue of s on V_{ω_1} . As $n \geq 2$, there are non-regular elements $s \in T$ such that $V_{\omega_1}^s = \{0\}$. For $G = B_n$, when p = 2, the non-zero weights of V_a are $\{\pm \varepsilon_i \pm \varepsilon_j \mid 1 \le i < j \le n\}$. Choose s such that $\varepsilon_1(s) = 1$, so s is not regular, but $(\varepsilon_i \pm \varepsilon_j)(s) \neq 1$ for all $i \neq j$.

3. Classical groups

In this section we prove Theorems 1.2, 1.4 and 1.5 for classical groups. In view of Theorem 1.6, we must examine the modules V_{ω} , for $\omega \in \Omega_p^+$, all of whose non-zero weights are of multiplicity 1. These are listed in Tables 2 and 3. Recall that we have fixed a maximal torus T of G, and we assume that our semisimple element belongs to T.

3.1. Groups of type A_n . Throughout this subsection, G is of type A_n .

Lemma 3.1. Let $\omega \in \Omega^+$ and $n \geq 2$.

- (1) If $\omega > 2\omega_1$ then $\omega \succeq \omega_1 + \omega_2 + \omega_n$.
- (2) If $\omega \succ \omega_1 + \omega_n$, then $\omega \succeq \omega_2 + \omega_{n-1}$ if $n \geq 3$ and $\omega \succeq \nu$, for some $\nu \in \{3\omega_1, 3\omega_2\}$ if n = 2.

Proof. Let $\omega > 2\omega_1$. By [14, Lemma 3.2], we can assume that $\omega = 2\omega_1 - \omega_{j-1} + \omega_j + \omega_k - \omega_{k+1}$ for some $j, k \in \{1, \ldots, n\}, j \leq k$. (We set $\omega_0 = 0 = \omega_{n+1}$.) It follows that k = n and $j \leq 2$, and in both cases (1) easily follows.

Now, let $\omega \succ \omega_1 + \omega_n$. As above, we can assume that $\omega = \omega_1 + \omega_n - \omega_{j-1} + \omega_j + \omega_k - \omega_{k+1}$. Then $j \in \{1,2\}$ and $k \in \{n-1,n\}$. If $n \geq 3$, then $\omega \in \{2\omega_1 + 2\omega_n, \omega_2 + \omega_{n-1}, \omega_2 + 2\omega_n, 2\omega_1 + \omega_{n-1}\}$. As $2\omega_1 + 2\omega_n \succ \omega_2 + 2\omega_n \succ \omega_2 + \omega_{n-1}$ and $2\omega_1 + \omega_{n-1} \succ \omega_2 + \omega_{n-1}$, the result follows. If n = 2 then $\omega \in \{2\omega_1 + 2\omega_2, 3\omega_2, 3\omega_1\}$ and $2\omega_1 + 2\omega_2 \succ 3\omega_i$, i = 1, 2, and the result of (2) follows.

Proposition 3.2. Let $G = A_n$, with $n \ge 1$, $s \in T_{reg}$ and let $\omega \in \Omega_p^+$.

- (1) Assume that $\omega \notin \{0, \omega_1, \dots, \omega_n, 2\omega_1, 2\omega_n\}$ and that s is almost cyclic on V_{ω} . Then s is strongly regular, unless (n, p) = (2, 3) and $\omega = \omega_1 + \omega_2$.
- (2) If $\omega \in \{2\omega_1, 2\omega_n\}$ and s is cyclic on V_{ω} , then s is strongly regular.
- (3) If (n,p)=(2,3) and $\omega=\omega_1+\omega_2$, with s cyclic on V_{ω} , then s is strongly regular.

Proof. Set $V = V_{\omega}$. Recall that the *T*-roots of *G* are $\varepsilon_i - \varepsilon_j$ for $i, j \in \{1, \dots, n+1\}, i \neq j$. Then *s* regular implies that $\alpha(s) \neq 1$ for all $\alpha \in \Phi(G)$.

For (1), suppose the contrary, that is, s is almost cyclic on V_{ω} and s is not strongly regular, so that $\alpha(s) = \beta(s)$ for some roots $\alpha \neq \beta$. As W acts transitively on the roots, we can assume that $\alpha = \alpha_1 = \varepsilon_1 - \varepsilon_2$.

If n = 1 then $\beta = -\alpha_1$ and $\alpha(s) = \beta(s)$ implies $(2\alpha_1)(s) = 1$, whence $\alpha_1(s) = -1$. Let $\omega = a\omega_1$, where by hypothesis $3 \le a \le p - 1$. Then the weights of V are $\omega - i\alpha_1$ for $0 \le i \le a$, and $(\omega - i\alpha_1)(s) = (-1)^i\omega(s)$, contradicting that s is almost cyclic on V.

Now let $n \geq 2$. Recall that W acts on $\varepsilon_1, \ldots, \varepsilon_{n+1}$ as the symmetric group S_{n+1} does. Since $\beta = \varepsilon_i - \varepsilon_j$ for some $i \neq j$, we can assume that $\beta \in \{-(\varepsilon_1 - \varepsilon_2), \pm(\varepsilon_1 - \varepsilon_3), \pm(\varepsilon_2 - \varepsilon_3), \varepsilon_3 - \varepsilon_4\}$ (where $\varepsilon_3 - \varepsilon_4$ occurs only if $n \geq 3$). As s is regular, $\alpha - \beta$ is not a root, so $\beta \notin \{\varepsilon_1 - \varepsilon_3, -\varepsilon_2 + \varepsilon_3\}$, and hence $\beta \in \{-(\varepsilon_1 - \varepsilon_2), -(\varepsilon_1 - \varepsilon_3), \varepsilon_2 - \varepsilon_3, \varepsilon_3 - \varepsilon_4\} = \{-\alpha_1, -\alpha_1 - \alpha_2, \alpha_2, \alpha_3\}$. In addition, as s is regular, if $\beta = -\alpha$ then $p \neq 2$ and $\alpha_1(s) = -1$.

By [12, Lemma 11] (see Definition 3, loc.cit.) and Theorem 2.5, there exists a weight $\mu \in \Omega(V) \cap \{\omega_1 + \omega_i, \omega_i + \omega_n : i = 1, ..., n\}$ and all weights subdominant to μ also lie in $\Omega(V)$. We consider in turn each possibility for the weight μ .

Suppose first that $\mu := \omega_1 + \omega_i$ for 1 < i < n. Then $\mu - \alpha_1 - \alpha_2 - \cdots - \alpha_i = \omega_{i+1} \in \Omega(V)$ and $\omega_{i+1} - \alpha$, $\omega_{i+1} - \beta \in W\mu$. In addition, $\omega_{i+1} - \alpha_{i+1} - \alpha$ and $\omega_{i+1} - \alpha_{i+1} - \beta \in \Omega(V)$, as long as $\beta \in \{-\alpha_1, \alpha_2\}$ when i = 2. Since s is almost cyclic on V, we deduce that either i = 2 and $\beta \in \{-\alpha_1 - \alpha_2, \alpha_3\}$ or $\omega_{i+1} - \alpha = \omega_{i+1} - \alpha - \alpha_{i+1}$, so $\alpha_{i+1}(s) = 1$, contradicting

the fact that s is regular. So it remains to consider the case where $i=2, \ \mu=\omega_1+\omega_2, \ n\geq 3, \ \text{and} \ \beta\in\{-\alpha_1-\alpha_2,\alpha_3\}.$ We give the details for the case $\beta=-\alpha_1-\alpha_2$. Here $\eta=\mu-\alpha_1-2\alpha_2-\alpha_3\in\Omega(V)$ and $\eta-\alpha,\eta-\beta\in\Omega(V)$. Hence we deduce as above that $\omega_{i+1}=\eta$ and so $(\alpha_2+\alpha_3)(s)=1$, contradicting that s is regular. The case i=2 and $\beta=\alpha_3$ is entirely similar.

Suppose now that $\mu = \omega_1 + \omega_n$. Then all roots are weights of V_{ω} , and since $\alpha(s) = \beta(s)$, we have $(-\alpha)(s) = (-\beta)(s)$, whence $\alpha(s) = \beta(s) = -1$ and $(\alpha + \beta)(s) = 1$. Then $\alpha + \beta$ is not a root as $s \in T_{reg}$. Therefore, $\beta \in \{-\alpha_1, \alpha_3\}$. If $\beta = \alpha_3$, so $n \geq 3$, then $\alpha_1 + \alpha_2 + \alpha_3$ is a root and $(\alpha_1 + \alpha_2 + \alpha_3)(s) = \alpha_2(s)$. As before, we deduce that $\alpha_2(s) = \alpha_1(s) = -1$, and then $(\alpha_1 + \alpha_2)(s) = 1$, whence a contradiction. So we are left with the case where $\beta = -\alpha$ and $\alpha(s) = (-\alpha)(s) = -1$. By Lemma 2.7 (and the hypothesis of (1)), we may assume that $\omega \neq \omega_1 + \omega_n$. By Lemma 3.1, $\omega \succeq \nu$, where $\nu \in \{3\omega_1, 3\omega_2\}$ if n = 2, and $\nu = \omega_2 + \omega_{n-1}$ if $n \geq 3$. As usual, Theorem 2.5 implies that all weights subdominant to ν lie in $\Omega(V)$.

Now if $n \geq 3$, so $\nu = \omega_2 + \omega_{n-1}$, then set $\eta = \nu - \alpha_1 - 2\alpha_2 - \alpha_3 - \cdots - \alpha_{n-1}$. Note that $\eta \pm \alpha_1$ and $\eta - \alpha_n \pm \alpha_1$ lie in $\Omega(V)$. But then we deduce that $\alpha_n(s) = 1$, giving the usual contradiction. If n = 2 and $\nu = 3\omega_1$, we argue similarly, setting $\eta = \nu - \alpha_1$, and deduce that $\alpha_2(s) = 1$. The case $\nu = 3\omega_2$ is analogous.

Note that the cases considered above cover also the weights $\mu = \omega_i + \omega_n$, for $2 \le i \le n$, by considering the dual V^* ; so (again invoking duality), it remains to consider the case $\mu = 2\omega_1$. Then Lemma 3.1 shows that $\omega \succeq \omega_1 + \omega_2 + \omega_n$, which then implies that $\omega \succ \omega_3 + \omega_n$, if $n \ge 3$, and $\omega \succeq \omega_1 + 2\omega_2$, if n = 2. In the former case the result follows from previously considered cases if $n \ge 4$. For the case n = 3 and $\omega \succeq \omega_1 + \omega_2 + \omega_3$, we consider the different possibilities for β , as before; recall that $\alpha = \alpha_1$ and $\beta \in \{-\alpha_1, -\alpha_1 - \alpha_2, \alpha_2, \alpha_3\}$. Setting $\nu = \omega_1 + \omega_2 + \omega_3$, and $\eta = \nu - \alpha_1 - \alpha_2 - \alpha_3$ and $\eta' = \nu - \alpha_1 - 2\alpha_2 - \alpha_3$, we have that $\eta - \alpha, \eta - \beta, \eta' - \alpha, \eta' - \beta \in \Omega(V)$ and deduce that $\eta(s) = \eta'(s)$ and so $\alpha_2(s) = 1$ giving the usual contradiction. Finally, let n = 2, and set $\nu = \omega_1 + 2\omega_2$ and assume that $\omega \succeq \nu$. We argue as in the previous cases. As before, since s is regular and $\alpha(s) = \beta(s)$, $\alpha - \beta$ is not a root. So taking $\alpha = \alpha_1$, we have that $\beta \in \{-\alpha_1, \alpha_2, -\alpha_1 - \alpha_2\}$. Set $\eta = \nu - \alpha_1 - \alpha_2$ and $\eta' = \nu - 2\alpha_1 - 2\alpha_2$. Then one checks that $\eta - \alpha, \eta - \beta, \eta' - \alpha, \eta' - \beta \in \Omega(V)$. So we deduce that $\eta(s) = \eta'(s)$ and so $(\alpha_1 + \alpha_2)(s) = 1$, a contradiction.

Turning to the proof of (2), we assume now that s is regular and cyclic on $V = V_{\omega}$, with $\omega \in \{2\omega_1, 2\omega_n\}$. By duality, we may assume that $\omega = 2\omega_1$; note that $p \neq 2$. Suppose the contrary, that is, s is not strongly regular, and let α, β be as in the proof of (1). Here, we only need to find distinct weights $\eta, \eta' \in \Omega(V)$ such that $\eta(s) = \eta'(s)$. It is straightforward to find such a pair for each choice of β (recall that $\alpha = \alpha_1$). For example, if $\beta = -\alpha_1$, then $2\alpha_1(s) = 0$ and so we may take $\eta = \omega$ and $\eta' = \omega - 2\alpha_1$. Or if $\beta = \alpha_2$ (so $n \geq 2$), we take $\eta = \omega - 2\alpha_1$ and $\eta' = \omega - \alpha_1 - \alpha_2$. The remaining cases are left to the reader.

To conclude, we turn to (3). Here, as $\Omega(V) = \{0\} \cup \Phi(G)$, and s is cyclic on V, we see that $\alpha(s) \neq \beta(s)$ for all $\alpha, \beta \in \Phi(G)$, $\alpha \neq \beta$; thus s is strongly regular as claimed.

Corollary 3.3. Theorem 1.2 and Theorem 1.4 hold for $G = A_n$.

Proof. For Theorem 1.2, we assume that $s \in T$ is cyclic on V_{ω} , for some $\omega \in \Omega_p^+$, $\omega \neq 0$. In addition, we assume that either p = 2 when n = 1 or $\omega \notin \{\omega_1, \ldots, \omega_n\}$. For n = 1, we have $\omega = \omega_1$ and $s \in T$ is cyclic on V_{ω} if and only if $s \neq 1$, if and only if s is strongly regular, as claimed. So we now assume $n \geq 2$; Theorem 1.3 implies that s is regular and then we use Proposition 3.2 to see that s is strongly regular.

For Theorem 1.4, we assume that $s \in T_{reg}$ is almost cyclic on V_{ω} , for some $\omega \in \Omega_p^+$, $\omega \neq 0$. We suppose further that either p=2 if n=1 or $\omega \notin \{2\omega_1, 2\omega_n, \omega_1, \ldots, \omega_n\}$. In addition, we assume that if n=2 and $\omega=\omega_1+\omega_2$, then $p\neq 3$. Now if n=1, with p=2 and so $\omega=\omega_1$, then s is almost cyclic on V_{ω} if and only if $s\neq 1$, if and only if s is cyclic on V_{ω} , if and only if s is strongly regular. For $n\geq 2$, Proposition 3.2 gives the result. \square

We now investigate the weights in Theorem 1.2(2) and Theorem 1.4(iii).

Lemma 3.4. Let $G = A_n$ and $\omega \in \Omega_p^+$, $\omega \neq 0$.

- (1) Assume that $p \neq 2$ if n = 1. If $\omega \in \{\omega_1, \ldots, \omega_n\}$, then there exists $s \in T$ with s not strongly regular and s cyclic on V_{ω} .
- (2) If $\omega \in \{2\omega_1, 2\omega_n\}$, then there exists $s \in T_{reg}$, with s not strongly regular and s almost cyclic on V_{ω} .
- *Proof.* (2) Here we have $p \neq 2$ and the result follows from Remark 2.8 when n = 1. So we now take $n \geq 2$. By duality, it suffices to consider $\omega = 2\omega_1$. Choose $T \leq \mathrm{SL}_{n+1}(F)$ to be the group of diagonal matrices $t = \mathrm{diag}(t_1, \ldots, t_{n+1})$ of determinant 1, and set $\varepsilon_i(t) = t_i$, for $i = 1, \ldots, n+1$.

Suppose first that $n \geq 3$. Let $s = \operatorname{diag}(a^3, a^2, a, b_1, \ldots, b_{n-2}) \in \operatorname{SL}_{n+1}(F)$ for some $a, b_1, \ldots, b_{n-2} \in F^{\times}$. Then $\alpha_1(s) = \alpha_2(s) = a$ so s is not strongly regular. The weights of V_{ω} are $2\varepsilon_i$ and $\varepsilon_i + \varepsilon_j$ for $i, j \in \{1, \ldots, n\}$, $i \neq j$. So the eigenvalues of s on V are a^6, a^4, a^2, b_i^2 for $i = 1, \ldots, n-2$ and $a^5, a^4, a^3, a^k b_i, b_i b_j$ for $1 \leq i < j \leq n-2, k = 1, 2, 3$. It is straightforward to see that there exist $a, b_1, \ldots, b_{n-2} \in F^{\times}$ such that a^4 is the only eigenvalue of s on V of multiplicity greater than 1, whence the result in this case.

- If n=2, then let $s=\mathrm{diag}(a,-a,-a^{-2})$. Then $\alpha_1(s)=(-\alpha_1)(s)=-1$ and the eigenvalues of s on $V_{2\omega_1}$ are $a^2,a^2,a^{-4},-a^2,-a^{-1},a^{-1}$, so s is not strongly regular and almost cyclic on V_{ω} for a suitable choice of $a \in F^{\times}$.
- (1) A similar construction can be used when $\omega \in \{\omega_1, \dots, \omega_n\}$. The assumption on the characteristic in case n = 1 is necessary as all non-identity semisimple elements in $SL_2(F)$ are strongly regular when char(F) = 2.

We note that Remark 2.8 and Lemma 3.4 establish the statements of Theorem 1.5 for the groups of type A_n .

3.2. Groups of type B_n , C_n and D_n . In this section, we prove the main results for classical groups not of type A_n . Recall that we assume that $n \geq 3$ when $G = B_n$, $n \geq 2$ when $G = C_n$, and $n \geq 4$ when $G = D_n$.

The weights of V_{ω_1} are $\pm \varepsilon_1, \ldots, \pm \varepsilon_n$ and, additionally, 0 when G is of type B_n and $p \neq 2$. So every element $s \in T$ can be written as $s = \text{diag}(a_1, \ldots, a_n, 1, a_n^{-1}, \ldots, a_1^{-1})$, with respect to some basis of V_{ω_1} , where 1 must be dropped if $G = B_n$ with p = 2.

Lemma 3.5. Let $G = B_n$, $n \ge 3$, or D_4 , $n \ge 4$, or $G = C_n$, $n \ge 3$ and p = 2. Let $\omega \in \Omega_p^+$, $\omega \ne 0$, and $s \in T_{reg}$. Suppose that s is almost cyclic on V_{ω} . Then one of the following holds:

- (i) s is strongly regular,
- (ii) $G = B_n$ and $\omega \in \{\omega_1, \omega_n\}$,
- (iii) $G = B_n$, p = 2 and $\omega = \omega_2$,
- (iv) $G = C_n$ and $\omega \in \{\omega_1, \omega_2, \omega_n\}$, or

(v) $G = D_n$ and $\omega \in \{\omega_1, \omega_{n-1}, \omega_n\}$.

Proof. By Theorem 1.6, all non-zero weights of V_{ω} are of multiplicity 1. The non-zero p-restricted irreducible G-modules with this property are listed in Tables 2 and 3. We find that $\omega \in \{0, \omega_1, 2\omega_1, \omega_2, \omega_n\}$ for $G = B_n$, $\omega \in \{0, \omega_1, 2\omega_1, \omega_2, \omega_{n-1}, \omega_n\}$ for $G = D_n$, and $\omega \in \{0, \omega_1, \omega_n, \omega_2\}$ for $G = C_n$. Lemma 2.7 then gives the result unless $G = B_n$ or D_n and $\omega = 2\omega_1$.

So suppose that $p \neq 2$, $\omega = 2\omega_1$ and s is almost cyclic on V_{ω} . Note that $\omega_2 = 2\omega_1 - \alpha_1$, so by Lemma 2.6(1), $\Omega(V_{\omega_2}) \subset \Omega(V_{2\omega_1})$. Moreover, the 0 weight of $V_{2\omega_1}$ has multiplicity strictly greater than 1. (See Table 3.) Finally, as $V_a = V_{\omega_2}$ and $p \neq 2$, we have $\Phi(G) \subset \Omega(V_{2\omega_1})$. Now suppose that s is not strongly regular; then Lemma 2.7 implies that s is not almost cyclic on V_{ω_2} , and since here as well the 0 weight occurs with multiplicity at least 2, we have that there exist $\alpha, \beta \in \Phi(G)$, $\alpha \neq \beta$ with $\alpha(s) = \beta(s) \neq 1$. But this then implies that s is not almost cyclic on $V_{2\omega_1}$, giving the result.

Corollary 3.6. Theorem 1.2 and Theorem 1.4 hold for $G = B_n$, $n \ge 3$, $G = D_n$, $n \ge 4$, and $G = C_n$ with $n \ge 3$ and p = 2.

Proof. For Theorem 1.2, we take $s \in T$ and $\omega \in \Omega_p^+$, $\omega \neq 0$, such that s is cyclic on V_ω . Then all weights of V_ω occur with multiplicity 1 and so ω is as in Table 2. As all of these weights appear in the conclusion of Theorem 1.2, the result holds.

For Theorem 1.4, we take ω as before and suppose that $s \in T_{reg}$, with s almost cyclic on V_{ω} . Then Lemma 3.5 gives the result.

We now turn to the consideration of the group $G = C_n$, when $p \neq 2$. Our first result gives some information about the associated weight lattice Ω .

Lemma 3.7. Let Φ be of type C_n and let $\omega \in \Omega^+$. Suppose that $\omega \notin \{0, 2\omega_1\}$ and ω is not a fundamental dominant weight. Then one of the following holds:

- (i) $\omega \succeq \omega_1 + \omega_2$,
- (ii) $n \geq 3$ and $\omega \succeq \omega_1 + \omega_3$, or
- (iii) n=2 and $\omega \succeq 2\omega_2$.

Proof. If ω is not radical (so $\omega \succ \omega_1$ by [12, Lemma 12]) then the result is available in [13, Lemma 2.2]. So suppose now that ω is radical and set $\omega = \sum c_i \omega_i$, with $\sum c_i \geq 2$. Consider first the case n = 2. If $c_2 \geq 2$ then $\omega - 2\omega_2$ is a radical dominant weight, so $\omega \succeq 2\omega_2$. If $c_2 = 1$ then c_1 is even and $c_1 \geq 2$ by hypothesis, so again $\omega \succ \omega - \alpha_1 = (c_1 - 2)\omega_1 + 2\omega_2 \succeq 2\omega_2$. So finally, suppose that $c_2 = 0$, so that $c_1 \geq 4$ is even. Then $\omega \succeq \omega - 2\alpha_1 = (c_1 - 4)\omega_1 + 2\omega_2 \succeq 2\omega_2$.

Consider now the case $n \geq 3$. If $c_1c_3 \neq 0$ then $\omega - \omega_1 - \omega_3$ is a dominant radical weight, so $\omega - \omega_1 - \omega_3 \succeq 0$, whence the result. Next suppose that $c_ic_j \neq 0$ for some odd i < j, $(i,j) \neq (1,3)$. As $\omega_i \succ \omega_{i-2}$ for $i \geq 2$, it follows that $\omega \succ \omega' = \omega + \omega_1 + \omega_3 - \omega_i - \omega_j$, and ω' is a dominant radical weight having non-zero coefficient of ω_1 and ω_3 . By the previous argument, $\omega' \succeq \omega_1 + \omega_3$, whence the result in this case. The same argument works if $c_i \geq 2$ for some odd $i \geq 3$, as $\omega_{i-2} = \omega_i - \beta$ for some $\beta \in \Phi^+$, and so $c_i\omega_i = (c_i - 1)\omega_i + \omega_i = (c_i - 1)\omega_i + \omega_{i-2} + \beta$, and we are in the previous case. We are left with the case where $c_i = 0$ for all odd i and either $c_i \geq 2$ for some even i or $c_ic_j \neq 0$ for some even i < j. In each case, it is a strightforward check to see that $\omega \succeq 2\omega_2 \succ \omega_1 + \omega_3$ as required.

Proposition 3.8. Let $G = C_n$, $n \ge 3$, $p \ne 2$, and $s \in T$. Let $\omega \in \Omega_p^+$, $\omega \ne 0$, and suppose that s is almost cyclic on V_ω . Then s is strongly regular unless one of the following holds:

- (1) $\omega \in \{\omega_1, \omega_2\}$;
- (2) n=3,4 and $\omega=\omega_n$; or
- (3) $n \ge 4$, p = 3, and $\omega \in \{\omega_{n-1}, \omega_n\}$.

Proof. We assume that $\omega \notin \{0, \omega_1\}$. Then by Theorem 1.3, s is regular, so $\alpha(s) \neq 1$ for every root α . By Theorem 1.6, ω occurs in Table 2 or 3. The case $\omega = 2\omega_1 = \omega_a$ is settled in Lemma 2.7, so we assume as well that $\omega \neq 2\omega_1$. The other entries of Tables 2 and 3 either appear in cases (1) - (3) of the statement or p > 3 and $\omega \in \{\omega_{n-1} + \frac{p-3}{2}\omega_n, \frac{p-1}{2}\omega_n\}$. So consider these remaining configurations. By Theorem 3.7, either $\omega \succeq \omega_1 + \omega_2$ or $\omega \succeq \omega_1 + \omega_3$. By Lemma 2.5, we have $\Omega(V_{\omega_1 + \omega_2}) \subseteq \Omega(V)$, respectively $\Omega(V_{\omega_1 + \omega_3}) \subseteq \Omega(V)$

Suppose that s is not strongly regular, so that we have $\alpha(s) = \beta(s)$ for some $\alpha, \beta \in \Phi(G)$, $\alpha \neq \beta$. The roots of G are $\pm 2\varepsilon_i, \pm \varepsilon_j \pm \varepsilon_k$ for $i, j, k \in \{1, ..., n\}$. As s is regular, we have $\varepsilon_i(s) \neq \pm 1$ and $(\varepsilon_j \pm \varepsilon_k)(s) \neq 1$, for all i and $j \neq k$. Using the Weyl group, we can assume that s satisfies one of:

- (i) $(2\varepsilon_1)(s) = (2\varepsilon_2)(s)$,
- (ii) $(2\varepsilon_1)(s) = (\varepsilon_2 \varepsilon_1)(s)$,
- (iii) $(2\varepsilon_1)(s) = (\varepsilon_2 \varepsilon_3)(s),$
- (iv) $n \ge 4$ and $(\varepsilon_1 \varepsilon_2)(s) = (\varepsilon_3 \varepsilon_4)(s)$, or
- (v) $(2\varepsilon_1)(s) = (-2\varepsilon_1)(s)$.

We now consider the two cases $\omega \succeq \omega_1 + \omega_2$ or $\omega \succeq \omega_1 + \omega_3$ separately.

Suppose first that $\omega \succeq \omega_1 + \omega_2 \succ \omega_1$. Then V_{ω} has weights $\pm \varepsilon_i$ $(i=1,\ldots,n)$ and $\pm 2\varepsilon_i \pm \varepsilon_j$, for $1 \le i \ne j \le n$. If (i) holds, $(2\varepsilon_1)(s) = (2\varepsilon_2)(s)$, so $(2\varepsilon_1 \pm \varepsilon_3)(s) = (2\varepsilon_2 \pm \varepsilon_3)(s)$. Since s is almost cyclic on V_{ω} , we have $(2\varepsilon_1 + \varepsilon_3)(s) = (2\varepsilon_1 - \varepsilon_3)(s)$, whence $\varepsilon_3(s) = \pm 1$, contradicting s regular. If (ii) holds, $(2\varepsilon_1)(s) = (\varepsilon_2 - \varepsilon_1)(s)$, so that $(2\varepsilon_1 - \varepsilon_2)(s) = (-\varepsilon_1)(s)$. As $\pm (2\varepsilon_1 - \varepsilon_2)$, $\pm \varepsilon_1$ are weights of V_{ω} , it follows that $(-\varepsilon_1)(s) = -1 = \varepsilon_1(s)$, again a contradiction. If (iii) holds, $(2\varepsilon_1)(s) = (\varepsilon_2 - \varepsilon_3)(s)$, so that $(2\varepsilon_1 - \varepsilon_2)(s) = (-\varepsilon_3)(s)$ and we get a contradiction as in (ii). Note that $\omega_1 + \omega_2 \succ \omega_3$, so V_{ω} contains the weights of V_{ω_3} . These include the weights $\pm (\varepsilon_1 - \varepsilon_3 + \varepsilon_4)$. If (iv) holds, then $(-\varepsilon_1 + \varepsilon_3 - \varepsilon_4)(s) = (-\varepsilon_2)(s)$, and $(+\varepsilon_1 - \varepsilon_3 + \varepsilon_4)(s) = \varepsilon_2(s)$; as usual, s almost cyclic on V_{ω} implies that $\varepsilon_2(s) = (-\varepsilon_2)(s)$, again a contradiction. Finally, if (v) holds, so that $(2\varepsilon_1)(s) = (-2\varepsilon_1)(s)$, then $(2\varepsilon_1 + \varepsilon_2)(s) = (-2\varepsilon_1 + \varepsilon_2)(s)$ and $(-2\varepsilon_1 + \varepsilon_3)(s) = (2\varepsilon_1 + \varepsilon_3)(s)$, which then implies that $\varepsilon_2(s) = \varepsilon_3(s)$ so $(\varepsilon_2 - \varepsilon_3)(s) = 1$, contradicting that s is regular.

Suppose now that $\omega \succeq \omega_1 + \omega_3 \succ 2\omega_1 \succ \omega_2$. Then $\Omega(V_\lambda) \subseteq \Omega(V_\omega)$, for $\lambda \in \{\omega_1 + \omega_3, 2\omega_1, \omega_2\}$; in particular, $\pm 2\varepsilon_i \pm \varepsilon_j \pm \varepsilon_k$ are weights of V_ω for i, j, k distinct, as are $\pm 2\varepsilon_i$ for $1 \le i \le n$. If (i) holds then $(2\varepsilon_1)(s) = (2\varepsilon_2)(s)$ and $(-2\varepsilon_1)(s) = (-2\varepsilon_2)(s)$. As usual, s almost cyclic on V_ω implies that $(2\varepsilon_1)(s) = (-2\varepsilon_2)(s)$, so that we have $(2\varepsilon_1 + \varepsilon_2 + \varepsilon_3)(s) = (\varepsilon_3 - \varepsilon_2)(s)$. Similarly, we find that $(-2\varepsilon_1 + \varepsilon_2 - \varepsilon_3)(s) = (-\varepsilon_2 - \varepsilon_3)(s)$ and we deduce that $\varepsilon_3(s) = (-\varepsilon_3)(s)$ contradicting that s is regular. If (ii) holds, $(2\varepsilon_1)(s) = (\varepsilon_2 - \varepsilon_1)(s)$, then $(2\varepsilon_1 + \varepsilon_2 + \varepsilon_3)(s) = (-\varepsilon_1 + 2\varepsilon_2 + \varepsilon_3)(s)$ and $(2\varepsilon_1 + \varepsilon_2 - \varepsilon_3)(s) = (-\varepsilon_1 + 2\varepsilon_2 - \varepsilon_3)(s)$. As $2\varepsilon_1 + \varepsilon_2 \pm \varepsilon_3$ and $-\varepsilon_1 + 2\varepsilon_2 \pm \varepsilon_3$ are weights of $V_{\omega_1 + \omega_3}$, it follows that $(2\varepsilon_1 + \varepsilon_2 + \varepsilon_3)(s) = (2\varepsilon_1 + \varepsilon_2 - \varepsilon_3)(s)$, whence $(2\varepsilon_3)(s) = 1$, a contradiction. If (iii) holds, $(2\varepsilon_1)(s) = (\varepsilon_2 - \varepsilon_3)(s)$, then $(2\varepsilon_1 - \varepsilon_2 + \varepsilon_3)(s) = 1$ and $(2\varepsilon_1 - \varepsilon_2 - \varepsilon_3)(s) = (-2\varepsilon_3)(s)$. The former equality means that the multiplicity of the eigenvalue 1 is greater than 1 and the second equality implies that $(-2\varepsilon_3)(s) = 1$, a contradiction. If (iv) holds, $(\varepsilon_1 - \varepsilon_2)(s) = (\varepsilon_3 - \varepsilon_4)(s)$, $n \ge 4$, then

 $(2\varepsilon_1 - \varepsilon_2 - \varepsilon_3)(s) = (\varepsilon_1 - \varepsilon_4)(s)$ and $(-\varepsilon_1 + \varepsilon_3 - 2\varepsilon_4)(s) = (-\varepsilon_2 - \varepsilon_4)(s)$. As s is almost cyclic on V_{ω} , we have $(\varepsilon_1 - \varepsilon_4)(s) = (-\varepsilon_2 - \varepsilon_4)(s)$, whence $\varepsilon_1(s) = (-\varepsilon_2)(s)$, giving the usual contradiction. Finally, if (v) holds, then $(2\varepsilon_1)(s) = (-2\varepsilon_1)(s)$, both of which occur in $V_{2\omega_1}$. We have as well $(2\varepsilon_1 + \varepsilon_2 + \varepsilon_3)(s) = (-2\varepsilon_1 + \varepsilon_2 + \varepsilon_3)(s)$. So we deduce that $(2\varepsilon_1 + \varepsilon_2 + \varepsilon_3)(s) = (2\varepsilon_1)(s)$ giving $(\varepsilon_2 + \varepsilon_3)(s) = 1$ and the final contradiction.

Corollary 3.9. Theorem 1.2 and Theorem 1.4 hold for $G = C_n$, $n \ge 3$ and $p \ne 2$.

Proof. We first let $s \in T$ and $\omega \in \Omega_p^+$, $\omega \neq 0$ such that s is cyclic on V_{ω} . Then all weights of V_{ω} occur with multiplicity 1 and so ω appears in Table 2. In addition, Proposition 3.8 applies. So either s is strongly regular or one of the following holds:

- (i) $\omega = \omega_1$,
- (ii) $\omega = \omega_2 \text{ and } (n, p) = (3, 3),$
- (iii) n = 3 and $\omega = \omega_3$, or
- (iv) $n \ge 4$, p = 3 and $\omega \in \{\omega_{n-1}, \omega_n\}$.

These are precisely the weights listed in the statement of Theorem 1.2.

Now for Theorem 1.4, we let ω be as above, but this time we take $s \in T_{reg}$ and suppose that s is almost cyclic on V_{ω} . Then Proposition 3.8 applies and we find that either s is strongly regular or ω is as listed in (i) - (iv) above, or $\omega = \omega_2$, for $(n, p) \neq (3, 3)$, or n = 4 and $\omega = \omega_4$ with $p \neq 3$. Hence we have the result of Theorem 1.4.

We now consider the group C_2 .

Lemma 3.10. Let $G = C_2$, and let $s \in T_{reg}$. Then

- (1) s is cyclic on V_{ω_1} and almost cyclic on V_{ω_2} . If s is not cyclic on V_{ω_2} then $p \neq 2$ and -1 is an eigenvalue of s on V_{ω_2} of multiplicity 2.
- (2) There exists $s' \in T_{reg}$, with s' is not strongly regular, and s' cyclic on V_{ω_1} and on V_{ω_2} .
- (3) Let $\omega \in \Omega_p^+$, $p \neq 2$, $\omega \notin \{0, \omega_1, \omega_2\}$. If s is almost cyclic on V_{ω} , then s is strongly regular.

Proof. The proof of (1) is straightforward and left to the reader. For (2), define s' by $\varepsilon_1(s') = a$, $\varepsilon_2(s') = a^3$, where $a \in F^{\times}$ and $a^{24} \neq 1$. Then s' is cyclic on V_{ω_1} (and hence regular) s' and is not strongly regular as $2\varepsilon_1(s') = (\varepsilon_2 - \varepsilon_1)(s')$. The eigenvalues of s' on V_{ω_2} are $1, a^{\pm 2}, a^{\pm 4}$ (where 1 is to be dropped if p = 2), so s' is cyclic on V_{ω_2} .

(3) As usual, Theorem 1.6 implies that ω appears in Table 2 or Table 3; we have to examine the cases with $p \neq 2$ and $\omega \in \{2\omega_2, \omega_1 + \frac{p-3}{2}\omega_2, \frac{p-1}{2}\omega_2\}$.

Suppose that s is not strongly regular. As in the proof of Proposition 3.8, we can assume that one of the following holds:

- (i) $(2\varepsilon_1)(s) = (2\varepsilon_2)(s)$,
- (ii) $(2\varepsilon_1)(s) = (\varepsilon_2 \varepsilon_1)(s)$, or
- (iii) $(2\varepsilon_1)(s) = (-2\varepsilon_1)(s)$.

By Lemma 3.7, either $\omega \succeq \omega_1 + \omega_2 \succ \omega_1$ or $\omega \succeq 2\omega_2$.

Suppose first that $\omega \succeq \omega_1 + \omega_2$. Note that $\pm 2\varepsilon_1 \pm \varepsilon_2, \pm \varepsilon_1 \pm 2\varepsilon_2, \pm \varepsilon_1, \pm \varepsilon_2 \in \Omega(V_{\omega_1 + \omega_2})$ If (i) holds, then $(2\varepsilon_1 - \varepsilon_2)(s) = \varepsilon_2(s)$ and $(-2\varepsilon_1 + \varepsilon_2)(s) = -\varepsilon_2(s)$. As s is almost cyclic on V_{ω} , it follows that $\varepsilon_2(s) = -\varepsilon_2(s)$, which is a contradiction as $2\varepsilon_2$ is a root. If (ii) holds then $(2\varepsilon_1)(s) = (\varepsilon_2 - \varepsilon_1)(s)$ implies $(2\varepsilon_1 - \varepsilon_2)(s) = (-\varepsilon_1)(s)$ and $(-2\varepsilon_1 + \varepsilon_2)(s) = \varepsilon_1(s)$. As s is almost cyclic on V_{ω} and $\pm (2\varepsilon_1 - \varepsilon_2), \pm \varepsilon_1 \in \Omega(V_{\omega_1 + \omega_2})$, it follows that $\varepsilon_1(s) = (-\varepsilon_1)(s)$, whence $(2\varepsilon_1)(s) = 1$, which contradicts s being regular. Finally, if (iii) holds, so that

 $(2\varepsilon_1)(s) = (-2\varepsilon_1)(s)$, then $(2\varepsilon_1 \pm \varepsilon_2)(s) = (-2\varepsilon_1 \pm \varepsilon_2)(s)$ which implies as before that $\varepsilon_2(s) = (-\varepsilon_2)(s)$, giving the usual contradiction.

Suppose that $\omega \succeq 2\omega_2$. As $2\omega_2 \succ 2\omega_1 \succ \omega_2$, we observe that $\pm 2\varepsilon_1 \pm 2\varepsilon_2$, $\pm 2\varepsilon_1$, $\pm 2\varepsilon_2$, $\pm \varepsilon_1 \pm \varepsilon_2$ are weights of V_ω . If (i) holds then $\pm (2\varepsilon_1 - 2\varepsilon_2)(s) = 1$ and hence the multiplicity of the eigenvalue 1 of s on V_ω is at least 2. In addition, $(2\varepsilon_1)(s) = (2\varepsilon_2)(s)$, so $(2\varepsilon_2)(s) = 1$ as s is almost cyclic on V_ω . This is a contradiction. If (ii) holds, that is, $(2\varepsilon_1)(s) = (\varepsilon_2 - \varepsilon_1)(s)$ then $(-2\varepsilon_1)(s) = (-\varepsilon_2 + \varepsilon_1)(s)$. So $(\varepsilon_2 - \varepsilon_1)(s) = -1$ and the multiplicity of -1 as an eigenvalue of s on V_ω is at least 2. As $\pm (2\varepsilon_1 - 2\varepsilon_2) \in \Omega(V_{2\omega_2})$ and $(2\varepsilon_1 - 2\varepsilon_2)(s) = 1$, we conclude that 1 is an eigenvalue of s on V_ω is of multiplicity at least 2. This is a contradiction, as s is almost cyclic on V_ω . Finally, if (iii) holds, then we have $(2\varepsilon_1)(s) = (-2\varepsilon_1)(s)$, and so $(2\varepsilon_1 + 2\varepsilon_2)(s) = (-2\varepsilon_1 + 2\varepsilon_2)(s)$ we deduce that $(2\varepsilon_2)(s) = 1$, a contradiction.

Corollary 3.11. Theorem 1.2 and Theorem 1.4 hold for $G = C_2$.

Proof. Recall that for $s \in T$ and $\omega \in \Omega_p^+$, $\omega \neq 0$, if s is almost cyclic on V_ω , then ω appears in Table 2 or Table 3. In particular, for p=2, the only weights which need to be considered are ω_1 and ω_2 and both of these are in the statements of the theorems. Henceforth we assume that $p \neq 2$. If $s \in T$ is cyclic on V_ω for some $\omega \in \Omega_p^+$, $\omega \neq 0$, then again consulting Table 2, we find that $\omega = \omega_1$ or $\omega = \omega_2$, as in the conclusion of Theorem 1.2. So finally, we are left to consider the case where $s \in T_{reg}$, $\omega \in \Omega_p^+$, and s is almost cyclic on V_ω . Then here Lemma 3.10(3) shows that either s is strongly regular or $\omega \in \{0, \omega_1, \omega_2\}$ as claimed in Theorem 1.4.

To complete our consideration of the classical type groups, we must establish the existence of elements as claimed in Theorem 1.5.

Lemma 3.12. Let $G = B_n$ with $n \ge 3$, or $G = C_n$, with p = 2 and $n \ge 3$. Then there exists $s \in T_{reg}$ such that s is not strongly regular, but cyclic on each of the modules V_{ω_1} and V_{ω_n} . Moreover, if $G = B_n$, then we may choose s so that s is cyclic on $V_{\omega_1} \oplus V_{\omega_n} \oplus \delta_{p,2}V_0$.

Proof. For $0 \le i \le n-3$, let $b_i \in F^{\times}$ be a primitive p_i -th root of unity where p_0, \ldots, p_{n-3} are distinct primes greater than 11. Set $X = \operatorname{diag}(b_0, (b_0)^3, (b_0)^5, b_1, \ldots, b_{n-3})$. Now set $t = \operatorname{diag}(X, 1, X^{-1})$, where the 1 is to be dropped if $G = C_n$. For a suitable choice of basis for the natural module for G, we have $t \in \operatorname{SO}_{2n+1}(F)$, respectively $t \in \operatorname{Sp}_{2n}(F)$, for $G = B_n$, respectively $G = C_n$.

We first note that t is cyclic on V_{ω_1} . In addition, $(\varepsilon_2 - \varepsilon_1)(t) = b_0^2 = (\varepsilon_3 - \varepsilon_2)(t)$, so t is not strongly regular.

Consider first the case $G=B_n$ and let $s\in G$ be a semisimple element acting on V_{ω_1} as t does. In particular, s is regular (since cyclic on V_{ω_1}), but s is not strongly regular. Now the weights of $V_{\omega_1}\oplus V_{\omega_n}\oplus \delta_{p,2}V_0$ are $0,\pm \varepsilon_i$ $(1\leq i\leq n),\frac{1}{2}(\pm \varepsilon_1\pm \cdots \pm \varepsilon_n)$. Hence, the remaining claim will follow if we show that the elements $\pm \varepsilon_i(s),1,(\frac{1}{2}(\pm \varepsilon_1\pm \cdots \pm \varepsilon_n)(s))$ are distinct. As |t| is odd, we may assume that |s| is odd as well. Hence, it suffices to show that the eigenvalues $2\varepsilon_i(s),1,(\pm \varepsilon_1\pm \cdots \pm \varepsilon_n)(s)$ are distinct. The elements $(\pm \varepsilon_1\pm \cdots \pm \varepsilon_n)(s)$ have the form $(b_0)^{\pm m}x$, for $m\in\{1,3,7,9\}$ and for some $x\in\{b_1^{\pm 1}\cdots b_{n-3}^{\pm 1}\}$. As $p_i>11$, all these elements are distinct, and differ from $1,(b_0)^{\pm 2},(b_0)^{\pm 6},\ b_0^{10},\ and\ b_1^{\pm 2},\ldots,b_{n-3}^{\pm 2},\ whence the result.$

The argument for $G = C_n$ is entirely similar as the weights of V_{ω_n} are $\pm \varepsilon_1 \pm \cdots \pm \varepsilon_n$. \square

Lemma 3.13. Let $G = D_n$, $n \ge 4$. Then there exists $s \in T_{reg}$ such that s is not almost cyclic on V_{ω_2} (equivalently, s is not strongly regular), but cyclic on $V_{\omega_1} \oplus V_{\omega_{n-1}} \oplus V_{\omega_n} \oplus V_0$.

Proof. As $\omega_a = \omega_2$, by Lemma 2.7, $s' \in T$ is almost cyclic on V_{ω_2} if and only if s' is strongly regular. We now exhibit the existence of the s as claimed in the statement.

The group $G=D_n$ is a maximal rank subgroup of B_n generated by root subgroups corresponding to long roots. The element s introduced in Lemma 3.12 is therefore contained in a subgroup $G=D_n$, and s is regular as an element of D_n , since it is regular in B_n . Moreover, we have seen in the proof of Lemma 3.12 that $(\varepsilon_2 - \varepsilon_1)(s) = (\varepsilon_3 - \varepsilon_2)(s)$; as $\varepsilon_2 - \varepsilon_1$ and $\varepsilon_3 - \varepsilon_2$ are long roots of B_n , they are roots of the subgroup D_n . So s is not strongly regular in D_n and by Lemma 2.7, it follows that s is not almost cyclic on V_{ω_2} . Furthermore, for the purposes of this proof, let η_1, \ldots, η_n denote the fundamental dominant weights of B_n . Then, the irreducible module for B_n with highest weight η_n restricts to D_n as $V_{\omega_{n-1}} \oplus V_{\omega_n}$. In addition, $V_{\eta_1}|_H = V_{\omega_1} \oplus (1 - \delta_{p,2})V_0$. Now Lemma 3.12 shows that s is cyclic on $V_{\eta_1} + V_{\eta_n} + \delta_{p,2}V_0$, and so s is cyclic on $V_{\omega_1} + V_{\omega_{n-1}} + V_{\omega_n} + V_0$, as claimed. \square

Remark 3.14. Note that the above two results provide a proof of Theorem 1.5(1) for $G = B_n$, $n \ge 3$, $G = D_n$ with $n \ge 4$, and $G = C_n$ for $n \ge 3$ and p = 2.

Lemma 3.15. Let $G = C_n$, $n \ge 2$. Then there exists $s \in T_{reg}$, not strongly regular, with s almost cyclic on V_{ω_2} .

Proof. Define s by $\varepsilon_1(s) = a$, $\varepsilon_2(s) = a^3$, $\varepsilon_i(s) = b_i$, for $a, b_1, \ldots, b_{n-2} \in F^{\times}$. Then s is not strongly regular as $(2\varepsilon_1)(s) = a^2 = (\varepsilon_2 - \varepsilon_1)(s)$, and $(2\varepsilon_1)(s)$, $\varepsilon_2 - \varepsilon_1$ are roots of G. Moreover, one may choose the a, b_i such that s is regular. The eigenvalues of s on V_{ω_2} are $1, a^{\pm 2}, a^{\pm 4}, a^{\pm 1}b_i^{\pm 1}, a^{\pm 3}b_i^{\pm 1}, b_i^{\pm 1}b_j^{\pm 1}, i \neq j$. We may choose $a, b_1, \ldots, b_{n-2} \in F^{\times}$ such that the eigenvalues different from 1 are distinct. Indeed, the non-zero weights of V_{ω_2} are roots and by choosing s regular, we know that the multiplicity of the eigenvalue 1 is precisely the multiplicity of the zero weight in V_{ω_2} , while all other weights have multiplicity 1. Hence s is almost cyclic on V_{ω_2} .

Remark 3.16. A similar argument works for $G = B_n$, with p = 2; for this case, set $\varepsilon_1(s) = a$ and $\varepsilon_2(s) = a^2$ and the remaining values as for C_n .

Lemma 3.17. Let $G = C_n$, $n \ge 2$. Then there exists $s \in T_{reg}$, s not strongly regular, which separates the weights of V_{ω_k} , for $k = 1, \ldots, n$.

Proof. Let $a, b_i \in F^{\times}$, for $1 \leq i \leq n$, and define $s \in T$ by $\varepsilon_i(s) = b_i$ for $1 \leq i \leq n$ and in addition $b_1 = a$ and $b_2 = a^3$. We first note that s is not strongly regular as $2\varepsilon_1(s) = (\varepsilon_2 - \varepsilon_1)(s)$. We now show that by choosing appropriate values for the b_i , the element s separates the weights of V_{ω_k} , as claimed. This will also show that s is regular as s is then cyclic on V_{ω_1} .

The non-zero weights of V_{ω_k} are $\pm \varepsilon_{i_1} \pm ... \pm \varepsilon_{i_j}$ for $j \leq k$ with k-j even and $1 \leq i_1 < ... < i_j \leq n$. This follows from the fact that V_{ω_j} occurs as a composition factor of $\wedge^j(V_{\omega_1})$, for all $1 \leq j \leq n$, and the subdominant weights of ω_k , for $k \geq 2$, are of the form ω_{k-2i} for $0 \leq i \leq \lfloor \frac{k}{2} \rfloor$. A weight $\pm \varepsilon_{i_1} \pm ... \pm \varepsilon_{i_j}$ takes the value $b_{i_1}^{\pm 1} \cdots b_{i_j}^{\pm 1} = xy$, on s, where $x = a^{\pm m}$ for some $0 \leq m \leq 4$ and y is a product of certain $b_i^{\pm 1}$, with i > 2. (In particular, if n = 2 then the weights of V_{ω_k} take values among $a^{\pm 1}, a^{\pm 3}, a^{\pm 2}, a^{\pm 4}, 1$ on s.) For a suitable choice of a and b_i , these values are distinct. For example, we can choose b_i to be a primitive p_i -th

root of unity for $3 \le i \le n$ for primes $p_3 > ... > p_n$, and a to be a primitive p_0 -th root of unity for a prime $p_0 > \max\{p_3, 7\}$. Therefore, s separates the weights of V_{ω_k} for every fixed k.

Proposition 3.18. Let $G = C_n, n \geq 2, p \neq 2$.

- (1) Let $\omega \in \Omega_n^+$ be as follows:
- (i) $\omega = \omega_1$,
- (ii) $\omega \in \{\omega_{n-1}, \omega_n\}, n \geq 3, p = 3, or$
- (iii) $\omega = \omega_3$, n = 3, $p \neq 3$.

Then there exists $s \in T_{reg}$, with s not strongly regular but cyclic on V_{ω} .

- (2) Let $\omega \in \Omega_p^+$ be one of:
- (i) $\omega = \omega_4$, n = 4 and $p \neq 3$, or
- (ii) $\omega = \omega_2$ with $(n, p) \neq (3, 3)$.

Then there exists $s \in T_{reg}$, with s not strongly regular but almost cyclic on V_{ω} .

Proof. This follows from Lemma 3.17 and Tables 2 and 3. Indeed, if s separates the weights of V then s is cyclic on V whenever all weights of V are of multiplicity 1, and almost cyclic if exactly one weight is of multiplicity greater than 1.

Lemma 3.19. Theorem 1.5 is true if G is of classical Lie type.

Proof. For G of type A_n this was established in Section 3.1. For Theorem 1.5(1), we refer to Remark 3.14 and note that it remains to consider $G = C_2$ and $G = C_n$ with $n \geq 3$ and $p \neq 2$. For $G = C_2$, we must establish the existence of $s \in T$, cyclic on V_{ω_1} and on V_{ω_2} , with s not strongly regular. This follows from Lemma 3.10(2).

For $G = C_n$, $n \ge 3$ and $p \ne 2$, we must establish the existence of $s \in T$, s not strongly regular, with s cyclic on

- (a) V_{ω_1} ,
- (b) V_{ω_3} when n=3, and
- (c) $V_{\omega_{n-1}}$ or V_{ω_n} , when p=3.

(If necessary, we may have different s for each of the modules.) This is precisely the statement of Proposition 3.18(1).

Now we turn to the list of modules to be treated for Theorem 1.5(2). For $n \geq 3$, $\omega = \omega_2$, and $G = C_n$, or $G = B_n$ with p = 2, the result follows from Lemma 3.15 and Remark 3.16. For $G = C_4$ with $\omega = \omega_4$, $p \neq 2, 3$, the result follows from Proposition 3.18(2).

4. Groups of exceptional types and Theorem 1.7

In this section we will establish all of the main results in case G is of exceptional type and deduce Theorem 1.7.

Lemma 4.1. Theorem 1.4 holds for G of exceptional type.

Proof. Let $\omega \in \Omega_p^+$ and $s \in T$ be almost cyclic on V_ω . Then Theorem 1.6 and Tables 2 and 3 imply that either $\omega = \omega_a$, or ω is the highest short root (in case $G = G_2$ or F_4), or ω is a miniscule weight. The miniscule weights appear in the conclusion of Theorem 1.4, as well as the highest short roots. Moreover, Lemma 2.7 shows that if s is almost cyclic on V_a , then either s is strongly regular or $(G, p) = (G_2, 3)$ or $(F_4, 2)$. The latter configurations are also in the list of exceptions in the statement of Theorem 1.4, hence the result.

Lemma 4.2. Theorem 1.2 holds for G of exceptional type.

Proof. Let $\omega \in \Omega_p^+$ and $s \in T$ be cyclic on V_ω . Then all weights of V_ω have multiplicity 1 and now comparing the statement of Theorem 1.2 with Table 2 gives the result.

We are left with proving Theorem 1.5; we treat each of the groups in turn.

Proposition 4.3. Let G be of type F_4 . There exists $s \in T_{reg}$ such that s is almost cyclic on V_{ω_4} but not strongly regular. If p = 3 then s can be chosen to be cyclic on V_{ω_4} .

Proof. Let $V = V_{\omega_4}$. Recall that G contains a subsystem subgroup H of type B_4 , generated by the long root subgroups of G, so $T \leq H$. For the purposes of this proof, let η_i , $1 \leq i \leq 4$ denote the fundamental dominant weights of $H = B_4$ (with respect to an appropriate choice of Borel subgroup of H). Then the non-zero weights of $V|_H$ are exactly the non-zero weights of the FH-module $V_{\eta_1} + V_{\eta_4}$. By comparison of the dimensions we conclude that the non-trivial composition factors of $V|_H$ are V_{η_1} and V_{η_4} , and there are $(1 - \delta_{p,3})(1 + \delta_{p,2})$ trivial composition factors.

Now let $s \in H$ be as in Lemma 3.12; then, $\alpha(s) = \beta(s)$, for α and β distinct roots of H, so s is not strongly regular in F_4 . Moreover, s is cyclic on $V_{\eta_1} + V_{\eta_4}$ and so almost cyclic on V. Then Theorem 1.3 implies that s is regular in G. If p = 3, then $V|_H = V_{\eta_1} + V_{\eta_4}$ and Lemma 3.12 shows that s is cyclic on V.

Remark 4.4. Let $G = F_4$, p = 2.

- (1) There exists an element $s \in T_{reg}$, s not strongly regular, which is almost cyclic on $V_a = V_{\omega_1}$ (and not almost cyclic on V_{ω_4}). Recall that, identifying T with $(F^{\times})^4$ and ε_i the i-th coordinate function for $1 \le i \le 4$, the roots of F_4 are $\pm \varepsilon_i, \frac{1}{2}(\pm \varepsilon_1 \pm \varepsilon_2 \pm \varepsilon_3 \pm \varepsilon_4)$ (short) and $\pm \varepsilon_i \pm \varepsilon_j$, $1 \le i < j \le 4$ (long). Let $a_i \in F^{\times}$ be a primitive p_i -root of unity, where $p_1 = 3, p_2 = 5, p_3 = 7$. Define $s \in T$ by $\varepsilon_1(s) = a_1, \varepsilon_2(s) = a_2, \varepsilon_3(s) = a_1a_2, \varepsilon_4(s) = a_3$. The non-zero weights of V_a are $\pm \varepsilon_i \pm \varepsilon_j$ ($1 \le i < j \le 4$) so s is almost cyclic on V_a . Then $(\varepsilon_1 + \varepsilon_2 \varepsilon_3 + \varepsilon_4)(s) = (-\varepsilon_1 \varepsilon_2 + \varepsilon_3 + \varepsilon_4)(s) = \varepsilon_4(s) = a_3$ so s is not almost cyclic on V_{ω_4} . Note that s is regular as $\alpha(s) \ne 1$ for every root α , but it is not strongly regular. This justifies one of the claims of Remark 2.8, and together with Proposition 4.3, completes the proof of Theorem 1.5 for the group $G = F_4$.
- (2) There exists a non-regular element $s \in T$ such that $V_a^s = V_a^T$. Indeed, let $\varepsilon_i(s) = t_i \in F^\times$, where $t_1 = 1$. Then s is not regular. The weights of $V_a = V_{\omega_1}$ are $0, \pm \varepsilon_i \pm \varepsilon_j$, so the eigenvalues of s on the non-zero weights of V_a are $1, (\pm \varepsilon_i \pm \varepsilon_j)(s)$, that is, $1, t_i^{\pm 1}, t_i^{\pm 1}t_j^{\pm 1}$, where $i, j = 2, 3, 4, i \neq j$. For an appropriate choice of $t_i \in F^\times$, these eigenvalues differ from 1, and the claim follows. This proves a claim in Remark 2.8 for the group $G = F_4$.

Proposition 4.5. Let $G = E_6$. Then there exists $s \in T$, not strongly regular, which is cyclic on V_{ω_1} and on V_{ω_6} .

Proof. It suffices to consider $V = V_{\omega_1}$. Let H < G be a subsystem subgroup of type D_5 . For the purposes of this proof, let η_i , for $1 \le i \le 5$, denote the fundamental dominant weights of H (with respect to an appropriate choice of maximal torus T_H and Borel subgroup of H). Then, we have that $V|_H = V_{\eta_1} \oplus V_{\eta_4} \oplus V_0$. (Here we have made a choice of the labeling of the roots of H.)

Now let $s \in H$ be as in Lemma 3.13. Then s is cyclic on $V_{\eta_1} + V_{\eta_4} + V_{\eta_5} + V_0$, so in particular on V. As well, s is not strongly regular in H and since roots of H are roots of G, s is not strongly regular in G.

Proposition 4.6. Let G be of type E_6 . Then there exists $s \in T_{reg}$ such that s is not strongly regular but almost cyclic on V_{ω_1} and on V_{ω_6} .

Proof. It suffices to establish the result for V_{ω_1} . Let $H \leq G$ be of type F_4 . Let s be as constructed in the proof of Proposition 4.3. For the purposes of this proof, let η_i , $1 \leq i \leq 4$ denote the fundamental dominant weights of H (with respect to a fixed choice of Borel subgroup and maximal torus). Then $V_{\omega_1}|_{H} = V_{\eta_4} \oplus (1 + \delta_{p,3})V_0$. Since s is almost cyclic on V_{η_4} , and cyclic if p = 3, and the zero weight has multiplicity 2 in V_{η_4} if $p \neq 3$, we see that s is almost cyclic on V_{ω_1} . We claim that s is regular but not strongly regular. Recall that $V_a = V_{\omega_2}$. We note that

$$V_{a|H} = \begin{cases} V_{\eta_1} \oplus V_{\eta_4}, & \text{if } p \neq 2, 3; \\ V_{\eta_1} \oplus V_{\eta_4} \oplus V_0, & \text{if } p = 3; \\ V_{\eta_1} \oplus V_{\eta_4} \oplus V_{\eta_4}, & \text{if } p = 2. \end{cases}$$

To see that s is regular, we need to show that the multiplicity of the eigenvalue 1 for s on V_a is equal to dim T=6. Note that s is regular in H and so the eigenvalue 1 on V_{η_1} has multiplicity $4-2\delta_{p,2}$. Also, the non-zero weights of V_{η_4} are roots of H and the zero weight has multiplicity $2-\delta_{p,3}$ in V_{η_4} ; so again the eigenvalue 1 has multiplicity $2-\delta_{p,3}$ in V_{η_4} . In each of the above decompositions, we find that the eigenvalue 1 for s on V_a has multiplicity 6.

Finally, to see that s is not strongly regular, we appeal to Lemma 2.7. It suffices to see that s is not almost cyclic on V_a . Recall that $\eta_4 \prec \eta_1$. Using Theorem 2.5 and the above decompositions of $V_a|_H$, we find that all non-zero weights of V_{η_4} occur with multiplicity 2 in V_a , whence the result.

We will use the following result when treating the group $G = E_7$.

Proposition 4.7. Let $G = A_n$, $n \ge 4$. Then there exists $s \in T_{reg}$, s not strongly regular, such that s is not almost cyclic on $V_{\omega_1+\omega_n}$ and cyclic on $V = V_{\omega_2} \oplus V_{\omega_{n-1}}$.

Proof. We start by considering a certain subvariety of F^m . Let $m \geq 4$. Set $X = \{(x_1,\ldots,x_m) \in F^m \mid x_1\cdots x_m=1\}$. Then X is isomorphic to the principal open subset $D(f) \subset F^{m-1}$ defined by the function $f \in F[T_1,\ldots,T_{m-1}], f=T_1\cdots T_{m-1}$. Indeed, the associated F-algebra of X is isomorphic to $F[T_1,\ldots,T_{m-1},(T_1\cdots T_{m-1})^{-1}]$ which is (isomorphic to) the localization of $F[T_1,\ldots,T_{m-1}]$ with respect to the multiplicative set $S = \{(T_1\cdots T_{m-1})^\ell \mid \ell \geq 0\}$. In particular, X is irreducible of dimension m-1. Now consider the closed subset Y of X, $Y = \{(x_1,\ldots,x_m) \in X \mid x_1^2 = x_2x_3\}$. Then the associated affine F-algebra is $F[X]/(T_1^2-T_2T_3)$. Since F[X] is a localization of $F[T_1,\ldots,T_{m-1}]$ and $T_1^2-T_2T_3$ is an irreducible polynomial function in $F[T_1,\ldots,T_{m-1}]$, Y is also an irreducible subvariety of F^m , of dimension m-2.

We now apply the above reasoning to a maximal torus of $G = \mathrm{SL}_{n+1}(F)$, namely the usual torus of diagonal matrices of determinant 1. Set $s = \mathrm{diag}(d_1, \ldots, d_{n+1}) \in T$, where $d_i \in F^{\times}$ and $\prod_{i=1}^{n+1} d_i = 1$. If we assume further that $d_1^2 = d_2 d_3$ then $(\varepsilon_1 - \varepsilon_2)(s) = (\varepsilon_3 - \varepsilon_1)(s)$ and so s is not almost cyclic on $V_{\omega_1 + \omega_n}$. Let $Y = \{s \in T \mid d_1^2 = d_2 d_3\}$ so that by the discussion

of the first paragraph, Y is an irreducible closed subvariety of T of dimension n-1. The module V has 1-dimensional T-weight spaces. For weights $\lambda, \mu \in \Omega(V)$, with $\lambda \neq \mu$, let $K_{\lambda\mu} = \ker(\lambda - \mu)$. The set $K := \bigcup_{\lambda,\mu} K_{\lambda\mu}$ is a proper closed subset of T. Since Y is irreducible, if $Y \subseteq K$, then there exist $\lambda, \mu \in \Omega(V)$, with $\lambda \neq \mu$, such that $Y \subseteq \ker(\lambda - \mu)^{\circ}$ and dimension considerations show that these two sets must be equal. But the weights of V are of the form $\pm(\varepsilon_i + \varepsilon_j)$, $1 \leq i < j \leq n+1$. So the possible weight differences $\lambda - \mu$ (up to a sign) are $\varepsilon_i + \varepsilon_j \pm (\varepsilon_k + \varepsilon_\ell)$, $\varepsilon_j - \varepsilon_\ell$, $2\varepsilon_i + \varepsilon_j + \varepsilon_\ell$ and $2(\varepsilon_i + \varepsilon_j)$, for i, j, k, ℓ distinct. So there are no distinct weights λ, μ with $Y = \ker(\lambda - \mu)^{\circ}$. Hence Y does not lie in K and the elements in $Y \setminus K$ are almost cyclic, indeed cyclic, on V and not almost cyclic on $V_{\omega_1 + \omega_n}$. So these elements are regular by Lemma 2.2 and not strongly regular by Lemma 2.7. \square

We now use the preceding result to establish the following

Proposition 4.8. Let $G = E_7$. There exists an element $s \in T_{reg}$, with s not strongly regular, and such that s is cyclic on V_{ω_7} .

Proof. Set $V = V_{\omega_7}$. Let $H \leq G$ be a maximal rank subgroup of type A_7 and for the purposes of this proof, let $\eta_i, 1 \leq i \leq 7$, denote the fundamental dominant weights of H (with respect to an appropriate choice of maximal torus T_H and Borel subgroup). Then $V|_H = V_{\eta_2} + V_{\eta_6}$. Let $s \in T_H$, be as in Proposition 4.7, so that s is cyclic on V and not strongly regular in H. Then s is not strongly regular in G as the roots of G. Finally, Lemma 2.2 shows that s is regular.

For the proof of Theorem 1.5, it remains to treat the group $G = G_2$. We investigate in detail the two modules corresponding to the fundamental dominant weights.

Proposition 4.9. Let $G = G_2$, $V_i = V_{\omega_i}$ for i = 1, 2.

- (1) Let $s \in T_{reg}$ and suppose that s is not cyclic on V_1 . Then $p \neq 2$ and the eigenvalues of s on V_1 are $\{1, -1, -1, -b, b^{-1}, b, -b^{-1}\}$, where $b \in F^{\times}$, $b^2 \neq 1$. In particular, s is almost cyclic on V_1 if and only if $b^2 \neq -1$.
- (2) Let $s \in T_{reg}$ and assume that p = 3. Then s is almost cyclic on V_2 and cyclic on V_1 . More precisely, if s is not cyclic on V_2 then, for some $a \in F^{\times}$, $a^4 \neq 1$, the eigenvalues of s on V_2 are $\{1, -1, -1, a^{\pm 3}, -a^{\pm 3}\}$, and those on V_1 are $\{1, a^{\pm 1}, -a^{\pm 1}, -(a^{\pm 2})\}$.
- (3) There exists $s \in T_{reg}$, not strongly regular, with s cyclic on V_1 .
- (4) Let p = 3. Then there exists $s \in T_{reg}$, not strongly regular, with s cyclic on V_2 .
- (5) Let p = 3. Then there exists a non-regular element $s \in T$, with $V_a^s = V_a^T$.

Proof. Set $\alpha_1(s) = a$, $\alpha_2(s) = b$, for some $a, b \in F^{\times}$.

(1) Here we suppose s to be regular and not cyclic on V_1 . The weights of V_1 are the short roots, and the weight 0 if $p \neq 2$, all with multiplicity 1. Then $(\alpha_1 + \alpha_2)(s) = ab$, $(2\alpha_1 + \alpha_2)(s) = a^2b$ so the eigenvalues of s on V_1 are 1 (if $p \neq 2$), $a^{\pm 1}$, $(ab)^{\pm 1}$, $(a^2b)^{\pm 1}$. Since s is not cyclic on V_1 , at least two of the eigenvalues are equal. As s is regular, $\alpha(s) \neq 1$ for every root α , in particular, $1 \notin \{a, ab, a^2b\}$. As W(G) is transitive on the short roots, we can assume that $a \in \{a^{-1}, (ab)^{\pm 1}, (a^2b)^{\pm 1}\}$. Now if $a \in \{(ab)^{\pm 1}, (a^2b)^{\pm 1}\}$, then $1 \in \{b, a^2b, ab, a^3b\}$. But $\alpha_2, 2\alpha_1 + \alpha_2$, $\alpha_1 + \alpha_2, 3\alpha_1 + \alpha_2$ are roots, so this contradicts s being regular. So we have $a = a^{-1}$. Then $a^2 = 1$ and $a = \alpha_1(s) \neq 1$, so $p \neq 2$ and a = -1. Then the eigenvalues of s on V_1 are $1, -1, -1, -b, -b^{-1}, b, b^{-1}$, where $b \neq \pm 1$ as s is regular, establishing the first claim. Finally, s is almost cyclic on V_1 if and only if $b^2 \neq -1$.

(2) Here we suppose s to be regular and p=3. The weights of V_2 are the long roots and the weight 0, so the eigenvalues of s on V_2 are $1, b^{\pm 1}, (a^3b)^{\pm 1}, (a^3b^2)^{\pm 1}$. Note that s regular implies that $b \neq 1$. As above, if s is not cyclic on V_2 , we can assume that $\alpha_2(s) \in \{-\alpha_2(s), \pm (3\alpha_1 + \alpha_2)(s), \pm (3\alpha_1 + 2\alpha_2)(s)\}$. As s is regular, $\alpha_2(s) \neq (-3\alpha_1 - \alpha_2)(s)$ and $\alpha_2(s) \neq (3\alpha_1 + 2\alpha_2)(s)$. If $\alpha_2(s) = (3\alpha_1 + \alpha_2)(s)$ then $(3\alpha_1)(s) = 1$, so $a^3 = 1$, and if $\alpha_2(s) = (-3\alpha_1 - 2\alpha_2)(s)$, then $(ab)^3 = 1$. But since p=3, this implies that a=1, respectively ab=1, contradicting that s is regular. So these two cases are ruled out, and we are left with the case where $\alpha_2(s) = -\alpha_2(s)$. So $b^2 = 1$, and hence b = -1, and $a^2 \neq 1$ since s is regular.

We now have that the eigenvalues of s on V_2 are $\{1, -1, -1, a^3, -a^3, a^{-3}, -a^{-3}\}$. Suppose for a contradiction that s is not almost cyclic on V_2 ; then either $a^3 = -1$ or $a^6 = -1$. In the first case, a = -1, and hence ab = 1 contradicting s regular. Thus $a^6 = -1$, equivalently, $a^2 = -1$. But then, $(2\alpha_1 + \alpha_2)(s) = a^2b = 1$, again contradicting that s is regular. Hence s is almost cyclic on V_2 as claimed. Note that $a^2 \neq -1$, else $(2\alpha_1 + \alpha_2)(s) = 1$. Now the eigenvalues of s on V_1 are $\{1, a, a^{-1}, -a, -a^{-1}, -a^2, -a^{-2}\}$, and hence s is cyclic on V_1 .

- (3) Let a = b, so that the eigenvalues of s on V_1 are $\{a^{\pm 3}, a^{\pm 2}, a^{\pm 1}, 1\}$, where 1 is to be dropped if p = 2; they are distinct if $a^k \neq 1$ for $k \leq 6$; for instance choose a to be a primitive 7-th root of unity. Then s is cyclic on V_1 (hence regular) and is not strongly regular as $\alpha_1(s) = \alpha_2(s) = a$.
- (4) Take again a = b, so that the eigenvalues of s on V_2 are $a^{\pm 1}, a^{\pm 4}, a^{\pm 5}, 1$. Then choosing a appropriately, we see that s is cyclic on V_2 , regular and not strongly regular, as in (3).

The final statement (5) is straightforward, as we may take a=1 so that s is not regular, while the non-zero weights of V_a are $\pm(3\alpha_1+2\alpha_2)$, $\pm(3\alpha_1+\alpha_2)$ and $\pm\alpha_2$; so that choosing b such that $b^2 \neq 1$, we have $V_a^s = V_a^T$.

We have now completed the proof of Theorem 1.5 for the exceptional groups. Finally, we conclude the article by proving Theorem 1.7:

Proof of Theorem 1.7. By Lemma 2.4, $\phi_i(s)$ is cyclic for every i. Therefore, by Theorem 1.2, s is strongly regular unless, possibly, for every i, (G, λ_i) is as in cases (2) - (10) there. Also, Theorem 1.6 implies that all non-zero weights of V_{ω} have multiplicity one, and we then apply [11, Theorem 2(2)] and [15, Proposition 2] to justify the conditions given in (i) - (iii) of the statement.

5. Tables

Tables 2 and 3 are taken from [11, Tables 1,2]. We recall here our convention for reading the tables in case $\operatorname{char}(F)=0$: for a natural number a the expressions $p>a,\ p\geq a$ or $p\neq a$ are to be interpreted as the absence of any restriction, that is, a is allowed to be any natural number. Note further that when a weight λ has coefficients expressed in terms of p, we are assuming that $\operatorname{char}(F)=p>0$.

G						
A_1	$a\omega_1, 1 \le a < p$					
$A_n, n > 1$	$a\omega_1, b\omega_n, \ 1 \le a, b < p$					
	$\omega_i, \ 1 < i < n$					
	$c\omega_i + (p-1-c)\omega_{i+1}, \ 1 \le i < n, \ 0 \le c < p$					
$B_n, n > 2$	$\omega_1, \ \omega_n$					
$C_n, n > 1, p = 2$	$\omega_1,\;\omega_n$					
$C_2, p > 2$	$\omega_1, \ \omega_2, \ \omega_1 + \frac{p-3}{2}\omega_2, \ \frac{p-1}{2}\omega_2$					
C_3	ω_3					
$C_n, n > 2, p > 2$	$\omega_1, \ \omega_{n-1} + \frac{p-3}{2}\omega_n, \ \frac{p-1}{2}\omega_n$					
$D_n, n > 3$	$\omega_1, \ \omega_{n-1}, \ \omega_n$					
E_6	$\omega_1,\;\omega_6$					
E_7	ω_7					
$F_4, p = 3$	ω_4					
$G_2, p \neq 3$	ω_1					
$G_2, p = 3$	$\omega_1,\;\omega_2$					

Table 2. Non-trivial irreducible p-restricted G-modules with all weights of multiplicity 1

G	conditions		weight 0 multiplicity
A_n ,	$n > 1, (n, p) \neq (2, 3)$	$\omega_1 + \omega_n$	$\begin{cases} n-1 & \text{if } p (n+1) \\ n & \text{if } p \not (n+1) \end{cases}$
A_3	$p > 3$ $n > 2, p \neq 2$	$2\omega_2$	2
B_n	$n > 2, p \neq 2$ n > 2, p = 2	ω_2 ω_2	$n - \gcd(2, n)$
	_	$2\omega_1$	$\begin{cases} n & \text{if } p (2n+1) \\ n+1 & \text{if } p \not (2n+1) \end{cases}$
C_n	n > 1	$2\omega_1$	n
	$n > 2, (n, p) \neq (3, 3)$	ω_2	$\begin{cases} n-2 & \text{if } p n\\ n-1 & \text{if } p \not n \end{cases}$
	$p \neq 5$	$2\omega_2$	2
C_4	$p \neq 2, 3$	ω_4	2
D_n	n > 3	$2\omega_1$	$\begin{cases} n-2 & \text{if } p n\\ n-1 & \text{if } p \not\mid n \end{cases}$
D	$n > 3, p \neq 2$	ω_2	n
D_n	n > 3, p = 2	ω_2	$n - \gcd(2, n)$
E_6		ω_2	$\begin{cases} 5 & \text{if } p = 3 \\ 6 & \text{if } p \neq 3 \end{cases}$
E_7		ω_1	$\begin{cases} 6 & \text{if } p = 2 \\ 7 & \text{if } p \neq 2 \end{cases}$
E_8		ω_8	8
F_4		ω_1	$\begin{cases} 2 & \text{if } p = 2\\ 4 & \text{if } p \neq 2 \end{cases}$
	$p \neq 3$ $p \neq 3$	ω_4	2
G_2	$p \neq 3$	ω_2	2

Table 3. Irreducible p-restricted G-modules with non-zero weights of multiplicity 1 and whose zero weight has multiplicity greater than 1.

$A_1 \\ p \neq 2$	A_n $n \ge 2$	B_n $n \ge 3$	C_2	C_n $n \ge 3, p = 2$	$C_3 \\ p \neq 2, 3$	C_n $n \ge 3, p = 3$
ω_1	ω_1,\ldots,ω_n	ω_1, ω_n	ω_1, ω_n	ω_1, ω_n	ω_1,ω_3	$\omega_1, \omega_{n-1}, \omega_n$
$D_n, n \ge 4$	E_6	E_7	$F_4, p = 3$	$G_2, p \neq 3$	$G_2, p = 3$	
$\omega_1, \omega_{n-1}, \omega_n$	ω_1,ω_6	ω_7	ω_4	ω_1	ω_1,ω_2	

Table 4. Highest weights of the non-trivial p-restricted modules in cases (2) - (10) of Theorem 1.2

References

- [1] N. Bourbaki, Groupes et algebres de Lie, Chapitres 4,5 et 6, Masson, Paris, 1981.
- [2] N. Bourbaki, Groupes et algebres de Lie, Chapitres 7 et 8, Hermann, Paris, 1975.
- [3] J. Humphreys, Modular representations of finite groups of Lie type, Cambridge Univ. Press, Cambridge, 2006.
- [4] F. Lübeck, Small degree representations of finite Chevalley groups in defining characteristic, *LMS J. Comput. Math.* 4(2001), 135 169.
- [5] F. Lübeck, Tables of weight multiplicities, on-line data http://www.math.rwth-aachen.de/~Frank.Luebeck/chev/WMSmall/index.html.
- [6] G. Malle and D. Testerman, Linear algebraic groups and finite groups of Lie type, Cambridge Univ. Press, Cambridge, 2011.
- [7] A.A. Premet, Weights of infinitesimally irreducible representations of Chevalley groups over a field of prime characteristic, *Math. USSR Sb.* **61** (1988), 167 183.
- [8] G.M. Seitz, The Maximal Subgroups of Classical Algebraic Groups, Memoirs Amer. Math. Soc., no. 365, Amer. Math. Soc., Providence, 1987.
- [9] G. M. Seitz, Bounds for dimensions of weight spaces of maximal tori, Contemporary Math. 153, 1993, 157 – 161.
- [10] T. Springer and R. Steinberg, Conjugacy classes, In: A. Borel et al., Seminar on algebraic groups and related finite groups, Lect. Notes in Math. vol. 131, Springer Verlag, Berlin, 1970, 167 – 266.
- [11] D. Testerman and A. Zalesski, Subgroups of simple algebraic groups containing maximal tori and representations with multiplicity 1 non-zero weights, *Transformation Groups* 20(2015), 831 861.
- [12] D. Testerman and A.E. Zalesski, Spectra of non-regular elements in irreducible representations of simple algebraic groups, *North-Western European J. Math.* 7(2021), 7(2021), 185 212.
- [13] A. Zalesski, Unisingular representations of finite symplectic groups, *Comm. Algebra* (appeared online)
- [14] A. Zalesski, Singer cycles in 2-modular representations of $GL_n(2)$, Archiv der Math. 110(2018), 433 446
- [15] A.E. Zalesski and I. Suprunenko, The representations of the dimensions $(p^n \mp 1)/2$ of the symplectic group of degree 2n over a field of characteristic p (in Russian), $Vesti\ AN\ BSSR,\ Ser.\ fiz-mat.\ navuk$ 1987, no.6, 9 15. (English translation: arXiv:2108.10650v2 [math.RT] 9 Oct 2021.)

Donna Testerman, Ecole Polytechnique Fédéral de Lausanne, Institute of Mathematics, Station 8, CH-1015 Lausanne, Switzerland E-mail: donna.testerman@epfl.ch

ALEXANDRE E. ZALESSKI, DEPARTMENT OF PHYSICS, MATHEMATICS AND INFORMATICS, ACADEMY OF SCIENCES OF BELARUS, 66 PROSPEKT NEZALEJNASTI, MINSK, 220000, BELARUS E-MAIL: ALEXANDRE.ZALESSKI@GMAIL.COM