

Large cycles in generalized Johnson graphs

Vladislav Kozhevnikov¹, Maksim Zhukovskii²

Abstract

We count cycles of an unbounded length in generalized Johnson graphs. Asymptotics of the number of such cycles is obtained for certain growth rates of the cycle length.

1 Introduction and new results

For integers $i \leq j$, everywhere below we denote $[i, j] := \{i, i+1, \dots, j\}$ and $[i] := [1, i]$. For integers n, r, s such that $0 \leq s < r < n$, a simple graph $G(n, r, s)$ with the set of vertices

$$V := V(G(n, r, s)) = \{x \subset [n] : |x| = r\}$$

and the set of edges

$$E := E(G(n, r, s)) = \{\{x, y\} : |x \cap y| = s\}$$

is called a *generalized Johnson graph*.

Unfortunately, there is no established term for graphs $G(n, r, s)$. In literature they appear as *generalized Johnson graphs* [1, 7, 33]; *uniform subset graphs* [9, 11, 41] and *distance graphs* [5, 6, 36, 45]. The family of $G(n, r, s)$ graphs was initially (to the best of our knowledge) considered in [9], where they are called “*uniform subset graphs*”. However, this name does not become widespread. In our opinion, the term “*generalized Johnson graph*” is preferred as the most comprehensible, since, if we set $s = r - 1$, then the definition of $G(n, r, s)$ turns into the definition of the well-known Johnson graph. Note that the Kneser graph is also a special case of $G(n, r, s)$ with $s = 0$. However, the term “*generalized Kneser graph*” is already used for another class of graphs [12, 16, 22, 25].

¹Moscow Institute of Physics and Technology (National Research University), Dolgoprudny, Moscow Region, Russia. Supported by Grant N NSh-775.2022.1.1 to support leading scientific schools of Russia

²Moscow Institute of Physics and Technology (National Research University), Dolgoprudny, Moscow Region, Russia; The Russian Presidential Academy of National Economy and Public Administration, Moscow, Russia; Moscow Center for Fundamental and Applied Mathematics, Moscow, Russia; Adyghe State University, Caucasus mathematical center, Maykop, Republic of Adyghea, Russia. Supported by the Ministry of Science and Higher Education of the Russian Federation (Goszadaniye No. 075-00337-20-03), project No. 0714-2020-0005.

On the one hand, as we mentioned above, graphs $G(n, r, s)$ generalize Johnson graphs $G(n, r, r - 1)$ [2, 14, 17, 18, 19] and Kneser graphs $G(n, r, 0)$ [4, 10, 30, 31, 34, 35, 43], which are themselves of interest in the graph theory. On the other hand, they are a special case of distance graphs in \mathbb{R}^n with the Euclidean metric, which are used to study problems of combinatorial geometry (Hadwiger–Nelson problem about the chromatic number $\chi(\mathbb{R}^n)$ [8, 13, 20, 21, 23, 28, 29], Borsuk problem about partitioning of a set in \mathbb{R}^n into subsets of a smaller diameter [27], and various generalizations of these problems [37, 38, 39]).

Throughout the paper we assume that r and s are constant and n approaches infinity. The total number of vertices in this graph is denoted by N :

$$N = |V| = \binom{n}{r} \sim \frac{n^r}{r!}.$$

From the definition of $G(n, r, s)$ it is evident that this graph is *vertex-transitive*, i.e. for any two vertices there exists an automorphism of the graph mapping the first vertex to the second one. In particular, $G(n, r, s)$ is regular. Let N_1 denote the degree of its vertex:

$$N_1 = \binom{r}{s} \binom{n-r}{r-s} \sim \binom{r}{s} \frac{n^{r-s}}{(r-s)!}.$$

In [9] it is proved that the graph $G(n, r, s)$ is Hamiltonian for $s \in \{r - 1, r - 2, r - 3\}$, arbitrary r and n as well as for $s \in \{0, 1\}$, arbitrary r , and sufficiently large n . Hamiltonian cycles has been extensively studied in Kneser graphs $G(n, r, 0)$. It is known that they are hamiltonian for $n \geq 2.62r$ [10] and for all r when $n \leq 27$ (except for the Petersen graph $G(5, 2, 0)$) [40]. Graphs $G(2r + 1, r, 0)$ are also known to be Hamiltonian for all $r \geq 3$ [34]. As for cycles of a constant length, the asymptotics of the number of their appearances in $G(n, r, s)$ is known for all constant r and s and given below in Theorem 1.

Let H and G be graphs. A map $\varphi : V(H) \rightarrow V(G)$ is called a *homomorphism* from H to G if, for any pair of vertices x, y of H , $\{x, y\} \in E(H) \Rightarrow \{\varphi(x), \varphi(y)\} \in E(G)$. If a homomorphism is injective, then it is called a *monomorphism*. Let $\text{hom}(H, G)$ and $\text{mon}(H, G)$ denote respectively the number of homomorphisms and monomorphisms from H to G . Throughout this paper we write simply $\text{hom}(H)$ and $\text{mon}(H)$ when $G = G(n, r, s)$.

Let C_t be a cycle on t vertices. The purpose of this paper is to find the asymptotic value of $\text{mon}(C_t)$ for different $t = t(n)$.

Burkin [5] found the asymptotics of $\text{mon}(C_t)$ for all $t = \text{const}$.

Theorem 1 (Burkin, 2016, [5]) *Let t be a fixed integer. Then*

$$\text{mon}(C_t) \sim N N_1 \left(\frac{N_1}{\binom{r}{s}} \right)^{t-2}. \quad (1)$$

We generalize this result to cycles of variable length, i.e. $t = t(n)$. It turns out that for slow enough (sublogarithmic) growth of $t(n)$ the asymptotics of $\text{mon}(C_t)$ remains the same as in (1). In contrast, for superlogarithmic $t(n) = o\left(\min\{\sqrt{N}, N_1\}\right)$ the asymptotics is different, namely, $\text{mon}(C_t) \sim N_1^t$. These results can be summarized in the following two theorems.

Theorem 2 *As $n \rightarrow +\infty$, $\text{mon}(C_t) \sim \text{hom}(C_t)$ iff $t = o\left(\min\{\sqrt{N}, N_1\}\right)$.*

Theorem 2 is the trickiest result of our paper. Asymptotics of $\text{hom}(C_t)$ (stated below in Theorem 3) is a more or less direct corollary (modulo technical asymptotical computations) of the well-known representation of $\text{hom}(C_t)$ in terms of eigenvalues of $G(n, r, s)$. Let us fix an arbitrarily small $\varepsilon > 0$ and consider the partition of \mathbb{N} obtained by excluding ε -neighborhoods of $\frac{\ln n}{\ln \binom{r-j}{s-j} - \ln \binom{r-j-1}{s-j-1}}$, $j \in [0, s]$, i.e. the intervals

$$I_j = \left[\left\lfloor \frac{(1+\varepsilon) \ln n}{\ln \binom{r-j}{s-j} - \ln \binom{r-j-1}{s-j-1}} \right\rfloor \right] \setminus \left[\left\lfloor \frac{(1-\varepsilon) \ln n}{\ln \binom{r-j+1}{s-j+1} - \ln \binom{r-j}{s-j}} \right\rfloor \right], \quad j \in [s-1],$$

$$I_s = \left[\left\lfloor \frac{(1-\varepsilon) \ln n}{\ln(r-s+1)} \right\rfloor \right], \quad I_0 = \left[\left\lfloor \frac{(1+\varepsilon) \ln n}{\ln \frac{r}{s}} \right\rfloor, \infty \right).$$

Theorem 3 *For arbitrary $t = t(n) \in \mathbb{N}$,*

$$\text{hom}(C_t) = N_1^t \left(1 + O\left(\frac{1}{n}\right) + \sum_{j=1}^s \frac{n^j}{j!} \left(\frac{\binom{r-j}{s-j}}{\binom{r}{s}} + O\left(\frac{1}{n}\right) \right)^t \right). \quad (2)$$

Moreover, for $j \in [0, s+1]$ and $t \in I_j$,

$$\text{hom}(C_t) \sim N_1^t \frac{n^j}{j!} \left(\frac{\binom{r-j}{s-j}}{\binom{r}{s}} \right)^t.$$

Note that for $t \in I_0$, $\text{hom}(C_t) \sim N_1^t$, while, for $t \in I_s$, $\text{hom}(C_t) \sim N_1^t \frac{n^s}{s!} \binom{r}{s}^{-t}$ i.e. (1) holds. Theorem 2 and Theorem 3 immediately yield asymptotics of the number of copies of C_t in $G(n, r, s)$ for all $t = o\left(\min\{\sqrt{N}, N_1\}\right)$ since it equals $\frac{1}{2t} \text{mon}(C_t)$.

The rest of the paper is organized as follows. First, in Section 2, we discuss general properties of random walks on graphs (Section 2.1) and more specific properties of random walks on $G(n, r, s)$ (Sections 2.2 and 2.3). Secondly, in Section 3, we prove Theorem 3.

Finally, in Sections 4 and 5 we prove that the condition in Theorem 2 is, respectively, sufficient and necessary.

The proof of the sufficiency provided in Section 4 uses exact expressions for the spectrum of $G(n, r, s)$. It should be noted that the proof in the case $r > 2s$ (in which $\sqrt{N} = o(N_1)$) as well as in the case $t = \omega(\ln N)$ can be considerably simplified by using a more general argument (which we omit in this paper) applicable to a wide subclass of spectral expanders (see Section 5). However, for an arbitrary N_1 -regular graph G on N vertices, the property $\text{mon}(C_t, G) \sim \text{hom}(C_t, G)$ does not necessarily hold when $t = O(\ln N)$ and $N_1 = O(\sqrt{N})$, even if G is a spectral expander. This fact can be demonstrated, for example, by considering the random regular graph $G(N, N_1)$ with $N_1 = \lfloor \ln^8 N \rfloor$, in which, for any $\varepsilon > 0$, the inequality $\text{mon}(C_t, G(N, N_1)) / \text{hom}(C_t, G(N, N_1)) < \varepsilon$ holds with probability approaching 1 as soon as $t = o(\ln N / \ln \ln N)$. This can be shown by translating the same property from the binomial random graph $G(N, (1 + o(1))N_1/N)$ to $G(N, N_1)$ using the sandwich conjecture, which is true for $N_1 = \omega(\ln^7 N)$ [24]. Note that $G(N, N_1)$ is a spectral expander [44]. For the definition and properties of binomial random graphs and regular random graphs see [26].

The proof of the necessity in Theorem 2 provided in Section 5 does not rely upon the whole spectrum of $G(n, r, s)$ but rather uses its spectral expansion property. The necessity of the condition $t = o(N_1)$ follows from the fact that a random walk starts backtracking with positive probability if $t > cN_1$ for a constant c , which is proved in Section 5.2 using almost solely the regularity of $G(n, r, s)$. The necessity of $t = o(\sqrt{N})$ is proved in Section 5.1 using a high convergence rate of a random walk on an expander, which is discussed in Section 2.1. Therefore, in Section 5 we formulate a generalization of Theorem 2 to a class of spectral expanders.

2 Random walks

Counting cycles in $G(n, r, s)$ can be reduced to analysing the distribution of a random walk on $G(n, r, s)$.

2.1 Distribution and adjacency matrix

Let G be an arbitrary regular connected graph on the vertex set $[N]$ with every vertex having degree N_1 . Let $A = (A_{i,j}, i, j \in [N])$ be its adjacency matrix ($A_{i,j} = 1$ if and only if i and j are adjacent in G). Moreover, let $\lambda_j, j \in [0, r]$, be all distinct eigenvalues of A , and let m_j be the multiplicity of λ_j .

Recall that a *random walk on G* is a discrete-time random process $(X_n, n \in \mathbb{Z}_+)$, where

X_0 is a vertex chosen uniformly at random from $[N]$, and, for every $n \in \mathbb{Z}_+$, X_{n+1} is chosen uniformly at random from the neighbors of X_n in G . For $x, y \in [N]$, let

$$P^t(x, y) := \mathbb{P}(X_t = y | X_0 = x)$$

and $P^t = (P^t(x, y), x, y \in [N])$ be the k -step transition probabilities matrix.

For a positive integer t , $\text{hom}(C_t, G)$ is exactly the trace of A^t . Since the trace of A^t equals the sum of its eigenvalues and the eigenvalues of A^t can be computed as the t th power of the eigenvalues of A (see, e.g., [32]), we get

$$\text{hom}(C_t, G) = \sum_{j=0}^r m_j \lambda_j^t. \quad (3)$$

If G is vertex-transitive, then, clearly, all $P^t(z, z)$, $z \in [N]$, are equal to each other. Then, for every $x \in [N]$,

$$NP^t(x, x) = \sum_{z \in [N]} P^t(z, z) = \frac{\text{hom}(C_t, G)}{N_1^t}. \quad (4)$$

Therefore, (3) implies that

$$P^t(x, x) = \frac{1}{N} \sum_{j=0}^r m_j \left(\frac{\lambda_j}{N_1} \right)^t. \quad (5)$$

Notice that (due to regularity of G) the distribution $\pi = (1/N, \dots, 1/N) \in \mathbb{R}^N$ is *stationary* meaning that $\pi P^1 = \pi$. Let us here assume that λ_0 is the largest eigenvalue and λ_1 is the largest in absolute value eigenvalue distinct from λ_0 . From the regularity of G it follows that $\lambda_0 = N_1$ and from its connectedness, that $m_0 = 1$ [3]. Let us also assume that $|\lambda_1| < \lambda_0$ (which is equivalent, for a connected graph, to the graph being non-bipartite [3]). Fix $v \in [N]$ and $\varepsilon > 0$. Let us recall that *the variation distance* at time $t \in \mathbb{Z}_+$ with initial state v is

$$\Delta_v(t) = \frac{1}{2} \sum_{u \in [N]} \left| P^t(v, u) - \frac{1}{N} \right|.$$

It is very well known [42] that *the mixing time* $\tau_v(\varepsilon) := \min\{t : \Delta_v(t') \leq \varepsilon \text{ for all } t' \geq t\}$ satisfies

$$\tau_v(\varepsilon) \leq \left(1 - \frac{|\lambda_1|}{N_1} \right)^{-1} \ln \frac{N}{\varepsilon}. \quad (6)$$

2.2 Eigenvalues of $G(n, r, s)$

Let A be the adjacency matrix of $G(n, r, s)$. The eigenvalues of A are known [15]. They are equal to

$$\lambda_j = \sum_{\ell=\max\{0, j-s\}}^{\min\{j, r-s\}} (-1)^\ell \binom{j}{\ell} \binom{r-j}{r-s-\ell} \binom{n-r-j}{r-s-\ell}, \quad j \in [0, r], \quad (7)$$

and the multiplicity of the eigenvalue λ_j equals (we let $\binom{n}{-1} = 0$)

$$m_j = \binom{n}{j} - \binom{n}{j-1}.$$

In order to prove Theorem 3, we need to analyze asymptotical behaviour of the expression to the right in (5).

Notice that $\lambda_0 = N_1$ and $m_0 = 1$. Also, for $j \in [1, s]$,

$$\frac{\lambda_j}{N_1} = \frac{\binom{r-j}{s-j}}{\binom{r}{s}} + O\left(\frac{1}{n}\right), \quad m_j = \frac{n^j}{j!} + O\left(\frac{1}{n}\right) \quad (8)$$

and, for $j \in [s+1, r]$,

$$\left| \frac{\lambda_j}{N_1} \right| \sim \frac{\binom{j}{s}(r-s)!}{\binom{r}{s}(r-j)!} n^{-(j-s)}, \quad m_j = \frac{n^j}{j!} + O\left(\frac{1}{n}\right). \quad (9)$$

Therefore, for $t \geq 2$ and $j \in [s+1, r]$,

$$\frac{m_j \lambda_j^t}{m_s \lambda_s^t} = (O(1))^t \cdot n^{-(j-s)(t-1)} = (O(n^{-(j-s)}))^{t-1} = O\left(\frac{1}{n}\right) \quad (10)$$

implying that

$$P^t(x, x) = \frac{1 + O\left(\frac{1}{n}\right)}{N} \sum_{j=0}^s m_j \left(\frac{\lambda_j}{N_1} \right)^t = \frac{1}{N} \left(1 + O\left(\frac{1}{n}\right) + \sum_{j=1}^s m_j \left(\frac{\binom{r-j}{s-j}}{\binom{r}{s}} + O\left(\frac{1}{n}\right) \right)^t \right). \quad (11)$$

2.3 Random walk on $G(n, r, s)$

Here, we consider the random walk $(X_n, n \in \mathbb{Z}_+)$ on $G(n, r, s)$. Since $G(n, r, s)$ is vertex-transitive, for any $x \in V$,

$$\text{mon}(C_t) = NN_1^t \mathbb{P}(X_t = x, X_0 \neq X_1 \neq \dots \neq X_{t-1} | X_0 = x).$$

In order to prove Theorem 2, we bound the deviation of $\frac{\text{mon}(C_t)}{\text{hom}(C_t)}$ from 1. For convenience, in this section, we express the bound in terms of the diagonal elements of P^t .

Due to (4), we get

$$\begin{aligned} 0 \leq \frac{\text{hom}(C_t) - \text{mon}(C_t)}{\text{hom}(C_t)} &= \frac{P^t(x, x) - \mathbb{P}(X_t = x, X_0 \neq X_1 \neq \dots \neq X_{t-1} | X_0 = x)}{P^t(x, x)} = \\ &= \frac{\mathbb{P}(X_t = x, \exists i, j \in [0, t-1] : i \neq j, X_i = X_j | X_0 = x)}{\mathbb{P}(X_t = x | X_0 = x)}. \end{aligned}$$

Note that the expression to the right is exactly the probability that the random walk meets itself somewhere on $[0, t-1]$ subject to $X_0 = x$ and $X_t = x$.

By the union bound,

$$\begin{aligned} \frac{\text{hom}(C_t) - \text{mon}(C_t)}{\text{hom}(C_t)} &\leq \sum_{0 \leq i < j < t} \frac{\mathbb{P}(X_j = X_i, X_t = x | X_0 = x)}{\mathbb{P}(X_t = x | X_0 = x)} \\ &= \sum_{0 \leq i < j < t} \frac{\sum_{z \in V} \mathbb{P}(X_t = x | X_j = z) \mathbb{P}(X_j = z | X_i = z) \mathbb{P}(X_i = z | X_0 = x)}{\mathbb{P}(X_t = x | X_0 = x)}. \end{aligned}$$

Due to vertex-transitivity of $G(n, r, s)$ the probabilities $\mathbb{P}(X_j = z | X_i = z)$ are equal for all z . Therefore,

$$\begin{aligned} \frac{\text{hom}(C_t) - \text{mon}(C_t)}{\text{hom}(C_t)} &\leq \sum_{0 \leq i < j < t} \frac{\sum_{z \in V} \mathbb{P}(X_{t-j+i} = x | X_i = z) P^{j-i}(x, x) \mathbb{P}(X_i = z | X_0 = x)}{\mathbb{P}(X_t = x | X_0 = x)} \\ &= \sum_{0 \leq i < j < t} \frac{P^{t-j+i}(x, x) P^{j-i}(x, x)}{P^t(x, x)} = \sum_{k=1}^{t-1} (t-k) \frac{P^k(x, x) P^{t-k}(x, x)}{P^t(x, x)} \quad (12) \\ &\leq t \sum_{k=1}^{t-1} \frac{P^k(x, x) P^{t-k}(x, x)}{P^t(x, x)} = t \sum_{k=2}^{t-2} \frac{P^k(x, x) P^{t-k}(x, x)}{P^t(x, x)}. \end{aligned}$$

3 Proof of Theorem 3

From (4) and (11), we get

$$\text{hom}(C_t) = N_1^t \left(1 + O\left(\frac{1}{n}\right) + \sum_{j=1}^s \frac{n^j}{j!} \left(\frac{\binom{r-j}{s-j}}{\binom{r}{s}} + O\left(\frac{1}{n}\right) \right)^t \right). \quad (13)$$

Let $j \in [s-1]$, $\varepsilon > 0$. If $t < (1 - \varepsilon) \frac{\ln n}{\ln \binom{r-j}{s-j} - \ln \binom{r-j-1}{s-j-1}}$, then

$$\left(\frac{\binom{r-j}{s-j}}{\binom{r}{s}} + o(1) \right)^t = o \left[n \left(\frac{\binom{r-j-1}{s-j-1}}{\binom{r}{s}} + o(1) \right)^t \right]$$

implying that the $(j+1)$ th term in the sum in (13) is asymptotically bigger than the j th term. Similarly, if $t > (1 + \varepsilon) \frac{\ln n}{\ln \binom{r-j}{s-j} - \ln \binom{r-j-1}{s-j-1}}$, then the $(j+1)$ th term in the sum in (13) is asymptotically smaller than the j th term. From this, we immediately get

$$\text{hom}(C_t) \sim \begin{cases} N_1^t \frac{n^s}{s!} \binom{r}{s}^{-t}, & t < (1 - \varepsilon) \frac{\ln n}{\ln \binom{r-s+1}{s-s+1} - \ln \binom{r-s}{s-s}}; \\ N_1^t \frac{n^j}{j!} \binom{r-j}{s-j}^t \binom{r}{s}^{-t}, & \frac{(1+\varepsilon) \ln n}{\ln \binom{r-j}{s-j} - \ln \binom{r-j-1}{s-j-1}} < t < \frac{(1-\varepsilon) \ln n}{\ln \binom{r-j+1}{s-j+1} - \ln \binom{r-j}{s-j}}, j \in [s-1]; \\ N_1^t, & t > (1 + \varepsilon) \frac{\ln n}{\ln \binom{r}{s} - \ln \binom{r-1}{s-1}}. \end{cases}$$

which proves Theorem 3. \square

4 Proof of sufficiency in Theorem 2

Here we prove that if $t = o(\min\{\sqrt{N}, N_1\})$, then

$$\frac{\text{hom}(C_t) - \text{mon}(C_t)}{\text{hom}(C_t)} \rightarrow 0, \quad n \rightarrow \infty.$$

Since this fraction is non-negative, it is sufficient to prove that the upper bound from (12) approaches 0. Clearly, we may assume that $t \geq 4$.

By (5), for every $x \in V$,

$$\begin{aligned} \sum_{k=2}^{t-2} P^k(x, x) P^{t-k}(x, x) &= \frac{1}{N^2} \sum_{k=2}^{t-2} \sum_{i,j=0}^r m_i m_j \left(\frac{\lambda_j}{N_1} \right)^k \left(\frac{\lambda_i}{N_1} \right)^{t-k} \\ &\leq \frac{t}{N^2} \sum_{i=0}^r m_i^2 \left| \frac{\lambda_i}{N_1} \right|^t + \frac{1}{N^2} \sum_{i \neq j} m_i m_j \left| \frac{\lambda_i}{N_1} \right|^t \sum_{k=2}^{t-2} \left| \frac{\lambda_j}{\lambda_i} \right|^k. \end{aligned} \tag{14}$$

Note that $|\lambda_i|^t \sum_{k=2}^{t-2} \left| \frac{\lambda_j}{\lambda_i} \right|^k = |\lambda_j|^t \sum_{k=2}^{t-2} \left| \frac{\lambda_i}{\lambda_j} \right|^k$. Therefore, for n large enough, due to (8) and (9), we get

$$\sum_{i \neq j} m_i m_j \left| \frac{\lambda_i}{N_1} \right|^t \sum_{k=2}^{t-2} \left| \frac{\lambda_j}{\lambda_i} \right|^k = 2 \sum_{0 \leq i < j \leq r} m_i m_j \left| \frac{\lambda_i}{N_1} \right|^t \sum_{k=2}^{t-2} \left| \frac{\lambda_j}{\lambda_i} \right|^k \leq 2 \sum_{0 \leq i < j \leq r} m_i m_j \left| \frac{\lambda_i}{N_1} \right|^t \frac{\left| \frac{\lambda_j}{\lambda_i} \right|^2}{1 - \left| \frac{\lambda_j}{\lambda_i} \right|}.$$

Moreover, from (8) and (9) we get that, for $j > i$ and $j > s$, $\frac{\lambda_j}{\lambda_i} = O\left(\frac{1}{n}\right)$, while, for $i < j \leq s$,

$$\frac{\lambda_j}{\lambda_i} = \frac{(s-i) \dots (s-j+1)}{(r-i) \dots (r-j+1)} \left(1 + O\left(\frac{1}{n}\right) \right).$$

The latter expression is less than $\frac{s}{r}$ if $j \neq 1$ and n is large enough. If $i = 0$, $j = 1$, then, from (7), we get

$$\frac{\lambda_1}{\lambda_0} = \frac{\binom{r-1}{r-s} \binom{n-r-1}{r-s} - \binom{r-1}{r-s-1} \binom{n-r-1}{r-s-1}}{\binom{r}{r-s} \binom{n-r}{r-s}} < \frac{\binom{r-1}{r-s} \binom{n-r-1}{r-s}}{\binom{r}{r-s} \binom{n-r}{r-s}} = \frac{s(n-2r+s)}{r(n-r)} < \frac{s}{r}.$$

Then, for n large enough,

$$\begin{aligned} \sum_{i \neq j} m_i m_j \left| \frac{\lambda_i}{N_1} \right|^t \sum_{k=2}^{t-2} \left| \frac{\lambda_j}{\lambda_i} \right|^k &\leq \frac{2}{1-s/r} \sum_{0 \leq i < j \leq r} m_i m_j \left| \frac{\lambda_i}{N_1} \right|^{t-2} \left| \frac{\lambda_j}{N_1} \right|^2 \\ &\leq \frac{2}{1-s/r} \sum_{i,j=0}^r m_i m_j \left| \frac{\lambda_i}{N_1} \right|^{t-2} \left| \frac{\lambda_j}{N_1} \right|^2 \\ &= \frac{2}{1-s/r} \left(\sum_{i=0}^r m_i \left| \frac{\lambda_i}{N_1} \right|^{t-2} \right) \left(\sum_{j=0}^r m_j \left| \frac{\lambda_j}{N_1} \right|^2 \right). \end{aligned}$$

By (11) and the definition of P^2 ,

$$\sum_{j=0}^r m_j \left| \frac{\lambda_j}{N_1} \right|^2 = NP^2(x, x) = \frac{N}{N_1}.$$

Moreover, by (8), (10) and (11),

$$\begin{aligned} \sum_{i=0}^r m_i \left| \frac{\lambda_i}{N_1} \right|^{t-2} &\sim \sum_{i=0}^s m_i \left| \frac{\lambda_i}{N_1} \right|^{t-2} \sim \sum_{i=0}^s m_i \left(\frac{\lambda_i}{N_1} \right)^{t-2} = O\left(\sum_{i=0}^s m_i \left(\frac{\lambda_i}{N_1} \right)^t \right) \\ &= O\left(\sum_{i=0}^r m_i \left(\frac{\lambda_i}{N_1} \right)^t \right) = O(NP^t(x, x)). \end{aligned}$$

It remains to estimate the first summand in the rightmost expression in (14). From (8) and (9), we get that, for $j \in [s+1, r]$,

$$\frac{m_j^2 \lambda_j^t}{m_s^2 \lambda_s^t} = (O(1))^t \cdot n^{-(j-s)(t-2)} = (O(n^{-(j-s)}))^{t-2} = o(1)$$

implying that

$$\sum_{i=0}^r m_i^2 \left| \frac{\lambda_i}{N_1} \right|^t \sim \sum_{i=0}^s m_i^2 \left(\frac{\lambda_i}{N_1} \right)^t.$$

Putting everything together and applying (11), we conclude that, for every $x \in V$,

$$t \sum_{k=2}^{t-2} \frac{P^k(x, x) P^{t-k}(x, x)}{P^t(x, x)} \leq \frac{t^2 \sum_{i=0}^s m_i^2 \left(\frac{\lambda_i}{N_1} \right)^t}{N \sum_{i=0}^s m_i \left(\frac{\lambda_i}{N_1} \right)^t} (1 + o(1)) + O\left(\frac{t}{N_1}\right).$$

We finish with proving that the condition $t = o(\sqrt{N})$ implies

$$\frac{t^2 \sum_{i=0}^s m_i^2 \left(\frac{\lambda_i}{N_1} \right)^t}{N \sum_{i=0}^s m_i \left(\frac{\lambda_i}{N_1} \right)^t} = o(1).$$

If $t > C \ln n$, where C is sufficiently large constant, then, by (8), for every $i \in [s]$, $m_i \left(\frac{\lambda_i}{N_1} \right)^t = o(1)$ and $m_i^2 \left(\frac{\lambda_i}{N_1} \right)^t = o(1)$. Therefore,

$$\frac{t^2 \sum_{i=0}^s m_i^2 \left(\frac{\lambda_i}{N_1} \right)^t}{N \sum_{i=0}^s m_i \left(\frac{\lambda_i}{N_1} \right)^t} \sim \frac{t^2}{N} = o(1).$$

If $t \leq C \ln n$, then, by (8),

$$\frac{t^2 \sum_{i=0}^s m_i^2 \left(\frac{\lambda_i}{N_1} \right)^t}{N \sum_{i=0}^s m_i \left(\frac{\lambda_i}{N_1} \right)^t} \leq \frac{t^2 m_s}{N} (1 + o(1)) = O\left(\frac{n^s \ln^2 n}{N}\right) = O\left(\frac{\ln^2 n}{n^{r-s}}\right) = o(1).$$

The sufficiency in Theorem 2 is proved. \square

5 Proof of necessity in Theorem 2

As was already noted, the necessity of the condition in Theorem 2 follows from a more general fact about spectral expanders. Consider a sequence of graphs $\{G_N, N \in \mathbb{N}\}$ such that $[N]$ is the set of vertices of G_N , and G_N is non-bipartite, connected and N_1 -regular (N_1 depends on N). Let $\lambda_1 = \lambda_1(N)$ be the second largest in absolute value eigenvalue of the adjacency matrix of G_N . We call the sequence $\{G_N, N \in \mathbb{N}\}$ a *spectral expander*, if there exists $\delta > 0$ such that, for all large enough N , $|\lambda_1|/N_1 < 1 - \delta$.

Theorem 4 *Let $\{G_N, N \in \mathbb{N}\}$ be a spectral expander such that $N_1 = \omega(\ln N)$. Then for any $c > 0$ there exists $\varepsilon > 0$ such that, for sufficiently large N , if $t > c \min\{\sqrt{N}, N_1\}$, then $\text{mon}(C_t, G_N) < (1 - \varepsilon)\text{hom}(C_t, G_N)$.*

Obviously, Theorem 4 implies the necessity of the condition $t = o(\min\{\sqrt{N}, N_1\})$ for $\text{mon}(C_t) \sim \text{hom}(C_t)$ stated in Theorem 2.

To prove Theorem 4, we introduce a random walk on G_N (see Section 2.1).

In Section 2.3, we note that the proportion of self-intersecting cycles in $G(n, r, s)$ is exactly the probability that the random walk meets itself somewhere on $[0, t - 1]$ subject to $X_0 = x$ and $X_t = x$. It is easy to see that (almost) the same is true for G_N since X_0 is chosen uniformly at random from $[N]$. Indeed,

$$\begin{aligned} \text{hom}(C_t, G_N) &= \sum_{x \in [N]} P^t(x, x) N_1^t = N N_1^t \sum_{x \in [N]} P^t(x, x) \mathbb{P}(X_0 = x) \\ &= N N_1^t \sum_{x \in [N]} \mathbb{P}(X_t = X_0 = x) = N N_1^t \mathbb{P}(X_t = X_0). \end{aligned}$$

In the same way, $\text{mon}(C_t, G) = N N_1^t \mathbb{P}(X_t = X_0, X_0 \neq X_1 \neq \dots \neq X_{t-1})$. Therefore,

$$\frac{\text{hom}(C_t, G) - \text{mon}(C_t, G)}{\text{hom}(C_t, G)} = \mathbb{P}(\exists i, j \in [0, t - 1] : i \neq j, X_i = X_j | X_t = X_0). \quad (15)$$

Further, we separately consider cases $\sqrt{N} = o(N_1)$ and $N_1 = O(\sqrt{N})$.

5.1 $\sqrt{N} = o(N_1)$

Let us first bound from above $P^t(x, y)$ for arbitrary x, y from $[N]$ and an integer $t \geq 1$. Since $P^t(x, y) = \sum_{v \in [N]} P^{t-1}(x, v) P^1(v, y)$, we get that

$$P^t(x, y) \leq \max_{v \in [N]} P^1(v, y) \leq \frac{1}{N_1}. \quad (16)$$

Let us also notice that from (6) it immediately follows that, for every $C > 0$, there exists κ such that, for all $t \geq \kappa \ln N$ and all $x, y \in [N]$, we have

$$\left| P^t(x, y) - \frac{1}{N} \right| < \frac{1}{N^2}. \quad (17)$$

Let us fix a positive $\tilde{c} < \min\{c, 1\}$ and prove that the random walk subject to $X_t = X_0$ intersects itself during the first $\tilde{c}\sqrt{N}$ steps with non-zero probability. Let

$$\mathcal{J} := \left\{ (i, j) \in [t]^2 : \kappa \ln N < i < j - 3\kappa \ln N < j < \tilde{c}\sqrt{N} \right\}.$$

Note that $|\mathcal{J}| = \frac{\tilde{c}^2 + o(1)}{2} N$ and that $j < t - \kappa \ln n$ for every $(i, j) \in \mathcal{J}$.

For all $(i, j) \in \mathcal{J}$, we have that

$$\frac{\mathbb{P}(X_i = X_j, X_t = X_0)}{\mathbb{P}(X_t = X_0)} = \sum_{x, u \in [N]} \frac{P^i(x, u) P^{j-i}(u, u) P^{t-j}(u, x)}{\sum_{y \in [N]} P^t(y, y)} = \frac{1 + o(1)}{N}$$

due to (17). Note that $o(1)$ in the expression above converges to 0 uniformly over all $(i, j) \in \mathcal{J}$. Therefore,

$$\sum_{(i, j) \in \mathcal{J}} \frac{\mathbb{P}(X_i = X_j, X_t = X_0)}{\mathbb{P}(X_t = X_0)} = \frac{\tilde{c}^2}{2} + o(1). \quad (18)$$

Let

$$\begin{aligned} \mathcal{J}_0 &:= \left\{ \{(i_1, j_1), (i_2, j_2)\} \in \binom{\mathcal{J}}{2} : (\forall \{i, j\} \subset \{i_1, i_2, j_1, j_2\} : |i - j| \geq \kappa \ln N) \right\}, \\ \mathcal{J}_1 &:= \left\{ \{(i_1, j_1), (i_2, j_2)\} \in \binom{\mathcal{J}}{2} : (\exists! \{i, j\} \subset \{i_1, i_2, j_1, j_2\} : |i - j| < \kappa \ln N) \right\}, \\ \mathcal{J}_2 &:= \left\{ \{(i_1, j_1), (i_2, j_2)\} \in \binom{\mathcal{J}}{2} : \max\{|i_1 - i_2|, |j_1 - j_2|\} < \kappa \ln N \right\}. \end{aligned}$$

It is clear from the definition of \mathcal{J} that $\mathcal{J}_0 \sqcup \mathcal{J}_1 \sqcup \mathcal{J}_2 = \binom{\mathcal{J}}{2}$ (recall that $j - i > 3\kappa \ln N$ for every $(i, j) \in \mathcal{J}$). We have

$$|\mathcal{J}_0| = \frac{1}{2} |\mathcal{J}^2| (1 + o(1)) = \frac{\tilde{c}^4 + o(1)}{8} N^2, \quad |\mathcal{J}_1| < 4\kappa \ln N (\tilde{c}\sqrt{N})^3, \quad |\mathcal{J}_2| < (2\kappa \ln N \tilde{c}\sqrt{N})^2.$$

As above, uniformly over all $\{(i_1, j_1), (i_2, j_2)\} \in \mathcal{J}_0$, (17) implies

$$\frac{\mathbb{P}(X_{i_1} = X_{j_1}, X_{i_2} = X_{j_2}, X_t = X_0)}{\mathbb{P}(X_t = X_0)} = \frac{1 + o(1)}{N^2}.$$

Uniformly over all $\{(i_1, j_1), (i_2, j_2)\} \in \mathcal{J}_1$, the relations (16) and (17) imply

$$\frac{\mathbb{P}(X_{i_1} = X_{j_1}, X_{i_2} = X_{j_2}, X_t = X_0)}{\mathbb{P}(X_t = X_0)} \leq \frac{1 + o(1)}{NN_1}.$$

Uniformly over all $\{(i_1, j_1), (i_2, j_2)\} \in \mathcal{J}_2$, (16) implies

$$\frac{\mathbb{P}(X_{i_1} = X_{j_1}, X_{i_2} = X_{j_2}, X_t = X_0)}{\mathbb{P}(X_t = X_0)} \leq \frac{1 + o(1)}{N_1^2}.$$

Summing up and recalling that, in the current case, $\sqrt{N} = o(N_1)$,

$$\begin{aligned} & \sum_{\{(i_1, j_1), (i_2, j_2)\} \in \binom{\mathcal{J}}{2}} \frac{\mathbb{P}(X_{i_1} = X_{j_1}, X_{i_2} = X_{j_2}, X_t = X_0)}{\mathbb{P}(X_t = X_0)} \\ & \leq \frac{\tilde{c}^4 + o(1)}{8} + \frac{(4\kappa\tilde{c}^3 + o(1))\sqrt{N} \ln N}{N_1} + \frac{(2\kappa\tilde{c} + o(1))^2 N \ln N}{N_1^2} = \frac{\tilde{c}^4 + o(1)}{8}. \end{aligned} \quad (19)$$

From (15), (18) and (19) we get

$$\begin{aligned} \frac{\text{hom}(C_t, G) - \text{mon}(C_t, G)}{\text{hom}(C_t, G)} & \geq \sum_{(i, j) \in \mathcal{J}} \frac{\mathbb{P}(X_i = X_j, X_t = X_0)}{\mathbb{P}(X_t = X_0)} \\ & \quad - \sum_{\{(i_1, j_1), (i_2, j_2)\} \in \binom{\mathcal{J}}{2}} \frac{\mathbb{P}(X_{i_1} = X_{j_1}, X_{i_2} = X_{j_2}, X_t = X_0)}{\mathbb{P}(X_t = X_0)} \\ & = \frac{\tilde{c}^2}{2} - \frac{\tilde{c}^4}{8} + o(1). \end{aligned}$$

Since $\frac{\tilde{c}^2}{2} - \frac{\tilde{c}^4}{8} > 0$, we conclude that $\text{mon}(C_t, G)/\text{hom}(C_t, G)$ is bounded away from 1 as needed.

5.2 $N_1 = O(\sqrt{N})$

W.l.o.g. we may assume that $c < 1$ and prove that the random walk subject to $X_t = X_0$ intersects itself during the first cN_1 steps with non-zero probability. In the same way as in Section 5.1, we use (15). However, here we consider all (i, j) such that i is even and $j = i + 2 \leq cN_1$. Let $\mathcal{J} := \{2i : i \in [0, \lfloor \frac{cN_1-2}{2} \rfloor]\}$. For every $i \in \mathcal{J}$, we have

$$\frac{\mathbb{P}(X_i = X_{i+2}, X_t = X_0)}{\mathbb{P}(X_t = X_0)} = \frac{1}{N} \sum_{x, u \in [N]} \frac{P^i(x, u) P^2(u, u) P^{t-i-2}(u, x)}{\mathbb{P}(X_t = X_0)} = \frac{1}{N_1} \frac{\mathbb{P}(X_{t-2} = X_0)}{\mathbb{P}(X_t = X_0)} \sim \frac{1}{N_1}$$

since $P^2(u, u) = \frac{1}{N_1}$ for all $u \in [N]$ and $\mathbb{P}(X_{t-2} = X_0) \sim \mathbb{P}(X_t = X_0) \sim 1/N$ due to (17). Thus,

$$\sum_{i \in \mathcal{J}} \frac{\mathbb{P}(X_i = X_{i+2}, X_t = X_0)}{\mathbb{P}(X_t = X_0)} = \frac{c}{2} + o(1).$$

Now, let $i_1, i_2 \in \mathcal{J}$, $i_1 < i_2$. Then, similarly,

$$\begin{aligned} & \frac{\mathbb{P}(X_{i_1} = X_{i_1+2}, X_{i_2} = X_{i_2+2}, X_t = X_0)}{\mathbb{P}(X_t = X_0)} = \\ &= \frac{1}{N} \sum_{x, u, v \in [N]} \frac{P^{i_1}(x, u) P^2(u, u) P^{i_2-i_1-2}(u, v) P^2(v, v) P^{t-i_2-2}(v, x)}{\mathbb{P}(X_t = X_0)} \\ &= \frac{1}{N} \sum_{x, u, v \in [N]} \frac{P^{i_1}(x, u) P^{i_2-i_1-2}(u, v) P^{t-i_2-2}(v, x)}{N_1^2 \mathbb{P}(X_t = X_0)} \\ &= \frac{\mathbb{P}(X_{t-4} = X_0)}{N_1^2 \mathbb{P}(X_t = X_0)} \sim \frac{1}{N_1^2}. \end{aligned}$$

Therefore,

$$\begin{aligned} \frac{\text{hom}(C_t, G) - \text{mon}(C_t, G)}{\text{hom}(C_t, G)} &\geq \sum_{i \in \mathcal{J}} \frac{\mathbb{P}(X_i = X_{i+2}, X_t = X_0)}{\mathbb{P}(X_t = X_0)} \\ &\quad - \sum_{i_1, i_2 \in \mathcal{J}: i_1 < i_2} \frac{\mathbb{P}(X_{i_1} = X_{i_1+2}, X_{i_2} = X_{i_2+2}, X_t = X_0)}{\mathbb{P}(X_t = X_0)} \\ &= \frac{c}{2} - \frac{c^2}{8} + o(1). \end{aligned}$$

Since $c/2 - c^2/8 > 0$, this finishes the proof of Theorem 4 and therefore of Theorem 2. \square

6 Acknowledgements

We would like to thank Andrey Kupavskii for very useful discussions.

References

- [1] L. A. Agong, C. Amarra, J. S. Caughman, A. J. Herman, T. S. Terada, *On the girth and diameter of generalized Johnson graphs*, Discrete Mathematics, **341**:1 (2018) pp. 138–142.
- [2] B. Alspach, *Johnson graphs are Hamilton-connected*, Ars Mathematica Contemporanea, **6**:1 (2012) pp. 21–23.
- [3] R. B. Bapat, *Graphs and Matrices*, Springer London (2014).
- [4] A. E. Brouwer, A. Schrijver, *Uniform Hypergraphs*, In Packing and Covering in Combinatorics. Mathematical Centre Tracts, **106** (1979) pp. 39–73.
- [5] A. V. Burkin, *Small Subgraphs in Random Distance Graphs*, Theory of Probability & Its Applications, **60**:3 (2016) pp. 367–382.
- [6] A. V. Burkin, M. E. Zhukovskii, *Small subgraphs and their extensions in a random distance graph*, Sbornik: Mathematics, **209**:2 (2018) pp. 163–186.
- [7] A. D. Cannon, J. Bamberg, C. E. Praeger, *A Classification of the Strongly Regular Generalised Johnson Graphs*, Annals of Combinatorics, **16**:3 (2012), pp.489–506.
- [8] K. Cantwell, *Finite Euclidean Ramsey theory*, Journal of Combinatorial Theory, Series A, **73**:2 (1996) pp. 273–285.
- [9] B.-L. Chen, K.-W. Lih, *Hamiltonian uniform subset graphs*, Journal of Combinatorial Theory, Series B, **42**:3 (1987) 257–263.
- [10] Y.-C. Chen, *Triangle-free Hamiltonian Kneser graphs*, Journal of Combinatorial Theory, Series B, **89**:1 (2003) pp. 1–16.
- [11] Y. Chen, W. Wang, *Diameters of uniform subset graphs*, Discrete Mathematics, **308**:24 (2008) 6645–6649.
- [12] Y. Chen, Y. Wang, *On the diameter of generalized Kneser graphs*, Discrete Mathematics, **308**:18 (2008) 4276–4279.
- [13] K. B. Chilakamarri, *On the chromatic number of rational five-space*, Aequationes Mathematicae, **39**:2-3 (1990) pp. 146–148.
- [14] M. Daven, C. A. Rodger, *The Johnson graph $J(v, k)$ has connectivity δ* , Congressus Numerantium, **139** (1999) pp. 123–128.

- [15] P. Delsarte, *An algebraic approach to the association schemes of coding theory*, Philips Res. Rep. Suppl. No. 10 (1973).
- [16] T. Denley, *The Odd Girth of the Generalised Kneser Graph*, European Journal of Combinatorics, **18**:6 (1997) pp. 607–611.
- [17] T. Etzion, *On the Nonexistence of Perfect Codes in the Johnson Scheme*, SIAM Journal on Discrete Mathematics, **9**:2 (1996) pp. 201–209.
- [18] T. Etzion, S. Bitan, *On the chromatic number, colorings, and codes of the Johnson graph*, Discrete Applied Mathematics, **70**:2 (1996) pp. 163–175.
- [19] T. Etzion, A. Brouwer, *Some new distance-4 constant weight codes*, Advances in Mathematics of Communications, **5**:3 (2011) 417–424.
- [20] G. Exoo, D. Ismailescu, M. Lim, *On the Chromatic Number of \mathbb{R}^4* , Discrete & Computational Geometry, **52**:2 (2014) pp. 416–423.
- [21] P. Frankl, *Extremal Problems and Coverings of the Space*, European Journal of Combinatorics, **1**:2 (1980) pp. 101–106.
- [22] P. Frankl, *On the chromatic number of the general Kneser-graph*, Journal of Graph Theory, **9**:2 (1985) pp. 217–220.
- [23] P. Frankl, R. M. Wilson, *Intersection theorems with geometric consequences*, Combinatorica, **1**:4 (1981) pp. 357–368.
- [24] P. Gao, *Kim-Vu’s sandwich conjecture is true for all $d = \omega(\log^7 n)$* , arXiv:2011.09449 (2021).
- [25] A. Jafari, M. J. Moghaddamzadeh, *On the chromatic number of generalized Kneser graphs and Hadamard matrices*, Discrete Mathematics, **343**:2 (2020) 111682.
- [26] S. Janson, T. Luczak, A. Ruciński, *Random Graphs*, Wiley, New York (2000).
- [27] J. Kahn, G. Kalai, *A Counterexample to Borsuk’s Conjecture*, Bulletin of the American Mathematical Society, **29**:1 (1993) pp. 60–63.
- [28] A. B. Kupavskii and A. M. Raigorodskii, *On the chromatic number of \mathbb{R}^9* , Journal of Mathematical Sciences, **163**:6 (2009) 720–731.
- [29] D. G. Larman, C. A. Rogers, *The realization of distances within sets in Euclidean space*, Mathematika, **19**:1 (1972) pp. 1–24.

- [30] L. Lovász, *Kneser's conjecture, chromatic number, and homotopy*, Journal of Combinatorial Theory, Series A, **25**:3 (1978) pp. 319–324.
- [31] J. Matoušek, *A Combinatorial Proof of Kneser's Conjecture*, Combinatorica, **24**:1 (2004) pp. 163–170.
- [32] C. D. Meyer, *Matrix analysis and applied linear algebra*, Society for Industrial and Applied Mathematics, Philadelphia (2000).
- [33] J. J. Moliterno, *Entries of the group inverse of the Laplacian matrix for generalized Johnson graphs*, Linear and Multilinear Algebra, **66**:6 (2017) pp. 1153–1170.
- [34] T. Mütze, J. Nummenpalo, B. Walczak, *Sparse Kneser graphs are Hamiltonian*, Journal of the London Mathematical Society **103**:4 (2021) 1253–1275.
- [35] S. Poljak, Z. Tuza, *Maximum bipartite subgraphs of Kneser graphs*, Graphs and Combinatorics, **3**:1 (1987) pp. 191–199.
- [36] M. M. Pyaderkin, *Independence numbers of random subgraphs of distance graphs*, Mathematical Notes, **99**:3-4 (2016) pp. 556–563.
- [37] A. M. Raigorodskii, *Borsuk's problem and the chromatic numbers of some metric spaces*, Russian Mathematical Surveys, **56**:1 (2001) pp. 103–139.
- [38] A. M. Raigorodskii, *Combinatorial Geometry and Coding Theory*, Fundamenta Informaticae, **145**:3 (2016) pp. 359–369.
- [39] A. M. Raigorodskii, *Coloring Distance Graphs and Graphs of Diameters*, Thirty Essays on Geometric Graph Theory ed. János Pach (2013) pp. 429–460.
- [40] I. Shields, C. Savage, *A Note on Hamilton Cycles in Kneser Graphs*, Bull. Inst. Combin. Appl. **40** (2002).
- [41] J. E. Simpson, *On uniform subset graphs*, Ars Combinatoria, **37** (1994) pp. 309–318.
- [42] A. Sinclair, *Improved bounds for mixing rates of Markov chains and multicommodity flow*, in: LATIN 1992. Lecture Notes in Computer Science, **583** (1992).
- [43] M. Valencia-Pabon, J.-C. Vera, *On the diameter of Kneser graphs*, Discrete Mathematics, **305**:1-3 (2005) pp. 383–385.
- [44] Y. Zhao, *Spectral Distributions of Random Graphs* (2012).

- [45] M. E. Zhukovskii, *On the probability of the occurrence of a copy of a fixed graph in a random distance graph*, Mathematical Notes, **92**:6 (2012) pp. 756–766.