

Numerable open covers and representability of topological stacks

Dmitri Pavlov

Department of Mathematics and Statistics, Texas Tech University
<https://dmitripavlov.org/>

Abstract. In this note, we establish an analogue of the Brown–Gersten property for numerable open covers of topological spaces: a simplicial presheaf on the site of topological spaces with numerable open covers satisfies the homotopy descent property for all open covers if and only if it satisfies it for covers with two elements and covers with pairwise disjoint elements. We apply this result to deduce a representability criterion for stacks on topological spaces similar to arXiv:1912.10544. We also use this result to establish new simple criteria for chain complexes of sheaves of abelian groups to satisfy the homotopy descent property.

Contents

1. Introduction
 - 1.4. Acknowledgments
2. Recollections on Grothendieck topologies
3. Numerable covers
4. Main theorem for topological spaces
5. Numerable sites
6. Homotopy descent for chain complexes of sheaves of abelian groups
7. Example: classical cohomology theories on smooth manifolds
8. Application: Classifying spaces for stacks over topological spaces
9. References

1 Introduction

The Brown–Gersten property allows one to verify the homotopy descent property for simplicial presheaves on Noetherian topological spaces of finite Krull dimension (the Zariski topology being the prime example) by verifying it for the empty cover and for covers consisting of two elements, see Brown–Gersten [1973, Theorem 4]. Morel–Voevodsky [1999, Proposition 1.16] extended this result to the Nisnevich topology, where the role of pairs of open subsets is assumed by Nisnevich squares. Voevodsky [2008.c] developed an abstract theory of completely decomposable topologies, which encompasses both cases.

These results allow one to drastically simplify the general homotopy descent condition, which involves homotopy limits of cosimplicial objects (also known as homotopy totalizations), by a much easier pair of conditions: $F(\emptyset)$ is weakly equivalent to the terminal object and the square

$$\begin{array}{ccc} F(U \cup V) & \longrightarrow & F(U) \\ \downarrow & & \downarrow \\ F(V) & \longrightarrow & F(U \cap V) \end{array}$$

is homotopy cartesian for any open U and V (more generally, for any distinguished square in the completely decomposable topology under consideration, see Voevodsky [2008.c]). This condition can be seen as the homotopy coherent version of the Mayer–Vietoris property. Apart from the fact that homotopy pullbacks are easier to manipulate, they also have better formal properties than homotopy totalizations. For example, filtered homotopy colimits of simplicial presheaves commute with homotopy pullbacks, but not with homotopy totalizations.

Recently, homotopy descent for the site of smooth manifolds (and related sites, such as the small site of a fixed smooth manifold or the site of manifolds with open embeddings as morphisms) has been gaining in importance in such fields as differential geometry and differential topology, with many examples, such as (higher) bundle gerbes, twists for vector bundles, Lie ∞ -groupoids, and Lie ∞ -algebroids, being formulated in this language, see, for example, the recent work of Sati–Schreiber–Stasheff [2008.b] and Fiorenza–Schreiber–Stasheff [2010].

One is naturally led to the question whether the Brown–Gersten property holds for the site of smooth manifolds. However, the most obvious analogue fails in a very simple situation. Fix a set A and consider the presheaf F of sets on the site of smooth manifolds that sends a smooth manifold M to the set of locally constant functions $M \rightarrow A$ with finitely many distinct values. This presheaf satisfies descent for all finite covers. However, it does not satisfy descent for the cover of a manifold with infinitely many connected components by its connected components. Indeed, the relevant descent object consists of all locally constant functions $M \rightarrow A$, which can have infinitely many distinct values.

The goal of this note is to show that the above situation is the worst thing that can happen: if we have homotopy descent for covers with two elements and covers of a disjoint union $\coprod_i U_i$ by its components U_i , then we have descent for all open covers. This is the content of the main theorem, which is proved in greater generality as Theorem 4.8:

Theorem 1.1. A simplicial presheaf on the site of smooth manifolds satisfies the homotopy descent property for all open covers if and only if it satisfies descent for covers with two elements and covers with pairwise disjoint elements.

Thus the general homotopy descent property is reduced to open covers with two elements (i.e., homotopy pullbacks) and to open covers with disjoint elements (i.e., infinite homotopy products), which are much easier to treat in practice than arbitrary homotopy totalizations.

More generally, Theorem 4.8 applies to arbitrary topological spaces as long as we work with numerable covers (Definition 3.9), defined as covers that admit a subordinate partition of unity. They are also known as normal covers. An even more general version for *numerable sites* (Definition 5.1) is given in Theorem 5.4. In particular, it works for the site of all smooth manifolds, where morphisms can be taken to be either arbitrary smooth maps, submersions, immersions, étale maps, or open embeddings. Instead of working with the site of all smooth manifolds we can restrict to the small site of some fixed smooth manifold. In particular, if we take open embeddings as morphisms we recover the usual notion of a sheaf on a manifold. The large site of a fixed smooth manifold can also be used. Smooth manifolds can be replaced with piecewise linear manifolds or topological manifolds.

Although the result appears to have a rather classical flavor, the author was unable to locate any reference in the literature that contains or readily implies it, and such a reference appears to be unknown to a nontrivial fraction of the mathematical community [2017.b]. Recent books like Amabel–Debray–Haine [2021, Theorem 3.6.1] only cite a weaker result (see, for example, Berwick–Evans–Boavida de Brito–Pavlov [2019, Theorem 5.1]), which shows that open covers with two elements and increasing chains of open embeddings generate the Grothendieck topology on smooth manifolds.

As an application, we leverage the existing work of Berwick–Evans–Boavida de Brito–Pavlov [2019, Theorem 5.1] to establish the following representability result for stacks on the site \mathbf{Top} of topological spaces with numerable open covers (Definition 3.16). See Theorem 8.5 for the full statement and details.

Theorem 1.2. Suppose

$$F: \mathbf{Top}^{\mathrm{op}} \rightarrow \mathbf{sSet}$$

is a simplicial presheaf on the site \mathbf{Top} . Define

$$\mathcal{C}F: \mathbf{Top}^{\mathrm{op}} \rightarrow \mathbf{sSet}, \quad \mathcal{C}F(X) = \operatorname{hocolim}_{n \in \Delta^{\mathrm{op}}} F(\Delta^n \times X),$$

where $\Delta^n \in \mathbf{Top}$ denotes the n -simplex as a topological space and the homotopy colimit is modeled by the diagonal of a bisimplicial set. If F satisfies the homotopy descent condition, then $\mathcal{C}F$ is representable in the following sense: the natural map of Definition 8.3

$$\mathcal{C}F(X) \rightarrow \operatorname{Map}(\operatorname{Sing} X, \mathcal{C}F(*))$$

is a weak equivalence whenever X is homotopy equivalent to a cofibrant topological space, e.g., X is a CW-complex, a topological manifold, or a polyhedron. In particular, we have a natural bijection of sets

$$F[X] \rightarrow [\operatorname{Sing} X, \mathcal{C}F(*)],$$

where $F[X]$ denotes concordance classes of the sheaf F over X .

The significance of this result is that it allows to easily establish Brown-type results about the existence of classifying spaces and representability of stacks. For example, it can be used to give a very short proof (Example 8.6) that for any topological group G the space BG classifies numerable principal G -bundles. Likewise, for any abelian topological group A , the Eilenberg–MacLane space $B^d A$ classifies numerable A -banded bundle $(d - 1)$ -gerbes (Example 8.7).

As another application, we give the following criterion for acyclicity of sheaves of abelian groups, see Corollary 6.2 for the proof. As shown in Corollary 6.7, the condition in this criterion is strictly more general than the usual notions of flabby, supple, and fine sheaves, yet it is easy to verify in practice.

Theorem 1.3. Suppose Site is a numerable site (Definition 5.1) and $F: \text{Site}^{\text{op}} \rightarrow \text{coCh}$ is a presheaf of cochain complexes such that in every cochain degree n the presheaf of abelian groups $F_n: \text{Site}^{\text{op}} \rightarrow \text{Ab}$ is a sheaf and for any $M \in \text{Site}$ and any covering family $\{U \rightarrow M, V \rightarrow M\}$ of M the map

$$F_n(U) \times F_n(V) \rightarrow F_n(U \cap V), \quad (u, v) \mapsto u|_{U \cap V} - v|_{U \cap V}$$

is surjective. Then F satisfies the homotopy descent property.

1.4. Acknowledgments

The author originally wrote up a proof of Theorem 4.8 for the paper Berwick-Evans–Boavida de Brito–Pavlov [2019], after polling on MathOverflow [2017.b] revealed that the result is not present in the literature. Subsequently, the authors of [2019] figured out a way to prove the main theorem of [2019] without this intermediate step, using a weaker lemma [2019, Theorem 5.1], which resulted in the original argument being split off as this note.

The author thanks Daniel Berwick-Evans and Pedro Boavida de Brito for helpful discussions.

2 Recollections on Grothendieck topologies

In this section we very briefly review the necessary definitions and facts about Grothendieck topologies. By Toën–Vezzosi [2002.b, Definition 3.3.1] (see also Lurie [2017.a, Remark 6.2.2.3]), on an ordinary category (as opposed to an ∞ -category) Grothendieck topologies in the sense of ∞ -categories coincide with ordinary Grothendieck topologies. The ∞ -categories that we consider are ordinary categories, so the usual notion of a Grothendieck topology suffices for our purposes.

Definition 2.1. (Johnstone [2002.a, Definition C.2.1.1].) A *coverage* on a category C is an assignment to every object $X \in C$ of a collection of families of morphisms (known as *generating covering families*) with codomain X such that for any such generating covering family $\{f_i: U_i \rightarrow X\}_{i \in I}$ and any morphism $g: Y \rightarrow X$ there is a generating covering family $\{h_j: V_j \rightarrow Y\}_{j \in J}$ such that for any $j \in J$ the morphism gh_j factors through the morphism f_i for some $i \in I$. A *site* is a category with a coverage. Used in 2.1, 2.3, 2.4, 2.5, 2.5*, 2.6, 2.7, 3.0*, 3.15, 3.16, 3.17, 4.1, 5.1, 5.8.

Definition 2.2. A *sieve* on an object X in a category C is a collection S of morphisms with codomain X such that $f \in S$ implies $fg \in S$ whenever fg is defined. Used in 2.3, 2.5, 5.8.

Definition 2.3. (Johnstone [2002.a, Definition C.2.1.8].) A *Grothendieck topology* (alias *Grothendieck coverage*) on a category C is an assignment to every object $X \in C$ of a collection of sieves on X (known as *covering sieves*) that together form a coverage on C , and the following two saturation conditions are satisfied: for any object $X \in C$ the maximal sieve (comprising all morphisms with codomain X) is a covering sieve, and if R is a covering sieve on X and S is a sieve on X such that $f^*S = \{g \mid fg \in S\}$ is a covering sieve for any $f \in R$, then S is also a covering sieve. Used in 2.3, 2.4, 2.5, 2.5*, 4.0*, 4.1, 4.8, 4.9, 5.4, 5.8.

Definition 2.4. The Grothendieck topology on C *generated* by a collection of families of morphisms $\{f_{k,i}: U_i \rightarrow X_k\}_{k \in K, i \in I_k}$ is the smallest Grothendieck topology (given by the intersection of all such topologies) for which the given families are contained in some covering sieves. In particular, we talk about the Grothendieck topology and covering sieves of a site. Used in 4.0*, 4.8, 4.9, 5.4.

Definition 2.5. A *covering family* for a site C is a family of morphisms $\{f_i: U_i \rightarrow X\}_{i \in I}$ in C such that the smallest sieve that contains this family is a covering sieve of C . Used in 2.5*, 2.6, 2.7, 4.0*, 4.2, 4.2*, 4.4, 4.4*, 4.5, 4.5*, 4.6, 4.6*, 4.7, 4.7*, 5.1, 5.2, 5.4, 5.4*, 5.8, 6.4.

The advantage of Grothendieck topologies is that passing from a Grothendieck topology to its category of sheaves defines an injective map from the collection of Grothendieck topologies on C to the collection of full subcategories of the category of presheaves on C . On the other hand, many different coverages can have the same category of sheaves. The advantage of coverages is that the given generating covering families can often be easier to work with than arbitrary covering sieves. For example, the coverage of **Man** (smooth manifolds) consisting of good open covers (covers whose finite intersections are either empty or diffeomorphic to \mathbf{R}^n) enjoys many special properties not shared by arbitrary covering sieves.

We spell out explicitly the particular formulation of the saturation properties for covering families that we will be using below.

Lemma 2.6. (cf. Johnstone [2002.a, Lemma C.2.1.6(i)].) Suppose C is a site. Given families of morphisms $\{f_i: U_i \rightarrow X\}_{i \in I}$ and $\{g_j: V_j \rightarrow X\}_{j \in J}$ in C such that any f_i factors through some g_j , if the family f is a covering family, then so is g . Used in 4.0*, 4.4*, 4.5*, 4.6*, 4.7*.

Lemma 2.7. (cf. Johnstone [2002.a, Lemma C.2.1.7(i)].) Suppose C is a site. Given a family $\{f_i: U_i \rightarrow X\}_{i \in I}$ and an I -indexed collection of families $\{g_{i,j}: V_{i,j} \rightarrow U_i\}_{j \in J_i}$ of morphisms in C , if the family f and all of g_i are covering families, then so is the family $\{f_i g_{i,j}: V_{i,j} \rightarrow X\}_{i \in I, j \in J_i}$. Used in 4.0*, 4.2*, 4.5*, 4.6*, 4.7*.

3 Numerable covers

In this section we review the definition of the site **Top** of topological spaces, continuous maps, and numerable open covers, i.e., open covers that admit a subordinate partition of unity. This site plays a fundamental role in defining bundle-like structures, such as principal bundles, vector bundles, (higher) bundle gerbes, etc., for which we say that a bundle is *numerable* if it is trivializable over a numerable open cover.

This is motivated by the fact that we typically expect or require that such bundle-like structures are classified by a map to some classifying space. Numerable bundles are stable under base change because numerable covers are stable under preimages (Tukey [1940, V.3.7]). Thus, if the universal bundle is numerable, then any bundle classified by a map to the classifying space (i.e., the base space of the universal bundle) must necessarily be numerable.

Indeed, the universal bundle over the classifying space of a topological group G such that G is a countable CW-complex is numerable because its base space is a CW-complex (Milnor [1956, Theorem 5.2]), CW-complexes are paracompact Hausdorff spaces (Miyazaki [1952.b]), and any open cover of a paracompact Hausdorff space is numerable (Michael [1953, Proposition 2]). The latter property of paracompact Hausdorff spaces is the reason why paracompact Hausdorff spaces are sometimes used in expositions related to classifying spaces. However, there is no need for such a restriction if one works with numerable covers, as we do below.

We start by recalling some properties of partitions of unity on topological spaces. It will be convenient to drop the normalization condition $\sum_{i \in I} f_i = 1$ and instead require that this sum exists and is a strictly positive continuous map. We refer to such families as positive partitions.

Definition 3.1. A *positive partition* on a topological space X is a family of continuous real-valued functions $\{f_i: X \rightarrow [0, \infty)\}_{i \in I}$ such that the map $\sum_{i \in I} f_i$ is everywhere defined and is a strictly positive continuous map $X \rightarrow (0, \infty)$. Used in 3.0*, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.7*, 3.8, 3.8*, 3.9, 3.10, 3.14*, 3.15*, 4.2*, 4.3*, 4.5*, 4.6*, 4.7*.

Remark 3.2. For any positive partition $\{f_i: X \rightarrow [0, \infty)\}_{i \in I}$ and for any $I' \subset I$ the sum $\sum_{i \in I'} f_i$ is everywhere defined and is a continuous map $X \rightarrow [0, \infty)$.

Definition 3.3. The *induced open cover* of a positive partition $\{f_i\}_{i \in I}$ on a topological space X is defined as the open cover $\{f_i^*(0, \infty)\}_{i \in I}$ of X . We say that a positive partition $\{f_i: X \rightarrow [0, \infty)\}_{i \in I}$ is *compatible* with an open cover $\{U_i\}_{i \in I}$ of X if $f_i^*(0, \infty) \subset U_i$ for all $i \in I$ and *subordinate* to U if the closure of $f_i^*(0, \infty)$ in X is a subset of U_i for all $i \in I$. Used in 1.1*, 3.0*, 3.4, 3.5, 3.6, 3.7*, 3.8, 3.8*, 3.9, 3.10, 3.12, 3.14*, 4.2*, 4.3*, 4.5*, 4.6*, 4.7*, 5.5, 6.4, 6.7*.

Remark 3.4. The above definition could be extended to the case when the indexing sets of positive partitions and open covers are different. For example, we could say that a positive partition $\{f_i: X \rightarrow [0, \infty)\}_{i \in I}$

is compatible with an open cover $\{U_j\}_{j \in J}$ of X if there is a map of sets $\alpha: I \rightarrow J$ such that $f_i^*(0, \infty) \subset U_{\alpha(i)}$ for all $i \in I$. However, in this case we can define a new positive partition $\{g_j: X \rightarrow [0, \infty)\}_{j \in J}$ by setting $g_j = \sum_{i: \alpha(i)=j} f_i$. The resulting positive partition has the same indexing set as the open cover $\{U_j\}_{j \in J}$ and is compatible with this open cover. This reindexing procedure works for all definitions and constructions below, e.g., a locally finite open cover could be reindexed by taking unions indexed by $\alpha^*\{j\}$, etc. Because of this, our constructions use a fixed indexing set for simplicity.

Definition 3.5. A positive partition $\{f_i\}_{i \in I}$ on a topological space X is *locally finite* if its induced open cover is a locally finite open cover, i.e., the collection of open subsets $V \subset X$ such that $f_i|_V = 0|_V$ for all but finitely many $i \in I$ forms an open cover of X . Used in 3.7*, 3.8, 3.8*, 3.10, 3.12, 3.14*, 5.5.

Remark 3.6. Occasionally, a weaker property is used. A positive partition $\{f_i\}_{i \in I}$ on a topological space X is *point-finite* if for any point $x \in X$ there is only finitely many $i \in I$ such that $f_i(x) \neq 0$, i.e., any $x \in X$ belongs only to finitely many elements of the induced open cover of f . We make no use of this notion below.

Definition 3.7. A *partition of unity* is a positive partition such that $\sum_{i \in I} f_i = 1$. Used in 1.1*, 3.0*, 3.7*, 3.8, 3.8*, 3.10, 3.12, 3.14*, 4.0*, 4.3*, 5.5, 6.5, 6.7*, 7.3*.

Often, the condition of local finiteness is included in the definition of a partition of unity. We will use both versions below, so for us it is important to be able to distinguish between them.

The following proposition is due to Michael R. Mather, see Dold [1972.a, Proposition A.2.8]. It shows that any positive partition f can be improved to a locally finite partition of unity g that is subordinate to the induced open cover of f . Dold's statement only claims compatibility, not subordination (i.e., does not mention closure), but our stronger version follows from his proof.

Proposition 3.8. For any positive partition $\{f_i: X \rightarrow [0, \infty)\}_{i \in I}$ on a topological space X we can find a locally finite partition of unity $\{g_i: X \rightarrow [0, \infty)\}_{i \in I}$ on X such that g is subordinate to the induced open cover of f . Used in 3.10, 3.13*.

Proof. We normalize f_i by dividing it by $\sum_{i \in I} f_i$, which converts $\{f_i\}_{i \in I}$ into a partition of unity. Set $\mu = \sup f_i: X \rightarrow (0, 1]$ and $g_i = \max(0, 2f_i - \mu): X \rightarrow [0, 1]$. By construction, the family $\{g_i\}_{i \in I}$ is a positive partition. The collection of open sets $V_J = h_J^*(0, \infty)$, where $h_J = \mu/2 - 1 + \sum_{i \in J} f_i$ for a finite subset $J \subset I$, forms an open cover of X because $\sum_{i \in I} f_i = 1$. We claim that the open cover $\{V_J\}_J$ exhibits the local finiteness of the positive partition $\{g_i\}_{i \in I}$. Indeed, for any finite subset $J \subset I$ and for any $i \in I \setminus J$ we have

$$f_i|_{V_J} \leq \sum_{k \in I \setminus J} f_k = 1 - \sum_{k \in J} f_k < \mu|_{V_J}/2,$$

so $g_i|_{V_J} = 0$, as desired.

Since $2f_i - \mu < 0$ on an open subset of X , we must have $2f_i - \mu \geq 0$ on the closure of $g_i^*(0, \infty)$. Since $\mu > 0$ everywhere, we have $f_i > 0$ on the closure of $g_i^*(0, \infty)$, so g is subordinate to the induced open cover of f . Replacing g_i with $g_i / \sum_{i \in I} g_i$, we obtain the desired locally finite partition of unity. ■

Thus, an open cover of a topological space admits a subordinate locally finite partition of unity if and only if it admits a compatible positive partition. Such covers play a fundamental role in general topology, where they are known as normal covers (introduced by John W. Tukey [1940, §V.2]), and in the theory of classifying spaces, where they are known as numerable covers (following Dold [1963, Definitions 2.1]).

Definition 3.9. An open cover $\{U_i\}_{i \in I}$ of a topological space X is *numerable* if it admits a compatible positive partition. Used in 1.1*, 3.0*, 3.8*, 3.10, 3.11, 3.13, 3.13*, 3.14, 3.14*, 3.15, 3.15*, 3.16, 3.17, 4.0*, 4.1, 4.2, 4.2*, 4.3, 4.3*, 4.4, 4.4*, 4.5, 4.5*, 4.6, 4.6*, 4.7, 4.7*, 4.8, 4.9, 5.5, 5.8.

Below we will also apply this definition to open covers of a subset X of some topological space Y , in which case we pass to the induced topology on X first.

The following proposition is a combination of results of Arthur H. Stone [1948], Ernest Michael [1953], and Kiiti Morita [1962, Theorem 1.2].

Recall that a *cozero set* is an (open) subset $U \subset X$ such that $U = f^*(0, \infty)$ for some continuous map $f: X \rightarrow [0, \infty)$. Recall also that an open cover $\{U_i\}_{i \in I}$ is a *star refinement* of an open cover $\{V_j\}_{j \in J}$ if the

open cover

$$\left\{ \bigcup_{k \in I: U_k \cap U_i \neq \emptyset} U_k \right\}_{i \in I}$$

refines V . A family $\{U_i\}_{i \in I}$ of open sets is *discrete* if there is an open cover $\{V_j\}_{j \in J}$ of X such that for every $j \in J$ we have $U_i \cap V_j \neq \emptyset$ for at most one $i \in I$. (Any discrete family is in particular disjoint.)

Proposition 3.10. (Ernest Michael, Kiiti Morita, Arthur H. Stone, see Morita [1962, Theorem 1.2]; also Derwent [1968].) For an open cover $\{U_i\}_{i \in I}$ of a topological space X the following properties are equivalent.

- U is a numerable cover, i.e., it admits a compatible positive partition;
- (Michael [1953, Proposition 2]; see Proposition 3.8.) U admits a subordinate locally finite partition of unity;
- (Michael [1953, Proposition 1], Hoshina [1989, Theorem 1.1 and 1.2].) U can be refined by an open cover given by the union of a countable collection of discrete families of cozero sets.
- (Michael [1953, Theorem 1], Morita [1964, Theorem 1.2].) U can be refined by an open cover given by the union of a countable collection of locally finite families of cozero sets.
- (Stone [1948, Theorems 1 and 2].) U is a *normal cover*, meaning there is a sequence $W_0 = U$, W_1 , W_2 , \dots of open covers of X such that W_{n+1} is a star refinement of W_n for all $n \geq 0$;
- U can be refined by the inverse image of an open cover of Y under some continuous map $X \rightarrow Y$, where Y is a metrizable topological space;
- (Mardešić–Segal [1982.b, Lemma I.6.1].) U can be refined by the inverse image of an open cover of Y under some continuous map $X \rightarrow Y$, where Y is an absolute neighborhood retract, i.e., a metrizable topological space Y such that any closed embedding $Y \rightarrow Z$ into a metrizable topological space Z factors through an open subset $U \subset Z$ such that there is a map $U \rightarrow Y$ for which the composition $Y \rightarrow U \rightarrow Y$ is identity;
- U can be refined by a locally finite normal open cover;
- U can be refined by a locally finite open cover consisting of cozero sets;

Used in 1.1*, 3.8*, 3.10, 3.13*, 3.16*.

Remark 3.11. We point out some incompatible definitions of normal and numerable covers in the literature. Hurewicz [1955, §5] defines normal covers as open covers consisting of cozero sets. Hurewicz uses this property in conjunction with local finiteness. Spanier [1966, the paragraph before Lemma 2.7.10] defines numerable covers as locally finite normal covers in the sense of Hurewicz. May [1975, §1] defines numerable covers as locally finite open covers consisting of cozero sets.

Remark 3.12. Normal spaces are precisely the topological spaces for which every finite open cover is normal (Tukey [1940, Theorem V.4.1]). Likewise, paracompact (i.e., every open cover has a locally finite refinement) Hausdorff spaces are precisely the topological spaces that satisfy the T_1 axiom and for which every open cover is normal (Stone [1948, Theorem 1]), or, equivalently, admits a locally finite compatible partition of unity (Michael [1953, Proposition 2]).

Remark 3.13. *Disjoint open covers* are always numerable: take f_i to be the characteristic function of U_i .

Used in 4.0*, 4.2, 4.2*, 4.4, 4.6, 4.7, 4.7*, 4.8, 4.9.

Most of Proposition 3.10 plays no role in what follows, except for the equivalence between first three properties, which we prove separately as Proposition 3.8 and Proposition 3.14. Our proof of Proposition 3.14 follows Husemöller–Joachim–Jurčo–Schottenloher [2008.a, Proposition 7.1.2]. Although Proposition 3.14 holds for discrete families instead of disjoint families (with the same proof), the latter case is sufficient for our purposes.

Proposition 3.14. (Michael [1953, Proposition 1]; Hoshina [1989, Theorem 1.1 and 1.2].) Any numerable open cover $\{U_i\}_{i \in I}$ of a topological space X can be refined by a numerable open cover $\{V_j\}_{j \in J}$ for which there is a map of sets $\lambda: J \rightarrow K$, where K is countable and for any $k \in K$ the family $\{V_j\}_{\lambda(j)=k}$ is disjoint.

Used in 3.13*, 3.16*, 4.0*, 4.7*.

Proof. Suppose $\{f_i\}_{i \in I}$ is a locally finite partition of unity subordinate to $\{U_i\}_{i \in I}$. Denote by J the collection of all finite nonempty subsets of I . Denote by K the set $\{1, 2, 3, \dots\}$ and by $\lambda: J \rightarrow K$ the map that computes

cardinality. Define an open cover $\{V_j\}_{j \in J}$ of X by

$$V_j = \{x \in X \mid f_k(x) > f_l(x) \text{ for all } k \in j \text{ and } l \in I \setminus j\},$$

where V_j is open because for any $x \in V_j$ we can find an open neighborhood W such that $f_l|_W \neq 0$ for only finitely many $l \in I \setminus j$, so the condition $f_k(x) > f_l(x)$ is an open condition. The family $\{V_j\}_{j \in J}$ is an open cover of X . Define a positive partition $\{g_j\}_{j \in J}$ exhibiting the numerability of V by

$$g_j(x) = \max \left(0, \min_{k \in j, l \in I \setminus j} (f_k(x) - f_l(x)) \right).$$

Again, the minimum exists and g_j is continuous because $\{f_i\}_{i \in I}$ is locally finite. Also, $V_j = g_j^*(0, \infty)$ and $V_j \cap V_{j'} = \emptyset$ whenever $j \setminus j' \neq \emptyset$ and $j' \setminus j \neq \emptyset$, by construction. In particular, for a fixed cardinality of j , all V_j are disjoint. Since $V_j \subset U_k$ for any $k \in j$, the open cover V refines the open cover U , which completes the proof. ■

Proposition 3.15. (Tukey [1940, V.3.7].) Numerable open covers form a coverage on the category of topological spaces and continuous maps.

Proof. Suppose X is a topological space, $\{U_i\}_{i \in I}$ is a numerable open cover of X , and $g: Y \rightarrow X$ is a continuous map. It suffices to show that $\{g^*U_i\}_{i \in I}$ is a numerable open cover of Y . Indeed, if $\{f_i: X \rightarrow [0, \infty)\}_{i \in I}$ is a positive partition on X such that $f_i^*(0, \infty) \subset U_i$ for all $i \in I$, then $\{f_i g: Y \rightarrow [0, \infty)\}_{i \in I}$ is a positive partition on Y such that $(f_i g)^*(0, \infty) = g^*(f_i^*(0, \infty)) \subset g^*U_i$. ■

Definition 3.16. Denote by **Top** the site of topological spaces and continuous maps, equipped with the coverage of numerable open covers. Used in 1.1*, 5.0*.

We finish this section by defining another site to which our main theorem is applicable, the site of locales and their morphisms. A *locale* is a poset that has finite infima and arbitrary suprema such that the map $b \mapsto \inf(a, b) = a \wedge b$ preserves suprema for any fixed a , and a *morphism of locales* is an order-preserving map of posets in the opposite direction that preserves finite infima and arbitrary suprema. These properties axiomatize the properties of the poset of open subsets of a topological space and the induced inverse image map on open subsets associated to a continuous map. In particular, we have a functor from topological spaces to locales. This functor becomes fully faithful on *sober spaces*, a large class of topological spaces that includes all Hausdorff spaces. The book by Picado–Pultr [2012] is a comprehensive introduction to locales, see, in particular, Theorem IX.2.3.4 there, which is a pointfree analogue of (a part of) Proposition 3.10. To keep the paper accessible to a larger audience, we formulate our lemmas and propositions using topological spaces. However, nothing in the statements or proofs depends on the availability of points, since all claims are formulated using open sets only. In particular, the use of points in the proof of Proposition 3.14 is merely a notational convenience.

Definition 3.17. Denote by **Loc** the site of locales and their morphisms, equipped with the coverage of numerable open covers. Used in 4.1, 5.0*.

4 Main theorem for topological spaces

In this section, we prove Theorem 4.8. The proof repeatedly uses Lemma 2.6 and Lemma 2.7 to show that more and more numerable covers are covering families in the Grothendieck topology generated by disjoint open covers (indexed by arbitrary sets) and numerable open covers with two elements. First, in Lemma 4.2 we show that ‘zigzag’ covers, i.e., covers P_0, P_1, \dots where the only nonempty intersections are $P_k \cap P_{k+1}$, are covering families. Next, using a trick with partitions of unity explained in Lemma 4.3, we can refine any countable cover of the form $U_0 \subset U_1 \subset \dots$ by a zigzag cover, which allows us to show that such increasing chains of open subsets are covering families (Lemma 4.4). Lemma 4.5 shows that finite numerable covers are covering families, by induction on the size of the cover. This is then used in Lemma 4.6 to show that arbitrary countable covers are covering families, whereas Lemma 4.7 uses Lemma 4.6 and Proposition 3.14 to show that covers of arbitrary cardinality are covering families. Together with Remark 3.13, Lemma 4.7 implies Theorem 4.8.

Definition 4.1. Denote by \mathbf{C} the category of topological spaces and continuous maps equipped with some arbitrary Grothendieck topology (not necessarily the Grothendieck topology generated by the coverage of numerable open covers). More generally, \mathbf{C} can denote the category of locales, as in Definition 3.17, but equipped with some arbitrary Grothendieck topology, not necessarily given by numerable open covers.

The following lemma and the underlying notion of a zigzag cover is essentially due to Greub–Halperin–Vanstone (see the proof of Proposition I.II in [1972.b]).

Lemma 4.2. If countable disjoint open covers and numerable open covers with two elements are covering families in \mathbf{C} , then so is any *zigzag cover*, i.e., a numerable open cover $\{P_0, P_1, P_2, \dots\}$ of X such that $P_k \cap P_l = \emptyset$ whenever $|k - l| > 1$. Used in 4.0*, 4.3, 4.4*.

Proof. The only nontrivial intersections of elements of P are $P_k \cap P_{k+1}$ for $k \geq 0$. Take

$$A = P_0 \cup P_2 \cup P_4 \cup \dots,$$

the (disjoint) union of even elements of P , and

$$B = P_1 \cup P_3 \cup P_5 \cup \dots,$$

the (disjoint) union of odd elements of P . We have $A \cup B = X$ and $A \cap B$ equals the (disjoint) union of $P_k \cap P_{k+1}$ for all $k \geq 0$. The open cover $\{A, B\}$ of X is numerable: if $f_i: X \rightarrow [0, 1]$ is a positive partition compatible with P , then $\sum_{i \geq 0} f_{2i}$ and $\sum_{i \geq 0} f_{2i+1}$ form a positive partition compatible with $\{A, B\}$. The numerable open cover $\{A, B\}$ of X is a covering family by assumption. The disjoint open covers $\{P_0, P_2, P_4, \dots\}$ of A and $\{P_1, P_3, P_5, \dots\}$ of B are covering families by assumption. By Lemma 2.7, $\{P_0, P_1, P_2, \dots\}$ is a covering family of X . ■

The following lemma constitutes the technical core of the proof. It shows that any increasing chain of open subsets can be refined by a zigzag cover.

Lemma 4.3. Suppose $U_0 \subset U_1 \subset U_2 \subset \dots$ is an increasing sequence of open subsets of a topological space X that forms a numerable open cover of X . Then there is a numerable open cover $P = \{P_0, P_1, P_2, \dots\}$ of X such that $P_i \subset U_i$ and $P_i \cap P_j = \emptyset$ whenever $|i - j| > 1$. (Thus, P is a zigzag cover of X in the sense of Lemma 4.2.) Used in 4.0*, 4.4*.

Proof. Choose a partition of unity $\{f_i: X \rightarrow [0, 1]\}_{i \geq 0}$ compatible with U . Define a sequence $\{g_i\}_{i \geq -2}$ of functions

$$g_i: X \rightarrow [0, 1], \quad g_i = \sum_{0 \leq k \leq i} f_k.$$

Fix some sequences α, β , and γ of real numbers such that

$$1 > \alpha_i > \beta_i > \gamma_i > \alpha_{i+1} > \beta_{i+1} > 0$$

for all $i \geq -2$. Now take

$$P_i = g_{i-2}^*[0, \beta_{i-2}) \cap g_i^*(\alpha_i, 1]$$

for $i \geq 0$. (As one can see from the formula, here the boundary -2 used above becomes important.) Observe that

$$P_i \subset g_i^*(\alpha_i, 1] \subset g_i^*(0, 1] \subset \bigcup_{0 \leq k \leq i} f_k^*(0, 1] \subset \bigcup_{0 \leq k \leq i} U_k = U_i.$$

Next we show that $X = \bigcup_{i \geq 0} P_i$, for which it suffices to show by induction on k that $\bigcup_{0 \leq i \leq k} P_i = g_k^*(\alpha_k, 1]$, since $\bigcup_{k \geq 0} g_k^*(\alpha_k, 1] = \bar{X}$. The claim holds for $k = -2$ and $k = -1$ because the union is empty and $g_k = 0$. Thus, suppose $\bigcup_{0 \leq i < k} P_i = g_{k-1}^*(\alpha_{k-1}, 1]$ for some $k \geq 0$. We have to show that $g_{k-1}^*(\alpha_{k-1}, 1] \cup P_k = g_k^*(\alpha_k, 1]$. One inclusion is trivial and the other boils down to

$$g_k^*(\alpha_k, 1] \subset g_{k-1}^*(\alpha_{k-1}, 1] \cup g_{k-2}^*[0, \beta_{k-2}),$$

which we strengthen to

$$g_k^*(\alpha_k, 1] \subset g_{k-1}^*(\alpha_{k-1}, 1] \cup g_{k-1}^*[0, \beta_{k-2}) = g_{k-1}^*((\alpha_{k-1}, 1] \cup [0, \beta_{k-2})) = g_{k-2}^*[0, 1] = X.$$

Now we demonstrate that $P_i \cap P_j = \emptyset$ for all $i \geq 0$ and $j \geq i + 2$. Once we restrict to $P_i \cap P_j$ we have the following inequalities: $g_i > \alpha_i$, $g_{i-2} < \beta_{i-2}$, $g_j > \alpha_j$, $g_{j-2} < \beta_{j-2}$. (Abusing notation, α_k and β_l denote constant functions on $P_i \cap P_j$ with indicated values.) Thus

$$\alpha_i < g_i \leq g_{j-2} < \beta_{j-2} < \alpha_i$$

must hold on $P_i \cap P_j$, which implies $P_i \cap P_j = \emptyset$.

To show that the open cover P of X is numerable, we construct a positive partition that is compatible with P . Take $h_i = 2^{-i} \cdot \max(0, \beta_{i-2} - g_{i-2}) \cdot \max(0, g_i - \alpha_i)$. By construction, $P_i = h_i^*(0, \infty)$. Thus, $\sum_{i \geq 0} h_i$ exists and is a strictly positive continuous map $X \rightarrow (0, 1]$ because P covers X and $h_i \leq 2^{-i}$. ■

Lemma 4.4. If countable disjoint open covers and numerable covers with two elements are covering families in \mathbb{C} , then so is any numerable open cover

$$\{U_0, U_1, U_2, \dots\}$$

of X such that $U_0 \subset U_1 \subset U_2 \subset \dots$. *Used in 4.0*, 4.6*.*

Proof. Using Lemma 4.3, we construct a numerable open cover $\{P_0, P_1, P_2, \dots\}$ of X such that $P_k \subset U_k$ and $P_k \cap P_l = \emptyset$ whenever $|k - l| > 1$, so the only nontrivial intersections are $P_k \cap P_{k+1}$ for $k \geq 0$. We have $P_k \subset U_k$ and P is a covering family by Lemma 4.2. By Lemma 2.6 U is also a covering family. ■

Lemma 4.5. If the empty cover and numerable covers with two elements are covering families in \mathbb{C} , then so is any finite numerable open cover. *Used in 4.0*, 4.6*.*

Proof. We prove by induction on n that numerable open covers with fewer than n elements are covering families. The empty cover (of the empty set) is a covering family by assumption, which establishes the base of the induction. Suppose all numerable open covers with fewer than $n > 0$ elements are covering families and $\{U_0, \dots, U_n\}$ is a numerable open cover of X with a positive partition $\{f_i: X \rightarrow [0, \infty] \mid 0 \leq i \leq n\}$. Set $A = (\sum_{i < n} f_i)^*(0, \infty) \cap \bigcup_{i < n} U_i$. The family $\{U_0 \cap A, \dots, U_{n-1} \cap A\}$ forms a numerable cover of A with a positive partition consisting of the restrictions of f_0, \dots, f_{n-1} because $\sum_{i < n} f_i > 0$ on A . This family has $n - 1$ elements, so it is a covering family by induction. The family $\{A, U_n\}$ is a numerable cover of X with a compatible positive partition $\sum_{i < n} f_i$ and f_n , hence a covering family by assumption. (The support of $\sum_{i < n} f_i$ need not be a subset of A , so we only require compatibility, not subordination.) By Lemma 2.7, $\{U_0 \cap A, \dots, U_{n-1} \cap A, U_n\}$ is a covering family of X . Hence $\{U_0, \dots, U_n\}$ is also a covering family of X by Lemma 2.6. ■

Lemma 4.6. If countable disjoint open covers and numerable covers with two elements are covering families in \mathbb{C} , then so is any countable numerable open cover. *Used in 4.0*, 4.7*.*

Proof. Suppose $\{V_i\}_{i \geq 0}$ is a countable numerable open cover of a topological space X with a compatible positive partition $\{f_i: X \rightarrow [0, \infty)\}_{i \geq 0}$. Set

$$U_i = V_0 \cup V_1 \cup \dots \cup V_i$$

and

$$g_i = f_0 + \cdots + f_i: X \rightarrow [0, \infty)$$

for all $i \geq 0$. Set $A_i = g_i^*(0, \infty)$. We have $A_0 \subset A_1 \subset A_2 \subset \cdots$ and $\bigcup_{i \geq 0} A_i = X$. Also $\{f_i\}_{i \geq 0}$ is a positive partition compatible with the open cover $\{A_i\}_{i \geq 0}$, so A is a numerable open cover of X . By Lemma 4.4, A is a covering family of X .

Next, $\{V_0 \cap A_i, V_1 \cap A_i, \dots, V_i \cap A_i\}$ is a numerable open cover of A_i with a positive partition given by the restrictions of f_0, \dots, f_i . By Lemma 4.5, $\{V_0 \cap A_i, V_1 \cap A_i, \dots, V_i \cap A_i\}$ is a covering family of A_i . Thus, by Lemma 2.7, $\{V_i \cap A_j\}_{0 \leq i \leq j}$ is a covering family of X . This covering family refines the open cover $\{V_i\}_{i \geq 0}$, so by Lemma 2.6, $\{V_i\}_{i \geq 0}$ is a covering family of X . ■

Lemma 4.7. If disjoint open covers indexed by arbitrary sets and numerable covers with two elements are covering families in \mathbf{C} , then so is any numerable open cover. Used in 4.0*, 4.7*.

Proof. By Proposition 3.14, any numerable open cover $\{U_i\}_{i \in I}$ of a topological space X can be refined by a numerable open cover $\{V_j\}_{j \in J}$ for which there is a map of sets $f: J \rightarrow K$, where K is countable and for any $k \in K$ the family $\{V_j\}_{f(j)=k}$ is disjoint. Choose a positive partition $\{h_j: X \rightarrow [0, \infty)\}_{j \in J}$ compatible with V . Set $W_k = \bigcup_{j: f(j)=k} V_j$ and $h_k = \sum_{j: f(j)=k} h_j$. Then $\{h_k\}_{k \in K}$ is a positive partition compatible with W , so W is a countable numerable open cover. By Lemma 4.6, W is a covering family of X . Furthermore, for every $k \in K$ the family $\{V_j\}_{f(j)=k}$ is a disjoint open cover of W_k , so also a covering family of W_k by assumption. Thus, by Lemma 2.7, $\{V_j\}_{j \in J}$ is a covering family of X that refines U . Hence, by Lemma 2.6, $\{U_i\}_{i \in I}$ is also a covering family of X . ■

Lemma 4.7 combined with Remark 3.13 immediately implies our main result.

Theorem 4.8. The Grothendieck topology of numerable open covers on \mathbf{Top} is generated by disjoint open covers (indexed by arbitrary sets) and numerable covers with two elements. Used in 1.0*, 1.1*, 1.4*, 4.0*, 4.8*, 5.0*, 6.0*, 6.1*, 8.2*.

All proofs in this section were carefully formulated to use only open sets, not individual points of X , and they work equally well without any modification in the following generalization of Theorem 4.8 to locales.

Theorem 4.9. The Grothendieck topology of numerable open covers on \mathbf{Loc} is generated by disjoint open covers (indexed by arbitrary sets) and numerable covers with two elements. Used in 5.4*.

5 Numerable sites

We would like to extend Theorem 4.8 to other sites, such as smooth manifolds. To this end we axiomatize the needed properties in the notion of a numerable site.

The most general notion of a numerable site uses the localic site \mathbf{Loc} (Definition 3.17). Readers not familiar with locales can substitute the site \mathbf{Top} (Definition 3.16) instead of \mathbf{Loc} in Definition 5.1 without losing any essential examples.

Definition 5.1. A *numerable site* is a site S equipped with a functor $\mathcal{U}: S \rightarrow \mathbf{Loc}$ such that \mathcal{U} admits cartesian lifts for all open inclusions and \mathcal{U} creates covering families, meaning the \mathcal{U} -image of a family f is a covering family (Definition 2.5) in \mathbf{Loc} if and only if f is a covering family in S . Used in 1.1*, 1.3, 5.0*, 5.2, 5.4, 5.4*, 5.5, 5.6, 5.7, 5.8, 5.9, 6.1, 6.3*, 6.4, 6.5, 6.6, 6.6*, 6.7.

Remark 5.2. The first condition in Definition 5.1 means that for any object X in S and any open inclusion $V \subset \mathcal{U}(X)$ there is a unique cartesian arrow $h: \hat{V} \rightarrow X$ that lifts $V \rightarrow \mathcal{U}(X)$. The latter means that for any morphism $f: W \rightarrow X$ in S such that the map $\mathcal{U}(f): \mathcal{U}(W) \rightarrow \mathcal{U}(X)$ factors through V , there is a unique morphism $g: W \rightarrow \hat{V}$ such that $hg = f$ and $\mathcal{U}(g)$ equals the map $\mathcal{U}(W) \rightarrow V$.

As an example, if $\{U \rightarrow M, V \rightarrow M\}$ is a covering family in S , then we can make define the intersection $U \cap V$ together with inclusions $U \cap V \rightarrow U$, $U \cap V \rightarrow V$ as the cartesian lift of $\mathcal{U}(U) \cap \mathcal{U}(V) \rightarrow \mathcal{U}(U)$, etc.

Used in 5.4*, 6.1, 6.7*.

Definition 5.3. A family $\{f_i: U_i \rightarrow X\}_{i \in I}$ of morphisms in a numerable site S is *disjoint* if the family $\{\mathcal{U}(f_i) \rightarrow \mathcal{U}(X)\}_{i \in I}$ factors through disjoint open inclusions in \mathbf{Loc} . Used in 5.4.

Theorem 5.4. If $\mathcal{U}: S \rightarrow \mathbf{Loc}$ is a numerable site, then the Grothendieck topology on S is generated by covering families with two elements and disjoint covering families indexed by a set. Used in 1.1*.

Proof. Using Remark 5.2, the proof of Theorem 4.9 and its dependencies continues to work for numerable sites: all constructions of covering families first take the \mathcal{U} -image in \mathbf{Loc} and construct the desired open covers in \mathbf{Loc} . Then, the above property allows us to lift the newly constructed open inclusions to S , and, furthermore, ensure that morphisms whose image factors through the new open inclusion themselves factor through the constructed lift. \blacksquare

Example 5.5. Normal Hausdorff topological spaces, continuous maps, and locally finite covers form a numerable site. Indeed, any locally finite open cover of a normal Hausdorff topological space admits a subordinate locally finite partition of unity (Bourbaki [1974, Proposition IX.4.3.3]), hence is a numerable cover.

Example 5.6. Paracompact Hausdorff topological spaces (Dieudonné [1944.b, §1]), continuous maps, and all open covers form a numerable site. Indeed, by Stone [1948, Theorem 1] the class of fully normal (i.e., every open cover is normal) Hausdorff spaces coincides with the class of paracompact (i.e., every open cover has a locally finite refinement) Hausdorff spaces.

Example 5.7. Paracompact Hausdorff smooth, PL, or topological manifolds together with smooth (respectively PL or continuous) maps, and all open covers form a numerable site.

Example 5.8. For a numerable site S and some object $B \in S$ the slice category S/B inherits a Grothendieck topology from S (a sieve is covering if and only if its image under the forgetful functor $S/B \rightarrow S$ is a covering family). The resulting site is known as the *big (gros) site* of B . It is numerable because the forgetful functor $S/B \rightarrow S$ creates covering families by construction, the functor $\mathcal{U}: S \rightarrow \mathbf{Loc}$ creates covering families by definition, hence the composition $S/B \rightarrow S \rightarrow \mathbf{Loc}$ also creates covering families. Cartesian lifts for $S/B \rightarrow \mathbf{Loc}$ can be constructed by lifting to S , which automatically provides the desired map to B .

Example 5.9. For a numerable site S and some object $B \in S$ the *little (petit) site* of B is the subcategory of the slice category S/B consisting of objects given by morphisms in S whose image under $\mathcal{U}: S \rightarrow \mathbf{Loc}$ is an open embedding and morphisms are commutative triangles in S whose image under \mathcal{U} is a commutative triangle of open embeddings. This subcategory is equipped with the induced Grothendieck topology from S/B . This is a numerable site since cartesian lifts of open inclusions by definition are morphisms in the little site.

6 Homotopy descent for chain complexes of sheaves of abelian groups

In this section, we use Theorem 4.8 to establish a simple sufficient condition (Theorem 6.1, Corollary 6.2) for a chain complex of sheaves of abelian groups to satisfy the homotopy descent property. The surjectivity condition in Corollary 6.2 below implies that all of the sheaves F_n are acyclic (Remark 6.3), so the theorem can be seen as an easy way to show that a sheaf of abelian groups is acyclic. On the other hand, the required property is strictly weaker than being flabby or fine (see Corollary 6.7), so our result does not follow from the classical theory.

Theorem 6.1. Suppose $\mathcal{U}: \mathbf{Site} \rightarrow \mathbf{Loc}$ is a numerable site (Definition 5.1) and $F: \mathbf{Site}^{\text{op}} \rightarrow \mathbf{coCh}$ is a presheaf of cochain complexes. Then F satisfies the homotopy descent property whenever the following conditions are satisfied.

- In every cochain degree n the presheaf of abelian groups $F_n: \mathbf{Site}^{\text{op}} \rightarrow \mathbf{Ab}$ sends disjoint unions to products.
- For any $M \in \mathbf{Site}$ and any covering family $\{U \rightarrow M, V \rightarrow M\}$ of M the map

$$F_n(U) \times F_n(V) \rightarrow F_n(U \cap V), \quad (u, v) \mapsto u|_{U \cap V} - v|_{U \cap V}$$

is surjective in every cochain degree n . (Here the map $U \cap V \rightarrow U$ is constructed as the cartesian lift of $\mathcal{U}(U) \cap \mathcal{U}(V) \rightarrow \mathcal{U}(U)$ and likewise for $U \cap V \rightarrow V$, see Remark 5.2.)

- For any $M \in \mathbf{Site}$ and any covering family $\{U \rightarrow M, V \rightarrow M\}$ of M the map

$$F(M) \rightarrow \ker(F(U) \times F(V) \rightarrow F(U \cap V))$$

is a quasi-isomorphism.

Used in 6.0*, 6.2*, 7.2*.

Proof. According to Theorem 4.8, the homotopy descent property for F is equivalent to the homotopy descent property for disjoint covers and covers with two elements. The homotopy descent property for disjoint covers holds because homotopy products of cochain complexes can be computed as ordinary products, and the latter are computed degreewise, so we conclude by the first property in the statement.

The case of covers with two elements means that for any $M \in \mathbf{Site}$ and a cover $\{U, V\}$ of M the restriction map

$$F(M) \rightarrow F(U) \times_{F(U \cap V)}^h F(V)$$

is a quasi-isomorphism. Computing the homotopy pullback as a homotopy equalizer, we reduce the problem to showing that in the sequence

$$F(M) \rightarrow F(U) \times F(V) \rightarrow F(U \cap V)$$

the left map (given by restrictions to U and V) is the homotopy fiber of the right map (given by $(u, v) \mapsto (u|_{U \cap V} - v|_{U \cap V})$). The right map is surjective by assumption. Degreewise surjections are fibrations in the projective model structure on cochain complexes, so the homotopy fiber can be computed as the kernel of the right map. Thus, it remains to show that the map from $F(M)$ to the kernel of the right map is a quasi-isomorphism, which holds by assumption. ■

Corollary 6.2. Suppose \mathbf{Site} is a numerable site and $F: \mathbf{Site}^{\text{op}} \rightarrow \mathbf{coCh}$ is a presheaf of cochain complexes such that in every cochain degree n the presheaf of abelian groups $F_n: \mathbf{Site}^{\text{op}} \rightarrow \mathbf{Ab}$ is a sheaf and for any $M \in \mathbf{Site}$ and any covering family $\{U \rightarrow M, V \rightarrow M\}$ of M the map

$$F_n(U) \times F_n(V) \rightarrow F_n(U \cap V), \quad (u, v) \mapsto u|_{U \cap V} - v|_{U \cap V}$$

is surjective. Then F satisfies the homotopy descent property. Used in 1.2*, 6.0*, 6.3.

Proof. We apply Theorem 6.1. The first property holds because F_n is a sheaf for any n . The second property holds by assumption. The third property holds because F_n is a sheaf for any n , so the map under consideration is an isomorphism. ■

Remark 6.3. In the language of classical abelian sheaf cohomology, Corollary 6.2 says that a sheaf of abelian groups $F: \mathbf{Site}^{\text{op}} \rightarrow \mathbf{Ab}$ is acyclic whenever for any $M \in \mathbf{Site}$ and any covering family $\{U \rightarrow M, V \rightarrow M\}$ of M the map

$$F_n(U) \times F_n(V) \rightarrow F_n(U \cap V), \quad (u, v) \mapsto u|_{U \cap V} - v|_{U \cap V}$$

is surjective for every n . Used in 6.0*, 6.6*, 6.7*.

The following definition generalizes the notion of a fine sheaf to numerable sites. Instead of using the original definition as a sheaf whose endomorphism sheaf is soft (which would require us to define soft sheaves on numerable sites), we use an equivalent (for paracompact spaces) characterization in terms of partitions of unity.

Definition 6.4. Suppose $\mathcal{U}: \mathbf{Site} \rightarrow \mathbf{Loc}$ is a numerable site (Definition 5.1). A sheaf

$$F: \mathbf{Site}^{\text{op}} \rightarrow \mathbf{Ab}$$

is *fine* if any covering family $\{U \rightarrow M, V \rightarrow M\}$ in \mathbf{Site} admits a *partition of unity*, defined as a pair of endomorphisms of F (as a sheaf of abelian groups):

$$p_U: F \rightarrow F, \quad p_V = \text{id} - p_U: F \rightarrow F$$

subordinate to the covering family $\{U \rightarrow M, V \rightarrow M\}$. The subordination condition means that there is an open inclusion $u' \subset \mathcal{U}(M)$ such that $\mathcal{U}(U) \cup u' = \mathcal{U}(M)$ and p_U vanishes when restricted along the cartesian lift $U' \rightarrow M$ of $u' \rightarrow \mathcal{U}(M)$. Likewise for p_V and V . Used in 6.5, 6.7, 6.7*, 7.3*.

Remark 6.5. Suppose $\mathcal{U}: \mathbf{Site} \rightarrow \mathbf{Loc}$ is a numerable site (Definition 5.1) and

$$F: \mathbf{Site}^{\text{op}} \rightarrow \mathbf{Ab}$$

is a sheaf that admits a module structure over the sheaf of rings $\mathbf{C} \circ \mathcal{U}$, where

$$\mathbf{C}: \mathbf{Loc}^{\text{op}} \rightarrow \mathbf{CRing}$$

sends a locale L to the commutative ring $\mathbf{C}(L, \mathbf{R})$ of real functions on L . Then F is a fine sheaf (Definition 6.4). Indeed, the endomorphisms p_U and p_V are given by multiplication by the elements of a partition of unity (Definition 3.7) subordinate to the open cover $\{\mathcal{U}(U), \mathcal{U}(V)\}$ of $\mathcal{U}(M)$. Analogous observations can be made using smooth functions, or other types of functions, provided that \mathcal{U} factors through smooth manifolds or other appropriate categories. Used in 6.7.

The following definition is due to Bengel–Schapira [1978]. See there for interesting examples of supple sheaves.

Definition 6.6. (Bengel–Schapira [1978].) Suppose $\mathcal{U}: \mathbf{Site} \rightarrow \mathbf{Loc}$ is a numerable site (Definition 5.1). A sheaf

$$F: \mathbf{Site}^{\text{op}} \rightarrow \mathbf{Ab}$$

is *supple* if for any $U \in \mathbf{Site}$, any $V_1, V_2 \rightarrow U$ whose images are open inclusions, and any $x \in F(U)$ such that $x|_{V_1 \cap V_2} = 0$, there are $y_1, y_2 \in F(U)$ such that $x = y_1 + y_2$, $y_1|_{V_1} = 0$, $y_2|_{V_2} = 0$. Used in 6.7, 6.7*.

We point out an important special case in which the additional condition of Remark 6.3 is guaranteed to hold. For topological spaces, this result is well known (see, for example, Bredon [1997, Theorems II.5.5, II.9.11, II.9.16]), the novelty here is the generalization to arbitrary numerable sites and a shorter proof.

Corollary 6.7. Suppose $\mathcal{U}: \mathbf{Site} \rightarrow \mathbf{Loc}$ is a numerable site (Definition 5.1). A sheaf $F: \mathbf{Site}^{\text{op}} \rightarrow \mathbf{Ab}$ is acyclic (i.e., $F[0]: \mathbf{Site}^{\text{op}} \rightarrow \mathbf{coCh}$ satisfies the homotopy descent property) whenever any of the following conditions hold.

- F is flabby (French: flasque), i.e., all restriction maps are surjections.
- F is supple (French: souple) in the sense of Definition 6.6.

- F is fine (French: fin) in the sense of Definition 6.4. (For example, F admits a module structure like in Remark 6.5.)

Used in 1.2*, 6.0*, 7.1*, 7.3*.

Proof. According to Remark 6.3, we have to show that the following property holds: for any $M \in \mathbf{Site}$ and any covering family $\{U \rightarrow M, V \rightarrow M\}$ of M , the map

$$F(U) \oplus F(V) \rightarrow F(U \cap V), \quad (u, v) \mapsto u|_{U \cap V} - v|_{U \cap V}$$

is degreewise surjective.

For a flabby sheaf this is true because the restriction map $F(U) \rightarrow F(U \cap V)$ is already surjective by definition.

For a supple sheaf, pick an arbitrary section

$$s \in F(U \cap V).$$

Pick a partition of unity $f_U, f_V: \mathbf{U}(M) \rightarrow [0, 1]$ subordinate to $\{\mathbf{U}(U), \mathbf{U}(V)\}$. Set $U' = f_U^*(2/3, 1]$ and $V' = f_V^*(2/3, 1]$. Denote by \hat{U}' and \hat{V}' the cartesian lifts of U' and V' (Remark 5.2). We have $U' \cap V' = \emptyset$ and $\{U', V', \mathbf{U}(U) \cap \mathbf{U}(V)\}$ is an open cover of $\mathbf{U}(M)$. By Definition 6.6, we can find $y_1, y_2 \in F(U \cap V)$ such that $s = y_1 + y_2$, $y_1|_{\hat{U}' \cap V} = 0$, $y_2|_{\hat{V}' \cap U} = 0$. Gluing y_1 and the zero section over \hat{U}' , we get $x_1 \in F(U)$. Likewise, gluing y_2 and the zero section over \hat{V}' , we get $x_2 \in F(V)$. By construction, $x_1|_{U \cap V} + x_2|_{U \cap V} = y_1 + y_2 = s$, as desired.

For a fine sheaf this is established using a standard argument with a partition of unity, whose existence is guaranteed by the definition of a fine sheaf. Throughout this entire proof, we only use those morphisms in \mathbf{Site} whose \mathbf{U} -images are open inclusions in \mathbf{Loc} . We adopt the convention that capital letters are used for objects of \mathbf{Site} , while small letters are used for their \mathbf{U} -images in \mathbf{Loc} . Thus, we write $u = \mathbf{U}(U)$. Sometimes, we construct an open inclusion $v \rightarrow w = \mathbf{U}(W)$, in which case its cartesian lift is denoted by $V \rightarrow W$. Also, $U \cap V \rightarrow M$ denotes the cartesian lift of $\mathbf{U}(U) \cap \mathbf{U}(V) \rightarrow \mathbf{U}(M)$. In particular, $\mathbf{U}(U \cap V) = \mathbf{U}(U) \cap \mathbf{U}(V)$.

Using Definition 6.4, choose a partition of unity

$$p_U: F \rightarrow F, \quad p_V = \text{id} - p_U: F \rightarrow F$$

subordinate to the covering family $\{U \rightarrow M, V \rightarrow M\}$. The subordination condition means that p_U and p_V are morphisms of sheaves of abelian groups and there is an open inclusion $u' \subset m$ such that $u \cup u' = m$ and p_U vanishes when restricted along the cartesian lift $U' \rightarrow M$; a similar condition is imposed on p_V and V .

Pick an arbitrary section

$$s \in F(U \cap V).$$

Denote by t the supremum of all opens $t \rightarrow m$ such that the restriction of s along the cartesian lift of $t \cap u \cap v \rightarrow m$ vanishes. Thus, the cartesian lift $T \rightarrow M$ can be seen as the complement of the support of s , and the restriction of s along $T \rightarrow M$ vanishes.

Now construct a section $q_U \in F(U)$ by gluing the zero section over $U \cap T$ and the section $p_V(s) \in F(U \cap V)$. These two sections are compatible because their restrictions to $U \cap T \cap V$ are both 0, by definition of T . Furthermore, $U \cap T \rightarrow U$ and $U \cap V \rightarrow U$ cover U because their \mathbf{U} -images satisfy $(u \cap t) \cup (u \cap v) = u \cap (t \cup v) = u \cap m = u$, since $t \cup v = m$ by definition of t and p_V (we can take $v' = t$ in the definition of p_V). Symmetrically, construct $q_V \in F(V)$ by gluing 0 over $V \cap T$ and $-p_U(s)$ over $U \cap V$.

The image of (q_U, q_V) under the map

$$F(U) \oplus F(V) \rightarrow F(U \cap V), \quad (u, v) \mapsto u|_{U \cap V} - v|_{U \cap V}$$

is s because q_U and q_V restrict to $p_V(s)$ respectively $-p_U(s)$ on $U \cap V$ by construction, and $p_V(s) - (-p_U(s)) = p_V(s) + p_U(s) = s$. ■

7 Example: classical cohomology theories on smooth manifolds

We illustrate the previous section by reproving the classical theorems on the equivalence of de Rham, singular, Alexander–Spanier, and sheaf cohomology on smooth manifolds. In the formalism of simplicial presheaves, the other classical cohomology theory, Čech cohomology, is equivalent to sheaf cohomology for trivial reasons: the Čech nerve of an open cover of a smooth manifold M is weakly equivalent to M in the local model structure on simplicial presheaves on the site of smooth manifolds, by definition.

The following theorem is due to Weil [1952.a], see also Bott–Tu [1982.a, Proposition 8.8]. Recall that $\Omega(M)$ denotes the de Rham complex of differential forms on a smooth manifold M .

Proposition 7.1. The presheaf of cochain complexes

$$\Omega: \mathbf{Man}^{\mathrm{op}} \rightarrow \mathbf{coCh}, \quad M \mapsto \Omega(M)$$

satisfies the homotopy descent condition on the site \mathbf{Man} . In particular, the integration map yields a natural weak equivalence

$$\Omega(M) \rightarrow \mathbf{C}^*(M, \mathbf{R})$$

where $\mathbf{C}^*(M, \mathbf{R})$ denotes the real smooth singular cochain complex of M .

Proof. Differential n -forms form a sheaf of abelian groups for any $n \geq 0$, which admits a module structure over smooth functions via multiplication, so the first result follows from Corollary 6.7 using smooth partitions of unity. For the second result, observe that both sides satisfy homotopy descent (see Proposition 7.2 for the right side) and are \mathbf{R} -invariant, i.e., send projection maps $M \times \mathbf{R} \rightarrow M$ to weak equivalences. Thus, it suffices to show the claim for the case $M = \mathbf{R}^0$, in which case both sides are $\mathbf{R}[0]$ and the map is identity. ■

Denote by $\mathbf{C}^*(M, A)$ the singular cochain complex with coefficients in an abelian group A on a smooth manifold M . Singular n -simplices can be taken to be either continuous or smooth maps $\Delta^n \rightarrow M$.

Proposition 7.2. For any abelian group A , the presheaf of cochain complexes

$$\mathbf{C}^*: \mathbf{Man}^{\mathrm{op}} \rightarrow \mathbf{coCh}, \quad M \mapsto \mathbf{C}^*(M, A)$$

satisfies the homotopy descent condition. In particular, singular cohomology is naturally isomorphic to sheaf cohomology. Used in 7.1*.

Proof. We invoke Theorem 6.1. The first property is satisfied because a singular cochain on a disjoint union is a collection of singular cochains on individual terms. The second property is satisfied because the presheaf $F = \mathbf{C}^*(-, A)$ is flabby: for any open embedding $W \rightarrow X$, the restriction map $\mathbf{C}^*(X, A) \rightarrow \mathbf{C}^*(W, A)$ is surjective because a singular cochain on W can be extended to a singular cochain on X by zeros.

The third property states that for any M and any cover $\{U, V\}$ of M the restriction map

$$r: F(M) \rightarrow \ker(F(U) \times F(V) \rightarrow F(U \cap V))$$

is a quasi-isomorphism. In our case, the kernel can be described as the cochain complex $\mathbf{C}^*(\{U, V\}, A)$ of singular cochains defined on singular simplices in X that factor through U or V . Recall (Eilenberg [1944.a, Chapter IV]) that the iterated barycentric subdivision construction on singular simplices defines a cochain map

$$\mathrm{Sd}: \mathbf{C}^*(\{U, V\}, A) \rightarrow \mathbf{C}^*(M, A)$$

such that $r \circ \mathrm{Sd} = \mathrm{id}$ and $\mathrm{id} - \mathrm{Sd} \circ r = h \circ d + d \circ h$ for a cochain homotopy h . Thus, the restriction map is a quasi-isomorphism, which completes the proof. ■

Denote by $\mathrm{AS}(M, A)$ the Alexander–Spanier cochain complex with coefficients in an abelian group A on a topological space M . In cochain degree n , these are given by germs of A -valued functions (not necessarily continuous) on U^{n+1} around the diagonal $U \rightarrow U^{n+1}$.

Proposition 7.3. The presheaf of cochain complexes

$$\mathbf{AS}: \mathbf{Man}^{\text{op}} \rightarrow \mathbf{coCh}, \quad M \mapsto \mathbf{AS}(M, A)$$

satisfies the homotopy descent condition. In particular, Alexander–Spanier cohomology is naturally isomorphic to sheaf cohomology.

Proof. The conditions of Definition 6.4 are satisfied: given an open cover $\{U, V\}$ of M , pick a partition of unity $f, g: M \rightarrow [0, 1]$ (so that $f + g = 1$) subordinate to $\{U, V\}$, set $u' = f^*[0, 1/3)$ and $v' = g^*[0, 1/3)$. Finally, define $p_U: \mathbf{AS}_n \rightarrow \mathbf{AS}_n$ by setting $p_U(s)$ to the germ of an A -valued function on M^{n+1} equal to s on $(f^*(1/2, 1))^{n+1}$ and 0 everywhere else. Likewise for p_V , using g^* instead of f^* . The pullback of p_U to $u' = f^*[0, 1/3)$ vanishes and $U \cup u' = M$. Likewise for p_V . By Corollary 6.7, we deduce that the sheaf \mathbf{AS}_n is acyclic for all n . Hence, \mathbf{AS} satisfies the homotopy descent condition. ■

8 Application: Classifying spaces for stacks over topological spaces

Definition 8.1. Suppose

$$F: \mathbf{Top}^{\text{op}} \rightarrow \mathbf{sSet}$$

is a simplicial presheaf on the site \mathbf{Top} . Define

$$\mathcal{C}F: \mathbf{Top}^{\text{op}} \rightarrow \mathbf{sSet}, \quad \mathcal{C}F(X) = \operatorname{hocolim}_{n \in \Delta^{\text{op}}} F(\Delta^n \times X),$$

where $\Delta^n \in \mathbf{Top}$ denotes the n -simplex as a topological space and the homotopy colimit is modeled by the diagonal of a bisimplicial set.

The importance of the presheaf $\mathcal{C}F$ lies in the fact that $\pi_0 \mathcal{C}F(X)$ is precisely the set of *concordance classes* of sections of F over X . Here two sections $a, b \in F(X)$ are *concordant* if there is a section $c \in F(\mathbf{R} \times X)$ such that $c|_{\{0\} \times X} \simeq a$ and $c|_{\{1\} \times X} \simeq b$, where \simeq means that the two given vertices of $F(X)$ are in the same connected component.

The following theorem can be seen as the analogue of Berwick–Evans–Boavida de Brito–Pavlov [2019, Theorem 1.2] for arbitrary topological spaces.

Theorem 8.2. Suppose

$$F: \mathbf{Top}^{\text{op}} \rightarrow \mathbf{sSet}$$

is a simplicial presheaf on the site \mathbf{Top} . If F satisfies the homotopy descent condition, then so does $\mathcal{C}F$. Used in 8.5*.

Proof. By Theorem 4.8, it suffices to show that $\mathcal{C}F$ satisfies the homotopy descent property for numerable open covers with two elements and for covers with disjoint elements. These two cases are proved in Berwick–Evans–Boavida de Brito–Pavlov [2019, Theorem 5.3 and Lemma 5.6]. Although the results are formulated for manifolds, the constructions in their proofs rely exclusively on the existence of partitions of unity, and continue to work unchanged in the setting of numerable open covers of topological spaces. ■

Definition 8.3. Suppose

$$F: \mathbf{Top}^{\text{op}} \rightarrow \mathbf{sSet}$$

is a simplicial presheaf on the site \mathbf{Top} . The natural map

$$\mathcal{C}F(X) \rightarrow \operatorname{Map}(\operatorname{Sing} X, \mathcal{C}F(*))$$

is defined as follows. Using the hom-product adjunction, it suffices to define a map

$$\mathcal{C}F(X) \times \operatorname{Sing} X \rightarrow \mathcal{C}F(*).$$

Unfolding the definition of \mathcal{C} as the diagonal of a bisimplicial set, it suffices to define a map of sets natural in $n \in \Delta$:

$$F(\Delta^n \times X)_n \times X^{\Delta^n} \rightarrow F(\Delta^n)_n.$$

Indeed, given a map $\Delta^n \rightarrow X$, we send the given element of $F(\Delta^n \times X)$ to an element of $F(\Delta^n \times \Delta^n)$ and then pull back along the diagonal map $\Delta^n \rightarrow \Delta^n \times \Delta^n$ to get an element of $F(\Delta^n)$. Used in 1.2, 8.5.

Remark 8.4. Below, we use the notion of a cofibrant topological space in the Serre–Quillen model structure on topological spaces. All CW-complexes, topological manifolds, and polyhedra are cofibrant topological spaces.

Theorem 8.5. Suppose

$$F: \mathbf{Top}^{\mathrm{op}} \rightarrow \mathbf{sSet}$$

is a simplicial presheaf on the site \mathbf{Top} . If F satisfies the homotopy descent condition, then $\mathcal{C}F$ is representable in the following sense: the natural map of Definition 8.3

$$\mathcal{C}F(X) \rightarrow \mathbf{Map}(\mathbf{Sing} X, \mathcal{C}F(*))$$

is a weak equivalence whenever X is homotopy equivalent to a cofibrant topological space. Furthermore, if F preserves weak equivalences, then the natural map

$$\mathcal{C}F(X) \rightarrow \mathbf{RMap}(\mathbf{Sing} X, \mathcal{C}F(*))$$

is a weak equivalence for an arbitrary topological space X . Used in 1.1*, 8.6, 8.7.

Proof. The statement about \mathbf{RMap} is an immediate consequence of the statement about \mathbf{Map} and the fact that every topological space X admits a weak equivalence to a cofibrant topological space.

By Theorem 8.2, the presheaf $\mathcal{C}F$ satisfies the homotopy descent condition. Furthermore, $\mathcal{C}F$ is homotopy invariant, as already observed by Morel–Voevodsky (see, for example, Berwick-Evans–Boavida de Brito–Pavlov [2019, Corollary 2.4]). The class of spaces X for which the natural map

$$\mathcal{C}F(X) \rightarrow \mathbf{Map}(X, \mathcal{C}F(*))$$

is a weak equivalence is closed under retracts and transfinite compositions because weak equivalences of simplicial sets are closed under retracts and transfinite compositions. It remains to show that for a map $X_{n-1} \rightarrow X_n$ that is a cobase change of the inclusion $S^{d-1} \rightarrow D^d$ for some $d \geq 0$, if X_{n-1} belongs to the class, then so does X_n . This follows from the descent property of $\mathcal{C}F$ with respect to the numerable open cover $\{U, V\}$ of X_n , where V is the interior of D^d inside X_n and U is the union of X_{n-1} and a small open band around S^{d-1} inside D^d . The spaces U , V , and $U \cap V$ are homotopy equivalent to X_{n-1} , D^d , and S^{d-1} , which implies the desired result. ■

Example 8.6. Suppose G is a topological group and consider the simplicial presheaf BG on the site \mathbf{Top} that sends a topological space X to the nerve of the groupoid of numerable principal G -bundles over X . By Theorem 8.5, concordance classes of numerable principal G -bundles over a cofibrant topological space X are classified by the space

$$\mathrm{hocolim}_{n \in \Delta^{\mathrm{op}}} BG(\Delta^n) \simeq B \mathrm{hocolim}_{n \in \Delta^{\mathrm{op}}} G(\Delta^n) \simeq B \mathbf{Sing}(G),$$

i.e., the classifying space of the topological group G . Taking π_0 on both sides, we recover the classification of numerable principal G -bundles up to concordance (in this case, isomorphism), for an arbitrary topological group G . Even better, Theorem 8.5 provides a statement on the level of spaces, not just π_0 : the space of maps $M \rightarrow B \mathbf{Sing}(G)$ is weakly equivalent to the space $(\mathcal{C}BG)(M)$ of numerable principal G -bundles on M up to concordance. Used in 1.2*.

Example 8.7. Suppose $d \geq 0$ and A is an abelian topological group. Consider the simplicial presheaf $B^d A$ on the site \mathbf{Top} that sends a topological space X to the simplicial set of numerable A -banded bundle $(d-1)$ -gerbes over X . The latter can be defined most easily as the fibrant replacement (in the Čech-local projective model structure on \mathbf{Top}) of the simplicial presheaf that sends X to $B^d C(X, A)$, where $C(X, A)$ denotes the discrete abelian group of continuous maps $X \rightarrow A$. By Theorem 8.5, concordance classes of such gerbes over a cofibrant topological space X are classified by the space

$$\mathrm{hocolim}_{n \in \Delta^{\mathrm{op}}} B^d A(\Delta^n) \simeq B^d \mathrm{hocolim}_{n \in \Delta^{\mathrm{op}}} A(\Delta^n) \simeq B^d \mathbf{Sing}(A),$$

i.e., the d -fold delooping of the topological group A . Taking π_0 on both sides, we recover the classification of numerable A -banded bundle $(d-1)$ -gerbes up to concordance (in this case, isomorphism), for an arbitrary abelian topological group A :

$$B^d A[M] \cong [M, B^d \text{Sing}(A)].$$

Even better, Theorem 8.5 provides a statement on the level of spaces, not just π_0 : the space of maps $M \rightarrow B^d \text{Sing}(A)$ is weakly equivalent to the space $(\mathcal{CB}^d A)(M)$ of numerable A -banded bundle $(d-1)$ -gerbes on M up to concordance. If A is a discrete abelian group, then $[M, B^d \text{Sing}(A)] \cong H^d(M, A)$ is the d th singular cohomology group of M with coefficients in A . This provides a geometric interpretation for singular cohomology groups of an arbitrary cofibrant topological space. Used in 1.2*.

9 References

- [1940] John W. Tukey. Convergence and Uniformity in Topology. *Annals of Mathematics Studies* 2 (1940). 3.0*, 3.8*, 3.12, 3.15.
- [1944.a] Samuel Eilenberg. Singular homology theory. *Annals of Mathematics (Second Series)* 45:3 (1944), 407–447. doi:10.2307/1969185. 7.2*.
- [1944.b] Jean Dieudonné. Une généralisation des espaces compacts. *Journal de Mathématiques Pures et Appliquées (Nouvelle Série)* 23 (1944), 65–76.
<https://gallica.bnf.fr/ark:/12148/bpt6k9704240z/f79.item>. 5.6.
- [1948] Arthur H. Stone. Paracompactness and product spaces. *Bulletin of the American Mathematical Society* 54:10 (1948), 977–983. doi:10.1090/s0002-9904-1948-09118-2. 3.9*, 3.10, 3.12, 5.6.
- [1952.a] André Weil. Sur les théorèmes de de Rham. *Commentarii Mathematici Helvetici* 26 (1952), 119–145. doi:10.1007/bf02564296, <https://eudml.org/doc/139040>. 7.0*.
- [1952.b] Hiroshi Miyazaki. The paracompactness of CW-complexes. *Tôhoku Mathematical Journal (Second Series)* 4:3 (1952), 309–313. doi:10.2748/tmj/1178245380. 3.0*.
- [1953] Ernest Michael. A note on paracompact spaces. *Proceedings of the American Mathematical Society* 4:5 (1953), 831–838. doi:10.1090/s0002-9939-1953-0056905-8. 3.0*, 3.9*, 3.10, 3.12, 3.14.
- [1955] Witold Hurewicz. On the concept of fiber space. *Proceedings of the National Academy of Sciences* 41:11 (1955), 956–961. doi:10.1073/pnas.41.11.956. 3.11.
- [1956] John Milnor. Construction of universal bundles, II. *Annals of Mathematics* 63:3 (1956), 430–436. doi:10.2307/1970012. 3.0*.
- [1962] Kiiti Morita. Paracompactness and product spaces. *Fundamenta Mathematicae* 50:3 (1962), 223–236. doi:10.4064/fm-50-3-223-236. 3.9*, 3.10.
- [1963] Albrecht Dold. Partitions of unity in the theory of fibrations. *Annals of Mathematics* 78:2 (1963), 223–255. doi:10.2307/1970341. 3.8*.
- [1964] Kiiti Morita. Products of normal spaces with metric spaces. *Mathematische Annalen* 154:4 (1964), 365–382. doi:10.1007/bf01362570. 3.10.
- [1966] Edwin H. Spanier. *Algebraic Topology*. McGraw–Hill, 1966. doi:10.1007/978-1-4684-9322-1. 3.11.
- [1968] John E. Derwent. A note on numerable covers. *Proceedings of the American Mathematical Society* 19:5 (1968), 1130–1130. doi:10.1090/s0002-9939-1968-0234461-x. 3.10.
- [1972.a] Albrecht Dold. *Lectures on Algebraic Topology*. Die Grundlehren der mathematischen Wissenschaften 200 (1972), Springer. Reprint: *Classics in Mathematics* (1995), Springer. doi:10.1007/978-3-662-00756-3. 3.7*.
- [1972.b] Werner Greub, Stephen Halperin, Ray Vanstone. *Connections, Curvature, and Cohomology*. Volume I. *De Rham Cohomology of Manifolds and Vector Bundles*. *Pure and Applied Mathematics* 47 (1972). doi:10.1016/S0079-8169(08)62849-4. 4.1*.
- [1973] Kenneth S. Brown and Stephen M. Gersten. Algebraic K-theory as generalized sheaf cohomology. *Lecture Notes in Mathematics* 341 (1973), 266–292. doi:10.1007/bfb0067062. 1.0*.
- [1974] Nicolas Bourbaki. *Topologie Générale*. Chapitres 5 à 10. *Actualités Scientifiques et Industrielles* 1235 (1974), Hermann. 5.5.

- [1975] J. Peter May. Classifying spaces and fibrations. *Memoirs of the American Mathematical Society* 1:155 (1975). doi:10.1090/memo/0155. 3.11.
- [1978] Gunter Bengel, Pierre Schapira. Décomposition microlocale analytique des distributions. *Journées Équations aux dérivées partielles* (1978), 1-2 doi:10.5802/jedp.176. 6.5*, 6.6.
- [1982.a] Raoul Bott, Loring W. Tu. *Differential Forms in Algebraic Topology*. Graduate Texts in Mathematics 82 (1982), Springer. doi:10.1007/978-1-4757-3951-0. 7.0*.
- [1982.b] Sibe Mardešić, Jack Segal. *Shape Theory*. North-Holland Mathematical Library 26 (1982). 3.10.
- [1989] Takao Hoshina. Extensions of mappings II. (Topics in General Topology.) North-Holland Mathematical Library 41 (1989), 41–80. 3.10, 3.14.
- [1997] Glen E. Bredon. *Sheaf Theory*. Second Edition. Graduate Texts in Mathematics 170 (1997), Springer. doi:10.1007/978-1-4612-0647-7. 6.6*.
- [1999] Fabien Morel and Vladimir Voevodsky. \mathbf{A}^1 -homotopy theory of schemes. *Publications mathématiques de l’I.H.É.S.* 90:1 (1999), 45–143. doi:10.1007/bf02698831. 1.0*.
- [2002.a] Peter T. Johnstone. Sketches of an Elephant. A Topos Theory Compendium. II. *Oxford Logic Guides* 44 (2002). 2.1, 2.3, 2.6, 2.7.
- [2002.b] Bertrand Toën, Gabriele Vezzosi. Segal topoi and stacks over Segal categories. arXiv:math/0212330v2. 2.0*.
- [2008.a] Dale Husemöller, Michael Joachim, Branislav Jurčo, Martin Schottenloher. *Basic Bundle Theory and K -Cohomology Invariants*. Lecture Notes in Physics 726 (2008), Springer. doi:10.1007/978-3-540-74956-1, https://www.mathematik.uni-muenchen.de/~schotten/Texte/978-3-540-74955-4_Book_LNP726.pdf. 3.13*.
- [2008.b] Hisham Sati, Urs Schreiber, Jim Stasheff. L_∞ -algebra connections and applications to String- and Chern-Simons n -transport. (Quantum Field Theory. Competitive Models.) Birkhäuser (2009), 303–424. arXiv:0801.3480v2, doi:10.1007/978-3-7643-8736-5_17. 1.0*.
- [2008.c] Vladimir Voevodsky. Homotopy theory of simplicial sheaves in completely decomposable topologies. *Journal of Pure and Applied Algebra* 214:8 (2010), 1384–1398. arXiv:0805.4578v1, doi:10.1016/j.jpaa.2009.11.004. 1.0*.
- [2010] Domenico Fiorenza, Urs Schreiber, Jim Stasheff. Čech cocycles for differential characteristic classes: an ∞ -Lie theoretic construction. *Advances in Theoretical and Mathematical Physics* 16:1 (2012), 149–250. arXiv:1011.4735v2, doi:10.4310/atmp.2012.v16.n1.a5. 1.0*.
- [2012] Jorge Picado, Aleš Pultr. *Frames and locales*. *Frontiers in Mathematics* (2012), Birkhäuser. 3.16*.
- [2017.a] Jacob Lurie. *Higher Topos Theory*. April 9, 2017. <https://www.math.ias.edu/~lurie/papers/HTT.pdf>. 2.0*.
- [2017.b] Dmitri Pavlov. Question 261195 (revision 1) on MathOverflow. February 2, 2017. <https://mathoverflow.net/revisions/261195/1>. 1.1*, 1.4*.
- [2019] Daniel Berwick-Evans, Pedro Boavida de Brito, Dmitri Pavlov. Classifying spaces of infinity-sheaves. arXiv:1912.10544v1. 1.1*, 1.4*, 8.1*, 8.2*, 8.5*.
- [2021] Araminta Amabel, Arun Debray, Peter J. Haine. *Differential Cohomology: Categories, Characteristic Classes, and Connections* arXiv:2109.12250v1. 1.1*.