

Orthogonal Stability ^{*}

Gabriele Nebe and Richard Parker

ABSTRACT. An ordinary character or a Brauer character χ of a finite group is called orthogonally stable, if all its indicator + constituents have even degree and there is a representation affording χ that fixes a non-degenerate quadratic form. In this case all non-degenerate quadratic forms in any representation affording χ have the same discriminant. This square class $\text{disc}(\chi) \in K^\times / (K^\times)^2$ of the character field K of χ is called the orthogonal discriminant of χ . The paper develops some character theoretic methods to compute orthogonal discriminants.

MSC: 20C15; 20C20; 11E12; 11E57.

KEYWORDS: orthogonal characters of finite groups; decomposition matrices; blocks with cyclic defect groups.

1 Introduction

Representation theory of finite groups studies linear actions of a finite group G on finite dimensional vector spaces V over a field K . Given such a group homomorphism $\rho : G \rightarrow \text{GL}(V)$ one may ask whether its image is contained in some other classical group, like the symplectic, a unitary or an orthogonal group and how to determine the isomorphism type of this classical group. In an ongoing project with Thomas Breuer and Jon Thackray we aim to answer this question for the absolutely irreducible representations ρ of all but the largest few groups G in the ATLAS of Brauer characters [6] over all finite fields K . The character tables in [6] and [2] yield enough information to determine the classical group with the only exception of the even degree orthogonal groups, where there are two possibilities, $O_{2m}^+(K)$ and $O_{2m}^-(K)$. The corresponding invariant quadratic forms are distinguished by their orthogonal discriminants $O+$ and $O-$ (see Definition 2.3). For the finitely many primes that divide the group order the determination of the orthogonal discriminants of the even degree absolutely irreducible indicator + Brauer characters is mostly a computational problem. With the meat-axe [13] we explicitly construct the corresponding representations and determine the orthogonal discriminants. Thanks to the main result of [11] the infinitely many other primes can be treated simultaneously by considering absolutely irreducible ordinary characters (see Section 5.2).

^{*}This is a contribution to Project-ID 286237555 – TRR 195 – by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation).

The central notion of the present paper is **orthogonal stability**. An ordinary character or a Brauer character χ of a finite group is called orthogonally stable if all its indicator + constituents have even degree and there is a representation affording χ that fixes a non-degenerate quadratic form. In this case all non-degenerate quadratic forms in any representation affording χ have the same discriminant, called the **orthogonal discriminant** $\text{disc}(\chi)$ of χ .

To compute $\text{disc}(\chi)$ it suffices to determine the orthogonal discriminants of all its indicator + constituents as shown in Theorem 4.13.

Assume that \mathcal{X} is an orthogonally stable ordinary character with character field K . Then K is a real number field and $\text{disc}(\mathcal{X})$ is a square class of the multiplicative group K^\times . We put

$$\Delta(\mathcal{X}) := K(\sqrt{\text{disc}(\mathcal{X})})/K, \text{ the discriminant field extension of } \mathcal{X}.$$

Theorem 5.5 shows that a prime ideal \wp of K is unramified in $\Delta(\mathcal{X})$ if the reduction of \mathcal{X} modulo \wp is orthogonally stable. For absolutely irreducible characters \mathcal{X} of p -defect one also the converse is true (see Theorem 5.10). This observation is enough to compute the orthogonal discriminant of all ordinary characters for the group J_1 from its decomposition matrices that are available in GAP [4]. Example 5.11 and Theorem 5.12 however show that the converse of Theorem 5.5 is not true in general.

For most groups, arguments based on cyclic defect are not enough to deduce all orthogonal discriminants. As the construction of the ordinary representations is computationally much harder than the one of the modular representations, Section 5.3 gives an algorithm to compute $\text{disc}(\mathcal{X})$ for absolutely irreducible indicator + ordinary characters \mathcal{X} of even degree by constructing enough modular representations affording the reduction of \mathcal{X} as a Brauer character and computing their orthogonal discriminants. We illustrate these methods computing the orthogonal discriminants of the irreducible ordinary characters of the Held group He from its p -modular orthogonal discriminants and the decomposition matrices for the primes p dividing $|He|$.

2 Quadratic forms

Let K be a field and V a vector space over K . A quadratic form Q is a map $Q : V \rightarrow K$ such that $Q(ax) = a^2Q(x)$ for all $a \in K$, $x \in V$ and such that its **polarisation**

$$B : V \times V \rightarrow K, B(x, y) := Q(x + y) - Q(x) - Q(y) \tag{1}$$

is a bilinear form. The determinant of Q is the determinant of a Gram matrix of its polarisation. This is well defined modulo squares. The quadratic form Q is called **non-degenerate**, if its polarisation is non-degenerate, i.e. $\det(Q) \neq 0$. We then also call (V, Q) a non-degenerate quadratic space.

Over number fields K two non-degenerate quadratic forms are isometric, if and only if they have the same dimension, n , Clifford invariant and **discriminant**, where the discriminant is

$$\text{disc}(Q) = (-1)^{\binom{n}{2}} \det(Q)(K^\times)^2.$$

For finite fields K the orthogonal discriminant (see Definition 2.3) distinguishes the two non-degenerate quadratic forms of given even dimension. Note that for even dimensional quadratic spaces, both, determinant and discriminant are multiplicative with respect to orthogonal direct sums:

Remark 2.1. *If Q and Q' are two quadratic forms of even dimension then*

$$\text{disc}(Q \perp Q') = \text{disc}(Q) \text{disc}(Q').$$

2.1 Restriction of scalars

Let (V, Q) be an n -dimensional quadratic space over the field K and let $F \leq K$ be a subfield of finite index, d . Then restriction of scalars turns V into an nd -dimensional space over F . If T denotes the trace of K/F then $T \circ Q : V \rightarrow F$ is a quadratic form that is non-degenerate if Q is non-degenerate and K/F is separable (the latter condition is always fulfilled if K is finite or of characteristic 0). Then

$$\det(T \circ Q) = N_{K/F}(\det(Q)) \text{disc}(K/F)^n$$

where $\text{disc}(K/F)$ is the field discriminant of K over F (see also [8, Lemma 2.2]).

Remark 2.2. *If K/F is separable and n is even, then $\text{disc}(T \circ Q) = N_{K/F}(\text{disc}(Q))$.*

2.2 Hyperbolic forms

For a finite dimensional K -vector space V we put

$$V^\vee := \text{Hom}_K(V, K) = \{f : V \rightarrow K \mid f \text{ is } K\text{-linear}\}$$

to denote the dual space of V . Any non-degenerate bilinear form B on V yields an isomorphism

$$\tilde{B} : V \rightarrow V^\vee, v \mapsto (x \mapsto B(v, x)). \quad (2)$$

The hyperbolic module $\mathbb{H}(V)$ is the quadratic K -space

$$(V \oplus V^\vee, Q) \text{ with } Q((v, f)) := f(v) \text{ for all } v \in V, f \in V^\vee.$$

Then $\dim(\mathbb{H}(V)) = 2 \dim(V)$ is even and $\text{disc}(\mathbb{H}(V)) = 1$.

2.3 Quadratic forms over finite fields

Let \mathbb{F}_q denote the field with q elements. It is well known that there are two non-degenerate quadratic forms of dimension 2 over \mathbb{F}_q , the hyperbolic plane $\mathbb{H}(\mathbb{F}_q)$ and the norm form $N(\mathbb{F}_q)$

of the quadratic extension of \mathbb{F}_q . The underlying space for $N(\mathbb{F}_q)$ is \mathbb{F}_{q^2} and the quadratic form is

$$Q(x) := x\bar{x} = xx^q = x^{q+1}.$$

On an \mathbb{F}_q -vector space of even dimension $2m$ there are two non-isometric non-degenerate quadratic forms. These two forms can be distinguished by many invariants such as discriminants resp. Arf invariants, Witt index, the number of isotropic vectors, and also the order of the orthogonal group (see for instance [7, Chapter IV]). To fix notation we denote these two forms by

$$Q_{2m}^+(q) := \mathbb{H}(\mathbb{F}_q)^m \text{ and } Q_{2m}^-(q) := \mathbb{H}(\mathbb{F}_q)^{m-1} \perp N(\mathbb{F}_q).$$

Definition 2.3. *The orthogonal discriminant of a $2m$ -dimensional non-degenerate quadratic space Q_{2m} over a finite field \mathbb{F}_q is $O+$ if $Q_{2m} \cong Q_{2m}^+(q)$ and $O-$ if $Q_{2m} \cong Q_{2m}^-(q)$. We denote this value $O+$ or $O-$ also by $\text{disc}(Q_{2m})$.*

We obtain the following analogue of Remark 2.1 on the multiplicativity of orthogonal discriminants:

Remark 2.4. *For $m, n \in \mathbb{N}$ we have $Q_{2m}^+(q) \perp Q_{2n}^+(q) \cong Q_{2(n+m)}^+(q)$, $Q_{2m}^+(q) \perp Q_{2n}^-(q) \cong Q_{2(n+m)}^-(q)$, and $Q_{2m}^-(q) \perp Q_{2n}^-(q) \cong Q_{2(n+m)}^-(q)$.*

Note that restriction of scalars does not change the orthogonal discriminant (see [17, Section 4]) where the key point is the following observation.

Proposition 2.5. *Let $T : \mathbb{F}_{q^2} \rightarrow \mathbb{F}_q$ denote the trace, $T(x) = x + x^q$. For $\epsilon = \pm$ we have $T \circ Q_{2m}^\epsilon(q^2) = Q_{4m}^\epsilon(q)$.*

Proof. It is enough to prove that $T(\mathbb{H}(\mathbb{F}_{q^2})) = \mathbb{H}(\mathbb{F}_q) \perp \mathbb{H}(\mathbb{F}_q)$ and $T(N(\mathbb{F}_{q^2})) = \mathbb{H}(\mathbb{F}_q) \perp N(\mathbb{F}_q)$. To see the first fact note that $\mathbb{H}(\mathbb{F}_{q^2})$ has a one dimensional isotropic \mathbb{F}_{q^2} -subspace. Restriction of scalars hence yields a 2-dimensional isotropic subspace in $T(\mathbb{H}(\mathbb{F}_{q^2}))$, so this form is hyperbolic. For the second form we count the number of non-zero isotropic vectors. So let $0 \neq y \in \mathbb{F}_{q^4}$ such that $T(N(y)) = 0$. Then

$$T(N(y)) = T(y^{q^2+1}) = y^{q^2+1} + (y^{q^2+1})^q = 0$$

and hence $y^{(q-1)(q^2+1)} = -1$. As the multiplicative group of \mathbb{F}_{q^4} is cyclic of order $(q+1)(q-1)(q^2+1)$, there are exactly $(q-1)(q^2+1)$ such elements $y \in \mathbb{F}_{q^4}$. By [7, Section IV] we hence get $T(N(\mathbb{F}_{q^2})) = Q_4^-(q)$. \square

So to obtain the analogous result as in Remark 2.2 for orthogonal discriminants $\in \{O+, O-\}$, the norm should be replaced by the identity.

Remark 2.6. *Any n -dimensional quadratic space (V, Q) over \mathbb{F}_q extends to an n -dimensional quadratic space $(V \otimes \mathbb{F}_{q^d}, Q)$ over \mathbb{F}_{q^d} by putting $Q(x \otimes a) := a^2Q(x)$ for $x \in V$, $a \in \mathbb{F}_{q^d}$. If d is even, the discriminant of Q becomes a square in \mathbb{F}_{q^d} and also any quadratic polynomial $X^2 + X + a \in \mathbb{F}_q[X]$ is reducible in $\mathbb{F}_{q^d}[X]$. Hence for even degree extensions this quadratic space over \mathbb{F}_{q^d} always has orthogonal discriminant $O+$. The orthogonal discriminant is, however, stable under odd degree field extensions.*

(a) If d is even then $Q_{2m}^+(q) \otimes \mathbb{F}_{q^d} \cong Q_{2m}^-(q) \otimes \mathbb{F}_{q^d} \cong Q_{2m}^+(q^d)$.

(b) If d is odd then $Q_{2m}^+(q) \otimes \mathbb{F}_{q^d} \cong Q_{2m}^+(q^d)$ and $Q_{2m}^-(q) \otimes \mathbb{F}_{q^d} \cong Q_{2m}^-(q^d)$.

2.4 Quadratic forms over discrete valuation rings

In this section we recall some results on regular quadratic forms over local rings for the special situation needed in this paper. So let R be a complete discrete valuation ring with uniformizer π and finite residue field $F := R/\pi R \cong \mathbb{F}_q$. We also assume that the field of fractions K of R has characteristic 0. A quadratic lattice (M, Q) is a free R -module M of finite rank together with an R -valued quadratic form $Q : M \rightarrow R$. The quadratic form is called **regular** if the map \tilde{B} from equation (2) restricts to an R -isomorphism between M and its dual module $M^\vee = \text{Hom}_R(M, R)$. This is equivalent to saying that the determinant of Q is a unit in R .

As shown in [7, Satz 15.6] there is a bijection between isometry classes of regular quadratic lattices of rank n over R and non-degenerate quadratic forms of dimension n over the residue field F . In particular in our situation there are exactly two isometry classes of regular quadratic R -lattices (M, Q) of even rank $2m$, corresponding to the two possible residue forms $(M/\pi M, \overline{Q}) \cong Q_{2m}^+(q)$ respectively $Q_{2m}^-(q)$. We denote these regular quadratic lattices by $Q_{2m}^+(R)$ respectively $Q_{2m}^-(R)$. Note that the norm form of the ring of integers in the unique unramified extension of degree 2 of K has as residue form $Q_2^-(q) = N(\mathbb{F}_q)$, so we get the following remark.

Remark 2.7. *The discriminant of $Q_{2m}^+(R)$ is 1. If $\delta \in \text{disc}(Q_{2m}^-(R))$ then $K[\sqrt{\delta}]$ is the unique unramified extension of degree 2 of K .*

The most important property of regular quadratic sublattices is that they split as orthogonal direct summands.

Lemma 2.8. *(see [7, Satz 1.6]) Let (M, Q) be a quadratic R -lattice and $N \leq M$ be some R -submodule such that $(N, Q|_N)$ is regular. Then*

$$(M, Q) = (N, Q|_N) \perp (N^\perp, Q|_{N^\perp}).$$

2.5 Hermitian forms

In this section we assume that we are given a Galois extension L/K of degree $[L : K] = 2$. Let $\bar{} \in \text{Gal}_K(L)$ denote the non-trivial Galois automorphism. A Hermitian form H on a non-zero L -vector space V is a map $H : V \times V \rightarrow L$ that is L -linear in the first argument and such that $H(y, x) = \overline{H(x, y)}$ for all $x, y \in V$. If $n := \dim_L(V)$ then restriction of scalars turns V into a K -vector space of dimension $2n$ and H defines a quadratic form $Q_H : V \rightarrow K$, $Q_H(x) := H(x, x)$ for all $x \in V$.

Proposition 2.9. *(see [16, page 350]) Let (V, H) be a non-degenerate Hermitian L -vector space.*

(a) Let $\text{char}(K) \neq 2$ and write $L = K[\sqrt{\delta}]$. Then $\text{disc}(Q_H) = \delta^n (K^\times)^2$.

(b) If $\text{char}(K) = 2$ and K is a finite field, then $Q_H \cong Q_{2n}^+$ if n is even and $Q_H \cong Q_{2n}^-$ if n is odd.

Note that [16] also gives the Clifford invariant of Q_H as well as the Arf invariant for general fields of characteristic 2.

3 Orthogonal representations

Let G be a finite group and K be a field. Any KG -module $V \neq \{0\}$ defines a group homomorphism $\rho : G \rightarrow \text{GL}(V)$. Then ρ is called a K -representation of G . Equivalence of representations is defined as isomorphism of KG -modules. The representation ρ is called **irreducible** if V is a simple KG -module, i.e. $\{0\}$ and V are the only G -invariant submodules of V . The **trivial representation** is the map $\rho : G \rightarrow \text{GL}(K)$, $\rho(g) = \text{id}_K$ for all $g \in G$.

For any KG -module V also the dual space V^\vee is a KG -module. The corresponding representation ρ^\vee is called the dual representation of ρ . The representation ρ is called **self-dual**, if ρ^\vee is equivalent to ρ .

Given a representation ρ or equivalently a KG -module V we put

$$\mathcal{Q}(V) = \mathcal{Q}(\rho) := \{Q : V \rightarrow K \mid Q \text{ is a quadratic form, } Q(x\rho(g)) = Q(x) \text{ for all } x \in V, g \in G\}$$

the space of G -invariant quadratic forms. We call ρ an **orthogonal representation**, if $\mathcal{Q}(\rho)$ contains a non-degenerate quadratic form Q . Then (V, Q) is also called an **orthogonal KG -module**. As the polarisation of Q defines a KG -isomorphism between V and its dual module V^\vee orthogonal representations are self-dual.

3.1 The anisotropic kernel

If the characteristic of K divides the group order, then not all representations are direct sums of irreducible ones. However, the theory of orthogonal representations is essentially a semi-simple one.

Lemma 3.1. *Let $\rho : G \rightarrow \text{GL}(V)$ be an irreducible representation and $Q \in \mathcal{Q}(\rho) \setminus \{0\}$ be a non-zero G -invariant quadratic form on V . Then either Q is non-degenerate or $\text{char}(K) = 2$, $Q : V \rightarrow K$ is an \mathbb{F}_2G -module homomorphism and ρ is the trivial representation.*

Proof. Assume that Q is degenerate. Since its polarisation B is G -invariant the radical

$$\text{rad}(B) = V^\perp = \{v \in V \mid B(v, w) = 0 \text{ for all } w \in V\}$$

is a KG -submodule of V . As V is simple and $\text{rad}(B) \neq 0$, we hence have $\text{rad}(B) = V$ and hence $B = 0$. Now $B(x, x) = 2Q(x)$ for all $x \in V$ so $Q \neq 0 = B$ implies that $\text{char}(K) = 2$.

Moreover the equation (1) shows that Q is a group homomorphism between the additive group of V and K . The G -invariance of Q implies that the kernel of this homomorphism is a submodule of the simple KG -module V , so Q is injective. Moreover for $x \in V$ and $g \in G$ we have $Q(x + xg) = Q(x) + Q(xg) = 2Q(x) = 0$, so $x + xg \in \ker(Q) = \{0\}$ and hence $x = xg$ for all $x \in V, g \in G$. This implies that V is the trivial KG -module. \square

Definition 3.2. *A submodule U of the orthogonal KG -module (V, Q) is called isotropic if $Q(U) = \{0\}$. An orthogonal KG -module (V, Q) is called anisotropic if it does not contain a non-zero isotropic submodule.*

Let $\rho : G \rightarrow \mathrm{GL}(V)$ be an orthogonal representation and $Q \in \mathcal{Q}(\rho)$ be a non-degenerate quadratic form. Let U be a maximal isotropic KG -submodule of V . Then Q defines a quadratic form \overline{Q} on U^\perp/U by $\overline{Q}(x + U) := Q(x)$ for all $x \in U^\perp$. The quadratic KG -module $(W, Q_0) := (U^\perp/U, \overline{Q})$ is uniquely determined by ρ and Q up to KG -isometry and called the anisotropic kernel of ρ (see for instance [3, Lemma 4.2 (8)]).

Theorem 3.3. *Assume that either $\mathrm{char}(K) \neq 2$ or the trivial representation is not a constituent of ρ . Then the anisotropic kernel (W, Q_0) of (V, Q) is the orthogonal sum of simple orthogonal KG -modules. Moreover $\mathrm{disc}(Q_0) = \mathrm{disc}(Q)$.*

Proof. If no simple submodule of V is isotropic, then put $W := V, Q_0 := Q$. Otherwise there is a simple isotropic submodule $S \leq V$. Then Q induces a quadratic form on S^\perp/S . Replacing V by this smaller dimensional module we finally arrive at a module (W, Q_0) for which the restriction of Q_0 to any simple submodule is non-zero. As we assumed that the trivial module is not a constituent of V (and hence of W) when the characteristic of K is 2, Lemma 3.1 shows that the restriction of Q_0 to any simple submodule S of W is non-degenerate. Hence S splits as a direct orthogonal summand $W = S \perp S^\perp$. So W is the orthogonal direct sum of simple quadratic KG -modules. \square

Remark 3.4. *The note [12] gives an explicit classification of all anisotropic orthogonal KG -modules for finite fields of characteristic 2. These have at most one indecomposable orthogonal direct summand R that is not simple; the socle and the top of R both are trivial KG -modules.*

4 Characters

We use capital letters \mathcal{X} only for ordinary characters, whereas χ may stand for both, an ordinary or a Brauer character.

4.1 Brauer characters

In characteristic 0, the character of a K -representation $\rho : G \rightarrow \mathrm{GL}(V)$ is the map $\mathcal{X}_\rho : G \rightarrow K, g \mapsto \mathrm{trace}(\rho(g))$ where the trace is the usual trace of an endomorphism. One important property here is that two representations are equivalent if and only if they have the same

character.

In positive characteristic, p , such a definition of character would waste information, for instance the character of the sum of p isomorphic modules would be identically 0. The definition of Brauer character can for instance be found in [9, Chapter 2] and [5, Chapter 15]. Given a field K of characteristic $p > 0$ and a K -representation ρ the Brauer character χ_ρ of ρ is a map from the p' -conjugacy classes of G to the field of complex numbers. For $g \in G$ of order n not divisible by p the eigenvalues of $\rho(g)$ (in an algebraic closure \hat{K} of K) are n -th roots of unity. Conway polynomials provide one canonical way to identify these roots of unity in \hat{K} with complex roots of unity (see for instance [6, Introduction]). Then the Brauer character of ρ assigns to g the sum of these complex roots of unity.

The Brauer character of ρ does in general not determine the representation ρ up to equivalence. However, it determines the multiset of composition factors of ρ up to isomorphism.

Definition 4.1. *Let \mathcal{X} be an ordinary character of G . The character field $K(\mathcal{X})$ is the abelian number field $\mathbb{Q}(\mathcal{X}(g) : g \in G)$ generated over the rationals by all character values of \mathcal{X} .*

For a p -Brauer character χ of G the character field $K(\chi)$ is obtained by reducing the ring of integers of the number field L generated by the Brauer character values of χ modulo any prime ideal that divides p . As L is Galois over \mathbb{Q} , the finite field $K(\chi)$ does not depend on the choice of the prime ideal.

Any ordinary character and any Brauer character is a unique sum of absolutely irreducible characters, its absolutely irreducible constituents.

4.2 Reduction of characters

Let K be a number field, O_K its ring of integers, \wp a non-zero prime ideal of O_K with residue field $F := O_K/\wp$. For a K -representation $\rho : G \rightarrow \mathrm{GL}(V)$ the reduction mod \wp is obtained by choosing a G -invariant O_K -lattice M in V , i.e. a finitely generated O_K -submodule M in V that contains a K -basis of V and such that $M\rho(g) = M$ for all $g \in G$. Then $\overline{V} := M/\wp M$ is a FG -module. The composition factors of \overline{V} are independent of the choice of the $O_K G$ -lattice M in V (see for instance [5, Theorem 15.6]). The corresponding representation $\overline{\rho} : G \rightarrow \mathrm{GL}(\overline{V})$ is called a reduction of ρ modulo \wp . Whereas the representation $\overline{\rho}$ depends on the choice of the lattice M , its Brauer character only depends on ρ and on \wp .

Remark 4.2. *Note that the character field of the Brauer character of $\overline{\rho}$ can be strictly contained in F . The smallest example in [2] are the two representations ρ of degree 4 of the group $2.S_5$, where the character field of ρ is $K = \mathbb{Q}(\sqrt{-3})$. The irrationality only occurs as the character value of the outer elements of order 6, thus the reduction of ρ modulo 2 has a rational Brauer character of degree 4.*

Given an ordinary character \mathcal{X} with character field K , the reduction of \mathcal{X} modulo \wp is the Brauer character of a reduction of a representation ρ (over a suitable field extension) affording \mathcal{X} . We can compute the reduction of \mathcal{X} modulo \wp directly from the character table:

Remark 4.3. For any finite abelian extension K of \mathbb{Q} the Conway polynomials from Section 4.1 identify a unique prime ideal \wp_0 of O_K that contains the prime p so that the reduction of \mathcal{X} modulo \wp_0 is the restriction of \mathcal{X} to the p' -classes of G .

For any prime ideal \wp of O_K that divides p , there is $\sigma \in \text{Gal}(K/\mathbb{Q})$ such that $\wp = \sigma^{-1}(\wp_0)$. Then the reduction of \mathcal{X} modulo \wp is obtained as the reduction of $\sigma(\mathcal{X})$ modulo \wp_0 .

4.3 Orthogonally simple characters

In the following the notion ‘‘character’’ denotes either an ordinary character or a Brauer character of the finite group G .

Definition 4.4. A character χ is called *orthogonal*, if there is an orthogonal representation with character χ . An orthogonal character χ is called *orthogonally simple*, if χ is not a proper sum of orthogonal characters.

The next definition recalls the Frobenius-Schur indicator:

Definition 4.5. Let χ be an absolutely irreducible character or Brauer character.

- (+) If χ is orthogonal then the indicator of χ is $+$.
- (+) In characteristic 2 also the trivial character is of indicator $+$.
- (o) If χ is not self-dual then its indicator is o .
- (-) In all other cases the indicator of χ is $-$.

In particular the absolutely irreducible characters of indicator $+$ are exactly those that are afforded by some representation ρ (over some extension field of the character field) for which $\mathcal{Q}(\rho) \neq \{0\}$.

Proposition 4.6. Let χ be an orthogonally simple character. Then one of the following holds:

- (a) χ is absolutely irreducible of indicator $+$.
- (b) $\chi = \psi + \psi^\vee$ for some absolutely irreducible character ψ of indicator o .
- (c) $\chi = 2\psi$ for some absolutely irreducible character ψ of indicator $-$.

Moreover for any orthogonal representation ρ affording the character χ the dimension of $\mathcal{Q}(\rho)$ is one.

Proof. Let ψ be an absolutely irreducible constituent of χ . As χ is self-dual, also ψ^\vee is a constituent of χ . If $\psi \neq \psi^\vee$ then $\psi + \psi^\vee$ is an orthogonal sub-character of χ and hence $\chi = \psi + \psi^\vee$. If $\psi = \psi^\vee$ and ψ has indicator $+$ then already ψ is orthogonal and hence $\chi = \psi$. If ψ is self-dual but not orthogonal, then all orthogonal characters containing ψ also contain 2ψ . As $2\psi = \psi + \psi^\vee$ is orthogonal we are in situation (c). \square

Corollary 4.7. *The only orthogonally simple characters of odd degree are the absolutely irreducible indicator + odd degree characters that are not the trivial Brauer character in characteristic 2.*

Definition 4.8. *Given an orthogonal ordinary character \mathcal{X} we say that “the orthogonal discriminant of \mathcal{X} is well defined” if there is a unique square class $\text{disc}(\mathcal{X}) \in K(\mathcal{X})^\times / (K(\mathcal{X})^\times)^2$ with the following property: Given an orthogonal representation ρ over some field L and with character \mathcal{X} and $Q \in \mathcal{Q}(\rho)$ a non-degenerate quadratic form. Then $\text{disc}(Q) = \text{disc}(\mathcal{X})(L^\times)^2$. The square class $\text{disc}(\mathcal{X}) \in K(\mathcal{X})^\times / (K(\mathcal{X})^\times)^2$ is then called the **orthogonal discriminant** of \mathcal{X} . For an orthogonal Brauer character χ “the orthogonal discriminant of χ is well defined” if all non-degenerate invariant quadratic forms in any orthogonal representation over the character field of χ have the same orthogonal discriminant $\in \{O+, O-\}$ which is then also called the **orthogonal discriminant** of χ .*

Theorem 4.9. *The orthogonal discriminant of an orthogonally simple character of even degree is well defined.*

Proof. We go through the three cases from Proposition 4.6:

- (a) Here χ is absolutely irreducible of indicator + and even degree. For number fields K the short note [11] shows that the orthogonal discriminant of χ is well defined. For finite fields K , there is an orthogonal representation ρ over K with character χ . As χ is absolutely irreducible the dimension of the K -space $\mathcal{Q}(\rho)$ is 1, so $\mathcal{Q}(\rho) = \{aQ \mid a \in K\}$ for some non-degenerate Q and $\text{disc}(\chi) = \text{disc}(Q)$.
- (b) If $K(\psi) = K(\chi)$, then Q is hyperbolic and hence its orthogonal discriminant is $O+$. Otherwise Q comes from a Hermitian form and $\text{disc}(\chi)$ can be read off from Proposition 2.9.
- (c) For number fields K [18, Theorem B] asserts that $\text{disc}(\chi) = 1$ in this case. If K is a finite field, then ψ is the character of a representation over K . If $\rho : G \rightarrow \text{GL}(V)$ is a representation affording the character 2ψ and $Q \in \mathcal{Q}(\rho)$ non-degenerate, then the restriction of the quadratic form Q to any simple submodule of V is identically zero. In particular the form Q is hyperbolic and hence of orthogonal discriminant $O+$.

□

Remark 4.10. *In characteristic 0 the orthogonally simple characters are exactly the characters of the irreducible \mathbb{R} -representations. In particular any orthogonally simple character is the character of a representation ρ over some real number field L . Moreover $B := \sum_{g \in G} \rho(g)\rho(g)^{\text{tr}}$ is an invariant symmetric bilinear form that is **totally positive definite**, i.e. for all embeddings $\epsilon : L \rightarrow \mathbb{R}$ the form $\epsilon(B)$ is positive definite. Hence the orthogonal discriminant of an orthogonally simple character of degree $2m$ is $(-1)^m d(K^\times)^2$ for some totally positive d in the character field K .*

For finite fields Proposition 2.5 tells us that restriction of scalars does not change the orthogonal discriminant of the orthogonally simple Brauer character. For ordinary characters we need to take the norm by Remark 2.2.

Proposition 4.11. *Let χ be an orthogonally simple ordinary character of even degree and let F be a subfield of $K := K(\chi)$. Then*

$$\psi := \sum_{\sigma \in \text{Gal}(K/F)} \chi^\sigma$$

has character field $K(\psi) = F$ and a well defined discriminant

$$\text{disc}(\psi) = N_{K/F}(\delta)(F^\times)^2 =: N_{K/F}(\text{disc}(\chi))$$

where $\delta \in \text{disc}(\chi)$ is any representative of the square class $\text{disc}(\chi) \in K^\times / (K^\times)^2$.

Proof. There is an extension field L of K and an L -representation ρ with character χ . Following [11] we put

$$E_-(\rho) = \langle \rho(g) - \rho(g^{-1}) \mid g \in G \rangle_{\mathbb{Q}}$$

to denote the space of skew adjoint elements in the central simple K -algebra $A = \langle \rho(g) \mid g \in G \rangle_{\mathbb{Q}}$ spanned by the matrices in $\rho(G)$. By [11]

$$\text{disc}(\chi) = (-1)^{\chi(1)/2} N_{\text{red}}(X)(K^\times)^2 \in K^\times / (K^\times)^2 \text{ for any invertible } X \in E_-(\rho)$$

where N_{red} is the reduced norm of A , the composition of the embedding of A into a matrix algebra of the same dimension with the determinant (see for instance [15, Section 9]). If $X = \sum_{g \in G} a_g(\rho(g) - \rho(g^{-1})) \in E_-(\rho)$ is such an invertible element then the direct sum of the representations ρ^σ maps X to the skew adjoint unit

$$\tilde{X} := \sum_{g \in G} a_g(\oplus_{\sigma \in \text{Gal}(K/F)} (\rho^\sigma(g) - \rho^\sigma(g^{-1}))) \in E_-(\oplus_{\sigma \in \text{Gal}(K/F)} \rho^\sigma).$$

As $N_{\text{red}}(X) \in K$ we compute $N_{\text{red}}(\tilde{X}) = N_{K/F}(N_{\text{red}}(X))$ and hence

$$\text{disc}(\psi) = (-1)^{(\psi(1)/2)} N_{\text{red}}(\tilde{X})(F^\times)^2 = N_{K/F}(\text{disc}(\chi)).$$

□

4.4 Orthogonally stable characters

An orthogonally stable character is the sum of even degree orthogonally simple characters:

Definition 4.12. *An orthogonal character or Brauer character χ is called orthogonally stable if all absolutely irreducible indicator + constituents of χ have even degree.*

Theorem 4.13. *Any orthogonally stable ordinary character \mathcal{X} has a well defined orthogonal discriminant. More precisely put $K = K(\mathcal{X})$. For an orthogonally simple constituent \mathcal{X}_j of \mathcal{X} we put $L_j := K \cdot K(\mathcal{X}_j)$ and $\mathcal{G}_j = \text{Gal}(L_j/K)$. If*

$$\mathcal{X} = \sum_{j=1}^n \sum_{\sigma \in \mathcal{G}_j} \mathcal{X}_j^\sigma$$

is the sum of orthogonally simple characters organised according to their Galois orbits then

$$\text{disc}(\mathcal{X}) = \prod_{j=1}^n N_{L_j/K}(\text{disc}(\mathcal{X}_j)).$$

Proof. Let ρ be an orthogonal representation with character \mathcal{X} and let $Q \in \mathcal{Q}(\rho)$ be non-degenerate. Then Q is the orthogonal sum of non-degenerate quadratic forms Q_j on the Galois orbits of the orthogonally simple constituents. By Proposition 4.11 the discriminant of Q_j is $N_{L_j/K} \text{disc}(\chi_j)$. So the theorem follows from the multiplicativity of the discriminants (cf. Remark 2.1). \square

From Proposition 2.5 and Remark 2.4 we get an analogous result for Brauer characters.

Remark 4.14. *Any orthogonally stable Brauer character χ has a well defined orthogonal discriminant.*

5 Reduction of orthogonal representations

In this section we fix a number field K with ring of integers O_K . Let $\rho : G \rightarrow \text{GL}(V)$ be an orthogonal K -representation of G and $Q \in \mathcal{Q}(\rho)$ be a non-degenerate G -invariant quadratic form on V with polarisation B .

Definition 5.1. *The discriminant field extension is $\Delta(Q) := K(\sqrt{\text{disc}(Q)})/K$. For an orthogonally stable ordinary character \mathcal{X} with character field K we put $\Delta(\mathcal{X}) := K(\sqrt{\text{disc}(\mathcal{X})})/K$.*

Note that the degree of $\Delta(Q)$ is 1 if $\text{disc}(Q) \in (K^\times)^2$ and the degree of $\Delta(Q)$ is 2 if $\text{disc}(Q) \notin (K^\times)^2$.

For a prime ideal \wp we denote by K_\wp the completion of K at \wp , by O_\wp its valuation ring and π a generator of the unique maximal ideal $\wp O_\wp = \pi O_\wp$. Put $F := O_\wp / \pi O_\wp \cong O_K / \wp$ to denote the residue field of O_\wp .

Clearly Q extends to a quadratic form on the completion $V_\wp = V \otimes_K K_\wp$. An O_\wp -lattice M in (V_\wp, Q) is called **even** if $Q(M) \subseteq O_\wp$, and **integral** if $B(M, M) \subseteq O_\wp$, i.e. M is contained in its dual lattice, $M^\# = \{x \in V_\wp \mid B(x, M) \subseteq O_\wp\}$. Clearly even lattices are integral and, as $2Q(v) = B(v, v)$, being integral and even is equivalent if 2 is a unit in O_\wp .

Assumption 5.2. Assume that either 2 is a unit in O_φ or the trivial representation is not a constituent of the reduction of ρ modulo φ .

Lemma 5.3. Under Assumption 5.2 all integral $O_\varphi G$ -lattices in V_φ are even.

Proof. The statement is clear, if 2 is a unit in O_φ . So assume that $\text{char}(F) = 2$ and that M is an integral $O_\varphi G$ -lattice in V_φ . Then $2Q(x) = B(x, x) \in O_\varphi$ and $Q(x+y) + O_\varphi = Q(x) + Q(y) + O_\varphi$ for all $x, y \in M$. Hence

$$Q : M \rightarrow \frac{1}{2}O_\varphi/O_\varphi$$

is a G -invariant homomorphism of abelian groups. Its kernel N is a $\rho(G)$ -invariant O_φ -sublattice of M and G acts trivially on the F -module $M/(N + \pi M)$. As the trivial module is not a constituent of the reduction of ρ modulo φ , we have $N + \pi M = M$, so $N = M$ by the well known Nakayama Lemma. This shows that $Q(M) \subseteq O_\varphi$ and hence M is even. \square

A maximal even $O_\varphi G$ -lattice is an even, $\rho(G)$ -invariant lattice M in V_φ such that no proper $\rho(G)$ -invariant overlattice of M is even.

Lemma 5.4. Assume Assumption 5.2 and let M be a maximal even $O_\varphi G$ -lattice. Then $\pi M^\# \subseteq M \subseteq M^\#$ and $(M^\#, \pi Q)$ is an even lattice. Reduction modulo φ yields two non-degenerate quadratic FG -modules $(M^\#/M, \overline{\pi Q})$ and $(M/\pi M^\#, \overline{Q})$.

Proof. Clearly M is integral, so $M \subseteq M^\#$. If $\pi M^\# \not\subseteq M$, then we put $X := \frac{1}{\pi}M \cap \pi M^\# + M$. This is clearly a proper $O_\varphi G$ -overlattice of M . Elementary computations show that X is integral, and hence even (by Lemma 5.3), contradicting the maximality of M . Similarly $(M^\#, \pi Q)$ is integral and hence even.

Moreover the radical of the reduction \overline{Q} of Q modulo φ on $M/\pi M$ is $\pi M^\#$, by definition. Similarly one concludes that $(M^\#/M, \overline{\pi Q})$ is non-degenerate. \square

5.1 Reduction of orthogonal characters

Theorem 5.5. Let \mathcal{X} be an orthogonal character, K its character field, φ a prime ideal of O_K and $F = O_K/\varphi$. Assume that the reduction χ of \mathcal{X} modulo φ is orthogonally stable. Then

- (a) \mathcal{X} is orthogonally stable.
- (b) φ is not ramified in $\Delta(\mathcal{X})$.
- (c) If φ is inert in $\Delta(\mathcal{X})$ and $\Delta(\mathcal{X})$ has degree 2 then χ has orthogonal discriminant $O-$.
- (d) Assume that $\dim_{K(\chi)}(F)$ is odd. If φ is split in $\Delta(\mathcal{X})$ or $\Delta(\mathcal{X})$ has degree 1 then χ has orthogonal discriminant $O+$.

Proof. (a) is clear.

(b) If the \wp -adic Schur index of \mathcal{X} is 1, then there is a K_\wp -representation $\rho : G \rightarrow \mathrm{GL}(V_\wp)$ affording the character \mathcal{X} . Otherwise there is a suitable ramified extension L of K_\wp and an L -representation ρ affording \mathcal{X} . Let O_\wp denote the ring of integers either in K_\wp or in L and π a generator of the maximal ideal of O_\wp . Then $F = O_\wp/\pi O_\wp$. Choose a non-degenerate $Q \in \mathcal{Q}(\rho)$.

The orthogonal stability of χ implies Assumption 5.2. Choose a maximal even $O_\wp G$ -lattice M in V_\wp . Then Lemma 5.4 gives a chain $\pi M^\# \subseteq M \subseteq M^\#$ of even $O_\wp G$ -lattices. Then $M^\#/M$ and $M/\pi M^\#$ are orthogonal FG -representations and hence orthogonally stable, in particular of even dimensions.

Take $v_1, \dots, v_s \in M$ such that their images form a basis of $M/\pi M^\#$ and put $N := \langle v_1, \dots, v_s \rangle_{O_\wp} \leq M$. Then $(N, Q|_N)$ is a regular quadratic O_\wp -lattice of discriminant, say, δ_1 . By Lemma 2.8 we get $M = N \perp N^\perp$. Lemma 5.4 yields that $(N^\perp, \frac{1}{\pi}Q|_{N^\perp})$ is regular. If δ_2 denotes its discriminant, then $\mathrm{disc}(Q) = \delta_1 \delta_2$. In particular Remark 2.7 says that $\Delta_\wp := K_\wp[\sqrt{\delta_1 \delta_2}]/K_\wp$ is not ramified. Moreover the orthogonal discriminant of the quadratic space

$$(N, Q|_N) \otimes F \oplus (N^\perp, \frac{1}{\pi}Q|_{N^\perp}) \otimes F$$

over F is $O+$ if Δ_\wp is of degree 1 and $O-$ if Δ_\wp has degree 2.

(c) Here Δ_\wp has degree 2 and hence by Remark 2.6 $\dim_{K(\chi)}(F)$ is odd and the orthogonal discriminant of χ is $O-$.

(d) When Δ_\wp has degree 1 then Remark 2.6 (a) shows that it is only possible to read off the orthogonal discriminant of χ from the computations over F if $\dim_{K(\chi)}(F)$ is odd. In this case the orthogonal discriminant of χ is $O+$. \square

In particular for dyadic primes, i.e. those that divide 2, Theorem 5.5 yields good restrictions on the discriminant. We highlight this for rational characters:

Corollary 5.6. *The discriminant of an orthogonally stable rational character \mathcal{X} with an orthogonally stable reduction modulo 2 is 1 modulo 4.*

Proof. Let $d \in \mathbb{Z}$ be square free so that $d(\mathbb{Q}^\times)^2 = \mathrm{disc}(\mathcal{X})$. As \mathcal{X} is orthogonally stable modulo 2, the prime 2 is either split in $\mathbb{Q}(\sqrt{d})$ (i.e. $d \equiv 1 \pmod{8}$) or inert in $\mathbb{Q}(\sqrt{d})$ (i.e. $d \equiv -3 \pmod{8}$). \square

5.2 Primes not dividing the group order

The Brauer character table modulo a prime not dividing the group order is exactly the ordinary character table. Given an absolutely irreducible ordinary character \mathcal{X} of even degree and indicator $+$ and some prime p not dividing the group order we aim to compute the orthogonal discriminant of the p -Brauer character $\chi = \mathcal{X}$. Let $K = K(\mathcal{X})$ be the character field of \mathcal{X} , O_K be its ring of integers. Then the Conway polynomials define a unique prime ideal \wp_0 of

O_K with $p \in \wp_0$. Put $F := O_K/\wp_0$. Then F is the character field of χ , χ is the reduction of \mathcal{X} modulo \wp_0 , and χ is orthogonally stable. So Theorem 5.5 implies the following corollary.

Corollary 5.7. *The prime \wp_0 is unramified in the discriminant field extension $\Delta(\mathcal{X})$.*

If \wp_0 is split in $\Delta(\mathcal{X})$ or $\text{degree}(\Delta(\mathcal{X})) = 1$ then $\text{disc}(\chi) = O+$.

If \wp_0 is inert in $\Delta(\mathcal{X})$ then $\text{disc}(\chi) = O-$.

5.3 An algorithm to compute the discriminant field extension

Theorem 5.5 can be applied to compute the orthogonal discriminant of an orthogonally stable ordinary character \mathcal{X} by computing enough orthogonally stable reductions.

To this aim, assume that we are given an orthogonally stable character \mathcal{X} of some finite group G . Let $K := K(\mathcal{X})$ denote its character field and O_K the ring of integers of K . Put

$$\mathcal{P} := \{ \wp \trianglelefteq_{\max} O_K \mid \text{reduction mod } \wp \text{ of } \mathcal{X} \text{ is not orthogonally stable} \}.$$

Then \mathcal{P} is contained in the set of prime ideals dividing $|G|O_K$. In particular it is a finite set.

Put $\epsilon := (-1)^{\mathcal{X}(1)/2}$, so that $\epsilon \text{disc}(\mathcal{X})$ is a totally positive square class in K (cf. Remark 4.10). Compute $\delta_1, \dots, \delta_t \in O_K$ such that the products $\prod_{j \in I} \delta_j$, for $I \subseteq \{1, \dots, t\}$ represent all totally positive square classes $\delta(K^\times)^2$ for which $K_\wp(\sqrt{\delta})$ is unramified for all $\wp \notin \mathcal{P}$. We assume that $(\delta_i(K^\times)^2 \mid i = 1, \dots, t)$ is \mathbb{F}_2 -linearly independent in $K^\times/(K^\times)^2$. For a non-dyadic prime $\wp \notin \mathcal{P}$ we define $a_\wp \in \mathbb{F}_2^t$ by $a_\wp(i) = 0$ if δ_i is a square in the residue field O_K/\wp and $a_\wp(i) = 1$ otherwise. Choose non-dyadic primes $\{\wp_1, \dots, \wp_t\}$ such that the matrix A with columns a_{\wp_i} , so

$$A = (a_{\wp_1}, \dots, a_{\wp_t}) \in \mathbb{F}_2^{t \times t}$$

has full rank t . Now compute the orthogonal discriminant ϵd_i of \mathcal{X} (mod \wp_i) for $i = 1, \dots, t$ and put $x_i := 1$ if $d_i = O-$ and $x_i := 0$ if $d_i = O+$. Put $y := xA^{-1}$ and $I := \{i \in \{1, \dots, t\} \mid y_i = 1\}$. Then

$$\delta := \epsilon \prod_{i \in I} \delta_i(K^\times)^2 = \text{disc}(\mathcal{X}).$$

Remark 5.8. *In practice we also use all dyadic primes \wp where \mathcal{X} has an orthogonally stable reduction χ as these yield even more restrictions than the non-dyadic primes. Instead of working only with $\delta_1, \dots, \delta_t$ we need to test the ramification behaviour of \wp in $K(\sqrt{\epsilon d})/K$ for all products d of the δ_i . By Theorem 5.5 we can exclude those d where \wp is ramified and the orthogonal discriminant of χ allows us to further exclude those d where \wp is inert (if $\text{disc}(\chi) = O+$) respectively split (if $\text{disc}(\chi) = O-$).*

Example 5.9. The group J_2 has two absolutely irreducible ordinary characters of degree 224 with character field $\mathbb{Q}(\sqrt{5})$ that are exchanged under an outer automorphism. So $\text{disc}(224a) = \text{disc}(224b) =: d(\mathbb{Q}(\sqrt{5})^\times)^2$. The decomposition matrices in GAP [4] tell us that both characters are orthogonally stable modulo 2 and 7. Constructing the representations we obtain that they yield embeddings into $O_{224}^+(\mathbb{F}_4)$ respectively $O_{224}^+(\mathbb{F}_{49})$. As $|J_2| = 2^7 3^3 5^2 7$ the only possibilities

for d are $1, 3, (5 + \sqrt{5})/2$, and $3(5 + \sqrt{5})/2$. Now 1 and 3 are squares in \mathbb{F}_{49} , whereas $(5 + \sqrt{5})/2$ is not a square. As the reduction mod 7 has orthogonal discriminant $+$, we are left with 1 or 3 as possible discriminants. The prime 2 is ramified in the extension $\mathbb{Q}(\sqrt{3}, \sqrt{5})/\mathbb{Q}(\sqrt{5})$, which excludes the possibility that $d = 3$ just using the fact that the reduction modulo 2 is orthogonally stable. We hence conclude that the discriminants are squares, $\text{disc}(224a) = \text{disc}(224b) = 1$.

5.4 Cyclic defect

In this section we assume that \mathcal{X} is an even degree absolutely irreducible orthogonal character in some p -block with cyclic defect group. Then the reduction χ of \mathcal{X} is multiplicity free. In characteristic 2 , the trivial character is the unique odd degree indicator $+$ irreducible Brauer character, so the fact that χ is multiplicity free and of even degree allows to conclude that χ is orthogonally stable and hence Theorem 5.5 applies. So the theory in this section is void for $p = 2$ and hence we may and will assume that $p \neq 2$. In particular we may consider bilinear forms instead of quadratic forms, all integral lattices are automatically even, and a prime ideal \wp is ramified in the discriminant field extension, if and only if there is a G -invariant maximal integral lattice L such that $L^\# / L$ has odd dimension over the residue field O_\wp / \wp .

For blocks of defect 1 the converse of Theorem 5.5 is true.

Theorem 5.10. *Let p be an odd prime and let $\mathcal{X} \in \text{Irr}_{\mathbb{C}}(G)$ be an orthogonally stable character in a p -block with defect 1 . Let K denote the character field of \mathcal{X} and \wp a prime ideal of O_K that contains p . Then the reduction χ of \mathcal{X} modulo \wp is orthogonally stable if and only if \wp is unramified (i.e. split or inert) in $\Delta(\mathcal{X})$.*

Proof. Let K_\wp be the completion of K at the prime \wp , L_\wp be some unramified extension of K_\wp with ring of integers O_\wp and $\rho : G \rightarrow \text{GL}_n(L_\wp)$ be some representation affording the character \mathcal{X} . Fix a non-degenerate quadratic form $Q \in \mathcal{Q}(\rho)$ and choose $M \leq L_\wp^n$ to be some maximal even $\rho(G)$ -invariant O_\wp -lattice. The assumption that the p -defect of \mathcal{X} is one implies that all $O_\wp G$ -lattices in L_\wp^n are linearly ordered by inclusion ([1, Theorem 11]). As $M^\# / M$ and $M / \wp M^\#$ are self-dual modules, their composition factors are ordered as $(S_1, S_2, \dots, S_s, S = S^\vee, S_s^\vee, \dots, S_2^\vee, S_1^\vee)$ for $M^\# / M$ and $(T_1, T_2, \dots, T_t, T = T^\vee, T_t^\vee, \dots, T_2^\vee, T_1^\vee)$ for $M / \wp M^\#$, possibly without the self-dual composition factors S, T . As the module $M^\# / \wp M^\#$ is multiplicity free, all these composition factors are pairwise non-isomorphic. The assumption that the reduction of \mathcal{X} modulo \wp is not orthogonally stable hence implies that both self-dual composition factors S and T do occur and are of odd dimension. In particular $M^\# / M$ has odd dimension so the \wp -adic valuation of the discriminant δ of Q is odd. As $p \neq 2$, this is equivalent to saying that \wp ramifies in $K(\sqrt{\delta})/K$. \square

Example 5.11. Let $G = \text{SL}_2(8)$ and \mathcal{X} be the rational absolutely irreducible character of degree $\mathcal{X}(1) = 8$. Then the reduction of \mathcal{X} modulo 3 is the sum of two characters of degree 1 and 7 , and hence not orthogonally stable. However, the discriminant of \mathcal{X} is 1 .

Example 5.11 is covered by the following general result for characters in blocks with cyclic defect groups.

Theorem 5.12. *Assume that \mathcal{X} is an indicator + absolutely irreducible ordinary character of even degree in a p -block of cyclic defect. Denote by K its character field, O_K the ring of integers and \wp a prime ideal of O_K containing the odd prime p .*

- *If \mathcal{X} belongs to the exceptional vertex then its reduction modulo \wp is not orthogonally stable, if and only if \wp is ramified in the discriminant field extension $\Delta(\mathcal{X})$.*
- *If \mathcal{X} does not belong to the exceptional vertex then \wp is ramified in the discriminant field extension $\Delta(\mathcal{X})$ if and only if the defect is odd and the reduction of $\mathcal{X} \bmod \wp$ is not orthogonally stable.*

In particular for even defect only the characters belonging to the exceptional vertex may have ramification at \wp in the discriminant field extension.

Proof. We keep the notation of Theorem 5.10. If \mathcal{X} belongs to the exceptional vertex, then by [14, Theorem (VIII.3)] the G -invariant O_\wp -lattices form a chain and we have the same situation as in the proof of Theorem 5.10 and hence the conclusion follows with the same argument.

Now assume that \mathcal{X} does not belong to the exceptional vertex and we also assume that its reduction mod \wp is not orthogonally stable. So there are exactly two constituents χ_S , χ_T of indicator + and odd degree. The paper [10] investigates the radical idealizer process, that starts with the O_\wp -order Λ spanned by the matrices in $\rho(G)$ and successively constructs involution invariant overorders until arriving at the hereditary order $H(\Lambda)$, the head order of Λ . As $H(\Lambda)$ is invariant under the canonical involution of G , its lattices hence form a chain that is invariant under dualization. The explicit form of $H(\Lambda)$ given in [10, Theorem 3.15] allows to conclude that χ_S and χ_T occur in the same constituent of $H(\Lambda) \bmod \wp$ if and only if the defect is even. The same argument as in the proof of Theorem 5.10 now applied to $H(\Lambda)$ allows to conclude the statement. \square

6 Two examples

6.1 The orthogonal discriminants of the orthogonally stable irreducible characters of J_1 .

Let J_1 denote the first Janko group of order $2^3 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 19$. Then there are seven indicator + absolutely irreducible ordinary characters of even degree: The two Galois conjugate characters $56a$, $56b$ with character field $\mathbb{Q}(\sqrt{5})$, two rational characters of degree 76 and three Galois conjugate characters $120a$, $120b$, $120c$ with character field $\mathbb{Q}(c_{19})$, the unique subfield of degree 3, index 6 of the 19th cyclotomic field. All these three character fields \mathbb{Q} , $\mathbb{Q}(\sqrt{5})$, and $\mathbb{Q}(c_{19})$

have narrow class number 1, which means that any ideal of their ring of integers has a totally positive generator that is unique up to multiplication by squares of units.

The reductions modulo 2, 3, 5 of all these seven characters stay absolutely irreducible, so the only primes that can divide the discriminants are those that divide $7 \cdot 11 \cdot 19$. With GAP [4] we compute the reduction modulo 7, 11, 19 as follows

\mathcal{X}	$56a$	$56b$	$76a$	$76b$	$120a$	$120b$	$120c$
(mod 7)	$56a$	$56b$	$1 + 75$	$31 + 45$	$45 + 75$	$31 + 89$	120
(mod 11)	56	$7 + 49$	$27 + 49$	$7 + 69$	$1 + 119$	$56 + 64$	$14 + 106$
(mod 19)	$1 + 55$	$22 + 34$	$76a$	$76b$	$43 + 77$	$43 + 77$	$43 + 77$

By Theorem 5.10 the determinant of both characters of degree 76 is $7 \cdot 11 = 77$.

For the non-rational characters the “reduction modulo 7,11,19” means the reduction modulo the prime ideal \wp_0 from Section 4.2. As in our cases, the residue fields are the prime fields, we can identify the ideal \wp_0 using the GAP-command

$$x := \text{Int}(\text{FrobeniusCharacterValue}(y, p));$$

where y is a name of the irrationality (here $y = ER(5)$ respectively $y = EC(19)$) and p the corresponding prime. We obtain

y	$\sqrt{5}$	$\sqrt{5}$	c_{19}	c_{19}
p	11	19	7	11
x	4	9	0	6

We get the factorization $11 = p_{11}p'_{11}$ where $p_{11} = 4 - \sqrt{5}$ and p'_{11} is its Galois conjugate $p'_{11} = 4 + \sqrt{5}$. Similarly $19 = p_{19}p'_{19}$ with $p_{19} = (9 - \sqrt{5})/2$, and hence

$$\text{disc}(56a) = p'_{11}p_{19}(\mathbb{Q}(\sqrt{5})^\times)^2 = (17 - 4\sqrt{5})(\mathbb{Q}(\sqrt{5})^\times)^2, \quad \text{disc}(56b) = (17 + 4\sqrt{5})(\mathbb{Q}(\sqrt{5})^\times)^2.$$

For the field $\mathbb{Q}(c_{19})$ we have $c_{19} = z + z^8 + z^{8^2} + z^{8^3} + z^{8^4} + z^{8^5}$ where z is a primitive 19th root of unity. A generator σ of the Galois group of $\mathbb{Q}(c_{19})$ acts on the cyclotomic field by squaring z and hence maps c_{19} to $c'_{19} := 4 - c_{19}^2$. Note that σ acts on the characters as the cyclic permutation $(120a, 120c, 120b)$. We compute

$$\alpha = 4 + c_{19} - c'_{19}, \quad \beta = 20 + 10c_{19} - 3c'_{19}, \quad \gamma = 7 + 3c_{19} - c'_{19}$$

as totally positive elements of norm 7, 11, respectively 19. Replacing c_{19} by 0 maps α onto $0 \in \mathbb{F}_7$ and similarly mapping c_{19} to 6 maps β to $0 \in \mathbb{F}_{11}$. So we get

$$\text{disc}(120a) = \alpha\sigma(\alpha)\beta\gamma(\mathbb{Q}(c_{19})^\times)^2 = (29 - 18c_{19} - 9c'_{19})(\mathbb{Q}(c_{19})^\times)^2$$

and $\text{disc}(120c) = \sigma(\text{disc}(120a))$, $\text{disc}(120b) = \sigma^2(\text{disc}(120a))$.

Theorem 6.1. *The discriminants of the ordinary absolutely irreducible orthogonal characters of J_1 are*

\mathcal{X}	$56a$	$56b$	$76a$	$76b$	$120a$	$120b$	$120c$
$\text{disc}(\mathcal{X})$	$17 - 4\sqrt{5}$	$17 + 4\sqrt{5}$	77	77	$29 - 18c_{19} - 9c'_{19}$	$47 + 9c_{19} + 18c'_{19}$	$38 + 9c_{19} - 9c'_{19}$

Remark 6.2. *The field $K := \mathbb{Q}(\sqrt{5})(\sqrt{\text{disc}(56a)})$ is not Galois over \mathbb{Q} , its normal closure $L = K(\sqrt{\text{disc}(56b)})$ has Galois group D_8 , the dihedral group of order 8.*

Similarly $K := \mathbb{Q}(c_{19})(\sqrt{\text{disc}(120a)})$ is not Galois over \mathbb{Q} , its normal closure is $L = K(\sqrt{\text{disc}(120b)}, \sqrt{\text{disc}(120c)})$ with Galois group $C_2 \wr C_3$ of order 24.

6.2 The Held group

The Held group He is a sporadic simple group of order $2^{10}3^35^27^317$. With the meat-axe [13] we construct all irreducible indicator + even degree modular representations for the prime divisors of the group order:

Characteristic 2

246	$O+$	246	$O+$	680	$O-$	1920	$O+$	2008	$O-$
4352	$O+$	4608	$O+$	21504	$O+$	21504	$O+$		

Characteristic 3

1920	$O-$	6172	$O-$	6272	$O-$	7650	$O-$	14400	$O-$
------	------	------	------	------	------	------	------	-------	------

Characteristic 5

104	$O+$	680	$O+$	1240	$O-$	4080	$O+$	4116	$O-$	6528	$O+$
7650	$O+$	9640	$O-$	10860	$O-$	11900	$O+$	14400	$O-$	22050	$O+$

Characteristic 7

50	$O+$	426	$O+$	798	$O+$	1072	$O-$	1700	$O+$
3654	$O+$	6154	$O-$	6272	$O-$	13720	$O-$	23324	$O+$

Characteristic 17

680	$O+$	4080	$O+$	4352	$O-$	6528	$O+$
7650	$O+$	10880	$O+$	11900	$O+$	23324	$O+$

The following table gives the information on the orthogonal discriminants in characteristic 0. The first column gives the number of the ordinary absolutely irreducible character \mathcal{X} followed by its degree $\mathcal{X}(1)$ and its discriminant $\text{disc}(\mathcal{X})$. For the prime divisors of the group order for which the reduction of \mathcal{X} is orthogonally stable we also give the orthogonal discriminant $O+$ or $O-$ of this reduction computed from the orthogonal discriminants in the

tables before. For characters in blocks with cyclic defect group such that the reduction mod p is not orthogonally stable we display p , to indicate that p is a prime divisor of the discriminant (see Theorem 5.10). All character fields are rational, except for the two algebraic conjugate characters of degree 21504, where the character field is $\mathbb{Q}(\sqrt{21})$. Here $\cdot, O-$ indicates that the reduction of the character number 30 modulo the prime ideal generated by $4 - \sqrt{21}$ is orthogonally stable of orthogonal discriminant $O-$ and the one modulo $4 + \sqrt{21}$ is not orthogonally stable.

<i>No</i>	<i>deg</i>	<i>disc</i>	(mod 2)	(mod 3)	(mod 5)	(mod 7)	(mod 17)
6	680	21	$O-$		$O+$		$O+$
12	1920	17	$O+$	$O-$	$O-$	$O-$	17
13	4080	1			$O+$	$O+$	$O+$
14	4352	105	$O+$				$O-$
15	6272	17	$O+$	$O-$		$O-$	17
16	6528	1	$O+$		$O+$		$O+$
19	7650	-1		$O-$	$O+$	$O-$	$O+$
22	10880	1	$O+$		$O+$	$O+$	$O+$
25	11900	21	$O-$		$O+$		$O+$
26	13720	17			$O-$	$O-$	17
27	14400	17		$O-$	$O-$		17
30	21504	17	$O+$		$\cdot, O-$		17
31	21504	17	$O+$		$O-, \cdot$		17
32	22050	-119		$O+$	$O+$	7	17
33	23324	1				$O+$	$O+$

For the characters of numbers $\notin \{30, 31, 33\}$ the proofs are all similar, so let us give the one for number 14. Here the possible discriminants are in $\{1, 3, 5, 7, 15, 21, 35, 105\}$. As the reduction mod 2 is orthogonally stable of discriminant $O+$, Corollary 5.6 tells us that the discriminant of \mathcal{X} is 1 mod 8, leaving 1 and 105 as the only possibilities. But the discriminant is not a square modulo 17, so $\text{disc}(\mathcal{X}) = 105$.

Characters number 30 and 31 are algebraic conjugate and so are their discriminants. We give the proof for number 30. Here the character field is $K = \mathbb{Q}(\sqrt{21})$. Its ring of integers has class number 1 and a totally positive unit u that is not a square, $u = (5 + \sqrt{21})/2$. Note that $3u = ((3 + \sqrt{21})/2)^2$ is a square in K . The prime ideals dividing $3 \cdot 5 \cdot 7 \cdot 17$ are generated by $p_3 := (3 + \sqrt{21})/2$ of norm -3 , $p_5 := 4 + \sqrt{21}$ and $p'_5 := 4 - \sqrt{21}$ of norm -5 , $p_7 = (7 + \sqrt{21})/2$ of norm 7 and 17 of norm 17^2 . By Theorem 5.10 we know that 17 divides the discriminant of \mathcal{X} . We also know that the reduction of \mathcal{X} modulo (p'_5) is orthogonally stable. So the discriminant of \mathcal{X} is in

$$\{17, 17u, 17p_7, 17up_7, 17p_3p_5, 17up_3p_5, 17p_7p_3p_5, 17up_7p_3p_5\}.$$

Only for $d = 17$ and $d = 17up_7p_3p_5$ the prime 2 is not ramified in $K(\sqrt{d})/K$. Now p'_5 is inert

in this extension for $d = 17$ and decomposed for $d = 17up_7p_3p_5$. As the reduction modulo p_5' of \mathcal{X} has orthogonal discriminant $O-$, we conclude that $d = 17$ is the only possibility.

For the last character, number 33, we use a completely different argument. This character extends in two ways to the group $He:2$. Take one of these extensions and restrict it to the maximal subgroup $S_4(4):4 \leq He:2$. This restriction is orthogonally stable and contains the following orthogonal simple characters:

<i>deg</i>	18 <i>ab</i>	68	50 <i>ab</i>	204	170	153 <i>ab</i>	816	900	1020	256 <i>ab</i>	256	680	680
<i>K</i>	$\mathbb{Q}(i)$	\mathbb{Q}	$\mathbb{Q}(\sqrt{-2})$	\mathbb{Q}	\mathbb{Q}	$\mathbb{Q}(i)$	\mathbb{Q}	\mathbb{Q}	\mathbb{Q}	$\mathbb{Q}(i)$	\mathbb{Q}	\mathbb{Q}	\mathbb{Q}
<i>mult</i>	1	1	1	2	1	1	4	7	6	3	1	2	5
<i>disc</i>	1	1	1	.	-1	-1	.	17	.	1	17	.	1

This table gives the degree of the orthogonal simple characters occurring in the restriction of the extension of \mathcal{X} to $S_4(4):4$. The small letters indicate pairs of complex conjugate characters, whose character fields are given in the next row. Their discriminant can hence be obtained by Proposition 2.9. The character of degree 68 is induced up from a rational character of degree 34 of the group $S_4(4):2$. Hence the invariant forms are orthogonal sums of two isometric forms and so the discriminant is a square. The same argument applies to the characters of degrees 170, 204, 816, 1020, and the two characters of degree 680. For the character of degree 900 we apply [11, Theorem 4.6] to conclude that the two characters of degree 450 of the group $S_4(4):2$ have discriminant $-(17 \pm 4\sqrt{17})$ respectively. So the discriminant of the character of degree 900 is their product, 17. The characters of degree 256 restrict irreducibly to the simple group $S_4(4)$. The restriction is irreducible modulo 2 and 3, not orthogonally stable modulo 17. So its discriminant is either 17 or 85. As the orthogonal discriminant of the reduction mod 2 is $O+$, we conclude that the discriminant is $\equiv 1 \pmod{8}$ (see Corollary 5.6) and hence it is 17.

References

- [1] Richard Brauer, Investigations on group characters, *Ann. of Math.* **42** (1941) 936-958.
- [2] John H. Conway, Robert T. Curtis, Simon P. Norton, Richard A. Parker and Robert A. Wilson, *Atlas of Finite Groups*. Oxford University Press (1985)
- [3] Andreas Dress, Induction and structure theorems for orthogonal representations of finite groups, *Annals of Mathematics, Second Series*, **102** (1975) 291-325.
- [4] The GAP Group, GAP – Groups, Algorithms, and Programming, Version 4.11.1; 2021.
- [5] Martin Isaacs, *Character Theory of Finite Groups*, Pure and Applied Mathematics, 69. New York-San Francisco-London: Academic Press (1976)
- [6] Christoph Jansen, Klaus Lux, Richard A. Parker, Robert A. Wilson, *An atlas of Brauer characters*. London Mathematical Society Monographs. Oxford: Clarendon Press (1995)

- [7] Martin Kneser, *Quadratische Formen*, Neu bearbeitet und herausgegeben in Zusammenarbeit mit Rudolf Scharlau. Springer (Berlin) (2002)
- [8] John Milnor, On isometries of inner product spaces. *Invent. Math.* **8** (1969) 83-97
- [9] Gabriel Navarro, *Characters and blocks of finite groups*. London Mathematical Society Lecture Note Series. 250. Cambridge: Cambridge University Press (1998).
- [10] Gabriele Nebe, On blocks with cyclic defect group and their head orders, *Comm. Alg.* **33** (2005) 689-708.
- [11] Gabriele Nebe, Orthogonal determinants of characters, Preprint.
- [12] Gabriele Nebe, Richard Parker, Equivariant Witt groups of quadratic forms in characteristic 2, Preprint.
- [13] Richard A. Parker, The computer calculation of modular characters. (The meat-axe). *Computational group theory, Proc. Symp., Durham/Engl. 1982 (1984)* 267-274
Highly improved partially parallel implementation by Richard Parker (2022)
- [14] Wilhelm Plesken, *Group rings of finite groups over p -adic integers*, Springer Lecture Notes in Mathematics **1026** Berlin etc.: Springer-Verlag (1983).
- [15] Irving Reiner, *Maximal Orders*. Lond. Math. Soc. Monogr. 5 (1975) Academic Press, London.
- [16] Winfried Scharlau, *Quadratic and Hermitian forms*, Grundlehren der Mathematischen Wissenschaften **270**. Berlin etc.: Springer-Verlag (1985).
- [17] Peter Sin, Wolfgang Willems, G -invariant quadratic forms, *J. Reine Angew. Math.* **420** (1991) 45-59.
- [18] Alexandre Turull, Schur index two and bilinear forms, *J. Algebra* **157** (1993) 562-572.