

# SOME RATIONAL HOMOLOGY COMPUTATIONS FOR DIFFEOMORPHISMS OF ODD-DIMENSIONAL MANIFOLDS

JOHANNES EBERT AND JENS REINHOLD

ABSTRACT. We calculate the rational cohomology of the classifying space of the diffeomorphism group of the manifolds  $U_{g,1}^n := \#^g(S^n \times S^{n+1}) \setminus \text{int}(D^{2n+1})$ , for large  $g$  and  $n$ , up to approximately degree  $n$ . The answer is that it is a free graded commutative algebra on an appropriate set of Miller–Morita–Mumford classes.

Our proof goes through the classical three-step procedure: (a) compute the cohomology of the homotopy automorphisms, (b) use surgery to compare this to block diffeomorphisms, (c) use pseudoisotopy theory and algebraic  $K$ -theory to get at actual diffeomorphism groups.

## CONTENTS

1. Introduction	2
1.1. Context: Madsen–Weiss type theorems	2
1.2. Main result	3
1.3. Relation to Hebestreit–Perlmutter’s work	4
1.4. Method of proof	4
1.5. Overview of the chapters	7
1.6. Acknowledgements	8
2. Characteristic classes of smooth and block bundles	8
2.1. Automorphism groups	8
2.2. Tautological classes	10
2.3. Some vanishing theorems for tautological classes	10
2.4. Borel classes	13
3. Rational cohomology of block diffeomorphism spaces: general theory	15
3.1. Some words about rational homotopy theory	15
3.2. Block diffeomorphisms versus tangential homotopy automorphisms	18
3.3. Cohomology of mapping spaces	20
3.4. Consequences for block diffeomorphism spaces	24
4. Homotopy calculations for the manifolds $U_{g,1}^n$	25
4.1. Homotopy groups	25
4.2. Mapping class groups	26
4.3. Some higher homotopy groups of the homotopy automorphisms	28
4.4. The spectral sequence for tangential homotopy automorphisms	31
5. A representation-theoretic calculation	35

---

*Date:* June 13, 2022.

The authors were supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 427320536 – SFB 1442, as well as under Germany’s Excellence Strategy EXC 2044 – 390685587, Mathematics Münster: Dynamics–Geometry–Structure.

5.1. Generalities	35
5.2. A special invariant calculation	37
6. The cohomology of the block diffeomorphism space	42
6.1. Using invariant theory	42
6.2. Using Borel’s vanishing theorem	46
7. The endgame: from block diffeomorphisms to actual diffeomorphisms	48
7.1. Getting the mapping class group under control	48
7.2. Computation of the cohomology	53
References	55

## 1. INTRODUCTION

**1.1. Context: Madsen–Weiss type theorems.** For a smooth compact manifold with boundary  $M$ , let  $\text{Diff}_\partial(M)$  denote the group of diffeomorphisms of a smooth compact manifold  $M$  which are equal to the identity near  $\partial M$ . One of the success stories of differential topology in the 21st century was a (partial) computation of the cohomology of the classifying space  $B\text{Diff}_\partial(M)$  for some even-dimensional manifolds, by Madsen–Weiss [50] (for surfaces) and by Galatius–Randal-Williams [25] [27] [26] (in the higher dimensional case). The simplest case of these results concerns the manifolds

$$W_{g,1}^n := \#^g(S^n \times S^n) \setminus \text{int}(D^{2n}),$$

the connected sum of  $g$  copies of  $S^n \times S^n$ , minus the interior of a disc, and are formulated in terms of the Madsen–Tillmann spectra<sup>1</sup>  $\text{MT}\theta_{2n}^n$ , the Thom spectrum of the additive inverse of the universal  $2n$ -dimensional vector bundle over the  $n$ -connected cover  $BO(2n)\langle n \rangle \rightarrow BO(2n)$ . There is a natural map  $\alpha_g : B\text{Diff}_\partial(W_{g,1}^n) \rightarrow \Omega_0^\infty \text{MT}\theta_{2n}^n$  to the unit component of the infinite loop space. These maps are compatible for varying value of  $g$ , and induce a map

$$\alpha_\infty : \text{hocolim}_{g \rightarrow \infty} B\text{Diff}_\partial(W_{g,1}^n) \rightarrow \Omega_0^\infty \text{MT}\theta_{2n}^n$$

in the limiting case, which is an integral homology equivalence (for  $n = 1$  by [50], for  $n \geq 3$  by [25] and for  $n = 2$  by [26]). This is complemented by homological stability theorems (unless  $n = 2$ ) due to [34] and [27], so that  $\alpha_g$  induces an isomorphism in homology in a range of degrees increasing with  $g$ .

The rational cohomology of  $\Omega_0^\infty \text{MT}\theta_{2n}^n$  (and more general Madsen–Tillmann spectra) is easily calculated using the standard tools from algebraic topology. The answer is that it is the free graded-commutative algebra generated by the vector space  $(s^{-2n} H^*(BO(2n)\langle n \rangle; \mathbb{Q}))_{>0}$ , the positive degree part of the desuspension of  $H^*(BO(2n)\langle n \rangle; \mathbb{Q})$ . Let  $\mu_c \in H^{k-2n}(\Omega_0^\infty \text{MT}\theta_{2n}^n; \mathbb{Q})$  be the element corresponding to  $c \in H^k(BO(2n)\langle n \rangle; \mathbb{Q})$ . The pullback  $\alpha_g^* \mu_c \in H^{k-2n}(B\text{Diff}_\partial(W_{g,1}^n); \mathbb{Q})$  is the tautological class  $\kappa_c$  of the universal bundle over  $B\text{Diff}_\partial(W_{g,1}^n)$ . Finally  $H^*(BO(2n)\langle n \rangle; \mathbb{Q})$  is the polynomial algebra generated by the Pontrjagin classes  $p_m$  with  $\frac{n+1}{4} \leq m \leq n-1$  and the Euler class. So altogether, in a range of degrees,  $H^*(B\text{Diff}_\partial(W_{g,1}^n); \mathbb{Q})$  is a polynomial algebra in certain tautological classes.

All these results are for *even-dimensional* manifolds. The construction of the map  $\alpha_g$  can be generalized to any manifold, and yields for oriented  $M$  of dimension

<sup>1</sup>We use the notation from [38] instead of that from [25].

$d$  a map  $\alpha_M : B\text{Diff}_\partial^+(M) \rightarrow \Omega_0^\infty \text{MT}SO(d)$  (the map  $\alpha_g$  above is a refinement of this construction, which exists as  $W_{g,1}^n$  is  $(n-1)$ -connected and  $n$ -parallelizable, i.e. the restriction of  $TW_{g,1}^n$  to the  $n$ -skeleton of  $W_{g,1}^n$  is trivial). It has been observed by the first named author [13] that the classes  $\alpha_M^* \mu_{L_m}$  associated to the components of the Hirzebruch  $L$ -class vanish, for each odd-dimensional  $d$ , though  $\mu_{L_m}$  is nonzero. Hence any naive generalization of say [50] or [25] will fail in odd dimensions.

**1.2. Main result.** Even though some substantial inroads into the odd-dimensional situation have been made recently [59], [7], [58], [38], it remains a mystery and there does not seem to be a convincing conjectural odd-dimensional analogue of the main result of [25]. Our modest hope in this work is that our main result, Theorem A below, might eventually serve as a piece of evidence which helps to formulate an odd-dimensional version of these results. Let us consider the manifolds

$$U_{g,1}^n := \#^g(S^n \times S^{n+1}) \setminus \text{int}(D^{2n+1})$$

which we view as an odd-dimensional variant of the manifolds  $W_{g,1}^n$ . Since  $U_{g,1}^n$  is  $(n-1)$ -connected and  $n$ -parallelizable, one obtains a map

$$\beta_g^n : B\text{Diff}_\partial(U_{g,1}^n) \rightarrow \Omega_0^\infty \text{MT}\theta_{2n+1}^n, \quad (1.1)$$

where the target is the Madsen–Tillmann spectrum of  $BO(2n+1)\langle n \rangle \rightarrow BO(2n+1)$ . Note that

$$H^*(BO(2n+1)\langle n \rangle; \mathbb{Q}) \cong \mathbb{Q}[L_m \mid \frac{n+1}{4} \leq m \leq n]$$

(there is no Euler class; and it is more convenient to use the components of the Hirzebruch  $L$ -class instead of the Pontrjagin classes as generators). Hence  $H^*(\Omega_0^\infty \text{MT}\theta_{2n+1}^n; \mathbb{Q})$  is the exterior algebra generated by the elements  $\{\mu_{L_{m_1} \cdots L_{m_r}}\}$ , where  $\frac{n+1}{4} \leq m_1 \leq \dots \leq m_r \leq n$ .

**Theorem A.** *Assume that  $n \geq 5$ . Then the map*

$$(\beta_g^n)^* : H^*(\Omega_0^\infty \text{MT}\theta_{2n+1}^n; \mathbb{Q}) \rightarrow H^*(B\text{Diff}_\partial(U_{g,1}^n); \mathbb{Q})$$

*is surjective in degrees  $*$   $\leq \min(\frac{g-2}{2}, n-4)$ , and in that range of degrees, the kernel is the ideal generated by the classes  $\mu_{L_m}$  (all  $m$ ) and by the linear subspace  $H^1(\Omega_0^\infty \text{MT}\theta_{2n+1}^n; \mathbb{Q})$ .*

*Remark 1.2.* (1) That  $\mu_{L_m}$  lies in the kernel of  $(\beta_g^n)^*$  is the main result of [13].

(2) The space  $H^1(\Omega_0^\infty \text{MT}\theta_{2n+1}^n; \mathbb{Q})$  is zero if  $n$  is even. If  $n \equiv 1 \pmod{4}$ , say  $n = 4k+1$ , one checks that

$$H^1(\Omega_0^\infty \text{MT}\theta_{8k+3}^{4k+1}; \mathbb{Q}) = \mathbb{Q}\{\mu_{L_{2k+1}}\},$$

and the triviality of  $(\beta_g^n)^*$  in degree 1 follows from [13]. If  $n \equiv 3 \pmod{4}$ , say  $n = 4k-1$ , we get a new relation. In that case, one checks that

$$H^1(\Omega_0^\infty \text{MT}\theta_{8k-1}^{4k-1}; \mathbb{Q}) = \mathbb{Q}\{\mu_{L_{2k}}, \mu_{L_k}^2\},$$

and the new relation is  $\kappa_{L_k}^2 = 0$ , which holds more generally for all stably parallelizable manifolds of those dimensions. We give the fairly elementary proof in Proposition 2.12 below; for the special manifolds  $U_{g,1}^n$  the relation comes out of the proof of Theorem A.

(3) The bound in  $g$  stems from a homological stability result due to Perlmutter [58, Corollary 1.3.2]: the stabilization map  $B\text{Diff}_\partial(U_{g,1}^n) \rightarrow B\text{Diff}_\partial(U_{g+1,1}^n)$  is homologically  $\frac{g-2}{2}$ -connected (with integral coefficients).

- (4) The bound in  $n$  comes from our method of proof which we describe informally in §1.4 below.

**1.3. Relation to Hebestreit–Perlmutter’s work.** Let us comment on the relationship of the present work with [38]. The disjoint union

$$BD := \coprod_g B\text{Diff}_\partial(U_{g,1}^n)$$

carries a natural structure of an algebra over the operad of little  $(2n + 1)$ -discs. Hence we can form its group completion  $\Omega B(BD)$  which is a  $(2n+1)$ -fold loop space. We clearly have  $\pi_0(\Omega B(BD)) \cong \mathbb{Z}$ , and an application of the group completion theorem shows that the homology of each of the components is  $H_*(\Omega_0 B(BD)) \cong H_*(\text{hocolim}_g B\text{Diff}_\partial(U_{g,1}^n))$ ; hence Theorem A also evaluates the rational cohomology of  $\Omega_0 B(BD)$  in a range of degrees. In [38], a larger  $E_{2n+1}$ -algebra is considered, namely

$$\mathcal{M}_{2n+1} := \coprod_{[W]} B\text{Diff}_\partial(W),$$

where  $[W]$  ranges through all diffeomorphism classes of  $(n - 1)$ -connected and  $n$ -parallizable  $(2n + 1)$ -manifolds  $W$  with boundary  $S^{2n}$ . The main result of [38] is that the group completion of  $\mathcal{M}_{2n+1}$  has the homotopy type of an infinite loop space if  $n \geq 4$  and  $n \neq 7$  (this is an odd-dimensional version of a theorem by Tillmann [72] for surfaces). This is done by showing that  $\Omega B\mathcal{M}_{2n+1}$  is homotopy equivalent to the infinite loop space of a spectrum denoted  $\text{MT}\mathcal{L}_{2n+1}$ . The latter is not a Madsen–Tillmann spectrum despite the notation, but rather obtained from a certain cobordism category of manifolds equipped with certain subspaces of their homology by using infinite loop space machinery. That cobordism category does not fit into the general theory of cobordism categories as in [28]; there is however a map  $\text{MT}\mathcal{L}_{2n+1} \rightarrow \text{MT}\theta_{2n+1}^n$  of spectra.

Clearly  $BD \subset \mathcal{M}_{2n+1}$  is a union of path components. However, while  $\pi_0(BD) \cong \mathbb{N}_0$ ,  $\pi_0(\mathcal{M}_{2n+1})$  is much larger; [38, Proposition 3.2.5] deduces a description of  $\pi_0(\mathcal{M}_{2n+1})$  from [77]. It is therefore not clear how to relate the group completions of  $BD$  and of  $\mathcal{M}_{2n+1}$ . As Fabian Hebestreit and Manuel Krannich pointed out to us, it seems conceivable that the map  $\Omega_0 B(BD) \rightarrow \Omega_0 B\mathcal{M}_{2n+1}$  is a rational homology equivalence. If that turns out to be true, Theorem A computes the rational homology of  $\Omega_0^\infty \text{MT}\mathcal{L}_{2n+1}$  in a range of degrees.

**1.4. Method of proof.** Having said that the methods of [25] must fail in the odd-dimensional case we need to say how we approach Theorem A. There is an established three-stage procedure to describe the topology of  $B\text{Diff}_\partial(M)$  for a high-dimensional manifold ( $d = \dim(M) \geq 5$ ) in a range depending on  $d$ . The *first step* is to get a hold on  $B\text{hAut}_\partial(M)$ , the classifying space of the homotopy automorphisms of  $M$ , relative to the boundary. The *second step* uses Quinn’s space-level version of the surgery exact sequence [63] to compare  $B\text{hAut}_\partial(M)$  with the classifying space  $B\widetilde{\text{Diff}}_\partial(M)$  of the *block diffeomorphism* group; the difference is in terms of the  $L$ -theory of the group ring of  $\pi_1(M)$ . The *third step* compares block diffeomorphisms to diffeomorphisms in a range of degrees, in terms of algebraic  $K$ -theory.

Our strategy in this paper is to first compute  $H^*(B\widetilde{\text{Diff}}_\partial(U_{g,1}^n); \mathbb{Q})$  in the same range of degrees as in Theorem A (the first and second step can largely be merged), and then to use the comparison from the third step (which holds in a larger range

of degrees) to arrive at  $H^*(B\text{Diff}_\partial(U_{g,1}^n); \mathbb{Q})$ . Let us first note that the map (1.1) does not extend to block diffeomorphisms, hence the spectrum  $\text{MT}\theta_{2n+1}^n$  does not play an important role in the calculation. However, the tautological classes can be extended by [15]. To state our results property, let us introduce some notation. Define a graded vector space  $Z(d)$  as follows:

$$Z(d)_i = \begin{cases} 0 & i \leq 0 \\ H^{d+i}(BO; \mathbb{Q}) & i > 0. \end{cases}$$

The element of  $Z(d)_i$  corresponding to  $c \in H^{d+i}(BO; \mathbb{Q})$  is denoted  $k_c$ .

For each  $d$ -manifold  $M$ , we therefore obtain maps

$$\Phi_M : \mathbb{F}(Z(d)) \rightarrow H^*(B\text{Diff}_\partial^+(M); \mathbb{Q})$$

and

$$\tilde{\Phi}_M : \mathbb{F}(Z(d)) \rightarrow H^*(B\widetilde{\text{Diff}}_\partial^+(M); \mathbb{Q})$$

from the free graded-commutative algebra generated by  $Z(d)$ . Sending  $k_c$  to  $\mu_c \in H^*(\Omega_0^\infty \text{MT}\theta_{2n+1}^n; \mathbb{Q})$  also gives an algebra map

$$\Psi : \mathbb{F}(Z(2n+1)) \rightarrow H^*(\Omega_0^\infty \text{MT}\theta_{2n+1}^n; \mathbb{Q})$$

which is surjective and whose kernel is the ideal generated by all  $k_{L_m c}$  with  $4m \leq n$  and  $c \in H^*(BO; \mathbb{Q})$ , and by all  $k_c$  with  $c \in \ker(H^*(BO; \mathbb{Q}) \rightarrow H^*(BO(2n+1); \mathbb{Q}))$ . Moreover  $(\beta_g^n)^* \circ \Psi = \Phi_{U_{g,1}^n}$ . Therefore Theorem A is equivalent to the following result.

**Theorem 1.3.** *Assume that  $n \geq 5$ . Then  $\Phi_{U_{g,1}^n}$  is surjective in degrees  $* \leq \min(\frac{n-2}{2}, n-4)$ , and in that range of degrees, the kernel is the ideal generated by the following list of elements:*

$$k_{L_m} \text{ all } m, \tag{1.4}$$

$$k_{L_m c} \text{ } 4m \leq n, c \in H^*(BO; \mathbb{Q}), \tag{1.5}$$

$$k_{L_{m_0} L_{m_1}} \text{ } 4(m_0 + m_1) = 2n + 2. \tag{1.6}$$

To state our result concerning  $B\widetilde{\text{Diff}}_\partial(U_{g,1}^n)$ , let  $B$  be the graded vector space

$$B := \bigoplus_{k \geq 1} \mathbb{Q}[4k + 1]$$

and recall that  $H^*(B\text{GL}_\infty(\mathbb{Z}); \mathbb{Q}) \cong \mathbb{F}(B)$  by Borel's famous calculation [5]. The action of the block diffeomorphism group on  $H_n(U_{g,1}^n; \mathbb{Z}) \cong \mathbb{Z}^g$  gives a map

$$\mathbb{F}(B) \cong H^*(B\text{GL}_\infty(\mathbb{Z}); \mathbb{Q}) \rightarrow H^*(B\text{GL}_g(\mathbb{Z}); \mathbb{Q}) \rightarrow H^*(B\widetilde{\text{Diff}}_\partial(U_{g,1}^n); \mathbb{Q}).$$

Combining this with  $\tilde{\Phi}_{U_{g,1}^n}$ , we obtain an algebra map

$$\Gamma : \mathbb{F}(B \oplus Z(2n+1)) \rightarrow H^*(B\widetilde{\text{Diff}}_\partial(U_{g,1}^n); \mathbb{Q}), \tag{1.7}$$

and we will prove the following result.

**Theorem 1.8.** *Let  $n \geq 5$ . The map  $\Gamma$  is surjective in degree  $* \leq n-4$  and for  $g \gg n$ , and the kernel is spanned by the same elements as given in Theorem 1.3.*

Grey [30, Theorem B] has shown a homological stability result for block diffeomorphism groups which applies to the manifolds  $U_{g,1}^n$  and can be used to give an explicit lower bound for  $g$  for which Theorem 1.8 holds (a worse bound can also be deduced from our proofs).

As already said, the proof of Theorem 1.8 uses Quinn's theory, which roughly expresses the homotopy fibre of the forgetful map  $B\widetilde{\text{Diff}}_\partial(M) \rightarrow B\text{hAut}_\partial(M)$  in terms of  $L$ -theory which is quite manageable for simply connected  $M$ . We refer to [2] for a more informative survey; more importantly, that paper contains a consequence of Quinn's theory [2, Theorem 1.1] which allows us to use surgery theory completely as a black box. The result is that the calculation of  $H^*(B\widetilde{\text{Diff}}_\partial(U_{g,1}^n); \mathbb{Q})$  is equivalent to the calculation of

$$H^*(\text{map}_*(U_{g,1}^n; BO_{\mathbb{Q}})^0 // \text{hAut}_\partial(U_{g,1}^n); \mathbb{Q}), \quad (1.9)$$

up to some smallprint that we shall ignore for the moment ( $\text{map}_*(U_{g,1}^n; BO_{\mathbb{Q}})^0$  is the component of the pointed mapping space containing the constant map). The computation of (1.9) fills the largest portion of this paper. The component group  $\pi_0(\text{hAut}_\partial(U_{g,1}^n))$  is very close to  $\text{GL}_g(\mathbb{Z})$  so that naturally Borel's work on the cohomology of arithmetic group enters. This is one of the reasons why  $g$  needs to be large in Theorem 1.8. The cohomology of mapping spaces such as  $\text{map}_*(U_{g,1}^n; BO_{\mathbb{Q}})^0$  is fairly easy to compute, but rather large. To get from there to Theorem 1.8, we employ Borel's vanishing theorem in a similar way to its use in [16] or [46], and a calculation in classical invariant theory.

While in principle the general theory allows us to compute the cohomology of block diffeomorphisms in arbitrary degrees, we ran into several difficulties which we could only resolve in the range of degrees of Theorem 1.8.

The main results of [2] about  $H^*(B\widetilde{\text{Diff}}_\partial(W_{g,1}^n); \mathbb{Q})$  show that  $H^*(B\widetilde{\text{Diff}}_\partial(W_{g,1}^n); \mathbb{Q})$  behaves very differently from  $H^*(B\text{Diff}_\partial(W_{g,1}^n); \mathbb{Q})$  in high degrees, in the sense that it contains contributions from graph homology. Recently, Stoll [69] computed  $H^*(B\text{hAut}_\partial(U_{g,1}^n); \mathbb{Q})$  for  $* \leq g-2$ , and he found a similar contribution from graph homology, which will also be present in  $H^*(B\widetilde{\text{Diff}}_\partial(U_{g,1}^n); \mathbb{Q})$ . We invite the curious and capable reader to figure out  $H^*(B\widetilde{\text{Diff}}_\partial(U_{g,1}^n); \mathbb{Q})$  in a larger range, at least up to degree  $2n-6$ , which is the range of degrees in which the third step, the comparison of diffeomorphisms and block diffeomorphisms, is valid.

Let us now turn to this step. The result is stated in detail as Theorem 7.1 below; we have formulated it in such a way that it is largely independent of our computation of  $H^*(B\widetilde{\text{Diff}}_\partial(U_{g,1}^n); \mathbb{Q})$ . One feature is that the Borel classes coming from the action on homology vanish on  $H^*(B\text{Diff}_\partial(U_{g,1}^n); \mathbb{Q})$ . We deduce this from the Dwyer–Weiss–Williams index theorem [12] in Proposition 2.16. In general, the comparison of diffeomorphisms and block diffeomorphisms is in terms of pseudoisotopy theory and algebraic  $K$ -theory, with the stable h-cobordism theorem [75] and Igusa's stability theorem [41] as the main points; the last one enforces a bound depending on the dimension of the manifold. An elaborate formulation of this step was given by Weiss–Williams in [78], but a simpler variant suffices for us. Let us describe briefly how the comparison is done. Recall the classical result by Farrell–Hsiang [20] stating that

$$\pi_k(B\text{Diff}_\partial(D^{2n+1})) \otimes \mathbb{Q} \cong \begin{cases} \mathbb{Q} & k \equiv 0 \pmod{4} \\ 0 & k \not\equiv 0 \pmod{4} \end{cases}$$

holds in a range of degrees. This range is nowadays known to be roughly  $2n$ , by recent work of Krannich [44] and Krannich–Randal-Williams [43]. An instance of Morlet’s lemma of disjunction states that the homotopy fibre of

$$B\text{Diff}_\partial(U_{g,1}^n) \rightarrow \widetilde{B\text{Diff}}_\partial(U_{g,1}^n) \tag{1.10}$$

is rationally equivalent to  $B\text{Diff}_\partial(D^{2n+1})$  up to degree approximately  $2n$ , and the main result of [14] (which is a consequence of [7] and [59]) says that the inclusion of the homotopy fibre into the total space of (1.10) induces the trivial map on rational homology in a range of degrees. To combine those facts, we use an elementary argument borrowed from Krannich’s paper [44] to show that the fibration (1.10) is “plus-constructible”.

In [44], the comparison map

$$B\text{Diff}_D(V_g^n) \rightarrow \widetilde{B\text{Diff}}_D(V_g^n) \tag{1.11}$$

is studied for the boundary connected sum  $V_g^n$  of  $g$  copies of  $S^n \times D^{n+1}$ , where  $D$  is a fixed disc in  $\partial V_g^n$ . The homotopy fibre of (1.11) is also rationally equivalent to  $B\text{Diff}_\partial(D^{2n+1})$ . Krannich used knowledge about  $B\text{Diff}_D(V_g^n)$  from [7] and  $\widetilde{B\text{Diff}}_D(V_g^n)$  from surgery theory to deduce knowledge about the homotopy fibre of (1.11) (and thereby extends the range in Farrell-Hsiang’s theorem to roughly  $2n$ ). In the present paper, the logic is reversed: we use knowledge about the base of (1.10) from surgery theory and the fibre from Farrell-Hsiang’s theorem (or Krannich’s improvement thereof) to deduce knowledge about the total space.

**1.5. Overview of the chapters.** To navigate the reader through this rather long paper, let us briefly describe the content of the chapters. §2: after setting up notation, we introduce the tautological classes and show in §2.3 the three vanishing theorems implied by Theorems A and 1.8 (one of the vanishing results is needed for the proof, the other two fall out as byproducts). In §2.4, we prove another vanishing result, namely Proposition 2.16 which says that the Borel classes on  $\widetilde{B\text{Diff}}_\partial(U_{g,1}^n)$  vanish when pulled back to  $B\text{Diff}_\partial(U_{g,1}^n)$ ; this is a fairly straightforward consequence of the Dwyer–Weiss–Williams theorem.

§3 is about the general theory behind the proof of Theorem 1.8. The goal is Proposition 3.25 which essentially gives a formula for the rational cohomology of  $B\text{Diff}_\partial(M)$  under some hypotheses on  $M$ , which are satisfied by  $U_{g,1}^n$ . The constituents for this formula are  $\text{hAut}_\partial(M_{\mathbb{Q}})$ , the homotopy automorphisms (relative to the boundary) of the rationalization of  $M$ , and the mapping space  $\text{map}_\partial(M; BO_{\mathbb{Q}})$ . This is derived from surgery theory, through a result of Berglund–Madsen [2]. §3.1 reviews some generalities on rational homotopy theory, and §3.3 describes the cohomology of mapping spaces of the above sort.

In §4, we begin to apply this theory to the manifolds  $U_{g,1}^n$ . Many of the results are exercises in standard homotopy theory (and already contained in [30]). The goal is Proposition 4.32, which computes the  $E_2$ -page of the spectral sequence of the fibration  $\text{map}_\partial(U_{g,1}^n; BO_{\mathbb{Q}})^0 // \text{hAut}_\partial(U_{g,1}^n)^{\text{id}} \rightarrow \text{BhAut}_\partial(U_{g,1}^n)^{\text{id}}$  (in a range of degrees, and including the  $d_2$ -differential).

Having determined the  $E_2$ -page of the spectral sequence, our tactical goal is to calculate the  $\pi_0(\text{hAut}_\partial(U_{g,1}^n))$ -invariant part of the  $E_\infty$ -page. For this, we need a calculation in classical invariant theory which we carry out in the purely algebraic §5. In §6, we eventually complete the proof of Theorem 1.8, with a use of Borel’s

vanishing theorem [6]. The rather short section §7 derives Theorem A from Theorem 1.8 along the lines sketched above.

*Notations.* We use the following standard notations in this paper. For two  $\mathbb{N}_0$ -graded algebras  $A$  and  $B$ ,  $A \otimes B$  always denotes the *graded* tensor product, to conform the conventions of homological algebra. The free graded-commutative algebra generated by a graded vector space  $V$  is denoted  $\mathbb{F}(V)$ . If  $V$  is a vector space and  $n \in \mathbb{N}_0$ , we let  $V[n]$  be the graded vector space which is  $V$  in degree  $n$  and 0 otherwise (and not the degree shift of a graded vector space). For a module  $N$  over a commutative ring  $R$ , we write  $N^\vee := \text{Hom}_R(N; R)$  for the dual.

**1.6. Acknowledgements.** The authors would like to thank Alexander Berglund, Fabian Hebestreit, Lutz Hille, Manuel Krannich, Sander Kupers, Ib Madsen, Thomas Nikolaus and Oscar Randal-Williams for helpful conversations about various aspects of this work. We also thank Andrea Bianchi for his careful reading of the first version of this paper. Last but not least, it is a pleasure to thank Jerzy Weyman. Without his help, we would not have been able to carry out the crucial invariant-theoretic calculation in Proposition 5.7, and would not have been able to get this project to a conclusion.

## 2. CHARACTERISTIC CLASSES OF SMOOTH AND BLOCK BUNDLES

**2.1. Automorphism groups.** Let us first establish some notation. Let  $M^d$  be a compact oriented smooth manifold with boundary. We write  $\text{Diff}(M)$  for the diffeomorphism group and  $\text{Diff}_\partial(M) \subset \text{Diff}(M)$  for the subgroup of diffeomorphisms which are the identity near  $\partial M$ . We furthermore let  $\text{Diff}^+(M) \subset \text{Diff}(M)$  and  $\text{Diff}_\partial^+(M) \subset \text{Diff}_\partial(M)$  the subgroups of orientation-preserving diffeomorphisms; note that  $\text{Diff}_\partial^+(M) = \text{Diff}_\partial(M)$  if the inclusion  $\partial M \rightarrow M$  is 0-connected.

In the present paper, we make heavy use of the *block diffeomorphism* group  $\widetilde{\text{Diff}}(M)$ . We won't repeat the definition here and refer instead to §1 and 2 of [44] for an up-to-date exposition. There are block analogues  $\widetilde{\text{Diff}}_\partial(M)$ ,  $\widetilde{\text{Diff}}^+(M)$ ,  $\widetilde{\text{Diff}}_\partial^+(M)$  of respective diffeomorphism groups. Let us also note that the natural map

$$I : \text{Diff}(M) \rightarrow \widetilde{\text{Diff}}(M)$$

is *by definition* 0-connected; the same holds for the decorated versions.

We shall need various flavours of homotopy automorphism groups. When forming mapping spaces, we secretly replace all spaces that occur by their singular simplicial set, and view the mapping space as a simplicial set. For a CW-pair  $(X, A)$  and a pointed space  $Y$ , we let  $\text{map}_A(X; Y)$  be the space of maps  $X \rightarrow Y$  whose restriction to  $A$  is the constant map to the basepoint in  $Y$  (or more formally the fibre of the restriction map  $\text{map}(X; Y) \rightarrow \text{map}(A; Y)$  over the constant map). For a CW-pair  $(X, A)$ , we let  $\text{hAut}_A(X)$  be the monoid of all homotopy self-equivalences of  $X$  which are the identity on  $A$ . For an oriented compact manifold, we let also  $\text{hAut}_\partial(M)^+ \subset \text{hAut}_\partial(M)$  the submonoid of those self-equivalences which preserve the fundamental homology class. There are natural maps

$$\widetilde{\text{Diff}}_\partial(M) \rightarrow \text{hAut}_\partial(M) \text{ and } \widetilde{\text{Diff}}_\partial^+(M) \rightarrow \text{hAut}_\partial^+(M)$$

(or rather a zig-zag, see the discussion in [2, p. 98 f] for more details).

Assume that  $V \rightarrow X$  is a vector bundle and that  $C \subset A$  is a subcomplex. We let  $\text{hAut}_A^C(V)$  be the monoid of all pairs  $(f, \hat{f})$  where  $f \in \text{hAut}_A(X)$  and  $\hat{f} : V \rightarrow V$  is a bundle map covering  $f$  which is fibrewise an isomorphism, and such that  $f|_{V|_C}$  is the identity, see [2, p. 107f] for more details. There is a stable version of that construction given on p. 110 loc.cit.; we define

$$\text{hAut}_A^C(V)^{\text{st}} := \text{colim}_k \text{hAut}_A^C(V \oplus \mathbb{R}^k).$$

All the monoids we just introduced are grouplike  $E_1$ -spaces and therefore admit classifying spaces. The classifying space  $\text{BhAut}_A^C(V)^{\text{st}}$  has a convenient description as follows. If  $V|_C$  is stably trivial and a stable trivialization is chosen, Proposition 4.13 of [2] provides a weak equivalence

$$\text{BhAut}_A^C(V)^{\text{st}} \simeq (\text{map}_C(X; BO) // \text{hAut}_A(X))_V, \quad (2.1)$$

where

$$(\text{map}_C(X; BO) // \text{hAut}_A(X))_V \subset \text{map}_C(X; BO) // \text{hAut}_A(X)$$

denotes the connected component determined by a fixed classifying map  $\lambda : X \rightarrow BO$  of  $V$  which extends the given trivialization of  $V|_C$ , viewed as a point in  $\text{map}_C(X; BO)$ . The map in (2.1) arises as follows: the total space of the universal fibration over  $\text{BhAut}_A^C(V)^{\text{st}}$  with fibre  $X$  carries a stable vector bundle which is built from  $V$ . For a detailed construction on the point-set level, we refer to [2]. When  $V$  is stably trivial, (2.1) can be reformulated as follows.

**Lemma 2.2.** *Assume that  $V \rightarrow X$  is stably trivial, and that a stable trivialization of  $V|_C$  is fixed. Then there is a weak equivalence*

$$\text{BhAut}_A^C(V)^{\text{st}} \simeq \text{map}_C(X; BO)^0 // \text{hAut}_A(X),$$

where  $\text{map}_C(X; BO)^0 \subset \text{map}_C(X; BO)$  denotes the component of the constant map.

*Proof.* According to (2.1), we must prove that

$$(\text{map}_C(X; BO) // \text{hAut}_A(X))_V \simeq \text{map}_C(X; BO)^0 // \text{hAut}_A(X)$$

if  $V$  is stably trivial. Now  $\pi_0(\text{map}_C(X; BO)) = KO^0(X, C)$  is an abelian group and  $\text{hAut}_A(X)$  acts by group automorphisms, and therefore fixes the neutral element. So  $\text{map}_C(X; BO)^0 // \text{hAut}_A(X)$  is the component of  $\text{map}_C(X; BO) // \text{hAut}_A(X)$  containing the constant map, which is exactly  $(\text{map}_C(X; BO) // \text{hAut}_A(X))_V$ .  $\square$

One important feature of block diffeomorphisms is the existence of the *derivative map*

$$D : \widetilde{\text{Diff}}_{\partial}(M) \rightarrow \text{hAut}_{\partial}^{\partial}(TM)^{\text{st}}, \quad (2.3)$$

which is a map of  $E_1$ -monoids and can therefore be delooped to a map

$$BD : B\widetilde{\text{Diff}}_{\partial}(M) \rightarrow \text{BhAut}_{\partial}^{\partial}(TM)^{\text{st}}. \quad (2.4)$$

The derivative map is constructed in [2, §4.3], see also [44, §1.9], and is an expression of the fact, first proven in [15] and expanded on in [37, §2.4], that block bundles have a stable vertical tangent bundle. By virtue of its definition, the derivative map is a map over  $\text{hAut}_{\partial}(M)$ . If  $M$  is stably parallelizable, we can use Lemma 2.2 to rewrite (2.4) in the form

$$BD : B\widetilde{\text{Diff}}_{\partial}(M) \rightarrow \text{map}_{\partial}(M; BO)^0 // \text{hAut}_{\partial}(M). \quad (2.5)$$

**2.2. Tautological classes.** Tautological classes (aka Miller–Morita–Mumford classes or  $\kappa$ -classes) for block bundles have been constructed in [15] and more systematically in [37]. The most streamlined construction can be given using the derivative map (2.4), and we sketch the definition briefly, in a level of generality that will prove to be useful for us later on.

*Construction 2.6.* Assume that  $M^d$  is a compact oriented smooth manifold with boundary. Consider the universal fibration pair  $(E, \partial E)$  with fibre  $(M, \partial M)$  over the space  $B := \text{map}_\partial(M; BO) // \text{hAut}_\partial^+(M)$ . The evaluation map

$$\text{ev} : \text{map}_\partial(M; BO) \times M \rightarrow BO$$

is  $\text{hAut}_\partial(M)$ -equivariant and maps  $\text{map}_\partial(M; BO) \times \partial M$  to the basepoint. As  $E = (\text{map}_\partial(M; BO) \times M) // \text{hAut}_\partial^+(M)$  and  $\partial E = (\text{map}_\partial(M; BO) \times \partial M) // \text{hAut}_\partial^+(M)$ ,  $\text{ev}$  induces a map  $\epsilon : E \rightarrow BO$  sending  $\partial E$  to the basepoint. We may think of  $\epsilon$  as a stable vector bundle on  $E$ , trivialized on  $\partial E$ . The Leray–Serre spectral sequence for the fibration pair  $(E, \partial E) \rightarrow B$  yields maps

$$\pi_! : H^k(E; \partial E) \rightarrow E_\infty^{k-d, d} \subset E_2^{k-d, d} \cong H^{k-d}(B; H^d(M; \partial M)) \rightarrow H^{k-d}(B)$$

(with coefficients in an arbitrary ring). Given a class  $c \in H^k(BO)$ , we can therefore form

$$\kappa_c := \pi_!(\epsilon^* c) \in H^{k-d}(\text{map}_\partial(M; BO) // \text{hAut}_\partial^+(M)).$$

*Construction 2.7.* Let  $M^d$  be an oriented compact smooth manifold with boundary and assume that  $TM|_{\partial M}$  is stably trivial. Combining the derivative map  $BD$  with (2.1) yields a map

$$\widetilde{BDiff}_\partial^+(M) \rightarrow B\text{hAut}_\partial^\partial(TM)^{\text{st}} \rightarrow \text{map}_\partial(M; BO) // \text{hAut}_\partial^+(M),$$

and we can pull back  $\kappa_c$  to a class, also denoted  $\kappa_c \in H^{k-d}(\widetilde{BDiff}_\partial^+(M))$ . Pulling this further back along  $BI : \widetilde{BDiff}_\partial(M) \rightarrow \widetilde{BDiff}_\partial^+(M)$ , we obtain the usual  $\kappa$ -classes on classifying spaces of diffeomorphism groups.

A map  $\gamma$  from a space  $X$  to one of the classifying spaces  $\widetilde{BDiff}_\partial^+(M)$ ,  $\widetilde{BDiff}_\partial^+(M)$  or  $\text{map}_\partial(M; BO) // \text{hAut}_\partial^+(M)$  classifies  $E \rightarrow X$ , which is a smooth fibre bundle / block bundle / fibration with a stable vector bundle on its total space. In such situations, we use the suggestive notation  $\kappa_c(E) := \gamma^* \kappa_c$ .

We now specialize to rational coefficients. Recall that  $H^*(BO; \mathbb{Q})$  is the polynomial algebra in the Pontrjagin classes. For our purposes, it is useful to observe that one can also write

$$H^*(BO; \mathbb{Q}) \cong \mathbb{Q}[L_1, L_2, \dots],$$

where  $L_m \in H^{4m}(BO; \mathbb{Q})$  is the  $m$ th component of the Hirzebruch  $L$ -class. The above is true by the well-known fact [40, p. 14] that the coefficient  $a_m$  of  $p_m$  in  $L_m$  is nonzero.

**2.3. Some vanishing theorems for tautological classes.** In this subsection, we review the three vanishing theorems that are entailed by Theorem A. Only one of them (Theorem 2.10) is actually used in the proof of our main theorem; the other two (Proposition 2.9 and Proposition 2.12) are only stated for sake of completeness; the fact that they are valid for the manifolds  $U_{g,1}^n$  is a byproduct of our computations below.

*An additivity property.* Let us begin with a fact which is well-known for diffeomorphism groups; the argument we give is essentially contained in [54], [56].

**Lemma 2.8.** *Let  $M$  and  $N$  be compact oriented  $d$ -manifolds with boundary with a common (closed) part  $\partial_0 \subset \partial M, \partial N$  of their boundary. Let*

$$\begin{aligned} \mu : \text{map}_\partial(M; BO) // \text{hAut}_\partial^+(M) \times \text{map}_\partial(N; BO) // \text{hAut}_\partial^+(N) &\rightarrow \\ &\rightarrow \text{map}_\partial(M \cup_{\partial_0} N; BO) // \text{hAut}_\partial^+(M \cup_{\partial_0} N) \end{aligned}$$

*be the obvious gluing map, let  $p_M$  be the projection to  $\text{map}_\partial(M; BO) // \text{hAut}_\partial^+(M)$ , and define  $p_N$  similarly. Then for each  $c \in H^k(BO)$  with  $k > d$ , we have*

$$\mu^* \kappa_c = p_M^* \kappa_c + p_N^* \kappa_c.$$

*Proof.* We must show the following: assume that  $\pi^E : E \rightarrow X$  and  $\pi^F : F \rightarrow X$  are two oriented fibrations with fibres  $M$  and  $N$  and trivialized boundaries, and that  $E$  and  $F$  contain a common part  $A \subset \partial E, \partial F$  of their boundary. Then for two vector bundles  $V \rightarrow E$  and  $W \rightarrow F$  of the same rank  $n$  which are trivialized over the respective boundaries and each  $c \in H^k(BO)$ , we have

$$\pi_1^{E \cup_A F}(c(V \cup W)) = \pi_1^E(c(V)) + \pi_1^F(c(W)) \in H^{k-d}(X),$$

provided that  $k - d > 0$ . We may suppose that  $X$  is a finite CW complex. It is an exercise in linear algebra to prove an isomorphism

$$(V \cup_A \underline{\mathbb{R}}_F^n) \oplus (\underline{\mathbb{R}}_E^n \cup_A W) \cong (V \cup_A W) \oplus (\underline{\mathbb{R}}_E^n \cup_A \underline{\mathbb{R}}_F^n)$$

of clutched bundles on  $E \cup_A F$  (hint: picking bundle maps  $f : \underline{\mathbb{R}}_F^n \rightarrow V$  and  $g : \underline{\mathbb{R}}_E^n \rightarrow W$  which are the identity on the boundary is a good start). Hence the classifying map  $\gamma_{E \cup_A F} : E \cup_A F \rightarrow BO$  of  $V \cup_A W$  can be factored as

$$E \cup_A F \rightarrow E/\partial \vee F/\partial \xrightarrow{\gamma^E \vee \gamma^F} BO \vee BO \xrightarrow{\text{fold}} BO.$$

The claim follows from the observation that

$$H^*(E, \partial E) \cong H^*(E \cup_A F, \partial E \cup F) \rightarrow H^*(E \cup_A F, \partial(E \cup_A F)) \xrightarrow{\pi_1^{E \cup_A F}} H^{*-d}(X)$$

is nothing else than  $\pi_1^E$  (and the similar fact for  $F$  in place of  $E$ ).  $\square$

The next result is fairly simple-minded; compare [25, Lemma 7.16] for a more elaborate, but closely related result about diffeomorphism groups.

**Proposition 2.9.** *Let  $M$  be a compact oriented  $d$ -manifold, assume that  $TM|_{\partial M}$  is stably trivial, that  $\partial M \rightarrow M$  is  $(k-1)$ -connected and assume that all rational Pontrjagin classes of  $TM$  up to degree  $k$  are zero. Then*

$$\kappa_c = 0 \in H^{|\kappa| - d}(\widetilde{BDiff}_\partial^+(M); \mathbb{Q})$$

*whenever  $c \in H^*(BO; \mathbb{Q})$  lies in the ideal generated by  $\bigoplus_{0 < j \leq k} H^j(BO; \mathbb{Q})$ .*

*Proof.* The cohomological Leray–Serre spectral sequence for the universal block bundle pair  $(E, \partial E)$  over  $(\widetilde{BDiff}_\partial^+(M), *)$  reads as follows:

$$E_2^{p,q} = H^p(\widetilde{BDiff}_\partial^+(M), *; H^q(M, \partial M; \mathbb{Q})) \Rightarrow H^{p+q}(E, M \cup \partial E; \mathbb{Q}).$$

The connectivity assumption on  $M$  prove that  $H^*(E, M \cup \partial E; \mathbb{Q}) = 0$  whenever  $* \leq k$ . Hence  $H^*(E; \mathbb{Q}) \rightarrow H^*(M \cup \partial E; \mathbb{Q})$  is injective for  $* \leq k$ . The Mayer–Vietoris sequence moreover proves that  $H^*(M \cup \partial E; \mathbb{Q}) \rightarrow H^*(M; \mathbb{Q}) \oplus H^*(\partial E; \mathbb{Q})$  is also injective for  $* \leq k$ . The Pontrjagin classes of  $T_v E$  go to  $(p(TM), \text{pr}_{\partial M}^* p(TM)) = 0$  and therefore vanish in degrees  $\leq k$ . Hence  $c(T_v E) = 0$ , and a fortiori  $\kappa_c = 0$ .  $\square$

*The family signature theorem.* The classical family signature theorem for smooth fibre bundles (which uses families of elliptic operators in its proof) holds more generally for block bundles, as shown by Randal–Williams in [66, Theorem 2.6]. It has two cases, the odd-dimensional and the even-dimensional case. The odd case reads as follows.

**Theorem 2.10.** *Let  $M^d$  be an odd-dimensional oriented manifold, and assume for simplicity that  $TM|_{\partial M}$  is stably trivial. Then*

$$\kappa_{L_m} = 0 \in H^{4k-d}(\widetilde{BDiff}_\partial^+ M; \mathbb{Q})$$

for each  $m \in \mathbb{N}$  (and hence the same is true for diffeomorphisms).

*References.* By Lemma 2.8, it is enough to show the theorem for closed  $M$ , and this is done in [66, Theorem 2.6] for block diffeomorphisms, and in [13] for diffeomorphisms.  $\square$

Let us state the even-dimensional case for sake of completeness, and only for the case where  $M$  is closed or  $\partial M$  is a sphere. The point is that in the case  $\dim(M) = 2n$ , the action of  $\mathrm{hAut}_\partial^+(M)$  on  $H_n(M; \mathbb{Q})$  preserves the (nondegenerate) intersection form  $I_M$ . This fact produces a map  $f : \mathrm{BhAut}_\partial^+(M) \rightarrow \mathrm{BAut}(H_n(M; \mathbb{Q}); I_M)$ ; the latter is the classifying space of a symplectic or an orthogonal group, depending on the parity of  $n$ . Randal–Williams defines classes  $\sigma_i \in H^i(\mathrm{BAut}(H_n(M; \mathbb{Q}); I_M); \mathbb{Q})$ , which live in degrees  $i \equiv 2 \pmod{4}$  if  $n$  is odd and  $i \equiv 0 \pmod{4}$  if  $n$  is even, and shows in [66, Theorem 2.6] that

$$\kappa_{L_m} = f^* \sigma_{4m-2n} \in H^{4m-2n}(\widetilde{BDiff}_\partial^+(M); \mathbb{Q}). \quad (2.11)$$

*Consequence of the family signature theorem.*

**Proposition 2.12.** *Let  $M$  be a compact oriented  $d$ -manifold, assume that  $TM|_{\partial M}$  is stably trivial, and suppose that all rational Pontrjagin classes of  $M$  are trivial. Then for each  $c \in H^k(BO; \mathbb{Q})$ ,  $k > \dim(M)$ , the homomorphism*

$$\pi_{k-d}(\widetilde{BDiff}_\partial^+(M)) \rightarrow \mathbb{Q}$$

given by

$$[f] \mapsto \langle f^* \kappa_c; [S^{k-d}] \rangle$$

is the zero map. In particular, if  $k = d + 1$ , then  $\kappa_c = 0 \in H^1(\widetilde{BDiff}_\partial^+(M); \mathbb{Q})$ .

*Proof.* For diffeomorphisms, this is a well-known fact, see e.g. [45, Proposition 13] or [33, Proposition 1.9]. To see that the proof also applies to block diffeomorphisms, we review the argument.

Firstly, the double  $M \cup_{\partial M} M$  has trivial rational Pontrjagin classes, by an argument given in the proof of [17, Theorem F]; hence by Lemma 2.8 it suffices to give the argument for closed  $M$ .

Let  $k = 4m$  and let  $\pi : E \rightarrow S^{4m-d}$  be an oriented block bundle with fibre  $M$ , classified by  $f : S^{4m-d} \rightarrow \widetilde{BDiff}_\partial^+(M)$ . The restriction of the stable vertical tangent bundle  $T_v E$  to the fibre  $M$  over  $*$  is stably isomorphic to  $TM$ . A brief inspection of the Leray–Serre spectral sequence of  $\pi$  proves that  $c(T_v E) = 0$  if  $c$  can be written as a product of classes in positive degrees. Hence we only have to prove that  $\kappa_{L_m}(E) = 0$ . This follows from [66, Theorem 3.1] (if  $d$  is even, we only need to

consider bundles over spheres of even dimension, which are simply-connected and (2.11) shows that  $\kappa_{L_m}$  is pulled back from an aspherical space).

The last sentence follows by the Hurewicz theorem.  $\square$

**2.4. Borel classes.** Let  $K(\mathbb{Z})$  denote the algebraic  $K$ -theory spectrum of the integers; recall that  $\pi_0(K(\mathbb{Z})) \cong \mathbb{Z}$  and that  $\Omega_0^\infty K(\mathbb{Z}) \simeq BGL_\infty(\mathbb{Z})^+$ . A celebrated result of Borel describes the rational cohomology of the latter space.

**Theorem 2.13** (Borel).

- (1) *The rational cohomology  $H^*(\Omega_0^\infty K(\mathbb{Z}); \mathbb{Q}) = H^*(BGL_\infty(\mathbb{Z}); \mathbb{Q})$  is the exterior algebra with generators  $\beta_{4k+1} \in H^{4k+1}(\Omega_0^\infty K(\mathbb{Z}); \mathbb{Q})$ ,  $k \geq 1$ . The classes  $\beta_{4k+1}$  are primitive.*
- (2) *The restriction maps*

$$H^p(BGL_\infty(\mathbb{Z}); \mathbb{Q}) \rightarrow H^p(BGL_g(\mathbb{Z}); \mathbb{Q}) \rightarrow H^p(BSL_g(\mathbb{Z}); \mathbb{Q})$$

*are isomorphisms provided that  $g \geq 2p + 2$ .*

- (3) *The group homomorphism  $\kappa : GL_g(\mathbb{Z}) \rightarrow GL_g(\mathbb{Z})$  given by  $\kappa(x) := (x^\top)^{-1}$  has the following effect on these classes:*

$$(B\kappa)^* \beta_{4k+1} = -\beta_{4k+1}.$$

*References.* (1) is of course a famous theorem of Borel [5] (he treats real cohomology which makes little difference as the cohomology spaces are all finite-dimensional by [62]). (2) A range in which the map from the stable cohomology to the unstable cohomology is an isomorphism is also determined in Borel's paper; the range as stated follows from Van der Kallen's homological stability theorem [74, Theorem 4.11], using that the Bass stable rank of the integers is 2; see [32, §4.1.11]. (3) can also easily be deduced from Borel's work. Since we do not know a reference, we shall indicate the proof here. It suffices to prove the statement for  $SL_g(\mathbb{Z})$  instead of  $GL_g(\mathbb{Z})$ , since the covering map  $BSL_g(\mathbb{Z}) \rightarrow BGL_g(\mathbb{Z})$  induces an injection in rational cohomology. We need to recall how the Borel classes are constructed. Let  $X$  be the symmetric space  $SL_g(\mathbb{R})/SO(g)$  and let  $\mathcal{A}^*(X)^{SL_g(\mathbb{R})}$  be the chain complex of invariant differential forms, which has trivial differential as each  $SL_g(\mathbb{R})$ -invariant differential form on  $X$  is closed, by a general fact about symmetric spaces. On the other hand,  $X$  is contractible and the  $SL_g(\mathbb{Z})$ -action is proper, so that there is a natural isomorphism

$$H^*(BSL_g(\mathbb{Z}); \mathbb{R}) \cong H^*(\mathcal{A}^*(X)^{SL_g(\mathbb{Z})}).$$

On  $X$ , there is the Cartan involution  $\tau : X \rightarrow X$ ,  $\tau(xSO(g)) := (x^\top)^{-1}SO(g)$ . It is easily verified that  $\tau$  induces an involution on  $\mathcal{A}^*(X)^{SL_g(\mathbb{R})}$ , and that the diagram

$$\begin{array}{ccc} \mathcal{A}^*(X)^{SL_g(\mathbb{R})} & \longrightarrow & H^*(BSL_g(\mathbb{Z}); \mathbb{R}) \\ \downarrow \tau^* & & \downarrow B\iota^* \\ \mathcal{A}^*(X)^{SL_g(\mathbb{R})} & \longrightarrow & H^*(BSL_g(\mathbb{Z}); \mathbb{R}) \end{array}$$

commutes. By definition, the Borel classes come from certain invariant forms on  $X$ . It is therefore enough to show that  $\tau^* : \mathcal{A}^p(X)^{SL_g(\mathbb{R})} \rightarrow \mathcal{A}^p(X)^{SL_g(\mathbb{R})}$  is multiplication by  $(-1)^p$ . The involution  $\tau$  fixes the basepoint  $o := SO(g) \in X$ , and since invariant forms on  $X$  are determined by their values at  $o$ , it is enough to check that  $T_o\tau = -1$ . But this is easily verified by a direct calculation.  $\square$

Now let  $X$  be a finite CW-complex. For each  $p$ , the action on  $H_p(X; \mathbb{Z})$  provides a map

$$B\rho_p : \text{BhAut}(X) \rightarrow \text{BGL}(H_p(X; \mathbb{Z})).$$

For each finitely generated abelian group  $A$  with torsion subgroup  $TA$ , there is a map

$$\iota : \text{BGL}(A) \rightarrow \text{BGL}(A/TA) \rightarrow \Omega^\infty K(\mathbb{Z})$$

well defined up to homotopy; it hits the component of  $\text{rank}(A) \in \mathbb{Z} = \pi_0(K(\mathbb{Z}))$ . Composing  $\iota$  and  $B\rho_p$  gives classes

$$\beta_{4k+1}^p := (\iota \circ B\rho_p)^* \beta_{4k+1} \in H^{4k+1}(\text{BhAut}(X); \mathbb{Q}).$$

Using the infinite loop space structure on  $\Omega^\infty K(\mathbb{Z})$ , we can form the alternating sum

$$\chi := \sum_{p \geq 0} (-1)^p \iota \circ B\rho_p : \text{BhAut}(X) \rightarrow \Omega_{\chi(X)}^\infty K(\mathbb{Z}) \simeq \Omega_0^\infty K(\mathbb{Z}),$$

the *algebraic K-theory Euler characteristic*. Because the Borel classes are primitive, the relation

$$\chi^* \beta_{4k+1} = \sum_{p \geq 0} (-1)^p \beta_{4k+1}^p \in H^{4k+1}(\text{BhAut}(X); \mathbb{Q})$$

holds.

**Theorem 2.14** (Dwyer–Weiss–Williams). *Let  $M$  be a smooth compact manifold, possibly with boundary. Then*

$$\chi^* \beta_{4k+1} = 0 \in H^{4k+1}(B\text{Diff}(M); \mathbb{Q}).$$

Let us remark that the analogue of Theorem 2.14 for block diffeomorphism groups is *false*; in fact our computation of  $H^*(B\widetilde{\text{Diff}}_\partial(U_{g,1}^n); \mathbb{Q})$  certifies its failure.

*References.* The Dwyer–Weiss–Williams index theorem [12, Corollary 8.12] shows that the map  $\chi$  factors through the Becker–Gottlieb transfer  $B\text{Diff}(M) \rightarrow QS^0$ , so that the result simply follows from Serre’s finiteness theorem. See also §2 of [14] for a more detailed summary.  $\square$

We now use Poincaré duality to deduce a sharper vanishing theorem from Theorem 2.14.

**Lemma 2.15.** *Let  $M^d$  be a connected smooth oriented manifold with boundary, and suppose that  $\partial M = S^{d-1}$ . Then*

$$\beta_{4k+1}^p = -\beta_{4k+1}^{d-p} \in H^{4k+1}(B\text{Diff}_\partial^+(M); \mathbb{Q})$$

for all  $p$ .

*Proof.* We can consider  $\hat{M} := M \cup_{\partial M} D^d$  instead, without changing the Borel classes. Poincaré duality, the universal coefficient theorem and Theorem 2.13 (3) proves the claim.  $\square$

**Proposition 2.16.** *Let  $M$  be as in Lemma 2.15. If  $d = 2n$ , then*

$$\beta_{4k+1}^n = 0 \in H^{4k+1}(B\text{Diff}_\partial^+(M); \mathbb{Q}).$$

If  $d = 2n + 1$ , then

$$\sum_{p=0}^n (-1)^p \beta_{4k+1}^p = 0 \in H^{4k+1}(B\text{Diff}_\partial^+(M); \mathbb{Q}).$$

*Proof.* For  $d = 2n$ , compute

$$\begin{aligned} 0 &\stackrel{(2.14)}{=} \sum_{p=0}^{n-1} \left( (-1)^p \beta_{4k+1}^p + (-1)^{2n-p} \beta_{4k+1}^{2n-p} \right) + (-1)^n \beta_{4k+1}^n = \\ &\stackrel{(2.15)}{=} \sum_{p=0}^{n-1} (-1)^p \left( \beta_{4k+1}^p - \beta_{4k+1}^p \right) + (-1)^n \beta_{4k+1}^n = (-1)^n \beta_{4k+1}^n. \end{aligned}$$

For  $d = 2n + 1$ , compute

$$\begin{aligned} 0 &\stackrel{(2.14)}{=} \sum_{p=0}^n \left( (-1)^p \beta_{4k+1}^p + (-1)^{2n+1-p} \beta_{4k+1}^{2n+1-p} \right) = \\ &\stackrel{(2.15)}{=} \sum_{p=0}^n \left( (-1)^p \beta_{4k+1}^p + (-1)^{2n+1-p+1} \beta_{4k+1}^p \right) = \\ &2 \sum_{p=0}^n (-1)^p \beta_{4k+1}^p. \end{aligned}$$

□

*Remark 2.17.* The even-dimensional case of Proposition 2.16 can be shown directly from Borel's work, without recourse to [12]. The point is that by Poincaré duality,  $B\rho_n$  factors through the symplectic group or through an orthogonal group of some signature, depending on the parity of  $n$ , and Borel also computed the stable rational cohomologies of such groups: they are concentrated in even degrees.

### 3. RATIONAL COHOMOLOGY OF BLOCK DIFFEOMORPHISM SPACES: GENERAL THEORY

We shall approach  $H^*(B\widetilde{\text{Diff}}_\partial(U_{g,1}^n); \mathbb{Q})$  by the surgery-theoretic approach to the topology of diffeomorphism groups which is due to Quinn [63]; a detailed exposition is available in [57]. For our purposes, work of Berglund and Madsen [2, §4] enables us to treat all the surgery theory as a black box. Our aim in this section is to reformulate the results of [2, §4] in a way which is readily applicable to  $U_{g,1}^n$ . Our goal is Proposition 3.25 below.

**3.1. Some words about rational homotopy theory.** Let us recall some notions and results from rational homotopy theory. For us, a *space* will be a Kan complex. The category of spaces is denoted  $\mathcal{S}$ , and the category of pointed spaces by  $\mathcal{S}_*$ . Recall that the category  $\mathcal{S}$  is enriched over itself.

We say that a space  $X$  is *finite* if the geometric realization  $|X|$  is homotopy equivalent to a finite CW complex.

A map  $f : X \rightarrow Y$  is a *HQ-equivalence* if the induced map  $f_* : H_*(X, \mathbb{Q}) \rightarrow H_*(Y, \mathbb{Q})$  is an isomorphism. When all path components of  $X$  and  $Y$  are nilpotent (e.g. simply connected or simple), this requirement is equivalent to saying that  $f_* : \pi_0(X) \rightarrow \pi_0(Y)$  is bijective and that  $f_* : \pi_k(X, x) \otimes \mathbb{Q} \rightarrow \pi_k(Y, f(x)) \otimes \mathbb{Q}$  is an isomorphism for all  $k \geq 1$  and all  $x \in X$  (for an arbitrary nilpotent group  $G$ , we use the notation  $G \otimes \mathbb{Q}$  for the  $\mathbb{Q}$ -localization of  $G$ , see [39, §I]). If source and target are nilpotent, we call an HQ-equivalence also a *rational homotopy equivalence*.

A space  $Z$  is  $\mathbb{Q}$ -local if for each  $H\mathbb{Q}$ -equivalence  $f : X \rightarrow Y$  and all choices of basepoints, the map

$$\_ \circ f : \text{map}_*(Y; Z) \rightarrow \text{map}_*(X; Z)$$

is a weak equivalence (equivalently the map induced by  $\_ \circ f$  on  $\pi_0$  is bijective for all such  $f$ ). If  $Z$  is nilpotent, this is equivalent to saying that all homotopy groups  $\pi_k(Z, z)$  for  $k \geq 2$  are  $\mathbb{Q}$ -vector spaces and that the fundamental groups  $\pi_1(Z, z)$  are  $\mathbb{Q}$ -local nilpotent groups in the sense of [39, p.4].

A map  $f : X \rightarrow Y$  is a  $\mathbb{Q}$ -localization if  $f$  is an  $H\mathbb{Q}$ -equivalence and  $Y$  is  $\mathbb{Q}$ -local. Such a map, if it exists, is unique up to weak equivalence. It was proven by Sullivan [70] that each nilpotent space admits a  $\mathbb{Q}$ -localization.

**Theorem 3.1.** *Let  $X$  and  $Y$  be connected pointed spaces and assume that  $X$  is finite. Then each component  $\text{map}_*(X; Y)^g \subset \text{map}_*(X; Y)$  of the pointed mapping space is nilpotent. Moreover, if  $Y$  is nilpotent and  $f : Y \rightarrow Z$  is a  $\mathbb{Q}$ -localization, the induced map*

$$\text{map}_*(X; Y)^g \rightarrow \text{map}_*(X; Z)^{f \circ g}$$

*is a  $\mathbb{Q}$ -localization and a rational homotopy equivalence, for each choice of  $g$ .*

The first sentence follows from Theorem 2.5 on p.64 of [39], and Theorem 3.11 on p. 77 of loc.cit. shows that the induced map is a  $\mathbb{Q}$ -localization, which is a rational homotopy equivalence as source and target are nilpotent.

For our purposes, it will be convenient to have a strictly functorial  $\mathbb{Q}$ -localization<sup>2</sup>.

**Theorem 3.2.** *There is an enriched functor  $(\_)_{\mathbb{Q}} : \mathcal{S} \rightarrow \mathcal{S}$ , together with an enriched natural transformation  $\eta : \text{id} \rightarrow (\_)_{\mathbb{Q}}$ , such that for each  $X \in \mathcal{S}$ , the map  $\eta_{\mathbb{Q}} : X \rightarrow X_{\mathbb{Q}}$  is a  $\mathbb{Q}$ -localization.*

We refer to [67, §3] for the vocabulary of enriched category theory. The statement that  $(\_)_{\mathbb{Q}}$  is enriched means that it comes along with natural maps

$$i_{X, Y} : \text{map}(X; Y) \rightarrow \text{map}(X_{\mathbb{Q}}; Y_{\mathbb{Q}}),$$

and the enriched natural transformation  $\eta$  is given by maps  $\eta_X : X \rightarrow X_{\mathbb{Q}}$  such that the composition

$$\text{map}(X; Y) \xrightarrow{i_{X, Y}} \text{map}(X_{\mathbb{Q}}; Y_{\mathbb{Q}}) \xrightarrow{-\circ \eta_X} \text{map}(X; Y_{\mathbb{Q}})$$

agrees with the map  $\eta_Y \circ \_$ . There is an induced enriched functor  $\mathcal{S}_* \rightarrow \mathcal{S}_*$  of pointed spaces.

*References for Theorem 3.2.* This is due to Bousfield; first in [8] without the word “enriched”, the enrichment is constructed in [9, §5].  $\square$

**Lemma 3.3.** *Let  $X$  and  $Z$  be connected pointed spaces and let  $g : X \rightarrow Z$  be a pointed map. Then if  $X$  is finite, the natural map*

$$i_{X, Z}^g : \text{map}_*(X; Z)^g \rightarrow \text{map}_*(X_{\mathbb{Q}}; Z_{\mathbb{Q}})^{g_{\mathbb{Q}}}$$

*is a rational homotopy equivalence of nilpotent spaces.*

<sup>2</sup>The only place in the paper where this enters essentially is in Observation 4.25.

*Proof.* The composition

$$\mathrm{map}_*(X; Z)^g \xrightarrow{i_{X,Z}^g} \mathrm{map}_*(X_{\mathbb{Q}}; Z_{\mathbb{Q}})^{g_{\mathbb{Q}}} \xrightarrow{-\circ\eta_X} \mathrm{map}_*(X; Z_{\mathbb{Q}})^{g_{\mathbb{Q}}\circ\eta_X}$$

is equal to  $\eta_Z \circ \_$ . The first and last space is nilpotent by Theorem 3.1, and the second map is a weak equivalence since  $Z_{\mathbb{Q}}$  is  $\mathbb{Q}$ -local and  $\eta_X$  is an  $H\mathbb{Q}$ -equivalence. So the middle space is nilpotent as well. The composition is a rational homotopy equivalence by Theorem 3.1, and so  $i_{X,Z}^g$  is a rational homotopy equivalence as well.  $\square$

A similar fact is true for homotopy automorphisms. Rationalization gives maps

$$j_X : \mathrm{hAut}_*(X) \rightarrow \mathrm{hAut}_*(X_{\mathbb{Q}})$$

and

$$j_{X,A} : \mathrm{hAut}_A(X) \rightarrow \mathrm{hAut}_{A_{\mathbb{Q}}}(X_{\mathbb{Q}})$$

when  $X$  is a pointed space or  $(X, A)$  is a space pair. We write  $\mathrm{hAut}_A(X)^{\mathrm{id}}$  for the unit component of  $\mathrm{hAut}_A(X)$ , and let moreover

$$\mathrm{hAut}_{A_{\mathbb{Q}}}(X_{\mathbb{Q}})_{\mathbb{Z}} \subset \mathrm{hAut}_{A_{\mathbb{Q}}}(X_{\mathbb{Q}})$$

be the union of all path components which are hit by  $j_{X,A}$ ; this are clearly grouplike submonoids.

**Lemma 3.4.** *Suppose that  $(X, A)$  is a pair of finite spaces. Then the natural map*

$$j_{X,A} : \mathrm{hAut}_A(X)^{\mathrm{id}} \rightarrow \mathrm{hAut}_{A_{\mathbb{Q}}}(X_{\mathbb{Q}})^{\mathrm{id}}$$

*is a rational homotopy equivalence.*

*Proof.* The case  $A = *$  is a special case of Lemma 3.3. In the general case, consider the commutative diagram

$$\begin{array}{ccc} \mathrm{hAut}_*(X)^{\mathrm{id}} & \xrightarrow{j_X} & \mathrm{hAut}_*(X_{\mathbb{Q}})^{\mathrm{id}} \\ \downarrow & & \downarrow \\ \mathrm{map}_*(A; X)^{\mathrm{inc}} & \xrightarrow{i_{A,X}^{\mathrm{inc}}} & \mathrm{map}_*(A_{\mathbb{Q}}; X_{\mathbb{Q}})^{\mathrm{inc}}, \end{array}$$

where the symbol *inc* is used to denote both inclusions  $A \rightarrow X$  and  $A_{\mathbb{Q}} \rightarrow X_{\mathbb{Q}}$ . The horizontal maps are rational homotopy equivalences by Lemma 3.3. It follows that the induced map on vertical homotopy fibres induces an isomorphism on all rational homotopy groups in degrees  $\geq 2$ . The same is true on fundamental groups, using [39, Proposition 1.10]. On the other hand, restricting the map on vertical fibres to the path component of the identity gives the map  $j_{A,X}$ .  $\square$

Having understood these matters, we usually abuse notation and write

$$\mathrm{hAut}_A(X_{\mathbb{Q}}) := \mathrm{hAut}_{A_{\mathbb{Q}}}(X_{\mathbb{Q}})$$

and use the notation  $\mathrm{hAut}_A(X_{\mathbb{Q}})_{\mathbb{Z}}$ ,  $\mathrm{hAut}_A(X)^{\mathrm{id}}$  similarly.

**3.2. Block diffeomorphisms versus tangential homotopy automorphisms.**  
We need to introduce some more notation.

**Notation 3.5.** (1) We write  $\text{Diff}_\partial(M)^{\sim\text{id}} \subset \text{Diff}_\partial(M)$  for the subgroup of diffeomorphisms which are homotopic to the identity (relative boundary), and define  $\widetilde{\text{Diff}}_\partial(M)^{\sim\text{id}} \subset \widetilde{\text{Diff}}_\partial(M)$  analogously.  
(2) Assume that  $C \subset A \subset X$  are subcomplexes and that  $V \rightarrow X$  is a vector bundle. In that situation, we denote by

$$\text{hAut}_A^C(V)^{\sim\text{id}} \subset \text{hAut}_A^C(V)$$

the preimage of  $\text{hAut}_A(X)^{\text{id}}$  under the forgetful map  $\text{hAut}_A^C(V) \rightarrow \text{hAut}_A(X)$ ; in other words the space of pairs  $(f, \hat{f})$  with  $f \sim \text{id}$  (relative  $A$ ). We define  $\text{hAut}_A^C(V)^{\text{st}, \sim\text{id}} \subset \text{hAut}_A^C(V)^{\text{st}}$  similarly.

The derivative map (2.3) is, by virtue of its definition, a map over  $\text{hAut}_\partial(M)$ , and hence it restricts to a map

$$D : \widetilde{\text{Diff}}_\partial(M)^{\sim\text{id}} \rightarrow \text{hAut}_\partial^\partial(TM)^{\text{st}, \sim\text{id}}.$$

If  $* \in \partial M$  is a basepoint, we can furthermore compose the derivative map with the forgetful map  $\text{hAut}_\partial^\partial(TM)^{\text{st}} \rightarrow \text{hAut}_\partial^*(TM)^{\text{st}}$ . Hence by restriction and taking classifying spaces, we obtain a map

$$BD : B\widetilde{\text{Diff}}_\partial(M)^{\sim\text{id}} \rightarrow B\text{hAut}_\partial^*(TM)^{\text{st}, \sim\text{id}}. \quad (3.6)$$

All the surgery theory we need enters the proof of the following result.

**Theorem 3.7.** [2, Theorem 1.1] *Assume that  $M^d$  is 1-connected,  $\partial M = S^{d-1}$  and that  $d \geq 5$ . Then the spaces  $B\widetilde{\text{Diff}}_\partial(M)^{\sim\text{id}}$  and  $B\text{hAut}_\partial^*(TM)^{\text{st}, \sim\text{id}}$  are nilpotent, and the map  $BD$  from (3.6) is a rational homotopy equivalence.*

In the rest of this subsection, we derive a version of Theorem 3.7 which involves the full block diffeomorphism group and not just  $\widetilde{\text{Diff}}_\partial(M)^{\sim\text{id}}$  and which is directly applicable to the manifolds  $U_{g,1}^n$ . Our goal is Theorem 3.11 below. We begin with the introduction of some more pieces of notation. We let

$$\text{hAut}_\partial(M)^{\cong} \subset \text{hAut}_\partial(M)$$

be the union of the components which are hit by the forgetful map  $\widetilde{\text{Diff}}_\partial(M) \rightarrow \text{hAut}_\partial(M)$  (or equivalently the forgetful map  $\text{Diff}_\partial(M) \rightarrow \text{hAut}_\partial(M)$ , as  $\text{Diff}_\partial(M) \rightarrow \widetilde{\text{Diff}}_\partial(M)$  is by definition 0-connected), and we define for a subcomplex  $C \subset \partial M$

$$\text{hAut}_\partial^C(TM)^{\text{st}, \cong} \subset \text{hAut}_\partial^C(TM)^{\text{st}}$$

as the preimage of  $\text{hAut}_\partial(M)^{\cong}$  under the forgetful map. Using this notation, the derivative map yields

$$BD : B\widetilde{\text{Diff}}_\partial(M) \rightarrow B\text{hAut}_\partial^\partial(TM)^{\text{st}, \cong} \rightarrow B\text{hAut}_\partial^*(TM)^{\text{st}, \cong}. \quad (3.8)$$

**Corollary 3.9.** *Let  $M$  be as in Theorem 3.7. Then the composition 3.8 is an  $H\mathbb{Q}$ -equivalence.*

*Proof.* There is a diagram

$$\begin{array}{ccc}
 B\widetilde{\text{Diff}}_{\partial}(M) \sim \text{id} & \xrightarrow{BD} & B\text{hAut}_{\partial}^*(TM)^{\text{st}, \sim \text{id}} \\
 \downarrow & & \downarrow \\
 B\widetilde{\text{Diff}}_{\partial}(M) & \xrightarrow{BD} & B\text{hAut}_{\partial}^*(TM)^{\text{st}, \cong} \\
 \downarrow & & \downarrow \\
 B\pi_0(\text{hAut}_{\partial}(M)^{\cong}) & \xlongequal{\quad} & B\pi_0(\text{hAut}_{\partial}(M)^{\cong})
 \end{array}$$

whose columns are fibre sequences, so the claim follows immediately from Theorem 3.7 and an application of the Leray–Serre spectral sequence.  $\square$

**Corollary 3.10.** *Assume that  $M$  is a 1-connected and stably parallelizable manifold of dimension  $d \geq 5$ , and that  $\partial M = S^{d-1}$ . Then the composition*

$$B\widetilde{\text{Diff}}_{\partial}(M) \rightarrow \text{map}_{\partial}(M; BO)^0 // \text{hAut}_{\partial}(M)^{\cong} \rightarrow \text{map}_*(M; BO)^0 // \text{hAut}_{\partial}(M)^{\cong}$$

*of the derivative map with the forgetful map is an  $H\mathbb{Q}$ -equivalence.*

*Proof.* Combine Corollary 3.9 with Lemma 2.2.  $\square$

Using the rationalization functor, we now give the variant of Theorem 3.7 which we shall eventually use. We let

$$\text{hAut}_{\partial}(M_{\mathbb{Q}})^{\cong} \subset \text{hAut}_{\partial}(M_{\mathbb{Q}})_{\mathbb{Z}}$$

the union of components which are hit by the rationalization map

$$\text{hAut}_{\partial}(M)^{\cong} \rightarrow \text{hAut}_{\partial}(M_{\mathbb{Q}})_{\mathbb{Z}}.$$

**Theorem 3.11.** *Let  $M$  be a simply connected manifold of dimension  $d \geq 5$ , assume that  $M$  is stably parallelizable and that  $\partial M = S^{d-1}$ . Then the composition*

$$B\widetilde{\text{Diff}}_{\partial}(M) \xrightarrow{BD} \text{map}_{\partial}(M; BO)^0 // \text{hAut}_{\partial}(M)^{\cong} \rightarrow \dots$$

$$\dots \rightarrow \text{map}_{\partial}(M_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \text{hAut}_{\partial}(M_{\mathbb{Q}})^{\cong} \rightarrow \text{map}_*(M_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \text{hAut}_{\partial}(M_{\mathbb{Q}})^{\cong}$$

*of the derivative map, the rationalization and the forgetful map is an  $H\mathbb{Q}$ -equivalence.*

*Proof.* The composition in question agrees with

$$B\widetilde{\text{Diff}}_{\partial}(M) \rightarrow \text{map}_*(M; BO)^0 // \text{hAut}_{\partial}(M)^{\cong} \rightarrow \text{map}_*(M_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \text{hAut}_{\partial}(M_{\mathbb{Q}})^{\cong},$$

so Corollary 3.10 leaves us with the task of proving that the second map is an  $H\mathbb{Q}$ -equivalence. This second map can clearly be factored as

$$\text{map}_*(M; BO)^0 // \text{hAut}_{\partial}(M)^{\cong} \rightarrow \text{map}_*(M_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \text{hAut}_{\partial}(M)^{\cong} \rightarrow \text{map}_*(M_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \text{hAut}_{\partial}(M_{\mathbb{Q}})^{\cong}.$$

The first of those maps is a  $H\mathbb{Q}$ -equivalence, by Lemma 3.3 and a straightforward spectral sequence argument. To prove that the second map is also an  $H\mathbb{Q}$ -equivalence, observe that there is a homotopy cartesian diagram

$$\begin{array}{ccc}
 \text{map}_*(M_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \text{hAut}_{\partial}(M)^{\cong} & \longrightarrow & B\text{hAut}_{\partial}(M)^{\cong} \\
 \downarrow & & \downarrow \\
 \text{map}_*(M_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \text{hAut}_{\partial}(M_{\mathbb{Q}})^{\cong} & \longrightarrow & B\text{hAut}_{\partial}(M_{\mathbb{Q}})^{\cong}.
 \end{array}$$

Therefore, it is sufficient to prove that the homotopy fibre  $F$  of the right vertical map is rationally acyclic.

The definition of  $B\mathrm{hAut}_\partial(M_\mathbb{Q})^\cong$  shows that  $F$  is connected, and Lemma 3.4 shows that  $\pi_k(F) \otimes \mathbb{Q} = 0$  for all  $k \geq 2$ . Hence the universal cover  $\tilde{F}$  is rationally acyclic, and the Leray–Serre spectral sequence of the fibre sequence  $\tilde{F} \rightarrow F \rightarrow B\pi_1(F)$  shows that

$$H_k(F; \mathbb{Q}) \cong H_k(B\pi_1(F); \mathbb{Q}). \quad (3.12)$$

The exact sequence

$$\pi_1(\mathrm{hAut}_\partial(M)^\cong) \rightarrow \pi_1(\mathrm{hAut}_\partial(M_\mathbb{Q})^\cong) \rightarrow \pi_1(F) \rightarrow \pi_0(\mathrm{hAut}_\partial(M)^\cong) \rightarrow \pi_0(\mathrm{hAut}_\partial(M_\mathbb{Q})^\cong)$$

yields a short exact sequence

$$0 \rightarrow T \rightarrow \pi_1(F) \rightarrow Q \rightarrow 1 \quad (3.13)$$

of groups, where

$$T := \mathrm{coker}\left(\pi_1(\mathrm{hAut}_\partial(M)^\cong) \rightarrow \pi_1(\mathrm{hAut}_\partial(M_\mathbb{Q})^\cong)\right)$$

is an abelian torsion group by Lemma 3.4 and

$$Q := \ker\left(\pi_0(\mathrm{hAut}_\partial(M)^\cong) \rightarrow \pi_0(\mathrm{hAut}_\partial(M_\mathbb{Q})^\cong)\right)$$

is finite by [18, Theorem 1.1]. The latter is a relative version of a Theorem of Sullivan [71, Theorem 10.2] which asserts that  $\pi_0(\mathrm{hAut}_*(X)) \rightarrow \pi_0(\mathrm{hAut}_*(X_\mathbb{Q}))$  has finite kernel whenever  $X$  is a finite and simply connected CW complex. As  $Q$  is finite, we get

$$H_k(\pi_1(F); \mathbb{Q}) \cong H_k(T; \mathbb{Q})_Q$$

from the Lyndon–Hochschild–Serre spectral sequence of the group extension (3.13). Since  $H_k(T; \mathbb{Q}) = 0$  for  $k \geq 1$ , being the colimit of  $H_k(H; \mathbb{Q})$ , where  $H$  runs through the finitely generated (and hence finite, as  $T$  is torsion) subgroups of  $T$ ,  $B\pi_1(F)$  is rationally acyclic. By (3.12),  $F$  is rationally acyclic as claimed.  $\square$

**3.3. Cohomology of mapping spaces.** Theorem 3.11 shows that we need to understand the cohomology of mapping spaces  $\mathrm{map}_*(M; BO_\mathbb{Q})^0$ . It turns out to be more convenient for our purposes to consider  $\mathrm{map}_\partial(M; BO_\mathbb{Q})^0$  instead. Since  $\mathrm{map}_\partial(M; BO_\mathbb{Q})^0 = \mathrm{map}_*(M/\partial M; BO_\mathbb{Q})^0$ , it suffices to consider pointed mapping spaces.

In what follows, all homology and cohomology groups are taken with coefficients in  $\mathbb{Q}$ . Though this is not needed for large parts of the section, it is all we shall need later on. To ease notation, we often write  $\alpha\beta := \alpha \cup \beta$  for the cup product of two cohomology classes.

*The slant product.* Let us first recall from [68, Chapter 6.1] the slant product

$$H^n(Y \times X) \otimes H_k(X) \rightarrow H^{n-k}(Y), \quad (\xi, x) \mapsto \xi/x,$$

which is related to the cohomology cross product and the Kronecker product by the formula

$$(\eta \times \zeta)/x = \eta\langle \zeta, x \rangle$$

for  $\eta \in H^*(Y)$ ,  $\zeta \in H^*(X)$  and  $x \in H_*(X)$ . Assume that  $X$  has finite type over  $\mathbb{Q}$  (i.e. each  $H_k(X)$  is finite-dimensional) and pick a homogeneous basis  $(b_i)_i$  of

$H_*(X)$  and let  $(\beta_i)_i$  be the dual basis of  $H^*(X)$ . An arbitrary  $\gamma \in H^*(Y \times X)$  can be written in the form  $\gamma = \sum_i \gamma_i \times \beta_i$  by the Künneth formula, and we get

$$\gamma/b_j = \sum_i (\gamma_i \times \beta_i)/b_j = \sum_i \gamma_i \langle \beta_i, b_j \rangle = \gamma_j.$$

or

$$\gamma = \sum_i (\gamma/b_i) \times \beta_i \quad (3.14)$$

The  $\lambda$ -classes.

**Definition 3.15.** Let  $X$  and  $Z$  be pointed spaces, with  $X$  of finite type over  $\mathbb{Q}$ . The evaluation map  $\text{ev} : \text{map}_*(X; Z) \times X \rightarrow Z$  and the slant product yields

$$\lambda : H^n(Z) \otimes H_k(X) \rightarrow H^{n-k}(\text{map}_*(X; Z)), \quad \xi \otimes x \mapsto \lambda_{x,\xi} := (\text{ev}^* \xi)/x.$$

This construction enjoys a naturality property, which is most concisely expressed by saying that  $\lambda$  is a natural transformation of functors

$$\mathcal{S}_* \times \mathcal{S}_*^{\text{op}} \rightarrow \mathbb{Q} - \text{Mod}$$

from  $(X, Z) \mapsto H^n(Z) \otimes H_k(X)$  to  $(X, Z) \mapsto H^{n-k}(\text{map}_*(X; Z))$ . The formula (3.14) leads to the equation

$$\text{ev}^* z = \sum_i (\text{ev}^* z/b_i) \times \beta_i = \sum_i \lambda_{b_i, z} \times \beta_i \in H^*(\text{map}_*(X; Z) \times X) \quad (3.16)$$

for each  $z \in H^*(Z)$ .

**Lemma 3.17.** Let  $c_{i,j}^k$  be the structure constants of the algebra  $H^*(X)$  with respect to the basis  $(\beta_i)$ , i.e.

$$\beta_i \beta_j = \sum_k c_{i,j}^k \beta_k.$$

Then

$$\lambda_{b_k, zy} = \sum_{i,j} (-1)^{|\beta_i|(|y|-|b_j|)} c_{i,j}^k \lambda_{b_i, z} \lambda_{b_j, y}$$

for all  $z, y \in H^*(Z)$ .

*Proof.* Using (3.16), one checks that firstly

$$\text{ev}^*(z) \text{ev}^*(y) = \text{ev}^*(zy) = \sum_k \lambda_{b_k, zy} \times \beta_k,$$

and secondly

$$\begin{aligned} \text{ev}^*(z) \text{ev}^*(y) &= \left( \sum_i \lambda_{b_i, z} \times \beta_i \right) \left( \sum_j \lambda_{b_j, y} \times \beta_j \right) = \\ &= \sum_{i,j} (-1)^{|\beta_i|(|y|-|b_j|)} (\lambda_{b_i, z} \lambda_{b_j, y}) \times (\beta_i \beta_j) = \\ &= \sum_{i,j,k} (-1)^{|\beta_i|(|y|-|b_j|)} c_{i,j}^k (\lambda_{b_i, z} \lambda_{b_j, y}) \times \beta_k. \end{aligned}$$

Comparing coefficients yields the result.  $\square$

**Lemma 3.18.** Assume that  $X$  is connected and that the cup length of  $X$  is  $\leq r-1$  (i.e. cup products of  $r$  elements of  $\check{H}^*(X)$  vanish). Then for  $z_1, \dots, z_r \in H^*(Z)$ , we have

$$\lambda_{b_k, z_1 \cdots z_r} = 0.$$

*Proof.* Using (3.16), we compute

$$\begin{aligned} \sum_k \lambda_{b_k, z_1 \cdots z_r} \times \beta_k &= \text{ev}^*(z_1 \cdots z_r) = \text{ev}^*(z_1) \cdots \text{ev}^*(z_r) = \\ &= \sum_{j_1, \dots, j_r} \epsilon_{j_1, \dots, j_r} (\lambda_{b_{j_1, z_1}} \cdots \lambda_{b_{j_r, z_r}}) \times (\beta_{j_1} \cdots \beta_{j_r}) = 0 \end{aligned}$$

for some signs  $\epsilon_{j_1, \dots, j_r} \in \{\pm 1\}$ . The claim follows immediately.  $\square$

*The case of an Eilenberg–Mac–Lane space.* Now we consider the case  $Z = K(\mathbb{Q}, m)$  and let  $u_m \in H^m(K(\mathbb{Q}, m); \mathbb{Q})$  be the fundamental class. We assume that  $X$  is connected, and we wish to calculate  $H^*(\text{map}_*(X; K(\mathbb{Q}, m))^0)$  in terms of the classes  $\lambda_{a_i, u^r}$ , where  $(a_i)$  is a homogeneous basis of  $H_*(X)$  and  $(\alpha_i)$  is dual basis.

There are a couple of relations between the classes  $\lambda_{a_i, u^r}$  which we state first.

**Lemma 3.19.** *Assume that  $X$  is connected. Then*

$$\lambda_{a_i, u_m^r} = 0 \in H^*(\text{map}_*(X; K(\mathbb{Q}, m))^0)$$

unless  $0 < |a_i| < rm$ .

*Proof.* For degree reasons, we get that  $\lambda_{a_i, u_m^r} = 0$  if  $|a_i| > rm$ . For the remaining cases, note that the evaluation map  $\text{map}_*(X; K(\mathbb{Q}, m))^0 \times X \rightarrow K(\mathbb{Q}, m)$  factors through the smash product  $\text{map}_*(X; K(\mathbb{Q}, m))^0 \wedge X$ . It follows that in the sum

$$\text{ev}^* u_m^r = \sum_i \lambda_{a_i, u_m^r} \times \alpha_i,$$

all terms in which one factor has degree 0 must vanish. This happens if  $|\alpha_i| = |a_i| = 0$  and  $rm - |a_i| = 0$ .  $\square$

**Proposition 3.20.** *Assume that  $X$  is connected and of finite type over  $\mathbb{Q}$ . Then the natural map*

$$\mathbb{F}\left(\bigoplus_{k=1}^{m-1} H_k(X)[m-k]\right) \rightarrow H^*(\text{map}_*(X; K(\mathbb{Q}, m))^0)$$

induced by the maps

$$H_k(X) \rightarrow H^{m-k}(\text{map}_*(X; K(\mathbb{Q}, m))^0), \quad a \mapsto \lambda_{a, u_m}$$

is an isomorphism. Hence  $H^*(\text{map}_*(X; K(\mathbb{Q}, m))^0)$  is the free graded commutative algebra generated by the elements  $\lambda_{a_i, u_m}$ , where  $a_i$  runs through a homogeneous basis for  $\tilde{H}_{* < m}(X)$ .

*Proof.* The space  $\text{map}_*(X; K(\mathbb{Q}, m))^0$  is a connected infinite loop space; in fact it is a generalized Eilenberg–Mac Lane space with homotopy groups

$$\pi_k(\text{map}_*(X; K(\mathbb{Q}, m))) \otimes \mathbb{Q} = [X, \Omega^k K(\mathbb{Q}, m)]_* = [X, K(\mathbb{Q}, m-k)]_* = \tilde{H}^{m-k}(X) \quad (3.21)$$

in positive degrees. It follows from the Milnor–Moore theorem [55] and the assumption that  $X$  is of finite type that  $H^*(\text{map}_*(X; K(\mathbb{Q}, m))^0)$  is isomorphic (as a graded algebra) to  $\mathbb{F}(\bigoplus_{k=1}^{m-1} H_k(X)[m-k])$ . The only issue is to verify that the map in question is indeed an isomorphism.

To achieve this, we use the following general principle. Assume that  $Y$  is a connected infinite loop space<sup>3</sup> of finite type, let  $V_*$  be a  $\mathbb{N}_0$ -graded, degreewise

<sup>3</sup>It would be enough to assume that  $Y$  is a connected homotopy-commutative H-space.

finite-dimensional  $\mathbb{Q}$ -vector space with  $V_0 = 0$  and let  $\sigma : V_* \rightarrow H^*(Y; \mathbb{Q})$  be a graded linear map. Then the induced algebra map

$$\mathbb{F}(\sigma) : \mathbb{F}(V_*) \rightarrow H^*(Y; \mathbb{Q})$$

from the free graded-commutative algebra on  $V$  to the cohomology of  $Y$  is an isomorphism if and only if the bilinear form

$$B_k : \pi_k(Y) \otimes \mathbb{Q} \times V_k \rightarrow \mathbb{Q}, (f, v) \mapsto \langle \sigma(v), \text{hur}_k^{\mathbb{Q}}(f) \rangle$$

is nondegenerate for each  $k$ . This principle is easily shown for  $Y = K(\mathbb{Q}, n)$ ; the general case follows from that and the fact that  $Y$  splits rationally as a product of Eilenberg–Mac-Lane spaces [21, §16].

We must therefore show that the bilinear form

$$B_{X,k} : \pi_{m-k}(\text{map}_*(X; K(\mathbb{Q}, m)^0)) \times \tilde{H}_k(X) \rightarrow \mathbb{Q}$$

given by

$$([f], a) \mapsto \langle \lambda_{a,u}, \text{hur}_{m-k}^{\mathbb{Q}}([f]) \rangle$$

is nondegenerate. If  $F : X \rightarrow Y$  is a map of pointed spaces, we denote by  $F^\sharp : \text{map}_*(Y; K(\mathbb{Q}, m))^0 \rightarrow \text{map}_*(X; K(\mathbb{Q}, m))^0$  the induced map. By the naturality of the slant product and the Kronecker product, we have

$$B_{Y,k}([f], F_*a) = B_{X,k}((F^\sharp)_*[f], a).$$

If  $F_* : H_k(X) \rightarrow H_k(Y)$  is an isomorphism then so is the induced map  $(F^\sharp)_* : \pi_{m-k}(\text{map}_*(Y; K(\mathbb{Q}, m))^0) \rightarrow \pi_{m-k}(\text{map}_*(X; K(\mathbb{Q}, m))^0)$  by (3.21), and it follows that nondegeneracy of  $B_{Y,k}$  is equivalent to nondegeneracy of  $B_{X,k}$  (in degree  $k$ ). Now there are maps  $X \rightarrow K(H_k(X; \mathbb{Q}), k)$  and  $\bigvee^g S^k \rightarrow K(H_k(X; \mathbb{Q}), k)$  inducing isomorphisms on  $H_k(-; \mathbb{Q})$ . This argument proves that it suffices to consider the case where  $X$  is a wedge of finitely many  $k$ -spheres. Using the naturality again, the form  $B_{\bigvee^g S^k, k}$  decomposes as the direct sum of  $k$  copies of the form  $B_{S^k, k}$ , and so we are left with the case  $X = S^k$ .

Both,  $\pi_{m-k}(\text{map}_*(S^k; K(\mathbb{Q}, m)^0)) \otimes \mathbb{Q}$  and  $\tilde{H}_k(S^k; \mathbb{Q})$  are 1-dimensional and  $\text{map}_*(S^k; K(\mathbb{Q}, m))^0 \simeq K(\mathbb{Q}, m-k)$  is  $(m-k-1)$ -connected. Hence (by the Hurewicz theorem) it is left to be proven that  $\lambda_{[S^k], u_m} \in H^{m-k}(\text{map}_*(S^k; K(\mathbb{Q}, m)))$  is nonzero. On the other hand, the map

$$H^m(K(\mathbb{Q}, m); \mathbb{Q}) \rightarrow H^{m-k}(\text{map}_*(S^k; K(\mathbb{Q}, m))) = H^{m-k}(\Omega^k K(\mathbb{Q}, m)), u \mapsto \text{ev}^*(u)/[S^k]$$

can be identified with the "transgression" map

$$H^m(K(\mathbb{Q}, m); \mathbb{Q}) = [K(\mathbb{Q}, m); K(\mathbb{Q}, m)] \xrightarrow{\Omega^k} [\Omega^k K(\mathbb{Q}, m); \Omega^k K(\mathbb{Q}, m)] = H^{m-k}(\Omega^k K(\mathbb{Q}, m))$$

which is well-known to be an isomorphism.  $\square$

Using that  $BO_{\mathbb{Q}} \simeq \prod_{m \geq 1} K(\mathbb{Q}; 4m)$  via the classes  $L_m$ , we deduce:

**Corollary 3.22.** *Let  $X$  be a connected space of finite type over  $\mathbb{Q}$ . Then the map*

$$\mathbb{F} \left( \bigoplus_{m \geq 1, 0 < k < 4m} H_k(X)[4m-k] \right) \rightarrow H^*(\text{map}_*(X; BO_{\mathbb{Q}})^0; \mathbb{Q})$$

which on the generators is the direct sum of the maps

$$H_k(X)[4m-k] \rightarrow H^{4m-k}(H^*(\text{map}_*(X; BO_{\mathbb{Q}})^0; \mathbb{Q})); a \mapsto \lambda_{a, L_m}$$

is an isomorphism. Thus  $H^*(\text{map}_*(X; BO_{\mathbb{Q}})^0; \mathbb{Q})$  is the free graded-commutative algebra with generators

$$\lambda_{a_i, L_m} \in H^{4m-|a_i|}(\text{map}_*(X; BO_{\mathbb{Q}})^0; \mathbb{Q})$$

where  $a_i$  runs through a homogeneous basis for  $\tilde{H}_*(X)$  and  $m$  through the natural numbers and  $4m - |a_i| > 0$ . The classes  $\lambda_{a_i, L_{m_1} \dots L_{m_r}}$  with  $m_1 \leq \dots \leq m_r$  are determined by the relation stated in Lemma 3.17.  $\square$

The naturality of the construction of the  $\lambda$ -classes, together with Lemma 3.3, has the following consequence which turns out to be helpful for us.

**Corollary 3.23.** *Let  $X$  be a connected space of finite type over  $\mathbb{Q}$ . Then the  $\pi_0(\text{hAut}_*(X))$ -action on  $H^*(\text{map}_*(X; BO_{\mathbb{Q}})^0; \mathbb{Q})$  is, under the isomorphism of Corollary 3.22, induced by the usual action of  $\pi_0(\text{hAut}_*(X))$  on the rational homology of  $X$ . Furthermore, the action extends to an action of  $\pi_0(\text{hAut}_*(X_{\mathbb{Q}}))$  on  $H^*(\text{map}_*(X; BO_{\mathbb{Q}})^0; \mathbb{Q})$  which is induced by the usual action of  $\pi_0(\text{hAut}_*(X_{\mathbb{Q}}))$  on  $H_*(X_{\mathbb{Q}}; \mathbb{Q}) \cong H_*(X; \mathbb{Q})$ .*

**3.4. Consequences for block diffeomorphism spaces.** We now collect some consequences of the above calculations. Let  $M^d$  be a smooth oriented compact manifold with boundary. In that case, there is a fundamental class  $[M] \in H_d(M/\partial M) \cong H_d(M, \partial M)$ .

**Lemma 3.24.** *Let  $M$  be a smooth oriented compact manifold with boundary, and denote by*

$$q : \text{map}_{\partial}(M; BO) \rightarrow \text{map}_{\partial}(M; BO) // \text{hAut}_{\partial}^+(M)$$

the fibre inclusion. Then

$$q^* \kappa_c = \lambda_{[M], c}$$

for each  $c \in H^*(BO; \mathbb{Q})$ .

*Proof.* This is a consequence of the definitions and the fact that for each space  $Y$ , the Gysin map of the trivial bundle  $\pi : Y \times (M/\partial M) \rightarrow Y$  is given by  $\pi_!(\xi) = \xi/[M]$ .  $\square$

**Proposition 3.25.** *Assume that  $M$  is a stably parallelizable 1-connected smooth manifold of dimension  $d \geq 5$  with  $\partial M = S^{d-1}$ . Then the algebra map*

$$H^*(\text{map}_{\partial}(M_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \text{hAut}_{\partial}(M_{\mathbb{Q}})^{\cong}) \rightarrow H^*(\widetilde{BDiff}_{\partial}(M))$$

is surjective. Moreover

- (1) If  $d$  is odd, the kernel is the ideal generated by the classes  $\kappa_{L_m}$ ,  $4m - d > 0$ .
- (2) If  $d = 2n$  is even, the kernel is the ideal generated by the classes  $\kappa_{L_m} - \sigma_{4m-d}$ ,  $4m - d > 0$ .

*Proof.* The composition

$$B\widetilde{Diff}_{\partial}(M) \rightarrow \text{map}_{\partial}(M_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \text{hAut}_{\partial}(M_{\mathbb{Q}})^{\cong} \rightarrow \text{map}_*(M_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \text{hAut}_{\partial}(M_{\mathbb{Q}})^{\cong}$$

is an  $H\mathbb{Q}$ -equivalence by Theorem 3.11, and so surjectivity is immediate. Theorem 2.10 and (2.11) prove that the classes listed in (1) and (2) lie in the kernel, and so it remains for us to show that the kernel is not larger. Consider the map

$$\Pi : \text{map}_{\partial}(M_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \text{hAut}_{\partial}^+(M_{\mathbb{Q}})^{\cong} \rightarrow \text{map}_*(M_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \text{hAut}_{\partial}^+(M_{\mathbb{Q}})^{\cong} \times \prod_{4m-d>0} K(\mathbb{Q}, 4m-d) \quad (3.26)$$

made out of the forgetful map and the classes  $\kappa_{L_m}$  or  $\kappa_{L_m} - \sigma_{4m-d}$  (for odd  $d$  or even  $d$ ). If we can prove that  $\Pi$  is a weak equivalence, the claim follows.

To prove this, note first that  $\Pi$  is a map over  $B\text{hAut}_\partial^+(M_{\mathbb{Q}})^\cong$ . The induced map on the homotopy fibres of the respective maps to  $B\text{hAut}_\partial^+(M_{\mathbb{Q}})^\cong$  is the map

$$\text{map}_\partial(M_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 \rightarrow \text{map}_*(M_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 \times \prod_{4m-d>0} K(\mathbb{Q}, 4m-d); \quad (3.27)$$

made out of the forgetful map and the classes  $\lambda_{L_m}$ . This follows from Lemma 3.24 and the fact that the classes  $\sigma_{4m-d}$  are pulled back from  $B\text{hAut}_\partial^+(M_{\mathbb{Q}})_{\mathbb{Z}}$ .

To see that (3.27) is a weak equivalence, one uses Corollary 3.22 for both mapping spaces and the naturality of the  $\lambda$ -classes.  $\square$

*Remark 3.28.* The proof suggests that one might extract a proof of Theorem 2.10 out of the arguments in [2, §4]. While surely true, it seems simpler to us to use the independent proof of Theorem 2.10, as it allows us to treat [2, §4] as a black box.

#### 4. Homotopy calculations for the manifolds $U_{g,1}^n$

**4.1. Homotopy groups.** Let us now focus our attention to the manifolds we are actually interested in, i.e.

$$U_{g,1}^n := \sharp^g(S^n \times S^{n+1}) \setminus \text{int}(D^{2n+1}). \quad (4.1)$$

Most of the following facts are also proven in [30], but as the setup in [30] is more general, we prefer to indicate the proofs here. Note that

$$U_{g,1}^n \simeq \bigvee^g S^n \vee \bigvee^g S^{n+1}.$$

Let us denote

$$N(g)_{\mathbb{Z}} := H_n(U_{g,1}^n; \mathbb{Z}) \cong \mathbb{Z}^g$$

and

$$N(g) := H_n(U_{g,1}^n; \mathbb{Q}) = N(g)_{\mathbb{Z}} \otimes \mathbb{Q}.$$

By Poincaré duality and the universal coefficient theorem, we have

$$H_{n+1}(U_{g,1}^n; \mathbb{Z}) \cong H^n(U_{g,1}^n; \mathbb{Z}) \cong N(g)_{\mathbb{Z}}^\vee.$$

These isomorphisms are natural with respect to the action of  $\text{hAut}_\partial(U_{g,1}^n)$ , and hold similarly for rational coefficients. The group  $\text{hAut}_\partial(U_{g,1}^n)$  acts trivially on  $H_{2n+1}(U_{g,1}^n, \partial; \mathbb{Z})$ ,  $H_{2n+1}(U_{g,1}^n, \partial; \mathbb{Q})$ ,  $H^{2n+1}(U_{g,1}^n, \partial; \mathbb{Z})$  and  $H^{2n+1}(U_{g,1}^n, \partial; \mathbb{Q})$ .

Let  $x_1, \dots, x_g \in \pi_n(U_{g,1}^n)$  be the elements represented by the inclusion of the  $g$  different  $S^n$ 's, and similarly let  $y_1, \dots, y_g \in \pi_{n+1}(U_{g,1}^n)$  be represented by the  $g$  copies of  $S^{n+1}$ . The inclusion of the boundary  $S^{2n} = \partial U_{g,1}^n \rightarrow U_{g,1}^n$  represents an element  $\omega \in \pi_{2n}(U_{g,1}^n)$  which agrees with the sum

$$\omega = \sum_{j=1}^g [x_j, y_j] \quad (4.2)$$

of Whitehead products of the generators, if the numbering and the signs of the generators are chosen appropriately. This is obvious from the definition of the Whitehead product when  $g = 1$ , and follows for higher  $g$ . We denote by

$$a_j := \text{hur}_n^{\mathbb{Z}}(x_j) \in H_n(U_{g,1}^n; \mathbb{Z}) \text{ and } b_j := \text{hur}_{n+1}^{\mathbb{Z}}(y_j) \in H_{n+1}(U_{g,1}^n; \mathbb{Z})$$

the images under the Hurewicz homomorphism, and use the same symbol for the image in rational homology.

We let  $(\alpha_1, \dots, \alpha_g)$  the basis of  $H^n(U_{g,1}^n; \mathbb{Z}) = H^n(U_{g,1}^n, \partial; \mathbb{Z})$  dual to  $(a_1, \dots, a_g)$  and let  $(\beta_1, \dots, \beta_g)$  be the basis of  $H^{n+1}(U_{g,1}^n; \mathbb{Z}) = H^{n+1}(U_{g,1}^n, \partial; \mathbb{Z})$  dual to  $(b_1, \dots, b_g)$ . Let  $\nu \in H^{2n+1}(U_{g,1}^n, \partial; \mathbb{Z})$  be dual to the fundamental class  $[U] = [U_{g,1}^n]$ . The cup product structure of  $H^*(U_{g,1}^n; \mathbb{Z})$  is given by

$$\alpha_i \beta_j = \delta_{ij} \nu; \quad (4.3)$$

all other cup products are zero for degree reasons. The cup length of  $U_{g,1}^n/\partial$  is therefore 2, and it follows from Lemma 3.17 that

$$\lambda_{[U], L_m L_k} = \sum_{j=1}^g \lambda_{a_j, L_m} \lambda_{b_j, L_k} + \sum_{j=1}^g \lambda_{b_j, L_m} \lambda_{a_j, L_k}, \quad (4.4)$$

and that

$$\lambda_{[U], L_{m_1} \dots L_{m_r}} = 0 \quad (4.5)$$

if  $r \geq 3$  and  $1 \leq m_1 \leq \dots \leq m_r$ .

Having understood the homological structure, let us turn to homotopy groups. The Hurewicz theorem shows that

$$\text{hur}_n^{\mathbb{Z}} : \pi_n(U_{g,1}^n) \rightarrow H_n(U_{g,1}^n; \mathbb{Z})$$

is an isomorphism, and that

$$\text{hur}_{n+1}^{\mathbb{Z}} : \pi_{n+1}(U_{g,1}^n) \rightarrow H_{n+1}(U_{g,1}^n; \mathbb{Z})$$

is surjective, if  $n \geq 2$ . The rational Hurewicz theorem implies that  $\text{hur}_{n+1}^{\mathbb{Z}}$  has finite kernel if  $n \geq 3$ .

The rational homotopy groups of  $U_{g,1}^n$  are known by a computation in rational homotopy theory. For a graded vector space  $V$ , we denote the desuspension by  $(s^{-1}V)_k := V_{k+1}$ . For a (simply connected) space  $X$ ,  $s^{-1}\pi_*(X_{\mathbb{Q}})$  is a graded Lie algebra under the Whitehead product, and since  $U_{g,1}^n$  is a wedge of spheres, we have

$$s^{-1}\pi_*((U_{g,1}^n)_{\mathbb{Q}}) \cong \mathbb{L}(\pi_n((U_{g,1}^n)_{\mathbb{Q}}) \oplus \pi_{n+1}((U_{g,1}^n)_{\mathbb{Q}})),$$

the free Lie algebra on the graded vector space which is  $\pi_n((U_{g,1}^n)_{\mathbb{Q}})$  in degree  $n-1$  and  $\pi_{n+1}((U_{g,1}^n)_{\mathbb{Q}})$  in degree  $n$ , by [21, Theorem 24.5]. It follows that

$$\pi_{n+k}(U_{g,1}^n) \otimes \mathbb{Q} = 0, \quad 2 \leq k \leq n-2, \quad n+2 \leq 2n-3. \quad (4.6)$$

## 4.2. Mapping class groups.

**Lemma 4.7.** *The map*

$$\pi_0(\text{hAut}_*(U_{g,1}^n)) \rightarrow \text{GL}(N(g)_{\mathbb{Z}}) \times \text{GL}(N(g)_{\mathbb{Z}})^{\vee}$$

*given by the action on integral homology is surjective when  $n \geq 1$ , and has finite kernel, provided that  $n \geq 3$ .*

*The analogous map*

$$\pi_0(\text{hAut}_*((U_{g,1}^n)_{\mathbb{Q}})) \rightarrow \text{GL}(N(g)) \times \text{GL}(N(g))^{\vee}$$

*is an isomorphism, which maps the subgroup  $\pi_0(\text{hAut}_*((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}}) \subset \pi_0(\text{hAut}_*((U_{g,1}^n)_{\mathbb{Q}}))$  onto the subgroup  $\text{GL}(N(g)_{\mathbb{Z}}) \times \text{GL}(N(g)_{\mathbb{Z}})^{\vee}$ .*

*Proof.* Since  $\mathrm{hAut}_*(\mathbb{V}^g S^n) \rightarrow \mathrm{GL}(H_n(\mathbb{V}^g S^n; \mathbb{Z}))$  is surjective, and similarly for a wedge of  $(n + 1)$ -spheres, surjectivity follows. To see that the kernel is finite, we note that  $\mathrm{hur}_{n+1}^{\mathbb{Z}}$  identifies  $N(g)_{\mathbb{Z}}^{\vee} = H_{n+1}(U_{g,1}^n; \mathbb{Z})$  with the torsionfree quotient of  $\pi_{n+1}(U_{g,1}^n)$ , which is why we can factor the map in question as

$$\pi_0(\mathrm{hAut}(U_{g,1}^n)) \rightarrow \mathrm{GL}(\pi_n(U_{g,1}^n)) \times \mathrm{GL}(\pi_{n+1}(U_{g,1}^n)) \rightarrow \mathrm{GL}(N(g)_{\mathbb{Z}}) \times \mathrm{GL}(N(g)_{\mathbb{Z}}^{\vee}). \quad (4.8)$$

The kernel of the second map can be identified with

$$\mathrm{Hom}(H_{n+1}(U_{g,1}^n; \mathbb{Z}); \ker(\mathrm{hur}_{n+1}^{\mathbb{Z}}))$$

and is therefore finite. The first map is injective: a pointed map  $f : U_{g,1}^n \rightarrow U_{g,1}^n$  which induces the identity on both,  $\pi_n$  and  $\pi_{n+1}$ , must be homotopic to the identity, as  $U_{g,1}^n$  is a wedge of spheres.

The rational case is analogous. The proof of surjectivity is similar (using elementary matrices to generate  $\mathrm{GL}_g(\mathbb{Q})$ ), and in the factorization analogous to (4.8), the second map is an isomorphism since  $\mathrm{hur}_{n+1}^{\mathbb{Q}}$  is an isomorphism.

The last statement follows easily from the others.  $\square$

We are interested in  $\mathrm{hAut}_{\partial}(U_{g,1}^n)$  rather than  $\mathrm{hAut}_*(U_{g,1}^n)$ . These two spaces are related by a fibre sequence

$$\mathrm{hAut}_{\partial}(U_{g,1}^n) \rightarrow \mathrm{hAut}_*(U_{g,1}^n) \rightarrow \Omega^{2n}U_{g,1}^n. \quad (4.9)$$

**Lemma 4.10.** *Let  $n \geq 3$ .*

(1) *The map*

$$\pi_0(\mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})) \rightarrow \mathrm{GL}(N(g))$$

*induced by the action on  $H_n(U_{g,1}^n; \mathbb{Q})$  is an isomorphism.*

(2) *It maps the subgroup  $\pi_0(\mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}})$  onto  $\mathrm{GL}(N(g)_{\mathbb{Z}})$ .*

(3) *Moreover  $\pi_0(\mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}}) \cong \pi_0(\mathrm{hAut}_*((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}})$ .*

*Proof.* (1) Consider the diagram

$$\begin{array}{ccc} \pi_0(\mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})) & \longrightarrow & \mathrm{GL}(N(g)) \\ \downarrow & & \downarrow \Delta \\ \pi_0(\mathrm{hAut}_*((U_{g,1}^n)_{\mathbb{Q}})) & \longrightarrow & \mathrm{GL}(N(g)) \times \mathrm{GL}(N(g)^{\vee}); \end{array} \quad (4.11)$$

the left vertical map is the obvious one, and the map  $\Delta$  sends  $h \in \mathrm{GL}(N(g)_{\mathbb{Z}})$  to  $(h, (h^{-1})^{\vee})$ . The horizontal maps are induced by the action on homology. The diagram commutes by Poincaré–Lefschetz duality.

The bottom horizontal map is an isomorphism by Lemma 4.7, and our task is to prove that the top horizontal map is an isomorphism as well. To prove that the top horizontal map in (4.11) is injective, it suffices to show that the left vertical map is injective, but that fits into an exact sequence

$$\pi_{2n+1}((U_{g,1}^n)_{\mathbb{Q}}) \xrightarrow{\delta} \pi_0(\mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})) \rightarrow \pi_0(\mathrm{hAut}_*((U_{g,1}^n)_{\mathbb{Q}}))$$

coming from the fibre sequence (4.9) or the rationalized version thereof. It is proven in [30] that the connecting homomorphism  $\delta$  is zero (see Proof of Proposition 6.6 and Remark 6.7 of the quoted paper).

To prove that the top horizontal map of (4.11) is surjective, we start with  $h \in \mathrm{GL}(N(g))$  and use Lemma 4.7 to find  $\varphi \in \mathrm{hAut}_*((U_{g,1}^n)_{\mathbb{Q}})$  which realizes  $\Delta(h)$  on  $H_n(U_{g,1}^n; \mathbb{Q}) \oplus H_{n+1}(U_{g,1}^n; \mathbb{Q})$ .

Let  $f = (f_{ij})$  and  $g = (g_{ij})$  be the matrices which describe the effect of  $\varphi$  on  $H_n(U_{g,1}^n; \mathbb{Q})$  and  $H_{n+1}(U_{g,1}^n; \mathbb{Q})$ , respectively, in terms of the bases  $(a_1, \dots, a_g)$  and  $(b_1, \dots, b_g)$ . By construction  $f^\top g = 1$  or  $\sum_i f_{ij} g_{ik} = \delta_{jk}$ . Using (4.2), we deduce that

$$\varphi_*(\omega) = \omega \in \pi_{2n}((U_{g,1}^n)_{\mathbb{Q}}).$$

So  $\varphi$  is homotopic relative to the basepoint to a map which is the identity on  $\partial U_{g,1}^n$ . So  $\varphi$  can be viewed as an element of  $\mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})$  which maps to  $h$  under the top horizontal map of (4.11).

To show (2) and (3), we consider the commutative diagram

$$\begin{array}{ccc} \pi_0(\mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})) \cong & & (4.12) \\ \downarrow & \searrow & \\ \pi_0(\mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}}))_{\mathbb{Z}} & \longrightarrow & \mathrm{GL}(N(g)_{\mathbb{Z}}) \\ \downarrow & & \downarrow \\ \pi_0(\mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})) & \xrightarrow{\cong} & \mathrm{GL}(N(g)); \end{array}$$

the bottom map is an isomorphism by item (1), and all maps pointing downwards are by definition injective.

Now every linear automorphism of  $H_n(U_{g,1}^n; \mathbb{Z})$  can be realized by a diffeomorphism fixing the boundary pointwise: it is explained in the proof of [30, Proposition 5.3] how to deduce this fact from [76, Lemma 17]. It follows that the diagonal map in (4.12) is surjective, hence both maps

$$\pi_0(\mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})) \cong \rightarrow \pi_0(\mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}}))_{\mathbb{Z}} \rightarrow \mathrm{GL}(N(g)_{\mathbb{Z}})$$

are isomorphisms; which shows (2) and (3).  $\square$

**4.3. Some higher homotopy groups of the homotopy automorphisms.** Let us turn to a description of some of the higher homotopy groups of  $\mathrm{hAut}_*(U_{g,1}^n)$  and  $\mathrm{hAut}_{\partial}(U_{g,1}^n)$ . Lemma 3.4 shows that the map

$$\pi_k(\mathrm{hAut}_*(U_{g,1}^n)) \otimes \mathbb{Q} \rightarrow \pi_k(\mathrm{hAut}_*((U_{g,1}^n)_{\mathbb{Q}})) \otimes \mathbb{Q} = \pi_k(\mathrm{hAut}_*((U_{g,1}^n)_{\mathbb{Q}}))$$

is an isomorphism when  $k \geq 1$ . The abelian groups  $\pi_k(\mathrm{hAut}_*((U_{g,1}^n)_{\mathbb{Q}}))$  are acted upon by the group  $\pi_0(\mathrm{hAut}_*((U_{g,1}^n)_{\mathbb{Q}})) \cong \mathrm{GL}(N(g)) \times \mathrm{GL}(N(g)^\vee)$ :  $\pi_0(\mathrm{hAut}_*((U_{g,1}^n)_{\mathbb{Q}})) = \pi_1(\mathrm{BhAut}_*((U_{g,1}^n)_{\mathbb{Q}}))$  acts on the higher homotopy groups of  $\mathrm{BhAut}_*((U_{g,1}^n)_{\mathbb{Q}})$ .

**Lemma 4.13.** *We have*

$$\dim_{\mathbb{Q}}(\pi_k(\mathrm{BhAut}_*(U_{g,1}^n)) \otimes \mathbb{Q}) = \begin{cases} g^2 & k = 2 \text{ and } n \geq 3, \\ 0 & 3 \leq k \leq n - 2. \end{cases}$$

*The same is true for  $(U_{g,1}^n)_{\mathbb{Q}}$  in place of  $U_{g,1}^n$ .*

*Proof.* For  $k \geq 1$ , we have  $\pi_k(\mathrm{hAut}_*(U_{g,1}^n)) \cong \pi_k(\mathrm{map}_*(U_{g,1}^n; U_{g,1}^n); \mathrm{id})$ . The inclusion

$$U_{g,1}^n \simeq \bigvee^g S^n \vee \bigvee^g S^{n+1} \rightarrow (S^n)^g \times (S^{n+1})^g$$

is the inclusion of the  $(2n - 1)$ -skeleton and hence  $(2n - 1)$ -connected. So the induced map

$$\text{map}_*(U_{g,1}^n; U_{g,1}^n) \rightarrow \text{map}_*(U_{g,1}^n; (S^n \times S^{n+1})^g)$$

is  $2n - 1 - (n + 1) = (n - 2)$ -connected. Because

$$\text{map}_*(U_{g,1}^n; (S^n \times S^{n+1})^g) \simeq (\Omega^n(S^n \times S^{n+1})^g)^g \times (\Omega^{n+1}(S^n \times S^{n+1})^g)^g,$$

we obtain an isomorphism

$$\begin{aligned} \pi_k(\text{map}_*(U_{g,1}^n; U_{g,1}^n)) &\cong \\ &(\pi_{n+k}(S^n))^{\oplus g^2} \oplus (\pi_{n+k}(S^{n+1}))^{\oplus g^2} \oplus (\pi_{n+1+k}(S^n))^{\oplus g^2} \oplus (\pi_{n+k+1}(S^{n+1}))^{\oplus g^2} \end{aligned}$$

for  $1 \leq k \leq n - 3$ . The claim follows by using the knowledge of the rational homotopy groups of the spheres.

The rational case is an immediate consequence of the integral one and Lemma 3.4.  $\square$

We need to know a precise description of  $\pi_1(\text{hAut}_*(U_{g,1}^n) \otimes \mathbb{Q}) \cong \pi_1(\text{hAut}_*((U_{g,1}^n)_{\mathbb{Q}}))$ , not merely its dimension which we just computed. We identify

$$\pi_1(\text{hAut}_*(U_{g,1}^n)) \cong \pi_2(\text{BhAut}_*(U_{g,1}^n)) = H_2((\text{BhAut}_*(U_{g,1}^n))^{\text{id}}; \mathbb{Z}).$$

An element  $\gamma \in \pi_2(\text{BhAut}_*(U_{g,1}^n))$  classifies a fibration  $E_\gamma \rightarrow S^2$ , together with a cross-section and a homotopy equivalence of the fibre over the basepoint with  $U_{g,1}^n$ .

The only potentially nonzero differential of the homological Leray–Serre spectral sequence of  $E_\gamma \rightarrow S^2$  is the map

$$d_{2,n}^2 : E_{2,n}^2 \rightarrow E_{0,n+1}^2,$$

which can be rewritten as

$$d(\gamma) : N(g)_{\mathbb{Z}} \cong H_2(S^2; N(g)_{\mathbb{Z}}) \rightarrow H_0(S^2; N(g)_{\mathbb{Z}}^{\vee}) \cong N(g)_{\mathbb{Z}}^{\vee}.$$

Therefore, assigning  $\gamma \mapsto d(\gamma)$  gives a map

$$d : \pi_2(\text{BhAut}_*(U_{g,1}^n)) \rightarrow \text{Hom}(N(g)_{\mathbb{Z}}; N(g)_{\mathbb{Z}}^{\vee}). \quad (4.14)$$

Similarly, we obtain

$$d_{\mathbb{Q}} : \pi_2(\text{BhAut}_*((U_{g,1}^n)_{\mathbb{Q}})) \rightarrow \text{Hom}(N(g); N(g)^{\vee}). \quad (4.15)$$

Source and target of (4.14) are  $\mathbb{Z}[\pi_0(\text{hAut}_*(U_{g,1}^n))]$ -modules; and source and target of (4.15) are  $\mathbb{Q}[\pi_0(\text{hAut}_*(U_{g,1}^n)_{\mathbb{Q}})]$ -modules.

**Lemma 4.16.** *The map  $d$  is a homomorphism of  $\mathbb{Z}[\pi_0(\text{hAut}_*(U_{g,1}^n))]$ -modules, and similarly  $d_{\mathbb{Q}}$  is a homomorphism of  $\mathbb{Q}[\pi_0(\text{hAut}_*(U_{g,1}^n)_{\mathbb{Q}})]$ -modules. When  $n \geq 4$ ,  $d_{\mathbb{Q}}$  is an isomorphism.*

*Proof.* The group  $\pi_0(\text{hAut}_*(U_{g,1}^n))$  acts on  $\pi_2(\text{BhAut}_*(U_{g,1}^n))$  by changing the identification of the fibres. Therefore, it is clear that  $d$  is  $\pi_0(\text{hAut}_*(U_{g,1}^n))$ -equivariant; and the same argument applies to  $d_{\mathbb{Q}}$ .

That  $d$  and  $d_{\mathbb{Q}}$  are additive requires a more detailed argument. The argument is the same in both cases, so we concentrate on  $d$ . Let  $\gamma_0, \gamma_1$  be two elements of  $\pi_2(\text{BhAut}_*(U_{g,1}^n))$ . Identification of the fibres over the basepoint gives a fibration over  $S^2 \vee S^2$  which we denote by  $E_{\gamma_0} \vee E_{\gamma_1} \rightarrow S^2 \vee S^2$ , abusing notation for simplicity. The fold map  $S^2 \vee S^2 \rightarrow S^2$  is covered by a fibrewise homotopy equivalence  $E_{\gamma_0} \vee$

$E_{\gamma_1} \rightarrow E_{\gamma_0 + \gamma_1}$ , and the inclusions  $S^2 \rightarrow S^2 \vee S^2$  are covered by fibrewise homotopy equivalences  $E_{\gamma_j} \rightarrow E_{\gamma_0} \vee E_{\gamma_1}$ . We obtain a commutative diagram

$$\begin{array}{ccccc} H_2(S^2; N(g)_{\mathbb{Z}}) \oplus H_2(S^2; N(g)_{\mathbb{Z}}) & \longrightarrow & H_2(S^2 \vee S^2; N(g)_{\mathbb{Z}}) & \longrightarrow & H_2(S^2; N(g)_{\mathbb{Z}}) \\ \downarrow d(\gamma_0) \oplus d(\gamma_1) & & \downarrow & & \downarrow d(\gamma_0 + \gamma_1) \\ H_0(S^2; N(g)_{\mathbb{Z}}^{\vee}) \oplus H_0(S^2; N(g)_{\mathbb{Z}}^{\vee}) & \longrightarrow & H_0(S^2 \vee S^2; N(g)_{\mathbb{Z}}^{\vee}) & \longrightarrow & H_0(S^2; N(g)_{\mathbb{Z}}^{\vee}) \end{array}$$

and the two horizontal composition maps are just the addition. Additivity of  $d$  follows. In a similar way, one proves that  $d_{\mathbb{Q}}$  is additive and hence  $\mathbb{Q}$ -linear.

For the second part of the proof, consider the diagram

$$\begin{array}{ccc} \pi_2(\mathrm{BhAut}_*(U_{g,1}^n)) & \xrightarrow{d} & \mathrm{Hom}(N(g)_{\mathbb{Z}}; N(g)_{\mathbb{Z}}^{\vee}) \\ \downarrow & & \downarrow \\ \pi_2(\mathrm{BhAut}_*((U_{g,1}^n)_{\mathbb{Q}})) & \xrightarrow{d_{\mathbb{Q}}} & \mathrm{Hom}(N(g); N(g)^{\vee}) \end{array} \quad (4.17)$$

which obviously commutes. We will show that

$$\ker(d) \subset \ker(\pi_2(\mathrm{BhAut}_*(U_{g,1}^n)) \rightarrow \pi_2(\mathrm{BhAut}_*((U_{g,1}^n)_{\mathbb{Q}}))). \quad (4.18)$$

Both vertical maps in (4.17) are rationalizations (the left one by Lemma 3.4), and so (4.18) shows that  $\ker(d)$  is finite. Hence  $d_{\mathbb{Q}}$  is injective, and Lemma 4.13, together with a dimension count, proves that  $d_{\mathbb{Q}}$  is an isomorphism.

To show (4.18), we must show that a fibration  $\pi : E \rightarrow S^2$  with fibre  $U_{g,1}^n$  and a cross-section whose Leray–Serre spectral sequence collapses at the  $E^2$ -stage is rationally fibre-homotopy equivalent to  $S^2 \times (U_{g,1}^n)_{\mathbb{Q}}$ . Since the spectral sequence collapses and since  $n \geq 3$ , the inclusion of the fibre  $U_{g,1}^n \rightarrow E$  induces isomorphisms on  $H_n$  and  $H_{n+1}$  and so gives isomorphisms  $N(g)_{\mathbb{Z}} \rightarrow H_n(E)$  and  $N(g)_{\mathbb{Z}}^{\vee} \rightarrow H_{n+1}(E)$ . From the inverses of those isomorphisms, we obtain a map

$$f : E \rightarrow K(N(g)_{\mathbb{Z}}, n) \times K(N(g)_{\mathbb{Z}}^{\vee}, n+1) \rightarrow K(N(g), n) \times K(N(g)^{\vee}, n+1)$$

inducing an isomorphism on rational homology in degrees  $n$  and  $n+1$ . The natural map  $(U_{g,1}^n)_{\mathbb{Q}} \rightarrow K(N(g), n) \times K(N(g)^{\vee}, n+1)$ , which induces the identity on homology in degrees  $n$  and  $n+1$ , is  $(2n-1) \geq (n+3)$ -connected, and since the homotopy dimension of  $E$  is  $n+3$ , we can deform  $f$  to a map  $g : E \rightarrow (U_{g,1}^n)_{\mathbb{Q}}$ , such that the composition with the inclusion of the fibre is the rationalization map  $U_{g,1}^n \rightarrow (U_{g,1}^n)_{\mathbb{Q}}$ . The map

$$h := (\pi, g) : E \rightarrow S^2 \times (U_{g,1}^n)_{\mathbb{Q}}$$

over  $S^2$  induces an isomorphism in rational homology up to degree  $n+1$ . Using a spectral sequence comparison argument, we obtain that  $h$  is a rational homology equivalence, and hence also a rational homology equivalence on the fibres.  $\square$

**Proposition 4.19.** *We have*

$$\pi_k(\mathrm{BhAut}_{\partial}(U_{g,1}^n)) \otimes \mathbb{Q} = 0$$

for  $3 \leq k \leq n-3$ . If  $n \geq 5$ , the map

$$\pi_2(\mathrm{BhAut}_{\partial}(U_{g,1}^n)) \otimes \mathbb{Q} \rightarrow \pi_2(\mathrm{BhAut}_*(U_{g,1}^n)) \otimes \mathbb{Q}$$

is injective and can be identified,  $\pi_0(\mathrm{hAut}_\partial((U_{g,1}^n)_\mathbb{Q}))$ -equivariantly, with the inclusion of

$$\begin{cases} S^2(N(g)^\vee) & n \text{ even} \\ \Lambda^2(N(g)^\vee) & n \text{ odd} \end{cases}$$

into  $N(g)^\vee \otimes N(g)^\vee = \mathrm{Hom}(N(g); N(g)^\vee)$ .

*Proof.* From the fibre sequence (4.9), we obtain an exact sequence

$$\pi_{2n+k}(U_{g,1}^n) \otimes \mathbb{Q} \pi_k(\mathrm{BhAut}_\partial(U_{g,1}^n)) \otimes \mathbb{Q} \rightarrow \pi_k(\mathrm{BhAut}_*(U_{g,1}^n)) \otimes \mathbb{Q}.$$

By (4.6), the leftmost group vanishes when  $2 \leq k \leq n-3$ . Together with Lemma 4.13, the first claim follows, as well as the injectivity on  $\pi_2$ .

For the remaining claim, consider the exact sequence

$$\begin{aligned} 0 \rightarrow \pi_2(\mathrm{BhAut}_\partial((U_{g,1}^n)_\mathbb{Q})) &\rightarrow \pi_2(\mathrm{BhAut}_*((U_{g,1}^n)_\mathbb{Q})) \rightarrow \pi_{2n+1}((U_{g,1}^n)_\mathbb{Q}) \rightarrow \\ &\rightarrow \pi_1(\mathrm{BhAut}_\partial((U_{g,1}^n)_\mathbb{Q})) \rightarrow \pi_1(\mathrm{BhAut}_*((U_{g,1}^n)_\mathbb{Q})) \rightarrow \pi_{2n}((U_{g,1}^n)_\mathbb{Q}) \end{aligned}$$

derived from (4.9). The proof of Lemma 4.10 shows that  $\pi_1(\mathrm{BhAut}_\partial((U_{g,1}^n)_\mathbb{Q})) \rightarrow \pi_1(\mathrm{BhAut}_*((U_{g,1}^n)_\mathbb{Q}))$  is injective, so we obtain a short exact sequence

$$0 \rightarrow \pi_2(\mathrm{BhAut}_\partial((U_{g,1}^n)_\mathbb{Q})) \rightarrow \pi_2(\mathrm{BhAut}_*((U_{g,1}^n)_\mathbb{Q})) \rightarrow \pi_{2n+1}((U_{g,1}^n)_\mathbb{Q}) \rightarrow 0. \quad (4.20)$$

The vector space  $\pi_{2n+1}((U_{g,1}^n)_\mathbb{Q})$  is generated by the Whitehead brackets  $[y_i, y_j]$ , modulo the relations that are universally satisfied by Whitehead brackets of two elements of degree  $n+1$ , that is

$$[y_i, y_j] = (-1)^{(n+1)^2} [y_j, y_i] = -(-1)^n [y_j, y_i].$$

It follows that

$$\pi_{2n+1}((U_{g,1}^n)_\mathbb{Q}) \cong \begin{cases} S^2(N(g)^\vee) & n \text{ odd} \\ \Lambda^2(N(g)^\vee) & n \text{ even.} \end{cases} \quad (4.21)$$

The sequence (4.20) is a sequence of  $\pi_0(\mathrm{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})) = \mathrm{GL}(N(g))$ -modules, and as  $S^2(N(g)^\vee)$ , as well as  $\Lambda^2(N(g)^\vee)$  are irreducible  $\mathrm{GL}(N(g))$ -modules, the second map in (4.20) must, up to multiplication by a nonzero constant, agree with the natural projection from  $\mathrm{Hom}(N(g); N(g)^\vee) = (N(g)^\vee)^{\otimes 2}$  to  $S^2(N(g)^\vee)$  or  $\Lambda^2(N(g)^\vee)$ . Therefore, the kernel of these maps are as asserted.  $\square$

**4.4. The spectral sequence for tangential homotopy automorphisms.** Lemma 4.10 (3) and Proposition 3.25 prove the following.

**Corollary 4.22.** *The algebra map*

$$H^*(\mathrm{map}_\partial((U_{g,1}^n)_\mathbb{Q}; \mathrm{BO}_\mathbb{Q})^0 // \mathrm{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})_{\mathbb{Z}}) \rightarrow H^*(\widetilde{\mathrm{BDiff}}_\partial(U_{g,1}^n))$$

is surjective if  $n \geq 3$ , and the kernel is the ideal generated by the classes  $\kappa_{L_m}$ ,  $m \in \mathbb{N}$ .  $\square$

Hence we need to calculate the cohomology of  $\mathrm{map}_\partial((U_{g,1}^n)_\mathbb{Q}; \mathrm{BO}_\mathbb{Q})^0 // \mathrm{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})_{\mathbb{Z}}$ , and we shall approach this through the spectral sequences for the two fibrations

$$\mathrm{map}_\partial((U_{g,1}^n)_\mathbb{Q}; \mathrm{BO}_\mathbb{Q})^0 // \mathrm{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\mathrm{id}} \rightarrow \mathrm{map}_\partial((U_{g,1}^n)_\mathbb{Q}; \mathrm{BO}_\mathbb{Q})^0 // \mathrm{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})_{\mathbb{Z}} \rightarrow \mathrm{BGL}(N(g)_{\mathbb{Z}}), \quad (4.23)$$

(where we identified the base space using  $\pi_0(\mathrm{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})_{\mathbb{Z}}) \cong \mathrm{GL}(N(g)_{\mathbb{Z}})$  which is valid when  $n \geq 3$  by Lemma 4.10), and

$$\mathrm{map}_\partial((U_{g,1}^n)_\mathbb{Q}; \mathrm{BO}_\mathbb{Q})^0 \rightarrow \mathrm{map}_\partial((U_{g,1}^n)_\mathbb{Q}; \mathrm{BO}_\mathbb{Q})^0 // \mathrm{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\mathrm{id}} \rightarrow \mathrm{BhAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\mathrm{id}}. \quad (4.24)$$

In this section, we are concerned with (4.24). Let us start with a crucial observation.

*Observation 4.25.* Consider generally a fibration  $p : E \rightarrow B$  over a 0-connected pointed  $B$  and let  $F := p^{-1}(b)$  be the fibre. Denote  $\Gamma := \pi_1(B, b)$ . The monodromy of  $p$  gives a homomorphism  $\mu : \Gamma \rightarrow \pi_0(\mathrm{hAut}(F))$ . Now consider the pullback

$$\begin{array}{ccc} \tilde{E} & \longrightarrow & E \\ \downarrow \tilde{p} & & \downarrow p \\ \tilde{B} & \longrightarrow & B \end{array}$$

along the universal covering  $\tilde{B} \rightarrow B$ ; the horizontal maps are  $\Gamma$ -Galois coverings. The rational cohomological spectral sequence  $\tilde{E}^{p,q}$  for  $\tilde{p}$  is a spectral sequence of  $\mathbb{Q}[\Gamma]$ -modules, just by naturality of the construction of the spectral sequence; in particular the isomorphism  $\tilde{E}_2^{p,0} \otimes \tilde{E}^{0,q} \rightarrow \tilde{E}_2^{p,q}$  given by the product structure is  $\Gamma$ -equivariant.

Now the usual identification  $H^q(F; \mathbb{Q}) \cong \tilde{E}_2^{0,q}$  is  $\Gamma$ -equivariant, where the action on the second group comes from the fact that  $\tilde{p}$  is  $\Gamma$ -equivariant, and the action on the first group comes from the monodromy of  $p$ .

We apply this general observation to the fibration  $p : \mathrm{map}_\partial((U_{g,1}^n)_\mathbb{Q}; \mathrm{BO}_\mathbb{Q})^0 // \mathrm{hAut}_\partial((U_{g,1}^n)_\mathbb{Q}) \rightarrow \mathrm{BhAut}_\partial((U_{g,1}^n)_\mathbb{Q})$ ; then  $\tilde{p}$  is just  $\mathrm{map}_\partial((U_{g,1}^n)_\mathbb{Q}; \mathrm{BO}_\mathbb{Q})^0 // \mathrm{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\mathrm{id}} \rightarrow \mathrm{BhAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\mathrm{id}}$ . We obtain that the spectral sequence of (4.24) is a spectral sequence of  $\mathrm{GL}(N(g))$ -representations. Moreover, the action on the  $E_2$ -term

$$E_2^{p,q} \cong H^p(\mathrm{BhAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\mathrm{id}}; \mathbb{Q}) \otimes H^q(\mathrm{map}_\partial((U_{g,1}^n)_\mathbb{Q}; \mathrm{BO}_\mathbb{Q})^0; \mathbb{Q}) \quad (4.26)$$

is the tensor product of the two actions. Corollary 3.22 computes the second factor in terms of the homology of  $U_{g,1}^n$ , and Corollary 3.23 states that the action is determined by the action of  $\mathrm{GL}(N(g))$  on the homology.

Combining our work so far, the  $E_2$ -term of the spectral sequence of (4.24) is readily computed in a range of degrees. Let us introduce some notation.

**Notation 4.27.** (1) We denote

$$L^2(N(g)) := \begin{cases} S^2(N(g)) & n \text{ even} \\ \Lambda^2(N(g)) & n \text{ odd.} \end{cases}$$

(2) We fix an integer  $M$ ; everything that matters is that  $M$  is large enough (see 4.31).

(3) Let  $V(n)$  be the graded vector space

$$V(n) := \bigoplus_{4m-2n-1 > 0, m \leq M} \mathbb{Q}[4m-2n-1],$$

(4) let  $U(n)$  be the graded vector space

$$U(n) := \bigoplus_{4m-n-1 > 0, m \leq M} \mathbb{Q}[4m-n-1]$$

(5) and let  $W(n)$  be the graded vector space

$$W(n) := \bigoplus_{4m-n > 0, m \leq M} \mathbb{Q}[4m-n].$$

**Definition 4.28.** Let  $n \geq 6$  be even. We define a map

$$S^*(S^2(N(g))) \otimes \Lambda^*(V(n)) \otimes S^*(N(g) \otimes W(n)) \otimes \Lambda^*(N(g)^\vee \otimes U(n)) \rightarrow E_2^{*,*} \quad (4.29)$$

of bigraded algebras as follows.

- On  $S^2(N(g))$ , which is in bidegree  $(2, 0)$ , it is the isomorphism  $S^2(N(g)) \cong H^2(\text{BhAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\text{id}})$  from Proposition 4.19.
- On  $V(n)_{4m-n-1} = \mathbb{Q}$ , which is in bidegree  $(0, 4m-n-1)$ , it is the map which sends 1 to the class  $\lambda_{[U_{g,1}^n], L_m} \in H^{4m-2n-1}(\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0) = E_2^{0, 4m-2n-1}$  (using the notations introduced in §4.1 and Definition 3.15).
- On  $N(g) \otimes W(n)_{4m-n} = N(g)$ , which is in bidegree  $(0, 4m-n)$ , it is the map which sends  $a \in N(g)$  to  $\lambda_{a, L_m} \in H^{4m-n}(\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0) = E_2^{0, 4m-n}$ .
- On  $N(g)^\vee \otimes U(n)_{4m-n-1} = N(g)^\vee$ , which is in bidegree  $(0, 4m-n-1)$ , it sends  $b \in N(g)^\vee$  to  $\lambda_{b, L_m} \in H^{4m-n-1}(\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0) = E_2^{4m-n-1, 0}$ .

We similarly define for odd  $n \geq 5$  a map

$$S^*(\Lambda^2(N(g))) \otimes \Lambda^*(V(n)) \otimes \Lambda^*(N(g) \otimes W(n)) \otimes S^*(N(g)^\vee \otimes U(n)) \rightarrow E_2^{*,*} \quad (4.30)$$

by the analogous formulas.

**Proposition 4.31.** *The maps (4.29) and (4.30) are  $\pi_0(\text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})) \cong \text{GL}(N(g))$ -equivariant. If  $n \geq 5$ , they are isomorphisms in bidegrees  $(p, q)$  with  $p \leq n-3$  and  $q \leq 4M-2n+2$ .*

*Proof.* The dual of the equivariant isomorphism  $L^2(N(g))^\vee \cong H_2(\text{BhAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\text{id}}; \mathbb{Q})$  from Proposition 4.19 extends to an equivariant algebra map

$$S^*(L^2(N(g))) \rightarrow H^*(\text{BBhAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\text{id}}; \mathbb{Q})$$

which is an isomorphism up to degree  $n-3$ , also by Proposition 4.19 and the well-known computation of the cohomology of rational Eilenberg–Mac-Lane spaces.

The other statements are clear from the naturality of the  $\lambda$ -classes that was recorded after Definition 3.15, and from Corollary 3.22 and Proposition 4.19.  $\square$

**Proposition 4.32.** *Assume  $n \geq 5$  and  $g \geq 2$ . With respect to the isomorphism of Proposition 4.31, the  $d_2$ -differential in the spectral sequence of the fibre sequence (4.24) is given on generators as follows.*

- (1) On  $L^2(N(g))$ , it is zero.
- (2) On  $V(n)$ , it is zero.
- (3) On  $N(g)^\vee \otimes U(n)$ , it is zero.
- (4) On  $N(g) \otimes W(n)$ , it is of the form

$$N(g) \otimes W(n) \xrightarrow{p \otimes S} L^2(N(g)) \otimes N(g)^\vee \otimes U(n)$$

where  $p : N(g) \rightarrow L^2(N(g)) \otimes N(g)^\vee$  is adjoint to the projection  $N(g)^{\otimes 2} \rightarrow L^2(N(g))$  and  $S : W(n) \rightarrow U(n)$  is a degree  $-1$  map whose restriction  $W(n)_{4m-n} \rightarrow U(n)_{4m-n-1}$  is an isomorphism unless  $4m-n-1 = 0$ .

*Proof.* Item (1) holds for degree reasons. For item (2), note that the inclusion map  $\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0 \rightarrow \text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0 // \text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\text{id}}$  pulls back  $\kappa_{L_m}$  to the class  $\lambda_{[U_{g,1}^n], L_m}$ . Therefore, the latter class is a permanent cycle, which verifies (2).

For the other two claims, we use the  $\text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})$ -equivariant homotopy equivalence

$$\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0 \simeq \prod_{m \geq 1} \text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; K(\mathbb{Q}, 4m))^0$$

given by the cohomology classes  $(L_m)_{m \geq 1}$ . Since the latter homotopy equivalence, combined with the projection onto the  $m$ th factor, pulls back  $\lambda_{a, u_m} \in H^*(\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; K(\mathbb{Q}, 4m))^0; \mathbb{Q})$  to  $\lambda_{a, L_m}$ , the proposition follows from the next Lemma.  $\square$

**Lemma 4.33.** *The  $E_2$ -term of the spectral sequence of the fibration*

$$\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; K(\mathbb{Q}, k))^0 // \text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\text{id}} \rightarrow \text{BhAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\text{id}} \quad (4.34)$$

is (in the columns up to degree  $n - 3$ , by Proposition 4.19) given by

$$\mathbb{F}(L^2(N(g)))[2, 0] \oplus N(g)[0, k - n] \oplus N(g)^\vee[0, k - n - 1]$$

(if  $k - n - 1 > 0$ ; if  $k - n - 1 = 0$ , the last summand is dropped and if  $k - n \leq 0$ , the last two summands are dropped). The differential  $d_2$  vanishes on  $N(g)^\vee[0, k - n - 1]$ , and on  $N(g)[0, k - n]$ , it is, up to a sign, the map adjoint to the projection  $N(g)^{\otimes 2} \rightarrow L^2(N(g))$ , if  $k - n - 1 > 1$ .

*Proof.* The differential vanishes on  $N(g)^\vee[0, k - n - 1]$  for degree reasons, and because the fibration (4.34) has a section. The claim about  $N(g)[0, k - n]$  is true for the same reason if  $k - n - 1 = 0$ , and so we may suppose  $k - n - 1 > 0$ . We pair the spectral sequence of (4.34) with the spectral sequence of the universal fibration

$$(U_{g,1}^n)_\mathbb{Q} // \text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\text{id}} \rightarrow \text{BhAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\text{id}}. \quad (4.35)$$

We write  $\tilde{E}_*^{*,*}$  for the rational cohomological spectral sequence of the latter. The  $E_2$ -term is of the form

$$\tilde{E}_2 = S^*(L^2 N(g)) \otimes H^*(U_{g,1}^n; \mathbb{Q})$$

(in the columns up to degree  $(n - 3)$ , by Proposition 4.19). Let  $x_{ij} \in L^2(N_g)$  the image of  $a_i \otimes a_j \in N_g^{\otimes 2}$  under the projection map; these elements satisfy  $x_{ji} = (-1)^n x_{ij}$ . By Proposition 4.19), the differential  $\tilde{d}_2 : \tilde{E}_2^{0, n+1} \rightarrow \tilde{E}_2^{2, n}$  is given on basis elements by the formulas

$$\tilde{d}_2(\beta_i) = \sum_j x_{ij} \otimes \alpha_j \quad (4.36)$$

and

$$\tilde{d}_2(\alpha_i) = 0. \quad (4.37)$$

To transfer this knowledge about  $\tilde{E}_*^{*,*}$  to information about  $E_*^{*,*}$ , observe that the evaluation map  $\text{ev} : U_{g,1}^n \times \text{map}_\partial(U_{g,1}^n; K(\mathbb{Q}, k))^0 \rightarrow K(\mathbb{Q}, k)$  is  $\text{hAut}_\partial(U_{g,1}^n)$ -equivariant and hence induces a map

$$(U_{g,1}^n \times \text{map}_\partial(U_{g,1}^n; K(\mathbb{Q}, k))^0) // \text{hAut}_\partial(U_{g,1}^n) \rightarrow K(\mathbb{Q}, k).$$

It follows that  $\text{ev}^*u_k \in H^k(U_{g,1}^n \times \text{map}_\partial(U_{g,1}^n; K(\mathbb{Q}, k))^0)$  lies in the image of the map induced by the inclusion

$$U_{g,1}^n \times \text{map}_\partial(U_{g,1}^n; K(\mathbb{Q}, k))^0 \rightarrow (U_{g,1}^n \times \text{map}_\partial(U_{g,1}^n; K(\mathbb{Q}, k))^0) // \text{hAut}_\partial(U_{g,1}^n)$$

of the fibre in the total space. This forces  $\text{ev}^*u_k \in \hat{E}_2^{0,k}$  to be a permanent cycle in the spectral sequence  $\hat{E}_*^{*,*}$  of

$$(U_{g,1}^n \times \text{map}_\partial(U_{g,1}^n; K(\mathbb{Q}, k))^0) // \text{hAut}_\partial((U_{g,1}^n)_{\mathbb{Q}})^{\text{id}} \rightarrow \text{BhAut}_\partial((U_{g,1}^n)_{\mathbb{Q}})^{\text{id}},$$

so  $\hat{d}_2(\text{ev}^*u_k) = 0$ . By formula (3.16)

$$\text{ev}^*u_k = \sum_{i=1}^g \alpha_i \times \lambda_{a_i, u_k} + \sum_{i=1}^g \beta_i \times \lambda_{b_i, u_k}.$$

Using (4.36) and (4.37), and that  $d_2(\lambda_{b_i, u_k}) = 0$  for degree reasons, we obtain

$$\begin{aligned} 0 &= \sum_{i=1}^g \hat{d}_2(\alpha_i \times \lambda_{a_i, u_k}) + \hat{d}_2(\beta_i \times \lambda_{b_i, u_k}) = \\ &= \sum_{i=1}^g (d_2\alpha_i) \times \lambda_{a_i, u_k} + \sum_{i=1}^g (-1)^n \alpha_i \times (d_2\lambda_{a_i, u_k}) + \sum_{i=1}^g (\tilde{d}_2\beta_i) \times \lambda_{b_i, u_k} + \sum_{i=1}^g (-1)^{n+1} \beta_i \times (\tilde{d}_2\lambda_{b_i, u_k}) = \\ &= \sum_{j=1}^g (-1)^n \alpha_j \times (d_2\lambda_{a_j, u_k}) + \sum_{i,j=1}^g x_{ij} \otimes (\alpha_j \times \lambda_{b_i, u_k}). \end{aligned}$$

Comparing coefficients leads to the identity

$$d_2\lambda_{a_j, u_k} = (-1)^{n+1} \sum_{i=1}^g x_{ij} \otimes \lambda_{b_i, u_k}$$

which finishes the proof.  $\square$

Propositions 4.31 and 4.32 were the goals of this section. Before we can use these to evaluate the two spectral sequences for (4.23) and (4.24), we need to switch gears and introduce some more algebraic background.

## 5. A REPRESENTATION-THEORETIC CALCULATION

### 5.1. Generalities.

*Rational representations.* Let  $\mathbb{K}$  be a field of characteristic 0, and let  $N$  be a finite-dimensional  $\mathbb{K}$ -vector space. Recall that a representation  $\rho : \text{GL}(N) \rightarrow \text{GL}(W)$  on some other  $\mathbb{K}$ -vector space is *rational* if the matrix entries (after choice of a basis) of  $\rho(g)$  are polynomial functions of the matrix entries of  $g$  and of  $\det(g)^{-1}$ . A similar definition applies to representations of  $\text{SL}(N)$ .

Let  $\mathbb{K} \subset \mathbb{L}$  be a field extension, let  $N$  and  $W$  be  $\mathbb{K}$ -vector spaces and let  $\rho : \text{GL}(N) \rightarrow \text{GL}(W)$  be rational. Then  $\rho$  extends to a rational representation  $\rho_{\mathbb{L}} : \text{GL}(N_{\mathbb{L}}) \rightarrow \text{GL}(W_{\mathbb{L}})$ . A similar statement is true for  $\text{SL}(N)$ .

If there is  $w \in \mathbb{Z}$  such that each element  $\lambda \in \mathbb{K}^\times \subset \text{GL}(N)$  in the centre acts by multiplication with  $\lambda^w$  on  $W$ , we say that  $\rho$  has *weight*  $w$ . We shall use the notation  $T^{k,l}(N) := N^{\otimes k} \otimes (N^\vee)^{\otimes l}$ .

The rational representations are described in terms of Schur functors, whose definition we briefly recall. Let  $\mathcal{P}_n$  be the set of partitions of  $n$ , thought of as Young diagrams. If  $\lambda$  is a partition of  $n$ , we also write  $|\lambda| = n$ . The *height*  $\text{ht}(\lambda)$

of  $\lambda \in \mathcal{P}_n$  is the number of rows of  $\lambda$ . To a partition  $\lambda$  of  $n$ , there is associated the Young symmetrizer  $c_\lambda \in \mathbb{Q}[\Sigma_n] \subset \mathbb{K}[\Sigma_n]$  and the associated irreducible  $\Sigma_n$ -representation  $M_\lambda$  (over  $\mathbb{K}$ ; it is irreducible since it is irreducible when the scalars are extended to the algebraic closure  $\overline{\mathbb{K}}$ ). The tensor power  $T^{n,0}(N)$  has a canonical  $\mathrm{GL}(N) \times \Sigma_n$ -representation, and the Schur functor  $S_\lambda(N)$  is defined as the  $\mathrm{GL}(N)$ -representation

$$S_\lambda(N) := c_\lambda \cdot T^{n,0}(N).$$

The theory of rational representations can be summarized in the following result.

**Theorem 5.1.** *Let  $N$  be a finite-dimensional  $\mathbb{K}$ -vector space.*

- (1) *As  $\mathrm{GL}(N) \times \Sigma_n$ -modules, we have*

$$T^{n,0}(N) = \bigoplus_{\lambda \in \mathcal{P}_n} S_\lambda(N) \otimes M_\lambda.$$

*If  $\mathrm{ht}(\lambda) > \dim(N)$ , then  $S_\lambda(N) = 0$ , and if  $\mathrm{ht}(\lambda) \leq \dim(N)$ ,  $S_\lambda(N)$  is nonzero and irreducible. The Schur functors  $S_\lambda(N)$  and  $S_\mu(N)$  are isomorphic only if they both vanish or if  $\lambda = \mu$ . The Schur functors remain irreducible when viewed as  $\mathrm{SL}(N)$ -modules.*

- (2) *After taking the tensor product with a suitable power of the determinant representation  $\det(N)$ , each  $\mathrm{GL}(N)$ -representation embeds into a  $\bigoplus_j T^{n_j,0}(N)$  for suitable  $n_j$ . The same is true for  $\mathrm{SL}(N)$ -representations instead.*
- (3) *Rational  $\mathrm{GL}(N)$ - and  $\mathrm{SL}(N)$ -representations are completely reducible. The Schur functors give a complete list of the irreducible  $\mathrm{SL}(N)$ -representations, and the Schur functors, tensorized with powers of the determinant representation, a complete list of the irreducible  $\mathrm{GL}(N)$ -representations.*
- (4) *Let  $N, V, W$  be  $\mathbb{K}$  vector spaces, let  $\mathrm{GL}(N) \rightarrow \mathrm{GL}(V)$  and  $\mathrm{GL}(N) \rightarrow \mathrm{GL}(W)$  be rational representations and let  $\mathbb{K} \subset \mathbb{L}$  be a field extension. Then  $W$  is irreducible if and only if  $W_{\mathbb{L}}$  is irreducible;  $V$  and  $W$  are isomorphic if and only if  $V_{\mathbb{L}}$  and  $W_{\mathbb{L}}$  are isomorphic, and furthermore*

$$(W^{\mathrm{GL}(N)})_{\mathbb{L}} = (W_{\mathbb{L}})^{\mathrm{GL}(N_{\mathbb{L}})}.$$

*The same is true for  $\mathrm{SL}$  in place of  $\mathrm{GL}$ .*

*References.* Statements (1)–(3) are well-known when  $\mathbb{K}$  is algebraically closed; [60] and [23] are convenient textbook references. We explain how to generalize the results to arbitrary  $\mathbb{K}$ .

(1) The decomposition is shown for algebraically closed fields (of characteristic 0) in [60, Theorem 9.3.1.4], but it exists over any ground field of characteristic 0 because the Young symmetrizer have rational coefficients. For a field extension  $\mathbb{K} \subset \mathbb{L}$ , we have  $S_\lambda(N_{\mathbb{L}}) \cong (S_\lambda(N))_{\mathbb{L}}$  as  $\mathrm{GL}(N_{\mathbb{L}})$ -representations. Hence the other statements follow from the algebraically closed case, which is done in [23, Theorem 6.3.1] and [60, Theorem 9.8.1].

(2) The proof of [60, Lemma 7.1.4] does not use that  $\mathbb{K}$  is algebraically closed.

(3) The first part follows from (2), since  $T^{n,0}(N)$  is completely reducible, and by generalities on completely reducible representations, e.g. [19, Proposition 3.1.4]. The second part follows from the algebraically closed case which is [60, Theorem 9.8.1].

(4) This is clear from the other items. □

**Definition 5.2.** Let  $\rho : \mathrm{SL}(N) \rightarrow \mathrm{GL}(W)$  be a rational representation. We say that  $\rho$  (or  $W$ ) has  $\mathrm{load} \leq n$  if each irreducible summand of  $W$  is a direct summand of  $T^{k,l}(N)$ , for some  $k, l$  with  $k + l \leq n$ .

*Invariants on tensor spaces.* We need the fundamental theorem of invariant theory of  $\mathrm{GL}(N)$ . We can identify  $T^{m,m}(N) \cong \mathrm{End}(N)^{\otimes m} \cong \mathrm{End}(N^{\otimes m})$ . There is a natural map

$$\sigma_{N,m} : \mathbb{K}[\Sigma_m] \rightarrow \mathrm{End}(N^{\otimes m})^{\mathrm{GL}(N)} \cong T^{m,m}(N)^{\mathrm{GL}(N)} \quad (5.3)$$

given by the  $\Sigma_m$ -action permuting the factors. In terms of a basis  $(a_1, \dots, a_g)$  of  $N$  and the dual basis  $(a^1, \dots, a^g)$  of  $N^\vee$ ,  $\sigma_{N,m}$  is given by the formula

$$s \mapsto \sum_{i_1, \dots, i_m} a_{i_1} \otimes \dots \otimes a_{i_m} \otimes a^{i_{s^{-1}(1)}} \otimes \dots \otimes a^{i_{s^{-1}(m)}} \in T^{m,m}(N). \quad (5.4)$$

**Theorem 5.5.** *Let  $N$  be a finite-dimensional  $\mathbb{K}$ -vector space. The map  $\sigma_{N,m}$  is surjective, and it is also injective if  $m \leq \dim(N)$ .*

Surjectivity is the content of the first fundamental theorem which is proven in e.g. [60, §9.1.2], [48, Theorem 9.1.2] or [29, Theorem 5.3.1] (and is a key ingredient for Theorem 5.1). Injectivity is the second fundamental theorem. The treatment of that result in [60] or [29] has a slightly different layout; the version as stated above is shown in [48, Theorem 9.1.3].

**Lemma 5.6.** *Let  $N$  be a finite-dimensional  $\mathbb{K}$ -vector space. Then*

- (1)  $T^{k,l}(N)^{\mathrm{GL}(N)} = 0$  unless  $k = l$ .
- (2)  $T^{k,l}(N)^{\mathrm{SL}(N)} = 0$  unless  $\dim(N)$  divides  $k - l$ .
- (3)  $T^{k,k}(N)^{\mathrm{SL}(N)} = T^{k,k}(N)^{\mathrm{GL}(N)}$ .

*Proof.* (1) is easy; just look at the action of a scalar matrix. (2) Use Theorem 5.1 (4) to replace  $\mathbb{K}$  by its algebraic closure. Then  $\mathrm{SL}(N)$  contains a primitive  $g$ th root of unity  $\zeta_g$ , where  $g = \dim(N)$ , which acts by  $\zeta_g^{k-l}$  on  $T^{k,l}(N)$ . Hence  $T^{k,l}(N)^{\mathrm{SL}(N)} = 0$  unless  $g$  divides  $k - l$ . (3) Use Theorem 5.1 (4) to replace  $\mathbb{K}$  by its algebraic closure. The canonical isomorphism  $T^{k,k}(N) \cong \mathrm{End}(N^{\otimes k})$  identifies  $T^{k,k}(N)^{\mathrm{GL}(N)}$  with the commutant algebra of the image of  $\rho_{\mathrm{GL}} : \mathbb{K}[\mathrm{GL}(N)] \rightarrow \mathrm{End}(N^{\otimes k})$ . Similarly,  $T^{k,k}(N)^{\mathrm{SL}(N)}$  is the commutant of the image of  $\rho_{\mathrm{SL}} : \mathbb{K}[\mathrm{SL}(N)] \rightarrow \mathrm{End}(N^{\otimes k})$ , so it is enough to prove that  $\rho_{\mathrm{GL}}$  and  $\rho_{\mathrm{SL}}$  have the same image. Each element  $A \in \mathrm{GL}(N)$  can be written as  $A = \lambda B$  with  $\lambda \in \mathbb{K}^\times$  and  $B \in \mathrm{SL}(V)$  (here we are using that  $\mathbb{K}$  is algebraically closed). It follows that  $\rho_{\mathrm{GL}}(A) = \lambda^k \rho_{\mathrm{SL}}(B) \in \mathrm{im}(\rho_{\mathrm{SL}})$ .  $\square$

**5.2. A special invariant calculation.** In this section, we carry out a representation-theoretic calculation that will be used later on. The main ideas for the proof were communicated to us by Jerzy Weyman, and we thank him for allowing to reproduce his argument here.

Let  $\mathbb{K}$  be a field of characteristic 0, and let  $N, W$  and  $U$  be finite-dimensional  $\mathbb{K}$ -vector spaces. We let  $g := \dim(N)$  and fix a basis  $(a_1, \dots, a_g)$  of  $N$ , with dual basis  $(a^1, \dots, a^g)$ . We consider the algebras

$$A := S^*(S^2(N)) \otimes S^*(N \otimes W) \otimes \Lambda^*(N^\vee \otimes U)$$

and

$$C := S^*(\Lambda^2(N)) \otimes \Lambda^*(N \otimes W) \otimes S^*(N^\vee \otimes U).$$

These algebras have an obvious trigrading and actions of the group  $\mathrm{GL}(N) \times \mathrm{GL}(W) \times \mathrm{GL}(U)$ . We want to determine the algebras  $A^{\mathrm{GL}(N)}$  and  $C^{\mathrm{GL}(N)}$  of  $\mathrm{GL}(N)$ -invariants. There are some obvious invariants. We define

$$\varphi_A : \Lambda^2(U) \rightarrow (S^1(S^2(N)) \otimes S^0(N \otimes W) \otimes \Lambda^2(N^\vee \otimes U))^{\mathrm{GL}(N)} = A_{1,0,2}^{\mathrm{GL}(N)}$$

by

$$u_1 \wedge u_2 \mapsto \sum_{i,j=1}^g (a_i \cdot a_j) \otimes 1 \otimes ((a^i \otimes u_1) \wedge (a^j \otimes u_2)),$$

and we define

$$\psi_A : W \otimes U \rightarrow (S^0(S^2(N)) \otimes S^1(N \otimes W) \otimes \Lambda^1(N^\vee \otimes U))^{\mathrm{GL}(N)} = A_{0,1,1}^{\mathrm{GL}(N)}$$

by

$$w \otimes u \mapsto \sum_{i=1}^g 1 \otimes (a_i \otimes w) \otimes (a^i \otimes u).$$

For  $x_0, x_1 \in \Lambda^2(U)$  and  $y_0, y_1 \in W \otimes U$ , the relations

$$\varphi_A(x_0)\varphi_A(x_1) = \varphi_A(x_1)\varphi_A(x_0),$$

$$\psi_A(y_0)\psi_A(y_1) = -\psi_A(y_1)\psi_A(y_0)$$

and

$$\varphi_A(x_0)\psi_A(x_1) = \psi_A(x_1)\varphi_A(x_0)$$

hold, and these imply that  $\varphi_A \oplus \psi_A$  extends to an algebra map

$$G : S^*(\Lambda^2(U)) \otimes \Lambda^*(W \otimes U) \rightarrow A^{\mathrm{GL}(N)}.$$

Similarly let us define

$$\varphi_C : \Lambda^2(U) \rightarrow (S^1(\Lambda^2(N)) \otimes \Lambda^0(N \otimes W) \otimes S^2(N^\vee \otimes U))^{\mathrm{GL}(N)} = C_{1,0,2}^{\mathrm{GL}(N)} \subset C^{\mathrm{GL}(N)}$$

by

$$u_1 \wedge u_2 \mapsto \sum_{1 \leq i < j \leq g} (a_i \wedge a_j) \otimes 1 \otimes ((a^i \otimes u_1) \cdot (a^j \otimes u_2))$$

and

$$\psi_C : W \otimes U \rightarrow (S^0(\Lambda^2(N)) \otimes \Lambda^1(N \otimes W) \otimes S^1(N^\vee \otimes U))^{\mathrm{GL}(N)} = C_{0,1,1}^{\mathrm{GL}(N)} \subset C^{\mathrm{GL}(N)}$$

by

$$w \otimes u \mapsto \sum_{i=1}^g 1 \otimes (a_i \otimes w) \otimes (a^i \otimes u).$$

A similar argument as above shows that  $\varphi_C \oplus \psi_C$  extends to an algebra map

$$H : S^*(\Lambda^2(U)) \otimes \Lambda^*(W \otimes U) \rightarrow C^{\mathrm{GL}(N)}.$$

**Proposition 5.7.** *We have  $A_{p,q,r}^{\mathrm{GL}(N)} = C_{p,q,r}^{\mathrm{GL}(N)} = 0$  unless  $2p + q - r = 0$ , and  $A_{p,q,r}^{\mathrm{SL}(N)} = C_{p,q,r}^{\mathrm{SL}(N)} = 0$  unless  $g := \dim(N)$  divides  $2p + q - r$ . The maps*

$$S^p(\Lambda^2(U)) \otimes \Lambda^q(W \otimes U) \xrightarrow{G} A_{p,q,2p+q}^{\mathrm{GL}(N)} \subset A_{p,q,2p+q}^{\mathrm{SL}(N)}$$

and

$$S^p(\Lambda^2(U)) \otimes \Lambda^q(W \otimes U) \xrightarrow{H} C_{p,q,2p+q}^{\mathrm{GL}(N)} \subset C_{p,q,2p+q}^{\mathrm{SL}(N)}$$

are surjective, and isomorphisms as long as  $2p + q \leq g$ .

We use Proposition 5.7 in conjunction with Proposition 5.8 below. To state it, let us assume that  $\mathbb{K} = \mathbb{Q}$  and that  $N$  has an *integral form*, i.e. a subgroup  $N_{\mathbb{Z}} \subset N$  such that  $N_{\mathbb{Z}} \otimes \mathbb{Q} = N$ . In that case, we have the subgroup  $\mathrm{SL}(N_{\mathbb{Z}}) \subset \mathrm{SL}(N)$  of automorphisms preserving  $N_{\mathbb{Z}}$ ; note that  $\mathrm{SL}(N_{\mathbb{Z}}) \cong \mathrm{SL}_g(\mathbb{Z})$ .

**Proposition 5.8.** *If  $\mathbb{K} = \mathbb{Q}$  and  $N$  has an integral form  $N_{\mathbb{Z}}$ , the inclusions*

$$A_{p,q,2p+q}^{\mathrm{SL}(N)} \subset A_{p,q,2p+q}^{\mathrm{SL}(N_{\mathbb{Z}})}$$

and

$$C_{p,q,2p+q}^{\mathrm{SL}(N)} \subset C_{p,q,2p+q}^{\mathrm{SL}(N_{\mathbb{Z}})}$$

are equalities.

*Proof of Proposition 5.7, surjectivity.* The first sentence is a straightforward application of Lemma 5.6. It also follows from Lemma 5.6 that  $A_{p,q,2p+q}^{\mathrm{GL}(N)} = A_{p,q,2p+q}^{\mathrm{SL}(N)}$  and  $C_{p,q,2p+q}^{\mathrm{GL}(N)} = C_{p,q,2p+q}^{\mathrm{SL}(N)}$ . So we must only show that  $G$  and  $H$  are isomorphism, and we start with surjectivity.

Consider the case of  $G$  first. We establish a commutative diagram

$$\begin{array}{ccc} \mathbb{Q}[\Sigma_{2p+q}] \otimes W^{\otimes q} \otimes U^{\otimes 2p+q} & \xrightarrow{F} & (N^{\otimes 2p+q} \otimes (N^{\vee})^{\otimes 2p+q} \otimes W^{\otimes q} \otimes U^{\otimes 2p+q})^{\mathrm{GL}(N)} \\ \downarrow Q & & \downarrow S \\ S^p(\Lambda^2(U)) \otimes \Lambda^q(W \otimes U) & \xrightarrow{G} & A_{p,q,2p+q}^{\mathrm{GL}(N)}, \end{array} \quad (5.9)$$

and show that  $F$  and  $S$  are surjective. The map  $S$  is the restriction of the  $(\mathrm{GL}(N)$ -equivariant) quotient map

$$N^{\otimes 2p+q} \otimes (N^{\vee})^{\otimes 2p+q} \otimes W^{\otimes q} \otimes U^{\otimes 2p+q} \cong N^{\otimes 2p} \otimes (N \otimes W)^{\otimes q} \otimes (N^{\vee} \otimes U)^{\otimes 2p+q} \rightarrow A_{q,p,2p+q}$$

to the invariant subspace. As the quotient map is surjective and as the category of rational representations of  $\mathrm{GL}(N)$  is semisimple,  $S$  is surjective.

The map  $F$  sends

$$s \otimes w_1 \otimes \dots \otimes w_q \otimes u_1 \otimes \dots \otimes u_{2p+q} \in \Sigma_{2p+q} \otimes W^{\otimes q} \otimes U^{\otimes 2p+q} \subset \mathbb{Q}[\Sigma_{2p+q}] \otimes W^{\otimes q} \otimes U^{\otimes 2p+q}$$

to

$$\sum_{i_1, \dots, i_{2p+q}} a_{i_1} \otimes \dots \otimes a_{i_{2p+q}} \otimes a^{i_{s-1}(1)} \otimes \dots \otimes a^{i_{s-1}(2p+q)} \otimes w_1 \otimes \dots \otimes w_q \otimes u_1 \otimes \dots \otimes u_{2p+q},$$

and is clearly  $\mathrm{GL}(N)$ -invariant. Formula (5.4) shows that (upon identification of its target)  $F$  is the tensor product of the map  $\sigma_{N,2p+q}$  defined in (5.3) and the identity on  $W^{\otimes q} \otimes U^{\otimes 2p+q}$ . Therefore, by Theorem 5.5,  $F$  is surjective.

We define the map  $Q$  on  $\Sigma_{2p+q} \otimes W^{\otimes q} \otimes U^{\otimes 2p+q}$  by

$$s \otimes w_1 \otimes \dots \otimes w_q \otimes u_1 \otimes \dots \otimes u_{2p+q} \mapsto$$

$$\mathrm{sgn}(s)(u_{s(1)} \wedge u_{s(2)} \cdots (u_{s(2p-1)} \wedge u_{s(2p)}) \otimes (w_1 \otimes u_{s(2p+1)}) \wedge \dots \wedge (w_q \otimes u_{s(2p+q)}).$$

It remains to prove that (5.9) commutes, but this follows from

$$SF(s \otimes w_1 \otimes \dots \otimes w_q \otimes u_1 \otimes \dots \otimes u_{2p+q}) =$$

$$\sum_{i_1, \dots, i_{2p+q}} (a_{i_1} \cdot a_{i_2}) \cdots (a_{i_{2p-1}} \cdot a_{i_{2p}}) \otimes (a_{i_{2p+1}} \otimes w_1) \cdots (a_{i_{2p+q}} \otimes w_q) \otimes (a^{i_{s-1}(1)} \otimes u_1) \wedge \dots \wedge (a^{i_{s-1}(2p+q)} \otimes u_{2p+q})$$

and

$$GQ(s \otimes w_1 \otimes \dots \otimes w_q \otimes u_1 \otimes \dots \otimes u_{2p+q}) =$$

$$\operatorname{sgn}(s) \sum_{i_1, \dots, i_{2p+q}} (a_{i_1} \cdot a_{i_2}) \cdots (a_{i_{2p-1}} \cdot a_{i_{2p}}) \otimes (a_{i_{2p+1}} \otimes w_1) \cdots (a_{i_{2p+q}} \otimes w_q) \otimes (a^{i_1} \otimes u_{s(1)}) \wedge \cdots \wedge (a^{i_{2p+q}} \otimes u_{s(2p+q)}).$$

This finishes the proof that  $G$  is surjective. The case of  $H$  is almost identical. In that case, we consider

$$\begin{array}{ccc} \mathbb{Q}[\Sigma_{2p+q}] \otimes W^{\otimes q} \otimes U^{\otimes 2p+q} & \xrightarrow{F} & (N^{\otimes 2p+q} \otimes (N^\vee)^{\otimes 2p+q} \otimes W^{\otimes q} \otimes U^{\otimes 2p+q})^{\operatorname{GL}(N)} \\ \downarrow Q' & & \downarrow S' \\ S^p(\Lambda^2(U)) \otimes \Lambda^q(W \otimes U) & \xrightarrow{G} & C_{p,q,2p+q}^{\operatorname{GL}(N)}. \end{array} \quad (5.10)$$

The map  $F$  is the same map as before. The map  $Q'$  is defined just as  $Q$ , the only difference being that the factor  $\operatorname{sgn}(s)$  in front of the definition of  $Q$  is dropped, and the map  $S'$  is again the quotient map.  $\square$

For the proof of injectivity of  $G$  and  $H$ , we need some classical results of invariant theory. Here is some notation: the conjugate partition to  $\lambda \in \mathcal{P}_n$  is denoted  $\tilde{\lambda} \in \mathcal{P}_n$ . By  $\mathcal{P}_{2p}^{\operatorname{er}}$ , we denote the set of partitions of  $2p$  with even rows, and by  $\mathcal{P}_{2p}^{\operatorname{ec}}$  the set of partitions of  $2p$  with even columns. The first ingredient we shall use are the Cauchy formulas which state that [60, §9.6.3, p. 271]

$$S^q(V \otimes W) = \bigoplus_{\lambda \in \mathcal{P}_q} S_\lambda(V) \otimes S_\lambda(W), \quad (5.11)$$

[60, §9.8.4]

$$\Lambda^r(V \otimes W) = \bigoplus_{\lambda \in \mathcal{P}_r} S_\lambda(V) \otimes S_{\tilde{\lambda}}(W) \quad (5.12)$$

and [60, §11.4.5]

$$S^p(S^2(V)) = \bigoplus_{\lambda \in \mathcal{P}_{2p}^{\operatorname{er}}} S_\lambda(V), \quad (5.13)$$

as well as [60, §11.4.5]

$$S^p(\Lambda^2(V)) = \bigoplus_{\lambda \in \mathcal{P}_{2p}^{\operatorname{ec}}} S_\lambda(V). \quad (5.14)$$

Furthermore, we need the formula [60, §12.5.1]

$$S_\lambda(V) \otimes S_\mu(V) = \bigoplus_{|\kappa|=|\lambda|+|\mu|} c_{\lambda,\mu}^\kappa S_\kappa(V). \quad (5.15)$$

The coefficients  $c_{\lambda,\mu}^\kappa \in \mathbb{N}_0$  are the well-known *Littlewood–Richardson coefficients*. These are the structure constants of the ring  $\Lambda$  of symmetric functions (over the integers) when one takes the Schur functions  $s_\lambda$ ,  $\lambda \in \mathcal{P}$ , as a basis. From this, it follows that the coefficients in (5.15) do not depend on  $\dim(V)$ . The symmetry

$$c_{\lambda,\mu}^\kappa = c_{\mu,\lambda}^\kappa;$$

of the Littlewood–Richardson coefficients is obvious; we also need to know the relation

$$c_{\lambda,\mu}^\kappa = c_{\tilde{\lambda},\tilde{\mu}}^{\tilde{\kappa}}. \quad (5.16)$$

To see (5.16), let  $\omega : \Lambda \rightarrow \Lambda$  be the involutive (ring) automorphism which is constructed in [49, p.21]; formula (3.8) on p.42 of [49] shows that  $\omega(s_\lambda) = s_{\tilde{\lambda}}$ . Since the Littlewood–Richardson coefficients are the structure constants with respect to

the Schur functions, (5.16) follows. An alternative proof of (5.16) can be found in [22, p. 62].

*Proof of Proposition 5.7, injectivity.* Since we already saw that  $G$  and  $H$  are surjective, it suffices to show that the dimensions of the two vector spaces agree (degree-wise, and in the range of degrees we claimed). It is therefore of no danger to write  $S = S'$  for isomorphic representations  $S$  and  $S'$ , and  $nS := S^{\oplus n}$ .

We first turn to the map  $G$ . Its components are maps

$$S^p(\Lambda^2 U) \otimes \Lambda^q(W \otimes U) \rightarrow (S^p(S^2(N)) \otimes S^q(N \otimes W) \otimes \Lambda^{2p+q}(N^\vee \otimes U))^{\mathrm{GL}(N)}.$$

As a  $\mathrm{GL}(N) \times \mathrm{GL}(W) \times \mathrm{GL}(U)$ -representation, we have by (5.13), (5.11) and (5.12)

$$\begin{aligned} & S^p(S^2(N)) \otimes S^q(N \otimes W) \otimes \Lambda^{2p+q}(N^\vee \otimes U) = \\ & \bigoplus_{\lambda \in \mathcal{P}_{2p}^{\mathrm{sr}}, |\mu|=q, |\nu|=2p+q} S_\lambda(N) \otimes S_\mu(N) \otimes S_\mu(W) \otimes S_\nu(N^\vee) \otimes S_{\bar{\nu}}(U). \end{aligned}$$

By (5.15), the latter is isomorphic to

$$\bigoplus_{\lambda \in \mathcal{P}_{2p}^{\mathrm{sr}}, |\mu|=q, |\nu|=|\kappa|=2p+q} c_{\lambda, \mu}^\kappa S_\kappa(N) \otimes S_\mu(W) \otimes S_\nu(N^\vee) \otimes S_{\bar{\nu}}(U).$$

Since the  $S_\lambda(N)$  are mutually nonisomorphic irreducible  $\mathrm{GL}(N)$ -representations or trivial, we have, with  $g := \dim(N)$ ,

$$(S_\kappa(N) \otimes S_\nu(N^\vee))^{\mathrm{GL}(N)} \cong \begin{cases} \mathbb{Q} & \kappa = \nu \text{ and } \mathrm{ht}(\nu) = \mathrm{ht}(\kappa) \leq g \\ 0 & \kappa \neq \nu \text{ or } \mathrm{ht}(\nu) > g \text{ or } \mathrm{ht}(\kappa) > g. \end{cases} \quad (5.17)$$

Therefore

$$(S_\kappa(N) \otimes S_\mu(W) \otimes S_\nu(N^\vee) \otimes S_{\bar{\nu}}(U))^{\mathrm{GL}(N)} = \begin{cases} S_\mu(W) \otimes S_{\bar{\nu}}(U) & \kappa = \nu \text{ and } \mathrm{ht}(\nu) = \mathrm{ht}(\kappa) \leq g \\ 0 & \kappa \neq \nu \text{ or } \mathrm{ht}(\nu) > g \text{ or } \mathrm{ht}(\kappa) > g, \end{cases}$$

and so

$$\begin{aligned} & (S^p(S^2(N)) \otimes S^q(N \otimes W) \otimes \Lambda^{2p+q}(N^\vee \otimes U))^{\mathrm{GL}(N)} \\ &= \bigoplus_{\lambda \in \mathcal{P}_{2p}^{\mathrm{sr}}, |\mu|=q, |\nu|=2p+q, \mathrm{ht}(\nu) \leq g} c_{\lambda, \mu}^\nu S_\mu(W) \otimes S_{\bar{\nu}}(U) \\ &= \bigoplus_{\lambda \in \mathcal{P}_{2p}^{\mathrm{sr}}, |\mu|=q, |\nu|=2p+q, \mathrm{ht}(\nu) \leq g} c_{\lambda, \bar{\mu}}^{\bar{\nu}} S_\mu(W) \otimes S_{\bar{\nu}}(U) \quad (\text{by (5.16)}). \end{aligned}$$

Under the hypothesis that  $g \geq 2p + q$ ,  $\mathrm{ht}(\nu) \leq g$  holds for all  $\nu \in \mathcal{P}_{2p+q}$ . Using (5.15) again, the latter is isomorphic to

$$\bigoplus_{\lambda \in \mathcal{P}_{2p}^{\mathrm{sr}}, |\mu|=q} S_\mu(W) \otimes S_{\bar{\lambda}}(U) \otimes S_{\bar{\mu}}(U),$$

and by (5.12), this agrees with

$$\begin{aligned} & \bigoplus_{\lambda \in \mathcal{P}_{2p}^{\mathrm{sr}}} \Lambda^q(W \otimes U) \otimes S_{\bar{\lambda}}(U) = \Lambda^q(W \otimes U) \otimes \bigoplus_{\lambda \in \mathcal{P}_{2p}^{\mathrm{sc}}} S_\lambda(U) = \\ &= \Lambda^q(W \otimes U) \otimes S^p(\Lambda^2(U)). \end{aligned}$$

The proof for  $H$  is almost identical. The components of  $H$  are maps

$$S^p(S^2 U) \otimes \Lambda^q(W \otimes U) \rightarrow (S^p(\Lambda^2(N)) \otimes \Lambda^q(N \otimes W) \otimes S^{2p+q}(N^\vee \otimes U))^{\mathrm{GL}(N)}.$$

We compute, by (5.14), (5.11) and (5.12) and (5.15),

$$\begin{aligned}
& S^p(\Lambda^2(N)) \otimes \Lambda^q(N \otimes W) \otimes S^{2p+q}(N^\vee \otimes U) = \\
&= \bigoplus_{\lambda \in \mathcal{P}_{2p}^{\text{sc}}, |\mu|=q, |\nu|=2p+q} S_\lambda(N) \otimes S_\mu(N) \otimes S_{\bar{\mu}}(W) \otimes S_\nu(N^\vee) \otimes S_\nu(U) = \\
&= \bigoplus_{\lambda \in \mathcal{P}_{2p}^{\text{sc}}, |\mu|=q, |\nu|=|\kappa|=2p+q} c_{\lambda, \mu}^\kappa S_\kappa(N) \otimes S_{\bar{\mu}}(W) \otimes S_\nu(N^\vee) \otimes S_\nu(U).
\end{aligned}$$

Taking  $\text{GL}(N)$ -invariants and using (5.17) yields

$$\begin{aligned}
& (S^p(\Lambda^2(N)) \otimes \Lambda^q(N \otimes W) \otimes S^{2p+q}(N^\vee \otimes U))^{\text{GL}(N)} = \\
&= \bigoplus_{\lambda \in \mathcal{P}_{2p}^{\text{sc}}, |\mu|=q, |\nu|=2p+q, \text{ht}(\nu) \leq g} c_{\lambda, \mu}^\nu S_{\bar{\mu}}(W) \otimes S_\nu(U).
\end{aligned}$$

If  $g \geq 2p + q$ , (5.15) shows that this is equal to

$$\begin{aligned}
& \bigoplus_{\lambda \in \mathcal{P}_{2p}^{\text{sc}}, |\mu|=q} S_{\bar{\mu}}(W) \otimes S_\lambda(U) \otimes S_\mu(U) = \\
& \left( \bigoplus_{\lambda \in \mathcal{P}_{2p}^{\text{sc}}} S_\lambda(U) \right) \otimes \left( \bigoplus_{|\mu|=q} S_{\bar{\mu}}(W) \otimes S_\mu(U) \right) \stackrel{(5.14), (5.12)}{=} \\
& S^p(\Lambda^2(U)) \otimes \Lambda^q(W \otimes U)
\end{aligned}$$

as claimed.  $\square$

Finally, we give the short proof of Proposition 5.8. This is an immediate consequence of a more general result.

**Lemma 5.18.** *Let  $N_{\mathbb{Z}}$  be a finitely generated free abelian group, write  $N := N_{\mathbb{Z}} \otimes \mathbb{Q}$  and let  $\rho : \text{SL}(N) \rightarrow \text{GL}(W)$  be a rational representation. Then*

$$W^{\text{SL}(N)} = W^{\text{SL}(N_{\mathbb{Z}})}.$$

*Proof.* Assume  $N = \mathbb{Z}^g$ . A special case of Borel's density theorem [4] states that  $\text{SL}_g(\mathbb{Z}) \subset \text{SL}_g(\mathbb{Q})$  is Zariski dense (a very short and elementary proof for the special linear group has been written down by Putman [61]). For  $v \in V^{\text{SL}_g(\mathbb{Z})}$  and  $\ell \in V^*$ , the function  $\text{SL}_g(\mathbb{Q}) \rightarrow \mathbb{Q}$ ,  $A \mapsto \ell(v - Av)$ , is polynomial and vanishes on  $\text{SL}_g(\mathbb{Z})$ , hence on  $\text{SL}_g(\mathbb{Q})$ , whence  $v = Av$  for all  $A \in \text{SL}_g(\mathbb{Q})$ .  $\square$

## 6. THE COHOMOLOGY OF THE BLOCK DIFFEOMORPHISM SPACE

**6.1. Using invariant theory.** In this section, we finish our partial evaluation of the spectral sequence of the fibration

$$\text{map}_{\partial}((U_{g,1}^n)_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 \rightarrow \text{map}_{\partial}((U_{g,1}^n)_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \text{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})^{\text{id}} \rightarrow \text{BhAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})^{\text{id}}. \quad (6.1)$$

Before we state the result, let us fix some bounds that the various parameters have to fulfil.

*Assumption 6.2.* (1) We assume throughout that  $n \geq 5$ .

(2) We pick  $M$  in (4.27) large enough so that

$$4M \geq 3n - 5.$$

(3) We furthermore choose  $g$  large enough to satisfy

$$g > n - 3,$$

which implies also that  $g \geq 3$ .

Using the number  $M$ , we define the graded vector spaces  $U(n)$ ,  $V(n)$  and  $W(n)$  as in (4.27). We let  $v_m \in V(n)_{4m-2n-1}$  and  $w_m \in W(n)_{4m-n}$  be the obvious generators, and let  $u_m := S(w_m) \in U(n)_{4m-n-1}$  be the image under the map  $S$  from Proposition 4.32. To formulate the result we are aiming at, some more notation is necessary.

**Definition 6.3.** We let  $K(n)$  be the following graded vector space. It has basis elements  $k_m \in K(n)_{4m-2n-1}$  for  $m \leq M$  and  $4m - 2n - 1 > 0$ , and it has basis elements  $k_{m_0, m_1} \in K(n)_{4(m_0+m_1)-2n-1}$  for  $m_0 \leq m_1 \leq M$ ,  $4m_0 \geq n + 1$ ,  $4(m_0 + m_1) - 2n - 1 > 0$ .

Note that all generators in  $K(n)$  are in odd degrees. We define a (degree-preserving) map

$$\xi : \Lambda^*(K(n)) \rightarrow H^*(\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0 // \text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})_{\mathbb{Z}}) \quad (6.4)$$

by

$$\xi(k_m) := \kappa_{L_m}, \quad \xi(k_{m_0, m_1}) := \kappa_{L_{m_0} L_{m_1}}.$$

The composition of  $\xi$  with the pullback map

$$H^*(\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0 // \text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})_{\mathbb{Z}}; \mathbb{Q}) \rightarrow H^*(\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0 // \text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\text{id}}; \mathbb{Q})$$

goes into the  $\text{SL}(N(g)_{\mathbb{Z}})$ -invariant part. Here is the goal of this subsection.

**Proposition 6.5.** *The map*

$$\xi : \Lambda^*(K(n)) \rightarrow H^*(\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0 // \text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\text{id}}; \mathbb{Q})^{\text{SL}(N(g)_{\mathbb{Z}})}$$

is an isomorphism in degrees  $* \leq (n - 4)$ , provided that  $M$  and  $g$  satisfy the bounds from (6.2).

Recall from Observation 4.25 that the spectral sequence  $E_*^{*,*}$  of (6.1) is a spectral sequence of  $\text{GL}(N(g)) \cong \pi_0(\text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q}))$ -modules. Let us elaborate this a little.

**Proposition 6.6.** *Let  $n \geq 5$  and  $g \geq 3$ .*

- (1) *For  $q \leq (n - 3)$ ,  $H^q(\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0 // \text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\text{id}}; \mathbb{Q})$  is a rational representation of  $\text{SL}(N(g)_{\mathbb{Z}})$ , of load  $\leq q$ .*
- (2) *Let  $E_*^{*,*}$  denote the spectral sequence of (6.1), and define*

$$\overline{E}_r^{p,q} := (E_r^{p,q})^{\text{SL}(N(g)_{\mathbb{Z}})}.$$

*Then  $\overline{E}_*^{*,*}$  is a spectral sequence, and it converges to  $H^*(\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0 // \text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\text{id}}; \mathbb{Q})^{\text{SL}(N(g)_{\mathbb{Z}})}$ .*

*Proof.* (1) We have to invoke a deep result by Bass–Milnor–Serre [1, Corollary 16.6]. The quoted result implies that a homomorphism  $\text{SL}(N(g)) \rightarrow \text{GL}(V)$ , where  $V$  is a finite-dimensional  $\mathbb{Q}$ -vector space and  $g \geq 3$ , is actually rational (see also the discussion on p. 63 f and p. 134 of loc.cit.). Hence  $H^q(\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0 // \text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\text{id}}; \mathbb{Q})$  is a rational  $\text{SL}(N(g))$ -representation.

From the description of the  $E_2$ -term given in Proposition 4.31, it is apparent that (in total degrees  $* \leq (n - 3)$ ) the  $\text{GL}(N(g))$ -representations in the  $E_2$ -term are all rational. Since subquotients of rational representations are again rational

(and the load does not increase), the representations occuring in  $E_\infty^{*,*}$  with total degree  $q$  are all rational, of load  $\leq q$ .

(2): Theorem 5.1 and Lemma 5.18 together imply that taking  $\mathrm{SL}(N(g)_\mathbb{Z})$ -invariants is an exact functor from rational  $\mathrm{GL}(N(g))$ -representations to  $\mathbb{Q}$ -vector spaces. Hence  $\overline{E}_*^{*,*}$  is also a spectral sequence. The claim about convergence follows from Theorem 6.17.  $\square$

Hence in order to prove Proposition 6.5, we can focus our attention completely on  $\overline{E}_*^{*,*}$ .

**Definition 6.7.** Let  $D^{*,*}$  be the bigraded algebra

$$D^{*,*} := \Lambda^*(V(n) \oplus W(n) \otimes U(n)) \otimes S^*(\mathbb{Q}[2, 0] \otimes \Lambda^2(U(n)))$$

where the graded vector spaces  $W(n), U(n)$  and  $V(n)$  are as in (4.27), but sit in bidegrees  $(0, *)$ . Let  $\delta : D^{*,*} \rightarrow D^{*,*}$  be the differential of degree  $(2, -1)$  which is a derivation, and is given on the generators as follows:

- (1)  $\delta|_{V(n)} = 0$ ,
- (2)  $\delta|_{W(n) \otimes U(n)}$  is the map  $W(n) \otimes U(n) \xrightarrow{S \otimes 1} U(n) \otimes U(n) \rightarrow \Lambda^2(U(n)) \cong \mathbb{Q}[2, 0] \otimes \Lambda^2(U(n))$ ,
- (3)  $\delta|_{\mathbb{Q}[2, 0] \otimes \Lambda^2(U(n))} = 0$ .

We define a graded algebra homomorphism

$$\eta : D^{*,*} \rightarrow \overline{E}_2^{*,*} = (E_2^{*,*})^{\mathrm{SL}(N_\mathbb{Z}(g))} \quad (6.8)$$

to the invariant part of the spectral sequence of (6.1), by sending

$$\begin{aligned} v_m &\mapsto \lambda_{[U_{g,1}^n], L_m} \in E_2^{0, 4m-2n-1} \\ w_{m_0} \otimes u_{m_1} &\mapsto \sum_i \lambda_{a_i, L_{m_0}} \lambda_{b_i, L_{m_1}} \in E_2^{0, 4m_0+4m_1-2n-1} \\ u_{m_0} \wedge u_{m_1} &\mapsto \sum_{ij} x_{ij} \otimes \lambda_{b_i, L_{m_0}} \lambda_{b_j, L_{m_1}} \in E_2^{2, 4m_0+4m_1-n-2}. \end{aligned}$$

Here  $x_{ij} \in E_2^{2,0}$  are the generators in  $E_2^{2,0} \cong L^2(N(g))$ .

**Proposition 6.9.** *The map  $\eta$  is a map of differential bigraded algebras, and it is an isomorphism in total degrees  $\leq (n-3)$ , provided that the bounds of 6.2 are satisfied.*

*Proof.* That  $\eta$  is an isomorphism in the indicated range of degrees follows from Propositions 4.31, 5.7 and 5.8. The first two conditions of 6.2 are needed for 4.31, and the third for 5.7.

Proposition 4.32 and the definition of the maps in Proposition 5.7 show that  $\eta$  is compatible with the differential.  $\square$

Let us next compute the cohomology of the differential graded algebras  $D^{*,*}$ . This is in terms of the following well-known construction.

**Definition 6.10.** Let  $F : Y \rightarrow X$  be a linear map of finite-dimensional  $\mathbb{Q}$ -vector spaces. The *Koszul complex* of the map  $F$  is the graded commutative differential graded algebra

$$D_F := \Lambda^*(Y) \otimes S^*(X),$$

where  $S^1(X)$  has degree 2,  $\Lambda^1(Y)$  has degree 1, and the differential  $d_F : D_F \rightarrow D_F$  is the derivation of degree +1 given by the condition that  $d_F|_{S^1(X)} = 0$  and that  $d_F : \Lambda^1(Y) \rightarrow S^1(X)$  is the map  $F$ .

Up to a different grading,  $D^{*,*}$  is  $D_F$ , where  $F$  is the map

$$V(n) \oplus W(n) \otimes U(n) \xrightarrow{0 \oplus S \otimes 1_{U(n)}} U(n)^{\otimes 2} \rightarrow \Lambda^2(U(n)). \quad (6.11)$$

In order to compute  $H^*(D^{*,*})$  and hence  $(E_3^{*,*})^{\text{SL}(N_{\mathbb{Z}}(g))}$  in a range of degrees, we compute  $H^*(D_F)$  in general. There are obvious linear maps

$$\ker(F) \cong \ker(d_F : \Lambda^1(Y) \rightarrow S^1(X)) \rightarrow H^1(D_F)$$

and

$$\text{coker}(F) \cong \text{coker}(d_F : \Lambda^1(Y) \rightarrow S^1(X)) \rightarrow H^2(D_F).$$

They give a map of graded commutative algebras

$$\eta_F : \Lambda^*(\ker(F)) \otimes S^*(\text{coker}(F)) \rightarrow H^*(D_F).$$

**Lemma 6.12.** *The map  $\eta_F$  is an isomorphism.*

*Proof.* It follows from the Künneth formula that if  $\eta_{F_0}$  and  $\eta_{F_1}$  are isomorphisms, then so is  $\eta_{F_0 \oplus F_1}$ .

By elementary linear algebra, we can write  $F$  as a direct sum of an isomorphism and a zero map, and so it suffices to treat these two cases separately. If  $F$  is a zero map, the claim is obvious. If  $F$  is an isomorphism, we can assume without loss of generality that  $F$  is an identity map. In that case,  $D_{\text{id}_V}$  is the Koszul complex of the vector space  $V$  which is well-known to be acyclic (since we are over a field of characteristic 0), see e.g. [31, §3.1].  $\square$

*Proof of Proposition 6.5.* The map in (6.11) is surjective. Lemma 6.12 and Proposition 6.9 prove that

$$\overline{E}_3^{p,q} = 0$$

if  $p + q \leq n - 3$  and  $p \neq 0$ . Hence, using Proposition 6.6, the natural map

$$A : H^q(\text{map}_{\partial}((U_{g,1}^n)_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \text{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})^{\text{id}}; \mathbb{Q})^{\text{SL}(N(g))_{\mathbb{Z}}} \rightarrow \overline{E}_{\infty}^{0,q} \subset \overline{E}_3^{0,q}$$

is an isomorphism if  $q \leq n - 4$ . It is therefore sufficient to show that  $A \circ \xi : \Lambda^*(K(n)) \rightarrow \overline{E}_3^{0,*}$  is an isomorphism in the indicated range of degrees. By the definition of  $\xi$  and by Lemma 3.24,  $A \circ \xi$  is given by

$$k_m \mapsto \lambda_{[U_{g,1}^n], L_m}$$

and

$$k_{m_0, m_1} \mapsto \lambda_{[U_{g,1}^n], L_{m_0} L_{m_1}}.$$

We must therefore check that  $\overline{E}_3^{0,*}$  is the free graded-commutative algebra on the listed generators. Lemma 6.12 tells us how to do that. We distinguish to cases.

First let  $n \not\equiv 3 \pmod{4}$ . Then the map  $S : W(n) \rightarrow U(n)$  is an isomorphism. We deduce that the following set is a basis for the kernel of (6.11):

$$\{v_m, (w_{m_0} \otimes u_{m_1} + w_{m_1} \otimes u_{m_0}) \mid 4m - 2n - 1 > 0, m \leq M, m_0 \leq m_1 \leq M, 4m_0 - n - 1 > 0\}. \quad (6.13)$$

Under the map  $\eta$  from (6.8), the element  $v_m$  is mapped to  $\lambda_{[U_{g,1}^n], L_m}$ . The element  $w_{m_0} \otimes u_{m_1} + w_{m_1} \otimes u_{m_0}$  is mapped to

$$\begin{aligned} & \sum_i \lambda_{a_i, L_{m_0}} \lambda_{b_i, L_{m_1}} + \lambda_{a_i, L_{m_1}} \lambda_{b_i, L_{m_0}} \\ &= \sum_i \lambda_{a_i, L_{m_0}} \lambda_{b_i, L_{m_1}} + \lambda_{b_i, L_{m_0}} \lambda_{a_i, L_{m_1}} \text{ (for degree reasons)} \\ &= \lambda_{[U_{g,1}^n], L_{m_0} L_{m_1}} \text{ (by (4.4)).} \end{aligned}$$

This completes the proof if  $n \not\equiv 3 \pmod{4}$ .

If  $n \equiv 3 \pmod{4}$ , the kernel of (6.11) is larger: one obtains a basis by adding the elements

$$w_{\frac{n+1}{4}} \otimes u_m, \quad 4m - n - 1 > 0, m \leq M$$

to the list of elements in (6.13). Using (6.8) and (4.4) again, these elements go to  $\lambda_{[U_{g,1}^n], L_{\frac{n+1}{4}} L_m}$ . So the proof is complete.  $\square$

**6.2. Using Borel's vanishing theorem.** We now look at the spectral sequence of the fibration

$$\mathrm{map}_{\partial}((U_{g,1}^n)_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})^{\mathrm{id}} \rightarrow \mathrm{map}_{\partial}((U_{g,1}^n)_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}} \rightarrow BGL(N(g)_{\mathbb{Z}}). \quad (6.14)$$

Recall the graded vector space  $K(n)$  from Definition 6.3 and from (6.4) the map

$$\xi : \Lambda^*(K(n)) \rightarrow H^*(\mathrm{map}_{\partial}((U_{g,1}^n)_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}}; \mathbb{Q}).$$

Let moreover  $B$  be the graded vector space  $B := \bigoplus_{k \geq 1} \mathbb{Q}[4k + 1]$ . Mapping the generators to the Borel classes gives furthermore a map

$$\beta : \Lambda^*(B) \rightarrow H^*(\mathrm{map}_{\partial}((U_{g,1}^n)_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}}; \mathbb{Q}).$$

**Proposition 6.15.** *Assume the bounds from (6.2) (which in particular means  $n \geq 5$ ), but strengthened by*

$$g \geq 2n - 4.$$

*Then the map*

$$\xi \otimes \beta : \Lambda^*(K(n) \oplus B) \rightarrow H^*(\mathrm{map}_{\partial}((U_{g,1}^n)_{\mathbb{Q}}; BO_{\mathbb{Q}})^0 // \mathrm{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}}; \mathbb{Q})$$

*is an isomorphism in degrees  $* \leq (n - 4)$ .*

When combined with Proposition 3.25, we obtain the following result as a corollary. This establishes Theorem 1.8 from the introduction.

**Corollary 6.16.** *Assume the bounds of Proposition 6.15. Then in degrees  $\leq (n - 4)$ ,  $H^*(B\widetilde{\mathrm{Diff}}_{\partial}(U_{g,1}^n); \mathbb{Q})$  is the exterior algebra on the Borel classes and on the tautological classes  $\kappa_{L_{m_0} L_{m_1}}$  with  $m_0 \leq m_1$ ,  $4m_0 - n > 0$  and  $4(m_0 + m_1) - 2n - 1 > 0$ .  $\square$*

The proof of Proposition 6.15 relies on Borel's vanishing theorem [6] that we shall state first.

**Theorem 6.17 (Borel).** *Let  $V$  be a rational representation of  $\mathrm{SL}_g(\mathbb{Q})$  of load at most  $n$ . Then the map*

$$H^p(\mathrm{SL}_g(\mathbb{Z}); \mathbb{Q}) \otimes V^{\mathrm{SL}_g(\mathbb{Z})} = H^p(\mathrm{SL}_g(\mathbb{Z}); \mathbb{Q}) \otimes V^{\mathrm{SL}_g(\mathbb{Q})} \rightarrow H^p(\mathrm{SL}_g(\mathbb{Z}); V^{\mathrm{SL}_g(\mathbb{Q})}) \rightarrow H^p(\mathrm{SL}_g(\mathbb{Z}); V)$$

*is an isomorphism, provided that  $2p + 2 \leq g - n$  (the first equation holds by Lemma 5.18).*

*References.* This is essentially due to Borel [6, Theorem 4.4], but the ranges are not made explicit in Borel's work, so some more words need to be said here. By complete reducibility, we can assume that  $V \subset T^{k,l}(\mathbb{Q}^g)$  with  $k+l \leq n$ , and finally suppose that  $V = T^{k,l}(\mathbb{Q}^g)$ . Moreover, it is enough to prove that statement for  $\mathbb{Q}$  replaced by  $\mathbb{R}$  and  $\mathbb{Q}^g$  replaced by  $\mathbb{R}^g$ .

Borel proved in loc.cit. that

$$H^p(\mathrm{SL}_g(\mathbb{Z}); \mathbb{R}) \otimes T^{k,l}(\mathbb{R}^g)^{\mathrm{SL}_g(\mathbb{R})} \rightarrow H^p(\mathrm{SL}_g(\mathbb{Z}); T^{k,l}(\mathbb{R}^g)) \quad (6.18)$$

is an isomorphism provided that  $p \leq \min(M(\mathrm{SL}_g(\mathbb{R}), (k, l)), C(\mathrm{SL}_g(\mathbb{R}), (k, l)))$ , where  $M(\mathrm{SL}_g(\mathbb{R}), (k, l))$  and  $C(\mathrm{SL}_g(\mathbb{R}), (k, l))$  are constants which can be read off from the root system of  $\mathfrak{sl}_g$  and the weights of  $T^{k,l}(\mathbb{Q}^g)$ . In loc.cit., Borel showed that  $M(\mathrm{SL}_g(\mathbb{R}), (k, l)) \geq g - 2$ , but left  $C(\mathrm{SL}_g(\mathbb{R}), (k, l))$  implicit. A relatively naive counting argument given in the proof of [43, Theorem 7.3] shows that  $C(\mathrm{SL}_g(\mathbb{R}), (k, l)) \geq \frac{1}{8}g^2 - \max(k, l) - 1$ , which implies that (6.18) is an isomorphism when  $p \leq f_{k,l}(g)$ , and  $f_{k,l}$  is a function with  $\lim_{g \rightarrow \infty} f_{k,l}(g) = \infty$ .

To get at the range claimed by us, we use Van der Kallen's work on homological stability, more precisely [74, Theorem 5.6]. The latter result implies that for  $2q+2 \leq g - (k+l)$  and all  $h \geq g$ , the map  $H_q(\mathrm{SL}_g(\mathbb{Z}); T^{k,l}(\mathbb{R}^g)) \rightarrow H_q(\mathrm{SL}_h(\mathbb{Z}); T^{k,l}(\mathbb{R}^h))$  is an isomorphism. This implies a cohomological statement by an instance of the universal coefficient theorem [65, Lemma 3.5], and we can pick  $h$  large enough so that  $f(h) \geq q$ .  $\square$

*Remark 6.19.* Tshishiku [73] showed by a careful analysis of root systems that the corresponding result is true for  $\mathrm{SO}_{g,g}(\mathbb{Z})$  and  $\mathrm{Sp}_{2g}(\mathbb{Z})$ , but in a range that only depends on  $g$ , not on the representation  $V$ . We have been informed by him that the analogous procedure for  $\mathrm{SL}_g(\mathbb{Z})$  does *not* lead to a range independent of  $V$ .

*Proof of Proposition 6.15.* Since Theorem 6.17 is about the special linear group rather than the general linear group, we modify the sequence a bit and look at

$$\mathrm{map}_\partial((U_{g,1}^n)_{\mathbb{Q}}; \mathrm{BO}_{\mathbb{Q}})^0 // \mathrm{hAut}_\partial((U_{g,1}^n)_{\mathbb{Q}})^{\mathrm{id}} \rightarrow \mathrm{map}_\partial((U_{g,1}^n)_{\mathbb{Q}}; \mathrm{BO}_{\mathbb{Q}})^0 // \mathrm{ShAut}_\partial((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}} \rightarrow \mathrm{BSL}(N(g)_{\mathbb{Z}}), \quad (6.20)$$

where  $\mathrm{ShAut}_\partial((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}} \subset \mathrm{hAut}_\partial((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}}$  consists of those homotopy automorphisms whose action on  $H_n(U_{g,1}^n)$  is by maps of determinant 1. The natural map

$$\mathrm{map}_\partial((U_{g,1}^n)_{\mathbb{Q}}; \mathrm{BO}_{\mathbb{Q}})^0 // \mathrm{ShAut}_\partial((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}} \rightarrow \mathrm{map}_\partial((U_{g,1}^n)_{\mathbb{Q}}; \mathrm{BO}_{\mathbb{Q}})^0 // \mathrm{hAut}_\partial((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}}$$

is, up to homotopy, a 2-fold covering. Hence it induces an injective map in rational cohomology by a general argument [35, Proposition 3G.1]. We may therefore show the Proposition with  $\mathrm{map}_\partial((U_{g,1}^n)_{\mathbb{Q}}; \mathrm{BO}_{\mathbb{Q}})^0 // \mathrm{hAut}_\partial((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}}$  replaced by  $\mathrm{map}_\partial((U_{g,1}^n)_{\mathbb{Q}}; \mathrm{BO}_{\mathbb{Q}})^0 // \mathrm{ShAut}_\partial((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}}$ .

Let  $E_*^{*,*}$  be the spectral sequence for (6.20). Its  $E_2$ -term is

$$E_2^{p,q} = H^p(\mathrm{SL}(N(g)_{\mathbb{Z}}); H^q(\mathrm{map}_\partial((U_{g,1}^n)_{\mathbb{Q}}; \mathrm{BO}_{\mathbb{Q}})^0 // \mathrm{hAut}_\partial((U_{g,1}^n)_{\mathbb{Q}})^{\mathrm{id}}; \mathbb{Q})).$$

By Proposition 6.6, the coefficient module is a rational representation of load  $\leq q$ .

Hence if  $2p + q + 2 \leq g$ , we may invoke Theorem 6.17 and see that

$$E_2^{p,q} = H^p(\mathrm{SL}(N(g)_{\mathbb{Z}}); \mathbb{Q}) \otimes H^q(\mathrm{map}_\partial((U_{g,1}^n)_{\mathbb{Q}}; \mathrm{BO}_{\mathbb{Q}})^0 // \mathrm{hAut}_\partial((U_{g,1}^n)_{\mathbb{Q}})^{\mathrm{id}}; \mathbb{Q})^{\mathrm{SL}(N(g)_{\mathbb{Z}})}. \quad (6.21)$$

Under our bound on  $g$ , this holds for all  $p, q$  with  $p + q \leq n - 4$ .

It follows from Proposition 6.5 and (6.4) that the map

$$H^*(\mathrm{map}_\partial((U_{g,1}^n)_{\mathbb{Q}}; \mathrm{BO}_{\mathbb{Q}})^0 // \mathrm{ShAut}_\partial((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}}; \mathbb{Q}) \rightarrow H^*(\mathrm{map}_\partial((U_{g,1}^n)_{\mathbb{Q}}; \mathrm{BO}_{\mathbb{Q}})^0 // \mathrm{hAut}_\partial((U_{g,1}^n)_{\mathbb{Q}})^{\mathrm{id}}; \mathbb{Q})^{\mathrm{SL}(N(g)_{\mathbb{Z}})}$$

is surjective if  $* \leq (n-4)$ , but that can be identified with the edge homomorphism of the spectral sequence, so that all differentials starting in the zeroth column vanish (in degrees  $* \leq n-4$ ). The isomorphism (6.21) implies that the map  $E_2^{*,0} \otimes E_2^{0,*} \rightarrow E_2^{*,*}$  of bigraded vector spaces is an isomorphism in total degrees  $\leq n-4$ . As the spectral sequence is a spectral sequence of algebras, we conclude that all differentials starting in a term  $E_r^{p,q}$  with  $p+q \leq (n-4)$  are trivial.

An application of Theorem 2.13 and the Leray–Hirsch theorem shows that the map  $\xi \otimes \beta$  is an isomorphism in degrees  $\leq (n-4)$ .  $\square$

## 7. THE ENDGAME: FROM BLOCK DIFFEOMORPHISMS TO ACTUAL DIFFEOMORPHISMS

In this rather short section, we compare the rational cohomologies of  $B\text{Diff}_\partial(U_{g,1}^n)$  and  $\widetilde{B\text{Diff}}_\partial(U_{g,1}^n)$  in a range of degrees and thereby complete the proof of Theorem A. We phrase the argument in a way that is largely independent of the calculations for  $H^*(\widetilde{B\text{Diff}}_\partial(U_{g,1}^n); \mathbb{Q})$  which we carried out in the previous sections. To formulate the result, let us introduce the following notations:

$$\mathcal{D}_g := \text{Diff}_\partial(U_{g,1}^n); \quad \widetilde{\mathcal{D}}_g := \widetilde{\text{Diff}}_\partial(U_{g,1}^n).$$

We let  $B\mathcal{D}_\infty := \text{hocolim}_{g \rightarrow \infty} B\mathcal{D}_g$  and define  $B\widetilde{\mathcal{D}}_\infty$  analogously. The aim of this section is the following result, which together with Theorem 1.8 and [58, Corollary 1.3.2] immediately implies Theorem A.

### Theorem 7.1.

- (1) *Let  $n \geq 3$ . Then  $H^*(B\widetilde{\mathcal{D}}_\infty; \mathbb{Q})$  is a free-graded commutative algebra which is degreewise finitely generated.*
- (2) *Let  $n \geq 4$ . Then  $H^*(B\mathcal{D}_\infty; \mathbb{Q})$  is a free-graded commutative algebra which is degreewise finitely generated.*
- (3) *Let  $n \geq 3$ . Then the map*

$$B\rho_n : B\widetilde{\mathcal{D}}_\infty \rightarrow B\text{GL}_\infty(\mathbb{Z})$$

*given by the action on  $H_n(-; \mathbb{Z})$  induces an injective map in rational cohomology, and  $H^*(B\widetilde{\mathcal{D}}_\infty)$  is isomorphic to the tensor product of  $H^*(B\text{GL}_\infty(\mathbb{Z}); \mathbb{Q})$  and a free graded-commutative algebra.*

- (4) *Let  $n \geq 5$ . Then the comparison map*

$$B\mathcal{I}_\infty : B\mathcal{D}_\infty \rightarrow B\widetilde{\mathcal{D}}_\infty$$

*induces a surjection in rational cohomology in degrees  $* \leq 2n-5$ , and in this range of degrees, the kernel of  $(B\mathcal{I}_\infty)^*$  is the ideal generated by the image of  $(B\rho_n)^*$ .*

**7.1. Getting the mapping class group under control.** For the proof of Theorem 7.1, we use that the spaces

$$B\mathcal{D} := \prod_{g \geq 0} B\mathcal{D}_g \quad \text{and} \quad B\widetilde{\mathcal{D}} := \prod_{g \geq 0} B\widetilde{\mathcal{D}}_g$$

carry  $E_{2n+1}$ -structures, analogous to the  $E_2$ -structure described in [24, §3] for diffeomorphisms of surfaces; we refrain from giving any more details here. As a consequence of May’s recognition principle [51], the group completions  $\Omega B\mathcal{D}$  and  $\Omega B\widetilde{\mathcal{D}}$  have the structures of  $(2n+1)$ -fold loop spaces. The comparison map

$BD \rightarrow B\tilde{D}$  is a map of  $E_{2n+1}$ -spaces, and hence the induced map on group completions is a map of  $(2n + 1)$ -fold loop spaces.

**Lemma 7.2.** *The natural maps*

$$\mathbb{Z} \times BD_\infty \rightarrow \Omega BB\mathcal{D} \tag{7.3}$$

and

$$\mathbb{Z} \times B\tilde{D}_\infty \rightarrow \Omega BB\tilde{D} \tag{7.4}$$

are acyclic (i.e. their homotopy fibres have the integral homology of a point). The commutator subgroups of  $\pi_1(B\mathcal{D}_\infty)$  and  $\pi_1(B\tilde{D}_\infty)$  are perfect. The maps (7.3) and (7.4) identify their targets with the Quillen plus construction on their source (here the Quillen plus construction is performed one component at a time, on the maximal perfect normal subgroup of the fundamental group).

*Proof.* A straightforward application of the group completion theorem [53] shows that the two maps are integral homology equivalences. An improved version of the group completion theorem, namely [64, Theorem 1.1], proves the (stronger) claim of acyclicity. Perfectness of the commutator subgroups follows from [64, Proposition 3.1], and the statement about the Quillen plus construction is a consequence: by [36, Proposition 3.1], acyclic maps out of a given space are classified up to homotopy equivalence by the kernels of their induced maps on fundamental groups.  $\square$

Our goal in this subsection is the following statement.

**Proposition 7.5.** *The diagram*

$$\begin{array}{ccc} BD_0 & \longrightarrow & BD_\infty^+ \\ \downarrow & & \downarrow \\ B\tilde{D}_0 & \longrightarrow & B\tilde{D}_\infty^+ \end{array}$$

is  $(2n - 5)$ -cartesian, i.e. the induced map on vertical homotopy fibres is  $(2n - 5)$ -connected.

The proof of Proposition 7.5 is a sequence of lemmas.

**Lemma 7.6.** *The diagram*

$$\begin{array}{ccc} BD_0 & \longrightarrow & BD_\infty \\ \downarrow & & \downarrow \\ B\tilde{D}_0 & \longrightarrow & B\tilde{D}_\infty \end{array}$$

is  $(2n - 4)$ -cartesian.

*Proof.* This is an immediate consequence of a version of the so-called *Morlet's lemma of disjunction*, more precisely [10, Corollary 3.2 on page 29]. In the case at hand, the quoted result says that

$$\begin{array}{ccc} BD_g & \longrightarrow & BD_{g+1} \\ \downarrow & & \downarrow \\ B\tilde{D}_g & \longrightarrow & B\tilde{D}_{g+1} \end{array} \tag{7.7}$$

is  $(2n - 4)$ -cartesian for each  $g$ , and the claim follows by passage to the colimit.  $\square$

The next step is to let

$$BD_g \rightarrow \mathcal{E}_g \rightarrow B\tilde{\mathcal{D}}_g$$

be the  $(2n - 4)$ th stage of the Moore–Postnikov tower of the natural map  $BD_g \rightarrow B\tilde{\mathcal{D}}_g$ . Using naturality of the Moore–Postnikov tower, we obtain the  $(2n - 4)$ -th Moore–Postnikov stage

$$BD_\infty \rightarrow \mathcal{E}_\infty \rightarrow B\tilde{\mathcal{D}}_\infty$$

in the colimit.

**Lemma 7.8.** *The diagrams*

$$\begin{array}{ccc} BD_g & \longrightarrow & \mathcal{E}_g \\ \downarrow & & \downarrow \\ B\tilde{\mathcal{D}}_g & \xlongequal{\quad} & B\tilde{\mathcal{D}}_g \end{array}$$

are  $(2n - 4)$ -cartesian for all  $0 \leq g \leq \infty$ . The diagrams

$$\begin{array}{ccc} \mathcal{E}_g & \longrightarrow & \mathcal{E}_{g+1} \\ \downarrow & & \downarrow \\ B\tilde{\mathcal{D}}_g & \longrightarrow & B\tilde{\mathcal{D}}_{g+1} \end{array}$$

are homotopy cartesian, for each  $0 \leq g < \infty$ .

*Proof.* If  $X \xrightarrow{f} Y \xrightarrow{g} Z$  are two maps and  $f$  is  $k$ -connected, then

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow g \circ f & & \downarrow g \\ Z & \xlongequal{\quad} & Z \end{array}$$

is  $k$ -cartesian. Apply this observation to the definition of  $\mathcal{E}_g$  to get the first claim. The second claim follows from the  $(2n - 4)$ -cartesianness of (7.7), together with the following general fact: if

$$\begin{array}{ccc} X_0 & \longrightarrow & X_1 \\ \downarrow f_0 & & \downarrow f_1 \\ Y_0 & \longrightarrow & Y_1 \\ \downarrow g_0 & & \downarrow g_1 \\ Z_0 & \longrightarrow & Z_1 \end{array}$$

is a commutative diagram of spaces, the maps  $f_i$  are  $k$ -connected, and the maps  $g_i$  are  $k$ -coconnected, and the large rectangle is  $k$ -cartesian, then the lower square is homotopy cartesian.  $\square$

**Lemma 7.9.**

(1) *The natural map*

$$BD_g \rightarrow B\tilde{\mathcal{D}}_g$$

*induces an isomorphism on fundamental groups.*

- (2) For each  $1 \leq g \leq \infty$ , the fibration  $\mathcal{E}_g \rightarrow \tilde{\mathcal{D}}_g$  is “very simple” in the following sense: if  $\mathcal{F}_g$  denotes its homotopy fibre, then the monodromy action  $\mu(\gamma) : \mathcal{F}_g \rightarrow \mathcal{F}_g$ , for each  $\gamma \in \pi_1(B\tilde{\mathcal{D}}_g)$ , is homotopic to the identity.

*Proof.* (1): Surjectivity of  $\pi_1(B\text{Diff}_\partial(M)) \rightarrow \pi_1(\widetilde{B\text{Diff}}_\partial(M))$  follows from the very definition of  $\widetilde{\text{Diff}}_\partial(M)$ , and injectivity follows from Cerf’s theorem [11, Théorème 0].

(2): We follow the outline of a very similar argument contained in [44, §5.3]. It is enough to deal with the case of finite  $g$ , the case  $g = \infty$  follows by passage to the colimit. We present the core argument first. Note that the homotopy fibre of  $B\tilde{\mathcal{D}}_g \rightarrow B\mathcal{D}_g$  is the homogeneous space  $\tilde{\mathcal{D}}_g/\mathcal{D}_g$ . Let

$$\sigma : \tilde{\mathcal{D}}_0/\mathcal{D}_0 \rightarrow \tilde{\mathcal{D}}_g/\mathcal{D}_g$$

be the stabilization map and let  $\gamma \in \tilde{\mathcal{D}}_g$ . Note that  $\sigma$  is given by gluing in (block) diffeomorphisms in a fixed disc, and note that  $\gamma$  can be isotoped to a diffeomorphism that fixes this disc. It follows that

$$L_\gamma \circ \sigma \sim \sigma : \tilde{\mathcal{D}}_0/\mathcal{D}_0 \rightarrow \tilde{\mathcal{D}}_g/\mathcal{D}_g,$$

where  $L_\gamma$  denotes the left translation by  $\gamma$  on  $\tilde{\mathcal{D}}_g/\mathcal{D}_g$ .

To turn this observation into an argument concerning the monodromy action, note that the monodromy action of  $\gamma$  on  $\tilde{\mathcal{D}}_g/\mathcal{D}_g$  is exactly  $L_\gamma$ , and note that the map  $q : \tilde{\mathcal{D}}_g/\mathcal{D}_g \rightarrow \mathcal{F}_g$  is the  $(2n - 4)$ th Postnikov truncation, and equivariant with respect to the two monodromy actions. Also, pick a CW model for  $\mathcal{F}_g$ .

Since  $\pi_k(\mathcal{F}_g) = 0$  if  $k \geq 2n - 4$  by construction, two maps  $f_0, f_1 : K \rightarrow \mathcal{F}_g$  from a CW complex are homotopic if and only if the restrictions  $f_j|_{K^{(2n-4)}}$  are homotopic. Apply this to  $f_0 = \text{id}_{\mathcal{F}_g}$  and  $f_1 = \mu(\gamma)$ . The inclusion of the  $(2n - 4)$ -skeleton  $\mathcal{F}_g^{(2n-4)} \rightarrow \mathcal{F}_g$  can be factored through maps

$$\mathcal{F}_g^{(2n-4)} \xrightarrow{h} \tilde{\mathcal{D}}_0/\mathcal{D}_0 \xrightarrow{\sigma} \tilde{\mathcal{D}}_g/\mathcal{D}_g \xrightarrow{q} \mathcal{F}_g.$$

Let  $\gamma \in \tilde{\mathcal{D}}_g$ . Then

$$\mu(\gamma)|_{\mathcal{F}_g^{(2n-4)}} \sim \mu(\gamma) \circ q \circ \sigma \circ h \sim q \circ L_\gamma \circ \sigma \circ h \sim q \circ \sigma \circ h = \text{id}|_{\mathcal{F}_g^{(2n-4)}},$$

so that  $\mu(\gamma) \sim \text{id}$ , as desired.  $\square$

**Lemma 7.10.** *The commutator subgroup of  $\pi_1(\mathcal{E}_\infty)$  is perfect, and the diagram*

$$\begin{array}{ccc} \mathcal{E}_\infty & \longrightarrow & \mathcal{E}_\infty^+ \\ \downarrow & & \downarrow \\ B\tilde{\mathcal{D}}_\infty & \longrightarrow & B\tilde{\mathcal{D}}_\infty^+ \end{array}$$

*is homotopy cartesian.*

*Proof.* First note that the composition

$$B\mathcal{D}_\infty \rightarrow \mathcal{E}_\infty \rightarrow B\tilde{\mathcal{D}}_\infty$$

induces an isomorphism on fundamental groups by Lemma 7.9 (1), and the first map is  $(2n - 4)$ -connected by definition, so that both maps induce isomorphisms on fundamental groups. By Lemma 7.2, the commutator subgroup of  $\mathcal{E}_\infty$  is hence perfect.

For the proof that the square is homotopy cartesian, we use a theorem by Berrick [3, Theorem 1.1]. Let  $\mathcal{F}_\infty$  be the homotopy fibre of  $\mathcal{E}_\infty \rightarrow B\tilde{\mathcal{D}}_\infty$ . The latter map induces an isomorphism on fundamental groups, which has three consequences:  $\mathcal{F}_\infty$  is connected;  $\pi_1(\mathcal{F}_\infty)$  is abelian (because it is a quotient of  $\pi_2(\widetilde{B\tilde{\mathcal{D}}_\infty})$ ); and  $\mathcal{F}_\infty$  is nilpotent ( $\mathcal{F}_\infty$  is also the homotopy fibre of the map  $\widetilde{\mathcal{E}_\infty} \rightarrow \widetilde{B\tilde{\mathcal{D}}_\infty}$  induced on universal coverings, so it is the homotopy fibre of a map of 1-connected spaces; it is a general fact that such homotopy fibres are nilpotent, see e.g. [52, Proposition 4.4.1]). Hence  $\mathcal{F}_\infty^+ = \mathcal{F}_\infty$  is nilpotent, which is part of hypothesis (b) of [3, Theorem 1.1]. Lemma 7.9 (2) shows that the commutator subgroup of  $\pi_1(B\tilde{\mathcal{D}}_\infty)$  acts trivially on the homology of  $\mathcal{F}_\infty$ . It now follows from [3, Theorem 1.1] that

$$\mathcal{F}_\infty \rightarrow \mathcal{E}_\infty^+ \rightarrow B\tilde{\mathcal{D}}_\infty^+$$

is a fibre sequence, which is exactly the statement that the square is homotopy cartesian.  $\square$

To finish the proof of Proposition 7.5, we need a general property of the Quillen plus construction.

**Lemma 7.11.** *Let  $f : X \rightarrow Y$  be a  $n$ -connected map of connected spaces,  $n \geq 2$ , let  $P \subset \pi_1(X) = \pi_1(Y)$  be a perfect normal subgroup of the common fundamental group, and let  $X \rightarrow X^+$ ,  $Y \rightarrow Y^+$  be the Quillen plus constructions on  $P$ . Then  $f^+ : X^+ \rightarrow Y^+$  is  $n$ -connected.*

*Proof.* If  $P = \pi_1(X)$ , there is not much say, besides quoting Hurewicz' theorem. In the general case, let  $\tilde{X} \rightarrow X$ ,  $\tilde{Y} \rightarrow Y$  be the coverings with fundamental group  $P$ . Now  $X^+$  can be realized as the homotopy pushout

$$\begin{array}{ccc} \tilde{X} & \longrightarrow & \tilde{X}^+ \\ \downarrow & & \downarrow \\ X & \longrightarrow & X^+ \end{array}$$

see e.g. [35, p.374], and the claim follows.  $\square$

*Proof of Proposition 7.5.* Lemma 7.6, Lemma 7.8 and Lemma 7.10 imply that the large rectangle in

$$\begin{array}{ccccccc} BD_0 & \longrightarrow & B\mathcal{D}_\infty & \longrightarrow & \mathcal{E}_\infty & \longrightarrow & \mathcal{E}_\infty^+ \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ B\tilde{\mathcal{D}}_0 & \longrightarrow & B\tilde{\mathcal{D}}_\infty & \xlongequal{\quad} & B\tilde{\mathcal{D}}_\infty & \longrightarrow & B\tilde{\mathcal{D}}_\infty^+ \end{array}$$

is  $(2n - 4)$ -cartesian. This is the same as the rectangle in

$$\begin{array}{ccc} BD_0 & \longrightarrow & B\mathcal{D}_\infty^+ \longrightarrow \mathcal{E}_\infty^+ \\ \downarrow & & \downarrow \quad \downarrow \\ B\tilde{\mathcal{D}}_0 & \longrightarrow & B\tilde{\mathcal{D}}_\infty^+ \xlongequal{\quad} B\tilde{\mathcal{D}}_\infty^+ \end{array}$$

The map  $B\mathcal{D}_\infty^+ \rightarrow \mathcal{E}_\infty^+$  is  $(2n - 4)$ -connected by Lemma 7.11, and so the right square is  $(2n - 4)$ -cartesian. Hence the left square is  $(2n - 5)$ -cartesian, as required.  $\square$

7.2. Computation of the cohomology.

**Lemma 7.12.** *Let  $\mathcal{G}$  be the homotopy fibre of the map  $\Omega BBD \rightarrow \Omega BB\tilde{D}$  at the basepoint (equivalently, the homotopy fibre of  $BD_\infty^+ \rightarrow B\tilde{D}_\infty^+$ ), so that there is a fibre sequence*

$$\mathcal{G} \xrightarrow{j} \Omega_0 BBD \xrightarrow{p} \Omega_0 BB\tilde{D}.$$

Then

- (1)  $\mathcal{G}$  has the homotopy type of a connected  $(2n + 1)$ -fold loop space, and its rational homotopy groups in degrees  $k \leq 2n - 5$  are given by

$$\pi_k(\mathcal{G}) \otimes \mathbb{Q} = \begin{cases} \mathbb{Q} & 1 \leq k \leq 2n - 5, k \equiv 0 \pmod{4} \\ 0 & 1 \leq k \leq 2n - 5, k \not\equiv 0 \pmod{4}. \end{cases}$$

- (2) The map  $j$  induces the zero map on (reduced) rational homology up to degree  $2n - 5$ .

*Proof.* It is clear that  $\mathcal{G}$  has the homotopy type of a connected loop space, so for (1) it suffices to calculate the dimensions of the rational homotopy groups in the indicated range. By Lemma 7.2 and Lemma 7.10, the homotopy groups of  $\mathcal{G}$  agree with the homotopy groups of the fibre of  $B\text{Diff}_\partial(D^{2n+1}) \rightarrow B\tilde{\text{Diff}}_\partial(D^{2n+1})$ , up to degree  $2n - 6$ . The homotopy group  $\pi_k(B\tilde{\text{Diff}}_\partial(D^{2n+1}))$  can be identified with the group  $\Theta_{2n+1+k}$  of homotopy spheres, which is of course finite by [42]. The rational homotopy groups of  $B\text{Diff}_\partial(D^{2n+1})$  are famously related to algebraic  $K$ -theory, originally by [20]; the range we need was given by Krannich in [44, Corollary B]. This establishes (1).

For (2), consider the diagram

$$\begin{array}{ccc} \tilde{\mathcal{D}}_0/\mathcal{D}_0 & \longrightarrow & \mathcal{G} \\ \downarrow & & \downarrow j \\ B\mathcal{D}_0 & \longrightarrow & B\mathcal{D}_\infty^+ \\ \downarrow & & \downarrow \\ B\tilde{\mathcal{D}}_0 & \longrightarrow & B\tilde{\mathcal{D}}_\infty^+. \end{array}$$

The first named author proved in [14, Theorem 1.7], based on [7] and [59] that  $B\mathcal{D}_0 \rightarrow B\mathcal{D}_\infty$  induces the zero map on (reduced) rational homology, up to degree  $2n - 3$ . The top map in the diagram is  $(2n - 5)$ -connected by Proposition 7.5, and as  $B\tilde{\mathcal{D}}_0$  has finite homotopy groups,  $\tilde{\mathcal{D}}_0/\mathcal{D}_0 \rightarrow B\mathcal{D}_0$  is a rational homotopy equivalence. Putting these facts together, triviality of  $H_*(j; \mathbb{Q})$  follows.  $\square$

*Proof of Theorem 7.1.* (1) We first prove that  $H^*(B\tilde{\text{Diff}}_\partial(U_{g,1}^n); \mathbb{Q})$  is degreewise finitely generated; this implies via the homological stability theorem of [30] that  $H^*(B\tilde{\mathcal{D}}_\infty; \mathbb{Q})$  is degreewise finitely generated. Theorem 3.11 reduces this to the question whether  $H^*(\text{map}_*((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0 // \text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\cong}; \mathbb{Q})$  is degreewise finitely generated. Using Lemma 4.10 (3), the spectral sequence for (4.23) and the well-known fact that  $H^*(BGL_g(\mathbb{Z}); V)$  is finite-dimensional for each finite-dimensional  $GL_g(\mathbb{Q})$ -representation  $V$ , one reduces the problem to the question whether  $H^*(\text{map}_\partial((U_{g,1}^n)_\mathbb{Q}; BO_\mathbb{Q})^0 // \text{hAut}_\partial((U_{g,1}^n)_\mathbb{Q})^{\text{id}}; \mathbb{Q})$  is degreewise finitely generated, but that is clear from Corollary 3.22 and Serre class theory.

By Lemma 7.2,  $H^*(B\tilde{\mathcal{D}}_\infty; \mathbb{Q}) \cong H^*(\Omega_0 BBD; \mathbb{Q})$ . As  $\Omega_0 BBD$  is a connected  $(2n+1)$ -fold loop space, it is rationally equivalent to a product of Eilenberg–MacLane spaces, from which the claim about the structure of  $H^*(B\tilde{\mathcal{D}}_\infty; \mathbb{Q})$  follows.

(2) Let  $U_g^n := U_{g,1}^n \cup_{\partial} D^{2n+1}$ . Kupers’ finiteness theorem<sup>4</sup> [47, Corollary C] shows that  $B\text{Diff}^+(U_g^n)$  is of homologically finite type if  $n \geq 4$ . There is a fibre sequence  $\text{Fr}^+(TU_g^n) \rightarrow B\text{Diff}_{\partial}(U_{g,1}^n) \rightarrow B\text{Diff}^+(U_g^n)$ , and a straightforward spectral sequence argument proves that  $B\text{Diff}_{\partial}(U_{g,1}^n)$  is of homologically finite type. Together with [58, Corollary 1.3.2], this establishes the finiteness claim. The rest of the proof is parallel to that of (1).

(3) For  $n \geq 5$  and in degrees  $\leq n-4$ , injectivity of  $(B\rho_n)^*$  is contained in the statement of Theorem 1.8 (this is all what is truly needed for the proof of Theorem A). To get the claim in full generality, we use recent work of Stoll [69]. Theorem A of loc.cit. implies that

$$\text{hocolim}_g \text{BhAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}} \rightarrow \text{BGL}_{\infty}(\mathbb{Z})$$

induces an injective map in rational cohomology and that  $H^*(\text{hocolim}_g \text{BhAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}}; \mathbb{Q})$  is a free  $H^*(\text{BGL}_{\infty}(\mathbb{Z}); \mathbb{Q})$ -module. On the other hand, the forgetful map

$$\text{map}_*(M_{\mathbb{Q}}; \text{BO}_{\mathbb{Q}})^0 // \text{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}}) \cong \rightarrow \text{BhAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}}) \cong$$

has a right inverse given by the constant map. Using that  $\text{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}})_{\mathbb{Z}} = \text{hAut}_{\partial}((U_{g,1}^n)_{\mathbb{Q}}) \cong$  (Lemma 4.10 (4)), the claim therefore follows from Theorem 3.11.

Now the map  $B\tilde{\mathcal{D}}_\infty^+ \rightarrow \text{BGL}_{\infty}(\mathbb{Z})^+$  is a map of  $H$ -spaces between connected  $(2n+1)$ -fold loop spaces of finite rational type. For such spaces, the rational cohomology is the free graded-commutative algebra on the dual space of the rational homotopy. From the injectivity in rational cohomology that we just established, we deduce surjectivity in rational homotopy, and this proves the claim about the structure of  $H^*(B\tilde{\mathcal{D}}_\infty; \mathbb{Q})$ .

(4) For the purpose of computing cohomology, we can replace  $BI_{\infty}$  by its plus-construction or equivalently by the map  $p : \Omega_0 BBD \rightarrow \Omega_0 BB\tilde{\mathcal{D}}$ . The three spaces in the fibre sequence

$$\mathcal{G} \xrightarrow{j} \Omega_0 BBD \xrightarrow{p} \Omega_0 BB\tilde{\mathcal{D}}$$

are connected double loop spaces. Hence by the Milnor–Moore theorem [55], their rational homology (with the Pontrjagin product) is the free graded commutative algebra on the rational homotopy. It follows from Lemma 7.12 that the map  $\pi_*(j) \otimes \mathbb{Q}$  is also the trivial map up to degree  $(2n-5)$ . Hence  $\pi_*(p) \otimes \mathbb{Q}$  is injective up to degree  $(2n-5)$ , so  $H_*(p; \mathbb{Q})$  is injective, and  $H^*(p; \mathbb{Q})$  is surjective, both in degrees  $\leq (2n-5)$ . Using the knowledge about the rational homotopy of  $\mathcal{G}$ , we see that  $\text{coker}(\pi_*(p) \otimes \mathbb{Q})$  is concentrated in degrees  $4k+1$ ,  $k \geq 1$  and in these degrees has dimension 1 (again in degrees  $\leq (2n-5)$ ).

Hence the kernel of  $H^*(p; \mathbb{Q})$  must, in degrees  $\leq (2n-5)$ , be an ideal generated by classes in each degree  $4k+1$ ,  $k \geq 1$ . However, we know by Proposition 2.16 that the ideal generated by the Borel classes lies in the kernel of  $H^*(p; \mathbb{Q})$ , and by a dimension count, justified by items (1),(2),(3), must be equal to the kernel. This finishes the evaluation of  $H^*(\Omega_0 BBD; \mathbb{Q})$ , which by Lemma 7.2 gives  $H^*(B\tilde{\mathcal{D}}_\infty; \mathbb{Q})$ .  $\square$

<sup>4</sup>All that is needed for the proof of Theorem A is that  $H^*(B\tilde{\mathcal{D}}_\infty; \mathbb{Q})$  is finite-dimensional for  $* \leq n-4$ , and this follows from (1) and the argument we shall give for (4). So the appeal to [47] is not really necessary.

## REFERENCES

- [1] H. Bass, J. Milnor, and J.-P. Serre. Solution of the congruence subgroup problem for  $SL_n$  ( $n \geq 3$ ) and  $Sp_{2n}$  ( $n \geq 2$ ). *Inst. Hautes Études Sci. Publ. Math.*, (33):59–137, 1967.
- [2] A. Berglund and I. Madsen. Rational homotopy theory of automorphisms of manifolds. *Acta Math.*, 224(1):67–185, 2020.
- [3] A. J. Berrick. The plus-construction and fibrations. *Quart. J. Math. Oxford Ser. (2)*, 33(130):149–157, 1982.
- [4] A. Borel. Density properties for certain subgroups of semi-simple groups without compact components. *Ann. of Math. (2)*, 72:179–188, 1960.
- [5] A. Borel. Stable real cohomology of arithmetic groups. *Ann. Sci. École Norm. Sup. (4)*, 7:235–272 (1975), 1974.
- [6] A. Borel. Stable real cohomology of arithmetic groups. II. In *Manifolds and Lie groups (Notre Dame, Ind., 1980)*, volume 14 of *Progr. Math.*, pages 21–55. Birkhäuser, Boston, Mass., 1981.
- [7] B. Botvinnik and N. Perlmutter. Stable moduli spaces of high-dimensional handlebodies. *J. Topol.*, 10(1):101–163, 2017.
- [8] A. K. Bousfield. The localization of spaces with respect to homology. *Topology*, 14:133–150, 1975.
- [9] A. K. Bousfield.  $K$ -localizations and  $K$ -equivalences of infinite loop spaces. *Proc. London Math. Soc. (3)*, 44(2):291–311, 1982.
- [10] D. Burghelea, R. Lashof, and M. Rothenberg. *Groups of automorphisms of manifolds*. Lecture Notes in Mathematics, Vol. 473. Springer-Verlag, Berlin-New York, 1975. With an appendix (“The topological category”) by E. Pedersen.
- [11] J. Cerf. La stratification naturelle des espaces de fonctions différentiables réelles et le théorème de la pseudo-isotopie. *Inst. Hautes Études Sci. Publ. Math.*, (39):5–173, 1970.
- [12] W. Dwyer, M. Weiss, and B. Williams. A parametrized index theorem for the algebraic  $K$ -theory Euler class. *Acta Math.*, 190(1):1–104, 2003.
- [13] J. Ebert. A vanishing theorem for characteristic classes of odd-dimensional manifold bundles. *J. Reine Angew. Math.*, 684:1–29, 2013.
- [14] J. Ebert. Diffeomorphisms of odd-dimensional discs, glued into a manifold. *arXiv e-prints, to appear in Algebraic & Geometric Topology*, page arXiv:2107.00903, July 2021.
- [15] J. Ebert and O. Randal-Williams. Generalised Miller-Morita-Mumford classes for block bundles and topological bundles. *Algebr. Geom. Topol.*, 14(2):1181–1204, 2014.
- [16] J. Ebert and O. Randal-Williams. Torelli spaces of high-dimensional manifolds. *J. Topol.*, 8(1):38–64, 2015.
- [17] J. Ebert and O. Randal-Williams. The positive scalar curvature cobordism category. *arXiv e-prints, to appear in Duke Math. Journal*, page arXiv:1904.12951, April 2019.
- [18] H. Espic and B. Saleh. On the group of homotopy classes of relative homotopy automorphisms. *arXiv e-prints*, page arXiv:2002.12083, February 2020.
- [19] P. Etingof, O. Golberg, S. Hensel, T. Liu, A. Schwendner, D. Vaintrob, and E. Yudovina. *Introduction to representation theory*, volume 59 of *Student Mathematical Library*. American Mathematical Society, Providence, RI, 2011. With historical interludes by Slava Gerovitch.
- [20] F. T. Farrell and W. C. Hsiang. On the rational homotopy groups of the diffeomorphism groups of discs, spheres and aspherical manifolds. In *Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 1*, Proc. Sympos. Pure Math., XXXII, pages 325–337. Amer. Math. Soc., Providence, R.I., 1978.
- [21] Y. Félix, S. Halperin, and J.-C. Thomas. *Rational homotopy theory*, volume 205 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2001.
- [22] W. Fulton. *Young tableaux*, volume 35 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1997. With applications to representation theory and geometry.
- [23] W. Fulton and J. Harris. *Representation theory*, volume 129 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1991. A first course, Readings in Mathematics.
- [24] S. Galatius, A. Kupers, and O. Randal-Williams.  $E_2$ -cells and mapping class groups. *Publ. Math. Inst. Hautes Études Sci.*, 130:1–61, 2019.
- [25] S. Galatius and O. Randal-Williams. Stable moduli spaces of high-dimensional manifolds. *Acta Math.*, 212(2):257–377, 2014.

- [26] S. Galatius and O. Randal-Williams. Homological stability for moduli spaces of high dimensional manifolds. II. *Ann. of Math. (2)*, 186(1):127–204, 2017.
- [27] S. Galatius and O. Randal-Williams. Homological stability for moduli spaces of high dimensional manifolds. I. *J. Amer. Math. Soc.*, 31(1):215–264, 2018.
- [28] S. Galatius, U. Tillmann, I. Madsen, and M. Weiss. The homotopy type of the cobordism category. *Acta Math.*, 202(2):195–239, 2009.
- [29] R. Goodman and N. R. Wallach. *Symmetry, representations, and invariants*, volume 255 of *Graduate Texts in Mathematics*. Springer, Dordrecht, 2009.
- [30] M. Grey. On rational homological stability for block automorphisms of connected sums of products of spheres. *Algebr. Geom. Topol.*, 19(7):3359–3407, 2019.
- [31] V. W. Guillemin and S. Sternberg. *Supersymmetry and equivariant de Rham theory*. Mathematics Past and Present. Springer-Verlag, Berlin, 1999. With an appendix containing two reprints by Henri Cartan [MR0042426 (13,107e); MR0042427 (13,107f)].
- [32] A. J. Hahn and O. T. O’Meara. *The classical groups and K-theory*, volume 291 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1989. With a foreword by J. Dieudonné.
- [33] B. Hanke, T. Schick, and W. Steimle. The space of metrics of positive scalar curvature. *Publ. Math. Inst. Hautes Études Sci.*, 120:335–367, 2014.
- [34] J. L. Harer. Stability of the homology of the mapping class groups of orientable surfaces. *Ann. of Math. (2)*, 121(2):215–249, 1985.
- [35] A. Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002.
- [36] J.-C. Hausmann and D. Husemoller. Acyclic maps. *Enseign. Math. (2)*, 25(1-2):53–75, 1979.
- [37] F. Hebestreit, M. Land, W. Lück, and O. Randal-Williams. A vanishing theorem for tautological classes of aspherical manifolds. *Geom. Topol.*, 25(1):47–110, 2021.
- [38] F. Hebestreit and N. Perlmutter. Cobordism categories and moduli spaces of odd dimensional manifolds. *Adv. Math.*, 353:526–590, 2019.
- [39] P. Hilton, G. Mislin, and J. Roitberg. *Localization of nilpotent groups and spaces*. North-Holland Mathematics Studies, No. 15. North-Holland Publishing Co., Amsterdam-Oxford; American Elsevier Publishing Co., Inc., New York, 1975.
- [40] F. Hirzebruch. *Neue topologische Methoden in der algebraischen Geometrie*. Ergebnisse der Mathematik und ihrer Grenzgebiete, (N.F.), Heft 9. Springer-Verlag, Berlin-Göttingen-Heidelberg, 1956.
- [41] K. Igusa. The stability theorem for smooth pseudoisotopies. *K-Theory*, 2(1-2):vi+355, 1988.
- [42] M. A. Kervaire and J. W. Milnor. Groups of homotopy spheres. I. *Ann. of Math. (2)*, 77:504–537, 1963.
- [43] M. Krannich and O. Randal-Williams. Diffeomorphisms of discs and the second Weiss derivative of  $B\text{Top}(-)$ . *arXiv e-prints*, page arXiv:2109.03500, September 2021.
- [44] Manuel Krannich. A homological approach to pseudoisotopy theory. I. *Invent. Math.*, 227(3):1093–1167, 2022.
- [45] M. Kreck. Cobordism of odd-dimensional diffeomorphisms. *Topology*, 15(4):353–361, 1976.
- [46] A. Kupers and O. Randal-Williams. On the cohomology of Torelli groups. *Forum Math. Pi*, 8:e7, 83, 2020.
- [47] Alexander Kupers. Some finiteness results for groups of automorphisms of manifolds. *Geom. Topol.*, 23(5):2277–2333, 2019.
- [48] J.-L. Loday. *Cyclic homology*, volume 301 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 1998. Appendix E by María O. Ronco, Chapter 13 by the author in collaboration with Teimuraz Pirashvili.
- [49] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, second edition, 1995. With contributions by A. Zelevinsky, Oxford Science Publications.
- [50] I. Madsen and M. Weiss. The stable moduli space of Riemann surfaces: Mumford’s conjecture. *Ann. of Math. (2)*, 165(3):843–941, 2007.
- [51] J. P. May. *The geometry of iterated loop spaces*. Lecture Notes in Mathematics, Vol. 271. Springer-Verlag, Berlin-New York, 1972.
- [52] J. P. May and K. Ponto. *More concise algebraic topology*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 2012. Localization, completion, and model categories.

- [53] D. McDuff and G. Segal. Homology fibrations and the “group-completion” theorem. *Invent. Math.*, 31(3):279–284, 1975/76.
- [54] E. Y. Miller. The homology of the mapping class group. *J. Differential Geom.*, 24(1):1–14, 1986.
- [55] J. W. Milnor and J. C. Moore. On the structure of Hopf algebras. *Ann. of Math. (2)*, 81:211–264, 1965.
- [56] S. Morita. Characteristic classes of surface bundles. *Invent. Math.*, 90(3):551–577, 1987.
- [57] A. J. Nicas. Induction theorems for groups of homotopy manifold structures. *Mem. Amer. Math. Soc.*, 39(267):vi+108, 1982.
- [58] N. Perlmutter. Homological stability for the moduli spaces of products of spheres. *Trans. Amer. Math. Soc.*, 368(7):5197–5228, 2016.
- [59] N. Perlmutter. Homological stability for diffeomorphism groups of high-dimensional handlebodies. *Algebr. Geom. Topol.*, 18(5):2769–2820, 2018.
- [60] C. Procesi. *Lie groups*. Universitext. Springer, New York, 2007. An approach through invariants and representations.
- [61] Putman, A. The Borel density theorem. Unpublished note, available at <https://www3.nd.edu/~andyp/notes/>, 2019.
- [62] D. Quillen. Finite generation of the groups  $K_i$  of rings of algebraic integers. In *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 179–198. Lecture Notes in Math., Vol. 341, 1973.
- [63] F. S. Quinn. *A geometric formulation of surgery*. ProQuest LLC, Ann Arbor, MI, 1970. Thesis (Ph.D.)–Princeton University.
- [64] O. Randal-Williams. ‘Group-completion’, local coefficient systems and perfection. *Q. J. Math.*, 64(3):795–803, 2013.
- [65] O. Randal-Williams. Cohomology of automorphism groups of free groups with twisted coefficients. *Selecta Math. (N.S.)*, 24(2):1453–1478, 2018.
- [66] Oscar Randal-Williams. The family signature theorem. *arXiv e-prints*, page arXiv:2204.11696, April 2022.
- [67] E. Riehl. *Categorical homotopy theory*, volume 24 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2014.
- [68] E. H. Spanier. *Algebraic topology*. McGraw-Hill Book Co., New York-Toronto, Ont.-London, 1966.
- [69] Robin Stoll. The stable cohomology of self-equivalences of connected sums of products of spheres, 2022.
- [70] D. Sullivan. Genetics of homotopy theory and the Adams conjecture. *Ann. of Math. (2)*, 100:1–79, 1974.
- [71] D. Sullivan. Infinitesimal computations in topology. *Inst. Hautes Études Sci. Publ. Math.*, (47):269–331 (1978), 1977.
- [72] U. Tillmann. On the homotopy of the stable mapping class group. *Invent. Math.*, 130(2):257–275, 1997.
- [73] B. Tshishiku. Borel’s stable range for the cohomology of arithmetic groups. *J. Lie Theory*, 29(4):1093–1102, 2019.
- [74] W. van der Kallen. Homology stability for linear groups. *Invent. Math.*, 60(3):269–295, 1980.
- [75] F. Waldhausen, B. Jahren, and J. Rognes. *Spaces of PL manifolds and categories of simple maps*, volume 186 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2013.
- [76] C. T. C. Wall. Classification problems in differential topology. II. Diffeomorphisms of handlebodies. *Topology*, 2:263–272, 1963.
- [77] C. T. C. Wall. Classification problems in differential topology. VI. Classification of  $(s - 1)$ -connected  $(2s + 1)$ -manifolds. *Topology*, 6:273–296, 1967.
- [78] M. Weiss and B. Williams. Automorphisms of manifolds and algebraic  $K$ -theory. I. *K-Theory*, 1(6):575–626, 1988.

*Email address:* johannes.ebert@uni-muenster.de

MATHEMATISCHES INSTITUT, WWU MÜNSTER, EINSTEINSTR. 62, 48149 MÜNSTER, GERMANY

*Email address:* jens.reinhold@posteo.de