THE CHARNEY-DAVIS CONJECTURE FOR SIMPLE THIN POLYOMINOES

MANOJ KUMMINI AND DHARM VEER

ABSTRACT. Let \mathcal{P} be a simple thin polyomino and \Bbbk a field. Let R be the toric \Bbbk -algebra associated to \mathcal{P} . Write the Hilbert series of R as $h_R(t)/(1-t)^{\dim(R)}$. We show that

$$(-1)^{\left\lfloor\frac{\deg h_R(t)}{2}\right\rfloor}h_R(-1) \ge 0$$

if *R* is Gorenstein. This shows that the Gorenstein rings associated to simple thin polyominoes satisfy the Charney-Davis conjecture.

1. INTRODUCTION

The Charney-Davis conjecture [CD95, Conjecture D] asserts that if h(t) is the *h*-polynomial of a flag simplicial homology (d - 1)-sphere, then $(-1)^{\lfloor \frac{d}{2} \rfloor}h(-1) \ge 0$. Stanley [Sta00, Problem 4] extended this conjecture to Gorenstein^{*} flag simplicial complexes. Generalizing it further, Reiner and Welker [RW05, Question 4.4] posed the following:

Question 1.1. Let \Bbbk be a field and *R* a standard graded Gorenstein Koszul \Bbbk -algebra. Write the Hilbert series of *R* as $h_R(t)/(1-t)^{\dim(R)}$. Is

$$(-1)^{\left\lfloor \frac{\deg h_R(t)}{2} \right\rfloor} h_R(-1) \ge 0$$
?

We say that a standard graded Gorenstein Koszul k-algebra R is CD if it gives an affirmative answer to the above question. In this article, we show that the toric k-algebras associated to simple thin polyominoes are CD, when they are Gorenstein.

Suppose that, in the notation of Question 1.1, deg $h_R(t)$ is odd. Then $h_R(-1) = 0$; see, e.g., [BH93, Corollary 4.4.6]. Therefore Question 1.1 is open only when deg $h_R(t)$ is even. See the bibliography of [RW05] and of [Sta00] for various classes of rings that are CD. A class of CD rings related to the ones we study in this paper are Gorenstein Hibi rings [Brä04, Corollary 4.3]. Recently, D'Alì and Venturello [1] proved that the answer to Question 1.1 is negative in general.

A cell in \mathbb{R}^2 is a set of the form $\{(x, y) \in \mathbb{R}^2 \mid a \le x \le a + 1, b \le y \le b + 1\}$ where $(a, b) \in \mathbb{Z}^2$. Let \mathcal{P} be a finite collection of cells. Then \mathcal{P} determines a unique topological subspace $\operatorname{sp}(\mathcal{P}) := \bigcup_{C \in \mathcal{P}} C$ of \mathbb{R}^2 . By abuse of terminology, we assign the topological attributes to \mathcal{P} that $\operatorname{sp}(\mathcal{P})$ has. We say that \mathcal{P} is a *polyomino* if \mathcal{P} is connected and does not have a finite cut-set [Sta97, 4.7.18] (i.e., $\operatorname{sp}(\mathcal{P})$ has these properties). There is a k-algebra $\Bbbk[\mathcal{P}]$ of finite-type associated to \mathcal{P} [Qur12].

We say that a polyomino \mathcal{P} is *simple* if $sp(\mathcal{P})$ is simply connected; it is *thin* if it does not have a 2 × 2 square such as the one shown in Figure 1. The S-property of simple thin polyominoes was introduced in [RR21] to characterise such polyominoes \mathcal{P} for which $\Bbbk[\mathcal{P}]$ is Gorenstein. If \mathcal{P} is simple then $\Bbbk[\mathcal{P}]$ is a Koszul algebra [QSS17, Corollary 2.3]. Therefore it is natural to ask whether $\Bbbk[\mathcal{P}]$ is CD if \mathcal{P} is a simple thin polyomino with the S-property. In this regard, we show the following:

Theorem 1.2. Let \mathcal{P} be a collection of cells such that its connected components are simple thin polyominoes with the *S*-property. Then $\Bbbk[\mathcal{P}]$ is CD.

Note that if $\mathcal{P}_1, \ldots, \mathcal{P}_m$ are the connected components of \mathcal{P} , then $\Bbbk[\mathcal{P}] \simeq \Bbbk[\mathcal{P}_1] \otimes_{\Bbbk} \cdots \otimes_{\Bbbk} \Bbbk[\mathcal{P}_m]$. Therefore $\Bbbk[\mathcal{P}]$ Gorenstein (respectively, Koszul) if and only if $\Bbbk[\mathcal{P}_i]$ is Gorenstein (respectively, Koszul) for each *i*.

Section 2 contains the definitions and preliminaries. Proof of the theorem is given in Section 3.

²⁰²⁰ Mathematics Subject Classification. 13D40 (Primary).

MK was partly supported by the grant CRG/2018/001592 from Science and Engineering Research Board, India and by an Infosys Foundation fellowship. DV was partly supported by an Infosys Foundation fellowship.



FIGURE 1. A 2×2 square polyomino

2. Preliminaries

Gorenstein rings, Koszul rings, Hilbert series, etc. Let *R* be a finitely generated k-algebra. Throughout this paper, we deal with only standard graded k-algebras *R*, i.e., *R* is generated as a k-algebra by homogeneous elements of degree 1. We say that *R* is *Gorenstein* if it is Cohen-Macaulay and $\operatorname{Ext}_{R}^{\dim R}(\Bbbk, R) \simeq \Bbbk$. We say that *R* is *Koszul* if k has a linear free resolution as an *R*-module.

The Hilbert series $H_R(t)$ of R is the formal power series $\sum_{i \in \mathbb{N}} \dim_{\mathbb{K}} R_i t^i$ where for each i, R_i is the finitedimensional \mathbb{k} -vector-space of the homogeneous elements of R of degree i. There exists a unique polynomial $h_R(t)$ such that

$$H_R(t) = \frac{h_R(t)}{(1-t)^{\dim R}}.$$

Question 1.1 asks whether $(-1)^{\lfloor \frac{\deg h_R(t)}{2} \rfloor} h_R(-1)$ is non-negative for all Gorenstein Koszul algebras R. Parenthetically, note that, if R is Cohen-Macaulay, deg $h_R(t)$ is the Castelnuovo-Mumford regularity of R.

Polyominoes. Let \mathcal{P} be a finite collection of cells. As mentioned in the introduction, we treat \mathcal{P} interchangeably with the topological space $\operatorname{sp}(\mathcal{P})$. Let $S = \Bbbk[\{x_{i,j} \mid (i,j) \in \mathcal{P} \cap \mathbb{Z}^2\}]$ be the standard graded polynomial ring in the variables $x_{i,j}$. Let $\mathfrak{a}_{\mathcal{P}}$ be the binomial ideal generated by the binomials $x_{i,j}x_{k,l} - x_{k,j}x_{i,l}$ for all $(i, j), (k, l) \in \mathcal{P} \cap \mathbb{Z}^2$ such that the rectangle with vertices (i, j), (k, l), (k, j) and (i, l) is a subset of $\operatorname{sp}(\mathcal{P})$. Define $\Bbbk[\mathcal{P}] = S/\mathfrak{a}_{\mathcal{P}}$ [Qur12]. If \mathcal{P} is a simple polyomino, then $\Bbbk[\mathcal{P}]$ is a Koszul Cohen-Macaulay integral domain [QSS17, Corollary 2.3].

For $k \in \mathbb{N}$, a *k*-rook configuration in \mathcal{P} is an arrangement of *k* rooks in pairwise non-attacking positions. The rook polynomial $r_{\mathcal{P}}(t)$ of \mathcal{P} is $\sum_{k \in \mathbb{N}} r_k t^k$ where r_k is the number of *k*-rook configurations in \mathcal{P} . The rook number $r(\mathcal{P})$ of \mathcal{P} is the degree of $r_{\mathcal{P}}(t)$, i.e., the largest *k* such that there is a *k*-rook configuration in \mathcal{P} . If \mathcal{P} is a simple thin polyomino, then $h_{k \mid \mathcal{P}_1}(t) = r_{\mathcal{P}}(t)$ [RR21, Theorem 1.1].

Let $C, D \in \mathcal{P}$. We say that *C* is a *neighbour* of *D* if $C \cap D$ is a line segment. A *path* from *C* to *D* is a sequence of cells $C = C_0, C_1, \ldots, C_m = D$ such that for all $i \neq j, C_i \neq C_j$ and for all $1 \leq i \leq m, C_i$ is a neighbour of C_{i-1} . If \mathcal{P} is a simple thin polyomino, then for all cells C, D of \mathcal{P} , there is a unique path from *C* to *D*.

A *inner interval* of \mathcal{P} is a subcollection I of \mathcal{P} such that sp(I) (which is a subspace of $sp(\mathcal{P})$) is a rectangle with vertices (i_1, j_1) , $(i_1 + 1, j_1)$, (i_1, j_2) and $(i_1 + 1, j_2)$ or a rectangle with vertices (i_1, j_1) , $(i_1, j_1 + 1)$, (i_2, j_1) and $(i_2, j_1 + 1)$ for some $i_1, i_2, j_1, j_2 \in \mathbb{Z}$ with $i_1 < i_2$ and $j_1 < j_2$. An inner interval of \mathcal{P} is *maximal* if it is maximal under inclusion.

Let \mathcal{P} be a simple thin polyomino. Observe that any cell of \mathcal{P} belongs to at most two maximal inner intervals. A cell *C* is said to be an *end-cell* of a maximal inner interval *I* if $C \in I$ and *C* has exactly one neighbour cell in *I*. A cell of \mathcal{P} is called *single* if it belongs to exactly one maximal inner interval of \mathcal{P} . We say that \mathcal{P} has the *S*-property if every maximal inner interval of \mathcal{P} has exactly one single cell. For a simple thin polyomino $\mathcal{P}, \mathbb{k}[\mathcal{P}]$ is Gorenstein if and only if \mathcal{P} has the *S*-property [RR21, Theorem 4.2].

3. Proof of the Theorem

We begin with an observation about how Hilbert series and rook polynomials behave in disjoint unions of polyominoes.




FIGURE 2. Collapse datum (cf. Definition 3.2)

Proposition 3.1. Let \mathcal{P} be a finite collection of cells. Write $\mathcal{P}_1, \ldots, \mathcal{P}_m$ for the connected components. Then:

$$h_{\mathbb{k}[\mathcal{P}]}(t) = \prod_{i=1}^{m} h_{\mathbb{k}[\mathcal{P}_i]}(t) \text{ and } r_{\mathcal{P}}(t) = \prod_{i=1}^{m} r_{\mathcal{P}_i}(t)$$

In particular, if \mathcal{P}_i is a simple thin polyomino for each *i*, then $h_{\mathbb{K}[\mathcal{P}]}(t) = r_{\mathcal{P}}(t)$.

Proof. Vertices of the \mathcal{P}_i are disjoint, so $\mathbb{k}[\mathcal{P}] \simeq \mathbb{k}[\mathcal{P}_1] \otimes_{\mathbb{k}} \cdots \otimes_{\mathbb{k}} \mathbb{k}[\mathcal{P}_m]$. Hence $H_{\mathbb{k}[\mathcal{P}]}(t) = \prod_{i=1}^m H_{\mathbb{k}[\mathcal{P}_i]}(t)$, from which it follows that $h_{\mathbb{k}[\mathcal{P}]}(t) = \prod_{i=1}^m h_{\mathbb{k}[\mathcal{P}_i]}(t)$. Let $k \in \mathbb{N}$. Then *k*-rook configurations in \mathcal{P} corresponds to independent choices of k_i -rook configurations in \mathcal{P}_i for each $1 \le i \le m$ and for each tuple $(k_1, \ldots, k_m) \in \mathbb{N}^m$ with $\sum_i k_i = k$. Hence $r_{\mathcal{P}}(t) = \prod_{i=1}^m r_{\mathcal{P}_i}(t)$. The final assertion now follows from noting that for each $i, h_{\mathbb{k}[\mathcal{P}_i]}(t) = r_{\mathcal{P}_i}(t)$ since \mathcal{P}_i is a simple thin polyomino [RR21, Theorem 1.1]. \Box

Let \mathcal{P} be a simple thin polyomino. In [RR21, Definition 3.4], Rinaldo and Romeo introduced a notion of collapsing \mathcal{P} in a maximal inner interval, and showed that if \mathcal{P} has at least two maximal inner intervals, then there exists a maximal inner interval in which \mathcal{P} is collapsible [RR21, Proposition 3.7]. We need a refinement of this result for simple thin polyominoes with the *S*-property, for which we rephrase [RR21, Definition 3.4] in a slightly different way.

Definition 3.2. Let \mathcal{P} be a simple thin polyomino. A *collapse datum* on \mathcal{P} is a tuple (I, J, \mathcal{P}^I) , where I and J are maximal inner intervals and \mathcal{P}^I is a sub-polyomino of \mathcal{P} satisfying the following conditions:

- (1) *J* is the only maximal inner interval of \mathcal{P} such that $I \cap J$ is a cell;
- (2) $\mathcal{P}^I \subseteq J$ and $I \cap J \notin \mathcal{P}^I$.
- (3) $\mathcal{P} \setminus (I \cup \mathcal{P}^I)$ is a non-empty sub-polyomino of \mathcal{P} .

Figure 2 gives an example of a collapse datum. Note that since \mathcal{P}^I is a sub-polyomino of \mathcal{P} and $\mathcal{P}^I \subseteq J$, it is an inner interval if it is non-empty. When \mathcal{P} has at least two maximal inner intervals, the maximal inner intervals *I* and *J* defined in the Definition 3.2 exist by [RR21, Lemma 3.6].

Discussion 3.3. Let \mathcal{P} be a simple thin polyomino with S-property. Assume that \mathcal{P} is not a cell. Then it has a collapse datum (I, J, \mathcal{P}^I) . Since, additionally, \mathcal{P} has the S-property, I has exactly two cells and \mathcal{P}^I is either empty or a cell. Write $I = \{C, D\}$ with C denoting the single cell of I and $\{D\} = I \cap J$. Denote the single cell of J by E. Let C_1, \ldots, C_k be the cells of J different from D and E. For $1 \leq i \leq k$, let B_i be the cell in \mathcal{P} such that $B_i \notin J$ and C_i is a neighbour cell of B_i ; such a B_i must exist, since C_i is not a single cell. We now consider the various cases.

(1) $\mathcal{P}^I \neq \emptyset$. Equivalently, $\mathcal{P}^I = \{E\}$. Then we may assume that C_k is an end-cell of J and for all $i \in \{1, \ldots, k - 1\}$, C_i and C_{i+1} are neighbours. So B_i and B_{i+1} can not be neighbours, since \mathcal{P} is thin. Hence for all $i \in \{1, \ldots, k\}$ whether B_i is above C_i or is below C_i determined by whether i is even or odd. Therefore in the neighbourhood of J, \mathcal{P} is as shown in Figure 3.




FIGURE 3. \mathcal{P}^I non-empty




FIGURE 4. \mathcal{P}^I is empty, *E* is a neighbour of C_k but not an end-cell of *J*




FIGURE 5. \mathcal{P}^I is empty, *E* is an end-cell of *J*

(2) $\mathcal{P}^I = \emptyset$, C_k is an end-cell of J and E is a neighbour cell of C_k . We may assume that for all $i \in \{1, \ldots, k-2\}$, C_i and C_{i+1} are neighbours, so B_i and B_{i+1} cannot be neighbours. Therefore, using the same considerations as in the above case, we see that \mathcal{P} is as shown in Figure 4 in the neighbourhood of J.

(3) $\mathcal{P}^I = \emptyset$ and *E* is an end-cell of *J*. Then, in the neighbourhood of *J*, \mathcal{P} is as shown in Figure 5.

(4) $\mathcal{P}^I = \emptyset$, C_k is an end-cell of J and E is not a neighbour cell of C_k . Then, in the neighbourhood of J, \mathcal{P} is one of the figures in Figure 6.

Discussion 3.4. By the *first end-cell* of *J*, we mean

$$\begin{cases} E, & \text{if } \mathcal{P}^I = \{E\}; \\ I \cap J, & \text{if } \mathcal{P}^I = \emptyset. \end{cases}$$

(Note that in both of the above cases, the cell in question is an end-cell of J.) We call the other end-cell of J the second end-cell of J. If E is the second end-cell of J, then E has exactly one neighbour cell. If C_k is the second end-cell of J and E is not a neighbour cell of C_k , then C_k has exactly two neighbour cells. If C_k is the second end-cell of J and E is a neighbour cell of C_k , then C_k has two or three neighbour cells; see Figures 3, 4, 5 and 6.




FIGURE 6. \mathcal{P}^{I} is empty, *E* is not a neighbour of C_{k}

The next lemma shows that simple thin polyominoes with the *S*-property have a special collapse datum. See Figure 7 for an example to illustrate that condition (2) of the next lemma cannot be deleted.

Lemma 3.5. Let \mathcal{P} be a simple thin polyomino with S-property. Assume that \mathcal{P} is not a cell. Then there exists a collapse datum (I, J, \mathcal{P}^I) of \mathcal{P} such that one of the following holds:

- (1) The second end-cell of *J* has at most two neighbour cells.
- (2) If the second end-cell of J has three neighbour cells, then one of its neighbour cells is both a single cell and an end-cell of the maximal inner interval containing it.

Proof. We proceed by induction on number of cells of \mathcal{P} . Let (I, J, \mathcal{P}^I) be a collapse datum of \mathcal{P} . If the neighbourhood of J in \mathcal{P} looks like the ones given in Figures 3, 5 or 6, then (1) holds. Therefore we may assume that we are in the situation of Figure 4. Let E be the single cell of J, C_k be the second end-cell of J. We may assume that B_k and B_{k+1} as marked in Figure 4 exist.

If B_k is both a single cell and an end-cell of the maximal inner interval that contains $\{B_k, C_k, B_{k+1}\}$ then (2) holds. Hence we may assume that either B_k is not a single cell or it is not an end-cell of the maximal inner interval that contains $\{B_k, C_k, B_{k+1}\}$. Similarly for B_{k+1} . Let


 $\mathcal{P}' = (\mathcal{P} \setminus \{A \in \mathcal{P} : \text{the (unique) path between } A \text{ and } E \text{ does not contain } C_k\}) \cup \{E\}.$

(E.g., in Figure 4, \mathcal{P}' is the sub-polyomino comprising E and the cells reachable from E through C_k .) Observe that \mathcal{P}' is a simple thin polyomino. We first show that \mathcal{P}' has the S-property. Let L be a maximal inner interval of \mathcal{P}' . Then L is a maximal inner interval of \mathcal{P} or $L = \{C_k, E\}$. In both cases, L has a unique single cell. Thus, \mathcal{P}' has the S-property. Also note that \mathcal{P}' is not a cell. The number of cells in \mathcal{P} are strictly less than the number of cells in \mathcal{P} . Therefore, by induction, there exists a collapse datum $(I', J', \mathcal{P}'^{I'})$ of \mathcal{P}' satisfying the assertion of the lemma for \mathcal{P}' .

Note that $I' \neq \{C_k, E\}$ and $J' \neq \{C_k, E\}$; therefore I' and J' are maximal inner intervals of \mathcal{P} . Let J_1 be a maximal inner interval of \mathcal{P} such that $J_1 \cap I'$ is a cell. Since \mathcal{P} is simple, $J_1 \subset \mathcal{P}'$, so J_1 is a maximal inner interval of \mathcal{P}' . Hence $J_1 = J'$. Note that $\mathcal{P}'^{I'} \subseteq J'$ and that $I' \cap J' \notin \mathcal{P}'^{I'}$. Moreover, since $J' \neq \{C_k, E\}$, it follows that $\{C_k, E\} \subseteq \mathcal{P}' \setminus (I' \cup \mathcal{P}'^{I'})$; hence $\mathcal{P} \setminus (I' \cup \mathcal{P}'^{I'})$ which equals

 $(\mathcal{P}' \smallsetminus (I' \cup \mathcal{P}'^{I'})) \cup \{A \in \mathcal{P} : \text{the path between } A \text{ and } E \text{ does not contain } C_k\}$

is a non-empty sub-polyomino of \mathcal{P} . Hence $(I', J', \mathcal{P'}^{I'})$ is a collapse datum of \mathcal{P} that satisfies the assertion of the lemma for \mathcal{P} .





k	<i>k</i> -rook configurations	number	<i>k</i> -rook configurations	number
	that have a rook at C		that do not have a rook at	
			С	
0	There are no 0-rook con-	0	Ø	1
	figurations that have a			
	rook at C			
1	$\{C\}$	1	${A}, {B}$	2
2	$\{C, A\}$	1	none	0
$k \ge 3$	none	0	none	0
	$r_{P,C}(t)$	$t + t^2$	$r_{P,\widehat{C}}(t)$	1 + 2t

TABLE 1. Calculation of $r_{P,C}(t)$ and $r_{P,\widehat{C}}(t)$



FIGURE 8. Polyomino from Example 3.8

Discussion 3.6. Let \mathcal{P} be a simple thin polyomino and C be a single cell in \mathcal{P} . Let $r_{\mathcal{P},C}(t)$ be the polynomial $\sum_{k \in \mathbb{N}} r_k t^k$, where r_k is the number of k-rook configurations in \mathcal{P} that have a rook at C. Let $r_{\mathcal{P},\widehat{C}}(t)$ be the polynomial $\sum_{k \in \mathbb{N}} r_k t^k$, where r_k is the number of k-rook configurations in \mathcal{P} that have no rook at C. Then,

$$r_{\mathcal{P}}(t) = r_{\varphi,\widehat{C}}(t) + r_{\mathcal{P},C}(t).$$

Let *I* be the maximal inner interval of \mathcal{P} such that $C \in I$. Let $r_{\mathcal{P},\hat{I}}(t)$ be the polynomial $\sum_{k \in \mathbb{N}} r_k t^k$, where r_k is the number of *k*-rook configurations in \mathcal{P} that has no rook at any cell of *I*. Note that $r_{\mathcal{P},C}(t) = r_{\mathcal{P},\hat{I}}(t)t$. Hence,

(3.7)
$$r_{\varphi}(t) = r_{\varphi \ \widehat{C}}(t) + r_{\varphi \ \widehat{I}}(t)t.$$

Example 3.8. We illustrate the above definitions now. Let \mathcal{P} be the polyomino as shown in the Figure 8. Note that *C* is a single cell in \mathcal{P} . The polynomials $r_{\mathcal{P},\widehat{C}}(t)$ and $r_{\mathcal{P},C}(t)$ are calculated in Table 1. The unique maximal inner interval *I* containing *C* is $\{B, C\}$. Hence $r_{\mathcal{P},\widehat{I}}(t) = 1 + t$, since this is the rook polynomial of the polyomino consisting of just the cell *A*. On the other hand, the number of *k*-rook configurations in \mathcal{P} for k = 0, 1, 2 are, respectively, 1, 3, 1; hence $r_{\mathcal{P}}(t) = 1 + 3t + t^2$. We thus see that

$$r_{\mathcal{P}}(t) = r_{\mathcal{P},\widehat{C}}(t) + r_{\mathcal{P},C}(t) = r_{\mathcal{P},\widehat{C}}(t) + r_{\mathcal{P},\widehat{I}}(t)t.$$

We now wish to express $r_{\mathcal{P},\widehat{C}}(t)$ and $r_{\mathcal{P},\widehat{I}}(t)$ as the rook polynomials of polyominoes when \mathcal{P} has the S-property.

Discussion 3.9. Let \mathcal{P} be a simple thin polyomino that has the S-property. Let (I, J, \mathcal{P}^I) be a collapse datum of \mathcal{P} satisfying the conclusion of Lemma 3.5. Let *C* and *D* be the cells of *I*, with *C* being the single cell. Let *E* be the single cell of *J*. Let $r_{\mathcal{P},\widehat{C}}(t)$ and $r_{\mathcal{P},\widehat{I}}(t)$ be as defined in Discussion 3.6.

Write $Q = \mathcal{P} \setminus \{C\}$. Then $r_{\mathcal{P},\widehat{C}}(t) = r_Q(t)$. If \mathcal{P}^I is empty, define \mathcal{R} to be the polyomino $\mathcal{P} \setminus I$. Otherwise, i.e. if \mathcal{P}^I is a cell E, define \mathcal{R} to be the polyomino $\mathcal{P} \setminus \{C, E\}$. Then $r_{\mathcal{P},\widehat{I}}(t) = r_{\mathcal{R}}(t)$. Thus (3.7) becomes

$$(3.10) r_{\mathcal{P}}(t) = r_{\mathcal{Q}}(t) + r_{\mathcal{R}}(t)t.$$

Note that Q does not have the S-property, so we cannot use an inductive argument to prove Theorem 1.2 directly. Hence we need to rewrite $r_Q(t)$ in terms of smaller polyominoes. To this end, we observe that D is a single cell of the maximal inner interval J inside Q. Therefore, by (3.7),

(3.11)
$$r_Q(t) = r_{Q,\widehat{D}}(t) + r_{Q,\widehat{I}}(t)t.$$

We note that

(3.12)

 $r_{\mathcal{O}\,\widehat{\mathcal{D}}}(t) = r_{\mathcal{R}}(t).$

We now find an expression for $r_{Q,\hat{J}}(t)$. Let E, C_1, \ldots, C_k be the other cells of J in \mathcal{P} . E denotes the single cell of J in \mathcal{P} . For $1 \leq i \leq k - 1$, let B_i be the cell in \mathcal{P} such that $B_i \notin J$ and C_i is a neighbour cell of B_i . (See Discussion 3.3 and Figures 3, 4, 5, and 6 for notational conventions.) When E is the second end-cell or C_k is an end-cell with two neighbour cells, let B_k be the cell in \mathcal{P} such that $B_k \notin J$ and C_k is a neighbour cell of B_k . When C_k is an end-cell with three neighbour cells, let B_k and B_{k+1} be the cells in \mathcal{P} such that $B_k, B_{k+1} \notin J$ and C_k is a neighbour cell of B_k and B_{k+1} is a neighbour cell of B_k and B_{k+1} be the cells in \mathcal{P} such that $B_k, B_{k+1} \notin J$ and C_k is a neighbour cell of B_k and B_{k+1} . In the case when C_k has three neighbour cells, by Lemma 3.5, we may assume that B_{k+1} is both a single cell and an end-cell of the maximal inner interval containing it.

Now for all $1 \le i \le k - 1$, define

 $Q_i := \{A \in Q : \text{the path between } A \text{ and } B_i \text{ does not contain } C_i\}.$

Also, define

 $\widetilde{Q_k} := \{A \in Q : \text{the path between } A \text{ and } B_k \text{ does not contain } C_k\}.$

When *E* is the second end-cell or C_k is an end-cell with two neighbour cells, define $Q_k = \widetilde{Q_k}$. When C_k is an end-cell with three neighbour cells, let $\{a, b, a', b'\}$ be the vertices of C_k where $a, b \in V(B_k)$ and $a', b' \in V(B_{k+1})$. We define Q_k as the polyomino obtained from $\widetilde{Q_k} \cup \{B_{k+1}\}$ by the identification of the vertices *a* and *b* of $V(B_k)$ with the vertices *a'* and *b'* of $V(B_{k+1})$, respectively, by translating the cell B_{k+1} .

Lemma 3.13. With notation as above, we have the following:

- (1) Q_1, \ldots, Q_k are precisely the connected components of $Q \setminus J$.
- (2) For each $1 \le i \le k$, Q_i is a simple thin polyomino with the S-property.
- (3) $r_{Q,\hat{I}}(t) = \prod_{i=1}^{k} r_{Q_i}(t)$ and $\sum_{i=1}^{k} r(Q_i) = r(\mathcal{P}) 2$.

Proof. (1): Each Q_i is connected, $Q_i \cap Q_j = \emptyset$ for all $i \neq j$ (since *P* is simple) and $Q \setminus J = Q_1 \cup \cdots \cup Q_k$.

(2): Since Q is simple thin, so is Q_i . Let L be a maximal inner interval of Q_i . Then, either L is a maximal inner interval of \mathcal{P} or $L = L' \setminus \{C_i\}$, where L' is a maximal inner interval of \mathcal{P} . In both cases, L has a unique single cell. Hence Q_i has the S-property.

(3): By (1) $r_{Q,\hat{J}}(t)$ is the rook polynomial of $Q \setminus J$. Thus, by Proposition 3.1, $r_{Q,\hat{J}}(t) = \prod_{i=1}^{k} r_{Q_i}(t)$. For any *k*-rook configuration α of $Q \setminus J$, we note that $\alpha \cup \{C, E\}$ is a (k + 2)-rook configuration of \mathcal{P} . Hence $\sum_{i=1}^{k} r(Q_i) \leq r(\mathcal{P}) - 2$. On the other hand, let β be a $r(\mathcal{P})$ -rook configuration of \mathcal{P} . Since \mathcal{P} has the S-property, β is the only $r(\mathcal{P})$ -rook configuration of \mathcal{P} and β is the collection of all single cells of \mathcal{P} . Then, $\beta \setminus \{C, E\}$ is a rook configuration of $Q \setminus J$. Therefore $\sum_{i=1}^{k} r(Q_i) \geq r(\mathcal{P}) - 2$.

We are now ready to prove Theorem 1.2.

Proof of the Theorem. By Proposition 3.1, we may assume that \mathcal{P} is a simple thin polyomino with the S-property. Let $h_{\Bbbk[\mathcal{P}]}/(1-t)^{\dim(\Bbbk[\mathcal{P}])}$ be the Hilbert series of $\Bbbk[\mathcal{P}]$. By [RR21, Theorem 1.1], $h_{\Bbbk[\mathcal{P}]}(t) = r_{\mathcal{P}}(t)$. We proceed by induction on the rook number $r(\mathcal{P})$. If $r(\mathcal{P})$ is odd (in particular if $r(\mathcal{P}) = 1$), then $r_{\mathcal{P}}(-1) = 0$. Hence we may assume that $r(\mathcal{P})$ is even. Let (I, J, \mathcal{P}^I) be a collapse datum of \mathcal{P} satisfying the conclusion of Lemma 3.5. Apply Discussion 3.9, adopting its notation. Let Q and \mathcal{R} be as in Discussion 3.9. Then, by (3.10)

$$r_{\mathcal{P}}(t) = r_{\mathcal{Q}}(t) + r_{\mathcal{R}}(t)t.$$

Now apply Discussion 3.6 to the single cell D of the maximal inner interval J of Q. By (3.11), (3.12) and Lemma 3.13, we see that

$$r_{\mathcal{P}}(t) = (1+t)r_{\mathcal{R}}(t) + t\prod_{i=1}^{k} r_{\mathcal{Q}_i}(t).$$

By Lemma 3.13 and induction hypothesis, $\mathbb{k}[Q_i]$ is CD for all $1 \le i \le k$. If $r(Q_i)$ is odd for some *i*, then $r_{\mathcal{P}}(-1) = 0$. Therefore we may assume that $r(Q_i)$ is even for all *i*.

$$(-1)^{\lfloor \frac{r(\mathcal{P})}{2} \rfloor} r_{\mathcal{P}}(-1) = (-1)^{\frac{r(\mathcal{P})}{2}+1} \prod_{i=1}^{k} r_{Q_{i}}(-1)$$
$$= (-1)^{\frac{r(\mathcal{P})-2}{2}} \prod_{i=1}^{k} r_{Q_{i}}(-1)$$
$$= \prod_{i=1}^{k} (-1)^{\frac{r(Q_{i})}{2}} r_{Q_{i}}(-1) \qquad \text{by Lemma 3.13}$$
$$\ge 0 \qquad \qquad \text{by induction.}$$

This completes the proof of the theorem.

References

- [BH93] W. Bruns and J. Herzog. Cohen-Macaulay rings, volume 39 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1993.
- [Brä04] P. Brändén. Sign-graded posets, unimodality of *W*-polynomials and the Charney-Davis conjecture. *Electron. J. Combin.*, 11(2):Research Paper 9, 15, 2004/06.
- [CD95] R. Charney and M. Davis. The Euler characteristic of a nonpositively curved, piecewise Euclidean manifold. *Pacific J. Math.*, 171(1):117–137, 1995.
 - [1] D'Alì, A. and Venturello, L. (2021). Koszul Gorenstein algebras from Cohen-Macaulay simplicial complexes. arXiv:2106.05051 [math.AC].
- [QSS17] A. A. Qureshi, T. Shibuta, and A. Shikama. Simple polyominoes are prime. J. Commut. Algebra, 9(3):413-422, 2017.
- [Qur12] A. A. Qureshi. Ideals generated by 2-minors, collections of cells and stack polyominoes. J. Algebra, 357:279–303, 2012.
- [RR21] G. Rinaldo and F. Romeo. Hilbert series of simple thin polyominoes. J. Algebraic Combin., 54:607 624, 2021.
- [RW05] V. Reiner and V. Welker. On the Charney-Davis and Neggers-Stanley conjectures. J. Combin. Theory Ser. A, 109(2):247–280, 2005.
- [Sta97] R. P. Stanley. *Enumerative combinatorics. Vol. 1*, volume 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1997. With a foreword by Gian-Carlo Rota, Corrected reprint of the 1986 original.
- [Sta00] R. P. Stanley. Positivity problems and conjectures in algebraic combinatorics. In *Mathematics: frontiers and perspectives*, pages 295–319. Amer. Math. Soc., Providence, RI, 2000.

CHENNAI MATHEMATICAL INSTITUTE, SIRUSERI, TAMILNADU 603103. INDIA *Email address*: mkummini@cmi.ac.in

CHENNAI MATHEMATICAL INSTITUTE, SIRUSERI, TAMILNADU 603103. INDIA *Email address*: dharm@cmi.ac.in