

REPRESENTATIONS OF THE LIE SUPERALGEBRA OF SUPERDERIVATIONS OF THE GRASSMANN ALGEBRA AT INFINITY

LUCAS CALIXTO AND CRYSTAL HOYT

ABSTRACT. The Lie superalgebra $W(\infty)$ is defined to be the direct limit of the simple finite-dimensional Cartan type Lie superalgebras $W(n)$ as n goes to infinity, where $W(n)$ denotes the Lie superalgebra of superderivations of the Grassmann algebra $\Lambda(n)$. The zeroth component of $W(\infty)$ in its natural \mathbb{Z} -grading is isomorphic to $\mathfrak{gl}(\infty)$.

In this paper, we study \mathbb{Z} -graded modules over $W(\infty)$, and introduce a category \mathbb{T}_W of $W(\infty)$ -modules that is closely related to the Koszul category $\mathbb{T}_{\mathfrak{sl}(\infty)}$ of tensor $\mathfrak{sl}(\infty)$ -modules introduced and studied by Dan-Cohen, Serganova and Penkov. We classify the simple objects of \mathbb{T}_W (up to isomorphism). We prove that each simple module in \mathbb{T}_W is isomorphic to the unique simple quotient of a module induced from a simple module in $\mathbb{T}_{\mathfrak{gl}(\infty)}$, and vice versa, which is analogous to the case for $W(n)$ studied by Serganova. We realize each simple module from \mathbb{T}_W as a module of tensor fields, generalizing work of Bernstein and Leites for $W(n)$. We prove that the category \mathbb{T}_W has enough injective objects, and for each simple module, we provide an explicit injective module in \mathbb{T}_W that contains it.

INTRODUCTION

The study of Lie algebras that can be obtained as a direct limit of classical finite-dimensional Lie algebras has been an active field of research over the last 20 years [PH22]. These Lie algebras are called locally finite Lie algebras. The program of understanding the structure and representation theory of such Lie algebras has been led by Ivan Penkov. The classical locally simple Lie algebras $\mathfrak{sl}(\infty)$, $\mathfrak{o}(\infty)$, $\mathfrak{sp}(\infty)$ are precisely the Lie algebras that appear in Baranov's classification of finitary locally simple Lie algebras over \mathbb{C} [Ba99]. These Lie algebras admit several natural categories of modules which turn out to have very rich and useful structure theory [PS11, DPS16, HPS19, PS19, CoP19]. For example, the category $\mathbb{T}_{\mathfrak{sl}(\infty)}$ of tensor modules over $\mathfrak{sl}(\infty)$ was introduced and studied in [DPS16], where it was shown to be a self-dual Koszul category [DPS16]. The category $\mathbb{T}_{\mathfrak{sl}(\infty)}$ is used in [FPS16] to categorify the Boson-Fermion Correspondence. In [HPS19], a category of integrable $\mathfrak{sl}(\infty)$ -modules is used to describe the induced $\mathfrak{sl}(\infty)$ -module structure on the complexified reduced Grothendieck groups arising from Brundan's categorical $\mathfrak{sl}(\infty)$ -action on the category $\mathcal{F}_{m|n}^{\mathbb{Z}}$ of finite-dimensional integral $\mathfrak{gl}(m|n)$ -modules and on the BGG category $\mathcal{O}_{m|n}^{\mathbb{Z}}$ (see [Br03]).

It is natural to consider the super analogs of locally finite Lie algebras and their categories of modules. The categories of tensor modules over $\mathfrak{gl}(\infty|\infty)$ and $\mathfrak{osp}(\infty|\infty)$ are described in [S14], and the category of integrable bounded weight modules over classical locally finite Lie superalgebras is studied in [CP22]. Although the study of the representation theory of locally finite Lie superalgebras

2010 *Mathematics Subject Classification.* 17B65, 17B10, 17B55.

Key words and phrases. locally finite Lie algebras, algebras of derivations, integrable modules, weight modules.

is still in its infancy, it has already proven to be very useful. In [S14], Serganova found a natural way to explain the equivalence between the categories of tensor modules over $\mathfrak{o}(\infty)$ and $\mathfrak{sp}(\infty)$ by means of the category of tensor modules over $\mathfrak{osp}(\infty|\infty)$.

Moving to non-classical Lie superalgebras, it is safe to say that the Lie superalgebra $W(n)$ of superderivations of the Grassmann algebra $\Lambda(n)$ is the most fundamental one. This Lie superalgebra appears in Kac's classification list of simple finite-dimensional Lie superalgebras [Kac77]. Bernstein and Leites described irreducible finite-dimensional $W(n)$ -modules and realized these representations as modules of tensor fields in [BL81, BL83]. Serganova studied the category of \mathbb{Z} -graded $W(n)$ -modules and described the structure of induced modules in [S05].

In this paper, we define the Lie superalgebra $W(\infty)$ to be the direct limit of the Lie superalgebras $W(n)$ as n goes to infinity, with respect to the natural embedding of $W(n)$ into $W(n+1)$. So by definition, $W(\infty)$ is a locally finite Lie superalgebra that is locally simple, and hence simple. The Lie superalgebra $W(\infty)$ has a natural \mathbb{Z} -grading such that the zeroth graded component is isomorphic to $\mathfrak{gl}(\infty)$. The Lie algebra $\mathfrak{gl}(\infty)$ has a corank 1 simple ideal $\mathfrak{sl}(\infty)$.

Our main goal in the current paper is to introduce and study categories \mathbb{T}_W , \mathbb{T}_W^{\geq} and \mathbb{T}_W^{\leq} of $W(\infty)$ -modules that are analogs of the category $\mathbb{T}_{\mathfrak{sl}(\infty)}$. The objects in these categories are \mathbb{Z} -graded $W(\infty)$ -modules that satisfy some nice properties that resemble those that finite-dimensional modules over $W(n)$ have. We prove that the simple objects of \mathbb{T}_W and \mathbb{T}_W^{\leq} coincide and that there is a one-to-one correspondence between such simple modules and simple tensor modules of $\mathfrak{gl}(\infty)$ (see Theorem 5.9 and Corollary 5.13). To obtain this bijection we rely on the fact that every simple object in \mathbb{T}_W is parabolically induced from a simple object in $\mathbb{T}_{\mathfrak{gl}(\infty)}$ (Proposition 5.6). This is analogous to the finite-dimensional situation where every simple finite-dimensional $W(n)$ -module can be obtained as a quotient of a parabolically induced simple finite-dimensional $\mathfrak{gl}(n)$ -module [S05]. As a corollary, we obtain that simple modules in \mathbb{T}_W are highest weight modules with respect to an appropriately chosen Borel subalgebra and that such modules can be parameterized by pairs of partitions.

In Section 6 we define modules of tensor fields for $W(\infty)$ and we prove that every simple module in \mathbb{T}_W is isomorphic to a submodule of a module of tensor fields (Theorem 6.7). This result gives a realization of all simple modules of \mathbb{T}_W and it is analogous to [BL81, Theorem 3]. In Section 7 we construct a functor Γ from the category $\mathfrak{g}\text{-mod}$ of all \mathfrak{g} -modules to the category \mathbb{T}_W , which allows us to prove that the category \mathbb{T}_W has enough injectives (Proposition 7.1). We show that combining Γ with a certain coinduction functor provides a way to construct injective objects of \mathbb{T}_W from injective objects in the category of integrable $\mathfrak{gl}(\infty)$ -modules (Corollary 7.5). Finally, we use the functor Γ along with Theorem 6.7 to explicitly construct, for every simple module in \mathbb{T}_W , an injective module of \mathbb{T}_W which contains it (Theorem 7.6).

Acknowledgements. We would like to thank Ivan Penkov and Vera Serganova for helpful discussions. The first author was supported by Fapemig Grant (APQ-02768-21) and by CNPq Grant (402449/2021-5). The second author was partially supported by ISF Grant 1221/17 and by ISF Grant 1957/21.

1. THE LIE ALGEBRAS $\mathfrak{gl}(\infty)$ AND $\mathfrak{sl}(\infty)$

Notation: The ground field is always assumed to be \mathbb{C} . We set $I := \{1, 2, 3, \dots\}$. If C is a category, then we write $M \in C$ whenever M is an object of C . If V is a \mathbb{Z}_2 -graded vector space, then \bar{v} denotes the parity of a homogeneous element $v \in V$. By convention, if we write \bar{v} then we assume that v is homogeneous. For a Lie superalgebra \mathfrak{a} , let $\mathbf{U}(\mathfrak{a})$ denote its universal enveloping superalgebra.

1.1. Construction and root system of $\mathfrak{gl}(\infty)$ and $\mathfrak{sl}(\infty)$. We begin with defining the Lie algebras $\mathfrak{gl}(\infty)$ and $\mathfrak{sl}(\infty)$ (see e.g. [PH22]). Let V, V_* be countable-dimensional vector spaces over \mathbb{C} . Fix a non-degenerate pairing $\langle \cdot, \cdot \rangle : V \otimes V_* \rightarrow \mathbb{C}$ and bases $\{\xi_i\}_{i \in I}$ for V , $\{\partial_i\}_{i \in I}$ for V_* such that $\langle \xi_i, \partial_j \rangle = \delta_{i,j}$ for all $i, j \in I$. Then $\mathfrak{gl}(\infty) := V \otimes V_*$ has a Lie algebra structure such that

$$[\xi_i \otimes \partial_j, \xi_k \otimes \partial_\ell] = \langle \xi_k, \partial_j \rangle \xi_i \otimes \partial_\ell - \langle \xi_i, \partial_\ell \rangle \xi_k \otimes \partial_j.$$

Moreover, V and V_* are the natural and conatural $\mathfrak{gl}(\infty)$ -modules, respectively, and they satisfy

$$(\xi_i \otimes \partial_j) \cdot \xi = \langle \xi, \partial_j \rangle \xi_i, \quad (\xi_i \otimes \partial_j) \cdot \partial = -\langle \xi_i, \partial \rangle \partial_j, \quad \text{for all } \xi, \xi_i \in V, \partial, \partial_j \in V_*.$$

In what follows, we write $\xi_i \partial_j$ for the element $\xi_i \otimes \partial_j$.

We can identify $\mathfrak{gl}(\infty)$ with the space of infinite matrices $(a_{ij})_{i,j \in I}$ with finitely many nonzero entries, where the vector $\xi_i \partial_j$ corresponds to the matrix $E_{i,j}$ with 1 in the i, j -position and zeros elsewhere. Under this identification, $\langle \cdot, \cdot \rangle$ is the trace map on $\mathfrak{gl}(\infty)$.

The Lie algebra $\mathfrak{sl}(\infty)$ is defined to be the kernel of $\langle \cdot, \cdot \rangle$. One can also realize $\mathfrak{sl}(\infty)$ as a direct limit of finite-dimensional Lie algebras $\varinjlim \mathfrak{sl}(n)$. So the Lie algebra $\mathfrak{sl}(\infty)$ is locally simple, and hence simple. Note however that the exact sequence

$$0 \rightarrow \mathfrak{sl}(\infty) \rightarrow \mathfrak{gl}(\infty) \rightarrow \mathbb{C} \rightarrow 0$$

does not split, since the center of $\mathfrak{gl}(\infty)$ is trivial.

We realize $\mathfrak{gl}(\infty)$ as the direct limit $\varinjlim \mathfrak{gl}(n)$ as follows. As a vector space, $\mathfrak{gl}(n) = V_n \otimes V_n^*$, where $\{\xi_1, \dots, \xi_n\}$ and $\{\partial_1, \dots, \partial_n\}$ are fixed dual bases for V_n and V_n^* , respectively. Then we have obvious embeddings $V_n \hookrightarrow V_{n+1}$, $V_n^* \hookrightarrow V_{n+1}^*$, $\mathfrak{gl}(n) \hookrightarrow \mathfrak{gl}(n+1)$ given on basis vectors by $\xi_i \rightarrow \xi_i$ and $\partial_j \rightarrow \partial_j$. Then $\mathfrak{gl}(\infty) = \cup_{n \in \mathbb{Z}_{\geq 2}} \mathfrak{gl}(n)$, and we identify $\mathfrak{gl}(n)$ with its image under the defining map.

We fix the Cartan subalgebras $\mathfrak{h}_{\mathfrak{gl}(n)} \subseteq \mathfrak{gl}(n)$, $\mathfrak{h}_{\mathfrak{gl}} \subseteq \mathfrak{gl}(\infty)$ and $\mathfrak{h}_{\mathfrak{sl}} \subseteq \mathfrak{sl}(\infty)$, which consist of diagonal matrices. Then $\mathfrak{h}_{\mathfrak{gl}(n)} = \mathfrak{h}_{\mathfrak{gl}} \cap \mathfrak{gl}(n)$ and $\mathfrak{h}_{\mathfrak{sl}} = \mathfrak{h}_{\mathfrak{gl}} \cap \mathfrak{sl}(\infty)$. If we choose the standard basis $\{\varepsilon_1, \varepsilon_2, \varepsilon_3, \dots\}$ of \mathfrak{h}_* , then the root system of $\mathfrak{a} = \mathfrak{gl}(\infty)$, $\mathfrak{sl}(\infty)$ is $\Delta = \{\varepsilon_i - \varepsilon_j \mid i \neq j \in I\}$, where the elements $\xi_i \partial_j = E_{i,j}$ are root vectors for $\varepsilon_i - \varepsilon_j$.

We let $\mathfrak{b}(\prec)^0$ be the Borel subalgebra of $\mathfrak{a} = \mathfrak{gl}(\infty)$, $\mathfrak{sl}(\infty)$ corresponding to the following linear order \prec on our index set I :

$$1 \prec 3 \prec 5 \prec \dots \prec 6 \prec 4 \prec 2, \quad (1.1)$$

that is, $\mathfrak{b}(\prec)^0 = \mathfrak{h} \oplus \mathfrak{n}^0$ where $\mathfrak{n}^0 = \bigoplus_{\alpha \in \Delta^+} \mathfrak{a}_\alpha$ and

$$\Delta^+ := \{\varepsilon_i - \varepsilon_j \mid i < j \text{ odd}\} \cup \{\varepsilon_i - \varepsilon_j \mid j < i \text{ even}\} \cup \{\varepsilon_i - \varepsilon_j \mid i \text{ odd}, j \text{ even}\}. \quad (1.2)$$

We let $\mathfrak{b}(\prec)_n^0$ be the Borel subalgebra of $\mathfrak{gl}(n)$ defined by $\mathfrak{b}(\prec)^0 \cap \mathfrak{gl}(n)$, using our fixed embedding $\mathfrak{gl}(n) \hookrightarrow \mathfrak{gl}(\infty)$. With respect to the Borel subalgebra $\mathfrak{b}(\prec)^0$, both the natural module V and

the conatural module V_* are highest weight modules, and they have highest weights ε_1 and $-\varepsilon_2$, respectively.

1.2. Tensor modules over $\mathfrak{gl}(\infty)$ and $\mathfrak{sl}(\infty)$. Let $\mathfrak{a} = \mathfrak{gl}(\infty), \mathfrak{sl}(\infty)$. An \mathfrak{a} -module is called a *tensor module* if it is isomorphic to a subquotient of a finite direct sum of modules of the form $V^{\otimes p_i} \otimes V_*^{\otimes q_i}$ for some $p_i, q_i \in \mathbb{Z}_{\geq 0}$ [DPS16]. The module $V^{\otimes p} \otimes V_*^{\otimes q}$ is not semisimple if $p, q > 0$. The simple subquotients of the modules $V^{\otimes p} \otimes V_*^{\otimes q}$, $p, q \in \mathbb{Z}_{>0}$ can be parameterized by two Young diagrams λ, μ , and we will denote them $V_{\lambda, \mu}$ (see Theorem 1.1).

By Schur–Weyl duality, we have the following isomorphism of $\mathfrak{gl}(\infty) \times (S_p \times S_q)$ -modules:

$$V^{\otimes p} \otimes V_*^{\otimes q} \simeq \bigoplus_{|\lambda|=p} \bigoplus_{|\mu|=q} (\mathbb{S}_\lambda(V) \otimes \mathbb{S}_\mu(V_*)) \otimes (Y_\lambda \boxtimes Y_\mu),$$

where \mathbb{S}_λ denotes the Schur functor corresponding to the Young diagram λ , the size of λ is $|\lambda| := \sum \lambda_i$, and $Y_\lambda \boxtimes Y_\mu$ is the outer tensor product of irreducible S_p - and S_q -modules.

Then $\tilde{V}_{\lambda, \mu} := (\mathbb{S}_\lambda(V) \otimes \mathbb{S}_\mu(V_*))$ are indecomposable \mathfrak{a} -modules, and their socle filtration was described by Penkov and Styrkas in [PSt11]. We recall that the *socle* of a module M , denoted $\text{soc } M$, is the largest semisimple submodule of M , and that the *socle filtration* of M is defined inductively by $\text{soc}^0 M := \text{soc } M$ and $\text{soc}^i M := p_i^{-1}(\text{soc}(M/(\text{soc}^{i-1} M)))$, where $p_i : M \rightarrow M/(\text{soc}^{i-1} M)$ is the natural projection. The layers of the socle filtration are denoted by $\overline{\text{soc}}^i M := \text{soc}^i M / \text{soc}^{i-1} M$.

The following theorem is from [PSt11, Theorem 2.3].

Theorem 1.1 (Penkov, Styrkas). *Let $\mathfrak{a} = \mathfrak{gl}(\infty), \mathfrak{sl}(\infty)$. The socle filtration of $\tilde{V}_{\lambda, \mu}$ has the following layers:*

$$\overline{\text{soc}}^k \tilde{V}_{\lambda, \mu} = \bigoplus_{|\gamma|=k} N_{\gamma, \lambda'}^\lambda N_{\gamma, \mu'}^\mu V_{\lambda', \mu'}, \quad (1.3)$$

where $N_{\gamma, \lambda'}^\lambda, N_{\gamma, \mu'}^\mu$ denotes the Littlewood–Richardson coefficients determined by the relation $s_\lambda s_\mu = \sum_\nu N_{\lambda, \mu}^\nu s_\nu$ for the Schur symmetric polynomials s_λ, s_μ, s_ν .

In particular, the indecomposable \mathfrak{a} -module $\tilde{V}_{\lambda, \mu}$ has an irreducible socle, denoted $V_{\lambda, \mu}$. For any partitions λ, μ , the \mathfrak{a} -module $V_{\lambda, \mu}$ is an irreducible highest weight module with respect to the Borel subalgebra $\mathfrak{b}(\prec)^0$ and has highest weight

$$\chi := \sum_{i \in \mathbb{Z}_{>0}} \lambda_i \varepsilon_{2i-1} - \sum_{j \in \mathbb{Z}_{>0}} \mu_j \varepsilon_{2j} \quad (1.4)$$

(see [PSt11, Theorem 2.1]).

One corollary of Theorem 1.1 is that each simple subquotient of $V^{\otimes p} \otimes V_*^{\otimes q}$ is isomorphic to a (simple) submodule of $V^{\otimes p'} \otimes V_*^{\otimes q'}$ for some p', q' .

1.3. The categories $\mathbb{T}_{\mathfrak{gl}(\infty)}$ and $\mathbb{T}_{\mathfrak{sl}(\infty)}$. A category of tensor modules over $\mathfrak{sl}(\infty)$ was introduced and studied in [DPS16]. Here we recall this construction, and define a similar category for $\mathfrak{gl}(\infty)$. First, we need the following definitions.

Let $\mathfrak{a} = \mathfrak{gl}(\infty), \mathfrak{sl}(\infty)$. An \mathfrak{a} -module M is called *integrable* if for every $x \in \mathfrak{a}$ and $m \in M$, we have

$$\dim \text{Span}_{\mathbb{C}}\{x^i m \mid i \in \mathbb{Z}_{>0}\} < \infty.$$

For each $n \in \mathbb{Z}_{>0}$, let \mathfrak{t}'_n be the subalgebra of $\mathfrak{gl}(\infty)$ generated by root vectors as follows:

$$\mathfrak{t}'_n := \langle \xi_i \partial_j = E_{i,j} \mid j, i \geq n \rangle. \quad (1.5)$$

Definition 1.2. We say that a subalgebra $\mathfrak{k} \subset \mathfrak{a}$ has *finite corank* in \mathfrak{a} if $(\mathfrak{t}'_n \cap \mathfrak{a}) \subset \mathfrak{k}$ for some $n \gg 0$.

Remark 1.3. For $\mathfrak{a} = \mathfrak{sl}(\infty)$, a subalgebra $\mathfrak{k} \subset \mathfrak{a}$ has finite corank if and only if \mathfrak{k} contains the commutator subalgebra of the centralizer of some finite-dimensional subalgebra of \mathfrak{a} .

We say that an \mathfrak{a} -module M satisfies the *large annihilator condition (l.a.c)* if, for every $m \in M$, the subalgebra $\text{Ann}_{\mathfrak{a}}(m)$ has finite corank.

Definition 1.4. The category $\mathbb{T}_{\mathfrak{gl}(\infty)}$ is the full subcategory of $\mathfrak{gl}(\infty)$ -mod consisting of modules M that satisfy the following conditions:

- (1) M has finite length;
- (2) M is integrable;
- (3) M satisfies the l.a.c.

The category $\mathbb{T}_{\mathfrak{sl}(\infty)}$ is defined similarly, by replacing $\mathfrak{gl}(\infty)$ with $\mathfrak{sl}(\infty)$.

Then $\mathbb{T}_{\mathfrak{gl}(\infty)}$ and $\mathbb{T}_{\mathfrak{sl}(\infty)}$ are abelian symmetric monoidal categories. It was proved in [DPS16, Corollary 4.6] that the category $\mathbb{T}_{\mathfrak{sl}(\infty)}$ consists of tensor modules. Moreover, the modules $V^{\otimes p} \otimes V_*^{\otimes q}$, $p, q \in \mathbb{Z}_{\geq 0}$ are injective in the category $\mathbb{T}_{\mathfrak{sl}(\infty)}$, and every indecomposable injective object of $\mathbb{T}_{\mathfrak{sl}(\infty)}$ is isomorphic to an indecomposable direct summand of $V^{\otimes p} \otimes V_*^{\otimes q}$ for some $p, q \in \mathbb{Z}_{\geq 0}$ [DPS16]. So by Theorem 1.1, an indecomposable injective module in $\mathbb{T}_{\mathfrak{sl}(\infty)}$ is isomorphic to $\tilde{V}_{\lambda, \mu} := (\mathbb{S}_{\lambda}(V) \otimes \mathbb{S}_{\mu}(V_*))$ for some λ, μ . Moreover, by [DPS16, Theorem 3.4], every module M in $\mathbb{T}_{\mathfrak{sl}(\infty)}$ is an $\mathfrak{h}_{\mathfrak{sl}}$ -weight module (since $\mathfrak{h}_{\mathfrak{sl}}$ is a “splitting” Cartan subalgebra).

Lemma 1.5. *The restriction functor*

$$\mathbf{R}_{\mathfrak{sl}} : \mathbb{T}_{\mathfrak{gl}(\infty)} \rightarrow \mathbb{T}_{\mathfrak{sl}(\infty)}$$

defined by restricting a module $M \in \mathbb{T}_{\mathfrak{gl}(\infty)}$ to $\mathfrak{sl}(\infty)$ is well-defined.

Proof. Let $\mathbf{R}_{\mathfrak{sl}}(M) := M'$ be the restriction of M to $\mathfrak{sl}(\infty)$. Trivially, M' satisfies conditions (2) and (3) of Definition 1.4. To show that (1) holds, it suffices to prove that M being simple implies that M' is simple. For this, let $m \in M$ be a non-zero weight vector. Then we have

$$M = \mathbf{U}(\mathfrak{gl}(\infty))m = \mathbf{U}(\mathfrak{sl}(\infty))m.$$

Thus M' is simple, and the claim follows. \square

Lemma 1.6. *If $M \in \mathbb{T}_{\mathfrak{gl}(\infty)}$, then M is an $\mathfrak{h}_{\mathfrak{gl}}$ -weight module. Moreover, $\mu, \lambda \in \text{Supp } M$ are equal if and only if $\mu|_{\mathfrak{h}_{\mathfrak{sl}}} = \lambda|_{\mathfrak{h}_{\mathfrak{sl}}}$.*

Proof. Since $\mathbf{R}_{\mathfrak{sl}}(M) \in \mathbb{T}_{\mathfrak{sl}(\infty)}$, we have that $\mathbf{R}_{\mathfrak{sl}}(M)$ is an $\mathfrak{h}_{\mathfrak{sl}}$ -weight module. By the l.a.c., for any $\mathfrak{h}_{\mathfrak{sl}}$ -weight vector m , we can find $E_{n,n} \in \mathfrak{h}_{\mathfrak{gl}}$ such that $E_{n,n}m = 0$. Since $\mathfrak{h}_{\mathfrak{gl}} = \mathfrak{h}_{\mathfrak{sl}} \oplus \mathbb{C}E_{n,n}$, we get that m is also an $\mathfrak{h}_{\mathfrak{gl}}$ -weight vector. Since M and $\mathbf{R}_{\mathfrak{sl}}(M)$ have the same underlying vector space, this completes the proof.

Now let $\mu, \lambda \in \text{Supp } M$ such that $\mu|_{\mathfrak{h}_{\mathfrak{sl}}} = \lambda|_{\mathfrak{h}_{\mathfrak{sl}}}$. Again by the l.a.c., for nonzero $m \in M_\mu$, we can choose $n \gg 0$ so that $0 = E_{n,n}m = \mu(E_{n,n})m$, and similarly for λ . Thus, we can choose $n \gg 0$, such that $\mu(E_{n,n}) = \lambda(E_{n,n}) = 0$. Since $\mathfrak{h}_{\mathfrak{gl}} = \mathfrak{h}_{\mathfrak{sl}} \oplus \mathbb{C}E_{n,n}$ this implies $\mu = \lambda$. \square

Proposition 1.7. *The categories $\mathbb{T}_{\mathfrak{gl}(\infty)}$ and $\mathbb{T}_{\mathfrak{sl}(\infty)}$ are equivalent.*

Proof. Let $M' \in \mathbb{T}_{\mathfrak{sl}(\infty)}$. Then, by [DPS16, Corollary 4.6], M' is isomorphic to a submodule of a finite direct sum of copies of the tensor algebra $T(V \oplus V_*)$, and so M' inherits a natural $\mathfrak{gl}(\infty)$ -module structure. The l.a.c. ensures that there is a unique way to extend the $\mathfrak{sl}(\infty)$ -module structure of M' to a $\mathfrak{gl}(\infty)$ -module structure in such way that the resulting $\mathfrak{gl}(\infty)$ -module is an object of $\mathbb{T}_{\mathfrak{gl}(\infty)}$. We let $M \in \mathbb{T}_{\mathfrak{gl}(\infty)}$ denote M' viewed as a $\mathfrak{gl}(\infty)$ -module endowed with such structure, and define a functor $F : \mathbb{T}_{\mathfrak{sl}(\infty)} \rightarrow \mathbb{T}_{\mathfrak{gl}(\infty)}$ by setting $F(M') = M$, and $F(f) = f$ for all $f \in \text{Hom}_{\mathbb{T}_{\mathfrak{sl}(\infty)}}(M, N)$. Then the functors F and $R_{\mathfrak{sl}}$ are inverse to each other. \square

We denote by $\widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$ the Grothendieck envelope of the category $\mathbb{T}_{\mathfrak{gl}(\infty)}$, that is, $\widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$ is the full subcategory of $\mathfrak{gl}(\infty)\text{-mod}$ with objects being arbitrary sums of objects in $\mathbb{T}_{\mathfrak{gl}(\infty)}$ (see [ChP17]). We let $\widetilde{\mathbb{T}}_{\mathfrak{sl}(\infty)}$ denote the Grothendieck envelope of $\mathbb{T}_{\mathfrak{sl}(\infty)}$. It follows that $\widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$ and $\widetilde{\mathbb{T}}_{\mathfrak{sl}(\infty)}$ are equivalent. Note that $\mathbb{T}_{\mathfrak{gl}(\infty)}$ and $\widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$ have the same simple objects.

We have the following characterization of $\widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$.

Proposition 1.8. *The category $\widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$ is the full subcategory of $\mathfrak{gl}(\infty)\text{-mod}$ consisting of modules M that satisfy the following conditions:*

- (1) M is an $\mathfrak{h}_{\mathfrak{gl}}$ -weight module;
- (2) M is integrable;
- (3) M satisfies the l.a.c.

Proof. First, if $M \in \widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$ then M satisfies (1)-(3), since $M = \sum M_\gamma$ where each $M_\gamma \in \mathbb{T}_{\mathfrak{gl}(\infty)}$ satisfies these conditions.

Next, suppose M is a $\mathfrak{gl}(\infty)$ -module satisfying conditions (1)-(3). We claim $M = M|_{\mathfrak{sl}(\infty)}$ lies in the category $\mathcal{OL}\mathcal{A}_{\mathfrak{b}}$, as defined in [PS19, Section 3]. To show this we need to verify the following conditions: (i) M satisfies the l.a.c.; (ii) M is \mathfrak{h} -semisimple; (iii) every $x \in \mathfrak{n}^0$ acts locally nilpotently on M . We get (i) and (ii) from the definition, while condition (iii) follows from integrability. Indeed, if $x \in (\mathfrak{n}(\prec)^0)_\alpha$ and $m \in M_\mu$, then integrability implies that $x^n m = 0$ for $n \gg 0$, since $x^k m \in M_{\mu+k\alpha}$. The claim follows.

Now M can be written as $M = \sum_\gamma M_\gamma$, where each M_γ is a finitely generated $\mathfrak{gl}(\infty)$ -module that is semisimple over $\mathfrak{h}_{\mathfrak{gl}}$ (and hence, when restricted to $\mathfrak{sl}(\infty)$, each M_γ is a finitely generated module in $\mathcal{OL}\mathcal{A}_{\mathfrak{b}}$). Thus by [PS19, Proposition 4.17], each M_γ is a finite-length $\mathfrak{sl}(\infty)$ -module (and hence finite-length $\mathfrak{gl}(\infty)$ -module). Therefore, each M_γ lies in $\mathbb{T}_{\mathfrak{gl}(\infty)}$, and we conclude that $M \in \widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$, since $\widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$ is closed under arbitrary sums. \square

The claim and proof of Proposition 1.8 hold if we replace $\mathfrak{gl}(\infty)$ by $\mathfrak{sl}(\infty)$ everywhere.

2. THE LIE SUPERALGEBRA $W(\infty)$

In this section, we define $W(\infty)$ to be the direct limit $W(\infty) := \varinjlim W(n)$ of the finite-dimensional Cartan type Lie superalgebras $W(n)$, with respect to the obvious embedding. The Lie superalgebra $W(\infty)$ is a proper subalgebra of $\text{Der}(\Lambda(\infty))$, the Lie superalgebra of all superderivations of the (infinite) Grassmann algebra $\Lambda(\infty)$. We also describe the root system of $W(\infty)$ with respect to our choice of Cartan subalgebra.

2.1. The Grassmann algebra $\Lambda(\infty)$. The Grassmann algebra $\Lambda(n)$ is by definition the free commutative (unital) superalgebra with n odd generators ξ_1, \dots, ξ_n , such that $\xi_i^2 = 0$ for $i \in \{1, 2, \dots, n\}$. We define the (infinite) Grassmann algebra $\Lambda(\infty)$ to be the free commutative superalgebra with countably many odd generators $\{\xi_i\}_{i \in I}$ satisfying $\xi_i^2 = 0$. Clearly, $\Lambda(\infty) = \varinjlim \Lambda(n)$, where $\Lambda(n) \hookrightarrow \Lambda(n+1)$ in the obvious way.

The Grassmann algebra $\Lambda(\infty)$ has a natural \mathbb{Z} -grading $\Lambda(\infty) = \bigoplus_{k \geq 0} \Lambda(\infty)^k$, obtained by setting $\deg \xi_i = 1$ for each generator ξ_i , $i \in I$, which is compatible with the superalgebra grading. We let

$$\mathcal{E} := \{\underline{e} : I \rightarrow \{0, 1\} \mid \text{Supp } \underline{e} < \infty\}$$

and, for any $\underline{e} \in \mathcal{E}$, we set

$$\xi^{\underline{e}} = \xi_{i_1} \cdots \xi_{i_n} := \xi_{i_1} \wedge \cdots \wedge \xi_{i_n},$$

where $\text{Supp } \underline{e} = \{i_1, \dots, i_n\}$ and $i_1 < \cdots < i_n$. Then $\deg \xi^{\underline{e}} = |\text{Supp } \underline{e}| = \sum_{i \in I} e(i)$ and the set $\{\xi^{\underline{e}} \mid \underline{e} \in \mathcal{E}\}$ is a \mathbb{C} -basis for $\Lambda(\infty)$. So $\Lambda(\infty)^k = \Lambda^{k+1}(V) = \langle \xi_{i_1} \cdots \xi_{i_k} \mid i_1 < \cdots < i_k \rangle_{\mathbb{C}}$ and $\Lambda(\infty)^0 = \mathbb{C}$.

2.2. The Lie superalgebra $W(\infty)$. Now let $W(n)$ denote the Lie superalgebra $\text{Der}(\Lambda(n))$ of superderivations of $\Lambda(n)$. An element of $W(n)$ can be written as $\sum_{i=1}^n P_i \partial_i$, where $P_i \in \Lambda(n)$ and ∂_i , for $i = 1, \dots, n$, is the derivation defined by

$$\partial_i(\xi_j) = \delta_{ij}.$$

Setting $\deg \partial_i = -1$ yields a \mathbb{Z} -grading

$$W(n) = \bigoplus_{k=-1}^{n-1} W(n)^k,$$

with the property that $W(n)^0$ is isomorphic to $\mathfrak{gl}(n)$ and $W(n)^{-1}$ is isomorphic to the conatural module of $\mathfrak{gl}(n)$.

We define the Lie superalgebra $W(\infty)$ to be the direct limit $W(\infty) := \varinjlim W(n)$, where the embedding $W(n) \hookrightarrow W(n+1)$ is defined in the obvious way, by sending the generators $\xi_1, \dots, \xi_n, \partial_1, \dots, \partial_n$ of $W(n)$ to the generators of $W(n+1)$ that have the same name. Then $W(\infty)$ is an infinite-dimensional locally finite Lie superalgebra, that is locally simple, and hence simple.

Let $\text{Der}(\Lambda(\infty)) \subseteq \text{End}_{\mathbb{C}}(\Lambda(\infty))$ be the Lie superalgebra of all superderivations of $\Lambda(\infty)$. The Lie superalgebra $\text{Der}(\Lambda(\infty))$ is quite large, and $W(\infty)$ can be identified with the subspace of $\text{Der}(\Lambda(\infty))$ spanned by all elements of the form $P(\xi)\partial_i$ with $P(\xi) \in \Lambda(\infty)$, $i \in I$.

The underlying super vector space of $W(\infty)$ is $\Lambda(V) \otimes V_*$, and the set

$$\{\underline{\xi}^{\underline{e}}\partial_j \mid \underline{e} \in \mathcal{E}, i \in I\}$$

is a \mathbb{C} -basis for $W(\infty)$. The element $\underline{\xi}^{\underline{e}}\partial_i$ has degree $(\sum \underline{e}(k)) - 1$. The bracket of homogeneous elements is given by

$$[\underline{\xi}^{\underline{a}}\partial_j, \underline{\xi}^{\underline{b}}\partial_\ell] = \underline{\xi}^{\underline{a}}\partial_j \circ \underline{\xi}^{\underline{b}}\partial_\ell - (-1)^{\deg(\underline{\xi}^{\underline{a}}\partial_j) \deg(\underline{\xi}^{\underline{b}}\partial_\ell)} \underline{\xi}^{\underline{b}}\partial_\ell \circ \underline{\xi}^{\underline{a}}\partial_j$$

and is of the form

$$[\underline{\xi}^{\underline{a}}\partial_j, \underline{\xi}^{\underline{b}}\partial_\ell] = \pm \delta_{\underline{b}(j),1} \underline{\xi}^{\underline{a}}\underline{\xi}^{\underline{b}-\underline{e}_j}\partial_\ell \pm \delta_{\underline{a}(\ell),1} \underline{\xi}^{\underline{b}}\underline{\xi}^{\underline{a}-\underline{e}_\ell}\partial_j, \quad (2.1)$$

where $\underline{a}, \underline{b} \in \mathcal{E}$ and $j, \ell \in \mathbb{Z}_{>0}$ (see e.g. [Gav15]). The \mathbb{Z} -grading on $W(\infty)$ is

$$W(\infty) = \bigoplus_{k \geq -1} W(\infty)^k, \text{ where } W(\infty)^k = \text{span}\{P_j\partial_j \mid P_j \in \Lambda(\infty)^{k+1}, j \in I\},$$

and is compatible with the superalgebra grading, i.e., $W(\infty)_0 = \bigoplus_{k \geq 0} W(\infty)^{2k}$.

Now the zeroth component $W(\infty)^0$ is isomorphic to $\mathfrak{gl}(\infty)$, and we will identify $W(\infty)^0$ with $\mathfrak{gl}(\infty)$ by sending $\xi_i\partial_j$ to $E_{i,j}$. Then $\mathfrak{gl}(\infty)$ acts on each graded component $W(\infty)^k$ by restricting the adjoint action to $W(\infty)^0$. For each $k \geq -1$, we have a $\mathfrak{gl}(\infty)$ -module isomorphism

$$W(\infty)^k \cong \Lambda^{k+1}(V) \otimes V_*,$$

while for $k \leq -2$, we have $W(\infty)^k = \{0\}$.

We fix a Cartan subalgebra \mathfrak{h} of $W(\infty)$ to be the subspace spanned by $\xi_i\partial_i$, for $i \in I$. Then \mathfrak{h} is also a Cartan subalgebra of $W(\infty)^0$ and, under our identification with $\mathfrak{gl}(\infty)$, we have that $\mathfrak{h} = \mathfrak{h}_{\mathfrak{gl}}$ is the space of diagonal matrices. The root system of $W(\infty)$ is

$$\Delta = \{\varepsilon_{i_1} + \cdots + \varepsilon_{i_k} - \varepsilon_j \mid i_1 < \cdots < i_k, j \neq i, k \geq 0\} \cup \{\varepsilon_{i_1} + \cdots + \varepsilon_{i_k} \mid i_1 < \cdots < i_k, k \geq 1\},$$

where $\xi_{i_1} \cdots \xi_{i_k} \partial_j \in W(\infty)_{\varepsilon_{i_1} + \cdots + \varepsilon_{i_k} - \varepsilon_j}$.

3. CATEGORIES OF $W(\infty)$ -MODULES

From now on let $\mathfrak{g} = W(\infty)$, let $\mathfrak{g}^k = W(\infty)^k$, and let \mathfrak{g}_n be the image of $W(n)$ in \mathfrak{g} under the natural inclusion so that $\mathfrak{g} = \bigcup \mathfrak{g}_n$. Then, under our identification, we have $\mathfrak{g}^0 = \mathfrak{gl}(\infty)$, $\mathfrak{g}_n^0 = \mathfrak{gl}(n)$ and $\mathfrak{g}^0 = \bigcup \mathfrak{g}_n^0$.

In this section, we introduce three categories of \mathfrak{g} -modules: \mathbb{T}_W , \mathbb{T}_W^{\geq} and \mathbb{T}_W^{\leq} . First, we need some definitions. For each $n \in \mathbb{Z}_{>0}$, let \mathfrak{t}_n be the subalgebra of \mathfrak{g} generated by root vectors as follows:

$$\mathfrak{t}_n := \langle \partial_j, \xi_{i_1} \cdots \xi_{i_k} \partial_j \mid j, i_t \geq n \rangle.$$

Then $\mathfrak{t}_n \cap \mathfrak{g}^0 = \mathfrak{t}'_n$ (see (1.5)).

Definition 3.1. Let \mathfrak{s} be a subalgebra of \mathfrak{g} . We say that a subalgebra $\mathfrak{k} \subset \mathfrak{g}$ has *finite corank relative to \mathfrak{s}* if for some $n \gg 0$ we have $(\mathfrak{t}_n \cap \mathfrak{s}) \subset \mathfrak{k}$.

Definition 3.2. Let M be a \mathfrak{g} -module, and let \mathfrak{s} be a subalgebra of \mathfrak{g} . We say that M satisfies the *large annihilator condition (l.a.c)* for \mathfrak{s} if, for every $m \in M$, the subalgebra $\text{Ann}_{\mathfrak{g}}(m)$ has finite corank relative to \mathfrak{s} .

For a \mathfrak{g} -module M , let $M^{\mathfrak{t}_n} = \{m \in M \mid \mathfrak{t}_n \cdot m = 0\}$.

Lemma 3.3. *M satisfies the l.a.c. for \mathfrak{g} if and only if $M = \bigcup_{i>0} M^{t_n}$.*

Proof. Now $m \in M^{t_n} \Leftrightarrow t_n \subseteq \text{Ann}_{\mathfrak{g}}(m)$. Hence, for each $m \in M$, the subalgebra $\text{Ann}_{\mathfrak{g}}(m)$ has finite corank in \mathfrak{g} if and only if $m \in M^{t_n}$ for some $n > 0$. \square

We consider the following subalgebras of \mathfrak{g} :

$$\mathfrak{g}^{\geq} := \bigoplus_{k \geq 0} \mathfrak{g}^k, \quad \mathfrak{g}^> := \bigoplus_{k > 0} \mathfrak{g}^k, \quad \mathfrak{g}^{\leq} := \mathfrak{g}^{-1} \oplus \mathfrak{g}^0, \quad \mathfrak{g}^{<} := \mathfrak{g}^{-1}. \quad (3.1)$$

Definition 3.4. We define $\widetilde{\mathbb{T}}_W$ (respectively, $\widetilde{\mathbb{T}}_W^{\geq}$, $\widetilde{\mathbb{T}}_W^{\leq}$) to be the full subcategory of \mathfrak{g} -mod consisting of modules M that satisfy the following conditions:

- (1) M has a \mathbb{Z} -grading $M = \bigoplus_{k \in \mathbb{Z}} M^k$ that is compatible with the superalgebra grading;
- (2) M is an \mathfrak{h} -weight module;
- (3) M is integrable over \mathfrak{g}^0 ;
- (4) M satisfies the l.a.c. for \mathfrak{g} (respectively, for \mathfrak{g}^{\geq} , \mathfrak{g}^{\leq}).

Let \mathbb{T}_W (respectively, \mathbb{T}_W^{\geq} , \mathbb{T}_W^{\leq}) be the full subcategory of $\widetilde{\mathbb{T}}_W$ (respectively, of $\widetilde{\mathbb{T}}_W^{\geq}$, $\widetilde{\mathbb{T}}_W^{\leq}$) consisting of finite-length \mathfrak{g} -modules (with respect to the Jordan–Hölder series).

Then \mathbb{T}_W and $\widetilde{\mathbb{T}}_W$ are abelian symmetric monoidal categories.

Remark 3.5. Clearly, \mathbb{T}_W is a subcategory of \mathbb{T}_W^{\leq} and \mathbb{T}_W^{\geq} . Moreover, \mathbb{T}_W is precisely the full subcategory of \mathfrak{g} -mod whose objects are in both \mathbb{T}_W^{\leq} and \mathbb{T}_W^{\geq} . Indeed, if $M \in \mathbb{T}_W^{\leq}$ and $M \in \mathbb{T}_W^{\geq}$, then for each $m \in M$ we can pick $n \gg 0$ so that both $t_n \cap \mathfrak{g}^{\geq}$ and $t_n \cap \mathfrak{g}^{\leq}$ are contained in $\text{Ann}_{\mathfrak{g}}(m)$, which implies that $t_n \subseteq \text{Ann}_{\mathfrak{g}}(m)$.

Example 3.6. Here are a few examples of modules in \mathbb{T}_W : the trivial module \mathbb{C} , the *natural module* $\Lambda(V)$, the simple natural module $\Lambda(V)_+ := \Lambda(V)/\mathbb{C}$, and the adjoint module \mathfrak{g} . It is clear that conditions (1) and (2) hold for these modules, while (3) follows from the next easy lemma. Condition (4) can be verified by direct computation. \circ

Lemma 3.7. *Suppose $M = \bigcup M_n$ is a \mathfrak{g} -module such that each M_n is a finite-dimensional \mathfrak{g}_n -module. Then M is integrable over \mathfrak{g}^0 .*

Proof. Let $m \in M$ and $x \in \mathfrak{g}^0$. Choose $N \gg 0$ such that $m \in M_n$ and $x \in \mathfrak{g}_n^0$. Then $\text{Span}_{\mathbb{C}}\{x^i m \mid i \in \mathbb{Z}_{>0}\} \subseteq M_n$, which is finite dimensional. \square

Remark 3.8. Note that integrability of a \mathfrak{g} -module over the even part \mathfrak{g}_0 implies integrability over \mathfrak{g} , since for any odd element $x \in \mathfrak{g}_1$, we have $x^2 = (1/2)[x, x] \in \mathfrak{g}_0$. However, integrability over the zeroth component \mathfrak{g}^0 does not imply integrability over \mathfrak{g} . Consider, for example, the induced module $K^-(X) = \text{Ind}_{\mathfrak{g}^{\leq}}^{\mathfrak{g}}(X)$ where $X \in \mathbb{T}_{\mathfrak{gl}(\infty)}$ and $\mathfrak{g}^{<}$ acts trivially on X . Then $K^-(X)$ is integrable over \mathfrak{g}^0 but not over \mathfrak{g} , since $\mathbf{U}(\mathfrak{g}^>)$ acts freely on it.

Lemma 3.9. *If $M \in \widetilde{\mathbb{T}}_W$, $\widetilde{\mathbb{T}}_W^{\geq}$, $\widetilde{\mathbb{T}}_W^{\leq}$, then $M|_{\mathfrak{g}^0} \in \widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$.*

Proof. We use the description of $\widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$ given in Proposition 1.8. By definition, M is integrable over \mathfrak{g}^0 , and M is a weight module for $\mathfrak{h} = \mathfrak{h}_{\mathfrak{gl}}$. So we only need to check l.a.c. for \mathfrak{g}^0 . Since $\mathfrak{g}^0 \subseteq \mathfrak{g}^{\geq} \cap \mathfrak{g}^{\leq}$, we have for any $m \in M$, that there exists $n \gg 0$ for which $t_n \cap \mathfrak{g}^0 \subseteq \text{Ann}(m)$. \square

Remark 3.10. If we replace condition (2) in Definition 3.4 with the requirement that each graded component M^k has finite length as a \mathfrak{g}^0 -module, then we get $(M^k)|_{\mathfrak{g}^0} \in \mathbb{T}_{\mathfrak{gl}(\infty)}$. This implies M^k is an \mathfrak{h} -weight module and $M|_{\mathfrak{g}^0}$ is in $\widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$. Hence, M is an \mathfrak{h} -weight module.

4. INDUCED MODULES

In this section, we study $W(\infty)$ -modules which are induced from $\mathfrak{gl}(\infty)$ -modules.

Definition 4.1. Let X be a module over $\mathfrak{g}^0 = \mathfrak{gl}(\infty)$, and let $\mathfrak{g}^>$ (respectively, $\mathfrak{g}^<$) act trivially on it. We define the induced \mathfrak{g} -modules

$$\begin{aligned} K^+(X) &:= \text{Ind}_{\mathfrak{g}^{\geq}}^{\mathfrak{g}}(X) = \mathbf{U}(\mathfrak{g}) \otimes_{\mathbf{U}(\mathfrak{g}^{\geq})} X, \\ K^-(X) &:= \text{Ind}_{\mathfrak{g}^{\leq}}^{\mathfrak{g}}(X) = \mathbf{U}(\mathfrak{g}) \otimes_{\mathbf{U}(\mathfrak{g}^{\leq})} X. \end{aligned}$$

Remark 4.2. It follows from the PBW Theorem that $\mathbf{U}(\mathfrak{g})$ is a free right $\mathbf{U}(\mathfrak{g}^{\geq})$ -module. Thus $K^+(-)$ is an exact functor. The same holds for $K^-(-)$.

It is standard to show that $K^{\pm}(X)$ admits a unique simple quotient if and only if X is a simple \mathfrak{g}^0 -module. Such a simple quotient will be denoted by $L^{\pm}(X)$.

We can define \mathfrak{g}_n^{\geq} , $\mathfrak{g}_n^>$, \mathfrak{g}_n^{\leq} and $\mathfrak{g}_n^<$ analogously to (3.1). Let X_n be a \mathfrak{g}_n^0 -module, and extend the action to $\mathfrak{g}_n^>$ (respectively, $\mathfrak{g}_n^<$) trivially. Similarly, we define the induced \mathfrak{g}_n -modules

$$K_n^+(X_n) := \text{Ind}_{\mathfrak{g}_n^{\geq}}^{\mathfrak{g}_n}(X_n), \quad K_n^-(X_n) := \text{Ind}_{\mathfrak{g}_n^{\leq}}^{\mathfrak{g}_n}(X_n),$$

and, for simple X_n , their simple quotients are denoted by $L_n^{\pm}(X_n)$.

We have the following result.

Proposition 4.3. *For each $n \geq 2$, let $f_n : X_n \hookrightarrow X_{n+1}$ be an embedding of \mathfrak{g}_n^0 -modules, and consider the natural embedding $f_n : K^{\pm}(X_n) \hookrightarrow K^{\pm}(X_{n+1})$ of \mathfrak{g}_n -modules, where $f_n(uv) = uf_n(v)$ for every $u \in \mathbf{U}(\mathfrak{g}_n)$ and $v \in X_n$.*

Then we have an isomorphism of \mathfrak{g} -modules

$$K^{\pm}(\varinjlim_n X_n) \cong \varinjlim_n K_n^{\pm}(X_n),$$

where the limits are taken over the family $\{f_n\}$.

Proof. Let $\varphi_{i,j} : X_i \hookrightarrow X_j$ be the family of embeddings defining $X := \varinjlim X_i$, and let $\varphi_i : X_i \hookrightarrow X$ such that $\varphi_i = \varphi_j \circ \varphi_{i,j}$ for all $i \leq j$. By definition, $\varinjlim K_i^{\pm}(X_i)$ is defined through the embeddings $\tilde{\varphi}_{i,j} := \iota_{i,j} \otimes \varphi_{i,j}$, where $\iota_{i,j} : \mathbf{U}(\mathfrak{g}_i) \hookrightarrow \mathbf{U}(\mathfrak{g}_j)$ are the canonical inclusions. Notice also that $\tilde{\varphi}_i = \iota_i \otimes \varphi_i$. Define

$$\phi_i : \mathbf{U}(\mathfrak{g}_i) \otimes X_i \hookrightarrow \mathbf{U}(\mathfrak{g}) \otimes X$$

by $\phi_i(u \otimes v) = u \otimes [v]$, where $[v]$ is a representative of the class of v in X . Notice that

$$\phi_j \circ \tilde{\varphi}_{i,j}(u \otimes v) = \phi_j(u \otimes \varphi_{i,j}(v)) = u \otimes [\varphi_{i,j}(v)] = u \otimes [v] = \phi_j(u \otimes v).$$

Thus, by the universal property of $\varinjlim K_i^{\pm}(X_i)$, there exists a unique homomorphism of \mathfrak{g} -modules

$$\sigma : \varinjlim K_i^{\pm}(X_i) \rightarrow K^{\pm}(\varinjlim X_i)$$

such that $\sigma \circ \tilde{\varphi}_i = \phi_i$ for every i . Then, for any $u \otimes [v] \in K^\pm(\varinjlim X_i)$, there is $i \gg 0$ for which $u \otimes [v] = \phi_i(u \otimes v) = \sigma \circ \tilde{\varphi}_i(u \otimes v)$. Hence σ is surjective.

On the other hand, let $w \in \ker \sigma$. Then we can write $w = \tilde{\varphi}_i(\sum u_i \otimes v_i)$ for linearly independent elements $u_i \in \mathbf{U}(\mathfrak{g}_i)$ for $i \gg 0$. In particular,

$$0 = \sigma(w) = \sigma \circ \tilde{\varphi}_i(\sum u_i \otimes v_i) = \phi_i(\sum u_i \otimes v_i) = \sum u_i \otimes [v_i],$$

and since the u_i are linearly independent, this implies $[v_i] = 0$ for every i . But φ_i is an embedding for every i , which implies $v_i = 0$, and hence $w = 0$. Therefore σ is an isomorphism. \square

We define a \mathbb{Z} -grading on $X \in \mathbb{T}_{\mathfrak{gl}(\infty)}$ as follows. Now X has a weight decomposition $X = \bigoplus_{\mu \in \mathfrak{h}^*} X_\mu$ and

$$\text{Supp } X \subset \left\{ \mu = \sum \mu_i \varepsilon_i \mid \mu_i \in \mathbb{Z} \text{ for all } i \in I, \text{ and } \mu_i = 0 \text{ for all but finitely many } i \right\}.$$

Indeed, this follows from the fact that the weights of the modules $V^{\otimes m} \otimes V_*^{\otimes n}$ are all of this form. For any such X , we define a \mathbb{Z} -grading $X = \bigoplus_{k \in \mathbb{Z}} X^k$ by letting

$$X^k := \bigoplus_{\mu \in \mathfrak{h}^*, |\mu| = k} X_\mu.$$

Notice that each X^k is invariant under the action of \mathfrak{g}^0 . In particular, if X simple, then $X = X^k$ for some $k \in \mathbb{Z}$, and in this case, we will write $|X| := k$.

Proposition 4.4. *Let $X \in \mathbb{T}_{\mathfrak{gl}(\infty)}$ such that $|X| = k$. The module $K^\pm(X)$ considered as a \mathfrak{g}^0 -module admits a decomposition*

$$K^+(X) = \bigoplus_{j \leq 0} T_+^j, \quad K^-(X) = \bigoplus_{j \geq 0} T_-^j \quad (4.1)$$

such that each T_\pm^j is in $\mathbb{T}_{\mathfrak{gl}(\infty)}$ and each simple subquotient Y of T_\pm^j satisfies $|Y| = k + j$. The decomposition in (4.1) is a \mathbb{Z} -grading of the \mathfrak{g} -module $K^\pm(X)$.

Proof. Let us first prove the claim for $K^-(X)$. Since $\mathfrak{g}^> = \bigoplus_{i > 0} \mathfrak{g}^i$ is graded, we can decompose the symmetric \mathfrak{g}^0 -module $S(\mathfrak{g}^>)$ as $S(\mathfrak{g}^>) = \bigoplus_{j > 0} S^j$, where S^j is the projection of $\bigoplus_{i_1 + \dots + i_n = j} \mathfrak{g}^{i_1} \otimes \dots \otimes \mathfrak{g}^{i_n}$ onto $S(\mathfrak{g}^>)$. So we have an isomorphism of \mathfrak{g}^0 -modules

$$K^-(X) \cong S(\mathfrak{g}^>) \otimes X \cong \bigoplus_{j \geq 0} (S^j \otimes X).$$

For each $j \geq 0$, set $T_-^j := S^j \otimes X$, and observe that T_-^j is in $\mathbb{T}_{\mathfrak{gl}(\infty)}$ since $S^j \in \mathbb{T}_{\mathfrak{gl}(\infty)}$. Notice now that any $\mu \in \text{Supp } S^j$ satisfies $|\mu| = \sum \mu_i = j$. Thus, for $N \gg 0$, the element $I_N := \sum_{i=1}^N \xi_i \partial_i$ acts on any weight vector of T_-^j (and hence on any weight vector of a subquotient of T_-^j) via multiplication by $k + j$.

Now for $K^+(X)$, we note that we have isomorphisms of \mathfrak{g}^0 -modules

$$K^+(X) \cong \Lambda(\mathfrak{g}^<) \otimes X \cong \bigoplus_{j \geq 0} \Lambda^j(V_*) \otimes X,$$

where each component $T_+^{-j} := \Lambda^j(V_*) \otimes X$ is in $\mathbb{T}_{\mathfrak{gl}(\infty)}$, and hence has finite length over \mathfrak{g}^0 . Moreover, for $N \gg 0$, the element I_N acts on any weight vector of $(\Lambda^j(V_*) \otimes X)$ via multiplication by $k - j$, and the claim follows for $K^+(X)$.

Now it is clear that $\mathfrak{g}^i T_{\pm}^j \subset T_{\pm}^{i+j}$, so the \mathbb{Z} -grading is compatible with the action of \mathfrak{g} . \square

Corollary 4.5. *If $X \in \mathbb{T}_{\mathfrak{gl}(\infty)}$ is simple, then the module $K := K^{\pm}(X)$ admits a \mathfrak{g}^0 -module filtration*

$$K = K_0 \supset K_1 \supset K_2 \supset K_3 \cdots,$$

where $\bigcap_i K_i = 0$ and K_{i-1}/K_i is simple for all $i > 0$. Moreover, the multiplicity of any simple subquotient of K is finite and does not depend on the choice of a filtration.

Proof. For each $j \geq 0$, let $T^j = F_j^0 \supset F_j^1 \supset \cdots \supset F_j^{n_j}$ be the Jordan–Hölder series of T^j provided by Proposition 4.4. Now, observe that

$$\begin{aligned} K_1 := F_1^1 \oplus \bigoplus_{j>1} T^j \supset K_2 := F_1^2 \oplus \bigoplus_{j>1} T^j \supset \cdots \supset K_{n_1} := F_1^{n_1} \oplus \bigoplus_{j>1} T^j \supset \\ \supset K_{n_1+1} := \bigoplus_{j>1} T_j \supset K_{n_1+2} := F_2^1 \oplus \bigoplus_{j>2} T_j \supset \cdots \end{aligned}$$

yields the required filtration. \square

Proposition 4.6. *If $X \in \mathbb{T}_{\mathfrak{gl}(\infty)}$, then any simple \mathfrak{g}^0 -module in the Jordan–Hölder series of the induced module $K^{\pm}(X)$, considered as a \mathfrak{g}^0 -module, has finite multiplicity.*

Proof. If X is simple, then any simple \mathfrak{g}^0 -subquotient M of $K^{\pm}(X)$ appears in a unique component T_{\pm}^j (see the proof of Proposition 4.4 for the definition of T_{\pm}^j) which is determined by $|M|$. Hence, M must have finite multiplicity since T_{\pm}^j has finite length.

Now we use induction on the length of X in its Jordan–Hölder series. Let Y be a maximal submodule of X and consider the short exact sequence

$$0 \rightarrow Y \rightarrow X \rightarrow X/Y \rightarrow 0$$

Applying the exact functor $K^{\pm}(-)$ on this sequence yields

$$0 \rightarrow K^{\pm}(Y) \rightarrow K^{\pm}(X) \rightarrow K^{\pm}(X/Y) \rightarrow 0.$$

By induction, a simple \mathfrak{g}^0 -subquotient M has finite multiplicity in $K^{\pm}(Y)$ and in $K^{\pm}(X/Y)$. Since the multiplicity of M is additive, the result follows. \square

Theorem 4.7. *If $X \in \mathbb{T}_{\mathfrak{gl}(\infty)}$, then $K^+(X)$ lies in $\widetilde{\mathbb{T}}_W^{\geq}$ and $K^-(X)$ lies in $\widetilde{\mathbb{T}}_W^{\leq}$.*

Proof. Write $X = \bigoplus_{k \in \mathbb{Z}} X^k$ where $|X^k| = k$. Then, it follows from Propositions 4.4 applied to each X^k that $K^{\pm}(X)$ satisfies (1)-(3) of Definition 3.4. So we just need to show that $K^+(X)$ satisfies l.a.c. for \mathfrak{g}^{\geq} and that $K^-(X)$ satisfies l.a.c. for \mathfrak{g}^{\leq} .

Let \mathfrak{k} be a subalgebra of \mathfrak{g} , and let $\mathbf{U}(\mathfrak{g})_{\text{ad } \mathfrak{k}}$ denote $\mathbf{U}(\mathfrak{g})$ viewed as a \mathfrak{k} -module via the adjoint action. Notice that \mathfrak{g} viewed as a \mathfrak{g} -module (via the adjoint action) satisfies l.a.c. for \mathfrak{g} (and hence for \mathfrak{g}^{\geq}). Thus $\mathbf{U}(\mathfrak{g})_{\text{ad } \mathfrak{g}^{\geq}}$, X (viewed as a \mathfrak{g}^{\geq} -module with $\mathfrak{g}^> \cdot X = 0$) and $\mathbf{U}(\mathfrak{g})_{\text{ad } \mathfrak{g}^{\geq}} \otimes_{\mathbb{C}} X$ also satisfy the l.a.c. for \mathfrak{g}^{\geq} . Consider the surjective map of vector spaces $\mathbf{U}(\mathfrak{g})_{\text{ad } \mathfrak{g}^{\geq}} \otimes_{\mathbb{C}} X \xrightarrow{\phi} K^+(X)$ given by $\phi(u \otimes x) = u \otimes x$ for all $x \in X$, $u \in \mathbf{U}(\mathfrak{g})$. This is a \mathfrak{g}^{\geq} -module homomorphism since for all $g \in \mathfrak{g}^{\geq}$ and $x \in X$, we have

$$\phi(g(u \otimes x)) = \phi([x, u] \otimes x + (-1)^{\bar{u}\bar{g}} u \otimes gx) = [x, u] \otimes x + (-1)^{\bar{u}\bar{g}} u \otimes gx$$

and $g\phi(u \otimes x) = gu \otimes x = [g, u] \otimes x + (-1)^{\bar{u}g} u \otimes gx$. Thus the result follows for $K^+(X)$. The proof for $K^-(X)$ is similar. \square

Remark 4.8. The induced \mathfrak{g} -module $K^+(X)$ is not in \mathbb{T}_W^{\leq} or \mathbb{T}_W , since it does not satisfy the l.a.c. for \mathfrak{g}^{\leq} . Similarly, $K^-(X)$ is not in \mathbb{T}_W^{\geq} or \mathbb{T}_W , since it does not satisfy the l.a.c. for \mathfrak{g}^{\geq} .

5. SIMPLE MODULES IN \mathbb{T}_W

In this section we prove that the simple modules of \mathbb{T}_W can be parameterized by two partitions, which we denote by $L_{\lambda, \mu}^-$, and that they are highest weight modules with respect to the Borel subalgebra $\mathfrak{b}(\prec)^{\min}$.

Given a Borel subalgebra \mathfrak{b}^0 of $\mathfrak{g}^0 = \mathfrak{gl}(\infty)$, we define Borel subalgebras for \mathfrak{g} by

$$\mathfrak{b}^{\max} := \mathfrak{b}^0 \oplus \mathfrak{g}^{\gt}, \quad \mathfrak{b}^{\min} := \mathfrak{b}^0 \oplus \mathfrak{g}^{\lt}.$$

Similarly, for a Borel subalgebra \mathfrak{b}_n^0 of $\mathfrak{g}_n^0 = \mathfrak{gl}(n)$, we define Borel subalgebras of \mathfrak{g}_n by $\mathfrak{b}_n^{\max} := \mathfrak{b}_n^0 \oplus \mathfrak{g}_n^{\gt}$ and $\mathfrak{b}_n^{\min} := \mathfrak{b}_n^0 \oplus \mathfrak{g}_n^{\lt}$.

For any weight $\gamma \in \mathfrak{h}^*$, we denote by $V_{\mathfrak{b}^0}(\gamma)$ the simple \mathfrak{b}^0 -highest weight \mathfrak{g}^0 -module with highest weight $\gamma \in \mathfrak{h}^*$. Similarly, $V_{\mathfrak{b}_n^0}(\gamma)$ denotes the simple \mathfrak{b}_n^0 -highest weight \mathfrak{g}_n^0 -module with highest weight $\gamma \in \mathfrak{h}_n^*$, where $\mathfrak{h}_n := \mathfrak{h} \cap \mathfrak{g}_n^0$.

For a Borel subalgebra $\mathfrak{b} \in \{\mathfrak{b}^{\max}, \mathfrak{b}^{\min}\}$ of \mathfrak{g} , let \mathbb{C}_γ denote the one-dimensional representation of \mathfrak{b} defined by $\gamma \in \mathfrak{h}^*$. The induced module $M_{\mathfrak{b}}(\gamma) := U(\mathfrak{g}) \otimes_{U(\mathfrak{b})} \mathbb{C}_\gamma$ is a \mathfrak{b} -highest weight \mathfrak{g} -module. We let $L_{\mathfrak{b}}(\gamma)$ denote the unique simple quotient of $M_{\mathfrak{b}}(\gamma)$.

Recall the Borel subalgebra $\mathfrak{b}(\prec)^0$ of \mathfrak{g}^0 corresponding to the linear order \prec on I given in (1.1). We let $\mathfrak{b}(\prec)^0 := \mathfrak{h} \oplus \mathfrak{n}(\prec)$ denote the standard Borel corresponding to the natural order on I . In what follows we set

$$\mathfrak{b}(\prec)_n^0 := \mathfrak{b}(\prec)^0 \cap \mathfrak{gl}(n), \quad \text{and} \quad \mathfrak{b}(\prec)_n^0 := \mathfrak{b}(\prec)^0 \cap \mathfrak{gl}(n).$$

Remark 5.1. Serganova studied induced $W(n)$ -modules in [S05], where she proved that the \mathfrak{g}_n -module $K_n^+(V_{\mathfrak{b}(\prec)_n^0}(\nu))$ is simple if and only if the weight ν is typical [S05, Theorem 6.3], and has length 2 when ν is atypical [S05, Corollary 7.6]. By Theorem 5.3 in [S05], the atypical weights for the standard Borel subalgebra $\mathfrak{b}(\prec)_n^{\max}$ are of the form $a\varepsilon_i + \varepsilon_{i+1} + \cdots + \varepsilon_n$, for some $a \in \mathbb{C}$.

Recall that $V_{\lambda, \mu}$ denotes the simple module in $\mathbb{T}_{\mathfrak{gl}(\infty)}$ corresponding to partitions λ, μ . By our convention, (0) represents the empty partition \emptyset .

Theorem 5.2. *The induced module $K^+(V_{\lambda, \mu})$ is locally simple (and hence simple) if and only if $(\lambda, \mu) \neq (\emptyset, (\mu_1))$, that is, if and only if $V_{\lambda, \mu} \not\cong S^k(V_*)$ for some $k \in \mathbb{Z}_{\geq 0}$.*

Proof. Recall that $\mathfrak{b}(\prec)^{\max} := \mathfrak{b}(\prec)^0 \oplus \mathfrak{g}^{\gt}$ and note that any simple $X \in \mathbb{T}_{\mathfrak{gl}(\infty)}$ is such that $X \cong V_{\lambda, \mu}$ for some partitions λ, μ [PSt11, DPS16]. In particular, $X \cong \varinjlim X_n$, where for $n \gg 0$, the \mathfrak{g}_n^0 -module X_n is the simple $\mathfrak{b}(\prec)_n^0$ -highest weight module

$$V_{\mathfrak{b}(\prec)_n^0} \left(\sum_{i \in \mathbb{Z}_{>0}} \lambda_i \varepsilon_{2i-1} - \sum_{i \in \mathbb{Z}_{>0}} \mu_i \varepsilon_{2i} \right).$$

In particular, using the Weyl group of \mathfrak{g}_n^0 , we conclude that

$$X_n \cong V_{\mathfrak{b}(\prec)_n^0}(\lambda_1 \varepsilon_1 + \cdots + \lambda_k \varepsilon_k - \mu_l \varepsilon_{n-l} - \cdots - \mu_1 \varepsilon_n).$$

It follows from Remark 5.1 that $K_n^+(X_n)$ is simple for $n > k + l$ if and only if its highest weight with respect to $\mathfrak{b}(\prec)_n^{\max}$ is not equal to $-\mu_1 \varepsilon_n$, that is, if $(\lambda, \mu) \neq (\emptyset, (\mu_1))$. Since $K^+(V_{\lambda, \mu}) \cong \varinjlim K_n^+(X_n)$, we conclude that $K^+(V_{\lambda, \mu})$ is locally simple when $(\lambda, \mu) \neq (\emptyset, (\mu_1))$. For the second claim, note that $V_{\emptyset, (k)} = S^k(V_*)$.

If $(\lambda, \mu) = (\emptyset, (k))$, then $V_{\emptyset, (k)} = S^k(V_*)$. The vector $1 \otimes \partial_2^k$ is the $\mathfrak{b}(\prec)^{\max}$ -highest weight vector of $K^+(S^k(V_*))$. We claim that the vector $\partial_2 \otimes \partial_2^k \in K^+(S^k(V_*))$ is $\mathfrak{b}(\prec)^{\max}$ -primitive and generates a proper submodule. Indeed, for any homogeneous element $x := \xi^{\underline{a}} \partial_j \in \mathfrak{g}^{\geq}$ we have

$$x(\partial_2 \otimes \partial_2^k) = [x, \partial_2] \otimes \partial_2^k + (-1)^{\bar{x}} \partial_2 \otimes x \partial_2^k,$$

and since ∂_2^k is $\mathfrak{b}(\prec)^{\max}$ -primitive, we just need to show that $[x, \partial_2] \otimes \partial_2^k$ is zero when $x \in \mathfrak{n}(\prec)^0$ and $x \in \mathfrak{g}^1$. By (2.1), $[\xi^{\underline{a}} \partial_j, \partial_2]$ is zero unless $\underline{a}(2) = 1$, and hence equals zero when $x \in \mathfrak{n}(\prec)^0$. Now consider $x_k := \xi^k \xi^2 \partial_k \in \mathfrak{g}_{\varepsilon_2}$, when $k \in \mathbb{Z}_{>0}$ and $k \neq 2$. Since $(\text{ad } \partial_2)^2 = 0$ by the Jacobi identity, we get $h_k := [x_k, \partial_2] \in \mathfrak{h}_{-\varepsilon_2}$ and $\varepsilon_2(h_k) = 0$. Hence, $[x_k, \partial_2] \otimes \partial_2^k = 1 \otimes h_k \partial_2^k = 0$, and the claim follows. \square

Remark 5.3. We can define a \mathfrak{g} -module homomorphism $K^+(S^{k+1}(V_*)) \rightarrow K^+(S^k(V_*))$ by $1 \otimes \partial_2^{k+1} \mapsto \partial_2 \otimes \partial_2^k$. It was proved for the finite-dimensional case in [S05], that the \mathfrak{g}_n -module $K^+(S^k((V_n)^*))$ has length 2 and the analogous maps give a resolution for $K^+(S^k((V_n)^*))$.

In contrast with Theorem 5.2 we have the following.

Proposition 5.4. *If X is a $\mathfrak{b}(\prec)^0$ -highest weight \mathfrak{g}^0 -module, then $K^-(X)$ is not simple. In particular, $K^-(V_{\lambda, \mu})$ is never simple for any simple $V_{\lambda, \mu} \in \mathbb{T}_{\mathfrak{gl}(\infty)}$.*

Proof. Let $v \in X$ be a nonzero $\mathfrak{b}(\prec)^0$ -highest weight vector, and let $x \in \mathfrak{g}_{\varepsilon_1 + \varepsilon_3 - \varepsilon_2} \subseteq \mathfrak{g}^1$ be a nonzero root vector. Since $\mathfrak{g}^1 \cong \Lambda^2(V) \otimes V_*$ as \mathfrak{g}^0 -modules, x is a $\mathfrak{b}(\prec)^0$ -highest weight vector of \mathfrak{g}^1 . We claim that $w := xv \in K^-(X)$ is a $\mathfrak{b}(\prec)^{\min}$ -highest weight vector. Indeed, since x and w are $\mathfrak{b}(\prec)^0$ -highest weight vectors, it is clear that w is a $\mathfrak{b}(\prec)^0$ -highest weight vector. Now let $y_i \in \mathfrak{g}_{-\varepsilon_i} \subseteq \mathfrak{g}^{-1}$ be a nonzero root vector, and observe that either $[y_i, x] \in \mathfrak{n}(\prec)^0$ or $[y_i, x] = 0$. In any case, we get $y_i w = 0$, and thus $\mathfrak{g}^{-1} w = 0$. This proves the claim, and the result. \square

Corollary 5.5. *If $X \in \mathbb{T}_{\mathfrak{gl}(\infty)}$, then $K^-(X)$ is not simple.*

Proof. If X is not simple, then $K^-(X)$ is not simple. If X is simple then $X \cong V_{\lambda, \mu}$ for some partitions λ, μ , so the claim follows from Proposition 5.4. \square

For a subalgebra \mathfrak{k} of \mathfrak{g} and a \mathfrak{g} -module M , we define $M^{\mathfrak{k}} = \{m \in M \mid \mathfrak{k} \cdot m = 0\}$. Recall that $\mathfrak{g}^< = \mathfrak{g}^{-1}$. We define the functor $\Psi : \mathbb{T}_{\tilde{W}}^{\leq} \rightarrow \tilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$ by

$$\Psi(M) := M^{\mathfrak{g}^<}.$$

Proposition 5.6. *If $M \in \mathbb{T}_{\tilde{W}}^{\leq}$, then $\Psi(M) \neq 0$. If $M \in \mathbb{T}_{\tilde{W}}^{\leq}$ is simple, then*

- (a) $\Psi(M)$ is a simple \mathfrak{g}^0 -module,
- (b) $\Psi(M)$ is in $\mathbb{T}_{\mathfrak{gl}(\infty)}$, and

(c) $M \cong L^-(\Psi(M))$.

Proof. For a nonzero vector $m \in M$ there exists n such that $(\mathfrak{t}_n \cap \mathfrak{g}^{\leq}) \subseteq \text{Ann}(m)$. Choose m' to be the longest non-zero element of the form $\partial_{i_1} \cdots \partial_{i_k} \cdot m$ such that $1 \leq i_1 < \cdots < i_k \leq n$. Then $\partial_j \cdot m' = 0$ for every $j \leq n$. Moreover, $\partial_j \cdot m' = 0$ for all $j > n$, since $\partial_j \in (\mathfrak{t}_n \cap \mathfrak{g}^{\leq}) \subseteq \text{Ann}(m)$ and $\partial_i \partial_j = -\partial_j \partial_i$ for any $i, j \in I$. This shows that $m' \in M^{\mathfrak{g}^{\leq}}$ and the first part is proved.

Assume now that $M \in \mathbb{T}_W$ is simple. The first part and the simplicity of M implies that there exist $n \in \mathbb{Z}$ such that $M = \bigoplus_{k \geq n} M^k$. Indeed, if $(M^{\mathfrak{g}^{\leq}})^n$ is nonzero, then $M = \mathbf{U}(\mathfrak{g})(M^{\mathfrak{g}^{\leq}})^n = \mathbf{U}(\mathfrak{g}^{\geq})(M^{\mathfrak{g}^{\leq}})^n$, which determines n . Moreover, we get $M^{\mathfrak{g}^{\leq}} = M^n$.

Now, if N is a proper \mathfrak{g}^0 -submodule of $M^{\mathfrak{g}^{\leq}}$, then $\mathbf{U}(\mathfrak{g})N = N \oplus \mathbf{U}(\mathfrak{g}^{\geq})^+ N$ and hence $M^{\mathfrak{g}^{\leq}} \cap \mathbf{U}(\mathfrak{g})N = N$. This implies that $\mathbf{U}(\mathfrak{g})N$ is a proper \mathfrak{g} -submodule of M , contradicting the irreducibility of M . Since $M^{\mathfrak{g}^{\leq}}$ is a simple \mathfrak{g}^0 -submodule of M , and $M|_{\mathfrak{g}^0} \in \widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$, we get $M^{\mathfrak{g}^{\leq}} \in \mathbb{T}_{\mathfrak{gl}(\infty)}$. Finally, we have $M \cong L^-(M^{\mathfrak{g}^{\leq}})$ since $M = \mathbf{U}(\mathfrak{g}^{\geq})M^{\mathfrak{g}^{\leq}}$. \square

Remark 5.7. The proof of Proposition 5.6 does not work if we replace \mathfrak{g}^{\leq} by \mathfrak{g}^{\geq} , since unlike $\mathbf{U}(\mathfrak{g}^{\leq})$, $\mathbf{U}(\mathfrak{g}^{\geq})$ is not a Grassmann algebra. In fact, $\Lambda(V)$, $\Lambda(V)_+$ and \mathfrak{g} are examples of modules in \mathbb{T}_W (and hence in \mathbb{T}_W^{\geq}) for which $M^{\mathfrak{g}^{\geq}} = 0$. Actually, we cannot have a simple module $M \in \mathbb{T}_W$ with $M^{\mathfrak{g}^{\geq}} \neq 0$. Indeed, if this were the case, then we could use the same arguments as those of Proposition 5.6 to conclude that $M^{\mathfrak{g}^{\geq}} \in \widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$, and hence that $M \cong K^+(M^{\mathfrak{g}^{\geq}})$ (by Theorem 5.2). But $K^+(M^{\mathfrak{g}^{\geq}}) \notin \mathbb{T}_W$, since it does not satisfy the l.a.c. for \mathfrak{g}^{\leq} .

Example 5.8. Note that $L^-(V)$ is isomorphic to $\Lambda(V)_+ := \Lambda(V)/\mathbb{C}$, since $(\Lambda(V)_+)^{\mathfrak{g}^{\leq}} = V$. And, $L^-(V_*)$ is isomorphic to the adjoint module of \mathfrak{g} , since it satisfies $\mathfrak{g}^{\mathfrak{g}^{\leq}} = \mathfrak{g}^{-1} \cong_{\mathfrak{g}^0} V_*$. A direct computation shows that the \mathfrak{g} -modules $\Lambda(V)_+$ and \mathfrak{g} are both objects in \mathbb{T}_W (see Example 3.6), and they have $\mathfrak{b}(\prec)^{\text{min}}$ -highest weights ε_1 and $-\varepsilon_2$, respectively. \circ

We set $L_{\lambda, \mu}^- := L^-(V_{\lambda, \mu})$. Then $L_{\lambda, \mu}^-$ is a highest weight \mathfrak{g} -module with respect to the Borel subalgebra $\mathfrak{b}(\prec)^{\text{min}}$ and has highest weight given by (1.4). It follows from Proposition 5.6 that $\Psi(L_{\lambda, \mu}^-) \cong V_{\lambda, \mu}$. The next result describes the simple objects of \mathbb{T}_W^{\leq} in terms of partitions.

Theorem 5.9. *The \mathfrak{g} -module $L_{\lambda, \mu}^-$ is in the category \mathbb{T}_W^{\leq} for any partitions λ and μ . Moreover, if $M \in \mathbb{T}_W^{\leq}$ is simple, then there exist partitions λ and μ for which $M \cong L_{\lambda, \mu}^-$.*

Proof. By Theorem 4.7, we have $K^-(V_{\lambda, \mu}) \in \widetilde{\mathbb{T}}_W^{\leq}$. It follows that $L_{\lambda, \mu}^- \in \mathbb{T}_W^{\leq}$, since $L_{\lambda, \mu}^-$ has finite length and $\widetilde{\mathbb{T}}_W^{\leq}$ is closed under taking quotients.

Now by Proposition 5.6, for any simple \mathfrak{g} -module M , we have that $\Psi(M)$ is a simple object of $\widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$, and therefore $\Psi(M) \cong V_{\lambda, \mu}$ for some partitions λ, μ . \square

Corollary 5.10. *Every simple module of \mathbb{T}_W^{\leq} (and hence of \mathbb{T}_W) is a highest weight module with respect to the Borel subalgebra $\mathfrak{b}(\prec)^{\text{min}}$.*

Proposition 5.11. *For any pair of partitions λ and μ , there exist $m, n \in \mathbb{Z}_{>0}$ such that the \mathfrak{g} -module $L_{\lambda, \mu}^-$ is a subquotient of $L^-(V)^{\otimes m} \otimes L^-(V_*)^{\otimes n}$. In particular, $L_{\lambda, \mu}^-$ lies in \mathbb{T}_W .*

Proof. Let $m = |\lambda|$ and $n = |\mu|$. Then $V^{\otimes m} \otimes V_*^{\otimes n}$ admits a $\mathfrak{b}(\prec)^0$ -highest weight vector $v \neq 0$ of highest weight $\sum_{i \in \mathbb{Z}_{>0}} \lambda_i \varepsilon_{2i-1} - \sum_{i \in \mathbb{Z}_{>0}} \mu_i \varepsilon_{2i} \in \mathfrak{h}^*$. The obvious embedding of \mathfrak{g}^0 -modules of

V into $K^-(V)$ gives rise to an embedding of \mathfrak{g}^0 -modules of V into $L^-(V)^{\mathfrak{g}^<}$, since V is simple. Therefore we have an embedding \mathfrak{g}^0 -modules

$$V^{\otimes m} \otimes V_*^{\otimes n} \hookrightarrow (L^-(V)^{\otimes m} \otimes L^-(V_*)^{\otimes n})^{\mathfrak{g}^<},$$

and we may assume that $v \in (L^-(V)^{\otimes m} \otimes L^-(V_*)^{\otimes n})^{\mathfrak{g}^<}$. Then v is a $\mathfrak{b}(\prec)^{\min}$ -highest weight vector, and $L_{\lambda,\mu}^-$ is a subquotient of $L^-(V)^{\otimes m} \otimes L^-(V_*)^{\otimes n}$. Since $L^-(V)$ and $L^-(V_*)$ are in \mathbb{T}_W (see Example 5.8), the result follows now from the fact that \mathbb{T}_W is closed under tensor products and subquotients. \square

Remark 5.12. Proposition 5.11 shows that the \mathfrak{g} -modules $L^-(V)$ and $L^-(V_*)$ play a similar role for the category \mathbb{T}_W as that of the \mathfrak{g}^0 -modules V and V_* for the category $\mathbb{T}_{\mathfrak{gl}(\infty)}$. In the category $\mathbb{T}_{\mathfrak{gl}(\infty)}$, all simple objects can be realized as subquotients of some $V^{\otimes m} \otimes V_*^{\otimes n}$.

We obtain the following corollary from Theorem 5.9 and 5.11.

Corollary 5.13. *The simple objects of \mathbb{T}_W and \mathbb{T}_W^{\leq} coincide.*

Remark 5.14. Note that Corollary 5.13 does not hold if we replace \mathbb{T}_W^{\leq} with \mathbb{T}_W^{\geq} . Indeed, by Theorem 4.7 and Theorem 5.2, the induced \mathfrak{g} -module $K^+(V_{\lambda,\mu})$ is simple when $(\lambda, \mu) \neq (\emptyset, (\mu_1))$; however, we have seen in Remark 4.8 that $K^+(V_{\lambda,\mu})$ is never in \mathbb{T}_W .

6. REALIZATION OF SIMPLE MODULES AS TENSOR FIELDS

In this section, we prove that the simple module $L_{\lambda,\mu}^-$ can be realized as a module of tensor fields. This is a natural generalization of the classical work of Bernstein and Leites for the finite-dimensional Cartan type Lie superalgebra $W(n)$ [BL81].

Let X be a $\mathfrak{gl}(\infty)$ -module. We define a \mathfrak{g} -module structure on the vector space

$$\mathcal{T}(X) := \Lambda(\infty) \otimes X$$

by linearly extending the following action:

$$\underline{\xi}^{\underline{e}} \partial_j \cdot (fv) = \underline{\xi}^{\underline{e}} \partial_j (f)v + (-1)^{p(\underline{\xi}^{\underline{e}} \partial_j)p(f)} \sum_{i \in I} \partial_i(\underline{\xi}^{\underline{e}}) f E_{i,j} v,$$

where $f \in \Lambda(\infty)$ and $v \in X$. This action is well defined since $\partial_i(\underline{\xi}^{\underline{e}})$ is nonzero only for finitely many $i \in I$. We call the module $\mathcal{T}(X)$ the module of tensor fields associated to X .

Remark 6.1. Notice that if $X \in \widetilde{\mathbb{T}}_{\mathfrak{gl}(\infty)}$, then $\mathcal{T}(X)$ lies in $\widetilde{\mathbb{T}}_W$. Indeed, the \mathfrak{g} -module $\mathcal{T}(X)$ is clearly an \mathfrak{h} -weight module. Moreover, since $X, \Lambda(\infty)$ are integrable over \mathfrak{g}^0 , we have that $\mathcal{T}(X)$ is also integrable over \mathfrak{g}^0 . Finally, $\mathcal{T}(X)$ satisfies the l.a.c. for \mathfrak{g} , since the action of \mathfrak{g} on $\Lambda(\infty)$ and the action of $\mathfrak{gl}(\infty)$ on X both satisfy the l.a.c..

Example 6.2. If V_* is the conatural $\mathfrak{gl}(\infty)$ -module, then the \mathfrak{g} -module $\mathcal{T}(V_*) = \Lambda(\infty) \otimes V_*$ is isomorphic to the adjoint module \mathfrak{g} (see Section 3). If \mathbb{C} is the trivial $\mathfrak{gl}(\infty)$ -module, then $\mathcal{T}(\mathbb{C}) \cong \Lambda(\infty)$ is the natural module. \circ

Let's consider X as a \mathfrak{g}^{\geq} -module by declaring $\mathfrak{g}^> \cdot X = 0$. Then we define

$$\text{Coind}_{\mathfrak{g}^{\geq}}^{\mathfrak{g}}(X) := \text{Hom}_{\mathfrak{g}^{\geq}}(\mathbf{U}(\mathfrak{g}), X).$$

An easy computation shows that if $t \in X \cong \mathbb{C} \otimes X \subseteq \mathcal{T}(X)$, then $\partial_i \cdot t = 0$ for all $i \in I$. For an arbitrary $t = \sum f_i v_i \in \mathcal{T}(X)$, we set

$$t(0) := \sum f_i(0)v_i \in X.$$

Now we have a natural embedding of \mathfrak{g} -modules $\varphi : \mathcal{T}(X) \hookrightarrow \text{Coind}_{\mathfrak{g}}^{\mathfrak{g}}(X)$, where for $t \in \mathcal{T}(X)$ and $u \in \mathbf{U}(\mathfrak{g})$,

$$\varphi(t)(u) := (-1)^{p(t)p(u)}(u \cdot t)(0).$$

Next we consider the finite-dimensional case. Suppose we have inclusions of \mathfrak{g}_n^0 -modules $X_n \subseteq X_{n+1}$ for every $n \in \mathbb{Z}_{>0}$, and consider the \mathfrak{g}^0 -module $X = \bigcup X_n$. We set $\mathcal{T}(X_n) := \Lambda(n) \otimes X_n$. Then we have $\mathcal{T}(X) = \bigcup \mathcal{T}(X_n)$, where we regard $\mathcal{T}(X_n) \subseteq \mathcal{T}(X_{n+1})$ in the obvious way. The restriction $\varphi_n := \varphi|_{\mathcal{T}(X_n)}$ gives embeddings of $\mathcal{T}(X_n)$ into $\text{Hom}_{\mathfrak{g}_n^{\geq}}(\mathbf{U}(\mathfrak{g}_n), X_n)$. Moreover we have the following statement from [BL81].

Lemma 6.3. $\mathcal{T}(X_n) \cong \text{Coind}_{\mathfrak{g}_n^{\geq}}^{\mathfrak{g}_n^{\geq}}(X_n)$, where the isomorphism is φ_n .

Proof. Since φ_n is an embedding, we just have to check that the dimensions match. This follows from the following

$$\begin{aligned} \mathcal{T}(X_n) &:= \Lambda(\xi_1, \dots, \xi_n) \otimes_{\mathbb{C}} X_n \\ &\cong \Lambda(\partial_1, \dots, \partial_n)^* \otimes_{\mathbb{C}} X_n \\ &\cong \text{Hom}_{\mathbb{C}}(\Lambda(\partial_1, \dots, \partial_n), X_n) \\ &\cong \text{Hom}_{\mathbb{C}}(\Lambda(\partial_1, \dots, \partial_n), \text{Hom}_{\mathfrak{g}_n^{\geq}}(\mathbf{U}(\mathfrak{g}_n^{\geq}), X_n)) \\ &\cong \text{Hom}_{\mathfrak{g}_n^{\geq}}(\Lambda(\partial_1, \dots, \partial_n) \otimes_{\mathbb{C}} \mathbf{U}(\mathfrak{g}_n^{\geq}), X_n) \\ &\cong \text{Hom}_{\mathfrak{g}_n^{\geq}}(\mathbf{U}(\mathfrak{g}_n), X_n). \quad \square \end{aligned}$$

Remark 6.4. In the infinite-dimensional case, the second isomorphism displayed above is just a natural embedding from $\Lambda(\partial)^* \otimes_{\mathbb{C}} X$ to $\text{Hom}_{\mathbb{C}}(\Lambda(\partial), X)$.

Consider the submodule $\bigcup \text{im } \varphi_n$ of $\text{Coind}_{\mathfrak{g}}^{\mathfrak{g}}(X)$. It follows from Lemma 6.3 that $\bigcup \text{im } \varphi_n \cong \bigcup \mathcal{T}(X_i)$ is a submodule of $\mathcal{T}(X)$. But since every element of $\mathcal{T}(X)$ lies in some $\mathcal{T}(X_n)$ for $n \gg 0$, we have the following:

$$\mathcal{T}(X) = \bigcup \mathcal{T}(X_n) \cong \bigcup \text{im } \varphi_n = \bigcup \text{Coind}_{\mathfrak{g}_n^{\geq}}^{\mathfrak{g}_n^{\geq}}(X_n) \subseteq \text{Coind}_{\mathfrak{g}}^{\mathfrak{g}}(X).$$

For any finite-dimensional \mathfrak{g}_n^0 -module X_n , we have

$$\begin{aligned} \mathcal{T}(X_n^*) &\cong \text{Hom}_{\mathfrak{g}_n^{\geq}}(\mathbf{U}(\mathfrak{g}_n), \text{Hom}_{\mathbb{C}}(X_n, \mathbb{C})) \\ &\cong \text{Hom}_{\mathbb{C}}(\mathbf{U}(\mathfrak{g}_n) \otimes_{\mathbf{U}(\mathfrak{g}_n^{\geq})} X_n, \mathbb{C}) = (\text{Ind}_{\mathfrak{g}_n^{\geq}}^{\mathfrak{g}_n^{\geq}} X_n)^* = K_n^+(X_n)^*. \end{aligned}$$

In particular, $\mathcal{T}(X_n) \cong K_n^+(X_n^*)^*$. Using this fact one can prove the following result.

Proposition 6.5. *If $(\lambda, \mu) \neq ((\lambda_1), \emptyset)$, then $\mathcal{T}(V_{\lambda, \mu})$ is a locally simple \mathfrak{g} -module, and hence simple.*

Proof. Let $\lambda = (\lambda_1 \geq \dots \geq \lambda_k)$ and $\mu = (\mu_1 \geq \dots \geq \mu_l)$. Recall that the simple \mathfrak{g}^0 -module $X := V_{\lambda, \mu}$ has a decomposition $X = \bigcup X_n$, where, for $n \gg 0$, X_n is the \mathfrak{g}_n^0 -module

$$X_n := V_{\mathfrak{b}(\langle \cdot \rangle)_n^0}(\lambda_1 \varepsilon_1 + \dots + \lambda_k \varepsilon_k - \mu_l \varepsilon_{n-l} - \dots - \mu_1 \varepsilon_n).$$

Now $\mathcal{T}(X_n) \cong K_n^+(X_n^*)^*$, where

$$X_n^* \cong V_{\mathfrak{b}(\prec)_n^0}(\mu_1\varepsilon_1 + \cdots + \mu_l\varepsilon_l - \lambda_k\varepsilon_{n-k} - \cdots - \lambda_1\varepsilon_n).$$

It follows from Remark 5.1 that the highest weight $\mu_1\varepsilon_1 + \cdots + \mu_l\varepsilon_l - \lambda_k\varepsilon_{n-k} - \cdots - \lambda_1\varepsilon_n$ of X_n^* is atypical if and only if $(\lambda, \mu) = ((\lambda_1), \emptyset)$. Hence, $\mathcal{T}(X_n) \cong K_n^+(X_n^*)^*$ is simple if and only if $(\lambda, \mu) \neq ((\lambda_1), \emptyset)$. Now the claim follows from the fact that $\mathcal{T}(X) \cong \bigcup \mathcal{T}(X_n)$. \square

The next corollary follows from Remark 6.1 and Proposition 6.5.

Corollary 6.6. *If $(\lambda, \mu) \neq ((\lambda_1), \emptyset)$, then $\mathcal{T}(V_{\lambda, \mu})$ is in \mathbb{T}_W .*

The following theorem gives a realization of the simple modules in \mathbb{T}_W as tensor fields.

Theorem 6.7. *For any (λ, μ) we have that $L_{\lambda, \mu}^-$ is isomorphic to a submodule of $\mathcal{T}(V_{\lambda, \mu})$. Moreover, if $(\lambda, \mu) \neq ((\lambda_1), \emptyset)$, then $L_{\lambda, \mu}^- \cong \mathcal{T}(V_{\lambda, \mu})$.*

Proof. For $n > k + l$, we denote by $V_{\lambda, \mu, n}$ the \mathfrak{g}_n^0 -module $V_{\mathfrak{b}(\prec)_n^0}(\nu)$ with highest weight $\nu := \sum_{i \in \mathbb{Z}_{>0}} \lambda_i \varepsilon_{2i-1} - \sum_{i \in \mathbb{Z}_{>0}} \mu_i \varepsilon_{2i}$. We notice that for all $n > k + l$ we have a short exact sequence of \mathfrak{g}_n -modules

$$0 \rightarrow Q_n(V_{\lambda, \mu, n}^*) \rightarrow K_n^+(V_{\lambda, \mu, n}^*) \rightarrow L_n^+(V_{\lambda, \mu, n}^*) \rightarrow 0$$

which yields the following short exact sequence of \mathfrak{g}_n -modules

$$0 \rightarrow L_n^+(V_{\lambda, \mu, n}^*)^* \rightarrow K_n^+(V_{\lambda, \mu, n}^*)^* \cong \mathcal{T}(V_{\lambda, \mu, n}) \rightarrow Q_n(V_{\lambda, \mu, n}^*)^* \rightarrow 0.$$

In particular, if v denotes the $\mathfrak{b}(\prec)_n^0$ -highest weight of $V_{\lambda, \mu, n}^*$, then $L_n^+(V_{\lambda, \mu, n}^*)^*$ is the submodule of $\mathcal{T}(V_{\lambda, \mu, n})$ generated by v . Since the weight of v in $\mathcal{T}(V_{\lambda, \mu, n}) \subset \mathcal{T}(V_{\lambda, \mu})$ is ν and $\mathcal{T}(V_{\lambda, \mu, n})_\nu = (V_{\lambda, \mu, n})_\nu = (V_{\lambda, \mu})_\nu$, we conclude that v is a $\mathfrak{b}(\prec)_n^0$ -highest weight of $V_{\lambda, \mu, n}$ and hence a $\mathfrak{b}(\prec)_n^{\min}$ -highest weight vector in $\mathcal{T}(V_{\lambda, \mu, n})$. Thus we have isomorphisms of \mathfrak{g}_n -modules $L_n^+(V_{\lambda, \mu, n}^*)^* \cong \mathbf{U}(\mathfrak{g}_n)v \cong L_n^-(V_{\lambda, \mu, n})$, for all $n \gg 0$. This implies that $\mathbf{U}(\mathfrak{g})v$ is locally simple and hence isomorphic to $L_{\lambda, \mu}^-$. Finally, the latter statement follows from Proposition 6.5. \square

7. INJECTIVE MODULES

In this section we prove that the category \mathbb{T}_W has enough injective objects. Moreover, for each simple module $L_{\lambda, \mu}^-$ in \mathbb{T}_W , we use Theorem 6.7 to provide an explicit injective module in \mathbb{T}_W that contains $L_{\lambda, \mu}^-$. Let \mathfrak{g} -mod (resp. \mathfrak{g}^0 -mod) be the category of all \mathfrak{g} -modules (resp. \mathfrak{g}^0 -mod), $\text{Int}_{\mathfrak{g}^0}$ be full subcategory of \mathfrak{g}^0 -mod consisting of integrable \mathfrak{g}^0 -modules, $\text{Int}_{\mathfrak{g}, \mathfrak{g}^0}$ be the full subcategory of \mathfrak{g} -mod consisting of \mathfrak{g}^0 -integrable modules, and $\text{Int}_{\mathfrak{g}, \mathfrak{g}^0}^{\text{wt}}$ be full subcategory of $\text{Int}_{\mathfrak{g}, \mathfrak{g}^0}$ consisting of \mathfrak{h} -weight modules. Define the functors $\Gamma_{\mathfrak{g}, \mathfrak{g}^0} : \mathfrak{g}$ -mod $\rightarrow \text{Int}_{\mathfrak{g}, \mathfrak{g}^0}$ where $\Gamma_{\mathfrak{g}, \mathfrak{g}^0}(M) = \{m \in M \mid \dim \text{Span}\{g^i m \mid i \geq 0\} < \infty, \forall g \in \mathfrak{g}^0\}$ is the largest submodule of M that is \mathfrak{g}^0 -integrable; $\Gamma_{\mathfrak{h}} : \text{Int}_{\mathfrak{g}, \mathfrak{g}^0} \rightarrow \text{Int}_{\mathfrak{g}, \mathfrak{g}^0}^{\text{wt}}$ where $\Gamma_{\mathfrak{h}}(M) := \bigoplus_{\mu \in \mathfrak{h}^*} M_\mu$ is the largest submodule of M that is \mathfrak{h} -semisimple; and $\Theta : \text{Int}_{\mathfrak{g}, \mathfrak{g}^0}^{\text{wt}} \rightarrow \mathbb{T}_W$ where $\Theta(M) = \bigcup_{n > 0} M^{t_n}$ is the largest submodule of M that satisfies the l.a.c. for \mathfrak{g} . Let $\Gamma : \mathfrak{g}$ -mod $\rightarrow \mathbb{T}_W$ denote the composition $\Phi \circ \Gamma_{\mathfrak{h}} \circ \Gamma_{\mathfrak{g}, \mathfrak{g}^0}$.

All three functors $\Gamma_{\mathfrak{g}, \mathfrak{g}^0}$, $\Gamma_{\mathfrak{h}}$ and Φ are left exact and are right adjoint to the respective inclusion functors $\text{Int}_{\mathfrak{g}, \mathfrak{g}^0} \subset \mathfrak{g}$ -mod, $\text{Int}_{\mathfrak{g}, \mathfrak{g}^0}^{\text{wt}} \subset \text{Int}_{\mathfrak{g}, \mathfrak{g}^0}$ and $\mathbb{T}_W \subset \text{Int}_{\mathfrak{g}, \mathfrak{g}^0}^{\text{wt}}$. Now we have the following result:

Proposition 7.1. *If $I \in \mathfrak{g}\text{-mod}$ is injective, then $\Gamma(I)$ is injective in \mathbb{T}_W . Moreover, the category \mathbb{T}_W has enough injectives.*

Proof. The former statement follows from the fact that Γ is right adjoint to the inclusion functor $\mathbb{T}_W \subset \mathfrak{g}\text{-mod}$. To prove the latter statement, take $M \in \mathbb{T}_W$ and notice that if I_M is the injective hull of M in $\mathfrak{g}\text{-mod}$, then the natural isomorphism

$$\text{Hom}_{\mathfrak{g}}(M, I_M) \cong \text{Hom}_{\mathfrak{g}}(M, \Gamma(I_M))$$

implies that the inclusion in $\text{Hom}_{\mathfrak{g}}(M, I_M)$ gives an inclusion in $\text{Hom}_{\mathfrak{g}}(M, \Gamma(I_M))$. \square

Corollary 7.2. *If $I \in \mathfrak{g}^0\text{-mod}$ is injective, then $\Gamma(\text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(I)) \in \mathbb{T}_W$ is injective.*

Proof. This is because the functor $\text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(-) : \mathfrak{g}^0\text{-mod} \rightarrow \mathfrak{g}\text{-mod}$ sends injectives to injectives [Kna88, Corollary 6.4]. \square

Proposition 7.3. *If $I \in \text{Int}_{\mathfrak{g}^0}$ is injective, then $\Gamma_{\mathfrak{g}, \mathfrak{g}^0}(\text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(I))$ is injective in $\text{Int}_{\mathfrak{g}, \mathfrak{g}^0}$.*

Proof. Let $M \in \text{Int}_{\mathfrak{g}^0}$. The adjunction between restriction and coinduction gives us that

$$\text{Ext}_{\mathfrak{g}}^1(M, \text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(I)) \cong \text{Ext}_{\mathfrak{g}^0}^1(M, I) = \text{Ext}_{\text{Int}_{\mathfrak{g}^0}}^1(M, I) = 0,$$

where we are using that $\text{Int}_{\mathfrak{g}^0}$ is closed under taking extensions [PS11]. Now consider the short exact sequence

$$0 \rightarrow \Gamma_{\mathfrak{g}, \mathfrak{g}^0}(\text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(I)) \rightarrow \text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(I) \rightarrow Q := \text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(I) / \Gamma_{\mathfrak{g}, \mathfrak{g}^0}(\text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(I)) \rightarrow 0$$

and the corresponding long exact sequence

$$\begin{aligned} 0 \rightarrow \text{Hom}_{\mathfrak{g}}(M, \Gamma_{\mathfrak{g}, \mathfrak{g}^0}(\text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(I))) &\rightarrow \text{Hom}_{\mathfrak{g}}(M, \text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(I)) \rightarrow \\ &\rightarrow \text{Hom}_{\mathfrak{g}}(M, Q) \rightarrow \text{Ext}_{\mathfrak{g}}^1(M, \Gamma_{\mathfrak{g}, \mathfrak{g}^0}(\text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(I))) \rightarrow \text{Ext}_{\mathfrak{g}}^1(M, \text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(I)) = 0 \end{aligned}$$

Since $M \in \text{Int}_{\mathfrak{g}^0}$, then $\text{Hom}_{\mathfrak{g}}(M, Q) = 0$. Thus $\text{Ext}_{\mathfrak{g}}^1(M, \Gamma_{\mathfrak{g}, \mathfrak{g}^0}(\text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(I))) = 0$, which proves the statement. \square

Let $\Gamma_{\mathfrak{g}^0} : \mathfrak{g}^0\text{-mod} \rightarrow \text{Int}_{\mathfrak{g}^0}$ be the functor defined by $\Gamma_{\mathfrak{g}^0}(M) = \{m \in M \mid \dim \text{Span}\{g^i m \mid i \geq 0\} < \infty, \forall g \in \mathfrak{g}^0\}$.

Corollary 7.4. *For every $M \in \text{Int}_{\mathfrak{g}^0}$, we have that $\Gamma_{\mathfrak{g}, \mathfrak{g}^0}(\text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(\Gamma_{\mathfrak{g}^0}(M^*)))$ is injective in $\text{Int}_{\mathfrak{g}, \mathfrak{g}^0}$.*

Proof. It follows from [PS11, Proposition 3.1]. \square

Corollary 7.5. *If $I \in \text{Int}_{\mathfrak{g}^0}$ is injective, then $\Gamma(\text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(I))$ is injective in \mathbb{T}_W . In particular, for every $I \in \text{Int}_{\mathfrak{g}^0}$ we have that $\Gamma(\text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(\Gamma_{\mathfrak{g}^0}(I^*)))$ is injective in \mathbb{T}_W .*

Proof. It follows from the fact that $\Gamma_{\mathfrak{h}}$ and Φ are right adjoint to the respective inclusion functors $\text{Int}_{\mathfrak{g}, \mathfrak{g}^0}^{\text{wt}} \subset \text{Int}_{\mathfrak{g}, \mathfrak{g}^0}$ and $\mathbb{T}_W \subset \text{Int}_{\mathfrak{g}, \mathfrak{g}^0}^{\text{wt}}$. \square

Proposition 7.6. *Each simple module $L_{\lambda, \mu}^-$ of \mathbb{T}_W is isomorphic to a submodule of the injective module $\Gamma(\text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}((V_{\lambda, \mu})^*))$.*

Proof. By Corollary 6.7 of [PS11], $((V_{\lambda,\mu})_*)^*$ is an injective hull of $V_{\lambda,\mu}$ in $\text{Int}_{\mathfrak{g}^0}$. Moreover, we have the following chain of inclusions

$$\text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(((V_{\lambda,\mu})_*)^*) \supset \text{Coind}_{\mathfrak{g}^{\geq}}^{\mathfrak{g}}(((V_{\lambda,\mu})_*)^*) \supset \mathcal{T}(((V_{\lambda,\mu})_*)^*) \supset \mathcal{T}(V_{\lambda,\mu}) \supset L_{\lambda,\mu}^-$$

where the latter inclusion follows from Theorem 6.7. Thus $L_{\lambda,\mu}^- \subset \Gamma\left(\text{Coind}_{\mathfrak{g}^0}^{\mathfrak{g}}(((V_{\lambda,\mu})_*)^*)\right)$. \square

REFERENCES

- [Ba99] A. Baranov, *Finitary simple Lie algebras*, J. Algebra, **219**(1), (1999) 299–329.
- [BL81] I. Bernstein, D. Leites, *Invariant differential operators and irreducible representations of Lie superalgebras of vector fields*, Selecta Math. Soviet **1**(2), (1981) 143–160.
- [BL83] I. Bernstein, D. Leites, *Irreducible representations of finite-dimensional Lie superalgebras of type W*, Selecta Math. Soviet, **3**(1), (1983) 63–68.
- [Br03] J. Brundan, *Kazhdan-Lusztig polynomials and character formulae for the Lie superalgebra $\mathfrak{gl}(m|n)$* , J. Amer. Math. Soc. **16** (2003), no. 1, 185–231.
- [ChP17] A. Chirvasitu, I. Penkov, *Representation categories of Mackey Lie algebras as universal monoidal categories*, Pure and Applied Mathematics Quarterly **13** (2017), 77–121.
- [CoP19] K. Coulembier, I. Penkov, *On an infinite limit of BGG categories \mathcal{O}* , Mosc. Math. J. **4** (2019), 655–693.
- [CP22] L. Calixto, I. Penkov, *Integrable bounded weight modules of classical Lie superalgebras at infinity*, J. Pure Appl. Algebra **226** (2022), 27 pp.
- [DPS16] E. Dan-Cohen, I. Penkov, V. Serganova, *A Koszul category of representations of finitary Lie algebras*, Advances in Mathematics **289**, (2016) 250–278.
- [FPS16] I. Frenkel, I. Penkov, V. Serganova, *A categorification of the boson-fermion correspondence via representation theory of $\mathfrak{sl}(\infty)$* , Comm. Math. Phys. **341** (2016) no.3, 911–931 .
- [Gav15] F. Gavarini, *Algebraic supergroups of Cartan type*, Forum Mathematicum **26**(5) (2014), 1473–1564.
- [HPS19] C. Hoyt, I. Penkov, V. Serganova, *Integrable $\mathfrak{sl}(\infty)$ -modules and Category \mathcal{O} for $\mathfrak{gl}(m|n)$* , J. Lond. Math. Soc. **99**(2), 403–427 (2019).
- [Kac77] V. Kac, *Lie superalgebras*, Adv. Math. **26** (1977), 8–96.
- [Kna88] A. Knapp, *Lie groups, Lie algebras, and cohomology*, Mathematical Notes, 34. Princeton University Press, Princeton, NJ, 1988. xii+510 pp. ISBN: 0-691-08498-X.
- [PH22] I. Penkov, C. Hoyt, *Classical Lie algebras at infinity*, Springer Monographs in Math., Springer International Publishing (2022).
- [PS11] I. Penkov, V. Serganova, *Categories of integrable $\mathfrak{sl}(\infty)$ -, $\mathfrak{o}(\infty)$ -, $\mathfrak{sp}(\infty)$ -modules*, Contemp. Math. **557** (2011), 335–357.
- [PS19] I. Penkov, V. Serganova, *Large annihilator category \mathcal{O} for $\mathfrak{sl}(\infty)$, $\mathfrak{so}(\infty)$, $\mathfrak{sp}(\infty)$* , J. of Algebra **532** (2019), 249–279.
- [PSt11] I. Penkov, K. Styrkas, *Tensor representations of infinite-dimensional root-reductive Lie algebras*, Developments and Trends in Infinite-Dimensional Lie Theory, Progress in Mathematics **288**, Birkhäuser (2011), 127–150.
- [S05] V. Serganova, *On representations of Cartan type Lie superalgebras*, Amer. Math. Soc. Transl. **213** (2005), 223–239.
- [S14] V. Serganova, *Classical Lie superalgebras at infinity*, Advances in Lie Superalgebras, Springer INdAM Ser. no. 7, Springer, Cham (2014), 181–201.

L. CALIXTO: DEPARTMENT OF MATHEMATICS, FEDERAL UNIVERSITY OF MINAS GERAIS, BRAZIL

Email address: lhcalixto@ufmg.br

C. HOYT: DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, RAMAT GAN 52900, ISRAEL

Email address: math.crystal@gmail.com