

A Unified View on the Functorial Nerve Theorem and its Variations

Ulrich Bauer* Michael Kerber[†] Fabian Roll*
Alexander Rolle*

August 2, 2022

Abstract

The nerve theorem is a basic result of algebraic topology that plays a central role in computational and applied aspects of the subject. In applied topology, one often needs a nerve theorem that is functorial in an appropriate sense, and furthermore one often needs a nerve theorem for closed covers, as well as for open covers. While the techniques for proving such functorial nerve theorems have long been available, there is unfortunately no general-purpose, explicit treatment of this topic in the literature. We address this by proving a variety of functorial nerve theorems. First, we show how one can use relatively elementary techniques to prove nerve theorems for covers by closed convex sets in Euclidean space, and for covers of a simplicial complex by subcomplexes. Then, we prove a more general, “unified” nerve theorem that recovers both of these, using standard techniques from abstract homotopy theory.

Contents

1. Introduction	2
2. Functorial Nerve Theorems via Homotopy Colimits	10
2.1. Homotopy Colimits and the Bar Construction	11
2.2. Functorial Nerve Theorems via the Bar Construction	14
3. Nerve Theorems for Closed Convex Covers	16
3.1. A Functorial Nerve Theorem for Closed Convex Covers	20
3.2. A One-Arrow Functorial Nerve Theorem for Pointed Covers	20
4. Nerve Theorems for Simplicial and Semi-Algebraic Covers	21
4.1. A Functorial Nerve Theorem for Simplicial Covers	22

*Technical University of Munich (TUM), Germany

[†]Graz University of Technology, Austria

4.2. A Functorial Nerve Theorem for Semi-Algebraic Covers	25
4.3. A Functorial Version of Björner’s Nerve Theorem	26
5. A Unified Nerve Theorem	27
5.1. Applications of the Unified Nerve Theorem	31
5.2. Simplicial Model Categories	33
5.3. Proof of the Unified Nerve Theorem	37
A. Auxiliary Lemmas about Geometric Simplicial Complexes	42
B. Discrete Morse Theory for Infinite Complexes	43
C. The Blowup Complex for Covers by Subcomplexes	46
D. The Blowup Complex for Open Covers	47

1. Introduction

Background If $\mathcal{U} = (U_i)_{i \in I}$ is a cover of a topological space X , then the *nerve* of \mathcal{U} , which dates back to Alexandroff [Ale28], is the simplicial complex $\text{Nrv}(\mathcal{U})$ whose simplices are the finite subsets $J \subseteq I$ such that the intersection $\bigcap_{i \in J} U_i$ is non-empty. The nerve of a cover played an important role in the development of homology and cohomology theory. In particular, Čech (co)homology is given by the (co)limit of the (co)homology groups of the nerves of a directed system of open covers ordered by refinement. A historical exposition can be found in [EH80, Chapter 2].

The *nerve theorem*, whose early versions are due to Leray [Ler45], Borsuk [Bor48], and Weil [Wei52], is a basic result in algebraic and combinatorial topology. Roughly speaking, it says that if every non-empty finite intersection of cover elements is contractible, then, subject to some further tameness conditions on X and \mathcal{U} , the space X is homotopy equivalent to the nerve of \mathcal{U} .

The literature on the nerve theorem is extensive but unfortunately hard to navigate. In part, this is because there are many different variants of the nerve theorem. There are many choices one can make for the “further tameness conditions” on the space and cover that yield a nerve theorem, and there are further choices one can make for the kind of equivalence one works with. For example, one could ask that the non-empty finite intersections are weakly homotopy equivalent to the one-point space, or to be acyclic. Many of the original results use concepts that are now obscure, and the many possible choices for hypotheses and proof techniques make it difficult to compare all the available nerve theorems.

Nowadays, the nerve theorem and the aspect of functoriality play a crucial role in applied topology. Nerves are the main way to replace a topological space, determined by the data points using geometric constructions, with a combinatorial model that is suitable for computations. Two prominent examples are the *Čech complex* and the *Delaunay complex*, which arise as nerves of a collection of closed balls and closed Voronoi

balls, respectively. Another important example is the *Vietoris–Rips complex*, which is not usually defined as the nerve of a cover, though it is isomorphic to a nerve [GM05; Cha+09]. Note that, while one can choose whether to use open or closed sets when defining the Čech and Vietoris–Rips complexes, the only standard way to define the Delaunay complex uses closed sets.




Figure 1.: A cover by closed balls (left) and closed Voronoi balls (right) together with the corresponding Čech and Delaunay complex.

These examples are typical, in that the topological spaces determined by data points usually depend on one or more parameters, leading to filtrations of topological spaces and covers. Now functoriality ensures that the corresponding nerves form a filtration as well. For example, if $X \subset \mathbb{R}^d$ is a finite set of points, the offset filtration O is the filtration of \mathbb{R}^d with $O_r = \cup_{x \in X} D_r(x)$ for $r > 0$, where $D_r(x)$ is the closed ball about x of radius r . The nerve of the cover $\mathcal{U}_r = (D_r(x))_{x \in X}$ is the Čech complex, and as r varies, these complexes form a filtration as well. In this case, the nerve theorem says that O_r is homotopy equivalent to the nerve of \mathcal{U}_r .

Going further, one wants a nerve theorem to provide homotopy equivalences that are somehow compatible with the inclusion maps in these two filtrations. This is necessary in particular if one is interested in persistent homology [EH10], which is an algebraic invariant of filtrations that encodes the homology of each step of the filtration, as well as the maps in homology induced by each inclusion. There are several ways in which the homotopy equivalences provided by a nerve theorem might be compatible with the inclusions in these two filtrations, as we will now explain.

In order to prove that the persistent homology of the offset filtration is isomorphic to the persistent homology of the associated Čech complex filtration, it suffices to have isomorphisms $H_n(O_r) \cong H_n(\text{Nrv}(\mathcal{U}_r))$ such that all the squares of the following form commute:

$$\begin{array}{ccc}
 H_n(O_r) & \xrightarrow{\cong} & H_n(\text{Nrv}(\mathcal{U}_r)) \\
 \downarrow & & \downarrow \\
 H_n(O_{r'}) & \xrightarrow{\cong} & H_n(\text{Nrv}(\mathcal{U}_{r'}))
 \end{array} \tag{1.1}$$

By Theorem B below, such isomorphisms can be constructed from the induced maps of homotopy equivalences $|\text{Nrv}(\mathcal{U}_r)| \rightarrow O_r$ between the nerves and the offsets such that all

squares of the following form commute:

$$\begin{array}{ccc}
O_r & \xleftarrow{\simeq} & |\mathrm{Nrv}(\mathcal{U}_r)| \\
\downarrow & & \downarrow \\
O_{r'} & \xleftarrow{\simeq} & |\mathrm{Nrv}(\mathcal{U}_{r'})|
\end{array} \tag{1.2}$$

The construction of these compatible homotopy equivalences relies on the fact that the cover elements of the offset filtration are convex and that the inclusions $O_r \hookrightarrow O_{r'}$ are affine linear.

For a more general filtration (X_r, \mathcal{A}_r) , with $X_r = \bigcup \mathcal{A}_r$, a similar strategy does not necessarily produce commuting diagrams as in 1.2. However, if one is only interested in the filtration after applying homology or some other homotopy-invariant functor, then it suffices to have homotopy equivalences $X_r \rightarrow |\mathrm{Nrv}(\mathcal{A}_r)|$ such that all squares of the following form commute *up to homotopy*:

$$\begin{array}{ccc}
X_r & \xrightarrow{\simeq} & |\mathrm{Nrv}(\mathcal{A}_r)| \\
\downarrow & \nearrow H & \downarrow \\
X_{r'} & \xrightarrow{\simeq} & |\mathrm{Nrv}(\mathcal{A}_{r'})|
\end{array} \tag{1.3}$$

In the diagram, H is a homotopy from the bottom route around the square to the top route. Nerve theorems with this structure are often used in the study of persistent homology (for references, see the end of this introduction).

However, in some homotopy-theoretic approaches to applied topology, we need a nerve theorem that is compatible with the inclusions $X_r \hookrightarrow X_{r'}$ on the nose, and not just up to homotopy. In this paper, we will prove nerve theorems that provide strictly commuting diagrams, at the cost of introducing an intermediary between the covered space and the nerve: we obtain a filtration Z_r and homotopy equivalences $Z_r \rightarrow X_r$ and $Z_r \rightarrow |\mathrm{Nrv}(\mathcal{A}_r)|$ such that all the diagrams of the following form commute:

$$\begin{array}{ccccc}
X_r & \xleftarrow{\simeq} & Z_r & \xrightarrow{\simeq} & |\mathrm{Nrv}(\mathcal{A}_r)| \\
\downarrow & & \downarrow & & \downarrow \\
X_{r'} & \xleftarrow{\simeq} & Z_{r'} & \xrightarrow{\simeq} & |\mathrm{Nrv}(\mathcal{A}_{r'})|
\end{array} \tag{1.4}$$

While one can avoid introducing intermediate objects in the special case of the offset filtration, this is not possible in general, as we explain below. Diagrams of the form 1.4 appear classically in the study of homotopy categories [GZ67]. More recently, they appear in Blumberg and Lesnick’s work on the *homotopy interleaving distance* [BL17], a distance on diagrams of spaces that is universal among stable and homotopy-invariant distances. The idea is to define an equivalence relation on filtered spaces such that F and F' are related if they can be connected via an intermediate filtration, as above, with the horizontal arrows weak homotopy equivalences. Then, filtered spaces F_1 and F_2 are homotopy interleaved if F_1 is related to some F'_1 , F_2 is related to some F'_2 , and F'_1 and F'_2 are interleaved. An important motivation for nerve theorems that provide diagrams of the form 1.4 is that they can be used in frameworks like the one of Blumberg–Lesnick.

Functorial Nerve Theorems In order to say precisely what we mean by a functorial nerve theorem, we need to explain how the nerve can be viewed as a functor. To this end, we will define the category of covered spaces.

To motivate our definition, we first briefly discuss a variant that is also common in the literature (see, e.g., [Bar02]). The objects in this category are of the form (X, \mathcal{U}) , where X is a topological space and \mathcal{U} is an unindexed cover of X . A map between covered spaces $f: (X, \mathcal{U}) \rightarrow (Y, \mathcal{V})$ is then given by a continuous map $f: X \rightarrow Y$ such that for any cover element $U \in \mathcal{U}$ there exists $V \in \mathcal{V}$ with $f(U) \subseteq V$. Choosing such a cover element $V \in \mathcal{V}$ for every element $U \in \mathcal{U}$ determines a simplicial map $\text{Nrv}(\mathcal{U}) \rightarrow \text{Nrv}(\mathcal{V})$ between the nerves. In general, different choices give different simplicial maps, but it will always be unique up to contiguity (see [Mun84, p. 67] for a definition). In particular, it follows that any two choices determine, up to homotopy, the same map on the geometric realization.

To avoid having to make choices, we work with indexed covers and record the choice of cover elements as above in a map between the indexing sets. This way, we circumvent the ambiguity of the induced map between the nerves up to homotopy.

Definition 1.1. Let X and Y be topological spaces, $(U_i)_{i \in I}$ a cover of X , and $(V_\ell)_{\ell \in L}$ a cover of Y . A map of indexed covers $\varphi: (U_i)_{i \in I} \rightarrow (V_\ell)_{\ell \in L}$ is defined formally as a map $\varphi: I \rightarrow L$ between the indexing sets. We say that a continuous map $f: X \rightarrow Y$ is carried by φ if for all $i \in I$ we have $f(U_i) \subseteq V_{\varphi(i)}$.

If f is carried by φ and g is carried by ψ , then $g \circ f$ is carried by $\psi \circ \varphi$ if the compositions are defined. Hence, we get the following category.

Definition 1.2. The *category of covered spaces* Cov has objects pairs of the form $(X, (U_i)_{i \in I})$, where X is a topological space and $(U_i)_{i \in I}$ is a cover of X . A morphism $(f, \varphi): (X, (U_i)_{i \in I}) \rightarrow (Y, (V_\ell)_{\ell \in L})$ between two covered spaces consists of a continuous map $f: X \rightarrow Y$ and a map of indexed covers $\varphi: (U_i)_{i \in I} \rightarrow (V_\ell)_{\ell \in L}$ such that f is carried by φ .

With this category in hand, we can define a functor $\text{Spc}: \text{Cov} \rightarrow \text{Top}$ by forgetting the cover: Spc takes a pair $(X, (U_i)_{i \in I})$ to X . By taking the geometric realization of the nerve of a cover, we obtain another such functor. We denote by Simp the category of simplicial complexes.

Definition 1.3. Let X be a topological space, and let $\mathcal{U} = (U_i)_{i \in I}$ be a cover of X . For any $J \subseteq I$, we write $U_J = \bigcap_{i \in J} U_i$. The *nerve* of \mathcal{U} is the simplicial complex $\text{Nrv}(\mathcal{U})$ with simplices

$$\{J \subseteq I \mid |J| < \infty \text{ and } U_J \neq \emptyset\}.$$

A morphism of covered spaces $(f, \varphi): (X, \mathcal{U}) \rightarrow (Y, \mathcal{V})$ induces a simplicial map between the nerves of the covers $\varphi_*: \text{Nrv}(\mathcal{U}) \rightarrow \text{Nrv}(\mathcal{V})$. Thus, the nerve can be seen to be a functor $\text{Nrv}: \text{Cov} \rightarrow \text{Simp}$. By postcomposing this with the geometric realization functor $|\cdot|: \text{Simp} \rightarrow \text{Top}$, we get the functor $|\text{Nrv}|: \text{Cov} \rightarrow \text{Top}$ that takes a pair (X, \mathcal{U}) to the geometric realization $|\text{Nrv}(\mathcal{U})|$.

Remark 1.4. There is a variant of this definition, where the vertex set is the set of cover elements, in contrast to our definition, where it is the indexing set. While these definitions yield different simplicial complexes in general, as the same subset can appear multiple times in the indexed cover, they are always homotopy equivalent. More precisely, if $\mathcal{U} = (U_i)_{i \in I}$ is an indexed cover and $U_j \subseteq U_l$, with $j \neq l$, are cover elements, then the inclusion $\text{Nrv}(\mathcal{V}) \hookrightarrow \text{Nrv}(\mathcal{U})$ is a homotopy equivalence, where $\mathcal{V} = (U_i)_{i \in I \setminus \{j\}}$: The link $\text{lk}(j) = \{\sigma \in \text{Nrv}(\mathcal{U}) \mid j \notin \sigma, \sigma \cup \{j\} \in \text{Nrv}(\mathcal{U})\}$ of the vertex j in $\text{Nrv}(\mathcal{U})$ is a cone with apex l , i.e., for all $\sigma \in \text{lk}(j)$ we have $\sigma \cup \{l\} \in \text{Nrv}(\mathcal{U})$. Therefore, there exists a collapse $\text{Nrv}(\mathcal{U}) \searrow \text{Nrv}(\mathcal{V})$ (see Appendix B for a definition).

Now that we can understand the covered space and the nerve as functors, we can consider natural transformations that relate them. In general, if F_1 and F_2 are functors from some category \mathcal{C} to \mathbf{Top} , and $\sigma: F_1 \Rightarrow F_2$ is a natural transformation, one says that σ is a pointwise homotopy equivalence if the component $\sigma_C: F_1(C) \rightarrow F_2(C)$ is a homotopy equivalence for all objects C of \mathcal{C} . Similarly one can consider pointwise weak homotopy equivalences, pointwise homology isomorphisms, et cetera. This paper is about nerve theorems that relate the covered space and the nerve through pointwise equivalences.

Most of these nerve theorems make use of a standard construction that is called the *blowup complex* by Zomorodian–Carlsson [ZC08], but goes back at least to Segal [Seg68]. It is a functor $\text{Blowup}: \mathbf{Cov} \rightarrow \mathbf{Top}$, along with natural transformations $\rho_S: \text{Blowup} \Rightarrow \text{Spc}$ and $\rho_N: \text{Blowup} \Rightarrow |\text{Nrv}|$. In particular, for any morphism of covered spaces $(f, \varphi): (X, \mathcal{U}) \rightarrow (Y, \mathcal{V})$ there exists a commuting diagram of the following form:

$$\begin{array}{ccccc} X & \xleftarrow{\rho_S} & \text{Blowup}(\mathcal{U}) & \xrightarrow{\rho_N} & |\text{Nrv}(\mathcal{U})| \\ f \downarrow & & \downarrow & & \downarrow |\varphi_*| \\ Y & \xleftarrow{\rho_S} & \text{Blowup}(\mathcal{V}) & \xrightarrow{\rho_N} & |\text{Nrv}(\mathcal{V})| \end{array} \quad (1.5)$$

We write $[n]$ for the set $\{0, \dots, n\}$. If $\mathcal{U} = (U_i)_{i \in [n]}$ is a finite cover of a space X , then the blowup complex is

$$\text{Blowup}(\mathcal{U}) = \bigcup_{J \in \text{Nrv}(\mathcal{U})} U_J \times \Delta^J \subseteq X \times \Delta^n,$$

where Δ^n is the standard topological n -simplex and Δ^J is a face of Δ^n determined by the inclusion $J \subseteq [n]$. The idea is that each piece of X expands according to the number of cover elements that contain it. See Section 2 for the definition for arbitrary covers.

To begin, we give three functorial nerve theorems (Theorems A, B and C below) whose proofs will be particularly attractive to students and newcomers to TDA. This is because the arguments are relatively elementary and use techniques that are interesting in their own right. In Section 3, we prove the following functorial nerve theorem:

Theorem A. *If $X \subset \mathbb{R}^d$, and $\mathcal{B} = (C_i)_{i \in [n]}$ is a cover by closed convex subsets, then the natural maps $\rho_S: \text{Blowup}(\mathcal{B}) \rightarrow X$ and $\rho_N: \text{Blowup}(\mathcal{B}) \rightarrow |\text{Nrv}(\mathcal{B})|$ are homotopy equivalences.*

The proof uses partitions of unity, and is similar to the strategy for open covers in Hatcher's textbook. In Section 3, we also prove a functorial nerve theorem for closed convex covers that does not require any intermediate object, subject to an additional assumption on the morphisms of covered spaces.

Before we state this theorem, we elaborate shortly on why such a functorial nerve theorem cannot exist in general. The reason is simple: there are no natural transformations between Spc and $|\text{Nrv}|$ in either direction. Consider the covered spaces $(*, (*))$, where $*$ is the one-point space, and $(Y, (Y))$, where $Y \neq *$ is any space. For any point $p \in Y$ the inclusion $\iota_p: * \hookrightarrow Y$ gives rise to a morphism of covered space $(\iota_p, * \mapsto Y)$. If there existed a natural transformation $|\text{Nrv}| \Rightarrow \text{Spc}$, then this would already fix a single inclusion $\iota_q: * \hookrightarrow Y$ as part of such a morphism of covered spaces, implying that $Y = \{q\}$ is a single point, yielding a contradiction. Similarly, consider any covered space $(Z, (U, V))$ with $p \in U \cap V$ any point. Consider the two morphisms of covered spaces $(\iota_p, * \mapsto U), (\iota_p, * \mapsto V): (*, (*)) \rightarrow (Z, (U, V))$. Then, these maps induce different simplicial maps on the nerves, implying that there exists no natural transformation $\text{Spc} \Rightarrow |\text{Nrv}|$.

Thus, in order to obtain a functorial nerve theorem that does not need an intermediate object, the map of indexed covers needs to have strong combinatorial control on the continuous map. To this end, we introduce the following notions.

Definition 1.5. A *pointed cover* $\mathcal{U}_* = (\mathcal{U} = (U_i)_{i \in I}, (u_\sigma)_{\sigma \in \text{Nrv}(\mathcal{U})})$ of a topological space X is a cover \mathcal{U} of X together with a point $u_\sigma \in U_\sigma$ for every $\sigma \in \text{Nrv}(\mathcal{U})$.

The *category of pointed covered spaces* Cov_* has objects tuples of the form (X, \mathcal{A}_*) , where X is a topological space and $\mathcal{A}_* = (\mathcal{A}, (a_\sigma)_{\sigma \in \text{Nrv}(\mathcal{A})})$ is a pointed cover of X . A morphism $(f, \varphi): (X, \mathcal{A}_*) \rightarrow (Y, \mathcal{B}_*)$ of pointed covered spaces is a morphism of covered spaces $(f, \varphi): (X, \mathcal{A}) \rightarrow (Y, \mathcal{B})$ that respects the basepoints, i.e., such that for any $\sigma \in \text{Nrv}(\mathcal{A})$ we have $f(a_\sigma) = b_{\varphi_*(\sigma)}$.

There is an obvious functor $\text{Cov}_* \rightarrow \text{Cov}$ that forgets the pointing, and hence we get, as for the category of covered space, the functors $\text{Spc}: \text{Cov}_* \rightarrow \text{Top}$ and $|\text{Sd Nrv}|: \text{Cov}_* \rightarrow \text{Top}$, where Sd Nrv is the subdivision of the nerve.

Now, we will describe a functorial nerve theorem that does not require an intermediate object. The subcategory ClConv_* of Cov_* consists of subsets of \mathbb{R}^d that are covered by finitely many closed convex sets. Further, we restrict to morphisms of pointed covered spaces whose underlying continuous maps are affine linear on each cover element. Many covers of interest in TDA are pointed.

Example 1.6. Let $\{x_0, \dots, x_n\} \subseteq \mathbb{R}^d$ be a finite set of points. Then, we can point the cover $\mathcal{U}_r = (D_r(x_i))_{i \in [n]}$ of the union of closed balls $O_r = \bigcup_{i=0}^n D_r(x_i)$ in the following way: For each non-empty subset $\sigma \subseteq [n]$ there exists a smallest real number r_σ such that the intersection $(\mathcal{U}_{r_\sigma})_\sigma$ is non-empty. We define the point p_σ to be the unique point in this intersection. This gives the pointed cover $(\mathcal{U}_r, (p_\sigma)_{\sigma \in \text{Nrv}(\mathcal{U}_r)})$ of O_r for each $r \in \mathbb{R}_{\geq 0}$. With this at hand, we see that the offset filtration is a functor $\mathbb{R}_{\geq 0} \rightarrow \text{ClConv}_*$.

Theorem B. *For every pointed covered space $(X, \mathcal{A}_*) \in \text{ClConv}_*$ there exists a homotopy equivalence*

$$\Gamma: |\text{Sd Nrv}(\mathcal{A})| \rightarrow X$$

that is natural with respect to the morphisms in ClConv_ .*

One says that a cover is *good* if all non-empty finite intersections of cover elements are contractible. As we have already mentioned, nerve theorems usually assume that the covers involved are good. Moreover, a cover is *locally finite* if every point has an open neighborhood that intersects only finitely many cover elements. In Section 4, we again use the blowup complex to prove a functorial nerve theorem for simplicial complexes:

Theorem C. *Let K be a simplicial complex and let $\mathcal{A} = (K_i \subseteq K)_{i \in I}$ be a locally finite good cover of K by subcomplexes. The natural maps $\rho_S: \text{Blowup}(|\mathcal{A}|) \rightarrow |K|$ and $\rho_N: \text{Blowup}(|\mathcal{A}|) \rightarrow |\text{Nrv}(\mathcal{A})|$ are homotopy equivalences.*

The proof is related to work of Björner [Bjö03], and uses elementary methods from combinatorial homotopy theory for constructing homotopy equivalences between simplicial complexes, together with discrete Morse theory. Combining this result with a well-known theorem on triangulations of semi-algebraic sets (Theorem 4.10), we obtain a nerve theorem for compact semi-algebraic sets that are covered by finitely many closed, semi-algebraic subspaces. Moreover, in Appendix C, we show that one can remove the assumption that the cover \mathcal{A} is locally finite by using contractible carriers for the proof.

Finally we use techniques from abstract homotopy theory to prove the following omnibus functorial nerve theorem. In particular, this result implies Theorems A and C.

In parts 1(b) and 2(b) of Theorem D, we restrict attention to compactly-generated spaces. This is a standard hypothesis in algebraic topology, as these spaces form a “convenient” subcategory of topological spaces that is suitable for developing the machinery of homotopy theory. In part 1(b), the intersection A_T and the *latching space* $L(T) = \bigcup_{T \subsetneq J} A_J \subseteq A_T$ are assumed to satisfy the homotopy extension property; for example, CW-pairs satisfy the homotopy extension property (cf. Remark 5.8). These assumptions on the latching spaces together with the assumption that the cover is locally finite dimensional allow for inductive arguments analogous to arguments that employ induction over the skeleton of a CW-complex.

At the beginning of Section 5 we introduce in detail all of the notions used in the statement of the theorem.

Theorem D. *Let X be a topological space and let $\mathcal{A} = (A_i)_{i \in I}$ be a cover of X .*

1. *Consider the natural map $\rho_S: \text{Blowup}(\mathcal{A}) \rightarrow X$.*
 - (a) *If \mathcal{A} is an open cover, then ρ_S is a weak homotopy equivalence. If furthermore X is paracompact, then ρ_S is a homotopy equivalence.*
 - (b) *Assume that X is compactly generated and that \mathcal{A} is a closed cover that is locally finite and locally finite dimensional. If for any $T \in \text{Nrv}(\mathcal{A})$ the latching space $L(T) \subseteq A_T$ is a closed subset and the pair $(A_T, L(T))$ satisfies the homotopy extension property, then ρ_S is a homotopy equivalence.*

2. Consider the natural map $\rho_N: \text{Blowup}(\mathcal{A}) \rightarrow |\text{Nrv}(\mathcal{A})|$.
 - (a) If \mathcal{A} is (weakly) good, then ρ_N is a (weak) homotopy equivalence.
 - (b) If for all $J \in \text{Nrv}(\mathcal{A})$ the space A_J is compactly generated and \mathcal{A} is homologically good with respect to a coefficient ring R , then ρ_N is an R -homology isomorphism.

We now summarize the ingredients that go into the proof. First of all, the blowup complex is closely related to another standard construction, called the bar construction. While the blowup complex has a natural map to the nerve of a cover, the bar construction has instead a natural map to the subdivision of the nerve, which is why we use the blowup complex in the statements of the theorems. However, the bar construction is in some ways easier to work with; in Section 2 we define the bar construction and explain how we can work with either the blowup complex or the bar construction, whichever is more convenient.

The statement in 1(a) about weak homotopy equivalences follows from work of Dugger–Isaksen [DI04]; we give a short proof in Appendix D. The statement in 1(a) about homotopy equivalences is proved in Hatcher’s textbook [Hat02, Proposition 4G.2]. Statement 1(b) follows from a standard argument using Reedy model structures, which is similar to the proof of [Dug, Corollary 14.17], for example. Both parts of 2(a) follow from the fact that the bar construction is homotopical, both for homotopy equivalences and weak homotopy equivalences. In the case of weak homotopy equivalences, this again uses Dugger–Isaksen [DI04]; see Proposition 5.42. We also give a short proof of this in Appendix D. Finally, 2(b) is proved using the bar construction in the setting of simplicial R -modules.

In summary, the proof of Theorem D is straightforward, given some powerful tools for studying diagrams of spaces from abstract homotopy theory. In Section 5 we provide an introduction to these tools. We hope this will be useful to interested members of the applied topology community.

The Literature on Nerve Theorems We now summarize the literature on the nerve theorem, with a particular focus on results that address functoriality.

The original work of Alexandroff [Ale28] on nerves, as well as the early nerve theorems of Leray [Ler45, Théorème 12] and Borsuk [Bor48, Corollary 3], considered closed covers, motivated in part by covers of polytopes by simplices. Open covers were considered by Weil [Wei52, Section 5], McCord [McC67], and Segal [Seg68]. There is renewed interest in the case of closed covers in applied topology, motivated by geometric constructions such as alpha shapes [EKS83].

A common way to relate the nerve of an open cover \mathcal{U} with the covered space X is by a partition of unity subordinate to the cover. Such a partition of unity defines a map from X to the nerve of \mathcal{U} in a straightforward way, which is a homotopy equivalence if \mathcal{U} is good. This idea appears in the work of Weil and Segal, and the textbook proofs of the nerve theorem by Hatcher [Hat02, Corollary 4G.3] and Kozlov [Koz08, Theorem 15.21]. Moreover, this map commutes up to homotopy with the maps on space and nerve

induced by a morphism of covered spaces. This was observed by Chazal–Oudot [CO08] (for certain inclusions of covered spaces) and by Bauer–Edelsbrunner–Jabłoński–Mrozek [Bau+20] (for general morphisms). See also Eilenberg–Steenrod [ES52, Ch. X.11], Lim–Mémoli–Okutan [LMO20] and Virk [Vir21].

In the case of open covers of paracompact spaces, Botnan–Spremann [BS15], following the approach that goes back at least to Segal, observe that the blowup complex provides a zigzag of natural transformations relating covered space and nerve.

Ferry–Mischaikow–Nanda [FMN14] consider covers by open and closed balls in Euclidean space; they also use the blowup complex as an intermediate object relating space and nerve, but use the Vietoris–Smale theorem on proper maps with contractible fibers to obtain a homotopy equivalence from nerve to space with control over the image of each simplex.

Bendich–Cohen–Steiner–Edelsbrunner–Harer–Morozov [Ben+07] give a nerve theorem for certain closed convex covers in Euclidean space; they define a map from the subdivision of the nerve of \mathcal{U} to the space by choosing a point in each non-empty intersection \mathcal{U}_J , mapping the vertex J to this point, and extending by piecewise linear interpolation. They show this map commutes up to homotopy with maps induced by certain morphisms of covered spaces.

Borel–Serre [BS73, Theorem 8.2.1] prove a nerve theorem for locally finite and closed covers whose nerve is finite dimensional and such that all finite non-empty intersections of cover elements are absolute retracts for metric spaces. Using similar techniques, Nagórko [Nag07] proves a nerve theorem for locally finite, locally finite dimensional, star-countable closed covers of normal spaces such that all non-empty intersections of cover elements are absolute extensors for metric spaces.

Björner [Bjö03] gives a proof of an n -connectivity version of the nerve theorem, which we discuss in Section 4.3. Given a good cover of a finite simplicial complex by subcomplexes, Barmak [Bar11] proves a related result, showing that the simplicial complex and the nerve have the same simple homotopy type.

2. Functorial Nerve Theorems via Homotopy Colimits

Theorems A, C, and D use the *blowup complex* $\text{Blowup}(\mathcal{U})$ of a cover \mathcal{U} as an intermediate object to relate the nerve of \mathcal{U} with the covered space. In this section, we define the blowup complex and its natural maps to the covered space and the nerve. The construction is not difficult, but there is an important point here: the blowup complex is closely related to the *bar construction*, and because of this, properties of the bar construction are used in many proofs of the nerve theorem, including Theorems A and D.

The bar construction is a standard model for the homotopy colimit: like the colimit, the homotopy colimit can be defined via a universal property, but since this universal property is phrased in terms of derived categories, it takes some work to define it precisely. A full discussion of the homotopy colimit is beyond the scope of this paper (see [Dug] for a nice introduction to the topic, or [Rie14, Part I] for a more abstract approach). However, in order to explain the properties of the bar construction that we

will use, we will at least describe the problem that the homotopy colimit addresses. So, in this section we will introduce a basic problem with the colimit of a diagram of topological spaces, give an idea of how the homotopy colimit addresses this problem, define the bar construction and the blowup complex, and explain how the properties of the bar construction can be used to prove functorial nerve theorems using the blowup complex.

First we establish some notation and recall some facts related to the barycentric subdivision of a simplicial complex. Write \mathbf{Po} for the category of posets, and \mathbf{Simp} for the category of simplicial complexes. Let $\text{Pos}: \mathbf{Simp} \rightarrow \mathbf{Po}$ be the functor that takes a simplicial complex to its poset of simplices (ordered by inclusion), and let $\text{Flag}: \mathbf{Po} \rightarrow \mathbf{Simp}$ be the functor that takes a poset P to the simplicial complex whose vertices are the elements of P and whose n -simplices are the chains $x_0 < \cdots < x_n$ of elements of P . Then the barycentric subdivision of a simplicial complex K is $\text{Sd}(K) = \text{Flag}(\text{Pos}(K))$. There is an affine linear homeomorphism $\alpha_K: |\text{Sd}(K)| \rightarrow |K|$ defined by the vertex map that sends a vertex σ of $\text{Sd}(K)$ to the barycenter of $|\sigma|$ in $|K|$. Note that, while the homeomorphism α_K is natural with respect to inclusions of simplicial complexes, it is not natural with respect to general simplicial maps.

2.1. Homotopy Colimits and the Bar Construction

While colimits are used everywhere in topology to construct new spaces, the colimit operation fails to respect homotopy equivalences, in the following sense. Take \mathcal{A} to be the category that looks like this:

$$\bullet \leftarrow \bullet \rightarrow \bullet$$

and consider the commutative diagram:

$$\begin{array}{ccccc} D^n & \longleftarrow & S^{n-1} & \longrightarrow & D^n \\ \downarrow & & \downarrow \text{id} & & \downarrow \\ * & \longleftarrow & S^{n-1} & \longrightarrow & * \end{array}$$

Here, the top maps are the boundary inclusions. We think of the rows as \mathcal{A} -shaped diagrams, and the vertical maps define a natural transformation between these two \mathcal{A} -shaped diagrams. Every component of this natural transformation is a homotopy equivalence, but the colimit of the top row is the sphere S^n , while the colimit of the bottom row is a one-point space $*$, so the induced map between the colimits cannot be a homotopy equivalence.

More generally, let \mathcal{C} be a small category, and write $\mathbf{Top}^{\mathcal{C}}$ for the category of functors $\mathcal{C} \rightarrow \mathbf{Top}$. One says that a functor $\Omega: \mathbf{Top}^{\mathcal{C}} \rightarrow \mathbf{Top}$ is *homotopical* if, given a natural transformation $\lambda: F \Rightarrow F'$ between \mathcal{C} -shaped diagrams F and F' that is a pointwise homotopy equivalence, the induced map $\Omega(F) \rightarrow \Omega(F')$ is also a homotopy equivalence. For any small category \mathcal{C} , the colimit defines a functor $\text{colim}: \mathbf{Top}^{\mathcal{C}} \rightarrow \mathbf{Top}$, and the previous example shows that this functor is not homotopical in general.

A homotopy colimit is a homotopical functor $\text{hocolim}: \mathbf{Top}^{\mathcal{C}} \rightarrow \mathbf{Top}$, together with a natural transformation $\text{hocolim} \Rightarrow \text{colim}$ that makes hocolim , in some sense, the

best possible homotopical approximation of the colimit functor. We now show how to construct a particular model for the homotopy colimit, called the bar construction, and we will see that it can be thought of as a “thickened” version of the colimit; see Example 2.5 for an illustration.

We write Δ^n for the standard topological n -simplex, and for $0 \leq i \leq n$, we write $d^i: \Delta^{n-1} \hookrightarrow \Delta^n$ for the inclusion of the face opposite the i^{th} vertex.

Definition 2.1. Let P be a poset, and let $F: P \rightarrow \mathbf{Top}$ be a diagram of topological spaces. The *bar construction* $\text{Bar}(P, F)$ of F is the quotient space

$$\text{Bar}(P, F) = \left(\bigsqcup_{\sigma=(v_0 < \dots < v_n)} F(v_0) \times \Delta^n \right) / \sim$$

where the disjoint union is taken over all chains in P , and the equivalence relation \sim is generated as follows. For a chain $\sigma = (v_0 < \dots < v_n)$ and $0 \leq i \leq n$, we write

$$\tau_i = (v_0 < \dots < \hat{v}_i < \dots < v_n) = (w_0 < \dots < w_{n-1}),$$

noting that w_0 equals v_0 for $i > 0$ and v_1 for $i = 0$. Now for any $x \in F(v_0)$ and $\alpha \in \Delta^{n-1}$, we identify $(x, d^i(\alpha))$ in the copy of $F(v_0) \times \Delta^n$ indexed by σ with $(F(v_0 \leq w_0)(x), \alpha)$ in the copy of $F(w_0) \times \Delta^{n-1}$ indexed by τ_i .

Example 2.2. Let $P = \{0 < 1\}$. Then a diagram $F: P \rightarrow \mathbf{Top}$ is just a map $F(0) \rightarrow F(1)$, and the bar construction $\text{Bar}(P, F)$ is the mapping cylinder of this map.

Definition 2.3. Let X be a topological space and $\mathcal{U} = (U_i)_{i \in I}$ a cover of X . Writing $P_{\mathcal{U}} = \text{Pos}(\text{Nrv}(\mathcal{U}))^{\text{op}}$, the *nerve diagram* of the cover \mathcal{U} is the functor $\mathcal{D}_{\mathcal{U}}: P_{\mathcal{U}} \rightarrow \mathbf{Top}$ with $\mathcal{D}_{\mathcal{U}}(J) = U_J$.

Remark 2.4. In many cases, the colimit of the diagram $\mathcal{D}_{\mathcal{U}}$ simply gives us back X : the inclusions $U_J \subseteq X$ induce a continuous map $\text{colim } \mathcal{D}_{\mathcal{U}} \rightarrow X$, which is in fact a bijection. If \mathcal{U} is an open cover, or if it is a closed cover that is locally finite (i.e., every point of X has an open neighborhood that intersects only finitely many cover elements), then this bijection is a homeomorphism.

Example 2.5. In this paper, we will mainly consider bar constructions of diagrams associated to a cover. For example, consider the following cover \mathcal{U} of the circle S^1 :





Figure 2.: A cover by three arcs (left) and the intersections of those (right).

If we label the three arcs $\{b, r, g\}$, the poset $P_{\mathcal{U}}$ associated to this cover has the following form:



By definition, the bar construction $\text{Bar}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}})$ of the nerve diagram $\mathcal{D}_{\mathcal{U}}: P_{\mathcal{U}} \rightarrow \text{Top}$ associated to the cover \mathcal{U} is built from pieces indexed by chains $v_0 < \dots < v_n$ in $P_{\mathcal{U}}$ and are of the form $\mathcal{D}_{\mathcal{U}}(v_0) \times \Delta^n$. More concretely, the bar construction in our example is built from the following pieces:



Figure 3.: Pieces indexed by chains in $P_{\mathcal{U}}$ of length zero (left) and of length one (right).

After making all identifications, the bar construction $\text{Bar}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}})$ is the following “thickened” version of $\text{colim } \mathcal{D}_{\mathcal{U}} \cong S^1$:




Figure 4.: The bar construction of the nerve diagram.

By $\text{Diag}_{\text{Po}}(\text{Top})$ we denote the category of diagrams over a poset: the objects are tuples (P, F) , where P is a poset and $F: P \rightarrow \text{Top}$ is a functor. A morphism $(g, \lambda): (P, F) \rightarrow (R, G)$ consists of a poset map $g: P \rightarrow R$ and a natural transformation $\lambda: F \Rightarrow G \circ g$. Then the bar construction defines a functor $\text{Bar}: \text{Diag}_{\text{Po}}(\text{Top}) \rightarrow \text{Top}$: a morphism (g, λ) induces a continuous map $\text{Bar}(P, F) \rightarrow \text{Bar}(R, G)$ defined by the maps

$$\lambda(v_0) \times |\text{Flag}(g)|: F(v_0) \times \Delta^n \rightarrow G(g(v_0)) \times \Delta^m,$$

where $|\text{Flag}(g)|: \Delta^n \rightarrow \Delta^m$ is the affine map that sends the vertex v_i to $g(v_i)$. Moreover, the projection maps $F(v_0) \times \Delta^n \rightarrow F(v_0)$ define a natural map $\text{Bar}(P, F) \rightarrow \text{colim } F$.

There are analogues of this quotient space construction in other settings, which are also called bar constructions. We will encounter some of these in Section 5. For a very general discussion of the bar construction, including a proof that it is a model of the homotopy colimit, see [Rie14, Chapters 4–5]. In particular, the bar construction is homotopical (see [Koz08, Theorem 15.12] for a proof in a slightly more general setting):

Proposition 2.6. *Let P be a poset, $F, F': P \rightarrow \text{Top}$ diagrams of topological spaces, and let $\lambda: F \Rightarrow F'$ be a natural transformation. If the component $\lambda(v)$ is a homotopy equivalence for all $v \in P$, then so is the induced map $\text{Bar}(P, F) \rightarrow \text{Bar}(P, F')$.*

2.2. Functorial Nerve Theorems via the Bar Construction

We can now explain how the bar construction can be used to prove functorial nerve theorems. For any poset P , if we write $*$ for the constant diagram $P \rightarrow \text{Top}$ to the one-point space, there is a canonical identification $\text{Bar}(P, *) \cong |\text{Flag}(P)|$. In particular, if $P_{\mathcal{U}}$ is the poset associated to a cover \mathcal{U} , we have $\text{Bar}(P_{\mathcal{U}}, *) \cong |\text{Flag}(P_{\mathcal{U}})| = |\text{Sd Nrv}(\mathcal{U})|$.

A morphism of covered spaces $(f, \varphi): (X, \mathcal{U}) \rightarrow (Y, \mathcal{V})$ induces a poset map $g: P_{\mathcal{U}} \rightarrow P_{\mathcal{V}}$ and a natural transformation $\lambda: \mathcal{D}_{\mathcal{U}} \Rightarrow \mathcal{D}_{\mathcal{V}} \circ g$. Thus, by what we have seen before, the operation $(X, \mathcal{U}) \mapsto \text{Bar}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}})$ defines a functor $\text{Cov} \rightarrow \text{Diag}_{\text{Po}}(\text{Top}) \rightarrow \text{Top}$. Moreover, the unique natural transformation $\mathcal{D}_{\mathcal{U}} \Rightarrow *$ induces a natural map

$$\pi_{\text{SdN}}: \text{Bar}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}}) \rightarrow \text{Bar}(P_{\mathcal{U}}, *) \cong |\text{Sd Nrv}(\mathcal{U})|.$$

If every non-empty finite intersection of cover elements happens to be contractible, then this map is a homotopy equivalence by Proposition 2.6. Using also the natural map from the bar construction to the colimit as mentioned at the end of Section 2.1, we get a diagram that is natural in morphisms of covered spaces:

$$X \leftarrow \text{colim } \mathcal{D}_{\mathcal{U}} \leftarrow \text{Bar}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}}) \rightarrow |\text{Sd Nrv}(\mathcal{U})|.$$

In Section 3 and Section 5 we will use this diagram to prove functorial nerve theorems by finding various sets of assumptions that make these natural maps equivalences of various kinds. This strategy – also employed in the well-known proof of the nerve theorem for open covers in Hatcher’s textbook [Hat02, Section 4.G] – relies on the well-known good properties of the bar construction. We exploit this established theory repeatedly in Section 5, where we use the fact that the bar construction is homotopical in several contexts, including homological algebra. In Section 4, we will prove a functorial nerve theorem using a bar construction constructed in the category of posets, rather than the topological construction.

However, for purposes of computational topology, we want a nerve theorem to relate the space X directly with the nerve of \mathcal{U} , not the much larger subdivision of the nerve. In order to obtain a functorial nerve theorem that works for morphisms of covered spaces as we have defined them, in which the map of indexed covers need not be an inclusion,

we cannot simply apply the usual homeomorphism $\alpha_K: |\mathrm{Sd}(K)| \rightarrow |K|$ defined for any simplicial complex K , as this map is natural only in inclusions of simplicial complexes. In the case of diagrams $\mathcal{D}_{\mathcal{U}}$ associated to a cover, the *blowup complex* is an efficient way to build a space homeomorphic to the bar construction, which comes with a natural map to $|\mathrm{Nrv}(\mathcal{U})|$ rather than $|\mathrm{Sd}\mathrm{Nrv}(\mathcal{U})|$. In the definition, for a non-empty finite set J , we write $|J|$ for the geometric realization of the full simplicial complex generated by J , which is homeomorphic to the standard topological simplex $\Delta^{|J|-1}$.

Definition 2.7. Let $\mathcal{U} = (U_i)_{i \in I}$ be a cover of a topological space X . The *blowup complex* $\mathrm{Blowup}(\mathcal{U})$ is the quotient space

$$\mathrm{Blowup}(\mathcal{U}) = \left(\bigsqcup_{J \in \mathrm{Nrv}(\mathcal{U})} U_J \times |J| \right) / \sim,$$

where the disjoint union is taken over all simplices $J \in \mathrm{Nrv}(\mathcal{U})$, and the equivalence relation \sim identifies, for all $J \subseteq J'$, the spaces $U_J \times |J|$ and $U_{J'} \times |J'|$ along their common subspace $U_{J'} \times |J|$.

Remark 2.8. If the cover $\mathcal{U} = (U_i)_{i \in [n]}$ is finite, the blowup complex can be defined as a subspace of the product $X \times \Delta^n$, as mentioned in the introduction. This is the approach used by Zomorodian and Carlsson [ZC08, Definition 3].

As before, the operation $(X, \mathcal{U}) \mapsto \mathrm{Blowup}(\mathcal{U})$ defines a functor $\mathrm{Cov} \rightarrow \mathrm{Top}$, and the projection maps $U_J \times |J| \rightarrow U_J$ define a natural map $\mathrm{Blowup}(\mathcal{U}) \rightarrow \mathrm{colim} \mathcal{D}_{\mathcal{U}}$, which gives us a natural map $\rho_S: \mathrm{Blowup}(\mathcal{U}) \rightarrow X$. But now the projection maps $U_J \times |J| \rightarrow |J|$ assemble to define a natural map $\rho_N: \mathrm{Blowup}(\mathcal{U}) \rightarrow |\mathrm{Nrv}(\mathcal{U})|$.

Example 2.9. As in Example 2.5, we consider the cover of the circle by three arcs.




Figure 5.: The bar construction (left) and the blowup complex (right).

Note that the blowup complex is combinatorially simpler.

We can use the homeomorphism $\alpha_K: |\mathrm{Sd}(K)| \rightarrow |K|$ defined for any simplicial complex K to construct a homeomorphism $\mathrm{Bar}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}}) \rightarrow \mathrm{Blowup}(\mathcal{U})$. For any simplex $J \in \mathrm{Nrv}(\mathcal{U})$, any flag $J \supset J_1 \supset \dots \supset J_n$ in $P_{\mathcal{U}}$ indexes a piece $U_J \times \Delta^n$ in $\mathrm{Bar}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}})$. The flags of this form glue together to give a copy of $U_J \times |\mathrm{Sd} J|$ inside $\mathrm{Bar}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}})$, where $\mathrm{Sd} J$ is the subdivision of the simplicial complex generated by J . Now for all $J \in \mathrm{Nrv}(\mathcal{U})$ we have a map

$$U_J \times |\mathrm{Sd} J| \xrightarrow{\alpha_J \times \mathrm{id}} U_J \times |J| \subset \mathrm{Blowup}(\mathcal{U}),$$

and assembling these maps gives the homeomorphism $\text{Bar}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}}) \rightarrow \text{Blowup}(\mathcal{U})$.

This homeomorphism is not natural in arbitrary morphisms of covered spaces, but it does fit into the following commutative diagram, where the solid arrows are natural in morphisms of covered spaces:

$$\begin{array}{ccccc}
 X & \xleftarrow{\pi_S} & \text{Bar}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}}) & \xrightarrow{\pi_{\text{Sd}N}} & |\text{Sd Nrv}(\mathcal{U})| \\
 \parallel & & \downarrow \cong & & \downarrow \cong \\
 X & \xleftarrow{\rho_S} & \text{Blowup}(\mathcal{U}) & \xrightarrow{\rho_N} & |\text{Nrv}(\mathcal{U})|
 \end{array} \tag{2.1}$$

The somewhat subtle point here is that, even though the homeomorphism $\text{Bar}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}}) \rightarrow \text{Blowup}(\mathcal{U})$ is not natural in arbitrary morphisms of covered spaces, we can use this homeomorphism and the good properties of the bar construction to prove functorial nerve theorems for the blowup complex: if some set of assumptions on the covered space (X, \mathcal{U}) imply that the top maps from $\text{Bar}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}})$ are equivalences of some kind, then the commutativity of the diagram 2.1 implies that the bottom maps from $\text{Blowup}(\mathcal{U})$ are equivalences of the same kind.

3. Nerve Theorems for Closed Convex Covers

In this section, we consider nerves of finite closed and convex covers of subsets of \mathbb{R}^d . The motivating instances are alpha complexes and Čech complexes of finite point sets. Our approach to prove a nerve theorem in this context is elementary and does not make explicit use of abstract homotopy-theoretic machinery. Nevertheless, it foreshadows the concepts that will take on a central role in Section 5.

We write $[n]$ for the set $\{0, \dots, n\}$. Let $\mathcal{C} = (C_i)_{i \in [n]}$ be a collection of closed convex subsets of \mathbb{R}^d , and let X be their union. We construct a continuous map $\Gamma: |\text{Sd Nrv}(\mathcal{C})| \rightarrow X$ and show that this map is a homotopy equivalence, establishing a nerve theorem for this setting. We then extend this result to prove the two functorial versions discussed in the introduction, Theorem A and Theorem B.

Each vertex $J \in \text{Sd Nrv}(\mathcal{C})$ represents a simplex in the nerve $\text{Nrv}(\mathcal{C})$, and hence we can choose a point p_J from the non-empty intersection $C_J = \bigcap_{j \in J} C_j$. By convexity of the cover elements in \mathcal{C} , this choice extends uniquely to a map $\Gamma: |\text{Sd Nrv}(\mathcal{C})| \rightarrow X$ that is affine linear on each simplex of the barycentric subdivision; see Fig. 6 for an illustration. Similar constructions can be found in the literature [BT82, Theorem 13.4] [Hau95, p. 179].




Figure 6.: Illustration of the map Γ .

Theorem 3.1. *The map Γ is a homotopy equivalence. In particular, $|\mathrm{Nrv}(\mathcal{C})|$ is homotopy equivalent to X .*

We prove the theorem by constructing a homotopy inverse Ψ to Γ . For this construction, we work with an open cover and a subordinate partition of unity, as in the familiar proof of the nerve theorem for open covers [Hat02, Proposition 4G.2]. To this end, we thicken the subsets C_i slightly so that the nerve remains unchanged. If the C_i are compact, it is possible to choose an ε such that the open ε -neighborhoods of the C_i have this desired property. More generally, we can choose such neighborhoods according to the following lemma.

Lemma 3.2. *Let $\mathcal{C} = (C_i)_{i \in [n]}$ be a collection of closed and convex subsets of \mathbb{R}^d . Then there exists a collection of open sets $\mathcal{G} = (U_i \supseteq C_i)_{i \in [n]}$ satisfying $\mathrm{Nrv}(\mathcal{C}) = \mathrm{Nrv}(\mathcal{G}_\varepsilon)$.*

Proof. Since \mathbb{R}^d is a normal space, every disjoint pair of closed sets C_i, C_j admits disjoint open neighborhoods $V_{i,j} \supseteq C_i, V_{j,i} \supseteq C_j$. Taking the finite intersection $U_i = \bigcap_{j: C_i \cap C_j = \emptyset} V_{i,j}$ for every i yields the desired open cover $(U_i)_{i \in [n]}$. \square

We choose \mathcal{G} according to Lemma 3.2. For each $i \in [n]$, there exists a Urysohn function $\varphi_i: \mathbb{R}^d \rightarrow [0, 1]$ that takes on the value 0 outside of U_i and the value 1 on C_i . For example, we may take

$$x \mapsto \frac{d(x, \mathbb{R}^d \setminus U_i)}{d(x, C_i) + d(x, \mathbb{R}^d \setminus U_i)}.$$

Normalizing these functions φ_i yields a partition of unity on X subordinate to the cover $(U_i \cap X)_{i \in [n]}$ of X : $\psi_i = \varphi_i / \sum_{j=0}^n \varphi_j: X \rightarrow [0, 1]$. We define the map $\Phi: X \rightarrow |\mathrm{Nrv}(\mathcal{C})|$ in barycentric coordinates for $|\mathrm{Nrv}(\mathcal{C})|$ as

$$\Phi: x \mapsto \sum_{i=0}^n \psi_i(x) \cdot |v_i|, \quad (3.1)$$

where $v_i = \{i\}$ is the vertex in $\mathrm{Nrv}(\mathcal{C})$ corresponding to i and $|v_i|$ is the corresponding point in the geometric realization. The map $\Psi: X \rightarrow |\mathrm{Sd} \mathrm{Nrv}(\mathcal{C})|$ is then given as the composite $\alpha_{\mathrm{Nrv}(\mathcal{C})} \circ \Phi$, where $\alpha_{\mathrm{Nrv}(\mathcal{C})}: |\mathrm{Nrv}(\mathcal{C})| \rightarrow |\mathrm{Sd} \mathrm{Nrv}(\mathcal{C})|$ is the standard homeomorphism from the nerve to its barycentric subdivision.

In order to show that Ψ is a homotopy inverse to Γ , we analyze more closely how these maps are related combinatorially. To this end, we use the following construction.

Definition 3.3. For every vertex v of a simplicial complex K , define the *closed barycentric star* as the subspace

$$\mathrm{bst} v = |\mathrm{ClSt}_{\mathrm{Sd} K} v| \subseteq |\mathrm{Sd} K|,$$

where $\mathrm{ClSt}_{\mathrm{Sd} K} v = \{\sigma \in \mathrm{Sd} K \mid \sigma \cup \{v\} \in \mathrm{Sd} K\}$ is the closure of the star of v in the barycentric subdivision of K .

We now state two lemmas about the closed barycentric stars, deferring the proofs to Appendix A.

Lemma 3.4. *Let K be a simplicial complex and let $\sigma \in K$ be a simplex. Then the intersection $\bigcap_{v \in \sigma} \text{bst } v$ is contractible. In particular, the collection of closed barycentric stars forms a good cover of $|K|$.*

It is not hard to see that the nerve of this cover is isomorphic to K . The following statement describes the closed barycentric stars in terms of barycentric coordinates.

Lemma 3.5. *Let K be a simplicial complex and let v be a vertex of K . The closed barycentric star $\text{bst } v$ consists of all points $x \in |K|$ that satisfy*

$$b_v(x) \geq b_w(x) \quad \text{for all } w \in \text{Vert } K, \quad (3.2)$$

where b_v denotes the barycentric coordinate with respect to the vertex v .

Proposition 3.6. *The pair of maps $(\Psi, \text{id}_{[n]})$ constitutes a morphism of covered spaces*

$$(X, \mathcal{C} = (C_i)_{i \in [n]}) \rightarrow (|\text{Sd Nrv}(\mathcal{C})|, (\text{bst } v_i)_{i \in [n]}).$$

Proof. Recall that $\Psi = \alpha_{\text{Nrv}(\mathcal{C})} \circ \Phi$, where $\alpha_{\text{Nrv}(\mathcal{C})}$ is the isomorphism $|\text{Nrv}(\mathcal{C})| \cong |\text{Sd Nrv}(\mathcal{C})|$ and $\Phi: X \rightarrow |\text{Nrv}(\mathcal{C})|$, $x \mapsto \sum_{i=0}^n \psi_i(x) \cdot |v_i|$. Note that if $x \in C_i$, then $\phi_i(x) = 1$ and thus $\psi_i(x)$ is maximal among the $\psi_j(x)$. Hence, by Lemma 3.5 we know that $\Psi(x) \in \text{bst}(v_i)$ and the claim follows. \square

Proposition 3.7. *The pair of maps $(\Gamma, \text{id}_{[n]})$ constitutes a morphism of covered spaces*

$$(|\text{Sd Nrv}(\mathcal{C})|, (\text{bst } v_i)_{i \in [n]}) \rightarrow (X, \mathcal{C} = (C_i)_{i \in [n]}).$$

Proof. By definition, the map Γ sends the vertices of a geometric simplex σ in $\text{bst } v_i$ to C_i . As the cover element C_i is convex and Γ is affine linear on σ , it follows that $\Gamma(\sigma)$ is also contained in C_i . This shows $\Gamma(\text{bst } v_i) \subseteq C_i$, proving the claim. \square

We will now show that Ψ is a homotopy inverse to Γ , which implies $|\text{Nrv}(\mathcal{C})| \cong |\text{Sd Nrv}(\mathcal{C})| \simeq X$.

Proof of Theorem 3.1. It follows from Proposition 3.7 and Proposition 3.6 that the pair of maps $(\Gamma \circ \Psi, \text{id}_{[n]})$ constitutes a morphism of covered spaces. Hence, $\Gamma \circ \Psi$ is carried by the identity on \mathcal{C} and thus it is homotopic to the identity id_X by a straight line homotopy: for every $x \in C_i$, we have $\Gamma \circ \Psi(x) \in C_i$, and since the C_i are convex, the line segment joining x and $\Gamma \circ \Psi(x)$ lies in C_i . Similarly, the pair of maps $(\Psi \circ \Gamma, \text{id}_{[n]})$ constitutes a morphism of covered spaces. That the composition $\Psi \circ \Gamma$ is homotopic to $\text{id}_{|\text{Sd Nrv}(\mathcal{C})|}$ now follows from Lemma 3.4 and the following Proposition 3.8. \square

Recall that any two maps into a contractible space are homotopic (to a constant map). The following statement generalizes this fact to good covers, where contractibility is only guaranteed locally. For a more general statement see Lemma C.1.

Proposition 3.8. *Let K be a finite simplicial complex and let Y be a topological space. Assume we have two morphisms of covered spaces with the same map of indexed covers*

$$(f, \varphi), (g, \varphi): (|K|, (|L_i|)_{i \in [n]}) \rightarrow (Y, (V_j)_{j \in J}),$$

where $(|L_i|)_{i \in [n]}$ is a cover by subcomplexes and $(V_j)_{j \in J}$ is a good cover. Then f is homotopic to g .

Proof. Let $I = [0, 1]$ denote the unit interval. We inductively construct homotopies $H^m: |\text{sk}_m K| \times I \rightarrow Y$ between $f|_{|\text{sk}_m K|}$ and $g|_{|\text{sk}_m K|}$ such that H^m is carried by the map of indexed covers $P^m: (|\text{sk}_m L_i| \times I)_{i \in [n]} \rightarrow (V_j)_{j \in J}$ induced by φ . If $m = n$ is the dimension of the simplicial complex, the map $H = H^n$ is the desired homotopy between f and g .

To establish the base case $m = 0$, let p be any vertex of K and let $i_0, \dots, i_k \in [n]$ be those indices with $|p| \in |L_{i_l}|$. By the assumption that f and g are carried by φ , we know that both $f(|p|)$ and $g(|p|)$ are contained in $S := \bigcap_{l=0}^k V_{\varphi(i_l)}$, which is contractible by assumption, and thus we can choose a path in S that connects these two points. This defines the desired homotopy H^0 .

To see that the map H^0 is carried by P^0 , let $(|p|, t) \in |p| \times I$ be a point. If $(|p|, t) \in |L_i| \times I$, then $i = i_l$ is one of the indices above. Thus, by construction, $H^0((|p|, t)) \in S \subseteq V_{\varphi(i_l)} = V_{\varphi(i)}$ and the claim is proven.

For the induction step from $(m-1)$ to m , let H^{m-1} be as above. Let σ be an m -simplex in $\text{sk}_m K$. Furthermore, let $i_0, \dots, i_k \in [n]$ be those indices with $|\partial\sigma| \subseteq |\sigma| \subseteq |L_{i_l}|$. Then we have

$$H^{m-1}(|\partial\sigma| \times I) \subseteq S := \bigcap_{l=0}^k V_{\varphi(i_l)}$$

by the induction hypothesis. As the space S is contractible, we can extend the homotopy $H^{m-1}|_{|\partial\sigma| \times I}$ to a homotopy $H^m|_{|\sigma| \times I}$ from $f|_{|\sigma|}$ to $g|_{|\sigma|}$:

$$\begin{array}{ccc} (|\partial\sigma| \times I) \cup (|\sigma| \times \{0, 1\}) \cong S^m & \xrightarrow{(H^{m-1}, (f, g))} & S \subseteq Y \\ \downarrow & \nearrow H^m|_{|\sigma| \times I} & \\ |\sigma| \times I \cong B^{m+1} & & \end{array}$$

Because the m -simplex σ was arbitrary, we can extend $H^{m-1}: |\text{sk}_{m-1} K| \times I \rightarrow Y$ to $H^m: |\text{sk}_m K| \times I \rightarrow Y$.

By construction, this map is carried by P^m : Let $(x, t) \in |\sigma| \times I$ be a point with x in the interior of $|\sigma|$. If $x \in |L_i|$, then already all of $|\sigma|$ lies in $|L_i|$, as L_i is a subcomplex of K . Hence, $i = i_l$ is one of the indices above, and by construction of H^m ,

$$H^m(x, t) \in H^m(|\sigma| \times I) \subseteq S \subseteq V_{\varphi(i_l)} = V_{\varphi(i)},$$

proving the claim. □

3.1. A Functorial Nerve Theorem for Closed Convex Covers

Now, we will discuss two ways of turning the result in Theorem 3.1 into a functorial nerve theorem. We start by giving a version that follows the strategy explained in Section 2. After this, we will give a version that is functorial on the nose but needs to use the concept of pointed covers.

Theorem 3.9. *If $X \subset \mathbb{R}^d$, and $\mathcal{C} = (C_i)_{i \in [n]}$ is a cover of X by closed convex subsets, then the natural maps $\rho_S: \text{Blowup}(\mathcal{C}) \rightarrow X$ and $\rho_N: \text{Blowup}(\mathcal{C}) \rightarrow |\text{Nrv}(\mathcal{C})|$, introduced in Section 2, are homotopy equivalences.*

Proof. As explained at the end of Section 2.2, it suffices to consider the (not necessarily commutative) diagram

$$\begin{array}{ccc} & \text{Bar}(P_{\mathcal{C}}, \mathcal{D}_{\mathcal{C}}) & \\ \pi_S \swarrow & & \searrow \pi_{\text{Sd}N} \\ X & \xleftarrow{\Gamma} & |\text{Sd Nrv}(\mathcal{C})| \end{array}$$

where Γ is as in Theorem 3.1, and show that π_S and $\pi_{\text{Sd}N}$ are homotopy equivalences.

By Proposition 2.6 and the fact that convex sets are contractible, we know that $\pi_{\text{Sd}N}$ is a homotopy equivalence. Every point $p \in \text{Bar}(P_{\mathcal{C}}, \mathcal{D}_{\mathcal{C}})$ can be described as a pair $p = (x, \alpha)$, where α is a point in $|\sigma|$, for some $\sigma = (J_n \subset \cdots \subset J_0) \in \text{Sd Nrv}(\mathcal{C})$, and $x \in C_{J_0}$. By construction, we have $\Gamma(\pi_{\text{Sd}N}(p)) = \Gamma(\alpha) \in C_{J_n}$ and $\pi_S(p) = x \in C_{J_0} \subseteq C_{J_n}$. Therefore, a straight line homotopy shows that the maps $\Gamma \circ \pi_{\text{Sd}N} \simeq \pi_S$ are homotopic. As Γ and $\pi_{\text{Sd}N}$ are homotopy equivalences the same is true for π_S . \square

3.2. A One-Arrow Functorial Nerve Theorem for Pointed Covers

We will now describe the second way of obtaining a functorial nerve theorem. Recall from Definition 1.5 and the paragraphs afterwards the definition of Cov_* and its subcategory ClConv_* . Before stating the functorial nerve theorem let us consider one more important example of a pointed covered space.

Example 3.10. Let K be a simplicial complex. The cover $(\text{bst}(v))_{v \in \text{Vert } K}$ of $|K|$ by the closed barycentric stars is pointed by the vertices $(|w|)_{w \in \text{Vert Sd } K}$.

Let $(X, \mathcal{A}_*) \in \text{ClConv}_*$ be a pointed covered space. Recall that the construction of Γ at the beginning of Section 3 requires many choices. Those choices can be made such that Γ is a morphism of pointed covered spaces, where the nerve is a pointed covered space as described in Example 3.10, and such that it is affine linear on each simplex of the barycentric subdivision of the nerve.

Theorem 3.11. *The homotopy equivalence $\Gamma: |\text{Sd Nrv}(\mathcal{A})| \rightarrow X$ is natural with respect to the morphisms in ClConv_* .*

Proof. To show naturality, let $(f, \varphi): (X, \mathcal{A}_*) \rightarrow (Y, \mathcal{C}_*)$ be a morphism in ClConv_* . Then we need to prove that the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \uparrow & & \uparrow \\ |\text{Sd Nrv}(\mathcal{A})| & \xrightarrow{|\text{Sd } \varphi_*|} & |\text{Sd Nrv}(\mathcal{C})| \end{array}$$

commutes. Both compositions are maps $|\text{Sd Nrv}(\mathcal{A})| \rightarrow Y$ that are affine linear on each simplex of the barycentric subdivision. Hence, they are completely determined by their values on the vertices, where both compositions coincide by construction. \square

4. Nerve Theorems for Simplicial and Semi-Algebraic Covers

One can prove a nerve theorem for simplicial complexes as a corollary of Quillen's Theorem A for posets. In this section, we use combinatorial arguments to prove a functorial version of this result. Using a well-known triangulation theorem for semi-algebraic sets, this functorial nerve theorem for simplicial complexes implies such a theorem for finite, closed, semi-algebraic covers of compact semi-algebraic sets. Finally, we use the same combinatorial methods to prove a functorial version of a nerve theorem of Björner.

Posets and Homotopy Theory We begin by reviewing some basic facts about posets, following Quillen [Qui78].

Recall that the flag complex $\text{Flag}(P)$ of a poset P is the simplicial complex whose vertices are the elements of P and whose n -simplices are the chains $x_0 < \dots < x_n$ of elements of P . We will sometimes say that a poset has a certain topological property if its flag complex has that property. For example, we say that a poset P is contractible if $|\text{Flag}(P)|$ is contractible, and we say a map $f: P \rightarrow Q$ of posets is a homotopy equivalence if the induced map $|\text{Flag}(P)| \rightarrow |\text{Flag}(Q)|$ is a homotopy equivalence. If P and Q are posets, then there is a canonical homeomorphism

$$|\text{Flag}(P \times Q)| \xrightarrow{\cong} |\text{Flag}(P)| \times |\text{Flag}(Q)| \quad (4.1)$$

induced by the projection maps. As explained in [Qui78], the product must be taken in the category of compactly generated spaces, Definition 5.1. However, if one of P or Q is finite, then this agrees with the usual product. It follows that if $f, g: P \rightarrow Q$ are maps of posets such that $f(x) \leq g(x)$ for all $x \in P$, then $|\text{Flag}(f)|, |\text{Flag}(g)|: |\text{Flag}(P)| \rightarrow |\text{Flag}(Q)|$ are homotopic. To see this, observe that the relation $f \leq g$ determines a map of posets $H: P \times \{0 < 1\} \rightarrow Q$, and $|\text{Flag}(\{0 < 1\})|$ is an interval.

The main result about posets that we need is Quillen's Theorem A [Qui73]. Given a map $f: P \rightarrow Q$ of posets and $y \in Q$, define the subposet of P : $f/y = \{x \in P \mid f(x) \leq y\}$.

Theorem 4.1 (Quillen's Theorem A). *If $f: P \rightarrow Q$ is a map of posets, and f/y is contractible for all $y \in Q$, then f is a homotopy equivalence.*

It should be said that Quillen's theorem is more general than this result, but this is what we will use. For a nice proof at this level of generality, see [Wal81] or [Bar11], where it is shown that for finite posets the map f is even a simple homotopy equivalence. We now use Quillen's Theorem A to prove a nerve theorem for covers of a simplicial complex by subcomplexes.

Proposition 4.2. *Let K be a simplicial complex and let $\mathcal{A} = (K_i \subseteq K)_{i \in I}$ be a locally finite good cover of K by subcomplexes. Then K is homotopy equivalent to $\text{Nrv}(\mathcal{A})$.*

Proof. Define a map of posets $f: \text{Pos}(K) \rightarrow \text{Pos}(\text{Nrv}(\mathcal{A}))^{\text{op}}$ by

$$f(\sigma) = \{i \in I \mid \sigma \in K_i\}.$$

As \mathcal{A} is locally finite, $f(\sigma)$ is finite for all $\sigma \in K$. We will show that f is a homotopy equivalence. As usual, for $J \subseteq I$, we write $K_J = \bigcap_{i \in J} K_i$. By Quillen's Theorem A, it suffices to show that, for all elements J of $\text{Pos}(\text{Nrv}(\mathcal{A}))^{\text{op}}$, the poset f/J is contractible. Unwinding the definition, f/J is the subposet of $\text{Pos}(K)$ with elements $\sigma \in \text{Pos}(K)$ such that $J \subseteq f(\sigma)$. By definition, $J \subseteq f(\sigma)$ if and only if $\sigma \in K_J$. So, $f/J = \text{Pos}(K_J)$. As the intersection K_J is nonempty, it is contractible by assumption, and so $|\text{Flag}(\text{Pos}(K_J))| \cong |K_J|$ is contractible. Thus, f is a homotopy equivalence. The homotopy equivalence of the proposition is the composition:

$$|K| \cong |\text{Sd}(K)| \xrightarrow{f_*} |\text{Sd}(\text{Nrv}(\mathcal{A}))| \cong |\text{Nrv}(\mathcal{A})|. \quad \square$$

4.1. A Functorial Nerve Theorem for Simplicial Covers

In order to prove a functorial version of Proposition 4.2, we now introduce a poset PoBar that is intermediate between the posets $\text{Pos}(K)$ and $\text{Pos}(\text{Nrv}(\mathcal{A}))^{\text{op}}$ that appeared in the proof. We use the notation PoBar because this construction can be seen as a bar construction taken in the category of posets, as we explain in Example 5.40.

Definition 4.3. If K is a simplicial complex and $\mathcal{A} = (K_i \subseteq K)_{i \in I}$ is a cover of K by subcomplexes, let $\text{PoBar}(\mathcal{A})$ be the poset with the underlying set

$$\text{PoBar}(\mathcal{A}) = \{(\sigma, J) \mid J \subseteq I \text{ finite}, \sigma \in K_J\}$$

where $(\sigma, J) \leq (\sigma', J')$ if and only if $\sigma \subseteq \sigma'$ and $J \supseteq J'$.

Since $\text{PoBar}(\mathcal{A})$ is a subposet of the product $\text{Pos}(K) \times \text{Pos}(\text{Nrv}(\mathcal{A}))^{\text{op}}$, it comes with projection maps $\lambda_S: \text{PoBar}(\mathcal{A}) \rightarrow \text{Pos}(K)$ and $\lambda_N: \text{PoBar}(\mathcal{A}) \rightarrow \text{Pos}(\text{Nrv}(\mathcal{A}))^{\text{op}}$. In the next lemma, f denotes the poset map defined in the proof of Proposition 4.2.

Lemma 4.4. *Let K be a simplicial complex and let $\mathcal{A} = (K_i \subseteq K)_{i \in I}$ be a locally finite good cover of K by subcomplexes. The following diagram of posets commutes up to homotopy after taking flag complexes, and all three maps are homotopy equivalences:*

$$\begin{array}{ccc} & \text{PoBar}(\mathcal{A}) & \\ \lambda_S \swarrow & & \searrow \lambda_N \\ \text{Pos}(K) & \xrightarrow{f} & \text{Pos}(\text{Nrv}(\mathcal{A}))^{\text{op}} \end{array}$$

Proof. We begin by showing that λ_S is a homotopy equivalence. Define a map of posets $\mu: \text{Pos}(K) \rightarrow \text{PoBar}(\mathcal{A})$ by $\mu(\sigma) = (\sigma, f(\sigma))$. The composition $\lambda_S \circ \mu$ is the identity on $\text{Pos}(K)$, and if $(\sigma, J) \in \text{PoBar}(\mathcal{A})$, then $(\mu \circ \lambda_S)(\sigma, J) = (\sigma, f(\sigma)) \leq (\sigma, J)$. It follows that $|\text{Flag}(\mu)| \circ |\text{Flag}(\lambda_S)|$ is homotopic to the identity on $|\text{Flag}(\text{PoBar}(\mathcal{A}))|$, so that λ_S is a homotopy equivalence.

If (σ, J) is in $\text{PoBar}(\mathcal{A})$, then $\lambda_N(\sigma, J) = J \subseteq f(\sigma) = (f \circ \lambda_S)(\sigma, J)$. So, we have $(f \circ \lambda_S)(\sigma, J) \leq \lambda_N(\sigma, J)$, which implies that $|\text{Flag}(f)| \circ |\text{Flag}(\lambda_S)|$ and $|\text{Flag}(\lambda_N)|$ are homotopic. We proved that f is a homotopy equivalence in the proof of Proposition 4.2. As λ_S and f are homotopy equivalences, so is λ_N . \square

The strategy now is to use what we have proved about the PoBar construction to show that the natural maps from the blowup complex to $|K|$ and $|\text{Nrv}(\mathcal{A})|$ (defined in Section 2) are homotopy equivalences.

To do this, we will identify a subcomplex $\text{T}(\mathcal{A}) \subseteq \text{Flag}(\text{PoBar}(\mathcal{A}))$ that is homeomorphic to the blowup complex after realization; we then show, using discrete Morse theory, that the inclusion $|\text{T}(\mathcal{A})| \hookrightarrow |\text{Flag}(\text{PoBar}(\mathcal{A}))|$ is a homotopy equivalence. In particular, it follows that the blowup complex is homotopy equivalent to $|\text{Flag}(\text{PoBar}(\mathcal{A}))|$.

Definition 4.5. Let K be a simplicial complex, and let $\mathcal{A} = (K_i \subseteq K)_{i \in I}$ be a cover of K by subcomplexes. Let $\text{T}(\mathcal{A})$ be the subcomplex of $\text{Flag}(\text{PoBar}(\mathcal{A}))$ consisting of the simplices $(\sigma_0, J_0) < \dots < (\sigma_m, J_m)$ such that $\sigma_m \in K_{J_0}$.

The letter T stands for “triangulation”, since $\text{T}(\mathcal{A})$ turns out to be a triangulation of the blowup complex.

Lemma 4.6. Let K be a simplicial complex, and let $\mathcal{A} = (K_i \subseteq K)_{i \in I}$ be a cover of K by subcomplexes; write $|\mathcal{A}| = (|K_i| \subseteq |K|)_{i \in I}$. There is a homeomorphism $\varphi: \text{Blowup}(|\mathcal{A}|) \rightarrow |\text{T}(\mathcal{A})|$ such that the following diagram commutes:

$$\begin{array}{ccccc}
|K| & \xleftarrow{\rho_S} & \text{Blowup}(|\mathcal{A}|) & \xrightarrow{\rho_N} & |\text{Nrv}(\mathcal{A})| \\
\cong \downarrow & & \downarrow \varphi & & \downarrow \cong \\
|\text{Sd}(K)| & \xleftarrow{\quad} & |\text{T}(\mathcal{A})| & \xrightarrow{\quad} & |\text{Sd}(\text{Nrv}(\mathcal{A}))| \\
& \swarrow \lambda_S & \downarrow & \searrow \lambda_N & \\
& & |\text{Flag}(\text{PoBar}(\mathcal{A}))| & &
\end{array} \tag{4.2}$$

Here, the vertical maps on the left and right are the standard homeomorphisms.

Proof. The blowup complex $\text{Blowup}(|\mathcal{A}|)$ is defined by glueing together pieces of the form $|K_J| \times |J|$ for $J \in \text{Nrv}(\mathcal{A})$. We abuse notation and write J also for the full simplicial complex on J . For any such piece, define φ by the composition

$$\begin{aligned}
|K_J| \times |J| &\cong |\text{Sd } K_J| \times |\text{Sd } J| \\
&= |\text{Flag}(\text{Pos}(K_J))| \times |\text{Flag}(\text{Pos}(J))| \\
&= |\text{Flag}(\text{Pos}(K_J))| \times |\text{Flag}(\text{Pos}(J)^{\text{op}})| \\
&\cong |\text{Flag}(\text{Pos}(K_J) \times \text{Pos}(J)^{\text{op}})| \subseteq |\text{T}(\mathcal{A})|
\end{aligned}$$

where the last homeomorphism is an instance of 4.1. As these maps respect the equivalence relation from the definition of the blowup complex, together they define a continuous map $\varphi: \text{Blowup}(|\mathcal{A}|) \rightarrow |\mathbb{T}(\mathcal{A})|$. By construction, the diagram 4.2 commutes.

To see that φ is a homeomorphism, we can construct its inverse. As J varies, the subcomplexes $\text{Flag}(\text{Pos}(K_J) \times \text{Pos}(J)^{\text{op}})$ cover $\mathbb{T}(\mathcal{A})$. For each J , we can reverse the homeomorphisms in the definition of φ to define φ^{-1} on $|\text{Flag}(\text{Pos}(K_J) \times \text{Pos}(J)^{\text{op}})|$. Since these maps agree on intersections, they glue together to define the inverse $\varphi^{-1}: |\mathbb{T}(\mathcal{A})| \rightarrow \text{Blowup}(|\mathcal{A}|)$. \square

Lemma 4.7. *Let K be a simplicial complex, and let $\mathcal{A} = (K_i \subseteq K)_{i \in I}$ be a cover of K by subcomplexes. Then the inclusion $|\mathbb{T}(\mathcal{A})| \hookrightarrow |\text{Flag}(\text{PoBar}(\mathcal{A}))|$ is a homotopy equivalence.*

Proof. We construct a discrete gradient vector field V on $\text{Flag}(\text{PoBar}(\mathcal{A}))$, such that the set of critical simplices is $\mathbb{T}(\mathcal{A})$. Then it will follow from Proposition B.2 that the inclusion $|\mathbb{T}(\mathcal{A})| \hookrightarrow |\text{Flag}(\text{PoBar}(\mathcal{A}))|$ is a homotopy equivalence.

First, define the discrete vector field V as follows: Take a simplex τ in $\text{Flag}(\text{PoBar}(\mathcal{A})) \setminus \mathbb{T}(\mathcal{A})$, i.e. a chain $(\sigma_0, J_0) < \dots < (\sigma_m, J_m)$ in $\text{PoBar}(\mathcal{A})$ such that $\sigma_m \notin K_{J_0}$. Then there exists a smallest index i with $\sigma_i \notin K_{J_0}$. If $J_{i-1} \neq J_i$, consider the chain

$$(\sigma_0, J_0) < \dots < (\sigma_{i-1}, J_{i-1}) < (\sigma_{i-1}, J_i) < (\sigma_i, J_i) < \dots < (\sigma_m, J_m)$$

and pair the corresponding simplex μ in $\text{Flag}(\text{PoBar}(\mathcal{A})) \setminus \mathbb{T}(\mathcal{A})$ with τ . Note that these pairs are disjoint and every simplex in $\text{Flag}(\text{PoBar}(\mathcal{A})) \setminus \mathbb{T}(\mathcal{A})$ appears in such a pairing. Hence, V is a discrete vector field.

Now, we prove that V is a discrete gradient vector field by showing that there are no non-trivial closed V -paths: Consider any V -path $\tau_0 \rightarrow \mu_0 \leftarrow \dots \rightarrow \mu_r \leftarrow \tau_{r+1}$ with $\{\tau_i, \mu_i\} \in V$ and $\tau_i \neq \tau_{i+1}$ for all i . We will show that it is not closed, i.e. $\tau_{r+1} \neq \tau_0$. Assume that τ_0 represents the chain $(\sigma_0, J_0) < \dots < (\sigma_m, J_m)$ and let i be the smallest index i with $\sigma_i \notin K_{J_0}$. We distinguish two cases: If τ_1 is either of the form

$$(\sigma_0, J_0) < \dots < \widehat{(\sigma_j, J_j)} \leq \dots \leq (\sigma_{i-1}, J_{i-1}) < (\sigma_{i-1}, J_i) < (\sigma_i, J_i) < \dots < (\sigma_m, J_m)$$

or of the form

$$(\sigma_0, J_0) < \dots < (\sigma_{i-1}, J_{i-1}) < (\sigma_{i-1}, J_i) < (\sigma_i, J_i) < \dots < \widehat{(\sigma_j, J_j)} < \dots < (\sigma_m, J_m),$$

then $r = 0$ and $\tau_{r+1} = \tau_1$, which means that the sequence is not closed. Moreover, after reindexing the chain to be of the form $(\tilde{\sigma}_0, \tilde{J}_0) < \dots < (\tilde{\sigma}_m, \tilde{J}_m)$, the smallest index i with $\tilde{\sigma}_i \notin K_{\tilde{J}_0}$ did not decrease. In the second case τ_1 is of the form

$$(\sigma_0, J_0) < \dots < (\sigma_{i-1}, J_{i-1}) < (\sigma_{i-1}, J_i) < \widehat{(\sigma_i, J_i)} < \dots < (\sigma_m, J_m).$$

Thus, after reindexing this chain to be of the form $(\tilde{\sigma}_0, \tilde{J}_0) < \dots < (\tilde{\sigma}_m, \tilde{J}_m)$, the smallest index i with $\tilde{\sigma}_i \notin K_{\tilde{J}_0}$ increased by one. This shows, after sufficiently many reapplications of these arguments, that τ_{r+1} is not equal to τ_0 , which means that the

sequence is not closed. Hence, as the sequence was arbitrary, we have proven that V is a discrete gradient vector field.

The reasoning above also implies that for every simplex in $\text{Flag}(\text{PoBar}(\mathcal{A}))$ its V -path height is finite and hence it follows from Lemma B.1 and Proposition B.2 that the inclusion $|\text{T}(\mathcal{A})| \hookrightarrow |\text{Flag}(\text{PoBar}(\mathcal{A}))|$ is a homotopy equivalence. \square

Theorem 4.8. *Let K be a simplicial complex and let $\mathcal{A} = (K_i \subseteq K)_{i \in I}$ be a locally finite good cover of K by subcomplexes. Then, the natural maps $\rho_S: \text{Blowup}(|\mathcal{A}|) \rightarrow |K|$ and $\rho_N: \text{Blowup}(|\mathcal{A}|) \rightarrow |\text{Nrv}(\mathcal{A})|$ are homotopy equivalences.*

Proof. Consider Diagram 4.2. By Lemma 4.4 the maps λ_S and λ_N are homotopy equivalences. By Lemma 4.7, the inclusion $|\text{T}(\mathcal{A})| \hookrightarrow |\text{Flag}(\text{PoBar}(\mathcal{A}))|$ is a homotopy equivalence, and by Lemma 4.6, φ is a homeomorphism. It follows that ρ_S and ρ_N are homotopy equivalences. \square

Remark 4.9. It is possible to remove the locally finiteness assumption from the theorem: From Proposition 2.6, the discussion at the beginning of Section 2.2 and Eq. (2.1) it follows that $\rho_N: \text{Blowup}(|\mathcal{A}|) \rightarrow |\text{Nrv}(\mathcal{A})|$ is a homotopy equivalence. Moreover, one can show by using contractible carriers that $\rho_S: \text{Blowup}(|\mathcal{A}|) \rightarrow |K|$ is a homotopy equivalence for any cover by subcomplexes; see Proposition C.3.

4.2. A Functorial Nerve Theorem for Semi-Algebraic Covers

As a corollary of Theorem 4.8, we get a functorial nerve theorem for finite, closed, semi-algebraic covers of compact semi-algebraic sets. For this, we need a well known theorem on the existence of triangulations of semi-algebraic sets [BCR98, Theorem 9.2.1], which we now state. For K a simplicial complex and σ a simplex of K , we write $\text{int}|\sigma| = |\sigma| \setminus |\partial\sigma| \subset |K|$ for the open simplex.

Theorem 4.10. *Let $S \subset \mathbb{R}^n$ be a compact semi-algebraic set, and let $(S_i)_{i=0}^q$ be a finite family of semi-algebraic subsets of S . There is a finite simplicial complex $K = \{\sigma_j\}_{j=0}^p$ and a homeomorphism $h: |K| \rightarrow S$, such that every S_i is the union of some images of simplices $h(\text{int}|\sigma_j|)$.*

Corollary 4.11. *Let $S \subset \mathbb{R}^n$ be a compact semi-algebraic set, and let $\mathcal{A} = (S_i)_{i=0}^q$ be a finite good cover of S such that each S_i is semi-algebraic and closed in S . The natural maps $\rho_S: \text{Blowup}(\mathcal{A}) \rightarrow S$ and $\rho_N: \text{Blowup}(\mathcal{A}) \rightarrow |\text{Nrv}(\mathcal{A})|$ are homotopy equivalences.*

Proof. By Theorem 4.10, there is a simplicial complex K , a homeomorphism $h: |K| \rightarrow S$, and a cover of K by subcomplexes $\mathcal{B} = (K_i \subseteq K)_{i=0}^q$ such that $h|_{K_i}$ is a homeomorphism between K_i and S_i . Then, h induces a homeomorphism $\text{Blowup}(|\mathcal{B}|) \rightarrow \text{Blowup}(\mathcal{A})$ such that the following diagram commutes:

$$\begin{array}{ccccc} |K| & \xleftarrow{\rho_S} & \text{Blowup}(|\mathcal{B}|) & \xrightarrow{\rho_N} & |\text{Nrv}(\mathcal{B})| \\ h \downarrow & & \downarrow & & \parallel \\ S & \xleftarrow{\rho_S} & \text{Blowup}(\mathcal{A}) & \xrightarrow{\rho_N} & |\text{Nrv}(\mathcal{A})| \end{array}$$

By Theorem 4.8 the top horizontal maps are homotopy equivalences, and the corollary follows. \square

4.3. A Functorial Version of Björner's Nerve Theorem

If K is a simplicial complex, and \mathcal{A} is a locally finite cover of K by subcomplexes, then we have a comparison map $|K| \rightarrow |\mathrm{Nrv}(\mathcal{A})|$ induced by the map of posets $f: \mathrm{Pos}(K) \rightarrow \mathrm{Pos}(\mathrm{Nrv}(\mathcal{A}))^{\mathrm{op}}$ defined in the proof of Proposition 4.2. In [Bjö03], Björner gives a detailed analysis of how the connectivity of this map is affected by the connectivity of the finite intersections of cover elements. For the final result of this section, we will use the PoBar construction and the blowup complex to prove a functorial version of Björner's theorem.

Definition 4.12. Let $k \geq 0$. A topological space X is k -connected if, for every $0 \leq r \leq k$, every map of the r -sphere into X is homotopic to a constant map.

Theorem 4.13 ([Bjö03, Theorem 6]). *Let K be a simplicial complex, let \mathcal{A} be a locally finite cover of K by subcomplexes, and let $k \geq 0$. Assume that every non-empty intersection $K_{i_1} \cap \cdots \cap K_{i_t}$ is $(k - t + 1)$ -connected, for all $t \geq 1$. Then f induces a bijection*

$$\pi_0(K) \cong \pi_0(\mathrm{Nrv}(\mathcal{A})),$$

and for all $1 \leq j \leq k$, and for all $x \in |K|$, f induces an isomorphism

$$\pi_j(|K|, x) \cong \pi_j(|\mathrm{Nrv}(\mathcal{A})|, f_*(x)).$$

In fact, the theorem of Björner deals with regular CW complexes, rather than simplicial complexes. The assumption that \mathcal{A} is locally finite is omitted from the original statement, but it is used in the proof. For convenience, Björner assumes that K is connected: Theorem 4.13 follows from Björner's theorem and the following lemma, which is easily proved.

Lemma 4.14. *Let K be a simplicial complex, and let $\mathcal{A} = (K_i)_{i \in I}$ be a locally finite cover of K by subcomplexes such that K_i is non-empty and connected for all $i \in I$. Then f induces a bijection $\pi_0(K) \cong \pi_0(\mathrm{Nrv}(\mathcal{A}))$. \square*

Using the PoBar construction and the blowup complex as before, we obtain the following functorial version of Björner's theorem.

Theorem 4.15. *Let $k \geq 0$, let K be a simplicial complex, and let $\mathcal{A} = (K_i)_{i \in I}$ be a locally finite cover of K by subcomplexes. Assume that every non-empty intersection $K_{i_1} \cap \cdots \cap K_{i_t}$ is $(k - t + 1)$ -connected, for all $t \geq 1$. The natural map $\rho_S: \mathrm{Blowup}(|\mathcal{A}|) \rightarrow |K|$ is a homotopy equivalence, and the natural map $\rho_N: \mathrm{Blowup}(|\mathcal{A}|) \rightarrow |\mathrm{Nrv}(\mathcal{A})|$ induces a bijection in path components, and for all $1 \leq j \leq k$, and for all $x \in \mathrm{Blowup}(|\mathcal{A}|)$, ρ_N induces an isomorphism $\pi_j(\mathrm{Blowup}(|\mathcal{A}|), x) \cong \pi_j(|\mathrm{Nrv}(\mathcal{A})|, \rho_N(x))$.*

Proof. Note that the proof of Lemma 4.4 shows that the poset map λ_S is a homotopy equivalence, and that the triangle commutes up to homotopy, without the assumption that \mathcal{A} is good. So, by Björner's Theorem 4.13, $\lambda_N: |\mathrm{Flag}(\mathrm{PoBar}(\mathcal{A}))| \rightarrow$

$|\mathrm{Sd}(\mathrm{Nrv}(\mathcal{A}))|$ induces a bijection in path components, and for all $1 \leq j \leq k$, and for all $x \in |\mathrm{Flag}(\mathrm{PoBar}(\mathcal{A}))|$, λ_N induces an isomorphism $\pi_j(|\mathrm{Flag}(\mathrm{PoBar}(\mathcal{A}))|, x) \cong \pi_j(|\mathrm{Nrv}(\mathcal{A})|, \lambda_N(x))$. The result follows from commutativity of Diagram 4.2. \square

5. A Unified Nerve Theorem

We now prove the unified nerve theorem (Theorem D in the introduction), which recovers Theorems 3.9 and 4.8. The proof has three main ingredients. First, as in Theorem 3.9, we use the connection between the blowup complex and the bar construction (explained in Section 2) to deduce statements about the blowup complex from the corresponding statements about the bar construction. Second, since the bar construction is a standard tool in homotopy theory, we can use well-known results to prove the statements we need about it. Finally, Hatcher [Hat02, Section 4.G] already proves one part of the theorem: if \mathcal{A} is an open cover of a paracompact space X , then the natural map $\mathrm{Blowup}(\mathcal{A}) \rightarrow X$ is a homotopy equivalence. We begin by introducing the various notions from topology we need to state the unified nerve theorem.

In order to avoid pathological behavior in the category \mathbf{Top} of all topological spaces, algebraic topologists often restrict attention to full subcategories that include all the spaces they are interested in (such as CW complexes), and that have nicer categorical properties. For example, it is often convenient to work in a category of topological spaces that is *cartesian closed*: roughly speaking, this means that for any spaces X and Y in the category, we have a “mapping space” Y^X in the category such that the set of maps $Z \rightarrow Y^X$ is in bijection with the set of maps $X \times Z \rightarrow Y$, and this bijection is natural in Y . Letting $Z = *$, we see that the points of Y^X are in bijection with continuous maps $X \rightarrow Y$. Such mapping spaces play an important role in algebraic topology, because they encode homotopy-theoretic information. For example, a path $\gamma: [0, 1] \rightarrow Y^X$ in the mapping space corresponds to a homotopy $H: X \times [0, 1] \rightarrow Y$. There is more than one standard choice for such a subcategory. We will consider the following:

Definition 5.1. A topological space X is *weak Hausdorff* if $g(K)$ is closed in X for every continuous map $g: K \rightarrow X$ with K compact Hausdorff. A subspace A of X is *compactly closed* if $g^{-1}(A)$ is closed in K for every continuous map $g: K \rightarrow X$ with K compact Hausdorff. A space X is a *k-space* if every compactly closed subspace of X is closed. A space X is *compactly generated* if it is a weak Hausdorff *k-space*. The full subcategory of \mathbf{Top} of compactly generated spaces is denoted by \mathbf{CGSp} .

Some authors refer to *k-spaces* as compactly generated spaces, and use the term “compactly generated weak Hausdorff spaces” for what we call compactly generated spaces. See [May99, Chapter 5] or [Str09] for basic facts about these spaces. For example, one can show that there exist functors that are adjoint to the inclusions

$$\mathbf{CGSp} \quad \begin{array}{c} \longleftarrow \\ \perp \\ \longrightarrow \end{array} \quad \mathbf{k-spaces} \quad \begin{array}{c} \longleftarrow \\ \top \\ \longrightarrow \end{array} \quad \mathbf{Top},$$

where *k-spaces* is the full subcategory of *k-spaces*.

Example 5.2. Many spaces are compactly generated:

- Every closed subset of a compactly generated space is compactly generated.
- Every CW-complex is compactly generated.
- Every locally compact Hausdorff space is compactly generated [Str09, Proposition 1.7]. In particular, \mathbb{R}^d is compactly generated.

A simplicial complex K is sometimes said to be *locally finite dimensional* if every vertex v of K has a finite dimensional star, i.e., $\sup\{\dim \sigma \mid v \in \sigma\} < \infty$. Following this usage, we say that a cover of a topological space is locally finite dimensional if the nerve of the cover is so. More explicitly, we have the following:

Definition 5.3. If X is a topological space, and $\mathcal{A} = (A_i)_{i \in I}$ is a cover, then \mathcal{A} is *locally finite dimensional* if for each cover element A_i there exists $k_i \in \mathbb{N}$ such that for any $J \subseteq I$ with $A_J \neq \emptyset$ and $i \in J$, we have $|J| \leq k_i$.

Definition 5.4. Let R be a commutative ring. We say that a continuous map f between topological spaces is an *R -homology isomorphism* if $H_n(f, R)$ is an isomorphism for all $n \geq 0$. We say that a cover $\mathcal{A} = (A_i)_{i \in I}$ is *homologically good* with respect to R if, for all non-empty $J \subseteq I$ such that $A_J \neq \emptyset$, the map $A_J \rightarrow *$ is an R -homology isomorphism, where $*$ is the one point space.

Definition 5.5. We say that a cover $\mathcal{A} = (A_i)_{i \in I}$ is *weakly good* if, for all non-empty $J \subseteq I$ such that $A_J \neq \emptyset$, the map $A_J \rightarrow *$ is a weak homotopy equivalence, where $*$ is the one point space.

Definition 5.6. Let X be a topological space, and let $\mathcal{A} = (A_i)_{i \in I}$ be a cover. For $T \in \text{Nrv}(\mathcal{A})$, the *latching space* is the subset

$$L(T) := \bigcup_{T \subsetneq J \subseteq I} A_J \subseteq A_T.$$

Finally, let us recall the homotopy extension property.

Definition 5.7. Let X be topological spaces and let A be a subset. We say that the pair (X, A) satisfies the *homotopy extension property* if for every commutative diagram of the following shape the dotted arrow exists

$$\begin{array}{ccc}
 A & \xrightarrow{\quad} & X \\
 \text{id}_A \times \{0\} \downarrow & & \text{id}_X \times \{0\} \downarrow \\
 A \times [0, 1] & \xrightarrow{\quad} & X \times [0, 1] \\
 & \searrow H & \downarrow \tilde{H} \\
 & & Y
 \end{array}$$

f (curved arrow from X to Y)
 H (curved arrow from $A \times [0, 1]$ to Y)

In words, the pair (X, A) has the homotopy extension property if for any map f every homotopy H on A can be extended to a homotopy \tilde{H} defined on all of X .

Remark 5.8. A large class of pairs has the homotopy extension property. For example, if X is a CW-complex and A a subcomplex, then (X, A) satisfies the homotopy extension property ([Hat02, Proposition 0.16] or [Koz08, Proposition 7.10]). We say more about the homotopy extension property in Section 5.1.

Theorem 5.9 (Unified Nerve Theorem). *Let X be a topological space and let $\mathcal{A} = (A_i)_{i \in I}$ be a cover of X .*

1. *Consider the natural map $\rho_S: \text{Blowup}(\mathcal{A}) \rightarrow X$.*
 - a) *If \mathcal{A} is an open cover, then ρ_S is a weak homotopy equivalence. If furthermore X is paracompact, then ρ_S is a homotopy equivalence.*
 - b) *Assume that X is compactly generated and that \mathcal{A} is a closed cover that is locally finite and locally finite dimensional. If for any $T \in \text{Nrv}(\mathcal{A})$ the latching space $L(T) \subseteq A_T$ is a closed subset and the pair $(A_T, L(T))$ satisfies the homotopy extension property, then ρ_S is a homotopy equivalence.*
2. *Consider the natural map $\rho_N: \text{Blowup}(\mathcal{A}) \rightarrow |\text{Nrv}(\mathcal{A})|$.*
 - a) *If \mathcal{A} is (weakly) good, then ρ_N is a (weak) homotopy equivalence.*
 - b) *If for all $J \in \text{Nrv}(\mathcal{A})$ the space A_J is compactly generated and \mathcal{A} is homologically good with respect to a coefficient ring R , then ρ_N is an R -homology isomorphism.*

We prove Theorem 5.9 in Section 5.3.

Remark 5.10. The compactly generated assumption in 2(b) is satisfied for example if X is a locally compact Hausdorff space and \mathcal{A} is an open cover. The assumption also holds if X is compactly generated and \mathcal{A} is a closed cover; this also includes the case of a cover of a CW-complex by subcomplexes. See Example 5.2.

Remark 5.11. If X is a regular CW-complex and \mathcal{A} is a cover of subcomplexes, then 2.b can also be proven by using spectral sequence techniques [Bro94, Chapter VII, Section 4]. Note that in loc. cit. the total complex associated to the double complex used is isomorphic to the cellular chain complex of $\text{Blowup}(\mathcal{A})$. Moreover, these techniques can also be used to prove an analogous statement to Theorem 4.13 for homology groups [Mes01, Theorem 2.1].

Remark 5.12. Most of the technical assumptions in the unified nerve theorem are necessary. To demonstrate this, we will now give some counterexamples.

- The classical nerve theorem for a good open cover of a paracompact space is proved by using 1(a) and 2(a). We will now give an example that shows that this paracompactness assumption is necessary in order to establish a homotopy equivalence between space and nerve. Consider the *long ray* L , which is constructed as follows: Take the first uncountable ordinal ω_1 , which is a well-ordered set and its elements are all countable ordinals, and insert a unit interval $(0, 1)$ between each countable ordinal α and its successor $\alpha + 1$, yielding a totally ordered set. The

topology on L is given by the order topology. The long line is a standard example for a non-paracompact space that is also not contractible [SS95; Jos83]. However, L is weakly contractible and for any point $p \in L$ the open set $L_{<p} = \{t \in L \mid t < p\} \subset L$ is homeomorphic to the interval $[0, 1)$. Thus, the open cover $\mathcal{A} = (L_{<p})_{p \in \omega_1}$ is a good cover of L and it follows from 1(a) and 2(a) that the nerve $|\mathrm{Nrv} \mathcal{A}|$ is weakly contractible and hence contractible by Whitehead's theorem [Hat02, Theorem 4.5]. This implies that the space L and the nerve $|\mathrm{Nrv} \mathcal{A}|$ are not homotopy equivalent.

- The finiteness conditions in 1(b) control the size of the cover. If \mathcal{A} is the cover of the circle S^1 by its points, then all conditions in 1(b) and 2(a) are satisfied except the locally finiteness assumption. As the nerve $|\mathrm{Nrv} \mathcal{A}|$ is a disjoint union of points, it is not homotopy equivalent to S^1 .
- Even if we are only interested in finite good and closed covers, the covered space does not need to be homotopy equivalent to the nerve of the cover. Consider the *double comb space* C and denote the two combs by A_1 and A_2 (see Fig. 7). Then, the nerve $|\mathrm{Nrv} \mathcal{A}|$ of the finite good and closed cover $\mathcal{A} = \{A_1, A_2\}$ of C is contractible. Hence, it can not be homotopy equivalent to C , because the latter is not contractible. In this example, the pairs $(A_1, A_1 \cap A_2)$ and $(A_2, A_1 \cap A_2)$ do not satisfy the homotopy extension property. This shows that the conditions on the latching spaces are crucial, as all other assumptions in 1(b) and 2(a) are satisfied.
- If \mathcal{A} is any homologically good open cover of a locally compact Hausdorff space X , then it follows from 1(a) and 2(b) that the space X and the nerve $\mathrm{Nrv} \mathcal{A}$ have isomorphic homology groups. This conclusion does not hold if one replaces the open cover by a closed cover. Consider the *Warsaw circle* $W \subseteq S^2$ that separates the sphere into two connected components U_1 and U_2 (see Fig. 7). The closed sets $A_1 = U_1 \cup W$ and $A_2 = U_2 \cup W$ cover the sphere and are contractible. Moreover, the intersection $A_1 \cap A_2 = W$ is acyclic and hence $\mathcal{A} = \{A_1, A_2\}$ is a homologically good closed cover of S^2 . Nevertheless, the space S^2 and the nerve $\mathrm{Nrv} \mathcal{A}$ do not have isomorphic homology groups, as $H_2(S^2) \cong \mathbb{Z}$ and $H_2(\mathrm{Nrv} \mathcal{A}) \cong 0$. Hence, the conditions on the latching spaces are crucial, as all other assumptions in 1(b) and 2(b) are satisfied. This counterexample also shows that the nerve of a weakly good closed cover is not necessarily weakly equivalent to the space it covers.




Figure 7.: The double comb space (left) and the Warsaw circle (right).

5.1. Applications of the Unified Nerve Theorem

The assumptions on the latching spaces in 1(b) of Theorem 5.9 might not be easy to check in all situations. We now give a reformulation, and a union theorem for pairs that satisfy the homotopy extension property, which help to verify these assumptions. We also show in this subsection that Theorem 5.9 implies the functorial nerve theorems Theorem 3.9 and Theorem 4.8.

To study pairs that satisfy the homotopy extension property it suffices to consider neighborhood deformation retracts.

Definition 5.13. A pair of topological spaces (X, A) is called an *NDR-pair* if there exist continuous maps $u: X \rightarrow [0, 1]$ and $h: X \times [0, 1] \rightarrow X$ such that

- (i) $A = u^{-1}(0)$
- (ii) $h(-, 0) = \text{id}_X$
- (iii) $h(a, -) = a$ for all $a \in A$
- (iv) $h(x, 1) \in A$ for all $x \in X$ with $u(x) < 1$.

Proposition 5.14 ([Koz08, Proposition 7.7]). *Let A be a closed subspace of X . Then (X, A) is an NDR-pair if and only if (X, A) satisfies the homotopy extension property.*

The following union theorem is due to Lillig [Lil73].

Proposition 5.15. *Let $A_0, \dots, A_n \subseteq X$ be closed subsets and assume that for all $J \subseteq [n]$ the pair (X, A_J) satisfies the homotopy extension property. Then $(X, \bigcup_{i=0}^n A_i)$ also satisfies the homotopy extension property.*

This proposition, together with 1(b) in Theorem 5.9, implies the following corollary, which does not involve the latching spaces.

Corollary 5.16. *Let X be a compactly generated topological space and $\mathcal{A} = (A_i)_{i \in [n]}$ a finite closed cover. Assume that for all $I \subseteq J \subseteq [n]$ the pair (A_I, A_J) satisfies the homotopy extension property. Then $\rho_S: \text{Blowup}(\mathcal{A}) \rightarrow X$ is a homotopy equivalence.*

We will now illustrate how these statements can be used to deduce the functorial nerve theorem for closed convex sets in \mathbb{R}^d (Theorem 3.9) from the unified nerve theorem (Theorem 5.9). The proof of the following lemma is elementary and left to the reader.

Lemma 5.17. *Let $K \subseteq \mathbb{R}^d$ be compact and convex. Assume that $\text{aff } K = \mathbb{R}^d$, where $\text{aff } K$ is the affine hull of K . Then we have $K = \overline{\text{int } K}$ and $\text{int } K$ is convex.*

Proposition 5.18. *Let K and K' be compact and convex sets in \mathbb{R}^d with $K \subseteq K'$. Then the pair (K', K) satisfies the homotopy extension property.*

Proof. Without loss of generality, we can assume that $\text{aff } K' = \mathbb{R}^d$ holds.

First of all, let us assume that K is the intersection of K' with an affine subspace. Now, choose a point x in K . By Lemma 5.17 and the proof of [Mun84, Lemma 1.1], we see that there exists a homeomorphism $\varphi: \mathbb{R}^d \rightarrow \mathbb{R}^d$ such that $\varphi(K') = \overline{\mathbb{B}}_1(0)$, $\varphi(x) = 0$ and $\varphi(K) = \overline{\mathbb{B}}_1(0) \cap \mathbb{R}^l \times \{0\}^{d-l}$, with $l = \dim \text{aff } K$. The pair

$$(\varphi(K'), \varphi(K)) = (\overline{\mathbb{B}}_1(0), \overline{\mathbb{B}}_1(0) \cap \mathbb{R}^l \times \{0\}^{d-l})$$

is a CW-pair and hence it satisfies the homotopy extension property (Remark 5.8).

Now, let $K \subseteq K'$ be such that $\text{aff } K = \text{aff } K' = \mathbb{R}^d$ holds. As before, choose a point $x \in K$ and let $\varphi: \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a homeomorphism with $\varphi(K) = \overline{\mathbb{B}}_1(0)$, $\varphi(x) = 0$ and $\varphi(K')$ star-shaped with respect to 0. It is easy to see that $(\varphi(K'), \varphi(K))$ is an NDR-pair and hence it follows from Proposition 5.14 that the pair (K', K) satisfies the homotopy extension property.

Finally, for arbitrary compact convex sets $K \subseteq K'$ we factor the inclusion as $K \hookrightarrow \text{aff } K \cap K' \hookrightarrow K'$. The claim now follows from the previous two cases together with the fact that if two pairs (X, Y) and (Y, Z) satisfy the homotopy extension property, then so does the pair (X, Z) . \square

By using a truncation argument, we extend this result to any pair of closed convex sets.

Corollary 5.19. *Let K and K' be closed and convex sets in \mathbb{R}^d with $K \subseteq K'$. Then the pair (K', K) satisfies the homotopy extension property.*

Proof. We verify Definition 5.7: Let $f: K' \rightarrow Y$ be any continuous map and $H: K \times [0, 1] \rightarrow Y$ be any homotopy with $H(-, 0) = f|_K$. We inductively construct an extension $\tilde{H}: K' \times [0, 1] \rightarrow Y$ of H with $\tilde{H}(-, 0) = f$. To this end, consider for every $n \in \mathbb{N}$ the compact convex sets $K_n = D_n(0) \cap K \subseteq K$ and $K'_n = D_n(0) \cap K' \subseteq K'$, where $D_n(0)$ is the closed ball of radius n centered at the origin. Denote by $f_n: K'_n \rightarrow Y$ and $H_n: K_n \times [0, 1] \rightarrow Y$ the restrictions of f and H , respectively.

By Proposition 5.18, the pair (K'_1, K_1) satisfies the homotopy extension property and hence there exists an extension $\tilde{H}_1: K'_1 \times [0, 1] \rightarrow Y$ of H_1 that satisfies $\tilde{H}_1(-, 0) = f_1$.

Let $n \in \mathbb{N}$ be arbitrary and consider the following diagram of inclusions:

$$\begin{array}{ccc}
 K'_n & \xrightarrow{\quad\quad\quad} & K'_{n+1} \\
 & \searrow & \nearrow \\
 & K'_n \cup K_{n+1} & \\
 & \nearrow & \searrow \\
 K_n & \xrightarrow{\quad\quad\quad} & K_{n+1}
 \end{array}$$

By Proposition 5.15 and Proposition 5.18, the pair $(K'_{n+1}, K'_n \cup K_{n+1})$ satisfies the homotopy extension property and hence we can extend the homotopy on $K'_n \cup K_{n+1}$ induced by (\tilde{H}_n, H_{n+1}) to a homotopy $\tilde{H}_{n+1}: K'_{n+1} \times [0, 1] \rightarrow Y$ that satisfies $\tilde{H}_{n+1}(-, 0) = f_{n+1}$. Taking to the colimit over all n yields the desired extension \tilde{H} . \square

Let $X \subset \mathbb{R}^d$ be a subset and let $\mathcal{B} = (C_i)_{i \in [n]}$ be a finite cover of X by closed convex sets. The previous corollary shows, together with the fact that any closed subset of \mathbb{R}^d is compactly generated (Example 5.2), that the assumptions in Corollary 5.16 are satisfied and hence, the map $\rho_S: \text{Blowup}(\mathcal{B}) \rightarrow X$ is a homotopy equivalence. As any cover by convex sets is good, it follows from 2(a) in Theorem 5.9 that the map $\rho_N: \text{Blowup}(\mathcal{B}) \rightarrow |\text{Nrv}(\mathcal{B})|$ is a homotopy equivalence, as well. This proves Theorem 3.9.

If in the functorial nerve theorem for covers by subcomplexes (Theorem 4.8) we additionally assume that the cover is locally finite dimensional, then this theorem also

follows readily from the unified nerve theorem (Theorem 5.9): the realization of a simplicial complex is compactly generated (Example 5.2); moreover, the latching spaces are subcomplexes and hence the homotopy extension properties are satisfied (Remark 5.8).

5.2. Simplicial Model Categories

In order to prove Theorem 5.9, we will need a generalization of the bar construction to other settings than the category of topological spaces. To make sense of the homotopy invariance property in other settings, we will need a general framework in which to understand the analogues of homotopy equivalences in other contexts. There are many choices for such frameworks: we will work with *model categories*, which have been a standard tool of abstract homotopy theory since they were introduced by Quillen in the 1960s. A thorough introduction to model categories is beyond the scope of this paper (see, e.g., [DS95] for a friendly introduction), but we will briefly introduce the aspects of model categories that are most relevant to this paper.

Model categories A model category is a category, together with three distinguished classes of morphisms, the *weak equivalences*, *fibrations*, and *cofibrations*, which are required to satisfy certain axioms. An admissible choice of these classes is called a *model structure* on the underlying category. The choice of morphisms define two distinguished classes of objects: an object X is *fibrant* if the unique map from X to the terminal object is a fibration, and it is *cofibrant* if the unique map from the initial object is a cofibration. Before we give the axioms, it is useful to have in mind some basic examples.

Example 5.20. There are several important model categories whose objects are topological spaces. Usually, one considers some subcategory of \mathbf{Top} in order to avoid pathological behavior. In the literature, there are a number of subcategories that algebraic topologists like to work with; we will choose the subcategory of compactly generated spaces. There is a model structure on the category of compactly generated spaces for which the weak equivalences are the homotopy equivalences, and the cofibrations are the *Hurewicz cofibrations*, which are the maps $i: A \rightarrow X$ that satisfy the homotopy extension property (see Definition 5.7, and replace the inclusion $A \subset X$ with i). This is called the *Hurewicz model structure*. It was originally established (on the category of all topological spaces) by Strøm [Str72]; see [MP12, Theorem 17.1.1] for an account in the setting of compactly generated spaces. Every space is both fibrant and cofibrant in the Hurewicz model structure, which is quite rare.

There is another model structure on the category of compactly generated spaces, called the *Quillen model structure*, for which the weak equivalences are the weak homotopy equivalences, i.e., a map that induces a bijection on path components and an isomorphism on homotopy groups for all choices of base point. This was first studied by Quillen in his original work on model categories [Qui67]; see [MP12, Theorem 17.2.2] for an account in our setting. Every space is fibrant in the Quillen model structure, and every CW complex is cofibrant.

Example 5.21. Model categories can be used to study homological algebra. Let R be a commutative ring. There is a model structure on the category of non-negatively

graded chain complexes of R -modules, for which the weak equivalences are the quasi-isomorphisms, and the cofibrations are those monomorphisms that have a degreewise-projective cokernel. In particular, the cofibrant objects are the degreewise-projective chain complexes. This is another of the original examples from [Qui67].

Definition 5.22. A *model category* \mathcal{M} is a category which is equipped with three subcategories of morphisms called *weak equivalences*, *fibrations* and *cofibrations* such that the following axioms hold:

1. The category \mathcal{M} has all small limits and colimits.
2. (2-of-3) If f and g are maps of \mathcal{M} such that $g \circ f$ is defined and two of the maps f , g , $g \circ f$ are weak equivalences, then so is the third.
3. If f is a retract of g and g is a weak equivalence, fibration, or cofibration, then so is f .
4. Given a commutative square

$$\begin{array}{ccc} A & \xrightarrow{f} & X \\ i \downarrow & & \downarrow p \\ B & \xrightarrow{g} & Y, \end{array}$$

where i is a cofibration and p is a fibration, then there is a map $h: B \rightarrow X$ such that $f = h \circ i$ and $g = p \circ h$ if either of i or p is a weak equivalence.

5. Any map f can be factored as (i) $f = p \circ i$, where i is a cofibration and p is a fibration and a weak equivalence, and (ii) $f = p' \circ i'$ where i' is a cofibration and a weak equivalence and p' is a fibration.

Remark 5.23. The definition of model category has evolved since it was first introduced. For example, we require all small limits and colimits, while Quillen originally required only all finite limits and colimits. For a discussion, see [Hov99, Chapter 1]

Remark 5.24. In a model category the weak equivalences together with the fibrations or the cofibrations determine the third subcategory; see, e.g., [Hov99, Lemma 1.1.10].

Many algebraic topologists prefer to work with certain kinds of combinatorial models of spaces, rather than with topological spaces themselves. These combinatorial models are called *simplicial sets*, and they are somewhat like simplicial complexes. While they are more complicated than simplicial complexes – for example, every simplicial set has simplices in every dimension, even the simplicial set that models the one-point space! – they have better categorical properties: for example, there is a geometric realization functor $|-|$ from the category of simplicial sets to the category of compactly generated topological spaces, and this functor preserves all small colimits, and all finite limits (we define this construction in Section 5.3). So, one can take limits and colimits in the category of simplicial sets, and these will model the corresponding limits and colimits of topological spaces. See, for example, [Fri12] for a friendly introduction to this topic.

Definition 5.25. Let Δ denote the *simplex category* that has as objects the finite ordinals $\{[n] = \{0, 1, \dots, n\} \mid n \geq 0\}$ and the morphisms are given by order preserving maps.

Definition 5.26. A *simplicial set* is a functor $X: \Delta^{\text{op}} \rightarrow \mathbf{Set}$, and a *morphism of simplicial sets* is a natural transformation. The set $X_n = X([n])$ is the set of *n-simplices* of X . The category of simplicial sets is denoted by \mathbf{sSet} . More generally, if \mathcal{C} is any category, a *simplicial object in \mathcal{C}* is a functor $\Delta^{\text{op}} \rightarrow \mathcal{C}$.

Example 5.27. The *Yoneda embedding* $Y: \Delta \hookrightarrow \mathbf{sSet}$, $[n] \mapsto \text{Hom}_{\Delta}(-, [n])$ gives rise to a family of simplicial sets. We denote $Y([n])$ by Δ^n and call it the *standard n-simplex*.

Example 5.28. Let X be a topological space. The *singular simplicial set of X* is the simplicial set $\text{Sing}(X)$ with

$$\text{Sing}(X)([n]) = \text{hom}(|\Delta^n|, X),$$

where $|\Delta^n|$ is the standard topological n -simplex, and $\text{hom}(-, -)$ denotes the set of continuous maps.

A fundamental fact about the relationship between simplicial sets and topological spaces is that the functor $\text{Sing}: \mathbf{CGSpc} \rightarrow \mathbf{sSet}$ is right adjoint to the geometric realization $|-|: \mathbf{sSet} \rightarrow \mathbf{CGSpc}$ mentioned above. This adjunction is what allows us to use simplicial sets as a model for spaces; we say more about this below.

Example 5.29. Let \mathcal{C} be a category. The *nerve of \mathcal{C}* is the simplicial set $N(\mathcal{C})$ such that

$$N(\mathcal{C})([n]) = \{v_0 \rightarrow v_1 \rightarrow \dots \rightarrow v_n \mid \text{string of composable morphisms in } \mathcal{C}\}.$$

If $\mathcal{A} = (A_i)_{i \in I}$ is a cover of a topological space, then the finite non-empty intersections of cover elements form a category with morphisms given by inclusion $A_J \hookrightarrow A_{J'}$ if $J' \subseteq J$. The geometric realization of the nerve of this category is homeomorphic to $|\text{Sd Nrv}(\mathcal{A})|$.

We can now introduce two more examples of model categories, both of which play a role in the proof of Theorem 5.9.

Example 5.30. The category of simplicial sets can be equipped with a model structure, also called the *Quillen model structure*, where the weak equivalences are those maps that become weak homotopy equivalences after applying the geometric realization functor and the cofibrations are the monomorphisms; see, for example, [GJ09, Chapter I]. From the definition of cofibration, it follows that every simplicial set is cofibrant.

Example 5.31. An alternative to the chain complexes of Example 5.21 is the category of simplicial R -modules. Here, R is a commutative ring as before, and a simplicial R -module is a simplicial object in the category $R\text{-Mod}$ of R -modules. We let $R: \mathbf{sSet} \rightarrow \mathbf{s}(R\text{-Mod})$ denote the functor that is induced by the free R -module functor $R: \mathbf{Set} \rightarrow R\text{-Mod}$; the forgetful functor $U: \mathbf{s}(R\text{-Mod}) \rightarrow \mathbf{sSet}$ is right adjoint to R . Then there is a model structure on $\mathbf{s}(R\text{-Mod})$ such that the weak equivalences and fibrations are exactly those morphisms whose underlying map of simplicial sets is a weak equivalence and fibration, respectively [GS07, Proposition 4.2]. Moreover, a continuous map $X \rightarrow Y$ is an R -homology isomorphism if and only if the induced map $R(\text{Sing}(X)) \rightarrow R(\text{Sing}(Y))$ is a weak equivalence of simplicial R -modules [Wei94, Dold-Kan Theorem 8.4.1].

We have now encountered two important adjunctions connecting model categories: the adjunction $(|-|, \text{Sing})$ relating spaces with simplicial sets, and (R, U) relating simplicial sets with simplicial R -modules. In general, a *Quillen adjunction* between model categories is an adjunction such that the left adjoint preserves cofibrations and trivial cofibrations, or equivalently, the right adjoint preserves fibrations and trivial fibrations. These adjunctions are the main way to relate model categories; both of the adjunctions just mentioned are Quillen adjunctions.

A Quillen adjunction (F, G) , where the left adjoint F is a functor $\mathcal{M} \rightarrow \mathcal{N}$, is a *Quillen equivalence* if, for all cofibrant X in \mathcal{M} and all fibrant Y in \mathcal{D} , a map $FX \rightarrow Y$ is a weak equivalence in \mathcal{N} if and only if the corresponding map $X \rightarrow GY$ is a weak equivalence in \mathcal{M} . The adjunction $(|-|, \text{Sing})$ is a Quillen equivalence when CGSpc is given the Quillen model structure; see for example [Hov99, Theorem 2.4.25, Theorem 3.6.7]. This simple definition has powerful consequences, and we now describe one that plays a role in the proof of Theorem 5.9. Since (F, G) is an adjunction, there is a natural map $\eta: X \rightarrow GFX$ called the *unit*, and another $\varepsilon: FGY \rightarrow Y$ called the *counit*. If (F, G) is a Quillen equivalence, then these maps are weak equivalences, subject to additional fibrancy and cofibrancy assumptions. See [Hov99, Proposition 1.3.13] for details. Once you know that $(|-|, \text{Sing})$ is a Quillen equivalence, then it follows immediately that the unit $K \rightarrow \text{Sing}(|K|)$ is a weak equivalence for every simplicial set K , and the counit $|\text{Sing}(Y)| \rightarrow Y$ is a weak equivalence for every compactly-generated space Y . The additional fibrancy and cofibrancy assumptions are vacuous in this case, as every simplicial set is cofibrant and every space is fibrant.

Simplicial model categories In the beginning of this section, we discussed the importance of mapping spaces; a simplicial model category \mathcal{M} is a model category equipped with additional structure that generalizes this feature of algebraic topology; see [Rie14, Definition 11.4.4] for a precise definition. For any two objects X and Y of a simplicial model category \mathcal{M} , we have a simplicial set $\mathbf{Hom}_{\mathcal{M}}(X, Y)$ that encodes homotopy-theoretic information about X and Y . Formally, one requires that the model category \mathcal{M} be enriched in simplicial sets, and tensored and cotensored. One then imposes an additional axiom that relates this structure to the model structure.

We will omit the axioms, since we will not use most of the structure explicitly. However, we will use the tensoring in order to define the bar construction in a simplicial model category, and we introduce this now. If \mathcal{M} is a simplicial model category, then for any object X of \mathcal{M} and any simplicial set K , there is an object $X \otimes K$ of \mathcal{M} , and this construction gives a functor $\mathcal{M} \times \mathbf{sSet} \rightarrow \mathcal{M}$. Furthermore, for any X , the functor $\mathbf{Hom}_{\mathcal{M}}(X, -): \mathcal{M} \rightarrow \mathbf{sSet}$ has a left adjoint given by $X \otimes -: \mathbf{sSet} \rightarrow \mathcal{M}$.

Example 5.32. The category CGSpc of compactly-generated spaces is enriched in simplicial sets, and tensored and cotensored. This makes CGSpc a simplicial model category with either the Hurewicz or Quillen model structures. If X and Y are compactly-generated spaces, $\mathbf{Hom}(X, Y)$ is the simplicial set with

$$\mathbf{Hom}(X, Y)_n = \text{hom}(X \times |\Delta^n|, Y).$$

So, the zero-simplices of $\mathbf{Hom}(X, Y)$ are maps from X to Y , the one-simplices are homotopies, the two-simplices are “homotopies between homotopies”, and so on. The operation \otimes is characterized by $X \otimes \Delta^n = X \times |\Delta^n|$.

Example 5.33. The Quillen model structure on \mathbf{sSet} gives a simplicial model category, and the operation \otimes is the cartesian product.

Example 5.34. The category $\mathbf{s}(R\text{-Mod})$ is a simplicial model category, with the model structure described in Example 5.31. If K is a simplicial set and M is a simplicial R -module, then $M \otimes K$ is the simplicial R -module with $(M \otimes K)_n = M_n \otimes_R R(K_n)$.

5.3. Proof of the Unified Nerve Theorem

We can now define the bar construction in the setting of a simplicial model category, generalizing the construction of Section 2.

Definition 5.35. Let P be a poset and let \mathcal{M} be a simplicial model category. The *simplicial bar construction* of a functor $F: P \rightarrow \mathcal{M}$ is the simplicial object

$$\mathrm{Bar}_\bullet(P, F): \Delta^{\mathrm{op}} \rightarrow \mathcal{M}$$

whose n -simplices $\mathrm{Bar}_n(P, F)$ are defined by the coproduct

$$\mathrm{Bar}_n(P, F) = \coprod_{v_0 \leq v_1 \leq \dots \leq v_n} F(v_0).$$

Equivalently, the coproduct is indexed by functors of the form $\gamma: [n] \rightarrow P$. For any map $\theta: [m] \rightarrow [n]$ in Δ , $\theta^*: \mathrm{Bar}_n(P, F) \rightarrow \mathrm{Bar}_m(P, F)$ takes the summand indexed by γ to the summand indexed by $\gamma \circ \theta$, via the map $F(\gamma(0)) \rightarrow F(\gamma(\theta(0)))$.

The identifications that were used to define the bar construction for topological spaces are achieved in this setting by the categorical notion of a *coend*:

Definition 5.36. Let \mathcal{C} be a small category, \mathcal{E} any category, and $H: \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathcal{E}$ a functor. The *coend* $\int^{\mathcal{C}} H$, sometimes denoted $\int^{c \in \mathcal{C}} H(c, c)$, is an object of \mathcal{E} equipped with arrows $H(c, c') \rightarrow \int^{\mathcal{C}} H$ for each $c \in \mathcal{C}$ that are collectively universal with the property that the diagram

$$\begin{array}{ccc} H(c', c) & \xrightarrow{f_*} & H(c', c') \\ f^* \downarrow & & \downarrow \\ H(c, c) & \longrightarrow & \int^{\mathcal{C}} H \end{array}$$

commutes for each $f: c \rightarrow c'$ in \mathcal{C} .

We can use this notion to define the geometric realization functor $|-|: \mathbf{sSet} \rightarrow \mathbf{CGSpc}$ mentioned in Section 5.2. Writing Δ for the simplex category as before, there is a functor $\Delta \rightarrow \mathbf{CGSpc}$ that takes $[n]$ to the standard n -simplex. If $X: \Delta^{\mathrm{op}} \rightarrow \mathbf{Set}$ is a simplicial

set, $|X|$ is the coend of the functor $\Delta^{\text{op}} \times \Delta \rightarrow \mathbf{CGSpc}$ that takes $([n], [m])$ to $X_n \times |\Delta^m|$, where X_n has the discrete topology. This is written

$$|X| = \int^{[n] \in \Delta} X_n \times |\Delta^n|.$$

We now apply the same idea to the simplicial bar construction:

Definition 5.37. Let P be a poset and let \mathcal{M} be a simplicial model category. The *bar construction* of a functor $F: P \rightarrow \mathcal{M}$ is the coend

$$\text{Bar}(P, F) = \int^{[n] \in \Delta} \text{Bar}_n(P, F) \otimes \Delta^n.$$

Note that there are canonical maps $\text{Bar}_n(P, F) \otimes \Delta^n \rightarrow \text{Bar}_n(P, F) \rightarrow \text{colim } F$ for all $n \geq 0$, which induce a map $\text{Bar}(P, F) \rightarrow \text{colim } F$ by the universal property of the coend.

Remark 5.38. Since the category \mathbf{CGSpc} of compactly generated spaces is a simplicial model category (with either the Hurewicz or Quillen model structures), we can consider the bar construction $\text{Bar}(P, F)$ of any functor $F: P \rightarrow \mathbf{CGSpc}$. Abusing notation, if $F: P \rightarrow \mathbf{Top}$ is a functor valued in the category of all topological spaces, we can define the bar construction $\text{Bar}(P, F)$ by first building the simplicial object $\text{Bar}_\bullet(P, F): \Delta^{\text{op}} \rightarrow \mathbf{Top}$ as in Definition 5.35, and then taking the coend

$$\text{Bar}(P, F) = \int^{[n] \in \Delta} \text{Bar}_n(P, F) \times |\Delta^n|$$

in \mathbf{Top} .

We now explain why, if $F: P \rightarrow \mathbf{CGSpc}$ is a functor valued in compactly generated spaces, then it does not matter whether we compute $\text{Bar}(P, F)$ in \mathbf{CGSpc} or \mathbf{Top} .

Following Dugger–Isaksen [DI04, Appendix A], given $F: P \rightarrow \mathbf{Top}$, we can compute $\text{Bar}(P, F)$ as a sequential colimit of pushouts, as follows. For $k = 0$, we define $\text{Bar}(P, F)(0) = \coprod_{v \in P} F(v)$, and for $k > 0$ we inductively define $\text{Bar}(P, F)(k)$ as the pushout

$$\begin{array}{ccc} \coprod_{v_0 < \dots < v_k} F(v_0) \times |\partial \Delta^k| & \longrightarrow & \text{Bar}(P, F)(k-1) \\ \downarrow & \lrcorner & \vdots f_{k-1} \\ \coprod_{v_0 < \dots < v_k} F(v_0) \times |\Delta^k| & \dashrightarrow & \text{Bar}(P, F)(k) \end{array}$$

where the top horizontal map is defined using the face maps $\text{Bar}_k(P, F) \rightarrow \text{Bar}_{k-1}(P, F)$. Then we have an isomorphism $\text{Bar}(P, F) \cong \text{colim}_k \text{Bar}(P, F)(k)$.

Now, if $F: P \rightarrow \mathbf{CGSpc}$ is a functor valued in compactly generated spaces, we can use this characterization of the bar construction computed in \mathbf{Top} to show that the result is in fact compactly generated, as follows. As the disjoint union of compactly generated spaces is compactly generated and because $|\partial \Delta^n|$ as well as $|\Delta^n|$ are locally compact Hausdorff, it follows from [Str09, Proposition 2.6. and Corollary 2.16] that the

spaces on the left hand side of the diagram and $\text{Bar}(P, F)(0)$ are compactly generated. Moreover, because the pushout of compactly generated spaces along a closed inclusion is again compactly generated [May99, p.40], it follows that the spaces $\text{Bar}(P, F)(k)$ are compactly generated. By [Str09, Proposition 2.35], the maps f_k are closed inclusions. Finally, it follows again from [May99, p.40] that $\text{Bar}(P, F) \cong \text{colim}_k \text{Bar}(P, F)(k)$ is compactly generated.

Since $\text{Bar}(P, F)$ computed in \mathbf{Top} is compactly generated, and it satisfies the universal property of the coend in \mathbf{Top} , it also satisfies this universal property in \mathbf{CGSpc} .

Furthermore, one can use this method for building the bar construction as a sequential colimit of pushouts to check that, given $F: P \rightarrow \mathbf{Top}$, the bar construction we consider in this section is naturally homeomorphic to the bar construction of Definition 2.1, which justifies using the same notation in both places.

Remark 5.39. The coend construction used to define $\text{Bar}(P, F)$ in Definition 5.37 is an example of the *geometric realization* of a simplicial space. We note that the blowup complex of Definition 2.7 can also be seen as the geometric realization of a simplicial space, the ordered Čech complex of [DI04, Section 2.5].

The map $\pi_{\text{Sd}N}: \text{Bar}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}}) \rightarrow |\text{Sd Nrv}(\mathcal{U})|$ is the geometric realization of a map $\text{Bar}_{\bullet}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}}) \rightarrow \text{Bar}_{\bullet}(P_{\mathcal{U}}, *)$ of simplicial spaces, and the map $\pi_S: \text{Bar}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}}) \rightarrow X$ is also the geometric realization of a map of simplicial spaces $\text{Bar}_{\bullet}(P_{\mathcal{U}}, \mathcal{D}_{\mathcal{U}}) \rightarrow X_{\bullet}$, where $X_n = X$ for all n . Analogously, the maps $\rho_N: \text{Blowup}(\mathcal{U}) \rightarrow |\text{Nrv}(\mathcal{U})|$ and $\rho_S: \text{Blowup}(\mathcal{U}) \rightarrow X$ can be seen as geometric realizations of maps of simplicial spaces.

Example 5.40. If we leave aside the requirement that we work in a simplicial model category, the PoBar construction from Section 4 is a bar construction. In more detail, let K be a simplicial complex, and let $\mathcal{A} = (K_i)_{i \in I}$ be a cover by subcomplexes. Let $\mathcal{D}_{\mathcal{A}}: P_{\mathcal{A}} \rightarrow \mathbf{Po}$ be the functor with $\mathcal{D}_{\mathcal{A}}(J) = U(\cap_{i \in J} K_i)$. There is a simplicial object $\text{Bar}_{\bullet}(P_{\mathcal{A}}, \mathcal{D}_{\mathcal{A}})$ in \mathbf{Po} defined as in Definition 5.35, and the inclusion $\Delta \subset \mathbf{Po}$ defines a cosimplicial object, i.e., a functor $\Delta \rightarrow \mathbf{Po}$. Then

$$\text{PoBar}(\mathcal{A}) = \int^{[n]} \text{Bar}_n(P_{\mathcal{A}}, \mathcal{D}_{\mathcal{A}}) \times [n].$$

Given a sufficiently well-behaved diagram $F: P \rightarrow \mathcal{M}$ in a simplicial model category \mathcal{M} , the bar construction of F computes the homotopy colimit of F . This appears as [Rie14, Corollary 5.1.3]. In the proof of Theorem 5.9, we will use two consequences of this result. The first says that the bar construction is homotopical for pointwise cofibrant diagrams:

Proposition 5.41. *Let \mathcal{M} be a simplicial model category, let P be a poset, and let $F, F': P \rightarrow \mathcal{M}$ be pointwise cofibrant diagrams. If $\sigma: F \Rightarrow F'$ is a natural transformation such that the component σ_x is a weak equivalence for all elements x of P , then the induced map $\text{Bar}(P, F) \rightarrow \text{Bar}(P, F')$ is a weak equivalence.*

In Proposition 2.6, we saw already that the bar construction respects pointwise homotopy equivalences between topological spaces (without any pointwise cofibrancy or compactly-generated assumptions!). There is a similar result for weak homotopy equivalences, which follows from work of Dugger–Isaksen [DI04]:

Proposition 5.42. *Let $F, G: P \rightarrow \mathbf{Top}$ be diagrams of topological spaces. Suppose there is a natural transformation $F \Rightarrow G$ that is a pointwise weak homotopy equivalence. Then the induced map $\mathrm{Bar}(P, F) \rightarrow \mathrm{Bar}(P, G)$ is a weak homotopy equivalence as well.*

Proof. The natural transformation $F \Rightarrow G$ induces a map $\mathrm{Bar}_\bullet(P, F) \rightarrow \mathrm{Bar}_\bullet(P, G)$ such that $\mathrm{Bar}_n(P, F) \rightarrow \mathrm{Bar}_n(P, G)$ is a weak homotopy equivalence for all $n \geq 0$. It is straightforward to check that the simplicial bar construction has free degeneracies in the sense of [DI04, Definition A.4], and therefore the induced map $\mathrm{Bar}(P, F) \rightarrow \mathrm{Bar}(P, G)$ is a weak homotopy equivalence by [DI04, Corollary A.6]. \square

We will soon use this last result to prove the “weak” version of Theorem 5.9 2(a). If we did not know this result, and applied only Proposition 5.41, we would need the additional assumption that the intersection A_J is cofibrant in the Quillen model structure on \mathbf{CGSpc} for all $J \in \mathrm{Nrv}(\mathcal{A})$.

For the proof of Theorem 5.9, we will also need a second consequence of the general fact that the bar construction computes the homotopy colimit. This is similar to [WZZ99, Lemma 4.5], for example. Recall that $\mathcal{D}_\mathcal{A}: P_\mathcal{A} \rightarrow \mathbf{Top}$ denotes the nerve diagram of the cover \mathcal{A} of a topological space X .

Proposition 5.43. *Let X be a compactly generated space, and let \mathcal{A} be a closed cover that is locally finite and locally finite dimensional. If for all $T \in \mathrm{Nrv}(\mathcal{A})$ the latching space $L(T) \subseteq A_T$ is closed and the pair $(A_T, L(T))$ satisfies the homotopy extension property, then the natural map $\mathrm{Bar}(P_\mathcal{A}, \mathcal{D}_\mathcal{A}) \rightarrow \mathrm{colim} \mathcal{D}_\mathcal{A} \cong X$ is a homotopy equivalence.*

Proof. A closed subspace of a compactly generated space is also compactly generated, so $\mathcal{D}_\mathcal{A}: P_\mathcal{A} \rightarrow \mathbf{Top}$ takes values in the subcategory \mathbf{CGSpc} of compactly generated spaces.

As \mathcal{A} is locally finite dimensional, the poset $P_\mathcal{A}$ is an upwards-directed Reedy category, with $\mathrm{deg}(J) = \sup\{(|J'| - |J|) \mid J' \in \mathrm{Nrv}(\mathcal{A}) \text{ with } J \subseteq J'\}$. Working with the Hurewicz model structure on \mathbf{CGSpc} , the Reedy model structure on the functor category $\mathrm{Fun}(P_\mathcal{A}, \mathbf{CGSpc})$ coincides with the projective model structure as $P_\mathcal{A}$ is upwards-directed. In particular, the projective model structure on $\mathrm{Fun}(P_\mathcal{A}, \mathbf{CGSpc})$ exists. See [Hir03, Chapter 15] for Reedy categories and the Reedy model structure; the relationship with the projective model structure is explained at [Dug, Proposition 13.12].

The condition on the latching spaces implies that all inclusions $L(T) \subseteq A_T$ are Hurewicz cofibrations, so that $\mathcal{D}_\mathcal{A}$ is Reedy cofibrant, i.e., projective cofibrant. As the bar construction $\mathrm{Bar}(P_\mathcal{A}, \mathcal{D}_\mathcal{A})$ computes the homotopy colimit of $\mathcal{D}_\mathcal{A}$ [Rie14, Corollary 5.1.3], the natural map $\mathrm{Bar}(P_\mathcal{A}, \mathcal{D}_\mathcal{A}) \rightarrow \mathrm{colim} \mathcal{D}_\mathcal{A}$ is a homotopy equivalence. As the cover is locally finite and X is compactly generated, it follows from [Str09, Corollary 2.23] that the colimit calculated in \mathbf{CGSpc} coincides with the one in \mathbf{Top} , and by Remark 2.4 this is naturally homeomorphic to X . \square

We are now ready to prove the unified nerve theorem.

Proof of Theorem 5.9. In Section 2.2, we explained how one can prove that the natural maps ρ_S and ρ_N from the blowup complex are equivalences by proving that the natural

maps π_S and $\pi_{\text{sd}N}$ from the bar construction are equivalences. So, we work with the bar construction in this proof.

The first part of 1(a) follows from work of Dugger and Isaksen [DI04, Theorem 2.1 and Proposition 2.7]; in Appendix D we give a short proof using their ideas. The second part of 1(a) is the content of [Hat02, Proposition 4G.2]. Statement 1(b) is the content of Proposition 5.43.

We now prove 2(a). By assumption, the unique natural transformation $\mathcal{D}_{\mathcal{A}} \Rightarrow *$ from the nerve diagram of the cover \mathcal{A} to the constant diagram on the one-point space is a pointwise (weak) homotopy equivalence. The results now follow from Proposition 2.6 and Proposition 5.42, respectively. Alternatively, see Appendix D for a short proof of the fact that ρ_N is a weak homotopy equivalence whenever \mathcal{A} is a weakly good cover.

We now prove 2(b). For every compactly generated space Z , there is a natural weak homotopy equivalence $|\text{Sing}(Z)| \rightarrow Z$, given by the counit of the adjunction $(|-|, \text{Sing})$ as explained in Section 5.2. So, there is a pointwise weak homotopy equivalence

$$|-| \circ \text{Sing} \circ \mathcal{D}_{\mathcal{A}} \Rightarrow \mathcal{D}_{\mathcal{A}},$$

that induces, by Proposition 5.42, a weak homotopy equivalence

$$\text{Bar}(P_{\mathcal{A}}, |-| \circ \text{Sing} \circ \mathcal{D}_{\mathcal{A}}) \rightarrow \text{Bar}(P_{\mathcal{A}}, \mathcal{D}_{\mathcal{A}}). \quad (5.1)$$

We work with the model structure on the category $\mathfrak{s}(R\text{-Mod})$ of simplicial R -modules described in Example 5.31; recall that a map $X \rightarrow Y$ of compactly-generated spaces is an R -homology isomorphism if and only if the induced map $\mathbf{R}(\text{Sing}(X)) \rightarrow \mathbf{R}(\text{Sing}(Y))$ is a weak equivalence of simplicial R -modules. By our assumption that the natural transformation $\mathcal{D}_{\mathcal{A}} \Rightarrow *$ is a pointwise R -homology isomorphism, the natural transformation $\mathbf{R} \circ \text{Sing} \circ \mathcal{D}_{\mathcal{A}} \Rightarrow \mathbf{R} \circ \text{Sing} \circ *$ is a pointwise weak equivalence of simplicial R -modules. As \mathbf{R} preserves cofibrant objects (since it is the left adjoint of a Quillen adjunction), and every simplicial set is cofibrant, both diagrams are pointwise cofibrant. So, by Proposition 5.41, the induced map $\text{Bar}(P_{\mathcal{A}}, \mathbf{R} \circ \text{Sing} \circ \mathcal{D}_{\mathcal{A}}) \rightarrow \text{Bar}(P_{\mathcal{A}}, \mathbf{R} \circ \text{Sing} \circ *)$ is a weak equivalence. Furthermore, for any poset P and any functor $F: P \rightarrow \mathfrak{s}\text{Set}$, there is a natural isomorphism $\text{Bar}(P, \mathbf{R} \circ F) \cong \mathbf{R}(\text{Bar}(P, F))$, using the definition of the tensor structure on $\mathfrak{s}(R\text{-Mod})$ and the fact that \mathbf{R} preserves colimits (as it is a left adjoint). So we have a commutative diagram:

$$\begin{array}{ccc} \text{Bar}(P_{\mathcal{A}}, \mathbf{R} \circ \text{Sing} \circ \mathcal{D}_{\mathcal{A}}) & \xrightarrow{\cong} & \mathbf{R}(\text{Bar}(P_{\mathcal{A}}, \text{Sing} \circ \mathcal{D}_{\mathcal{A}})) \\ \simeq \downarrow & & \downarrow \\ \text{Bar}(P_{\mathcal{A}}, \mathbf{R} \circ \text{Sing} \circ *) & \xrightarrow{\cong} & \mathbf{R}(\text{Bar}(P_{\mathcal{A}}, \text{Sing} \circ *)) \end{array}$$

and hence, the morphism on the right is a weak equivalence by 2-of-3.

For any simplicial set K , the unit map $K \rightarrow \text{Sing}(|K|)$ is a natural weak equivalence, as explained in Section 5.2. The functor \mathbf{R} preserves all weak equivalences, as every simplicial set is cofibrant [Hov99, Lemma 1.1.12 (Ken Brown's lemma)]. So we have a

commutative square, in which the indicated maps are weak equivalences:

$$\begin{array}{ccc}
\mathbf{R}(\mathrm{Bar}(P_{\mathcal{A}}, \mathrm{Sing} \circ \mathcal{D}_{\mathcal{A}})) & \xrightarrow{\cong} & \mathbf{R}(\mathrm{Sing}(|\mathrm{Bar}(P_{\mathcal{A}}, \mathrm{Sing} \circ \mathcal{D}_{\mathcal{A}})|)) \\
\cong \downarrow & & \downarrow \\
\mathbf{R}(\mathrm{Bar}(P_{\mathcal{A}}, \mathrm{Sing} \circ *)) & \xrightarrow[\cong]{} & \mathbf{R}(\mathrm{Sing}(|\mathrm{Bar}(P_{\mathcal{A}}, \mathrm{Sing} \circ *)|))
\end{array}$$

It follows, by 2-of-3, that the fourth map in the square is a weak equivalence as well, and so $|\mathrm{Bar}(P_{\mathcal{A}}, \mathrm{Sing} \circ \mathcal{D}_{\mathcal{A}})| \rightarrow |\mathrm{Bar}(P_{\mathcal{A}}, \mathrm{Sing} \circ *)|$ is an R -homology isomorphism.

For any poset P and any functor $F: P \rightarrow \mathbf{sSet}$, there is a natural isomorphism $\mathrm{Bar}(P, | - | \circ F) \cong |\mathrm{Bar}(P, F)|$, again using the definitions and the fact that geometric realization preserves colimits, being a left adjoint. So we have the following commutative diagram:

$$\begin{array}{ccc}
|\mathrm{Bar}(P_{\mathcal{A}}, \mathrm{Sing} \circ \mathcal{D}_{\mathcal{A}})| & \xrightarrow{\sim_R} & |\mathrm{Bar}(P_{\mathcal{A}}, \mathrm{Sing} \circ *)| \\
\downarrow \cong & & \downarrow \cong \\
\mathrm{Bar}(P_{\mathcal{A}}, | - | \circ \mathrm{Sing} \circ \mathcal{D}_{\mathcal{A}}) & \longrightarrow & \mathrm{Bar}(P_{\mathcal{A}}, | - | \circ \mathrm{Sing} \circ *) \\
& \searrow & \uparrow \\
& & \mathrm{Bar}(P_{\mathcal{A}}, \mathcal{D}_{\mathcal{A}})
\end{array}$$

Recall that that the map $\mathrm{Bar}(P_{\mathcal{A}}, | - | \circ \mathrm{Sing} \circ \mathcal{D}_{\mathcal{A}}) \rightarrow \mathrm{Bar}(P_{\mathcal{A}}, \mathcal{D}_{\mathcal{A}})$ from 5.1 is a weak equivalence. Together with the fact that weak homotopy equivalences are also R -homology isomorphisms [Hat02, Proposition 4.21], we get that the canonical map

$$\pi_{\mathrm{Sd}N}: \mathrm{Bar}(P_{\mathcal{A}}, \mathcal{D}_{\mathcal{A}}) \rightarrow |\mathrm{Sd} \mathrm{Nrv}(\mathcal{A})| \cong \mathrm{Bar}(P_{\mathcal{A}}, *) \cong \mathrm{Bar}(P_{\mathcal{A}}, | - | \circ \mathrm{Sing} \circ *)$$

is, once more by 2-of-3, an R -homology isomorphism as well. \square

A. Auxiliary Lemmas about Geometric Simplicial Complexes

Lemma A.1. *Let $\sigma = \{v_0, \dots, v_k\} \in K$ be a simplex and consider the subcomplex $L = \{\tau_0 \subset \dots \subset \tau_m \mid \sigma \subseteq \tau_0\} \subseteq \mathrm{Sd} K$. Then $\bigcap_{i=0}^k \mathrm{bst} v_i = |L|$.*

Proof of Lemma A.1. First, let $\phi = (\tau_0 \subset \dots \subset \tau_m) \in L$ be a simplex. By definition, ϕ is contained in the simplex $\sigma \subseteq \tau_0 \subset \dots \subset \tau_m$ of $\mathrm{Sd} K$. Thus, the realization of ϕ is contained in $\mathrm{bst} v_i$ for all i , and so we have $|L| \subseteq \bigcap_{i=0}^k \mathrm{bst} v_i$.

Now, let $|\phi = (\tau_0 \subset \dots \subset \tau_m)| \subseteq \bigcap_{i=0}^k \mathrm{bst} v_i$. Since for all i we have $|\tau_0 \subset \dots \subset \tau_m| \subseteq \mathrm{bst} v_i$, we know that $v_i \in \tau_0$. Thus, the simplex σ is also contained in τ_0 . Therefore, $\phi \in L$ and so we have $\bigcap_{i=0}^k \mathrm{bst} v_i \subseteq |L|$. \square

Proof of Lemma 3.4. By Lemma A.1, every (geometric) simplex in $\bigcap_{v \in \sigma} \mathrm{bst} v \subseteq |\mathrm{Sd} K|$ has a coface in this intersection with $z(|\sigma|)$ as a vertex, where $z(|\sigma|)$ is the barycenter of $|\sigma|$. Thus, $\bigcap_{v \in \sigma} \mathrm{bst} v$ is star-shaped with respect to $z(|\sigma|)$ and hence contractible. \square

The following two lemmas are straightforward calculations (compare [ES52, p.62]).

Lemma A.2. *Let K be a simplicial complex and let $x \in |K|$. Write x in barycentric coordinates of K as*

$$x = \sum_{j=0}^m \nu_j \cdot |w_j|$$

with $w_i \in \text{Vert } K$, $\nu_i > 0$ and $\sum_{j=0}^m \nu_j = 1$ as well as $\nu_0 \geq \nu_1 \geq \dots \geq \nu_m$. Then, by using the (geometric) simplices

$$|\tau_i| = \text{conv}\{|w_0|, \dots, |w_i|\} \quad \text{for all } i \in \{0, \dots, m\} \quad (\text{A.1})$$

in the realization $|K|$ and by writing $z(|\tau_i|)$ for the barycenter of $|\tau_i|$, we have $x \in \text{conv}\{z(|\tau_0|), \dots, z(|\tau_m|)\}$. Specifically, writing x in barycentric coordinates of $\text{Sd } K$ as $x = \sum_{j=0}^m \mu_j z(|\tau_j|)$, we have

$$\begin{aligned} \mu_i &= (i+1) \left(\nu_i(x) - \nu_{i+1}(x) \right) \quad \text{for } i = 0, \dots, m-1 \\ \mu_m &= (m+1) \nu_m(x). \end{aligned}$$

Lemma A.3. *Let $x \in |\text{Sd } K|$, written in barycentric coordinates as $x = \sum_{j=0}^m \mu_j z(|\tau_j|)$ for some flag of simplices $\tau_0 \subset \dots \subset \tau_m$ in K , where*

$$|\tau_i| = \text{conv}\{|w_0|, \dots, |w_i|\} \quad \text{for all } i \in \{1, \dots, m\}$$

and $w_i \in \text{Vert } K$. Then we have $x \in |\tau_m| = \text{conv}\{|w_0|, \dots, |w_m|\}$. Specifically, the barycentric coordinates ν_i of x in K with respect to $|w_0|, \dots, |w_m|$ take the form

$$\nu_i = \sum_{j=i}^m \frac{1}{j+1} \mu_j. \quad (\text{A.2})$$

Proof of Lemma 3.5. Let $x \in |K|$ be a point satisfying Eq. (3.2). It suffices to show that x is contained in a simplex of $|\text{Sd } K|$ having $|v|$ as a vertex. Let v_0, \dots, v_m be the vertices in K with $b_{v_i}(x) > 0$ in descending order of barycentric coordinates. By Eq. (3.2) we may choose $v_0 = v$. Now, by Lemma A.2, we know that the point x is contained in $\text{conv}\{|v| = z(|\tau_0|), \dots, z(|\tau_m|)\}$ for the simplices $|\tau_i| \subseteq |K|$ specified as in Eq. (A.1). Hence, by definition the point x is contained in $\text{bst } v$.

Conversely, let $x \in \text{bst } v$ for some vertex $v \in \text{Vert } K$. Then there exists a simplex $\tau \in \text{Sd } K$ with v as a vertex such that $x \in |\tau|$ and that τ corresponds to a flag $v = \tau_0 \subset \dots \subset \tau_m$ of simplices in K . From Lemma A.3, or more specifically Eq. (A.2), we deduce that the barycentric coordinate $\nu_0 = b_v(x)$ of x in K with respect to v is maximal. \square

B. Discrete Morse Theory for Infinite Complexes

Discrete Morse theory for finite cell complexes was introduced by Forman [For95; For02] and since then found its way into algorithms, applications and underwent reformulations

as well as generalizations. For an extension to infinite cell complexes that is similar to the results presented here, see [Bat02].

Let K be a simplicial complex and $\sigma \in K$ a simplex. We call a facet $\tau \subseteq \sigma$ a *free facet* if σ is the only proper coface of τ . An *elementary collapse* $K \searrow K \setminus \{\tau, \sigma\}$ is the removal of a pair of simplices, where τ is a free facet of σ . A *collapse* $K \searrow L$ onto a subcomplex L is a sequence of elementary collapses starting in K and ending in L . Moreover, an elementary collapse can be realized continuously by a strong deformation retract and therefore collapses preserve the homotopy type.

By a *discrete vector field* on K we mean a partition V of K into singletons $\{\sigma\}$, σ is then called a *critical simplex*, and pairs $\{\sigma, \tau\}$ corresponding to arcs (σ, τ) in the Hasse diagram $\mathcal{H}(K)$ of the face poset, i.e. the directed graph whose nodes are the simplices and whose arcs are the pairs (σ, τ) in which σ is a facet of τ . We call the discrete vector field V a *discrete gradient vector field* if the graph $\mathcal{H}(K, V)$ that is obtained from $\mathcal{H}(K)$ by reversing all the arcs (σ, τ) for which $\{\sigma, \tau\} \in V$ is acyclic. Note that it suffices to check that there are no non-trivial closed V -paths, i.e. that there are no undirected paths in $\mathcal{H}(K)$ that are of the form $\tau_0 \rightarrow \mu_0 \leftarrow \cdots \rightarrow \mu_r \leftarrow \tau_{r+1}$ with $\{\tau_i, \mu_i\} \in V$, $\tau_i \neq \tau_{i+1}$ and $\tau_0 = \tau_{r+1}$.

Given a discrete gradient vector field V on a simplicial complex K , we can define a poset structure on V as follows: For two elements $A, B \in V$ we define $A \leq_V B$ if and only if there exists a sequence $A = C_0, C_1, \dots, C_n = B$ in V such that for every i there exist elements $x_{i-1} \in C_{i-1}, x_i \in C_i$ with x_{i-1} a face of x_i .

Moreover, for any element $A \in V$ we define its *height* to be

$$\text{ht}(A) = \sup\{n \in \mathbb{N} \mid \exists A = B_0 > \cdots > B_n \text{ in } V\}.$$

The following lemma is useful in practice.

Lemma B.1. *The height $\text{ht}(A)$ is finite for every $A \in V$ if and only if for every simplex $\sigma \in K$ its V -path height*

$$\text{ht}_V(\sigma) = \sup\{n \in \mathbb{N} \mid \exists V\text{-path } \sigma = \tau_0 \rightarrow \mu_0 \leftarrow \cdots \rightarrow \mu_{n-1} \leftarrow \tau_n\}$$

is finite.

Proof. Every V -path $\tau_0 \rightarrow \mu_0 \leftarrow \cdots \rightarrow \mu_{n-1} \leftarrow \tau_n$ induces a descending chain $\{\tau_0, \mu_0\} > \cdots > \{\tau_{n-1}, \mu_{n-1}\}$. Hence, if the height is finite for every $A \in V$ this implies that the V -path height is finite for every simplex in K .

For the converse, we employ induction over the dimension $\dim A = \dim \min A$ of an element $A \in V$. If $\dim A = 0$, then $\text{ht}(A) = \text{ht}_V(\min A)$ and this is finite by assumption. For the induction step, consider the set $F_{\min A}$ of all V -paths starting in $\min A$. For a gradient path $\gamma = \tau_0 \rightarrow \mu_0 \leftarrow \cdots \rightarrow \mu_{n-1} \leftarrow \tau_n$ write $\text{end } \gamma = \tau_n$ and $\text{length } \gamma = n$. Then we can bound $\text{ht}(A)$ from above as follows:

$$\text{ht}(A) \leq \max\{\text{length } \gamma + 1 + \text{ht}(B) \mid \gamma \in F_{\min A}, \sigma \subsetneq \text{end } \gamma, \sigma \in B \in V\}.$$

To complete the induction step, note that for every B as above we have $\dim B < \dim A$ and hence $\text{ht}(B)$ is finite by the induction assumption. Thus, it suffices to show that

the set $F_{\min A}$ is finite. This can be seen as follows: Given a gradient path γ as above, then the path ends in τ_i or there are $\dim \mu_i = \dim \tau_0 + 1$ choices for τ_{i+1} once τ_i is fixed. Hence, the cardinality of $F_{\min A}$ is bounded from above by $(\dim(\sigma) + 2)^{\text{ht}_V(\sigma)}$. \square

The essential ideas for the proof of the following proposition can be found already in the proof of [Bro92, Proposition 1], which predates Forman's papers.

Proposition B.2. *Let $L \subseteq K$ be a pair of simplicial complexes and let V be a discrete gradient vector field on K such that for every element $A \in V$ its height $\text{ht}(A)$ is finite. Moreover, assume that $K \setminus L$ is the union of pairs in V . Then the inclusion $|L| \hookrightarrow |K|$ is a homotopy equivalence.*

Before we prove Proposition B.2 we need one small lemma.

Lemma B.3. *Let K be a simplicial complex and $K_0 \subseteq K_1 \subseteq \dots \subseteq K$ a filtration of subcomplexes whose union is K such that for all $i \in \mathbb{N}$ the inclusion $|K_i| \hookrightarrow |K_{i+1}|$ is a homotopy equivalence. Then the inclusion $|K_0| \hookrightarrow |K|$ is also a homotopy equivalence.*

Proof. By Whitehead's theorem [Hat02, Theorem 4.5], it suffices to show that for all $n \in \mathbb{N}$ the induced morphism on homotopy groups $g_n: \pi_n(|K_0|) \rightarrow \pi_n(|K|)$ is an isomorphism. For any map $f: S^n \rightarrow |K|$ its image is compact and hence contained in some $|K_i|$. As $|K_0| \rightarrow |K_i|$ is a homotopy equivalence, it follows that the homotopy class $[f]$ is in the image of the composite $\pi_n(|K_0|) \rightarrow \pi_n(|K_i|) \rightarrow \pi_n(|K|)$ and hence also in the image of g_n . This shows surjectivity. A similar argument applied to any homotopy $h: S^n \times [0, 1] \rightarrow |K|$ shows that g_n is injective. This proves the lemma. \square

Proof of Proposition B.2. Without loss of generality, we can assume that L is the union of critical simplices.

Consider the filtration $L = K_0 \subseteq K_1 \subseteq \dots \subseteq K$ of K , where K_i is the subcomplex

$$K_i = L \cup \bigcup_{A \in V, \text{ht}(A) \leq i} A.$$

We show that for every $i \in \mathbb{N}$ the inclusion $|K_i| \hookrightarrow |K_{i+1}|$ is a homotopy equivalence and the proposition then follows from Lemma B.3: Let $\{\tau, \sigma\} = A \in V$ be any element, with τ a facet of σ , such that $\text{ht}(A) = i + 1$. Then, τ is a free facet of σ in K_{i+1} , as otherwise there would exist a pair $B \in V$ with $\text{ht}(B) \leq i + 1$ and $B > A$. But this cannot be true, because then the last property implies that the height of B satisfies $\text{ht}(B) \geq \text{ht}(A) + 1 = i + 2$, contradicting the construction. A similar argument shows that σ is not properly contained in any simplex of K_{i+1} . Therefore, the complement $K_{i+1} \setminus K_i$ is partitioned by pairs in V of height $i + 1$ and the corresponding simplices to different pairs can only possibly intersect in the subcomplex K_i . Thus, executing the elementary collapses that are encoded by those pairs simultaneously induces a strong deformation retract

$$|K_{i+1}| \rightarrow \left| K_{i+1} \setminus \bigcup_{A \in V, \text{ht}(A)=i+1} A \right| = |K_i|$$

and hence the inclusion $|K_i| \hookrightarrow |K_{i+1}|$ is a homotopy equivalence. \square

C. The Blowup Complex for Covers by Subcomplexes

In this section we show that for any simplicial complex K and any cover \mathcal{A} by subcomplexes the natural map $\rho_S: \text{Blowup}(|\mathcal{A}|) \rightarrow |K|$ is a homotopy equivalence. As explained in Remark 4.9, this proves that the locally finiteness assumption in Theorem 4.8 can be dropped.

In essence, the proof of the following lemma is the same as the proof of [Wal81, Lemma 2.1] and proved by induction over the skeleton, similarly to Proposition 3.8. Therefore, it is left to the reader.

Lemma C.1. *Let (K, L) be a pair of regular CW-complexes. Further, let Y be a topological space and let $\mathcal{U} = (U_j)_{j \in J}$ be a cover of Y . Assume that there exists a map of indexed covers $\varphi: \text{cells}(K) \rightarrow \mathcal{U}$ such that for any pair of cells $\tau \subseteq \sigma$ in K we have $\varphi(\tau) \subseteq \varphi(\sigma)$ and that $\varphi(\sigma)$ is contractible. If $(f, \varphi|_{\text{cells}(L)}): (L, \text{cells}(L)) \rightarrow (Y, \mathcal{U})$ is a morphism of covered spaces, then f extends to a continuous map $F: K \rightarrow Y$ such that $(F, \varphi): (K, \text{cells}(K)) \rightarrow (Y, \mathcal{U})$ is also a morphism of covered spaces.*

Remark C.2. In particular, by considering the pair $(K \times [0, 1], K \times \{0, 1\} \cup L \times [0, 1])$ this lemma implies that any two such extensions are homotopic relative to L . Moreover, it also implies Proposition 3.8.

Let K be a simplicial complex and let $\mathcal{A} = (A_i)_{i \in I}$ be a cover of K by subcomplexes. Note that for each $J \in \text{Nrv}(\mathcal{A})$ the product $|A_J| \times |J|$ carries the product CW-structure and so, by recalling Definition 2.7, we see that the blowup complex $\text{Blowup}(|\mathcal{A}|)$ is a regular CW-complex. Moreover, in Lemma 4.6 we constructed a triangulation for it.

Proposition C.3. *The natural map $\rho_S: \text{Blowup}(|\mathcal{A}|) \rightarrow |K|$ is a homotopy equivalence.*

Proof. Let us construct a homotopy inverse $g: |K| \rightarrow \text{Blowup}(|\mathcal{A}|)$ to ρ_S . Define the function

$$\Gamma: \{\sigma \mid \sigma \in K\} \rightarrow \mathcal{P}(I), \quad \sigma \mapsto \{i \in I \mid \sigma \in A_i\};$$

for $\tau \subseteq \sigma$ we have $\Gamma(\sigma) \subseteq \Gamma(\tau)$. Note that $\Gamma(\sigma)$ is not a simplex in $\text{Nrv}(\mathcal{A})$ if σ is contained in infinitely many cover elements. Nevertheless, all of its non-empty finite subsets are and by abuse of notation we write $|\Gamma(\sigma)| = \bigcup\{|\tau| \mid \tau \in \mathcal{P}(\Gamma(\sigma)) \cap \text{Nrv}(\mathcal{A})\} \subseteq |\text{Nrv} \mathcal{A}|$. For every simplex $\mu \in K$ define

$$B_\mu = \text{im}(|\mu| \times |\Gamma(\mu)| \rightarrow \text{Blowup}(|\mathcal{A}|)).$$

We introduce the bijective map of indexed covers

$$\varphi: \mathcal{S} := \{|\sigma| \mid \sigma \in K\} \rightarrow \mathcal{U} := \{B_\sigma \cup \bigcup_{\tau \in \partial\sigma} B_\tau \mid \sigma \in K\}, \quad |\sigma| \mapsto B_\sigma \cup \bigcup_{\tau \in \partial\sigma} B_\tau,$$

where the union is taken over all proper faces of σ . Note that $(\rho_S, \varphi^{-1}): (\text{Blowup}(|\mathcal{A}|), \mathcal{U}) \rightarrow (|K|, \mathcal{S})$ is a morphism of covered spaces.

Let us show that for every simplex $\sigma \in K$ the subspace $\varphi(|\sigma|)$ is contractible: For any face $\tau \subseteq \sigma$ we have $\Gamma(\sigma) \subseteq \Gamma(\tau)$ and hence $\varphi(|\sigma|)$ deformation retracts for any

chosen point $p_\sigma \in |\Gamma(\sigma)|$ onto $|\sigma| \times \{p_\sigma\}$ by a straight line homotopy. Therefore, $\varphi(|\sigma|)$ is contractible.

It follows from Lemma C.1 with $L = \emptyset$ that there exists a continuous map $g: |K| \rightarrow \text{Blowup}(|\mathcal{A}|)$ such that $(g, \varphi): (|K|, \mathcal{S}) \rightarrow (\text{Blowup}(|\mathcal{A}|), \mathcal{U})$ is a morphism of covered spaces. By composition, we get the morphisms of covered spaces $(\rho_S \circ g, \text{id}): (|K|, \mathcal{S}) \rightarrow (|\mathcal{A}|, \mathcal{S})$ and $(g \circ \rho_S, \text{id}): (\text{Blowup}(|\mathcal{A}|), \mathcal{U}) \rightarrow (\text{Blowup}(|\mathcal{A}|), \mathcal{U})$. It follows from Lemma C.1 and Remark C.2 that the map $\rho_S \circ g$ is homotopic to the identity on $|K|$; more explicitly, $\rho_S \circ g$ is homotopic to $\text{id}_{|K|}$ by a straight line homotopy. The same reasoning applied to the morphism of covered spaces

$$(g \circ \rho_S, \psi): (\text{Blowup}(|\mathcal{A}|), \text{cells}(\text{Blowup}(|\mathcal{A}|))) \rightarrow (\text{Blowup}(|\mathcal{A}|), \mathcal{U}), \quad \psi: (|\sigma| \times |\tau|) \mapsto \varphi(|\sigma|)$$

shows that $g \circ \rho_S$ is homotopic to the identity on $\text{Blowup}(|\mathcal{A}|)$. Therefore, ρ_S is a homotopy equivalence. \square

D. The Blowup Complex for Open Covers

The parts of 1(a) and 2(a) in Theorem 5.9 that establish weak homotopy equivalences follow from work of Dugger and Isaksen [DI04]. In this section, we adapt their proof strategy to give a more direct proof of the fact that the natural map $\rho_S: \text{Blowup}(\mathcal{A}) \rightarrow X$ is a weak homotopy equivalence whenever \mathcal{A} is an open cover of X . Moreover, we use the same ideas to give a short proof of the fact that $\rho_N: \text{Blowup}(\mathcal{A}) \rightarrow |\text{Nrv}(\mathcal{A})|$ is a weak homotopy equivalence whenever \mathcal{A} is a weakly good cover.

By the following lemma, a map is a weak homotopy equivalence if it is so locally.

Lemma D.1 ([Gra75, Lemma 16.24]; [Die08, Theorem 6.7.11]). *Let $f: Y \rightarrow X$ be a continuous map and let $\mathcal{A} = (A_i)_{i \in I}$ be an open cover of X . If for every $\sigma \in \text{Nrv}(\mathcal{A})$ the restricted map $f^{-1}(A_\sigma) \rightarrow A_\sigma$ is a weak homotopy equivalence, then so is f .*

In order to apply this lemma to ρ_S , we need to determine the preimages of the finite intersections of cover elements in $\mathcal{A} = (A_i)_{i \in I}$. Recall from Definition 2.7 that the blowup complex is defined as

$$\text{Blowup}(\mathcal{A}) = \left(\bigsqcup_{J \in \text{Nrv}(\mathcal{A})} A_J \times |J| \right) / \sim$$

and ρ_S is induced by the projections of the products $A_J \times |J|$ onto the first coordinate. For any $\sigma \in \text{Nrv}(\mathcal{A})$ the preimage is the subspace

$$\rho_S^{-1}(A_\sigma) = \left(\bigsqcup_{J \in \text{St}_{\text{Nrv}(\mathcal{A})}(\sigma)} A_J \times |J| \right) / \sim,$$

where $\text{St}_{\text{Nrv}(\mathcal{A})}(\sigma) = \{J \in \text{Nrv}(\mathcal{A}) \mid \sigma \subseteq J\}$ is the star of σ in $\text{Nrv}(\mathcal{A})$. This can be seen as follows: Whenever $A_J \cap A_\sigma = A_{J \cup \sigma}$ is non-empty for some J , the union $J \cup \sigma$ is a simplex in $\text{St}_{\text{Nrv}(\mathcal{A})}(\sigma)$ and so

$$(A_J \cap A_\sigma) \times |J| \subseteq A_{J \cup \sigma} \times |J \cup \sigma|$$

is contained in the right hand side of the equality above. Conversely, $A_J \subseteq A_\sigma$ for every $J \in \text{St}_{\text{Nrv}(\mathcal{A})}(\sigma)$, and so the above equality holds.

Proposition D.2. *Let $\mathcal{A} = (A_i)_{i \in I}$ be an open cover of the topological space X . Then the natural map $\rho_S: \text{Blowup}(\mathcal{A}) \rightarrow X$ is a weak homotopy equivalence.*

Proof. By Lemma D.1 it suffices to prove that for every $\sigma \in \text{Nrv}(\mathcal{A})$ the restricted map $\rho_S^{-1}(A_\sigma) \rightarrow A_\sigma$ is a weak homotopy equivalence. We show that this map is in fact a homotopy equivalence.

Choose any point $z \in |\sigma|$ and consider the following subspace

$$A_\sigma \times \{z\} \hookrightarrow A_\sigma \times |\sigma| \hookrightarrow \rho_S^{-1}(A_\sigma).$$

Note that the space $|\text{St}_{\text{Nrv}(\mathcal{A})}(\sigma)|$ is star-shaped with respect to z and hence it deformation retracts onto this point. As $A_\tau \subseteq A_\sigma$ for every $\tau \in \text{St}_{\text{Nrv}(\mathcal{A})}(\sigma)$, this lifts to a deformation retraction of $\rho_S^{-1}(A_\sigma)$ onto $A_\sigma \times \{z\}$. Therefore, we get the following commutative diagram, where the horizontal maps are homotopy equivalences:

$$\begin{array}{ccc} A_\sigma \times \{z\} & \xrightarrow{\simeq} & \rho_S^{-1}(A_\sigma) \\ \pi_1 \downarrow & & \downarrow \rho_S \\ A_\sigma & \xrightarrow{=} & A_\sigma \end{array}$$

Obviously, π_1 is a homotopy equivalence and hence so is the map on the right. This proves that $\rho_S: \text{Blowup}(\mathcal{A}) \rightarrow X$ is weak homotopy equivalence. \square

Recall now that ρ_N is induced by the projections of the products $A_J \times |J|$ onto the second coordinate. To apply Lemma D.1 to ρ_N , we cover $|\text{Nrv}(\mathcal{A})|$ by the open simplex stars $(S_\sigma)_{\sigma \in \text{Nrv}(\mathcal{A})}$, where

$$S_\sigma = \bigcup \{ \text{int } |J| \mid J \in \text{St}_{\text{Nrv}(\mathcal{A})}(\sigma) \}.$$

Note that this cover is closed under finite intersections. Hence, it suffices to consider for any $\sigma \in \text{Nrv}(\mathcal{A})$ the preimage

$$\rho_N^{-1}(S_\sigma) = \left(\bigsqcup_{J \in \text{St}_{\text{Nrv}(\mathcal{A})}(\sigma)} A_J \times \text{int } |J| \right) / \sim.$$

Proposition D.3. *Let $\mathcal{A} = (A_i)_{i \in I}$ be a weakly good cover of the topological space X . Then the natural map $\rho_N: \text{Blowup}(\mathcal{A}) \rightarrow |\text{Nrv}(\mathcal{A})|$ is a weak homotopy equivalence.*

Proof. By Lemma D.1 it suffices to prove that for every $\sigma \in \text{Nrv}(\mathcal{A})$ the restricted map $\rho_N^{-1}(S_\sigma) \rightarrow S_\sigma$ is a weak homotopy equivalence.

Similarly to the the proof of Proposition D.2, we get the following commutative diagram, where $z \in \text{int } |\sigma|$ is any point and the horizontal maps are homotopy equivalences:

$$\begin{array}{ccc} A_\sigma \times \{z\} & \xrightarrow{\cong} & \rho_N^{-1}(S_\sigma) \\ \pi_2 \downarrow & & \downarrow \rho_N \\ \{z\} & \xrightarrow{\cong} & S_\sigma \end{array}$$

As \mathcal{A} is a weakly good cover, the map π_2 is a weak homotopy equivalence and hence so is the map on the right. This proves that $\rho_N: \text{Blowup}(\mathcal{A}) \rightarrow |\text{Nrv}(\mathcal{A})|$ is a weak homotopy equivalence. \square

Acknowledgements This research has been supported by the DFG Collaborative Research Center SFB/TRR 109 *Discretization in Geometry and Dynamics*, and Austrian Science Fund (FWF) grants P 29984-N35 and P 33765-N.

References

- [Ale28] Paul Alexandroff. “Über den allgemeinen Dimensionsbegriff und seine Beziehungen zur elementaren geometrischen Anschauung”. In: *Math. Ann.* 98.1 (1928), pp. 617–635. ISSN: 0025-5831. DOI: [10.1007/BF01451612](https://doi.org/10.1007/BF01451612).
- [Bar11] Jonathan Ariel Barmak. “On Quillen’s Theorem A for posets”. In: *J. Combin. Theory Ser. A* 118.8 (2011), pp. 2445–2453. ISSN: 0097-3165. DOI: [10.1016/j.jcta.2011.06.008](https://doi.org/10.1016/j.jcta.2011.06.008).
- [Bar02] Michael Barr. *Acyclic models*. Vol. 17. CRM Monograph Series. American Mathematical Society, Providence, RI, 2002, pp. xii+179. ISBN: 0-8218-2877-0. DOI: [10.1090/crmm/017](https://doi.org/10.1090/crmm/017).
- [Bat02] Ekkehard Batzies. *Discrete Morse Theory for Cellular Resolutions*. Philipps-Universität Marburg. 2002. DOI: [10.17192/z2002.0115](https://doi.org/10.17192/z2002.0115).
- [Bau+20] U. Bauer, H. Edelsbrunner, G. Jabłoński, and M. Mrozek. “Čech–Delaunay gradient flow and homology inference for self-maps”. In: *Journal of Applied and Computational Topology* 4 (Dec. 2020), pp. 455–480. DOI: [10.1007/s41468-020-00058-8](https://doi.org/10.1007/s41468-020-00058-8).
- [Ben+07] Paul Bendich, David Cohen-Steiner, Herbert Edelsbrunner, John Harer, and Dmitriy Morozov. “Inferring Local Homology from Sampled Stratified Spaces”. In: *48th Annual IEEE Symposium on Foundations of Computer Science (FOCS 2007), October 20-23, 2007, Providence, RI, USA, Proceedings*. IEEE Computer Society, 2007, pp. 536–546. DOI: [10.1109/FOCS.2007.45](https://doi.org/10.1109/FOCS.2007.45).
- [Bjö03] Anders Björner. “Nerves, fibers and homotopy groups”. In: *J. Combin. Theory Ser. A* 102.1 (2003), pp. 88–93. ISSN: 0097-3165. DOI: [10.1016/S0097-3165\(03\)00015-3](https://doi.org/10.1016/S0097-3165(03)00015-3).

- [BL17] Andrew J. Blumberg and Michael Lesnick. *Universality of the Homotopy Interleaving Distance*. 2017. arXiv: [1705.01690](https://arxiv.org/abs/1705.01690) [[math.AT](#)].
- [BCR98] Jacek Bochnak, Michel Coste, and Marie-Françoise Roy. *Real algebraic geometry*. Vol. 36. *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)* [Results in Mathematics and Related Areas (3)]. Translated from the 1987 French original, Revised by the authors. Springer-Verlag, Berlin, 1998, pp. x+430. ISBN: 3-540-64663-9. DOI: [10.1007/978-3-662-03718-8](https://doi.org/10.1007/978-3-662-03718-8).
- [BS73] A. Borel and J.-P. Serre. “Corners and arithmetic groups”. In: *Comment. Math. Helv.* 48 (1973), pp. 436–491. ISSN: 0010-2571. DOI: [10.1007/BF02566134](https://doi.org/10.1007/BF02566134).
- [Bor48] Karol Borsuk. “On the imbedding of systems of compacta in simplicial complexes”. eng. In: *Fundamenta Mathematicae* 35.1 (1948), pp. 217–234. URL: <http://eudml.org/doc/213158>.
- [BS15] Magnus Bakke Botnan and Gard Spreemann. “Approximating persistent homology in Euclidean space through collapses”. In: *Applied Algebra in Engineering, Communication and Computing* 26.1-2 (2015), pp. 73–101. DOI: [10.1007/s00200-014-0247-y](https://doi.org/10.1007/s00200-014-0247-y).
- [BT82] Raoul Bott and Loring W. Tu. *Differential forms in algebraic topology*. Vol. 82. Graduate Texts in Mathematics. Springer-Verlag, New York-Berlin, 1982, pp. xiv+331. ISBN: 0-387-90613-4.
- [Bro94] Kenneth S. Brown. *Cohomology of groups*. Vol. 87. Graduate Texts in Mathematics. Corrected reprint of the 1982 original. Springer-Verlag, New York, 1994, pp. x+306. ISBN: 0-387-90688-6.
- [Bro92] Kenneth S. Brown. “The geometry of rewriting systems: a proof of the Anick-Groves-Squier theorem”. In: *Algorithms and classification in combinatorial group theory (Berkeley, CA, 1989)*. Vol. 23. Math. Sci. Res. Inst. Publ. Springer, New York, 1992, pp. 137–163. DOI: [10.1007/978-1-4613-9730-4_6](https://doi.org/10.1007/978-1-4613-9730-4_6).
- [Cha+09] Frédéric Chazal, David Cohen-Steiner, Leonidas J. Guibas, Facundo Mémoli, and Steve Oudot. “Gromov-Hausdorff Stable Signatures for Shapes using Persistence”. In: *Comput. Graph. Forum* 28.5 (2009), pp. 1393–1403. DOI: [10.1111/j.1467-8659.2009.01516.x](https://doi.org/10.1111/j.1467-8659.2009.01516.x).
- [CO08] Frédéric Chazal and Steve Oudot. “Towards Persistence-Based Reconstruction in Euclidean Spaces”. In: *Proceedings of the Annual Symposium on Computational Geometry* (Jan. 2008). DOI: [10.1145/1377676.1377719](https://doi.org/10.1145/1377676.1377719).
- [Die08] Tammo tom Dieck. *Algebraic topology*. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2008, pp. xii+567. ISBN: 978-3-03719-048-7. DOI: [10.4171/048](https://doi.org/10.4171/048).
- [Dug] Daniel Dugger. *A primer on homotopy colimits*. URL: <https://pages.uoregon.edu/ddugger/hocolim.pdf> (visited on 05/13/2020).

- [DI04] Daniel Dugger and Daniel Isaksen. “Topological hypercovers and \mathbb{A}^1 -realizations”. In: *Mathematische Zeitschrift* 246 (Apr. 2004), pp. 667–689. DOI: [10.1007/s00209-003-0607-y](https://doi.org/10.1007/s00209-003-0607-y).
- [DS95] W. Dwyer and J. Spalinski. “Homotopy theories and model categories”. In: *Handbook of Algebraic Topology*. Ed. by I.M. James. 1995, pp. 73–126.
- [EH10] H. Edelsbrunner and J. Harer. *Computational Topology. An Introduction*. American Mathematical Society, 2010. ISBN: 0-8218-4925-5.
- [EKS83] H. Edelsbrunner, D. Kirkpatrick, and R. Seidel. “On the shape of a set of points in the plane”. In: *IEEE Transactions on Information Theory* 29.4 (1983), pp. 551–559. DOI: [10.1109/TIT.1983.1056714](https://doi.org/10.1109/TIT.1983.1056714).
- [EH80] David A. Edwards and Harold M. Hastings. “Čech theory: its past, present, and future”. In: *Rocky Mountain J. Math.* 10.3 (1980), pp. 429–468. ISSN: 0035-7596. DOI: [10.1216/RMJ-1980-10-3-429](https://doi.org/10.1216/RMJ-1980-10-3-429).
- [ES52] Samuel Eilenberg and Norman Steenrod. *Foundations of algebraic topology*. Princeton University Press, Princeton, New Jersey, 1952, pp. xv+328.
- [FMN14] Steve Ferry, Konstantin Mischaikow, and Vidit Nanda. “Reconstructing functions from random samples”. In: *Journal of Computational Dynamics* 1.2 (2014), pp. 233–248. DOI: [10.3934/jcd.2014.1.233](https://doi.org/10.3934/jcd.2014.1.233).
- [For95] Robin Forman. “A discrete Morse theory for cell complexes”. In: *Geometry, topology, & physics*. Conf. Proc. Lecture Notes Geom. Topology, IV. Int. Press, Cambridge, MA, 1995, pp. 112–125.
- [For02] Robin Forman. “A user’s guide to discrete Morse theory”. In: *Sém. Lothar. Combin.* 48 (2002), Art. B48c, 35.
- [Fri12] Greg Friedman. “Survey article: An elementary illustrated introduction to simplicial sets”. In: *Rocky Mountain J. Math.* 42.2 (2012), pp. 353–423. ISSN: 0035-7596. DOI: [10.1216/RMJ-2012-42-2-353](https://doi.org/10.1216/RMJ-2012-42-2-353).
- [GZ67] P. Gabriel and M. Zisman. *Calculus of fractions and homotopy theory*. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 35. Springer-Verlag New York, Inc., New York, 1967, pp. x+168.
- [GM05] R. Ghrist and A. Muhammad. “Coverage and hole-detection in sensor networks via homology”. In: *IPSN 2005. Fourth International Symposium on Information Processing in Sensor Networks, 2005*. 2005, pp. 254–260. DOI: [10.1109/IPSN.2005.1440933](https://doi.org/10.1109/IPSN.2005.1440933).
- [GS07] Paul Goerss and Kristen Schemmerhorn. “Model categories and simplicial methods”. In: *Interactions between homotopy theory and algebra*. Vol. 436. Contemp. Math. Amer. Math. Soc., 2007, pp. 3–49.
- [GJ09] Paul G. Goerss and John F. Jardine. *Simplicial homotopy theory*. Modern Birkhäuser Classics. Reprint of the 1999 edition [MR1711612]. Birkhäuser Verlag, Basel, 2009, pp. xvi+510. ISBN: 978-3-0346-0188-7. DOI: [10.1007/978-3-0346-0189-4](https://doi.org/10.1007/978-3-0346-0189-4).

- [Gra75] Brayton Gray. *Homotopy theory*. Pure and Applied Mathematics, Vol. 64. An introduction to algebraic topology. Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1975, pp. xiii+368.
- [Hat02] Allen Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002, pp. xii+544. ISBN: 9780521795401.
- [Hau95] Jean-Claude Hausmann. “On the Vietoris-Rips complexes and a cohomology theory for metric spaces”. In: *Prospects in topology (Princeton, NJ, 1994)*. Vol. 138. Ann. of Math. Stud. Princeton Univ. Press, Princeton, NJ, 1995, pp. 175–188.
- [Hir03] Philip S. Hirschhorn. *Model categories and their localizations*. Vol. 99. Mathematical Surveys and Monographs. American Mathematical Society, 2003.
- [Hov99] Mark Hovey. *Model categories*. Vol. 63. Mathematical Surveys and Monographs. American Mathematical Society, 1999.
- [Jos83] K. D. Joshi. *Introduction to general topology*. A Halsted Press Book. John Wiley & Sons, Inc., New York, 1983, pp. xii+412. ISBN: 0-470-27556-1.
- [Koz08] Dmitry Kozlov. *Combinatorial algebraic topology*. Vol. 21. Algorithms and Computation in Mathematics. Springer, Berlin, 2008, pp. xx+389. ISBN: 978-3-540-71961-8. DOI: [10.1007/978-3-540-71962-5](https://doi.org/10.1007/978-3-540-71962-5).
- [Ler45] Jean Leray. “Sur la forme des espaces topologiques et sur les points fixes des représentations”. In: *J. Math. Pures Appl. (9)* 24 (1945), pp. 95–167. ISSN: 0021-7824.
- [Lil73] Joachim Lillig. “A union theorem for cofibrations”. In: *Arch. Math. (Basel)* 24 (1973), pp. 410–415. ISSN: 0003-889X. DOI: [10.1007/BF01228231](https://doi.org/10.1007/BF01228231).
- [LMO20] Sunhyuk Lim, Facundo Mémoli, and Osman Berat Okutan. *Vietoris–Rips Persistent Homology, Injective Metric Spaces, and The Filling Radius*. 2020. arXiv: [2001.07588](https://arxiv.org/abs/2001.07588) [math.AT].
- [May99] J.P. May. *A Concise Course in Algebraic Topology*. The University of Chicago Press, 1999.
- [MP12] J.P. May and K. Ponto. *More Concise Algebraic Topology: Localization, Completion, and Model Categories*. The University of Chicago Press, 2012.
- [McC67] Michael C. McCord. “Homotopy type comparison of a space with complexes associated with its open covers”. In: *Proc. Amer. Math. Soc.* 18 (1967), pp. 705–708. ISSN: 0002-9939. DOI: [10.2307/2035443](https://doi.org/10.2307/2035443).
- [Mes01] Roy Meshulam. “The clique complex and hypergraph matching”. In: *Combinatorica* 21.1 (2001), pp. 89–94. ISSN: 0209-9683. DOI: [10.1007/s004930170006](https://doi.org/10.1007/s004930170006).
- [Mun84] James R. Munkres. *Elements of algebraic topology*. Addison-Wesley Publishing Company, Menlo Park, CA, 1984, pp. ix+454. ISBN: 0-201-04586-9.
- [Nag07] Andrzej Nagórko. “Carrier and nerve theorems in the extension theory”. In: *Proc. Amer. Math. Soc.* 135.2 (2007), pp. 551–558. ISSN: 0002-9939. DOI: [10.1090/S0002-9939-06-08477-2](https://doi.org/10.1090/S0002-9939-06-08477-2).

- [Qui73] Daniel Quillen. “Higher algebraic K -theory. I”. In: *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*. 1973, 85–147. Lecture Notes in Math., Vol. 341.
- [Qui67] Daniel Quillen. *Homotopical Algebra*. Vol. 43. Lecture Notes in Mathematics. Springer, 1967.
- [Qui78] Daniel Quillen. “Homotopy properties of the poset of nontrivial p -subgroups of a group”. In: *Adv. in Math.* 28.2 (1978), pp. 101–128. ISSN: 0001-8708. DOI: [10.1016/0001-8708\(78\)90058-0](https://doi.org/10.1016/0001-8708(78)90058-0).
- [Rie14] Emily Riehl. *Categorical homotopy theory*. Vol. 24. New Mathematical Monographs. Cambridge University Press, Cambridge, 2014, pp. xviii+352. ISBN: 978-1-107-04845-4. DOI: [10.1017/CB09781107261457](https://doi.org/10.1017/CB09781107261457).
- [Seg68] Graeme Segal. “Classifying spaces and spectral sequences”. In: *Inst. Hautes Études Sci. Publ. Math.* 34 (1968), pp. 105–112. ISSN: 0073-8301.
- [SS95] Lynn Arthur Steen and J. Arthur Seebach Jr. *Counterexamples in topology*. Reprint of the second (1978) edition. Dover Publications, Inc., Mineola, NY, 1995, pp. xii+244. ISBN: 0-486-68735-X.
- [Str09] N. P. Strickland. *The Category of CGWH Spaces*. 2009. URL: <https://ncatlab.org/nlab/files/StricklandCGHWSpaces.pdf>.
- [Str72] Arne Strøm. “The homotopy category is a homotopy category”. In: *Archiv der Mathematik* 23 (1972), pp. 435–441.
- [Vir21] Žiga Virk. “Rips Complexes as Nerves and a Functorial Dowker-Nerve Diagram”. In: *Mediterranean Journal of Mathematics* 18 (2021). DOI: [10.1007/s00009-021-01699-4](https://doi.org/10.1007/s00009-021-01699-4).
- [Wal81] James W. Walker. “Homotopy type and Euler characteristic of partially ordered sets”. In: *European J. Combin.* 2.4 (1981), pp. 373–384. ISSN: 0195-6698. DOI: [10.1016/S0195-6698\(81\)80045-5](https://doi.org/10.1016/S0195-6698(81)80045-5).
- [Wei94] Charles A. Weibel. *An introduction to homological algebra*. Vol. 38. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1994, pp. xiv+450. ISBN: 0-521-43500-5; 0-521-55987-1. DOI: [10.1017/CB09781139644136](https://doi.org/10.1017/CB09781139644136).
- [Wei52] André Weil. “Sur les théorèmes de de Rham”. In: *Comment. Math. Helv.* 26 (1952), pp. 119–145. ISSN: 0010-2571. DOI: [10.1007/BF02564296](https://doi.org/10.1007/BF02564296).
- [WZŽ99] Volkmar Welker, Günter M. Ziegler, and Rade T. Živaljević. “Homotopy colimits – comparison lemmas for combinatorial applications”. In: *Journal für die reine und angewandte Mathematik* 509 (1999), pp. 117–149. DOI: [10.1515/crll.1999.509.117](https://doi.org/10.1515/crll.1999.509.117).
- [ZC08] Afra Zomorodian and Gunnar Carlsson. “Localized homology”. In: *Comput. Geom.* 41.3 (2008), pp. 126–148. ISSN: 0925-7721. DOI: [10.1016/j.comgeo.2008.02.003](https://doi.org/10.1016/j.comgeo.2008.02.003).