ON THE LS-CATEGORY OF HOMOMORPHISMS

ALEXANDER DRANISHNIKOV AND NURSULTAN KUANYSHOV

ABSTRACT. We prove the equality $\operatorname{cat}(\phi) = \operatorname{cd}(\phi)$ for homomorphisms $\phi : \Gamma \to \Lambda$ of a torsion free finitely generated nilpotent groups Γ to an arbitrary group Λ . We construct an epimorphism $\psi : G \to H$ between geometrically finite groups with $\operatorname{cat}(\psi) > \operatorname{cd}(\psi)$.

1. INTRODUCTION

Definition 1.1. The (reduced) Lusternik-Schnirelmann category (LS-category), cat X, of an ANR space X is the minimal number k such that X admits an open cover by k + 1 sets U_0, U_1, \ldots, U_k such that each U_i is contractible in X.

The Lusternik-Schnirelmann category is an important invariant, since it gives a lower bound on the number of critical points for a smooth real-valued function on a closed manifold [LS]. Since it is a homotopy invariant, it can be defined for discrete groups Γ as cat Γ = cat $B\Gamma$ where $B\Gamma = K(\Gamma, 1)$ is a classifying space. Eilenberg and Ganea [EG] proved that the LS-category of a discrete group equals its cohomological dimension, cat $(\pi) = cd(\pi)$. We recall that the cohomological dimension of a group Γ is defined as follows,

 $cd(\Gamma) = \max\{k \mid H^k(\Gamma, M) \neq 0\}$

where the maximum is taken over all $\mathbb{Z}\Gamma$ -modules M.

Theorem 1.2 ([Sch, DR]). For the cohomological dimension of a discrete group Γ ,

$$cd(\Gamma) = \max\{k \mid (\beta_{\Gamma})^k \neq 0\}$$

where $\beta_{\Gamma} \in H^1(\Gamma, I(\Gamma))$ is the Berstein-Schwarz class of Γ .

The LS-category of the map $f: X \to Y$, cat f, is the minimal number k such that X admits an open cover by k + 1 open sets $U_0, U_2, ..., U_k$ with nullhomotopic restrictions $f|_{U_i}: U_i \to Y$ for all i. The LS-category cat ϕ of a group homomorphism $\phi: \Gamma \to \pi$ is defined as cat f where the map $f: B\Gamma \to B\pi$ induces the homomorphism ϕ for the fundamental groups.

The cohomological dimension $cd(\phi)$ of a group homomorphism $\phi: \Gamma \to \pi$ was defined by Mark Grant about 10 years ago on Mathoverflow [Gr] as maximum of k such that there is a π -module M with the nonzero induced homomorphism $\phi^*: H^k(\pi, M) \to H^k(\Gamma, M)$. In view of universality of the Berstein-Schwarz class [DR] for any homomorphism $\phi: \Gamma \to \pi$

$$\operatorname{cd}(\phi) = \max\{k \mid \phi^*(\beta_\pi)^k \neq 0\}.$$

This brings the inequality $cd(\phi) \leq cat \phi$ for all homomorphisms.

In view of the equity $cd(\Gamma) = cat \Gamma$, the following conjecture seems to be natural:

²⁰⁰⁰ Mathematics Subject Classification. Primary 55M30; Secondary 55M25, 57R65, 57R67.

Key words and phrases. Lusternik-Schnirelmann category, group homomorphism.

Conjecture 1.3. For any group homomorphism $\phi: \Gamma \to \pi$ always

$$\operatorname{cat} \phi = \operatorname{cd}(\phi).$$

In [Sc] Jamie Scott considered this conjecture for geometrically finite groups and he proved it for monomorphisms of any groups and for homomorphisms of free and free abelian groups. In [Gr] Tom Goodwillie gave an example of an epimorphism of an infinitely generated group $\phi: G \to \mathbb{Z}^2$ with $cd(\phi) = 1$ that disproves the conjecture.

In the first part of this paper we prove Conjecture 1.3 for finitely generated torsion free nilpotent groups. In the second part we present a finitely generated counterexample by constructing a map between aspherical manifolds $f: M \to N$ with

$$\operatorname{cat} f > \operatorname{cd}(f_{\#} : \pi_1(M) \to \pi_1(N)).$$

2. Preliminaries

2.1. Nilpotent groups. The upper central series of a group Γ is a chain of subgroups

$$e = Z_0 \leqslant Z_1 \leqslant \dots \leqslant Z_n \leqslant \dots$$

where $Z_1 = Z(\Gamma)$ is the center of the group, and Z_{i+1} is the preimage under the canonical epimorphism $\Gamma \to \Gamma/Z_i$ of the center of Γ/Z_i . A group Γ is *nilpotent* if $Z_n = \Gamma$ for some n. The least such n is called *the nilpotency class* of Γ , denoted nil(Γ). Note that the groups with the nilpotency class one are exactly abelian groups.

The lower central series of a group Γ is a chain of subgroups

$$\Gamma = \gamma_0(\Gamma) \ge \gamma_1(\Gamma) \ge \gamma_2(\Gamma) \ge \dots$$

defined as $\gamma_{i+1}(\Gamma) = [\gamma_i(\Gamma), \Gamma]$. It's known that for nilpotent groups Γ the nilpotency class nil(Γ) equals the least *n* for which $\gamma_n(\Gamma) = 1$.

Proposition 2.1. (1) Let $\phi : \Gamma \to \Gamma'$ be an epimorphism. Then $\phi(Z(\Gamma)) \subset Z(\Gamma')$ and $\phi(\gamma_i(\Gamma)) = \gamma_i(\Gamma')$ for all *i*.

(2) For any finitely generated torsion free nilpotent group Γ , any $z \in \Gamma$, and any $n \in \mathbb{N}$ the condition $z^n \in Z(\Gamma)$ implies $z \in Z(\Gamma)$.

Proof. (1) Straightforward (see for example [B], Theorem 5.1.3).

(2) This is Mal'cev's Theorem 1 in [Ma2]. We note that the proof there is not selfcontained. This statement follows from the fact that z and z^n have the same centralizers. The latter can be proven using Mal'cev theorem about embedding Γ into the group of unipotent upper triangular matrices [Ra] and the fact that z belongs to Zariski closure of z^n .

Corollary 2.2. For any finitely generated torsion free nilpotent group Γ the group $\Gamma/Z(\Gamma)$ is torsion free finitely generated nilpotent group.

Proof. The torsion free part follows from (2). The rest follows from (1). \Box

We note that $\operatorname{nil}(\Gamma/Z(\Gamma)) < \operatorname{nil}(\Gamma)$.

Suppose that G is a connected, simply connected nilpotent Lie group and $\Gamma \subset G$ is a uniform lattice. Then G is the universal cover for Γ and $N = G/\Gamma$ is an aspherical manifold, called a *nilmanifold*.

Mal'cev Theorem. [Ma1] Every torsion free finitely generated nilpotent group Γ can be realized as the fundamental group of some nilmanifold.

The corresponding simply connected Lie groups G is obtained as the Mal'cev completion of Γ .

2.2. Ganea-Schwarz's approach to cat. Recall that an element of an iterated join $X_0 * X_1 * \cdots * X_n$ of topological spaces is a formal linear combination $t_0x_0 + \cdots + t_nx_n$ of points $x_i \in X_i$ with $\sum t_i = 1, t_i \ge 0$, in which all terms of the form $0x_i$ are dropped. Given fibrations $f_i : X_i \to Y$ for i = 0, ..., n, the fiberwise join of spaces $X_0, ..., X_n$ is defined to be the space

 $X_0 *_Y X_1 *_Y \cdots *_Y X_n = \{ t_0 x_0 + \cdots + t_n x_n \in X_0 * \cdots * X_n \mid f_0(x_0) = \cdots = f_n(x_n) \}.$

The fiberwise join of fibrations $f_0, ..., f_n$ is the fibration

 $f_0 *_Y * \cdots *_Y f_n : X_0 *_Y X_1 *_Y \cdots *_Y X_n \longrightarrow Y$

defined by taking a point $t_0x_0 + \cdots + t_nx_n$ to $f_i(x_i)$ for any *i* such that $t_i \neq 0$.

When $X_i = X$ and $f_i = f : X \to Y$ for all *i* the fiberwise join of spaces is denoted by $*_Y^{n+1}X$ and the fiberwise join of fibrations is denoted by $*_Y^{n+1}f$.

For a path connected space X, we turn an inclusion of a point $* \to X$ into a fibration $p_0^X : G_0(X) \to X$, whose fiber is known to be the loop space ΩX . The *n*-th Ganea space of X is defined to be the space $G_n(X) = *_X^{n+1} G_0(X)$, while the *n*-th Ganea fibration $p_n^X : G_n(X) \to X$ is the fiberwise join $*_X^{n+1} p_0^X$. Then the fiber of p_n^X is $*^{n+1} \Omega X$.

The following theorem give the Ganea-Shwarz characterization of cat [Sch],[CLOT]:

Theorem 2.3. If X is a connected ANR, then $\operatorname{cat} X \leq n$ if and only if the fibration $p_n^X : G_n(X) \to X$ admits a section.

This characterization can be extendedd to maps:

Theorem 2.4. If $f : X \to Y$ is a map between connected ANRs, then $\operatorname{cat} f \leq n$ if and only if there is a lift of f with respect to the fibration $p_n^Y : G_n(Y) \to Y$ admits a section.

2.3. Berstein-Schwarz cohomology class. The Berstein-Schwarz class of a discrete group π is the first obstruction β_{π} to a lift of $B\pi = K(\pi, 1)$ to the universal covering $E\pi$. Note that $\beta_{\pi} \in H^1(\pi, I(\pi))$ where $I(\pi)$ is the augmentation ideal of the group ring $\mathbb{Z}\pi$ [Be],[Sch].

Theorem 2.5 (Universality [DR],[Sch]). For any cohomology class $\alpha \in H^k(\pi, L)$ there is a homomorphism of π -modules $I(\pi)^k \to L$ such that the induced homomorphism for cohomology takes $(\beta_{\pi})^k \in H^k(\pi; I(\pi)^k)$ to α where $I(\pi)^k = I(\pi) \otimes \cdots \otimes I(\pi)$ and $(\beta_{\pi})^k = \beta_{\pi} \smile \cdots \smile \beta_{\pi}$.

In the paper we use notations $H^*(\Gamma, A)$ for cohomology of a group Γ with coefficient in Γ -module A. The cohomology groups of a space X with the fundamental group Γ we denote as $H^*(X; A)$. Thus, $H^*(\Gamma, A) = H^*(B\Gamma; A)$ where $B\Gamma = K(\Gamma, 1)$.

3. Reduction to epimorphisms

Lemma 3.1. Let $\pi \subset \Lambda$ be a subgroup, $j : B\pi \to B\Lambda$ be a map generated by this inclusion, and $p' : j^* E\Lambda \to B\pi$ be the pull-back of the universal covering $p_\Lambda : E\Lambda \to B\Lambda$. Then p' has a lift with respect to the universal covering $p_\pi : E\pi \to B\pi$. Proof. Clearly, for each path component C of $j^*E\Lambda$ the restriction $p'|_C : C \to B\pi$ of a covering p' is a covering. Easy diagram chasing shows that C is simply connected. Hence, each path component C of $j^*E\Lambda$ is homeomorphic to $E\pi$ and the restriction $p'|_C : C \to B\pi$ is the universal covering. We may assume that $j^*E\Lambda$ is a CW complex. This would imply that each path component C is open. Thus, the lift $j^*E\Lambda \to E\pi$ can be defined independently on each path component.

Given a homomorphism $\phi : \Gamma \to \Lambda$, by $\phi' : \Gamma \to im(\phi)$ we denote the restriction of ϕ from the codomain to its range. The following theorem is a generalization of Theorem 6.11 from [Sc].

Theorem 3.2. For any group homomorphism $\phi : \Gamma \to \Lambda$, $\operatorname{cat} \phi = \operatorname{cat} \phi'$.

Proof. Clearly, $\cot \phi \leq \cot \phi'$. We show that $\cot \phi \geq \cot \phi'$. Let $\cot \phi = k$. This means that there is a lift of the corresponding map $f: B\Gamma \to B\Lambda$ with respect to the Ganea fibration $p_k: G_k(B\Lambda) \to B\Lambda$. Since the path fibration $p: P_0(B\pi) \to B\pi$ is fiber-wise homotopy equivalent to the universal covering $E\pi \to B\pi$ for any discrete group π , the k-th Ganea fibration $G_k(B\pi) \to B\pi$ is fiberwise homotopy equivalent to the iterated fiberwise join $*_{B\pi}^{k+1}E\pi$ of the universal covering. Then f admits a lift with respect to

$$*^{k+1}_{B\Lambda}p_{\Lambda}:*^{k+1}_{B\Lambda}E\Lambda \to B\Lambda.$$

The map f factors as $f = j \circ f'$ where $f' : B\Gamma \to B\pi$, $j : B\pi \to B\Lambda$ and $\pi = im(\phi)$. Hence, the map f' admits a lift to the pull-back

$$\bar{f}': B\Gamma \to j^*(*^{k+1}_{B\Lambda}E\Lambda)$$

By Lemma 3.1 there is a lift s of $*^{k+1}_{B\pi}p': j^*(*^{k+1}_{B\Lambda}E\Lambda) \to B\pi$ with respect to

$$*^{k+1}_{B\pi}p_{\pi}:*^{k+1}_{B\pi}E\pi \to B\pi.$$

Then the composition $s \circ \bar{f}'$ is a lift of f' with respect to $*_{B\pi}^{k+1}p_{\pi}$. By Theorem 2.4 cat $f' \leq k$ and, hence, cat $\phi' \leq k$.

Theorem 3.3. For any group homomorphism $\phi: \Gamma \to \Lambda$ and $\phi': \Gamma \to im(\phi) = \pi$ as above

$$\operatorname{cd}(\phi) = \operatorname{cd}(\phi').$$

Proof. Clearly, $\operatorname{cd}(\phi') \ge \operatorname{cd}(\phi)$. We show that $\operatorname{cd}(\phi') \le \operatorname{cd}(\phi)$. Let $\operatorname{cd}(\phi') = k$. Then ϕ'^* : $H^k(\pi, M) \to H^k(\Gamma, M)$ is not zero for some π -module M. Let

$$\alpha: \operatorname{Coind}_{\pi}^{\Lambda} M = Hom_{\pi}(\mathbb{Z}\Lambda, M) \to M$$

denote the canonical π -homomorphism, defined for $f : \Lambda \to M$ as $\alpha(f) = f(1)$. Consider the commutative diagram

where α^* is the coefficient homomorphism generated by α . By Shapiro Lemma [Br] the top row through homomorphism is an isomorphism. Therefore, the homomorphism

$$\phi^* = \phi'^* j^* : H^k(\Lambda, \operatorname{Coind}_{\pi}^{\Lambda} M) \to H^k(\Gamma, \operatorname{Coind}_{\pi}^{\Lambda} M)$$

is nonzero. Hence, $cd(\phi) \ge k$.

Theorem 3.2 and Theorem 3.3 imply the following:

Corollary 3.4. Suppose that Conjecture 1.3 holds true for all epimorphisms $\phi : \Gamma \to \pi$ for some class of groups. Then it holds for all homomorphisms for groups from that class.

4. Homomorphisms of nilpotent groups

Lemma 4.1. Let Γ be π are finitely generated, torsion free nilpotent groups. Then every epimorphism $\phi: \Gamma \to \pi$ can be realized as a locally trivial bundle of nilmanifolds with the fiber a nilmanifold.

Proof. We prove it by induction on n = s + t where $s = \operatorname{nil}(\Gamma)$ and $t = \operatorname{nil}(\pi)$ are the nilpotency classes of Γ and π . The base of induction is the case of abelian groups where any epimorphism $\phi : \mathbb{Z}^{k+\ell} \to \mathbb{Z}^k$ is the projection onto a factor, since it is a split surjection. Clearly, the projection ϕ can be realized as a trivial fiber bundle of tori $T^{k+\ell} \to T^k$ with the fiber a torus T^{ℓ} .

Suppose that n > 2 and the statement of the lemma holds true for s + t < n. We denote by $B = Z(\Gamma) \cap \operatorname{Ker} \phi$ and by $A = Z(\Gamma)/B$, where $Z(\Gamma)$ is the center of Γ . Note that B is a direct summand in $Z(\Gamma)$ or, equivalently, A is free. For the later we claim that if $z^n \in B$, then $z \in B$. Indeed, If $z^n \in \operatorname{Ker} \phi$ then $z \in \operatorname{Ker} \phi$, since π is torsion free. If $z^n \in Z(\Gamma)$ then $z \in Z(\Gamma)$ by Proposition 2.1. Thus, $Z(\Gamma) \cong B \oplus A$. Let \overline{A} be the direct summand of $Z(\pi)$ that contains $\phi(Z(\Gamma)) \cong A$ as a finite index subgroup. In view of Corollary 2.2, π/\overline{A} is a torsion free nilpotent group.

We consider the nilpotent group $\Gamma' = \Gamma/B$ and the epimorphism $\phi' : \Gamma' \to \pi$. induced by ϕ . In view of the principal fiber bundle $B\Gamma \to B\Gamma'$ with the fiber a torus, it suffices to show that ϕ' can be realized as a fiber bundle of nilmanifolds. We consider the commutative diagram

$$\begin{array}{ccc} \Gamma' & \longrightarrow & \Gamma'/A \\ \phi' \downarrow & & \bar{\phi} \downarrow \\ \pi & \longrightarrow & \pi/\bar{A}. \end{array}$$

Note that $\Gamma'/A = \Gamma/Z(\Gamma)$ and, hence, $\bar{\phi}$ is an epimorphism with $\operatorname{nil}(\Gamma'/A) + \operatorname{nil}(\pi/\bar{A}) < s + t$. The homomorphism ϕ' factors through the pull-back $\Lambda = \bar{\phi}^* \pi$, $\phi' = \bar{\phi}' \circ \xi$ with respect to $\bar{\phi}$. Since $\phi'|_A : A \to \bar{A}$ is an embedding of a finite index subgroup, the homomorphism $\xi : \Gamma' \to \Lambda$ in the commutative diagram of short exact sequences generated by ϕ' and the pull-back

$$1 \longrightarrow A \longrightarrow \Gamma' \longrightarrow \Gamma/Z(\Gamma) \longrightarrow 1$$
$$\downarrow^{\subset} \qquad \xi \downarrow^{\subset} \qquad \downarrow^{=}$$
$$1 \longrightarrow \bar{A} \longrightarrow \Lambda \longrightarrow \Gamma/Z(\Gamma) \longrightarrow 1$$
$$\downarrow^{=} \qquad \bar{\phi}' \downarrow \qquad \bar{\phi} \downarrow$$
$$1 \longrightarrow \bar{A} \longrightarrow \pi \longrightarrow \pi/\bar{A} \longrightarrow 1$$

is an embedding of a finite index subgroup.

By induction assumption applied to $\overline{\phi}$ there is a fiber bundle between nilmanifolds

$$\bar{f}: B(\Gamma/Z(\Gamma)) \to B(\pi/\bar{A})$$

with the fiber a nilmanifold F. Consider the pull-back diagram

$$B\Lambda \xrightarrow{p'} B(\Gamma/Z(\Gamma))$$
$$\bar{f}' \downarrow \qquad \bar{f} \downarrow$$
$$B\pi \xrightarrow{p} B(\pi/\bar{A}).$$

Let $q : M \to B\Lambda$ be a covering that corresponds to the subgroup $\xi(\Gamma') \subset \Lambda = \pi_1(B\Lambda)$. Since $\xi(\Gamma')$ is of finite index, M is a closed aspherical manifold with $\pi_1(M) = \Gamma'$. Thus, the composition $f' = \bar{f}' \circ q : M \to B\pi$ realizes the homomorphism ϕ' as a fiber bundle. The fiber F' of f' is homeomorphic to the total space of a fiber bundle $F' \to F$ with a finite fiber. The homotopy exact sequence of the fibration $F' \to B\Gamma' \xrightarrow{f'} B\pi$ and the fact that ϕ' is surjective imply that F' is connected. Hence, F' is a nilmanifold.

Lemma 4.2. For every locally trivial bundle of closed aspherical manifolds $f : M^m \to N^n$ with compact connected fiber F the induced homomorphism

$$f^*: H^n(N; \mathbb{Z}\pi) \to H^n(M; \mathbb{Z}\pi)$$

is nonzero where $\pi = \pi_1(N)$.

Proof. Consider the pull-back diagram:

$$\widetilde{M} \xrightarrow{p_2} f^* \widetilde{N} \xrightarrow{f} \widetilde{N} \xrightarrow{p_M} p_1 \qquad \qquad \downarrow^{p_N} \\ M \xrightarrow{p_M} M \xrightarrow{f} N$$

where p_M and p_N are he universal covering maps. Since $f^*(\widetilde{N})$ is a covering of M and \widetilde{M} is the universal covering, p_M factors through covering maps p_2 and p_1 .

We recall [Br, Lemma 7.4] that for every left π -module M there is a natural isomorphism of right modules

$$\Theta: Hom_{\pi}(M, \mathbb{Z}\pi) \to Hom_{c}(M, \mathbb{Z})$$

defined by the formula $F \mapsto f_1$ where $F: M \to \mathbb{Z}\pi$, $F(m) = \sum_{\gamma \in \pi} f_{\gamma}(m)\gamma$. We consider the following commutative diagrams of cochain groups:

is defined as follows. For each simplex σ in X we fix a lift $\tilde{\sigma}$ in \widetilde{M} . We define

$$\Phi(F)(\sigma) = F(\widetilde{\sigma})$$

for $F \in Hom_{\Gamma}(C_*(\widetilde{M}), \mathbb{Z}\pi)$ and simplex σ in X.

Claim 1. $\Phi(F) \in Hom_{\pi}(C_*(X), \mathbb{Z}\pi).$

Proof. Since the fiber F is connected, it follows that f induces an epimorphism of the fundamental groups. Let $g \in \pi$ and let $f_*(\bar{g}) = g$. Then $\tilde{g}\tilde{\sigma}$ and $\bar{g}\tilde{\sigma}$ both cover $g\sigma$. Hence there is $\gamma \in Kerf_*$ such that $\gamma(\tilde{g}\tilde{\sigma}) = \bar{g}\tilde{\sigma}$. Then Φ is π -equivariant:

$$\Phi(F)(g\sigma) = F(\widetilde{g}\widetilde{\sigma}) = F(\gamma\widetilde{g}\widetilde{\sigma}) = F(\bar{g}\widetilde{\sigma}) = gF(\widetilde{\sigma}) = g\Phi(F)(\sigma).$$

Claim 2. $\Phi \circ p_2^* = 1$.

Proof.
$$\Phi \circ p_2^*(F')(\sigma) = \Phi(p_2^*(F'))(\sigma) = (p_2^*F')(\tilde{\sigma}) = F'(p_2(\tilde{\sigma})) = F'(\sigma)$$

Since \tilde{N} is contractible, the bottom row in the diagram above gives us an isomorphism of the cohomology groups $\Theta^* : H^*(N;\mathbb{Z}\pi) \to H^*_c(\tilde{N};\mathbb{Z})$. The commutative diagram on the cochain level produces the commutative diagram for cohomology:

$$H^{n}(M; \mathbb{Z}\pi) \xrightarrow{\Psi^{*}} H^{n}_{c}(X; \mathbb{Z})$$

$$f^{*}_{c} \uparrow \qquad (\bar{f})^{*} \uparrow$$

$$H^{n}(N; \mathbb{Z}\pi) \xrightarrow{\Theta^{*}} H^{n}_{c}(\tilde{N}; \mathbb{Z}) = \mathbb{Z}$$

Since \widetilde{N} is contractible, $X = f^*(\widetilde{N})$ is a trivial fiber bundle with the fiber F. Since $X \cong \widetilde{N} \times F$ admits a proper retraction onto \widetilde{N} , we obtain that $(\overline{f}_c)^*$ is a monomorphism. Hence, the homomorphism $f^* : \mathbb{Z} = H^n(N; \mathbb{Z}\pi) \to H^n(M; \mathbb{Z}\pi)$ is not trivial. \Box

Theorem 4.3. For a homomorphism $\phi : \Gamma \to \pi$ of finitely generated torsion free nilpotent groups $\operatorname{cat} \phi = \operatorname{cd}(\phi)$.

Proof. In view of Theorem 3.2 and Theorem 3.3, it suffices to prove the theorem when ϕ is surjective. By Lemma 4.1 there is a fiber bundle of nilmanifolds $M^m = B\Gamma \rightarrow B\pi = N^n$ with a compact fiber. By Lemma 4.2, $\operatorname{cd}(\phi) = n$. The inequalities $\operatorname{cd}(\phi) \leq \operatorname{cat} \phi \leq \dim N = n$ complete the proof.

5. Counterexample

Answering a question of M. Grant, T. Goodwillie gave an example of an epimorphism $\phi: G \to \Gamma$ satisfying the inequality $\operatorname{cd}(\phi) < \min\{\operatorname{cd}(G), \operatorname{cd}(\Gamma)\}$. We present here a slightly modified version of his example. Let $\Gamma = \mathbb{Z}^2$ and $\phi: G \to \Gamma$ be the epimorphism defined by the extension of Γ by the abelian group $I(\Gamma)^2 = I(\Gamma) \otimes_{\mathbb{Z}} I(\Gamma)$ that corresponds to the square of the Berstein-Schwarz cohomology class $(\beta_{\Gamma})^2 \in H^2(\Gamma, I(\Gamma)^2)$. Then $\phi^*(\beta_{\Gamma})^2 = 0$. Hence, $\operatorname{cd}(\phi) < 2$. If $\operatorname{cat} \phi \leq 1$, then the induced map $\hat{\phi}: BG \to B\Gamma$ can be lifted to the Ganea's space $G_1(B\Gamma)$ which is homotopy equivalent to 1-dimensional complex $\Sigma\Gamma$, the suspension of Γ . [CLOT]. Then the epimorphism ϕ factors through a free group F via epimorphisms $G \to F \to \Gamma$. Hence $\operatorname{rank} F \geq 2$. Therefore, G contains a free group of $\operatorname{rank} \geq 2$. We note that G is amenable as an abelian-by-abelian extension and hence G cannot contain F. Therefore, $\operatorname{cat} \phi \geq 2$. Thus, ϕ is an infinitely generated counterexample to Conjecture 1.3.

Remark 5.1. We note there are simpler examples of epimorphisms $\phi: G \to \Gamma$ satisfying

$$\operatorname{cd}(\phi) < \min\{\operatorname{cd}(G), \operatorname{cd}(\Gamma)\}.$$

Namely, the homomorphism $\phi : F_n * \mathbb{Z}^n \to \mathbb{Z}^n$ defined by the abelianization of the first factor and by the projection onto $\mathbb{Z}^n \to \mathbb{Z} \subset \mathbb{Z}^n$ has $\operatorname{cd}(\phi) = 1$ and $\operatorname{cd}(F^n * \mathbb{Z}^n) = \operatorname{cd} \mathbb{Z}^n = n$. Here F_n denotes the free group on n generators.

Let S denote the sphere spectrum. We recall that every stably parallelizable manifold is S-orientable [Sw]. The S-cohomology groups of X are exactly the stable cohomotopy groups $\pi_s^*(X)$. Then Lemma 3.5 of [Ru] in the case of the spectrum S can be stated as follows:

Lemma 5.2. Suppose that $f: W \to M$ is a map of degree one between closed stably paralleliable manifolds. Then the induced map $f^*: \pi_s^*(M) \to \pi_s^*(W)$ is injective.

We note that the natural map $[X, S^n] \to \pi_s^n(X)$ is a bijection when dim $X \leq 2n-2$ [Hu].

5.1. Bolotov's example. Answering a question of Gromov [Gro1], D. Bolotov constructed [Bo] a closed 4-manifold M with the fundamental group $\pi = \mathbb{Z} * \mathbb{Z}^3$ whose cohomological dimension $cd(\pi) = 3$ such that a classifying map $u_M : M \to B\pi$ cannot be deformed to the 2skeleton $B\pi^{(2)}$. His manifold is defined as the pull-back $M = g^*(S^3 \times S^1)$ of the S^1 -bundle $h \times 1 : S^3 \times S^1 \to S^2 \times S^1$, where h is the Hopf fiber bundle, with respect to the collapsing map $g : N = (S^2 \times S^1) \# T^3 \to S^2 \times S^1$. Here T^3 is a 3-dimensional torus. The pull-back bundle is denoted by $p : M \to N$.

Proposition 5.3. Bolotov's manifold has the following properties:

(a) The map p induces an isomorphism of the fundamental groups.

(b) The homomorphism $u_M^*: H^3(B\pi; A) \to H^3(M; A)$ is trivial for any π -module A.

(c) The through map $M \xrightarrow{u_M} B\pi = S^1 \vee T^3 \xrightarrow{q_2} T^3 \xrightarrow{q_1} S^3$ is essential where q_2 collapses S^1 to the wedge point, and q_1 is a map of degree one.

(d) The manifold M is stably parallelizable.

Proof. (a) This is straightforward ([Bo]). Thus, the classifying map u_M factors through $p: M \to N, u_M = q_3 q_4 p$ where $q_4: N \to (S^2 \times S^1) \vee T^3$ collapses the connected sum to the wedge and $q_3: (S^2 \times S^1) \vee T^3 \to S^1 \vee T^3$ projects $S^2 \times S^1$ onto the factor S^1 .

(b) Let $a \in H^3(B\pi; A)$. By the Poincare Duality for local coefficients there is 1-dimensional class $\bar{\beta}$ in M such that $u_M^*(a) \cup \bar{\beta} \neq 0$. In view of universality of the Berstein-Schwarz class β_{π} and the fact that u_M induces an isomorphism of 1-cohomology, we may assume that $\bar{\beta} = u_M^*(\beta_{\pi})$. Then $0 \neq u_M^*(a) \cup \bar{\beta} = u_M^*(a \cup \beta_{\pi}) = 0$. The last equality is due to dimensional reason.

(c) This was proven in [Bo]. Here we present a simplified proof. Let $q = q_1 q_2 q_3 q_4 : N \to S^3$:

$$N = (S^2 \times S^1) \# T^3 \xrightarrow{q_4} (S^2 \times S^1) \vee T^3 \xrightarrow{q_3} S^1 \vee T^3 \xrightarrow{q_2} T^3 \xrightarrow{q_1} S^3.$$

Let a be a generator in $H^3(K(\mathbb{Z},3))$ and let a_0 be its restriction to $S^3 = K(\mathbb{Z},3)^{(4)}$. Part (b) implies that $p^*(\alpha_0) = 0$ where $\alpha_0 = q^*(a_0)$. The exact sequence of pair (C_p, N) implies $(j^*)^{-1}(\alpha_0) \neq \emptyset$ where $j: N \to C_p$ is the inclusion of the codomain of p into the mapping cone. It suffices to show that $q: N \to S^3$ does not extend to C_p . Every extension ψ' of q to the 4-skeleton $C_p^{(4)}$ can be extended to a map $\psi: C_p \to K(\mathbb{Z},3)$ defining an element $\alpha \in (j^*)^{-1}(\alpha_0)$. We show that such ψ cannot be deformed to $S^3 \subset K(\mathbb{Z},3)$. Here we assume that $K(\mathbb{Z},3)^{(5)} = S^3 \cup_{\nu} D^5$.

For a cohomology class x we denote by \bar{x} its mod 2 reduction. Since q is a map of degree one, $\bar{\alpha}_0 \neq 0$. We recall that the obstruction to retraction of $S^3 \cup_{\nu} D^5$ to S^3 is the Steenrod square $Sq^2\bar{a}$. By the naturality of primary obstructions the obstruction to deform the map $\psi: C_p \to K(\mathbb{Z}, 3)$ to S^3 is $\psi^*(Sq^2\bar{a}) = Sq^2\bar{\alpha}$. We show that $Sq^2\bar{\alpha} \neq 0$. Let E be the disk bundle associated with the circle budle p. For the use of cohomology we can identify N with E. Then $M = \partial E$ and the pair (E, M) can be identified with the mapping cone C_p . In view of the Thom isomorphism there is $x \in H^1(N; \mathbb{Z}_2)$ such that $x \cup u = \bar{\alpha}$ where u is the mod 2 Thom class. Since $j^*(\bar{\alpha}) = \bar{\alpha}_0$ the naturality of the cup product

$$\begin{array}{ccc} H^{3}(N,\partial N;\mathbb{Z}_{2}) \times H^{2}(N,\partial N;\mathbb{Z}_{2}) & \xrightarrow{\cup} & H^{5}(N,\partial N;\mathbb{Z}_{2}) \\ & & & & & \\ j^{*} \times 1 & & & & 1 \\ H^{3}(N;\mathbb{Z}_{2}) \times H^{2}(N,\partial N;\mathbb{Z}_{2}) & \xrightarrow{\cup} & H^{5}(N,\partial N;\mathbb{Z}_{2}) \end{array}$$

implies that $\bar{\alpha}_0 \cup u = \bar{\alpha} \cup u$. Then

$$0 \neq \bar{\alpha}_0 \cup u = \bar{\alpha} \cup u = x \cup u \cup u = x \cup Sq^2u = Sq^2(x \cup u) = Sq^2(\bar{\alpha}).$$

Here we use the equality $Sg^2u = u \cup u$ and the Cartan formula for Steenrod squares.

(d) The manifold M is stably parallelizable as the total space of an orientable S^1 -bundle over a stably parallelizable manifold N.

We recall that for every closed manifold M there is a hyperbolization $f: W \to M$ which a map of degree one of a closed aspherical manifold which is surjective on the fundamental groups. Moreover, the map f induces an isomorphism between stable tangent bundles of Mand W [DJ],[CD],[Gro2].

Theorem 5.4. There is a map of a closed aspherical 4-manifold $g: W \to T^3$ onto a 3-torus that induces an epimorphism of the fundamental groups $g_{\#}: \pi_1(W) \to \mathbb{Z}^3$ such that $\operatorname{cat} g_{\#} = 3$ and $\operatorname{cd}(g_{\#}) < 3$.

Proof. Let M be the Bolotov's example and let $f: W \to M$ be the hyperbolization of M. Let $g = q_2 \circ u_M \circ f$. Since Bolotov's example is stably parallelizable and the hyperbolization is a tangential map, by Lemma 5.2 we obtain that $f^*: \pi_s^3(M) \to \pi_s^3(W)$ is injective. By dimensional reason $[M, S^3] = \pi_s^3(M)$ and $[W, S^3] = \pi_s^3(W)$. Thus, since by Proposition 5.3 (c) the map $q_1q_2u_M: M \to S^3$ is essential, the map $q_1 \circ g: W \to S^3$ is essential as well. Hence, the map g cannot be deformed to the 2-skeleton. Therefore, $\operatorname{cat} g > 2$. Clearly, $\operatorname{cat} g = \operatorname{cat} g_{\#} = 3$.

By Proposition 5.3 (a) the homomorphism $(q_1 \circ u_M)^*$ is trivial on 3-dimensional cohomology. Hence, so is g^* . This means that $\operatorname{cd}(g_{\#}) < 3$.

Acknowledgments

The first author was supported by the Simons Foundation Grant.

References

- [B] H. Bechtell, The theory of groups, Addison-Wesley, 1971.
- [Be] I. Berstein, On the Lusternik-Schnirelmann category of Grassmannians. Math. Proc. Camb. Philos. Soc. 79 (1976) 129-134.
- [Bo] D. Bolotov, Gromov's macroscopic dimension conjecture, AGT 6 (2006), 1669-1676.
- [Br] K. Brown, Cohomology of Groups. Graduate Texts in Mathematics, 87 Springer, New York Heidelberg Berlin, 1994.
- [CD] R. M. Charney, M. W. Davis, Strict hyperbolization, Topology 34 (1995), no.2, 329-350.
- [CLOT] O. Cornea, G. Lupton, J. Oprea, D. Tanre, Lusternik-Schnirelmann Category, AMS, 2003.
 [DJ] M. W. Davis and T. Januszkiewicz, Hyperbolization of polyhedra, J. Differential Geom. 34 (1991),
- no.2, 347-388.
 [DR] A. Dranishnikov, Yu. Rudyak, On the Berstein-Svarc theorem in dimension 2. Math. Proc. Cambridge Philos. Soc. 146 (2009), no. 2, 407-413.
- [EG] S. Eilenberg, T. Ganea, On the Lusternik-Schnirelmann Category of Abstract Groups. Annals of Mathematics, 65, (1957), 517-518.
- [Gr] M. Grant, https://mathoverflow.net/questions/89178/cohomological-dimension-of-a-homomorphism
- [Gro1] M. Gromov Positive curvature, macroscopic dimension, spectral gaps and higher signatures, Functional analysis on the eve of the 21st century. Vol. II, Birkhauser, Boston, MA, 1996.
- [Gro2] M. Gromov, Hyperbolic groups, Essays in group theory, Math. Sci. Res. Inst. Publ., 8, Springer, New York, (1987) 75–263.
- [Hu] S.-T. Hu, Homotopy theory, Acad. Press, 1959.
- [LS] L. Lusternik, L. Schnirelmann, "Sur le probleme de trois geodesiques fermees sur les surfaces de genre 0", Comptes Rendus de l'Academie des Sciences de Paris, 189: (1929) 269-271.
- [Ma1] A. I. Mal'tsev, On a class of homogeneous spaces, Izvestiya Rossiiskoi Akademii Nauk. Seriya Matematicheskaya, vol. 13 (1949), no 3, 201-212.
- [Ma2] A. I. Mal'tsev, Nilpotent groups without torsion, Izvestiya Rossiiskoi Akademii Nauk. Seriya Matematicheskaya, vol. 13 (1949), no 1, 9–32.
- [Ra] M. S. Raghunathan, Discrete subgroups of Lie groups, Springer, 1972.
- [Ru] Yu. Rudyak, On category weight and its applications. Topology 38 (1999) no. 1, 37–55.
- [Sch] A. Schwarz, The genus of a fibered space. Trudy Moscov. Mat. Obsc. 10, 11 (1961 and 1962), 217-272, 99-126.
- [Sc] J. Scott, On the Topological Complexity of Maps, preprint arXiv:2011.10646 2020.
- [Sw] R. Switzer, Algebraic ¹opologyĐHomotopy and Homology. Springer, Berlin, 1975.

Alexander Dranishnikov, Department of Mathematics, University of Florida, 358 Little Hall, Gainesville, FL 32611-8105, USA

Email address: dranish@math.ufl.edu

NURSULTAN KUANYSHOV, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, 358 LITTLE HALL, GAINESVILLE, FL 32611-8105, USA

Email address: kuanyshov@math.ufl.edu