

DIFFERENTIAL OPERATOR APPROACH TO \imath QUANTUM GROUPS AND THEIR OSCILLATOR REPRESENTATIONS

ZHAOBING FAN, JICHENG GENG, AND SHAOLONG HAN

ABSTRACT. For a quasi-split Satake diagram, we define a modified q -Weyl algebra, and show that there is an algebra homomorphism between it and the corresponding \imath quantum group. In other words, we provide a differential operator approach to \imath quantum groups. Meanwhile, the Oscillator representations are obtained. The crystal basis of the irreducible subrepresentations of these Oscillator representations are constructed.

1. INTRODUCTION

Quantum groups were introduced independently by Drinfeld and Jimbo in [7, 18]. To date, the theory of quantum groups has played important roles in representation theory, low dimensional topology and mathematical physics etc. From the perspective of the development of quantum groups, mathematicians have extended various theories of quantum groups to more and more generalized cartan data, such as from semi-simple cases to Kac-Moody cases, and then to Borcherds cases. Another direction to develop quantum groups is so called \imath quantum groups, which are the main objects studied in the current paper.

Let \mathfrak{g} be a Kac-Moody algebra and θ be an involution on \mathfrak{g} . The pair $(\mathfrak{g}, \mathfrak{g}^\theta)$ is called a symmetric pair, where \mathfrak{g}^θ is the θ -stable subalgebra of \mathfrak{g} . The quantization of $(\mathfrak{g}, \mathfrak{g}^\theta)$ is denoted by $(\mathbf{U}_q(\mathfrak{g}), \mathbf{U}_q(\mathfrak{g}^\theta))$ which is called the quantum symmetric pair. In particular, the algebra $\mathbf{U}_q(\mathfrak{g}^\theta)$ is called the \imath quantum group, which is a coideal subalgebra of $\mathbf{U}_q(\mathfrak{g})$. In general, the algebra $\mathbf{U}_q(\mathfrak{g}^\theta)$ is not a Hopf algebra. The \imath quantum groups are classified by Satake diagrams. Moreover, \imath quantum groups can be degenerated to ordinary quantum groups if the Satake diagram is quasi-split of diagonal type.

In 2013, Bao and Wang initiated the \imath -program in [3]. The purpose of this program is to generalize various results for ordinary quantum groups to \imath quantum groups. From geometric side, in [1], Bao, Kujawa, Li and Wang gave a geometric realization of \mathbf{U}^j and \mathbf{U}^i , the \imath quantum groups of type B/C , by using equivariant function on double flag varieties. This generalizes the classic work of Beilinson, Lusztig and MacPherson for ordinary quantum group in [4]. Later on, the first author and his collaborators further gave a geometric realization of the \imath quantum groups of affine type C in [11] by using a similar approach. Moreover, the first author, Ma and Xiao provided another geometric realization of the \imath quantum groups \mathbf{U}^j in [14] by using equivariant K -group of corresponding Steinberg variety. This fills Ginzburg and Vasserot's work in [15] into \imath -program.

From algebraic side, q -Schur duality, Hall algebra and categorification are important theory related to quantum groups. In the framework of \imath -program, these theories have been generalized to certain \imath quantum groups. The Hecke-algebraic approach to the \imath quantum groups was given in [12], which was a generalization of affine type A construction in [8]. The three-parameter q -Schur duality between \imath quantum groups and Hecke algebras was studied in [13], which can be degenerated to q -Schur dualities of affine types B , C and D by various specialization of parameters. In [25, 5], Ringel and Bridgeland gave a realization of the half part and

Key words and phrases. \imath quantum group, differential operator, Oscillator representation, crystal basis.

entire of quantum groups by using Hall algebras, respectively. The \imath Hall algebra approach toward \imath quantum groups was developed by Lu and Wang in series papers [20, 21, 22, 23]. The highest weight theory and crystal (global) basis theory of \imath quantum groups \mathbf{U}^j of type AIII was studied in [26] and [27]. The categorification of the modified \imath quantum group $\check{\mathbf{U}}^j$ was developed in [2]. In [9, 10], Du and Wu presented a new realization of the \imath quantum groups \mathbf{U}^j and \mathbf{U}^i in terms of a BLM type basis.

In this paper, we provide a differential operator realization of \imath quantum groups ${}^iU(\mathcal{S})$ of type AIII. This realization can be regarded as an application of Hayashi's approach [16] to \imath quantum groups. To achieve this goal, we introduce the modified q -Weyl algebra $\mathbf{A}_q(\mathcal{S})$ associated to Satake diagrams, generated by certain linear operators on the polynomial ring $\mathbb{Q}(q)[X_0, X_1, \dots, X_{r+1}]$, and show that there is an algebra homomorphism from ${}^iU(\mathcal{S})$ to $\mathbf{A}_q(\mathcal{S})$. Thus we obtain naturally a differential operator realization of ${}^iU(\mathcal{S})$ on the q -Weyl algebra $\mathbf{A}_q(A_{r+1})$ and representations of ${}^iU(\mathcal{S})$ on the polynomial ring which are called the Oscillator representations. Following the framework of crystal basis theory in [19] and [17, Chapter 4], we construct the crystal basis of irreducible modules arising from the Oscillator representations of \imath quantum groups ${}^iU(A_{2r+1})$, ${}^iU(A_{2r+1}^{(1)})$ and ${}^iU(A_1^{(1)})$.

The paper is organized as follows. In Section 2, we recall the differential operator realization of the quantum group $\mathbf{U}_q(\mathfrak{sl}_{r+2})$ and its Oscillator representation. In Section 3, we define the modified q -Weyl algebras $\mathbf{A}_q(\mathcal{S})$ associated with Satake diagrams and study their representations. In Section 4, we construct the differential operator realizations of \imath quantum groups ${}^iU(\mathcal{S})$. In Section 5, we study the Oscillator representations of ${}^iU(\mathcal{S})$ and the crystal basis of irreducible modules over \imath quantum groups ${}^iU(A_{2r+1})$, ${}^iU(A_{2r+1}^{(1)})$ and ${}^iU(A_1^{(1)})$.

Acknowledgements. Z. Fan was partially supported by the NSF of China grant 11671108, the NSF of Heilongjiang Province grant JQ2020A001, and the Fundamental Research Funds for the central universities.

Notations. Let

$$[a] = \frac{q^a - q^{-a}}{q - q^{-1}}, \quad [a]^{k!} = [ka][k(a-1)] \cdots [2k][k], \quad X^{(a)_k} = \frac{X^a}{[a]^{k!}} \quad (a, k \in \mathbb{Z}),$$

$$(a; x)_0 = 1, \quad (a; x)_n = (1-a)(1-ax) \cdots (1-ax^{n-1}), \quad n \geq 1,$$

$$\begin{bmatrix} n \\ d \end{bmatrix} = \begin{cases} \frac{[n][n-1] \cdots [n-d+1]}{[d]!}, & \text{if } d > 0, \\ 1, & \text{if } d = 0, \\ 0, & \text{if } d < 0. \end{cases}$$

2. DIFFERENTIAL OPERATOR REALIZATION OF $\mathbf{U}_q(\mathfrak{sl}_{r+2})$

Let $r \geq 0$ and $\mathbb{I} = \{0, 1, \dots, r+1\}$. The q -Weyl algebra $\mathbf{A}_q(A_{r+1})$ associated to Dynkin diagram of type A_{r+1} is generated by $\mathcal{D}_i, \mathcal{X}_i, \mathcal{M}_i^{\pm 1}$ ($i \in \mathbb{I}$) over $\mathbb{Q}(q)$ subject to the following relations:

$$(2.1) \quad \mathcal{M}_i \mathcal{M}_i^{-1} = \mathcal{M}_i^{-1} \mathcal{M}_i = 1, \quad \mathcal{M}_i \mathcal{M}_j = \mathcal{M}_j \mathcal{M}_i,$$

$$(2.2) \quad \mathcal{D}_i \mathcal{M}_j = \mathcal{M}_j \mathcal{D}_i, \quad \mathcal{X}_i \mathcal{M}_j = \mathcal{M}_j \mathcal{X}_i, \quad \mathcal{D}_i \mathcal{X}_j = \mathcal{X}_j \mathcal{D}_i, \quad \text{if } i \neq j,$$

$$(2.3) \quad \mathcal{D}_i \mathcal{D}_j = \mathcal{D}_j \mathcal{D}_i, \quad \mathcal{X}_i \mathcal{X}_j = \mathcal{X}_j \mathcal{X}_i,$$

$$(2.4) \quad \mathcal{D}_i \mathcal{M}_i = q \mathcal{M}_i \mathcal{D}_i, \quad \mathcal{X}_i \mathcal{M}_i = q^{-1} \mathcal{M}_i \mathcal{X}_i,$$

$$(2.5) \quad \mathcal{D}_i \mathcal{X}_i = \frac{q \mathcal{M}_i - q^{-1} \mathcal{M}_i^{-1}}{q - q^{-1}}, \quad \mathcal{X}_i \mathcal{D}_i = \frac{\mathcal{M}_i - \mathcal{M}_i^{-1}}{q - q^{-1}}.$$

Let $\mathbb{P} = \mathbb{Q}(q)[X_0, X_1, \dots, X_{r+1}]$ be the polynomial ring over $\mathbb{Q}(q)$. We define $\mathbf{A}_q(A_{r+1})$ acting on \mathbb{P} , for any $f \in \mathbb{P}$, as follows

$$(2.6) \quad \begin{aligned} \mathcal{D}_i f(X_0, \dots, X_{r+1}) &= \frac{f(X_0, \dots, qX_i, \dots, X_{r+1}) - f(X_0, \dots, q^{-1}X_i, \dots, X_{r+1})}{qX_i - q^{-1}X_i}, \\ \mathcal{X}_i f(X_0, \dots, X_i, \dots, X_{r+1}) &= X_i f(X_0, \dots, X_i, \dots, X_{r+1}), \\ \mathcal{M}_i^{\pm 1} f(X_0, \dots, X_i, \dots, X_{r+1}) &= f(X_0, \dots, q^{\pm 1}X_i, \dots, X_{r+1}). \end{aligned}$$

Theorem 2.1 ([16, Proposition 2.1]). *The polynomial ring \mathbb{P} is an irreducible $\mathbf{A}_q(A_{r+1})$ -module.*

The generators $\mathcal{D}_i, \mathcal{X}_i, \mathcal{M}_i$ ($i \in \mathbb{I}$) of the algebra $\mathbf{A}_q(A_{r+1})$ can be seen as linear operators on the polynomial ring \mathbb{P} . Moreover, the operator \mathcal{D}_i is called *q-differentiation operator*, which satisfying the *q*-analog of the Leibniz rule as follows:

$$\begin{aligned} &\mathcal{D}_i(f(X_0, \dots, X_i, \dots, X_{r+1})g(X_0, \dots, X_i, \dots, X_{r+1})) \\ &= \mathcal{D}_i f(X_0, \dots, X_i, \dots, X_{r+1})g(X_0, \dots, q^{-1}X_i, \dots, X_{r+1}) \\ &\quad + f(X_0, \dots, qX_i, \dots, X_{r+1})\mathcal{D}_i g(X_0, \dots, X_i, \dots, X_{r+1}). \end{aligned}$$

Let $c_{ij} = 2\delta_{ij} - \delta_{i,j+1} - \delta_{i+1,j}$ for $i, j \in \mathbb{Z}$. Recall that the quantum group $\mathbf{U}_q(\mathfrak{sl}_{r+2})$ is generated by $E_i, F_i, K_i^{\pm 1}$ ($0 \leq i \leq r$) satisfying the following relations:

$$\begin{aligned} K_i K_i^{-1} &= K_i^{-1} K_i = 1, K_i K_j = K_j K_i, \\ K_i E_j K_i^{-1} &= q^{c_{ij}} E_j, K_i F_j K_i^{-1} = q^{-c_{ij}} F_j, \\ E_i F_j - F_j E_i &= \delta_{ij} \frac{K_i - K_i^{-1}}{q - q^{-1}}, \\ E_i E_j &= E_j E_i, F_i F_j = F_j F_i, \quad \text{if } |i - j| > 1, \\ E_i^2 E_j - (q + q^{-1}) E_i E_j E_i + E_j E_i^2 &= 0, \quad \text{if } |i - j| = 1, \\ F_i^2 F_j - (q + q^{-1}) F_i F_j F_i + F_j F_i^2 &= 0, \quad \text{if } |i - j| = 1. \end{aligned}$$

Theorem 2.2 ([16, Theorem 3.2]). *There is a $\mathbb{Q}(q)$ -algebra homomorphism $\chi_r : \mathbf{U}_q(\mathfrak{sl}_{r+2}) \rightarrow \mathbf{A}_q(A_{r+1})$ such that*

$$E_i \mapsto \mathcal{X}_i \mathcal{D}_{i+1}, \quad F_i \mapsto \mathcal{X}_{i+1} \mathcal{D}_i, \quad K_i \mapsto \mathcal{M}_i \mathcal{M}_{i+1}^{-1}.$$

By Theorems 2.1 and 2.2, the polynomial ring \mathbb{P} is a $\mathbf{U}_q(\mathfrak{sl}_{r+2})$ -module via pull-back map. The polynomial ring \mathbb{P} is called the Oscillator representation of $\mathbf{U}_q(\mathfrak{sl}_{r+2})$. Moreover, we have

Theorem 2.3 ([16, Theorem 4.1]). *Let $s \geq 0$, the subspace of \mathbb{P} spanned by the set*

$$\{X_0^{a_0} X_1^{a_1} \cdots X_{r+1}^{a_{r+1}} \mid a_0 + a_1 + \cdots + a_{r+1} = s\}$$

is an irreducible $\mathbf{U}_q(\mathfrak{sl}_{r+2})$ -module.

3. MODIFIED *q*-WEYL ALGEBRA

In this section, we shall introduce the modified *q*-Weyl algebra associated to a given Satake diagram. Let \mathbf{I} be a finite set and assume $|\mathbf{I}| \geq r + 2$. Denote by D the Dynkin diagram of type $A_{|\mathbf{I}|-1}$ or $A_{|\mathbf{I}|-1}^{(1)}$. The nodes in D are labeled by the elements in \mathbf{I} . For $i, j \in \mathbf{I}$, we define the Cartan datum (\mathbf{I}, \cdot) on D as follows:

$$i \cdot j = \begin{cases} 2, & \text{if } i = j, \\ -k, & \text{if there are } k \text{ lines connected between } i \text{ and } j, \\ 0, & \text{otherwise.} \end{cases}$$

Let $(Y, X, \langle \cdot, \cdot \rangle)$ be a root datum of type (\mathbf{I}, \cdot) ; cf. [24, Section 2.1]. Let τ be an involution of the Cartan datum (\mathbf{I}, \cdot) and W be the Weyl group generated by simple reflections s_i for $i \in \mathbf{I}$. Given a subset $\mathbf{I}_\bullet \subset \mathbf{I}$, let $W_{\mathbf{I}_\bullet} = \langle s_i \mid i \in \mathbf{I}_\bullet \rangle$ be the parabolic subgroup of W with longest element w_\bullet , and let ρ_\bullet (resp. ρ_\bullet^\vee) be the half sum of all positive roots (resp. coroots) in the root system $R_{\mathbf{I}_\bullet}$ (resp. $R_{\mathbf{I}_\bullet}^\vee$).

Let $\mathbf{I}_\circ = \mathbf{I} \setminus \mathbf{I}_\bullet$. A Satake diagram \mathcal{S} is the pair $(\mathbf{I} = \mathbf{I}_\circ \cup \mathbf{I}_\bullet, \tau)$ such that $\tau(\mathbf{I}_\bullet) = \mathbf{I}_\bullet$, the actions of τ and $-w_\bullet$ on \mathbf{I}_\bullet coincide, and $\langle \rho_\bullet^\vee, j' \rangle \in \mathbb{Z}$ if $\tau j = j \in \mathbf{I}_\circ$. By abuse of notations, denote by \mathbb{I} the set of τ -orbits of \mathbf{I} .

In this paper, we only consider the case $\mathbf{I} = \mathbf{I}_\circ$. We use natural numbers $\{0, 1, 2, \dots\}$ to label the elements in \mathbf{I} .

Definition 3.1. *Given a Satake diagram \mathcal{S} , the associated modified q -Weyl algebra, $\mathbf{A}_q(\mathcal{S})$, is generated by $\mathfrak{d}_i, \mathfrak{r}_i, \mathfrak{m}_i$ ($i \in \mathbb{I}$) over $\mathbb{Q}(q)$ subject to the following relations:*

$$(3.1) \quad \mathfrak{m}_i \mathfrak{m}_i^{-1} = \mathfrak{m}_i^{-1} \mathfrak{m}_i = 1, \quad \mathfrak{m}_i \mathfrak{m}_j = \mathfrak{m}_j \mathfrak{m}_i,$$

$$(3.2) \quad \mathfrak{d}_i \mathfrak{m}_j = \mathfrak{m}_j \mathfrak{d}_i, \quad \mathfrak{r}_i \mathfrak{m}_j = \mathfrak{m}_j \mathfrak{r}_i, \quad \mathfrak{d}_i \mathfrak{r}_j = \mathfrak{r}_j \mathfrak{d}_i, \quad \text{if } i \neq j,$$

$$(3.3) \quad \mathfrak{d}_i \mathfrak{d}_j = \mathfrak{d}_j \mathfrak{d}_i, \quad \mathfrak{r}_i \mathfrak{r}_j = \mathfrak{r}_j \mathfrak{r}_i,$$

$$(3.4) \quad \mathfrak{d}_i \mathfrak{m}_i = q^{\xi_i} \mathfrak{m}_i \mathfrak{d}_i, \quad \mathfrak{r}_i \mathfrak{m}_i = q^{-\xi_i} \mathfrak{m}_i \mathfrak{r}_i,$$

$$(3.5) \quad \mathfrak{d}_i \mathfrak{r}_i = \frac{q^{\xi_i} \mathfrak{m}_i - q^{-\xi_i} \mathfrak{m}_i^{-1}}{q - q^{-1}}, \quad \mathfrak{r}_i \mathfrak{d}_i = \frac{\mathfrak{m}_i - \mathfrak{m}_i^{-1}}{q - q^{-1}},$$

where $\xi_i = 1 - i \cdot \tau i$.

In the rest of the paper, we shall consider the following Satake diagrams.

Diagram I : Satake diagram of type A_{2r+3} .




Diagram II: Satake diagram of type $A_{2r+2,1}$.




Diagram III: Satake diagram of type $A_{2r+3}^{(1)}$.







Diagram IV: Satake diagram of type $A_{2r+2,1}^{(1)}$.Diagram V: Satake diagram of type $A_{1,2r+2}^{(1)}$.Diagram VI: Satake diagram of type $A_{2r+1,2}^{(1)}$.

Proposition 3.1. *Let \mathcal{S} be the Satake diagram in Diagram I-VI, then there is a $\mathbb{Q}(q)$ -algebra homomorphism $\iota : \mathbf{A}_q(\mathcal{S}) \rightarrow \mathbf{A}_q(A_{r+1})$ which sends*

$$\begin{aligned} \mathfrak{d}_i &\mapsto \mathcal{D}_i\left(\sum_{k=0}^{\xi_i-1} \mathcal{M}_i^{\xi_i-1-2k}\right), & \mathfrak{x}_i &\mapsto \mathcal{X}_i, & \mathfrak{m}_i^{\pm 1} &\mapsto \mathcal{M}_i^{\pm \xi_i}, & \text{if } \xi_i > 0, \\ \mathfrak{d}_i &\mapsto -\mathcal{D}_i\left(\sum_{k=0}^{-\xi_i-1} \mathcal{M}_i^{-\xi_i-1-2k}\right), & \mathfrak{x}_i &\mapsto \mathcal{X}_i, & \mathfrak{m}_i^{\pm 1} &\mapsto \mathcal{M}_i^{\pm \xi_i}, & \text{if } \xi_i < 0. \end{aligned}$$

Proof. Since \mathfrak{d}_i , \mathfrak{x}_i and \mathfrak{m}_i are generators of $\mathbf{A}_q(\mathcal{S})$, it's enough to show that the map ι is well-defined. The verification of relations (3.1)-(3.4) is straightforward. We shall only verify the relations in (3.5). If $\xi_i > 0$, by using the relations (2.4) and (2.5), we have

$$\begin{aligned} \iota(\mathfrak{d}_i \mathfrak{x}_i) &= \mathcal{D}_i\left(\sum_{k=0}^{\xi_i-1} \mathcal{M}_i^{\xi_i-1-2k}\right) \mathcal{X}_i = \mathcal{D}_i \mathcal{X}_i \left(\sum_{k=0}^{\xi_i-1} q^{\xi_i-1-2k} \mathcal{M}_i^{\xi_i-1-2k}\right) \\ &= \frac{q\mathcal{M}_i - q^{-1}\mathcal{M}_i^{-1}}{q - q^{-1}} \left(\sum_{k=0}^{\xi_i-1} q^{\xi_i-1-2k} \mathcal{M}_i^{\xi_i-1-2k}\right) = \frac{q^{\xi_i} \mathcal{M}_i^{\xi_i} - q^{-\xi_i} \mathcal{M}_i^{-\xi_i}}{q - q^{-1}} \\ &= \frac{q^{\xi_i} \iota(\mathfrak{m}_i) - q^{-\xi_i} \iota(\mathfrak{m}_i^{-1})}{q - q^{-1}}, \end{aligned}$$

on the other hand,

$$\iota(\mathfrak{x}_i \mathfrak{d}_i) = \mathcal{X}_i \mathcal{D}_i\left(\sum_{k=0}^{\xi_i-1} \mathcal{M}_i^{\xi_i-1-2k}\right) = \frac{\mathcal{M}_i - \mathcal{M}_i^{-1}}{q - q^{-1}} \left(\sum_{k=0}^{\xi_i-1} \mathcal{M}_i^{\xi_i-1-2k}\right)$$

$$\begin{aligned}
&= \frac{\sum_{k=0}^{\xi_i-1} \mathcal{M}_i^{\xi_i-2k} - \sum_{k=0}^{\xi_i-1} \mathcal{M}_i^{\xi_i-2-2k}}{q - q^{-1}} = \frac{\sum_{k=0}^{\xi_i-1} \mathcal{M}_i^{\xi_i-2k} - \sum_{k=1}^{\xi_i} \mathcal{M}_i^{\xi_i-2k}}{q - q^{-1}} \\
&= \frac{\mathcal{M}_i^{\xi_i} - \mathcal{M}_i^{-\xi_i}}{q - q^{-1}} = \frac{\iota(\mathbf{m}_i) - \iota(\mathbf{m}_i^{-1})}{q - q^{-1}}.
\end{aligned}$$

The case for $\xi_i < 0$ can be shown similiary. \square

Remark 3.1. In Diagram I, if there is no line connected between the nodes $r+1$ and $r+2$, the diagram is called the Satake diagram of diagonal type. In this case, the modified q -Weyl algebra $\mathbf{A}_q(\mathcal{S})$ is exactly the q -Weyl algebra $\mathbf{A}_q(A_{r+1})$.

Let $\mathbf{X} = (X_0, \dots, X_{r+1})$ and $\mathbf{a} = (a_i)_{i \in \mathbb{I}} \in \mathbb{Z}_{\geq 0}^{r+2}$. We denote $\mathbf{X}^{\mathbf{a}} = X_0^{a_0} X_1^{a_1} \dots X_{r+1}^{a_{r+1}}$. Let \mathbf{e}_i be the $(r+2)$ -tuple such that the i -th entry is 1 and 0 otherwise. We define a lexicographic order $>_{lex}$ on the set $\mathbb{Z}_{\geq 0}^{r+2}$ as follows.

For any $\mathbf{a} = (a_0, \dots, a_{r+1})$, $\mathbf{b} = (b_0, \dots, b_{r+1}) \in \mathbb{Z}_{\geq 0}^{r+2}$, $\mathbf{a} >_{lex} \mathbf{b}$ if the leftmost nonzero entry of $\mathbf{a} - \mathbf{b} \in \mathbb{Z}^{r+2}$ is positive.

Theorem 3.2. The polynomial ring \mathbb{P} is a $\mathbf{A}_q(\mathcal{S})$ -module which is irreducible if q is not a root of unity.

Proof. By Theorem 2.1 and Proposition 3.1, \mathbb{P} is a $\mathbf{A}_q(\mathcal{S})$ -module via pull-back. More precisely, the generators \mathfrak{d}_i , \mathfrak{r}_i and \mathfrak{m}_i in $\mathbf{A}_q(\mathcal{S})$ act on \mathbb{P} as follows:

$$(3.6) \quad \mathfrak{d}_i \mathbf{X}^{\mathbf{a}} = [\xi_i a_i] \mathbf{X}^{\mathbf{a} - \mathbf{e}_i}, \quad \mathfrak{r}_i \mathbf{X}^{\mathbf{a}} = \mathbf{X}^{\mathbf{a} + \mathbf{e}_i}, \quad \mathfrak{m}_i \mathbf{X}^{\mathbf{a}} = q^{\xi_i a_i} \mathbf{X}^{\mathbf{a}},$$

where $\mathfrak{d}_i \mathbf{X}^{\mathbf{a}} = 0$ if $a_i = 0$.

Let \mathbb{P}' be a nonzero submodule of \mathbb{P} . We choose a non-zero vector $w = \sum_{\mathbf{a}} c_{\mathbf{a}} \mathbf{X}^{\mathbf{a}} \in \mathbb{P}'$ with $c_{\mathbf{a}} \in \mathbb{Q}(q)$. Let $\mathbf{k} = (k_i)_{0 \leq i \leq r+1}$ be the maximum element in the set $\{\mathbf{a} \in \mathbb{Z}_{\geq 0}^{r+2} | c_{\mathbf{a}} \neq 0\}$ with respect to the order $>_{lex}$. It follows that

$$\mathfrak{d}_0^{k_0} \mathfrak{d}_1^{k_1} \dots \mathfrak{d}_{r+1}^{k_{r+1}} w = c_{\mathbf{k}} \prod_{i=0}^{r+1} [k_i]^{\xi_i} \mathbf{X}^{\mathbf{0}}.$$

Since the coefficient of $\mathbf{X}^{\mathbf{0}}$ is not zero, we have $\mathbf{X}^{\mathbf{0}} \in \mathbb{P}'$. For any element $v = \mathbf{X}^{\mathbf{a}} \in \mathbb{P}$ with $\mathbf{a} = (a_i)_{0 \leq i \leq r+1}$, we have $\mathfrak{r}_0^{a_0} \mathfrak{r}_1^{a_1} \dots \mathfrak{r}_{r+1}^{a_{r+1}} \mathbf{X}^{\mathbf{0}} = v \in \mathbb{P}'$, i.e., $\mathbb{P} \subseteq \mathbb{P}'$. Hence $\mathbb{P} = \mathbb{P}'$. Since \mathbb{P}' is arbitrary, \mathbb{P} is irreducible. \square

4. DIFFERENTIAL OPERATOR APPROACH TO ι QUANTUM GROUPS

In this section, we shall show that there is an algebra homomorphism from ${}^{\iota}U(\mathcal{S})$ to $\mathbf{A}_q(\mathcal{S})$ for a quasi-split Satake diagram \mathcal{S} . In other words, this provides a differential operator approach to ${}^{\iota}U(\mathcal{S})$. Let's first recall the definition of ${}^{\iota}U(\mathcal{S})$ from [6].

Definition 4.1. Given a Satake diagram \mathcal{S} , the associated ι quantum groups ${}^{\iota}U(\mathcal{S})$ is the $\mathbb{Q}(q)$ -algebra generated by B_i, H_i ($i \in \mathbf{I}$), subjecting to the following relations

$$(4.1) \quad H_i H_{\tau i} = 1, \quad H_i H_j = H_j H_i,$$

$$(4.2) \quad H_j B_i - q^{i \cdot \tau j - i \cdot j} B_i H_j = 0,$$

$$(4.3) \quad B_i B_j - B_j B_i = 0, \quad \text{if } i \cdot j = 0 \text{ and } \tau i \neq j,$$

$$(4.4) \quad \sum_{n=0}^{1-i \cdot j} (-1)^n B_i^{(n)} B_j B_i^{(1-i \cdot j - n)} = 0, \quad \text{if } j \neq \tau i \neq i,$$

$$(4.5) \quad \sum_{n=0}^{1-i\cdot\tau i} (-1)^{n+i\cdot\tau i} B_i^{(n)} B_{\tau i} B_i^{(1-i\cdot\tau i-n)} \\ = \frac{1}{q-q^{-1}} \left(q^{i\cdot\tau i+3(\delta_{i,0}-\delta_{i,\tau 0})\delta_{0,\tau 0,-1}} (q^{-2}; q^{-2})_{-i\cdot\tau i \varsigma_{\tau i}} B_i^{(-i\cdot\tau i)} H_i \right. \\ \left. - q^{3(\delta_{i,\tau 0}-\delta_{i,0})\delta_{0,\tau 0,-1}} (q^2; q^2)_{-i\cdot\tau i \varsigma_i} B_i^{(-i\cdot\tau i)} H_{\tau i} \right), \quad \text{if } \tau i \neq i,$$

$$(4.6) \quad \sum_{n=0}^{1-i\cdot j} (-1)^n \begin{bmatrix} 1-i\cdot j \\ n \end{bmatrix} B_i^n B_j B_i^{1-i\cdot j-n} = \delta_{i,j,-1} q \varsigma_i B_j, \quad \text{if } \tau i = i \neq j.$$

For the Diagram I-VI, let

$$(\varsigma_i, \varsigma_{\tau i}) = \begin{cases} (q^{-1}, q^{-1}), & \text{if } i \cdot \tau i = 2, \\ (1, 1), & \text{if } i \cdot \tau i = 0, \\ (q, 1), & \text{if } i \cdot \tau i = -1, \\ (q, -q^{-3}), & \text{if } i \cdot \tau i = -2. \end{cases}$$

For $i \in \mathbb{I}$, let

$$f_i = B_i, \quad e_i = B_{\tau i}, \quad k_i = H_i, \quad k_i^{-1} = H_{\tau i}, \quad \text{if } \tau i \neq i \text{ and } t_i = B_i, \quad \text{if } \tau i = i.$$

Theorem 4.1. *Given a Satake diagram S in Diagram I-VI, there exists a $\mathbb{Q}(q)$ -algebra homomorphism $\Phi_S : {}^i U(S) \rightarrow \mathbf{A}_q(S)$.*

Proof. For each case of S , we shall give an explicit construction of Φ_S , and show that it is an algebra homomorphism.

(1) If S is A_{2r+1} ($r \geq 0$) in Diagram I, the map Φ_S is given by

$$e_i \mapsto \mathfrak{r}_i \mathfrak{d}_{i+1}, \quad f_i \mapsto \mathfrak{r}_{i+1} \mathfrak{d}_i, \quad k_i \mapsto \mathfrak{m}_i \mathfrak{m}_{i+1}^{-1}.$$

(2) If S is $A_{2r+2,1}$ ($r \geq 0$) in Diagram II, the map Φ_S is given by

$$e_i \mapsto \mathfrak{r}_i \mathfrak{d}_{i+1}, \quad f_i \mapsto \mathfrak{r}_{i+1} \mathfrak{d}_i, \quad k_i \mapsto (-1)^{\delta_{i,r}} \mathfrak{m}_i \mathfrak{m}_{i+1}^{-(-1)^{\delta_{i,r}}}, \quad t_{r+1} \mapsto \mathfrak{r}_{r+1} \mathfrak{d}_{r+1}.$$

(3) If S is $A_{2r+1}^{(1)}$ ($r \geq 1$) in Diagram III, the map Φ_S is given by

$$e_i \mapsto \mathfrak{r}_i \mathfrak{d}_{i+1}, \quad f_i \mapsto \mathfrak{r}_{i+1} \mathfrak{d}_i, \quad k_i \mapsto q^{-2\delta_{i,0}} \mathfrak{m}_i \mathfrak{m}_{i+1}^{-1}.$$

(4) If S is $A_1^{(1)}$ in Diagram III, the map Φ_S is given by

$$e_0 \mapsto \mathfrak{r}_0 \mathfrak{d}_1, \quad f_0 \mapsto \mathfrak{r}_1 \mathfrak{d}_0, \quad k_0 \mapsto q^{-1} \mathfrak{m}_0 \mathfrak{m}_1^{-1}.$$

(5) If S is $A_{2r+2,1}^{(1)}$ ($r \geq 0$) in Diagram IV, the map Φ_S is given by

$$e_i \mapsto \mathfrak{r}_i \mathfrak{d}_{i+1}, \quad f_i \mapsto \mathfrak{r}_{i+1} \mathfrak{d}_i, \quad k_i \mapsto (-1)^{\delta_{i,r}} q^{-2\delta_{i,0}} \mathfrak{m}_i \mathfrak{m}_{i+1}^{-(-1)^{\delta_{i,r}}}, \quad t_{r+1} \mapsto \mathfrak{r}_{r+1} \mathfrak{d}_{r+1}.$$

(6) If S is $A_{1,2r+2}^{(1)}$ ($r \geq 0$) in Diagram V, the map Φ_S is given by

$$e_i \mapsto \mathfrak{r}_{i-1} \mathfrak{d}_i, \quad f_i \mapsto \mathfrak{r}_i \mathfrak{d}_{i-1}, \quad k_i \mapsto (-1)^{\delta_{i,1}} \mathfrak{m}_{i-1}^{(-1)^{\delta_{i,1}}} \mathfrak{m}_i^{-1}, \quad t_0 \mapsto \mathfrak{r}_0 \mathfrak{d}_0.$$

(7) If S is $A_{2r+1,2}^{(1)}$ ($r \geq 1$) in Diagram VI, the map Φ_S is given by

$$e_i \mapsto \mathfrak{r}_i \mathfrak{d}_{i+1}, \quad f_i \mapsto \mathfrak{r}_{i+1} \mathfrak{d}_i, \quad k_i \mapsto (-1)^{\delta_{i,r}} \mathfrak{m}_i \mathfrak{m}_{i+1}^{-(-1)^{\delta_{i,r}}}, \quad t_0 \mapsto \mathfrak{r}_1 \mathfrak{d}_1, \quad t_{r+1} \mapsto \mathfrak{r}_{r+1} \mathfrak{d}_{r+1}.$$

We shall only prove the cases (1)-(4), the other cases can be shown similarly.

(1) By the relations (4.2) and (4.5), we have

$$(4.7) \quad k_i e_j k_i^{-1} = q^{c_{ij} + \delta_{i,r} \delta_{j,r}} e_j, \quad k_i f_j k_i^{-1} = q^{-c_{ij} - \delta_{i,r} \delta_{j,r}} f_j,$$

$$(4.8) \quad e_r^2 f_r + f_r e_r^2 = (q + q^{-1})(e_r f_r e_r - e_r (q k_r + q^{-1} k_r^{-1})),$$

$$(4.9) \quad f_r^2 e_r + e_r f_r^2 = (q + q^{-1})(f_r e_r f_r - (q k_r + q^{-1} k_r^{-1}) f_r).$$

By the involution τ in Diagram I, it follows that $\xi_i = 2^{\delta_{i,r+1}}$. According to the definition of homomorphism Φ_S and relations (3.1)-(3.5), for $i = j = r$ in the relations (4.7), we have

$$\begin{aligned} \Phi_S(k_r e_r k_r^{-1}) &= \mathbf{m}_r \mathbf{m}_{r+1}^{-1} \mathfrak{r}_r \mathfrak{d}_{r+1} \mathbf{m}_r^{-1} \mathbf{m}_{r+1} = q^3 \mathfrak{r}_r \mathfrak{d}_{r+1} = q^3 \Phi_S(e_r), \\ \Phi_S(k_r f_r k_r^{-1}) &= \mathbf{m}_r \mathbf{m}_{r+1}^{-1} \mathfrak{r}_r \mathfrak{d}_{r+1} \mathfrak{d}_r \mathbf{m}_r^{-1} \mathbf{m}_{r+1} = q^{-3} \mathfrak{r}_{r+1} \mathfrak{d}_r = q^{-3} \Phi_S(f_r), \end{aligned}$$

and the verification of other cases in the relations (4.7) is similar.

For the relation (4.8), we have

$$\begin{aligned} &\Phi_S(e_r^2 f_r) + \Phi_S(f_r e_r^2) = \mathfrak{r}_r \mathfrak{d}_{r+1} \mathfrak{r}_r \mathfrak{d}_{r+1} \mathfrak{r}_{r+1} \mathfrak{d}_r + \mathfrak{r}_{r+1} \mathfrak{d}_r \mathfrak{r}_r \mathfrak{d}_{r+1} \mathfrak{r}_r \mathfrak{d}_{r+1} \\ &= (q - q^{-1})^{-2} \mathfrak{r}_r \mathfrak{d}_{r+1} ((\mathbf{m}_r - \mathbf{m}_r^{-1})(q^2 \mathbf{m}_{r+1} - q^{-2} \mathbf{m}_{r+1}^{-1}) + (q^2 \mathbf{m}_r - q^{-2} \mathbf{m}_r^{-1})(q^{-2} \mathbf{m}_{r+1} - q^2 \mathbf{m}_{r+1}^{-1})) \\ &= (q - q^{-1})^{-2} \mathfrak{r}_r \mathfrak{d}_{r+1} ((q^2 + 1) \mathbf{m}_r \mathbf{m}_{r+1} + (q^{-2} + 1) \mathbf{m}_r^{-1} \mathbf{m}_{r+1}^{-1} \\ &\quad - (q^{-2} + q^4) \mathbf{m}_r \mathbf{m}_{r+1}^{-1} - (q^2 + q^{-4}) \mathbf{m}_r \mathbf{m}_{r+1}^{-1}) \\ &= (q + q^{-1})(q - q^{-1})^{-2} \mathfrak{r}_r \mathfrak{d}_{r+1} (q \mathbf{m}_r \mathbf{m}_{r+1} + q^{-1} \mathbf{m}_r^{-1} \mathbf{m}_{r+1}^{-1} \\ &\quad - (q + q(q - q^{-1})^2) \mathbf{m}_r \mathbf{m}_{r+1}^{-1} - (q^{-1} + q^{-1}(q - q^{-1})^2) \mathbf{m}_r^{-1} \mathbf{m}_{r+1}) \\ &= (q + q^{-1}) \mathfrak{r}_r \mathfrak{d}_{r+1} ((q - q^{-1})^{-2} (\mathbf{m}_{r+1} - \mathbf{m}_{r+1}^{-1})(q \mathbf{m}_r - q^{-1} \mathbf{m}_r^{-1}) - q \mathbf{m}_r \mathbf{m}_{r+1}^{-1} - q^{-1} \mathbf{m}_r^{-1} \mathbf{m}_{r+1}) \\ &= (q + q^{-1}) (\mathfrak{r}_r \mathfrak{d}_{r+1} \mathfrak{r}_{r+1} \mathfrak{d}_r \mathfrak{r}_r \mathfrak{d}_{r+1} - q \mathfrak{r}_r \mathfrak{d}_{r+1} \mathbf{m}_r \mathbf{m}_{r+1}^{-1} - q^{-1} \mathfrak{r}_r \mathfrak{d}_{r+1} \mathbf{m}_r^{-1} \mathbf{m}_{r+1}) \\ &= (q + q^{-1}) (\Phi_S(e_r f_r e_r) - q \Phi_S(e_r k_r) - q^{-1} \Phi_S(e_r k_r^{-1})). \end{aligned}$$

(2) For $i, j \in \{r, r+1\}$, the relations (4.4) and (4.6) can be converted to

$$(4.10) \quad e_r^2 t_{r+1} + t_{r+1} e_r^2 = [2] e_r t_{r+1} e_r,$$

$$(4.11) \quad f_r^2 t_{r+1} + t_{r+1} f_r^2 = [2] f_r t_{r+1} f_r$$

$$(4.12) \quad e_i t_{r+1} = t_{r+1} e_i, \quad f_i t_{r+1} = t_{r+1} f_i, \quad \text{if } i \neq r,$$

$$(4.13) \quad t_{r+1}^2 e_r + e_r t_{r+1}^2 = [2] t_{r+1} e_r t_{r+1} + e_r,$$

$$(4.14) \quad t_{r+1}^2 f_r + f_r t_{r+1}^2 = [2] t_{r+1} f_r t_{r+1} + f_r.$$

By the involution τ in Diagram II, we have $\xi_i = (-1)^{\delta_{i,r+1}}$. For the relations (4.10) and (4.13), we have

$$\begin{aligned} &\Phi_S(e_r^2 t_{r+1}) + \Phi_S(t_{r+1} e_r^2) \\ &= \mathfrak{r}_r^2 \mathfrak{d}_{r+1}^2 \frac{\mathbf{m}_{r+1} - \mathbf{m}_{r+1}^{-1}}{q - q^{-1}} + \mathfrak{r}_r^2 \mathfrak{d}_{r+1}^2 \frac{q^2 \mathbf{m}_{r+1} - q^{-2} \mathbf{m}_{r+1}^{-1}}{q - q^{-1}} \\ &= (q + q^{-1}) \mathfrak{r}_r^2 \mathfrak{d}_{r+1}^2 \frac{q \mathbf{m}_{r+1} - q^{-1} \mathbf{m}_{r+1}^{-1}}{q - q^{-1}} = [2] \Phi_S(e_r t_{r+1} e_r), \\ &\Phi_S(t_{r+1}^2 e_r) + \Phi_S(e_r t_{r+1}^2) \\ &= (q - q^{-1})^{-2} ((\mathbf{m}_{r+1} - \mathbf{m}_{r+1}^{-1})^2 + (q^{-1} \mathbf{m}_{r+1} - q \mathbf{m}_{r+1}^{-1})^2) \mathfrak{r}_r \mathfrak{d}_{r+1} \end{aligned}$$

$$\begin{aligned}
&= (q - q^{-1})^{-2} ((1 + q^{-2})\mathfrak{m}_{r+1}^2 + (1 + q^2)\mathfrak{m}_{r+1}^{-2} - 4)\mathfrak{r}_r \mathfrak{d}_{r+1} \\
&= (q - q^{-1})^{-2} (q + q^{-1})(\mathfrak{m}_{r+1} - \mathfrak{m}_{r+1}^{-1})(q^{-1}\mathfrak{m}_{r+1} - q\mathfrak{m}_{r+1}^{-1})\mathfrak{r}_r \mathfrak{d}_{r+1} + \mathfrak{r}_r \mathfrak{d}_{r+1} \\
&= [2]\Phi_{\mathcal{S}}(t_{r+1}e_r t_{r+1}) + \Phi_{\mathcal{S}}(e_r).
\end{aligned}$$

(3) By the relation (4.5), we have

$$(4.15) \quad e_i f_j - f_j e_i = \delta_{ij} \frac{k_i - k_i^{-1}}{q - q^{-1}}, \quad \text{if } (i, j) \neq (0, 0), (r, r),$$

$$(4.16) \quad e_0^2 f_0 + f_0 e_0^2 = (q + q^{-1})(e_0 f_0 e_0 - (qk_0 + q^{-1}k_0^{-1})e_0),$$

$$(4.17) \quad e_r^2 f_r + f_r e_r^2 = (q + q^{-1})(e_r f_r e_r - e_r(qk_r + q^{-1}k_r^{-1})),$$

$$(4.18) \quad f_0^2 e_0 + e_0 f_0^2 = (q + q^{-1})(f_0 e_0 f_0 - f_0(qk_0 + q^{-1}k_0^{-1})),$$

$$(4.19) \quad f_r^2 e_r + e_r f_r^2 = (q + q^{-1})(f_r e_r f_r - (qk_r + q^{-1}k_r^{-1})f_r).$$

We provide a detailed argument for the relation (4.16). By the involution τ in Diagram III, we have $\xi_i = 2^{\delta_{i,0} + \delta_{i,r+1}}$. Then

$$\begin{aligned}
&\Phi_{\mathcal{S}}(e_0^2 f_0) + \Phi_{\mathcal{S}}(f_0 e_0^2) = \mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{r}_1 \mathfrak{d}_0 + \mathfrak{r}_1 \mathfrak{d}_0 \mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{r}_0 \mathfrak{d}_1 \\
&= (q - q^{-1})^{-2} \mathfrak{r}_0 \mathfrak{d}_1 ((\mathfrak{m}_0 - \mathfrak{m}_0^{-1})(q\mathfrak{m}_1 - q^{-1}\mathfrak{m}_1^{-1}) + (q^{-1}\mathfrak{m}_1 - q\mathfrak{m}_1^{-1})(q^4\mathfrak{m}_0 - q^{-4}\mathfrak{m}_0^{-1})) \\
&= (q - q^{-1})^{-2} \mathfrak{r}_0 \mathfrak{d}_1 ((q + q^3)\mathfrak{m}_0 \mathfrak{m}_1 - (q^{-1} + q^5)\mathfrak{m}_0 \mathfrak{m}_1^{-1} - (q + q^{-5})\mathfrak{m}_0^{-1} \mathfrak{m}_1 + (q^{-1} + q^{-3})\mathfrak{m}_0^{-1} \mathfrak{m}_1^{-1}) \\
&= (q - q^{-1})^{-2} (q + q^{-1}) \mathfrak{r}_0 \mathfrak{d}_1 (q^2 \mathfrak{m}_0 \mathfrak{m}_1 + q^{-2} \mathfrak{m}_0^{-1} \mathfrak{m}_1^{-1} \\
&\quad - (q^2 + (q - q^{-1})^2 q^2) \mathfrak{m}_0 \mathfrak{m}_1^{-1} - (q^{-2} + (q - q^{-1})^2 q^{-2}) \mathfrak{m}_0^{-1} \mathfrak{m}_1) \\
&= (q + q^{-1}) \mathfrak{r}_0 \mathfrak{d}_1 ((q - q^{-1})^{-2} (\mathfrak{m}_1 - \mathfrak{m}_1^{-1})(q^2 \mathfrak{m}_0 - q^{-2} \mathfrak{m}_0^{-1}) - (q^2 \mathfrak{m}_0 \mathfrak{m}_1^{-1} + q^{-2} \mathfrak{m}_0^{-1} \mathfrak{m}_1)) \\
&= (q + q^{-1}) (\mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{r}_1 \mathfrak{d}_1 \mathfrak{r}_0 \mathfrak{d}_1 - q^2 \mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{m}_0 \mathfrak{m}_1^{-1} - q^{-2} \mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{m}_0^{-1} \mathfrak{m}_1) \\
&= (q + q^{-1}) (\Phi_{\mathcal{S}}(e_0 f_0 e_0) - q \Phi_{\mathcal{S}}(k_0 e_0) - q^{-1} \Phi_{\mathcal{S}}(k_0^{-1} e_0)).
\end{aligned}$$

(4) It follows by the relations (4.1), (4.2) and (4.5) that

$$(4.20) \quad k_0 k_0^{-1} = 1, \quad k_0 e_0 = q^4 e_0 k_0, \quad k_0 f_0 = q^{-4} f_0 k_0,$$

$$(4.21) \quad e_0^3 f_0 - [3]e_0^2 f_0 e_0 + [3]e_0 f_0 e_0^2 - f_0 e_0^3 = [3]!(q - q^{-1})e_0(k_0 - k_0^{-1})e_0,$$

$$(4.22) \quad f_0^3 e_0 - [3]f_0^2 e_0 f_0 + [3]f_0 e_0 f_0^2 - e_0 f_0^3 = -[3]!(q - q^{-1})f_0(k_0 - k_0^{-1})f_0.$$

We need to make slight modifications for the algebra $\mathbf{A}_q(\mathcal{S})$ defined in Definition 3.1. Let $r = 0$ and $\xi_0 = 1$, $\xi_1 = 3$. We only verify the relation (4.21) and we have

$$\begin{aligned}
&\Phi_{\mathcal{S}}(e_0^3 f_0) - [3]\Phi_{\mathcal{S}}(e_0^2 f_0 e_0) + [3]\Phi_{\mathcal{S}}(e_0 f_0 e_0^2) - \Phi_{\mathcal{S}}(f_0 e_0^3) \\
&= \mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{r}_1 \mathfrak{d}_0 - [3]\mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{r}_1 \mathfrak{d}_0 \mathfrak{r}_0 \mathfrak{d}_1 + [3]\mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{r}_1 \mathfrak{d}_0 \mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{r}_0 \mathfrak{d}_1 - \mathfrak{r}_1 \mathfrak{d}_0 \mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{r}_0 \mathfrak{d}_1 \mathfrak{r}_0 \mathfrak{d}_1 \\
&= (q - q^{-1})^{-2} \mathfrak{r}_0^2 \mathfrak{d}_1^2 ((\mathfrak{m}_0 - \mathfrak{m}_0^{-1})(q^3 \mathfrak{m}_1 - q^{-3} \mathfrak{m}_1^{-1}) - [3](\mathfrak{m}_1 - \mathfrak{m}_1^{-1})(q\mathfrak{m}_0 - q^{-1}\mathfrak{m}_0^{-1}) \\
&\quad + [3](q^2 \mathfrak{m}_0 - q^{-2} \mathfrak{m}_0^{-1})(q^{-3} \mathfrak{m}_1 - q^3 \mathfrak{m}_1^{-1}) - (q^{-6} \mathfrak{m}_1 - q^6 \mathfrak{m}_1^{-1})(q^3 \mathfrak{m}_0 - q^{-3} \mathfrak{m}_0^{-1})) \\
&= (q - q^{-1})^{-2} \mathfrak{r}_0^2 \mathfrak{d}_1^2 (((q^3 - q^{-3}) + [3](q^{-1} - q))\mathfrak{m}_0 \mathfrak{m}_1 - ((q^{-3} - q^9) + [3](q^5 - q))\mathfrak{m}_0 \mathfrak{m}_1^{-1} \\
&\quad - ((q^3 - q^{-9}) + [3](q^{-5} - q^{-1}))\mathfrak{m}_1 \mathfrak{m}_0^{-1} - ((q^{-3} - q^3) + [3](q - q^{-1}))\mathfrak{m}_0^{-1} \mathfrak{m}_1^{-1}) \\
&= [3]!(q - q^{-1}) \mathfrak{r}_0^2 \mathfrak{d}_1^2 (q^3 \mathfrak{m}_0 \mathfrak{m}_1^{-1} - q^{-3} \mathfrak{m}_0^{-1} \mathfrak{m}_1) = [3]!(q - q^{-1}) \mathfrak{r}_0 \mathfrak{d}_1 (q^{-1} \mathfrak{m}_0 \mathfrak{m}_1^{-1} - q \mathfrak{m}_0^{-1} \mathfrak{m}_1) \mathfrak{r}_0 \mathfrak{d}_1 \\
&= [3]!(q - q^{-1}) \Phi_{\mathcal{S}}(e_0(k_0 - k_0^{-1})e_0).
\end{aligned}$$

The other relations in ${}^{\iota}U(\mathcal{S})$ can be verified in a similar way. \square

5. CRYSTAL BASIS OF THE OSCILLATOR REPRESENTATIONS

In this section, we study the irreducible modules of \imath quantum groups and construct the crystal basis of these irreducible modules over \imath quantum groups ${}^{\imath}U(A_{2r+1})$, ${}^{\imath}U(A_{2r+1}^{(1)})$ and ${}^{\imath}U(A_1^{(1)})$.

5.1. Irreducible modules of \imath quantum groups. By Proposition 3.1 and Theorem 4.1, we have

Corollary 5.1. *There exists a $\mathbb{Q}(q)$ -algebra homomorphism $\iota \circ \Phi_S : {}^{\imath}U(S) \rightarrow \mathbf{A}_q(A_{r+1})$.*

From Theorems 3.2 and 4.1, it's easy to see that \mathbb{P} is a ${}^{\imath}U(S)$ -module. The ${}^{\imath}U(S)$ -action is given as follows.

1) For the \imath quantum group ${}^{\imath}U(A_{2r+1})$ ($r \geq 0$), the action is given by

$$e_i \mathbf{X}^{\mathbf{a}} := [2^{\delta_{i,r}} a_{i+1}] \mathbf{X}^{\mathbf{a}+\mathbf{e}_i-\mathbf{e}_{i+1}}, \quad f_i \mathbf{X}^{\mathbf{a}} := [a_i] \mathbf{X}^{\mathbf{a}-\mathbf{e}_i+\mathbf{e}_{i+1}}, \quad k_i \mathbf{X}^{\mathbf{a}} := q^{a_i-2^{\delta_{i,r}} a_{i+1}} \mathbf{X}^{\mathbf{a}}.$$

2) For the \imath quantum group ${}^{\imath}U(A_{2r+2,1})$ ($r \geq 0$), the action is given by

$$e_i \mathbf{X}^{\mathbf{a}} := [(-1)^{\delta_{i,r}} a_{i+1}] \mathbf{X}^{\mathbf{a}+\mathbf{e}_i-\mathbf{e}_{i+1}}, \quad f_i \mathbf{X}^{\mathbf{a}} := [a_i] \mathbf{X}^{\mathbf{a}-\mathbf{e}_i+\mathbf{e}_{i+1}}, \\ k_i \mathbf{X}^{\mathbf{a}} := (-1)^{\delta_{i,r}} q^{a_i-a_{i+1}} \mathbf{X}^{\mathbf{a}}, \quad t_{r+1} \mathbf{X}^{\mathbf{a}} := -[a_{r+1}] \mathbf{X}^{\mathbf{a}}.$$

3) For the \imath quantum group ${}^{\imath}U(A_{2r+1}^{(1)})$ ($r \geq 1$), the action is given by

$$e_i \mathbf{X}^{\mathbf{a}} := [2^{\delta_{i,r}} a_{i+1}] \mathbf{X}^{\mathbf{a}+\mathbf{e}_i-\mathbf{e}_{i+1}}, \quad f_i \mathbf{X}^{\mathbf{a}} := [2^{\delta_{i,0}} a_i] \mathbf{X}^{\mathbf{a}-\mathbf{e}_i+\mathbf{e}_{i+1}}, \\ k_i \mathbf{X}^{\mathbf{a}} := q^{2^{\delta_{i,0}} a_i-2^{\delta_{i,r}} a_{i+1}-2\delta_{i,0}} \mathbf{X}^{\mathbf{a}},$$

4) For the \imath quantum group ${}^{\imath}U(A_1^{(1)})$, the action is given by

$$e_0 \mathbf{X}^{\mathbf{a}} := [3a_1] \mathbf{X}^{\mathbf{a}+\mathbf{e}_0-\mathbf{e}_1}, \quad f_0 \mathbf{X}^{\mathbf{a}} := [a_0] \mathbf{X}^{\mathbf{a}-\mathbf{e}_0+\mathbf{e}_1}, \quad k_0 \mathbf{X}^{\mathbf{a}} := q^{a_0-3a_1-1} \mathbf{X}^{\mathbf{a}}.$$

5) For the \imath quantum group ${}^{\imath}U(A_{2r+2,1}^{(1)})$ ($r \geq 0$), the action is given by

$$e_i \mathbf{X}^{\mathbf{a}} := (-1)^{\delta_{i,r}} [a_{i+1}] \mathbf{X}^{\mathbf{a}+\mathbf{e}_i-\mathbf{e}_{i+1}}, \quad f_i \mathbf{X}^{\mathbf{a}} := [2^{\delta_{i,0}} a_i] \mathbf{X}^{\mathbf{a}-\mathbf{e}_i+\mathbf{e}_{i+1}}, \\ k_i \mathbf{X}^{\mathbf{a}} := (-1)^{\delta_{i,r}} q^{2^{\delta_{i,0}} a_i-a_{i+1}-2\delta_{i,0}} \mathbf{X}^{\mathbf{a}}, \quad t_{r+1} \mathbf{X}^{\mathbf{a}} := -[a_{r+1}] \mathbf{X}^{\mathbf{a}}.$$

6) For the \imath quantum group ${}^{\imath}U(A_{1,2r+2}^{(1)})$ ($r \geq 0$), the action is given by

$$e_i \mathbf{X}^{\mathbf{a}} := [2^{\delta_{i,r+1}} a_i] \mathbf{X}^{\mathbf{a}+\mathbf{e}_i-\mathbf{e}_{i+1}}, \quad f_i \mathbf{X}^{\mathbf{a}} := [(-1)^{\delta_{i,1}} a_{i-1}] \mathbf{X}^{\mathbf{a}-\mathbf{e}_i+\mathbf{e}_{i+1}}, \\ k_i \mathbf{X}^{\mathbf{a}} := (-1)^{\delta_{i,1}} q^{a_{i-1}-2^{\delta_{i,r+1}} a_i} \mathbf{X}^{\mathbf{a}}, \quad t_0 \mathbf{X}^{\mathbf{a}} := -[a_0] \mathbf{X}^{\mathbf{a}}.$$

7) For the \imath quantum group ${}^{\imath}U(A_{2r+1,2}^{(1)})$ ($r \geq 1$), the action is given by

$$e_i \mathbf{X}^{\mathbf{a}} := (-1)^{\delta_{i,r}} [a_{i+1}] \mathbf{X}^{\mathbf{a}+\mathbf{e}_i-\mathbf{e}_{i+1}}, \quad f_i \mathbf{X}^{\mathbf{a}} := [a_i] \mathbf{X}^{\mathbf{a}-\mathbf{e}_i+\mathbf{e}_{i+1}}, \\ k_i \mathbf{X}^{\mathbf{a}} := (-1)^{\delta_{i,r}} q^{a_i-a_{i+1}} \mathbf{X}^{\mathbf{a}}, \quad t_0 \mathbf{X}^{\mathbf{a}} := [a_1] \mathbf{X}^{\mathbf{a}}, \quad t_{r+1} \mathbf{X}^{\mathbf{a}} := -[a_{r+1}] \mathbf{X}^{\mathbf{a}}.$$

The ${}^{\imath}U(S)$ -module \mathbb{P} is called *the Oscillator representation of ${}^{\imath}U(S)$* . The following diagram explains the relations between ${}^{\imath}U(S)$, $\mathbf{A}_q(S)$, $\mathbf{A}_q(A_{r+1})$ and \mathbb{P} .

$$\begin{array}{ccccc} {}^{\imath}U(S) & \xrightarrow{\Phi_S} & \mathbf{A}_q(S) & \xrightarrow{\iota} & \mathbf{A}_q(A_{r+1}) \\ & \searrow & \downarrow & \swarrow & \\ & & \text{End}(\mathbb{P}) & & \end{array}$$

For $s \geq 0$, let $\Lambda_s = \{\mathbf{a} \in \mathbb{Z}_{\geq 0}^{r+2} \mid \sum_{i=0}^{r+1} a_i = s\}$. The polynomial ring \mathbb{P} has the following direct sum decomposition

$$\mathbb{P} = \bigoplus_{s=0}^{\infty} \mathbb{P}_s, \quad \mathbb{P}_s = \bigoplus_{\mathbf{a} \in \Lambda_s} \mathbb{K}\mathbf{X}^{\mathbf{a}}.$$

Since the actions of e_i, f_i, k_i and t_j on \mathbb{P} preserve the degree of arbitrary monomial in \mathbb{P}_s , the subspace \mathbb{P}_s is a ${}^{\iota}U(\mathcal{S})$ -module.

Theorem 5.1. *The ${}^{\iota}U(\mathcal{S})$ -module \mathbb{P}_s is irreducible if q is not a root of unity.*

Proof. Let \mathbb{P}'_s be a non-zero submodule of \mathbb{P}_s . We fix a non-zero element $\mathbf{X}^{\mathbf{a}} \in \mathbb{P}'_s$. It follows that

$$\left(\prod_{i=0}^r e_i^{\sum_{j=i}^r a_{j+1}} \right) \mathbf{X}^{\mathbf{a}} = \left(\prod_{i=0}^r (\mathfrak{r}_i \mathfrak{d}_{i+1})^{\sum_{j=i}^r a_{j+1}} \right) \mathbf{X}^{\mathbf{a}} = \left(\prod_{i=1}^{r+1} [a_i + a_{i+1} + \cdots + a_{r+1}]^{\xi_i!} \right) X_0^s \in \mathbb{P}'_s.$$

Since the coefficient is not zero, we get $X_0^s \in \mathbb{P}'_s$. Let $\mathbf{X}^{\mathbf{b}}$ be any monomial in \mathbb{P} , where $\mathbf{b} = (b_i)_{0 \leq i \leq r+1}$. It follows that

$$\left(\prod_{i=0}^r f_i^{s - \sum_{j=0}^i b_j} \right) X_0^s = \left(\prod_{i=0}^r (\mathfrak{r}_{i+1} \mathfrak{d}_i)^{s - \sum_{j=0}^i b_j} \right) X_0^s = \left(\frac{[s]^{\xi_0!}}{[b_0]^{\xi_0!}} \prod_{i=1}^r \frac{[s - (\sum_{j=0}^{i-1} b_j)]^{\xi_i!}}{[b_i]^{\xi_i!}} \right) \mathbf{X}^{\mathbf{b}} \in \mathbb{P}'_s.$$

The relation $\mathbb{P}_s = \mathbb{P}'_s$ holds automatically due to the non-zero coefficient of $\mathbf{X}^{\mathbf{b}}$. Hence the ${}^{\iota}U(\mathcal{S})$ -module \mathbb{P}_s is irreducible. \square

5.2. Crystal basis of irreducible module. In this subsection, we only consider the ι quantum groups ${}^{\iota}U(A_{2r+1})$, ${}^{\iota}U(A_{2r+1}^{(1)})$ and ${}^{\iota}U(A_1^{(1)})$.

Let $\mathbf{X}^{(\mathbf{a})} = \prod_{i=0}^{r+1} X_i^{(a_i)\xi_i}$ for a sequence $\mathbf{a} \in \Lambda_s$. It's easy to verify:

$$\mathbf{X}^{(\mathbf{a})} = f_i^{(a_{i+1})\xi_{i+1}} \mathbf{X}^{(\mathbf{a} + a_{i+1}(\mathbf{e}_i - \mathbf{e}_{i+1}))}.$$

Let $\mathbb{P}_s = \bigoplus_{\mathbf{a} \in \Lambda_s} \mathbb{P}_s^{\mathbf{a}}$, where $\mathbb{P}_s^{\mathbf{a}} = \mathbb{Q}(q)\mathbf{X}^{(\mathbf{a})}$. We define Kashiwara operators \tilde{e}_i and \tilde{f}_i ($0 \leq i \leq r$) on \mathbb{P}_s as follows:

$$(5.1) \quad \tilde{e}_i \mathbf{X}^{(\mathbf{a})} = f_i^{(a_{i+1}-1)\xi_{i+1}} \mathbf{X}^{(\mathbf{a} + a_{i+1}(\mathbf{e}_i - \mathbf{e}_{i+1}))}, \quad \tilde{f}_i \mathbf{X}^{(\mathbf{a})} = f_i^{(a_{i+1}+1)\xi_{i+1}} \mathbf{X}^{(\mathbf{a} + a_{i+1}(\mathbf{e}_i - \mathbf{e}_{i+1}))}.$$

Let $\mathbb{A}_0 = \{f/g \in \mathbb{Q}(q) \mid f, g \in \mathbb{Q}[q], g(0) \neq 0\}$. The crystal lattice and crystal basis of the irreducible module \mathbb{P}_s are defined as follows.

Definition 5.2. *Let $\mathcal{L}(s)$ be an \mathbb{A}_0 -submodule of \mathbb{P}_s . We say that $\mathcal{L}(s)$ is a crystal lattice of \mathbb{P}_s if*

- (L1) $\mathcal{L}(s)$ is a free \mathbb{A}_0 -module of rank $\dim_{\mathbb{Q}(q)} \mathbb{P}_s$, and $\mathbb{Q}(q) \otimes_{\mathbb{A}_0} \mathcal{L}(s) = \mathbb{P}_s$,
- (L2) $\mathcal{L}(s) = \bigoplus_{\mathbf{a} \in \Lambda_s} \mathcal{L}(s)_{\mathbf{a}}$, where $\mathcal{L}(s)_{\mathbf{a}} := \mathcal{L}(s) \cap \mathbb{P}_s^{\mathbf{a}}$,
- (L3) $\tilde{f}_i(\mathcal{L}(s)) \subset \mathcal{L}(s)$ and $\tilde{e}_i(\mathcal{L}(s)) \subset \mathcal{L}(s)$.

Definition 5.3. A crystal basis of the module \mathbb{P}_s is a pair $(\mathcal{L}(s), \mathcal{B}(s))$ such that

- (B1) $\mathcal{L}(s)$ is a crystal lattice of \mathbb{P}_s ,
- (B2) $\mathcal{B}(s)$ is a \mathbb{Q} -basis of $\mathcal{L}(s)/q\mathcal{L}(s)$,
- (B3) $\mathcal{B}(s) = \bigsqcup_{\lambda \in \Lambda_s} \mathcal{B}(s)_{\lambda}$, where $\mathcal{B}(s)_{\lambda} := \mathcal{B}(s) \cap (\mathcal{L}(s)_{\lambda}/q\mathcal{L}(s)_{\lambda})$,
- (B4) $\tilde{f}_i(\mathcal{B}(s)) \subset \mathcal{B}(s) \sqcup \{0\}$ and $\tilde{e}_i(\mathcal{B}(s)) \subset \mathcal{B}(s) \sqcup \{0\}$,
- (B5) for each $b, b' \in \mathcal{B}(s)$, one has $\tilde{f}_i(b) = b'$ if and only if $b = \tilde{e}_i(b')$.

From Theorem 5.1, \mathbb{P}_s is an irreducible ${}^iU(\mathcal{S})$ -module with $\mathbb{Q}(q)$ -basis $\{\mathbf{X}^{(\mathbf{a})} | \mathbf{a} \in \Lambda_s\}$. We define

$$(5.2) \quad \mathcal{L}^*(s) = \bigoplus_{\mathbf{a} \in \Lambda_s} \mathcal{L}^*(s)_{\mathbf{a}}, \quad \mathcal{B}^*(s) = \{\mathbf{X}^{(\mathbf{a})} + q\mathcal{L}^*(s) | \mathbf{a} \in \Lambda_s\},$$

where $\mathcal{L}^*(s)_{\mathbf{a}} = \mathbb{A}_0\mathbf{X}^{(\mathbf{a})}$.

Theorem 5.2. *The pair $(\mathcal{L}^*(s), \mathcal{B}^*(s))$ is a crystal basis of \mathbb{P}_s .*


Proof. It's straightforward to verify that the conditions (L1), (L2) and (L3) hold in Definition 5.2 due to the definition of Kashiwara operators in (5.1). Hence $\mathcal{L}^*(s)$ is a crystal lattice of \mathbb{P}_s . From the definition of $\mathcal{B}^*(s)$ in (5.2), it's easy to see $\mathcal{B}^*(s)$ is a \mathbb{Q} -basis of $\mathcal{L}^*(s)/q\mathcal{L}^*(s) \cong \mathbb{Q} \otimes_{\mathbb{A}_0} \mathcal{L}^*(s)$. The conditions (B3) and (B4) are easy to verify. Now let's verify (B5):

Let $b = \mathbf{X}^{(\mathbf{a})}$ and $b' = f_i^{(a_{i+1}+1)\xi_{i+1}} \mathbf{X}^{(\mathbf{a}+a_{i+1}(\mathbf{e}_i-\mathbf{e}_{i+1}))}$. Then we have $\tilde{f}_i b = b'$.

$$\begin{aligned} \tilde{e}_i b' &= \tilde{e}_i f_i^{(a_{i+1}+1)\xi_{i+1}} \mathbf{X}^{(\mathbf{a}+a_{i+1}(\mathbf{e}_i-\mathbf{e}_{i+1}))} = \tilde{e}_i \frac{f_i}{[\xi_{i+1}(a_{i+1}+1)]} f_i^{(a_{i+1})\xi_{i+1}} \mathbf{X}^{(\mathbf{a}+a_{i+1}(\mathbf{e}_i-\mathbf{e}_{i+1}))} \\ &= \tilde{e}_i \frac{\mathbf{x}_{i+1} \mathbf{d}_i}{[\xi_{i+1}(a_{i+1}+1)]} \mathbf{X}^{(\mathbf{a})} = \tilde{e}_i \mathbf{X}^{(\mathbf{a}-\mathbf{e}_i+\mathbf{e}_{i+1})} = f_i^{(a_{i+1})\xi_{i+1}} \mathbf{X}^{(\mathbf{a}-\mathbf{e}_i+\mathbf{e}_{i+1}+(a_{i+1}+1)(\mathbf{e}_i-\mathbf{e}_{i+1}))} \\ &= \frac{f_i^{a_{i+1}}}{[a_{i+1}]^{\xi_{i+1}!}} \mathbf{X}^{(\mathbf{a}+a_{i+1}(\mathbf{e}_i-\mathbf{e}_{i+1}))} = \mathbf{X}^{(\mathbf{a})} = b. \end{aligned}$$

Hence $(\mathcal{L}^*(s), \mathcal{B}^*(s))$ is a crystal basis of \mathbb{P}_s . \square

Example 5.3. *Let $r = 1$ and $s = 3$, we use $(a_0 a_1 a_2)$ to represent $X_0^{(a_0)\xi_0} X_1^{(a_1)\xi_1} X_2^{(a_2)\xi_2} + q\mathcal{L}^*(3)$. We use the red arrow to represent the action of \tilde{f}_0 and the blue arrow to represent the action of \tilde{f}_1 . The crystal graph is described as follows:*



REFERENCES

- [1] H. Bao, J. Kujawa, Y. Li and W. Wang, *Geometric Schur duality of classical type*, Transform. Groups **23** (2018), 329–389.

- [2] H. Bao, P. Shan, W. Wang and B. Webster, *Categorification of quantum symmetric pairs I*, Quantum Topology **9** (2018), 643–714.
- [3] H. Bao and W. Wang, *A new approach to Kazhdan-Lusztig theory of type B via quantum symmetric pairs*, Astérisque **402** (2018), vii+134pp.
- [4] A. Beilinson, G. Lusztig and R. McPherson, *A geometric setting for the quantum deformation of GL_n* , Duke Math. J. **61** (1990), 655–677.
- [5] T. Bridgeland, *Quantum groups via Hall algebras of complexes*, Ann. Math. **177** (2013), 739–759.
- [6] X. Chen, M. Lu and W. Wang, *A Serre presentation for the ι -quantum groups*, Transform. Groups **26** (2021), 827–857.
- [7] V. Drinfeld, *Quantum groups*, Proc. Int. Congr. Math. Berkeley **1** (1987), 798–820.
- [8] J. Du, and Q. Fu, *Quantum affine \mathfrak{gl}_n via Hecke algebras*, Adv. Math. **282** (2015), 23–46.
- [9] J. Du and Y. Wu, *A new realisation of the ι -quantum groups $\mathbf{U}^j(n)$* , J. Pure Appl. Algebra **226** (2022), 106793.
- [10] J. Du and Y. Wu, *The ι -quantum group $\mathbf{U}^i(n)$* , arXiv:2111.08898.
- [11] Z. Fan, C. Lai, Y. Li, L. Luo and W. Wang, *Affine flag varieties and quantum symmetric pairs*, Mem. Amer. Math. Soc. **265** (2020), no. 1285, v+123pp.
- [12] Z. Fan, C. Lai, Y. Li, L. Luo and W. Wang, *Affine Hecke algebras and quantum symmetric pairs*, Mem. Amer. Math. Soc. (to appear), arXiv:1609.06199v2.
- [13] Z. Fan, C. Lai, Y. Li, L. Luo, W. Wang and H. Watanabe, *Quantum Schur duality of affine type C with three parameters*, Math. Res. Lett. **27** (2020), 79–114.
- [14] Z. Fan, H. Ma and H. Xiao, *Equivariant K-theory approach to ι -quantum groups*, Publ. Res. Inst. Math. Sci.(to appear), 2020, arXiv:1911.00851.
- [15] V. Ginzburg and E. Vasserot, *Langlands reciprocity for affine quantum groups of type A_n* , Internat. Math. Res. Notices **3** (1993), 67–85.
- [16] T. Hayashi, *q -Analogues of Clifford and Weyl algebras—spinor and oscillator representations of quantum enveloping algebras*, Commun. Math. Phys. **127** (1990), 129–144.
- [17] J. Hong and S.-J. Kang, *Introduction to Quantum Groups and Crystal Bases*, Grad. Stud. Math. **42**, Amer. Math. Soc. 2002.
- [18] M. Jimbo, *A q -difference analogue of $\mathbf{U}(\mathfrak{g})$, Hecke algebra and the Yang Baxter equation*, Lett. Math. **11** (1986), 247–252.
- [19] M. Kashiwara, *On crystal bases of the q -analogue of the universal enveloping algebra*, Duke Math. J. **63** (1991), 465–516.
- [20] M. Lu and W. Wang, *Hall algebras and quantum symmetric pairs I: foundations*, Proc. Lond. Math. Soc. **124** (2022), 1–82.
- [21] M. Lu and W. Wang, *Hall algebras and quantum symmetric pairs II: reflection functors*, Commun. Math. Phys. **381** (2021), 799–855.
- [22] M. Lu and W. Wang, *Hall algebras and quantum symmetric pairs III: quiver varieties*, Adv. Math. **393** (2021), 108071.
- [23] M. Lu and W. Wang, *Hall algebras and quantum symmetric pairs of Kac-Moody type*, arXiv:2006.06904.
- [24] G. Lusztig, *Introduction to quantum groups*, Modern Birkhäuser Classics, Reprint of the 1993 Edition, Birkhäuser, Boston, 2010.
- [25] C.M. Ringel, *Hall algebras and quantum groups*, Invent. Math. **101** (1990), 583–591.
- [26] H. Watanabe, *Crystal basis theory for a quantum symmetric pair $(\mathbf{U}, \mathbf{U}^j)$* , Int. Math. Res. Not. **22** (2020), 8292–8352.
- [27] H. Watanabe, *Global crystal bases for integrable modules over a quantum symmetric pair of type AIII*, Represent. Theory **25** (2021), 27–66.

HARBIN ENGINEERING UNIVERSITY, HARBIN, CHINA
 Email address: fanzhaobing@hrbeu.edu.cn

HARBIN ENGINEERING UNIVERSITY, HARBIN, CHINA
 Email address: jcgeng@hrbeu.edu.cn

HARBIN ENGINEERING UNIVERSITY, HARBIN, CHINA
 Email address: algebra@hrbeu.edu.cn