# TETRAVALENT VERTEX-TRANSITIVE GRAPHS OF ORDER 6p

### MAJID AREZOOMAND, MOHSEN GHASEMI\* AND MOHAMMAD A. IRANMANESH

ABSTRACT. A graph is vertex-transitive if its automorphism group acts transitively on vertices of the graph. A vertex-transitive graph is a Cayley graph if its automorphism group contains a subgroup acting regularly on its vertices. In this paper, the tetravalent vertex-transitive non-Cayley graphs of order 6p are classified for each prime p.

## 1. INTRODUCTION

In this paper we consider undirected finite connected graphs without loops or multiple edges. For a graph X we use V(X), E(X), A(X) and Aut(X) to denote its vertex set, edge set, arc set and its full automorphism group, respectively. For  $u, v \in V(X), u \sim v$ represents that u is adjacent to v, and is denoted by  $\{u, v\}$  the edge incident to u and v in X, and  $N_X(u)$  is the neighborhood of u in X, that is, the set of vertices adjacent to u in X. A graph X is said to be G-vertex-transitive, G-edge-transitive and G-arctransitive (or G-symmetric) if  $G \leq \operatorname{Aut}(X)$  acts transitively on V(X), E(X) and A(X), respectively. In the special case, if  $G = \operatorname{Aut}(X)$  then X is said to be vertex-transitive. edge-transitive and arc-transitive (or symmetric). An s-arc in a graph X is an ordered (s+1)-tuple  $(v_0, v_1, \dots, v_s)$  of vertices of X such that  $v_{i-1}$  is adjacent to  $v_i$  for  $1 \le i \le s$ , and  $v_{i-1} \neq v_{i+1}$  for  $1 \leq i \leq s$ ; in other words, a directed walk of length s which never includes a backtracking. A graph X is said to be s-arc-transitive if Aut(X) is transitive on the set of s-arcs in X. A subgroup of the automorphism group of a graph X is said to be s-regular if it acts regularly on the set of s-arcs of X. Recall that a permutation group G acting on a set  $\Omega$  is called semiregular if the stabilizer of  $\alpha \in G$ ,  $G_{\alpha} = 1$  for all  $\alpha \in G$ and is called regular if it is semiregular and transitive.

Let G be a finite group and S be a subset of G such that  $1 \notin S$  and  $S = S^{-1}$  where  $S^{-1} = \{s^{-1} \mid s \in S\}$ . The Cayley graph  $X = \operatorname{Cay}(G, S)$  on G with respect to S is defined as the graph with vertex set V(X) = G and edge set  $E(X) = \{\{g, sg\} \mid g \in G, s \in S\}$ . The automorphism group  $\operatorname{Aut}(X)$  of X contains the right regular representation R(G) of G, the acting group of G by right multiplication, as a subgroup, and R(G) is regular on V(X). A Cayley graph,  $\operatorname{Cay}(G, S)$  is said to be normal if R(G) is normal in  $\operatorname{Aut}(\operatorname{Cay}(G, S))$ . By [36, Proposition 3.1],  $N_{\operatorname{Aut}(\operatorname{Cay}(G,S))}(R(G)) = R(G)\operatorname{Aut}(G, S)$ , where  $\operatorname{Aut}(G, S) = \{\alpha \in C, S\}$ .

MSC(2010): Primary: 20B20, Secondary: 05C25.

Keywords: Tetravalent graph, vertex transitive graph, non-Cayley graph.

<sup>\*</sup>Corresponding author.

Aut(G) |  $S^{\alpha} = S$ }. Note that Aut(X) is normal if and only if Aut(G, S) = Aut(X)\_v for some  $v \in V(X)$ .

It is well known that a vertex-transitive graph is a Cayley graph if and only if its automorphism group contains a subgroup acting regularly on its vertex set (see, for example, [32, Lemma 4]). There are vertex-transitive graphs which are not Cayley graphs and the smallest one is the well-known Petersen graph. Such a graph will be called a *vertex-transitive non-Cayley graph*, or a VNC-graph for short.

Many publications have been put into service of investigation the  $\mathcal{VNC}$ -graphs from different perspectives. For example, in [19], Marušič asked for a determination of the set  $\mathcal{NC}$  of non-Cayley numbers, that is, those numbers n for which there exists a  $\mathcal{VNC}$ -graph of order n, and to settle this question, a lot of  $\mathcal{VNC}$ -graphs were constructed in [12, 14, 15, 20, 21, 22, 23, 24, 25, 27, 30]. In [8], Feng considered the question to determine the smallest valency for  $\mathcal{VNC}$ -graphs of a given order and it was solved for the graphs of odd prime power order. From [6, 18] all cubic or tetravalent  $\mathcal{VNC}$ -graphs of order 2p are known for each prime p. In [38, 39, 40], all cubic  $\mathcal{VNC}$ -graphs of order 2pq were classified for any primes p and q. Also tetravalent  $\mathcal{VNC}$ -graphs of order  $2p^2$  and 4p were classified for any prime p (see [5, 37]). We know that all components of vertex-transitive graphs are isomorphic. So without loss of generality we may assume that X is connected.

A graph X is said to be a *bi-Cayley graph* over a group  $G \leq Aut(X)$  if it admits G as a semiregular automorphism group with two orbits of equal size. Note that every bi-Cayley graph admits the following concrete realization. Let R, L and S be subsets of a group G such that  $R = R^{-1}$ ,  $L = L^{-1}$  and  $R \cup L$  does not contain the identity element of G. Let  $g_0 = (g, 0)$  and  $g_1 = (g, 1)$  where  $g \in G$ . Define the graph BiCay(G, R, L, S)to have vertex set the union of the right part  $G_0 = \{g_0 \mid g \in G\}$  and the left part  $G_1 = \{g_1 \mid g \in G\}$  and edge set the union of the right edges  $\{\{h_0, g_0\} \mid gh^{-1} \in R\}$ , the left edges  $\{\{h_1, g_1\} \mid gh^{-1} \in L\}$  and the spokes  $\{\{h_0, g_1\} \mid gh^{-1} \in S\}$ . Now we introduce the coset graph constructed from a finite group G relative to a subgroup Hof G and a union D of some double cosets of H in G such that  $D^{-1} = D$ . The coset graph  $\operatorname{Cos}(G, H, D)$  of G with respect to H and D is defined to have vertex set [G: H], the set of right cosets of H in G, and edge set  $\{\{Hg, Hdg\} \mid g \in G, d \in D\}$ . The graph  $\cos(G, H, D)$  has valency |D|/|H| and is connected if and only if D generates the group G. The action of G on  $V(\cos(G, H, D))$  by right multiplication induces a vertex-transitive automorphism group, which is arc-transitive if and only if D is a single double coset. Clearly,  $\operatorname{Cos}(G, H, D) \cong \operatorname{Cos}(G, H^{\alpha}, D^{\alpha})$  for every  $\alpha \in \operatorname{Aut}(G)$ . Conversely, let X be a graph and let A be a vertex-transitive subgroup of Aut(X). By [31], the graph X is isomorphic to a coset graph Cos(A, H, D), where  $H = A_u$  is the stabilizer of  $u \in V(X)$  in

A and D consists of all elements of A which map u to one of its neighbors. It is easy to show that  $\operatorname{Core}_A(H) = 1$  and that D is a union of some double cosets of H in A satisfying  $D = D^{-1}$ . Recall that a graph  $\Gamma = (V, E)$  is called an (m, n)-metacirculant, where m, nare positive integers, if  $\Gamma$  is of order |V| = mn and has two automorphisms  $\rho, \sigma$  such that

- (a)  $\langle \rho \rangle$  is semiregular and has m orbits on V,
- (b)  $\sigma$  cyclically permutes the *m* orbits of  $\langle \rho \rangle$  and normalizes  $\langle \rho \rangle$ , and
- (c)  $\sigma^m$  fixes at least one vertex of  $\Gamma$ .

A graph  $\widetilde{X}$  is called a *covering* of a graph X with projection  $\wp : \widetilde{X} \to X$  if there is a surjection  $\wp : V(\widetilde{X}) \to V(X)$  such that  $\wp|_{N_{\widetilde{X}}(\widetilde{v})} : N_{\widetilde{X}}(\widetilde{v}) \to N_X(v)$  is a bijection for any vertex  $v \in V(X)$  and  $\widetilde{v} \in \wp^{-1}(v)$ . A covering  $\widetilde{X}$  of X with a projection  $\wp$  is said to be *regular* (or K-covering) if there is a semiregular subgroup K of the automorphism group Aut( $\widetilde{X}$ ) such that the graph X is isomorphic to the quotient graph  $\widetilde{X}/K$ , say by  $\Im$ , and the quotient map  $\widetilde{X} \to \widetilde{X}/K$  is the composition  $\wp \Im$  of  $\wp$  and  $\Im$  (for the purpose of this paper, all functions are composed from left to right). If K is cyclic or elementary abelian then  $\widetilde{X}$  is called a *cyclic* or an *elementary abelian covering* of X, and if  $\widetilde{X}$  is connected K becomes the covering transformation group. The *fibre* of an edge or a vertex is its preimage under  $\wp$ . An automorphism of  $\widetilde{X}$  is said to be *fibre-preserving* if it maps a fibre to a fibre, while every covering transformation maps a fibre on to itself. All of fibre-preserving automorphisms form a group called the *fibre-preserving group*.

Let  $\tilde{X}$  be a *K*-covering of *X* with a projection  $\wp$ . If  $\alpha \in \operatorname{Aut}(X)$  and  $\tilde{\alpha} \in \operatorname{Aut}(\tilde{X})$ satisfy  $\tilde{\alpha}\wp = \wp \alpha$ , we call  $\tilde{\alpha}$  a *lift* of  $\alpha$ , and  $\alpha$  the *projection* of  $\tilde{\alpha}$ . Concepts such as a lift of a subgroup of  $\operatorname{Aut}(X)$  and the projection of a subgroup of  $\operatorname{Aut}(\tilde{X})$  are self-explanatory. The lifts and the projections of such subgroups are of course subgroups in  $\operatorname{Aut}(\tilde{X})$  and  $\operatorname{Aut}(X)$ , respectively.

Let  $m_1, m_2 > 1$  be two odd integers such that  $(m_1, m_2) = 1$ . Let  $1 \leq t \leq m_2$ ,  $(t, m_2) = 1$  be such that  $t^2 \equiv -1 \pmod{m_2}$ . Let  $H = \langle r \rangle \times \langle s \rangle \cong \mathbb{Z}_{m_1} \times \mathbb{Z}_{m_2} = \mathbb{Z}_{m_1 m_2}$ . Set  $R = \{r, r^{-1}\}, L = \{r^t, r^{-t}\}$  and  $S = \{1, s\}$  and  $X_{m_1, m_2, t} = \operatorname{BiCay}(H, R, L, S)$ . Then we have the following result.

**Proposition 1.** [29, Theorem 4.3] Let X be a connected tetravalent vertex-transitive bi-Cayley graph over a cyclic group of odd order n. Then X is a  $\mathcal{VNC}$ -graph if and only if  $X \cong X_{m_1,m_2,t}$  for some integers  $m_1, m_2$  and t.

The following theorem is the main result of this paper.

**Theorem 2.** Let X be a connected tetravalent vertex-transitive graph of order 6p, where p is a prime. Then X is a  $\mathcal{VNC}$ -graph if and only if  $X \cong X_{3,p,t}$ , where  $1 \le t \le p-1$  and  $t^2 \equiv -1 \pmod{p}$  or X is one of the nine specified graphs.

## 2. Preliminaries

In this section, we introduce some notation and definitions as well as some preliminary results which will be used later in the paper.

For a regular graph X, use d(X) to represent the valency of X, and for any subset B of V(X), the subgraph of X induced by B will be denoted by X[B]. Let X be a connected vertex-transitive graph, and let  $G \leq \operatorname{Aut}(X)$  be vertex-transitive on X. For a G-invariant partition  $\Omega$  of V(X), the quotient graph  $X_{\Omega}$  is defined as the graph with vertex set  $\Omega$  such that, for any two vertices  $B, C \in \Omega$ , B is adjacent to C if and only if there exist  $u \in B$  and  $v \in C$  which are adjacent in X. Let N be a normal subgroup of G. Then the set  $\Omega$  of orbits of N in V(X) is a G-invariant partition of V(X). In this case, the symbol  $X_{\Omega}$  will be replaced by  $X_N$ .

For two groups M and N, by  $N \rtimes M$  we denote the semidirect product of N by M. For a group G, the largest solvable normal subgroup of G is called the *solvable radical* of G and is denoted by Sol(G). The outer automorphism group of the group G, is the quotient, Aut(G)/Inn(G), where Aut(G) is the automorphism group of G and Inn(G) is the subgroup consisting of inner automorphisms. The outer automorphism group is usually denoted by Out(G). Also a lexicographic product of two graphs X and Y which is denoted by X[Y] is defined as the graph with vertex set  $V(X) \times V(Y)$  such that for any two vertices  $u = (x_1, y_1)$  and  $v = (x_2, y_2)$  in X[Y], u is adjacent to v in X[Y], whenever either  $\{x_1, x_2\} \in E(X)$  or  $x_1 = x_2$  and  $\{y_1, y_2\} \in E(Y)$ . Note that the lexicographic product of two Cayley graphs is a Cayley graph. For group and graph-theoretic terms not defined here we refer the reader to [4] and [35], respectively. Now we state the following well known result.

**Proposition 3.** Let X be a G-vertex-transitive graph. Then X is symmetric if and only if each vertex stabilizer  $G_v$  acts transitively on the set of vertices adjacent to v, where  $v \in V(X)$ .

A finite simple group G is said to be a  $K_3$ -group if its order has exactly three distinct prime divisors. By [11, Pages 12-14], G is isomorphic to one of the following groups:

(1) 
$$A_5, A_6, PSL_2(7), PSL_2(8), PSL_2(17), PSL_3(3), U_3(3), U_4(2).$$

The *socle* of a group G is the subgroup generated by the set of all minimal normal subgroups of G, it is denoted by soc(G). Also a group G is said to be *almost simple* if  $T \leq G \leq \operatorname{Aut}(T)$ , where T is a non-abelian simple group. It is well known that G is an almost simple group if and only if  $\operatorname{soc}(G) = T$  for some non-abelian simple group T.

Now we prove the following lemma.

**Lemma 4.** Let G be an almost simple group and soc(G) = A, where A is a non-abelian simple  $K_3$ -group. Then

- (i) If  $A \cong A_5$  then  $G \cong A_5$  or  $S_5$ .
- (*ii*) If  $A \cong A_6$  then  $G \cong A_6$  or  $S_6$  or  $S_6 \rtimes \mathbb{Z}_2$ .
- (iii) If  $A \cong PSL_2(p)$ , where  $p \in \{7, 17\}$ , then  $G \cong PSL_2(p)$  or  $PGL_2(p)$ .
- (iv) If  $A \cong PSL_2(8)$  then  $G \cong PSL_2(8)$  or  $G/PSL_2(8) \cong \mathbb{Z}_3$ .
- (v) If  $A \cong PSL_3(3)$  then  $G \cong PSL_3(3)$  or  $G/PSL_3(3) \cong \mathbb{Z}_2$ .
- (vi) If  $A \cong U_3(3)$  then  $G \cong U_3(3)$  or  $G/U_3(3) \cong \mathbb{Z}_2$ .
- (vii) If  $A \cong U_4(2)$  then  $G \cong U_4(2)$  or  $G/U_4(2) \cong \mathbb{Z}_2$ .

Proof. Since G is an almost simple group and  $\operatorname{soc}(G) = A$ , it implies that  $A \leq G \leq \operatorname{Aut}(A)$ . If  $A \cong A_5$ , then  $A_5 \leq G \leq \operatorname{Aut}(A_5)$ . Now since  $|\operatorname{Out}(A_5)| = 2$ , it follows that  $G \cong A_5$ or  $S_5$  and (i) holds. Also if A is isomorphic to one of  $\operatorname{PSL}_2(7)$ ,  $\operatorname{PSL}_2(17)$ ,  $\operatorname{PSL}_3(3)$ ,  $\operatorname{U}_3(3)$ or  $\operatorname{U}_4(2)$  then  $|\operatorname{Out}(A)| = 2$  and the assertions in (iii), (v), (vi) and (vii) hold. If  $A \cong A_6$ then  $|\operatorname{Out}(A_6)| = 4$  and  $\operatorname{Aut}(A_6) \cong S_6 \rtimes \mathbb{Z}_2$ . Thus  $G \cong A_6$  or  $S_6$  or  $S_6 \rtimes \mathbb{Z}_2$  and (ii) holds. Finally if  $A \cong \operatorname{PSL}_2(8)$  then  $|\operatorname{Out}(\operatorname{PSL}_2(8))| = 3$ . Hence  $G \cong \operatorname{PSL}_2(8)$  or  $G/\operatorname{PSL}_2(8) \cong \mathbb{Z}_3$ and (iv) holds.

### 3. Main Results

In this section we classify all connected tetravalent  $\mathcal{VNC}$ -graphs of order 6p where p is a prime. To do this we prove the following propositions.

**Proposition 5.** Let X be a graph,  $A = \operatorname{Aut}(X)$  and N be the normal subgroup of A. Also let  $\Omega$  be the set of orbits of N on V(X) and  $X_N$  be a Cayley graph on subgroup of A/K, where K is the kernel of action N on  $\Omega$ . Then X is a Cayley graph on some subgroup of A.

Proof. Suppose that  $X_N = \operatorname{Cay}(H, S)$ , where  $H \leq A/K$ . Thus we may suppose that H = T/K, where  $T \leq A$ . Let x, y be two arbitrary elements of V(X). Set  $X = x^N$  and  $Y = y^N$ . If X = Y then there is  $\alpha \in N$  so that  $x^{\alpha} = y$ . Thus we may suppose that  $X \neq Y$ . By our assumption there is an element  $K\beta \in T/K$  such that  $(x^N)^{K\beta} = (y^N)$  and so  $(x^N)^{\beta} = (y^N)$ . Now  $x^{\beta} = y^n$  for some  $n \in N$ . Now  $x^{(\beta n^{-1})} = y$ , where  $\beta n^{-1} \in T$ . Thus T acts transitively on V(X). Suppose that  $x^t = x$ , where  $t \in T$ . Now  $(x^N)^{Kt} = (x^N)$ , a contradiction. Thus T acts regularly on V(X) and X is a Cayley graph on T.

**Proposition 6.** Let X be a connected tetravalent vertex-transitive graph of order 3p, where p is a prime. Then X is a VNC-graph if and only if X is isomorphic to  $L(O_3)$ , the line graph of the Petersen graph.

Proof. Suppose that X is a connected tetravalent  $\mathcal{VNC}$ -graph of order 3p. By [26], all tetravalent vertex-transitive graphs of order 6, 9 and 21 are Cayley graphs. Also a tetravalent vertex-transitive graph of order 15 is not a Cayley graph and is isomorphic to  $L(O_3)$ . Thus we may assume that  $p \geq 11$ . If X is symmetric then by [34, Page 215], is isomorphic to one of  $L_3(2)_{21}^4$  or G(3p, 2), which are Cayley graphs. Thus in the following we may assume that X is not symmetric. Let  $A = \operatorname{Aut}(X)$  and  $v \in V(X)$ . Since  $A_v$  is a  $\{2, 3\}$ -group we have  $|A| = 2^s 3^t p$ , for some non-negative integers s, t. We claim that A is not solvable.

Suppose by contrary that A is solvable and let N be a minimal normal subgroup of A. Then N is an elementary abelian r-group, where  $r \in \{2,3,p\}, N = T^k$ , and T is a simple group. Let  $\Omega$  be the set of orbits of N on V(X) and K be the kernel of the action A on the set of orbits of N. By considering the order of graph, we see that either r=3 or r=p. Assume first that N is an elementary abelian 3-group. Then  $X_N$  has order p and has valency 2 or 4. It is easy to see that K acts faithfully on each orbits and therefore  $K \leq S_3$ , which implies that |N| = 3. Let P be a Sylow p-subgroup of A. Then |P| = p and since N is a normal subgroup of A, it follows that the orbits of N form a complete block system of A. Moreover  $P \times N$  acts regularly on V(X) which implies that X is a Cayley graph, a contradiction. Next assume that N is an elementary abelian *p*-group. Then  $X_N$  has order 3 and its valency is 2. Hence  $\operatorname{Aut}(X_N) \cong D_6$ . Suppose that  $\Omega = \{\Delta_0, \Delta_1, \Delta_2\}$  and  $\Delta_i \sim \Delta_{i+1}$  where the subscripts are taken modulo 3. If  $\Delta_i$  has an edge, then  $X[\Delta_i] \cong C_p$  and it is easy to see that  $|K_v| \leq 2$  where  $v \in V(X)$ . Thus we have  $|K| \leq 2p$ . In case  $\Delta_i$  has no edge, then since  $|V(X_N)| = 3$  and  $p \geq 11$  we conclude that  $X[\Delta_i \cup \Delta_{i+1}] \cong C_{2p}$ . Let  $K^*$  be the kernel of the action of K on  $X[\Delta_i \cup \Delta_{i+1}]$ . Now the connectivity of X and transitivity of A/K on  $V(X_N)$  imply that  $K^* = 1$  and consequently,  $K \leq \operatorname{Aut}(X[\Delta_i \cup \Delta_{i+1}]) \cong D_{4p}$ . Since K fixes  $\Delta_i$  it follows that  $|K| \leq 2p$ . Since A/Kis transitive on  $V(X_N)$  it follows that 3 divides |A/K| and A/K has an element of order 3, say  $K\alpha$ , where  $\alpha \in A$ . Therefore  $\alpha^3 \in K$  and hence  $o(\alpha^3) = 1, 2, p$  or 2p. If  $\alpha^3 = 1$ then  $\langle \alpha \rangle$ . P acts transitively and so regularly on V(X), a contradiction. Also if  $o(\alpha^3) = 2$ , p or 2p then  $\alpha^2$ ,  $\alpha^p$  and  $\alpha^{2p}$  have order 3. In each case there is an element of order 3, say  $\beta$ , in A such that  $\langle \beta \rangle P$  acts regularly on V(X), another contradiction. Hence A is not solvable.

Let L = Sol(A),  $\Sigma$  be the set of orbits of L on V(X) and K be the kernel of the action of A on the set of orbits of L. Assume first that L is not trivial. Since  $K_v$  is a  $\{2, 3\}$ -group, it implies that  $K_v$  is solvable and by considering  $K = LK_v$  we see that K is solvable. By the

definition of solvable radical we conclude that K = L and  $A/L \leq \operatorname{Aut}(X_L)$ . If deg $(X_L) = 0$ then A/L is solvable which implies that A is solvable, a contradiction. Since |X| = 3p, this implies that deg $(X_L) \neq 1, 3$ . In case deg $(X_L) = 2$  we have Aut $(X_L) \cong D_{2n}$ , where  $|V(X_L)| = n$  and  $n \in \{3, p\}$ . Now A/L is solvable and so A is solvable, which is impossible. Thus we may assume that deg $(X_L) = 4$  and  $L \leq S_3$ . Thus  $X_L$  is a circulant graph and  $X_L = \operatorname{Cay}(G, S)$ , where  $G \cong \mathbb{Z}_p = \langle a \rangle$ ,  $S = \{a^r, a^{-r}, a^s, a^{-s}\}$  where (r, p) = (s, p) = 1. By [3, Theorem 1.2],  $X_L$  is normal. Hence Aut $(G, S) = \operatorname{Aut}(X_L)$ . Since  $\sigma : a \mapsto a^r$  is an automorphism of G we have  $S = \{a, a^{-1}, a^i, a^{-i}\}$  where (i, p) = 1.

Set  $B = \operatorname{Aut}(X_L)$  and  $Y = X_L$ . Let  $B_1^*$  be the subgroup of  $B_1$  which fixes  $\{1\} \cup S$  and  $Y_2(1)$  be the subgraph of Y with vertex set

$$\{1\} \cup S \cup \{a^2, a^{-2}, a^{i+1}, a^{-i+1}, a^{2i}, a^{-2i}, a^{-i-1}, a^{i-1}\}.$$

Also

$$N_Y(a) = \{1, a^2, a^{i+1}, a^{-i+1}\}, N_Y(a^{-1}) = \{1, a^{-2}, a^{-i-1}, a^{i-1}\}, N_Y(a^i) = \{1, a^{i+1}, a^{2i}, a^{i-1}\}$$
  
and  $N_Y(a^{-i}) = \{1, a^{-i+1}, a^{-2i}, a^{-i-1}\}$ . By a tedious computation we find that  $B_1^* = 1$  and  $B_1$  have no any element of order 4. Thus  $B_1$  acts faithfully on  $S$  and so  $B_1 \leq S_4$ . Since  $B_1$  has no any element of order 4, by considering the subgroups of  $S_4$  we see that either  $B_1 \cong \mathbb{Z}_2 \times \mathbb{Z}_2$  or  $|B_1| \in \{1, 2, 6\}$ . Since  $|B| = |B_1||G|$  we see that  $|B| \in \{p, 2p, 4p, 6p\}$ .  
Moreover  $\alpha : a \to a^{-1}$  is an element of  $\operatorname{Aut}(G, S)$  and  $o(\alpha) = 2$ . Hence  $|B_1| = 2$  or 4 or 6. From the fact that  $A/L$  acts transitively on  $V(Y)$  we have  $p \mid |A/L|$  and  $|A/L| \in \{p, 2p, 4p, 6p\}$ . By considering the order of  $B$  we see that  $G \leq A/L$ . So  $G = T/L$  where  $T \leq A$  and by Proposition 5,  $X$  is a Cayley graph, a contradiction.

Assume now that L is trivial. Then N is not solvable, where N is the minimal normal subgroup of A. If N is semiregular then |N| divides 3p and N is solvable, a contradiction. Therefore N is not semiregular and the valency of quotient graph  $X_N$  is not 4. By considering the order of graph we see that  $X_N$  is not cubic. Thus  $\deg(X_N) = 0$  or 2 and since  $A/K \leq \operatorname{Aut}(X_N)$  we conclude that A/K is solvable. Moreover, since  $K = K_vN$ and  $K_v$  is a  $\{2,3\}$ -group we conclude that K/N is solvable. From the fact that  $A/K \cong$ (A/N)/(K/N) we conclude that A/N is solvable. We claim that N is the unique minimal normal subgroup of A. Suppose by contrary that  $M \neq N$  is another minimal normal subgroup of A. Since  $MN/N \cong M/(M \cap N) \leq A/N$  we see that  $M/(M \cap N)$  is solvable. But  $M \cap N = 1$  and so M is solvable, a contradiction. Hence N is the only minimal normal subgroup of A. Also since |A| is not divided by  $p^2$  we conclude that N is a simple group. Therefore A is an almost simple group and  $\operatorname{soc}(A) = N$ . From the fact that A is a  $\{2,3,p\}$ -group, we conclude that N is a  $K_3$ -group. Also since  $p \geq 11$  it follows that Nis isomorphic to either  $\operatorname{PSL}_2(17)$  or  $\operatorname{PSL}_3(3)$ . First assume that  $N \cong \text{PSL}_2(17)$ . Then by part (*iii*) of Lemma 4,  $A \cong \text{PSL}_2(17)$  or  $\text{PGL}_2(17)$ . If  $A \cong \text{PSL}_2(17)$  then  $|A : A_v| = 51$ . By [7], we see that A has not any subgroup of index 51, a contradiction. Also if  $A \cong \text{PGL}_2(17)$ , then by using GAP [10], A has no any subgroup of index 51, another contradiction.

Assume now that  $N \cong PSL_3(3)$ . In this case, by part (v) of Lemma 4,  $A \cong PSL_3(3)$  or  $A = Aut(PSL_3(3))$ . In the first case, by using GAP [10], and the fact that A is a transitive permutation group of degree 39, we see that A has a regular subgroup which means that X is a Cayley graph, a contradiction. However the later case is impossible, because there is no transitive permutation representation of  $Aut(PSL_3(3))$  of degree 39.

Let X be a connected tetravalent vertex-transitive graph of order 6p. If p = 2 or 3, then by [26], we see that X is a Cayley graph. Also if p = 5 or p = 7 then by [13] and using [28] we see that X is isomorphic to one of the graphs in the following lemma.

**Lemma 7.** A connected tetravalent vertex-transitive non-Cayley graph of order at most 42 are listed below:

- (i) The Coxeter graph;
- (*ii*) the Desargues graph;
- (*iii*) the dodecahedron graph;
- (iv) one of the six graphs in Figure 1.




FIGURE 1. Six graphs illustrated in Lemma 7

**Proof of the Theorem 2.** Let X be a connected tetravalent vertex-transitive graph of order 6p. If  $p \leq 7$ , then X is one of the nine specified graphs in Lemma 7. So we may

assume that  $p \ge 11$ . Suppose that  $X \cong X_{3,p,t}$ , then by Proposition 1, X is a  $\mathcal{VNC}$ -graph. Assume that X is a  $\mathcal{VNC}$ -graph. Let  $A = \operatorname{Aut}(X)$  and  $v \in V(X)$ . Since X is connected and tetravalent,  $A_v$  is a  $\{2,3\}$ -group and so  $|A| = 2^s 3^t p$  for some integers  $s, t \ge 1$ . Assume that N is a minimal normal subgroup of A. We know that  $N = T^k$ , where T is a finite simple group. Let  $\Omega$  be the set of orbits of N on V(X) and K be the kernel of the action of A on  $\Omega$ . Now we consider two cases:

Case I. A is solvable.

In this case N is an elementary abelian r-group, where  $r \in \{2, 3, p\}$ . First suppose that N is an elementary abelian 2-group. In this case  $X_N$  has order 3p and valency 2 or 4. If  $X_N$  has valency 2 then  $X_N \cong C_{3p}$  and  $\operatorname{Aut}(X_N) \cong D_{6p}$ . Suppose that  $\Omega =$  $\{\Delta_0, \Delta_1, \Delta_2, \dots, \Delta_{3p-1}\}$ , where the subscripts are taken modulo 3p. Also suppose that  $\Delta_i \sim \Delta_{i+1}$  in  $X_N$ . If for some *i*, the induced graph  $X[\Delta_i]$  has an edge then by considering the valency of X, we see that  $X[\Delta_{i-1} \cup \Delta_i] \cong C_4$  and  $X[\Delta_i \cup \Delta_{i+1}] \cong K_4$  where  $1 \leq i \leq$ 3p - 1. Now by considering the induced graph  $X[\Delta_0 \cup \Delta_{3p-1}]$ , we have a contradiction. Thus we may assume that each  $X[\Delta_i]$  has no any edge. Thus for each  $i, X[\Delta_i \cup \Delta_{i+1}] \cong C_4$ . It is easy to see that in this case  $X \cong C_{3p}[2K_1]$ , which is a Cayley graph, a contradiction.

Assume now that  $X_N$  has valency 4. Then  $K_v = 1$  for each vertex v. By Proposition 6,  $X_N$  is a Cayley graph, unless the case where  $X_N$  is isomorphic to the line graph of the Petersen graph. Since  $p \ge 11$ ,  $X_N$  is not isomorphic to the line graph of the Petersen graph. Hence  $X_N$  is a Cayley graph and by [21, Proposition 1.2], is a (3, p)-metacirculant graph. Thus  $X_N$  has an automorphism  $\sigma$  of order p such that  $\langle \sigma \rangle$  is semiregular on the vertex set of  $X_N$ . Moreover  $X_N$  has an automorphism say  $\tau$  such that  $\tau$  normalizing  $\langle \sigma \rangle$  and cyclically permutes the 3 orbits of  $\langle \sigma \rangle$  and has a cycle of size 3 in its cycle decomposition. Thus, for some positive integer i, the group  $\langle \sigma, \tau^i \rangle$ , has order 3p and acts regularly on  $V(X_N)$ . Therefore we may suppose that  $X_N = \operatorname{Cay}(H, S)$  where  $H = \langle \sigma, \tau^i \rangle$ . First suppose that  $H \cong F_{3p}$ . If  $X_N$  is normal then  $\operatorname{Aut}(X_N)_1 = \operatorname{Aut}(H, S)$ . Since  $X_N$  is connected and Aut(H, S) acts faithfully on S we conclude that Aut( $X_N$ )<sub>1</sub>  $\leq S_4$ . Moreover  $S = \{x, x^{-1}, y, y^{-1}\}$  where x and y have not the same order. It is easy to see that Aut(H, S) has no any element of order 3 or 4. Thus by considering the subgroups of  $S_4$ we see that  $\operatorname{Aut}(X_N)_1 = 1$  or  $\mathbb{Z}_2$  or  $\mathbb{Z}_2 \times \mathbb{Z}_2$ . Therefore  $|\operatorname{Aut}(X_N)| \leq 12p$  and so |A/K|divides 12p. By considering the order of  $\operatorname{Aut}(X_N)$  we see that  $H \leq A/K$  and so H = T/Kfor some  $T \leq A$ . Now |T| = 6p and T acts regularly on V(X), a contradiction. Thus we may suppose that  $X_N$  is not normal. If  $X_N$  is not primitive then by [17, Theorem 3.1] we see that the valency of  $X_N$  is not equal to 4. Also if  $X_N$  is primitive then again by [34, Lemma 2.1] we have a contradiction. Therefore  $X_N$  is a Cayley graph on  $\mathbb{Z}_{3p}$ .

Thus  $X_N$  is a circulant graph and  $X_N = \operatorname{Cay}(H, S)$ , where  $S = \{a^i, a^{-i}, a^j, a^{-j}\}$  and either (i, 3p) = 1 or (j, 3p) = 1 or (i - j, 3p) = 1. Set  $B = \operatorname{Aut}(X_N)$  and  $Y = X_N$ . Let  $B_1^*$ be the subgroup of  $B_1$  which fixes  $\{1\} \cup S$  and  $Y_2(1)$  be the subgraph of Y with vertex set  $\{1\} \cup S \cup (S^2 - (1 \cup S))$ . We see that

$$S^2 - (1 \cup S) = \{a^{2i}, a^{i+j}, a^{i-j}, a^{-2i}, a^{-i+j}, a^{-i-j}, a^{2j}, a^{-2j}\}.$$

Also

$$N_Y(a^i) = \{1, a^{2i}, a^{i+j}, a^{i-j}\}, N_Y(a^{-i}) = \{1, a^{-2i}, a^{-i+j}, a^{-i-j}\}, N_Y(a^j) = \{1, a^{i+j}, a^{-i+j}, a^{2j}\}$$

and  $N_Y(a^{-j}) = \{1, a^{i-j}, a^{-i-j}, a^{-2j}\}$ . By a tedious computation we have  $B_1^* = 1$  and  $B_1$  has no any element of order 4. Thus  $B_1$  acts faithfully on S and so  $B_1 \leq S_4$ . Since  $B_1$  has no any element of order 3, and the fact that  $\alpha : a \to a^{-1}$  is an element of  $\operatorname{Aut}(H, S)$  and  $o(\alpha) = 2$ , it implies that  $B_1 \cong \mathbb{Z}_2$  or  $\mathbb{Z}_2 \times \mathbb{Z}_2$  and so  $|\operatorname{Aut}(X_N)| \in \{6p, 12p\}$ . Since A/N acts transitively on V(Y) we have that 3p divides |A/N|. Thus  $|A/N| \in \{3p, 6p, 12p\}$ . By considering the order of B, we see that  $H \leq A/N$  and so H = T/N where  $T \leq A$ . Now by Proposition 5, X is a Cayley graph, a contradiction.

Now suppose that N is an elementary abelian 3-group and  $\Omega = \{\Delta_0, \Delta_1, \Delta_2, \cdots, \Delta_{2p-1}\},\$ where the subscripts are taken modulo 2p. Then  $X_N$  has order 2p and so  $X_N$  has valency 2, 3 or 4. First suppose that  $X_N$  has valency 2. Then  $X_N \cong C_{2p}$  and  $\operatorname{Aut}(X_N) \cong D_{4p}$ . Suppose that  $\Delta_i \sim \Delta_{i+1}$  in  $X_N$ . If  $\Delta_i$  has an edge then by considering the valency of X we see that N acts faithfully on  $X[\Delta_i]$ . Hence |N| = 3 and by considering the valency of X we see that  $X[\Delta_{i-1} \cup \Delta_i] \cong 3K_2, |K_v| \le 2$  and  $|K| \le 6$ . Since A/K is transitive on  $V(X_N)$ it follows that 2p divides |A/K|. Hence  $A/K \cong \mathbb{Z}_{2p}$  or  $A/K \cong D_{2p}$  or  $A/K \cong D_{4p}$ . If  $A/K \cong \mathbb{Z}_{2p}$  or  $A/K \cong D_{4p}$ , then A/K has an element of order 2p, say  $K\alpha$ , where  $\alpha \in A$ . Therefore  $\alpha^{2p} \in K$  and hence  $o(\alpha^{2p}) = 1, 2, 3$  or 6. If  $\alpha^{2p} = 1$  then  $\langle \alpha \rangle \times \langle \gamma \rangle$ , where  $\gamma$  is an element of order 3 in K, acts transitively and so regularly on V(X), a contradiction. Also if  $o(\alpha^{2p}) = 2$ , 3 or 6 then  $\alpha^2$ ,  $\alpha^3$  and  $\alpha^6$  have order 2p. Thus in each case there is an element of order 2p, say  $\beta$ , in A such that  $\langle \beta \rangle \times \langle \gamma \rangle$  acts transitively and so regularly on V(X), a contradiction. Thus we may suppose that  $A/K \cong D_{2p}$ . Thus A/K has an element of order p, say  $K\alpha$ . Clearly,  $K\alpha$  does not fix any element of  $V(X_N)$  and so  $\alpha$ does not. Clearly,  $\alpha^p \in K$  and so  $o(\alpha^p) = 1$  or 2 or 3 or 6. In each case we can find an element of order p, say  $\beta$  such that  $\langle \beta \rangle$  acts semiregularly on V(X). Set  $H = \langle \beta \rangle \times \langle \gamma \rangle$ , where  $\gamma$  is an element of order 3 in K, acts semiregularly on V(X) and has two orbits on V(X), which means that X is a bi-Cayley graph over  $H \cong \mathbb{Z}_{3p}$ . Then, by Proposition 1,  $X \cong X_{3,p,t}$ , where  $1 \le t \le p-1$ , and  $t^2 \equiv -1 \pmod{p}$ .

Suppose that  $\Delta_i$  has no edge. Then it is easy to see that either  $X[\Delta_0 \cup \Delta_{2p-1}] = X[\Delta_{i-1} \cup \Delta_i] \cong C_6$   $(1 \le i \le 2p-1)$  or  $X[\Delta_i \cup \Delta_{i+1}] \cong 3K_2$   $(1 \le i \le 2p-1, i \text{ is odd})$  and  $X[\Delta_i \cup \Delta_{i+1}] \cong K_{3,3}$   $(0 \le i \le 2p-1, i \text{ is even}).$ 

First suppose that  $X[\Delta_0 \cup \Delta_{2p-1}] = X[\Delta_{i-1} \cup \Delta_i] \cong C_6$   $(1 \le i \le 2p-1)$ . Then we consider the action of K in  $X[\Delta_i]$   $(0 \le i \le 2p-1)$ . Now it is easy to see that K acts faithfully on  $X[\Delta_i]$  and so  $K \le S_3$ . Since 2p divides |A/K| and |A/K| divides 4p with a similar arguments as above we get a contradiction. Now suppose that  $X[\Delta_i \cup \Delta_{i+1}] \cong 3K_2$  $(1 \le i \le 2p-1, i \text{ is odd})$  and  $X[\Delta_i \cup \Delta_{i+1}] \cong K_{3,3}$   $(0 \le i \le 2p-2, i \text{ is even})$ .

Let  $V(\Delta_i) = \{a_i, b_i, c_i\}$   $(0 \le i \le 2p-1)$ . Without loss of generality we may assume that  $a_i \sim a_{i+1}, b_i \sim b_{i+1}$  and  $c_i \sim c_{i+1}$ . Now it is easy to see that  $\alpha : a_i \mapsto b_i, b_i \mapsto c_i, c_i \mapsto a_i$  and  $\beta : a_i \mapsto a_{i+2}, b_i \mapsto b_{i+2}, c_i \mapsto c_{i+2}$  are automorphisms of X of orders 3 and p, respectively, where  $(0 \le i \le 2p-1)$ . Clearly  $\langle \alpha \rangle \times \langle \beta \rangle \cong \mathbb{Z}_{3p}$  acts semiregularly and has two orbits on V(X). Therefore X is a bi-Cayley graph over the group  $\mathbb{Z}_{3p}$  and again, by Proposition 1,  $X = X_{3,p,t}$ , where  $1 \le t \le p-1$  and  $t^2 \equiv -1 \pmod{p}$ .

Suppose that  $X_N$  has valency either 3 or 4. If  $Aut(X_N)$  is primitive then by [18, Corollary 6.6],  $p \ge 313$ ,  $2p = m^2 + 1$  for some composite integer m. Also  $\Gamma$  has valency either  $\frac{m(m-1)}{2}$  or  $\frac{m(m+1)}{2}$ , a contradiction. Thus we may suppose that  $Aut(X_N)$  is imprimitive. Now by [21, Theorem],  $X_N$  is metacirculant. If  $X_N$  has valency 3 then by [33, Main Theorem],  $X_N$  is isomorphic to generalized Petersen graph GP(p, k). By [9, Theorems 1, 2], we see that  $|\operatorname{Aut}(X_N)| = 4p$ . If  $\Delta_i$  has an edge then  $X[\Delta_i] \cong C_3$   $(0 \le i \le 2p-1)$ . We see that  $X_N$  can not have valency 3, a contradiction. Thus we may suppose that  $\Delta_i$  $(0 \le i \le 2p-1)$  has no edges. Let  $V(\Delta_i) = \{a_i, b_i, c_i\}$   $(0 \le i \le 2p-1)$ . Let  $\Delta_{i+1}, \Delta_{i+2}$ and  $\Delta_{i+3}$  are the vertices adjacent to  $\Delta_i$ . Since  $X_N$  has valency 3, we may suppose that  $X[\Delta_i \cup \Delta_{i+1}] \cong C_6, X[\Delta_i \cup \Delta_{i+2}] \cong 3K_2 \text{ and } X[\Delta_i \cup \Delta_{i+3}] \cong 3K_2.$  Now it is easy to see that  $\alpha : a_i \mapsto b_i, b_i \mapsto c_i, c_i \mapsto a_i$  is an automorphisms of X of order 3. Also K acts faithfully on  $X[\Delta_i]$  and so  $K = N \cong \mathbb{Z}_3$ . By [1, Theorem 1.1], we see that the generalized Petersen graph is a normal bi-Cayley graph on  $H = \mathbb{Z}_p$ . Clearly, H acts semiregularly on  $V(X_N)$ . If H is not a subgroup of A/K then  $p^2 \mid |\operatorname{Aut}(X_N)|$ , a contradiction. Thus  $H \leq A/K$  and A/K has a semiregular element of order p, say  $K\alpha$ . Now  $\alpha^p \in K$  and so  $o(\alpha^p) = 1, 2$  or 3. In each case it is easy to find an element of order p, say  $\beta$ , which is a power of  $\alpha$ . If  $\beta$  fixes an element then it is easy to see that  $K\alpha$  or  $K\alpha^2$  or  $K\alpha^3$  fixes at least one  $\Delta_i$ , a contradiction. Now  $\langle \beta \rangle \times \langle \gamma \rangle \cong \mathbb{Z}_{3p}$  acts semiregularly and has two orbits on V(X) where  $\gamma \in K$  and  $O(\gamma) = 3$ . Therefore X is a bi-Cayley graph over group  $\mathbb{Z}_{3p}$ and by Proposition 1,  $X = X_{3,p,t}$ , where  $1 \le t \le p-1$  and  $t^2 \equiv -1 \pmod{p}$ . Also if  $X_N$ has valency 4 then N = K. Since  $A_v$  is  $\{2, 3\}$ -group we conclude that A has a semiregular element of order p, say  $\delta$ . Since  $N_A(N)/C_A(N)$  is a subgroup of  $\operatorname{Aut}(N) \cong \mathbb{Z}_2$  we conclude that  $\delta$  or  $\delta^2 \in C_A(N)$ . Also if  $N = \langle \gamma \rangle$  then  $\langle \delta, \gamma \rangle$  or  $\langle \delta^2, \gamma \rangle$  is a semiregular subgroup of  $\operatorname{Aut}(X)$  which is isomorphic to  $\mathbb{Z}_{3p}$ . Thus X is a bi-Cayley graph over  $H \cong \mathbb{Z}_{3p} = \langle a \rangle$  and by Proposition 1,  $X = X_{3,p,t}$ , where  $1 \leq t \leq p-1$  and  $t^2 \equiv -1 \pmod{p}$ .

Finally, suppose that N is an elementary abelian p-group. By considering the order of A we see that |N| = p. Suppose that  $N_v \neq 1$  for some  $v \in V(X)$  and w is an another arbitrary vertex of X. Since X is transitive, there exists  $\alpha \in \operatorname{Aut}(X)$  such that  $w^{\alpha} = v$  and so  $N_v = N_{w^{\alpha}} = \alpha^{-1}N_w\alpha = N_w$ . Thus N is trivial, a contradiction. Therefore N acts semiregularly on V(X) and so X is an N-regular cover of  $X_N$ . By [26, Page 173] we know that  $X_N$  is a Cayley graph. Thus  $\operatorname{Aut}(X_N)$  has a subgroup which acts regularly on V(X), say G. Suppose that  $\tilde{G}$  is a lift of G. Now it is easy to see that  $\tilde{G}$  acts regularly on V(X) and so X is a Cayley graph, a contradiction.

**Case II.** A is not solvable.

Let L = Sol(A) be the solvable radical of A and  $\Omega$  be the set of orbits of L on V(X). First suppose that L is not trivial. Let K be the kernel of A on  $\Omega$ . Since  $K_v$  is a  $\{2,3\}$ -group it implies that  $K_v$  is solvable and so by considering  $K = LK_v$  we see that K is solvable. Now by the definition of L we conclude that K = L and  $A/L \leq \operatorname{Aut}(X_L)$ . If  $\deg(X_L) = 0$ then A/L is solvable and so A is solvable, a contradiction. Also if deg $(X_L) = 2$  then Aut $(X_L) \cong D_{2n}$ , where  $|V(X_L)| = n$  and  $n \in \{3, 6, p, 2p, 3p\}$ . Now A/L is solvable and so A is solvable, a contradiction. Thus we may suppose that  $deg(X_L) = 3$  or 4. If  $\deg(X_L) = 3$  then  $n \in \{6, 2p, 3p\}$ . By [16, Theorem 1.1], we see that  $\operatorname{Aut}(X_L)$  is solvable and so A/L is solvable. Now A is solvable, a contradiction. Also if deg $(X_L) = 4$  then L = K and K acts faithfully on each orbits of K. Moreover, in this case  $n \in \{6, p, 2p, 3p\}$ . If n = 6, then  $X_L$  is isomorphic to the octahedral graph and so  $|\operatorname{Aut}(X_L)| = 48$ . Thus  $Aut(X_L)$  is solvable which implies that A is solvable, a contradiction. If n = p then by |V(X)| = |L|n we see that |L| = 6 and  $L \cong S_3$  or  $\mathbb{Z}_6$ . Also since  $A_v$  is a  $\{2,3\}$ -group, it implies that A has a semiregular element of order p, say  $\alpha$ . If L as subgroup of S<sub>6</sub> is isomorphic to  $S_3$  then since L is semiregular we conclude that  $S_3 \times \langle \alpha \rangle$  acts regularly on V(X), a contradiction. Similarly if L as subgroup of  $S_6$  is isomorphic to  $\mathbb{Z}_6 = \langle \beta \rangle$  then  $\langle \alpha \rangle \times \langle \beta \rangle$  acts regularly on V(X), a contradiction. Also if  $|V(X_L)| = 2p$  then |L| = 3. Since  $A_v$  is  $\{2,3\}$ -group we conclude that A has a semiregular element, say  $\alpha$ , of order p, Since  $N_A(L)/C_A(L)$  is a subgroup of Aut(L) we conclude that  $\alpha \in C_A(L)$ . Also if  $\beta \in L$  and  $\beta$  has order 3 then  $\langle \alpha, \beta \rangle$  is a semiregular subgroup of Aut(X) which is isomorphic to  $\mathbb{Z}_{3p}$ . Thus X is a bi-Cayley graph over H, where  $H \cong \mathbb{Z}_{3p}$ . By Proposition 1,  $X = X_{3,p,t}$ , where  $1 \le t \le p-1$  and  $t^2 \equiv -1 \pmod{p}$ . Finally if  $|V(X_L)| = 3p$ then |L| = 2. By Proposition 6,  $X_N$  is a Cayley graph, unless the case where  $X_N$  is isomorphic to the line graph of the Petersen graph. Since  $p \ge 11$ , it follows that  $X_N$  is

not isomorphic to the line graph of the Petersen graph. Also if  $X_N$  is a Cayley graph then  $\operatorname{Aut}(X_N)$  has a subgroup say H which acts regularly on  $V(X_N)$ . Thus |H| = 3p and by [21, Proposition 1.2],  $X_N$  is a (3, p)-metacirculant graph. Thus  $X_N$  has an automorphism  $\sigma$  of order p such that  $\langle \sigma \rangle$  is semiregular on the vertex set of  $X_N$ , and an automorphism  $\tau$ normalizing  $\langle \sigma \rangle$  and cyclically permuting the 3 orbits of  $\langle \sigma \rangle$  such that  $\tau$  has a cycle of size 3 in its cycle decomposition. Thus  $\langle \sigma, \tau^i \rangle$  for some positive integer *i*, has order 3*p* and acts regularly on  $V(X_N)$ . Therefore we may suppose that  $X_N = \operatorname{Cay}(H, S)$ . First suppose that  $H = \langle \sigma, \tau^i \rangle \cong F_{3p}$ . If  $X_N$  is normal then  $\operatorname{Aut}(X_N)_1 = \operatorname{Aut}(H, S)$ . Since  $\operatorname{Aut}(H, S)$  acts faithfully on S we conclude that  $\operatorname{Aut}(X_N)_1 \leq S_4$ . Also since  $X_N$  is connected we conclude that  $S = \{x, x^{-1}, y, y^{-1}\}$  and x and y have not the same orders. Now it is easy to see that Aut(H, S) has no any element of order 3 or 4. Thus by considering the subgroups of  $S_4$  we see that  $\operatorname{Aut}(X_N)_1 = 1$  or  $\mathbb{Z}_2$  or  $\mathbb{Z}_2 \times \mathbb{Z}_2$ . Therefore  $|\operatorname{Aut}(X_N)| \leq 12p$  and so |A/K| divides 12p. Now by considering the order of  $\operatorname{Aut}(X_N)$  we see that  $H \leq A/K$  and so H = T/Kfor some  $T \leq A$ . Now |T| = 6p and T acts regularly on V(X), a contradiction. Thus we may suppose that  $X_N$  is not normal. If  $X_N$  is not primitive then by [17, Theorem 3.1] we see that the valency of  $X_N$  is not equal to 4. Also if  $X_N$  is primitive then again by [34, Lemma 2.1] we have a contradiction. Therefore  $X_N$  is a Cayley graph on  $\mathbb{Z}_{3p}$ . Now similar to the case where A is solvable, we conclude that X is a Cayley graph, a contradiction.

Suppose that L is trivial. By considering the order of A, we see that N is a  $K_3$ -group. Since  $p \ge 11$ , it follows that  $N \cong PSL_2(17)$  or  $PSL_3(3)$ . If N has more than two orbits then  $n \in \{3, 6, 2p, 3p\}$ . First assume that n = 2p or 3p. We know that A/K acts transitively on  $V(X_N)$  and  $K = K_{\alpha}N$ , where  $v \in V(X)$ . Thus  $p^2$  divides |A|, a contradiction. Thus we may suppose that n = 3 or 6. By using GAP [10], it is easy to see that  $PSL_2(17)$  and  $PSL_3(3)$  does not have any orbit of order p or 2p, where  $p \in \{13, 17\}$ . Therefore N has at most two orbits. Now  $Aut(X_N)$  is solvable and so A/K is solvable. Also since  $K_v$  is  $\{2, 3\}$ group and  $K = K_v N$ , it follows that K/N is solvable. Since  $A/K \cong (A/N)/(K/N)$  we conclude that A/N is also solvable. Suppose that M is another minimal normal subgroup of A. Thus by the fact that  $MN/N \cong M/(M \cap N) \le A/N$  we see that  $M/(M \cap N)$  is solvable. But  $M \cap N = 1$  and so M is solvable, a contradiction. Hence N is the only minimal normal subgroup of A. Since |A| is not divisible by  $p^2$  we conclude that N is a simple group. Therefore A is an almost simple group and soc(A) = N. Since A is  $\{2, 3, p\}$ group, it follows that N is a  $K_3$ -group. Also since  $p \ge 11$  we conclude that  $N \cong PSL_2(17)$ or  $N \cong PSL_3(3)$ .

First assume that  $N \cong \text{PSL}_2(17)$ . Then  $A \cong \text{PSL}_2(17)$  or  $\text{PGL}_2(17)$ . If  $A \cong \text{PSL}_2(17)$ then  $|A : A_v| = 102$ . By [7], we see that A has two subgroups of index 102 and both of them have an element of order 4. Also we see that  $A_v$  acts transitively on the set of vertices adjacent to v. By Proposition 3, X is a symmetric graph. If X is one-regular then  $|A| = 2^3.3.17$ , a contradiction. By [2, Theorem 1], we see that  $X \cong C_{3p}[2K_1]$  and so X is a Cayley graph, a contradiction. Also if  $A \cong PGL_2(17)$  then by using GAP [10], A has no any subgroup of index 102, a contradiction.

Assume that  $N \cong \mathrm{PSL}_3(3)$ . Then  $A \cong \mathrm{PSL}_3(3)$  or  $A = \mathrm{Aut}(\mathrm{PSL}_3(3))$ . Let  $A = \mathrm{Aut}(\mathrm{PSL}_3(3))$ . Since  $A \leq S_{78}$  is a transitive permutation group of degree 78, there exists a core-free subgroup H of A of index 78 such that A is isomorphic to the permutation representation of the action of A on the right cosets of H in G by right multiplication. One can check, by using GAP [10], that in this case A has a regular subgroup of order 78 which means that X is a Cayley graph, a contradiction. Hence  $A = \mathrm{PSL}_3(3)$  and  $X = \mathrm{Cos}(\mathrm{PSL}_3(3), H, D)$ , is a coset graph, where  $H \cong (\mathbb{Z}_3 \times \mathbb{Z}_3) \rtimes Q_8$  and D is a union of double cosets of H in G. Since X is 4-valent, we have |D|/|H| = 4. By using GAP [10], there are 15 possibilities for D and in each case  $\langle D \rangle$  is a proper subgroup of  $\mathrm{PSL}_3(3)$  which means that X is disconnected, a contradiction. This complete the proof.

#### References

- M. Arezoomand, M. Ghasemi, Normality of one-matching semi-Cayley graphs over finite abelian groups with maximum degree three, *Contrib. Discrete. Math.* 15(3) (2020) 75–87.
- K. Berčič, M. Ghasemi, Tetravalent arc-transitive graphs of order twice a product of two primes, Discrete Math. 312 (2012) 3643–3648.
- [3] Y.G. Baik, Y.-Q. Feng, H.S. Sim, M.Y. Xu, On the normality of Cayley graphs of abelian groups, Algebra Collog. 5 (1998) 297–304.
- [4] N.L. Biggs, Algebraic Graph Theory, 2nd edn., Cambridge Mathematica Library, Cambridge University Press, Camberidge, 1986.
- [5] H.W. Cheng, M. Ghasemi, S. Qiao, Tetravalent vertex-transitive graphs of order twice a prime square, Graph. Combin. 32 (2016) 1763–1771.
- [6] Y. Cheng, J. Oxley, On weakly symmetric graphs of order twice a prime, J. Combin. Theory B 42 (1987) 196–211.
- [7] J.H. Conway, R.T. Curties, S.P. Norton, R.A. Parker, R.A. Wilson, Atlas of Finite Groups, Clarendon Press, Oxford, 1985.
- [8] Y.-Q. Feng, On vertex-transitive graphs of odd prime-power order, *Discrete Math.* 248 (2002) 265–269.
- [9] R. Frucht, J. E. Graver, M.E. Watkins, The groups of the generalized Petersen graphs, Math. Proc. Camb. Phil. Soc. 70 (1971) 211–218.
- [10] GAP Team, Group, GAP-Groups, Algorithms, and Programming, Version 4.5.5, 2012, http://www.gapsystem.org.
- [11] D. Gorenstein, *Finite simple groups* Plenum Press, New York, 1982.
- [12] A. Hassani, M. A. Iranmanesh, C. E. Praeger, On vertex-imprimitive graphs of order a product of three distinct odd primes, J. Combin. Math. Combin. Comput. 28 (1998) 187–213.
- [13] D. Holt, G. Royle, G. Tracey, The transitive groups of degree 48 and some applications, accepted to J. Algebra

- [14] M.A. Iranmanesh, C.E. Praeger, On non-Cayley vertex-transitive graphs of order a product of three primes, J. Combin. Theory B 81 (2001) 1–19.
- [15] C.H. Li, A. Seress, On vertex-transitive non-Cayley graphs of square-free order, Designs, Codes and Cryptography. 34 (2005) 265–281.
- [16] C.H. Li, Z.P. Lu, G.X. Wang, Vertex-transitive cubic graphs of square-free order, J. Graph Theory. 75 (2014) 1–19.
- [17] Z.P. Lu, M.Y. Xu, On the normality of Cayley graphs of order pq, Austral. J. Combin. 27 (2003) 81–93.
- [18] D. Marušič, On vertex symmetric digraphs, *Discrete Math.* **36** (1981) 69–81.
- [19] D. Marušič, Cayley properties of vertex symmetric graphs, Ars Combin. 16B (1983) 297–302.
- [20] D. Marušič, Vertex transitive graphs and digraphs of order p<sup>k</sup>, Ann. Discrete Math. 27 (1985) 115– 128.
- [21] D. Marušič, R. Scapellato, Characterizing vertex-transitive pq-graphs with an imprimitive automorphism subgroup, J. Graph Theory. 16 (1992) 375–387.
- [22] D. Marušič, R. Scapellato, Classifying vertex-transitive graphs whose order is a product of two primes, *Combinatorica*. 14 (1994) 187–201.
- [23] D. Marušič, R. Scapellato, B. Zgrablič, On quasiprimitive pqr-graphs, Algebra Colloq. 2 (1995) 295– 314.
- [24] B.D. McKay, C.E. Praeger, Vertex-transitive graphs which are not Cayley graphs I, J. Austral. Math. Soc. 56 (1994) 53–63.
- [25] B.D. McKay, C.E. Praeger, Vertex-transitive graphs which are not Cayley graphs II, J. Graph Theory 22 (1996) 321–334.
- [26] B.D. McKay, G. F. Royle, The Transitive Graphs with at Most 26 Vertices. Ars Combin. 30 (1990) 161–176.
- [27] A.A. Miller, C.E. Praeger, Non-Cayley vertex-transitive graphs of order twice the product of two odd primes, J. Algebraic Combin. 3 (1994) 77–111.
- [28] Private communication with Holt and Royle.
- [29] S. Qiao, J. X. Zhou, On tetravalent vertex-transitive bicirculants, Indian J. Pure Appl. Math. 51(1) (2020) 277-288.
- [30] A. Seress, On vertex-transitive non-Cayley graphs of order pqr, Discrete Math. 182 (1998) 279–292.
- [31] B.O. Sabidussi, Vertex-transitive graphs, Monash Math. 68 (1964) 426–438.
- [32] G. Sabidussi, On a class of fix-point-free graphs, Proc. Amer. Math. Soc. 9 (1958) 800-804.
- [33] N.D. Tan, Cubic (m, n)-metacirculant graphs Cayley graphs which are not Cayley graphs, Discrete Math. 154 (1996) 237–244.
- [34] R.J. Wang, M.Y. Xu, A classification of symmetric graphs of order 3p, J. Combin. Theory B 58 (1993) 197–216.
- [35] H. Wielandt, Finite Permutation Groups, Acadamic Press, New York 1964.
- [36] M.Y. Xu, Automorphism groups and isomorphisms of Cayley digraphs, Discrete Math. 82 (1998) 309–319.
- [37] J.-X. Zhou, Tetravalent vertex-transitive graphs of order 4p, J. Graph Theory 71 (2012) 402–415.
- [38] J.-X. Zhou, Cubic vertex-transitive graphs of order 4p (Chinese), J. Sys. Sci. Math. Sci. 28 (2008) 1245–1249.
- [39] J.-X. Zhou, Cubic vertex-transitive graphs of order  $2p^2$  (Chinese), Advance in Math. **37** (2008) 605–609.

## 16 MAJID AREZOOMAND, MOHSEN GHASEMI\* AND MOHAMMAD A. IRANMANESH

[40] J.-X. Zhou, Y.-Q. Feng, Cubic vertex-transitive graphs of order 2pq, J. Graph Theory 65 (2010) 285–302.

### M. Arezoomand

University of Larestan, 74317-16137, Larestan, I. R. Iran Email: arezoomand@lar.ac.ir

# M. Ghasemi

Department of Mathematics, Urmia University, 57135, Urmia, I. R. Iran Email: m.ghasemi@urmia.ac.ir

# Mohammad A. Iranmanesh

Department of Mathematics, Yazd University, 89195-741, Yazd, I. R. Iran Email: iranmanesh@yazd.ac.ir