ON S-1-ABSORBING PRIMARY SUBMODULES

MOHAMMED ISSOUAL, NAJIB MAHDOU, NESLIHAN AYSEN OZKIRISCI, AND ECE YETKIN CELIKEL

ABSTRACT. In this work, we introduce the notion of S-1-absorbing primary submodule as an extension of 1-absorbing primary submodule. Let S be a multiplicatively closed subset of a ring R and M be an R-module. A submodule N of M with $(N :_R M) \cap S = \emptyset$ is said to be S-1-absorbing primary if whenever $abm \in N$ for some non-unit $a, b \in R$ and $m \in M$, then either $sab \in (N :_R M)$ or $sm \in M$ -rad(N). We examine several properties of this concept and provide some characterizations. In addition, S-1-absorbing primary avoidance theorem and S-1-absorbing primary property for idealization and amalgamation are presented.

1. INTRODUCTION

In this article, we assume all rings are commutative with non-zero identity and all modules are unital. We will start with certain definitions and notations that will be used throughout the paper. Let I be an ideal of a ring R and K, L be two submodules of R-module M. The residual K by L and I is defined as $(K :_R L) =$ $\{r \in R \mid rL \subseteq K\}$ and $(K :_M I) = \{m \in M \mid Im \subseteq K\}$, respectively. If I is a principal ideal of R generated by an element $r \in R$, then $(K :_M r)$ is preferred instead of $(K :_M Rr)$. By M-rad(N), we denote the M-radical of N which is the intersection of all prime submodules containing the submodule N.

Consider a multiplicatively closed subset S of a ring R. Recently, S-versions and their features of prime and primary ideals have been thoroughly researched in various studies. Moreover, absorbing-types of such ideals have been proposed and investigated. From the literature, we have gathered the following definitions that we are interested in:

- (1) A submodule N of M with $(N :_R M) \cap S = \emptyset$ is said to be S-1-absorbing prime, if there exists a fixed element $s \in S$ whenever $abm \in N$ for some non-unit elements $a, b \in R$ and $m \in M$, then $sab \in (N :_R M)$ or $sm \in N$, [16].
- (2) A proper ideal I of R is called 1-absorbing primary if whenever $abc \in I$ for some non-unit $a, b, c \in R$, then either $ab \in I$ or $c \in \sqrt{I}$, [7].
- (3) A proper submodule N of M is called 1-absorbing primary if whenever $abm \in N$ for some non-unit $a, b \in R$ and $m \in M$, then either $ab \in (N :_R M)$ or $m \in M$ -rad(N), [27].

²⁰⁰⁰ Mathematics Subject Classification. Primary 13A15, 13A99, 13C13.

Key words and phrases. S-1-absorbing primary submodule, 1-absorbing primary submodule, S-2-absorbing primary submodule, S-primary ideal.

This paper is in final form and no version of it will be submitted for publication elsewhere.

- (4) A submodule N of M with $(N :_R M) \cap S = \emptyset$ is said to be S-primary, if there exists a fixed element $s \in S$ such that whenever $am \in N$ for some $a \in R$ and $m \in M$, then $sa \in \sqrt{(N :_R M)}$ or $sm \in N$, [3].
- (5) A submodule N of M with $(N :_R M) \cap S = \emptyset$ is said to be S-2-absorbing primary, if there exists a fixed element $s \in S$ such that whenever $abm \in N$ for some $a, b \in R$ and $m \in M$, then $sam \in N$ or $sbm \in N$ or $sab \in \sqrt{(N :_R M)}$, [24].

Our purpose in this paper is to extend the notion of S-1-absorbing primary ideals and introduce the concept of S-1-absorbing primary submodules. This study is composed of five sections; in the second section, the concept of S-1-absorbing primary submodule is defined and examined in detail. Several characterizations of S-1-absorbing primary submodules are also presented. We exemplify the differences between S-1-absorbing primary and S-1-absorbing prime submodules, as well as 1-absorbing primary submodules. We provide some theorems to examine the relations between S-1-absorbing primary submodules and the other notions such as S-primary and 1-absorbing primary submodules.

In [20], it has been proposed and investigated the prime avoidance theorem for modules. This theorem says that if a submodule N of M is contained in a union of finitely many prime submodules P_i 's with $(P_j: M) \not\subseteq (P_k: M)$ whenever $j \neq k$, then N is contained in P_i for some i. This theorem has also been extended to primary submodules in various ways [4, 13]. In the third section, we provide S-1-absorbing primary avoidance theorem under the assumption that M is finitely generated multiplication.

In section 4, we examine the S-1-absorbing primary property for idealization of M in R. The idealization of M in R is the ring $R(+)M = \{(r,m) \mid r \in R, m \in M\}$ with component-wise addition and multiplication defined as $(r_1, m_1)(r_2, m_2) = (r_1r_2, r_1m_2 + r_2m_1)$. For more information and the ideal structure of an idealization of M, the reader may see [2].

Let (A, B) be a pair of commutative rings with unity, $f : A \to B$ be a ring homomorphism and J be an ideal of B. In this setting, we can consider the following subring of $A \times B$:

$$A \bowtie^{f} J := \{(a, f(a) + j) \mid a \in A, j \in J\}$$

called the amalgamation of A with B along J with respect to f. This construction is defined and examined by D'Anna et al. [11] and then extensively investigated in different papers [12, 25, 15]. In the last section, we study S-1-absorbing primary ideals of the amalgamation $A \bowtie^f J$.

2. Characterization of S-1-absorbing primary submodules

Definition 1. Let S be a multiplicatively closed subset of a ring R and M an Rmodule. We call a submodule N of M with $(N :_R M) \cap S = \emptyset$ an S-1-absorbing primary if there exists an (fixed) $s \in S$ such that for all non-unit $a, b \in R$ and $m \in M$ if $abm \in N$, then either sab $\in (N :_R M)$ or $sm \in M$ -rad(N). This fixed element $s \in S$ is called an S-element of N.

In particular, if one take M = R as an *R*-module, then we have the definition of *S*-1–absorbing primary ideal of *R* as following:

Definition 2. Let S be a multiplicatively closed subset of a ring R. We call a proper ideal I of R with $I \cap S = \emptyset$ an S-1-absorbing primary if there exists an

(fixed) $s \in S$ such that whenever $abc \in I$ for some non-unit $a, b, c \in R$, then either $sab \in I$ or $sc \in \sqrt{I}$. This fixed element $s \in S$ is called an S-element of I.

Let N be a proper submodule of M. It is clear that any S-1-absorbing prime submodule is a S-1-absorbing primary submodule and that two concepts coincide if M-rad(N) = N. However, this implication is not reversible in general, see the following example.

Example 1. (S-1-absorbing primary submodule that is not S-1-absorbing prime) Consider the \mathbb{Z} -module $M = \mathbb{Z} \times \mathbb{Z}$ and the submodule $N = 3\mathbb{Z} \times 8\mathbb{Z}$ of M. For the multiplicatively closed subset $S = \{3^n : n > 0\}$, N is an S-1-absorbing primary submodule which is not S-1-absorbing prime.

Proof. Suppose that $ab(m_1, m_2) \in N$ for some integers m_1, m_2 and $a, b \neq \pm 1$. If m_2 is even, then $sm \in 3\mathbb{Z} \times 2\mathbb{Z} = M\text{-}rad(N)$ for all $s \in S$. If m_2 is odd, then since $abm_2 \in 8\mathbb{Z}$, we conclude that $ab \in 8\mathbb{Z}$, and so $sab \in 24\mathbb{Z} = (N :_R M)$ for all $s \in S$. Thus N is an S-1-absorbing primary submodule. As $2 \cdot 3 \cdot (1, 4) \in N$ but neither $6s \in (N :_R M) = 24\mathbb{Z}$ nor $s(1, 4) \in N$ for any $s \in S$, N is not S-1-absorbing prime. \Box

Let S be a multiplicatively closed subset of a ring R. In the following proposition, by S^* , we denote the saturation set of S defined by $\{r \in R : \frac{r}{1} \in U(S^{-1}R)\}$. It is clear that $S \subseteq S^*$ and S^* is also a multiplicatively closed subset of R. In case of $S = S^*$, then S is said to be a saturated set, [17]. For an R-module M, by $U_M(R)$, we denote the set $\{r \in R : rM = M\}$ which contains U(R) is a saturated multiplicatively closed subset of R.

Proposition 1. Let S be a multiplicatively closed subset of a ring R, and N be a submodule of an R-module M.

- (1) Let $T \subseteq S$ be two multiplicatively closed subsets of R. If N is an T-1absorbing primary submodule such that $(N :_R M) \cap S = \emptyset$, then N is also an S-1-absorbing primary submodule. Conversely, if for each $s \in S$, there is an element $s' \in T$ such that $ss' \in T$ and N is a S-1-absorbing primary submodule of M, then N is an T-1-absorbing primary submodule of M.
- (2) Every 1-absorbing primary submodule N with $(N :_R M) \cap S = \emptyset$, is S-1absorbing primary. However, it is clear that these two classes of 1-absorbing primary and S-1-absorbing primary submodules coincide if $S \subseteq U(R)$.
- (3) N is an S-1-absorbing primary submodule of M if and only if N is S^{\star} -1-absorbing primary.
- (4) Suppose that M = Rm is a cyclic *R*-module and $S \subseteq U_M(R)$. Then *N* is a 1-absorbing primary submodule if and only if *N* is an *S*-1-absorbing primary submodule.

Proof. (1) It is clear.

(2) The claim follows by taking $T = \{1\}$ in (1).

(3) Suppose that N is S-1-absorbing primary submodule. It is clear that $(N :_R M) \cap S^* = \emptyset$. Since $S \subseteq S^*$, then by (1), N is S^* -1-absorbing primary. For the converse, let $abm \in N$ for some non-unit elements $a, b \in R$ and $m \in M$. Since N is an S^* -1-absorbing primary submodule, there exists an $s \in S^*$ such that $sab \in (N :_R M)$ or $sm \in M$ -rad(N). Since $\frac{s}{1}$ is a unit of $S^{-1}R$, $sur \in S$ for

some $u \in S$ and $r \in R$. Say $t := sur \in S$. Hence, we conclude $tm \in M$ -rad(N) or $tab \in (N :_R M)$, as needed.

(4) Suppose that N is an S-1-absorbing primary submodule. Let $abm' \in N$ for some non-unit $a, b \in R$ and $m' = rm \in M$. Then there exists an $s \in S$ such that $sab \in (N :_R M)$ or $srm \in M$ -rad(N). If $srm \in M$ -rad(N), then since $S \subseteq U_M(R)$ we have $srRm = rM \subseteq M$ -rad(N) which implies $m' = rm \in M$ -rad(N). If $sab \in (N :_R M)$, then since M is a multiplication module and $S \subseteq U_M(R)$, we conclude $abM = sabM \subseteq (N :_R M)M \subseteq M$ -rad(N). Thus, N is a 1-absorbing primary submodule. The converse part follows from (2).

Example 2. (S-1-absorbing primary submodule that is not 1-absorbing primary) Consider the multiplicatively closed subset $S = \{2^k 3^t : k, t \ge 0\}$ of \mathbb{Z} . Then the \mathbb{Z} -submodule N = (0) of $M = \mathbb{Q} \times \mathbb{Z}_2 \times \mathbb{Z}_3$ is an S-1-absorbing primary submodule which is not 1-absorbing primary.

Proof. Suppose that $ab\left(\frac{c}{d}, \overline{x}, \overline{y}\right) \in N$ for some $\pm 1 \neq a, b \in \mathbb{Z}$ and $\left(\frac{c}{d}, \overline{x}, \overline{y}\right) \in M$. Consider the element $6 \in S$. Here, abc = 0 which implies a = 0 or b = 0 or c = 0. If a = 0 or b = 0, then clearly $sab \in (N :_{\mathbb{Z}} M)$. If c = 0, then $6\left(\frac{c}{d}, \overline{x}, \overline{y}\right) = (0, \overline{0}, \overline{0}) \in M$ -rad(N) = (0), and thus N is an S-1-absorbing primary submodule. However, it is not 1-absorbing primary as $2 \cdot 3 \cdot (0, \overline{1}, \overline{1}) \in N$ but neither $2 \cdot 3 \in (N :_{\mathbb{Z}} M) = 0$ nor $(0, \overline{1}, \overline{1}) \in M$ -rad(N) = (0).

Theorem 1. Let S be a multiplicatively closed subset of a ring R. For a submodule N of an R-module M satisfying $(N :_R M) \cap S = \emptyset$, the following are equivalent.

- (1) N is an S-1-absorbing primary submodule of M.
- (2) There exists an $s \in S$ such that whenever $Ibm \subseteq N$ for some proper ideal I of $R, b \in R \setminus U(R)$ and $m \in M$ implies either $sIb \subseteq (N :_R M)$ or $sm \in M$ -rad(N).
- (3) There exists an $s \in S$ such that whenever $IJm \subseteq N$ for some proper ideals I, J of R and $m \in M$ implies either $sIJ \subseteq (N :_R M)$ or $sm \in M$ -rad(N).
- (4) There exists an $s \in S$ such that whenever $IJK \subseteq N$ for some proper ideals I, J of R and a submodule K of M implies either $sIJ \subseteq (N:_R M)$ or $sK \subseteq M$ -rad(N).

Proof. $(1) \Rightarrow (2)$. Let s be an S-element of N. Suppose that $Ibm \subseteq N$ and $sIb \notin (N :_R M)$. Then $sxb \notin (N :_R M)$ for some non-unit $x \in I$. Since $xbm \in N$ and $sxb \notin (N :_R M)$, we have $sm \in M$ -rad(N), as needed.

 $(2) \Rightarrow (3)$. Let s be the fixed element in (2). Suppose that $IJm \subseteq N$ but $sIJ \notin (N:_R M)$ for some proper ideals I, J of R and $m \in M$. Then $sIb \notin (N:_R M)$ for some non-unit $b \in J$. From our assumption (2), we conclude $sm \in M$ -rad(N).

 $(3) \Rightarrow (4)$. Let s be the fixed element in (3). Suppose $IJK \subseteq N$ but $sK \not\subseteq N$ for some proper ideals I, J of R and a submodule K of M. Hence $sm \notin M\text{-}rad(N)$ for some $m \in K$ which implies $sIJ \subseteq (N :_R M)$ by (3).

 $(4) \Rightarrow (1)$. Let $a, b \in R$ be non-unit elements, $m \in M$ and $abm \in N$. Put I = aR, J = bR and K = Rm in (4), then we are done.

In the following, we characterize S-1-absorbing primary ideals of a ring R by taking M = R as an R-module in Theorem 1.

Theorem 2. Let S be a multiplicatively closed subset of a ring R and I an ideal of R with $I \cap S = \emptyset$. Then I is an S-1-absorbing primary ideal of R if and only if

there exists an $s \in S$ such that whenever $I_1I_2I_3 \subseteq I$ for some proper ideals I_1, I_2 and I_3 of R implies either $sI_1I_2 \subseteq I$ or $sI_3 \subseteq \sqrt{I}$.

Recall from [22, Lemma 2.4] that if M is a finitely generated multiplication R-module, then $\sqrt{(N:_R M)} = (M \cdot rad(N):_R M)$ for any submodule N of M

Next, we give a characterization for S-1-absorbing primary submodules in terms of submodules of a finitely generated multiplication modules.

Theorem 3. Let S be a multiplicatively closed subset of a ring R and M be a finitely generated multiplication R-module. A proper submodule N of M with $(N :_R M) \cap S = \emptyset$ is an S-1-absorbing primary submodule of M if and only if there exists an $s \in S$ such that whenever $N_1N_2N_3 \subseteq N$ for some proper submodules N_1, N_2, N_3 of M implies either $sN_1N_2 \subseteq N$ or $sN_3 \subseteq M$ -rad(N).

Proof. Suppose that $N_1N_2N_3 \subseteq N$ for some proper submodules $N_1 = I_1M$, $N_2 = I_2M$ and $N_3 = I_3M$ where I_1 , I_2 , I_3 are proper ideals of R. Hence $I_1I_2I_3 \subseteq (N:_R M)$ and $(N:_R M)$ is a S-1-absorbing primary ideal of R by Lemma 4. This yields that there exists $s \in S$ such that $sI_1I_2 \subseteq (N:_R M)$ or $sI_3 \subseteq \sqrt{(N:_R M)}$ by Theorem 2. Thus we conclude either $sN_1N_2 = sIJM \subseteq (N:_R M)M = N$ or $sN_3 \subseteq \sqrt{(N:_R M)M} = M$ -rad(N), as desired. For the converse, suppose that $IJm \subseteq N$ for some proper ideals I, J of R and $m \in M$. Put $N_1 = IM$, $N_2 = JM$ and $N_3 = Rm$. This follows that $sIJ \subseteq (N:_R M)$ or $sm \in sN_3 \subseteq M$ -rad(N), thus we are done by Theorem 1.

Now, we are ready for a characterization of S-1-absorbing primary submodules of finitely generated faithful multiplication modules in terms of presentation ideals of them.

Theorem 4. Let M be a finitely generated faithful multiplication R-module and S be a multiplicatively closed subset of R. For a submodule N of M provided $(N :_R M) \cap S = \emptyset$, the following statements are equivalent.

- (1) N is a S-1-absorbing primary submodule of M.
- (2) N = IM for some S-1-absorbing primary ideal I of R.
- (3) $(N:_R M)$ is an S-1-absorbing primary ideal of R.

Proof. (1) \Rightarrow (2). Let N be a S-1-absorbing primary submodule of M. Then $(N :_R M)$ is an S-1-absorbing primary ideal of R by Proposition 4. Since M is multiplication, put $I = (N :_R M)$, so the claim is clear.

 $(2) \Rightarrow (3)$. Since M is cancellation, that is, $IM = JM \Leftrightarrow I = J$ for every ideal I, J of R which means that $(N :_R M) = (IM :_R M) = I$ is an S-1-absorbing primary ideal of R.

 $(3) \Rightarrow (1)$. Suppose that $0 \neq IJK \subseteq N$ for some proper ideals I, J of R and a submodule K of M. Since M is a multiplication module, K = LM for some ideal L of R. Then we have $IJLM \subseteq (N :_R M)$. We may assume that L is a proper ideal of R. Since $(N :_R M)$ is an S-1-absorbing primary ideal of R, so it is an S-1-absorbing primary submodule of an R-module R and there exists a fixed $s \in S$ such that $sIJ \subseteq (N :_R M)$ or $sL \subseteq \sqrt{(N :_R M)} = (M \cdot rad(N) :_R M)$ by Theorem 2. This yields that $sIJ \subseteq (N :_R M)$ or $sK = LM \subseteq M \cdot rad(N)$ and N is S-1-absorbing primary submodule of M by Theorem 1. \Box

Every S-primary submodule of a finitely generated faithful multiplication module is S-1-absorbing primary, see the following.

Proposition 2. Let S be a multiplicatively closed subset of a ring R and M be a finitely generated faithful multiplication R-module. If N is an S-primary submodule of M, then it is S-1-absorbing primary.

Proof. Let N be an S-primary submodule of M. Then $(N :_R M)$ is a S-primary ideal of R by [3, Proposition 2.11 (1)] and hence $(N :_R M)$ is a S-1-absorbing primary ideal of R. Thus N is an S-1-absorbing primary submodule of M by Theorem 4.

Lemma 1. Let S be a multiplicatively closed subset of R. If I is an S-1-absorbing primary ideal, then \sqrt{I} is an S-prime ideal of R.

Proof. Let s be an S-element of I, $a, b \in R$ and $ab \in \sqrt{I}$. If a or b is unit, then the claim is clear. So, assume that a, b are non-unit elements. Then $a^k b^k = a^t a^t b^k \in I$ for some k = 2t > 1 which implies that $sa^k \in I$ or $sb^k \in \sqrt{I}$. Thus $sa \in \sqrt{I}$ or $sb \in \sqrt{I}$ and \sqrt{I} is an S-prime ideal of R.

Proposition 3. Let S be a multiplicatively closed subset of R. If N is an S-1absorbing primary submodule of M, then M-rad(N) is an S-prime submodule of M.

Proof. Follows from Theorem 4, Lemma 1 and [26, Proposition 2.9 (2)]. \Box

Let S be a multiplicatively closed subset of a ring R. Then S is said to satisfy the maximal multiple condition if there exists $s \in S$ such that t divides s for each $t \in S$, [1]. For instance, if S is finite, then S satisfies the maximal multiple condition. Also, U(R) is a multiplicatively closed set satisfying the maximal multiple condition.

Proposition 4. Let S be a multiplicatively closed subset of a ring R, and N be a submodule of an R-module M.

- (1) If N is an S-1-absorbing primary submodule of M, then $S^{-1}N$ is a 1-absorbing primary $S^{-1}R$ -submodule of $S^{-1}M$.
- (2) If S satisfies the maximal multiple condition, and $S^{-1}N$ is a 1-absorbing primary $S^{-1}R$ -submodule of $S^{-1}M$, then N is an S-1-absorbing primary submodule of M.

Proof. (1) Let $\frac{a}{s_1}, \frac{b}{s_2}$ be non-unit elements of $S^{-1}R$ and $\frac{m}{s_3} \in S^{-1}M$ with $\frac{a}{s_1}, \frac{b}{s_2}, \frac{m}{s_3} \in S^{-1}N$. Then $uabm \in N$ for some $u \in S$. Hence, there exists an $s \in S$ such that $suab \in (N :_R M)$ or $sm \in M$ -rad(N). This yields either $\frac{a}{s_1}, \frac{b}{s_2} = \frac{suab}{sus_1s_2} \in S^{-1}(N :_R M) \subseteq (S^{-1}N :_{S^{-1}R}, S^{-1}M)$ or $\frac{m}{s_3} = \frac{sm}{ss_3} \in S^{-1}(M$ -rad $(N)) = S^{-1}M$ -rad $(S^{-1}N)$ and $S^{-1}N$ is a 1-absorbing primary submodule of $S^{-1}M$.

(2) Let $s_{\max} \in S$ is the element which has the maximal multiple condition. Suppose that $abm \in N$ for some non-unit elements $a, b \in R$ and $m \in M$. Then $\frac{a}{1}\frac{b}{1}\frac{m}{1} \in S^{-1}N$ which implies $\frac{a}{1}\frac{b}{1} \in (S^{-1}N :_{S^{-1}R} S^{-1}M) \subseteq S^{-1}(N :_R M)$ or $\frac{m}{1} \in S^{-1}M$ -rad $(S^{-1}N) = S^{-1}(M$ -rad(N)). This follows that $uab \in (N :_R M)$ or $vm \in M$ -rad(N). Since u and v divide s_{\max} , we get $s_{\max}ab \in (N :_R M)$ or $s_{\max}m \in M$ -rad(N) as needed.

Proposition 5. Let S be a multiplicatively closed subset of R, and M be a multiplication R-module. If $\{N_i\}_{i=1}^k$ is a family of S-1-absorbing primary submodules of

M with the same M-radical, then $\bigcap_{i=1}^{k} N_i$ is also S-1-absorbing primary submodule of M.

Proof. Let $s_i \in S$ be an *S*-element of N_i for each i = 1, ..., k. First, note that $\left(\bigcap_{i=1}^k N_i\right) \cap S = \emptyset$. Put $s = s_1 s_2 ... s_k$. Suppose that $abm \in \bigcap_{i=1}^k N_i$ but $sab \notin \left(\bigcap_{i=1}^k N_i :_R M\right)$ for some non-unit elements $a, b \in R$ and $m \in M$. Then $sab \notin (N_j :_R M)$ for some $j \in \{1, ..., k\}$. Since N_j is *S*-1-absorbing primary and $abm \in N_j$, we have $s_jm \in M$ -rad (N_j) and so $sm \in M$ -rad $(N_j) = M$ -rad $\left(\bigcap_{i=1}^k N_i\right)$ by [22, Proposition 2.14 (3)], we are done. □

Lemma 2. Let $f : M_1 \to M_2$ be an *R*-module epimorphism and *S* a multiplicatively closed subset of *R*.

- (1) If N is a submodule of M_1 , then $(N :_R M_1) \subseteq (f(N) :_R M_2)$, [27].
- (2) If K is a submodule of M_2 , then $(K :_R M_2) \subseteq (f^{-1}(K) :_R M_1), [27].$
- (3) If N is a submodule of M_1 and ker $(f) \subseteq N$, then $f(M_1 \operatorname{rad}(N)) = M_2 \operatorname{rad}(f(N))$, [19].
- (4) If K is a submodule of M_2 , then $f^{-1}(M_2 \operatorname{rad}(K)) = M_1 \operatorname{rad}(f^{-1}(K)),$ [19].

Proposition 6. Let $f : M_1 \to M_2$ be an *R*-module epimorphism and *S* be a multiplicatively closed subset of *R*.

- (1) Suppose that $(f(N) :_R M_2) \cap S = \emptyset$. If N is an S-1-absorbing primary submodule of M_1 containing Ker(f), then f(N) is an S-1-absorbing primary submodule of R_2 .
- (2) If K is an S-1-absorbing primary submodule of M_2 , then $f^{-1}(K)$ is an S-1-absorbing primary submodule of M_1 .

Proof. (1) Let a, b be non-unit elements of R and $m \in M$ with $abm \in f(N)$. Say $m = f(m_1)$ for some $m_1 \in M_1$. Since $abf(m_1) \in f(N)$ and $Ker(f) \subseteq I$, we have $abm_1 \in N$ which implies that there exists an $s \in S$ such that $sab \in (N :_R M_1)$ or $sm \in M_1$ -rad(N). Thus, $sab \in (N :_R M_1) \subseteq f((N) :_R M_2)$ or $sf(m) \in f(N) \subseteq M_2$ -rad(f(N)) by Lemma 2.

(2) First, we show that $(f^{-1}(K):_R M_1) \cap S = \emptyset$. Assume that $r \in (f^{-1}(K):_R M_1) \cap S$. S. Then $rM_1 \subseteq f^{-1}(K)$ implies $rM_2 = rf(M_1) \subseteq f(f^{-1}(K)) \subseteq K$, and hence $r \in (K:_R M_2)$, a contradiction. Let *a* and *b* be non-unit elements of *R* and $m \in M_1$ with $abm \in f^{-1}(K)$. Then $abf(m) \in K$ which implies that there exists $s \in S$ such that $sab \in (K:_R M_2)$ or $sf(m) \in M_2$ -rad(K). Thus, we conclude $sab \in (K:_R M_2) \subseteq (f^{-1}(K):_R M_1)$ or $sm \in f^{-1}(M_2$ -rad $(K)) = M_1$ -rad $(f^{-1}(K))$ by Lemma 2. Therefore, $f^{-1}(K)$ is an S-1-absorbing primary submodule of M_1 . \Box

As a consequence of Theorem 6, we have the following result.

Corollary 1. Let S be a multiplicatively closed subset of R and N, K be submodules of R-module M.

- (1) Let $K \subseteq N$. Then N is an S-1-absorbing primary submodule of M if and only if N/K is an S-1-absorbing primary submodule of M/K.
- (2) If N is an S-1-absorbing primary submodule of M with $(N :_R K) \cap S = \emptyset$, then $N \cap K$ is an S-1-absorbing primary submodule of K.

Proof. (1) It is clear that $(N/K :_R M/K) \cap S = \emptyset$. Consider the canonical epimorphism $\pi : M \to M/K$ in Proposition 6 (1). Then N/K is an S-1-absorbing primary submodule of M/K. For the converse part, suppose that $abm \in N$ for some non-unit elements a and b of R, $m \in M$. Hence $ab(m + K) \in N/K$ which yields that there exists a fixed $s \in S$ such that $sab \in (N/K :_R M/K)$ or $s(m + K) \in M/K$ -rad(N/K) = M-rad(N)/K. Therefore $sab \in (N :_R M)$ or $sm \in M$ -rad(N), so we are done.

(2) Consider that the injection $i: K \to M$ defined by i(k) = k for all $k \in K$. It is easy to see that $(i^{-1}(N): K) \cap S = \emptyset$. Then $i^{-1}(N) = N \cap K$ is an S-1-absorbing primary submodule of K by Proposition 6 (2).

Next, we show that if a ring R admits an S-1-absorbing primary ideal that is not an S-primary ideal, then R is a quasilocal ring. First, recall the following lemma.

Lemma 3. [7] Let R be a ring. Suppose that for every non-unit element w of R and for every unit element u of R, we have w + u is a unit element of R. Then R is a quasilocal ring.

Theorem 5. Let R be a ring. If there is an S-1-absorbing primary ideal which is not an S-primary, then R is a quasilocal ring.

Proof. Let $s' \in S$ be an S-element of J which is not S-primary ideal of R. Then there exist non-unit elements $a, b \in R$ with $ab \in J$ satisfying $sa \notin J$ and $sy \notin \sqrt{J}$ for all $s \in S$. Choose $r \in R \setminus U(R)$. Hence $rab \in J$ which yields $s'ra \in J$. We show that $r + u \in U(R)$ for any $u \in U(R)$. Let $u \in U(R)$. Assume that $r + u \in R \setminus U(R)$. Then $(r + u)ab \in J$ and $s'y \notin \sqrt{I}$ implies that $s'(r + u)a \in I$. Now, since $s'ra \in I$, we get $s'ua \in I$, and so $s'a \in J$, a contradiction. Therefore, $r + u \in U(R)$ and R is a quasilocal ring by Lemma 3.

As a direct result of Theorem 5, we have the following corollary.

Corollary 2. Let I be an ideal of a non-quasi local ring R with $I \cap S = \emptyset$. Then I is an S-1-absorbing primary ideal of R if and only if I is an S-primary ideal of R.

Lemma 4. Let S be a multiplicatively closed subset of R. If N is an S-1-absorbing primary submodule of a finitely generated multiplication R-module M. Then the following statements hold.

- (1) $(N:_R M)$ is an S-1-absorbing primary ideal of R.
- (2) If $(N:_R m) \cap S = \emptyset$ for some $m \in M \setminus \mathfrak{P}$ where \mathfrak{P} is any prime submodule of M containing N, then $(N:_R m)$ is an S-1-absorbing primary ideal of R.
- (3) If $((N:_M r):_R M) \cap S = \emptyset$ for some $r \in R \setminus (N:_R M)$, then $(N:_M r)$ is an S-1-absorbing primary submodule of M.

Proof. (1) Let $a, b, c \in R \setminus U(R)$ and $abc \in (N :_R M)$. Then $ab(cM) \subseteq N$ implies that there exists an $s \in S$ such that $sab \in (N :_R M)$ or $scM \subseteq M$ -rad(N) by Theorem 1. Thus we have $sab \in (N :_R M)$ or $sc \in (M$ - $rad(N) :_R M) = \sqrt{(N :_R M)}$ and $(N :_R M)$ is an S-1-absorbing primary ideal of R.

(2) Let $a, b, c \in R \setminus U(R)$ and $abc \in (N :_R m)$. Then we have $abcm \in N$. As N is an S-1-absorbing primary submodule, there exists an $s \in S$ such that $sab \in (N :_R M)$ or $scm \in M$ -rad(N). This indicates that $sab \in (N :_R m)$ or $sc \in (M$ -rad $(N) :_R m)$. For the rest of the proof, we only need to show that (M-rad $(N) :_R m) = \sqrt{(N :_R m)}$. The inclusion $\sqrt{(N :_R m)} \subseteq (M$ -rad $(N) :_R m)$ is satisfied by ([21], Lemma 2.1). For the reverse inclusion, let $a \in (M$ -rad $(N) :_R m)$ is satisfied by ([21], Lemma 2.1). For the reverse inclusion, let $a \in (M$ -rad $(N) :_R m)$ which implies $am \in M$ -rad $(N) = \bigcap_{N \subseteq P} P$ where P is any prime submodule of M that contains N. Since M is multiplication, there exists an ideal I of R such that N = IM and a prime ideal Q of R such that P = QM. It follows that $am \in \bigcap_{N \subseteq P} P = \bigcap_{IM \subseteq QM} QM$. Then, we obtain $am \in QM$ for all prime ideals Q containing I by using ([14], Theorem 3.2). This means that $a \in Q$ or $m \in QM$ for all primes Q containing I by ([14], Lemma 2.10) and hence $a \in \bigcap_{I \subseteq Q} Q = \sqrt{I}$ which implies $a^n \in I$ for some $n \in \mathbb{N}$. We conclude that $a^n m \in IM = N$, namely $a \in \sqrt{(N :_R m)}$, which completes the proof.

(3) Suppose that $abm \in (N :_M r)$ for some non-unit $a, b \in R$ and $m \in M$. Then $abrm \in N$ which implies that there exists an $s \in S$ such that $sab \in (N :_R M) \subseteq ((N :_M r) :_R M)$ or $srm \in M$ - $rad(N) \subseteq M$ - $rad((N :_M r))$. Thus $(N :_M r)$ is an S-1-absorbing primary submodule of M.

Theorem 6. Let M be a finitely generated multiplication R-module where R is a non-quasi local ring and N be a submodule of M with $(N :_R M) \cap S = \emptyset$. Then N is an S-1-absorbing primary submodule of M if and only if N is an S-primary submodule of M.

Proof. Suppose that N is a S-1-absorbing primary submodule of M. Then $(N :_R M)$ is an S-1-absorbing primary ideal of R by Lemma 4 and $(N :_R M)$ is S-primary ideal of R by Corollary 2. Hence, N is an S-primary submodule of M by [3, Theorem 2.17]. Conversely, if N is an S-primary submodule of M, then $(N :_R M)$ is an S-primary ideal of R by [3, Theorem 2.17]. Since every S-primary ideal with $(N :_R M) \cap S = \emptyset$ is an S-1-absorbing primary ideal, $(N :_R M)$ is an S-1-absorbing primary ideal of R. Thus, N is an S-1-absorbing primary submodule of M by Theorem 1. □

In view of Corollary 2, we have the following result.

Theorem 7. Let S_i be multiplicatively closed subset of ring R_i for each i = 1, ..., nand $S = S_1 \times \cdots \times S_n$. For an ideal $I = I_1 \times \cdots \times I_n$ of $R = R_1 \times \cdots \times R_n$, the following are equivalent.

- (1) I is an S-1-absorbing primary ideal of R.
- (2) I is an S-primary ideal of R.
- (3) I_j is an S_j -primary ideal of R_j for some $j \in \{1, \ldots, n\}$ and $I_k \cap S_k \neq \emptyset$ for all $k \in \{1, \ldots, n\} \setminus \{j\}$.

Proof. (1) \Leftrightarrow (2) Since R is not quasi-local, S-1-absorbing primary ideals and S-primary ideals coincide by Corollary 2. (2) \Leftrightarrow (3) Follows from [3, Theorem 2.21] considering $M_i = R_i$ as an R_i -module for all $i \in \{1, \ldots, n\}$.

Proposition 7. Let S be a multiplicatively closed subset of R, M_1 , M_2 be R-modules and $M = M_1 \times M$ and N_1 be a proper submodule of M_1 . If $N = N_1 \times M_2$ is an $S \times 0$ -1-absorbing primary submodule of R-module M, then N_1 is an S-1-absorbing primary submodule of M_1 .

Proof. Suppose that $N = N_1 \times M_2$ is an $S \times 0$ -1-absorbing primary submodule of M. Then $N_1 \cong N/(\{0\} \times M_2)$ is an S-1-absorbing primary submodule of $M/(\{0\} \times M_2) \cong M_1$ by Corollary 1.

Theorem 8. Let S be a multiplicatively close subset of R, M_1 , M_2 be R-modules and $M = M_1 \times M_2$ and N_1 , N_2 be proper submodules respectively of M_1 and M_2 . Let $M = M_1 \times M_2$, $N = N_1 \times M_2$ and $N' = M_1 \times N_2$. We suppose that $(N_i : M_i) \cap S = \emptyset$, where i = 1, 2.

- (1) N_1 is an S-1-absorbing primary submodule of M_1 if and only if N is an $S \times \{1\}$ -1-absorbing primary submodule of M.
- (2) N_2 is an S-1-absorbing primary submodule of M_2 if and only if N' is a $\{1\} \times S$ -1-absorbing primary submodule of M.

Proof. (1) Suppose that N is an S-1-absorbing primary submodule of M, let (s, 1) be $S \times \{1\}$ -element of N. Let a, b be non-nunit elements of R and $m_1 \in M_1$ with $abm_1 \in N_1$, so $(a, 1)(b, 1)(m_1, 0) = (abm_1, 0) \in N$. Hence $(s, 1)(a, 1)(b, 1) \in (N_1 \times M_2 : M_1 \times M_2)$ or $(s, 1)(m_1, 0) \in M$ -rad $(N_1 \times M_2)$. If $(s, 1)(a, 1)(b, 1) \in (N_1 \times M_2 : M_1 \times M_2)$, it is clear that $sab \in (N_1 : M_1)$. Now suppose $(s, 1)(m_1, 0) \in M$ -rad $(N_1 \times M_2)$. By [5, Lemma 2.10], we have $(s, 1)(m_1, 0) \in M_1$ -rad $(N_1) \times M_2$, and thus $sm_1 \in M_1$ -rad (N_1) . We conclude that N_1 is an S-1-absorbing primary submodule. Now assume that N_1 is an S-1-absorbing primary submodule. Now assume that N_1 is an S-1-absorbing primary submodule of M_1 and let $s \in S$ be an S-element of N_1 . Let (a, a'), (b, b') be non-units of $R \times R$ and $(m, m') \in M$ with $(a, a')(b, b')(m, m') \in N$. Then $abm \in N_1$ and so either $sab \in (N_1 : M_1)$ or $sm_1 \in M_1$ -rad(N). If $sab \in N_1$, we conclude that $(s, 1)(a, a')(b, b') = (sab, a'b') \in (N_1 \times M_2 : M_1 \times M_2)$, now if $sab \in M_1$ -rad (N_1) , we get $(s, 1)(a, a')(m, m') = (sam, sa'm') \in M_1$ -rad $(N_1) \times M_2 = M$ -rad $(N_1 \times M_2)$ by [5, Lemma 2.10]. Hence, N is an $S \times \{1\}$ -1-absorbing primary submodule of M.

(2) The similar proof of (1).

Lemma 5. Let N be a submodule of M and S be a multiplicatively closed subset of R. If $(P : M) \cap S = \emptyset$ for all prime submodule P containing N, then (M - rad(N) : s) = M - rad(N) = M - rad(N : s) for any $s \in S$.

Proof. The inclusions M-rad $(N) \subseteq M$ -rad(N : s) and M-rad $(N) \subseteq (M$ -rad(N) : s) are clear. Let $m \in (N : s)$ and suppose that P is a prime submodule of M containing N. Then $sm \in N \subseteq P$ implies directly $m \in P$ due to our assumption. Thus, $(N : s) \subseteq P$ and we conclude M-rad(N) = M-rad(N : s). Now, let $m \in (M$ -rad(N) : s). Then $sm \in M$ -rad(N) which means that $sm \in P$ for all prime submodules P containing N. Again we have $m \in P$ by our assumption. It follows that $m \in M$ -rad(N) and consequently we get (M-rad $(N) : s) \subseteq M$ -rad(N). \Box

In the following proposition, we denote $Z_I(R)$ by the set $\{x \in R \mid xy \in I \text{ for some } y \in R \setminus I\}$.

Proposition 8. Let N be a submodule of M and S be a multiplicatively closed subset of R provided $(P: M) \cap S = \emptyset$ for all prime submodule P containing N. If $Z_{(N:M)}(R) \cap S = \emptyset$, then (N:s) is a 1-absorbing primary submodule of M for all $s \in S$.

Proof. Suppose that $abm \in (N:s)$ for some non-unit $a, b \in R$ and $m \in M$. Hence, $sabm \in M$ and there exists $s' \in S$ such that either $s'sab \in (N:M)$ or $s'm \in M$ rad(N). If $s'sab \in (N:M)$, then we conclude $ab \in (N:M) \subseteq ((N:s):M)$ since $Z_{(N:M)}(R) \cap S = \emptyset$. Now, assume that $s'm \in M$ -rad(N). Then we have $m \in (M$ rad(N):s') = M-rad(N) = M-rad(N:s) by Lemma 5, so we are done. \Box

3. S-1-Absorbing Primary Avoidance Theorem

In this section, S-1-absorbing primary avoidance theorem is proved. Throughout this section, let M be a finitely generated multiplication R-module, N, N_1, N_2, \ldots, N_n be submodules of M and S be a multiplicatively closed subset of R. Recall that a covering $N \subseteq N_1 \cup N_2 \cup \cdots \cup N_n$ is said to be efficient if no N_k is unnecessary. Besides, if none of the N_k may be excluded, the union $N = N_1 \cup N_2 \cup \cdots \cup N_n$ is efficient [20]. We first state some theorems that we will require.

Lemma 6. ([18], Corollary 1) Let R be a ring, $S \subseteq R$ a multiplicatively closed set and P an ideal of R disjoint with S. Then P is S-prime if and only if there exists $s \in S$, such that for all I_1, \ldots, I_n ideals of R, if $I_1 \ldots I_n \subseteq P$, then $sI_j \in P$ for some $j \in \{1, \ldots, n\}$.

Lemma 7. ([18], Proposition 4) Let R be a ring, $S \subseteq R$ a multiplicatively closed set and P an ideal of R disjoint with S. Then P is S-prime if and only if there exists $s \in S$ such that for all $x_1, \ldots, x_n \in R$, if $x_1 \ldots x_n \in P$, then $sx_j \in P$ for some $j \in \{1, \ldots, n\}$.

Theorem 9. Let $N \subseteq N_1 \cup N_2 \cup \cdots \cup N_n$ be an efficient covering of submodules N_1, N_2, \ldots, N_n of M where n > 2. Assume that $\sqrt{(N_k : m)}$ is an ideal of R disjoint with S for all $1 \le k \le n$ and all $m \in M \setminus P$ where P is any prime submodule of M containing N_k . If $\sqrt{(N_j : M)} \nsubseteq (\sqrt{(N_k : m)} : s)$ for all $s \in S$ whenever $j \ne k$, then no N_k is an S-1-absorbing primary submodule of M.

Proof. It can be easily seen that $N = (N \cap N_1) \bigcup (N \cap N_2) \bigcup \cdots \bigcup (N \cap N_n)$ is an efficient union since $N \subseteq N_1 \cup N_2 \cup \cdots \cup N_n$ is efficient. Thus, there exists an element $m_k \in N \setminus N_k$ for all $k \leq n$. From ([20], Lemma 2.2), it is known that $\bigcap_{i\neq k} (N \cap N_j) \subseteq N \cap N_k$. Now assume that N_k is an S-1-absorbing primary submodule of M for some $k \in \{1, ..., n\}$. Then, $(N_k : m)$ is an S-1-absorbing primary ideal of R by Lemma 4. Furthermore, by Lemma 1, $\sqrt{(N_k:m)}$ is an Sprime ideal of R. By our hypothesis, $s\sqrt{(N_j:M)} \not\subseteq (\sqrt{(N_k:m)})$ and then there exists a non-unit element $a_j \in \sqrt{(N_j : M)}$ such that $sa_j \notin \sqrt{(N_k : m)}$ for all $s \in S$ and all $m \in M \setminus N_k$. (If a_j is unit, then $\sqrt{(N_j : M)} = R$ and hence $N_j = M$ which contradicts with our assumption that the covering is efficient.) So, there exists a positive integer n_j such that $a_j^{n_j} \in (N_j : M)$ where $j \neq k$. From Lemma 6, we also have $\prod_{j \neq k} \sqrt{(N_j:M)} \notin \sqrt{(N_k:m)}$ due to $\sqrt{(N_k:m)}$ is S-prime. Now let us say $a = \prod_{j \neq k} a_j$ and $t = max\{n_j\}_{j \neq k}$. Then we get $a^t \in (N_j : M)$ and it follows that $a^t m_k \in N \cap N_j$ for every $j \neq k$. However, $a \notin \sqrt{(N_k : m)}$ by Lemma 7 and this implies that $a^t m \notin N_k$ for all $m \in M \setminus N_k$. As a result, $a^t m_k \notin N_k$ because $m_k \notin N_k$ as well. This shows that $a^t m_k \in \bigcap_{i \neq k} (N \cap N_j) \setminus N \cap N_k$, which is a contradiction.

Theorem 10 (S-1-absorbing primary avoidance theorem). Let N, N_1, N_2, \ldots, N_n be submodules of M such that $N \subseteq N_1 \cup N_2 \cup \cdots \cup N_n$ where at most two of N_1, N_2, \ldots, N_n are not S-1-absorbing primary submodules. Suppose that $\sqrt{(N_k : m)} \cap S = \emptyset$ for all $m \in M \setminus N_k$ where $1 \le k \le n$. If $\sqrt{(N_j : M)} \nsubseteq (\sqrt{(N_k : m)} : s)$ for all $s \in S$ whenever $j \ne k$, then $N \subseteq N_k$ for some k.

Proof. We can assume that the covering $N \subseteq N_1 \cup N_2 \cup \cdots \cup N_n$ is efficient because any cover including submodules of M can be reduced to an efficient covering by eliminating any superfluous terms and therefore $n \neq 2$. Since there is no S-1absorbing primary submodule N_k from Theorem 9, n < 2. Hence we obtain $N \subseteq N_k$ for some $k \in \{1, \ldots, n\}$.

Corollary 3. Let N be a submodule of M. If the S-1-absorbing primary avoidance theorem holds for M, then the theorem holds also for M/N.

Proof. Let $K/N, N_1/N, N_2/N, ..., N_n/N$ be submodules of M/N such that at most two of $N_1/N, N_2/N, ..., N_n/N$ are not S-1-absorbing primary and $K/N \subseteq N_1/N \cap$ $N_2/N \cap \cdots \cap N_n/N$ where $n \geq 2$. Thus, $K \subseteq N_1 \cap N_2 \cap \cdots \cap N_n$ and at most two of $N_1, N_2, ..., N_n$ are not S-1-absorbing primary by Corollary 1. Assume that $\sqrt{N_j/N : M/N} \notin (\sqrt{N_k/N : m + N} : s)$ for all $s \in S$ and all $m + N \in M/N \setminus N_k/N$ whenever $j \neq k$. Then $\sqrt{(N_j : M)} \notin (\sqrt{(N_k : m)} : s)$ for all $s \in S$. Since the S-1absorbing primary avoidance theorem is satisfied by the submodules of M, the rest of the proof is immediate. □

4. Idealization of S-1-absorbing primary submodules

Let M be a unitary R-module. Recall from [23] the idealization R(+)M of M is the direct sum $R(+)M = R \oplus M$ which is a commutative ring with componentwise addition and multiplication $(r_1, m_1)(r_2, m_2) = (r_1r_2, r_1m_2 + r_2m_1)$. It is well known that if N is a submodule of M with $IM \subset N$, then I(+)N is an ideal of R(+)M.

Proposition 9. Let S be a multiplicatively closed subset of R, I be an ideal of R with $I \cap S = \emptyset$. Then the following statements are equivalent:

- (1) I is an S-1-absorbing primary ideal of R.
- (2) I(+)M is an S(+)0-1-absorbing primary ideal of R(+)M.
- (3) I(+)M is an S(+)M-1-absorbing primary ideal of R(+)M.

Proof. (1)⇒ (2) It is clear that $(I(+)N) \cap (S(+)0) = \emptyset$ if and only if $S \cap I = \emptyset$. Suppose that *I* is an *S*-1-absorbing primary ideal of *R* and *s* be an *S*-element of *I*. Let $(a,m)(b,n)(c,p) \in I(+)M$ where (a,m), (b,n), (c,p) are non-unit elements of R(+)M. Since U(R(+)M) = U(R)(+)M, we conclude that a, b, c are non-units in *R* and $abc \in I$. Then either $sab \in I$ or $sc \in \sqrt{I}$. If $sab \in I$, then $(s,0)(a,m)(b,n) = (sab, san + sbm) \in I(+)M$. If $sc \in \sqrt{I}$, then $(s,0)(c,p) = (sc,0) \in \sqrt{I(+)M}$ due to $\sqrt{I(+)M} = \sqrt{I}(+)M$. Thus I(+)M is an S(+)0-1-absorbing primary ideal of R(+)M.

 $(2) \Rightarrow (3)$ Is clear since $S(+)0 \subset S(+)M$.

 $(3) \Rightarrow (1)$ Assume that I(+)M is an S(+)M-1-absorbing primary ideal of R(+)M. Let (s,m) be an S(+)M-element of I(+)M and let $abc \in I$ for non-unit elements a, b, c of R. Then (a, 0), (b, 0), (c, 0) are also non-unit elements of R(+)M since U(R(+)M) = U(R)(+)M. Besides, we have $(a, 0)(b, 0)(c, 0) \in I(+)M$. So, either $(s,m)(a, 0)(b, 0) = (sab, abm) \in I(+)M$ or $(s,m)(c, 0) \in \sqrt{I(+)M}$ which yields $sab \in I$ or $sc\sqrt{I}$. Hence I is an S(+)M-1-absorbing primary ideal of R. \Box **Proposition 10.** Let S be a multiplicatively closed subset of R and I be an ideal of R with $I \cap S = \emptyset$. Suppose N is a proper submodule of M such that $IM \subset N$. Then $(1) \Longrightarrow (2) \Longrightarrow (3)$ is satisfied for the below statements:

- (1) I(+)N is an S(+)0-1-absorbing primary ideal of R(+)M.
- (2) I(+)N is an S(+)M-1-absorbing primary ideal of R(+)M.
- (3) I is an S-1-absorbing primary ideal of R.

Proof. (1) \Longrightarrow (2) It is clear since $S(+)0 \subset S(+)M$.

 $(2) \Longrightarrow (3)$ Let $abc \in I$ for non-unit elements $a, b, c \in R$. We have $(a, 0)(b, 0)(c, 0) \in I(+)N$ which is S(+)M-1-absorbing primary ideal of R(+)M where (a, 0), (b, 0), (c, 0) are non-units in R(+)M. Then either

$$(s,m)(a,0)(b,0) = (sab,abm) \in I(+)N$$

or

$$(s,m)(c,0) = (sv,cm) \in \sqrt{I(+)N} = \sqrt{I}(+)M.$$

where (s,m) is an S(+)M-element of I(+)N. Then either $sab \in I$ or $sc \in \sqrt{I}$. Hence I is an S-1-absorbing primary ideal of R.

Example 3. The implication $(3) \implies (1)$ is not true in general case. If I is an S-1-absorbing primary ideal, then I(+)N (where N a proper submodule of M) is not necessarily S(+)0-1-absorbing primary ideal of R(+)M. Take $R = \mathbb{Z} = M$ and $S = \{\mp 1\}$. Then $I = \{0\}$ is an S-1-absorbing primary ideal of \mathbb{Z} , since $(0(+)6\mathbb{Z} : (0,1)) = 6\mathbb{Z}(+)\mathbb{Z}$ is not an S(+)0-prime ideal. We conclude that $0(+)6\mathbb{Z}$ is not an S(+)0-1-absorbing primary ideal of $\mathbb{Z}(+)\mathbb{Z}$ by Lemma 1.

5. Amalgamation of S-1-absorbing primary submodules

Let A and B be commutative rings with unity, let J be an ideal of B and let $f: A \to B$ be a ring homomorphism. In this setting, we can consider the following subring of $A \times B$:

$$A \bowtie^{f} J := \{(a, f(a) + j) \mid a \in A, j \in J\}$$

called the *amalgamation of* A with B along J with respect to f. This construction is a generalization of the amalgamated duplication of a ring along an ideal (cf., for instance [6, 8, 9, 10, 11, 12]).

It is clear that a subset S of A is a multiplicatively closed subset of A if and only if $S \bowtie^f J := \{(s, f(s) + j) \mid s \in S; j \in J\}$ is a multiplicatively closed subset of $A \bowtie^f J$.

Lemma 8. If I is an ideal of A, then

$$\sqrt{I \bowtie^f J} = \sqrt{I} \bowtie^f J.$$

Proof. Let $(a, f(a) + j) \in A \bowtie^f J$ such that $(a, f(a) + j) \in \sqrt{I \bowtie^f J}$. Then there exists a nonzero integer n such that $(a, f(a) + j)^n = (a^n, f(a^n) + j') \in I \bowtie^f J$ where $j' = \sum_{k=0}^{n-1} \binom{n}{k} f(a^k) j^{n-k} \in J$. Then $a^n \in I$, so $a \in \sqrt{I}$ and $(a, f(a) + j) \in \sqrt{I} \bowtie^f J$. Hence, we conclude that $\sqrt{I \bowtie^f J} \subset \sqrt{I} \bowtie^f J$. Conversely, let $(a, f(a) + j) \in \sqrt{I} \bowtie^f J$. Then there exists $n \in \mathbb{N}^*$ such that $a^n \in I$. On the other

hand $(a, f(a) + j)^n = (a^n, f(a^n) + j')$. Then $(a, f(a) + j) \in \sqrt{I \bowtie^f J}$, which yields $\sqrt{I} \bowtie^f J \subset \sqrt{I \bowtie^f J}$. Finally, we conclude $\sqrt{I \bowtie^f J} = \sqrt{I} \bowtie^f J$. \Box

Proposition 11. Let S be a multiplicatively closed subset and I, J be ideals of A. The following statements are equivalent:

- (1) I is an S-1-absorbing primary ideal of A.
- (2) $I \bowtie^f J$ is an $S \bowtie^f J$ -1-absorbing ideal of $A \bowtie^f J$.

Proof. Assume that $I \bowtie^f J$ is an $S \bowtie^f J$ -1-absorbing primary ideal of $A \bowtie^f J$ and let (s, f(s) + j) be an $S \bowtie^f J$ -element of $I \bowtie^f J$. Take $abc \in I$ for non-unit elements a, b, c of A. Then (a, f(a)), (b, f(b)), (c, f(c)) are also non-unit elements of $A \bowtie^f J$ and $(a, f(a))(b, f(b))(c, f(c)) \in I \bowtie^f J$. Since $I \bowtie^f J$ is an $S \bowtie^f J$ -1absorbing primary submodule, we get either $(s, f(s) + j)(a, f(a))(b, f(b)) \in I \bowtie^f J$ or $(s, f(s) + j)(c, f(c)) \in \sqrt{I \bowtie^f J} = \sqrt{I} \bowtie^f J$ by Lemma 8. Then either $sab \in I$ or $sc \in \sqrt{I}$. Thus I is an S-1-absorbing primary ideal of A.

Conversely, suppose that I is an S-1-absorbing primary ideal of A and let $s \in S$ be an S-element of I. Let (a, f(a)+j), (b, f(b)+k), (c, f(c)+l) be non-unit elements of $A \bowtie^f J$ such that $(a, f(a)+j)(b, f(b)+k)(c, f(c)+l) \in I \bowtie^f J$. Then $abc \in I$. It is necessary that one of the a, b, c is non-unit, otherwise, abc is unit and $abc \in I$, a contradiction. Since I is an S-1-absorbing primary submodule of A, we get either $sab \in I$ or $sc \in \sqrt{I}$ which gives either $(s, f(s))(a, f(a) + j)(b, f(b) + k) \in I \bowtie^f J$ or $(s, f(s))(c, f(c) + l) \in \sqrt{I} \bowtie^f J = \sqrt{I \bowtie^f J}$ by Lemma 8. Hence $I \bowtie^f J$ is an $S \bowtie^f J$ -1-absorbing primary ideal of $A \bowtie^f J$.

Corollary 4. Let R be a ring and I, K be ideals of R with $K \cap S = \emptyset$. Then the following statements are equivalent:

- (1) K is an S-1-absorbing primary ideal of R.
- (2) $K \bowtie I$ is an $S \bowtie I$ -1-absorbing primary ideal of $R \bowtie I$.

Proof. It is easy to see that $(K \bowtie I) \cap (S \bowtie I) = \emptyset$. The rest of the proof follows from the previous proposition.

Let S_2 be a multiplicatively closed subset of the ring f(A) + J. Consider the set

$$\overline{S_2}^f := \{ (r, f(r) + j) \mid j \in J, r \in A, f(r) + j \in S_2 \}$$

It is clear that $\overline{S_2}^f$ is a multiplicatively closed subset of $A \bowtie^f J$.

If K is an ideal of f(A) + J, then the set

$$\overline{K}^{f} := \{ (a, f(a) + j) \mid a \in A, j \in J, f(a) + j \in K \}$$

is an ideal of $A \bowtie^f J$.

Lemma 9. Under the above notations, we have

$$\sqrt{\overline{K}^f} = \overline{\sqrt{K}}^f.$$

Proposition 12. Let $f : A \longrightarrow B$ be a ring homomorphism, J, K be ideals of B and S_2 be a multiplicatively closed subset of f(A) + J with $K \cap S_2 = \emptyset$. Let

$$\overline{K}^{f} = \{(a, f(a) + j) \mid a \in A, j \in J, f(a) + j \in K\}.$$

Then the following statements are equivalent:

(1) K is an S₂-1-absorbing primary ideal of f(A) + J.

(2) \overline{K}^f is an \overline{S}_2^f -1-absorbing primary ideal of $A \bowtie^f J$.

Proof. Assume that \overline{K}^f is an \overline{S}^f -1-absorbing primary ideal of $A \bowtie^f J$ and let (s, f(s) + p) be an \overline{S}^{f} -element of \overline{K}^{f} . Let $(f(a) + j)(f(b) + k)(f(c) + l) \in K$ for non-unit elements f(a) + j, f(b) + k, $f(c) + l \in f(A) + J$. Then (a, f(a) + j), (b, f(b) + j)k), (c, f(c)+l) are non-unit elements of $A \bowtie^f J$ and (a, f(a)+j)(b, f(b)+k)(c, f(c)+k)(c, f(c $l \in \overline{K}^{f}$. Thus, we have either $(s, f(s)+p)(a, f(a)+j)(b, f(b)+k) \in \overline{K}^{f}$ or $(s, f(s)+p)(a, f(a)+j)(b, f(b)+k) \in \overline{K}^{f}$ $p)(c, f(c)+l) \in \sqrt{\overline{K}^f}$. In the first case, we have $(f(s)+p)(f(a)+j)(f(b)+k) \in K$. In the second case, we have $(f(s)+p)(f(c)+k) \in \sqrt{\overline{K}^f} = \overline{\sqrt{K}^f}$ by Lemma 9 and hence $(f(s)+p)(f(c)+k) \in \sqrt{K}$. We conclude that K is an S₂-1-absorbing primary ideal of B. Conversely, suppose that K is an S_2 -1-absorbing primary ideal of B and let f(s)+p an S_2 -element of K. Let $(a, f(a)+j)(b, f(b)+k)(c, f(c)+l) \in \overline{K}^f$ for non-unit elements $(a, f(a) + j), (b, f(b) + k), (c, f(c) + l) \in A \bowtie^f J$. Then we get $(abc, (f(a) + b)) \in A \bowtie^f J$. $j)(f(b)+k)(f(c)+l)) \in \overline{K}^{f}$. Now suppose that f(a)+j, f(b)+k, f(c)+l are non-unit elements of f(A) + J. Since K is an S₂-1-absorbing primary ideal of f(A) + J, we conclude that either $(f(s)+p)(f(a)+j)(f(b)+k) \in K$ or $(f(s)+p)(f(c)+l) \in \sqrt{K}$. In the first case, we have $(s, f(s) + p)(a, f(a) + j)(b, f(b) + k) \in \overline{K}^{f}$. In the second case, we have $(s, f(s)+p)(c, f(c)+l) \in \sqrt{\overline{K}^f}$. Now, if one of f(a)+j, f(b)+k, f(c)+lis unit, then the claim is clear. Hence \overline{K}^f is an $\overline{S_2}^f$ -1-absorbing primary ideal of $A \bowtie^f J.$ \square

An *R*-module can be regarded as an $R \bowtie I$ -module via the natural projection $R \bowtie I \longrightarrow R; (r, r + i) \longmapsto r$, and an $R \bowtie I$ -module can also be regarded as an *R*-module via the natural injection $\iota : R \longrightarrow R \bowtie I; r \longmapsto (r, r)$.

Let R be a ring, I an ideal of R, M an R-module and set

$$M \bowtie I := \{ (m, m') \in M \times M \mid m - m' \in IM \},\$$

which is an $R \bowtie I$ -module with the multiplication given by

$$(r, r+i)(m, m') = (rm, (r+i)m')$$

where $r \in R$, $i \in I$ and $(m, m') \in M \bowtie I$. Then the set $M \bowtie I$ is called the *duplication of the R-module M along the ideal I*.

Let N be a submodule of M. It is clear that

$$N \bowtie I : \{(n,m) \in N \times M \mid n-m \in IM\}$$

and

$$N := \{ (m, n) \in M \times N \mid n - m \in IM \}$$

are submodules of $M \bowtie I$. If S is a multiplicatively closed subset of R, then it is obvious that the sets $S \bowtie I := \{(s, s+i) \mid s \in S, i \in I\}$ and $\overline{S} := \{(s, s+i) \mid i \in I, s+i \in S\}$ are multiplicatively closed subsets of $R \bowtie I$.

Lemma 10. Let N be a submodule of M. Then

 $(N \bowtie I :_{R \bowtie I} M \bowtie I) = (N :_R M) \bowtie I.$

Proof. Let $(a, a + i) \in (N \bowtie I :_{R \bowtie I} M \bowtie I)$. Then $(a, a + i)(M \bowtie I) \subset N \bowtie I$. Clearly, $aM \subset N$ and thus $(a, a + i) \in (N :_R M) \bowtie I$. Conversely, suppose that $(a, a + i) \in (N :_R M) \bowtie I$ and take $(m, m') \in M \bowtie I$. We have (a, a + i)(m, m') = $(am, (a+i)m') \in M \times M$ and $a(m-m') - im' \in IM$ since $m-m' \in IM$. Hence $(a, a+i) \in (N \bowtie I :_{R\bowtie I} M \bowtie I)$. Thus, we conclude that $(N \bowtie I :_{R\bowtie I} M \bowtie I) = (N :_{R} M) \bowtie I$.

Lemma 11. Let L be a submodule of M. Then the following statements are equivalent:

- (1) L is a prime submodule of M.
- (2) $L \bowtie I$ is a prime submodule of $M \bowtie I$.

Proof. Assume that $L \bowtie I$ is a prime submodule of $M \bowtie I$. Let $am \in L$ for $a \in R$ and $m \in M$. Then $(a, a)(m, m) \in L \bowtie I$. Since $L \bowtie I$ is a prime submodule of $M \bowtie I$, we get either $(a, a) \in (L \bowtie I :_{R\bowtie I} M \bowtie I)$ which implies $a \in (N : M)$ by Lemma 10, or $(m, m) \in L \bowtie I$ which follows $m \in L$. Hence L is a prime submodule of M. Conversely, assume that L is a prime submodule of M and let $(a, a + i)(m, m') \in L \bowtie I$ for $(a, a + i) \in R \bowtie I$ and $(m, m') \in M \bowtie I$. Thus $(am, (a + i)m') \in L \bowtie I$ and we conclude that $am \in L$. Due to the fact that L is a prime submodule, we obtain either $a \in (N : M)$ or $m \in L$. In the first case, we have $(a, a + i) \in (N : M) \bowtie I = (N \bowtie I :_{R\bowtie I} M \bowtie I)$. In the second case, we have $(m, m') \in L \bowtie I$, since $m - m' \in IM$. Thus, $L \bowtie I$ is a prime submodule of $M \bowtie I$.

Lemma 12. Let N be a submodule of M. Then the prime submodule which contains $N \bowtie I$ are exactly of the form $L \bowtie I$ where L is a prime submodule containing N. Moreover, $M \bowtie I \operatorname{-rad}(N \bowtie I) = M \operatorname{-rad}(N) \bowtie I$.

Proof. Let $\varphi : M \bowtie I \longrightarrow M; (m, m') \longmapsto m$ be an $R \bowtie I$ -epimorphism with ker $\varphi = 0 \times IM$. Then $M \bowtie I/0 \times IM \simeq M$. So, if L is a submodule of $M \bowtie I$ such that $0 \times IM \subset I$, then there exists a submodule K of M such that $L/0 \times IM \simeq K$, which implies $L = K \bowtie I$. From Lemma 11, we conclude that L is a prime submodule of $M \bowtie I$ if and only if K is a prime submodule of M. Now, if L is a prime submodule of $M \bowtie I$ such that $N \bowtie I \subset L$, then $0 \times IM \subset L$, and so there exists a prime submodule K of M with $L = K \bowtie I$.

Now, it is clear that $\bigcap_i (K_i \bowtie I) = (\bigcap_i K_i) \bowtie I$ where K_i 's are the submodules of M and consequently we have $M \bowtie I \operatorname{-rad}(N \bowtie I) = M \operatorname{-rad}(N) \bowtie I$. \Box

Theorem 11. Let N be a submodule of M with $(N :_R M) \cap S = \emptyset$. Then the following statements are equivalent:

- (1) N is an S-1-absorbing primary submodule of M.
- (2) $N \bowtie I$ is an $S \bowtie I$ -1-absorbing primary submodule of $M \bowtie I$.

Proof. From Lemma 10, we clearly note that $(N \bowtie I :_{R\bowtie I} M \bowtie I) \cap S \bowtie I = \emptyset$ if and only if $(N :_R M) \cap S = \emptyset$. Now suppose that $N \bowtie I$ is an $S \bowtie I$ -1absorbing primary submodule of $M \bowtie I$. Let (s, s + i) be an $S \bowtie I$ -element of $N \bowtie I$ and let $abm \in N$ for non-unit elements $a, b \in R$ and $m \in M$. Then $(a, a)(b, b)(m, m) \in N \bowtie I$. Since (a, a), (b, b) are non-units and $N \bowtie I$ is an $S \bowtie I$ -1-absorbing primary submodule of $M \bowtie I$, we have either

$$(s, s+i)(a, a)(b, b) \in (N \bowtie I : M \bowtie I)$$

or

$$(s, s+i)(m, m) \in M \bowtie I - \operatorname{rad}(N \bowtie I).$$

In the first case, we have $sab \in (N:M)$ by Lemma 10. In the second case, $M \bowtie I$ - $rad(N \bowtie I) = M$ - $rad(N) \bowtie I$ by Lemma 12 and we conclude that $(sm, (s+i)m) \in M$ - $rad(N) \bowtie I$, and so $sm \in M$ -rad(N). If such an N does not exists, it is well known that M-rad(N) = M. Then N is an S-1-absorbing primary submodule of M. Conversely, assume that N is an S-absorbing primary submodule of M. Let $s \in S$ be an S-element of N and let $(a, a+i)(b, b+i)(m, m') \in N \bowtie I$ for non-unit elements $(a, a + i), (b, b + i) \in R \bowtie I$ and $(m, m') \in M \bowtie I$. Then $abm \in N$. If a or b is unit, the claim is clear. Now we can assume a and b are non-unit elements of R. Due to the fact that N is an S-1-absorbing primary submodule of R, we get either $sab \in (N : M)$ or $sm \in M$ -rad(N). In the first case, we conclude that $(s, s)(a, a + i)(b, b + i) \in (N : M) \bowtie I = (N \bowtie I : M \bowtie I)$. In the second case, we have $(sm, sm' + im') \in M$ - $rad(N) \bowtie I$, since $r(m - m') - im' \in IM$. Thus, the claim follows by M- $rad(N) \bowtie I = M \bowtie I$ - $rad(N \bowtie I)$.

References

- Anderson DD, Arabaci T, Tekir U and Koc S (2020) On S-multiplication modules. Comm. Algebra 48(8): 3398-3407
- [2] Anderson DD, Winders M (2009) Idealization of a module. J. Commut. Algebra 1(1): 3-56
- [3] Ansari-Toroghy H, Pourmortazavi SS (2022), On S-primary Submodules. Int. Electron. J. Algebra 31(31): 74-89
- [4] Atani SE, Tekir U (2006) On the primary avoidance theorem for modules over commutative rings. Int J Pure Appl Math. 31(2): 203-207
- [5] Atani SE and Khalil Saraei FE (2007) Modules which Satisfy the Radical Formula. Int. J. Contemp. Math. Sci. 2(1): 13-18
- [6] Aung PP (2016) Gorenstein dimensions over some rings the form $R \oplus C$. J. Algebra Appl. 15(3): 1650043
- [7] Badawi A, Yetkin Celikel E (2020) On 1-absorbing primary ideals of commutative rings. J. Algebra Its Appl. 19(06): 2050111
- [8] Bagheri S, Salimi M, Tavasoli E, Yassemi S (2012) A construction of quasi-Gorenstein rings. J. Algebra. Appl. 11: 12500131-1-1250013-9
- [9] Bouba F, Mahdou N, Tamekkante M (2018) Duplication of a module along an ideal. Acta Math. Hungar. 154(1): 29-42
- [10] Chhiti M, Jerrar M, Kabbaj S, Mahdou N (2015) Prüfer conditions in an amalgamated duplication of a ring along an ideal. Comm. Algebra 41: 249-261
- [11] D'anna M, Fontana M (2007) An amalgamated duplication of a ring along an ideal: the basic properties. J. Algebra Appl. 6: 443-459
- [12] D'anna M, Finocchiaro CA, Fontana M (2016) New algebraic properties of an amalgamated algebra along an ideal. Comm. Alg. 44: 1836-1851
- [13] Dubey MK, Sarohe P (2017) On prime and primary avoidance theorem for subsemimodules. Quasigr. Relat. Syst. 25: 221-229
- [14] El-Bast ZA and Smith PF (1988) Multiplication Modules. Comm. Algebra 16(4): 755-779
- [15] El Khalfaoui R, Mahdou N, Sahandi P, Shirmohammadi N (2021) Amalgamated modules along an ideal. Commun. Korean Math. Soc. 36: 1-10
- [16] Farzalipour F and Ghiasvand P (2021) On S-1-absorbing prime submodules. J. Algebra Appl. https://doi.org/10.1142/S0219498822501158
- [17] Gilmer R (1968) Multiplicative Ideal Theory. Queen's Papers in Pure and Appl. Math.
- [18] Hamed A and Malek A (2020) S-prime ideals of a commutative ring. Beitr Algebra Geom, 61: 533–542
- [19] Lu CP (1989) M-radicals of submodules in modules. Math. Japonica, 34(2): 211-219
- [20] Lu CP (1997) Unions of prime submodules. Houston J. Math, 23 (2): 203-213
- [21] Moghimi HF, Rashedi F, Samiei M (2014) A Note on primary-like submodules of multiplication modules. J. Algebra Relat. Top. 2(2): 37-41

MOHAMMED ISSOUAL, NAJIB MAHDOU, NESLIHAN AYSEN OZKIRISCI, AND ECE YETKIN CELIKEL

- [22] Mostafanasab H, Yetkin Celikel E, Tekir U and Darani AY (2016) On 2-absorbing primary submodules of modules over commutative rings. An. St. Univ. Ovidius Constanta 24(1): 335-351
- [23] Nagata M (1962) Local rings. Interscience Publishers, New York.
- [24] Naji OA (2021) On S-2-absorbing primary submodules. Filomat 35(5): 1477-1487
- [25] Salimi M, Tavasoli E, Yassemi S (2013) The amalgamated duplication along a semidualizing ideal. Rendiconti del Semin. Mat. dell Univ. di Padovo 129: 115-127
- [26] Sevim ES, Arabaci T, Tekir U and Koc S (2019) On S-prime submodules. Turk. J. Math. 43(2): 1036-1046
- [27] Yetkin Celikel E (2021) 1-absorbing primary submodules. Analele Stiint. ale Univ. Ovidius Constanta Ser. Mat. 29(3): 285-296

CRMEF MEKNES, MOROCCO. Email address: issoual2@yahoo.fr

LABORATORY OF MODELING AND MATHEMATICAL STRUCTURES, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY OF FEZ, BOX 2202, UNIVERSITY S.M. BEN ABDELLAH FEZ, MOROCCO.

Email address: mahdou@hotmail.com

DEPARTMENT OF MATHEMATICS, YILDIZ TECHNICAL UNIVERSITY, DAVUTPASA CAMPUS, ESENLER, 34210, ISTANBUL, TURKEY

Email address: aozk@yildiz.edu.tr

DEPARTMENT OF ELECTRICAL-ELECTRONICS ENGINEERING, FACULTY OF ENGINEERING, HASAN KALYONCU UNIVERSITY, GAZIANTEP, TURKEY

Email address: ece.celikel@hku.edu.tr, yetkinece@gmail.com