

THE ONE-ROW COLORED \mathfrak{sl}_3 JONES POLYNOMIALS FOR PRETZEL LINKS

KOTARO KAWASOE

ABSTRACT. The colored \mathfrak{sl}_3 Jones polynomial $J_{(n_1, n_2)}^{\mathfrak{sl}_3}(L; q)$ are given by a link and an (n_1, n_2) -irreducible representation of \mathfrak{sl}_3 . In general, it is hard to calculate $J_{(n_1, n_2)}^{\mathfrak{sl}_3}(L; q)$ for an oriented link L . However, we calculate the one-row \mathfrak{sl}_3 colored Jones polynomials $J_{(n, 0)}^{\mathfrak{sl}_3}(P(\alpha, \beta, \gamma); q)$ for three-parameter families of oriented pretzel links $P(\alpha, \beta, \gamma)$ by using Kuperberg's linear skein theory by setting $n_2 = 0$. Furthermore, we show the existence of the tails of $J_{(n, 0)}^{\mathfrak{sl}_3}(P(2\alpha + 1, 2\beta + 1, 2\gamma); q)$ for the alternating pretzel knots $P(2\alpha + 1, 2\beta + 1, 2\gamma)$.

1. INTRODUCTION

The quantum invariant of a link is obtained by Lie algebra \mathfrak{g} and its irreducible representation. In particular, if $\mathfrak{g} = \mathfrak{sl}_2$ and $V = \mathbb{C}^2$, we call the quantum invariant of a link Jones polynomial [Jon85]. Kauffman [Kau87] reformulated the Jones polynomial of a link L by using Kauffman bracket skein relation. Furthermore, as a generalization of the Jones polynomial, we constitute the colored \mathfrak{sl}_2 Jones polynomial $J_N^{\mathfrak{sl}_2}(L; q)$ by an $(N + 1)$ -dimensional irreducible representation of \mathfrak{sl}_2 . For a link L , the graphical techniques with Kauffman bracket and Jones-Wenzel projector [Wen87] for a link diagram give $J_N^{\mathfrak{sl}_2}(L; q)$. Jones-Wenzel projector is an element of A_1 web space and plays a crucial role in $J_N^{\mathfrak{sl}_2}(L; q)$. Linear skein theory is this method of calculating $J_N^{\mathfrak{sl}_2}(L; q)$. There are many examples of $J_N^{\mathfrak{sl}_2}(L; q)$ calculated by using linear skein theory. We can see one example in [Lic97]. The explicit formulae in $J_N^{\mathfrak{sl}_2}(L; q)$ are helpful when we consider the property that there exists the limit of the colored Jones polynomial called the tail, and conjectures related to quantum invariants of links such as the volume conjecture [Kas97], the Jones slope conjecture [Gar11], Etc.

As regards the tail of links, first, Dasbach and Lin [DL06] showed that the first two coefficients and the last two coefficients of $J_{N+1}^{\mathfrak{sl}_2}(K; q)$ do not depend on N for alternating knots K . They also showed that the third and the third to last coefficients of $J_N^{\mathfrak{sl}_2}(K; q)$ on N for alternating knots do not depend on N provided $N \geq 3$. This result led them to predict that the coefficients of $J_N^{\mathfrak{sl}_2}(K; q)$ up to the k -th do not depend on N provided $N \geq k$; for alternating knots K , there exists a q series $T^{\mathfrak{sl}_2}(K; q)$ such that

$$T^{\mathfrak{sl}_2}(K; q) - \hat{J}_N^{\mathfrak{sl}_2}(K; q) \in q^{N+1} \mathbb{Z}[[q]]$$

where $\hat{J}_N^{\mathfrak{sl}_2}(L; q)$ is normalization with the minimum degree of $J_N^{\mathfrak{sl}_2}(L; q)$ and $T^{\mathfrak{sl}_2}(L; q)$ is called the tails of $J_N^{\mathfrak{sl}_2}(K; q)$. Armond [Arm13] showed the existence of the tails of the colored Jones \mathfrak{sl}_3 polynomials for adequate knots containing alternating knots. Garoufalidis and Lê [GL15] gave proof of the existence of more general stability of coefficients of $J_N^{\mathfrak{sl}_2}(K; q)$ for alternating knots. Armond and Dasbach [AD11] gave explicit tails for $(2, 2m+1)$ -torus knots and $(2, 2m)$ -torus links, Elhamdadi, Hajij [EH17], and Beirne [Bei19] gave explicit tails for pretzel knots, Osburn, Kcilthy and Beirne [KO17] [BO17] gave explicit tails for knots with small crossing numbers.

Meanwhile, the colored \mathfrak{sl}_3 Jones polynomial $J_{(n_1, n_2)}^{\mathfrak{sl}_3}(L; q)$ is obtained by a link and an (n_1, n_2) -irreducible representation of \mathfrak{sl}_3 . However, due to the complexity of the calculation, the explicit formulae for the colored \mathfrak{sl}_3 Jones polynomial only exists for trefoil knot [Law03], $(2, 2m+1)$ -torus knots [GV17][GMV13], two bridge links [Yua17], and $(2, 2m)$ -torus links [Yua17][Yua18a][Yua21a]. For trefoil knot and $(2, 2m+1)$ -torus knot, the explicit formulae are given by Kuperberg's linear skein theory [Kup96].

1)-torus knots, representation theory methods gave the colored \mathfrak{sl}_3 Jones polynomials. In the colored \mathfrak{sl}_3 Jones polynomial, the A_2 bracket is the same role as the Kauffman bracket, and the A_2 clasp is the same role as the Jones-Wenzel projector. For $(2, 2m)$ -torus links and two bridge links, graphical techniques with A_2 bracket and the A_2 claps gave the one-row colored \mathfrak{sl}_3 Jones polynomials obtained by an $(n, 0)$ -irreducible representation of \mathfrak{sl}_3 . Kuperberg's linear skein theory [Kup96] is these graphical techniques. Besides, we can consider the tail concerning $J_{(n_1, n_2)}^{\mathfrak{sl}_3}(L; q)$. Garoufalidis and Young [GV17] showed stability of coefficients of $J_{(n_1, n_2)}^{\mathfrak{sl}_3}(L; q)$ for torus knots. Yuasa [Yua17] [Yua18a] [Yua21a] [Yua21b] gave the explicit tails of $J_{(n, 0)}^{\mathfrak{sl}_3}(L; q)$ for $(2, 2m)$ -torus links and proved that the existence of the tails of the one-row colored \mathfrak{sl}_3 Jones polynomials for minus adequate oriented links.

In this paper, we will give explicitly the one-row \mathfrak{sl}_3 colored Jones polynomials for all three-parameter family of pretzel links $P(\alpha, \beta, \gamma)$ except for those where α, β, γ are all odd. These pretzel links include 8_5 and 8_{19} . 8_5 and 8_{19} are not $(2, 2m)$ -torus link and three-bridge knot. Furthermore, we prove the exsistace of the tail of alternating pretzel knots $P(2\alpha + 1, 2\beta + 1, 2\gamma)$. For instance, the one-row colored \mathfrak{sl}_3 Jones polynomial for $P(3, 3, 2) = 8_5$ multiplied by q^{3n^2+8n} is

$$\begin{aligned} n=1 : & 1 - q + q^2 - 2q^4 + q^5 - 2q^6 + q^7 + q^8 + q^{10} \\ n=2 : & 1 - q + q^3 - 2q^4 + 2q^6 - 2q^7 - 2q^8 + 4q^9 + q^{10} + \dots \\ n=3 : & 1 - q + q^4 + q^5 + q^7 - 4q^8 - q^9 + 8q^{10} + \dots \\ n=4 : & 1 - q - 2q^4 + 2q^5 + q^6 - 2q^8 - 4q^9 + 4q^{10} \dots \\ n=5 : & 1 - q - 2q^4 + q^5 + 2q^6 + q^7 - 3q^8 - 4q^9 + q^{10} + \dots \\ n=6 : & 1 - q - 2q^4 + q^5 + q^6 + 2q^7 - 2q^8 - 2q^9 + 3q^{10} + \dots \\ n=7 : & 1 - q - 2q^4 + q^5 + q^6 + q^7 - q^8 - q^9 + 2q^{10} + \dots \end{aligned}$$

In Appendix, we also give the one-row \mathfrak{sl}_3 colored Jones polynomials for four and five parameter pretzel knots $8_{10}, 8_{15}, 8_{20}$, and 8_{21} . As a result, the one-row \mathfrak{sl}_3 colored Jones polynomials are determined for all knots with eight or fewer crossings except $8_{16}, 8_{17}, 8_{18}$.

This paper is organized as follows. In section 2, we introduce the Kurperberg's linear skein theory and the formulae for the A_2 clasp given by Ohtsuki and Yamada [OY97], Kim [Kim06] [Kim07] and Yuasa [Yua17] [Yua18b]. In section 3, we derive a formula for m times-half twists where two strands with the same directions for Kurperberg's web space. In section 4, we give explicit formulae for the one-row \mathfrak{sl}_3 colored Jones polynomial of three-parameter family of pretzel links $P(\alpha, \beta, \gamma)$. Moreover, for alternating pretzel knots $P(2\alpha + 1, 2\beta + 1, 2\gamma)$, we give the proof that the tail of $J_{(n, 0)}^{\mathfrak{sl}_3}(P(2\alpha + 1, 2\beta + 1, 2\gamma); q)$ exists. In Appendix, we give the one-row \mathfrak{sl}_3 colored Jones Polynomials for $8_{10}, 8_{15}, 8_{20}$ and 8_{21} .

2. PRELIMINARIES

We use the following q-integer notations.

$$\{n\}_q = \{q^{\frac{n}{2}} - q^{-\frac{n}{2}}\}_q, \quad \{n\}_q! = \{n\}_q \{n-1\}_q \cdots \{1\}_q, \quad [n]_q = \frac{\{n\}_q}{\{1\}_q}, \quad [n]_q! = [n]_q [n]_q \cdots [1]_q$$

where n is a non-negative integer. A q-Pochhammer symbol is difined by

$$(q)_k = \prod_{i=1}^k (1 - q^i).$$


A q-binomial coefficient is defined by

$${n \brack k}_q = \frac{[n]_q!}{[k]_q! [n-k]_q!}.$$

Lemma 2.1 ([OY97]). For any integer a, b and c ,

- $$(1) \quad [a]_q[b]_q = \sum_{i=1}^a [a+b-(2i-1)]_q = \sum_{i=1}^b [a+b-(2i-1)]_q,$$
- $$(2) \quad [a]_q[b]_q - [a-c]_q[b-c]_q = [a+b-c]_q[c]_q,$$
- $$(3) \quad [a]_q[b-c]_q + [c]_q[a-c]_q = [b]_q[a-c]_q.$$


We first define the A_2 web spaces. Let D be a disk with signed marked point (P, ϵ) on its boundary where P is a finite set such that $P \subset \partial D$ and $\epsilon : P \rightarrow \{+, -\}$ is a map. A bipartite uni-trivalent graph on D is an immersion of a directed graph such that every vertex is either univalent or trivalent and is divided into sink or source.



A tangled bipartite uni-trivalent graph diagram on D is a bipartite uni-trivalent graph graph on D satisfying:

- (1) the set of univalent vertices coincide with P ,
- (2) every crossing point is a transverse double point of two edges with under or over crossing data.

For tangled uni-trivalent graph diagrams G and G' , we call G regular isotopic to G' on D if G is related to G' by a finite sequence of boundary-fixing isotopies and Reidemeister moves (R1) – (R4) with some directions of edges.



The tangled uni-trivalent graphs on D are regular isotopy classes of tangled trivalent graph diagrams on D . We denote $T(P, \epsilon)$ the set of tangled uni-trivalent graphs on D . The A_2 basis web is the boundary-fixing isotopy class of bipartite uni-trivalent graphs with crossing on D without internal 0, 2, 4-gons. Let $B(P, \epsilon)$ be the set of A_2 basis web. The A_2 web space $W(P, \epsilon)$ is a $\mathbb{Q}(q^{\frac{1}{6}})$ -vector space spanned by $B(P, \epsilon)$. An element in $W(P, \epsilon)$ is called web.

Definition 2.2 (the A_2 bracket [Kup96]). We define a $\mathbb{Q}(q^{\frac{1}{6}})$ linear map $\langle \cdot \rangle_3 : \mathbb{Q}(q^{\frac{1}{6}})T(P, \epsilon) \rightarrow W(P, \epsilon)$ by the following:

$$\begin{aligned} \left\langle \begin{array}{c} \text{circle with two crossing arcs} \\ \text{one arc goes up-right, one down-left} \end{array} \right\rangle_3 &= q^{\frac{1}{3}} \left\langle \begin{array}{c} \text{circle with two arcs} \\ \text{one arc goes up-right, one down-left} \end{array} \right\rangle_3 - q^{-\frac{1}{6}} \left\langle \begin{array}{c} \text{circle with two arcs} \\ \text{one arc goes up-right, one down-left} \end{array} \right\rangle_3, \\ \left\langle \begin{array}{c} \text{circle with two crossing arcs} \\ \text{one arc goes up-left, one down-right} \end{array} \right\rangle_3 &= q^{-\frac{1}{3}} \left\langle \begin{array}{c} \text{circle with two arcs} \\ \text{one arc goes up-left, one down-right} \end{array} \right\rangle_3 - q^{\frac{1}{6}} \left\langle \begin{array}{c} \text{circle with two arcs} \\ \text{one arc goes up-left, one down-right} \end{array} \right\rangle_3, \\ \left\langle \begin{array}{c} \text{circle with four edges} \\ \text{top-left edge goes up-right, top-right down-left, bottom-left up-left, bottom-right down-right} \end{array} \right\rangle_3 &= \left\langle \begin{array}{c} \text{circle with two arcs} \\ \text{one arc goes up-right, one down-left} \end{array} \right\rangle_3 + \left\langle \begin{array}{c} \text{circle with two arcs} \\ \text{one arc goes up-left, one down-right} \end{array} \right\rangle_3, \\ \left\langle \begin{array}{c} \text{circle with one edge} \\ \text{edge goes right} \end{array} \right\rangle_3 &= [2]_q \left\langle \begin{array}{c} \text{circle with no edges} \end{array} \right\rangle_3, \\ \left\langle G \cup \begin{array}{c} \text{circle with one edge} \\ \text{edge goes right} \end{array} \right\rangle_3 &= [3]_q G \end{aligned}$$

where G is a bipartite uni-tri valent graph.

We can easily show that this linear map is invariant under the regular isotopy. We next introduce A_2 clasps of type $(n, 0)$ according to [Kup96] and [OY97]. Let P be the set $\{p_1, p_2, \dots, p_{2n}\}$ where p_1, p_2, \dots, p_{2n} are elements of P aligned counterclockwise on ∂D from p_1 in order of decreasing subscript of p_i , and ϵ is defined by

$$(2.1) \quad \epsilon(p_i) = \begin{cases} + & (1 \leq i \leq n) \\ - & (n+1 \leq i \leq 2n) \end{cases}.$$

We denote $W(P; \epsilon)$ as $W_{n^+ + n^-}$.

Definition 2.3. Let n be a positive integer. We define the A_2 clasps of type $(n, 0)$ by

$$\begin{aligned} \left\langle \begin{array}{c} \text{rectangle with one edge} \\ \text{edge goes up} \end{array} \right\rangle_3 &= \left\langle \begin{array}{c} \text{vertical line segment} \\ \text{top edge goes up} \end{array} \right\rangle_3 \in W_{1^+ + 1^-}, \\ \left\langle \begin{array}{c} \text{rectangle with one edge} \\ \text{edge goes up} \end{array} \right\rangle_3 &= \left\langle \begin{array}{c} \text{rectangle with one edge} \\ \text{edge goes up} \end{array} \right\rangle_3 - \frac{[n-1]_q}{[n]_q} \left\langle \begin{array}{c} \text{rectangle with two edges} \\ \text{top edge goes up, bottom edge goes down} \end{array} \right\rangle_3 \in W_{n^+ + n^-}. \end{aligned}$$

The following properties hold for A_2 clasps.

Lemma 2.4 ([Kup96]). For any positive integer n , we have

$$(2.2) \quad \left\langle \begin{array}{c} n \\ \text{---} \\ | \end{array} \right\rangle_3 = \left\langle \begin{array}{c} n \\ \text{---} \\ | \\ \text{---} \\ | \end{array} \right\rangle_3,$$

$$(2.3) \quad \left\langle \begin{array}{c} n-k-2 \\ \text{---} \\ | \\ \text{---} \\ | \\ k \\ \text{---} \\ | \end{array} \right\rangle_3 = 0 \quad (k = 0, 1, \dots, n-2).$$

Here, a strand labeled by the number n implies the n parallel copies of the strand.

We also introduce the A_2 clasp of type (n_1, n_2) according to [OY97].

Definition 2.5 (the A_2 clasp of type (n_1, n_2) [OY97]). Let n_1 and n_2 be non-negative integers. We define the A_2 clasp of type (n_1, n_2) by

$$\left\langle \begin{array}{c} n_1 \\ \text{---} \\ | \\ \text{---} \\ | \\ n_2 \\ \text{---} \\ | \end{array} \right\rangle_3 = \sum_{i=0}^{\min\{n_1, n_2\}} (-1)^i \frac{[n_1]_q [n_2]_q}{[n_1+n_2+1]_q} \left\langle \begin{array}{c} n_1 \\ \text{---} \\ | \\ \text{---} \\ | \\ n_2 \\ \text{---} \\ | \end{array} \right\rangle_3 \in W_{n_1^+ + n_2^- + n_1^- + n_2^+}.$$

Ohtsuki, Yamada [OY97] and Yuasa [Yua17] gave formulae for A_2 clasps of type $(n, 0)$.

Lemma 2.6 ([OY97][Yua17]). For $k = 0, 1, \dots, n$, we have

$$(2.4) \quad \left\langle \begin{array}{c} n-k \\ \text{---} \\ | \\ \text{---} \\ | \\ k \\ \text{---} \\ | \end{array} \right\rangle_3 = \frac{[n+1]_q [n+2]_q}{[n-k+1]_q [n-k+2]_q} \left\langle \begin{array}{c} n-k \\ \text{---} \\ | \\ \text{---} \\ | \end{array} \right\rangle_3$$

$$(2.5) \quad \left\langle \begin{array}{c} k \\ \text{---} \\ | \\ \text{---} \\ | \\ n-k \\ \text{---} \\ | \end{array} \right\rangle_3 = q^{\frac{k(n-k)}{3}} \left\langle \begin{array}{c} n \\ \text{---} \\ | \\ \text{---} \\ | \end{array} \right\rangle_3, \quad \left\langle \begin{array}{c} k \\ \text{---} \\ | \\ \text{---} \\ | \\ n-k \\ \text{---} \\ | \end{array} \right\rangle_3 = q^{-\frac{k(n-k)}{3}} \left\langle \begin{array}{c} n \\ \text{---} \\ | \\ \text{---} \\ | \end{array} \right\rangle_3$$


$$(2.6) \quad \left\langle \begin{array}{c} n \\ \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \\ | \end{array} \right\rangle_3 = q^{\frac{n^2+3n}{3}} \left\langle \begin{array}{c} n \\ \text{---} \\ | \\ \text{---} \\ | \end{array} \right\rangle_3, \quad \left\langle \begin{array}{c} n \\ \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \\ | \end{array} \right\rangle_3 = q^{-\frac{n^2+3n}{3}} \left\langle \begin{array}{c} n \\ \text{---} \\ | \\ \text{---} \\ | \end{array} \right\rangle_3$$

$$(2.7) \quad \Delta(n, 0) = \left\langle \begin{array}{c} n \\ \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \\ | \end{array} \right\rangle_3 = \frac{[n+1]_q [n+2]_q}{[2]_q}$$

$$(2.8) \quad \left\langle \begin{array}{c} n \\ \text{---} \\ | \\ \text{---} \\ | \\ n-k \\ \text{---} \\ | \\ \text{---} \\ | \\ k \\ \text{---} \\ | \\ n-k \\ \text{---} \\ | \\ n-k \\ \text{---} \\ | \end{array} \right\rangle_3 = q^{-(n-k)} \frac{(1-q^{n+1})(1-q^{n+2})}{(1-q^{k+1})(1-q^{k+2})} \Delta(n, 0)$$

We use the following two types of web, stair-step and triangle, which we can see in [Kim06], [Kim07], [Yua17], Etc.


Definition 2.7. For positive integers n and m , stair-step



– is defined by

$$\begin{array}{c}
 \text{Diagram 1:} \\
 \begin{array}{ccc}
 \text{Left:} & \xrightarrow{\quad n \quad} & \boxed{\text{Block}} \xrightarrow{\quad m \quad} \\
 \downarrow & & \downarrow \\
 \boxed{\text{Block}} \xrightarrow{\quad n \quad} & = & \text{Right:} \\
 & & \left\{ \begin{array}{c} \xrightarrow{\quad n \quad} \\ \vdots \\ \xrightarrow{\quad n \quad} \end{array} \right. \xrightarrow{\quad m-1 \quad} \xrightarrow{\quad 1 \quad} \xrightarrow{\quad m \quad}
 \end{array}
 \end{array}$$

For positive integers n , triangle is defined by



$$\begin{array}{c} \text{Diagram A: } \begin{array}{c} \diagdown \quad \diagup \\ \backslash \quad / \\ \triangle \end{array} \quad \text{Diagram B: } \begin{array}{c} \diagdown \quad \diagup \\ \backslash \quad / \\ = \end{array} \\ \text{Diagram C: } \begin{array}{c} \diagdown \quad \diagup \\ \backslash \quad / \\ \text{Y-shaped tree} \end{array} \quad , \end{array}$$

$$n \begin{array}{c} \diagdown \\ \diagup \end{array} n = \begin{array}{c} 1 \rightarrow \leftarrow 1 \\ n-1 \rightarrow \leftarrow \end{array} \quad (n > 1).$$

The opposite direction is defined in the same way.

The following formulae help calculation of the \mathfrak{sl}_3 colored Jones polynomial. Yuasa [Yua17], [Yua18b], [Yua21b] and Kim [Kim06] gave them. We sometimes omit directions of edges of A_2 webs.

Lemma 2.8 ([Yua17] [Yua18b] [Yua21b] [Kim06]). For positive integer m and n , we have

$$(2.9) \quad \left\langle \begin{array}{c} n \rightarrow \square \\ \square \rightarrow n \\ n \rightarrow \square \\ \square \rightarrow n \end{array} \right\rangle_3 = \sum_{k=0}^n (-1)^k q^{\frac{2n^2 - 6nk + 3k^2}{6}} \frac{(q)_n}{(q)_k (q)_{n-k}} \left\langle \begin{array}{c} n \rightarrow \square \\ \square \rightarrow n \\ n \rightarrow \square \\ \square \rightarrow n \\ \square \rightarrow n \\ \square \rightarrow n \\ n-k \\ k \\ k \\ n-k \end{array} \right\rangle_3,$$

$$(2.10) \quad \left\langle \begin{array}{c} n \rightarrow \square \\ \square \rightarrow n \\ n \rightarrow \square \\ \square \rightarrow n \end{array} \right\rangle_3 = \sum_{k=0}^n (-1)^k q^{\frac{-2n^2+3k^2}{6}} \frac{(q)_n}{(q)_k (q)_{n-k}} \left\langle \begin{array}{c} n \rightarrow \square \quad n-k \\ \square \rightarrow n \quad \square \rightarrow n \\ k \quad k \\ n-k \end{array} \right\rangle_3,$$

$$(2.11) \quad \left\langle \begin{array}{c} n \\ | \\ \square \end{array} \begin{array}{c} \triangle \\ | \\ \square \end{array} \begin{array}{c} n \\ | \\ \square \end{array} \right\rangle_3 = \left\langle \begin{array}{c} n \\ | \\ \square \end{array} \begin{array}{c} \triangle \\ | \\ \square \end{array} \begin{array}{c} n \\ | \\ \square \end{array} \right\rangle_3,$$

$$(2.12) \quad \left\langle \begin{array}{c} n \\ | \\ \square \end{array} \begin{array}{c} \square \\ | \\ \square \end{array} \begin{array}{c} n \\ | \\ \square \end{array} \right\rangle_3 = \left\langle \begin{array}{c} n \\ | \\ \square \end{array} \begin{array}{c} \triangle \\ | \\ \square \end{array} \begin{array}{c} n \\ | \\ \square \end{array} \right\rangle_3,$$

$$(2.13) \quad \left\langle \begin{array}{c} n \\ | \\ \square \end{array} \begin{array}{c} \triangle \\ | \\ \square \end{array} \begin{array}{c} n \\ | \\ \square \end{array} \right\rangle_3 = \left\langle \begin{array}{c} k \\ | \\ \square \end{array} \begin{array}{c} k \\ | \\ \square \end{array} \begin{array}{c} n-k \\ | \\ \square \end{array} \begin{array}{c} n-k \\ | \\ \square \end{array} \begin{array}{c} n-k \\ | \\ \square \end{array} \right\rangle_3,$$

$$(2.14) \quad \left\langle \begin{array}{c} n \\ | \\ \square \end{array} \begin{array}{c} \triangle \\ | \\ \square \end{array} \begin{array}{c} n \\ | \\ \square \end{array} \right\rangle_3 = \left\langle \begin{array}{c} n \\ | \\ \square \end{array} \begin{array}{c} \square \\ | \\ \square \end{array} \begin{array}{c} 1 \\ | \\ \square \end{array} \begin{array}{c} n-1 \\ | \\ \square \end{array} \begin{array}{c} n-1 \\ | \\ \square \end{array} \right\rangle_3 - \frac{[n-1]_q}{[n]_q} \left\langle \begin{array}{c} n \\ | \\ \square \end{array} \begin{array}{c} \square \\ | \\ \square \end{array} \begin{array}{c} 1 \\ | \\ \square \end{array} \begin{array}{c} n-1 \\ | \\ \square \end{array} \begin{array}{c} n-1 \\ | \\ \square \end{array} \right\rangle_3,$$

$$(2.15) \quad \left\langle \begin{array}{c} n+m \\ | \\ \square \end{array} \begin{array}{c} \square \\ | \\ \square \end{array} \begin{array}{c} n \\ | \\ \square \end{array} \begin{array}{c} m \\ | \\ \square \end{array} \begin{array}{c} m \\ | \\ \square \end{array} \right\rangle_3 = \left\langle \begin{array}{c} n+m \\ | \\ \square \end{array} \begin{array}{c} \square \\ | \\ \square \end{array} \begin{array}{c} n \\ | \\ \square \end{array} \begin{array}{c} m \\ | \\ \square \end{array} \begin{array}{c} m \\ | \\ \square \end{array} \right\rangle_3,$$

$$(2.16) \quad \left\langle \begin{array}{c} n \\ | \\ \square \end{array} \begin{array}{c} \square \\ | \\ \square \end{array} \begin{array}{c} n \\ | \\ \square \end{array} \right\rangle_3 = [n+1]_q \sum_{k=0}^n \left\langle \begin{array}{c} k \\ | \\ \square \end{array} \right\rangle_3,$$

$$(2.17) \quad \left\langle \begin{array}{c} n \\ | \\ \square \end{array} \begin{array}{c} \square \\ | \\ \square \end{array} \begin{array}{c} n \\ | \\ \square \end{array} \right\rangle_3 = \sum_{k=0}^n \left\langle \begin{array}{c} n-k \\ | \\ \square \end{array} \begin{array}{c} k \\ | \\ \square \end{array} \begin{array}{c} n-k \\ | \\ \square \end{array} \right\rangle_3,$$

$$(2.18) \quad \left\langle \begin{array}{c} n \\ | \\ \square \end{array} \begin{array}{c} k \\ | \\ \square \end{array} \begin{array}{c} n-k \\ | \\ \square \end{array} \begin{array}{c} n-k \\ | \\ \square \end{array} \right\rangle_3 = \frac{[n+1]_q[n+2]_q}{[n-k+2]_q} \Delta(n,0) \quad (0 \leq k \leq n),$$

$$(2.19) \quad \left\langle \begin{array}{c} n \\ | \\ \square \end{array} \right\rangle_3 = \sum_{i=0}^{n-1} (-1)^i \frac{[n-i]}{[n]_q} \left\langle \begin{array}{c} i \\ | \\ \square \end{array} \begin{array}{c} 1 \\ | \\ \square \end{array} \begin{array}{c} n-i-1 \\ | \\ \square \end{array} \right\rangle_3 \in W_{1^+1^-},$$

(2.20)

$$\left\langle \begin{array}{c} n \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ n \end{array} \right\rangle_3 = (-1)^n q^{-\frac{n^2+3n}{6}} \left\langle \begin{array}{c} n \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ n \end{array} \right\rangle_3, \quad \left\langle \begin{array}{c} n \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ n \end{array} \right\rangle_3 = (-1)^n q^{\frac{n^2+3n}{6}} \left\langle \begin{array}{c} n \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ n \end{array} \right\rangle_3,$$

(2.21)

$$\left\langle \begin{array}{c} m \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ m \end{array} \right\rangle_3 = (-1)^{mn} q^{-\frac{mm}{6}} \left\langle \begin{array}{c} m \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ m \end{array} \right\rangle_3, \quad \left\langle \begin{array}{c} m \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ m \end{array} \right\rangle_3 = (-1)^{mn} q^{\frac{mm}{6}} \left\langle \begin{array}{c} m \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ m \end{array} \right\rangle_3,$$

(2.22)

$$\left\langle \begin{array}{c} m \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ m \end{array} \right\rangle_3 = (-1)^{mn} q^{-\frac{mm}{6}} \left\langle \begin{array}{c} m \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ m \end{array} \right\rangle_3, \quad \left\langle \begin{array}{c} m \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ m \end{array} \right\rangle_3 = (-1)^{mn} q^{\frac{mm}{6}} \left\langle \begin{array}{c} m \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ m \end{array} \right\rangle_3.$$

In [Yua17], Yuasa gave the m times full twists formula for two strands with opposite directions and the A_2 bracket bubble skein expansion formula. We use these formulae to calculate the one-row \mathfrak{sl}_3 colored Jones polynomial for pretzel links in Section 4.

Theorem 2.9 (m times full twist formula [Yua17]). For a positive integer n , we have

$$\left\langle \begin{array}{c} n \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ n \end{array} \right\rangle_3 = \sum_{0 \leq k_m \leq k_{m-1} \leq \dots \leq k_1 \leq n} \phi(n, k_1, k_2, \dots, k_m) q^{\epsilon_m} \left\langle \begin{array}{c} n-k_m \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ n-k_m \end{array} \right\rangle_3$$

m times full twists

where

$$\phi(n, k_1, k_2, \dots, k_m) q^{\epsilon_m} = \frac{(q^{\epsilon_m})^{-\frac{2m}{3}(n^2+3n)} (q^{\epsilon_m})^{n-k_m} (q^{\epsilon_m})^{\sum_{i=1}^m (k_i^2 + 2k_i)} (q^{\epsilon_m})_n^2}{(q^{\epsilon_m})_{n-k_1} (q^{\epsilon_m})_{k_1-k_2} \cdots (q^{\epsilon_m})_{k_{m-1}-k_m} (q^{\epsilon_m})_{k_m}^2}.$$

Theorem 2.10 (the A_2 bracket bubble skein expansion formula [Yua17]). Let n be a positive integer and k, l non-negative integers. For $n \geq k, l$, we have

$$\left\langle \begin{array}{c} n-k \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ n-k \end{array} \right\rangle_3 = \sum_{t=\max\{k,l\}}^{\min\{k+l,n\}} \psi(n, t, k, l) \left\langle \begin{array}{c} n-t \\ \nearrow \nwarrow \\ \text{bubble} \\ \nearrow \nwarrow \\ n-t \end{array} \right\rangle_3$$

where

$$\psi(n, t, k, l) = \frac{q^{(t+1)(t-k-l)+kl} (q)_k (q)_l (q)_{n-k}^2 (q)_{n-l}^2 (q)_{2n-t+2}}{(q)_n^2 (q)_{n-t}^2 (q)_{t-k} (q)_{t-l} (q)_{2n-k-l+2} (q)_{-t+k+l}}.$$

3. THE FORMULAE FOR THE A_2 WEB SPACE

We give m times half twists formulae for two strands with the same directions for A_2 bracket.

Proposition 3.1 (m times half twists formula). Let n be a positive integer and $k_0 = n$. For a positive integer m , we have

$$(3.1) \quad \left\langle \begin{array}{c} \text{Diagram of } m \text{ times half twists} \\ \text{with strands } n \text{ and } n \end{array} \right\rangle_3 = \sum_{0 \leq k_m \leq k_{m-1} \leq \dots \leq k_1 \leq n} \chi_1(n, k_1, k_2, \dots, k_m) \left\langle \begin{array}{c} \text{Diagram with strands } n, n-k_m, n-k_m, k_m, k_m \\ \text{and a crossing} \end{array} \right\rangle_3$$

where

$$\chi_1(n, k_1, k_2, \dots, k_m) = (-1)^{nm} \frac{q^{-\frac{1}{6}(n^2+3n)m} q^{\frac{1}{2}(n-k_m)} (-1)^{\sum_{i=1}^m k_i} q^{\sum_{i=1}^m \frac{1}{2}(k_i^2+k_i)} (q)_n}{\prod_{i=1}^m (q)_{k_{i-1}-k_i} (q)_{k_m}}.$$

For a negative integer m , we have

$$(3.2) \quad \left\langle \begin{array}{c} \text{Diagram of } m \text{ times-half twists} \\ \text{with strands } n \text{ and } n \end{array} \right\rangle_3 = \sum_{0 \leq k_m \leq k_{m-1} \leq \dots \leq k_1 \leq n} \chi_{-1}(n, k_1, k_2, \dots, k_m) \left\langle \begin{array}{c} \text{Diagram with strands } n, n-k_m, n-k_m, k_m, k_m \\ \text{and a crossing} \end{array} \right\rangle_3$$

where

$$\chi_{-1}(n, k_1, k_2, \dots, k_m) = (-1)^{nm} \frac{q^{\frac{1}{6}(n^2+3n)m} q^{-\frac{1}{2}(n-k_m)} (-1)^{\sum_{i=1}^m k_i} q^{\sum_{i=1}^m \frac{1}{2}(k_i^2-k_i)} q^{\sum_{i=1}^m k_{i-1} k_i} (q)_n}{\prod_{i=1}^m (q)_{k_{i-1}-k_i} (q)_{k_m}}.$$

To prove Proposition 3.1, we prepare the following lemma.

Lemma 3.2. Let n be a positive integer and k non-negative integer. For $0 \leq k \leq n$, we have

$$(3.3) \quad \left\langle \begin{array}{c} \text{Diagram with strands } n, n-k, k \\ \text{and a crossing} \end{array} \right\rangle_3 = \left\langle \begin{array}{c} \text{Diagram with strands } n, n-k, k \\ \text{and a crossing} \end{array} \right\rangle_3.$$

Proof. By definition of A_2 clasp of type (n_1, n_2) , we obtain

$$\left\langle \begin{array}{c} n \\ \nearrow \quad \searrow \\ \text{---} \\ \text{---} \\ \nearrow \quad \searrow \\ n-k \end{array} \right\rangle_3 = \sum_{i=0}^{\min\{n-k,k\}} (-1)^i \frac{\begin{bmatrix} n-k \\ i \end{bmatrix}_q \begin{bmatrix} k \\ i \end{bmatrix}_q}{\begin{bmatrix} n+i \\ i \end{bmatrix}_q} \left\langle \begin{array}{c} n \\ \nearrow \quad \searrow \\ \text{---} \\ \text{---} \\ \nearrow \quad \searrow \\ n-k \\ \nearrow \quad \searrow \\ i \\ \nearrow \quad \searrow \\ i \end{array} \right\rangle_3$$

The right-hand side of the above equation is zero by (2.11), (2.2), and (2.3) except for $i = 0$.

Proof of Proposition 3.1. We proceed by induction on m . We can see that (3.1) holds by (2.9) for $m = 1$. Assume (3.1) holds when $m = l$ for some integers $l \geq 1$.

$$\begin{aligned}
& \left\langle \begin{array}{c} n \\ \diagup \quad \diagdown \\ \vdots \\ n \end{array} \right\rangle_3 \\
& m \text{ times-half twists} \\
& = \sum_{0 \leq k_l \leq k_{l-1} \leq \dots \leq k_1 \leq n} \chi_1(n, k_1, k_2, \dots, k_l) \left\langle \begin{array}{c} n-k_l \\ \diagup \quad \diagdown \\ n-k_l \\ n-k_l \\ \diagup \quad \diagdown \\ n-k_l \\ n-k_l \end{array} \right\rangle_3 = \sum_{0 \leq k_l \leq k_{l-1} \leq \dots \leq k_1 \leq n} \chi_1(n, k_1, k_2, \dots, k_l) \left\langle \begin{array}{c} n-k_l \\ \diagup \quad \diagdown \\ n-k_l \\ n-k_l \\ \diagup \quad \diagdown \\ n-k_l \\ n-k_l \end{array} \right\rangle_3 \\
& = \sum_{0 \leq k_l \leq k_{l-1} \leq \dots \leq k_1 \leq n} \chi_1(n, k_1, k_2, \dots, k_l) (-1)^{n-k_l} q^{-\frac{1}{6}(n-k_l)^2 - \frac{1}{2}(n-k_l)} q^{-\frac{2}{3}(n-k_l)k_l} \left\langle \begin{array}{c} n-k_l \\ \diagup \quad \diagdown \\ n-k_l \\ n-k_l \\ \diagup \quad \diagdown \\ n-k_l \\ n-k_l \end{array} \right\rangle_3 \\
& = \sum_{0 \leq k_l \leq k_{l-1} \leq \dots \leq k_1 \leq n} \chi_1(n, k_1, k_2, \dots, k_l) (-1)^{n-k_l} q^{-\frac{1}{6}(n-k_l)^2 - \frac{1}{2}(n-k_l)} q^{-\frac{2}{3}(n-k_l)k_l} \left\langle \begin{array}{c} n-k_l \\ \diagup \quad \diagdown \\ n-k_l \\ n-k_l \\ \diagup \quad \diagdown \\ n-k_l \\ n-k_l \end{array} \right\rangle_3 \\
& \text{(Lemma 3.2)}
\end{aligned}$$

$$\begin{aligned}
& \stackrel{\text{(Lemma 3.2, (2.21))}}{=} \sum_{0 \leq k_l \leq k_{l-1} \leq \dots \leq k_1 \leq n} \chi_1(n, k_1, k_2, \dots, k_l) (-1)^{n-k_l} q^{-\frac{1}{6}(n-k_l)^2 - \frac{1}{2}(n-k_l)} q^{-\frac{2}{3}(n-k_l)k_l} q^{\frac{1}{3}(n-k_l)k_l} \left\langle \begin{array}{c} n-k_l \\ n-k_l \end{array} \right\rangle_l \\
& = \stackrel{\text{(Lemma 3.2)}}{=} \sum_{0 \leq k_l \leq k_{l-1} \leq \dots \leq k_1 \leq n} \chi_1(n, k_1, k_2, \dots, k_l) (-1)^{n-k_l} q^{-\frac{1}{6}(n-k_l)^2 - \frac{1}{2}(n-k_l)} q^{-\frac{1}{3}(n-k_l)k_l} \left\langle \begin{array}{c} n-k_l \\ n-k_l \end{array} \right\rangle_3 \\
& \stackrel{\text{(2.21))}}{=} \sum_{0 \leq k_l \leq k_{l-1} \leq \dots \leq k_1 \leq n} \chi_1(n, k_1, k_2, \dots, k_l) (-1)^{n-k_l} q^{-\frac{1}{6}(n-k_l)^2 - \frac{1}{2}(n-k_l)} q^{-\frac{2}{3}(n-k_l)k_l} \left\langle \begin{array}{c} n-k_l \\ n-k_l \end{array} \right\rangle_3 \\
& \stackrel{\text{(2.9))}}{=} \sum_{0 \leq k_{l+1} \leq k_l \leq \dots \leq k_1 \leq n} \chi_1(n, k_1, k_2, \dots, k_{l+1}) \left\langle \begin{array}{c} n-k_l \\ n-k_l \end{array} \right\rangle_3
\end{aligned}$$

We can see that

$$(3.4) \quad \left\langle \begin{array}{c} n \\ \nearrow k_l \\ \text{---} \\ n-k_l \\ \swarrow \\ \text{---} \\ n-k_l \\ \nearrow k_{l+1} \\ \text{---} \\ k_l-k_{l+1} \end{array} \right\rangle_3 \text{ (Lemma 3.2, (2.22))} = \left\langle \begin{array}{c} n \\ \nearrow k_l \\ \text{---} \\ n-k_l \\ \swarrow \\ \text{---} \\ n-k_l \\ \nearrow k_{l+1} \\ \text{---} \\ k_l-k_{l+1} \end{array} \right\rangle_3 \text{ (Definition 2.7)} = \left\langle \begin{array}{c} n \\ \nearrow k_{l+1} \\ \text{---} \\ n-k_l \\ \swarrow \\ \text{---} \\ n-k_l \\ \nearrow k_{l+1} \\ \text{---} \\ k_l-k_{l+1} \end{array} \right\rangle_3$$

(2.12) $\left\langle \begin{array}{c} n \\ \nearrow k_{l+1} \\ \text{---} \\ n-k_{l+1} \\ \swarrow \\ \text{---} \\ n-k_{l+1} \\ \nearrow k_{l+1} \\ \text{---} \\ k_l-k_{l+1} \end{array} \right\rangle_3 \text{ (Lemma 3.2, (2.22))} = \left\langle \begin{array}{c} n \\ \nearrow k_{l+1} \\ \text{---} \\ n-k_{l+1} \\ \swarrow \\ \text{---} \\ n-k_{l+1} \\ \nearrow k_{l+1} \\ \text{---} \\ k_l-k_{l+1} \end{array} \right\rangle_3$

By using (3.4), we have

$$(3.5) \quad \left\langle \begin{array}{c} \text{Diagram showing two configurations of a link component with labels } n, k_l, n-k_l, k_{l+1}, n-k_{l+1}, k_{l+2}, n-k_{l+2} \end{array} \right\rangle_3 = \left\langle \begin{array}{c} \text{Diagram showing two configurations of a link component with labels } n, k_{l+1}, n-k_{l+1}, k_{l+1}, n-k_{l+1}, k_{l+2}, n-k_{l+2} \end{array} \right\rangle_3$$

We remove the A_2 clasps by using Lemma 3.2 and (3.4) in (RHS of (3.5)). Therefore, (3.1) holds for $n = l + 1$. We can see (3.2) in a similar way. \square

We use the same definition for the one-row \mathfrak{sl}_3 colored Jones polynomial as we do in [Yua18a].

Definition 3.3. Let L be an oriented link, and \bar{L} a diagram of L . The one-row \mathfrak{sl}_3 colored Jones polynomial for L is defined by

$$J_{(n,0)}^{\mathfrak{sl}_3}(L; q) = (q^{\frac{n^2+3n}{3}})^{-w(\bar{L})} \langle L(n, 0) \rangle_3 / \Delta(n, 0)$$

where $w(\bar{L})$ is the writhing number of \bar{L} and $L(n, 0)$ replaces a part of \bar{L} with A_2 clasp of type $(n, 0)$. For example,

$$T(2,3)(n,0) = \begin{array}{c} \text{Diagram of a torus knot } T(2,3) \text{ with a } A_2 \text{ clasp of type } (n,0) \end{array}$$

Let a map $sign : \mathbb{Z} \rightarrow \{-1, 0, 1\}$ be defined by

$$sign(x) = \begin{cases} 1 & (x > 0) \\ 0 & (x = 0) \\ -1 & (x < 0) \end{cases}$$

Corollary 3.4. The one-row colored \mathfrak{sl}_3 Jones polynomials for $(2, m)$ -torus knots $T(2, m)$ are the following:

$$(3.6) \quad J_{(n,0)}^{\mathfrak{sl}_3}(T(2, m); q) = (q^{\frac{n^2+3n}{3}})^{-(m)} \sum_{0 \leq k_{|m|} \leq k_{|m|-1} \leq \dots \leq k_1 \leq n} \chi_{sign(m)} q^{\frac{n+k_{|m|}}{2}} \frac{(1-q^{n+1})(1-q^{n+2})}{(1-q)(1-q^{n-k_{|m|}+2})}.$$

Proof. we can obtain (3.6) from Definition 3.3, Proposition 3.1 and (2.18) easily. \square

Remark 3.5. For $m = 3, 5, 7$, $n=1, 2, \dots, 10$, Corollary 3.4 are equal to Theorem 2.1 in [GMV13] and Theorem 5.7 in [Yua17] for $T(2, m)$ by using Mathematica. For example,

$$\begin{aligned} J_{(4,0)}^{\mathfrak{sl}_3}(T(2,3); q) = & q^{-42} - q^{-40} - q^{-39} - q^{-38} + q^{-36} + q^{-35} + 2q^{-31} + q^{-30} - q^{-28} - q^{-27} + q^{-25} - q^{-23} - 2q^{-22} - q^{-21} \\ & + q^{-20} + q^{-19} - q^{-17} + q^{-14} + q^{-13} + q^{-8} \end{aligned}$$

Proposition 3.6. Let n be a positive integer and k, l non-negative integers. For $n \geq k, l$, we have

$$\left\langle \begin{array}{c} \text{Diagram showing two configurations of a link component with labels } n, n-k, n-l, k, l, n-k, n-l \end{array} \right\rangle_3 = \sum_{t=\max\{k,l\}}^{\min\{k+l,n\}} \sum_{a=t}^n \Omega(n, t, k, l) \left\langle \begin{array}{c} \text{Diagram showing two configurations of a link component with labels } n, n-t, n-t, t, t, n-t \end{array} \right\rangle_3$$

where

$$\Omega(n, k, l, t) = \frac{q^{-\frac{k+l}{2}+t} q^{(t+1)(t-k-l)+kl} (1 - q^{n+1-k}) (1 - q^{n+1-l}) (q)_k (q)_l (q)_{n-k}^2 (q)_{n-l}^2 (q)_{2n-t+2}}{(1 - q^{n+1-t})^2 (q)_n^2 (q)_{n-t}^2 (q)_{t-k} (q)_{t-l} (q)_{2n-k-l+2} (q)_{-t+k+l}}.$$

We first prove the following lemma.

Lemma 3.7. Let n and a be positive integers. We have

$$\left\langle \begin{array}{c} n \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} \square \xrightarrow{\quad} \square \\ \square \xleftarrow{\quad} \square \end{array} \xrightarrow{\quad} \begin{array}{c} n \\ \square \end{array} \right\rangle_3 = \frac{[n+1]_q}{[n-a+1]_q} \left\langle \begin{array}{c} n \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} \square \xrightarrow{\quad} \square \\ \square \xleftarrow{\quad} \square \end{array} \xrightarrow{\quad} \begin{array}{c} a \\ \square \end{array} \right\rangle_3$$

Proof. First, we prove that it holds for $a = 1$.

$$(3.7) \quad \left\langle \begin{array}{c} n \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} \square \xrightarrow{\quad} \square \\ \square \xleftarrow{\quad} \square \end{array} \xrightarrow{\quad} \begin{array}{c} n \\ \square \end{array} \right\rangle_3^1 = \frac{[n+1]_q}{[n]_q} \left\langle \begin{array}{c} n \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} \square \xrightarrow{\quad} \square \\ \square \xleftarrow{\quad} \square \end{array} \xrightarrow{\quad} \begin{array}{c} 1 \\ \square \end{array} \right\rangle_3$$

The equation (3.7) is the same lemma in [Yua21b], but we prove it in a different way. We proceed by induction on n . We can see that (3.7) holds by an easy calculation for $n = 2$. Assume (3.7) holds when $n \leq k-1$ for some integer $k \geq 3$. We obtain

$$\begin{aligned} & \left\langle \begin{array}{c} k \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} \square \xrightarrow{\quad} \square \\ \square \xleftarrow{\quad} \square \end{array} \xrightarrow{\quad} \begin{array}{c} k \\ \square \end{array} \right\rangle_3^1 \\ &= ((2.14)) \left\langle \begin{array}{c} k \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} \square \xrightarrow{\quad} \square \\ \square \xleftarrow{\quad} \square \end{array} \xrightarrow{\quad} \begin{array}{c} k-1 \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} k-2 \\ \square \end{array} \right\rangle_3 - \frac{[k-1]_q}{[k]_q} \left\langle \begin{array}{c} k \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} \square \xrightarrow{\quad} \square \\ \square \xleftarrow{\quad} \square \end{array} \xrightarrow{\quad} \begin{array}{c} k-1 \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} k-2 \\ \square \end{array} \right\rangle_3 \\ &= (\text{Definition 2.7, (2.19)}) \frac{[k]_q}{[k-1]_q} \left\langle \begin{array}{c} k \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} \square \xrightarrow{\quad} \square \\ \square \xleftarrow{\quad} \square \end{array} \xrightarrow{\quad} \begin{array}{c} k-1 \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} k-2 \\ \square \end{array} \right\rangle_3 - (-1)^{k-1} \frac{1}{[k]_q} \left\langle \begin{array}{c} k \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} \square \xrightarrow{\quad} \square \\ \square \xleftarrow{\quad} \square \end{array} \xrightarrow{\quad} \begin{array}{c} k-1 \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} k-2 \\ \square \end{array} \right\rangle_3 \end{aligned}$$

$$\begin{aligned}
 &= \frac{[k]_q}{[k-1]_q} \left\langle \begin{array}{c} k \\ \downarrow \\ -k \end{array} \right\rangle_3 - (-1)^{k-1} \frac{1}{[k]_q} \left\langle \begin{array}{c} k \\ \downarrow \\ -k \\ \downarrow \\ k-1 \\ \downarrow \\ k-1 \end{array} \right\rangle_3
 \end{aligned}$$

For the second term,

$$\begin{aligned}
 &\left\langle \begin{array}{c} k \\ \downarrow \\ k-1 \\ \downarrow \\ k-2 \\ \downarrow \\ k-1 \\ \downarrow \\ k-1 \end{array} \right\rangle_3 \\
 &= \frac{[k-2]_q}{[k-1]_q} \left\langle \begin{array}{c} k \\ \downarrow \\ k-1 \\ \downarrow \\ k-3 \\ \downarrow \\ k-1 \\ \downarrow \\ k-1 \end{array} \right\rangle_3 - \frac{[k-2]_q}{[k-1]_q} \left\langle \begin{array}{c} k \\ \downarrow \\ k-1 \\ \downarrow \\ k-3 \\ \downarrow \\ k-1 \\ \downarrow \\ k-1 \end{array} \right\rangle_3 \\
 &= -\frac{[k-2]_q}{[k-1]_q} \left\langle \begin{array}{c} k \\ \downarrow \\ k-3 \\ \downarrow \\ k-2 \\ \downarrow \\ k-1 \\ \downarrow \\ k-1 \end{array} \right\rangle_3 \stackrel{\text{(Definition 2.7)}}{=} -\frac{[k-2]_q}{[k-1]_q} \left\langle \begin{array}{c} k \\ \downarrow \\ k-3 \\ \downarrow \\ k-2 \\ \downarrow \\ k-1 \\ \downarrow \\ k-1 \end{array} \right\rangle_3 \\
 &\stackrel{\text{(Definition 2.7)}}{=} -\frac{[k-2]_q}{[k-1]_q} \left\langle \begin{array}{c} k \\ \downarrow \\ k-2 \\ \downarrow \\ k-2 \\ \downarrow \\ k-1 \\ \downarrow \\ k-1 \end{array} \right\rangle_3 \stackrel{\text{(Definition 2.7, (2.19))}}{=} -(-1)^{k-2} \frac{1}{[k-1]_q} \left\langle \begin{array}{c} k \\ \downarrow \\ k-2 \\ \downarrow \\ k-2 \\ \downarrow \\ k-1 \\ \downarrow \\ k-1 \end{array} \right\rangle_3 \\
 &\stackrel{\text{(Definition 2.7)}}{=} -(-1)^{k-2} \frac{1}{[k-1]_q} \left\langle \begin{array}{c} k \\ \downarrow \\ k-2 \\ \downarrow \\ k-2 \\ \downarrow \\ k-1 \\ \downarrow \\ k-1 \end{array} \right\rangle_3 \stackrel{\text{(induction hypothesis)}}{=} -(-1)^{k-2} \frac{1}{[k-1]_q} \left\langle \begin{array}{c} k \\ \downarrow \\ -k \end{array} \right\rangle_3
 \end{aligned}$$

Regarding second equality, we have

$$\left\langle \begin{array}{c} k \\ \downarrow \\ k-1 \\ \downarrow \\ k-3 \\ \downarrow \\ k-1 \\ \downarrow \\ k-1 \end{array} \right\rangle_3 = 0$$

by using Definition 2.7 and (2.3). Hence,

$$\left\langle \begin{array}{c} k \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} \square \\ \searrow \swarrow \end{array} \xleftarrow{\quad} \begin{array}{c} k \\ \square \end{array} \right\rangle_3 = \frac{[k]_q^2 - 1}{[k]_q[k-1]_q} \left\langle \begin{array}{c} k \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} \square \\ \swarrow \searrow \end{array} \xleftarrow{\quad} \begin{array}{c} k \\ \square \end{array} \right\rangle_3$$

We can check easily the following equation by Lemma 2.1.

$$(3.8) \quad \frac{[k]_q^2 - 1}{[k]_q[k-1]_q} = \frac{[k+1]_q}{[k]_q}$$

Therefore, (3.7) holds for $n = k$. (3.8) is not defined for $n = 1$. However (3.7) holds for $n = 1$ by definition of A_2 bracket. We can prove easily that if we use (3.8) repeatedly, then Lemma 3.7 holds. \square

Proof of Proposition 3.6.

$$\begin{aligned} & \left\langle \begin{array}{c} n \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} n-k \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} n-l \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} l \\ \square \end{array} \right\rangle_3 \stackrel{\text{(Theorem 2.10)}}{=} \sum_{t=\max\{k,l\}}^{\min\{k+l,n\}} \psi(n,t,k,l) \left\langle \begin{array}{c} n \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} n-k \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} n-l \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} l \\ \square \end{array} \right\rangle_3 \\ & \stackrel{\text{(2.2), Lemma 3.7}}{=} \sum_{t=\max\{k,l\}}^{\min\{k+l,n\}} \frac{[n-k+1]_q[n-l+1]_q}{[n+1-t]_q^2} \psi(n,t,k,l) \left\langle \begin{array}{c} n \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} n-t \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} t \\ \square \end{array} \right\rangle_3 \\ & \stackrel{\text{((2.17))}}{=} \sum_{t=\max\{k,l\}}^{\min\{k+l,n\}} \sum_{a=t}^n \Omega(n,t,k,l) \left\langle \begin{array}{c} n \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} n-a \\ \square \end{array} \xrightarrow{\quad} \begin{array}{c} a \\ \square \end{array} \right\rangle_3 \end{aligned}$$

\square

4. COMPUTING THE ONE-ROW \mathfrak{sl}_3 COLORED JONES POLYNOMIAL FOR PRETZEL LINKS

In this section, we compute the one-row \mathfrak{sl}_3 colored Jones polynomials for pretzel links. We consider the three-parameter family of pretzel links $P(\alpha, \beta, \gamma)$ for integers α, β, γ .

$$P(\alpha, \beta, \gamma) = \boxed{\alpha \quad \beta \quad \gamma}$$

where a box marked with the letter α represents a right-handed (resp. left-handed) $|\alpha|$ -twist if $\alpha > 0$ (resp. $\alpha < 0$). Boxes with other letters represent the same thing. The one-row colored \mathfrak{sl}_3 Jones polynomial of a link depends on the direction of the link. Therefore, it is necessary to distinguish links with different directions. For example, we consider a knot with the directions using the arrow symbol as follows.

$$P(\downarrow 3 \downarrow, \uparrow 3 \uparrow, \downarrow 2 \uparrow) = \text{Diagram of a trefoil knot with arrows indicating direction.}$$

Theorem 4.1. Let α, β, γ be integers. The one-row colored \mathfrak{sl}_3 Jones polynomials for pretzel links $P(\alpha, \beta, \gamma)$ are the following:

$$(4.1) \quad J_{(n,0)}^{\mathfrak{sl}_3}(P(\downarrow 2\alpha \uparrow, \downarrow 2\beta \uparrow, \downarrow 2\gamma \uparrow); q) = \sum_{0 \leq k_{|\alpha|} \leq k_{|\alpha|-1} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{|\beta|} \leq l_{|\beta|-1} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_{|\gamma|} \leq m_{|\gamma|-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{|\alpha|}, l_{|\beta|}\}}^{\min\{k_{|\alpha|}+l_{|\beta|}, n\}} \sum_{t=\max\{s, m_{|\gamma|}\}}^{\min\{s+m_{|\gamma|}, n\}} (q^{\frac{n^2+3n}{3}})^{-(2\alpha+2\beta+2\gamma)} \\ \phi(n, k_1, k_2, \dots, k_{|\alpha|}) q^{\epsilon_\alpha} \phi(n, l_1, l_2, \dots, l_{|\beta|}) q^{\epsilon_\beta} \phi(n, m_1, m_2, \dots, m_{|\gamma|}) q^{\epsilon_\gamma} \psi(n, s, k_{|\alpha|}, l_{|\beta|}) \psi(n, t, s, m_{|\gamma|}) q^{-(n-t)} \frac{(1-q^{n+1})(1-q^{n+2})}{(1-q^{t+1})(1-q^{t+2})},$$

$$(4.2) \quad J_{(n,0)}^{\mathfrak{sl}_3}(P(\downarrow 2\alpha \downarrow, \uparrow 2\beta \uparrow, \downarrow 2\gamma \uparrow); q) = \sum_{0 \leq k_{2|\alpha|} \leq k_{2|\alpha|-1} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2|\beta|} \leq l_{2|\beta|-1} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_{|\gamma|} \leq m_{|\gamma|-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{2|\alpha|}, l_{2|\beta|}\}}^{\min\{k_{2|\alpha|}+l_{2|\beta|}, n\}} \sum_{a=0}^{n-s} \sum_{t=\max\{a, m_{|\gamma|}\}}^{\min\{a+t+m_{|\gamma|}, n\}} (q^{\frac{n^2+3n}{3}})^{2\alpha+2\beta-2\gamma} \\ \chi_{sign(2\alpha)}(n, k_1, k_2, \dots, k_{|\alpha|}) \chi_{sign(2\beta)}(n, l_1, l_2, \dots, l_{2|\beta|}) \phi(n, m_1, m_2, \dots, m_{|\gamma|}) q^{\epsilon_\gamma} \Omega(n, s, k_{2|\alpha|}, l_{2|\beta|}) \psi(n, t, a, m_{|\gamma|}) q^{-(n-t)} \frac{(1-q^{n+1})(1-q^{n+2})}{(1-q^{t+1})(1-q^{t+2})},$$

$$(4.3) \quad J_{(n,0)}^{\mathfrak{sl}_3}(P(\downarrow 2\alpha+1 \downarrow, \uparrow 2\beta+1 \uparrow, \downarrow 2\gamma \uparrow); q) = \sum_{0 \leq k_{|2\alpha+1|} \leq k_{|2\alpha|} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{|2\beta+1|} \leq l_{|2\beta|} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_{|\gamma|} \leq m_{|\gamma|-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{|2\alpha+1|}, l_{|2\beta+1|}\}}^{\min\{k_{|2\alpha+1|}+l_{|2\beta+1|}, n\}} \sum_{a=s}^n \sum_{t=\max\{a, s\}}^{\min\{a+s, n\}} (q^{\frac{n^2+3n}{3}})^{-(2\alpha+2\beta-2\gamma+2)} \\ \chi_{sign(2\alpha+1)}(n, k_1, k_2, \dots, k_{|2\alpha+1|}) \chi_{sign(2\beta+1)}(n, l_1, l_2, \dots, l_{|2\beta+1|}) \phi(n, m_1, m_2, \dots, m_{|\gamma|}) q^{\epsilon_\gamma} \\ \Omega(n, s, k_{|2\alpha+1|}, l_{|2\beta+1|}) \psi(n, a, m_{|\gamma|}, t) q^{-(n-t)} \frac{(1-q^{n+1})(1-q^{n+2})}{(1-q^{t+1})(1-q^{t+2})},$$

(4.4)

$$\begin{aligned}
& J_{(n,0)}^{\mathfrak{sl}_3}(P(\downarrow 2\alpha+1 \downarrow, \uparrow 2\beta \uparrow, \downarrow 2\gamma \uparrow); q) \\
&= \sum_{0 \leq k_{|2\alpha+1|} \leq k_{|2\alpha|} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{|\beta|} \leq l_{|\beta|-1} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_{|\gamma|} \leq m_{|\gamma|-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{|2\alpha+1|}, l_{|\beta|}\}}^{\min\{k_{|2\alpha+1|} + l_{|\beta|}\}} \sum_{a=0}^{n-s} \sum_{t=\{a,s\}}^{\min\{a+s,n\}} (q^{\frac{n^2+3n}{3}})^{(-2\alpha-2\beta+2\gamma-1)} \\
&\quad \chi_{\text{sign}(2\alpha+1)}(n, k_1, k_2, \dots, k_{|2\alpha+1|}) \chi_{\text{sign}(2\beta)}(n, l_1, l_2, \dots, l_{|\beta|}) \phi(n, m_1, m_2, \dots, m_{|\gamma|})_{q^{\epsilon_{|\gamma|}}} \Omega(n, k_{|2\alpha+1|}, l_{|\beta|}, t) \\
&\quad \psi(n, t, a, m_{|\gamma|}) q^{-(n-t)} \frac{(1-q^{n+1})(1-q^{n+2})}{(1-q^{t+1})(1-q^{t+2})},
\end{aligned}$$

(4.5)

$$\begin{aligned}
& J_{(n,0)}^{\mathfrak{sl}_3}(P(\downarrow 2\alpha+1 \downarrow, \uparrow 2\beta \downarrow, \uparrow 2\gamma \uparrow); q) \\
&= \sum_{0 \leq k_{|2\alpha+1|} \leq k_{|2\alpha|} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{|\beta|} \leq l_{|\beta|-1} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_{|\gamma|} \leq m_{|\gamma|-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{|2\alpha|}, l_{|\beta|}\}}^{\min\{k_{|2\alpha|} + l_{|\beta|}\}} \sum_{a=0}^{n-s} \sum_{t=\{a,s\}}^{\min\{a+s,n\}} (q^{\frac{n^2+3n}{3}})^{(-2\alpha-2\beta+2\gamma-1)} \\
&\quad \chi_{\text{sign}(2\alpha+1)}(n, k_1, k_2, \dots, k_{|2\alpha+1|}) \phi(n, l_1, l_2, \dots, l_{|\beta|})_{q^{\epsilon_\beta}} \chi_{\text{sign}(2\gamma)}(n, m_1, m_2, \dots, m_{|\gamma|})_{q^{\epsilon_{|\gamma|}}} \Omega(n, k_{|2\alpha|}, l_{|\beta|}, t) \\
&\quad \psi(n, t, a, m_{|\gamma|}) q^{-(n-t)} \frac{(1-q^{n+1})(1-q^{n+2})}{(1-q^{t+1})(1-q^{t+2})}.
\end{aligned}$$

Proof. We first prove (4.2).

$$\begin{aligned}
& J_{(n,0)}^{\mathfrak{sl}_3}(P(\downarrow 2\alpha+1 \downarrow, \uparrow 2\beta+1 \uparrow, \downarrow 2\gamma \uparrow); q) \\
&= (\text{Definition 3.3}) (q^{\frac{n^2+3n}{3}})^{-(2\alpha+2\beta-2\gamma+2)} \left\langle \begin{array}{c} n \\ \swarrow \uparrow \downarrow \uparrow \downarrow \uparrow \downarrow \\ 2\alpha+1 & 2\beta+1 & 2\gamma \\ \uparrow \downarrow \uparrow \downarrow \uparrow \downarrow \uparrow \downarrow \\ \searrow \end{array} \right\rangle_3 / \Delta(n, 0) \\
&= (\text{Proposition 3.1}) (q^{\frac{n^2+3n}{3}})^{-(2\alpha+2\beta-2\gamma+2)} \sum_{0 \leq k_{|2\alpha+1|} \leq k_{|2\alpha|} \leq \dots \leq k_1 \leq n} \chi_{\text{sign}(2\alpha+1)}(n, k_1, k_2, \dots, k_{|2\alpha+1|}) \\
&\quad \left\langle \begin{array}{c} k_{|2\alpha+1|} \\ \swarrow \uparrow \downarrow \uparrow \downarrow \uparrow \downarrow \\ 2\beta+1 & 2\gamma \\ \uparrow \downarrow \uparrow \downarrow \uparrow \downarrow \uparrow \downarrow \\ \searrow \end{array} \right\rangle_3 / \Delta(n, 0) \\
&= (\text{Proposition 3.1, Theorem 2.9}) (q^{\frac{n^2+3n}{3}})^{-(2\alpha+2\beta-2\gamma+2)} \sum_{0 \leq k_{|2\alpha+1|} \leq k_{|2\alpha|} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{|2\beta+1|} \leq l_{|2\beta|} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{|\gamma|-1} \leq \dots \leq m_1 \leq n} \\
&\quad \times \chi_{\text{sign}(2\alpha+1)}(n, k_1, k_2, \dots, k_{|2\alpha+1|}) \chi_{\text{sign}(2\beta+1)}(n, l_1, l_2, \dots, l_{|2\beta+1|}) \phi(n, m_1, m_2, \dots, m_\gamma)_{q^{\epsilon_\gamma}}
\end{aligned}$$

$$\begin{aligned}
& \times \left\langle k_{|2\alpha+1|}, l_{|2\beta+1|}, m_{|\gamma|} \right\rangle_3 / \Delta(n, 0) \\
& \stackrel{(2.2)}{=} (q^{\frac{n^2+3n}{3}})^{-(2\alpha+2\beta-2\gamma+2)} \sum_{0 \leq k_{|2\alpha+1|} \leq k_{|2\alpha|} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{|2\beta+1|} \leq l_{|2\beta|} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \\
& \quad \times \chi_{sign(2\alpha+1)}(n, k_1, k_2, \dots, k_{|2\alpha+1|}) \chi_{sign(2\beta+1)}(n, l_1, l_2, \dots, l_{|2\beta+1|}) \phi(n, m_1, m_2, \dots, m_\gamma) q^{\epsilon_\gamma} \\
& \quad \times \left\langle k_{|2\alpha+1|}, l_{|2\beta+1|}, m_{|\gamma|} \right\rangle_3 / \Delta(n, 0) \\
& \stackrel{(Propositon 3.6)}{=} (q^{\frac{n^2+3n}{3}})^{-(2\alpha+2\beta-2\gamma+2)} \sum_{0 \leq k_{|2\alpha+1|} \leq k_{|2\alpha|} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{|2\beta+1|} \leq l_{|2\beta|} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{|2\alpha+1|}, l_{|2\beta+1|}\}}^{\min\{k_{|2\alpha+1|}+l_{|2\beta+1|}, n\}} \sum_{a=s}^n \\
& \quad \times \chi_{sign(2\alpha+1)}(n, k_1, k_2, \dots, k_{|2\alpha+1|}) \chi_{sign(2\beta+1)}(n, l_1, l_2, \dots, l_{|2\beta+1|}) \phi(n, m_1, m_2, \dots, m_\gamma) q^{\epsilon_\gamma} \left\langle \begin{array}{c} n \\ a \\ n-a \\ n-m_{|\gamma|} \end{array} \right\rangle_3 / \Delta(n, 0) \\
& \stackrel{(2.2)}{=} (q^{\frac{n^2+3n}{3}})^{-(2\alpha+2\beta-2\gamma+2)} \sum_{0 \leq k_{|2\alpha+1|} \leq k_{|2\alpha|} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{|2\beta+1|} \leq l_{|2\beta|} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{|2\alpha+1|}, l_{|2\beta+1|}\}}^{\min\{k_{|2\alpha+1|}+l_{|2\beta+1|}, n\}} \sum_{a=s}^n \\
& \quad \times \chi_{sign(2\alpha+1)}(n, k_1, k_2, \dots, k_{|2\alpha+1|}) \chi_{sign(2\beta+1)}(n, l_1, l_2, \dots, l_{|2\beta+1|}) \phi(n, m_1, m_2, \dots, m_\gamma) q^{\epsilon_\gamma} \\
& \quad \times \Omega(n, s, k_{|2\alpha+1|}, l_{|2\beta+1|}) \left\langle \begin{array}{c} n \\ a \\ n-a \\ n-m_{|\gamma|} \end{array} \right\rangle_3 / \Delta(n, 0) \\
& \stackrel{(Theorem 2.10)}{=} (q^{\frac{n^2+3n}{3}})^{-(2\alpha+2\beta-2\gamma+2)} \sum_{0 \leq k_{|2\alpha+1|} \leq k_{|2\alpha|} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{|2\beta+1|} \leq l_{|2\beta|} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{|2\alpha+1|}, l_{|2\beta+1|}\}}^{\min\{k_{|2\alpha+1|}+l_{|2\beta+1|}, n\}}
\end{aligned}$$

$$\begin{aligned}
& \sum_{a=s}^n \sum_{t=\max\{s, m_\gamma\}}^{\min\{s+m_\gamma, n\}} \chi_{sign(2\alpha+1)}(n, k_1, k_2, \dots, k_{|2\alpha+1|}) \chi_{sign(2\beta+1)}(n, l_1, l_2, \dots, l_{|2\beta+1|}) \phi(n, m_1, m_2, \dots, m_\gamma) q^{e_\gamma} \\
& \times \Omega(n, s, k_{|2\alpha+1|}, l_{|2\beta+1|}) \psi(n, a, m_\gamma, t) \left\langle \begin{array}{c} \text{Diagram} \\ \text{with } n-t \text{ segments} \end{array} \right\rangle_3 / \Delta(n, 0) \\
& \stackrel{(2.8)}{=} \sum_{0 \leq k_{2|\alpha|} \leq k_{2|\alpha|-1} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2|\beta|} \leq l_{2|\beta|-1} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{2|\alpha|}, l_{2|\beta|}\}}^{\min\{k_{2|\alpha|}+l_{2|\beta|}, n\}} \sum_{a=0}^{n-s} \sum_{t=\max\{a, m_{|\gamma|}\}}^{\min\{a+t+m_{|\gamma|}, n\}} (q^{\frac{n^2+3n}{3}})^{2\alpha+2\beta-2\gamma} \\
& \times \chi_{sign(2\alpha)}(n, k_1, k_2, \dots, k_{|\alpha|}) \chi_{sign(2\beta)}(n, l_1, l_2, \dots, l_{|\beta|}) \phi(n, m_1, m_2, \dots, m_{|\gamma|}) q^{e_\gamma} \Omega(n, s, k_{2|\alpha|}, l_{2|\beta|}) \psi(n, t, a, m_{|\gamma|}) \\
& \times q^{-(n-t)} \frac{(1-q^{n+1})(1-q^{n+2})}{(1-q^{t+1})(1-q^{t+2})}
\end{aligned}$$

We can prove (4.1), (4.3) and (4.5) in a similar way to (4.2). \square

Remark 4.2. For $n=1, 2, \dots, 10$, Theorem 3.1 is equal to Theorem 5.7 in [Yua17] for $4_1, 6_2$ and 8_2 by using Mathematica. For example,

$$\begin{aligned}
J_{(3,0)}^{\mathfrak{sl}_3}(4; q) = & q^{15} - q^{13} - q^{12} - q^{11} + 2q^{10} + q^9 - 2q^7 - q^6 + 4q^5 + 2q^4 - 2q^3 - 4q^2 + 5 - 4q^{-2} - 2q^{-3} \\
& + 2q^{-4} + 4q^{-5} - q^{-6} - 2q^{-7} + q^{-9} + 2q^{-10} - q^{-11} - q^{-12} - q^{-13} + q^{-15}.
\end{aligned}$$

5. THE TAILS OF THE ONE-ROW COLORED \mathfrak{sl}_3 JONES POLYNOMIALS FOR $P(2\alpha+1, 2\beta+1, 2\gamma)$ PRETZEL KNOTS

In this section, we use Theorem 4.1 to show the existence of the tails of the one-row colored \mathfrak{sl}_3 Jones polynomials for alternating pretzel knots $P(2\alpha+1, 2\beta+1, 2\gamma)$.

Definition 5.1. For any formal Laurent series $f(q)$, we define $\hat{f}(q)$ by

$$\hat{f}(q) = \pm q^{-\text{mindeg}(f(q))} f(q) = \sum_{i=0}^{\infty} a_i q^i \in \mathbb{Z}[[q]].$$

In the above normalization, we determine $a_0 > 0$.

Corollary 5.2. For positive integers α, β and γ ,

$$\text{mindeg}(J_{(n,0)}^{\mathfrak{sl}_3}(P(\downarrow 2\alpha+1 \downarrow, \uparrow 2\beta+1 \uparrow, \downarrow 2\gamma \uparrow); q)) = -(\alpha+\beta+1)n^2 - (3\alpha+3\beta+2)n$$

Proof.

$$\begin{aligned}
& \text{mindeg}(J_{(n,0)}^{\text{sl}_3}(P(\downarrow 2\alpha+1 \downarrow, \uparrow 2\beta+1 \uparrow, \downarrow 2\gamma \uparrow); q)) \\
&= \text{mindeg} \left(\sum_{0 \leq k_{2\alpha+1} \leq k_{2\alpha} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2\beta+1} \leq l_{2\beta} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1} + l_{2\beta+1}, n\}} \sum_{a=s}^n \sum_{t=\max\{a, m_\gamma\}}^{\min\{a+m_\gamma, n\}} \right. \\
&\quad \left. (q^{\frac{n^2+3n}{3}})^{-(2\alpha+2\beta-2\gamma+2)} q^{-\frac{1}{6}(n^2+3n)(2\alpha+1)} q^{\frac{1}{2}(n-k_{2\alpha+1})} q^{\sum_{i=1}^{2\alpha+1} \frac{1}{2}(k_i^2+k_i)} (q)_n \times q^{-\frac{1}{6}(n^2+3n)(2\beta+1)} q^{\frac{1}{2}(n-l_{2\beta+1})} q^{\sum_{i=1}^{2\beta+1} \frac{1}{2}(l_i^2+l_i)} (q)_n \right. \\
&\quad \left. \times (q)^{-\frac{2\gamma}{3}(n^2+3n)} (q)^{n-m_\gamma} (q)^{\sum_{i=1}^\gamma (m_i^2+2m_i)} (q)_n^2 \right. \\
&\quad \left. \times \left(\prod_{i=1}^\gamma (q)_{m_{i-1}-m_i} \right) (q)_{m_\gamma}^2 \right. \\
&\quad \left. \times \frac{q^{-\frac{k_{2\alpha+1}+l_{2\beta+1}}{2}+s} q^{(s+1)(s-k_{2\alpha+1}-l_{2\beta+1})+k_{2\alpha+1}l_{2\beta+1}} (1-q^{n+1-k_{2\alpha+1}})(1-q^{n+1-l_{2\beta+1}})(q)_{k_{2\alpha+1}}(q)_{l_{2\beta+1}} (q)_{n-k_{2\alpha+1}}^2 (q)_{n-l_{2\beta+1}}^2 (q)_{2n-s+2}}{(1-q^{n+1-s})^2 (q)_{n-s}^2 (q)_{s-k_{2\alpha+1}} (q)_{s-l_{2\beta+1}} (q)_{2n-k_{2\alpha+1}-l_{2\beta+1}+2} (q)_{-s+k_{2\alpha+1}+l_{2\beta+1}} \right. \\
&\quad \left. \times \frac{q^{(t+1)(t-a-m_\gamma)+am_\gamma} (q)_a (q)_{m_\gamma} (q)_{n-a}^2 (q)_{n-m_\gamma}^2 (q)_{2n-t+2}}{(q)_n^2 (q)_{n-t}^2 (q)_{t-a} (q)_{t-m_\gamma} (q)_{2n-a-m_\gamma+2} (q)_{-t+a+m_\gamma}} q^{-(n-t)} \frac{(1-q^{n+1})(1-q^{n+2})}{(1-q^{t+1})(1-q^{t+2})} \right) \\
&= \text{mindeg}(q^{-(\alpha+\beta+1)n^2-(3\alpha+3\beta+2)n} \sum_{0 \leq k_{2\alpha+1} \leq k_{2\alpha} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2\beta+1} \leq l_{2\beta} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1} + l_{2\beta+1}, n\}} \\
&\quad \sum_{a=s}^n \sum_{t=\max\{a, m_\gamma\}}^{\min\{a+m_\gamma, n\}} q^{\frac{1}{2}(-k_{2\alpha+1})} q^{\sum_{i=1}^{2\alpha+1} \frac{1}{2}(k_i^2+k_i)} q^{\frac{1}{2}(-l_{2\beta+1})} q^{\sum_{i=1}^{2\beta+1} \frac{1}{2}(l_i^2+l_i)} q^{-m_\gamma} q^{\sum_{i=1}^\gamma (m_i^2+2m_i)} q^{-\frac{k_{2\alpha+1}+l_{2\beta+1}}{2}+s} \\
&\quad \times q^{(s+1)(s-k_{2\alpha+1}-l_{2\beta+1})+k_{2\alpha+1}l_{2\beta+1}} q^{(t+1)(t-a-m_\gamma)+am_\gamma} q^t) \\
&= \text{mindeg}(q^{-(\alpha+\beta+1)n^2-(3\alpha+3\beta+2)n} \sum_{0 \leq k_{2\alpha+1} \leq k_{2\alpha} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2\beta+1} \leq l_{2\beta} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1} + l_{2\beta+1}, n\}} \\
&\quad \sum_{a=s}^n \sum_{t=\max\{a, m_\gamma\}}^{\min\{a+m_\gamma, n\}} q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2+k_i)} q^{\sum_{i=1}^{2\beta} \frac{1}{2}(l_i^2+l_i)} q^{m_\gamma^2+m_\gamma} q^{\sum_{i=1}^{\gamma-1} (m_i^2+2m_i)} q^{\frac{k_{2\alpha+1}^2+l_{2\beta+1}^2-(k_{2\alpha+1}+l_{2\beta+1})}{2}} \\
&\quad \times q^{s+(s+1)(s-k_{2\alpha+1}-l_{2\beta+1})+k_{2\alpha+1}l_{2\beta+1}} q^{t+(t+1)(t-a-m_\gamma)+am_\gamma} \\
&= -(\alpha+\beta+1)n^2 - (3\alpha+3\beta+2)n
\end{aligned}$$

We use $\frac{1}{1-q} = 1 + q + q^2 + \dots$ in the second equality. If $l_{2\beta+1} \leq k_{2\alpha+1} \leq s$, then

$$\begin{aligned}
& s + (s+1)(s - (k_{2\alpha+1} + l_{2\beta+1})) + k_{2\alpha+1}l_{2\beta+1} \\
(5.1) \quad &= s^2 + s(2 - (k_{2\alpha+1} + l_{2\beta+1}) - k_{2\alpha+1} - l_{2\beta+1} + k_{2\alpha+1}l_{2\beta+1}) \\
&= (s - k_{2\alpha+1} + 1)(s - l_{2\beta+1} + 1) - 1 \\
&\geq (k_{2\alpha+1} - l_{2\beta+1} + 1) - 1 \geq 0
\end{aligned}$$

The same is true for $k_{2\alpha+1} \leq l_{2\beta+1} \leq s$. In the above last inequality, we have equality if and only if $l_{2\beta+1} = k_{2\alpha+1} = s$ holds. By the same method as the above, if $a \leq m_\gamma \leq t$ and $m_\gamma \leq a \leq t$, then we can obtain

$$(5.2) \quad t + (t+1)(t - (a + m_\gamma)) + am_\gamma \geq 0.$$

The equality sign is valid for $a = m_\gamma = t$ and $m_\gamma = a = t$. By (5.1), (5.2), $k_{2\alpha+1}^2 - k_{2\alpha+1} \geq 0$ and $\frac{2}{2\beta+1} - l_{2\beta+1} \geq 0$, the fourth equality holds. \square

Theorem 5.3. Let α, β , and γ be positive integers. For oriented pretzel knots $P(\downarrow 2\alpha + 1 \downarrow, \uparrow 2\beta + 1 \uparrow, \downarrow 2\gamma \uparrow)$, there exists $\mathcal{T}^{\mathfrak{sl}_3}(P(\downarrow 2\alpha + 1 \downarrow, \uparrow 2\beta + 1 \uparrow, \downarrow 2\gamma \uparrow); q)$ in $\mathbb{Z}[[q]]$ such that

$$\mathcal{T}^{\mathfrak{sl}_3}(P(\downarrow 2\alpha + 1 \downarrow, \uparrow 2\beta + 1 \uparrow, \downarrow 2\gamma \uparrow); q) - J_{(n,0)}^{\hat{\mathfrak{sl}}_3}(P(\downarrow 2\alpha + 1 \downarrow, \uparrow 2\beta + 1 \uparrow, \downarrow 2\gamma \uparrow); q) \in q^{n+1}\mathbb{Z}[[q]].$$

Remark 5.4. For a simple Lie algebra \mathfrak{g} , Le[Le00] proved integrality theorem for a quantum \mathfrak{g} invariant. This says $\hat{J}_{(n,0)}^{\mathfrak{sl}_3}(L; q)$ belongs to $\mathbb{Z}[[q]]$.

Definition 5.5. For two Laurent series $\hat{f}(q), \hat{g}(q) \in \mathbb{Z}[[q]]$, we define

$$f\hat{g}(q) \equiv_n g\hat{f}(q)$$

by $f\hat{g}(q) = g\hat{f}(q)$ in $\mathbb{Z}[[q]]/q^{n+1}\mathbb{Z}[[q]]$ for all n .

Proof of Theorem 5.3. There exists $\mathcal{T}^{\mathfrak{sl}_3}(P(\downarrow 2\alpha + 1 \downarrow, \uparrow 2\beta + 1 \uparrow, \downarrow 2\gamma \uparrow); q)$ if and only if

$$(5.3) \quad \hat{J}_{(n,0)}^{\mathfrak{sl}_3}(P(\downarrow 2\alpha + 1 \downarrow, \uparrow 2\beta + 1 \uparrow, \downarrow 2\gamma \uparrow); q) \equiv_{n+1} J_{(n+1,0)}^{\hat{\mathfrak{sl}}_3}(P(\downarrow 2\alpha + 1 \downarrow, \uparrow 2\beta + 1 \uparrow, \downarrow 2\gamma \uparrow); q))$$

for all n . First, we can easily obtain the following.

$$(5.4) \quad \begin{aligned} & \hat{J}_{(n,0)}^{\mathfrak{sl}_3}(P(\downarrow 2\alpha + 1 \downarrow, \uparrow 2\beta + 1 \uparrow, \downarrow 2\gamma \uparrow); q)) \\ & \equiv_{n+1} \sum_{0 \leq k_{2\alpha}+1 \leq k_{2\alpha} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2\beta}+1 \leq l_{2\beta} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1}+l_{2\beta+1}, n\}} \sum_{a=s}^n \sum_{t=\max\{a, m_\gamma\}}^{\min\{a+m_\gamma, n\}} \\ & \quad \frac{(-1)^{\sum_{i=1}^{2\alpha+1} k_i} q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2+k_i)}}{\left(\prod_{i=2}^{2\alpha+1} (q)_{k_{i-1}-k_i}\right)(q)_{k_{2\alpha+1}}} \times \frac{(-1)^{\sum_{i=1}^{2\beta+1} k_i} q^{\sum_{i=1}^{2\beta} \frac{1}{2}(l_i^2+l_i)}}{\left(\prod_{i=2}^{2\beta+1} (q)_{l_{i-1}-l_i}\right)(q)_{l_{2\beta+1}}} \times \frac{(q)_{m_\gamma^2+m_\gamma} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2+2m_i)} (q)_n}{\left(\prod_{i=2}^{\gamma} (q)_{m_{i-1}-m_i}\right)(q)_{m_\gamma^2}} \\ & \quad \times \frac{q^{\frac{(k_{2\alpha+1}^2+l_{2\beta+1}^2)-(k_{2\alpha+1}+l_{2\beta+1})}{2}}}{(q)_{s-k_{2\alpha+1}} (q)_{s-l_{2\beta+1}} (q)_{-s+k_{2\alpha+1}+l_{2\beta+1}}} q^{s+(s+1)(s-k_{2\alpha+1}-l_{2\beta+1})+k_{2\alpha+1}l_{2\beta+1}} (q)_{k_{2\alpha+1}} (q)_{l_{2\beta+1}} \\ & \quad \times \frac{q^{t+(t+1)(t-a-m_\gamma)+am_\gamma}}{(q)_{t-a} (q)_{t-m_\gamma} (q)_{-t+a+m_\gamma}} \frac{1}{(1-q^{t+1})(1-q^{t+2})} \end{aligned}$$

Here,

$$(5.5) \quad \begin{aligned} \frac{q^s}{1-q^{n+2-s}} &= q^s (1 + q^{n+2-s} + q^{2(n+2-s)} + \dots) \\ &= q^s + q^{n+2} + (\text{higher order terms}) \\ &\equiv_{n+1} 0 \end{aligned}$$

We use a deformation similar to the above equation repeatedly. We next consider $\hat{J}_{(n+1,0)}^{\text{sl}_3}(P(\downarrow 2\alpha+1 \downarrow, \uparrow 2\beta+1 \uparrow, \downarrow 2\gamma \uparrow); q))$.

$$\begin{aligned}
& \hat{J}_{(n+1,0)}^{\text{sl}_3}(P(\downarrow 2\alpha+1 \downarrow, \uparrow 2\beta+1 \uparrow, \downarrow 2\gamma \uparrow); q)) \\
= & \sum_{0 \leq k_{2\alpha+1} \leq k_{2\alpha} \leq \dots \leq k_1 \leq n+1} \sum_{0 \leq l_{2\beta+1} \leq l_{2\beta} \leq \dots \leq l_1 \leq n+1} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n+1} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1}+l_{2\beta+1}, n+1\}} \sum_{a=s}^{n+1} \sum_{t=\max\{a, m_\gamma\}}^{\min\{a+m_\gamma, n+1\}} \\
& \times \frac{(-1)^{\sum_{i=1}^{2\alpha+1} k_i} q^{\frac{1}{2}k_{2\alpha+1}^2} q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2+k_i)} (q)_{n+1}}{\left(\prod_{i=1}^{2\alpha+1} (q)_{k_{i-1}-k_i}\right) (q)_{n+1-k_1} (q)_{k_{2\alpha+1}}} \times \frac{(-1)^{\sum_{i=1}^{2\beta+1} l_i} q^{\frac{1}{2}l_{2\beta+1}^2} q^{\sum_{i=1}^{2\beta} \frac{1}{2}(l_i^2+l_i)} (q)_{n+1}}{\left(\prod_{i=1}^{2\beta+1} (q)_{l_{i-1}-l_i}\right) (q)_{n+1-l_1} (q)_{l_{2\beta+1}}} \times \frac{(q)^{m_\gamma^2+m_\gamma} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2+2m_i)} (q)_{n+1}^2}{\left(\prod_{i=1}^{\gamma} (q)_{m_{i-1}-m_i}\right) (q)_{n+1-m_1} (q)_{m_\gamma}^2} \\
& \times \frac{q^{-\frac{k_{2\alpha+1}+l_{2\beta+1}}{2}+s} q^{(s+1)(s-k_{2\alpha+1}-l_{2\beta+1})+k_{2\alpha+1}l_{2\beta+1}} (1-q^{n+2-k_{2\alpha+1}}) (1-q^{n+2-l_{2\beta+1}}) (q)_{k_{2\alpha+1}} (q)_{l_{2\beta+1}} (q)_{n+1-k_{2\alpha+1}}^2 (q)_{n+1-l_{2\beta+1}}^2}{(1-q^{n+2-s})^2 (q)_{n+1-s}^2 (q)_{n+1-s}^2 (q)_{s-k_{2\alpha+1}} (q)_{s-l_{2\beta+1}} (q)_{2(n+1)-k_{2\alpha+1}-l_{2\beta+1}+2} (q)_{-s+k_{2\alpha+1}+l_{2\beta+1}}} \\
& \times \frac{q^{(t+1)(t-a-m_\gamma)+am_\gamma} (q)_a (q)_{m_\gamma} (q)_{n+1-a}^2 (q)_{n+1-m_\gamma}^2 (q)_{2(n+1)-t+2}}{(q)_{2(n+1)-s+2} (q)_{n+1}^2 (q)_{n+1-t}^2 (q)_{t-a} (q)_{t-m_\gamma} (q)_{2(n+1)-a-m_\gamma+2} (q)_{-t+a+m_\gamma}} q^{-(n+1-t)} \frac{(1-q^{n+2}) (1-q^{n+3})}{(1-q^{t+1}) (1-q^{t+2})} \\
\equiv_{n+1} & \sum_{0 \leq k_{2\alpha+1} \leq k_{2\alpha} \leq \dots \leq k_1 \leq n+1} \sum_{0 \leq l_{2\beta+1} \leq l_{2\beta} \leq \dots \leq l_1 \leq n+1} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n+1} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1}+l_{2\beta+1}, n+1\}} \sum_{a=s}^{n+1} \sum_{t=\max\{a, m_\gamma\}}^{\min\{a+m_\gamma, n+1\}} \\
& \frac{(-1)^{\sum_{i=1}^{2\alpha+1} k_i} q^{\frac{1}{2}k_{2\alpha+1}^2} q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2+k_i)} }{\left(\prod_{i=2}^{2\alpha+1} (q)_{k_{i-1}-k_i}\right) (q)_{k_{2\alpha+1}}} \times \frac{(-1)^{\sum_{i=1}^{2\beta+1} l_i} q^{\frac{1}{2}l_{2\beta+1}^2} q^{\sum_{i=1}^{2\beta} \frac{1}{2}(l_i^2+l_i)} }{\left(\prod_{i=2}^{2\beta+1} (q)_{l_{i-1}-l_i}\right) (q)_{l_{2\beta+1}}} \times \frac{(q)^{m_\gamma^2+m_\gamma} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2+2m_i)} (q)_n}{\left(\prod_{i=2}^{\gamma} (q)_{m_{i-1}-m_i}\right) (q)_{m_\gamma}^2} \\
& \times \frac{q^{-\frac{k_{2\alpha+1}+l_{2\beta+1}}{2}+s} q^{(s+1)(s-k_{2\alpha+1}-l_{2\beta+1})+k_{2\alpha+1}l_{2\beta+1}} (q)_{k_{2\alpha+1}} (q)_{l_{2\beta+1}}}{(q)_{s-k_{2\alpha+1}} (q)_{s-l_{2\beta+1}} (q)_{-s+k_{2\alpha+1}+l_{2\beta+1}}} \times \frac{q^{t+(t+1)(t-a-m_\gamma)+am_\gamma} (q)_a (q)_{m_\gamma}}{(q)_{t-a} (q)_{t-m_\gamma} (q)_{-t+a+m_\gamma}} \frac{1}{(1-q^{t+1}) (1-q^{t+2})} \\
= & \sum_{0 \leq k_{2\alpha+1} \leq k_{2\alpha} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2\beta+1} \leq l_{2\beta} \leq \dots \leq l_1 \leq n+1} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n+1} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1}+l_{2\beta+1}, n+1\}} \sum_{a=s}^{n+1} \sum_{t=\max\{a, m_\gamma\}}^{\min\{a+m_\gamma, n+1\}} \\
& \frac{(-1)^{\sum_{i=1}^{2\alpha+1} k_i} q^{\frac{1}{2}k_{2\alpha+1}^2} q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2+k_i)} }{\left(\prod_{i=2}^{2\alpha+1} (q)_{k_{i-1}-k_i}\right) (q)_{k_{2\alpha+1}}} \times \frac{(-1)^{\sum_{i=1}^{2\beta+1} l_i} q^{\frac{1}{2}l_{2\beta+1}^2} q^{\sum_{i=1}^{2\beta} \frac{1}{2}(l_i^2+l_i)} }{\left(\prod_{i=2}^{2\beta+1} (q)_{l_{i-1}-l_i}\right) (q)_{l_{2\beta+1}}} \times \frac{(q)^{m_\gamma^2+m_\gamma} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2+2m_i)} (q)_n}{\left(\prod_{i=2}^{\gamma} (q)_{m_{i-1}-m_i}\right) (q)_{m_\gamma}^2} \\
& \times \frac{q^{-\frac{k_{2\alpha+1}+l_{2\beta+1}}{2}+s} q^{(s+1)(s-k_{2\alpha+1}-l_{2\beta+1})+k_{2\alpha+1}l_{2\beta+1}} (q)_{k_{2\alpha+1}} (q)_{l_{2\beta+1}}}{(q)_{s-k_{2\alpha+1}} (q)_{s-l_{2\beta+1}} (q)_{-s+k_{2\alpha+1}+l_{2\beta+1}}} \times \frac{q^{t+(t+1)(t-a-m_\gamma)+am_\gamma} (q)_a (q)_{m_\gamma}}{(q)_{t-a} (q)_{t-m_\gamma} (q)_{-t+a+m_\gamma}} \frac{1}{(1-q^{t+1}) (1-q^{t+2})} \\
& + q^{n+1+\frac{1}{2}(n^2+n)} \sum_{0 \leq k_{2\alpha+1} \leq k_{2\alpha} \leq \dots \leq k_2 \leq n+1} \sum_{0 \leq l_{2\beta+1} \leq l_{2\beta} \leq \dots \leq l_1 \leq n+1} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n+1} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1}+l_{2\beta+1}, n+1\}} \\
& \sum_{a=s}^{n+1} \sum_{t=\max\{a, m_\gamma\}}^{\min\{a+m_\gamma, n+1\}} \frac{(-1)^{\sum_{i=2}^{2\alpha+1} k_i} q^{\sum_{i=2}^{2\alpha} \frac{1}{2}(k_i^2+k_i)} }{\left(\prod_{i=2}^{2\alpha+1} (q)_{k_{i-1}-k_i}\right) (q)_{k_{2\alpha+1}}} \times \frac{(-1)^{\sum_{i=1}^{2\beta+1} l_i} q^{\sum_{i=1}^{2\beta} \frac{1}{2}(l_i^2+l_i)} }{\left(\prod_{i=2}^{2\beta+1} (q)_{l_{i-1}-l_i}\right) (q)_{l_{2\beta+1}}} \times \frac{(q)^{m_\gamma^2+m_\gamma} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2+2m_i)} (q)_n}{\left(\prod_{i=2}^{\gamma} (q)_{m_{i-1}-m_i}\right) (q)_{m_\gamma}^2}
\end{aligned}$$

$$\begin{aligned} & \times \frac{q^{\frac{(k_{2\alpha+1}^2 + l_{2\beta+1}^2) - (k_{2\alpha+1} + l_{2\beta+1})}{2}} q^{s+(s+1)(s-k_{2\alpha+1}-l_{2\beta+1}) + k_{2\alpha+1}l_{2\beta+1}} (q)_{k_{2\alpha+1}} (q)_{l_{2\beta+1}}}{(q)_{s-k_{2\alpha+1}} (q)_{s-l_{2\beta+1}} (q)_{-s+k_{2\alpha+1}+l_{2\beta+1}}} \\ & \times \frac{q^{t+(t+1)(t-a-m_\gamma)+am_\gamma} (q)_a (q)_{m_\gamma}}{(q)_{t-a} (q)_{t-m_\gamma} (q)_{-t+a+m_\gamma}} \frac{1}{(1-q^{t+1})(1-q^{t+2})} \end{aligned}$$

By (5.1), (5.2), $k_{2\alpha+1}^2 - k_{2\alpha+1} \geq 0$ and $\frac{1}{2}l_{2\beta+1} - l_{2\beta+1} \geq 0$,

$$\begin{aligned} & q^{n+1+\frac{1}{2}(n^2+n)} \sum_{0 \leq k_{2\alpha+1} \leq k_{2\alpha} \leq \dots \leq k_2 \leq n+1} \sum_{0 \leq l_{2\beta+1} \leq l_{2\beta} \leq \dots \leq l_1 \leq n+1} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n+1} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1}+l_{2\beta+1}, n+1\}} \\ & \sum_{a=s}^{n+1} \sum_{t=\max\{a, m_\gamma\}}^{\min\{a+m_\gamma, n+1\}} \frac{(-1)^{\sum_{i=1}^{2\alpha} k_i} q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2 + k_i)}}{\left(\prod_{i=2}^{2\alpha} (q)_{k_{i-1}-k_i}\right) (q)_{k_{2\alpha+1}}} \times \frac{(-1)^{\sum_{i=1}^{2\beta+1} l_i} q^{\sum_{i=1}^{2\beta+1} \frac{1}{2}(l_i^2 + l_i)}}{\left(\prod_{i=2}^{2\beta+1} (q)_{l_{i-1}-l_i}\right) (q)_{l_{2\beta+1}}} \times \frac{(q)^{m_\gamma^2 + m_\gamma} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2 + 2m_i)} (q)_n}{\left(\prod_{i=2}^{\gamma} (q)_{m_{i-1}-m_i}\right) (q)_{m_\gamma}^2} \\ & \times \frac{q^{\frac{(k_{2\alpha+1}^2 + l_{2\beta+1}^2) - (k_{2\alpha+1} + l_{2\beta+1})}{2}} q^{s+(s+1)(s-k_{2\alpha+1}-l_{2\beta+1}) + k_{2\alpha+1}l_{2\beta+1}} (q)_{k_{2\alpha+1}} (q)_{l_{2\beta+1}}}{(q)_{s-k_{2\alpha+1}} (q)_{s-l_{2\beta+1}} (q)_{2(n+1)-k_{2\alpha+1}-l_{2\beta+1}+2} (q)_{-s+k_{2\alpha+1}+l_{2\beta+1}}} \\ & \times \frac{q^{t+(t+1)(t-a-m_\gamma)+am_\gamma} (q)_a (q)_{m_\gamma}}{(q)_{t-a} (q)_{t-m_\gamma} (q)_{-t+a+m_\gamma}} \frac{1}{(1-q^{t+1})(1-q^{t+2})} \\ & = q^{n+1+\frac{1}{2}(n^2+n)} + (\text{high order terms}) \\ & \equiv_{n+1} 0 \end{aligned}$$

We consider $l_1 = n+1$ and $m_1 = n+1$ in the same way. Therefore,

$$\begin{aligned} (5.6) \quad & \hat{J}_{(n+1,0)}^{\mathfrak{sl}_3}(P(\downarrow 2\alpha+1 \downarrow, \uparrow 2\beta+1 \uparrow, \downarrow 2\gamma \uparrow; q)) \\ & \equiv_{n+1} \sum_{0 \leq k_{2\alpha+1} \leq k_{2\alpha} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2\beta+1} \leq l_{2\beta} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1}+l_{2\beta+1}, n+1\}} \sum_{a=s}^{n+1} \sum_{t=\max\{a, m_\gamma\}}^{\min\{a+m_\gamma, n+1\}} \\ & \frac{(-1)^{\sum_{i=1}^{2\alpha} k_i} q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2 + k_i)}}{\left(\prod_{i=2}^{2\alpha} (q)_{k_{i-1}-k_i}\right) (q)_{k_{2\alpha+1}}} \times \frac{(-1)^{\sum_{i=1}^{2\beta+1} l_i} q^{\sum_{i=1}^{2\beta+1} \frac{1}{2}(l_i^2 + l_i)} (q)_n}{\left(\prod_{i=2}^{2\beta+1} (q)_{l_{i-1}-l_i}\right) (q)_{l_{2\beta+1}}} \times \frac{(q)^{m_\gamma^2 + m_\gamma} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2 + 2m_i)} (q)_n}{\left(\prod_{i=2}^{\gamma} (q)_{m_{i-1}-m_i}\right) (q)_{m_\gamma}^2} \\ & \times \frac{q^{\frac{(k_{2\alpha+1}^2 + l_{2\beta+1}^2) - (k_{2\alpha+1} + l_{2\beta+1})}{2}} q^{s+(s+1)(s-k_{2\alpha+1}-l_{2\beta+1}) + k_{2\alpha+1}l_{2\beta+1}} (q)_{k_{2\alpha+1}} (q)_{l_{2\beta+1}}}{(q)_{s-k_{2\alpha+1}} (q)_{s-l_{2\beta+1}} (q)_{-s+k_{2\alpha+1}+l_{2\beta+1}}} \\ & \times \frac{q^{t+(t+1)(t-a-m_\gamma)+am_\gamma} (q)_a (q)_{m_\gamma}}{(q)_{t-a} (q)_{t-m_\gamma} (q)_{-t+a+m_\gamma}} \frac{1}{(1-q^{t+1})(1-q^{t+2})} \end{aligned}$$

If $k_{2\alpha+1} + l_{2\beta+1} \geq n + 1$, then

(RHS of (5.6))

$$\begin{aligned}
&= \sum_{0 \leq k_{2\alpha+1} \leq k_{2\alpha} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2\beta+1} \leq l_{2\beta} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^n \sum_{a=s}^{n+1} \sum_{t=\max\{a, m_\gamma\}}^{\min\{a+m_\gamma, n+1\}} \\
&\times \frac{(-1)^{\sum_{i=1}^{2\alpha+1} q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2+k_i)}}}{(\prod_{i=2}^{2\alpha+1} (q)_{k_{i-1}-k_i})(q)_{k_{2\alpha+1}}} \times \frac{(-1)^{\sum_{i=1}^{2\beta+1} q^{\sum_{i=1}^{2\beta} \frac{1}{2}(l_i^2+l_i)}}}{(\prod_{i=2}^{2\beta+1} (q)_{l_{i-1}-l_i})(q)_{l_{2\beta+1}}} \times \frac{(q)^{m_\gamma^2+m_\gamma} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2+2m_i)} (q)_n}{(\prod_{i=2}^\gamma (q)_{m_{i-1}-m_i})(q)_{m_\gamma}^2} \\
&\times \frac{q^{\frac{(k_{2\alpha+1}^2+l_{2\beta+1}^2)-(k_{2\alpha+1}+l_{2\beta+1})}{2}}}{(q)_{s-k_{2\alpha+1}} (q)_{s-l_{2\beta+1}} (q)_{-s+k_{2\alpha+1}+l_{2\beta+1}}} q^{s+(s+1)(s-k_{2\alpha+1}-l_{2\beta+1})+k_{2\alpha+1}l_{2\beta+1}} (q)_{k_{2\alpha+1}} (q)_{l_{2\beta+1}} \\
&\times \frac{q^{t+(t+1)(t-a-m_\gamma)+am_\gamma} (q)_a (q)_{m_\gamma}}{(q)_{t-a} (q)_{t-m_\gamma} (q)_{-t+a+m_\gamma}} \frac{1}{(1-q^{t+1})(1-q^{t+2})} \\
&+ \sum_{0 \leq k_{2\alpha+1} \leq k_{2\alpha} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2\beta+1} \leq l_{2\beta} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \frac{(-1)^{\sum_{i=1}^{2\alpha+1} q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2+k_i)}}}{(\prod_{i=2}^{2\alpha+1} (q)_{k_{i-1}-k_i})(q)_{k_{2\alpha+1}}} \times \frac{(-1)^{\sum_{i=1}^{2\beta+1} q^{\sum_{i=1}^{2\beta} \frac{1}{2}(l_i^2+l_i)}}}{(\prod_{i=2}^{2\beta+1} (q)_{l_{i-1}-l_i})(q)_{l_{2\beta+1}}} \\
&\times \frac{(q)^{m_\gamma^2} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2+2m_i)} (q)_n}{(\prod_{i=2}^\gamma (q)_{m_{i-1}-m_i})(q)_{m_\gamma}^2} \times \frac{q^{\frac{(k_{2\alpha+1}^2+l_{2\beta+1}^2)-(k_{2\alpha+1}+l_{2\beta+1})}{2}}}{(q)_{n+1-k_{2\alpha+1}} (q)_{n+1-l_{2\beta+1}} (q)_{-(n+1)+k_{2\alpha+1}+l_{2\beta+1}}} \times \frac{q^{(n+1)} (q)_{n+1}}{(q)_{n+1-m_\gamma}} \\
&\times \frac{1}{(1-q^{n+2})(1-q^{n+3})}
\end{aligned}$$

Here, by (5.1), (5.2), $k_{2\alpha+1}^2 - k_{2\alpha+1} \geq 0$ and $\frac{1}{2\beta+1} - l_{2\beta+1} \geq 0$,

$$\begin{aligned}
&\sum_{0 \leq k_{2\alpha+1} \leq k_{2\alpha} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2\beta+1} \leq l_{2\beta} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \frac{(-1)^{\sum_{i=1}^{2\alpha+1} q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2+k_i)}}}{(\prod_{i=2}^{2\alpha+1} (q)_{k_{i-1}-k_i})(q)_{k_{2\alpha+1}}} \times \frac{(-1)^{\sum_{i=1}^{2\beta+1} q^{\sum_{i=1}^{2\beta} \frac{1}{2}(l_i^2+l_i)}}}{(\prod_{i=2}^{2\beta+1} (q)_{l_{i-1}-l_i})(q)_{l_{2\beta+1}}} \\
&\times \frac{(q)^{m_\gamma^2} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2+2m_i)} (q)_n}{(\prod_{i=2}^\gamma (q)_{m_{i-1}-m_i})(q)_{m_\gamma}^2} \times \frac{q^{\frac{(k_{2\alpha+1}^2+l_{2\beta+1}^2)-(k_{2\alpha+1}+l_{2\beta+1})}{2}}}{(q)_{n+1-k_{2\alpha+1}} (q)_{n+1-l_{2\beta+1}} (q)_{-(n+1)+k_{2\alpha+1}+l_{2\beta+1}}} \times \frac{q^{(n+1)} (q)_{n+1}}{(q)_{n+1-m_\gamma}} \\
&\times \frac{1}{(1-q^{n+2})(1-q^{n+3})} \\
&= q^{n+1} + ((\text{high order terms})) \\
&\equiv_{n+1} 0
\end{aligned}$$

If $a = n + 1$, then terms greater than the n -th power of q vanish in the above way. Thus,

$$(5.7) \quad \begin{aligned} & \hat{J}_{(n+1,0)}^{\mathfrak{sl}_3}(P(\downarrow 2\alpha+1 \downarrow, \uparrow 2\beta+1 \uparrow, \downarrow 2\gamma \uparrow; q)) \\ & \equiv_{n+1} \sum_{0 \leq k_{2\alpha}+1 \leq k_{2\alpha} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2\beta}+1 \leq l_{2\beta} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1}+l_{2\beta+1}, n\}} \sum_{a=s}^n \sum_{t=\max\{a, m_\gamma\}}^{\min\{a+m_\gamma, n+1\}} \\ & \quad \frac{q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2+k_i)}}{\left(\prod_{i=2}^{2\alpha+1} (q)_{k_{i-1}-k_i}\right)(q)_{k_{2\alpha+1}}} \times \frac{q^{\sum_{i=1}^{2\beta} \frac{1}{2}(l_i^2+l_i)}}{\left(\prod_{i=2}^{2\beta+1} (q)_{l_{i-1}-l_i}\right)(q)_{l_{2\beta+1}}} \times \frac{(q)_{m_\gamma^2+m_\gamma} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2+2m_i)}}{\left(\prod_{i=2}^{\gamma} (q)_{m_{i-1}-m_i}\right)(q)_{m_\gamma}^2} \\ & \quad \times \frac{q^{\frac{(k_{2\alpha+1}^2+l_{2\beta+1}^2)-(k_{2\alpha+1}+l_{2\beta+1})}{2}}}{(q)_{s-k_{2\alpha+1}} (q)_{s-l_{2\beta+1}} (q)_{-s+k_{2\alpha+1}+l_{2\beta+1}}} \\ & \quad \times \frac{q^{t+(t+1)(t-a-m_\gamma)+am_\gamma} (q)_a (q)_{m_\gamma}}{(q)_{t-a} (q)_{t-m_\gamma} (q)_{-t+a+m_\gamma}} \frac{1}{(1-q^{t+1})(1-q^{t+2})} \end{aligned}$$

If $n+1 \leq a+m_\gamma$, then

$$\begin{aligned} & (\text{RHS of (5.7)}) \\ & = \sum_{0 \leq k_{2\alpha}+1 \leq k_{2\alpha} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2\beta}+1 \leq l_{2\beta} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1}+l_{2\beta+1}, n\}} \sum_{a=s}^n \sum_{t=\max\{a, m_\gamma\}}^n \\ & \quad \frac{q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2+k_i)}}{\left(\prod_{i=2}^{2\alpha+1} (q)_{k_{i-1}-k_i}\right)(q)_{k_{2\alpha+1}}} \times \frac{q^{\sum_{i=1}^{2\beta} \frac{1}{2}(l_i^2+l_i)}}{\left(\prod_{i=2}^{2\beta+1} (q)_{l_{i-1}-l_i}\right)(q)_{l_{2\beta+1}}} \times \frac{(q)_{m_\gamma^2+m_\gamma} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2+2m_i)}}{\left(\prod_{i=2}^{\gamma} (q)_{m_{i-1}-m_i}\right)(q)_{m_\gamma}^2} \\ & \quad \times \frac{q^{\frac{(k_{2\alpha+1}^2+l_{2\beta+1}^2)-(k_{2\alpha+1}+l_{2\beta+1})}{2}}}{(q)_{s-k_{2\alpha+1}} (q)_{s-l_{2\beta+1}} (q)_{-s+k_{2\alpha+1}+l_{2\beta+1}}} \\ & \quad \times \frac{q^{t+(t+1)(t-a-m_\gamma)+am_\gamma} (q)_a (q)_{m_\gamma}}{(q)_{t-a} (q)_{t-m_\gamma} (q)_{-t+a+m_\gamma}} \frac{1}{(1-q^{t+1})(1-q^{t+2})} \\ & + \sum_{0 \leq k_{2\alpha}+1 \leq k_{2\alpha} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2\beta}+1 \leq l_{2\beta} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1}+l_{2\beta+1}, n\}} \sum_{a=s}^n \\ & \quad \frac{q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2+k_i)}}{\left(\prod_{i=2}^{2\alpha+1} (q)_{k_{i-1}-k_i}\right)(q)_{k_{2\alpha+1}}} \times \frac{q^{\sum_{i=1}^{2\beta} \frac{1}{2}(l_i^2+l_i)}}{\left(\prod_{i=2}^{2\beta+1} (q)_{l_{i-1}-l_i}\right)(q)_{l_{2\beta+1}}} \times \frac{(q)_{m_\gamma^2+m_\gamma} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2+2m_i)}}{\left(\prod_{i=2}^{\gamma} (q)_{m_{i-1}-m_i}\right)(q)_{m_\gamma}^2} \\ & \quad \times \frac{q^{\frac{(k_{2\alpha+1}^2+l_{2\beta+1}^2)-(k_{2\alpha+1}+l_{2\beta+1})}{2}}}{(q)_{s-k_{2\alpha+1}} (q)_{s-l_{2\beta+1}} (q)_{-s+k_{2\alpha+1}+l_{2\beta+1}}} \\ & \quad \times \frac{q^{n+1+(n+1+1)(n+1-a-m_\gamma)+am_\gamma} (q)_a (q)_{m_\gamma}}{(q)_{t-a} (q)_{t-m_\gamma} (q)_{-t+a+m_\gamma}} \frac{1}{(1-q^{n+2})(1-q^{n+3})} \end{aligned}$$

$$\begin{aligned}
& \equiv_{n+1} \sum_{0 \leq k_{2\alpha}+1 \leq k_{2\alpha} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2\beta}+1 \leq l_{2\beta} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1}+l_{2\beta+1}, n\}} \sum_{a=s}^n \sum_{t=\max\{a, m_\gamma\}}^n \\
& \quad \times \frac{q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2+k_i)}}{\left(\prod_{i=2}^{2\alpha+1} (q)_{k_{i-1}-k_i}\right)(q)_{k_{2\alpha+1}}} \times \frac{q^{\sum_{i=1}^{2\beta} \frac{1}{2}(l_i^2+l_i)}}{\left(\prod_{i=2}^{2\beta+1} (q)_{l_{i-1}-l_i}\right)(q)_{l_{2\beta+1}}} \times \frac{(q)^{m_\gamma^2+m_\gamma} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2+2m_i)} (q)_n}{\left(\prod_{i=2}^{\gamma} (q)_{m_{i-1}-m_i}\right)(q)_{m_\gamma}^2} \\
& \quad \times \frac{q^{\frac{(k_{2\alpha+1}^2+l_{2\beta+1}^2)-(k_{2\alpha+1}+l_{2\beta+1})}{2}}}{(q)_{s-k_{2\alpha+1}} (q)_{s-l_{2\beta+1}} (q)_{-s+k_{2\alpha+1}+l_{2\beta+1}}} q^{s+(s+1)(s-k_{2\alpha+1}-l_{2\beta+1})+k_{2\alpha+1}l_{2\beta+1}} (q)_{k_{2\alpha+1}} (q)_{l_{2\beta+1}} \\
& \quad \times \frac{q^{t+(t+1)(t-a-m_\gamma)+am_\gamma} (q)_a (q)_{m_\gamma}}{(q)_{t-a} (q)_{t-m_\gamma} (q)_{-t+a+m_\gamma}} \frac{1}{(1-q^{t+1})(1-q^{t+2})}
\end{aligned}$$

The second congruence is obtained by the fact that

$$\begin{aligned}
& n + 1 + (n + 1 + 1)(n + 1 - a - m_\gamma) + am_\gamma + m_\gamma \\
& = (n + 1 - a + 1)(n + 1 - m_\gamma + 1) - 1 + m_\gamma \\
& \geq 2(n + 1 - m_\gamma + 1) - 1 + m_\gamma \\
& = n + 1 + n - m_\gamma + 4 \geq n + 5
\end{aligned}$$

Therefore,

$$\begin{aligned}
& (5.8) \quad \hat{J}_{(n+1,0)}^{\text{sl}_3}(P(\downarrow 2\alpha+1 \downarrow, \uparrow 2\beta+1 \uparrow, \downarrow 2\gamma \uparrow); q)) \\
& \equiv_{n+1} \sum_{0 \leq k_{2\alpha}+1 \leq k_{2\alpha} \leq \dots \leq k_1 \leq n} \sum_{0 \leq l_{2\beta}+1 \leq l_{2\beta} \leq \dots \leq l_1 \leq n} \sum_{0 \leq m_\gamma \leq m_{\gamma-1} \leq \dots \leq m_1 \leq n} \sum_{s=\max\{k_{2\alpha+1}, l_{2\beta+1}\}}^{\min\{k_{2\alpha+1}+l_{2\beta+1}, n\}} \sum_{a=s}^n \sum_{t=\max\{a, m_\gamma\}}^n \\
& \quad \times \frac{q^{\sum_{i=1}^{2\alpha} \frac{1}{2}(k_i^2+k_i)}}{\left(\prod_{i=2}^{2\alpha+1} (q)_{k_{i-1}-k_i}\right)(q)_{k_{2\alpha+1}}} \times \frac{q^{\sum_{i=1}^{2\beta} \frac{1}{2}(l_i^2+l_i)} (q)_n}{\left(\prod_{i=2}^{2\beta+1} (q)_{l_{i-1}-l_i}\right)(q)_{l_{2\beta+1}}} \times \frac{(q)^{m_\gamma^2+m_\gamma} (q)^{\sum_{i=1}^{\gamma-1} (m_i^2+2m_i)} (q)_n}{\left(\prod_{i=2}^{\gamma} (q)_{m_{i-1}-m_i}\right)(q)_{m_\gamma}^2} \\
& \quad \times \frac{q^{\frac{(k_{2\alpha+1}^2+l_{2\beta+1}^2)-(k_{2\alpha+1}+l_{2\beta+1})}{2}}}{(q)_{s-k_{2\alpha+1}} (q)_{s-l_{2\beta+1}} (q)_{-s+k_{2\alpha+1}+l_{2\beta+1}}} q^{s+(s+1)(s-k_{2\alpha+1}-l_{2\beta+1})+k_{2\alpha+1}l_{2\beta+1}} (q)_{k_{2\alpha+1}} (q)_{l_{2\beta+1}} \\
& \quad \times \frac{q^{t+(t+1)(t-a-m_\gamma)+am_\gamma} (q)_a (q)_{m_\gamma}}{(q)_{t-a} (q)_{t-m_\gamma} (q)_{-t+a+m_\gamma}} \frac{1}{(1-q^{t+1})(1-q^{t+2})}
\end{aligned}$$

According to (5.4) and (5.8),

$$\hat{J}_{(n,0)}^{\text{sl}_3}(P(\downarrow 2\alpha+1 \downarrow, \uparrow 2\beta+1 \uparrow, \downarrow 2\gamma \uparrow); q)) \equiv_{n+1} \hat{J}_{(n+1,0)}^{\text{sl}_3}(P(\downarrow 2\alpha+1 \downarrow, \uparrow 2\beta+1 \uparrow, \downarrow 2\gamma \uparrow); q))$$

□

6. APPENDIX

The knots $8_{10}, 8_{15}, 8_{20}$ and 8_{21} are four or five parameters pretzel knots. For these knots, we can calculate the one-row colored sl_3 Jones polynomials by combining the similar method as three-parameter pretzel knots and links.

Theorem 6.1.

The one-row colored \mathfrak{sl}_3 Jones polynomials for $P(-3, -2, 3, -1) = 8_{10}$, $P(3, -1, -2, -1, 3) = 8_{15}$, $P(3, -2, -3, 1) = 8_{20}$ and $P(3, 3, -1, 2) = 8_{21}$ are the following:

- (1) $J_{(n,0)}^{\mathfrak{sl}_3}(P(-3, -2, 3, -1); q)$

$$= \sum_{0 \leq k_3 \leq k_2 \leq k_1 \leq n} \sum_{0 \leq l_2 \leq l_1 \leq n} \sum_{0 \leq p_3 \leq p_2 \leq p_1 \leq n} \sum_{m=0}^n \sum_{t=\max\{k_3, l_2\}}^{\min\{k_3+l_2, n\}} \sum_{a=t}^n \sum_{s=\max\{p_3, m\}}^{\min\{p_3+m, n\}} \sum_{b=s}^n \sum_{u=\max\{a, b\}}^{\min\{a+b, n\}} (q^{\frac{n^2+3n}{3}})^3$$

$$\chi_{-1}(n, k_0, k_1, k_2) \chi_{-1}(n, l_0, l_1) q^{-1} \chi_1(n, p_1, p_2, p_3) \chi_{-1}(n, m) \Omega(n, t, k_3, l_2) \Omega(n, s, p_3, m) \psi(n, u, a, b)$$

$$q^{-(n-u)} \frac{(1-q^{n+1})(1-q^{n+2})}{(1-q^{u+1})(1-q^{u+2})},$$
- (2) $J_{(n,0)}^{\mathfrak{sl}_3}(P(3, -1, -2, -1, 3); q)$

$$= \sum_{0 \leq k_3 \leq k_2 \leq k_1 \leq n} \sum_{0 \leq l \leq n} \sum_{0 \leq m_2 \leq m_1} \sum_{0 \leq p \leq n} \sum_{0 \leq r_3 \leq r_2 \leq r_1 \leq n} \sum_{t=\max\{k_3, l\}}^{\min\{k_3+l, n\}} \sum_{a=t}^n \sum_{s=\max\{p, r_3\}}^{\min\{p+r_3, n\}} \sum_{b=s}^n$$

$$\sum_{u=\max\{a, m_1\}}^{\min\{a+m_2, n\}} \sum_{v=\max\{b, m_2\}}^{\min\{b+m_2, n\}} (q^{\frac{n^2+3n}{3}})^{-2} \chi_1(n, k_1, k_2, k_3) \chi_{-1}(n, l) \phi(n, m_1, m_2) \chi_{-1}(n, p) \chi_1(n, r_1, r_2, r_3) \Omega(n, t, k_3, l)$$

$$\Omega(n, s, p, r_3) \psi(n, u, a, m_2) \psi(n, v, u, b) q^{-(n-v)} \frac{(1-q^{n+1})(1-q^{n+2})}{(1-q^{v+1})(1-q^{v+2})}$$
- (3) $J_{(n,0)}^{\mathfrak{sl}_3}(P(3, -2, -3, 1); q),$

$$= \sum_{0 \leq k_3 \leq k_2 \leq k_1 \leq n} \sum_{0 \leq l_2 \leq l_1 \leq n} \sum_{0 \leq p_3 \leq p_2 \leq p_1 \leq n} \sum_{m=0}^n \sum_{t=\max\{k_3, l_2\}}^{\min\{k_3+l_2, n\}} \sum_{a=t}^n \sum_{s=\max\{p_3, m\}}^{\min\{p_3+m, n\}} \sum_{b=s}^n \sum_{u=\max\{a, b\}}^{\min\{a+b, n\}} q^{\frac{n^2+3n}{3}}$$

$$\chi_1(n, k_1, k_2, k_3) \chi_{-1}(n, l_1, l_2) q^{-1} \chi_{-1}(n, p_1, p_2, p_3) \chi_1(n, m) \Omega(n, t, n-k_3, l_2) \Omega(n, s, n-p_2, n-m)$$

$$\psi(n, u, t+a, s+b) q^{-(n-u)} \frac{(1-q^{n+1})(1-q^{n+2})}{(1-q^{u+1})(1-q^{u+2})}$$
- (4) $J_{(n,0)}^{\mathfrak{sl}_3}(P(3, 3, -1, -2); q),$

$$= \sum_{0 \leq k_3 \leq k_2 \leq k_1 \leq n} \sum_{0 \leq l_3 \leq l_2 \leq l_1 \leq n} \sum_{0 \leq p \leq n} \sum_{0 \leq m_2 \leq m_1 \leq n} \sum_{t=\max\{k_3, l_3\}}^{\min\{k_3+l_3, n\}} \sum_{a=t}^n \sum_{s=\max\{p, m\}}^{\min\{p+m, n\}} \sum_{b=s}^n \sum_{u=\max\{a, b\}}^{\min\{t+s, n\}}$$

$$(q^{\frac{n^2+3n}{3}})^{-3} \chi_1(n, k_1, k_2, k_3) \chi_1(n, l_1, l_2, l_3) \chi_{-1}(n, p) \chi_{-1}(n, m_1, m_2)$$

$$\Omega(n, t, k_3, l_3) \Omega(n, s, p, m_2) \psi(n, u, a, b) q^{-(n-u)} \frac{(1-q^{n+1})(1-q^{n+2})}{(1-q^{u+1})(1-q^{u+2})}.$$

Remark 6.2. According to Theorem 6.1, Theorem 4.1 and known results, the one-row \mathfrak{sl}_3 colored Jones polynomial is determined for all knots with eight or fewer crossings except $8_{16}, 8_{17}, 8_{18}$.

Acknowledgment. I would like to express my sincere gratitude to my advisor, Professor Masaaki Suzuki, for his constructive suggestions.

REFERENCES

- [Arm13] C. Armond, The head and tail conjecture for alternating knots, Algebr. Geom. Topol. 13 (2013), no. 5, 2809–2826. MR 3116304
- [AD11] C. Armond and O. T. Dasbach, Rogers-Ramanujan type identities and the head and tail of the colored Jones polynomial, arXiv:1106.3948 (2011).

- [Bei19] P. Beirne, On the 2-head of the colored Jones polynomial for pretzel knots, arXiv:1902.07061(2019)
- [BO17] P. Beirne and R. Osburn, q-series and tails of colored Jones polynomials, *Indagationes Mathematicae* 28, no. 1, (2017), 247–260
- [DL06] O. T. Dasbach and X. S. Lin, On the head and the tail of the colored Jones polynomial, *Compos. Math.* 142 (2006), no. 5, 1332–1342. MR 2264669
- [EH17] Mohamed Elhamdadi and Mustafa Hajij, Pretzel knots and q-series, *Osaka J. Math.* 54 (2017), no. 2, 363–381. MR 3657236
- [Gar11] S. Garoufalidis, The Jones slopes of a knot, *Quantum Topol.* 2 (2011), 43–69. MR 2763086
- [GL15] S. Garoufalidis and T. T. Q. Lê, Nahm sums, stability, and the colored Jones polynomial, *Res. Math. Sci.* 2 (2015), Art. 1, 55. MR 337565
- [GMV13] S. Garoufalidis, H. Morton, and T. Vuong, The SL_3 colored Jones polynomial of the trefoil, *Proc. Amer. Math. Soc.* 141 (2013), no. 6, 2209–2220. MR 3034446
- [GV17] S. Garoufalidis and T. Vuong, A stability conjecture for the colored Jones polynomial, *Topology Proc.* 49 (2017), 211–249. MR 3570390
- [Haj14] M. Hajij, The Bubble skein element and applications, *J. Knot Theory Ramifications* 23 (2014), no. 14, 1450076, 30. MR 3312619
- [Jon85] V. F. R. Jones, A polynomial invariant for knots via von Neumann algebras, *Bull. Amer. Math. Soc. (N.S.)* 12 (1985), no. 1, 103–111. MR 766964
- [Kas97] R. M. Kashaev, The hyperbolic volume of knots from the quantum dilogarithm, *Lett. Math. Phys.* 39 (1997), no. 3, 269–275. MR 1434238
- [Kau87] L. H. Kauffman, State models, and the Jones polynomial, *Topology* 26 (1987), no. 3, 395–407. MR 899057
- [KL94] L. H. Kauffman and Sóstenes L. Lins, Temperley-Lieb recoupling theory and invariants of 3-manifolds, *Annals of Mathematics Studies*, vol. 134, Princeton University Press, Princeton, NJ, 1994. MR 1280463
- [Kim06] D. Kim, Trihedron coefficients for $U_q(sl(3, \mathbb{C}))$, *J. Knot Theory Ramifications* 15 (2006), no. 4, 453–469. MR 2221529
- [Kim07] ———, Jones-Wenzl idempotents for rank two simple Lie algebras, *Osaka J. Math.* 44 (2007), no. 3, 691–722. MR 23609
- [KO17] A. Keilthy and R. Osburn, Rogers-Ramanujan type identities for alternating knots, *J. Number Theory* 161 (2016), 255–280. MR 3435728
- [Kup96] G. Kuperberg, Spiders for rank 2 Lie algebras, *Comm. Math. Phys.* 180 (1996), no. 1, 109–151. MR 1403861
- [Law03] R. Lawrence, The $\text{PSU}(3)$ invariant of the Poincaré homology sphere, *Proceedings of the Pacific Institute for the Mathematical Sciences Workshop “Invariants of Three-Manifolds”* (Calgary, AB, 1999), vol. 127, 2003, pp. 153–168. MR 1953324
- [Le00] T. T. Q. Le, Integrality and symmetry of quantum link invariants, *Duke Math. J.* 102 (2000), no. 2, 273–306. MR 1749439
- [Lic93] W. B. R. Lickorish, The skein method for three-manifold invariants, *J. Knot Theory Ramifications* 2 (1993), no. 2, 171–194. MR 1227009
- [Lic97] ———, An introduction to knot theory, *Graduate Texts in Mathematics*, vol. 175, Springer-Verlag, New York, 1997. MR 1472978
- [OY97] T. Ohtsuki and S. Yamada, Quantum $SU(3)$ invariant of 3-manifolds via linear skein theory, *J. Knot Theory Ramifications* 6 (1997), no. 3, 373–404. MR 1457194
- [RT91] N. Reshetikhin and V. G. Turaev, Invariants of 3-manifolds via link polynomials and quantum groups, *Invent. Math.* 103 (1991), no. 3, 547–597. MR 1091619
- [Wen87] H. Wenzl, On sequences of projections, *C. R. Math. Rep. Acad. Sci. Canada* 9 (1987), no. 1, 5–9. MR 873400
- [Yua17] W. Yuasa, The \mathfrak{sl}_3 colored Jones polynomials for 2-bridge links, *J. Knot Theory Ramifications* 26(2017), no. 7, 1750038, 37. MR 3660093
- [Yua18a] ———, A q -series identity via the \mathfrak{sl}_3 colored Jones polynomials for the $(2, 2m)$ -torus link, *Proc. Amer. Math. Soc.* 146 (2018), no. 7, 3153–3166. MR 3787374
- [Yua18b] ———, A_2 colored polynomials of rigid vertex graphs, *New York J. Math.* (2018), vol. 24, 355–374. MR 3829741
- [Yua21a] ———, Twist formulas for one-row colored A_2 webs and \mathfrak{sl}_3 tails of $(2, 2m)$ -torus links, *Acta Math. Vietnam.* 46 (2021), no. 2, 369–387. MR 4264242
- [Yua21b] ———, The zero stability for the one-row colored \mathfrak{sl}_3 Jones polynomial, arXiv:2007.15621(2021)