

Parameterized Algorithms for Upward Planarity

Steven Chaplick ✉ 

Maastricht University, Maastricht, The Netherlands

Emilio Di Giacomo ✉ 

Università degli Studi di Perugia, Perugia, Italy

Fabrizio Frati ✉ 

Roma Tre University, Rome, Italy

Robert Ganian ✉ 

Technische Universität Wien, Wien, Austria

Chrysanthi N. Raftopoulou ✉ 

National Technical University of Athens, Athens, Greece

Kirill Simonov ✉

Technische Universität Wien, Wien, Austria

Abstract

We obtain new parameterized algorithms for the classical problem of determining whether a directed acyclic graph admits an upward planar drawing. Our results include a new fixed-parameter algorithm parameterized by the number of sources, an XP-algorithm parameterized by treewidth, and a fixed-parameter algorithm parameterized by treedepth. All three algorithms are obtained using a novel framework for the problem that combines SPQR tree-decompositions with parameterized techniques. Our approach unifies and pushes beyond previous tractability results for the problem on series-parallel digraphs, single-source digraphs and outerplanar digraphs.

2012 ACM Subject Classification Theory of computation → Parameterized complexity and exact algorithms; Human-centered computing → Graph drawings

Keywords and phrases Upward planarity, parameterized algorithms, SPQR trees, treewidth, treedepth

Digital Object Identifier 10.4230/LIPIcs...

Related Version An extended abstract of this manuscript was accepted for publication in the proceedings of the 38th International Symposium on Computational Geometry (SoCG 2022).

Funding *Emilio Di Giacomo*: MIUR, grant 20174LF3T8, Dip. Ing. - UNIPG, grants RICBA19FM and RICBA20EDG

Fabrizio Frati: MIUR, grant 20174LF3T8

Robert Ganian: Austrian Science Fund (FWF) Project Y1329

Chrysanthi N. Raftopoulou: NTUA research program IIEBE 2020

Kirill Simonov: Austrian Science Fund (FWF) Project P31336

Acknowledgements The authors thank Fabrizio Montecchiani and Giuseppe Liotta for fruitful discussions on the topic of upward planarity. This research was initiated at Dagstuhl Seminar 21293: Parameterized Complexity in Graph Drawing [22].

1 Introduction

A digraph is called *upward planar* if it admits an upward planar drawing, that is, a planar drawing where all edges are oriented upward. The problem of upward planarity testing (UPWARD PLANARITY) and constructing an associated upward planar drawing arises, among others, in the context of visualization of hierarchical network structures; application domains include project management, visual languages and software engineering [2]. Upward planarity



© S. Chaplick, E. Di Giacomo, F. Frati, R. Ganian, C. N. Raftopoulou, K. Simonov;
licensed under Creative Commons License CC-BY 4.0

Leibniz International Proceedings in Informatics

LIPICS Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

is the most prominent notion of planarity that is inherently directed, and also has classical connections to the theory of ordered sets: the orders arising from the transitive closure of upward planar single-source digraphs have bounded dimension [38].

Since the introduction of the notion, UPWARD PLANARITY has become the focus of extensive theoretical research. The problem has been shown to be NP-complete more than 25 years ago [25, 26], but the first polynomial-time algorithms for restricted variants of UPWARD PLANARITY have been published even earlier [31, 32]. Among others, the problem is known to be polynomial-time tractable when G is provided with a planar embedding [3] (which also implies polynomial-time tractability for triconnected DAGs, since these admit a single planar embedding), or when restricted to the class of outerplanar DAGs [35], DAGs whose underlying graph is series-parallel [15], and most prominently single-source DAGs [2, 7, 32].

In spite of the number of results on UPWARD PLANARITY that analyze the classical complexity of the problem on specific subclasses of instances, the problem was up to now mostly unexplored from the more fine-grained perspective of parameterized complexity analysis¹ [12, 18]. In particular, while it was known that UPWARD PLANARITY is fixed-parameter tractable when parameterized by the cyclomatic number of the input DAG (or, equivalently, the feedback edge number of the underlying undirected graph) [10], by the number of triconnected components and cut vertices [28], or the number of triconnected components plus the maximum diameter of a split component [15], the complexity of the problem under classical structural parameterizations has remained completely open.

Contribution. We develop a novel algorithmic framework for solving UPWARD PLANARITY which combines parameterized dynamic programming with the SPQR-tree decompositions of planar graphs [14, 27, 29]. In essence, our framework uses a characterization of the “shapes” of faces in an upward planar drawing that is inspired by earlier work on the notion of spirality [3, 15] and reduces UPWARD PLANARITY to the task of handling the “rigid” nodes in these decompositions. Informally, the task that needs to be handled there can be stated as follows: what are all the possible ways to combine the possible shapes of the children of a rigid node to obtain an upward planar drawing for the node itself? The framework is formalized in the form of a general “Interface Lemma” (Lemma 24) which can be complemented with numerous parameterizations as well as other algorithmic approaches.

In the remainder of this article, we use this framework to push the boundaries of tractability for UPWARD PLANARITY. Our first result in this direction is a fixed-parameter algorithm for UPWARD PLANARITY parameterized by the number of sources in the input graph². This result generalizes the polynomial-time tractability of the single-source case [2, 7] and answers an open question from a recent Dagstuhl seminar [22]. On a high level, we use the Interface Lemma to reduce the problem to a case where almost all children of a rigid node have a simple shape, and we show how this can be handled via a flow network approach.

Having established the tractability of instances with few sources, we turn towards understanding which structural properties of the underlying undirected graph can be used to solve UPWARD PLANARITY efficiently. In this context, there is a well-known hierarchy of fundamental structural parameters of graphs (see, e.g., [1, Figure 1]) and the study of their algorithmic applications is a prominent research direction that has been pursued not only for a number of problems in computational geometry [9, 30], but also in many other fields such as in artificial intelligence [4, 23] and constraint satisfaction [37, 24]. Yet,

¹ A basic introduction to parameterized complexity is provided in the preliminaries.

² It is worth noting that this immediately also implies tractability parameterized by the number of sinks: inverting all arcs results in an equivalent instance.

apart from the fixed-parameter tractability of UPWARD PLANARITY parameterized by the feedback edge number [10] (which is one of the by far most restrictive parameterizations in the aforementioned hierarchy), nothing was known about whether the more widespread “decompositional” parameters can be used to solve the problem. The parameters that will be of interest here are *treewidth* [36], the most prominent structural graph parameter, and *treedepth* [34], the arguably best known parameter that lies below treewidth in the parameter hierarchy.

To obtain new boundaries of tractability for UPWARD PLANARITY with respect to these two parameters, we first show that the problem posed by the Interface Lemma can be restated as a purely combinatorial problem on a suitable combinatorization of the embedding of the graph represented by the rigid node, and—crucially—that a bound on the input graph’s treewidth also implies a bound for the treewidth of this combinatorization. Once that is done, we design a non-trivial dynamic program that exploits this treewidth bound to handle the rigid nodes, which together with the Interface Lemma allows us to solve UPWARD PLANARITY. This yields an XP-algorithm for UPWARD PLANARITY parameterized by the treewidth of the underlying undirected graph—a result which unifies and generalizes the polynomial-time tractability of UPWARD PLANARITY on outerplanar as well as series-parallel graphs [15, 35]. Furthermore, a more detailed analysis of the dynamic program reveals that the same algorithm runs in fixed-parameter time when parameterized by treedepth.

Paper Organization. After introducing the necessary preliminaries in Section 2, we introduce a characterization of components of the SPQR trees used by our framework (Section 3), provide the bulk of our framework for digraphs whose underlying undirected graph is 2-connected (Section 4), and then complete the description of the framework by handling general digraphs (Section 5). Sections 6 and 7 are then dedicated to applying the framework to UPWARD PLANARITY parameterized by the number of sources and treewidth along with treedepth, respectively.

2 Preliminaries

We refer to the classical books for basic graph and graph drawing terminology [16, 13]. We use $N_G(v)$ to denote the set of vertices adjacent to a vertex v in a graph G .

2.1 Upward planar drawings and embeddings

A *drawing* of a graph maps each vertex to a point in the plane and each edge to a Jordan arc between the end-points of the edge. A drawing is *planar* if no two edges intersect, except at common end-points. A planar drawing partitions the plane into regions, called *faces*. The bounded faces are called *internal*, while the unbounded face is the *outer face*. Two planar drawings of a graph are *equivalent* if: (1) they have the same *rotation system*, that is, for each vertex v , the clockwise order of the edges incident to v is the same in both drawings; and (2) their outer faces are delimited by the same walk, that is, the order of the edges encountered when clockwise traversing the boundary of the outer face is the same in both drawings. A *planar embedding* of a graph is an equivalence class of planar drawings of that graph.

Thus, a planar embedding of a graph consists of a rotation system and a choice for the walk delimiting the outer face. We often talk about a *face of a planar embedding*, meaning a face of any planar drawing that respects the planar embedding. The *flip* of a planar embedding is the planar embedding obtained by reversing the clockwise order of the edges

incident to each vertex and by reversing the order of the edges encountered when clockwise traversing the boundary of the outer face.

Throughout the paper, we use the term *digraph* as short for “directed graph”. A digraph is *acyclic* if it contains no directed cycle. An acyclic digraph is usually called *DAG*, for short. A vertex in a digraph is a *source* if it is only incident to outgoing edges and it is a *sink* if it is only incident to incoming edges. A vertex in a digraph is a *switch* if it is a source or a sink, and it is a *non-switch* otherwise. The *underlying graph* of a digraph is the undirected graph obtained from the digraph by ignoring the edge directions. A *plane digraph* is a digraph together with a prescribed planar embedding for its underlying graph.

A drawing of a digraph is *upward* if every edge is represented by a Jordan arc monotonically increasing from the source to the sink of the edge, and it is *upward planar* if it is both upward and planar. A digraph is *upward planar* if it admits an upward planar drawing; we use UPWARD PLANARITY to denote the problem of determining whether a digraph is upward planar; w.l.o.g., we assume that the input digraph is connected.

Consider an upward planar drawing Γ of a digraph G . An *angle* α of a face f of Γ is a triple (e_1, v, e_2) , where e_1 and e_2 are two edges of G that are incident to the vertex v , that are incident to the face f , and that are consecutive in the order of the edges encountered when clockwise traversing the boundary of f . We say that α is *flat* if one between e_1 and e_2 is incoming v and the other one is outgoing v , otherwise α is a *switch angle*. Then Γ defines an *angle assignment*, which assigns the value -1 , 0 , and 1 to each small, flat, and large angle, respectively, in every face of Γ . The angle assignment, together with the planar embedding of the underlying graph of G in Γ , constitutes an *upward planar embedding* of G .

A switch angle at a vertex v is hence delimited by two outgoing or by two incoming edges for v . Each switch angle is further classified as *large* or *small* as follows. Consider a switch angle $\alpha = (e_1, v, e_2)$ at a vertex v in a face f delimited by two outgoing edges (resp. by two incoming edges) and consider a disk D centered at v , sufficiently small so that its boundary has a single intersection with every edge incident to v . The edges e_1 and e_2 divide D into two regions, one of which contains part of f and contains no portion of any edge incident to v in its interior; call D' this region. Then we say that α is *large* if D' contains a suitably short vertical segment that has v as its highest (resp. lowest) end-point, it is *small* otherwise.

An upward planar drawing hence defines an *angle assignment*, which is an assignment of the value -1 , 0 , and 1 to each small, flat, and large angle, respectively, in every face of Γ . This angle assignment, together with the planar embedding of the underlying graph of G in Γ , constitutes an *upward planar embedding* of G . An *upward plane digraph* is a digraph together with a prescribed upward planar embedding.

The angle assignments that enhance a planar embedding into an upward planar embedding have been characterized by Didimo et al. [15], building on the work by Bertolazzi et al. [3]. Note that, once the planar embedding \mathcal{E} of a digraph G is specified, then so are the angles of the faces of \mathcal{E} ; in particular, whether an angle is flat or switch only depends on \mathcal{E} . Consider an angle assignment for \mathcal{E} . If v is a vertex of G , we denote by $n_i(v)$ the number of angles at v that are labeled i , with $i \in \{-1, 0, 1\}$. If f is a face of G , we denote by $n_i(f)$ the number of angles of f that are labeled i , with $i \in \{-1, 0, 1\}$. The cited characterization is as follows.

► **Theorem 1** ([3, 15]). *Let G be a digraph, \mathcal{E} be a planar embedding of the underlying graph of G , and λ be an assignment of each angle of each face in \mathcal{E} to a value in $\{-1, 0, 1\}$. Then \mathcal{E} and λ define an upward planar embedding of G if and only if the following properties hold:*

UP0 *If α is a switch angle, then $\lambda(\alpha) \in \{-1, 1\}$, and if α is a flat angle, then $\lambda(\alpha) = 0$.*

UP1 *If v is a switch vertex of G , then $n_1(v) = 1$, $n_{-1}(v) = \deg(v) - 1$, $n_0(v) = 0$.*

UP2 If v is a non-switch vertex of G , then $n_1(v) = 0$, $n_{-1}(v) = \deg(v) - 2$, $n_0(v) = 2$.

UP3 If f is a face of G , then $n_1(f) = n_{-1}(f) - 2$ if f is an internal face and $n_1(f) = n_{-1}(f) + 2$ if f is the outer face.

Theorem 1 has the following algorithmic consequence.

► **Theorem 2** ([3, 6]). *Let G be an n -vertex digraph and \mathcal{E} be a planar embedding of the underlying graph of G . It is possible to test in $\mathcal{O}(n \log^3 n)$ time whether there exists an angle assignment λ such that \mathcal{E} and λ define an upward planar embedding of G . Consequently, it can be tested in $\mathcal{O}(n \log^3 n)$ time whether G admits an upward planar drawing Γ which respects \mathcal{E} ; in the positive case, Γ can be constructed within the same time bound.*

Theorem 2 was originally proved in [3] with an $\mathcal{O}(n^2)$ running time. The main idea for its proof is the following. Construct a *planar bipartite network* $\mathcal{N}(S, T, A)$, where:

- S is a set of sources; there is a source s_w for each vertex w of G ; each source can supply a single unit of flow;
- T is a set of sinks; there is a sink t_f for each face f of \mathcal{E} ; each sink t_f demands a number of units of flow equal to $n_f/2 - 1$ or $n_f/2 + 1$, depending on whether f is an internal face or the outer face of \mathcal{E} , where n_f is the number of switch angles incident to f ; and
- A is a set of arcs from the sources to the sinks; there is an arc from a source s_w to a sink t_f if the vertex w corresponding to s_w is incident to the face f corresponding to t_f ; each arc a has a capacity c_a of a single unit of flow.

It was proved in [3] that there exists an angle assignment λ such that \mathcal{E} and λ define an upward planar embedding of G if and only if the bipartite network \mathcal{N} admits a flow whose value is the sum of the demands of the sinks in T (or the sum of the supplies of the sources in S). A *flow* is an assignment of a value $\phi_a \leq c_a$ to each arc a such that the sum of the values assigned to the arcs outgoing each node s_w does not exceed the supply of s_w and the sum of the values assigned to the arcs incoming each node t_f does not exceed the demand of t_f . The *value* of a flow is the sum of the values assigned to its arcs.

Borradaile et al. [6] presented an $\mathcal{O}(n \log^3 n)$ algorithm to find a maximum flow in a planar network with multiple sources and sinks. This result, plugged into the described framework by Bertolazzi et al. [3], gives us a proof of Theorem 2.

2.2 Parameterized Complexity

In parameterized complexity [12, 18], the complexity of a problem is studied not only with respect to the input size, but also with respect to some problem parameter(s). The core idea behind parameterized complexity is that the combinatorial explosion resulting from the NP-hardness of a problem can sometimes be confined to certain structural parameters that are small in practical settings. We now proceed to the formal definitions.

A *parameterized problem* Q is a subset of $\Omega^* \times \mathbb{N}$, where Ω is a fixed alphabet. Each instance of Q is a pair (I, κ) , where $\kappa \in \mathbb{N}$ is called the *parameter*. A parameterized problem Q is *fixed-parameter tractable* (FPT) [18, 12], if there is an algorithm, called a *fixed-parameter algorithm*, that decides whether an input (I, κ) is a member of Q in time $f(\kappa) \cdot |I|^{\mathcal{O}(1)}$, where f is a computable function. The class FPT denotes the class of all fixed-parameter tractable parameterized problems. A weaker notion of tractability is that of XP: a parameterized problem Q is in the class XP if there is an algorithm that decides whether an input (I, κ) is a member of Q in time $(|I| + 1)^{f(\kappa)}$, where f is a computable function.

2.3 Treewidth and Treedepth

Here we consider the treewidth and treedepth of the underlying graphs³. A *tree-decomposition* \mathcal{T} of a graph $G = (V, E)$ is a pair (T, χ) , where T is a tree (whose vertices we call *nodes*) rooted at a node r and χ is a function that assigns each node t a set $\chi(t) \subseteq V$ such that the following holds:

- For every $uv \in E$ there is a node t such that $u, v \in \chi(t)$.
- For every vertex $v \in V$, the set of nodes t satisfying $v \in \chi(t)$ forms a nonempty subtree of T .

A tree-decomposition is *nice* if the following two conditions are also satisfied:

- $|\chi(\ell)| = 1$ for every leaf ℓ of T and $|\chi(r)| = 0$.
- There are only three kinds of non-leaf nodes in T :

Introduce node: a node t with exactly one child t' such that $\chi(t) = \chi(t') \cup \{v\}$ for some vertex $v \notin \chi(t')$.

Forget node: a node t with exactly one child t' such that $\chi(t) = \chi(t') \setminus \{v\}$ for some vertex $v \in \chi(t')$.

Join node: a node t with two children t_1, t_2 such that $\chi(t) = \chi(t_1) = \chi(t_2)$.

The *width* of a tree-decomposition (T, χ) is the size of a largest set $\chi(t)$ minus 1, and the *treewidth* of the graph G , denoted $tw(G)$, is the minimum width of a tree-decomposition of G . It is known that a tree-decomposition can be transformed into a nice tree-decomposition of the same width in linear time. Efficient fixed-parameter algorithms are known for computing a nice tree-decomposition of near-optimal width [5, 33].

► **Proposition 3 ([5]).** *There exists an algorithm which, given an n -vertex graph G and an integer k , in time $2^{\mathcal{O}(k)} \cdot n$ either outputs a tree-decomposition of G of width at most $5k + 4$ and $\mathcal{O}(n)$ nodes, or determines that $tw(G) > k$.*

We let T_t denote the subtree of T rooted at a node t , and we use $\chi(T_t)$ to denote the set $\bigcup_{t' \in V(T_t)} \chi(t')$. In the context of dynamic programming, the set $past(t) = \chi(T_t) \setminus \chi(t)$ is called the *past* while the set $past(t) = V \setminus \chi(T_t)$ is called the *future*.

The second structural parameter that we will be considering here is the *treedepth* of a graph G , denoted $td(G)$ [34]. A useful way of thinking about graphs of bounded treedepth is that they are (sparse) graphs with no long paths.

A *rooted forest* is a disjoint union of rooted trees. For a vertex x in a tree T of a rooted forest, the *height* (or *depth*) of x in the forest is the number of vertices in the path from the root of T to x . The *height of a rooted forest* is the maximum height of a vertex of the forest.

► **Definition 4 (Treedepth).** *Let the closure of a rooted forest \mathcal{F} be the graph $clos(\mathcal{F}) = (V_c, E_c)$ with the vertex set $V_c = \bigcup_{T \in \mathcal{F}} V(T)$ and the edge set $E_c = \{xy : x \text{ is an ancestor of } y \text{ in some } T \in \mathcal{F}\}$. A treedepth decomposition of a graph G is a rooted forest \mathcal{F} such that $G \subseteq clos(\mathcal{F})$. The treedepth $td(G)$ of a graph G is the minimum height of any treedepth decomposition of G .*

We will later use T_x to denote the vertex set of the subtree of T rooted at a vertex x of T . The following properties of treedepth will be crucial for our considerations.

► **Proposition 5 ([34]).** *For every path of length d in a graph G , it holds that $td(G) \leq d \leq 2^{td(G)}$. Moreover, $tw(G) \leq td(G)$.*

³ Directed alternatives to treewidth exist, but are typically not well-suited for algorithmic applications [21].

2.4 Expansion

In our algorithms, we will employ a linear-time preprocessing step to simplify the input digraphs so that every vertex has at most one incoming edge or at most one outgoing edge; this operation, which was introduced by Bertolazzi et al. [2], is called *expansion* and is defined as follows. We begin by marking each non-switch vertex in the digraph $G = (V, E)$ as *unprocessed*. We then loop over each non-switch unprocessed vertex v , for which we create two new *processed* vertices v_1 and v_2 and add the following edges to the digraph: $\{v_1v_2\} \cup \{(av_1 \mid av \in E) \cup \{v_2b \mid vb \in E\}$, and then we delete v . In an expanded graph, that is, a graph that has gone through the expansion operation, every vertex has at most one outgoing or at most one incoming edge; in the former case we say it is a *bottom* vertex, while in the latter case it is a *top* vertex. The only outgoing (incoming) edge of a bottom (top) vertex of an expanded digraph is the *special edge* of that vertex. Since a non-switch vertex of degree 2 has exactly one incoming and one outgoing edge, it could be considered both a top and a bottom vertex. To avoid this ambiguity, we treat degree-2 vertices as top vertices.

A digraph expansion preserves upward planarity [2]; furthermore, it preserves biconnectivity and the number of sources, and at most doubles the number of vertices. Moreover, below we show that applying expansion on graphs of bounded treedepth results in graphs which also have bounded treedepth. Hence, in the remainder, we will assume, without loss of generality, that each digraph for which we aim to test upward planarity has been expanded and thus each of its vertices has at most one incoming edge or at most one outgoing edge.

► **Observation 6.** *Let G be a digraph and G' be its expansion. The treedepth and treewidth of G' are at most twice the treedepth and treewidth of G , respectively. Moreover, G and G' have the same number of sources.*

Proof. Consider a treedepth decomposition \mathcal{F} of G , replace every vertex of this decomposition with an edge to obtain a rooted forest \mathcal{F}' . We claim that \mathcal{F}' is a treedepth decomposition of G' : For a vertex v of G corresponding to a vertex u in \mathcal{F} , associate the respective vertices v_1, v_2 of G' with the endpoints of the edge in \mathcal{F}' that replaced u . It can be easily seen that no ancestor-descendant relation is violated, and that the height of \mathcal{F}' is at most twice the height of \mathcal{F} .

For treewidth, it is possible to simply transform the original tree-decomposition of G into one of G' by replacing each vertex by the two vertices that replaced it, resulting in a new tree-decomposition of G' with at most twice the width of the original. Regarding the number of sources, it can be observed that each vertex that is not a source is replaced by two vertices which are not sources, and each source is replaced by two vertices of which precisely one is a source. ◀

2.5 SPQR-tree decomposition

Let G be a biconnected undirected graph. A pair of vertices is a *separation pair* if its removal disconnects G . A *split pair* is either a separation pair or a pair of adjacent vertices. A *split component* of G with respect to a split pair $\{u, v\}$ is either an edge (u, v) or a maximal subgraph $G_{uv} \subset G$ such that $\{u, v\}$ is not a split pair of G_{uv} . A split pair $\{s', t'\}$ of G is *maximal* with respect to a split pair $\{s, t\}$ of G , if for every other split pair $\{s^*, t^*\}$ of G , there is a split component that includes the vertices s', t', s and t .

An *SPQR-tree* T of G with respect to an edge e^* is a rooted tree that describes a recursive decomposition of G induced by its split pairs [14]. In what follows, we call *nodes* the vertices of T , to distinguish them from the vertices of G . The nodes of T are of four types S, P, Q,

and R. Each node μ of T is associated with a split pair $\{u, v\}$ of G , where u and v are the *poles* of μ , with a subgraph G_μ of G , called the *pertinent graph* of μ , which consists of one or more split components of G with respect to $\{u, v\}$, and with a multigraph $\text{sk}(\mu)$, called the *skeleton* of μ , which represents the arrangement of such split components in G_μ . The edges of $\text{sk}(\mu)$ are called *virtual edges*. Each node μ of T whose pertinent graph is not a single edge has some children, each corresponding to a split components of G in G_μ . Each of these children is the root of a subtree of T .

Formally, T is defined as follows. The root ρ of T is a Q-node corresponding to the edge $e^* = (u, v)$. The child σ of ρ has u and v as poles, and its pertinent graph is defined as $G_\sigma = G \setminus e^*$. Now consider a node μ of T with poles s and t and a pertinent graph G_μ ; assume that G_μ is *st-biconnectible*, i.e., it is either biconnected or it becomes so if the edge (s, t) is added to it. We distinguish some cases.

Base case: G_μ consists of a single edge e between s and t . Then, μ is a Q-node whose skeleton is the edge (s, t) . The node μ is a leaf of T .

Series case: G_μ is not biconnected; since G_μ is *st-biconnectible*, each cut-vertex of G_μ (i.e., each vertex whose removal disconnects G_μ) lies on any path from s to t . Then, μ is an S-node. Let v_1, \dots, v_{k-1} , where $k \geq 2$, be the cut vertices of G_μ , in the order in which they are encountered in any path from s to t . The skeleton of μ is a path consisting of the virtual edges e_1, \dots, e_k ; for $i = 1, \dots, k$, we have $e_i = (v_{i-1}, v_i)$, where $v_0 = s$ and $v_k = t$. Further, μ has k children ν_1, \dots, ν_k ; for $i = 1, \dots, k$, the poles of ν_i are v_{i-1} and v_i , and the pertinent graph G_{ν_i} of ν_i is the union of all the split components with respect to $\{v_{i-1}, v_i\}$ that do not contain both s and t . The decomposition recurs on the nodes ν_1, \dots, ν_k .

Parallel case: G_μ is not biconnected and $\{s, t\}$ is a split pair of G_μ . Then μ is a P-node. If the split pair $\{s, t\}$ defines k maximal split components of G_μ , then the skeleton of μ is a set of k parallel edges e_1, \dots, e_k between s and t . Further, μ has k children ν_1, \dots, ν_k ; for $i = 1, \dots, k$, the poles of ν_i are s and t , and the pertinent graph G_{ν_i} of ν_i is a maximal split component with respect to $\{s, t\}$. The decomposition recurs on the nodes ν_1, \dots, ν_k .

Rigid case: None of the other cases is applicable. Let $\{s_1, t_1\}, \dots, \{s_k, t_k\}$ be the maximal split pairs of G with respect to $\{s, t\}$ ($k \geq 1$) such that $\{s_i, t_i\}$ belongs to G_μ , for $i = 1, \dots, k$. Then μ is an R-node whose skeleton is the graph whose vertex set is $\{s_1, t_1\} \cup \dots \cup \{s_k, t_k\}$ and whose edge set is $\{(s_1, t_1), \dots, (s_k, t_k)\}$. Then μ has k children ν_1, \dots, ν_k ; for $i = 1, \dots, k$, the poles of ν_i are s_i and t_i , and the pertinent graph G_{ν_i} of ν_i is the union of all the split components with respect to $\{s_i, t_i\}$ that do not contain both s and t . The decomposition recurs on the nodes ν_1, \dots, ν_k .

Note that each virtual edge e_i in the skeleton of a node μ of T *corresponds* to the pertinent graph G_{ν_i} of a child ν_i of μ . We say that G_{ν_i} is a *component* of G_μ . Figs. 1a and 1b show a planar graph and its SPQR-tree. To simplify our algorithms, we assume that every S-node of T has two children. If this is not the case, we can modify T to achieve this property (see Fig. 1c). An SPQR-tree T of an n -vertex planar graph has $\mathcal{O}(n)$ Q-, S-, P-, and R-nodes. Also, the total number of vertices of the skeletons for the nodes in T is $\mathcal{O}(n)$ [14].

When talking about an SPQR-tree T of a biconnected directed graph G , we mean an SPQR-tree of its underlying graph. Let μ be a node of T with poles u and v . A *uv-external upward planar embedding* of G_μ is an upward planar embedding of G_μ such that u and v are incident to the outer face. In our algorithms, when testing the upward planarity of a digraph G , the fact that its SPQR-tree T is rooted at an edge e^* of G corresponds to the requirement that e^* is incident to the outer face of the upward planar embedding \mathcal{E} of G we are looking for. For each node μ of T , the restriction of \mathcal{E} to the vertices and edges of the pertinent graph G_μ of μ is a *uv-external upward planar embedding* of G_μ .





Figure 1 (a) A planar DAG G . (b) An SPQR-tree of G . For each node that is not a Q-node, the skeleton is depicted together with a dashed edge to represent the rest of the graph; for each Q-node, the corresponding edge is shown. (c) An SPQR-tree of G whose S-nodes have exactly two children.

3 The Shapes of Components

Let G be a biconnected DAG, let T be an SPQR-tree of G rooted at an edge e^* , let μ be a node of T with poles u and v , and let \mathcal{E}_μ be a uv -external upward planar embedding of G_μ . Let λ be the angle assignment defined by \mathcal{E}_μ . The poles u and v identify two paths on the boundary of the outer face f_0 of \mathcal{E}_μ : the *left outer path* $P_l = \langle v_0 = u, v_1, \dots, v_k = v \rangle$ is the path that leaves f_0 on the left when walking from u to v ; the *right outer path* $P_r = \langle w_0 = u, w_1, \dots, w_h = v \rangle$ of \mathcal{E}_μ is the path that leaves f_0 on the right when walking from u to v ; see Fig. 2. Notice that P_l and P_r may share some vertices other than u and v if G_μ is not biconnected. For $i = 0, 1, \dots, k$, let α_i denote the angle at v_i inside f_0 and, for $i = 0, 1, \dots, h$, let β_i denote the angle at w_i inside f_0 . The *left-turn-number* $\tau_l(\mathcal{E}_\mu, u, v)$ of \mathcal{E}_μ is defined as $\sum_{i=1}^{k-1} \lambda(\alpha_i)$, while the *right-turn-number* $\tau_r(\mathcal{E}_\mu, u, v)$ of \mathcal{E}_μ is $\sum_{i=1}^{h-1} \lambda(\beta_i)$. Note that $\alpha_0 = \beta_0$ and $\alpha_k = \beta_h$ are the angles at u and v inside f_0 , respectively. The values $\lambda(\alpha_0)$ and $\lambda(\alpha_k)$ are also denoted by $\lambda(\mathcal{E}_\mu, u)$ and $\lambda(\mathcal{E}_\mu, v)$, respectively. Finally, given a vertex $w \in \{u, v\}$, let $\rho_l(\mathcal{E}_\mu, w)$ denote the orientation of the edge e_l of P_l incident to w , that is, $\rho_l(\mathcal{E}_\mu, w) = in$ if e_l is an incoming edge for w , $\rho_l(\mathcal{E}_\mu, w) = out$ otherwise. Analogously, let $\rho_r(\mathcal{E}_\mu, w)$ denote the orientation of the edge e_r of P_r incident to w . The *shape description* of \mathcal{E}_μ is the tuple $\langle \tau_l(\mathcal{E}_\mu, u, v), \tau_r(\mathcal{E}_\mu, u, v), \lambda(\mathcal{E}_\mu, u), \lambda(\mathcal{E}_\mu, v), \rho_l(\mathcal{E}_\mu, u), \rho_r(\mathcal{E}_\mu, u), \rho_l(\mathcal{E}_\mu, v), \rho_r(\mathcal{E}_\mu, v) \rangle$; see Fig. 2.



■ **Figure 2** An upward planar embedding of a split component G_μ with poles u and v and shape description $\langle 3, 0, 0, -1, out, in, out, out \rangle$. The left (right) outer path is shown in green (orange).

We note that there are some dependencies between the values of a shape description. First, $\rho_l(\mathcal{E}_\mu, u)$ and $\rho_r(\mathcal{E}_\mu, u)$ have the same value if $\lambda(\mathcal{E}_\mu, u) \in \{-1, 1\}$, while they have different values if $\lambda(\mathcal{E}_\mu, u) = 0$. Similarly, $\rho_r(\mathcal{E}_\mu, v)$ is implied by the values $\rho_l(\mathcal{E}_\mu, v)$ and $\lambda(\mathcal{E}_\mu, v)$. Furthermore, the values of $\tau_l(\mathcal{E}_\mu, u, v)$ and $\rho_l(\mathcal{E}_\mu, u)$ imply the value of $\rho_l(\mathcal{E}_\mu, v)$; indeed, if $\tau_l(\mathcal{E}_\mu, u, v)$ is even, then $\rho_l(\mathcal{E}_\mu, u)$ and $\rho_l(\mathcal{E}_\mu, v)$ are different, while if $\tau_l(\mathcal{E}_\mu, u, v)$ is odd, then $\rho_l(\mathcal{E}_\mu, u)$ and $\rho_l(\mathcal{E}_\mu, v)$ are the same. Finally, $\lambda(\mathcal{E}_\mu, v)$ is implied by $\tau_l(\mathcal{E}_\mu, u, v)$, $\tau_r(\mathcal{E}_\mu, u, v)$, and $\lambda(\mathcal{E}_\mu, u)$. This is a consequence of the following observation, which adopts the notation of Theorem 1 and follows from the definition of the involved parameters.

► **Observation 7.** *We have $\tau_l(\mathcal{E}_\mu, u, v) + \tau_r(\mathcal{E}_\mu, u, v) + \lambda(\mathcal{E}_\mu, u) + \lambda(\mathcal{E}_\mu, v) = n_1(f_0) - n_{-1}(f_0)$.*

Observation 7, together with Property **UP3** of Theorem 1, implies the following.


► **Corollary 8.** *We have $\tau_l(\mathcal{E}_\mu, u, v) + \tau_r(\mathcal{E}_\mu, u, v) + \lambda(\mathcal{E}_\mu, u) + \lambda(\mathcal{E}_\mu, v) = 2$.*

Recall that if u is a top or bottom vertex of G , then it has at most one incoming edge or at most one outgoing edge, respectively, which is called the *special edge* of u . If G_μ contains this edge, then G_μ is a *special component* for u , otherwise we say that G_μ is a *normal component* for u . Note that, if u is a source or a sink of G , then it has no special component. The following lemma bounds the right-turn number of a uv -external upward planar embedding \mathcal{E}_μ of G_μ with respect to its left-turn number.

► **Lemma 9.** *Let G_μ be a split component of an upward plane digraph G with respect to a split pair $\{u, v\}$. Let \mathcal{E}_μ be the induced uv -external upward planar embedding of G_μ with shape description $\langle \tau_l(\mathcal{E}_\mu, u, v), \tau_r(\mathcal{E}_\mu, u, v), \lambda(\mathcal{E}_\mu, u), \lambda(\mathcal{E}_\mu, v), \rho_l(\mathcal{E}_\mu, u), \rho_r(\mathcal{E}_\mu, u), \rho_l(\mathcal{E}_\mu, v), \rho_r(\mathcal{E}_\mu, v) \rangle$. If $\tau_l(\mathcal{E}_\mu, u, v) = c$ is the left turn-number of \mathcal{E}_μ , then the right turn-number $\tau_r(\mathcal{E}_\mu, u, v)$ is $-c + h$ where $h \in \{0, 1, 2, 3, 4\}$. In particular, if G_μ is a normal component for both u and v , then $\{\lambda(\mathcal{E}_\mu, u), \lambda(\mathcal{E}_\mu, v)\} \subseteq \{-1, 1\}$ and*

1. $\tau_r(\mathcal{E}_\mu, u, v) = -c$, if $\lambda(\mathcal{E}_\mu, u) = \lambda(\mathcal{E}_\mu, v) = 1$ (see Fig. 3a);
2. $\tau_r(\mathcal{E}_\mu, u, v) = -c + 2$, if $\{\lambda(\mathcal{E}_\mu, u), \lambda(\mathcal{E}_\mu, v)\} = \{-1, 1\}$ (see Figs. 3b and 3c); and
3. $\tau_r(\mathcal{E}_\mu, u, v) = -c + 4$, if $\lambda(\mathcal{E}_\mu, u) = \lambda(\mathcal{E}_\mu, v) = -1$ (see Fig. 3d).

Proof. If G_μ is a normal component for u , then either all the edges incident to u are incoming u or they are all outgoing u , hence the angle incident to u in the outer face f_0 of \mathcal{E}_μ is a switch angle and $\lambda(\mathcal{E}_\mu, u) \in \{-1, 1\}$. On the other hand, if G_μ is a special component for u , then either $\lambda(\mathcal{E}_\mu, u) \in \{-1, 0\}$, if G_μ contains at least one incoming and at least one outgoing edge of u , or $\lambda(\mathcal{E}_\mu, u) = 1$, if G_μ contains only the special



■ **Figure 3** Illustrations of the various cases of Lemma 9 for normal components.

edge of u . Similar considerations hold true for v . The rest of the statement follows by Corollary 8. Indeed, we have that $\tau_l(\mathcal{E}_\mu, u, v) + \tau_r(\mathcal{E}_\mu, u, v) + \lambda(\mathcal{E}_\mu, u) + \lambda(\mathcal{E}_\mu, v) = 2$, hence $\tau_r(\mathcal{E}_\mu, u, v) = 2 - \tau_l(\mathcal{E}_\mu, u, v) - \lambda(\mathcal{E}_\mu, u) - \lambda(\mathcal{E}_\mu, v)$, from which the statements follow for $\tau_l(\mathcal{E}_\mu, u, v) = c$ and by substituting the values of $\lambda(\mathcal{E}_\mu, u)$ and $\lambda(\mathcal{E}_\mu, v)$ of each case. ◀

The following two lemmata establish some useful properties for the shape descriptions of components that are parallel compositions of sub-components. Let G_μ be the pertinent digraph of a P-node μ . Let $\nu'_1, \nu'_2, \dots, \nu'_{k'}$ be a subset of the children of μ in T and let G'_μ be the subgraph of G_μ consisting of the union of $G_{\nu'_1}, G_{\nu'_2}, \dots, G_{\nu'_{k'}}$. Assume that the upward planar embedding \mathcal{E} is such that the pertinent graphs $G_{\nu'_1}, G_{\nu'_2}, \dots, G_{\nu'_{k'}}$ appear in this order from left to right around u . Denote by \mathcal{E}_ν the upward planar embedding of G_ν in \mathcal{E} . Then the outer boundary of G'_μ in \mathcal{E} is formed by the left outer path of $\mathcal{E}_{\nu'_1}$ and the right outer path of $\mathcal{E}_{\nu'_{k'}}$. Let f'_i be the face bounded by the right outer path of $\mathcal{E}_{\nu'_i}$ and by the left outer path of $\mathcal{E}_{\nu'_{i+1}}$, for $i = 1, 2, \dots, k' - 1$, and let f'_0 be the outer face of G'_μ in \mathcal{E} .

► **Lemma 10.** *Suppose f'_i is an internal face of G'_μ with respect to an upward planar embedding \mathcal{E} where c is the right-turn-number of $\mathcal{E}_{\nu'_i}$, that is $c = \tau_r(\mathcal{E}_{\nu'_i}, u, v)$, and $\alpha_u(f'_i)$ and $\alpha_v(f'_i)$ are the angles at vertices u and v inside f'_i . Then the left-turn-number $\tau_l(\mathcal{E}_{\nu'_{i+1}}, u, v)$ of $G_{\nu'_{i+1}}$ satisfies:*

1. $\tau_l(\mathcal{E}_{\nu'_{i+1}}, u, v) = -c$, if $\lambda(\alpha_u(f'_i)) = \lambda(\alpha_v(f'_i)) = 1$
2. $\tau_l(\mathcal{E}_{\nu'_{i+1}}, u, v) = -c - 2$, if $\{\lambda(\alpha_u(f'_i)), \lambda(\alpha_v(f'_i))\} = \{-1, 1\}$
3. $\tau_l(\mathcal{E}_{\nu'_{i+1}}, u, v) = -c - 4$, if $\lambda(\alpha_u(f'_i)) = \lambda(\alpha_v(f'_i)) = -1$.

Proof. By UP3 we have that $n_1(f'_i) + n_{-1}(f'_i) = -2$. Since the boundary of f'_i consists of the right outer path of $\mathcal{E}_{\nu'_i}$ and the right outer path of $\mathcal{E}_{\nu'_{i+1}}$, it follows that $n_1(f'_i) + n_{-1}(f'_i) = \tau_r(\mathcal{E}_{\nu'_i}, u, v) + \tau_l(\mathcal{E}_{\nu'_{i+1}}, u, v) + \lambda(\alpha_u(f'_i)) + \lambda(\alpha_v(f'_i)) = -2$. The lemma's statements follow by replacing $\tau_r(\mathcal{E}_{\nu'_i}, u, v)$ with c and substituting the considered values of $\lambda(\alpha_u(f'_i))$ and $\lambda(\alpha_v(f'_i))$. ◀

Let ν be a child of μ and let G_ν be the corresponding pertinent graph with poles u and v . Suppose that G_ν admits a uv -external upward planar embedding \mathcal{E}_ν such that the shape description of G_ν is s . We say that s is a *thin shape description* if it satisfies the following conditions, which we call *thin conditions*:

- $\tau_r(\mathcal{E}_\nu, u, v) = -\tau_l(\mathcal{E}_\nu, u, v)$;
- $\lambda(\mathcal{E}_\nu, u) = \lambda(\mathcal{E}_\nu, v) = 1$;
- $\rho_l(\mathcal{E}_\nu, u) = \rho_r(\mathcal{E}_\nu, u)$;
- $\rho_l(\mathcal{E}_\nu, v) = \rho_r(\mathcal{E}_\nu, v)$.

Let N_μ be the set of children $\nu_1, \nu_2, \dots, \nu_k$ of μ . For any subset N of N_μ , we denote by G_N the subgraph of G_μ that consists of all component G_ν with $\nu \in N$. Consider two subsets N_1 and N_2 of N such that $N_2 \subseteq N_1$ and assume that G_{N_1} and G_{N_2} are upward planar. We say that the upward planar embeddings \mathcal{E}_{N_1} and \mathcal{E}_{N_2} of G_{N_1} and G_{N_2} respectively, *form an equivalence pair* if the components of G_{N_2} appear in the same order in G_{N_1} around u starting from the outer face, and for every $\nu \in N_2$ component G_ν has the same shape description in both \mathcal{E}_{N_1} and \mathcal{E}_{N_2} . Let s be a thin shape description, and let N_s be a subset of N_μ such that the pertinent graph G_ν , for $\nu \in N_s$ admits an upward planar embedding with shape description s . For $N \subseteq N_\mu$ with $N \cap N_s \neq \emptyset$, we say that an upward planar embedding \mathcal{E}_N of G_N *respects shape description s for the subset N_s* if the components G_ν for $\nu \in N \cap N_s$ are consecutive around u when starting from the outer face of \mathcal{E}_N , and all have shape description s in \mathcal{E}_N .

► **Lemma 11.** *Let N_s be a subset of N_μ such that the pertinent graph G_ν , for $\nu \in N_s$ admits an upward planar embedding with thin shape description s . Let also N_1 and N_2 be two subsets of N with $N_2 \subseteq N_1$ and $N_1 \setminus N_2 \subseteq N_s$. Then, G_{N_1} has an upward planar embedding \mathcal{E}_{N_1} that respects s for N_s if and only if G_{N_2} has an upward planar embedding \mathcal{E}_{N_2} that respects s for N_s , and embeddings \mathcal{E}_{N_1} and \mathcal{E}_{N_2} form an equivalence pair. Furthermore, the shape descriptions of G_{N_1} and G_{N_2} in \mathcal{E}_{N_1} and \mathcal{E}_{N_2} , respectively, are the same.*

Proof. Suppose first that G_{N_1} has an upward planar embedding \mathcal{E}_{N_1} that respects s for N_s . The restriction of \mathcal{E}_{N_1} to G_{N_2} is an upward planar embedding \mathcal{E}_{N_2} of G_{N_2} that respects s for N_s . Moreover the components shared by G_{N_1} and G_{N_2} have the same shape description and appear in the same order around u , that is, \mathcal{E}_{N_1} and \mathcal{E}_{N_2} form an equivalence pair.

Suppose now that G_{N_2} has an upward planar embedding \mathcal{E}_{N_2} that respects s for N_s . Let $G_{\nu'_1}, G_{\nu'_2}, \dots, G_{\nu'_{k'}}$ be the sequence of the components of G_{N_2} in the left to right order defined by \mathcal{E}_{N_2} . Denote by \mathcal{E}_ν^2 the upward planar embedding of component G_ν in \mathcal{E}_{N_2} , for $\nu \in N_2$. Let f'_i be the face bounded by the right outer path of $\mathcal{E}_{\nu'_i}^2$ and by the left outer path of $\mathcal{E}_{\nu'_{i+1}}^2$, for $i = 1, 2, \dots, k' - 1$, and let f'_0 be the outer face of \mathcal{E}_{N_2} . As \mathcal{E}_{N_2} respects N_s , there exist consecutive components $G_{\nu'_\ell}, G_{\nu'_{\ell+1}}, \dots, G_{\nu'_{\ell+j}}$ of G_{N_2} such that $\nu'_{\ell+i} \in N_s$ and $\mathcal{E}_{\nu'_{\ell+i}}^2$ has shape description s , for $i = 0, \dots, j$. We now change the upward planar embedding \mathcal{E}_{N_2} of G_{N_2} to an upward planar embedding \mathcal{E}_{N_1} of G_{N_1} as follows. Note that, since $N_2 \subseteq N_1 \cup N_s$, if component G_ν of G_{N_1} does not belong to G_{N_2} then $\nu \in N_s$. We replace components $G_{\nu'_{\ell+i}}$ ($i = 0, \dots, j$) with a sequence consisting of all the components G_ν of G_{N_1} where $\nu \in N_s$. Let \mathcal{G}_s denote the subgraph of G_{N_1} that contains components G_ν where $\nu \in N_1 \cap N_s$. To completely define \mathcal{E}_{N_1} we need to define the upward planar embedding of all the components in \mathcal{G}_s , the labels of the angles at u and v inside the faces formed by consecutive elements of \mathcal{G}_s , and the labels of the angles at u and v inside the faces $f'_{\ell-1}$ and $f'_{\ell+j}$ (notice that the boundary of both faces is changed). For each component of \mathcal{G}_s we choose an upward planar embedding such that the resulting shape description is s . We label -1 all the angles at u and v inside the faces formed by consecutive elements of \mathcal{G}_s . For the angles at u and v inside $f'_{\ell-1}$ and $f'_{\ell+j}$, we preserve the labels they had in \mathcal{E}_{N_2} . We denote by \mathcal{E}_ν^1 the upward planar embedding of G_ν in \mathcal{E}_{N_1} for $\nu \in N_1$. Note that $\mathcal{E}_\nu^1 = \mathcal{E}_\nu^2$ for every $\nu \in N_1 \setminus N_s$. In order to prove that the resulting labeling is a valid angle assignment we need to prove that properties **UP0**–**UP3** hold. Consider an angle α at a vertex w . If α is an angle that exists also in \mathcal{E}_{N_2} , then **UP0** holds for α in \mathcal{E}_{N_1} as well; if α is an angle inside an internal face of \mathcal{E}_ν^1 of a component $G_\nu \in \mathcal{G}_s$ then **UP0** holds for α in \mathcal{E}_{N_1} because it holds in the upward planar embedding chosen for G_ν . If α is an angle inside the outer face of \mathcal{E}_ν^1 at a vertex different from u and v , then again, **UP0** holds in \mathcal{E}_{N_1} because it holds

in the upward planar embedding chosen for G_ν . If α is an angle at u or at v inside a face f_j formed by consecutive elements G_ν and $G_{\nu'}$ of \mathcal{G}_s then the two edges defining α have the same orientation because the two upward planar embeddings \mathcal{E}_ν^1 and $\mathcal{E}_{\nu'}^1$ have the same shape description; since α is labeled 1, **UP0** holds. Finally, if α is an angle at u or at v inside $f'_{\ell-1}$ or $f'_{\ell+j}$, then the two edges defining α have the same orientation with the two edges defining the corresponding angle in \mathcal{E}_{N_1} (because the leftmost component $G_{\nu'_\ell}$ of G_{N_2} that is replaced by the sequence of components in \mathcal{G}_s has the same shape description in \mathcal{E}_{N_2} as the leftmost component of \mathcal{G}_s in \mathcal{E}_{N_1} , and the rightmost component $G_{\nu'_{\ell+j}}$ of G_{N_2} has the same shape description in \mathcal{E}_{N_2} with the rightmost component of \mathcal{G}_s in \mathcal{E}_{N_1}). Since the label at α is preserved, **UP0** holds because it holds in \mathcal{E}_{N_2} . Concerning **UP1** and **UP2**, we observe that for each vertex different from u and v **UP1** or **UP2** holds in \mathcal{E}_{N_1} because it either holds in \mathcal{E}_{N_2} or it holds in the upward planar embedding chosen for the components of \mathcal{G}_s . So, the only vertices for which we have to prove **UP1** or **UP2** are u and v . To this aim we observe that either **UP1** or **UP2** holds for u and v in \mathcal{E}_{N_2} and that the angles at u and at v that are in \mathcal{E}_{N_1} but not in \mathcal{E}_{N_2} are all labeled -1 ; thus, they cannot create a violation of **UP1** or of **UP2**. Concerning **UP3**, consider a face f of \mathcal{E}_{N_1} . If f is also a face of \mathcal{E}_{N_2} then **UP3** holds for f in \mathcal{E}_{N_1} because it holds in \mathcal{E}_{N_2} . If f is an internal face of \mathcal{E}_ν^1 of a component $G_\nu \in \mathcal{G}_s$, then **UP3** holds for \mathcal{E}_{N_1} because it holds for the upward planar embedding chosen for G_ν . If f is a face f_j formed by consecutive elements G_ν and $G_{\nu'}$ of \mathcal{G}_s , then its boundary is formed by the right outer path of \mathcal{E}_ν^1 and by the left outer path of $\mathcal{E}_{\nu'}^1$. Since G_ν and $G_{\nu'}$ have the same shape description that satisfies the thin conditions, we have $\tau_r(\mathcal{E}_\nu^1, u, v) = -\tau_l(\mathcal{E}_{\nu'}^1, u, v)$; moreover, the two angles at u and v inside f_j are both labeled -1 . Thus $n_1(f) - n_{-1}(f) = \tau_r(\mathcal{E}_\nu^1, u, v) + \tau_l(\mathcal{E}_{\nu'}^1, u, v) - 2 = -2$ and **UP3** holds. Consider now the case when f is the face $f'_{\ell-1}$ and assume that it is an internal face (a similar proof holds when it is the outer face). The boundary of $f'_{\ell-1}$ in \mathcal{E}_{N_2} consists of the right outer path of $\mathcal{E}_{\nu'_{\ell-1}}^2$ and of the left outer path of $\mathcal{E}_{\nu'_\ell}^2$; since **UP3** holds for $f'_{\ell-1}$ in \mathcal{E}_{N_2} , we have $n_1(f) - n_{-1}(f) = \tau_r(\mathcal{E}_{\nu'_{\ell-1}}^2, u, v) + \tau_l(\mathcal{E}_{\nu'_\ell}^2, u, v) + \alpha_u + \alpha_v = -2$, where α_u and α_v are the angles at u and v inside $f'_{\ell-1}$, respectively. The boundary of $f'_{\ell-1}$ in \mathcal{E}_{N_1} consists of the right outer path of $\mathcal{E}_{\nu'_{\ell-1}}^1$ and of the left outer path of \mathcal{E}_ν^1 , where G_ν is the first component of the set \mathcal{G}_s . Thus, we have $n_1(f) - n_{-1}(f) = \tau_r(\mathcal{E}_{\nu'_{\ell-1}}^1, u, v) + \tau_l(\mathcal{E}_\nu^1, u, v) + \alpha_u + \alpha_v$; since $\mathcal{E}_{\nu'_{\ell-1}}^2$ and \mathcal{E}_ν^1 have the same shape description we have $\tau_l(\mathcal{E}_{\nu'_{\ell-1}}^2, u, v) = \tau_l(\mathcal{E}_\nu^1, u, v)$ and therefore **UP3** holds for $f'_{\ell-1}$ in \mathcal{E}_{N_1} . If f is the face $f'_{\ell+j}$, the proof is analogous to the one for $f'_{\ell-1}$. This concludes the proof that properties **UP0–UP3** hold for \mathcal{E}_{N_1} . Now, since all \mathcal{G}_s are consecutive in \mathcal{E}_{N_1} and have the same shape description s , it follows that \mathcal{E}_{N_1} respects s for N_s . Additionally \mathcal{E}_{N_1} and \mathcal{E}_{N_2} form an equivalence pair. Hence we proved that if G_{N_2} has an upward planar embedding \mathcal{E}_{N_2} that respects s for N_s , then G_{N_1} has an upward planar embedding \mathcal{E}_{N_1} that respects s for N_s and the two embeddings form an equivalence pair.

It remains to prove that the shape descriptions of G_{N_1} in \mathcal{E}_{N_1} and of G_{N_2} in \mathcal{E}_{N_2} are the same. In general, the shape description of a component in an upward planar embedding only depends on the outer boundary of that component. Let f_0^1 and f_0^2 be the outer faces of the two upward planar embeddings \mathcal{E}_{N_1} and \mathcal{E}_{N_2} described above. The boundary of f_0^2 consists of the left outer boundary of $\mathcal{E}_{\nu'_1}^2$ and by the right outer boundary of $\mathcal{E}_{\nu'_{k'}}^2$. If neither ν'_1 nor $\nu'_{k'}$ belong to N_s , then the boundary of f_0^1 coincides with that of f_0^2 , they have the same labeling and therefore \mathcal{E}_{N_1} and \mathcal{E}_{N_2} have the same shape description. If this is not the case then either $\nu'_1 \in N_s$ or $\nu'_{k'} \in N_s$. If $\nu'_1 \in N_s$ the left boundary of f_0^2 consists of the left outer path of $\mathcal{E}_{\nu'_1}^2$ and the left boundary of f_0^1 consists of the left outer path of \mathcal{E}_ν^1 , where G_ν is a component of \mathcal{G}_s . Since $\mathcal{E}_{\nu'_1}^2$ and \mathcal{E}_ν^1 have the same shape description, they have the same

left-turn-number. Similarly, if $\nu'_{k'} \in N_s$ the right outer paths of f_0^2 and f_0^1 have the same right-turn-numbers. Since the angles at u and v inside f_0^2 are the same as those inside f_0^1 , \mathcal{E}_{N_1} and \mathcal{E}_{N_2} have the same shape description. \blacktriangleleft

4 General Algorithm

Let G be an n -vertex biconnected expanded DAG whose underlying undirected graph is planar and let T be an SPQR-tree decomposition of G . Let also τ_{\min} and τ_{\max} be two integers such that $\tau_{\min} \leq \tau_{\max}$. In this section, we present a general algorithm to compute the shape descriptions of all possible upward planar embeddings of G with respect to T , and such that the left- and right-turn numbers of the induced upward planar embeddings of all pertinent graphs of T are within the range $[\tau_{\min}, \tau_{\max}]$. In case G is not upward planar under these restrictions, the algorithm returns an empty set. We visit the nodes of T bottom-up and we compute for each node μ the set \mathcal{F}_μ of the shape descriptions of all possible upward planar embeddings of the pertinent graph G_μ of μ . We call \mathcal{F}_μ the *feasible set* of μ . The feasible set \mathcal{F}_μ is computed starting from the feasible sets $\mathcal{F}_{\nu_1}, \mathcal{F}_{\nu_2}, \dots, \mathcal{F}_{\nu_k}$, where $\nu_1, \nu_2, \dots, \nu_k$ are the children of μ in T . If at any point the feasible set \mathcal{F}_μ of node μ is empty, we conclude that G is not upward planar (under the above restrictions) and the process returns an empty set, otherwise we continue the traversal of T .

Storing feasible sets

For each node μ of T we associate a matrix $M(\mu)$ of size $(\tau_{\max} - \tau_{\min} + 1) \times 5$ where the element $M(\mu)[i, j]$ of the matrix contains all shape descriptions for G_μ with left-turn-number $\tau_l = \tau_{\min} + i$ and right-turn-number $\tau_r = -\tau_l + j$. Note that by Lemma 9, the right-turn-number of G_μ can only take values in $[-c, -c + 4]$, where c is the left-turn-number τ_l . A shape description within $M(\mu)[i, j]$ is stored as a pair $\langle \lambda(\mathcal{E}_\mu, u), \rho_l(\mathcal{E}_\mu, u) \rangle$ as its first two values are implied by indices i and j and the other four values of the shape description can be derived from the indices and the stored values (recall that the values of a shape description are not all independent). The following lemma summarizes the space requirements and run-time of basic operations for feasible sets.

► **Lemma 12.** *Let $\tau = \tau_{\max} - \tau_{\min} + 1$. Then:*

1. *creating matrix $M(\mu)$ for all nodes μ of T takes $\mathcal{O}(|T|\tau)$ time and $\mathcal{O}(|T|\tau)$ space;*
2. *retrieving all shape descriptions of $M(\mu)$ takes $\mathcal{O}(\tau)$ time;*
3. *adding, deleting or finding a shape description in $M(\mu)$ requires $\mathcal{O}(1)$ time;*
4. *there are at most 18 shape descriptions with given left- or right-turn-number, and finding them takes $\mathcal{O}(1)$ time.*

Proof. The first part of the lemma is obvious. We now argue that for specific values of $\tau_l(\mathcal{E}_\mu, u, v)$ and $\tau_r(\mathcal{E}_\mu, u, v)$ there is a bounded number of possible shape descriptions with the given left- and right-turn-numbers. Indeed, for the pair $\langle \lambda(\mathcal{E}_\mu, u), \rho_l(\mathcal{E}_\mu, u) \rangle$, $\lambda(\mathcal{E}_\mu, u)$ takes a value from $\{-1, 0, 1\}$, while $\rho_l(\mathcal{E}_\mu, u) \in \{in, out\}$. Hence the possible values for the pairs are in total six. Hence, each element $M(\mu)[i, j]$ of matrix $M(\mu)$ contains at most six pairs, and parts 2 and 3 of the lemma follows. For the last part of the lemma, we first argue that the number of shape descriptions with given left- or right-turn-number are at most 18. Indeed, if we specify the values $\lambda(\mathcal{E}_\mu, u)$, $\lambda(\mathcal{E}_\mu, v)$, $\rho_l(\mathcal{E}_\mu, u)$, and either the left- or the right-turn-number, then the remaining values of a shape description are uniquely determined. As there are $3^2 \cdot 2 = 18$ ways to specify all of $\lambda(\mathcal{E}_\mu, u)$, $\lambda(\mathcal{E}_\mu, v)$ and $\rho_l(\mathcal{E}_\mu, u)$, it follows that

there are at most 18 shape descriptions with specific left- or right-turn-number. Now, the shape descriptions with left-turn-number equal to τ_l are stored in row $i = \tau_l - \tau_{\min}$ of $M(\mu)$, which has length 5. On the other hand, if the right-turn-number τ_r is given, by Lemma 9 the right-turn-number can only take values in the interval $[\tau_r, -\tau_r + 4]$, and therefore it suffices to consider the pairs of at most five elements of the matrix. In both cases the shape descriptions with given left- or right-turn-number can be found in $\mathcal{O}(1)$ time. \blacktriangleleft

We describe how to compute the feasible set \mathcal{F}_μ of a node μ of T depending on its type.

Q-node

In this case, the pertinent graph G_μ is a single directed edge connecting poles u and v . If the edge is directed from u to v then the feasible set consists of the tuple $\langle 0, 0, 1, 1, out, out, in, in \rangle$, while if the edge is directed from v to u then it consists of the tuple $\langle 0, 0, 1, 1, in, in, out, out \rangle$.

► **Lemma 13.** *Let μ be a Q-node of T . The feasible set \mathcal{F}_μ can be computed in $\mathcal{O}(1)$ time.*

S-node

Recall that node μ has exactly two children in T , ν_1 and ν_2 . Let u_1, v_1 be the poles of G_{ν_1} and u_2, v_2 be the poles of G_{ν_2} . The poles of G_μ are u_1 and v_2 , that is $v_1 = u_2$. Let $\langle \tau_l(\mathcal{E}_{\nu_1}, u_1, v_1), \tau_r(\mathcal{E}_{\nu_1}, u_1, v_1), \lambda(\mathcal{E}_{\nu_1}, u_1), \lambda(\mathcal{E}_{\nu_1}, v_1), \rho_l(\mathcal{E}_{\nu_1}, u_1), \rho_r(\mathcal{E}_{\nu_1}, u_1), \rho_l(\mathcal{E}_{\nu_1}, v_1), \rho_r(\mathcal{E}_{\nu_1}, v_1) \rangle$ be a tuple in \mathcal{F}_{ν_1} and let $\langle \tau_l(\mathcal{E}_{\nu_2}, u_2, v_2), \tau_r(\mathcal{E}_{\nu_2}, u_2, v_2), \lambda(\mathcal{E}_{\nu_2}, u_2), \lambda(\mathcal{E}_{\nu_2}, v_2), \rho_l(\mathcal{E}_{\nu_2}, u_2), \rho_r(\mathcal{E}_{\nu_2}, u_2), \rho_l(\mathcal{E}_{\nu_2}, v_2), \rho_r(\mathcal{E}_{\nu_2}, v_2) \rangle$ be a tuple in \mathcal{F}_{ν_2} .


Let α_l be the angle at the common pole that is created by the two left outer paths of \mathcal{E}_{ν_1} and \mathcal{E}_{ν_2} (see Fig. 4). Similarly, let α_r be the angle at the common pole that is created by the two right outer paths of \mathcal{E}_{ν_1} and \mathcal{E}_{ν_2} . We assign the labels λ_l and λ_r to α_l and α_r respectively as follows: $\lambda_l = 0$ if $\rho_l(\mathcal{E}_\mu, v_1) \neq \rho_l(\mathcal{E}_\mu, u_2)$ otherwise $\lambda_l \in \{-1, 1\}$, and $\lambda_r = 0$ if $\rho_r(\mathcal{E}_\mu, v_1) \neq \rho_r(\mathcal{E}_\mu, u_2)$ otherwise $\lambda_r \in \{-1, 1\}$. Note that we exclude the case $\lambda_l = \lambda_r = 1$ as this would imply that two angles on the same vertex of G are large, which is not possible. Given the two values for λ_l and λ_r , we construct a candidate tuple $\langle \tau_l(\mathcal{E}_\mu, u_1, v_2), \tau_r(\mathcal{E}_\mu, u_1, v_2), \lambda(\mathcal{E}_\mu, u_1), \lambda(\mathcal{E}_\mu, v_2), \rho_l(\mathcal{E}_\mu, u_1), \rho_r(\mathcal{E}_\mu, u_1), \rho_l(\mathcal{E}_\mu, v_2), \rho_r(\mathcal{E}_\mu, v_2) \rangle$ with

- $\tau_l(\mathcal{E}_\mu, u_1, v_2) = \tau_l(\mathcal{E}_{\nu_1}, u_1, v_1) + \tau_l(\mathcal{E}_{\nu_2}, u_2, v_2) + \lambda_l$,
- $\tau_r(\mathcal{E}_\mu, u_1, v_2) = \tau_r(\mathcal{E}_{\nu_1}, u_1, v_1) + \tau_r(\mathcal{E}_{\nu_2}, u_2, v_2) + \lambda_r$,
- $\lambda(\mathcal{E}_\mu, u_1) = \lambda(\mathcal{E}_{\nu_1}, u_1)$,
- $\lambda(\mathcal{E}_\mu, v_2) = \lambda(\mathcal{E}_{\nu_2}, v_2)$,
- $\rho_l(\mathcal{E}_\mu, u_1) = \rho_l(\mathcal{E}_{\nu_1}, u_1)$,
- $\rho_r(\mathcal{E}_\mu, u_1) = \rho_r(\mathcal{E}_{\nu_1}, u_1)$,
- $\rho_l(\mathcal{E}_\mu, v_2) = \rho_l(\mathcal{E}_{\nu_2}, v_2)$,
- $\rho_r(\mathcal{E}_\mu, v_2) = \rho_r(\mathcal{E}_{\nu_2}, v_2)$.

We accept the candidate tuple if and only if $\tau_l(\mathcal{E}_\mu, u_1, v_2) + \tau_r(\mathcal{E}_\mu, u_1, v_2) + \lambda(\mathcal{E}_\mu, u_1) + \lambda(\mathcal{E}_\mu, v_2) = 2$ holds, that is if Corollary 8 holds for the outer face of \mathcal{E}_μ .

► **Lemma 14.** *Let μ be an S-node of T with children ν_1 and ν_2 . The feasible set \mathcal{F}_μ can be computed in $\mathcal{O}(\tau + |\mathcal{F}_{\nu_1}| \cdot |\mathcal{F}_{\nu_2}|)$ time.*

Proof. By Lemma 12, getting the shape descriptions of each feasible set \mathcal{F}_{ν_1} and \mathcal{F}_{ν_2} takes $\mathcal{O}(\tau)$ time. There are $|\mathcal{F}_{\nu_1}| \cdot |\mathcal{F}_{\nu_2}|$ pairs of shape descriptions for \mathcal{E}_{ν_1} and \mathcal{E}_{ν_2} , and for each pair, the candidate tuple is constructed in $\mathcal{O}(1)$ time. Finally, by Lemma 12 storing each accepted



■ **Figure 4** Series composition. The resulting shape description is $\langle 2, 2, -1, -1, in, in, out, out \rangle$



■ **Figure 5** Different shape descriptions corresponding to the shape sequence $[s_1, s_2]$, with $s_1 = \langle 0, 0, 1, 1, out, out, in, in \rangle$ (gray shaded) and $s_2 = \langle -2, 2, 1, 1, out, out, in, in \rangle$ (orange shaded). The shape descriptions are (a) $\langle 0, 2, 1, -1, out, out, in, in \rangle$ and (b) $\langle 0, 2, -1, 1, out, out, in, in \rangle$.

tuple requires $\mathcal{O}(1)$ time. Hence, the feasible set \mathcal{F}_μ can be computed in $\mathcal{O}(\tau + |\mathcal{F}_{\nu_1}| \cdot |\mathcal{F}_{\nu_2}|)$ time, as claimed. ◀

P-node

Let μ be a P-node with poles u and v and children $\nu_1, \nu_2, \dots, \nu_k$. Let $\nu'_1, \nu'_2, \dots, \nu'_{k'}$ be a subset of the children of μ and let G'_μ be the subgraph of G_μ consisting of the union of $G_{\nu'_1}, G_{\nu'_2}, \dots, G_{\nu'_{k'}}$. Consider an upward planar embedding \mathcal{E}'_μ of G'_μ and assume that the pertinent graphs $G_{\nu'_1}, G_{\nu'_2}, \dots, G_{\nu'_{k'}}$ appear in this order from left to right around u . Denote by $\mathcal{E}'_{\nu'_i}$ the uv -external upward planar embedding of $G_{\nu'_i}$, for $i = 1, 2, \dots, k'$. Then the outer boundary of \mathcal{E}'_μ is formed by the left outer path of $\mathcal{E}'_{\nu'_1}$ and the right outer path of $\mathcal{E}'_{\nu'_{k'}}$. Denote by f'_i the face bounded by the right outer path of $\mathcal{E}'_{\nu'_i}$ and by the left outer path of $\mathcal{E}'_{\nu'_{i+1}}$, for $i = 1, 2, \dots, k' - 1$, and let f'_0 be the outer face of G'_μ . Denote by S' the sequence of shape descriptions of the pertinent digraphs of $\nu'_1, \nu'_2, \dots, \nu'_{k'}$ in the order $G_{\nu'_1}, G_{\nu'_2}, \dots, G_{\nu'_{k'}}$. The sequence S' is the *shape sequence* of G'_μ with respect to the upward planar embedding \mathcal{E}'_μ . To describe the sequence S' we use the following notation: a^* denotes a subsequence of S' consisting of 0 or more elements equal to a ; a^+ denotes a subsequence of S' consisting of 1 or more elements equal to a . Not all sequences of shape descriptions of $\mathcal{E}'_{\nu'_1}, \mathcal{E}'_{\nu'_2}, \dots, \mathcal{E}'_{\nu'_{k'}}$ are shape sequences of G'_μ . For G'_μ we say that a shape description s' *corresponds* to a shape sequence S' if there exists an upward planar embedding of G'_μ with shape description s' and whose shape sequence is S' . Notice that, the same shape sequence might correspond to different upward planar embeddings of G'_μ , even if the order of $G_{\nu'_1}, G_{\nu'_2}, \dots, G_{\nu'_{k'}}$ is the same (see Fig. 5). Let S be a sequence of shape descriptions; the *reduced sequence* of S is the sequence of shape descriptions obtained from S by replacing each maximal subsequence a^+ of S with the single element a . We define the *size* of S as the number of elements in its reduced sequence.

► **Lemma 15.** *Let S be a sequence of shape descriptions from the feasible sets of $G_{\nu'_1}, G_{\nu'_2},$*

$\dots, G_{\nu_{k'}}'$. We can decide whether S is a shape sequence of G'_μ and compute the corresponding shape descriptions of G'_μ in $\mathcal{O}(r^3)$ time, where r is the size of S . Furthermore there are $\mathcal{O}(r^2)$ computed shape descriptions of G'_μ .

Proof. Assume first that S is reduced. We want to use **UP0-UP3** in order to compute all possible corresponding shape descriptions of G'_μ . Let \mathcal{E}'_μ be an upward planar embedding of G'_μ with shape sequence S . Let $G_{\nu'_i}$ be a component of G'_μ . Denote by $\mathcal{E}_{\nu'_i}$ its uv -external upward embedding induced by \mathcal{E}'_μ and let $s_i = \langle \tau_l(s_i), \tau_r(s_i), \lambda_u(s_i), \lambda_v(s_i), \rho_l(u, s_i), \rho_r(u, s_i), \rho_l(v, s_i), \rho_r(v, s_i) \rangle$ be its shape description in S . For all faces f'_i of \mathcal{E}'_μ , **UP3** gives $\tau_r(s_i) + \tau_l(s_{i+1}) + \lambda(\alpha_u(f'_i)) + \lambda(\alpha_v(f'_i)) = -2$ holds if $i \neq 0$, and $\tau_r(s_{k'}) + \tau_l(s_1) + \lambda(\alpha_u(f'_0)) + \lambda(\alpha_v(f'_0)) = 2$ for the outer face f'_0 . For the labels at the angles $\alpha_u(f'_i)$ and $\alpha_v(f'_i)$, we have that $\lambda(\alpha_w(f'_i)) = 0$ (i.e. $\alpha_w(f'_i)$ is a flat angle) if and only if $\rho_r(w, s_i) \neq \rho_l(w, s_{i+1})$, where w is a pole of μ . Hence the label of $\alpha_w(f'_i)$ is not determined only if it is a switch angle and its label can be either -1 or 1 . Note that by **UP2** there is exactly one angle with label 1 . Then, if u is a switch vertex, the possible labelings of all angles $\alpha_u(f'_i)$ are $r + 1$; namely there are r faces f'_i that can have the unique label 1 , or none of them has label 1 . Similarly if v is a switch vertex, the possible labelings of all angles $\alpha_v(f'_i)$ are $r + 1$. This implies that there exist in total at most $(r + 1)^2$ different labelings for the angles at u and v inside all faces f'_i of G'_μ . For each labeling, we want to test whether **UP3** holds for all faces f'_i of G'_μ . This can be done in constant time for each face, and thus in $\mathcal{O}(r)$ time for all faces. Still we need to guarantee that **UP1** and **UP2** also hold.

Assume that pole w is a switch vertex of G'_μ . Note that if the shape description for a component $G_{\nu'_i}$ has label at pole w equal to -1 , that is $\lambda(w, \mathcal{E}_{\nu'_i}) = -1$, then the large angle of w is inside an interior face of $\mathcal{E}_{\nu'_i}$. Then $\lambda(\alpha_w(f'_i)) = -1$ must hold for all faces f'_i of \mathcal{E}'_μ , and $\lambda(w, \mathcal{E}_{\nu'_j}) = 1$ for all components $G_{\nu'_j}$ different from $G_{\nu'_i}$. Otherwise, $\lambda(w, \mathcal{E}_{\nu'_i}) = 1$ holds for all components $G_{\nu'_i}$, and the large angle of w must be inside a face f'_i of \mathcal{E}'_μ , that is $\lambda(\alpha_w(f'_i)) = 1$ and $\lambda(\alpha_w(f'_j)) = -1$ for all other faces f'_j . Note that all shape descriptions used for the components of G'_μ belong to their feasible sets. Then all angles of w in the interior of all components (except possibly for one angle in the interior of one $\mathcal{E}_{\nu'_i}$) have label -1 . This implies that the above restrictions are sufficient to test whether **UP1** holds for w , and this can be done in $\mathcal{O}(r)$ time for each labeling. Now, assume that pole w is a non-switch vertex of G'_μ . Note that if the shape description for a component $G_{\nu'_i}$ has label at pole w equal to -1 , that is $\lambda(w, \mathcal{E}_{\nu'_i}) = -1$, then both flat angles of u are inside $\mathcal{E}_{\nu'_i}$, while if the label equals 0 that is $\lambda(w, \mathcal{E}_{\nu'_i}) = 0$, then one flat angle of u is inside $\mathcal{E}_{\nu'_i}$. In the first case, we have that $\lambda(\alpha_w(f'_i)) = -1$ must hold for all faces f'_i , and $\lambda(w, \mathcal{E}_{\nu'_j}) = 1$ for all components $G_{\nu'_j}$ different from $G_{\nu'_i}$. In the second case, $G_{\nu'_i}$ is the unique special component of pole w , and $\lambda(w, \mathcal{E}_{\nu'_j}) = 1$ must hold for all other components $G_{\nu'_j}$, while $\lambda(\alpha_w(f'_i)) = 1$ must hold for all faces f'_i , except for one that must have label 0 . In both cases, since all shape descriptions used for the components of G'_μ belong to their feasible sets, the switch angles of w in the interior of all components have label -1 . Hence the above restrictions are sufficient to test whether **UP2** holds for w , and this can be done in $\mathcal{O}(r)$ time for each labeling.

Overall, there exist $\mathcal{O}(r^2)$ possible labelings for the angles $\alpha_u(f'_i)$ and $\alpha_v(f'_i)$ of u and v respectively inside all faces f'_i of \mathcal{E}'_μ . Testing whether **UP1-UP3** holds for each of them requires $\mathcal{O}(r)$ time. If this is the case, by Theorem 1 G'_μ is upward planar and the shape description of \mathcal{E}'_μ can be computed in constant time from the first element of S , its last element, and the labels at u and v chosen for the outer face f'_0 . Thus, the overall complexity of deciding whether a sequence S is a shape sequence for G'_μ is $\mathcal{O}(r^3)$. In the same time all $\mathcal{O}(r^2)$ shape descriptions for G'_μ that correspond to S are computed.

Assume now that S is not reduced. We describe how this case can be brought down to

the previous one. Let s^+ be a maximal subsequence of S . Let $G_{\nu'_j}, G_{\nu'_{j+1}}, \dots, G_{\nu'_{j+\ell}}$ be the components of G'_μ that correspond to the subsequence s^+ and let G_μ^+ be the subgraph of G'_μ consisting of these components. Let S^- be the sequence obtained from S by replacing s^+ with s and let G_μ^- be the graph obtained from G'_μ by replacing G_μ^+ with $G_{\nu'_j}$.

First we check under which conditions s^+ is a shape sequence for G_μ^+ . Let $G_{\nu'_i}$ and $G_{\nu'_{i+1}}$ be two consecutive components of G'_μ that both have shape description s in S . The internal face f'_i of \mathcal{E}'_μ is defined by the right outer path of $\mathcal{E}_{\nu'_i}$ and the left outer path of $\mathcal{E}_{\nu'_{i+1}}$. Let $\alpha_u(f'_i)$ and $\alpha_v(f'_i)$ be the angles at u and v respectively inside f'_i . If S is a shape sequence of G'_μ , **UP3** must be satisfied for f'_i , that is $\tau_r(s) + \tau_l(s) + \lambda(\alpha_u(f'_i)) + \lambda(\alpha_v(f'_i)) = -2$. Recall that by Lemma 9 the right-turn-number of s satisfies $\tau_r(s) = -\tau_l(s) + h$ with $h = 0, 1, 2, 3, 4$. Also the labels $\lambda(\alpha_u(f'_i))$ and $\lambda(\alpha_v(f'_i))$ take values in $\{-1, 0, 1\}$. The previous equation gives $h + \lambda(\alpha_u(f'_i)) + \lambda(\alpha_v(f'_i)) = -2$ which is only satisfied if $h = 0$ and $\lambda(\alpha_u(f'_i)) = \lambda(\alpha_v(f'_i)) = -1$. Hence the shape description $s = \langle \tau_l(s), \tau_r(s), \lambda_u(s), \lambda_v(s), \rho_l(u, s), \rho_r(u, s), \rho_l(v, s), \rho_r(v, s) \rangle$ must satisfy the thin conditions presented in Section 3. Checking whether s satisfies the thin conditions can be done in constant time. Let N_1 be the set $\{\nu'_1, \dots, \nu'_{k'}\}$ of children of μ , $N_s = \{\nu'_j, \dots, \nu'_{j+\ell}\}$ and $N_2 = N_1 \setminus N_s \cup \{\nu'_j\}$. Note that G'_μ and G_μ^- are the graphs G_{N_1} and G_{N_2} of Lemma 11. Also, an upward embedding \mathcal{E}_{N_1} of G_{N_1} with shape sequence S , and an upward embedding \mathcal{E}_{N_2} of G_{N_2} with shape sequence S^- , both respect s for the set N_s , and they form an equivalence pair. Hence, by Lemma 11, if s satisfies the thin conditions, then every shape description for G'_μ that corresponds to S is also a shape description for G_μ^- that corresponds to S^- and vice versa.

We try to replace every maximal subsequence s^+ of S with s . If this is not possible, that is s does not satisfy the thin conditions, then S is not a shape sequence of G'_μ . Otherwise, we have a subgraph G_μ^- of G'_μ and a sequence S^- which is reduced. By using the procedure described at the beginning of the proof, we can compute the shape descriptions of G_μ^- that corresponds to S^- in $\mathcal{O}(r^3)$ time. ◀

Let S' be the shape sequence of G'_μ with respect to \mathcal{E}'_μ . Let ν be a child of G_μ that is not in the set $\{\nu'_1, \nu'_2, \dots, \nu'_{k'}\}$, let s be a shape description from the feasible set of G_ν , and let G''_μ be the subgraph of G_μ consisting of the union of G_ν and G'_μ . We say that S' can be extended with s to a shape sequence S'' of G''_μ if S'' is a shape sequence of G''_μ , s belongs to S'' , and removing s from S'' we obtain S' . Notice that s can be either the first, or the last, or an intermediate element of S'' .


► **Lemma 16.** *Let S' be a shape sequence of G'_μ . Given a shape description s from the feasible set of G_ν , we can decide whether S' can be extended with s to a shape sequence S'' of G''_μ and compute the corresponding shape descriptions for G''_μ in $\mathcal{O}(r^4)$ time, where r is the size of S' .*

Proof. The shape description s can be placed either at the beginning, at the end or between two consecutive elements of S' . In the first two cases, $S'' = [s, S']$ or $S'' = [S', s]$. The shape descriptions that correspond to S'' can be computed using Lemma 15 in $\mathcal{O}(r^3)$ time each. For the last case, let a and b be the two consecutive elements of S' where s is placed. If $a \neq b$ there are at most $r - 1$ possible placements for s , that is, $r - 1$ different sequences S'' . Note that if $s = a$ or $s = b$, then S'' might be the same as S' . On the other hand, if $a = b$ there are at most r such sequences S'' . In fact, arguing similarly with the first part of the proof of Lemma 15, if $s \neq a$ the sequence S'' is not a shape sequence of G''_μ and if $s = a$ then S'' is the same sequence with S' . By Lemma 15, we can test in $\mathcal{O}(r^3)$ time if each produced sequence S'' is a shape sequence of G''_μ or not. The total require time is $\mathcal{O}(r^4)$ as claimed. ◀

Suppose that G_μ is upward planar and consider an upward planar embedding \mathcal{E}_μ of G_μ . We first remove the special components of u and v and the normal components G_ν whose uv -external upward planar embedding \mathcal{E}_ν induced by \mathcal{E}_μ has shape description with $\lambda(\mathcal{E}_\nu, u) = -1$ or $\lambda(\mathcal{E}_\nu, v) = -1$. Denote by G'_μ the subgraph of G_μ obtained after this removal. We say that G'_μ is the *thin subgraph* of G_μ with respect to \mathcal{E}_μ . Denote by \mathcal{E}'_μ the upward planar embedding of G'_μ . Let $\nu'_1, \nu'_2, \dots, \nu'_{k'}$ be the children of μ corresponding to the remaining components, that is, normal components of both u and v where $\lambda(\mathcal{E}_{\nu'_i}, u) = \lambda(\mathcal{E}_{\nu'_i}, v) = 1$ for $i = 1, \dots, k'$. Assume that the pertinent graphs $G_{\nu'_1}, G_{\nu'_2}, \dots, G_{\nu'_{k'}}$ appear in this order from left to right around u , such that the outer boundary of \mathcal{E}'_μ is formed by the left outer path of $\mathcal{E}_{\nu'_1}$ and the right outer path of $\mathcal{E}_{\nu'_{k'}}$. We first present few technical lemmata that will be used later in this section.

► **Lemma 17.** *The thin subgraph of G_μ with respect to an upward planar embedding \mathcal{E}_μ contains k' components of G_μ , where $k - 2 \leq k' \leq k$.*

Proof. As G_μ has k components, the thin subgraph G'_μ contains $k' \leq k$ components of G_μ . In order to prove that $k' \geq k - 2$ we argue that there are most two components of G_μ that don't belong to the thin subgraph G'_μ . Note that if component G_{ν_i} does not belong to G'_μ , then either G_{ν_i} is a special component of u or v , or it is a normal component for both poles u and v , with $\lambda(\mathcal{E}_{\nu_i}, u) = -1$ or $\lambda(\mathcal{E}_{\nu_i}, v) = -1$. We say that G_{ν_i} is *associated* with pole w if G_{ν_i} is either a special component of w or it is a normal component for both poles with $\lambda(\mathcal{E}_{\nu_i}, w) = -1$. Note that if G_{ν_i} is a special component of w then w is a non-switch vertex of G_μ . On the other hand, if it is a normal component of w with $\lambda(\mathcal{E}_{\nu_i}, w) = -1$, then there is a large angle at w inside \mathcal{E}_{ν_i} and w is a switch vertex of G_μ . Assume for a contradiction that there exist at least three components that don't belong to the thin subgraph G'_μ . Then at least two of them are associated with the same pole, say w . If w is a non-switch vertex, then both components are special components of w ; contradiction, since there is at most one special component of w . Hence w is a switch vertex and both components are normal components with a large angle at w in their interior; contradiction to **UP2**, since, in any upward planar embedding, each switch vertex has exactly one large angle. ◀




■ **Figure 6** A graph G_μ that is a parallel composition of seven components; the thin subgraph is shown in gray. The component $G_{\nu_{j1}}$ (shown in orange) is a normal component for u and v with $\lambda(G_{\nu_{j1}}, v) = -1$; the component $G_{\nu_{j2}}$ (shown in green) is a special component of u .

In the next lemma, if $w \in \{u, v\}$ is a top vertex then $\rho_w = out$, otherwise $\rho_w = in$.

► **Lemma 18.** *Let μ be a P -node such that G_μ is upward planar and let \mathcal{E}_μ be an upward planar embedding of G_μ such that the left-turn-number of G'_μ is c . Then the shape sequence*

of G'_μ with respect to \mathcal{E}_μ is $[s_1^+, s_2^*, s_3^*]$, with $s_1 = \langle c, -c, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$, $s_2 = \langle c - 2, -c + 2, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$, $s_3 = \langle c - 4, -c + 4, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$.

Proof. Since all components of G'_μ are normal components of both poles u and v , it follows that u and v are switch vertices of G'_μ . Hence, for every $i = 1, \dots, k'$, $\rho_l(\mathcal{E}_{\nu'_i}, u) = \rho_r(\mathcal{E}_{\nu'_i}, u) = \rho_u$ and $\rho_l(\mathcal{E}_{\nu'_i}, v) = \rho_r(\mathcal{E}_{\nu'_i}, v) = \rho_v$. Additionally, at each pole u and v there is exactly one large angle. By our assumptions $\lambda(\mathcal{E}_{\nu'_i}, u) = \lambda(\mathcal{E}_{\nu'_i}, v) = 1$ for $i = 1, \dots, k'$, hence the large angles at u and v lie inside two (not necessarily distinct) faces of \mathcal{E}'_μ . Let f_i and f_j be the two faces where the large angle at u and v lie in respectively, where $0 \leq i, j < k'$. We distinguish some cases depending on whether $i = 0$ or $j = 0$ holds.



■ **Figure 7** Illustration for Lemma 18. Components with shape description s_1 , s_2 and s_3 are shaded with gray, orange and green respectively.

Case 1: $i = j = 0$. The large angles at u and v are both inside the outer face of \mathcal{E}'_μ , that is, $\lambda(G'_\mu, u) = \lambda(G'_\mu, v) = 1$ (see Fig. 7a). In this case, by Lemma 9, the left and the right turn number of \mathcal{E}'_μ are the same in absolute value, that is $\tau_l(\mathcal{E}'_\mu, u, v) = -\tau_r(\mathcal{E}'_\mu, u, v)$. Since the angles at u and v inside the face f_h is small, it follows from Lemma 10 that $\tau_r(\mathcal{E}_{\nu'_h}, u, v) = -\tau_l(\mathcal{E}_{\nu'_{h+1}}, u, v)$, for each $h = 1, 2, \dots, k - 1$. Also, since the angles at u and v inside the internal faces of $\mathcal{E}_{\nu'_h}$ are small, it follows that $\lambda(\mathcal{E}_{\nu'_h}, u) = \lambda(\mathcal{E}_{\nu'_h}, v) = 1$, which implies by Lemma 9 that $\tau_l(\mathcal{E}_{\nu'_h}, u, v) = -\tau_r(\mathcal{E}_{\nu'_h}, u, v)$. Combining these equalities we have that all \mathcal{E}_{ν_h} (gray components in Fig. 7a) have the same shape description $\langle c, -c, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$, where $c = \tau_l(\mathcal{E}'_\mu, u, v)$. Hence the shape sequence is $[\langle c, -c, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle^+]$.

Case 2: $i = 0$ and $j > 0$. The large angle at u is inside the outer face of \mathcal{E}'_μ while the large angle at v is inside face f'_j , with $1 \leq j < k'$, that is, $\lambda(\mathcal{E}'_\mu, u) = 1$ and $\lambda(\mathcal{E}'_\mu, v) = -1$ (see Fig. 7b). By Lemma 10 we have that $\tau_r(\mathcal{E}_{\nu'_i}, u, v) = -\tau_l(\mathcal{E}_{\nu'_{i+1}}, u, v) - 2$. For all $h \neq j$, the angles at u and v inside the face f'_h are small, hence from Lemma 10 it follows that $\tau_r(\mathcal{E}_{\nu'_h}, u, v) = -\tau_l(\mathcal{E}_{\nu'_{h+1}}, u, v)$. Also, since the angles at u and v inside the internal faces of $\mathcal{E}_{\nu'_h}$ are small, it follows that $\lambda(\mathcal{E}_{\nu'_h}, u) = \lambda(\mathcal{E}_{\nu'_h}, v) = 1$, which implies by Lemma 9 that $\tau_l(\mathcal{E}_{\nu'_h}, u, v) = -\tau_r(\mathcal{E}_{\nu'_h}, u, v)$. Combining these equations, we conclude that all \mathcal{E}_{ν_h} with $h \leq j$ (gray components in Fig. 7b) have the same shape description $\langle c, -c, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$, where $c = \tau_l(\mathcal{E}'_\mu, u, v)$, while all \mathcal{E}_{ν_h} with $h > j$ (orange components in Fig. 7b) have the same, yet different, shape description $\langle c - 2, -c + 2, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$. Then the shape sequence for \mathcal{E}'_μ is $[\langle c, -c, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle^+, \langle c - 2, -c + 2, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle^+]$.

Case 3: $i > 0$ and $j = 0$. The large angle at v is inside the outer face of \mathcal{E}'_μ while the large angle at u is inside face f'_i , with $1 \leq i < k'$, that is, $\lambda(\mathcal{E}'_\mu, u) = -1$ and $\lambda(\mathcal{E}'_\mu, v) = 1$ (see

Fig. 7c). Symmetrically to the previous case, we can prove that all $\mathcal{E}_{\nu'_h}$ with $h \leq i$ (gray components in Fig. 7c) have the same shape description $\langle c, -c, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$, where $c = \tau_l(\mathcal{E}'_\mu, u, v)$, while all $\mathcal{E}_{\nu'_h}$ with $h > i$ (orange components in Fig. 7c) have the same, yet different, shape description $\langle c - 2, -c + 2, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$. As before, the shape sequence for \mathcal{E}'_μ is $[\langle c, -c, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle^+, \langle c - 2, -c + 2, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle^+]$.

Case 4: $i, j > 0$. None of the large angles at u and v is inside the outer face of \mathcal{E}'_μ , that is, $\lambda(\mathcal{E}'_\mu, u) = \lambda(\mathcal{E}'_\mu, v) = -1$. The large angle at u is inside face f'_i , with $1 \leq i < k'$ and the large angle at v is inside face f'_j , with $1 \leq j < k'$. Assume first that $i < j$ holds. For each $h \neq i, j$, we have that the angles at u and v inside the face f'_h and inside the internal faces of $\mathcal{E}_{\nu'_h}$ are small. By Lemmata 10 and 9 it follows that $\tau_r(\mathcal{E}_{\nu'_h}, u, v) = -\tau_l(\mathcal{E}_{\nu'_{h+1}}, u, v)$ and $\tau_l(\mathcal{E}_{\nu'_h}, u, v) = -\tau_r(\mathcal{E}_{\nu'_h}, u, v)$. This means that all $\mathcal{E}_{\nu'_h}$ with $h \leq i$ (gray components in Fig. 7d) have the same shape description $\langle c, -c, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$, where $c = \tau_l(\mathcal{E}'_\mu, u, v)$, all $\mathcal{E}_{\nu'_h}$ with $i < h \leq j$ (orange components in Fig. 7d) have the same shape description $\langle c - 2, -c + 2, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$, and all $\mathcal{E}_{\nu'_h}$ with $j < h$ (green components in Fig. 7d) have the same shape description $\langle c - 4, -c + 4, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$; see Fig. 7d. The shape sequence defined is $[\langle c, -c, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle^+, \langle c - 2, -c + 2, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle^+, \langle c - 4, -c + 4, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle^+]$. Note that the shape sequence is the same when $i > j$, while for $i = j$ it becomes $[\langle c, -c, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle^+, \langle c - 4, -c + 4, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle^+]$; see Fig. 7e. \blacktriangleleft

Based on Lemma 18, our algorithm computes the shape descriptions of all possible upward planar embeddings of G_μ that match some fixed left- and right-turn-numbers in three steps. In the first step, we decide which sequences S' of shape descriptions given in Lemma 18 are shape sequences for some maximal subgraph G'_μ of G_μ . If S' is such a sequence, then, by Lemma 17, there are at most two children whose shape descriptions are not in S' . So, in the second step we compute all possible extensions of S' with the missing shape descriptions to shape sequences of G_μ . In the last step, an extended sequence S of G_μ is modified by appropriately changing the shape descriptions for at most two components of G'_μ in order to match the given left- and right-turn-numbers.

We now describe in detail how to compute the feasible set \mathcal{F}_μ of a P-node μ . For each possible value c_l of $\tau_l(\mathcal{E}_\mu, u, v)$ and c_r of $\tau_r(\mathcal{E}_\mu, u, v)$, that is $c_l, c_r \in [\tau_{min}, \tau_{max}]$, we compute the shape descriptions for all possible upward planar embeddings of G_μ with the given left- and right-turn-numbers. All normal components of G_μ have uv -external upward planar embeddings with left- and right-turn-number odd if both u and v are either top or bottom vertices, and even if one of them is a top vertex and the other one is a bottom vertex. Then any uv -external upward planar embedding of every subgraph G'_μ that contains only normal components has a maximal left-turn-number c'_l that is either equal to c_l or to $c_l - 1$ depending on the parity of both c_l and c'_l . For the first step of the algorithm we consider all sequences of shape descriptions $S' = [s_1^*, s_2^*, s_3^*]$ where $s_1 = \langle c'_l, -c'_l, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$, $s_2 = \langle c'_l - 2, -c'_l + 2, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$, $s_3 = \langle c'_l - 4, -c'_l + 4, 1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$. For each of them we identify a maximal subgraph G'_μ of G_μ such that S' is a shape sequence of G'_μ . To this aim, we check whether it is possible to choose a shape description from the feasible set of the pertinent graph G_{ν_i} of each child ν_i of μ (with $i = 1, 2, \dots, k$) and to sort these pertinent graphs so that the resulting shape sequence is S' . We observe that S' contains at most three different shape descriptions, namely s_1, s_2 and s_3 which can occur multiple times in the sequence. For each child ν_i of μ (with $i = 1, 2, \dots, k$) we check whether the feasible set \mathcal{F}_{ν_i} contains one of the three shape descriptions of S' in the order that they appear in S' ; if so, we choose it for G_{ν_i} . Note that the shape description assigned to a child is the first one among s_1, s_2 and s_3 that is contained in its feasible set. However, this does

not necessarily produce the desired sequence S' for G'_μ , especially if S' has two or three elements. Thus, we check whether it is possible to modify the assigned shape descriptions so as to produce the sequence S' . If S' consists of two elements and all components of G'_μ are assigned to the first one, we check whether one of them can be reassigned the second one. If this is not possible, then no subgraph G'_μ has S' as its shape sequence. If S' consists of three elements, that is $S' = [s_1^+, s_2^+, s_3^+]$, then we first make sure that s_2 has at least one assigned component by reassigning a component from s_1 , as in the previous case. If this is not possible, then no subgraph G'_μ has S' as its shape sequence. Otherwise, if s_3 has at least one assigned component we are done. If not, we check whether a component from s_1 or s_2 can be reassigned to s_3 . If such a component exists, and both s_1 and s_2 have still at least one element, we are done. Otherwise if s_1 is empty, then no subgraph G'_μ has S' as its shape sequence; if s_2 is empty we check whether a component from s_1 can be reassigned to s_2 .

For the second step of the algorithm, assume that for some sequence S' we have identified a maximal subgraph G'_μ of G_μ such that S' is a shape sequence of G'_μ . Consider now those children of μ where no shape description could be assigned for their pertinent graph. Recall that by Lemma 17 there can be at most two such children of μ , say ν and ν' . If this is not true, we conclude that G_μ does not admit an upward drawing whose thin subgraph has S' as its shape sequence. Otherwise, let s_ν ($s_{\nu'}$) be a shape description in the feasible set of ν (ν' , respectively), if it exists. Using Lemma 16 we compute all possible extensions of S' with s_ν to shape sequences of G''_μ , where G''_μ is the subgraph of G_μ consisting of G'_μ and G_ν . If ν' does not exist, then $G''_\mu = G_\mu$ and each computed shape sequence S'' is a shape sequence of G_μ . Otherwise, using the same lemma, for each computed shape sequence S'' , we further compute all possible extensions of S'' with $s_{\nu'}$ to shape sequences of G_μ . For every computed shape sequence S of G_μ we check whether it matches the given left- and right-turn-numbers, that is, the left-turn-number of the first element of S equals c_l and the right-turn-number of its last element equals c_r . If this is the case, then we add all shape descriptions for G_μ that correspond to S to the feasible set \mathcal{F}_μ of μ . Otherwise, we proceed to the third step of the algorithm with S .

For the last step of our algorithm, let S be a shape sequence of G_μ given by the previous step. That is, S is an extension of S' and contains s_ν and $s_{\nu'}$ for children ν and ν' (if they exist) at specific positions, and such that the given left-turn-number and/or the right-turn-numbers are not matched. We consider cases depending on whether only one or both of the two turn-numbers are not matched. Assume first the the first element of S has left-turn-number different from c_l , while the last element of S has right-turn-number equal to c_r . Then the first element has left-turn-number smaller than c_l . Our goal is to find a component $G_{\nu'_i}$ of the thin subgraph G'_μ , remove its current shape description from S and use another one from its feasible set at the beginning of the sequence in order to match the desired left-turn-number. Note that this is possible only if c_l and c'_l have the same parity. So, consider component $G_{\nu'_i}$ of the thin subgraph G'_μ and let $s_i \in \{s_1, s_2, s_3\}$ be the shape description assigned to $G_{\nu'_i}$ in S . Let s'_i be another shape description in the feasible set of $G_{\nu'_i}$ that is different from s_1 , s_2 and s_3 . We denote by S^- the sequence S after removing s_i . Before proceeding, we want to assure that S^- contains the same elements with S . This is not true if $G_{\nu'_i}$ was the only component assigned to s_i during the first step of the algorithm. Note that all components of G'_μ are normal components for both u and v and the shape descriptions in their feasible sets have maximum left-turn-number equal to c'_l . Hence s_1 does not belong to S , that is, the shape sequence of G'_μ is $S' = [s_2^*, s_3^*]$ and $G_{\nu'_i}$ is initially assigned either to s_2 or to s_3 . If $G_{\nu'_i}$ was the only component of G'_μ assigned to s_2 , then s_2 does not belong to the feasible set of the other components of G'_μ and S^- can't

be modified to contain all elements of S' . If $G_{\nu'_i}$ was the only component of G'_μ assigned to s_3 , then we check whether s_2 belongs to the feasible set of the other components of G'_μ . If we can't find such a component, then S^- can't be modified to contain all elements of S' , otherwise, we reassign that component to s_2 and S^- contains the same elements with S . Using Lemma 15, we can compute the shape descriptions of all upward planar embeddings of G_μ that correspond to the sequence $[s'_i, S^-]$. The computed shape descriptions have the desired left- and right-turn-number and are added to the feasible set \mathcal{F}_μ of μ .

For the case where S matches the left-turn-number c_l but the last element of S has right-turn-number different from c_r , we follow a similar approach. In this case, s_3 does not belong to S and for each component $G_{\nu'_i}$ of the thin subgraph G'_μ we first remove its current shape description s_i from S . If the new sequence S^- does not contain all elements of S' , then $s_i = s_2$ and we check if we can reassign another component of G'_μ from s_1 to s_2 . If this is not possible, then S^- can't be modified to contain all elements of S' . Otherwise, by Lemma 15, we can compute all possible shape descriptions for G_μ that correspond to the sequence $[S^-, s'_i]$. The computed shape descriptions match the desired left- and right-turn-number and are added to the feasible set \mathcal{F}_μ of μ .

In the special case where both the left- and right-turn-number of S do not match c_l and c_r respectively, then S does not contain neither s_1 nor s_3 . Hence, the initial sequence for the thin subgraph G'_μ is $S' = [s_2^+]$. Now, the shape sequence of a normal component for both u and v that has left-turn-number equal to c'_l , right-turn-number at most $-c'_l + 2$ and at least one label of the angles at u or v is -1 , is either $t_l^u = \langle c'_l, -c'_l + 2, -1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$ or $t_l^v = \langle c'_l, -c'_l + 2, 1, -1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$ (see Lemma 9). Similarly the shape sequence of such a component with left-turn-number at most $c'_l - 2$ and right-turn-number $-c'_l + 4$ is either $t_r^u = \langle c'_l - 2, -c'_l + 4, -1, 1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$ or $t_r^v = \langle c'_l - 2, -c'_l + 4, 1, -1, \rho_u, \rho_u, \rho_v, \rho_v \rangle$. In order to match both the given left- and right-turn-numbers, S needs to be extended with t_l^u or t_l^v at the beginning, and with t_r^u or t_r^v at the end. As t_l^u and t_r^u both have $\lambda(u) = -1$, and t_l^v and t_r^v both have $\lambda(v) = -1$, the possible extensions of S are $[t_l^u, S, t_r^v]$ and $[t_l^v, S, t_r^u]$. We consider all components $G_{\nu'_i}$ of the thin subgraph G'_μ and check whether t_l^u , t_l^v , t_r^u and t_r^v are contained in their feasible set. Let T_l^u be the set of components $G_{\nu'_i}$ that contain t_l^u in their feasible set; sets T_l^v , T_r^u and T_r^v are defined similarly. For the sequence $[t_l^u, S, t_r^v]$ we want to select one component from T_l^u and a different one from T_r^v and reassign their shape descriptions from s_2 to t_l^u and t_r^v respectively. This is not possible only if either one of T_l^u or T_r^v is empty, or if T_l^u is the same as T_r^v and they both contain one element. If this is the case, S can't be extended to $[t_l^u, S, t_r^v]$, otherwise, we use Lemma 15 to decide whether this is a shape sequence of G_μ and to compute the corresponding shape descriptions for \mathcal{F}_μ . Similarly, S can't be extended to the second sequence $[t_l^v, S, t_r^u]$ if either one of T_l^v and T_r^u is empty or if they both have size one and contain the same element, otherwise the shape descriptions for \mathcal{F}_μ that correspond to $[t_l^v, S, t_r^u]$ are computed with Lemma 15 and added to the feasible set \mathcal{F}_μ of μ .

► **Lemma 19.** *Let μ be an P -node of T . The described algorithm computes correctly the feasible set \mathcal{F}_μ of μ .*

Proof. Our algorithm computes sequences of shape descriptions for G_μ based on the feasible sets of the pertinent graphs of the children of μ . For each computed sequence Lemma 15 tests whether it is a shape sequence of G_μ and the corresponding shape descriptions are added in the feasible set \mathcal{F}_μ of μ . Hence, whenever a shape description s is computed for a sequence S , there exists an upward planar embedding \mathcal{E}_μ of G_μ with shape sequence S and whose shape description is s .

It remains to prove that if s_μ is a shape description of G_μ , then our algorithm finds a shape sequence S such that s_μ corresponds to S . Let \mathcal{E}_μ be an upward planar embedding of G_μ with shape sequence S_μ and such that the shape description of G_μ is s_μ . Let $G'_{\mathcal{E}_\mu}$ be the thin subgraph of G_μ with respect to \mathcal{E}_μ and let S'_μ be its shape sequence.

For the left- and right-turn-numbers of s_μ , the first step of the algorithm computes a maximal thin subgraph G'_μ for S'_μ assigning each child of G_μ to the first element of S'_μ that is contained in its feasible set. If an element of S'_μ is not assigned, the algorithm tries to reassign some components of G'_μ . This is not possible only if an element of S'_μ is not contained in the feasible sets of all components of G_μ . Note that the children of μ whose pertinent graphs belong to the thin subgraph $G'_{\mathcal{E}_\mu}$ contain at least one element of S'_μ in their feasible set. Hence, the algorithm will successfully compute a maximal thin subgraph G'_μ for S'_μ . Additionally, the computed maximal thin subgraph G'_μ contains all components of $G'_{\mathcal{E}_\mu}$, although they may be assigned to different elements of S'_μ .

If $G'_{\mathcal{E}_\mu} = G_\mu$, that is $S_\mu = S'_\mu$, then all components of G_μ are assigned to elements of S'_μ , the second step can't extend S'_μ , and the left- and right-turn-numbers of S'_μ match the given values. Hence, during the third step of the algorithm all shape descriptions for G_μ that correspond to S'_μ will be computed, including s_μ . So, assume that there is at least one component of G_μ , say G_ν , whose shape description s_ν in S_μ is not an element of S'_μ . If G_ν is a special component of one of the poles, or a normal component for both poles whose feasible set contains no element of S'_μ , then the second step of the algorithm tests whether S'_μ can be extended with all shape descriptions in the feasible set of G_ν , including s_ν . Since S_μ is a shape sequence of G_μ , the algorithm will extend S'_μ with s_ν to all possible shape sequences of the subgraph consisting of G'_μ and G_ν . In particular the sequence where s_ν is placed between the same elements of S'_μ as in S_μ , will be one of the computed shape sequences. If there is another component $G_{\nu'}$ that is either a special component of one of the poles, or a normal component for both poles whose feasible set contains no element of S'_μ , then S'_μ will be further extended to contain the shape description $s_{\nu'}$ of $G_{\nu'}$ at the correct position. At the end of the second step, either the computed shape sequence S is the same with S_μ , or all elements of S appear in S_μ in the same order from left to right.

If S has the same left- and right-turn-numbers with S_μ , then the set of shape descriptions that correspond to S will include s_μ , since all elements of S are contained in S'_μ . So, we examine the case where the left- and/or the right-turn-numbers of S do not match those of S_μ . Assume first that the left-turn-number is different from S_μ , while the right-turn-number is the same. Let s_1 be the first element of S_μ that corresponds to component G_{ν_1} of G_μ with respect to \mathcal{E}_μ . Then, G_{ν_1} is a component $G_{\nu'_1}$ of the maximal thin subgraph G'_μ computed in the first step, and its shape description in S belongs to S'_μ . During the third step of the algorithm, it will be tested whether the sequence $[s_1, S]$ is a shape sequence for G_μ . Note that since all elements of $[s_1, S]$ belong to S_μ in the same order, and the left- and right-turn-numbers of the two sequences are the same, the shape description s_μ of G_μ corresponds to both S_μ and $[s_1, S]$ and will be therefore computed by the algorithm. Similarly, if S_μ and S have the same left-turn-number and different right-turn-number, the third step of the algorithm will identify s_μ as a shape description corresponding to the sequence $[S, s_k]$, where s_k is the last element of S_μ and it is the shape description used for the last component G_{ν_k} of G_μ with respect to \mathcal{E}_μ . Hence it remains to consider the case where S and S_μ have different left- and right-turn-numbers. In this case S_μ is one of $[t_l^u, S'_\mu, t_r^v]$ or $[t_l^v, S'_\mu, t_r^u]$, and $S'_\mu = [s_2^+]$. The last part of the algorithm tests whether it is possible to reassign two components of G_μ from s_2 , one to t_l^u and the other to t_r^v for the first sequence, or one to t_l^v and the other to t_r^u for the second one. Hence S_μ will be successfully identified as a shape sequence of G_μ ,

concluding the lemma. ◀

► **Lemma 20.** *Let μ be an P -node of T . The feasible set \mathcal{F}_μ can be computed in $\mathcal{O}(\tau \cdot k)$ time, where k is the number of children of μ .*

Proof. We prove that the time complexity of the algorithm described above is $\mathcal{O}(\tau \cdot k)$. The number of choices for the left-turn-number c_l of a uv -external upward planar embedding of G_μ is τ , and for each value of c_l by Lemma 9 the right-turn-number c_r can only take values $-c_l + h$ for $h = 0, 1, 2, 3, 4$. Hence there are 5τ possible pairs for c_l and c_r . For each pair, we compute the shape descriptions of all possible uv -external upward planar embeddings of G_μ with left-turn-number equal to c_l and right-turn-number equal to c_r . For given c_l and c_r , first we retrieve the feasible sets of the children of μ . For each feasible set we are only interested in those shape descriptions that have left-turn-number in $[c_l - 4, c_l]$ and right-turn-number in $[c_r - 4, c_r]$. Let $|\mathcal{F}|$ denote the size of the largest feasible set under these constraints. By Lemma 12 the shape descriptions with specific left- and right-turn-numbers are at most six and can be fetched in $\mathcal{O}(1)$ time. Hence $|\mathcal{F}|$ is a constant and all feasible sets can be computed in $\mathcal{O}(k|\mathcal{F}|)$ time.

For the first step of the algorithm, we consider all possible sequences $S' = [s_1^*, s_2^*, s_3^*]$. There are eight such sequences depending on whether each of s_1, s_2 and s_3 belongs to the sequence or not. Let S' be one of them. For $i = 1, \dots, k$ we check if \mathcal{F}_{ν_i} contains one of the at most three elements of S' ; by Lemma 12 each check takes $\mathcal{O}(1)$ time, and we store which nodes among ν_i are assigned to s_1, s_2 and s_3 (if they belong to S'). This step takes $\mathcal{O}(k)$ time and the maximal thin subgraph G'_μ is identified. If an element of S' has no nodes assigned to it, at most two components of G'_μ are reassigned. Each reassignment process takes $\mathcal{O}(k)$ time, since a specific shape description is looked for in the feasible sets of the components of G'_μ . Hence the maximal thin subgraph G'_μ with shape sequence S' is identified in $\mathcal{O}(k)$ time or the algorithm concludes that there is no subgraph of G_μ with shape sequence S' .

The second step considers children ν and ν' of μ (if they exist). For each $s_\nu \in \mathcal{F}_\nu$ and $s_{\nu'} \in \mathcal{F}_{\nu'}$, shape sequence S' is extended in all possible ways with s_ν and $s_{\nu'}$ to a shape sequence S for G_μ . Each extension uses Lemma 16 and requires $\mathcal{O}(1)$ time, since S' has size at most three. Hence, computing all possible sequences S for G_μ that extend a specific shape sequence S' of the first step can be done in $\mathcal{O}(|\mathcal{F}|^2)$ time.

Checking whether the left- or right-turn-number of a sequence is equal to a given number takes $\mathcal{O}(1)$ time. If both the left- and right-turn-numbers are matched by S , the shape descriptions that correspond to S are computed with Lemma 15 in $\mathcal{O}(1)$ time (as S has size at most five). Otherwise the algorithm proceeds with the last step. We consider two cases depending on whether one of c_l and c_r are not matched or both. Consider first the case where one of them is not matched. Assume that the first element of S has left-turn-number different from c_l and the right-turn-number equals c_r ; the case where S matches c_l and not c_r is similar. There are at most k choices for selecting component $G_{\nu'_i}$. Computing the sequence S^- takes constant time since its size is at most five. For at most one component, say $G_{\nu'_j}$, S^- might not contain all elements of S' ; for this component only, we reassign in $(k' - 1)\mathcal{O}(|\mathcal{F}|)$ time another component from s_2 to s_3 . We check whether all elements of S' have at least one node assigned to them in $\mathcal{O}(1)$ time. If this is the case, then the new sequence S^- contains all elements of S' . For each shape description s'_i of the feasible set of $G_{\nu'_i}$ with left-turn-number equal to c_l we compute all possible shape descriptions for G_μ that correspond to the sequence $[s'_i, S^-]$ in $\mathcal{O}(1)$ time by Lemma 15. As the computed shape descriptions match the desired left- and right-turn-number, they are added to the feasible set \mathcal{F}_μ of μ in $\mathcal{O}(1)$ time each. The process for a component $G_{\nu'_i}$ requires $\mathcal{O}(k|\mathcal{F}|^2)$ time if the

sequence S^- does not contain all elements of S' , and $\mathcal{O}(|\mathcal{F}|)$ time otherwise. Since this can occur for at most one component of G'_μ , the third step runs in $\mathcal{O}(k|\mathcal{F}|^2)$ time, for the case where one of c_l or c_r are not matched.

In the special case where both the left- and right-turn-number of S do not match c_l and c_r respectively, then the four sets T_l^u, T_l^v, T_r^u and T_r^v are computed, by checking whether t_l^u, t_l^v, t_r^u or t_r^v respectively belong to the feasible set of all components of G'_μ . By Lemma 12 each check takes $\mathcal{O}(1)$ time, hence the sets are computed in $\mathcal{O}(k)$ time. There are two possible sequences for G_μ , namely $[t_l^u, S, t_r^v]$ and $[t_l^v, S, t_r^u]$. For each of them, we check if any of the corresponding sets is empty, or if they have both size one and contain the same element. If this is the case S can't be extended to match the given left- and right-turn-numbers, otherwise, by Lemma 15 the corresponding shape descriptions for \mathcal{F}_μ can be computed in $\mathcal{O}(1)$ time; each shape description is added to \mathcal{F}_μ in $\mathcal{O}(1)$ time. Hence, if S matches none of c_l and c_r the last step of the algorithm requires $\mathcal{O}(k)$ time.

The overall time complexity is $5\tau(\mathcal{O}(k|\mathcal{F}|) + 8 \cdot (\mathcal{O}(k) + \mathcal{O}(|\mathcal{F}|^2) \cdot (\mathcal{O}(k|\mathcal{F}|^2) + \mathcal{O}(k))))$, where $|\mathcal{F}| = \mathcal{O}(1)$, that is $\mathcal{O}(\tau \cdot k)$, as claimed. \blacktriangleleft

R-node

While for S-, P- and Q-nodes we were able to provide polynomial-time procedures to compute the feasible sets, the NP-hardness of UPWARD PLANARITY prevents the existence of such a polynomial-time procedure for R-nodes. Indeed, obtaining a suitable algorithm that can compute the feasible sets for an R-node given the feasible sets for all of its children will be the task of our parameterized algorithms presented in Sections 6 and 7. In order to complete the description of our framework and also present it in a modular fashion which can be reused by future work, here we formalize the notion of an *R-node subprocedure* which, intuitively, is an algorithm that can compute a feasible set of an R-node.

Formally, an *R-node subprocedure* is an algorithm which takes as input an R-node μ of T with skeleton H , a mapping \mathcal{S}_μ which assigns each child of μ to its feasible set, runs in time at most $\alpha(G_\mu, \mathcal{S}_\mu)$, and computes the feasible set \mathcal{F}_μ . For a digraph G with SPQR-tree T , let $\alpha(G)$ be $\sum_{\text{R-node } \mu} \alpha(G_\mu, \mathcal{S}_\mu)$. We say that $\alpha(G)$ is the *total time complexity* of the R-node subprocedure for graph G .

Root node

We now focus on the root node of the SPQR-tree T . The root node r corresponds to an edge $e = (u, v)$ of G that lies on its outer face. Node r has only one child node μ with poles u and v . Let s be a shape description in the feasible set \mathcal{F}_μ of μ . Then, there exists a uv -external upward planar embedding \mathcal{E}_μ of the pertinent graph G_μ of node μ , such that G_μ has shape description s , and poles u and v are on the outer face of G_μ . Note that $G_\mu = G \setminus e$. Hence, r can be treated as a P-node with poles u and v and two children; one of them is μ and the other one is a Q-node, say μ' for the edge (u, v) . Then the feasible set of μ' contains only one shape description, namely $s' = \langle 0, 0, 1, 1, \text{out}, \text{out}, \text{in}, \text{in} \rangle$. By Lemma 20 we can compute in $\mathcal{O}(\tau)$ time the feasible set of r . Since r has only two children, any shape sequence S of r contains s' as its first or last element. This implies that for any upward planar embedding \mathcal{E} of G_r with shape sequence S , either the left or the right outer path of $G_{\mu'}$ belongs to the outer face of G_r with respect to \mathcal{E} . As $G_{\mu'}$ consists only of edge (u, v) , it follows that (u, v) belongs to the outer face of G in all possible upward planar embeddings.

► **Lemma 21.** *The feasible set \mathcal{F}_r of the root node r of T can be computed in $\mathcal{O}(\tau)$ time.*

Time complexity

From the discussion above and in particular from Lemmata 13, 14, 20 and 21, we obtain the following lemma.

► **Lemma 22.** *Let G be a biconnected DAG with n vertices and let T be an SPQR-tree of G rooted at a Q -node corresponding to an edge $e = (u, v)$. Let τ_{\min} and τ_{\max} be two given integer values, and let $\tau = \tau_{\max} - \tau_{\min} + 1$. Given an R -node subprocedure with total time complexity $\alpha(G)$, it is possible to compute in time $\mathcal{O}(\alpha(G) + \tau^2 \cdot n)$ the shape descriptions of every upward planar embedding with e on the outer face, such that the left- and right-turn-numbers of the pertinent graph of every node of T are in the range $[\tau_{\min}, \tau_{\max}]$.*

5 Extension to the Single-Connected Case

In this section, we show that Lemma 22 can be used as a blackbox to extend our results from the biconnected case to arbitrary graphs. Our goal is to prove Interface Lemma 24, which reduces the task of solving UPWARD PLANARITY to the one of obtaining an R -node subprocedure, for *all* graphs including single-connected ones. First, in the next definition we encapsulate the property that the shapes occurring in SPQR-tree decompositions of a graph have a bounded range of left- and right-turn-numbers, as in Sections 6 and 7 we obtain efficient algorithms precisely in this regime.

► **Definition 23.** *Let G be a digraph and τ_{\min}, τ_{\max} be two integers. We call G $[\tau_{\min}, \tau_{\max}]$ -turn-bounded if the following holds for every upward planar embedding \mathcal{E} of G . For any biconnected component B of G and every SPQR-tree decomposition T of B that is rooted at a Q -node corresponding to an edge that lies on the outer face of B in the embedding induced by \mathcal{E} , left- and right-turn-numbers of the pertinent graph of every node of T are in range $[\tau_{\min}, \tau_{\max}]$.*

We now give the precise statement of our Interface Lemma.

► **Lemma 24 (Interface Lemma).** *Let G be an n -vertex digraph, and τ_{\min}, τ_{\max} be integers such that G is $[\tau_{\min}, \tau_{\max}]$ -turn-bounded. Given an R -node subprocedure with total time complexity $\alpha(G)$, it is possible to determine whether G admits an upward planar embedding in time $\mathcal{O}(n(\alpha(G) + \tau^2 \cdot n))$ where $\tau = \tau_{\max} - \tau_{\min} + 1$.*

Note that, for a single-connected graph G , we define the total time complexity $\alpha(G)$ of an R -node subprocedure to be the sum of $\alpha(B)$ over all biconnected components B of G .

To give an intuition of the proof, we start by guessing the root of the block-cut tree of G , which is assumed to see the outer face in the desired upward planar embedding of G . The core of the proof is the following lemma, which states that leaf components can always be added on top of an upward planar embedding, up to simple conditions on the cut-vertex.

► **Lemma 25.** *Consider a rooted block-cut tree of G , its cut vertex v that is adjacent to leaf blocks B_1, \dots, B_ℓ , and the parent block P . Denote by G^P the corresponding component of $G - \{v\}$. Any upward planar embedding of G^P where the root block sees the outer face can be extended to an embedding of G with the same property if the following holds:*

1. *Each B_i has an upward-planar embedding with v on the outer face.*
2. *If v is a non-switch vertex in P , each B_i has an upward-planar embedding with v on the outer face where the large angle of v is on the outer face.*


As the conditions of Lemma 25 turn out to be necessary as well, our algorithm simply proceeds upwards along the block-cut tree, checking every time the conditions of the lemma and removing the respective leaf components. If at some point the conditions are not satisfied, we conclude that G is a no-instance; otherwise, any upward planar embedding of the root block can be iteratively extended to an embedding of the whole graph. Finally, finding upward planar embeddings of the blocks is performed via Lemma 22; it can also be used to verify conditions of Lemma 25.

Proof of Lemma 25. Fix an upward planar embedding \mathcal{E}^P of G^P , and upward planar embeddings $\mathcal{E}^{B_1}, \dots, \mathcal{E}^{B_\ell}$ of B_1, \dots, B_ℓ subject to the conditions of the lemma. W.l.o.g. assume that the leaf blocks are indexed in such a way that for the first p blocks, the angle of v at the outer face is small or flat, for some $0 \leq p \leq \ell$, and for the remaining blocks the outer angle at v is large. If $p > 0$, it follows from the conditions of the lemma that v is a switch vertex in P and in all the blocks B_1, \dots, B_ℓ except maybe one; if there is a block where v is a non-switch vertex w.l.o.g. let that block be B_p .

We first consequently deal with the blocks B_1, \dots, B_p , that is, we construct an upward planar embedding \mathcal{E}_i of $G_i = G[V(G_p) \cup \bigcup_{j=1}^i V(B_j)]$ for each $i \in [p]$ from an upward planar embedding \mathcal{E}_{i-1} of G_{i-1} , starting with the embedding $\mathcal{E}_0 = \mathcal{E}^P$ of $G_0 = G^P$. For each vertex of G_i that is not v , we infer both the circular order of the incident edges and the angle assignment from \mathcal{E}_{i-1} or \mathcal{E}^{B_i} , depending on whether the vertex belongs to B_i or not. To define a circular order on edges incident to v , split the ordering on neighbors of v in G_{i-1} at a large angle of v (according to \mathcal{E}_{i-1}), which necessarily exists as v is a switch vertex of G_{i-1} , and join it with the ordering of neighbors of v in B_i split at the outer angle of v in \mathcal{E}^{B_i} . This results in two new angles, for each of them, we assign a value of -1 if it is a switch angle, and 0 if it is a flat angle. All the other angles around v keep their value from either \mathcal{E}_{i-1} or \mathcal{E}^{B_i} . We now verify that the conditions **UP0–UP3** from Theorem 1 hold for the vertex v and the new face of the constructed embedding. **UP0–UP2** hold by construction, next we verify **UP3** by considering two cases. In what follows, f denotes the newly formed face. We assume that **UP3** is already verified for \mathcal{E}_{i-1} .

Case 1. The large angle at v is inside the outer face of \mathcal{E}_{i-1} . This necessarily implies $i = 1$, as each B_i for $i \in [p]$ is embedded with a small angle at v on the outer face. Since we embed B_1 in the outer angle of v , the newly formed face is the outer face. Observe that the set of angles at f is precisely the union of the angles of G_0 and the angles of B_1 on the outer faces in the respective embeddings, except for the angles at v . The angle at v on the outer face is necessarily 1 in \mathcal{E}_0 , and there are two cases for the value in \mathcal{E}^{B_1} . If the corresponding angle is a switch angle, then the value is -1 , and the two newly formed angles at v in f by construction receive the value -1 (see Fig. 8a). If the angle is flat in B_1 , its value is 0 and one of the newly formed angles is also flat, receiving a value of 0 , while the other is a switch angle, receiving a value of -1 (see Fig. 8b). In either case, the sum of angle values on f at v is two less than the sum of two outer angles at v in G_0 and B_1 . Thus, summing up all angle values along f we obtain two (sum of angles on the outer face of G_0) plus two (sum of angles on the outer face of B_1 minus two (difference in the angles at v), which satisfies **UP3** since f is the outer face.

Case 2. The large angle at v is inside an inner face of \mathcal{E}_{i-1} . In this case, f is also an inner face in \mathcal{E}_i . Analogously to Case 1, summing up the angles on f in G_i is equivalent to summing angles on the corresponding faces in G_{i-1} and B_i accounting for the change of angles at v . By the same arguments, the change at v results in minus two, while the sum in G_{i-1} is minus two (inner face) and the sum in B_i is two (outer face) by **UP3** in \mathcal{E}_{i-1} and \mathcal{E}^{B_i} , respectively. This sums up to minus two, fulfilling **UP3** for f in \mathcal{E}_i .



■ **Figure 8** Illustrations for the proof of Lemma 25.

It remains to extend the resulting upward planar embedding of G_p by accounting for the block B_{p+1}, \dots, B_ℓ for which there is an upward planar embedding with a large angle at v on the outer face. Again, we consequently extend the embedding one block at a time, constructing an upward planar embedding \mathcal{E}_i of $G_i = G[V(G_p) \cup \bigcup_{j=1}^i V(B_j)]$ for each $i \in [p+1..\ell]$. Fix B_i , v is a switch vertex in B_i since it has an adjacent large angle. We first define the embedding on every vertex except v , by repeating the respective circular order and angle assignment from \mathcal{E}_{i-1} or \mathcal{E}^{B_i} . Now, consider a flat or large angle α at v in G_{i-1} as given by \mathcal{E}_{i-1} , at least one necessarily exists by Theorem 1. Split the neighbor list of v in G_{i-1} at α , split the neighbor list of v in B_i at the large angle, then join these lists to obtain a circular order of neighbors of v in G_i . Consider the two newly formed angles. If α is flat, one of them is flat and the other is a switch angle, assign 0 to the former and -1 to the latter (see Fig. 8c). If α is a large angle, either the edges at v are oriented differently in G_{i-1} and B_i , in which case both new angles are flat (see Fig. 8d), or v is a switch vertex in G_i , then assign 1 to one of the newly formed edges and -1 to the other (arbitrarily) (see Fig. 8e). By construction, the conditions **UP0**–**UP2** of Theorem 1 are satisfied at v , it remains to verify **UP3** for the newly formed face f . Analogously to the case $i \leq p$, the sum of angles on f can be computed as the sum of three parts: the sum of angles in G_{i-1} and B_i on the corresponding faces, and the change in the angle values at v . The first part is two or minus two, depending on whether the large angle at v is on the outer face of G_{i-1} or not, but it matches exactly the target value for f . The second part is always two since v is on the outer face of B_i . It can be easily seen that the third part is exactly minus two, in each of the three cases above. Thus, **UP3** is satisfied for f , and \mathcal{E}_i is a required upward planar embedding. Setting $i = \ell$ finishes the proof. ◀

By proceeding upwards along the block-cut tree of G , Lemma 24 follows shortly from Lemma 25.

Proof of Lemma 24. The algorithm proceeds as follows. First, compute the block-cut tree T of G . Then, branch on the choice of the root block B^R of T . The algorithm now verifies whether there is an upward planar embedding of G where at least one edge of B^R is incident to the outer face. To this end, the algorithm iteratively checks the conditions of Lemma 25 in the bottom-up fashion. Specifically, the algorithm picks a cut vertex v in T , such that all its adjacent blocks are leaves in T , except for the parent block P . For each of the leaf blocks B_1, \dots, B_ℓ , the algorithm uses Lemma 22 to compute possible shape descriptions for every upward planar drawing where v lies on the outer face. That is, for every edge e incident to v in B_i , the algorithm constructs an SPQR-tree decomposition of B_i rooted at a Q-node corresponding to e , and runs Lemma 22 on this decomposition.

If some of the blocks does not admit any feasible shape at this stage, the algorithm

immediately concludes that G is a no-instance. Additionally, if v is a non-switch vertex in P , the algorithm checks whether for each $i \in [\ell]$ there exists a feasible shape description which assigns a large angle to the outer face at v . Since for each $e = (v, u) \in E(B_i)$, we have computed the shape descriptions of every upward planar embedding of B_i with edge e on the outer face via Lemma 22, this information is readily available to the algorithm. If for some component the condition does not hold, the algorithm finishes with reporting that G is a no-instance. Otherwise, the algorithm removes the blocks B_1, \dots, B_ℓ from the graph and proceeds. The algorithm continues in this fashion until only the root block B^R is left, where it finally uses Lemma 22 to check whether B^R admits an upward planar drawing. That is, for every edge $e \in E(B^R)$, the algorithm constructs an SPQR-tree decomposition of B^R rooted at e and runs Lemma 22 on this decomposition. If for each e this results in an empty set of feasible shapes, the algorithm concludes that G is a no-instance. Otherwise, the algorithm returns that G is a yes-instance.

Correctness. First, assume that the algorithm concluded that G admits an upward planar embedding. Then, there is at least one feasible shape description of B^R returned by the algorithm of Lemma 22, which means that there is an upward planar embedding of B^R . By going through the sequence of considered cut vertices v in the reverse order, the embedding can be extended to all children blocks containing v by Lemma 25. Specifically, fix a cut vertex v that splits the graph into the leaf blocks B_1, \dots, B_ℓ and the subgraph G^P that contains the parent block, and assume that the existence of an upward planar embedding for G^P is already shown. Now, by Lemma 25 this embedding extends to an upward planar embedding of $G[V(G^P) \cup V(B_1) \cup \dots \cup V(B_\ell)]$. When the whole block-cut tree T is traversed back in this way, we obtain an upward planar embedding of G .

In the other direction, we argue that if G has an upward planar embedding \mathcal{E}^* , the algorithm necessarily returns one as well (not necessarily the same). In \mathcal{E}^* , there is an edge e that is incident to the outer face, consider the block B^R that contains e and the branch of the algorithm where B^R is set to be the root block of the block-cut tree T of G . Since G admits an upward planar embedding, each block B in T admits one as well. Here we assume that a particular upward planar drawing of G is fixed as well, and an upward planar drawing for its subgraph is induced naturally by removing the respective part of the drawing of G . We now show that no step of the algorithm can result with reporting a no-instance, this immediately implies that the algorithm returns an upward planar embedding of G .

Fix a graph G' that is obtained from G by removing a sequence of leaf blocks (as during the work of our algorithm), \mathcal{E}^* induces an upward planar embedding of G' as well. If G' consists of a single biconnected block, our algorithm successfully decides whether there exists an upward planar embedding of this block by Lemma 22. Otherwise, consider a cut-vertex v with adjacent children leaf blocks B_1, \dots, B_ℓ and the parent block P , denote the component of $G' - v$ that contains P by G_P . For each B_1, \dots, B_ℓ , G_P fix an upward planar embedding that is induced by \mathcal{E}^* . Since G_P contains B^R , there is an edge e in G_P that is incident to the outer face. This immediately implies that for each B_i , v is on the outer face in its embedding — otherwise the whole G_P together with e lies in an inner face of an embedding of B_i , thus contradicting the choice of \mathcal{E}^* and e . This verifies the first condition of Lemma 25, we now move to the second condition. Let v be a non-switch vertex in P , then by the expansion procedure v is a switch vertex in all B_1, \dots, B_ℓ . Assume there is a block B_i where v has a small angle on the outer face, in its fixed embedding. Then by the condition UP1 of Theorem 1, v has a large angle in one of the inner faces of B_i . However, since by the discussion above P is embedded in the outer face of B_i , the flat angles at v will end up being inside the small angle on the outer face of B_i . This contradicts the fact that \mathcal{E}^* is an

upward planar embedding. Thus, at every intermediate step of the algorithm Lemma 25 is invoked successfully.

Running time. For a fixed rooting of the block-cut tree T , on each non-root block B our algorithm runs the algorithm of Lemma 22 on B for each edge incident to v , where v is the parent cut-vertex of this block. Additionally, on the root block B^R Lemma 22 is invoked once for every edge of B^R . Thus, over all possible rootings of T , Lemma 22 is run once on each block per edge of the block, and once for each edge incident to a cut vertex. The latter holds since if our algorithm queries Lemma 22 several times with the same block and the same root edge e , there is no need to recompute the set of admissible shape descriptions as it only depends on the block itself. Therefore, the running time is dominated by invocations of Lemma 22, where the number of invocations on a block is proportional to the number of vertices of the block, and that takes time

$$\sum_{B \in T} \mathcal{O}(|B|(\alpha(B) + \tau^2 \cdot |B|)),$$

Since the blocks of T intersect only in cut-vertices, and the amount of intersections is linear in n , the running time can be upper-bounded by $n(\alpha(G) + \tau^2 \cdot n)$, as required (here we also use that by definition $\alpha(B) \leq \alpha(G)$ for each block B of G). Every other part of the algorithm (e.g. constructing the block-cut tree) takes linear time. ◀

6 An Algorithm Parameterized by the Number of Sources

Let G be an n -vertex digraph whose underlying graph is planar and let σ be the number of sources of G . First, we expand G . As discussed in Section 2, this operation does not alter whether G is upward planar, it preserves the number of sources and at most doubles the number of vertices of G . In view of Lemma 24, in order to obtain an algorithm for UPWARD PLANARITY parameterized by σ , it suffices to devise an R-node subprocedure whose runtime depends on σ and, polynomially, on n . We hence consider a single biconnected component B of G and observe the following.

► **Observation 26.** *The digraph B is expanded and it has at most σ sources.*

Proof. First, B is expanded since G is expanded and since each vertex of B has a set of incident edges which is a subset of the set of edges incident to the same vertex in G .

We prove that B has at most σ sources by induction on the number b_G of biconnected components of G . Indeed, if $b_G = 1$, then $B = G$ and the statement is trivial. If $b_G > 1$, remove from G a biconnected component $B' \neq B$ that has a single cut-vertex c ; let G' be the resulting graph. We prove that G' has at most σ sources. This is obvious if c is a source of G or if c is not a source of G' . Otherwise, suppose that c is not a source of G and it is a source of G' . The two things imply that c is not a source of B' ; since G is acyclic, it follows that B' contains a source $s \neq c$ of B' ; note that s is also a source of G . Thus, although G' contains the “new” source c , it does not contain the source s , hence G' has at most σ sources. Since G' has one less biconnected component than G , the inductive hypothesis applies and we can conclude that B has σ sources. ◀

For the sake of familiarity of notation, in the following we denote by G the considered biconnected digraph, by n the number of its vertices, and by σ the number of its sources. Let e^* be any edge of G ; we compute an SPQR-tree T of G rooted at the Q-node representing e^* in $\mathcal{O}(n)$ time [14, 27]. We will show how to compute, for each R-node μ of T , the feasible set \mathcal{F}_μ , assuming that the feasible sets for the children of μ have been already computed. However, before doing that, we need to establish some combinatorial lemmata.

6.1 Turn-Numbers and Size of Feasible Sets

We start by proving that a bounded number of sources implies bounded left- and right-turn-numbers. Let μ be any node of T , let u and v be the poles of μ , let σ_μ be the number of sources of G_μ different from its poles, and let \mathcal{E}_μ be any uv -external upward planar embedding of G_μ . We have the following.

► **Lemma 27.** *The left- and right-turn-numbers of \mathcal{E}_μ are in the interval $[-2\sigma_\mu - 1, 2\sigma_\mu + 1]$.*

Proof. Let $\langle v_0 = u, v_1, v_2, \dots, v_k = v \rangle$ and $\langle w_0 = u, w_1, w_2, \dots, w_h = v \rangle$ be the left and right outer paths of \mathcal{E}_μ , respectively. For $i = 0, 1, \dots, k$, denote by α_i the angle at v_i inside the outer face of \mathcal{E}_μ and, for $i = 0, 1, \dots, h$, by β_i the angle at w_i inside the outer face of \mathcal{E}_μ .

We first prove that the left-turn-number of \mathcal{E}_μ is at most $2\sigma_\mu + 1$. This is done by proving, for every $m = 1, \dots, k - 1$, the following claim. Let σ_m be the number of sources of G_μ among v_1, \dots, v_m and let $p_m \leq m$ be the largest index such that α_{p_m} is a large angle, where p_m is undefined if each angle among $\alpha_1, \dots, \alpha_m$ is small or flat. Note that, if p_m is defined, then v_{p_m} is either a source or a sink of G_μ , by properties **UP1-UP2** of Theorem 1. Then the claim is that:

- $\sum_{i=1}^m \lambda(\alpha_i)$ is smaller than or equal to $2\sigma_m + 1$, if the index p_m is defined and v_{p_m} is a sink; and
- $\sum_{i=1}^m \lambda(\alpha_i)$ is smaller than or equal to $2\sigma_m$, if the index p_m is undefined or v_{p_m} is a source.

Observe that the claim, applied with $m = k - 1$, implies that the left-turn-number of \mathcal{E}_μ is at most $2\sigma_\mu + 1$, since $\sigma_{k-1} \leq \sigma_\mu$.

The proof is by induction on m . In the base case $m = 1$. We distinguish some cases, according to the type of α_1 .

- If α_1 is a non-large angle, then $\lambda(\alpha_1) \leq 0 = 2\sigma_1$.
- If α_1 is a large angle and v_1 is a source of G_μ , then $\lambda(\alpha_1) = 1 < 2 = 2\sigma_1$.
- If α_1 is a large angle and v_1 is a sink of G_μ , then $\lambda(\alpha_1) = 1 = 2\sigma_1 + 1$.

For the inductive case, suppose that the claim is true for some $m \in \{1, \dots, k - 2\}$, we prove that the claim is true for $m + 1$ as well. We distinguish some cases, depending on the type of α_{m+1} and on whether p_m is defined or not.

- First, suppose that α_{m+1} is a non-large angle and that p_m is undefined; then p_{m+1} is undefined, $\sigma_{m+1} = \sigma_m$, and $\lambda(\alpha_{m+1}) \leq 0$. It follows that $\sum_{i=1}^{m+1} \lambda(\alpha_i) \leq \sum_{i=1}^m \lambda(\alpha_i) \leq 2\sigma_m = 2\sigma_{m+1}$, where the second inequality uses the inductive hypothesis.
- Second, suppose that α_{m+1} is a non-large angle, that p_m is defined, and that v_{p_m} is a source; then $v_{p_{m+1}} = v_{p_m}$ is a source, $\sigma_{m+1} = \sigma_m$, and $\lambda(\alpha_{m+1}) \leq 0$. As in the previous case, it follows that $\sum_{i=1}^{m+1} \lambda(\alpha_i) \leq \sum_{i=1}^m \lambda(\alpha_i) \leq 2\sigma_m = 2\sigma_{m+1}$.
- Third, suppose that α_{m+1} is a non-large angle, that p_m is defined, and that v_{p_m} is a sink; then $v_{p_{m+1}} = v_{p_m}$ is a sink, $\sigma_{m+1} = \sigma_m$, and $\lambda(\alpha_{m+1}) \leq 0$. It follows that $\sum_{i=1}^{m+1} \lambda(\alpha_i) = \sum_{i=1}^m \lambda(\alpha_i) \leq 2\sigma_m + 1 = 2\sigma_{m+1} + 1$.
- Fourth, suppose that α_{m+1} is a large angle and that v_{m+1} is a source; then $\sigma_{m+1} = \sigma_m + 1$ and $\lambda(\alpha_{m+1}) = 1$. It follows that $\sum_{i=1}^{m+1} \lambda(\alpha_i) = 1 + \sum_{i=1}^m \lambda(\alpha_i) \leq 1 + (2\sigma_m + 1) = 2\sigma_{m+1} + 1$.
- Fifth, suppose that α_{m+1} is a large angle, that v_{m+1} is a sink, and that p_m is undefined; then $\lambda(\alpha_{m+1}) = 1$. Further, each angle among $\alpha_1, \dots, \alpha_m$ is small or flat, hence $\sum_{i=1}^m \lambda(\alpha_i) \leq 0$. It follows that $\sum_{i=1}^{m+1} \lambda(\alpha_i) \leq 1 \leq 2\sigma_{m+1} + 1$.

- Sixth, suppose that α_{m+1} is a large angle, that v_{m+1} is a sink, that p_m is defined, and that v_{p_m} is a source; then $\sigma_{m+1} = \sigma_m$ and $\lambda(\alpha_{m+1}) = 1$. It follows that $\sum_{i=1}^{m+1} \lambda(\alpha_i) = 1 + \sum_{i=1}^m \lambda(\alpha_i) \leq 1 + 2\sigma_m = 2\sigma_{m+1} + 1$.
- Finally, suppose that α_{m+1} is a large angle, that v_{m+1} is a sink, that p_m is defined, and that v_{p_m} is a sink; then $\sigma_{m+1} = \sigma_m$ and $\lambda(\alpha_{m+1}) = 1$.

With respect to the previous cases, here we need to prove the following observation: there exists (at least) one angle among $\alpha_{p_m+1}, \alpha_{p_m+2}, \dots, \alpha_m$ that is small. Indeed, by definition of p_m , each angle among $\alpha_{p_m+1}, \alpha_{p_m+2}, \dots, \alpha_m$ is either small or flat. Further, since v_{p_m} is a sink, the edge between v_{p_m} and v_{p_m+1} is incoming v_{p_m} . Now consider the smallest index $i \geq 1$ such that the edge between v_{p_m+i} and v_{p_m+i+1} is outgoing v_{p_m+i} ; such an index exists, as otherwise v_{m+1} would not be a sink. Then the edges connecting v_{p_m+i} to v_{p_m+i-1} and v_{p_m+i+1} are both outgoing v_{p_m+i} ; since α_{p_m+i} is not large, it is small, which proves the claimed observation.

It follows that $\sum_{i=1}^{m+1} \lambda(\alpha_i) = \sum_{i=1}^{p_m} \lambda(\alpha_i) + \sum_{i=p_m+1}^m \lambda(\alpha_i) + \lambda(\alpha_{m+1}) \leq (2\sigma_m + 1) - 1 + 1 = 2\sigma_{m+1} + 1$.

This completes the induction, hence proves the claim and that the left-turn-number of \mathcal{E}_μ is at most $2\sigma_\mu + 1$. An analogous proof shows that the right-turn-number of \mathcal{E}_μ is also at most $2\sigma_\mu + 1$. From this, it follows that $\sum_{i=0}^k \lambda(\alpha_i) \leq 2\sigma_\mu + 3$ and $\sum_{i=0}^h \lambda(\beta_i) \leq 2\sigma_\mu + 3$ (the two sums are the left- and right-turn-number of \mathcal{E}_μ , respectively, increased by the angles at u and v inside the outer face of \mathcal{E}_μ). By property **UP3** of Theorem 1, we have that $\sum_{i=1}^{k-1} \lambda(\alpha_i) + \sum_{i=0}^h \lambda(\beta_i) = 2$, hence $\sum_{i=1}^{k-1} \lambda(\alpha_i) = 2 - \sum_{i=0}^h \lambda(\beta_i) \geq 2 - (2\sigma_\mu + 3) = -2\sigma_\mu - 1$, hence the left-turn-number of \mathcal{E}_μ is at least $-2\sigma_\mu - 1$. Analogously, the right-turn-number of \mathcal{E}_μ is at least $-2\sigma_\mu - 1$, which concludes the proof of the lemma. ◀

A direct consequence of Lemma 27 is that we can bound the size of a feasible set \mathcal{F}_μ of μ .


► **Lemma 28.** *The size of the feasible set \mathcal{F}_μ of μ is at most $72\sigma_\mu + 54$.*

Proof. Note that G_μ contains at most σ_μ sources other than u and v . By Lemma 27, the shape description of \mathcal{E}_μ has the value $\tau_l(\mathcal{E}_\mu, u, v)$ in $[-2\sigma_\mu - 1, 2\sigma_\mu + 1]$, hence $\tau_l(\mathcal{E}_\mu, u, v)$ can have one of $4\sigma_\mu + 3$ possible values. By Lemma 12 there are at most 18 shape descriptions with given $\tau_l(\mathcal{E}_\mu, u, v)$. Hence, \mathcal{F}_μ has at most $(4\sigma_\mu + 3) \cdot 18 = 72\sigma_\mu + 54$ possible shape descriptions. ◀

6.2 An R-node Subprocedure

In the following, we show an algorithm running in $\mathcal{O}(\sigma^{1.45} \cdot k \log^3 k)$ time that computes the feasible set \mathcal{F}_μ of an R-node μ of T with k children. Let u and v be the poles of μ . Let σ_μ be the number of sources of G_μ different from its poles. Furthermore, let ν_1, \dots, ν_k be the children of μ and, for $i = 1, \dots, k$, let u_i and v_i be the poles of ν_i and e_i be the virtual edge representing ν_i in the skeleton $\text{sk}(\mu)$ of μ . We assume that we have the feasible set \mathcal{F}_{ν_i} for each child ν_i of μ .

We first introduce two classifications of the components of G_μ (that is, $G_{\nu_1}, \dots, G_{\nu_k}$). First, a component G_{ν_i} is *boring* if it does not contain any sources except, possibly, for its poles, while it is *interesting* if it contains at least one source different from its poles. Because G_μ has σ_μ sources, there are at most σ_μ interesting components among $G_{\nu_1}, \dots, G_{\nu_k}$, while there can be any number of boring components. Second, a component G_{ν_i} is *extreme* if e_i is incident to a pole of μ and is incident to the face of the (unique) planar embedding of $\text{sk}(\mu)$ containing u and v , it is *non-extreme* otherwise. Note that there are four extreme components



■ **Figure 9** Shape descriptions of boring components.

among $G_{\nu_1}, \dots, G_{\nu_k}$. Clearly, an $\mathcal{O}(n)$ -time pre-processing of G allows us to equip every virtual edge e in the skeleton of every R-node τ of T with a label, indicating whether the component of G_τ corresponding to e is boring, interesting, extreme, or non-extreme. With such a labeling, we can produce in $\mathcal{O}(k)$ time (and thus in $\mathcal{O}(n)$ time over all the R-nodes of T) an ordering of the components of G_μ such that all the extreme or interesting components precede all the non-extreme boring components. We assume, w.l.o.g., that $G_{\nu_1}, \dots, G_{\nu_k}$ is such an order, where $G_{\nu_1}, \dots, G_{\nu_h}$ are the extreme or interesting components.

The outline of our algorithm to compute the feasible set \mathcal{F}_μ of μ is as follows. For each shape description $s = \langle \tau_l, \tau_r, \lambda^u, \lambda^v, \rho_l^u, \rho_r^u, \rho_l^v, \rho_r^v \rangle$, where $\tau_l \in [-2\sigma_\mu - 1, 2\sigma_\mu + 1]$, $\tau_r \in [-\tau_l, -\tau_l + 4]$, $\lambda^u \in \{-1, 0, 1\}$, $\lambda^v \in \{-1, 0, 1\}$, $\rho_l^u \in \{\text{in}, \text{out}\}$, $\rho_r^u \in \{\text{in}, \text{out}\}$, $\rho_l^v \in \{\text{in}, \text{out}\}$, and $\rho_r^v \in \{\text{in}, \text{out}\}$, we independently test whether G_μ admits a uv -external upward planar embedding with shape description s ; then \mathcal{F}_μ contains the shape descriptions for which the test was successful. In order to test whether G_μ admits a uv -external upward planar embedding with shape description s , we consider all possible combinations of shape descriptions for the extreme and interesting components of G_μ . For each of these combinations, we create a flow network that admits a flow of a certain size if and only if G_μ admits a uv -external upward planar embedding whose shape description is s and in which the shape descriptions of the extreme and interesting components are those in the considered combination.

6.2.1 Boring components

Despite their name, boring components play an important role in our algorithm. In the following we discuss their features.

► **Lemma 29.** *If G_{ν_i} is a boring component, then the shape description of any $u_i v_i$ -external upward planar embedding of G_{ν_i} is one of the following:*

1. $\langle 0, 0, 1, 1, \text{out}, \text{out}, \text{in}, \text{in} \rangle$ (see Fig. 9a);
2. $\langle 0, 0, 1, 1, \text{in}, \text{in}, \text{out}, \text{out} \rangle$ (see Fig. 9a with u_i and v_i inverted);
3. $\langle 0, 1, 1, 0, \text{out}, \text{out}, \text{in}, \text{out} \rangle$ (see Fig. 9b);
4. $\langle 1, 0, 0, 1, \text{out}, \text{in}, \text{out}, \text{out} \rangle$ (see Fig. 9b with u_i and v_i inverted);
5. $\langle 1, 0, 1, 0, \text{out}, \text{out}, \text{out}, \text{in} \rangle$ (see Fig. 9c);
6. $\langle 0, 1, 0, 1, \text{out}, \text{in}, \text{out}, \text{out} \rangle$ (see Fig. 9c with u_i and v_i inverted);
7. $\langle -1, 1, 1, 1, \text{out}, \text{out}, \text{out}, \text{out} \rangle$ (see Fig. 9d);
8. $\langle 1, -1, 1, 1, \text{out}, \text{out}, \text{out}, \text{out} \rangle$ (see Fig. 9d with u_i and v_i inverted);
9. $\langle 1, 1, 1, -1, \text{out}, \text{out}, \text{out}, \text{out} \rangle$ (see Fig. 9e); and
10. $\langle 1, 1, -1, 1, \text{out}, \text{out}, \text{out}, \text{out} \rangle$ (see Fig. 9e with u_i and v_i inverted);

Proof. Consider any uv -external upward planar embedding \mathcal{E}_{ν_i} of G_{ν_i} . Let s_{ν_i} be the shape description of \mathcal{E}_{ν_i} . Since G_{ν_i} is a boring component, only u_i and v_i can be sources of G_{ν_i} .

Then, by Lemma 27, the left- and right-turn-numbers $\tau_l(\mathcal{E}_{\nu_i}, u_i, v_i)$ and $\tau_r(\mathcal{E}_{\nu_i}, u_i, v_i)$ of \mathcal{E}_{ν_i} can only take the values -1 , 0 , and 1 .

- Assume first that both u_i and v_i are sources of G_{ν_i} . It follows that $\{\lambda(\mathcal{E}_{\nu_i}, u_i), \lambda(\mathcal{E}_{\nu_i}, v_i)\} \subseteq \{-1, 1\}$, that the last four values of s_{ν_i} are all out, and that $\tau_l(\mathcal{E}_{\nu_i}, u_i, v_i)$ and $\tau_r(\mathcal{E}_{\nu_i}, u_i, v_i)$ are both odd. By Corollary 8, we have that $\tau_l(\mathcal{E}_{\nu_i}, u_i, v_i) + \tau_r(\mathcal{E}_{\nu_i}, u_i, v_i) + \lambda(\mathcal{E}_{\nu_i}, u_i) + \lambda(\mathcal{E}_{\nu_i}, v_i) = 2$. Hence:
 - If $\lambda(\mathcal{E}_{\nu_i}, u_i) = \lambda(\mathcal{E}_{\nu_i}, v_i) = 1$, then one between $\tau_l(\mathcal{E}_{\nu_i}, u_i, v_i)$ and $\tau_r(\mathcal{E}_{\nu_i}, u_i, v_i)$ is -1 and the other one is 1 (see Fig. 9d, possibly with u_i and v_i inverted).
 - If $\{\lambda(\mathcal{E}_{\nu_i}, u_i), \lambda(\mathcal{E}_{\nu_i}, v_i)\} = \{-1, +1\}$, then $\tau_l(\mathcal{E}_{\nu_i}, u_i, v_i)$ and $\tau_r(\mathcal{E}_{\nu_i}, u_i, v_i)$ are both 1 (see Fig. 9e, possibly with u_i and v_i inverted).
 - Note that $\lambda(\mathcal{E}_{\nu_i}, u_i)$ and $\lambda(\mathcal{E}_{\nu_i}, v_i)$ cannot be both -1 , as otherwise $\tau_l(\mathcal{E}_{\nu_i}, u_i, v_i) + \tau_r(\mathcal{E}_{\nu_i}, u_i, v_i) + \lambda(\mathcal{E}_{\nu_i}, u_i) + \lambda(\mathcal{E}_{\nu_i}, v_i) = 2$ could not be satisfied, as $\tau_l(\mathcal{E}_{\nu_i}, u_i, v_i) + \tau_r(\mathcal{E}_{\nu_i}, u_i, v_i) \leq 2$.
- Assume next that u_i is a source of G_{ν_i} and v_i is not. It follows that the fifth and sixth values of s_{ν_i} are both out.
 - If v_i is a sink of G_{ν_i} , then the seventh and eighth values of s_{ν_i} are both in. Hence, $\tau_l(\mathcal{E}_{\nu_i}, u_i, v_i)$ and $\tau_r(\mathcal{E}_{\nu_i}, u_i, v_i)$ are even and thus they are both 0 . By Corollary 8, we have that $\tau_l(\mathcal{E}_{\nu_i}, u_i, v_i) + \tau_r(\mathcal{E}_{\nu_i}, u_i, v_i) + \lambda(\mathcal{E}_{\nu_i}, u_i) + \lambda(\mathcal{E}_{\nu_i}, v_i) = 2$, hence $\lambda(\mathcal{E}_{\nu_i}, u_i) = \lambda(\mathcal{E}_{\nu_i}, v_i) = 1$ (see Fig. 9a).
 - If v_i is a non-switch vertex of G_{ν_i} , then $\lambda(\mathcal{E}_{\nu_i}, v_i) = 0$; further, either the seventh value of s_{ν_i} is in and the eight value is out, or vice versa. In the former case (see Fig. 9b), $\tau_l(\mathcal{E}_{\nu_i}, u_i, v_i)$ is even (and thus it is 0) and $\tau_r(\mathcal{E}_{\nu_i}, u_i, v_i)$ is odd (and thus it is 1 , as if it were -1 , then $\tau_l(\mathcal{E}_{\nu_i}, u_i, v_i) + \tau_r(\mathcal{E}_{\nu_i}, u_i, v_i) + \lambda(\mathcal{E}_{\nu_i}, u_i) + \lambda(\mathcal{E}_{\nu_i}, v_i) = 2$ could not be satisfied), thus $\lambda(\mathcal{E}_{\nu_i}, u_i) = 1$. In the latter case (see Fig. 9c), $\tau_r(\mathcal{E}_{\nu_i}, u_i, v_i)$ is even (and thus it is 0) and $\tau_l(\mathcal{E}_{\nu_i}, u_i, v_i)$ is odd (and thus it is 1 , as if it were -1 , then $\tau_l(\mathcal{E}_{\nu_i}, u_i, v_i) + \tau_r(\mathcal{E}_{\nu_i}, u_i, v_i) + \lambda(\mathcal{E}_{\nu_i}, u_i) + \lambda(\mathcal{E}_{\nu_i}, v_i) = 2$ could not be satisfied), thus $\lambda(\mathcal{E}_{\nu_i}, u_i) = 1$.

The case in which v_i is a source of G_{ν_i} and u_i is not is symmetric to the previous one (see Figs. 9a–9c with u_i and v_i inverted). Finally, one of u_i and v_i is a source, given that G_{ν_i} is acyclic. ◀

In the following, we call (inverted-) *sausage* the shape description in item 1 (resp. item 2) of Lemma 29, (inverted-) *right-wing* the shape descriptions in item 3 (resp. item 4) of Lemma 29, (inverted-) *left-wing* the shape description in item 5 (resp. item 6) of Lemma 29, (inverted-) *hat* the shape descriptions in item 7 (resp. item 8) of Lemma 29, and (inverted-) *heart* the shape descriptions in item 9 (resp. item 10) of Lemma 29. The next lemma describes the combinations of shape descriptions that can appear in the feasible set of a non-extreme boring component and which shape descriptions are “better” than others.

► **Lemma 30.** *For every $i = 1, \dots, h$, the following statements hold true:*

- S1** *If \mathcal{F}_{ν_i} contains the sausage or the inverted-sausage, then $|\mathcal{F}_{\nu_i}| = 1$, i.e., \mathcal{F}_{ν_i} contains no other shape description.*
- S2** *If \mathcal{F}_{ν_i} contains the right-wing (the left-wing), then it contains the left-wing (resp. the right-wing), it might contain the heart, and it does not contain any other shape description.*
- S3** *If \mathcal{F}_{ν_i} contains the inverted-right-wing (the inverted-left-wing), then it contains the inverted-left-wing (the inverted-right-wing), it might contain the inverted-heart, and it does not contain any other shape description.*

- S4** If \mathcal{F}_{ν_i} contains the hat (the inverted-hat), then it contains the inverted-hat (resp. the hat), it might contain the heart, it might contain the inverted-heart, and it does not contain any other shape description.
- S5** Suppose that G_μ admits a uv -external upward planar embedding \mathcal{E}_μ with shape description s , in which the shape description of the $u_i v_i$ -external upward planar embedding \mathcal{E}_{ν_i} of G_{ν_i} is the heart. Suppose also that G_{ν_i} admits a $u_i v_i$ -external upward planar embedding \mathcal{E}'_{ν_i} whose shape description is the right-wing, the left-wing, the hat, or the inverted-hat. Then replacing \mathcal{E}_{ν_i} with \mathcal{E}'_{ν_i} in \mathcal{E}_μ results in a uv -external upward planar embedding \mathcal{E}'_μ of G_μ with shape description s .
- S6** Suppose that G_μ admits a uv -external upward planar embedding \mathcal{E}_μ with shape description s , in which the shape description of the $u_i v_i$ -external upward planar embedding \mathcal{E}_{ν_i} of G_{ν_i} is the inverted-heart. Suppose also that G_{ν_i} admits a $u_i v_i$ -external upward planar embedding \mathcal{E}'_{ν_i} whose shape description is the inverted-right-wing, the inverted-left-wing, the hat, or the inverted-hat. Then replacing \mathcal{E}_{ν_i} with \mathcal{E}'_{ν_i} in \mathcal{E}_μ results in a uv -external upward planar embedding \mathcal{E}'_μ of G_μ with shape description s .


Proof. We prove statement S1. Suppose that \mathcal{F}_{ν_i} contains the sausage; the case in which \mathcal{F}_{ν_i} contains the inverted-sausage is analogous. Let \mathcal{E}_{ν_i} be a $u_i v_i$ -external upward planar embedding of G_{ν_i} whose shape description is the sausage. Since both the edges incident to v_i and to the outer face of \mathcal{E}_{ν_i} are incoming v_i (as the seventh and eighth values in the sausage are both in) and since the angle at v_i incident to the outer face of \mathcal{E}_{ν_i} is large (as the fourth value in the sausage is 1), it follows that v_i contains no outgoing edge. In all other shape descriptions of a boring component, the seventh and eighth values contain at least one out (see items 2–10 in Lemma 29), hence they require at least one outgoing edge for v_i . Hence, the sausage is the only shape description in the feasible set of μ .

We prove statement S2. Suppose that \mathcal{F}_{ν_i} contains the right-wing; the case in which \mathcal{F}_{ν_i} contains the left-wing is analogous. Let \mathcal{E}_{ν_i} be a $u_i v_i$ -external upward planar embedding of G_{ν_i} whose shape description is the right-wing. Then a $u_i v_i$ -external upward planar embedding of G_{ν_i} whose shape description is the left-wing can be simply obtained by flipping \mathcal{E}_{ν_i} . By statement S1, we have that \mathcal{F}_{ν_i} does not contain the (possibly inverted-) sausage. Further, v_i contains both incoming and outgoing edges (as the seventh and eighth values in the right-wing are one in and one out). On the other hand, each shape description among the inverted-right-wing, the inverted-left-wing, the hat, the inverted-hat, and the inverted-heart requires v_i to only have outgoing edges (as, in each of such shape descriptions, the seventh and eighth values are both out and the fourth value is 1, see items 4, 6, 7, 8, and 10 in Lemma 29). Hence, \mathcal{F}_{ν_i} does not contain any of these shape descriptions.

The proof of statement S3 is analogous to the one of statement S2.

We prove statement S4. Suppose that \mathcal{F}_{ν_i} contains the hat; the case in which \mathcal{F}_{ν_i} contains the inverted-hat is analogous. Let \mathcal{E}_{ν_i} be a $u_i v_i$ -external upward planar embedding of G_{ν_i} whose shape description is the hat. Then a $u_i v_i$ -external upward planar embedding of G_{ν_i} whose shape description is the inverted-hat can be simply obtained by flipping \mathcal{E}_{ν_i} . By statements S1–S3, we have that \mathcal{F}_{ν_i} contains neither the (possibly inverted-) sausage, nor the (possibly inverted-) right-wing, nor the (possibly inverted-) left-wing.

We finally prove statement S5; the proof of statement S6 is analogous. Suppose that the shape description of \mathcal{E}_{ν_i} is the heart and that the one of \mathcal{E}'_{ν_i} is the right-wing; the other cases are analogous. Let f (g) be the face of \mathcal{E}_μ that is to the left (resp. to the right) of \mathcal{E}_{ν_i} when traversing the left (resp. right) outer path of \mathcal{E}_{ν_i} from u_i to v_i . Similarly, let f' (g') be the face of \mathcal{E}'_μ that is to the left (resp. to the right) of \mathcal{E}'_{ν_i} when traversing the left (resp. right) outer path of \mathcal{E}'_{ν_i} from u_i to v_i .



■ **Figure 10** (a) The upward planar embedding \mathcal{E}_μ , where only \mathcal{E}_{ν_i} and the faces f and g are shown. The internal faces of \mathcal{E}_{ν_i} are gray. The angles whose values are different from -1 are shown. (b) The upward planar embedding \mathcal{E}'_μ , where only \mathcal{E}'_{ν_i} and the faces f' and g' are shown. The internal faces of \mathcal{E}'_{ν_i} are gray. The angles whose values are different from -1 are shown.

Let us specify how the replacement of \mathcal{E}_{ν_i} with \mathcal{E}'_{ν_i} in \mathcal{E}_μ defines the angle assignment of \mathcal{E}'_μ ; refer to Fig. 10. First, every face h' of \mathcal{E}'_μ different from f' and g' is also a face of \mathcal{E}_μ or of \mathcal{E}'_{ν_i} , hence the value in $\{-1, 0, 1\}$ assigned to a vertex incident to h' in \mathcal{E}'_μ coincides with the one in \mathcal{E}_μ . Second, for each vertex $w \notin \{u_i, v_i\}$ incident to f' (resp. to g'), the value in $\{-1, 0, 1\}$ assigned to w in f' (resp. in g') coincides with the value assigned to w in the outer face of \mathcal{E}'_{ν_i} . Third, the value in $\{-1, 0, 1\}$ assigned to u_i in f' (resp. in g') coincides with the one assigned to u_i in f (resp. in g). Finally, the values assigned to v_i in f' and in g' are 0 and -1 , respectively.

In order to prove that \mathcal{E}'_μ is a uv -external upward planar embedding of G_μ , we need to prove that Properties **UP0**–**UP3** of Theorem 1 are satisfied by \mathcal{E}'_μ .

UP0 Consider the angle $\alpha = \langle e'_1, w, e'_2 \rangle$ at a vertex w in a face h' of \mathcal{E}'_μ .

If $h' \notin \{f', g'\}$ or if $w \notin \{u_i, v_i\}$, then α is also an angle of \mathcal{E}_μ or of \mathcal{E}'_{ν_i} and, by construction, the value assigned to α in \mathcal{E}'_μ is the same as the one assigned to α in \mathcal{E}_μ or of \mathcal{E}'_{ν_i} . Hence, from the fact that \mathcal{E}_μ and \mathcal{E}'_{ν_i} are upward planar embeddings, it follows that the value assigned to α is in $\{-1, 1\}$ if α is a switch angle, and 0 if α is a flat angle.

If $h' \in \{f', g'\}$, say $h' = f'$, and $w = u_i$, then e'_1 is the edge incident to u_i in the left outer path of \mathcal{E}'_{ν_i} and e'_2 is the edge incident to u_i and different from e'_1 in the boundary of f' . Let $\beta = \langle e_1, w, e_2 \rangle$ be the angle at w in f , where e'_1 is the edge incident to u_i in the left outer path of \mathcal{E}_{ν_i} and e_2 is the edge incident to u_i and different from e_1 in the boundary of f . Note that $e'_2 = e_2$. Further, since the shape descriptions of \mathcal{E}_{ν_i} and \mathcal{E}'_{ν_i} both have the fifth value equal to out (see items 9 and 3 in Lemma 29, respectively), e_1 and e'_1 are both outgoing u_i . Hence, α is a switch angle or a flat angle if and only if β is a switch angle or a flat angle, respectively. By construction, the value assigned to α in \mathcal{E}'_μ coincides with the value assigned to β in \mathcal{E}_μ . Since \mathcal{E}_μ is an upward planar embedding, the value assigned to α is in $\{-1, 1\}$ if α is a switch angle, and 0 if α is a flat angle.

If $h' = f'$ and $w = v_i$, then e'_1 is the edge incident to v_i in the left outer path of \mathcal{E}'_{ν_i} and e'_2 is the edge incident to v_i and different from e'_1 in the boundary of f' . Since the shape description of \mathcal{E}'_{ν_i} has the seventh value equal to in (see item 3 in Lemma 29), the edge e'_1 is incoming v_i . Further, we have that e'_2 is outgoing v_i . Indeed, since the shape description of \mathcal{E}_{ν_i} has the fourth value equal to -1 (see item 9 in Lemma 29), all the edges that are incident to v_i in G_μ and that do not belong to G_{ν_i} are outgoing v_i . It follows that α is a flat angle; by construction, α is assigned the value 0.

If $h' = f'$ and $w = v_i$, then e'_1 is the edge incident to v_i in the right outer path of \mathcal{E}'_{ν_i} and e'_2 is the edge incident to v_i and different from e'_1 in the boundary of g' . Since the shape description of \mathcal{E}'_{ν_i} has the eighth value equal to out (see item 3 in Lemma 29), the edge e'_1 is outgoing v_i . Further, we have that e'_2 is outgoing v_i , as all the edges that are incident to v_i in G_μ and that do not belong to G_{ν_i} are outgoing v_i . It follows that α is a small angle; by construction, α is assigned the value -1 .

UP1 Consider a switch vertex w of G_μ and let $n_1(w)$, $n_0(w)$, and $n_{-1}(w)$ be the number of angles labeled 1, 0, and -1 in \mathcal{E}'_μ , respectively.

First, suppose that $w \notin \{u_i, v_i\}$ of G_μ and suppose that w belongs to G_{ν_i} (to $G_\mu \setminus G_{\nu_i}$). Then we have $n_1(w) = 1$, $n_0(w) = 0$, and $n_{-1}(w) = \deg(w) - 1$, because the same equalities hold true in \mathcal{E}'_{ν_i} (resp. in \mathcal{E}_μ) and because the number of angles incident to w assigned with the value 1, 0, and -1 in \mathcal{E}'_μ coincides with the number of angles incident to w assigned with the value 1, 0, and -1 , respectively, in \mathcal{E}'_{ν_i} (resp. in \mathcal{E}_μ).

Second, suppose that $w = u_i$. Since the shape descriptions of \mathcal{E}_{ν_i} and \mathcal{E}'_{ν_i} have the third value equal to 1 (see items 9 and 3 in Lemma 29, respectively), all the angles incident to u_i in the internal faces of \mathcal{E}_{ν_i} and \mathcal{E}'_{ν_i} are assigned the value -1 . Furthermore, the angle incident to u_i in f' (in g') coincides with the angle incident to u_i in f (resp. in g); finally, every face h' of \mathcal{E}'_μ that is incident to u_i and that is not incident to edges of \mathcal{E}'_{ν_i} is also a face of \mathcal{E}_μ , hence the value assigned to the angle at u_i in h' coincides with the one in \mathcal{E}_μ . Thus, we have $n_1(w) = 1$, $n_0(w) = 0$, and $n_{-1}(w) = \deg(w) - 1$ in \mathcal{E}'_μ as the same equalities hold true in \mathcal{E}_μ .

Finally, we have that $w \neq v_i$. Indeed, the shape description of \mathcal{E}'_{ν_i} has the seventh and eighth values equal to in and out, respectively (see item 3 in Lemma 29), hence v_i has at least one outgoing and one incoming edge, thus it is not a switch vertex.

UP2 Consider a non-switch vertex w of G_μ and let $n_1(w)$, $n_0(w)$, and $n_{-1}(w)$ be the number of angles labeled 1, 0, and -1 in \mathcal{E}'_μ , respectively. If $w \neq v_i$, then the proof that $n_1(w) = 0$, $n_0(w) = 2$, and $n_{-1}(w) = \deg(w) - 2$ is analogous to the proof that $n_1(w) = 1$, $n_0(w) = 0$, and $n_{-1}(w) = \deg(w) - 1$ in the case in which w is a switch vertex. We prove that $n_1(v_i) = 0$, $n_0(v_i) = 2$, and $n_{-1}(v_i) = \deg(v_i) - 2$. First, by construction, the angle at v_i in f' has value 0, while the one at v_i in g' has value -1 . Second, since v_i is a non-switch vertex of \mathcal{E}'_{ν_i} , there are two angles labeled 0 and incident to v_i in \mathcal{E}'_{ν_i} ; one of them is the angle in the outer face of \mathcal{E}'_{ν_i} (which is not a face of \mathcal{E}'_μ) and the other one is an angle in an internal face of \mathcal{E}'_{ν_i} (this is also an angle labeled 0 in \mathcal{E}'_μ). Every other angle incident to v_i in an internal face of \mathcal{E}'_{ν_i} has label -1 . Finally, since the shape description of \mathcal{E}_{ν_i} has the fourth value equal to -1 (see item 9 in Lemma 29), all the angles incident to v_i inside a face of \mathcal{E}_μ that lies outside \mathcal{E}_{ν_i} and that is different from f' and g' have label -1 .

UP3 Consider any face h' of \mathcal{E}'_μ . Denote by $n_1(h')$ and $n_{-1}(h')$ the number of angles of h' labeled 1 and -1 , respectively. If $h' \notin \{f', g'\}$, then h' is also a face of \mathcal{E}_μ or of \mathcal{E}'_{ν_i} ; further, by construction, the value assigned to each angle of h' in \mathcal{E}'_μ is the same as the one assigned to it in \mathcal{E}_μ or in \mathcal{E}'_{ν_i} . Hence, we have $n_1(h') = n_{-1}(h') - 2$ if h' is an internal face of \mathcal{E}'_μ or $n_1(h') = n_{-1}(h') + 2$ if h' is the outer face of \mathcal{E}'_μ , because the same is true in \mathcal{E}_μ or in \mathcal{E}'_{ν_i} .

We now prove that $n_1(f') = n_{-1}(f') - 2$ if f' is an internal face of \mathcal{E}'_μ or $n_1(f') = n_{-1}(f') + 2$ if f' is the outer face of \mathcal{E}'_μ ; the proof that $n_1(g') = n_{-1}(g') - 2$ if g' is an internal face of \mathcal{E}'_μ or $n_1(g') = n_{-1}(g') + 2$ if g' is the outer face of \mathcal{E}'_μ is analogous. Let $(u_i = w_1, w_2, \dots, w_x = v_i)$ and $(u_i = w'_1, w'_2, \dots, w'_{x'} = v_i)$ be the left outer paths of \mathcal{E}_{ν_i} and \mathcal{E}'_{ν_i} , respectively. Further, let z_1, \dots, z_y be the vertices of G_μ incident to f (and to f') and not in G_{ν_i} . For $i = 1, \dots, x$, let $\alpha(w_i)$ be angle incident to w_i inside f . Furthermore,

for $i = 1, \dots, x'$, let $\beta(w'_i)$ be angle incident to w'_i inside f' . Similarly, for $i = 1, \dots, y$, let $\alpha(z_i)$ the angle incident to z_i inside f and f' .

Denote by λ the angle assignment of \mathcal{E}_μ . Since \mathcal{E}_μ is an upward planar embedding, we have $\sum_{i=2}^{x-1} \lambda(\alpha(w_i)) + \sum_{i=1}^y \lambda(\alpha(z_i)) + \lambda(\alpha(u_i)) + \lambda(\alpha(v_i)) = -2$ if f is an internal face of \mathcal{E}_μ , or $\sum_{i=2}^{x-1} \lambda(\alpha(w_i)) + \sum_{i=1}^y \lambda(\alpha(z_i)) + \lambda(\alpha(u_i)) + \lambda(\alpha(v_i)) = 2$ if f is the outer face of \mathcal{E}_μ . The sum $\sum_{i=2}^{x-1} \lambda(\alpha(w_i))$ is the left-turn-number of \mathcal{E}_{ν_i} , hence it is equal to 1 (see the first value in the shape description at item 9 of Lemma 29). Further, the value $\lambda(\alpha(v_i))$ is equal to -1 , given that the shape description of \mathcal{E}_{ν_i} has the third value equal to -1 (see again item 9 of Lemma 29). It follows that $\sum_{i=1}^y \lambda(\alpha(z_i)) + \lambda(\alpha(u_i)) = -2$ if f is an internal face of \mathcal{E}_μ or $\sum_{i=1}^y \lambda(\alpha(z_i)) + \lambda(\alpha(u_i)) = 2$ if f is the outer face of \mathcal{E}_μ .

Denote by λ' the angle assignment of \mathcal{E}'_μ . We need to prove that $\sum_{i=2}^{x'-1} \lambda'(\beta(w'_i)) + \sum_{i=1}^y \lambda'(\alpha(z_i)) + \lambda'(\beta(u_i)) + \lambda'(\beta(v_i)) = -2$ if f' is an internal face of \mathcal{E}'_μ , or $\sum_{i=2}^{x'-1} \lambda'(\beta(w'_i)) + \sum_{i=1}^y \lambda'(\alpha(z_i)) + \lambda'(\beta(u_i)) + \lambda'(\beta(v_i)) = 2$ if f' is the outer face of \mathcal{E}'_μ . First, note that f is the outer face of \mathcal{E}_μ if and only if f' is the outer face of \mathcal{E}'_μ . Second, by construction we have $\lambda'(\beta(u_i)) = \lambda(\alpha(u_i))$, we have $\lambda'(\beta(v_i)) = 0$, and, for $i = 1, \dots, y$, we have $\lambda'(\alpha(z_i)) = \lambda(\alpha(z_i))$. Third, the sum $\sum_{i=2}^{x'-1} \lambda'(\beta(w'_i))$ is the left-turn-number of \mathcal{E}'_{ν_i} , hence it is equal to 0 (see the first value in the shape description at item 3 of Lemma 29). It follows that the sum $\sum_{i=2}^{x'-1} \lambda'(\beta(w'_i)) + \sum_{i=1}^y \lambda'(\alpha(z_i)) + \lambda'(\beta(u_i)) + \lambda'(\beta(v_i))$ is equal to $\sum_{i=1}^y \lambda(\alpha(z_i)) + \lambda(\alpha(u_i))$, hence it is equal to -2 if f' is an internal face of \mathcal{E}'_μ or to 2 if f' is the outer face of \mathcal{E}'_μ , as required.

This concludes the proof that \mathcal{E}'_μ is a uv -external upward planar embedding of G_μ .

It remains to prove that the shape description of \mathcal{E}'_μ is s . Since G_{ν_i} is a non-extreme component, the replacement of \mathcal{E}_{ν_i} with \mathcal{E}'_{ν_i} does not alter the last six values of the shape description of the uv -external upward planar embedding of G_μ . We prove that the left-turn-number of \mathcal{E}'_μ is equal to the one of \mathcal{E}_μ ; the proof that the right-turn-number of \mathcal{E}'_μ is equal to the one of \mathcal{E}_μ . First, if the virtual edge e_i of the skeleton of μ corresponding to G_{ν_i} does not belong to the left outer path of the planar embedding of the skeleton of μ in \mathcal{E}_μ , the replacement of \mathcal{E}_{ν_i} with \mathcal{E}'_{ν_i} does not alter the left-turn-number of the uv -external upward planar embedding of G_μ . Otherwise, the contribution of \mathcal{E}_{ν_i} to the left-turn-number of \mathcal{E}_μ is given by the label of u_i in the outer face of \mathcal{E}_μ , plus the left-turn-number of \mathcal{E}_{ν_i} , plus the label of v_i in the outer face of \mathcal{E}_μ . Similarly, the contribution of \mathcal{E}'_{ν_i} to the left-turn-number of \mathcal{E}'_μ is given by the label of u_i in the outer face of \mathcal{E}'_μ , plus the left-turn-number of \mathcal{E}'_{ν_i} , plus the label of v_i in the outer face of \mathcal{E}'_μ . The label of u_i in the outer face of \mathcal{E}_μ coincides with the label of u_i in the outer face of \mathcal{E}'_μ . Further, the left-turn-number of \mathcal{E}_{ν_i} is 1, while the left-turn-number of \mathcal{E}'_{ν_i} is 0. Finally, the label of v_i in the outer face of \mathcal{E}_μ is -1 , while the label of v_i in the outer face of \mathcal{E}'_μ is 0. Hence, the contribution of \mathcal{E}_{ν_i} to the left-turn-number of \mathcal{E}_μ is equal to the contribution of \mathcal{E}'_{ν_i} to the left-turn-number of \mathcal{E}'_μ . It follows that the left-turn-number of \mathcal{E}_μ is equal to the one of \mathcal{E}'_μ , hence the shape description of \mathcal{E}'_μ is s . ◀

Based on Lemma 30, we can associate to each boring component ν_i a *preferred set* \mathcal{P}_{ν_i} , which is a subset of its feasible set \mathcal{F}_{ν_i} defined as follows.

- If \mathcal{F}_{ν_i} contains the sausage or the inverted-sausage, then \mathcal{P}_{ν_i} coincides with \mathcal{F}_{ν_i} ; see statement S1 of Lemma 30.
- If \mathcal{F}_{ν_i} contains the left-wing, then \mathcal{P}_{ν_i} contains the left-wing and the right-wing; see statement S2 of Lemma 30.
- If \mathcal{F}_{ν_i} contains the inverted-left-wing, then \mathcal{P}_{ν_i} contains the inverted-left-wing and the inverted-right-wing; see statement S3 of Lemma 30.

- If \mathcal{F}_{ν_i} contains the hat, then \mathcal{P}_{ν_i} contains the hat and the inverted-hat; see statement S4 of Lemma 30.
- If \mathcal{F}_{ν_i} does not satisfy any of the previous conditions, then, by statements S1–S4 of Lemma 30, it contains only the heart, or the inverted-heart, or both. Then we let \mathcal{P}_{ν_i} coincide with \mathcal{F}_{ν_i} .

By Lemma 30, if there exists a uv -external upward planar embedding of G_μ with some shape description s , then there exists a uv -external upward planar embedding of G_μ with some shape description s in which each boring component G_{ν_i} has a shape description in its preferred set.

6.2.2 The Algorithm

We now present an algorithm to construct the feasible set \mathcal{F}_μ of μ .

Consider a shape description $s = \langle \tau_l, \tau_r, \lambda^u, \lambda^v, \rho_l^u, \rho_r^u, \rho_l^v, \rho_r^v \rangle$, where $\tau_l \in [-2\sigma_\mu - 1, 2\sigma_\mu + 1]$, $\tau_r \in [-\tau_l, -\tau_l + 4]$, $\lambda^u \in \{-1, 0, 1\}$, $\lambda^v \in \{-1, 0, 1\}$, $\rho_l^u \in \{\text{in}, \text{out}\}$, $\rho_r^u \in \{\text{in}, \text{out}\}$, $\rho_l^v \in \{\text{in}, \text{out}\}$, and $\rho_r^v \in \{\text{in}, \text{out}\}$; we show how to test whether $s \in \mathcal{F}_\mu$.

6.2.2.1 Checks

In order to test whether $s \in \mathcal{F}_\mu$, we start by performing some preliminary checks, which might let us conclude that $s \notin \mathcal{F}_\mu$.

We start with a *coherence check*. Recall that only four out of the eight values in s are independent. Thus, we check:

- whether ρ_l^u and ρ_r^u have the same value if $\lambda^u \in \{-1, 1\}$;
- whether ρ_l^u and ρ_r^u have different values if $\lambda^u = 0$;
- whether ρ_l^v and ρ_r^v have the same value if $\lambda^v \in \{-1, 1\}$;
- whether ρ_l^v and ρ_r^v have different values if $\lambda^v = 0$;
- whether ρ_l^u and ρ_l^v have the same value if τ_l is odd;
- whether ρ_l^u and ρ_l^v have different values if τ_l is even; and
- whether $\tau_l + \tau_r + \lambda^u + \lambda^v = 2$ (see Corollary 8).

If any of these checks fails, we conclude that \mathcal{F}_μ does not contain s , otherwise we proceed.

Second, for each extreme or interesting component G_{ν_i} of G_μ , we select a shape description $s_i \in \mathcal{F}_{\nu_i}$. Clearly, the number ℓ of ways this selection can be done is the product of the cardinalities of the sets \mathcal{F}_{ν_i} of the extreme and interesting components of G_μ ; that is, $\ell = \prod_{i=1}^h |\mathcal{F}_{\nu_i}|$ (recall that $G_{\nu_1}, G_{\nu_2}, \dots, G_{\nu_h}$ are the components of G_μ that are extreme or interesting). We will argue later that ℓ can be bounded by a simply-exponential function of σ (and, in particular, is independent of the number of vertices of G_μ). For now, assume that shape descriptions $s_1 \in \mathcal{F}_{\nu_1}, \dots, s_h \in \mathcal{F}_{\nu_h}$ have been fixed. We also fix \mathcal{S}_μ to be a planar embedding of the skeleton $\text{sk}(\mu)$ of μ in which u and v are incident to the outer face. Since μ is an R-node, there are two such planar embeddings, which are one the flip of the other one. The goal now becomes the one of testing whether G_μ admits a uv -external upward planar embedding \mathcal{E}_μ such that:

- P1** the shape description of \mathcal{E}_μ is s ;
- P2** for $i = 1, \dots, h$, the $u_i v_i$ -external upward planar embedding \mathcal{E}_{ν_i} of G_{ν_i} in \mathcal{E}_μ has shape description s_i ; and
- P3** the planar embedding of $\text{sk}(\mu)$ induced by \mathcal{E}_μ is \mathcal{S}_μ .

Then we have that s belongs to \mathcal{F}_μ if and only if this test is successful for at least one selection of the shape descriptions s_1, \dots, s_h and of the planar embedding \mathcal{S}_μ .

We start by performing an *extreme-edge check*. The rationale here is that, since the shape descriptions of the extreme components of G_μ have been already selected, the direction of the edges incident to the outer face of any uv -external upward planar embedding \mathcal{E}_μ of G_μ satisfying Property P2 is already determined. These directions have to be coherent with the last four values in s in order to satisfy Property P1. Let e_i be the virtual edge of $\text{sk}(\mu)$ that is incident to u and that belongs to the left outer path of \mathcal{S}_μ ; assume, w.l.o.g., that $u_i = u$. Then we check whether the fifth value of s_i coincides with the fifth value of s . Three more checks are performed for the other edges incident to u and v and incident to the outer face of \mathcal{E}_μ . If any of these checks fails, then there is no uv -external upward planar embedding of G_μ satisfying Properties P1–P3. Otherwise, we proceed (while being sure that, if a uv -external upward planar embedding \mathcal{E}_μ of G_μ is found satisfying Property P2, then the last four values of the shape description of \mathcal{E}_μ are those specified in s).

We next perform an *angle check*. This verifies whether the shape descriptions s_1, \dots, s_h of the extreme or interesting components and the feasible sets $\mathcal{F}_{\nu_{h+1}}, \dots, \mathcal{F}_{\nu_k}$ of the non-extreme boring components allow for an angle assignment that satisfies Properties **UP0-UP2** of Theorem 1. Consider each vertex w of $\text{sk}(\mu)$.


We first deal with the case in which w is a switch vertex of G_μ . Let l_w be a counter, initially set to 0. The counter l_w is going to represent, in any uv -external upward planar embedding \mathcal{E}_μ of G_μ satisfying Property P2, the number of angles incident to w that are required to be large inside some component G_{ν_i} of G_μ .

- For each extreme or interesting component G_{ν_i} such that $w = u_i$ (or $w = v_i$) and such that the third (resp. the fourth) value of s_i is -1 , increase l_w by 1. Namely, if the third (resp. the fourth) value of s_i is equal to -1 , there is a large angle at w in an internal face of every $u_i v_i$ -external upward planar embedding of G_{ν_i} whose shape description is s_i .
- For each non-extreme boring component G_{ν_i} such that $w = v_i$ (or $w = u_i$), if \mathcal{F}_{ν_i} contains only the heart (resp. the inverted-heart), then increase l_w by 1. Namely, if \mathcal{F}_{ν_i} contains only the heart (resp. the inverted-heart), there is a large angle at w in an internal face of every $u_i v_i$ -external upward planar embedding of G_{ν_i} whose shape description is the heart (resp. the inverted-heart).

We now check whether $l_w \leq 1$. If the check fails, we conclude that there is no uv -external upward planar embedding of G_μ satisfying Properties P1–P3. Indeed, in order to satisfy Property P2, there should be at least two large angles incident to w , which is not possible by Property **UP1**. If $l_w = 1$, then we label w as *unavailable* (meaning that w is already incident to a large angle inside some component of G_μ , hence it cannot be incident to another large angle), whereas if $l_w = 0$, then we label w as *available* (meaning that w still has to be assigned a large angle).

We next deal with the case in which w is a non-switch vertex of G_μ . Recall that G is expanded, hence G_μ contains a unique edge incoming w or a unique edge outgoing w . Let G_{ν_i} be the component of G_μ containing such an edge. Then we check whether the angles at w in the internal faces of every other component are all small. Thus, for each component G_{ν_j} of G_μ with $j \neq i$ such that $w = u_j$ (or $w = v_j$), we check whether the third value of s_j (resp. the fourth value of s_j) is equal to 1. If any of these checks fails, we conclude that there is no uv -external upward planar embedding of G_μ satisfying Properties P1–P3.

There is a final angle check that needs to be performed on the poles of μ . Namely, if $\lambda^u = 1$ ($\lambda^v = 1$) and either u (resp. v) is not a switch vertex or it is an unavailable



■ **Figure 11** (a) Sources for switch vertices. (b) Sources for non-switch vertices. (c)–(d) Sources for extreme or interesting components. (e) Sources for boring hats. (f) Sources for boring hearts.

switch vertex, then we conclude that there is no uv -external upward planar embedding of G_μ satisfying Properties P1–P3.

6.2.2.2 Flow network

We are now going to borrow ideas from the algorithm presented by Bertolazzi et al. in [3] for testing the upward planarity of a plane digraph. As described in Section 2.1, Bertolazzi et al. reduce the problem of testing whether a digraph with a fixed planar embedding \mathcal{E} admits an upward planar embedding to the one of testing whether the vertex-face incidence network \mathcal{N} of \mathcal{E} (in which the capacities of the arcs of \mathcal{N} and the supplies and demands of the nodes of \mathcal{N} are suitably set) admits a flow with a certain value. The skeleton $\text{sk}(\mu)$ of our R-node μ indeed has a fixed planar embedding \mathcal{S}_μ . However, the edges of $\text{sk}(\mu)$ are not actual edges, but rather virtual edges that correspond to components of G_μ . This is not really problematic for the extreme or interesting components, as we already fixed their shape description. In fact, as the shape description of a component G_{ν_i} contains all the relevant information that describes how G_{ν_i} interacts with the rest of G_μ , we do not even need to represent G_{ν_i} in the flow network, but rather we just need to take into account the constant amount of information in its shape description. Handling non-extreme boring components is more challenging. According to Lemma 29, each boring component has $\mathcal{O}(1)$ shape descriptions in its feasible set, however the number of such components is not, in general, bounded by a function of σ only, hence we cannot try all possible combinations for their shape descriptions, as we are doing for the interesting components. Lemma 30 comes here to the rescue in order to restrict the possible shape descriptions of each boring component to at most two, namely those in its preferred set; the choice between these two shape descriptions can be plugged into the flow network.

We now define a bipartite network $\mathcal{N}(S, T, A)$, where S is a set of sources, T is a set of sinks, and A is a set of arcs from the sources to the sinks such that G_μ admits a uv -external upward planar embedding satisfying Properties P1–P3 if and only if \mathcal{N} admits a flow of a certain size. Intuitively, and following the approach in [3], each unit of flow going from a source s to a sink t represents the choice that a vertex of G_μ (represented by s) has a large angle in a face (represented by t) of the planar embedding of G_μ we construct. The number of such large angles has to be equal to $\frac{n_t}{2} - 1$, where n_t is the number of switch angles in t .

The network \mathcal{N} is formally defined as follows. We start by defining the set S of sources.

Switch vertices. For each available switch vertex w of G_μ that is a vertex of $\text{sk}(\mu)$, there is a source s_w in S which can supply a single unit of flow; see Figure 11a.

Non-switch vertices. For each non-switch vertex w of G_μ that is a vertex of $\text{sk}(\mu)$, there is a source s_w in S which can supply a single unit of flow if and only if the following three conditions are satisfied (see Figure 11b):

1. The special component for w is a non-extreme boring component G_{ν_i} of G_μ .
2. Either $w = v_i$ and the preferred set \mathcal{P}_{ν_i} of ν_i consists of the left-wing and the right-wing, or $w = u_i$ and \mathcal{P}_{ν_i} consists of the inverted-left-wing and the inverted-right-wing.
3. The special edge for w is outgoing w .


Extreme or interesting components. Consider each extreme or interesting component G_{ν_i} .

If the shape description s_i has the first value (i.e., the left-turn-number) τ_l^i greater than 0, then there is a source z_l^i in S which can supply τ_l^i units of flow. Further, if s_i has the second value (i.e., the right-turn-number) τ_r^i greater than 0, then there is a source z_r^i in S which can supply τ_r^i units of flow; see Figures 11c and 11d.

Non-extreme boring hats. For each non-extreme boring component G_{ν_i} whose preferred set \mathcal{P}_{ν_i} consists of the hat and of the inverted-hat, there is a source b^i which can supply a single unit of flow; see Figure 11e.

Non-extreme boring hearts. For each non-extreme boring component G_{ν_i} whose preferred set \mathcal{P}_{ν_i} consists of the heart, or of the inverted-heart, or of both, there are two sources b_1^i and b_2^i , each of which can supply a single unit of flow; see Figure 11f.

The set T contains the following sinks. For a virtual edge e_i of $\text{sk}(\mu)$, denote by $f_l(e_i)$ and $f_r(e_i)$ the faces of S_μ which are respectively to the left and to the right of e_i when traversing such an edge from u_i to v_i .



■ **Figure 12** Non-switch vertices w that count for the value of n_f . The special edge for w is red. (a) G_{ν_i} is not the special component for w . In (b)–(e) G_{ν_i} is the special component for w , in (b)–(c) it is an extreme or interesting component, while in (d)–(e) it is non-extreme and boring. (b) The fifth value of s_i is in and the special edge for w is outgoing w . (c) The fifth value of s_i is out and the special edge for w is incoming w . (d) \mathcal{P}_{ν_i} consists of the inverted-heart. (e) \mathcal{P}_{ν_i} consists of the inverted-right-wing and the special edge is outgoing w .

Internal faces. For each internal face f of S_μ , there is a sink t_f in T that demands a number d_f of units of flow equal to $\frac{n_f}{2} - 1$, where n_f is equal to the sum of the following numbers (roughly speaking, n_f represents the number of switch angles incident to f which are not yet “paired” with large angles incident to f ; what follows allows us to count the number of such switch angles although a planar embedding of G_μ is not specified):

- For each virtual edge e_i incident to f such that G_{ν_i} is an extreme or interesting component and such that $f = f_l(e_i)$ ($f = f_r(e_i)$), the absolute value of τ_l^i (resp. of τ_r^i).
- The number of virtual edges e_i incident to f such that G_{ν_i} is a non-extreme boring component whose preferred set consists of the hat and of the inverted-hat, or contains the heart and/or the inverted-heart.
- The number of switch vertices of $\text{sk}(\mu)$ incident to f .

- The number of non-switch vertices w of $\text{sk}(\mu)$ incident to f that satisfy one of the following three conditions. Let e_i and e_j be the virtual edges incident to w and f . Assume, w.l.o.g. up to relabeling of e_i with e_j , that G_{ν_j} is not the special component for w . Further, assume that $w = u_i$; the case $w = v_i$ is analogous. Finally, assume that $f = f_l(e_i)$; the case $f = f_r(e_i)$ is analogous. The conditions are as follows.
 1. G_{ν_i} is not the special component for w (see Figure 12a); or
 2. G_{ν_i} is the special component for w , it is an extreme or interesting component, and the fifth value of s_i is in if and only if the special edge for w is outgoing w (see Figures 12b and 12c); note that fifth value of s_i determines the direction of the edge of G_{ν_i} that is incident to u_i and to f in any $u_i v_i$ -external upward planar embedding of G_{ν_i} whose shape description is s_i ; or
 3. G_{ν_i} is the special component for w , it is a non-extreme boring component, and either (a) the preferred set \mathcal{P}_{ν_i} of G_{ν_i} consists of the inverted-heart (see Figure 12d), or (b) \mathcal{P}_{ν_i} consists of the inverted-right-wing and the special edge is outgoing w (see Figure 12e).

Left- and right-turn-number. There are two sinks t_l and t_r which demand $\frac{\tau_l + n_f^l}{2}$ and $\frac{\tau_r + n_f^r}{2}$ units of flow, respectively, where τ_l and τ_r are the left- and right-turn-numbers of the shape description s whose membership in \mathcal{F}_μ we are establishing, and where n_f^l (n_f^r) is defined as n_f in the previous item, however the face f is the outer face, and the virtual edges and the vertices to be considered in the count for n_f^l (resp. n_f^r) are only those in the left (resp. right) outer path of \mathcal{S}_μ and different from u and v .

Heart or inverted-heart. For each non-extreme boring component G_{ν_i} whose preferred set consists of the heart and of the inverted-heart, there is a sink t_i which demands a single unit of flow; see Figure 12c. Note that either an angle at u_i or an angle at v_i inside the upward planar embedding of G_{ν_i} has to be large.

Angle at u . If $\lambda^u = 1$, there is a sink t^u which demands a single unit of flow.

Angle at v . If $\lambda^v = 1$, there is a sink t^v which demands a single unit of flow.

Finally, the set A contains the following arcs.

Switch vertices to faces. For each available switch vertex $w \notin \{u, v\}$ in $\text{sk}(\mu)$, there is an arc with capacity 1 from the corresponding source $s_w \in S$ to each sink t_f corresponding to a face f of \mathcal{S}_μ incident to w (among these sinks there is t_l or t_r , if w belongs to the left or to the right outer path of \mathcal{S}_μ , respectively).

Non-switch vertices to faces. For each non-switch vertex w of G_μ that corresponds to a source s_w in S , let e_i be the virtual edge of $\text{sk}(\mu)$ corresponding to the special component for w . Then there is an arc with capacity 1 from s_w to each of the two sinks corresponding to the two faces $f_l(e_i)$ and $f_r(e_i)$ of \mathcal{S}_μ incident to e_i (one of these sinks is t_l or t_r if e_i belongs to the left or to the right outer path of \mathcal{S}_μ , respectively).

Extreme or interesting components to faces on the left. For each extreme or interesting component G_{ν_i} such that the first value τ_l^i of s_i is greater than 0, there is an arc with capacity τ_l^i from z_l^i to the sink t_f corresponding to the face $f = f_l(e_i)$ of \mathcal{S}_μ ; note that $t_f = t_l$ or $t_f = t_r$ if e_i belongs to the left or to the right outer path of \mathcal{S}_μ , respectively, and f is the outer face of \mathcal{S}_μ .

Extreme or interesting components to faces on the right. For each extreme or interesting component G_{ν_i} such that the second value τ_r^i of s_i is greater than 0, there is an arc with capacity τ_r^i from z_r^i to the sink t_f corresponding to the face $f = f_r(e_i)$ of \mathcal{S}_μ ; note that $t_f = t_l$ or $t_f = t_r$ if e_i belongs to the left or to the right outer path of \mathcal{S}_μ , respectively, and f is the outer face of \mathcal{S}_μ .


Non-extreme boring hats to faces. For each non-extreme boring component \mathcal{G}_{ν_i} whose preferred set \mathcal{P}_{ν_i} consists of the hat and of the inverted-hat, there is an arc with capacity 1 from b_i^1 to each of the two sinks corresponding to the faces $f_l(e_i)$ and $f_r(e_i)$ of \mathcal{S}_μ incident to e_i (one of these sinks is t_l or t_r if e_i belongs to the left or to the right outer path of \mathcal{S}_μ , respectively).

Non-extreme boring hearts to faces. For each non-extreme boring component \mathcal{G}_{ν_i} whose preferred set \mathcal{P}_{ν_i} consists of the heart, or of the inverted-heart, or of both, there is an arc with capacity 1 from b_i^1 to the sink corresponding to the face $f_l(e_i)$ of \mathcal{S}_μ and there is an arc with capacity 1 from b_i^2 to the sink corresponding to the face $f_r(e_i)$ of \mathcal{S}_μ (one of these sinks is t_l or t_r if e_i belongs to the left or right outer path of \mathcal{S}_μ , respectively).

Switch vertices to non-extreme boring hearts and inverted-hearts. For each non-extreme boring component G_{ν_i} whose preferred set contains the heart and the inverted-heart, there is an arc with capacity 1 from each pole of G_{ν_i} that is an available switch vertex to the sink t_i .

First pole. If $\lambda^u = 1$, then there is an arc with capacity 1 from the source s_u to the sink t^u . If u is a switch vertex and $\lambda^u \in \{-1, 0\}$, there is an arc with capacity 1 from s_u to each sink t_f corresponding to an internal face f of \mathcal{S}_μ incident to u .

Second pole. If $\lambda^v = 1$, then there is an arc with capacity 1 from the source s_v to the sink t^v . If v is a switch vertex and $\lambda^v \in \{-1, 0\}$, there is an arc with capacity 1 from s_v to each sink t_f corresponding to an internal face f of \mathcal{S}_μ incident to v .



■ **Figure 13** Illustration for the construction of a flow network \mathcal{N} that allows us to determine whether a shape description $s = \langle 1, 0, -1, 0, \text{in}, \text{out}, \text{in}, \text{in} \rangle$ belongs to \mathcal{F}_μ . (a) shows the input, which contains of a shape description s_i for each extreme or interesting component G_{ν_i} of G_μ and the preferred set \mathcal{P}_{ν_i} for each non-extreme boring component G_{ν_i} of G_μ . (b) shows the corresponding flow network; arc capacities are not shown (each of them is equal to the supply of the source of the arc). (c) shows a flow for \mathcal{N} whose value coincides with the total number of units of flow that are demanded by the sinks of \mathcal{N} ; each shown arc is traversed by a flow equal to its capacity. (d) shows a uv -external upward planar embedding of G_μ with shape description s corresponding to the flow.

This concludes the construction of the network \mathcal{N} . Note that \mathcal{N} has $\mathcal{O}(k)$ nodes and arcs,

as $\text{sk}(\mu)$ has $\mathcal{O}(k)$ vertices and virtual edges. We define F as the total number of units of flow that are demanded by the sinks in T . We test whether every sink has a non-negative demand and whether \mathcal{N} admits a flow with value F . The latter can be done in $\mathcal{O}(k \log^3 k)$ time by means of an algorithm by Borradaile et al. [6]. We conclude that G_μ admits a uv -external upward planar embedding satisfying Properties P1–P3 if and only if the tests are successful.

Figure 13 for an example of the described algorithm to construct \mathcal{N} . In this example, the demand of each sink x is the following.

- For the sink corresponding to the internal face x incident to u , a , and b , the demand is 1, as $n_x = 4$: One unit comes from the switch vertex a , one from the non-switch vertex u , one from the boring component corresponding to the virtual edge (u, b) , and one from the interesting component corresponding to the virtual edge (b, a) .
- For the sink corresponding to the internal face x incident to u , b , and c , the demand is 1, as $n_x = 4$: Three units come from the non-switch vertices u , b , and c , and one from the boring component corresponding to the virtual edge (u, b) .
- For the sink corresponding to the internal face x incident to a , b , e , and f , the demand is 1, as $n_x = 4$: Two units come from the switch vertices a and f , one from the non-switch vertex b , and one from the interesting component corresponding to the virtual edge (a, f) .
- For the sink corresponding to the internal face x incident to b , c , d , and e , the demand is 1, as $n_x = 4$: One unit comes from the switch vertex d , two from the non-switch vertices b and c , and one from the boring component corresponding to the virtual edge (d, e) .
- For the sink corresponding to the internal face x incident to d , e , and g , the demand is 2, as $n_x = 6$: Two units come from the switch vertices d and g , one from the non-switch vertex e , and three from the boring components corresponding to the virtual edges (d, e) , (d, g) , and (e, g) .
- For the sink corresponding to the internal face x incident to e , f , g , and v , the demand is 2, as $n_x = 6$: Three units come from the switch vertices f , g , and v , one from the non-switch vertex e , one from the boring component corresponding to the virtual edge (e, g) , and one from the interesting component corresponding to the virtual edge (f, v) .
- For the sink t_l , the demand is 4, as τ_l is 1 and $n_x^l = 7$, where x is the outer face: Two units come from the switch vertices a and f , three units from the interesting component corresponding to the virtual edge (u, a) , two units from the interesting components corresponding to the virtual edges (a, f) and (f, v) .
- For the sink t_r , the demand is 2, as τ_r is 0 and $n_x^r = 4$, where x is the outer face: Two units come from the switch vertices d and g , one from the non-switch vertex c , and one from the boring component corresponding to the virtual edge (d, g) .
- For the sink corresponding to the boring component whose preferred set consists of the heart and the inverted-heart, the demand is 1.
- For the sink t^v , the demand is 1, as $\lambda^v = 1$.

6.2.2.3 Correctness and Running Time

In Lemma 32 below, we argue about the correctness and the running time of our algorithm. For the latter, we need the following auxiliary lemma.

► **Lemma 31.** *We have $\ell \in \mathcal{O}(1.45^\sigma)$.*

Proof. For $i = 1, \dots, h$, let σ_i be the number of sources of G_{ν_i} different from u_i and v_i and note that $\sum_{i=1}^h \sigma_i =: \bar{\sigma} \leq \sigma$. By Lemma 28, the size of \mathcal{F}_{ν_i} is at most $72\sigma_i + 54$. Hence, we have that $\ell \leq \prod_{i=1}^h (72\sigma_i + 54)$.

Recall that each component among $G_{\nu_1}, \dots, G_{\nu_h}$ is extreme or interesting, and that at most four components are extreme; each extreme component is either boring or interesting. Let $0 \leq m \leq 4$ be the number of boring components among $G_{\nu_1}, \dots, G_{\nu_h}$ and assume, w.l.o.g., that such components come first in the order $G_{\nu_1}, \dots, G_{\nu_h}$. Hence, we have $\sigma_i = 0$ for $i = 1, \dots, m$, and $\sigma_i > 0$ for $i = m+1, \dots, h$. Then we have $\ell \leq (\prod_{i=1}^m 54) \cdot \left(\prod_{i=m+1}^h (72\sigma_i + 54) \right) \leq 54^4 \cdot 73 \prod_{i=m+1}^h \sigma_i$. We prove that $\prod_{i=m+1}^h \sigma_i \leq e^{\sigma/e}$.

First, we have $\prod_{i=m+1}^h \sigma_i \leq (\bar{\sigma}/x)^x$, where $x := h - m$; that is, $\prod_{i=m+1}^h \sigma_i$ is maximum when all the values σ_i are equal. Indeed, if there is a value $\sigma_i > \bar{\sigma}/x$ and a value $\sigma_j < \bar{\sigma}/x$, then $\sigma_i \cdot \sigma_j < (\bar{\sigma}/x) \cdot (\sigma_i + \sigma_j - \bar{\sigma}/x)$, as can be proved by substituting $\sigma_i = \bar{\sigma}/x + y$ and $\sigma_j = \bar{\sigma}/x - y'$. Hence, the values σ_i can be modified into $\bar{\sigma}/x$ one by one without altering the sum $\sum_{i=m+1}^h \sigma_i = \bar{\sigma}$ and while increasing the product $\prod_{i=m+1}^h \sigma_i$.

Second, we have that $f(x) := (\bar{\sigma}/x)^x$ has its maximum when $x = \bar{\sigma}/e$, where e is the base of the natural logarithm. Indeed, we have $\frac{\partial f(x)}{\partial x} = (\bar{\sigma}/x)^x (\ln(\bar{\sigma}/x) - 1)$, which is positive for $x \in (0, \bar{\sigma}/e)$, null for $x = \bar{\sigma}/e$, and negative for $x \in (\bar{\sigma}/e, \bar{\sigma}]$. The value of $f(x)$ at its maximum is hence $e^{\bar{\sigma}/e} < 1.45^{\bar{\sigma}}$. It follows that $\ell \leq 54^4 \cdot 73 \prod_{i=m+1}^h \sigma_i \in \mathcal{O}(1.45^\sigma)$. ◀

We are now ready to prove the following.

► **Lemma 32.** *Let μ be an R -node of T . The feasible set \mathcal{F}_μ of μ can be computed in $\mathcal{O}(\sigma 1.45^\sigma \cdot k \log^3 k)$ time, where k is the number of children of μ in T and σ is the number of sources of G .*

Proof. First, we argue about the running time of the algorithm we presented. The number of shape descriptions that are checked for membership in \mathcal{F}_μ is in $\mathcal{O}(\sigma)$, by Lemma 28; this determines the multiplicative factor of σ in the running time. For each shape description s , we consider ℓ possible choices for the shape descriptions of the extreme and interesting components of G_μ and two possible choices for the planar embedding of $\text{sk}(\mu)$. By Lemma 31, we have $\ell \in \mathcal{O}(1.45^\sigma)$; this determines the multiplicative factor of 1.45^σ in the running time. Finally, for each choice for the shape descriptions of the extreme and interesting components of G_μ and for each choice for the planar embedding of $\text{sk}(\mu)$, we construct in $\mathcal{O}(k)$ time the planar bipartite network \mathcal{N} and determine in $\mathcal{O}(k \log^3 k)$ time its maximum flow, by means of the algorithm by Borradaile et al. [6].

We now prove the correctness of the algorithm we presented for computing \mathcal{F}_μ . Namely, we prove that, for any shape description s , the graph G_μ admits a uv -external upward planar embedding \mathcal{E}_μ if and only if our algorithm determines that $s \in \mathcal{F}_\mu$.

(\implies) Suppose first that G_μ admits a uv -external upward planar embedding \mathcal{E}_μ with shape description s . We prove that our algorithm determines that $s \in \mathcal{F}_\mu$. First, for each non-extreme boring component G_{ν_i} of G_μ , we replace the $u_i v_i$ -external upward planar embedding of G_{ν_i} contained in \mathcal{E}_μ with a $u_i v_i$ -external upward planar embedding of G_{ν_i} whose shape description is one in the preferred set of G_{ν_i} . By Lemma 30, this modification preserves the property that \mathcal{E}_μ is a uv -external upward planar embedding with shape description s .

By Lemma 27, the first value τ_l of s (i.e., the left-turn-number of \mathcal{E}_μ) is an integer in the interval $[-2\sigma - 1, 2\sigma + 1]$. By Lemma 9, the second value τ_r of s (i.e., the right-turn-number of \mathcal{E}_μ) is a value in $[-\tau_l, -\tau_l + 4]$. Further, by definition, the third and fourth values λ^u and λ^v of s are in the set $\{-1, 0, 1\}$ and the fifth, sixth, seventh, and eighth values $\rho_l^u, \rho_r^u, \rho_l^v$, and ρ_r^v of s are in the set $\{\text{in}, \text{out}\}$. It follows that our algorithm considers s as a possible shape description which is to be tested for membership in \mathcal{F}_μ .

As discussed after the definition of shape description, we have that ρ_l^u and ρ_r^u have the same value if and only if $\lambda^u \in \{-1, 1\}$, that ρ_l^v and ρ_r^v have the same value if and only

if $\lambda^v \in \{-1, 1\}$, that ρ_l^u and ρ_l^v have the same value if and only if τ_l is odd, and that $\tau_l + \tau_r + \lambda^u + \lambda^v = 2$. Hence, s does not fail the coherence check.

Let \mathcal{S}_μ be the planar embedding of $\text{sk}(\mu)$ in \mathcal{E}_μ . Further, for each component G_{ν_i} of G_μ , let \mathcal{E}_{ν_i} be the $u_i v_i$ -external upward planar embedding of G_{ν_i} contained in \mathcal{E}_μ , where u_i and v_i are the poles of ν_i , and let s_i be the shape description of \mathcal{E}_{ν_i} .

Recall that $G_{\nu_1}, \dots, G_{\nu_h}$ are the extreme or interesting components of G_μ . For $i = 1, \dots, h$, since a $u_i v_i$ -external upward planar embedding of G_{ν_i} with shape description s_i exists, we have that $s_i \in \mathcal{F}_{\nu_i}$. Hence, our algorithm tests whether G_μ admits a uv -external upward planar embedding whose shape description is s , whose restriction to G_{ν_i} is a $u_i v_i$ -external upward planar embedding with shape description s_i , for $i = 1, \dots, h$, and such that the corresponding planar embedding of $\text{sk}(\mu)$ is \mathcal{S}_μ . We now prove that, when considering the shape descriptions s_1, \dots, s_h for $G_{\nu_1}, \dots, G_{\nu_h}$, respectively, and the planar embedding \mathcal{S}_μ for $\text{sk}(\mu)$, our algorithm indeed concludes that $s \in \mathcal{F}_\mu$.

The extreme-edge check does not fail, as the shape description s_i of each extreme component G_{ν_i} of G_μ is the one of a $u_i v_i$ -external upward planar embedding contained in a uv -external upward planar embedding of G_μ with shape description s .

The angle check does not fail either. Indeed, consider any vertex w of $\text{sk}(\mu)$. Suppose first that w is a switch vertex of G_μ . Since \mathcal{E}_μ is an upward planar embedding, by Property **UP1** of Theorem 1, there is one angle incident to w in \mathcal{E}_μ which is labeled 1. Hence, at most one component G_{ν_i} of G_μ is such that the angle at w in one of the internal faces of \mathcal{E}_{ν_i} is large; hence, the counter l_w is at most 1. Suppose next that w is a non-switch vertex of G_μ . Since \mathcal{E}_μ is an upward planar embedding, by Property **UP2** of Theorem 1, no angle incident to w in \mathcal{E}_μ is labeled 1. Hence, for every non-special component G_{ν_i} of G_μ , the angle at w in every internal face of \mathcal{E}_{ν_i} is small, hence the angle check does not fail.

The final check at u and v also does not fail. Namely, if $\lambda^u = 1$ ($\lambda^v = 1$), then by Properties **UP1** and **UP2** of Theorem 1, we have that u (resp. v) is a switch vertex; further, since the angle incident to u (resp. to v) in the outer face of \mathcal{E}_μ is large, it follows that every angle at u (resp. at v) inside every internal face of \mathcal{E}_μ , and consequently inside every internal face of \mathcal{E}_{ν_i} , is small, hence u (resp. v) is available.

It remains to prove that the network $\mathcal{N}(S, T, A)$ admits a flow whose value is equal to the sum of the demands of the sinks in T . We compute such a flow as follows.

Switch vertices to faces. For each available switch vertex w in $\text{sk}(\mu)$, if the large angle incident to w in \mathcal{E}_μ lies in the interior of a face of \mathcal{E}_μ corresponding to a face f of \mathcal{S}_μ , we let one unit of flow pass from s_w to the sink t_f corresponding to f (this sink is t_l or t_r , if w belongs to the left or to the right outer path of \mathcal{S}_μ and the large angle incident to w lies in the outer face of \mathcal{E}_μ).

Non-switch vertices to faces. For each non-switch vertex w of G_μ that corresponds to a source s_w in S , let e_i be the virtual edge of $\text{sk}(\mu)$ corresponding to the special component for w . Then we let one unit of flow pass from s_w to the sink corresponding to $f_l(e_i)$ if $w = v_i$ and the shape description of \mathcal{E}_{ν_i} is the left-wing or if $w = u_i$ and the shape description of \mathcal{E}_{ν_i} is the inverted-right-wing, or we let one unit of flow pass from s_w to the sink corresponding to $f_r(e_i)$ if $w = v_i$ and the shape description of \mathcal{E}_{ν_i} is the right-wing or if $w = u_i$ and the shape description of \mathcal{E}_{ν_i} is the inverted-left-wing.

Extreme or interesting components to faces on the left. For each extreme or interesting component G_{ν_i} such that the first value τ_l^i of s_i is greater than 0, we let τ_l^i units of flow pass from z_l^i to the sink t_f corresponding to the face $f = f_l(e_i)$ of \mathcal{S}_μ .

Extreme or interesting components to faces on the right. For each extreme or interesting component G_{ν_i} such that the second value τ_r^i of s_i is greater than 0, we let τ_r^i units of

flow pass from z_r^i to the sink t_f corresponding to the face $f = f_r(e_i)$ of \mathcal{S}_μ .

Non-extreme boring hats to faces. For each non-extreme boring component \mathcal{G}_{ν_i} whose preferred set \mathcal{P}_{ν_i} consists of the hat and of the inverted-hat, we let one unit of flow pass from b^i to the sink corresponding to the face $f_r(e_i)$ or to the face $f_l(e_i)$ of \mathcal{S}_μ , depending on whether the shape description of \mathcal{E}_{ν_i} is the hat or the inverted-hat, respectively.

Non-extreme boring hearts to faces. For each non-extreme boring component \mathcal{G}_{ν_i} whose preferred set \mathcal{P}_{ν_i} consists of the heart, or of the inverted-heart, or of both, we let one unit of flow pass from b_1^i to the sink corresponding to the face $f_l(e_i)$ of \mathcal{S}_μ and unit of flow pass from b_2^i to the sink corresponding to the face $f_r(e_i)$ of \mathcal{S}_μ .

Switch vertices to non-extreme boring hearts and inverted-hearts. For each non-extreme boring component G_{ν_i} whose preferred set contains the heart and the inverted-heart, we let one unit of flow pass from v_i to the sink t_i or from u_i to the sink t_i , depending on whether the shape description of \mathcal{E}_{ν_i} is the heart or the inverted-heart, respectively (note that this satisfies the demand of the sink t_i).

First pole. If $\lambda^u = 1$, then we let one unit of flow pass from the source s_u to the sink t^u (note that this satisfies the demand of the sink t^u).

Second pole. If $\lambda^v = 1$, then we let one unit of flow pass from the source t_u to the sink t^v (note that this satisfies the demand of the sink t^v).

It is easy to verify that each source in S sends a number of units of flow equal to the maximum number of units of flow it can supply. The only interesting sources to consider are:

- The source s_u (similar arguments deal with s_v); note that s_u exists only if $\lambda^u = 1$. If $\lambda^u = 1$, then s_u sends one unit of flow to t^u and it does not send flow to any sink corresponding to an internal face of \mathcal{S}_μ , as the angle at u in the outer face of \mathcal{E}_μ is large, hence the angle at u in any internal face of \mathcal{E}_μ is small.
- Each source s_w that sends one unit of flow to a sink t_i corresponding to a component G_{ν_i} of G_μ whose preferred set contains the heart and the inverted-heart. By the definition of the flow, the angle at the vertex w corresponding to s_w in the outer face of \mathcal{E}_{ν_i} is small (as either $w = v_i$ and the shape of \mathcal{E}_{ν_i} is the heart or $w = u_i$ and the shape of \mathcal{E}_{ν_i} is the inverted-heart). It follows that the angle at w in any internal face of \mathcal{E}_μ that is not an internal face of \mathcal{E}_{ν_i} is small, hence s_w does not send further units of flow.

We now prove that each sink receives a number of units of flow equal to its demand. This was already observed during the construction of the flow for the sinks t^u and t^v , as well as for each sink corresponding to a non-extreme boring component of G_μ whose preferred set contains the heart and the inverted-heart. We now consider the sinks corresponding to internal faces of \mathcal{S}_μ and for the sinks t_l and t_r .

Consider any sink t_f corresponding to an internal face f of \mathcal{S}_μ . First, recall that the demand of t_f is equal to $\frac{n_f}{2} - 1$, by construction.

For each component G_{ν_i} of G_μ , let s_i be the shape description of \mathcal{E}_{ν_i} , and let τ_l^i and τ_r^i be the left- and right-turn-number of \mathcal{E}_{ν_i} , respectively. Let E_1 (E_2) be the set that contains every virtual edge e_i of $\text{sk}(\mu)$ such that $f = f_l(e_i)$ (resp. $f = f_r(e_i)$). Let E'_1 (E'_2) be the subset of E_1 (resp. E_2) that only contains the virtual edges e_i such that $\tau_l^i > 0$ (resp. $\tau_r^i > 0$). Let $f_\mathcal{E}$ be the face of \mathcal{E}_μ corresponding to f . Furthermore, let $n'_1(f_\mathcal{E})$ and $n'_{-1}(f_\mathcal{E})$ be the number of large and small angles of $f_\mathcal{E}$, respectively, incident to vertices of $\text{sk}(\mu)$. Finally, let g be the number of *ignored* virtual edges; a virtual edge e_i is ignored if: (a) $e_i \in E'_1$, s_i is the left-wing, and the special edge for v_i is incoming v_i ; or (b) $e_i \in E'_2$, s_i is the right-wing, and the special edge for v_i is incoming v_i ; or (c) $e_i \in E'_1$, s_i is the inverted-right-wing, and

the special edge for u_i is incoming u_i ; or (d) $e_i \in E'_2$, s_i is the inverted-left-wing, and the special edge for u_i is incoming u_i .

We now prove a sequence of statements.

▷ **Claim 33.** The amount of flow sent to t_f is equal to $\sum_{e_i \in E'_1} \tau_l^i + \sum_{e_i \in E'_2} \tau_r^i + n'_1(f_{\mathcal{E}}) - g$.

Proof. The number of units of flow that are sent to t_f from the sources in S corresponding to available switch vertices of $\text{sk}(\mu)$ is equal to $n'_1(f_{\mathcal{E}})$. Indeed, if a vertex of $\text{sk}(\mu)$ is incident to a large angle in $f_{\mathcal{E}}$, it is an available switch vertex, by Property **UP1** of Theorem 1.

We now deal with the flow sent to t_f from the sources in S corresponding to components of G_{μ} . We need to prove that the total amount of such flow is $\sum_{e_i \in E'_1} \tau_l^i + \sum_{e_i \in E'_2} \tau_r^i - g$.

- Each extreme or interesting component G_{ν_i} such that $e_i \in E'_1$ (such that $e_i \in E'_2$) sends to t_f a number of units of flow equal to τ_l^i (resp. to τ_r^i).
- Each non-extreme boring component G_{ν_i} sends one unit of flow to t_f if $e_i \in E'_1$ and s_i is the left-hat, the heart, or the inverted-heart (in each case we have $\tau_l^i = 1$), or if $e_i \in E'_2$ and s_i is the right-hat, the heart, or the inverted-heart (in each case we have $\tau_r^i = 1$).
- A non-extreme boring component G_{ν_i} such that $e_i \in E'_1$ ($e_i \in E'_2$), such that e_i is not ignored, and such that s_i is the left-wing or the inverted-right-wing (resp. the right-wing or the inverted-left-wing) does not send any unit of flow to t_f , while we have $\tau_l^i = 1$ (resp. $\tau_r^i = 1$). However, this “missing” unit of flow is “compensated” by the non-switch vertex v_i or u_i , for which G_{ν_i} is the special component, which sends one unit of flow to t_f .
- Finally, a non-extreme boring component G_{ν_i} such that $e_i \in E'_1$ ($e_i \in E'_2$), such that e_i is ignored, and such that s_i is the left-wing or the inverted-right-wing (resp. the right-wing or the inverted-left-wing) does not send any unit of flow to t_f , while we have $\tau_l^i = 1$ (resp. $\tau_r^i = 1$). This “missing” unit of flow is not “compensated” by any non-switch vertex, which results in the g term subtracted from the amount of flow sent to t_f .

This concludes the proof of the claim. ◀

▷ **Claim 34.** We have $n_f = \sum_{e_i \in E_1} |\tau_l^i| + \sum_{e_i \in E_2} |\tau_r^i| + n'_1(f_{\mathcal{E}}) + n'_{-1}(f_{\mathcal{E}}) - 2g$.

Proof. By Properties **UP1** and **UP2** of Theorem 1, every large angle of $f_{\mathcal{E}}$ is incident to a switch vertex. Denote by $p'_{-1}(f_{\mathcal{E}})$ and $q'_{-1}(f_{\mathcal{E}})$ the number of small angles of $f_{\mathcal{E}}$ incident to switch and non-switch vertices of $\text{sk}(\mu)$, respectively. Then we have $n'_{-1}(f_{\mathcal{E}}) = p'_{-1}(f_{\mathcal{E}}) + q'_{-1}(f_{\mathcal{E}})$. By construction, each switch vertex of $\text{sk}(\mu)$ incident to f contributes 1 to n_f . Since a switch vertex does not have incident flat angles, the total contribution of the switch vertices to n_f is $n'_1(f_{\mathcal{E}}) + p'_{-1}(f_{\mathcal{E}})$.

Each non-switch vertex incident to a small angle in $f_{\mathcal{E}}$ contributes 1 to n_f , except for the non-switch vertices w such that the special component G_{ν_i} for w corresponds to an ignored virtual edge e_i . By definition, the number of such non-switch vertices is g , hence the total contribution of the non-switch vertices to n_f is $q'_{-1}(f_{\mathcal{E}}) - g$.

We now deal with the contributions to n_f stemming from the components of G_{μ} .

- Each extreme or interesting component G_{ν_i} such that $e_i \in E_1$ ($e_i \in E_2$) contributes to n_f by an amount of $|\tau_l^i|$ (resp. $|\tau_r^i|$).
- Similarly, each non-extreme boring component G_{ν_i} such that $e_i \in E_1$ ($e_i \in E_2$) and such that s_i is the hat, the inverted-hat, the heart, or the inverted-heart contributes to n_f by an amount of $|\tau_l^i| = 1$ (resp. $|\tau_r^i| = 1$).
- Further, each non-extreme boring component G_{ν_i} such that $e_i \in E_1$ and such that s_i is the sausage, the inverted-sausage, the right-wing, or the inverted-left-wing contributes to n_f by an amount of $|\tau_l^i| = 0$.

- Similarly, each non-extreme boring component G_{ν_i} such that $e_i \in E_2$ and such that s_i is the sausage, the inverted-sausage, the left-wing, or the inverted-right-wing contributes to n_f by an amount of $|\tau_r^i| = 0$.
- Each non-extreme boring component G_{ν_i} such that $e_i \in E_1$ ($e_i \in E_2$), such that e_i is not ignored, and such that s_i is the left-wing or the inverted-right-wing (resp. the right-wing or the inverted-left-wing) contributes by an amount of 0 to n_f , while we have $\tau_l^i = 1$ (resp. $\tau_r^i = 1$). However, this “missing” contribution to n_f is “compensated” by the non-switch vertex v_i or u_i , for which G_{ν_i} is the special component, which contributes 1 to n_f while being incident to a flat angle in $f_{\mathcal{E}}$.
- Finally, each non-extreme boring component G_{ν_i} such that $e_i \in E'_1$ ($e_i \in E'_2$), such that e_i is ignored, and such that s_i is the left-wing or the inverted-right-wing (resp. the right-wing or the inverted-left-wing) contributes by an amount of 0 to n_f , while we have $\tau_l^i = 1$ (resp. $\tau_r^i = 1$). This “missing” contribution to n_f is not “compensated” by any non-switch vertex, which results in a second g term subtracted from the value of n_f .

This concludes the proof of the claim. \blacktriangleleft

▷ **Claim 35.** We have $\sum_{e_i \in E'_1} \tau_l^i - \sum_{e_i \in E_1 \setminus E'_1} |\tau_l^i| + \sum_{e_i \in E'_2} \tau_r^i - \sum_{e_i \in E_2 \setminus E'_2} |\tau_r^i| + n'_1(f_{\mathcal{E}}) - n'_{-1}(f_{\mathcal{E}}) = -2$.

Proof. By definition, for each component G_{ν_i} of G_{μ} such that $e_i \in E_1$ ($e_i \in E_2$), the value τ_l^i (resp. τ_r^i) is equal to the number of non-pole vertices of G_{ν_i} incident to large angles in $f_{\mathcal{E}}$ minus the number of non-pole vertices of G_{ν_i} incident to small angles in $f_{\mathcal{E}}$. Hence, the sum $\sum_{e_i \in E_1} \tau_l^i + \sum_{e_i \in E_2} \tau_r^i + n'_1(f_{\mathcal{E}}) - n'_{-1}(f_{\mathcal{E}}) = \sum_{e_i \in E'_1} \tau_l^i - \sum_{e_i \in E_1 \setminus E'_1} |\tau_l^i| + \sum_{e_i \in E'_2} \tau_r^i - \sum_{e_i \in E_2 \setminus E'_2} |\tau_r^i| + n'_1(f_{\mathcal{E}}) - n'_{-1}(f_{\mathcal{E}})$ is equal to the number of vertices of G_{μ} incident to large angles in $f_{\mathcal{E}}$ minus the number of vertices of G_{μ} incident to small angles in $f_{\mathcal{E}}$. By Property **UP3** of Theorem 1, this difference is equal to -2 . \blacktriangleleft

By Claim 35, we have $\sum_{e_i \in E_1 \setminus E'_1} |\tau_l^i| + \sum_{e_i \in E_2 \setminus E'_2} |\tau_r^i| = \sum_{e_i \in E'_1} \tau_l^i + \sum_{e_i \in E'_2} \tau_r^i + n'_1(f_{\mathcal{E}}) - n'_{-1}(f_{\mathcal{E}}) + 2$. This can be replaced into the equality $n_f = \sum_{e_i \in E'_1} \tau_l^i + \sum_{e_i \in E_1 \setminus E'_1} |\tau_l^i| + \sum_{e_i \in E'_2} \tau_r^i + \sum_{e_i \in E_2 \setminus E'_2} |\tau_r^i| + n'_1(f_{\mathcal{E}}) + n'_{-1}(f_{\mathcal{E}}) - 2g$ given by Claim 34, in order to get that $n_f = 2 \sum_{e_i \in E'_1} \tau_l^i + 2 \sum_{e_i \in E'_2} \tau_r^i + 2n'_1(f_{\mathcal{E}}) - 2g + 2$. Hence, the demand of t_f is equal to $\frac{n_f}{2} - 1 = \sum_{e_i \in E'_1} \tau_l^i + \sum_{e_i \in E'_2} \tau_r^i + n'_1(f_{\mathcal{E}}) - g$, which by Claim 33 is equal to the flow sent to t_f .

Consider now the sink t_l (the arguments for the sink t_r are analogous). First, recall that the demand of t_l is equal to $\frac{\tau_l + n'_l}{2}$. We need to prove that the amount of flow which is sent to t_l is equal to the same amount. The proof is very similar to the one for the sink t_f corresponding to an internal face f of \mathcal{S}_{μ} , and is hence only sketched here.

For each component G_{ν_i} of G_{μ} , let τ_l^i and τ_r^i be defined as before. Let E_1 (E_2) be the set that contains every virtual edge e_i of $\text{sk}(\mu)$ that belongs to the left outer path of \mathcal{S}_{μ} and such that the outer face of \mathcal{S}_{μ} is $f_l(e_i)$ (resp. $f_r(e_i)$). Let E'_1 (E'_2) be the subset of E_1 (resp. E_2) that only contains the virtual edges e_i such that $\tau_l^i > 0$ (resp. $\tau_r^i > 0$). Let $f_{\mathcal{E}}$ be the outer face of \mathcal{E}_{μ} . Further, let $n'_1(f_{\mathcal{E}})$ and $n'_{-1}(f_{\mathcal{E}})$ be the number of large and small angles, respectively, incident to $f_{\mathcal{E}}$ and to non-pole vertices of the left outer path of \mathcal{S}_{μ} . Finally, let g be defined as before.

As in the proof for the sink t_f corresponding to an internal face f of \mathcal{S}_{μ} , we need three statements. The first one is the analogue of Claim 33: The amount of flow which is sent to t_l is equal to $\sum_{e_i \in E'_1} \tau_l^i + \sum_{e_i \in E'_2} \tau_r^i + n'_1(f_{\mathcal{E}}) - g$; the proof is identical to the one of Claim 33. The second statement is the analogue of Claim 34: $n'_l = \sum_{e_i \in E_1} |\tau_l^i| + \sum_{e_i \in E_2} |\tau_r^i| + n'_1(f_{\mathcal{E}}) +$

$n'_{-1}(f_{\mathcal{E}}) - 2g$; again, the proof is identical to the one of Claim 34. Finally, the third statement is similar to Claim 35: $\sum_{e_i \in E'_1} \tau_l^i - \sum_{e_i \in E_1 \setminus E'_1} |\tau_l^i| + \sum_{e_i \in E'_2} \tau_r^i - \sum_{e_i \in E_2 \setminus E'_2} |\tau_r^i| + n'_1(f_{\mathcal{E}}) - n'_{-1}(f_{\mathcal{E}}) = \tau_l$; again, its proof follows the same lines of the proof of Claim 35, while using that the left-turn-number of \mathcal{E}_{μ} is τ_l in place of Property **UP3** of Theorem 1.

The three statements imply that the amount of flow which is sent to t_l is equal to $\frac{\tau_l + n'_f}{2}$, as requested. Indeed, by the third statement we have $\sum_{e_i \in E_1 \setminus E'_1} |\tau_l^i| + \sum_{e_i \in E_2 \setminus E'_2} |\tau_r^i| = \sum_{e_i \in E'_1} \tau_l^i + \sum_{e_i \in E'_2} \tau_r^i + n'_1(f_{\mathcal{E}}) - n'_{-1}(f_{\mathcal{E}}) - \tau_l$. This can be replaced into the equality $n'_f = \sum_{e_i \in E'_1} \tau_l^i + \sum_{e_i \in E_1 \setminus E'_1} |\tau_l^i| + \sum_{e_i \in E'_2} \tau_r^i + \sum_{e_i \in E_2 \setminus E'_2} |\tau_r^i| + n'_1(f_{\mathcal{E}}) + n'_{-1}(f_{\mathcal{E}}) - 2g$ given by the second statement, in order to get that $n'_f = 2 \sum_{e_i \in E'_1} \tau_l^i + 2 \sum_{e_i \in E'_2} \tau_r^i + 2n'_1(f_{\mathcal{E}}) - 2g - \tau_l$. Hence, the demand of t_l is equal to $\frac{n'_f + \tau_l}{2} = \sum_{e_i \in E'_1} \tau_l^i + \sum_{e_i \in E'_2} \tau_r^i + n'_1(f_{\mathcal{E}}) - g$, which by the first statement is equal to the flow sent to t_l .

(\Leftarrow) Suppose next that our algorithm determines that $s \in \mathcal{F}_{\mu}$. We prove that G_{μ} admits a uv -external upward planar embedding \mathcal{E}_{μ} with shape description s .

Our algorithm determines that $s \in \mathcal{F}_{\mu}$ when examining some shape descriptions s_1, \dots, s_h for the extreme and interesting components $G_{\nu_1}, \dots, G_{\nu_h}$ of G_{μ} , together with a planar embedding \mathcal{S}_{μ} for the skeleton $\text{sk}(\mu)$ of μ . If the corresponding flow network \mathcal{N} admits a flow in which each sink is supplied with a number of units of flow equal to its demand, then our algorithm indeed concludes that $s \in \mathcal{F}_{\mu}$.

The uv -external upward planar embedding \mathcal{E}_{μ} is constructed starting from \mathcal{S}_{μ} , by selecting a $u_i v_i$ -external upward planar embedding \mathcal{E}_{ν_i} for each component G_{ν_i} of G_{μ} , and by assigning a label to each vertex of $\text{sk}(\mu)$ inside the faces of \mathcal{S}_{μ} . Selecting a $u_i v_i$ -external upward planar embedding \mathcal{E}_{ν_i} for each component G_{ν_i} of G_{μ} implies that the label of each angle at a vertex $w \notin \{u_i, v_i\}$ of G_{ν_i} in \mathcal{E}_{μ} is the same as in \mathcal{E}_{ν_i} ; further, the label of each angle in an internal face of \mathcal{E}_{ν_i} at u_i or v_i in \mathcal{E}_{μ} is the same as in \mathcal{E}_{ν_i} . Hence, after this selection, the only missing labels for an angle assignment are those for the vertices of $\text{sk}(\mu)$ inside the faces of \mathcal{S}_{μ} .

For $i = 1, \dots, h$, we select \mathcal{E}_{ν_i} to be any $u_i v_i$ -external upward planar embedding of G_{ν_i} with shape description s_i . This embedding exists, as our algorithm only consider shape descriptions in \mathcal{F}_{ν_i} for the component G_{ν_i} . It remains to select a $u_i v_i$ -external upward planar embedding \mathcal{E}_{ν_i} for each non-extreme boring component G_{ν_i} of G_{μ} and a label for each vertex of $\text{sk}(\mu)$ inside each face of \mathcal{S}_{μ} . This is done based on the solution for the flow network \mathcal{N} .

Let γ be a function that assigns to each arc $a \in A$ a value $\gamma(a)$ equal to the amount of flow traversing a in the solution for \mathcal{N} . By the integral flow theorem, we can assume that $\gamma(a)$ is a non-negative integer, for each arc $a \in A$. Similarly to [3], it can be proved that the total demand of the sinks in T is equal to the total amount of units of flow that can be supplied from the sources in S . This implies that each source sends out the maximum number of units of flow it can supply. Since the flow is integral and the only sources that can supply more than one unit of flow have a single incident arc, it follows that each source sends its flow on a single incident arc. This allows us to complete the construction of \mathcal{E}_{μ} as follows. Consider each non-extreme boring component G_{ν_i} of G_{μ} and its preferred set \mathcal{P}_{ν_i} .

- If \mathcal{P}_{ν_i} contains a single shape description, then we select \mathcal{E}_{ν_i} to be any $u_i v_i$ -external upward planar embedding of G_{ν_i} with that shape description.
- If \mathcal{P}_{ν_i} contains the hat and the inverted-hat, we select \mathcal{E}_{ν_i} to be any $u_i v_i$ -external upward planar embedding of G_{ν_i} with the former or the latter shape description, depending on whether the source b^i sends its unit of flow to the sink corresponding to the face $f_r(e_i)$ or to the sink corresponding to the face $f_l(e_i)$, respectively.
- If \mathcal{P}_{ν_i} contains both the heart and the inverted-heart, we select \mathcal{E}_{ν_i} to be any $u_i v_i$ -external upward planar embedding of G_{ν_i} with the former or the latter shape description,

depending on whether the unit of flow demanded by the sink t_i is supplied by a source corresponding to v_i or by a source corresponding to u_i , respectively.

- If \mathcal{P}_{ν_i} contains both the left-wing and the right-wing (the case in which it contains both the inverted-left-wing and the inverted-right-wing is analogous), then we further distinguish two cases.

v_i is a bottom vertex: then we select \mathcal{E}_{ν_i} to be any $u_i v_i$ -external upward planar embedding of G_{ν_i} whose shape description is the left-wing or the right-wing (the choice on whether to use one or the other shape description is arbitrary);

v_i is a top vertex: then we select \mathcal{E}_{ν_i} to be any $u_i v_i$ -external upward planar embedding of G_{ν_i} whose shape description is the left-wing or the right-wing, depending on whether the source corresponding to the non-switch vertex v_i sends its unit of flow to the sink corresponding to $f_l(e_i)$ or to the sink corresponding to $f_r(e_i)$, respectively.

We now have a planar embedding (which is not yet an upward planar embedding) of G_μ . For each vertex w of $\text{sk}(\mu)$ that is a non-switch vertex of G_μ , we label each angle at w in a face of \mathcal{S}_μ incident to precisely one incoming edge of w with the value 0, and we label every other angle at w in a face of \mathcal{S}_μ with the value -1 . For each vertex w of $\text{sk}(\mu)$ that is an unavailable switch vertex of G_μ , we label each angle at w in a face of \mathcal{S}_μ with the value -1 . For each vertex w of $\text{sk}(\mu)$ that is an available switch vertex of G_μ , let f be the unique face of \mathcal{S}_μ such that the source corresponding to w sends its unit of flow to the sink corresponding to f ; then we label the angle at w in f with the value 1 and we label every other angle at w in a face of \mathcal{S}_μ with the value -1 .

This completes the construction of the upward planar embedding \mathcal{E}_μ of G_μ . Since u and v are incident to the outer face of \mathcal{S}_μ , by construction, it follows that \mathcal{E}_μ is uv -external. We now prove that \mathcal{E}_μ is actually an upward planar embedding of G_μ , i.e., it satisfies Properties **UP0**–**UP3** of Theorem 1. Let λ be the angle assignment in \mathcal{E}_μ ; for each vertex w of G_μ and each face f of \mathcal{E}_μ , we define $n_{-1}(w)$, $n_0(w)$, $n_1(w)$, $n_{-1}(f)$, and $n_1(f)$ as before Theorem 1.

Property **UP0** (if α is a switch angle of \mathcal{E}_μ , then $\lambda(\alpha) \in \{-1, 1\}$, and if α is a flat angle, then $\lambda(\alpha) = 0$) is true for each angle α internal to \mathcal{E}_{ν_i} and for each angle α at a non-pole vertex of G_{ν_i} , for any $i \in \{1, \dots, k\}$, since \mathcal{E}_{ν_i} is an upward planar embedding. Further, it is true for each angle α at a vertex of $\text{sk}(\mu)$ inside a face of \mathcal{S}_μ , by construction.

Property **UP1** (if w is a switch vertex of G_μ , then $n_1(w) = 1$, $n_{-1}(w) = \deg(w) - 1$, $n_0(w) = 0$) is true for each non-pole switch vertex of G_{ν_i} , for any $i \in \{1, \dots, k\}$, since \mathcal{E}_{ν_i} is an upward planar embedding. Further, Property **UP1** is true for each available switch vertex w of $\text{sk}(\mu)$, as the only angle at w which is labeled 1 is the one in the face f of \mathcal{S}_μ such that the source in S corresponding to w sends its unit of flow to the sink in T corresponding to f ; indeed, every other angle at w in a face of \mathcal{S}_μ is labeled -1 , by construction, and every other angle at w in an internal face of an embedding \mathcal{E}_{ν_i} is labeled -1 since w is available. Moreover, Property **UP1** is true for each unavailable switch vertex w of $\text{sk}(\mu)$, as every angle at w in a face of \mathcal{S}_μ is labeled -1 , by construction, and there is exactly one angle at w in an internal face of an embedding \mathcal{E}_{ν_i} that is labeled 1, while all the others are labeled -1 , since w is unavailable. Finally, observe that a switch vertex is either available or unavailable, since the angle check does not fail.

Property **UP2** (if w is a non-switch vertex of G_μ , then $n_1(w) = 0$, $n_{-1}(w) = \deg(w) - 2$, $n_0(w) = 2$) is true for each non-switch vertex w of G_μ . Indeed, by Property **UP1** and since G is expanded, there are two angles at w labeled 0. Moreover, every other angle α at w is labeled -1 . This is by construction if α is in a face of \mathcal{S}_μ ; further, it is since \mathcal{E}_{ν_i} satisfies Property **UP2** and the angle check did not fail if α is in an internal face of \mathcal{E}_{ν_i} .

We prove Property **UP3** (if $f_{\mathcal{E}}$ is an internal face of \mathcal{E}_{μ} , then $n_1(f_{\mathcal{E}}) = n_{-1}(f_{\mathcal{E}}) - 2$, and if $f_{\mathcal{E}}$ is the outer face of \mathcal{E}_{μ} , then $n_1(f_{\mathcal{E}}) = n_{-1}(f_{\mathcal{E}}) + 2$). Consider an internal face $f_{\mathcal{E}}$ of \mathcal{E}_{μ} ; the outer face of \mathcal{E}_{μ} can be dealt with similarly. If $f_{\mathcal{E}}$ is also an internal face of \mathcal{E}_{ν_i} , for some $i \in \{1, \dots, k\}$, then we have $n_1(f_{\mathcal{E}}) = n_{-1}(f_{\mathcal{E}}) - 2$, given that \mathcal{E}_{ν_i} is an upward planar embedding. Otherwise, $f_{\mathcal{E}}$ corresponds to an internal face f of \mathcal{S}_{μ} . Let $E_1, E_2, E'_1, E'_2, n'_1(f_{\mathcal{E}}), n'_{-1}(f_{\mathcal{E}}), \tau_l^i, \tau_r^i$, and g be defined as in the proof of necessity.

A key observation is the following.

▷ **Claim 36.** If the number of small angles in $f_{\mathcal{E}}$ at non-pole vertices of the components G_{ν_i} of G_{μ} is χ , for some $\chi \geq 0$, then the number of large angles in $f_{\mathcal{E}}$ at non-pole vertices of the components G_{ν_i} of G_{μ} is equal to $\chi + \sum_{e_i \in E_1} \tau_l^i + \sum_{e_i \in E_2} \tau_r^i$.

Proof. For every edge $e_i \in E_1$, we have that, if the number of small angles in $f_{\mathcal{E}}$ at non-pole vertices of G_{ν_i} is equal to x_i , for some integer $x_i \geq 0$, then the number of large angles in $f_{\mathcal{E}}$ at non-pole vertices of G_{ν_i} is equal to $\tau_l^i + x_i$; this just comes from the definition of the left-turn-number of \mathcal{E}_{ν_i} . A similar relationship (with τ_r^i in place of τ_l^i) holds true for every edge $e_i \in E_2$. Then the claim follows with $\chi = \sum_{e_i \in E_1 \cup E_2} x_i$. ◀

By Claim 36, the number of large angles in $f_{\mathcal{E}}$ is equal to $\sum_{e_i \in E_1} \tau_l^i + \sum_{e_i \in E_2} \tau_r^i + \chi + n'_1(f_{\mathcal{E}})$, while the number of small angles in $f_{\mathcal{E}}$ is equal to $\chi + n'_{-1}(f_{\mathcal{E}})$. Hence, we want to prove that $\sum_{e_i \in E_1} \tau_l^i + \sum_{e_i \in E_2} \tau_r^i + n'_1(f_{\mathcal{E}}) - n'_{-1}(f_{\mathcal{E}}) = -2$.

The rest of the proof proceeds similarly to the proof of sufficiency. First, we have that the amount of flow sent to the sink t_f corresponding to f is equal to $\sum_{e_i \in E'_1} \tau_l^i + \sum_{e_i \in E'_2} \tau_r^i + n'_1(f_{\mathcal{E}}) - g$ (as in Claim 33). This is because:

- (a) we make the angle at a switch vertex of $\text{sk}(\mu)$ large if and only if the source corresponding to the switch vertex sends one unit of flow to t_f ;
- (b) t_f receives τ_l^i (τ_r^i) units of flow from an extreme or interesting component G_{ν_i} such that $e_i \in E_1$ (resp. $e_i \in E_2$) if and only if $\tau_l^i > 0$ (resp. $\tau_r^i > 0$);
- (c) for each edge $e_i \in E_1$ such that G_{ν_i} is non-extreme, boring, and non-ignored, we choose \mathcal{E}_{ν_i} in such a way that $\tau_l^i = 1$ if and only if t_f receives one unit of flow from a source corresponding to G_{ν_i} or corresponding to a non-switch vertex for which G_{ν_i} is the special component;
- (d) for each edge $e_i \in E_2$ such that G_{ν_i} is non-extreme, boring, and non-ignored, we choose \mathcal{E}_{ν_i} in such a way that $\tau_r^i = 1$ if and only if t_f receives one unit of flow from a source corresponding to G_{ν_i} or corresponding to a non-switch vertex for which G_{ν_i} is the special component; and
- (e) for each edge $e_i \in E_1$ ($e_i \in E_2$) such that G_{ν_i} is non-extreme, boring, and ignored, we have $\tau_l^i = 1$ (resp. $\tau_r^i = 1$), although this does not correspond to any unit of flow received by t_f ; this determines the $-g$ term in the amount of flow received by t_f .

Similar considerations on the value of n_f establish that $n_f = \sum_{e_i \in E_1} |\tau_l^i| + \sum_{e_i \in E_2} |\tau_r^i| + n'_1(f_{\mathcal{E}}) + n'_{-1}(f_{\mathcal{E}}) - 2g$ (as in Claim 34). Since the demand of t_f is equal to $\frac{n_f}{2} - 1$ and since the amount of flow received by t_f is equal to its demand, we get $\sum_{e_i \in E'_1} \tau_l^i + \sum_{e_i \in E'_2} \tau_r^i + n'_1(f_{\mathcal{E}}) - g = \frac{\sum_{e_i \in E_1} |\tau_l^i| + \sum_{e_i \in E_2} |\tau_r^i| + n'_1(f_{\mathcal{E}}) + n'_{-1}(f_{\mathcal{E}}) - 2g}{2} - 1$, which is $2 \sum_{e_i \in E'_1} \tau_l^i + 2 \sum_{e_i \in E'_2} \tau_r^i + 2n'_1(f_{\mathcal{E}}) = \sum_{e_i \in E'_1} \tau_l^i - \sum_{e_i \in E_1 \setminus E'_1} \tau_l^i + \sum_{e_i \in E'_2} \tau_r^i - \sum_{e_i \in E_2 \setminus E'_2} \tau_r^i + n'_1(f_{\mathcal{E}}) + n'_{-1}(f_{\mathcal{E}}) - 2$. This gives us $\sum_{e_i \in E_1} \tau_l^i + \sum_{e_i \in E_2} \tau_r^i + n'_1(f_{\mathcal{E}}) - n'_{-1}(f_{\mathcal{E}}) = -2$, as desired.

Finally, we need to prove that the shape description of \mathcal{E}_{μ} is s . The proof that the left-turn-number of \mathcal{E}_{μ} coincides with the first value of s , say τ_l , is very similar to the one we

just presented that $n_1(f_\varepsilon) = n_{-1}(f_\varepsilon) - 2$ (the main difference is that the demand of the sink t_l is equal to $\frac{n_f + \tau_l}{2}$ and this value is used in place of $\frac{n_f}{2} - 1$); this proof is hence omitted here. Likewise for the proof that the right-turn-number of \mathcal{E}_μ coincides with the second value of s . That the label of u in the outer face of \mathcal{E}_μ coincides with the third value of s , say λ^u , comes from the requirement that the source s_u corresponding to u must send one unit of flow to the sink t^u , if $\lambda^u = 1$, comes from the requirement that s_u must send one unit of flow to an internal face of \mathcal{S}_μ incident to u , if $\lambda^u = -1$, and comes from the extreme-edge check otherwise. A similar argument shows that the label of v in the outer face of \mathcal{E}_μ coincides with the fourth value of s . Finally, that the last four values in the shape description of \mathcal{E}_μ coincide with the last four values of s directly comes from the extreme-edge check. ◀

We are finally ready to state the main theorem of this section.

► **Theorem 37.** *It is possible to solve UPWARD PLANARITY in $\mathcal{O}(\sigma 1.45^\sigma \cdot n^2 \log^3 n)$ time for a digraph with n vertices and σ sources.*

Proof. Let G be an n -vertex digraph whose underlying graph is planar and let σ be the number of sources of G . First, we expand G , which is done in $\mathcal{O}(n)$ time. As discussed in Section 2, an expansion does not alter whether G is upward planar, it preserves the number of sources, and at most doubles the number of vertices of G . By Observation 26, each biconnected component B of G is expanded and has at most σ sources. For any edge e^* of B , the SPQR-tree T of B can be computed in $\mathcal{O}(n)$ time. By Lemma 27, for any R-node μ of T , the left- and right-turn-numbers of any uv -external upward planar embedding of the pertinent graph of μ are in $\mathcal{O}(\sigma)$. Further, by Lemma 32, there exists an R-node subprocedure whose total time complexity is $\mathcal{O}(\sigma 1.45^\sigma \cdot n \log^3 n)$. Hence, by Lemma 24, UPWARD PLANARITY can be solved in $\mathcal{O}(n((\sigma 1.45^\sigma \cdot n \log^3 n) + (\sigma^2 \cdot n))) \subseteq \mathcal{O}(\sigma 1.45^\sigma \cdot n^2 \log^3 n)$ time. ◀

7 An Algorithm Parameterized by Treewidth


The aim of this section is to provide an R-node subprocedure which yields parameterized algorithms for UPWARD PLANARITY when parameterized by treewidth and treedepth. The idea behind this is to obtain a combinatorization of the task that is asked in the subprocedure. This will be done by extending the skeleton of the R-node with additional information, notably via a so-called *embedding graph*. The R-node subprocedure is then obtained by performing dynamic programming over the embedding graph. However, to obtain the desired runtime, we will first have to show that the embedding graph has bounded treewidth.

It is worth noting that the notion of embedding graph used here shares similarities with the embedding graphs that have been recently used to solve several drawing extension problems [19, 20]. However, the notions are not the same due to problem-specific differences, and the use of the combinatorizations differ as well: while in the two aforementioned papers the combinatorizations were used in conjunction with Courcelle's Theorem [11], this approach cannot be used here due to the fact that we will need to make the turn numbers add up to -2 for each non-outer face.

7.1 A Combinatorial Representation of the Skeleton

Let G be a connected graph with a planar embedding \mathcal{G} , and let F be the set of faces of \mathcal{G} . Let G^- be the graph obtained from G by subdividing each edge e once, creating the vertex v_e . We define the *embedding graph* \tilde{G} of G as the graph obtained from G^- by adding a vertex f for each face in F , and connecting f to each vertex in G^- incident to f . Formally,

$V(\tilde{G}) = V(G) \cup F \cup \{v_e \mid e \in E(G)\} = V^v \cup V^f \cup V^e$, and $E(\tilde{G}) = E(G^-) \cup \{fv \mid v \in V(G) \wedge v \text{ is incident to } f\} \cup \{fv_e \mid e \in E(G) \wedge \text{both endpoints of } e \text{ are incident to } f\}$. Observe that \tilde{G} is tripartite, and we call the three sets of vertices that occur in the definition of $V(\tilde{G})$ the *true vertices*, *face-vertices* and *edge-vertices* of \tilde{G} , respectively. An illustration is shown in Fig. 14.



■ **Figure 14** (a) A planar graph G . (b) The embedding graph \tilde{G} of G . True-, face-, and edge-vertices are shown in black, green, and orange, respectively.

Furthermore, note that each edge-vertex is adjacent to at most two face-vertices.

It is well known (and easy to show) that if the treewidth of a graph G is at most k , then the treewidth of its subdivision is at most k [12, 18]. Our aim in this section, however, will be to show that if the treewidth of an embedded graph is bounded, then the treewidth of its embedding graph is also bounded. We begin by identifying a small number of faces that are, in some sense, “relevant” for a bag in a tree decomposition of G^- . We will then show that it is possible to construct a bounded-width decomposition of the embedding graph by adding the face-vertices of these relevant faces to the appropriate bags.

Given a tree-decomposition (T, χ) of G^- , a face $f \in F$ is called *crossing* for a node $t \in T$ if f is incident to at least one vertex in $fut(t)$ and to at least one vertex in $past(t)$. Our first task is to bound the number of crossing faces in a bag.

► **Lemma 38.** *If G has treewidth k , then there exists a tree-decomposition (T, χ) of width k such that there are at most $2k$ crossing faces for each node t of T .*

Proof. The lemma follows from previous work on so-called *geometric tree-decompositions* [17]. In particular, Dorn [17, Theorem 1] showed that if a plane graph G has treewidth k , there exists a tree-decomposition (T, χ) of G of width at most k with the additional property that for each node $t \in V(T)$, the bag $\chi(t)$ contains at most two vertex-subsets $S_1, S_2 \subseteq \chi(t)$ and there is a partitioning of $past(t)$ into $past_1(t), past_2(t)$ with the following properties:

- S_1, S_2 are minimal separators in G that separate $fut(t)$ from $past_1(t)$ and $past_2(t)$, respectively, and
- there are Jordan curves C_1, C_2 which intersect the drawing of G precisely in the vertices of S_1, S_2 , respectively, and separate $fut(t)$ from $past_1(t)$ and $past_2(t)$, respectively.

Given the above, consider such a geometric tree-decomposition (T, χ) of an arbitrary drawing that realizes the embedding of G^- . Each crossing face must, by definition, be intersected by at least one of the Jordan curves C_1, C_2 . Since each of these curves intersects at most k vertices and no edges, the total number of faces intersecting at least one of these curves is upper-bounded by $2k$. ◀

Let us now proceed with a tree-decomposition (T, χ) satisfying Lemma 38, and let $F(t)$ be the set of crossing faces for t . We say that a face is *inside* a bag t if every vertex on its boundary lies in t , and denote the set of faces inside t as $I(t)$.

► **Observation 39.** $|I(t)| \leq 2k - 4$.

Proof. Let H be the subgraph of G induced on $\chi(t)$. Since H is planar, by Euler's formula we obtain that it contains at most $2k - 4$ faces. Moreover, $I(t)$ is a subset of the faces of H . ◀

Finally, in order to obtain a valid tree-decomposition for \tilde{G} , we will need to add the set $F(t)$ of crossing faces also to all nodes adjacent to t (this will be important in cases where t is a forget node or when t 's parent is an introduce node). Formally, let the *propagation faces* $P(t)$ for $t \in T$ simply be defined as $\bigcup_{\{p \mid pt \in E(T)\}} F(p)$. Since a nice tree-decomposition has degree at most 3, it follows that $|P(t)| \leq 3 \cdot 2k$.

Let $\psi(t) = \chi(t) \cup F(t) \cup I(t) \cup P(t)$. While Lemma 38 together with Observation 39 means that adding the face-vertices for the crossing, inside and propagation faces to the relevant bags (i.e., replacing $\chi(t)$ with $\psi(t)$) would preserve a bound on the bag size, we need to show that this construction results in a valid tree-decomposition for \tilde{G} .

► **Lemma 40.** $(T, \psi(t))$ is a tree-decomposition of \tilde{G} .

Proof. To prove the lemma, it suffices to verify that (1) every edge incident to F occurs in some bag, and (2) for each face-vertex $f \in F$, the set of nodes of T satisfying $f \in \chi(t)$ forms a nonempty subtree of T . But first, we'll establish the following subclaim: for each face-vertex f , there exists a node t such that $f \in \psi(t)$.

Indeed, if all vertices adjacent to f in \tilde{G} occur together in some bag, say $\psi(t)$, then $f \in \psi(t)$ and the subclaim holds. Otherwise, let us consider a forget node t with a child p such that $\chi(p) \setminus \chi(t) = \{z\}$ where $zf \in E(\tilde{G})$ with the additional property that t achieves maximum distance from the root r (i.e., t is “as far away” from the root as possible). In particular, this implies that no other forget node which forgets a vertex adjacent to f occurs in the subtree rooted at t . Since $\chi(p)$ does not contain all the neighbors of f and no neighbor of f was forgotten in the subtree rooted at t , there must exist some neighbor of f , say y , in $\text{fut}(t)$. Since $z \in \text{past}(t)$, we have that $f \in F(t)$ and hence $f \in \psi(t)$, and the subclaim holds.

Let us now turn to establishing property (1). Again, if all vertices adjacent to f in \tilde{G} occur in some bag, then the property trivially holds. Otherwise, let v be an arbitrary vertex such that $vf \in E(\tilde{G})$; our aim is to show that $\{v, f\}$ occur together in some bag of (T, ψ) . Recall that there must exist some node in T where f occurs as a crossing vertex; let us pick one such node, say u . Consider the node t where v was forgotten. We distinguish three cases.

- u is a descendant of t : There must exist some vertex $w \in \text{past}(u)$ adjacent to f . Consider the unique u - t path in T , and let s' be the unique node on this path such that its parent s' introduces v . Since $v \in \text{fut}(s')$ and $w \in \text{past}(s')$, it follows that $f \in F(s')$. But then $\{f, v\} \subseteq P(s')$, as desired.
- u is an ancestor of t : There must exist some vertex $w \in \text{fut}(u)$ adjacent to f . But then $w \in \text{fut}(t)$ as well, and since $v \in \text{past}(t)$ we obtain that $f \in F(t)$. Then $\{f, v\} \subseteq P(t')$ where t' is the unique child of t , as desired.
- u is neither an ancestor nor a descendant of t : There must exist some vertex $w \in \text{past}(u)$ adjacent to f . But then $w \in \text{fut}(t)$, and since $v \in \text{past}(t)$ we obtain that $f \in F(t)$. Then $\{f, v\} \subseteq P(t')$ where t' is the unique child of t , as desired.

Finally, let us establish property (2). In particular, let t, u be nodes of T such that $f \in \psi(t) \cap \psi(u)$, and it suffices to show that the bag of every node on the t - u path in T contains f . Let us first consider the case where $f \in I(t)$. Since the intersection of connected subtrees on T is itself a connected subtree and each vertex v adjacent to f occurs on a connected subtree of T , it follows that the set of bags where $f \in I(t)$ also forms a connected subtree. Moreover, this implies that in this case there can be no node u such that $f \in F(u)$, and so the claim holds.

On the other hand, assume that $f \in F(t) \cup P(t)$. If $f \in P(t)$, then there exists a neighbor t' of t such that $f \in F(t')$ and the t' - u path contains the t - u path; hence it suffices to establish the case where $f \in F(t)$. Moreover, the previous paragraph established that in this case $f \in F(u) \cup P(u)$, and hence by the same argument as in the previous sentence it also suffices to consider the case where $f \in F(u)$. We now distinguish three cases.

- u is a descendant of t : By the definition of crossing faces, there must exist a vertex $a \in \text{past}(u)$ and a vertex $b \in \text{fut}(t)$ where $a \neq b$. But then each bag p on the u - t path in T satisfies that $a \in \text{past}(p)$ and $b \in \text{fut}(p)$, and in particular $f \in F(p)$.
- t is a descendant of u : This case is completely symmetrical to the previous one.
- u is neither an ancestor nor a descendant of t : By the definition of crossing faces, there must exist a vertex $a \in \text{past}(u)$ and a vertex $b \in \text{past}(t)$ where $a \neq b$. Let s be the least common ancestor of t and u . Each node p on the unique t - s path except for s satisfies that $b \in \text{past}(p)$ and $a \in \text{fut}(p)$, meaning that $f \in F(p)$. The same claim then holds symmetrically for the unique u - s path. Finally, since we showed that s has a neighbor p' satisfying $f \in F(p')$, we obtain $f \in P(s)$ as well. In particular, the bag of every node on the t - u path contains f . ◀

Lemma 38, Observation 39 and Lemma 40 together with the known fact that subdividing edges does not increase the treewidth [12, 18] immediately imply the following result, which we believe may be of general interest.

► **Theorem 41.** *Let G be a graph with a planar embedding of treewidth k where $k \geq 1$. Then the embedding graph \tilde{G} has treewidth at most $11k - 4 \in \mathcal{O}(k)$.*

7.2 Problem Reformulation

While the embedding graph defined in Subsection 7.1 is well-suited for our purposes, we still need to formulate the problem we will be solving on these embedding graphs. First of all, the R-node subprocedure required by Lemma 24 can be straightforwardly reduced to the task of checking whether a specific shape description ψ can be achieved at the R-node. This reduction takes time at most $\mathcal{O}(\tau)$ by Lemma 12. At this point, the input consists of (1) an R-node μ of T with skeleton H , (2) a mapping \mathcal{S}_μ which assigns each virtual edge in H to its feasible set, (3) a bound κ on the treewidth of the embedding graph \tilde{H} obtained from H , and (4) a target shape description ψ .

The main difficulty here—and the task of this subsection—will be to express the question of whether ψ is in the feasible set of μ as a combinatorial problem over the embedding graph \tilde{H} of H : Determine if there exists a *valid* angle mapping α and a shape selector β . We will need some additional terminology to elaborate on the meaning of “valid”, “angle mapping” and “shape selector”. Denote the poles of μ by \tilde{u}, \tilde{v} . Fix a $\tilde{u}\tilde{v}$ -external planar embedding \mathcal{E}_μ^H of H . Let \tilde{H} be the corresponding embedding graph, where $V(\tilde{H}) = V^v \cup V^e \cup V^f$. Let $F(H)$ denote the set of faces defined by \mathcal{E}_μ^H , we may assume that the outer face $f_H \in F(H)$ is fixed. For each edge $e \in E(H)$, denote by ν^e the respective child of the node μ in the used

SPQR decomposition, and recall that G_{ν^e} is then the corresponding pertinent graph. Let the set A_μ of active switch vertices be defined as the set of all switch vertices in G_μ which are contained in $V(H)$. Then:

- an angle mapping α maps each active switch vertex $v \in A_\mu$ to a vertex in $N_{\tilde{H}}(v)$; intuitively, this describes the placement of the large angle at v in the upward planar embedding of the pertinent graph (which could either be in a face between two virtual edges—in which case α maps v to the corresponding face vertex—or in a virtual edges—in which case α maps v to that virtual edge).
- a shape selector β maps each vertex v_e in V^e obtained from virtual edge e of H to a shape description that occurs in a feasible set in the range of $\mathcal{S}_\mu(e)$.

It remains to formalize the meaning of “valid”. For a true vertex $v \in V^\nu$ and an edge-vertex $e \in V^e$ adjacent to v , denote by $\lambda_\beta(v, v_e)$ the value assigned to the outer angle of v in G_{ν^e} by the shape description $\beta(v_e)$. For a true vertex $v \in V^\nu$ and an adjacent face vertex $v_f \in V^f$, denote by $\lambda_\alpha(v, v_f)$ the value that α induces for the angle at v inside v_f . That is, if $v \in A_\mu$, $\lambda_\alpha(v, v_f)$ is equal to 1 if $\alpha(v) = v_f$, and -1 otherwise. If $v \notin A_\mu$, $\lambda_\alpha(v, v_f)$ is equal to 0 if the corresponding angle is flat angle, and -1 otherwise, independently of α . For a face vertex $v_f \in V^f$ and an adjacent edge-vertex $v_e \in V^e$, denote by $\text{turn}_\beta(v_f, v_e)$ the turn number that $\beta(v_e)$ induces on the respective part of f . Specifically, if v_e corresponds to the component (G_{ν^e}, u, v) , then $\text{turn}_\beta(v_f, v_e)$ is the left-turn-number $\tau_l(G_{\nu^e}, u, v)$ if u appears right before v in the cyclic order induced by \mathcal{E}_μ^H on the vertices of H adjacent to v_f , and the right-turn-number $\tau_r(G_{\nu^e}, u, v)$ otherwise, where both $\tau_l(G_{\nu^e}, u, v)$ and $\tau_r(G_{\nu^e}, u, v)$ are defined by the shape $\beta(v_e)$. Analogously, for v adjacent to v_e define $\rho_\beta(v_f, v_e, v)$ as the value $\rho_l(G_{\nu^e}, v)$ or $\rho_r(G_{\nu^e}, v)$.

The pair (α, β) is valid if the following three Validity Conditions hold:

1. For each face vertex $f \in V^f$, we have that

$$\sum_{v_e \in V^e \cap N_{\tilde{H}}(v_f)} \text{turn}_\beta(v_f, v_e) + \sum_{v \in V^\nu \cap N_{\tilde{H}}(v_f)} \lambda_\alpha(v, v_f) = \begin{cases} 2, & \text{if } f = f_H \\ -2, & \text{otherwise.} \end{cases}$$

2. For each true vertex $v \in V^\nu$, the following conditions hold on the pole angles in the shapes selected by β for the incident edges:

- a. If v is a switch vertex in G_μ , for each $v_e \in V^e \cap N_{\tilde{H}}(v) \setminus \{\alpha(v)\}$, $\lambda_\beta(v, v_e) = 1$, and $\lambda_\beta(v, v_e) = -1$ if $v_e = \alpha(v)$. That is, in accordance with α , at most one of the virtual edges “takes” a large angle inside, meaning that the outside angle of this component at v is small.
- b. If v is a non-switch vertex in G_μ , for each $v_e \in V^e \cap N_{\tilde{H}}(v)$, we distinguish two cases. If the interior of G_{ν^e} does not contain a flat angle at v , then $\lambda_\beta(v, v_e) = 1$. Note that otherwise (i.e., if there is both an ingoing edge and an outgoing edge at v in G_{ν^e}), $\lambda_\beta(v, v_e) \in \{-1, 0\}$.

3. (α, β) agrees with the target shape ψ , which denotes the tuple $(\tau_l(G_\mu, \tilde{u}, \tilde{v}), \tau_r(G_\mu, \tilde{u}, \tilde{v}), \lambda(G_\mu, \tilde{u}), \lambda(G_\mu, \tilde{v}), \rho_l(G_\mu, \tilde{u}), \rho_r(G_\mu, \tilde{u}), \rho_l(G_\mu, \tilde{v}), \rho_r(G_\mu, \tilde{v}))$.

The circular order on the vertices and edges incident to f_H induces a right path $v'_0 = \tilde{u}$, $e'_1, v'_1, \dots, e'_h, v'_h = \tilde{v}$ and a left path $v_0 = \tilde{u}$, $e_1, v_1, \dots, e_\ell, v_\ell = \tilde{v}$ from \tilde{u} to \tilde{v} , where the left path is defined as the reverse of the corresponding path from \tilde{v} to \tilde{u} in the circular order. Let L_e and L_v contain all intermediate edge-vertices and true vertices on the left path, respectively. That is, $L_e = \{e_1, e_2, \dots, e_\ell\}$, $L_v = \{v_1, v_2, \dots, v_{\ell-1}\}$. For the right path, define analogously R_e and R_v . The following holds:

a. Left-turn and right-turn numbers match, that is,

$$\begin{aligned} \sum_{v_e \in L_e} \text{turn}_\beta(f_H, v_e) + \sum_{v_v \in L_v} \lambda_\alpha(v_v, f_H) &= \tau_l(G_\mu, \tilde{u}, \tilde{v}), \\ \sum_{v_e \in R_e} \text{turn}_\beta(f_H, v_e) + \sum_{v_v \in R_v} \lambda_\alpha(v_v, f_H) &= \tau_r(G_\mu, \tilde{u}, \tilde{v}). \end{aligned}$$

b. Angles at the poles match,

$$\begin{aligned} \lambda_\alpha(\tilde{u}, f_H) &= \lambda(G_\mu, \tilde{u}), \\ \lambda_\alpha(\tilde{v}, f_H) &= \lambda(G_\mu, \tilde{v}). \end{aligned}$$

c. Boundary edge directions at the poles match,

$$\begin{aligned} \rho_\beta(f_H, e_1, \tilde{u}) &= \rho_l(G_\mu, \tilde{u}), \\ \rho_\beta(f_H, e'_1, \tilde{u}) &= \rho_r(G_\mu, \tilde{u}), \\ \rho_\beta(f_H, e_\ell, \tilde{v}) &= \rho_l(G_\mu, \tilde{v}), \\ \rho_\beta(f_H, e'_\ell, \tilde{v}) &= \rho_r(G_\mu, \tilde{v}). \end{aligned}$$

We are now ready to prove that the problem of determining whether there exists a valid pair (α, β) is equivalent to checking whether ψ is in the feasible set of μ .

► **Lemma 42.** *There is an upward planar embedding of G_μ with the shape description ψ if and only if there is a valid pair (α, β) .*

Proof. In the forward direction, take the upward planar embedding \mathcal{E}_μ of G_μ that has shape description ψ and produce the corresponding (α, β) . For each children node ν of μ , consider the upward planar embedding \mathcal{E}_ν induced by \mathcal{E}_μ on G_ν which is a subgraph of G_μ . Let e be the edge of H that corresponds to ν , assign $\beta(e)$ to be the shape in $\mathcal{S}_\mu(e)$ that is defined by the embedding \mathcal{E}_ν . For each active switch vertex $v \in A_\mu$, define $\alpha(v)$ as follows. By Theorem 1, there is precisely one large angle at v in \mathcal{E}_μ . If this angle falls inside an inner face of G_{ν^e} for some virtual edge e , $\alpha(v)$ is set to be v_e . Otherwise, this angle belongs to one of the faces induced by a face f of H ; let $\alpha(v)$ be the corresponding vertex v_f . It remains to verify that α and β defined in this way form a valid pair.

1. By the condition **UP3** of Theorem 1, in \mathcal{E}_μ the angles belonging to a face sum up to two if the face is the outer face, and to -2 otherwise. In particular, this holds for any face that is induced by a face f of H . The sum of angle values on f is precisely the left part of the Validity Condition 1, since for each $v \in V(H)$, $\lambda_\alpha(v, v_f)$ is by construction equal to the value of this angle in \mathcal{E}_μ , and for each $e \in E(H)$ incident to v , $\text{turn}_\beta(v_f, v_e)$ is precisely the sum of angle values along the boundary walk of G_{ν^e} lying on f , by definition of a shape description.
2. Validity Condition 2 follows immediately from conditions **UP0–UP2** of Theorem 1 applied to \mathcal{E}_μ .
3. Validity Condition 3 boils down to verifying that the constructed pair (α, β) is set in accordance with the embedding \mathcal{E}_μ of G_μ that has shape description ψ : 3a is shown analogously to Validity Condition 1, 3b is immediate by the construction of α , in 3c the values of ρ on both sides are taken from the same embedding and thus necessarily match.

In the other direction, consider a valid pair (α, β) . First, fix a planar embedding \mathcal{E}_μ of G_μ based on (α, β) together with \mathcal{E}_μ^H : For each children component (G_{ν^e}, u, v) fix a

uv -external upward planar embedding that has the shape description given by $\beta(v_e)$ on the corresponding virtual edge v_e . In particular that fixes a planar embedding \mathcal{E}_{ν^e} of G_{ν^e} , and an angle assignment λ_{ν^e} . It remains to fix the circular order of the edges incident to the vertices of the skeleton H and the walk defining the boundary of the outer face. For the former, the embedding \mathcal{E}_μ^H defines a circular order on the set of virtual edges adjacent to each vertex v of H . On the other hand, the order of the edges incident to v inside each corresponding children component G_{ν^e} is given by the fixed planar embedding of G_{ν^e} . Together, this defines a circular order on all the edges of G_μ incident to v . Finally, the boundary walk of the outer face is also defined unambiguously from the outer face in \mathcal{E}_μ^H and the embeddings of the children components. Specifically, consider a sequence of virtual edges $v_{e_1}, \dots, v_{e_\ell}$ forming the boundary of the outer face in \mathcal{E}_μ^H . Obtain a walk in G_μ by replacing each v_{e_i} in the above sequence by the left- or right-walk from u_i to v_i around the outer face in the fixed embedding $\mathcal{E}_{\nu^{e_i}}$ of the component $(G_{\nu^{e_i}}, u_i, v_i)$, where the left-walk is taken if u_i appears right before v_i in the clockwise boundary walk of the outer face in $\mathcal{E}_{\nu^{e_i}}$, and the right-walk is taken otherwise. Clearly, \mathcal{E}_μ defined in this way is a valid planar embedding of G_μ .

Next, from (α, β) we construct an angle assignment λ for G_μ with the embedding \mathcal{E}_μ that satisfies Theorem 1, while simultaneously achieving the shape description ψ for $(G_\mu, \tilde{u}, \tilde{v})$. In what follows we specify how λ is defined for each pair of a vertex v and an incident face f . We distinguish the cases where $v \in V(H)$ and $v \notin V(H)$, and also the cases where f lies inside G_{ν^e} for some $e \in E(H)$, or f is induced by a face of H .

Case 1: If f lies inside G_{ν^e} for some $e \in E(H)$, $\lambda(v, f) = \lambda_{\nu^e}(v, f)$, that is, the value of the angle assignment is copied from the corresponding angle assignment on G_{ν^e} that yields shape description $\beta(v_e)$.

Case 2: If f is induced by a face f' of H , and v is a vertex of G_{ν^e} that does not belong to $V(H)$, we put $\lambda(v, f) = \lambda_{\nu^e}(v, f_e)$, where f_e is the external face in the embedding \mathcal{E}_{ν^e} of G_{ν^e} .

Case 3: If f is induced by a face f' of H and v is a vertex of H , we consider two cases. If v is an active vertex, we put $\lambda(v, f) = 1$ if $\alpha(v) = v_{f'}$ and $\lambda(v, f) = -1$ otherwise. If v is not an active vertex, $\lambda(v, f) = 0$ if the corresponding angle is flat, and $\lambda(v, f) = -1$ otherwise.

Next, we verify that λ defined in this way satisfies the conditions of Theorem 1. If $v \in V(G_{\nu^e})$, but not in $V(H)$, $\lambda(v, f)$ is either $\lambda_{\nu^e}(v, f)$ or $\lambda_{\nu^e}(v, f_e)$, and $\lambda_{\nu^e}(v, \cdot)$ satisfies **UP0** since $(\mathcal{E}_{\nu^e}, \lambda_{\nu^e})$ is an upward planar embedding of G_{ν^e} . Moreover, **UP1** and **UP2** are also automatically satisfied for v since the angle assignment around v is identical to that of λ_{ν^e} . Analogously, if f is a face inside G_{ν^e} , the condition **UP3** holds since the angle assignment for the angles of f is identical to that of λ_{ν^e} , which is a part of an upward planar embedding and thus satisfies **UP3**.

It remains to verify **UP0–2** for $v \in V(H)$ and **UP3** for each face f induced by a face of H . **UP0** follows immediately from the definition of λ , see **Case 3**. If v is a switch vertex of G_μ , v is an active vertex, and the mapping α picks a certain value for v . If $\alpha(v) = v_{f'}$ for some face f' of H , then $\lambda(v, f) = 1$ for the corresponding face f of G_μ , and $\lambda(v, f) = -1$ for all other faces of G_μ that are induced by the faces of H . Moreover, by Validity Condition 2a, for each virtual edge e incident to v in H , the outer angle of v in G_{ν^e} is large, thus every angle of v inside G_{ν^e} is small, meaning that every angle at v in G_μ is small, except for the angle at the face f . If $\alpha(v) = v_{e'}$ for some edge e' of H , by **Case 3** all the angles of form (v, f) , where f is a face induced by H , are small. Analogously, for each virtual edge e incident to v in H , the outer angle of v in G_{ν^e} is large, meaning that all angles at v inside

G_{ν^e} are small. Finally, for the angles at v inside $G_{\nu^{e'}}$, exactly one of them is assigned to large, as the angle assignment follows that of λ_{ν^e} , which satisfies **UP1**. Thus, **UP1** holds for v .

If v is in H and not a switch vertex of G_μ , by **Case 3** among the angles induced by faces of H , every flat angle is assigned 0, and every switch angle is assigned -1 . Consider a virtual edge e incident to v in H . If the corresponding component contains only edges of the same direction from v , by Validity condition 2b the outer angle of G_{ν^e} at v is large, and thus each angle at v inside G_{ν^e} is small. If, on the other hand, G_{ν^e} contains a flat angle at v , then either both flat angles are inside G_{ν^e} , the outer angle of G_{ν^e} at v is large, and all the other angles at v in G_{ν^e} (and thus in G_μ) are small. Or, the second flat angle at v occurs in a face adjacent to G_{ν^e} , the outer angle of G_{ν^e} is flat, and all the other angles are small. In either case, the Validity Condition 2b together with the fact that the angles assigned in G_{ν^e} by λ_{ν^e} satisfy **UP0–2**, ensure the fact that the condition **UP2** at v is satisfied.

For each face f of G_μ induced by a face f' of H , Validity Condition 1 of a valid pair immediately implies that **UP3** is satisfied. Specifically, for a vertex v in H , the angle of v is set exactly to $\lambda_\alpha(v, v_f)$. For a vertex v that is not in H , but inside a children component G_{ν^e} , the value of its angle is a part of the respective left- or right-turn number of G_{ν^e} , $\text{turn}_\beta(v_{f'}, v_e)$. Thus, the sum of angle values along f is precisely encoded in the left part of Validity Condition 1.

Finally, it remains to see that the $\tilde{u}\tilde{v}$ -external upward planar embedding $(\mathcal{E}_\mu, \lambda)$ of G_μ has shape description ψ . Analogously to verifying **UP3**, the left parts in Validity Condition 3a can be seen to encode the sum of angle values along the left and right boundary walk of G_μ , thus the turn numbers of G_μ are as required by ψ . Validity Condition 2b explicitly ensures that the outer pole angles of G_μ are in accordance with ψ . Finally, Validity Condition 3c verifies that the edges incident to the poles on the left and tight outer walk have the direction as specified by ψ . ◀

7.3 Finding Valid Pairs Using Treewidth

Subsections 7.1 and 7.2 together result in a combinatorial problem that we can solve in order to provide an R-node subprocedure. To provide a more careful analysis of the running time, we introduce a new measure ζ of the graph which, intuitively, bounds the maximum amount of spiraling that can occur for any face of the pertinent graph. Formally, ζ is the maximum over $n_1(f)$ and $n_{-1}(f)$ (see Theorem 1), over all faces f of all possible planar embeddings of the pertinent graph G_μ of μ . Recalling Proposition 5, we obtain:

► **Observation 43.** $\zeta \leq V(G_\mu)$, and moreover $\zeta \leq 2^{td(G_\mu)}$.

We can now design a dynamic program that solves the task at hand. The program computes sets of records for each node of a tree-decomposition in a leaf-to-root fashion, where each record is a tuple of the form **(angle, shape, score, left, right)** where **angle** and **shape** contain snapshots of α and β in the given bag, respectively; **score** keeps track of the sum of large and small angles for each face in the given bag; and **left, right** store information about the left- and right-turn-numbers of the outer face.

► **Lemma 44.** *There is an algorithm that runs in time $\zeta^{\mathcal{O}(tw(H))} \cdot (|V(H)| + |\mathcal{S}_\mu|)$ and either computes a valid pair, or correctly determines that no such pair exists.*

Proof. As our first step, we compute the embedding graph \tilde{H} of H and then use the well-known 5-approximation algorithm [5] to obtain a nice tree-decomposition (T, χ) of the embedding graph of width at most $5tw(H)$, which is in $\mathcal{O}(tw(H))$ by Lemma 41. We assume

that (T, χ) is rooted at the node r and that $\chi(r) = \emptyset$. Without loss of generality and for ease of presentation, we will alter this decomposition by adding the face-vertex v_H of the outer face to every bag (other than $\chi(r)$) and apply the standard conversion to nice tree-decompositions in order to obtain a nice tree-decomposition with the property that v_H is the first vertex introduced on each leaf and the last vertex forgotten just below the root. Let the width of the resulting tree-decomposition (T, χ) be k . To complete the proof, we provide a dynamic programming algorithm \mathcal{A} that processes T in a leaf-to-root fashion.

We begin by formalizing the records used by \mathcal{A} . For each node $t \in V(T)$, let V_t^v, V_t^e, V_t^f denote the intersection of $\chi(t)$ with V^v, V^e, V^f , respectively. At each node $t \in V(T)$, we will store a record set R_t which will be a set of tuples of the form $(\text{angle}, \text{shape}, \text{score}, \text{left}, \text{right})$ where:

- **angle** maps each switch vertex in $V_t^v \cap A_\mu$ to an element of $(V_t^e \cup V_t^f \cup \{\text{assigned}, \text{todo}\})$,
- **shape** maps each edge-vertex $w \in V_t^e$ to a shape description in a feasible set in $\mathcal{S}_\mu(w)$,
- **score** maps each face-vertex in V_t^f to an integer z such that $-\zeta \leq z \leq \zeta$, and
- **left, right** are integers whose absolute value is at most ζ .

Intuitively, for each node $t \in V(T)$ we will use R_t to store all the relevant combinatorial information about the behavior of a valid pair (α, β) of G w.r.t. the target shape description ψ in the subgraph induced on $\chi(T_t)$; let us call this subgraph H_t . To formalize the semantics of our records, we will first introduce a suitable projection of pairs to H_t : a t -pair is the restriction of pairs to $\chi(T_t)$, i.e., a pair of the form (α_t, β_t) where the domain (but not the range) of both mappings α_t and β_t is restricted to $\chi(T_t) = \text{past}(t) \cup \chi(t)$. A t -pair is *valid* if Validity Conditions 1, 2, 3b and 3c are satisfied for all vertices in $\text{past}(t)$ —in particular, (1) each face-vertex in $\text{past}(t)$ achieves the correct sum, (2) each true vertex in $\text{past}(t)$ satisfies the condition on its pole angles, and (3b,c) the behavior of the poles and the edges on the outer face incident to the poles in $\text{past}(t)$ match ψ . Note that Validity Condition 3a is not included in this definition, since it is a global condition that we will keep track of separately and check at the end.

Our records for each node t will capture information about valid t -pairs as follows: for each tuple $(\text{angle}, \text{shape}, \text{score}, \text{left}, \text{right})$ of the form described above, $(\text{angle}, \text{shape}, \text{score}, \text{left}, \text{right}) \in R_t$ if and only if there exists a valid t -pair (α_t, β_t) with the following properties:

- For each switch vertex $v \in V_t^v \cap A_\mu$ such that $\alpha_t(v) = w$,
 - **angle**(v) = w if and only if $w \in \chi(t)$,
 - **angle**(v) = **assigned** if and only if $w \in \text{past}(t)$, and
 - **angle**(v) = **todo** otherwise, i.e., if and only if $w \in \text{fut}(t)$.
- For each $v_e \in V_t^e$, **shape**(v_e) = $\beta_t(v_e)$.
- For each $v_f \in V_t^f$ representing a face f of H , **score**(v_f) is the intermediate sum for the term in Validity Condition 1; formally, **score**(v) = $\sum_{v_e \in V^e \cap N_{\tilde{H}}(v_f) \cap \chi(T_t)} \text{turn}_\beta(v_f, v_e) + \sum_{v \in V^v \cap N_{\tilde{H}}(v_f) \cap \chi(T_t)} \lambda_\alpha(v, v_f)$.
- **right** and **left** contain the intermediate sums of Validity Condition 3a, formally:

$$\begin{aligned} \sum_{v_e \in L_e \cap \chi(T_t)} \text{turn}_\beta(f_H, v_e) + \sum_{v_v \in L_v \cap \chi(T_t)} \lambda_\alpha(v_v, f_H) &= \text{left}, \\ \sum_{v_e \in R_e \cap \chi(T_t)} \text{turn}_\beta(f_H, v_e) + \sum_{v_v \in R_v \cap \chi(T_t)} \lambda_\alpha(v_v, f_H) &= \text{right}. \end{aligned}$$

Observe that since $H_r = H$ and $\chi(r) = \emptyset$, the set of valid r -pairs is precisely the set of pairs for the instance satisfying all Validity Conditions except for 3a. Hence, in view of Lemma 42, in order to determine whether there exists a valid pair it suffices to verify whether there is a valid r -pair satisfying Validity Condition 3a—i.e., whether there R_r contains a record such that $\text{left} = \tau_l(G_\mu, \tilde{u}, \tilde{v})$ and $\text{right} = \tau_r(G_\mu, \tilde{u}, \tilde{v})$.

Having defined the syntax and semantics of the records, we proceed to a description of the dynamic programming steps themselves—in particular, we describe how one can compute the record set at each of the four kinds of nodes in a nice tree-decomposition.

Leaf node. If $t \neq r$ is a leaf in T with $\chi(t) = \{v_H\}$, we recall that each leaf contains a single vertex v_H and we simply set R_t to $\{(\emptyset, \emptyset, v_H \mapsto 0, 0, 0)\}$.

Introduce node. If t is an introduce node with child t_{old} and $\chi(t) \setminus \chi(t_{\text{old}}) = \{v\}$, we compute R_t as follows.

If $v \in V^v$, then for each tuple $(\text{angle}, \text{shape}, \text{score}, \text{left}, \text{right}) \in R_{t_{\text{old}}}$ and each $w \in (N(v) \cap \chi(t)) \cup \{\text{todo}\}$, we construct the tuple $(\text{angle} \cup \{v \mapsto w\}, \text{shape}, \text{score}', \text{left}', \text{right}')$ and add it to R_t where

- **score'** is obtained from **score** by updating the values for each face $v_f \in \chi(t)$, i.e., by adding the value of $\lambda_\alpha(v, v_f)$ to each $\text{score}(v_f)$. Notably, if $v \in A_\mu$ then every face $f \in \chi(t)$ such that $\text{angle}(v) \neq f$ has its score decreased by 1 while the face $\text{angle}(v)$ has its score increased by 1 (the latter only applies if $\text{angle}(v) \neq \text{todo}$). On the other hand, if $v \notin A_\mu$ then every face in $\chi(t)$ has its score decreased by 1 for every small angle it has at v (which can be verified directly from \tilde{H} , regardless of the records).
- **left'** and **right'** are obtained from **left** and **right**, respectively, by updating the values of the intermediate sums on the left side in Validity Condition 3a based on the impact on the newly introduced vertex v . This update is only carried out if $v \in L_v$ or R_v , respectively, and is done by simply adding $\lambda_\alpha(v, f_H)$ to **score**.

If $v \in V^e$, then for each tuple $(\text{angle}, \text{shape}, \text{score}, \text{left}, \text{right}) \in R_{t_{\text{old}}}$ and each $\iota \in \mathcal{S}_\mu$, we construct the tuple $(\text{angle}', \text{shape} \cup \{v \mapsto \iota\}, \text{score}', \text{left}', \text{right}')$ where **score'**, **left'** and **right'** are updated based on the value of turn_β of the newly introduced shape description ι in the same way as in the previous case. Moreover, for each $v_v \in V_t^v \cap N(v)$ such that $\text{angle}(v_v) = \text{todo}$, we perform an additional branching to determine whether $\text{angle}'(v_v) = \text{todo}$ or $\text{angle}'(v_v) = v$; we set $\text{angle}' = \text{angle}$ for all vertices not satisfying the condition of this sentence. In each case, we verify whether Validity Condition 2 holds for the pair v_v, v_e ; if not, we discard the non-compliant choice of **angle'**, and otherwise we add the tuple $(\text{angle}', \text{shape} \cup \{v \mapsto \iota\}, \text{score}', \text{left}', \text{right}')$ to $R_{t_{\text{old}}}$.

If $v \in V^f$, then for each tuple $(\text{angle}, \text{shape}, \text{score}, \text{left}, \text{right}) \in R_{t_{\text{old}}}$ and each $v_v \in V_t^v \cap N(v)$ such that $\text{angle}(v_v) = \text{todo}$, we branch to determine whether $\text{angle}'(v_v) = \text{todo}$ or $\text{angle}'(v_v) = v$. In each branch, we construct the tuple $(\text{angle}', \text{shape}, \text{score}', \text{left}, \text{right})$ where **score'** matches **score** on all vertices other than v , and maps v to the intermediate sum of v obtained in an analogous way as in the previous two cases (in particular, we compute the sum $\text{turn}_\beta(v, v_e)$ over all $v_e \in V_t^e$ plus the sum $\lambda_\alpha(v_v, v)$ over all $v_v \in V_t^v$ based on the information in **angle'** and **shape**).

Join node. If t is a join node with children t_1 and t_2 , we compute R_t as follows. For each $\rho_1 = (\text{angle}_1, \text{shape}_1, \text{score}_1, \text{left}_1, \text{right}_1) \in R_{t_1}$ and each $\rho_2 = (\text{angle}_2, \text{shape}_2, \text{score}_2, \text{left}_2, \text{right}_2) \in R_{t_2}$, we perform two consistency checks:

- Check the consistency of **angle**₁ and **angle**₂ for each $v_v \in V_t^v \cap A_\mu$: if $\text{angle}_1(v_v) \in \chi(t)$ then $\text{angle}_1(v_v) = \text{angle}_2(v_v)$, if $\text{angle}_1(v_v) = \text{assigned}$ then $\text{angle}_2(v_v) = \text{todo}$, and

if $\text{angle}_1(v_v) = \text{todo}$ then $\text{angle}_2(v_v) \in \{\text{todo}, \text{assigned}\}$. We perform the analogous checks also for $\text{angle}_2(v_v)$. If any of these checks fail, we discard the current pair of (ρ_1, ρ_2) .

- Check the consistency of shape_1 and shape_2 for each $v_e \in V_t^e$: it must hold that $\text{shape}_1(v_e) = \text{shape}_2(v_e)$. If any of these checks fail, we discard the current pair of (ρ_1, ρ_2) .

If the above checks succeeded, we construct a new entry $(\text{angle}, \text{shape}, \text{score}, \text{left}, \text{right})$ and add it to R_t . This is carried out as follows:

1. For each $v_v \in V_t^v \cap A_\mu$ such that $\text{angle}_1(v_v) = \text{angle}_2(v_v)$, set $\text{angle}(v_v) = \text{angle}_1(v_v)$. For each $v_v \in V_t^v \cap A_\mu$ such that $\text{angle}_1(v_v) \neq \text{angle}_2(v_v)$, based on the previous checks we know that one was set to **assigned** while the other to **todo**; we hence set $\text{angle}(v_v) = \text{assigned}$.
2. For each $v_e \in V_t^e$, set $\text{shape}(v_e) = \text{shape}_1(v_e)$.
3. For each $v_f \in V_t^f$, set $\text{score}(v_f) = \text{score}_1(v_f) + \text{score}_2(v_f) - \text{corr}(v_f)$ where $\text{corr}(v_f)$ nullifies the double-counting of vertices in $V_t^v \cup V_t^e$ and is computed as $\sum_{v_e \in V_t^e \cap N_{\tilde{H}}(v_f)} \text{turn}_\beta(v_f, v_e) + \sum_{v \in V_t^v \cap N_{\tilde{H}}(v_f)} \lambda_\alpha(v, v_f)$ (where turn_β and λ_α are obtained from **shape** and **angle** in the same way as for the introduce nodes).
4. Similarly as for **score**, set $\text{left} = \text{left}_1 + \text{left}_2 - \text{leftcorr}$ and $\text{right} = \text{right}_1 + \text{right}_2 - \text{rightcorr}$. Here, leftcorr nullifies the double-counting of vertices in $\chi(t) \cap (L_v \cup L_e)$ and is equal to $\sum_{v_e \in L_e \cap \chi(t)} \text{turn}_\beta(f_H, v_e) + \sum_{v_v \in L_v \cap \chi(t)} \lambda_\alpha(v_v, f_H)$. The value of rightcorr is defined and computed analogously.

Forget node. If t is a forget node with child t_{old} and $\chi(t_{\text{old}}) \setminus \chi(t) = \{v\}$, we compute R_t as follows.

If $v \in V^v$, for each tuple $(\text{angle}, \text{shape}, \text{score}, \text{left}, \text{right}) \in R_{t_{\text{old}}}$ we construct a new tuple $(\text{angle}', \text{shape}, \text{score}, \text{left}, \text{right})$ where angle' is obtained from **angle** by omitting v from the domain. If v is a pole of H , we also check that Validity Conditions 3b and 3c are satisfied for each neighbor of v , and if not we discard the computed tuple. Otherwise, we add the tuple to R_t .

If $v \in V^e$, for each tuple $(\text{angle}', \text{shape}, \text{score}, \text{left}, \text{right}) \in R_{t_{\text{old}}}$ we construct a new tuple $(\text{angle}, \text{shape}', \text{score}, \text{left}, \text{right})$ where shape' is obtained from **shape** by omitting v from the domain and angle' is obtained from **angle** by mapping every vertex that **anglemapped** to v to **assigned** instead. If v is incident to a pole of H , we also check that Validity Condition 3c is satisfied. If it is, we add the tuple to R_t .

If $v \in V^f$, for each tuple $(\text{angle}, \text{shape}, \text{score}, \text{left}, \text{right}) \in R_{t_{\text{old}}}$ we construct a new tuple $(\text{angle}', \text{shape}, \text{score}', \text{left}, \text{right})$ where score' is obtained from **score** by omitting v from the domain and angle' is obtained from **angle** by mapping every vertex that **anglemapped** to v to **assigned** instead. If Validity Condition 1 is satisfied (i.e., if $\text{score}(v) = -2$ or, in the case of $v = f_h$, $+2$), we add the resulting tuple to R_t , and discard it otherwise.

Using the above procedures, we compute the record set of all nodes up to R_r , for which we check if there is a record such that $\text{left} = \tau_l(G_\mu, \tilde{u}, \tilde{v})$ and $\text{right} = \tau_r(G_\mu, \tilde{u}, \tilde{v})$. If yes, the algorithm outputs “Yes”, and otherwise it outputs “No”. This concludes the description of the algorithm.

Regarding the running time, the size of each record set is upper-bounded by $\zeta^{\mathcal{O}(tw)}$, and the procedures described in the computation of each type of node can be completed in time at most $\zeta^{\mathcal{O}(tw)}$ per node. The resulting runtime can hence be upper-bounded by $\zeta^{\mathcal{O}(tw)} \cdot V(H)$.

Correctness follows from the correctness of the computation of the record set for each node $t \in V(T)$, which can be verified directly from the definitions and the semantics of R_t . ◀

As an consequence of Lemma 44 together with Observation 43, Lemma 42 and Theorem 41, we obtain an R-node subprocedure that runs in XP-time parameterized by treewidth and fixed-parameter time parameterized by treedepth. Combining this with the Interface Lemma 24, we conclude:

► **Theorem 45.** *It is possible to solve UPWARD PLANARITY in time $n^{\mathcal{O}(tw(G))}$ and time $2^{\mathcal{O}(td(G)^2)} \cdot n^2$, where n is the number of vertices of the input digraph G .*

8 Concluding Remarks

The presented results show that the combination of SPQR-trees with parameterized techniques is a promising algorithmic tool for geometric graph problems. Indeed, for the case of upward planarity, our framework allows us to reduce the general problem to a similar one on 3-connected graphs, at which point it is possible to use parameter-specific approaches such as dynamic programming or flow networks to obtain a solution. We believe not only that the framework developed here can help obtain other algorithms for UPWARD PLANARITY, but that the idea behind the framework can be adapted to solve other problems of interest as well—a candidate problem in this regard would be constrained level planarity testing [8].

All algorithms and arguments given within this paper are constructive and can be extended to output an upward planar drawing for each yes-instance of UPWARD PLANARITY. An open problem is whether UPWARD PLANARITY is W[1]-hard when parameterized by treewidth, or fixed-parameter tractable. It seems that improving the current dynamic programming algorithm to a fixed-parameter one would require non-trivial and non-obvious insights, but the problem has so far also proven resilient to our attempts at obtaining a W[1]-hardness reduction. Another question is whether the fixed-parameter tractability of UPWARD PLANARITY parameterized by the number of sources can be lifted to parameterizing by the maximum turn number of a face in the final drawing.

References

- 1 Rémy Belmonte, Eun Jung Kim, Michael Lampis, Valia Mitsou, and Yota Otachi. Grundy distinguishes treewidth from pathwidth. In *28th Annual European Symposium on Algorithms, ESA 2020*, volume 173 of *LIPIcs*, pages 14:1–14:19. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- 2 Paola Bertolazzi, Giuseppe Di Battista, Carlo Mannino, and Roberto Tamassia. Optimal upward planarity testing of single-source digraphs. *SIAM J. Comput.*, 27(1):132–169, 1998.
- 3 Paola Bertolazzi, Giuseppe Di Battista, Giuseppe Liotta, and Carlo Mannino. Upward drawings of triconnected digraphs. *Algorithmica*, 12(6):476–497, 1994.
- 4 Bernhard Bliem, Michael Morak, Marius Moldovan, and Stefan Woltran. The impact of treewidth on grounding and solving of answer set programs. *J. Artif. Intell. Res.*, 67:35–80, 2020.
- 5 Hans L. Bodlaender, Pål Grønås Drange, Markus S. Dregi, Fedor V. Fomin, Daniel Lokshtanov, and Michal Pilipczuk. A $c^k n$ 5-approximation algorithm for treewidth. *SIAM J. Comput.*, 45(2):317–378, 2016.
- 6 Glencora Borradaile, Philip N. Klein, Shay Mozes, Yahav Nussbaum, and Christian Wulff-Nilsen. Multiple-source multiple-sink maximum flow in directed planar graphs in near-linear time. *SIAM J. Comput.*, 46(4):1280–1303, 2017.

- 7 Guido Brückner, Markus Himmel, and Ignaz Rutter. An SPQR-tree-like embedding representation for upward planarity. In Daniel Archambault and Csaba D. Tóth, editors, *27th International Symposium on Graph Drawing and Network Visualization, GD 2019*, volume 11904 of *Lecture Notes in Computer Science*, pages 517–531. Springer, 2019.
- 8 Guido Brückner and Ignaz Rutter. Partial and constrained level planarity. In Philip N. Klein, editor, *28th Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2017*, pages 2000–2011. SIAM, 2017.
- 9 Sergio Cabello and Christian Knauer. Algorithms for graphs of bounded treewidth via orthogonal range searching. *Comput. Geom.*, 42(9):815–824, 2009.
- 10 Hubert Y. Chan. A parameterized algorithm for upward planarity testing. In Susanne Albers and Tomasz Radzik, editors, *12th Annual European Symposium on Algorithms, ESA 2004*, volume 3221 of *Lecture Notes in Computer Science*, pages 157–168. Springer, 2004.
- 11 Bruno Courcelle. The monadic second-order logic of graphs. I. Recognizable sets of finite graphs. *Inf. Comput.*, 85(1):12–75, 1990.
- 12 Marek Cygan, Fedor V. Fomin, Lukasz Kowalik, Daniel Lokshantov, Dániel Marx, Marcin Pilipczuk, Michal Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015.
- 13 Giuseppe Di Battista, Peter Eades, Roberto Tamassia, and Ioannis G. Tollis. *Graph Drawing: Algorithms for the Visualization of Graphs*. Prentice-Hall, 1999.
- 14 Giuseppe Di Battista and Roberto Tamassia. On-line planarity testing. *SIAM J. Comput.*, 25(5):956–997, 1996.
- 15 Walter Didimo, Francesco Giordano, and Giuseppe Liotta. Upward spirality and upward planarity testing. *SIAM J. Discret. Math.*, 23(4):1842–1899, 2009.
- 16 Reinhard Diestel. *Graph Theory, 4th Edition*, volume 173 of *Graduate texts in mathematics*. Springer, 2012.
- 17 Frederic Dorn. Dynamic programming and planarity: Improved tree-decomposition based algorithms. *Discret. Appl. Math.*, 158(7):800–808, 2010.
- 18 Rodney G. Downey and Michael R. Fellows. *Fundamentals of Parameterized Complexity*. Texts in Computer Science. Springer, 2013.
- 19 Eduard Eiben, Robert Ganian, Thekla Hamm, Fabian Klute, and Martin Nöllenburg. Extending partial 1-planar drawings. In Artur Czumaj, Anuj Dawar, and Emanuela Merelli, editors, *47th International Colloquium on Automata, Languages, and Programming, ICALP 2020*, volume 168 of *LIPIcs*, pages 43:1–43:19. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- 20 Robert Ganian, Thekla Hamm, Fabian Klute, Irene Parada, and Birgit Vogtenhuber. Crossing-optimal extension of simple drawings. In Nikhil Bansal, Emanuela Merelli, and James Worrell, editors, *48th International Colloquium on Automata, Languages, and Programming, ICALP 2021*, volume 198 of *LIPIcs*, pages 72:1–72:17. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.
- 21 Robert Ganian, Petr Hliněný, Joachim Kneis, Daniel Meister, Jan Obdržálek, Peter Rossmanith, and Somnath Sikdar. Are there any good digraph width measures? *J. Comb. Theory, Ser. B*, 116:250–286, 2016.
- 22 Robert Ganian, Fabrizio Montecchiani, Martin Nöllenburg, and Meirav Zehavi. Parameterized Complexity in Graph Drawing (Dagstuhl Seminar 21293). *Dagstuhl Reports*, 11(6):82–123, 2021.
- 23 Robert Ganian and Sebastian Ordyniak. The complexity landscape of decompositional parameters for ILP. *Artif. Intell.*, 257:61–71, 2018.
- 24 Robert Ganian, M. S. Ramanujan, and Stefan Szeider. Combining treewidth and backdoors for CSP. In Heribert Vollmer and Brigitte Vallée, editors, *34th Symposium on Theoretical Aspects of Computer Science, STACS 2017, March 8-11, 2017, Hannover, Germany*, volume 66 of *LIPIcs*, pages 36:1–36:17. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2017.
- 25 Ashim Garg and Roberto Tamassia. On the computational complexity of upward and rectilinear planarity testing. In Roberto Tamassia and Ioannis G. Tollis, editors, *DIMACS International*

- Workshop on Graph Drawing, GD '94*, volume 894 of *Lecture Notes in Computer Science*, pages 286–297. Springer, 1994.
- 26 Ashim Garg and Roberto Tamassia. On the computational complexity of upward and rectilinear planarity testing. *SIAM J. Comput.*, 31(2):601–625, 2001.
 - 27 Carsten Gutwenger and Petra Mutzel. A linear time implementation of SPQR-trees. In Joe Marks, editor, *8th International Symposium on Graph Drawing, GD '00*, volume 1984 of *Lecture Notes in Computer Science*, pages 77–90. Springer, 2000.
 - 28 Patrick Healy and Karol Lynch. Two fixed-parameter tractable algorithms for testing upward planarity. *Int. J. Found. Comput. Sci.*, 17(5):1095–1114, 2006.
 - 29 John E. Hopcroft and Robert Endre Tarjan. Dividing a graph into triconnected components. *SIAM J. Comput.*, 2(3):135–158, 1973.
 - 30 Kristóf Huszár and Jonathan Spreer. 3-manifold triangulations with small treewidth. In Gill Barequet and Yusu Wang, editors, *35th International Symposium on Computational Geometry, SoCG 2019, June 18–21, 2019, Portland, Oregon, USA*, volume 129 of *LIPIcs*, pages 44:1–44:20. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2019.
 - 31 Michael D. Hutton and Anna Lubiw. Upward planar drawing of single source acyclic digraphs. In Alok Aggarwal, editor, *2nd Annual ACM/SIGACT-SIAM Symposium on Discrete Algorithms, SODA 1991*, pages 203–211. ACM/SIAM, 1991.
 - 32 Michael D. Hutton and Anna Lubiw. Upward planar drawing of single-source acyclic digraphs. *SIAM J. Comput.*, 25(2):291–311, 1996.
 - 33 T. Kloks. *Treewidth: Computations and Approximations*. Berlin, 1994.
 - 34 Jaroslav Nešetřil and Patrice Ossona de Mendez. *Sparsity - Graphs, Structures, and Algorithms*, volume 28 of *Algorithms and combinatorics*. Springer, 2012.
 - 35 Achilleas Papakostas. Upward planarity testing of outerplanar dags. In Roberto Tamassia and Ioannis G. Tollis, editors, *DIMACS International Workshop on Graph Drawing, GD '94*, volume 894 of *Lecture Notes in Computer Science*, pages 298–306. Springer, 1994.
 - 36 Neil Robertson and Paul D. Seymour. Graph minors. III. Planar tree-width. *J. Comb. Theory, Ser. B*, 36(1):49–64, 1984.
 - 37 Marko Samer and Stefan Szeider. Constraint satisfaction with bounded treewidth revisited. *J. Comput. Syst. Sci.*, 76(2):103–114, 2010.
 - 38 William T. Trotter and John I. Moore Jr. The dimension of planar posets. *J. Comb. Theory, Ser. B*, 22(1):54–67, 1977.