Asymptotic dimension of Artin groups and a new upper bound for Coxeter groups.

Panagiotis Tselekidis

March 14, 2022

Abstract

If $A_{\Gamma}(W_{\Gamma})$ is the Artin (Coxeter) group with defining graph Γ we denote by $Sim(\Gamma)$ the number of vertices of the largest clique in Γ . We show that $asdimA_{\Gamma} \leq Sim(\Gamma)$, if $Sim(\Gamma) = 2$. We conjecture that the inequality holds for every Artin group. We prove that if for all free of infinity Artin (Coxeter) groups the conjecture holds, then it holds for all Artin (Coxeter) groups. As a corollary, we show that $asdimW_{\Gamma} \leq Sim(\Gamma)$ for all Coxeter groups, which is the best known upper bound for the asymptotic dimension of Coxeter Groups. As a further corollary, we show that the asymptotic dimension of any Artin group of large type with $Sim(\Gamma) = 3$ is exactly two.

Contents

1	Introduction	1
2	Preliminaries.	4
3	The case $Sim(\Gamma) = 2$. 3.1 Graph-Groups	5 7
4	The main results.	8

1 Introduction

Artin groups are easily defined via presentations, like Coxeter groups, but they are often very poorly understood. There are numerous conjectures about them, for example, it is not known in general whether they are torsion free. While Artin groups remain much more mysterious than their Coxeter relatives, a driving theme behind current research has been to show that these groups are as well behaved as Coxeter groups.

Some examples of Artin groups are the free non-abelian groups, free abelian groups, and RAAGs. In recent years, Artin and Coxeter groups proved to be a very rich source of examples and counterexamples of interesting phenomena in geometric group theory.

In this paper we are interested in the geometry of Artin and Coxeter groups and in particular in the asymptotic dimension of these groups. The asymptotic dimension achieved particular prominence in geometric group theory after a paper of Guoliang Yu, (see [23]) which proved the Novikov higher signature conjecture for manifolds whose fundamental group has finite asymptotic dimension. It is known for many classes of groups that they have finite asymptotic dimension, for instance, hyperbolic, relative hyperbolic, hierarchically hyperbolic, mapping class groups of surfaces, and one relator groups have finite asymptotic dimension (see [5], [18], [1], [7], [17], [20]).

However, there are still many open questions regarding asymptotic dimension, for example,

Do all Artin groups have finite asymptotic dimension? [8] Do all $Out(F_n)$ groups have finite asymptotic dimension, when n > 2? [8]

Recently, it was proved in [14] that Artin groups being both of large and of hyperbolic type are hierarchically hyperbolic. As a consequence, these Artin groups have finite asymptotic dimension. However, no specific bound was obtained.

The purpose of this paper is to give some new results about the asymptotic dimension of Artin and Coxeter groups. To be more precise, let Γ be a finite simplicial labeled graph, we denote by A_{Γ} (W_{Γ}) the Artin group (Coxeter group) associated to the graph Γ . We set

 $Sim(\Gamma) = max\{n \mid \Gamma \text{ contains the 1-skeleton of the standard}$ $(n-1)\text{-simplex } \Delta^{n-1}\}.$

We conjecture that:

Conjecture 1.1. $asdim A_{\Gamma} \leq Sim(\Gamma)$, for all Artin groups.

Conjecture 1.2. $asdimW_{\Gamma} \leq Sim(\Gamma)$, for all Coxeter groups.

It is known that if A_{Γ} is a RAAG, then $asdimA_{\Gamma} = Sim(\Gamma)$ see ([20] and [22]). Since when $Sim(\Gamma) = 1$ the group A_{Γ} is a free group, $asdim(A_{\Gamma}) = 1$. We prove that:

Theorem 1.1. Let A_{Γ} be an Artin group with $Sim(\Gamma) = 2$. Then

 $asdimA_{\Gamma} = 2.$

Theorem 1.2. Let A_{Γ} be an Artin group of large type with $Sim(\Gamma) = 3$. Then

$$asdimA_{\Gamma} = 2.$$

It is known by an isometric embedding theorem of Januszkiewicz (see [21]) that Coxeter groups have finite asymptotic dimension. In particular, Januszkiewicz's theorem shows that for any Coxeter group W_{Γ} with defining graph Γ we have the following upper bound: $asdimW_{\Gamma} \leq \sharp V(\Gamma)$.

Dranishnikov notes that this can be improved to the following: $asdimW_{\Gamma} \leq ch(N(W_{\Gamma}))$. Where $N(W_{\Gamma})$ is the nerve of W_{Γ} and $ch(N(W_{\Gamma}))$ is the chromatic number of the 1-skeleton of $N(W_{\Gamma})$, we note that the 1-skeleton of $N(W_{\Gamma})$ is Γ . In general, $Sim(\Gamma) \leq ch(\Gamma)$.

Using theorem 1.1 we prove that:

Theorem 1.3. If for all free of infinity Artin (Coxeter) groups the conjecture 1.1 (conjecture 1.2) holds, then it holds for all Artin (Coxeter) groups.

We recall that an Artin (Coxeter) group is *free of infinity* if the corresponding graph is complete. Using theorem 1.3 we show that conjecture 1.2 is true.

Theorem 1.4. Let W_{Γ} be a Coxeter group. Then

 $asdimW_{\Gamma} \leq Sim(\Gamma).$

Since $Sim(\Gamma) \leq ch(N(W_{\Gamma}))$ and there are graphs such that the inequality is strict, theorem 1.4 improves the previously best known upper bound.

By a finite simplicial *G*-labeled graph we mean a finite simplicial graph Γ such that every edge [a, b] is labeled by a cyclically reduced word $r_{a,b}$ (in terms of letters-verices a, b). The word $r_{a,b}$ is neither of the form $b^k a b^{\lambda}$ nor $a^k b a^{\lambda}$.

The *Graph group* associated to a finite simplicial G-labeled graph Γ is the group G_{Γ} given by the following presentation:

 $G_{\Gamma} = \langle a, b | r_{a,b}$ when a, b are connected by an edge \rangle .

Obviously, Graph groups generalize Artin groups.

Proposition 1.5. Let Γ be a finite simplicial *G*-graph such that $Sim(\Gamma) = 2$. We further assume that $H_1(\Gamma, \mathbb{Z}) = 0$. Then

asdim $G_{\Gamma} \leq 2$.

2 Preliminaries.

The asymptotic dimension asdim X of a metric space X is defined as follows: $asdim X \leq n$ if and only if for every R > 0 there exists a uniformly bounded covering \mathcal{U} of X such that the R-multiplicity of \mathcal{U} is smaller than or equal to n + 1 (i.e. every R-ball in X intersects at most n + 1 elements of \mathcal{U}).

There are many equivalent ways to define the asymptotic dimension of a metric space. It turns out that the asymptotic dimension of an infinite tree is 1 and the asymptotic dimension of \mathbb{E}^n is n.

By a finite simplicial labeled graph we mean a finite simplicial graph Γ such that every edge [a, b] is labeled by a natural number $m_{ab} > 1$. The Artin group associated to Γ is the group A_{Γ} given by the following presentation:

$$A_{\Gamma} = \langle a, b \in V(\Gamma) | \underbrace{ab_{\cdots}}_{m_{ab}} = \underbrace{ba_{\cdots}}_{m_{ab}} \text{ when a, b are connected by an edge } \rangle.$$

Convention. We will use the same symbol m_{ab} both for the label of the edge [a, b] and the relation $\underbrace{ab \dots}_{m_{ab}} \cdot \underbrace{a^{-1}b^{-1}\dots}_{m_{ab}}$ of the group.

We call an Artin group Right-angled Artin group (RAAG) if $m_{ab} = 2$ when a, b are connected by an edge. We call an Artin group free of infinity if the corresponding graph is complete. An Artin group is of large type if every label m_{ab} is at least three.

The Coxeter group associated to Γ is the group W_{Γ} given by the following presentation:

 $W_{\Gamma} = \langle V(\Gamma) | a^2$ for all $a \in V(\Gamma)$ and $\underbrace{ab \dots}_{\substack{m_{ab} \\ m_{ab}}} = \underbrace{ba \dots}_{\substack{m_{ab} \\ m_{ab}}}$ when a, b are connected by an edge \rangle .

A Coxeter group is called *Right-angled Coxeter group* (RACG) if $m_{ab} = 2$ when a, b are connected by an edge. We call a Coxeter group *free of infinity* if the corresponding graph is complete.

We say a simplicial graph is *complete* or equivalently a *clique* if any two vetrices are connected by an edge. An *n*-*clique* is the complete graph on n vertices. There is another equivalent way to define $Sim(\Gamma)$, in particular, it can be viewed as the larger natural number *n* such that Γ contains a complete subgraph Γ' with *n* vertices.

We recall that the *full subgraph* defined by a subset V of the vertices of a graph Γ is a subgraph of Γ formed from V and from all of the edges of Γ which have both endpoints in the subset V.

The following theorem is proved by the author in [20]

Theorem 2.1. Let (\mathbb{G}, Y) be a finite graph of groups with vertex groups $\{G_v \mid v \in Y^0\}$ and edge groups $\{G_e \mid e \in Y^1_+\}$. Then the following inequality holds:

 $asdim \, \pi_1(\mathbb{G}, Y, \mathbb{T}) \leq max_{v \in Y^0, e \in Y^1_{\perp}} \{asdim G_v, asdim \, G_e + 1\}.$

We also need a theorem for free products from [6], see also [3].

Theorem 2.2. Let A, B be two finitely generated groups. Then:

 $asdim A * B = max\{asdim A, asdim B, 1\}.$

We recall also the following (see [20])

Theorem 2.3. Let G be a finitely generated one relator group. Then one of the following is true:

(i) G is finite cyclic, and asdim G = 0

(ii) G is a nontrivial free group or a free product of a nontrivial free group and a finite cyclic group, and asdim G = 1

(iii) G is an infinite freely indecomposable not cyclic group or a free product of a nontrivial free group and an infinite freely indecomposable not cyclic group, and asdim G = 2.

3 The case $Sim(\Gamma) = 2$.

Let Γ be a finite simplicial labeled graph, such that $Sim(\Gamma) = 2$. We distinguish two cases, either the graph contains a cycle or not. We will handle these cases separately.

Proposition 3.1. Let Γ be a finite simplicial labeled graph, such that $Sim(\Gamma) = 2$. We further assume that $H_1(\Gamma, \mathbb{Z}) = 0$, then

asdim
$$A_{\Gamma} = 2$$
 and asdim $W_{\Gamma} \leq 1$.

Proof. (Artin groups) We observe that if Γ is not connected, then $A_{\Gamma} = *_{i=1}^{n} A_{\Gamma_i}$, where Γ_i are the connected components of Γ. By the previous observation and theorem 2.2 we may consider only connected graphs. So we assume that Γ is connected, thus Γ is a tree.

We observe that if Γ is a tree, then A_{Γ} is the fundamental group of graph of groups where the vertex groups are one-relator parabolic subgroups of A_{Γ} and the edge groups are infinite cyclic parabolic subgroups. By theorem 2.1 and theorem 2.3 we have that $asdim A_{\Gamma} \leq 2.$.

Suppose that A_{Γ} is the artin group defined by a single edge. Obviously, A_{Γ} is one relator group. Since A_{Γ} is not a free group and it is torsion free (see [16] page 200), by theorem 2.3 we obtain that $asdim A_{\Gamma} = 2$. Since every Artin group with $Sim(\Gamma) = 2$ contains an one relator parabolic subgroup we conclude that the proposition is true.

(*Coxeter groups*) If Γ is a tree, then W_{Γ} is the fundamental group of graph of finite parabolic subgroups. By theorem 2.1 we have that $asdim W_{\Gamma} \leq 1$.

In the second case we have that the graph contains a cycle, however, it is possible to contain only cycles of length larger than or equal to four.

Theorem 3.2. Let Γ be a finite simplicial labeled graph, such that $Sim(\Gamma) = 2$. Then

asdim
$$A_{\Gamma} = 2$$
 and asdim $W_{\Gamma} \leq 2$.

Proof. Again we may assume that Γ is connected. We will use induction on the number n_C of cycles contained in the graph. If $n_C = 0$, then by proposition 3.1 the statement of this theorem is true. We assume that the statement is true for all $n_C \leq n$. Let Γ be a finite simplicial labeled graph such that $n_C = n + 1$. By the previous observation we know that the graph contains a cycle γ of length $m + 1 \geq 4$.

Let $\{v_1, \ldots, v_{m+1}\}$ be the vertex set of γ and $\{e_1, \ldots, e_{m+1}\}$ be the edges of the cycle where $e_i = [v_i, v_{i+1}]$. We set $\Gamma_1 = \Gamma \setminus \{v_1\}$, obviously this graph contains at least one cycle less than Γ , thus asdim $A_{\Gamma_1} = 2$ (asdim $W_{\Gamma_1} \leq 2$).

Let $U = \{u_1, \ldots, u_k\}$ be all the vertices of Γ such that for each u_j there exists an edge connecting it with v_1 . It is easy to observe that the

full subgraph Γ_U of Γ formed by U contains no edge. Thus the parabolic subgroup $A_{\Gamma_U}(W_{\Gamma_U})$ of $A_{\Gamma}(W_{\Gamma})$ is the free group with k generators F_k (is the virtually free group $*_{i=1}^k \mathbb{Z}_2$).

Let Γ_{v_1} be the full subgraph of Γ formed by $U \cup \{v_1\}$, we observe that contains no cycle. It is not hard to see that Γ_{v_1} contains at least an edge, thus $asdim A_{\Gamma_{v_1}} = 2$ (asdim $W_{\Gamma_{v_1}} \leq 1$.).

It is easy to check that the Artin (Coxeter) group $A_{\Gamma}(W_{\Gamma})$ is an amalgamated product of its parabolic subgroups $A_{\Gamma_1}(W_{\Gamma_1})$, $A_{\Gamma_U}(W_{\Gamma_U})$ and $A_{\Gamma_{v_1}}(W_{\Gamma_{v_1}})$. To be more precise, the following equalities are a direct consequence of the presentation of A_{Γ} and W_{Γ} :

$$A_{\Gamma} = A_{\Gamma_1} \underset{A_{\Gamma_U}}{*} A_{\Gamma_{v_1}} \text{ and } W_{\Gamma} = W_{\Gamma_1} \underset{W_{\Gamma_U}}{*} W_{\Gamma_{v_1}}$$

Then by theorem 2.1 (or using Dranishnikov's theorem (see [10])) we conclude that the theorem holds.

3.1 Graph-Groups.

In fact we may obtain a more general result than proposition 3.1.

By a finite simplicial *G*-labeled graph we mean a finite simplicial graph Γ such that every edge [a, b] is labeled by a cyclically reduced word $r_{a,b}$ (in terms of letters-verices a, b). The word $r_{a,b}$ is neither of the form $b^k a b^{\lambda}$ nor $a^k b a^{\lambda}$.

The *Graph group* associated to a finite simplicial G-labeled graph Γ is the group G_{Γ} given by the following presentation:

 $G_{\Gamma} = \langle a, b | r_{a,b}$ when a, b are connected by an edge \rangle .

Obviously, Artin groups are Graph groups.

Proposition 3.3. Let Γ be a finite simplicial G-graph, such that $Sim(\Gamma) = 2$. We further assume that $H_1(\Gamma, \mathbb{Z}) = 0$, then

$$asdim G_{\Gamma} \leq 2.$$

Proof. We assume that Γ is connected, thus Γ is a tree. Suppose that G_{Γ} is a graph group defined by a single edge. Obviously, G_{Γ} is one relator group. By theorem 2.3 we obtain that $\operatorname{asdim} G_{\Gamma} \leq 2$.

It is important to note that if a is a vertex of Γ , then $\langle a \rangle$ is an infinite cyclic subgroup of G_{Γ} . This follows by the Freiheitssatz (see [23], thm 5.1, page 198) and induction on the number of vertices. Since $Sim(\Gamma) = 2$, the graph contains at least an edge.

We observe that if Γ is a tree, then G_{Γ} is the fundamental group of graph of groups where the vertex groups are one-relator Graph subgroups of G_{Γ} and the edge groups are infinite cyclic subgroups. By theorem 2.1 and theorem 2.3 we have that $asdim G_{\Gamma} \leq 2.$

4 The main results.

Theorem 4.1. If for all free of infinity Artin (Coxeter) groups the conjecture 1.1 (conjecture 1.2) holds, then it holds for all Artin (Coxeter) groups.

Proof. We will show the theorem only for Artin groups since the proof for Coxeter groups is similar to that for Artin groups.

We observe that the free of infinity Artin groups are those defined by a complete graph. By theorem 2.2 it suffices to prove the statement only for Artin groups defined by a connected graph.

Let Γ be a finite simplicial connected labeled graph. We will use induction on $Sim(\Gamma)$. We distinguish two cases.

Case 1. The graph has at most two vertices.

Then A_{Γ} is either a free group or an one relator group. In the first subcase the asymptotic dimension is one and in the second subcase by theorem 3.2 the asymptotic dimension is two. In particular in both subcases the conjecture 1.1 holds.

Case 2. The graph has at least three vertices.

For $Sim(\Gamma) = 1$ or 2 the conjecture 1.1 true, we assume that the statement of this theorem is true for any $Sim(\Gamma) \leq n$, where $n \geq 2$. Let Γ be a finite simplicial connected labeled graph with $Sim(\Gamma) = n + 1$.

It is vital to use another induction here, in particular, we use induction on $n_{V(\Gamma)} = \sharp V(\Gamma)$. If $n_{V(\Gamma)} = Sim(\Gamma)$, then the graph is a complete graph thus the group A_{Γ} is free of infinity, so $asdimA_{\Gamma} \leq n + 1$. We assume that $asdimA_{\Gamma} \leq n + 1$ for every graph with $n_{V(\Gamma)} < N + 1$, we note that $N \geq Sim(\Gamma)$. Let Γ be a finite simplicial connected labeled graph with $Sim(\Gamma) = n + 1$ and $n_{V(\Gamma)} = N + 1$.

Let $\{C_1, \ldots, C_k\}$ be the complete subgraphs of Γ such that $Sim(C_i) = Sim(\Gamma)$, we will call these subgraphs maximal complete subgraphs. We set $n_C = \sharp\{C_1, \ldots, C_k\}$. We further distinguish two subcases.

Subcase 2.1 $n_C = 1$.

Let C be the only maximal complete subgraph of Γ . Since $n_{V(\Gamma)} > Sim(\Gamma)$, there exist vertices $v_1, v_2 \in C$ and $x \notin C$ such that there is no edge connecting v_1 with x. We set $\Gamma_1 = \Gamma \setminus \{v_1\}, \Gamma_x = \Gamma \setminus \{x\}$ and $\Gamma_{1x} = \Gamma \setminus \{v_1, x\}$

and we observe that all these graphs form the following parabolic subgroups of A_{Γ} , $A_{\Gamma_1} A_{\Gamma_x}$ and $A_{\Gamma_{1x}}$. As a direct consequence of the presentation of A_{Γ} we have that

$$A_{\Gamma} = A_{\Gamma_1} \underset{A_{\Gamma_{1x}}}{*} A_{\Gamma_x}. \tag{1}$$

It is easy to see that $Sim(\Gamma_1), Sim(\Gamma_{1x}) < n+1$, while $Sim(\Gamma_{1x}) = n+1$ but $n_{V(\Gamma_{1x})} < n_{V(\Gamma)} = N+1$. Thus $asdim A_{\Gamma_1}, asdim A_{\Gamma_{1x}} < n+1$ while $asdim A_{\Gamma_x} \le n+1$. Applying theorem 2.1 we conclude that $asdim A_{\Gamma} \le n+1$.

Subcase 2.2 $n_C > 1$.

This case is a bit more complicate than the previous, the goal is to write the A_{Γ} as an amalgamatated product, similar to the equality (1).

Recall that $S_C = \{C_1, \ldots, C_{n_C}\}$ is the set of the maximal complete subgraphs of Γ . Observe that there exists a vertex v_2 of C_2 and an edge $e = [v_0, v_1]$ of C_1 such that there is no edge between v_2 and v_1 . We will divide S_C into two disjoint subsets S_1 and S_2 , the first set consists of the maximal subgraphs containing v_1 and the second set containing the rest of the elements. After possibly relabeling the set S_C we may assume that $S_2 = \{C_2, \ldots, C_{\sigma}\}$. We fix an element v_i for every $C_i \in S_2$ such that there is no edge between v_1 and v_i (such elements exist and they are not equal to v_0).

We set $\Gamma_1 = \Gamma \setminus \{v_1\}$, $\Gamma_{\sigma} = \Gamma \setminus \{v_2, \ldots, v_{\sigma}\}$ and $\Gamma_{1\sigma} = \Gamma \setminus \{v_1, \ldots, v_{\sigma}\}$. We observe that all these graphs form the following parabolic subgroups of A_{Γ} , $A_{\Gamma_1} A_{\Gamma_{\sigma}}$ and $A_{\Gamma_{1\sigma}}$. As a direct consequence of the presentation of A_{Γ} we have that

$$A_{\Gamma} = A_{\Gamma_1} \underset{A_{\Gamma_1\sigma}}{*} A_{\Gamma_{\sigma}}.$$
 (2)

Observe that $Sim(\Gamma_{1\sigma}) < n+1$, while $Sim(\Gamma_{1}) = Sim(\Gamma_{\sigma}) = n+1$, however, $n_{V(\Gamma_{\sigma})}, n_{V(\Gamma_{1})} < n_{V(\Gamma)} = N+1$. So we obtain $asdim A_{\Gamma_{1\sigma}} < n+1$ while $asdim A_{\Gamma_{\sigma}}, asdim A_{\Gamma_{1}} \leq n+1$. By theorem 2.1 we conclude that $asdim A_{\Gamma} \leq n+1$.

This completes the proof that $asdim A_{\Gamma} \leq n+1$ in any subcase when $Sim(\Gamma) = n+1$, thus by induction the theorem holds.

Theorem 4.2. Let Γ be a finite simplicial labeled graph, and let W_{Γ} be the Coxeter group defined by Γ . Then

$$asdimW_{\Gamma} \leq Sim(\Gamma).$$

Proof. By an isometric embedding theorem of Januszkiewicz (see [21]) we have that $asdimW_{\Gamma} \leq \sharp V(\Gamma)$. Observe that if Γ is complete graph, then $\sharp V(\Gamma) = Sim(\Gamma)$. This means that the conjecture 1.2 is true for complete graphs.

Thus by theorem 4.1 we conclude that conjecture 1.2 is true for any Coxeter group.

As a corollary of theorem 4.1 we have:

Proposition 4.3. Let A_{Γ} be an Artin group of large type with $Sim(\Gamma) = 3$. Then

$$asdimA_{\Gamma} = 2$$

Proof. By [15] every Artin group of large type defined by a triangle splits as a graph of free groups, thus by theorem 2.1 $asdimA_{\Gamma} = 2$. By the proof of theorem 4.1 and the fact that every parabolic subgroup of a large type Artin group is also a large type Artin group, we conclude that the proposition is true.

References

- Jason Behrstock, Mark F. Hagen, and Alessandro Sisto. Asymptotic dimension and smallcancellation for hierarchically hyperbolic spaces and groups. Proc. Lond. Math. Soc. (3), 114(5):890–926, 2017
- [2] G Bell, A Dranishnikov, On asymptotic dimension of groups, Algebr. Geom. Topol. 1 (2001) 57-71 MR1808331
- [3] G.Bell, and A.Dranishnikov, On asymptotic dimension of groups acting on trees. Geometriae Dedicata, 103 (2004), 89-101
- G. C. Bell and A. N. Dranishnikov, Asymptotic dimension in Bedlewo, (2005) ArXiv:math.GR/0507570
- [5] G. Bell and A. Dranishnikov. Asymptotic dimension. Topology Appl., 155(12):1265-1296, 2008.
- [6] G. C. Bell, A. N. Dranishnikov, and J. E. Keesling, On a formula for the asymptotic dimension of free products, Fundamenta Mathematicae 183.1 (2004): 39-45.

- [7] Bestvina, Mladen, Ken Bromberg, and Koji Fujiwara, *Constructing group actions on quasi-trees and applications tomapping class groups*, Publications mathematiques de l'IHES 122.1 (2015): 1-64.
- [8] Site: https://sites.google.com/a/scu.edu/rscott/pggt, Problems in Geometric Group Theory, Asymptotic Dimension – Dranishnikov.
- [9] A. Dranishnikov, Cohomological approach to asymptotic dimension. Geom Dedicata 141, 59 (2009). https://doi.org/10.1007/s10711-008-9343-0
- [10] A. Dranishnikov, On asymptotic dimension of amalgamated products and right-angled Coxeter groups, Algebr. Geom. Topol. 8 (2008), no 3, 1281-1293
- [11] Elements de mathematique. Fasc. XXXIV. Groupes et algebres de Lie. Chapitre IV: Groupes de Coxeter et systemes de Tits. Chapitre V: Groupes engendres par des reflexions. Chapitre VI: Systemes de racines. Actualites Scientifiques et Industrielles, No. 1337, Hermann, Paris, 1968.
- [12] E.Godelle, L.paris. $K(\pi, 1)$ and word problem, for infinite type Artin-Tits groups, and applications to virtual braid groups. Preprint, arXiv:1007.1365.
- [13] M.Gromov, Asymptotic invariants of infinite groups, in Geometric Group Theory, v. 2, 1-295, London Math. Soc. Lecture Note Ser., vol. 182, Cambridge Univ. Press, Cambridge, 1993.
- [14] M.Hagen, A.Martin, A.Sisto, Extra-large type Artin groups are hierarchically hyperbolic (2021)arXiv:2109.04387
- [15] Kasia Jankiewicz, Residual finiteness of certain 2-dimensional Artin groups (2020) arXiv:2006.04717
- [16] R. Lyndon and P. Schupp, Combinatorial group theory, Springer-Verlag, 1977.
- [17] D. Matsnev, Asymptotic dimension of one relator groups, (2006) ArXiv:
- [18] Denis Osin, Asymptotic dimension of relatively hyperbolic groups, International Mathematics Research Notices, 2005, no. 35, 2143-2161

- [19] L. Paris. Universal cover of Salvetti's complex and topology of simplicial arrangements of hyperplanes. Trans. Amer. Math. Soc. 340 (1993), no. 1, 149–178.
- [20] Panagiotis Tselekidis, Asymptotic Dimension of Graphs of Groups and One Relator Groups, (2019), Arxiv: 1905.07925
- [21] Tadeusz Januszkiewicz, For Coxeter groups $z^{|g|}$ is a coefficient of a uniformly bounded representation, Fund. Math. 174 (2002), no. 1, 79–86. MR MR1925487 (2003f:20061)
- [22] N.Wright, Finite asymptotic dimension for CAT(0) cube complexes, Geom. Topol. 16 (2012), 527–554.
- [23] Guoliang Yu, The Novikov conjecture for groups with finite asymptotic dimension, Ann. of Math. (2) 147 (1998), no. 2, 325-355

E-mail: panagiotis.tselekidis@queens.ox.ac.uk *Address:* Mathematical Institute, University of Oxford, Andrew Wiles Building, Woodstock Rd, Oxford OX2 6GG, U.K.