

# Adjunctions between Eilenberg-Moore categories and a PBW-type theorem

Mamta Balodi\*

Abhishek Banerjee†

Anita Naolekar‡

## Abstract

Recently, Dotsenko and Tamaroff have shown that a morphism of  $T \rightarrow S$  of monads over a category  $\mathcal{C}$  satisfies the PBW-property if and only if it makes  $S$  into a free right  $T$ -module. We consider an adjunction  $\Psi = (G, F)$  between categories  $\mathcal{C}, \mathcal{D}$ , a monad  $S$  on  $\mathcal{C}$  and a monad  $T$  on  $\mathcal{D}$ . We show that a morphism  $\phi : (\mathcal{C}, S) \rightarrow (\mathcal{D}, T)$  that is well behaved with respect to the adjunction  $\Psi$  has a PBW-property if and only if it makes  $S$  satisfy a certain freeness condition with respect to  $T$ -modules with values in  $\mathcal{C}$ .

**MSC(2020) Subject Classification:** 16D90, 18C20

**Keywords:** Monads, PBW-theorem, Eilenberg-Moore category

## 1 Introduction

Let  $\mathcal{C}$  be a category and let  $\phi : T \rightarrow S$  be a morphism of monads on  $\mathcal{C}$ . Let  $\mathcal{C}^S$  and  $\mathcal{C}^T$  denote respectively the Eilenberg-Moore categories with respect to  $S$  and  $T$ . Then, if  $x \in \mathcal{C}$  carries the structure of an  $S$ -algebra, given by a structure map  $\lambda_x : Sx \rightarrow x$ , the composition  $Tx \xrightarrow{\phi x} Sx \xrightarrow{\lambda_x} x$  gives  $x$  the structure of a  $T$ -algebra. This determines a functor  $\phi_* : \mathcal{C}^S \rightarrow \mathcal{C}^T$ . Moreover, this functor  $\phi^*$  has a left adjoint

$$\phi^! : \mathcal{C}^T \rightarrow \mathcal{C}^S \quad (1.1)$$

which may be viewed as a “universal enveloping algebra.” In [3], Dotsenko and Tamaroff prove a striking result; they say that a morphism of monads on  $\mathcal{C}$  has the PBW property if the underlying object of the universal enveloping algebra of an object in  $\mathcal{C}^T$  does not depend on its  $T$ -algebra structure. Then, Dotsenko and Tamaroff [3] show that, if the Eilenberg-Moore categories contain reflexive coequalizers,  $\phi : T \rightarrow S$  satisfies the PBW-property if and only if  $\phi$  makes  $S$  into a free right  $T$ -module.

In our situation, we start with an adjunction  $\Psi = (G, F)$  between categories  $\mathcal{C}$  and  $\mathcal{D}$ . We consider a monad  $S$  on  $\mathcal{C}$  and a monad  $T$  on  $\mathcal{D}$ . By a  $\Psi$ -morphism  $\phi = (\phi_G, \phi_F) : (\mathcal{C}, S) \rightarrow (\mathcal{D}, T)$ , we mean a pair of natural transformations  $\phi_G : GT \rightarrow TS$  and  $\phi_F : TF \rightarrow FS$  satisfying certain condition that make the “morphism of monads” compatible with the adjunction  $\Psi = (G, F)$ . We then define a PBW-property similar to that of [3]. Our aim is to show that a  $\Psi$ -morphism has the PBW-property if and only if it satisfies a certain freeness condition with respect to  $T$ -modules with values in  $\mathcal{C}$ .

---

\*Stat-Math Unit, Indian Statistical Institute, Bangalore. Email: mamta.balodi@gmail.com

†Department of Mathematics, Indian Institute of Science, Bangalore. Email: abhishekbanerjee1313@gmail.com

‡Stat-Math Unit, Indian Statistical Institute, Bangalore. Email: anita.naolekar@gmail.com

## 2 Adjunctions between Eilenberg-Moore categories

We begin by fixing some notation. Suppose that  $(G, F)$  is a pair of adjoint functors between categories  $\mathcal{A}$  and  $\mathcal{B}$ . Accordingly, for objects  $a \in \mathcal{A}$  and  $b \in \mathcal{B}$ , we have an isomorphism

$$\mathcal{B}(Ga, b) \cong \mathcal{A}(a, Fb) \quad (2.1)$$

For a morphism  $g \in \mathcal{B}(Fa, b)$ , we denote by  $g^R$  the corresponding element in  $\mathcal{A}(a, Gb)$ . Similarly, for any  $f \in \mathcal{A}(a, Fb)$ , we will denote by  $f^L$  the corresponding element in  $\mathcal{B}(Ga, b)$ .

A monad on a category  $\mathcal{C}$  consists of an endofunctor  $S : \mathcal{C} \rightarrow \mathcal{C}$  along with natural transformations  $\mu(S) : S^2 \rightarrow S$  and  $\eta(S) : 1 \rightarrow S$  satisfying the usual associativity and unit conditions. We will denote this monad as a pair  $(\mathcal{C}, S)$ . A map of such pairs  $(\mathcal{C}, S) \rightarrow (\mathcal{D}, T)$  consists of a functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  along with a natural transformation  $\phi_F : TF \rightarrow FS$  such that the following diagrams commute

$$\begin{array}{ccc} T^2F & \xrightarrow{T\phi_F} TFS & \xrightarrow{\phi_FS} FS^2 \\ \mu(T)F \downarrow & & \downarrow F\mu(S) \\ TF & \xrightarrow{\phi_F} & FS \end{array} \quad \begin{array}{ccc} & F & \\ \eta(T)F \swarrow & & \searrow F\eta(S) \\ TF & \xrightarrow{\phi_F} & FS \end{array} \quad (2.2)$$

An  $S$ -algebra, or more precisely, an algebra over the monad  $(\mathcal{C}, S)$  consists of an object  $x \in \mathcal{C}$  along with a morphism  $\lambda_x : Sx \rightarrow x$  which satisfies associativity and unit conditions with respect to the action of  $S$ , i.e.,  $\lambda_x \circ \mu(S)x = \lambda_x \circ (S\lambda_x)$  and  $1_x = \lambda_x \circ \eta(S)x$ . The category of algebras over the monad  $(\mathcal{C}, S)$  is the Eilenberg-Moore category which we denote by  $\mathcal{C}^S$ . A morphism  $f : (x, \lambda_x) \rightarrow (x', \lambda_{x'})$  in  $\mathcal{C}^S$  is given by  $f : x \rightarrow x'$  such that  $\lambda_{x'} \circ Sf = f \circ \lambda_x$ . Given a monad  $\mathcal{C}^S$ , we also denote by  $\rho(S) : \mathcal{C} \rightarrow \mathcal{C}^S$  the free functor which is left adjoint to the forgetful functor  $\pi(S) : \mathcal{C}^S \rightarrow \mathcal{C}$ . The following result is well known.

**Lemma 2.1.** *Let  $(F, \phi_F) : (\mathcal{C}, S) \rightarrow (\mathcal{D}, T)$  be a morphism of monads. Then, there is a functor  $\hat{\phi}_F : \mathcal{C}^S \rightarrow \mathcal{D}^T$  that fits into the following commutative diagram*

$$\begin{array}{ccc} \mathcal{C}^S & \xrightarrow{\hat{\phi}_F} & \mathcal{D}^T \\ \pi(S) \downarrow & & \downarrow \pi(T) \\ \mathcal{C} & \xrightarrow{F} & \mathcal{D} \end{array} \quad (2.3)$$

*Proof.* We consider  $(y, \lambda_y) \in \mathcal{C}^S$ . It may be verified that the composition

$$\lambda_{Fy} : TFy \xrightarrow{\phi_F(y)} FSy \xrightarrow{F\lambda_y} Fy \quad (2.4)$$

gives  $Fx$  the structure of an algebra over  $(\mathcal{D}, T)$ . This defines a functor  $\hat{\phi}_F : \mathcal{C}^S \rightarrow \mathcal{D}^T$  that makes the diagram (2.3) commute.  $\square$

We now fix a pair  $\Psi = (G : \mathcal{D} \rightarrow \mathcal{C}, F : \mathcal{C} \rightarrow \mathcal{D})$  of adjoint functors between  $\mathcal{D}$  and  $\mathcal{C}$ . Our aim is to construct a commutative diagram of left adjoints corresponding to the commutative square of right adjoints in (2.3).

**Definition 2.2.** *Let  $(\mathcal{C}, S)$  and  $(\mathcal{D}, T)$  be monads and  $\Psi = (G, F)$  be an adjunction between  $\mathcal{D}$  and  $\mathcal{C}$ . A  $\Psi$ -morphism of monads is a pair  $\phi = (\phi_G, \phi_F)$  such that:*

(1)  $\phi_G : GT \rightarrow SG$  is a natural transformation which satisfies

$$\begin{array}{ccc} GT^2 & \xrightarrow{\phi_GT} SGT & \xrightarrow{S\phi_G} S^2G \\ G\mu(T) \downarrow & & \downarrow \mu(S)G \\ GT & \xrightarrow{\phi_G} & SG \end{array} \quad \begin{array}{ccc} G & \xrightarrow{G\eta(T)} & GT \\ \eta(S)G \searrow & & \downarrow \phi_G \\ & & SG \end{array} \quad (2.5)$$

(2)  $\phi_F : TF \longrightarrow FS$  is a natural transformation which satisfies

$$\begin{array}{ccc} T^2F & \xrightarrow{T\phi_F} & TFS \xrightarrow{\phi_F S} FS^2 \\ \mu(T)F \downarrow & & \downarrow F\mu(S) \\ TF & \xrightarrow{\phi_F} & FS \end{array} \quad \begin{array}{ccc} & F & \\ \eta(T)F \swarrow & & \searrow F\eta(S) \\ TF & \xrightarrow{\phi_F} & FS \end{array} \quad (2.6)$$

(3) For any morphism  $f : Gx \longrightarrow y$  in  $\mathcal{C}$ , the following two equivalent conditions are satisfied

$$\begin{array}{ccc} GTx & \xrightarrow{GT(f^R)} & GTFy \\ \phi_G(x) \downarrow & & \downarrow \phi_F(y)^L \\ SGx & \xrightarrow{Sf} & Sy \end{array} \Leftrightarrow \begin{array}{ccc} Tx & \xrightarrow{T(f^R)} & TFy \\ \phi_G(x)^R \downarrow & & \downarrow \phi_F(y) \\ FSGx & \xrightarrow{FSf} & FSy \end{array} \quad (2.7)$$

**Lemma 2.3.** Let  $\phi = (\phi_G, \phi_F) : (\mathcal{C}, S) \longrightarrow (\mathcal{D}, T)$  be a  $\Psi$ -morphism of monads. Let  $(x, \lambda_x) \in \mathcal{D}^T$ . Then, the following two compositions are identity

$$SGx \xrightarrow{SG\eta(T)x} SGTx \xrightarrow[\text{SG}\lambda_x]{(\mu(S)Gx) \circ (S\phi_G(x))} SGx \quad (2.8)$$

*Proof.* From the condition (2.5), we observe that

$$\begin{array}{ccc} Gx & \xrightarrow{G\eta(T)(x)} & GTx \\ & \searrow \eta(S)G(x) & \downarrow \phi_G(x) \\ & & SGx \end{array} \Rightarrow \begin{array}{ccc} SGx & \xrightarrow{SG\eta(T)(x)} & SGTx \\ & \searrow S\eta(S)G(x) & \downarrow S\phi_G(x) \\ & & SSGx \end{array} \quad (2.9)$$

Since  $(\mu(S)Gx) \circ (S\eta(S)Gx) = id$ , it follows from (2.9) that  $(\mu(S)Gx) \circ (S\phi_G(x)) \circ (SG\eta(T)x) = id$ . Additionally, since  $(x, \lambda_x)$  is an algebra over  $(\mathcal{D}, T)$ , we have  $(SG\lambda_x) \circ (SG\eta(T)x) = SG(\lambda_x \circ \eta(T)x) = id$ .  $\square$

From now onwards, we assume that the Eilenberg-Moore categories  $\mathcal{C}^S$  and  $\mathcal{D}^T$  both contain reflexive coequalizers. For more on this condition, we refer the reader to [1], [2, § 9.3] and [4].

**Lemma 2.4.** Let  $\phi = (\phi_G, \phi_F) : (\mathcal{C}, S) \longrightarrow (\mathcal{D}, T)$  be a  $\Psi$ -morphism of monads. Let  $(x, \lambda_x) \in \mathcal{D}^T$ . Then, the coequalizer

$$Coeq \left( SGTx \xrightarrow[\text{SG}\lambda_x]{(\mu(S)Gx) \circ (S\phi_G(x))} SGx \right) \quad (2.10)$$

in  $\mathcal{C}$  is equipped with the structure of an  $S$ -algebra. This determines a functor  $\hat{\phi}_G : \mathcal{D}^T \longrightarrow \mathcal{C}^S$ .

*Proof.* We observe that each of the objects in (2.8) is canonically equipped with the structure of an  $S$ -algebra and each arrow in (2.8) is a morphism of  $S$ -algebras. Applying Lemma 2.3, it follows that the coequalizer in (2.10) is a reflexive coequalizer. By assumption,  $\mathcal{C}^S$  contains reflexive coequalizers and the result follows.  $\square$

**Proposition 2.5.** Let  $\Psi = (G : \mathcal{D} \longrightarrow \mathcal{C}, F : \mathcal{C} \longrightarrow \mathcal{D})$  be a pair of adjoint functors between  $\mathcal{D}$  and  $\mathcal{C}$ . Let  $\phi = (\phi_G, \phi_F) : (\mathcal{C}, S) \longrightarrow (\mathcal{D}, T)$  be a  $\Psi$ -morphism of monads. Then,  $\hat{\phi}_G : \mathcal{D}^T \longrightarrow \mathcal{C}^S$  is left adjoint to the functor  $\hat{\phi}_F$  and fits into the following commutative diagram

$$\begin{array}{ccc} \mathcal{C}^S & \xleftarrow{\hat{\phi}_G} & \mathcal{D}^T \\ \rho(S) \uparrow & & \uparrow \rho(T) \\ \mathcal{C} & \xleftarrow{G} & \mathcal{D} \end{array} \quad (2.11)$$

*Proof.* We consider  $(x, \lambda_x) \in \mathcal{D}^T$ ,  $(y, \lambda_y) \in \mathcal{C}^S$  and a morphism  $\beta : x \rightarrow Fy$  which is a map of  $T$ -algebras. In particular,  $\beta \in \mathcal{D}(x, Fy)$ . Accordingly, we have a morphism  $\beta^L : Gx \rightarrow y$  in  $\mathcal{C}$ . From the coequalizer in (2.10) we see that in order to construct a morphism  $\alpha : \hat{\phi}_G(x) \rightarrow y$  we must show that

$$\lambda_y \circ S\beta^L \circ (\mu(S)Gx) \circ (S\phi_G(x)) = \lambda_y \circ S\beta^L \circ (SG\lambda_x) \quad (2.12)$$

Since  $\beta : x \rightarrow Fy$  is a morphism of  $T$ -algebras, the following diagrams commute

$$\begin{array}{ccc} Tx & \xrightarrow{T\beta} & TFy \xrightarrow{\phi_F(y)} FSy \\ \downarrow \lambda_x & & \swarrow F\lambda_y \\ x & \xrightarrow{\beta} & Fy \end{array} \quad \Rightarrow \quad \begin{array}{ccccc} GTx & \xrightarrow{GT\beta} & GTFy & \xrightarrow{\phi_F(y)^L} & Sy \\ \downarrow G\lambda_x & & & & \swarrow \lambda_y \\ Gx & \xrightarrow{\beta^L} & y & & \end{array} \quad (2.13)$$

Here the right hand side diagram follows from the left hand side diagram using the adjointness of  $(G, F)$ . Applying the functor  $S$  to the right hand side diagram in (2.13) and composing with  $\lambda_y$ , we obtain

$$\begin{array}{ccccccc} SGTx & \xrightarrow{SGT\beta} & SGTfy & \xrightarrow{S(\phi_F(y)^L)} & SSy & \xrightarrow{\mu(S)y} & Sy \\ \downarrow SG\lambda_x & & & & \downarrow S\lambda_y & & \downarrow \lambda_y \\ SGx & \xrightarrow{S\beta^L} & Sy & \xrightarrow{\lambda_y} & y & & \end{array} \quad (2.14)$$

Using the compatibility condition in (2.7) in Definition 2.2, we also have the following commutative diagram

$$\begin{array}{ccccc} SGTx & \xrightarrow{SGT\beta} & SGTfy & & \\ \downarrow S\phi_G(x) & & \downarrow S(\phi_F(y)^L) & & \\ SSGx & \xrightarrow{SS\beta^L} & SSy & & \\ \downarrow \mu(S)Gx & & \downarrow \mu(S)y & & \\ SGx & \xrightarrow{S\beta^L} & Sy & \xrightarrow{\lambda_y} & y \end{array} \quad (2.15)$$

Combining (2.14) and (2.15), the equality in (2.12) follows. It may be verified directly that  $\alpha : \hat{\phi}_G(x) \rightarrow y$  is a morphism of  $S$ -algebras. Conversely, if we have an  $S$ -algebra morphism  $\hat{\phi}_G(x) \rightarrow y$ , we have an induced map  $Gx \xrightarrow{\eta(S)Gx} SGx \rightarrow y$ , which corresponds to a morphism  $x \rightarrow Fy$ . Again, it may be verified directly that this is a  $T$ -algebra morphism and these two associations are inverses of each other.

Accordingly, we now have  $\hat{\phi}_G : \mathcal{D}^T \rightarrow \mathcal{C}^S$  which is left adjoint to  $\hat{\phi}_F : \mathcal{C}^S \rightarrow \mathcal{D}^T$ . Then,  $\hat{\phi}_G \circ \rho(T)$  is left adjoint to  $\pi(T) \circ \hat{\phi}_F$ . By Lemma 2.1, we have  $\pi(T) \circ \hat{\phi}_F = F \circ \pi(S)$  and hence the diagram (2.11) of left adjoints must commute.  $\square$

### 3 The Main theorem

We begin with the following definition.

**Definition 3.1.** Let  $\mathcal{C}, \mathcal{D}$  be categories and let  $(\mathcal{D}, T)$  be a monad. By a  $T$ -module taking values in  $\mathcal{C}$ , we will mean a pair  $(P, \nu(P))$  such that

(a)  $P : \mathcal{D} \rightarrow \mathcal{C}$  is a functor

(b)  $\nu(P) : PT \longrightarrow P$  is a natural transformation satisfying

$$\begin{array}{ccc} PT^2 & \xrightarrow{P\mu(T)} & PT \\ \nu(P)T \downarrow & & \downarrow \nu(P) \\ PT & \xrightarrow{\nu(P)} & P \end{array} \quad \begin{array}{ccc} P & \xrightarrow{P\eta(T)} & PT \\ & \searrow id & \downarrow \nu(P) \\ & & P \end{array} \quad (3.1)$$

A morphism  $\alpha : (P, \nu(P)) \longrightarrow (P', \nu(P'))$  of such  $T$ -modules consists of a natural transformation  $\alpha : P \longrightarrow P'$  of functors satisfying  $\nu(P') \circ (\alpha T) = \alpha \circ \nu(P)$ . We will denote by  $Mod_T^{\mathcal{C}}$  the category of  $T$ -modules taking values in  $\mathcal{C}$ .

**Proposition 3.2.** Let  $\mathcal{C}, \mathcal{D}$  be categories and let  $(\mathcal{D}, T)$  be a monad. There is a canonical functor

$$\Sigma_T^{\mathcal{C}} : Fun(\mathcal{D}, \mathcal{C}) \longrightarrow Mod_T^{\mathcal{C}} \quad P \mapsto (PT, P\mu(T)) \quad (3.2)$$

which is left adjoint to the forgetful functor  $Mod_T^{\mathcal{C}} \longrightarrow Fun(\mathcal{D}, \mathcal{C})$ .

*Proof.* Since  $(\mathcal{D}, T)$  is a monad, it is easy to verify that for any  $P \in Fun(\mathcal{D}, \mathcal{C})$ , the pair  $(PT, P\mu(T))$  satisfies the conditions in (3.1) for being a  $T$ -module with values in  $\mathcal{C}$ . It remains to show that for any  $(P', \nu(P')) \in Mod_T^{\mathcal{C}}$ , we have isomorphisms

$$Mod_T^{\mathcal{C}}((PT, P\mu(T)), (P', \nu(P'))) \cong Fun(\mathcal{D}, \mathcal{C})(P, P') \quad (3.3)$$

Indeed, given  $\alpha : P \longrightarrow P'$  in  $Fun(\mathcal{D}, \mathcal{C})$ , we have  $\alpha^L : (PT, P\mu(T)) \longrightarrow (P', \nu(P'))$  defined by setting

$$\alpha^L : PT \xrightarrow{\alpha T} P'T \xrightarrow{\nu(P')} P' \quad (3.4)$$

Conversely, for  $\beta : (PT, P\mu(T)) \longrightarrow (P', \nu(P'))$  in  $Mod_T^{\mathcal{C}}$ , we have the transformation  $\beta^R : P \longrightarrow P'$  given by

$$\beta^R : P \xrightarrow{P\eta(T)} PT \xrightarrow{\beta} P' \quad (3.5)$$

Because  $(\mathcal{D}, T)$  is a monad, it is clear that these two associations are inverse to each other.  $\square$

Following Proposition 3.2, we will say that  $(PT, P\mu(T))$  is the free  $T$ -module in  $\mathcal{C}$  associated to the functor  $P : \mathcal{D} \longrightarrow \mathcal{C}$ .

**Lemma 3.3.** Let  $\Psi = (G, F)$  be an adjunction and let  $\phi = (\phi_G, \phi_F) : (\mathcal{C}, S) \longrightarrow (\mathcal{D}, T)$  be a  $\Psi$ -morphism of monads. Then,  $SG : \mathcal{D} \longrightarrow \mathcal{C}$  is canonically equipped with the structure of a  $T$ -module taking values in  $\mathcal{C}$ .

*Proof.* We set  $\nu' : SGT \xrightarrow{S\phi_G} S^2G \xrightarrow{\mu(S)G} SG$ . In order to show that this determines a  $T$ -module, we note that the following diagram commutes.

$$\begin{array}{ccccc} SGT^2 & \xrightarrow{S\phi_G T} & SSGT & \xrightarrow{\mu(S)GT} & SGT \\ \downarrow SG\mu(T) & & \downarrow S\phi_G & & \downarrow S\phi_G \\ & & SSSG & \xrightarrow{\mu(S)SG} & SSG \\ & & \downarrow S\mu(S)G & & \downarrow \mu(S)G \\ SGT & \xrightarrow{S\phi_G} & S^2G & \xrightarrow{\mu(S)G} & SG \end{array} \quad (3.6)$$

In (3.6), the left hand square is obtained by applying  $S$  to the commutative square in (2.5). Using the triangle in (2.5), we also have the commutative diagram

$$\begin{array}{ccccc} & & SG & & \\ & \swarrow S\eta(T) & \downarrow S\eta(S)G & \searrow 1 & \\ SGT & \xrightarrow{S\phi_G} & SSG & \xrightarrow{\mu(S)G} & SG \end{array} \quad (3.7)$$

This proves the result.  $\square$

**Definition 3.4.** Let  $\Psi = (G, F)$  be an adjunction and let  $\phi = (\phi_G, \phi_F) : (\mathcal{C}, S) \longrightarrow (\mathcal{D}, T)$  be a  $\Psi$ -morphism of monads. We will say that  $\phi$  satisfies the PBW-property if there exists a functor  $Q : \mathcal{D} \longrightarrow \mathcal{C}$  which fits into the following commutative diagram

$$\begin{array}{ccc} \mathcal{D}^T & \xrightarrow{\hat{\phi}_G} & \mathcal{C}^S \\ \pi(T) \downarrow & & \downarrow \pi(S) \\ \mathcal{D} & \xrightarrow{Q} & \mathcal{C} \end{array} \quad (3.8)$$

**Proposition 3.5.** Let  $\Psi = (G, F)$  be an adjunction and let  $\phi = (\phi_G, \phi_F) : (\mathcal{C}, S) \longrightarrow (\mathcal{D}, T)$  be a  $\Psi$ -morphism of monads. Then, if  $\phi = (\phi_G, \phi_F)$  has the PBW property,  $\nu' : SGT \xrightarrow{S\phi_G} S^2G \xrightarrow{\mu(S)G} SG$  gives  $SG$  the structure of a free  $T$ -module in  $\mathcal{C}$ .

*Proof.* Since  $\phi$  has the PBW property, we have  $Q : \mathcal{D} \longrightarrow \mathcal{C}$  which fits into the commutative diagram (3.8). We first claim that  $SG = QT$ . Using (2.11) and (3.8), we have the following commutative diagram.

$$\begin{array}{ccccc} \mathcal{D} & \xrightarrow{\rho(T)} & \mathcal{D}^T & \xrightarrow{\pi(T)} & \mathcal{D} \\ G \downarrow & & \hat{\phi}_G \downarrow & & \downarrow Q \\ \mathcal{C} & \xrightarrow{\rho(S)} & \mathcal{C}^S & \xrightarrow{\pi(S)} & \mathcal{C} \end{array} \quad (3.9)$$

From (3.9), it is clear that for any  $x \in \mathcal{D}$ , we have  $QTx = SGx \in \mathcal{C}$ . It remains to show that  $QT = SG$  as objects of  $Mod_T^{\mathcal{C}}$ . For this, we note the following commutative diagram

$$\begin{array}{ccccccc} & & SGTy & & & & \\ & \nearrow \mu(S)GTy & & \searrow S\phi_Gy & & & \\ SSGTy & \xrightarrow{SS\phi_Gy} & SSSGy & \xrightarrow{\mu(S)SGy} & SSGy & & \\ \uparrow S\phi_GTy & & \downarrow S\mu(S)Gy & & \downarrow \mu(S)Gy & & \\ SGTTY & \xrightarrow{SG\mu(T)y} & SGTy & \xrightarrow{S\phi_G(T)y} & SSGy & \xrightarrow{\mu(S)Gy} & SGy \end{array} \quad (3.10)$$

which gives for any  $y \in \mathcal{D}$  the morphism from the coequalizer  $QTy = Q\pi(T)(Ty, \lambda_Ty) = \pi(S)\hat{\phi}_G(Ty, \lambda_Ty) \longrightarrow SGy$ . The condition (2.5) gives us the commutative diagram

$$\begin{array}{ccccccc} SGTTTY & \xrightarrow{SG\mu(T)Ty} & SGTTY & \xrightarrow{S\phi_GTy} & SSGTy & \xrightarrow{\mu(S)GTy} & SGTy \\ \downarrow SGT\mu(T)y & & \downarrow SG\mu(T)y & & \downarrow SS\phi_Gy & & \downarrow S\phi_Gy \\ & & & & SSSGy & \xrightarrow{\mu(S)SGy} & SSGy \\ & & & & \downarrow S\mu(S)Gy & & \downarrow \mu(S)Gy \\ SGTTy & \xrightarrow{SG\mu(T)y} & SGTy & \xrightarrow{S\phi_Gy} & SSGy & \xrightarrow{\mu(S)Gy} & SGy \end{array} \quad (3.11)$$

and also

$$\begin{array}{ccccccc} SGTTTY & \xrightarrow{S\phi_GTTY} & SSGTTY & \xrightarrow{\mu(S)GTTY} & SGTTY & \xrightarrow{S\phi_GTy} & SSGTy & \xrightarrow{\mu(S)GTy} & SGTy \\ \downarrow SGT\mu(T)y & & \downarrow SSG\mu(T)y & & \downarrow SG\mu(T)y & & \downarrow SS\phi_Gy & & \downarrow S\phi_Gy \\ & & & & & & SSSGy & \xrightarrow{\mu(S)SGy} & SSGy \\ & & & & & & \downarrow S\mu(S)Gy & & \downarrow \mu(S)Gy \\ SGTTy & \xrightarrow{S\phi_GTy} & SSGTy & \xrightarrow{\mu(S)GTy} & SGTy & \xrightarrow{S\phi_Gy} & SSGy & \xrightarrow{\mu(S)Gy} & SGy \end{array} \quad (3.12)$$

Putting together (3.11) and (3.12) and considering the coequalizers defining  $\hat{\phi}_G(Ty, \lambda_{Ty})$  and  $\hat{\phi}_G(TTy, \lambda_{TTy})$ , we see that  $QT = SG$  is compatible with the right action of  $T$  on both sides. This proves the result.  $\square$

**Proposition 3.6.** *Let  $\Psi = (G, F)$  be an adjunction and let  $\phi = (\phi_G, \phi_F) : (\mathcal{C}, S) \rightarrow (\mathcal{D}, T)$  be a  $\Psi$ -morphism of monads. Then,  $SG \in \text{Mod}_T^{\mathcal{C}}$  is isomorphic to a free  $T$ -module taking values in  $\mathcal{C}$ .*

*Proof.* We take  $Q : \mathcal{D} \rightarrow \mathcal{C}$  so that  $SG \cong QT$  in  $\text{Mod}_T^{\mathcal{C}}$ . We claim that  $Q\pi(T) = \pi(S)\hat{\phi}_G$ . Using the definition of  $\hat{\phi}_G$  in (2.10), this means that we must show that the coequalizer

$$\text{Coeq} \left( SGTx \xrightarrow[\text{SG}\lambda_x]{(\mu(S)Gx) \circ (S\phi_G(x))} SGx \right) \cong Q(x) \quad (3.13)$$

in  $\mathcal{C}$  for any  $T$ -algebra  $(x, \lambda_x) \in \mathcal{D}^T$ . Since  $SG \cong QT$  as  $T$ -modules, it follows that the composition  $SGT \xrightarrow{S\phi_G} SSG \xrightarrow{\mu(S)G} SG$  giving the  $T$ -module structure on  $SG$  corresponds to the composition  $QTT \xrightarrow{Q\mu(T)} QT$  giving the  $T$ -module structure on  $QT$ . Accordingly, we can prove (3.13) by showing that

$$\text{Coeq} \left( QTTx \xrightarrow[\text{QT}\lambda_x]{Q\mu(T)x} QT x \right) \cong Q(x) \quad (3.14)$$

We will prove this by showing that the following is a split coequalizer diagram in  $\mathcal{C}$ :

$$QT x \xrightarrow{\epsilon = Q\eta(T)Tx} QTTx \xrightarrow[\delta = QT\lambda_x]{\gamma = Q\mu(T)x} QT x \xleftarrow[\alpha = Q\lambda_x]{\beta = Q\eta(T)x} Qx \quad (3.15)$$

Since  $(x, \lambda_x)$  is a  $T$ -algebra, we see that  $\alpha\gamma = \alpha\delta$ . Since  $T$  is a monad, we get  $\alpha\beta = id$  and  $\gamma\epsilon = id$ . We also observe directly that  $\delta\epsilon = \beta\alpha$ . We conclude that  $Qx = \text{Coeq}(\gamma, \delta)$ , which proves the result.  $\square$

**Theorem 3.7.** *Let  $\Psi = (G, F)$  be an adjunction and let  $\phi = (\phi_G, \phi_F) : (\mathcal{C}, S) \rightarrow (\mathcal{D}, T)$  be a  $\Psi$ -morphism of monads. Then, the following are equivalent.*

(1) *There exists a functor  $Q : \mathcal{D} \rightarrow \mathcal{C}$  such that there are isomorphisms in  $\mathcal{C}$*

$$\hat{\phi}_G(x, \lambda_x) = Q(x) \quad \forall (x, \lambda_x) \in \mathcal{D}^T \quad (3.16)$$

*natural with respect to morphisms in  $\mathcal{D}^T$ .*

(2) *The functor  $SG$ , along with the map  $\nu' : SGT \xrightarrow{S\phi_G} S^2G \xrightarrow{\mu(S)G} SG$ , is isomorphic to a free  $T$ -module with values in  $\mathcal{C}$ .*

*Proof.* (1)  $\Rightarrow$  (2) follows from Proposition 3.5 and (2)  $\Rightarrow$  (1) follows from Proposition 3.6.  $\square$

## References

- [1] J. Adámek and V. Koubek, *Are colimits of algebras simple to construct?*, J. Algebra **66** (1980), no. 1, 226–250.
- [2] M. Barr and C. Wells, *Toposes, triples and theories*, Grundlehren der mathematischen Wissenschaften, vol. 278, Springer-Verlag, New York, 1985.
- [3] V. Dotsenko and P. Tamaroff, *Endofunctors and Poincaré-Birkhoff-Witt theorems*, Int. Math. Res. Not. **16** (2021), 12670–12690.
- [4] F. E. J. Linton, *Coequalizers in categories of algebras*, Sem. on Triples and Categorical Homology Theory (ETH, Zürich, 1966/67), Springer, Berlin, 1969, pp. 75–90.