

Finite temperature quantum condensations in the space of states

Massimo Ostilli¹ and Carlo Presilla^{2,3}

¹*Instituto de Física, Universidade Federal da Bahia, Salvador, 40210-340, BA, Brazil*

²*Dipartimento di Fisica, Sapienza Università di Roma, Piazzale A. Moro 2, Roma 00185, Italy*

³*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, Roma 00185, Italy*

(Dated: October 6, 2022)

If an isolated quantum system admits a partition of its space of states \mathcal{H} into two subspaces, $\mathcal{H}_{\text{cond}}$ and $\mathcal{H}_{\text{norm}}$, such that, in the thermodynamic limit, $\dim \mathcal{H}_{\text{cond}} / \dim \mathcal{H} \rightarrow 0$ and the ground state energies of the system restricted to these subspaces cross each other for some value of the Hamiltonian parameters, then, the system undergoes a first-order quantum phase transition driven by that parameter. A proof of this general class of phase transitions, which represent a condensation in the space of quantum states, was recently provided at zero temperature. It is reasonable to extend the above condensation quantum phase transitions to finite temperature by substituting the ground state energies with the corresponding free energies. Here, we illustrate this criterion in two different systems. We derive, analytically, the phase diagram of the paradigmatic Grover model and, numerically, that of a system of free fermions in a one-dimensional inhomogeneous lattice, where the condensation realizes as a spatial condensation. These phase diagrams are structurally similar, in agreement with the universal features of the present class of phase transitions. Finally, we suggest an experimental realization of the fermionic system in terms of heterostructure superlattices.

Quantum phase transitions (QPTs), i.e. the thermodynamic singularities emerging at zero temperature ($T = 0$) driven by some Hamiltonian parameter of the system, originate from quantum fluctuations, consequence of Heisenberg's uncertainty principle. However, an isolated system at $T = 0$ represents an abstract limit and understanding the finite temperature counterpart of a QPT (if any) is of paramount importance. Such an aim represents a quite challenging issue, from both the theoretical and experimental viewpoints since, above zero temperature, quantum and thermal fluctuations may compete in an intricate manner. Put in simple terms, in general, two kinds of scenarios are expected [1–6]: the ordered phase exists only at zero temperature; the ordered phase exists also at finite temperature, below some critical value T_c which, in turn, might signal a purely classical phase transition when the Hamiltonian parameters are set to a value rendering the system classical (e.g., the Ising model in a transverse field becomes the classical Ising model when the transverse field is set to zero). This second scenario is particularly appealing for potential applications to quantum computing protocols aimed at finding the ground state (GS) by working at finite temperature, the GS of the ordered phase being the solution of some (classical) combinatorial problem of interest, possibly hard [7–9]. Intuitively, first-order QPTs, a much less explored field when compared to second-order QPTs, might favor this scenario since, at the transition, the order parameter jumps between two very different quantum states.

Recently, we have introduced a class of first-order QPTs taking place via condensation in the space of states [10]. Let us consider a system described by the Hamiltonian

$$H = \Gamma K + JV, \quad (1)$$

where K and V are two noncommuting dimensionless Hermitian operators, and Γ and J parameters with energy dimensions. Representing H in the eigenbasis of V , it is natural to call V potential operator, K hopping operator, Γ hopping pa-

rameter and J potential strength. We will use, equivalently, Γ or J as the control parameter of the supposed QPT. Since phase transitions occur in the thermodynamic limit (TDL), we need a fair competition between K and V in this limit. By this we mean that, supposing that H describes a system of N particles, the eigenvalues of K and V both scale linearly with N . An important family of models to bear in mind is a collection of N qubits (spins). For these systems, the space of states \mathcal{H} can be identified with the space spanned by the $M = 2^N$ spin states indicated by $|\mathbf{n}\rangle = |n_1\rangle|n_2\rangle \dots |n_N\rangle$, where $|n_i\rangle = |\pm\rangle$ is an eigenstate of the Pauli matrix σ_i^z relative to the qubit $i = 1, \dots, N$. The potential V is a diagonal operator in the states $|\mathbf{n}\rangle$, namely, $V = \sum_{\mathbf{n}} V_{\mathbf{n}} |\mathbf{n}\rangle \langle \mathbf{n}|$. The hopping operator K is chosen as the sum of single-flip operators $K = -\sum_{i=1}^N \sigma_i^x$. A simple yet non trivial example of this family of systems is the Grover Hamiltonian, emulating a benchmark model for quantum search [11–15], in which $V_{\mathbf{n}} = -JN\delta_{\mathbf{n},\mathbf{n}_1}$, $J > 0$, and \mathbf{n}_1 represents the target of a totally unstructured (worst case scenario) search [16].

The general result proven in [10], valid at $T = 0$, is as follows. If we can find a partition of the space of states \mathcal{H} of the system into two subspaces, $\mathcal{H} = \mathcal{H}_{\text{cond}} \oplus \mathcal{H}_{\text{norm}}$, such that, in the TDL, $\dim \mathcal{H}_{\text{cond}} / \dim \mathcal{H} \rightarrow 0$ and the ground state energies of H restricted to these subspaces cross each other at a finite value of Γ (or J), then the system undergoes a first-order QPT driven by this parameter. In fact, crossing this point the ground state of the system steeply changes from being a superposition in the *normal* subspace, as large as \mathcal{H} , to a superposition in the *condensed* subspace, with vanishing relative dimension. We thus have a condensation QPT.

In the present paper, we show how to naturally extend the above $T = 0$ quantum condensations to finite temperature. For systems at canonical equilibrium at temperature T , we have just to replace the ground state energies with the free energies. While a detailed proof is given in [17], here we illustrate the new criterion with two examples. As first exam-

ple, we derive analytically the phase diagram of the paradigmatic Grover model. Then we consider a system of relevant physical interest, namely, free fermions in a one-dimensional (1D) inhomogeneous lattice. Also in this case we obtain, numerically, a phase diagram, structurally similar to that of the Grover model. This phase-diagram structure can be proved to be a universal feature of the condensation QPTs [17] and thus the above similarity strongly supports our numerical findings. We conclude suggesting an experimental realization of condensation QPTs in semiconductor superlattices.

Before recalling the $T = 0$ case, two comments are in order about the nature of the condensation QPTs. i) They are intrinsically first-order, for they can be driven by using even one single Hamiltonian parameter. In contrast, as for the classical case, jumps of the order parameter can result when crossing the coexistence line of two different phases that originate from the critical point of a second-order QPT. Notice that, for such a scenario to take place at zero temperature, one needs that the Hamiltonian depends on at least two independent parameters (think to the 1D Ising model in the presence of both a transverse and a longitudinal magnetic field [18, 19]); ii) Condensation QPTs are far from being exotic. As we have recently shown, the renowned Wigner crystallization belongs to this class of QPTs [20].

Normal and condensed subspaces. We start by defining a proper partition of the space of states. Consider a system with Hamiltonian (1), and let $\{|\mathbf{n}_k\rangle\}_{k=1}^M$ be a complete orthonormal set of eigenstates of V , the configurations: $V|\mathbf{n}_k\rangle = V_k|\mathbf{n}_k\rangle$, $k = 1, \dots, M$. We assume ordered potential values $V_1 \leq \dots \leq V_M$. For a system of N qubits, for instance, the set of the configurations may correspond to the set of $M = 2^N$ product states of N spins along some direction, as stated above. Given an integer M_{cond} with $1 \leq M_{\text{cond}} < M$, we make a partition of the set of the configurations as $\{|\mathbf{n}_k\rangle\}_{k=1}^M = \{|\mathbf{n}_k\rangle\}_{k=1}^{M_{\text{cond}}} \cup \{|\mathbf{n}_k\rangle\}_{k=M_{\text{cond}}+1}^M$. Correspondingly, the Hilbert space of the system, $\mathcal{H} = \text{span}\{|\mathbf{n}_k\rangle\}_{k=1}^M$, equipped with standard complex scalar product $\langle u|v\rangle$, is decomposed as the direct sum of two mutually orthogonal subspaces, denoted condensed and normal, $\mathcal{H} = \mathcal{H}_{\text{cond}} \oplus \mathcal{H}_{\text{norm}}$, where $\mathcal{H}_{\text{cond}} = \text{span}\{|\mathbf{n}_k\rangle\}_{k=1}^{M_{\text{cond}}}$, and $\mathcal{H}_{\text{norm}} = \text{span}\{|\mathbf{n}_k\rangle\}_{k=M_{\text{cond}}+1}^M = \mathcal{H}_{\text{cond}}^\perp$. Finally, we define $E = \inf_{|u\rangle \in \mathcal{H}} \langle u|H|u\rangle / \langle u|u\rangle$, $E_{\text{cond}} = \inf_{|u\rangle \in \mathcal{H}_{\text{cond}}} \langle u|H|u\rangle / \langle u|u\rangle$ and $E_{\text{norm}} = \inf_{|u\rangle \in \mathcal{H}_{\text{norm}}} \langle u|H|u\rangle / \langle u|u\rangle$, which are the GS eigenvalues, respectively, of H and of H restricted to the condensed and normal subspaces. According to the scaling properties assumed for K and V , we have that E , E_{cond} and E_{norm} increase linearly with N (at least in the TDL).

Quantum phase transitions at $T = 0$. The Hilbert space dimension M generally diverges exponentially with N , while the dimension M_{cond} , may or may not be a growing function of N . In [10] we have shown that:

$$\text{if } \lim_{N \rightarrow \infty} \frac{M_{\text{cond}}}{M} = 0, \quad (2)$$

$$\text{then } \lim_{N \rightarrow \infty} \frac{E}{N} = \lim_{N \rightarrow \infty} \min \left\{ \frac{E_{\text{cond}}}{N}, \frac{E_{\text{norm}}}{N} \right\}. \quad (3)$$

For finite sizes, up to corrections $O(1)$, Eq. (3) reads

$$E \simeq \begin{cases} E_{\text{cond}}, & \text{if } E_{\text{cond}} < E_{\text{norm}}, \\ E_{\text{norm}}, & \text{if } E_{\text{norm}} < E_{\text{cond}}. \end{cases} \quad (4)$$

As a consequence of Eq. (3), by varying some parameter of the Hamiltonian we obtain a QPT, necessarily of first order, whenever a crossing takes place between E_{cond} and E_{norm} . In the TDL, the space of states splits at the quantum critical point (QCP) defined by

$$\lim_{N \rightarrow \infty} \frac{E_{\text{cond}}}{N} = \lim_{N \rightarrow \infty} \frac{E_{\text{norm}}}{N}, \quad (5)$$

and, in correspondence with Eq. (4), for the GS $|E\rangle$ we have either $|E\rangle \in \mathcal{H}_{\text{cond}}$ or $|E\rangle \in \mathcal{H}_{\text{norm}}$.

Apart from the necessary condition (2), M_{cond} should be properly chosen so that Eq. (5) admits a solution [20]. In addition, if we look for protocols aimed at capturing the GS of (1) for $\Gamma = 0$, we have to set M_{cond} as the degeneracy of the GS of V .

Grover model at $T = 0$. Here $V_1 = -JN$, with $J > 0$, and $V_k = 0$, for $k = 2, 3, \dots, M = 2^N$. We can assume $M_{\text{cond}} = 1$ independent of N . We find $|E_{\text{cond}}\rangle = |\mathbf{n}_1\rangle$ and $E_{\text{cond}} = V_1$. Up to a correction $O(N/M)$, we also have $E_{\text{norm}} = -\Gamma N$ [10, 21]. Therefore Eq. (4) becomes

$$E \simeq \begin{cases} -JN, & \text{if } \Gamma < J, \\ -\Gamma N, & \text{if } \Gamma > J. \end{cases} \quad (6)$$

According to Eq. (5), we thus have a critical value $\Gamma_c = J$ if Γ is the moving parameter, or $J_c = \Gamma$ if J is varied. For $\Gamma > J$ the GS of the model coincides with the GS of the hopping operator K , while for $\Gamma < J$ the system stays locked in the configuration $|\mathbf{n}_1\rangle$. We thus have a QPT that corresponds to a condensation in the space of states.

Order parameter. The interpretation of the above class of QPTs in terms of a condensation in the space of states holds in general, even when $\mathcal{H}_{\text{cond}}$ contains many eigenstates of V [20]. At zero temperature, the probability for the condensed subspace to be occupied is

$$p_{\text{cond}} = \sum_{k=1}^{M_{\text{cond}}} |\langle \mathbf{n}_k | E \rangle|^2. \quad (7)$$

On the other hand, in the TDL, since it is either $|E\rangle \in \mathcal{H}_{\text{cond}}$ or $|E\rangle \in \mathcal{H}_{\text{norm}}$, we find either $p = 1$ or $p = 0$, respectively (we assume $|E\rangle$ normalized). In other words, p_{cond} represents an order parameter of these QPTs.

Finite temperature quantum condensations. Our aim is to extend the above class of condensation QPTs to finite temperature. We suppose that the system, in contact with a heat bath, is at canonical equilibrium at temperature $T = 1/(k_B\beta)$, i.e., it is in the state described by the Gibbs density matrix operator $\rho = e^{-\beta H} / \text{tr } e^{-\beta H}$.

Analogously to the $T = 0$ case, we proceed by defining the Gibbs free energies associated to the spaces \mathcal{H} , $\mathcal{H}_{\text{cond}}$, $\mathcal{H}_{\text{norm}}$,

$$\begin{aligned} e^{-\beta F} &= \text{tr} e^{-\beta H} = \sum_{|\mathbf{n}\rangle \in \mathcal{H}} \langle \mathbf{n} | e^{-\beta H} | \mathbf{n} \rangle, \\ e^{-\beta F_{\text{cond}}} &= \text{tr}_{\text{cond}} e^{-\beta H_{\text{cond}}} = \sum_{|\mathbf{n}\rangle \in \mathcal{H}_{\text{cond}}} \langle \mathbf{n} | e^{-\beta H_{\text{cond}}} | \mathbf{n} \rangle, \\ e^{-\beta F_{\text{norm}}} &= \text{tr}_{\text{norm}} e^{-\beta H_{\text{norm}}} = \sum_{|\mathbf{n}\rangle \in \mathcal{H}_{\text{norm}}} \langle \mathbf{n} | e^{-\beta H_{\text{norm}}} | \mathbf{n} \rangle, \end{aligned}$$

where H_{cond} and H_{norm} are the restrictions of H to the condensed and normal subspaces [22]. It is natural to investigate whether Eq. (3) can be generalized to finite temperature just by substituting the energies E , E_{cond} , E_{norm} with the free energies F , F_{cond} , F_{norm} , which scale linearly with N too.

For any partition $\mathcal{H} = \mathcal{H}_{\text{cond}} \oplus \mathcal{H}_{\text{norm}}$, we can prove that (X stands for either cond or norm and Y for its complement)

$$1 \leq \frac{\langle \mathbf{n} | e^{-\beta H} | \mathbf{n} \rangle}{\langle \mathbf{n} | e^{-\beta H_X} | \mathbf{n} \rangle} \leq e^{\beta \Gamma \min\{A_X^{(\text{out})}, A_Y^{(\text{out})}\}}, \quad |\mathbf{n}\rangle \in \mathcal{H}_X, \quad (8)$$

$$F \leq \min\{F_{\text{cond}}, F_{\text{norm}}\}, \quad (9)$$

$$F \geq \min\{F_{\text{cond}}, F_{\text{norm}}\} - \min\{A_{\text{cond}}^{(\text{out})}, A_{\text{norm}}^{(\text{out})}\} \Gamma, \quad (10)$$

where $A_X^{(\text{out})} = \sup_{|\mathbf{n}\rangle \in \mathcal{H}_X} \sum_{|\mathbf{n}'\rangle \in \mathcal{H}_Y} |\langle \mathbf{n} | K | \mathbf{n}' \rangle|$ represents the maximum number of outgoing links (nonzero matrix elements of K) from \mathcal{H}_X to \mathcal{H}_Y . The product $\min\{A_X^{(\text{out})}, A_Y^{(\text{out})}\} \Gamma$ determines the rate of convergence to 1 of the probability for crossing the boundary between \mathcal{H}_X and \mathcal{H}_Y along the quantum dynamics. Hereafter, we assume $\min\{A_{\text{cond}}^{(\text{out})}, A_{\text{norm}}^{(\text{out})}\} = A_{\text{norm}}^{(\text{out})}$. In the Grover model, e.g., $A_{\text{norm}}^{(\text{out})} = 1$ while $A_{\text{cond}}^{(\text{out})} = N$. The important point is that, in most of the systems of interest, the conditions $M_{\text{cond}}/M \rightarrow 0$ or $A_{\text{norm}}^{(\text{out})}/N \rightarrow 0$ are equivalent [17] and, under any of these conditions, Eqs. (9) and (10), up to a $o(N)$ term, provide the natural generalization of Eq. (4)

$$F \simeq \begin{cases} F_{\text{cond}}, & \text{if } F_{\text{cond}} < F_{\text{norm}}, \\ F_{\text{norm}}, & \text{if } F_{\text{norm}} < F_{\text{cond}}. \end{cases} \quad (11)$$

Equation (11) extends the $T = 0$ QPT to finite temperature. The crossing between F_{cond} and F_{norm} gives rise to a first-order QPT controlled by Hamiltonian parameters and temperature, the equation for the critical surface being

$$\lim_{N \rightarrow \infty} \frac{F_{\text{cond}}}{N} = \lim_{N \rightarrow \infty} \frac{F_{\text{norm}}}{N}. \quad (12)$$

The proof of Eqs. (8) is detailed in [17] and is based on the exact probabilistic representation of the quantum evolution introduced in [23] used at an imaginary time identified with the inverse temperature β . Equations (9)-(10) are easily derived from Eqs. (8).

The probability for the condensed subspace to be occupied represents an order parameter also at finite temperature and




Figure 1. Phase diagram J - T for the Grover model at thermal equilibrium, the solid line separating the two phases is drawn according to Eq. (15).

the phase transition can be interpreted as a first-order condensation in the space of states. In fact, due to Eqs. (8)

$$p_{\text{cond}} = \sum_{|\mathbf{n}\rangle \in \mathcal{H}_{\text{cond}}} \langle \mathbf{n} | \rho | \mathbf{n} \rangle \simeq \frac{1}{1 + e^{-\beta(F_{\text{norm}} - F_{\text{cond}})}}, \quad (13)$$

where the equality holds in the TDL with $p_{\text{cond}} = 1$ in the condensed phase $F_{\text{cond}} < F_{\text{norm}}$ and $p_{\text{cond}} = 0$ in the normal one $F_{\text{norm}} < F_{\text{cond}}$. At the critical surface separating the two phases we have $p_{\text{cond}} = 1/2$.

Grover Model at $T > 0$. We illustrate the mechanism of the finite temperature condensation in the exactly solvable Grover model. Since $M_{\text{cond}} = 1$, we have $-\beta F_{\text{cond}} = -\beta V_1$ with $V_1 = -JN$. Up to corrections exponentially small in N , the free energy of the normal subspace coincides with that of the hopping operator K whose levels are $-\Gamma(N - 2j)$, $j = 0, \dots, N$, and have degeneracy $N!/(j!(N-j)!)$,

$$e^{-\beta F_{\text{norm}}} = \text{tr} e^{-\beta K} = \sum_{j=0}^N \binom{N}{j} e^{-\beta(-\Gamma(N-2j))}, \quad (14)$$

which yields $-\beta F_{\text{norm}} = N \log(2 \cosh(\beta \Gamma))$. The critical surface defined by Eq. (12) is thus

$$J = k_B T \log(2 \cosh(\Gamma/k_B T)), \quad (15)$$

also found in Ref. [15] via approximate methods. Note that Eq. (15) has solution only for $J \geq \Gamma$. A parametric plot of the critical surface is shown in Fig. 1. For any fixed Γ , at high temperature Eq. (15) provides the asymptotic slope $T = J/(k_B \log 2)$, while at low temperature the slope becomes infinite in correspondence of the QCP $J = \Gamma$ of the $T = 0$ transition. We can prove that the above two features are a universal fingerprint of the condensation QPTs at finite temperature [17].

Free fermions in a 1D inhomogeneous lattice. Finite temperature condensation QPTs can be observed in a variety of

physically relevant systems. Here, we study the case of N_p spinless fermions moving in a 1D lattice with $N \geq N_p$ sites. Some sites, the first consecutive $N_i < N_p$, for simplicity, differ from the others by the presence of an attractive potential so that the Hamiltonian of the system is

$$H = -\eta \sum_{l=1}^{N-1} (c_l^\dagger c_{l+1} + c_{l+1}^\dagger c_l) - g \sum_{l=1}^{N_i} c_l^\dagger c_l, \quad (16)$$

where c_l is the fermionic annihilation operator on site l and we choose open boundary conditions. The hopping parameter η and the attractive strength g are positive constants. Note that we are considering a system of noninteracting particles, nevertheless H is the sum of two noncommuting operators as in Eq. (1) and we can look for a condensation QPT by varying the parameter g .

At first, we define the subspace $\mathcal{H}_{\text{cond}}$. Besides satisfying the necessary condition (2), this subspace should be large enough for the free energies restricted to $\mathcal{H}_{\text{cond}}$ and $\mathcal{H}_{\text{norm}}$ to cross each other at some finite value g_c of the parameter g . The latter condition is equivalent to have $F_{\text{norm}}(g=0) < F_{\text{cond}}(g=0)$ and $\lim_{g \rightarrow \infty} F_{\text{cond}}(g)/g < \lim_{g \rightarrow \infty} F_{\text{norm}}(g)/g$ (see [20] for the $T=0$ analog). In the present system and for $N_i = N_p$, the above two inequalities are satisfied if $M_{\text{cond}} = 1 + N_p^2$, i.e., if the subspace $\mathcal{H}_{\text{cond}}$ consists of the ground state of V , in which all the N_p fermions are in the N_i attractive sites, and of the N_p^2 first excited states of V , in which $N_p - 1$ fermions are in the N_i attractive sites and 1 is in the remaining $N - N_i$ sites. Other choices of M_{cond} are possible, but all lead to the same TDL. Remarkably, in this system, the nature of $\mathcal{H}_{\text{cond}}$ shows that a condensation in the space of states corresponds to an actual space localization in the attractive sites.

Second, we evaluate the order parameter (13) as

$$p_{\text{cond}} = \frac{1}{\sum_{j=1}^M e^{-\beta E_j}} \sum_{j=1}^M e^{-\beta E_j} \sum_{k=1}^{M_{\text{cond}}} |\langle \mathbf{n}_k | E_j \rangle|^2, \quad (17)$$

where E_j and $|E_j\rangle$ are the N_p -particle eigenvalues and eigenvectors of H . These are easily obtained by Pauli's principle combining the single-particle eigenvalues and eigenvectors of H calculated by numerically diagonalizing the $N \times N$ tridiagonal matrix with non zero elements $A_{l+1,l} = A_{l,l+1} = -\eta$, for $l = 1, \dots, N-1$, and $A_{l,l} = -g$, for $l = 1, \dots, N_i$. The computation of E_j and $|E_j\rangle$ is a simple task which requires a time $O(N^2)$. Also the computation of the sum of the squared scalar products in Eq. (17) requires an affordable time $O(N^2 N_p^2)$ [24]. It's the sum over M which limits the computation of p_{cond} to a relatively small number of particles, in fact, M grows as $N!/(N_p!(N - N_p)!)$.

In Fig. 2 we show the behavior of p_{cond} obtained as a function of the attractive strength g for a number of particles $N_p = 4, 6, 8, \dots, 16$. As indicated above, we chose $N_p = N_i = N/2$ and $M_{\text{cond}} = 1 + N_p^2$. The equilibrium temperature is $k_B T/\eta = 0.2$, similar plots at different temperatures are provided in [25]. Whereas the order param-




Figure 2. Order parameter p_{cond} versus g/η for the fermionic system (16) with $N_p = N_i = N/2 = 4, 6, 8, 10, 12, 14, 16$, symbols left to right. Here $M_{\text{cond}} = 1 + N_p^2$ and the system is at canonical equilibrium at temperature $k_B T/\eta = 0.2$. Inset: value of g/η at $p_{\text{cond}} = 1/2$ for different N_p . Data (circles) compare quite well ($\chi^2 \simeq 10^{-4}$) with two fitting models $a + b/N_p + c/N_p^2$ (black solid line, $a = 4.424$, $b = 9.685$ and $c = 13.760$) and $a + b \log N_p + c \log(\log N_p)$ (red dashed line, $a = 2.261$, $b = -2.268$ and $c = 7.627$). The second model, however, must be rejected because in the TDL gives $g_c \rightarrow -\infty$.

ter exhibits a clear tendency toward the step-like behavior expected for a first-order QPT, we are still far from the TDL and no classical supercomputer could allow to reach much larger values of N_p . However, we can estimate the critical value g_c of the QPT, namely, the value of g at which $p_{\text{cond}} = 1/2$ in the TDL, by a fit-and-extrapolate procedure. The validity of this procedure in the case of the Grover model is illustrated in [25]. We fit the curve $a + b/N_p + c/N_p^2$ to the values of g at which $p_{\text{cond}} = 1/2$ for the available N_p and extrapolate $g_c = a$. A different fit model which provides a slowly diverging g_c , namely, $a + b \log N_p + c \log(\log N_p)$, must be rejected as we obtain a pointless negative value of b , see Fig. 2.

In Fig. 3 we plot the values of g_c derived as explained above for different temperatures. The result is a phase diagram in the plane g - T sharing the same universal features [17] of the phase diagram of Fig. 1. This corroborates the existence of a condensation QPT for the present fermionic system.

Conclusions. Under the same conditions valid at $T=0$ but replacing the ground state energies of the system and of its restrictions to the condensed and normal subspaces, with the corresponding free energies, the condensation QPTs can be extended to $T > 0$. Even if the proof is rather involved [17], the result is natural and, as we have shown, describes accurately the behavior of two rather different systems: the Grover model, analytically solvable, and the case of free fermions in a 1D inhomogeneous lattice.

The two studied systems, despite being very different, share similar phase diagrams, with a critical temperature asymptotically proportional to the potential strength and having an infinite slope at the QCP. In fact, these are universal features of




Figure 3. Phase diagram g - T for the fermionic system (16). The solid line separating the two phases is evaluated by the fitting procedure described in Fig. 2

the condensation QPTs [17].

The 1D fermionic system considered here could be investigated experimentally by using superlattices, grown with reliable technologies in the well established 2D semiconductor heterojunctions [30] or in 1D nanowires [31]. In both cases, the alternance of nano-layers of different materials forms effective 1D lattices made of wells and barriers obtained by the corresponding band-gap energies. By varying the barrier widths and heights one can tune the hopping coefficients between neighbouring wells, and by using different/doped materials adjust the bottom level of the wells, i.e., create local attractive potentials. The number of electrons in the nanocells of the superlattice can be fixed by photoexcitation and their dislocation at thermal equilibrium detected by photoluminescence spectroscopy [32]. In this way, one could have direct access to our order parameter and, for large superlattices in which the TDL has been effectively reached, to the critical surface of the phase diagram.

We are grateful to Antonio Polimeni for helpful discussions on the experimental implementation of our model. Grant CNPq 307622/2018-5 - PQ (Brazil) is acknowledged. M. O. thanks the Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, and the Department of Physics of Sapienza University of Rome for financial support and hospitality.

-
- [1] S. L. Sondhi, S. M. Girvin, J. P. Carini, and D. Shahar, “Continuous quantum phase transitions”, *Rev. Mod. Phys.* **69**, 315 (1997).
 [2] T. R. Kirkpatrick and D. Belitz, “Quantum phase transitions in electronic systems”, in *Electron Correlations in the Solid State* ed. by N. H. March, (Imperial College Press, London 1999).
 [3] T. Vojta, “Quantum phase transitions in electronic systems”, *Ann. Phys. (Leipzig)* **9**, 403 (2000).

- [4] S. Sachdev, *Quantum Phase Transitions* (Cambridge University Press, Cambridge 2000).
 [5] L. D. Carr, “Understanding Quantum Phase Transitions”, (CRC Press, Taylor & Francis 2010).
 [6] A. Plastino and E. M. F. Curado, “Finite temperature approach to quantum phase transitions”, *International Journal of Bifurcation and Chaos* **20**, 397 (2010).
 [7] A. B. Finilla, M. A. Gomez, C. Sebenik, and D. J. Doll, “Quantum annealing: A new method for minimizing multidimensional functions”, *Chem. Phys. Lett.* **219**, 343 (1994).
 [8] T. Kadowaki and H. Nishimori, “Quantum annealing in the transverse Ising model”, *Phys. Rev. E* **58**, 5355 (1998).
 [9] G. E. Santoro and E. Tosatti, “Optimization using quantum mechanics: quantum annealing through adiabatic evolution”, *J. Phys. A* **39**, R393 (2006).
 [10] M. Ostilli and C. Presilla, “First-order quantum phase transitions as condensations in the space of states”, *J. Phys. A: Math. Theor.* **54**, 055005 (2021).
 [11] L. K. Grover, “A fast quantum-mechanical search algorithm for database search”, *Proceedings, 28th Annual ACM Symposium on the Theory of Computing (STOC)*, May 1996, pages 212-219; “Quantum Mechanics Helps in Searching for a Needle in a Haystack”, *Phys. Rev. Lett.* **79**, 325 (1997); “From Schrödinger’s equation to the quantum search algorithm”, *Am. J. Phys.* **69**, 769 (2001).
 [12] E. Farhi and S. Gutmann, “Analog analogue of a digital quantum computation”, *Phys. Rev. A* **57**, 2403 (1998).
 [13] J. Roland and N. J. Cerf, “Quantum search by local adiabatic evolution”, *Phys. Rev. A*, **65**, 042308 (2002).
 [14] T. Jörg, F. Krzakala, J. Kurchan, and A. C. Maggs, “Simple glass models and their quantum annealing”, *Phys. Rev. Lett.* **101**, 147204 (2008).
 [15] T. Jörg, F. Krzakala, J. Kurchan, A. C. Maggs, and J. Pujos, “Energy gaps in quantum first-order mean-field-like transitions: The problems that quantum annealing cannot solve”, *Europhysics Letters* **89**, 40004 (2010).
 [16] In contrast, structured searches correspond to potentials having a smooth minimum around the target and, therefore, benefit from the application of gradient-descent based methods like, e.g., the Ising model where, however, the corresponding QPTs are second-order.
 [17] M. Ostilli and C. Presilla, “Finite temperature quantum condensations in the space of states: general proof”, arXiv:2209.13244
 [18] M. A. Continentino and A. S. Ferreira, “First-order quantum phase transitions”, *J. Magnetism and Magnetic Materials* **310**, 828 (2007).
 [19] M. Campostrini, J. Nespolo, A. Pelissetto, and E. Vicari, “Finite-Size Scaling at First-Order Quantum Transitions” *Phys. Rev. Lett.* **113**, 070402 (2014).
 [20] M. Ostilli and C. Presilla, “Wigner crystallization of electrons in a one-dimensional lattice: a condensation in the space of states”, *Phys. Rev. Lett.* **127**, 040601 (2021).
 [21] C. Presilla and M. Ostilli, “Phase transitions and gaps in quantum random energy models”, *Physica A* **515**, 57 (2019).
 [22] In the representation of the eigenstates of V , H_{cond} corresponds to a null matrix except for the block $\langle \mathbf{n}_k | H_{\text{cond}} | \mathbf{n}_{k'} \rangle = \langle \mathbf{n}_k | H | \mathbf{n}_{k'} \rangle$, $k, k' = 1, \dots, M_{\text{cond}}$. Analogously, H_{norm} corresponds to a null matrix except for the block $\langle \mathbf{n}_k | H_{\text{norm}} | \mathbf{n}_{k'} \rangle = \langle \mathbf{n}_k | H | \mathbf{n}_{k'} \rangle$, $k, k' = M_{\text{cond}} + 1, \dots, M$. Note that $H_{\text{cond}} + H_{\text{norm}} \neq H$.
 [23] M. Beccaria, C. Presilla, G. F. De Angelis, and G. Jona-Lasinio, “An exact representation of the fermion dynamics in terms of Poisson processes and its connection with Monte Carlo algorithms”, *Europhys. Lett.* **48**, 243 (1999).

- [24] The evaluation of the scalar product between the antisymmetrized states $|n_k\rangle$ and $|E_j\rangle$ can be reduced to the evaluation of the determinant of the $N \times N$ matrix whose elements are the projections of the single-particle eigenvectors of H in the basis of the eigenvectors of V [28]. The calculation of this determinant requires a time $O(N^2)$.
- [25] Supplemental Material.
- [26] H. Lan and Y. Ding, “Ordering, positioning and uniformity of quantum dot arrays”, *Nano Today*, **7**, 94 (2012).
- [27] I. Piquero-Zulaica, et al. “Precise engineering of quantum dot array coupling through their barrier widths”, *Nat. Commun.* **8**, 787 (2017).
- [28] J. P. Blaizot and G. Ripka, *Quantum Theory of Finite Systems* (The MIT Press, Cambridge, Massachusetts 1986).
- [29] T. Kato, “On the adiabatic theorem of quantum mechanics”, *J. Phys. Soc. Jpn.* **5**, 435 (1950).
- [30] Qian Wang, Lin Zhang, Xuejuan Liu and Sha Li, “Two-Dimensional Semiconductor Heterojunctions for Optoelectronics and Electronics”, *Front. Energy Res.*, 9:802055 (2021).
- [31] O. Arif, V. Zannier, F. Rossi, D. De Matteis, K. Kress, M. De Luca, I. Zardo and L. Sorba, “GaAs/GaP superlattice nanowires: growth, vibrational and optical properties”, *Nanoscale*, 2022, Advance Article.
- [32] D. Tedeschi, M. De Luca, A. Polimeni, “Photoluminescence Spectroscopy Applied to Semiconducting Nanowires: A Valuable Probe for Assessing Lattice Defects, Crystal Structures, and Carriers’ Temperature”, pages 289–306 in *Fundamental Properties of Semiconductor Nanowires*, N. Fukata and R. Riccardio editors (Springer, Singapore 2021).

Supplemental Material for “Finite temperature quantum condensations in the space of states”

Massimo Ostilli and Carlo Presilla

Grover model

In the following, we provide two plots of the order parameter p_{cond} evaluated numerically in the Grover Model as a function of Γ/J for different values of N . Note that the higher is the temperature the slower is the convergence of the data to the $N \rightarrow \infty$ limit of p_{cond} represented by the step-like solid line. Nevertheless, the fit of $a + b/N + c/N^2$ to the values of Γ/J at $p_{\text{cond}} = 0.5$ provides an asymptotic value of $\Gamma_c/J = a$ in good agreement with the exact value.




Figure S1. Left panel. Order parameter p_{cond} versus Γ/J for the Grover model at thermal equilibrium at temperature $k_B T/J = 0.5$ for $N = 7, 9, 11, 13$ (open symbols). The step-like solid line is the asymptotic $N \rightarrow \infty$ exact value. Right panel. Value of Γ/J at $p_{\text{cond}} = 1/2$ for different N (open circles). The function $a + b/N + c/N^2$ fits quite well the data (black solid line, $a = 0.96$, $b = 1.88$ and $c = -4.18$) and predicts the asymptotic exact value $\Gamma_c/J = 0.99$ (dashed red line) with a 3% error.




Figure S2. As in figure S1 at temperature $k_B T/J = 1.2$. The fit parameters ($a = 0.67$, $b = 0.79$ and $c = -0.75$) predict the asymptotic exact value $\Gamma_c/J = 0.65$ with a 3% error.

Free fermions in a 1D inhomogeneous lattice

In the following figures, we report the values of p_{cond} obtained numerically for the system (16) at canonical equilibrium at different temperatures T . In each plot, the parameter a of the fit is taken as the critical value $g_c(T)$ shown in Fig. 3.




Figure S3. Order parameter p_{cond} versus g/η for the fermionic system (16) with $N_p = N_i = N/2 = 4, 6, 8, 10, 12$ (symbols from left to right). We put $M_{\text{cond}} = 1 + N_p^2$ and the system is at canonical equilibrium at temperature $k_B T/\eta = 0.02$. Inset: value of g/η at $p_{\text{cond}} = 1/2$ for different N_p . Data (circles) compare quite well with the fitting function $a + b/N_p + c/N_p^2$ (solid line, $a = 4.277$, $b = -6.860$ and $c = -26.841$).




Figure S4. As in Fig. S3 at temperature $k_B T/\eta = 0.1$. The fit gives $a = 4.303$, $b = -7.058$ and $c = -22.568$.




Figure S5. As in Fig. S3 at temperature $k_B T/\eta = 0.3$. The fit gives $a = 4.622$, $b = -11.987$ and $c = -7.506$.




Figure S6. As in Fig. S3 at temperature $k_B T/\eta = 0.4$. The fit gives $a = 4.922$, $b = -15.144$ and $c = 0.4423$.




Figure S7. As in Fig. S3 at temperature $k_B T/\eta = 0.5$. The fit gives $a = 5.277$, $b = -18.517$ and $c = 8.364$.




Figure S8. As in Fig. S3 at temperature $k_B T / \eta = 0.7$. The fit gives $a = 6.060$, $b = -24.931$ and $c = 21.991$.




Figure S9. As in Fig. S3 at temperature $k_B T / \eta = 1.0$. The fit gives $a = 7.359$, $b = -33815$ and $c = 38.458$.




Figure S10. As in Fig. S3 at temperature $k_B T / \eta = 1.3$. The fit gives $a = 8.774$, $b = -42.407$ and $c = 52.764$.