

COMPUTING EQUIVALENCE CLASSES OF FINITE GROUP ACTIONS ON ORIENTABLE SURFACES: A DYNAMIC SURVEY

JÁN KARABÁŠ, ROMAN NEDELA, AND MÁRIA SKYVOVÁ

ABSTRACT. This paper focuses on the classification of classes of topological equivalence of finite group actions on Riemann surfaces. By the Riemann-Hurwitz bound, there are just finitely many groups that act conformally on a closed orientable surface $\mathcal{S}_g$ of genus $g \geq 2$. With each such action of a group G on $\mathcal{S}_g$ one can associate the fundamental group $\Gamma = \pi(\mathcal{O})$ of the quotient orbifold $\mathcal{O} = \mathcal{S}_g/G$, isomorphic to a Fuchsian group determined completely by orbifold's signature. The Riemann existence theorem reduces the problem of the existence of an action of G on $\mathcal{S}_g$ to a purely group-theoretical problem of deciding whether there is a smooth epimorphism mapping the Fuchsian group Γ onto the group G . Using computer algebra systems such as MAGMA or GAP, together with the library of small groups, the generation of all finite group actions on a surface of fixed small genus $g \geq 2$ becomes almost a routine procedure. The difficult part is to determine the classes of these actions with respect to topological equivalence. To achieve this, one needs to investigate the action of the automorphism group of a Fuchsian group on the set of finite group actions on $\mathcal{S}_g$ with the corresponding signature. In this paper we derive several results on the topological equivalence of finite group actions on Riemann surfaces. As an application, we derive complete lists of finite group actions of genus $g \leq 9$ distinguished up to the topological equivalence. A summary of the actions can be found in Appendix; the reader interested in more details is referred to the web page <https://www.savbb.sk/~karabas/science/discactions.html>. It is expected that we will be able to extend the list to higher genera, refreshed partial results are available on the web page. The following text is an extended version of the paper [23].

1. INTRODUCTION

The investigation of surface symmetries is a long-standing project started in ancient Greece. By the beginning of 20th century the classification of the spherical and wallpaper groups, which are the groups of isometries of the sphere and of the Euclidean plane, had been established. Groups acting on the projective plane, the torus, and the Klein bottle are understood as well, since these are quotients of the spherical and wallpaper groups. In contemporary mathematics, the symmetries of surfaces of higher genera are studied within the frame of the theory of Riemann surfaces and of algebraic curves. With each orientation-preserving action of a group G on a orientable surface $\mathcal{S}_g$ of genus g , one can associate the *signature* $(\gamma; m_1, m_2, \dots, m_r)$, where γ is the genus of the quotient surface, r is the number of singular orbits, and m_i , $i = 1, \dots, r$, are the orders

Date: February 22, 2024; A dynamic survey, version 2.

2020 Mathematics Subject Classification. 30F10, 20F34.

Key words and phrases. group action, Riemann surface, Fuchsian group, mapping class group.

of the point stabilizers in the corresponding singular orbits (the point stabilizers are necessarily cyclic). The signature, genus g , and order of the group G are related by the Riemann-Hurwitz equation (2.2). As a consequence, the order of G is related via the Hurwitz bound, $|G| \leq 84(g-1)$ for $g > 1$, see e.g. [21, Theorem 5.11.1]. It follows that for genera greater than one, there are only finitely many finite groups acting on these surfaces. Thus, determining all of them is a finite problem. On the other hand, groups that are small with respect to Hurwitz bound have usually a huge number of actions even on surfaces of small genera. Hence it makes sense to consider proper equivalence relations and classify the actions up to the chosen equivalence. An excellent account on equivalence relations of group actions on surfaces, together with many open problems, can be found in recent papers [8, 9].

One of the most important relations defined on actions of groups on orientable surfaces is the topological equivalence defined as follows. For an orientable surface $\mathcal{S}$ denote by $\text{Hom}^+(\mathcal{S})$ the group of orientation-preserving homeomorphisms. Two actions of a group G , given by embeddings $\varepsilon_i: G \rightarrow \text{Hom}^+(\mathcal{S}_i)$, $i = 1, 2$, on possibly different surfaces $\mathcal{S}_1, \mathcal{S}_2$ of the same genus, are *topologically equivalent* if there is an intertwining, orientation preserving homeomorphism $h: \mathcal{S}_1 \rightarrow \mathcal{S}_2$ and an automorphism $a \in \text{Aut}(G)$ such that $\varepsilon_2(g) = h\varepsilon_1(a(g))h^{-1}$, for every $g \in G$. As it is noted in [8, p.23], “topological equivalence of actions is important since the action classes are directly linked to conjugacy classes of finite subgroups of the mapping class group, in a 1-1 fashion, and to the strata of the branch locus of moduli space in an almost 1-1 fashion”.

Problem 1. *For a fixed integer $g > 1$, derive the list of actions of finite groups on an orientable surface of genus g , distinguished up to topological equivalence.*

A complete solution of Problem 1 is known only up to genus 4, see Broughton [7] for genera 2 and 3, and Bogopolsky [3] for genus four. Incomplete lists for genera 4 and 5 were derived by Kimura [24], and by Kuribayashi and Kimura [26]. To determine the classes of topological equivalence of finite groups of genus g seems to be a hard problem, both from theoretical and practical point of view. Broughton in [8] discusses several related equivalences of group actions which may be used to approximate the topological equivalence. In particular, Breuer in [5] explains how one can employ computations with group characters to determine the equivalence relation $\sim_{\mathcal{H}^1}$ (following the notation from [8]). The numbers of $\sim_{\mathcal{H}^1}$ -classes are computed up to genus 48, see [5, p.95]. By Broughton [8, p.28], two actions that are topologically equivalent are $\sim_{\mathcal{H}^1}$ -equivalent as well. Unfortunately, in many instances of Problem 1 the topological equivalence differs significantly from $\sim_{\mathcal{H}^1}$ -equivalence, see discussion in Section 8 for details. Paulhus in [36] introduces a database of group actions, however, the actions are not distinguished up to topological equivalence. Moreover, we have checked that the database [35] is not complete, see Section 8 for some details.

An action of a finite group G on (an orientable) surface of genus g determines the quotient *orbifold* $\mathcal{S}_g/G$ with the fundamental group Γ . On the other hand, Γ acts as a group of orientation-preserving symmetries of the universal cover $\mathcal{U}$ of the surface $\mathcal{S}_g$. In this setting, Γ is known under the name *Fuchsian group*. The surface $\mathcal{S}_g$ and the group G can be obtained as quotients $\mathcal{U}/K$ and Γ/K by some normal subgroup $K \triangleleft \Gamma$, respectively. The group K is a surface group isomorphic to $\pi_1(\mathcal{S}_g)$. The Fuchsian group Γ is completely determined by the signature $(\gamma; m_1, \dots, m_r)$, therefore, we use

the notation $\Gamma = F(\gamma; m_1, \dots, m_r)$. By the Riemann existence theorem (Theorem 2.1), an action of an abstract finite group G on $\mathcal{S}_g$ with a prescribed signature exists if and only if there is a smooth epimorphism $\eta: \Gamma \rightarrow G$. An epimorphism is *smooth* if for each element x of finite order $|\eta(x)| = |x|$. Equivalently, smooth epimorphisms are called *surface kernel* epimorphisms [30]. In view of the above, the Riemann existence theorem yields that the classification of discrete groups of a given genus can be seen as a purely group-theoretical problem. More precisely, Lloyd proved in [30] that two actions of a group G with given signature, given by associated epimorphisms $\eta_1, \eta_2: \Gamma \rightarrow G$, are topologically equivalent if and only if there exist $a \in \text{Aut}(G)$ and orientation-preserving automorphism $\alpha \in \text{Aut}^+(\Gamma)$ such that $\eta_2 = a\eta_1\alpha$. It follows that in order to determine the equivalence classes it is necessary to study the action of the *group of orientation-preserving automorphisms* $\text{Aut}^+(\Gamma)$ of the Fuchsian group Γ on the set of all smooth epimorphisms $\Gamma \rightarrow G$. A particular subproblem is to find a proper finite set of generators of $\text{Aut}^+(\Gamma)$. The problem of determining a finite generating sets of $\text{Aut}^+(\Gamma)$ is related to the problem of determining the generating sets of mapping class groups of two-dimensional orbifolds, a topic intensely studied within the framework of geometric topology. In the cases we are interested in, the mapping class group is isomorphic to the outer automorphism group of Γ . Therefore, finding a finite generating set for $\text{Aut}^+(\Gamma)$ and finding the generating set for the corresponding mapping class group are equivalent problems; see [6, 9] for details. Various generating sets of mapping class groups of surfaces are determined in [28, 31, 20, 19, 39, 25, 37]. In [16, 27] one can find a finite set of generators for mapping class groups of punctured surfaces.

A significant part of the paper deals with symmetries of Fuchsian groups with planar signatures. Following Zieschang et al. [41], a Fuchsian group is *planar* if its signature is of the form $(0; m_1, m_2, \dots, m_r)$. Finite sets of generators and presentations of $\text{Aut}^+(\Gamma)$, for planar Γ and $r \leq 6$ can be found in [38]. In Section 2 we introduce basic concepts and relevant general results. In Section 3 we collect and organise known results on the automorphism group of a planar Fuchsian group derived by Zieschang [41], Birman [2], and Chow [10]. As a consequence, in Section 4 we are able to formulate and prove Theorem 4.1 characterising the action of $\text{Aut}^+(\Gamma)$ on the *sets of smooth epimorphisms* $\text{Epi}_o(\Gamma, G)$ for Fuchsian groups Γ with planar signatures. Based on the theoretical part of the paper we were able to implement an algorithm determining representatives of equivalence classes of finite groups with planar signatures for a fixed genus. Further, employing the generating set of the automorphism group of a surface group derived by McCool in [31] we implemented the algorithm computing equivalence classes of actions with signatures of type $(\gamma; -)$, $\gamma \geq 1$. The two algorithms are complemented by an approximation algorithm producing an equivalence relation refining the topological equivalence for actions with nonplanar signature. The approximation uses automorphisms of Fuchsian groups introduced in [18]. Actual outputs of programs and other related information can be found online [22]. In particular, in this paper we deal with actions of finite groups of genera $2 \leq g \leq 9$, see Section 8 and Appendix for more details.

2. FINITE GROUP ACTIONS ON SURFACES

The following set of transformations, known as the *group of Möbius transformations*, maps the upper half-plane $\mathcal{U} = \{z \in \mathbb{C} \mid \operatorname{Im}(z) > 0\}$ onto itself by taking $z \mapsto w(z) = \frac{az+b}{cz+d}$ for $a, b, c, d \in \mathbb{R}$, $ad - bc = 1$. The group, known as $\operatorname{LF}(2, \mathbb{R})$ or $\operatorname{PSL}(2, \mathbb{R})$, can be topologised by taking the usual metric topology in $\mathbb{R}^4$ and identifying w with the equivalence class of $(a, b, c, d) \in \mathbb{R}^4$, $(a, b, c, d) \sim \lambda(a, b, c, d)$ for all $\lambda \in \mathbb{R} \setminus \{0\}$. A discrete subgroup of $\operatorname{PSL}(2, \mathbb{R})$ is called a Fuchsian group. Fuchsian groups for which the orbit space is compact will be of interest here, so henceforth ‘Fuchsian group’ will mean ‘Fuchsian group with compact orbit space’.

A torsion-free Fuchsian group is a *surface group*. Surface groups are isomorphic to fundamental groups of compact connected orientable surfaces. It is known that a Fuchsian group Γ always contains a normal surface subgroup K of finite index [34]. On the other hand, every normal torsion-free subgroup of finite index in Γ is a surface group. Recall that a discrete group of automorphisms of a connected compact closed surface is finite. By Zieschang [40], every such group is a quotient of a subgroup of the group of Möbius transformations of the upper half-plane $\mathcal{U}$.

Let $\mathcal{S}_g$ be an orientable surface of the genus g , and let $\operatorname{Hom}^+(\mathcal{S}_g)$ be its group of orientation-preserving homeomorphisms. We say that the group G acts on the surface $\mathcal{S}_g$ if there is a monomorphism $\varepsilon: G \rightarrow \operatorname{Hom}^+(\mathcal{S}_g)$. Every action of G on $\mathcal{S}_g$ is determined by the pair of Fuchsian groups $K \triangleleft \Gamma < \operatorname{PSL}(2, \mathbb{R})$ that act discontinuously on $\mathcal{U}$ and by an epimorphism $\eta: \Gamma \rightarrow G$ with kernel K , where K is a surface group. Such an epimorphism is smooth by definition. The epimorphism η is constructed from ε and from a homeomorphism of $\mathcal{U}/K \cong \mathcal{S}_g$. More precisely, let G be a finite group. The smooth epimorphism $\eta: \Gamma \rightarrow G$ determines a discrete action of the group G on the quotient Riemann surface $\mathcal{U}/\ker \eta$. For every finite group G acting as a group of automorphisms of a compact orientable surface $\mathcal{S}$ there is a Fuchsian group Γ and an smooth epimorphism $\eta: \Gamma \rightarrow G$, such that the action $(G, \mathcal{S})$ is equivalent to the action $(\Gamma/\ker \eta, \mathcal{U}/\ker \eta)$.

The group $\Gamma = F(\gamma; m_1, m_2, \dots, m_r)$ has the (canonical) presentation

$$(2.1) \quad \langle x_1, x_2, \dots, x_r, a_1, \dots, a_\gamma, b_1, \dots, b_\gamma \mid x_1^{m_1} = \dots = x_r^{m_r} = 1, \prod_{i=1}^{\gamma} [a_i, b_i] \prod_{j=1}^r x_j = 1 \rangle,$$

where $1 < m_1 \leq m_2 \leq \dots \leq m_r$ are integers, $0 \leq \gamma \leq g$ is the genus of the quotient surface $\mathcal{S}_g/\eta(\Gamma)$, and m_i are the branch indices of the (regular) branched covering $\mathcal{S}_g \rightarrow \mathcal{S}_g/\eta(\Gamma)$, such that m_i divides $|G|$ for $i = 1, 2, \dots, r$. All the integer parameters in the presentation (2.1) are related with the genus g of the surface $\mathcal{S}_g$ by the Riemann-Hurwitz equation

$$(2.2) \quad 2 - 2g = |G| \left(2 - 2\gamma - \sum_{i=1}^r \left(1 - \frac{1}{m_i} \right) \right);$$

$$m_i \geq 2, \quad m_i \mid |G|, \quad i = 1, \dots, r.$$

As a consequence, one finds that if $g \geq 2$, then the order of the group G is bounded by the Hurwitz bound, $|G| \leq 84(g-1)$. It follows that there are finitely many groups acting on the surface $\mathcal{S}_g$ of genus $g \geq 2$. The integers $\gamma, m_1, \dots, m_r$ determined by the action of G , usually written as vector $(\gamma; m_1, \dots, m_r)$, will be called the *signature* or the *branch data*. Consider the inverse problem: Given the genus $g \geq 2$, let $|G|$ and a signature $(\gamma; m_1, \dots, m_r)$ satisfy the Riemann-Hurwitz equation. Is there a group G of order $|G|$ that acts on the orientable surface $\mathcal{S}_g$ of the genus g such that the Riemann-Hurwitz equation is satisfied for the signature $(\gamma; m_1, m_2, \dots, m_r)$? It turns out that this is not always true. A complete answer is given by the following theorem.

Theorem 2.1 (Riemann existence theorem [7, 17]). *Group G acts on a surface $\mathcal{S}_g$ of genus g with branch data $(\gamma; m_1, \dots, m_r)$ if and only if there is a smooth epimorphism $F(\gamma; m_1, \dots, m_r) \rightarrow G$.*

Recall that the epimorphism $\eta: \Gamma \rightarrow G$ is smooth if and only if $|\eta(x_i)| = m_i = |x_i|$, for $i = 1, 2, \dots, r$. We say that a pair (Γ, G) , where G is a finite group and $\Gamma = F(\gamma; m_1, \dots, m_r)$ is a Fuchsian group, is *numerically g -admissible* if the corresponding parameters satisfy (2.2). If $\text{Epi}_o(\Gamma, G)$ is non-empty, then we say that (Γ, G) is *g -admissible*.

The following theorem gives an algebraic criterion for the equivalence of actions in terms of the associated smooth epimorphisms.

Theorem 2.2 ([30, Theorem 4]). *Let Γ be a Fuchsian group. Given a finite group G , let $\eta_1, \eta_2 \in \text{Epi}_o(\Gamma, G)$ be smooth epimorphisms, and let $(G, \mathcal{S})$ and $(G, \mathcal{S}')$ be the corresponding actions. Then $(G, \mathcal{S})$ and $(G, \mathcal{S}')$ are topologically equivalent if and only if there exists $\alpha \in \text{Aut}^+(\Gamma)$ and $a \in \text{Aut}(G)$ such that $\eta_2 = a\eta_1\alpha$.*

The smooth epimorphisms η_1, η_2 satisfying the conditions in Theorem 2.2 will be called *equivalent*, in short, we write $\eta_1 \sim \eta_2$. If $\eta_2 = a\eta_1$ for some $a \in \text{Aut}(G)$, then $\eta_1 \cong \eta_2$ are *$\text{Aut}(G)$ -equivalent*.

Let Γ be a Fuchsian group given by presentation (2.1) and G be a finite group. Every epimorphism $\eta \in \text{Epi}_o(\Gamma, G)$ is determined by the images of the $r + 2\gamma$ generators from presentation (2.1) written as a vector

$$\vec{\eta} = (\bar{x}_1, \dots, \bar{x}_r, \bar{a}_1, \dots, \bar{a}_\gamma, \bar{b}_1, \dots, \bar{b}_\gamma),$$

where $\bar{x}_i = \eta(x_i)$, $\bar{a}_j = \eta(a_j)$, $\bar{b}_j = \eta(b_j)$, for $i = 1, \dots, r$ and $j = 1, \dots, \gamma$. Hence the set $\text{Epi}_o(\Gamma, G)$ of smooth epimorphisms $\Gamma \rightarrow G$ can be identified with the set of above vectors of length $r + 2\gamma$. Let

$$GEN = (x_1, \dots, x_r, a_1, \dots, a_\gamma, b_1, \dots, b_\gamma)$$

be a vector (treated as an array) of generators of a Fuchsian group Γ . Then we can express a modified version of Theorem 2.2 as follows.

Corollary 2.3 ([8, p.22]). *The classes of topological equivalence are in one-to-one correspondence with the orbits of $\text{Aut}(G) \times \text{Aut}^+(\Gamma)$ on $\text{Epi}_o(\Gamma, G)$ given by $(a, \alpha) \cdot \vec{\eta}_1 = \vec{\eta}_2$, where the k -th entry of $\vec{\eta}_2$ is $\vec{\eta}_2[k] = a(\eta_1(\alpha(GEN[k])))$, for $k = 1, \dots, r + 2\gamma$.*

Let $\eta \in \text{Epi}_o(\Gamma, G)$. Write the multi-set $\{\eta(x_i) \mid i = 1, \dots, r\}$ in the form $\{y_1^{n_1}, \dots, y_k^{n_k}\}$, where $n_1 \leq n_2 \leq \dots \leq n_k$, where every n_i is the multiplicity of an element $y \in \{\eta(x_i) \mid i =$

$1, \dots, r\}$. Clearly, $r = n_1 + n_2 + \dots + n_k$. Denote $s(\eta) = (n_1, \dots, n_k)$. We show that $s(\eta)$ is an invariant of a class of topological equivalence provided G is abelian.

Lemma 2.4. *Let (Γ, G) be a g -admissible pair and let G be abelian. If $\eta_1, \eta_2 \in \text{Epi}_o(\Gamma, G)$ are topologically equivalent, then $s(\eta_1) = s(\eta_2)$.*

Proof. By Theorem 2.2 there exist $\alpha \in \text{Aut}^+(\Gamma)$ and $a \in \text{Aut}(G)$ such that $\eta_2 = a\eta_1\alpha$. By [21, Theorem 5.8.6] there exists a permutation $\mu \in \text{Sym}(r)$ and $g_i \in \Gamma$ such that $\alpha(x_i) = g_i x_{\mu(i)} g_i^{-1}$, for $i = 1, \dots, r$. It follows that

$$\eta_2(x_i) = a(\eta_1(\alpha(x_i))) = a(\eta_1(g_i x_{\mu(i)} g_i^{-1})) = a(\eta_1(g_i) \eta_1(x_{\mu(i)}) \eta_1(g_i)^{-1}) = a(\eta_1(x_{\mu(i)})).$$

Since $\{\eta_2(x_1), \dots, \eta_2(x_r)\} = \{a\eta_1(x_{\mu(1)}), \dots, a\eta_1(x_{\mu(r)})\}$, we have $s(\eta_1) = s(\eta_2)$. $\square$

Lemma 2.5. *Let (Γ, G) be a g -admissible pair. If $\eta_1, \eta_2 \in \text{Epi}_o(\Gamma, G)$ are topologically equivalent, then there exists $a \in \text{Aut}(G)$ such that $N_G(\langle \eta_1(x_1), \dots, \eta_1(x_r) \rangle) = aN_G(\langle \eta_2(x_1), \dots, \eta_2(x_r) \rangle)$, where $N_G(H)$ denotes the normaliser of $H \leq G$ in G .*

Proof. By Theorem 2.2 there exist $\alpha \in \text{Aut}^+(\Gamma)$ and $a \in \text{Aut}(G)$ such that $\eta_2 = a\eta_1\alpha$. Set $K = \langle x_1, \dots, x_r \rangle \leq \Gamma$. By [21, Theorem 5.8.6] we have $\alpha(N_\Gamma(K)) = N_\Gamma(K)$.

It follows that

$$\eta_2(N_\Gamma(K)) = a(\eta_1(\alpha(N_\Gamma(K)))) = a(\eta_1(N_\Gamma(K))) = aN_G(\langle \eta_1(x_1), \dots, \eta_1(x_r) \rangle).$$

On the other hand, $\eta_2(N_\Gamma(K)) = N_G(\langle \eta_2(x_1), \dots, \eta_2(x_r) \rangle)$, and we are done. $\square$

3. AUTOMORPHISM GROUP OF A PLANAR FUCHSIAN GROUP

This section is aimed to derive a finite generating set of $\text{Aut}^+(\Gamma)$ for a *planar Fuchsian group*. By a planar Fuchsian group $\Gamma = F(0; m_1, m_2, \dots, m_r)$ we mean the group with presentation

$$(3.1) \quad \langle x_1, x_2, \dots, x_r \mid x_1^{m_1} = \dots = x_r^{m_r} = 1, \prod_{j=1}^r x_j = 1 \rangle, \quad 2 + \sum_{i=1}^r \frac{1}{m_i} < r.$$

In what follows, in Sections 3 and 4, the symbol Γ will denote a planar Fuchsian group.

Denote by $\text{Sym}(r)$ the symmetric group that acts on the set $\{1, 2, \dots, r\}$ and by $\mathcal{F}_r = \mathcal{F}(\tilde{x}_1, \dots, \tilde{x}_r)$ the free group of rank r generated by the free generators $\tilde{x}_1, \dots, \tilde{x}_r$. Let $B_r = \langle \sigma_1, \dots, \sigma_{r-1} \rangle$, be the *braid group* with the defining relations $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$, for $i = 1, \dots, r-1$, and $\sigma_i \sigma_j = \sigma_j \sigma_i$, for $|i-j| \geq 2$. Let $\nu: B_r \rightarrow \text{Sym}(r)$ be the homomorphism defined by $\nu(\sigma_i) = (i, i+1)$. Since B_r acts on $\text{Sym}(r)$ through a transitive homomorphism ν and for $r \geq 3$ the image contains a 3-cycle, the image $\nu(B_r)$ is $\text{Sym}(r)$. Hence, ν is a group epimorphism $B_r \rightarrow \text{Sym}(r)$. The kernel $\ker(\nu)$ is called the *pure braid group* P_r . It is known that P_r is generated by the elements

$$A_{i,j} = \sigma_{j-1}^{-1} \sigma_{j-2}^{-1} \dots \sigma_{i+1}^{-1} \sigma_i^2 \sigma_{i+1} \dots \sigma_{j-1},$$

for $1 \leq i < j \leq r$, see [10].

Denote by $\delta: B_r \hookrightarrow \text{Aut}(\mathcal{F}_r)$ the embedding determined by $\sigma_i \mapsto c_i$, where

$$c_i: \tilde{x}_i \mapsto \tilde{x}_i \tilde{x}_{i+1} \tilde{x}_i^{-1}, \quad \tilde{x}_{i+1} \mapsto \tilde{x}_i \quad \text{and} \quad \tilde{x}_j \mapsto \tilde{x}_j \quad \text{for } j \notin \{i, i+1\}.$$

We now consider the following natural question: How do we recognize the image $\delta(B_r)$ in $\text{Aut}(\mathcal{F}_r)$? This was answered by Birman 1974 [2] and Chow 1948 [10] as follows.

Theorem 3.1 ([10]). *An automorphism $\beta \in \delta(B_r)$ if and only if there exists a permutation $\mu \in \text{Sym}(r)$ and elements $\lambda_i \in \mathcal{F}_r$ such that*

- (i) $\beta(\tilde{x}_i) = \lambda_i \tilde{x}_{\mu(i)} \lambda_i^{-1}$, $i = 1, 2, \dots, r$, and
- (ii) $\beta(\tilde{x}_1 \tilde{x}_2 \dots \tilde{x}_r) = \tilde{x}_1 \tilde{x}_2 \dots \tilde{x}_r$.

By [40, Theorem 3], every automorphism of a Fuchsian group Γ is induced by an automorphism of the free group $\mathcal{F}_r$. The following theorem characterizing the lift $\tilde{A}(\Gamma) \leq \text{Aut}(\mathcal{F}_r)$ of $\text{Aut}^+(\Gamma)$ is a restricted version of [40, Theorem 3].

Theorem 3.2 ([40, Theorem 8]). *If an automorphism $\tilde{\phi} \in \tilde{A}(\Gamma) \leq \text{Aut}(\mathcal{F}_r)$, then there exists a permutation $\mu \in \text{Sym}(r)$ satisfying $|x_{\mu(i)}| = m_i = |x_i|$ and elements $\lambda_i, \lambda \in \mathcal{F}_r$, for $i = 1, \dots, r$, such that*

- (i) $\tilde{\phi}(\tilde{x}_i) = \lambda_i \tilde{x}_{\mu(i)} \lambda_i^{-1}$, $i = 1, 2, \dots, r$, and
- (ii) $\tilde{\phi}(\tilde{x}_1 \tilde{x}_2 \dots \tilde{x}_r) = \lambda \tilde{x}_1 \tilde{x}_2 \dots \tilde{x}_r \lambda^{-1}$,
- (iii) $\omega(\lambda) = 1$, where $\omega: \mathcal{F}_r \rightarrow \Gamma$ is the natural projection.

Furthermore, given $\tilde{\phi} \in \tilde{A}(\Gamma)$ determined by $\mu \in \text{Sym}(r)$, λ and λ_i ($i = 1, \dots, r$) projects to a unique $\phi \in \text{Aut}^+(\Gamma)$.

The following lemma is proved in [38].

Lemma 3.3 ([38, Lemma 2.1]). *The lift $\tilde{I}(\Gamma) \leq \text{Aut}(\mathcal{F}_r)$ of the group of inner automorphisms $\text{Inn}(\Gamma)$ is a subgroup of $\delta(P_r)$.*

Using the aforementioned statements, we can prove the following proposition.

Proposition 3.4. *Let Γ be a planar Fuchsian group. Then $\text{Aut}^+(\Gamma)$ acts on the set $\{1, 2, \dots, r\}$ by a homomorphism $\nu^*: \text{Aut}^+(\Gamma) \rightarrow \text{Sym}(r)$ defined by setting $\nu^*(\phi) = \nu(\delta^{-1}(\tilde{\phi}))$, where $\tilde{\phi}$ is a lift of $\phi \in \text{Aut}^+(\Gamma)$.*

Proof. It is sufficient to prove that ν^* is well defined. According to Lemma 3.3, the group $\tilde{I}(\Gamma)$ of lifts of inner automorphisms of Γ is a subgroup of $\delta(P_r) = \delta(\ker(\nu))$. By Theorem 3.2 and Lemma 3.3 we have $\tilde{A}(\Gamma) \leq \tilde{I}(\Gamma) \cdot \delta(B_r) = \delta(B_r)$. It follows that two lifts $\tilde{\phi}_j$, $j = 1, 2$, of $\phi \in \text{Aut}^+(\Gamma)$ are described by the elements $\mu_j \in \text{Sym}(r)$ and $\lambda_{j,i} \in \mathcal{F}_r$, $i = 1, \dots, r$. Since $\tilde{\phi}_j$ are lifts of the same automorphism ϕ , the product $\kappa = \tilde{\phi}_1 \cdot \tilde{\phi}_2^{-1}$ is a lift of identity. Hence $\mu_1 = \mu_2$ and $\kappa \in \delta(P_r)$. It follows that $\tilde{\phi}_1 = \kappa \tilde{\phi}_2$. Since $\delta^{-1}(\kappa) \in \ker(\nu)$, we have $\nu \delta^{-1}(\tilde{\phi}_1) = \nu \delta^{-1}(\tilde{\phi}_2)$. It follows that $\nu^*(\phi)$ does not depend on the choice of the lift $\tilde{\phi}$. $\square$

Let $\mathcal{P}$ be the partition of the index set $\{1, 2, \dots, r\}$ given by the equivalence $i \cong j$ iff $m_i = m_j$. We assume that the integers m_i are ordered in non-decreasing order. Assume that there are $\ell \geq 1$ classes of $\mathcal{P}$. Then there exist integers $0 = r_0 < r_1 < r_2 < \dots < r_\ell = r$ such that each part of $\mathcal{P}$ has form

$$\mathcal{P}_j = \{r_{j-1} + 1, \dots, r_j\},$$

where $r_j - r_{j-1} = |\mathcal{P}_j|$, for some j , $\ell \geq j \geq 1$. We say that a permutation $\mu \in \text{Sym}(r)$ is $\mathcal{P}$ -invariant if $\mu(\mathcal{P}_j) = \mathcal{P}_j$, for $j = 1, \dots, \ell$. The set of all $\mathcal{P}$ -invariant permutations forms a subgroup $\text{Sym}(r, \mathcal{P}) \leq \text{Sym}(r)$ isomorphic to the direct product $\prod_{j=1}^{\ell} \text{Sym}(r_j - r_{j-1})$.

Theorem 3.5. *Let Γ be a planar Fuchsian group. Let $\omega: \tilde{A}(\Gamma) \rightarrow \text{Aut}^+(\Gamma)$ be the natural projection. Then for the epimorphism $\nu^*: \text{Aut}^+(\Gamma) \rightarrow \text{Sym}(r)$ the following statements hold:*

- (i) *The kernel $\ker \nu^*$ is the image $P_r^* = \omega\delta(P_r)$ of the pure braid group;*
- (ii) *The image $\nu^*(\text{Aut}^+(\Gamma)) = \text{Sym}(r, \mathcal{P}) \cong \prod_{j=1}^{\ell} \text{Sym}(r_j - r_{j-1})$ is the group of $\mathcal{P}$ -invariant permutations;*
- (iii) *$\text{Aut}^+(\Gamma) = \omega\delta\nu^{-1}(\text{Sym}(r, \mathcal{P}))$.*

Proof. By definition, a generator c_i , $i = 1, 2, \dots, r-1$, of $\delta(B_r)$ is in $\tilde{A}(\Gamma)$ exactly when $i \neq r_j$ for some j . Therefore, $\text{Sym}(r, \mathcal{P}) = \nu^*(\text{Aut}^+(\Gamma))$.

Let $\phi \in \ker(\nu^*)$. Then $\nu(\delta^{-1}(\tilde{\phi})) = 1$. It follows that $\tilde{\phi} \in \delta(P_r)$, and consequently, $\tilde{\phi} \in \tilde{I}(\Gamma) \cdot \delta(P_r)$, where $\tilde{I}(\Gamma)$ is the lift of the inner automorphisms (see Lemma 3.3). By [38, Lemma 2.1] we have $\tilde{I}(\Gamma) \cdot \delta(P_r) = \delta(P_r)$, and we have $\ker(\nu^*) \leq \omega\delta(P_r) = P_r^*$. Since $P_r^* \leq \ker(\nu^*)$ holds trivially, $P_r^* = \ker(\nu^*)$.

By Theorem 3.2 and by [38, Lemma 2.1] we have $\tilde{A}(\Gamma) = \tilde{I}(\Gamma) \cdot \delta\nu^{-1}(\text{Sym}(r, \mathcal{P})) = \delta\nu^{-1}(\text{Sym}(r, \mathcal{P}))$. It follows that $\text{Aut}^+(\Gamma) = \omega\delta\nu^{-1}(\text{Sym}(r, \mathcal{P}))$. $\square$

Corollary 3.6. *For a permutation $\mu \in \text{Sym}(r, \mathcal{P})$ choose an element $x_\mu \in \text{Aut}^+(\Gamma)$ with the property $\nu^*(x_\mu) = \mu$. Then $\text{Aut}^+(\Gamma) = \cup_\mu x_\mu P_r^*$. In particular, the index $[\text{Aut}^+(\Gamma) : P_r^*] = \prod_{j=1}^{\ell} (r_j - r_{j-1})!$.*

An explicit set of generators of the group $P_r^* \leq \text{Aut}^+(\Gamma)$ can be derived from the generating set $\{A_{s,t}\}$ of the pure braid group; see [38]. For $1 \leq s < t \leq r$ we set

$$(3.2) \quad \mathcal{A}_{s,t}(x_i) = \begin{cases} x_i, & i < s \text{ or } t < i, \\ x_i^{(x_s x_t)} = x_s x_t x_i x_t^{-1} x_s^{-1}, & s = i, \\ x_i^{x_s} = x_s x_i x_s^{-1}, & t = i, \\ x_i^{[x_s, x_t]} = x_s x_t x_s^{-1} x_t^{-1} x_i x_t x_s x_t^{-1} x_s^{-1}, & s < i < t \end{cases}$$

Corollary 3.7. *Let $\omega: \mathcal{F}_r \rightarrow \Gamma$ be the natural projection $\tilde{x}_i \rightarrow x_i$, $i = 1, \dots, r$. Then the group $\text{Aut}^+(\Gamma)$ is generated by the set*

$$\{\mathcal{A}_{s,t} : 1 \leq s < t \leq r\} \cup \{\omega c_i : i \in \{1, 2, \dots, r\} \setminus \{r_1, \dots, r_\ell\}\}.$$

In what follows, we set $\mathcal{H}_i = \omega c_i \in \text{Aut}^+(\Gamma)$, for $i \in \{1, 2, \dots, r\} \setminus \{r_1, \dots, r_\ell\}$.

4. ACTION OF $\text{Aut}^+(\Gamma)$ ON $\text{Epi}_o(\Gamma, G)$

Let $\Gamma = F(0; m_1, m_2, \dots, m_r)$ be a planar Fuchsian group, and let G be a finite group. The action of $\text{Aut}^+(\Gamma)$ on $\text{Epi}_o(\Gamma, G)$ is given by $\eta \mapsto \eta\alpha$, for $\eta \in \text{Epi}_o(\Gamma, G)$ and $\alpha \in \text{Aut}^+(\Gamma)$. In what follows we use the representation of epimorphisms from $\text{Epi}_o(\Gamma, G)$ by the vectors of images of the generators of Γ introduced in Section 2.

We call *vertical action* the action of the image $P_r^* \leq \text{Aut}^+(\Gamma)$ of the pure braid group on $\text{Epi}_o(\Gamma, G)$. It is generated by the permutations $\alpha_{s,t}$ corresponding to the generators $\mathcal{A}_{s,t}$:

$$(4.1) \quad \alpha_{s,t} : (\eta(x_1), \dots, \eta(x_r)) \mapsto (\eta(\mathcal{A}_{s,t}(x_1)), \dots, \eta(\mathcal{A}_{s,t}(x_r))), \text{ for } 1 \leq s < t \leq r.$$

The images $\mathcal{A}_{s,t}(x_i)$ are defined by (3.2). It follows from Theorem 3.5(iii) that the action of $\text{Aut}^+(\Gamma)$ on E is generated by permutations $\alpha_{s,t}$ and by permutations $\gamma_i \in \text{Sym}(E)$ corresponding to the elements $\mathcal{H}_i$ of $\text{Aut}^+(\Gamma)$ satisfying $\nu^*(\mathcal{H}_i) \in \text{Sym}(r, \mathcal{P})$. In particular,

$$(4.2) \quad \gamma_i: (y_1, \dots, y_i, y_{i+1}, \dots, y_r) \mapsto (y_1, \dots, y_i y_{i+1} y_i^{-1}, y_i, \dots, y_r),$$

where $i \in \{1, 2, \dots, r\} \setminus \{r_1, \dots, r_\ell\}$ and $y_i = \eta(x_i)$, $i = 1, \dots, r$.

Now we are ready to describe the action of $\text{Aut}^+(\Gamma)$ on the set $\text{Epi}_0(\Gamma, G)$.

Theorem 4.1. *Let Γ be a planar Fuchsian group with presentation (3.1) and let G be a finite group. For $\eta \in \text{Epi}_0(\Gamma, G)$ set $\vec{\eta} = (\eta(x_1), \dots, \eta(x_r))$. Two epimorphisms $\eta_1, \eta_2 \in \text{Epi}_0(\Gamma, G)$ are equivalent if and only if there exist $a \in \text{Aut}(G)$, $\pi \in \langle \alpha_{s,t} \rangle$, $1 \leq s < t \leq r$, and $\gamma \in \langle \gamma_i \rangle$, $i \in \{1, 2, \dots, r\} \setminus \{r_1, \dots, r_\ell\}$, such that $a(\vec{\eta}_1) = \gamma \pi(\vec{\eta}_2)$.*

Proof. By Theorem 2.2, η_1 is equivalent to η_2 if and only if there exist $a \in \text{Aut}(G)$ and $\alpha \in \text{Aut}^+(\Gamma)$ such that $a(\vec{\eta}_1) = \vec{\eta}_2 \alpha$. Since $P_r^* \triangleleft \text{Aut}^+(\Gamma)$, the group decomposes into a disjoint union of cosets $\text{Aut}^+(\Gamma) = \cup_{\mu \in \text{Sym}(r, \mathcal{P})} x_\mu P_r^*$, where $x_\mu \in (\nu^*)^{-1}(\mu)$. Therefore, the action of α is determined by the action of $\bar{x}_\mu \cdot \pi$, for some $\mu \in \text{Sym}(r)$ and $\pi \in \langle \alpha_{s,t} \rangle$, $1 \leq s < t \leq r$. By $\bar{x}_\mu$ we have denoted a transformation of $\text{Epi}_0(\Gamma, G)$ that corresponds to x_μ . Denote by $\tau_i = (i, i+1) \in \text{Sym}(r)$ the transposition swapping the points i and $i+1$. Each $\mu \in \text{Sym}(r, \mathcal{P})$ can be expressed as a composition $\mu = \tau_{i_1} \tau_{i_2} \dots \tau_{i_k}$, where $\{i_1, i_2, \dots, i_k\} \subseteq \{1, 2, \dots, r\} \setminus \{r_1, \dots, r_\ell\}$. Recall that $\omega: \tilde{\mathcal{F}}_r \rightarrow \Gamma$ is the natural projection taking $\tilde{x}_i \mapsto x_i$, for $i = 1, 2, \dots, r$. Then x_μ can be expressed as the product $\prod_{j=1}^k \mathcal{H}_{i_j}$. Setting $\bar{x}_\mu = \gamma = \prod_{j=1}^k \gamma_{i_j}$ we obtain the statement. $\square$

Example 4.2. Let

$$\Gamma = \langle x_1, x_2, x_3 \mid x_1^{m_1} = x_2^{m_2} = x_3^{m_3} = x_1 x_2 x_3 = 1 \rangle$$

be a triangle group with signature $(0; m_1, m_2, m_3)$. We claim that the vertical action on $\text{Epi}_0(\Gamma, G)$ coincides with the action of the group $\text{Inn}(\Gamma)$ of inner automorphisms. Indeed, the vertical action is generated by the automorphisms $\mathcal{A}_{1,2}$, $\mathcal{A}_{1,3}$, and $\mathcal{A}_{2,3}$, where the images of the generators are defined by the following table:

	x_1	x_2	x_3
$\mathcal{A}_{1,2}$	$x_1^{(x_1 x_2)}$	$x_2^{x_1}$	x_3
$\mathcal{A}_{1,3}$	$x_1^{(x_1 x_3)}$	$x_2^{[x_1, x_3]}$	$x_3^{x_1}$
$\mathcal{A}_{2,3}$	x_1	$x_2^{(x_2 x_3)}$	$x_3^{x_2}$

TABLE 1. Automorphisms of a triangle group

For a generator $x_i \in \Gamma$, denote by I_{x_i} the inner automorphism determined by conjugation by x_i . Checking the Table 1, we see that $\mathcal{A}_{1,2} = I_{x_1} I_{x_2}$, $\mathcal{A}_{1,3} = I_{x_1} I_{x_3}$, $\mathcal{A}_{2,3} = I_{x_2} I_{x_3}$. It follows that the action of the outer automorphism group $\text{Out}^+(\Gamma)$ on the set of smooth epimorphisms $\Gamma \rightarrow G$ is isomorphic to the action of a subgroup of $\text{Sym}(3)$. An epimorphic image of Γ by a smooth epimorphism in a finite group G can be identified with a regular hypermap $\mathcal{M}$ of type $\{m_1, m_2, m_3\}$. In particular, there is a bijection between

the kernels of epimorphisms in $\text{Epi}_0(\Gamma, G)$ and the isomorphism classes of hypermaps of type $\{m_1, m_2, m_3\}$. Regular hypermaps up to genus 101 were determined by Conder [14, 12, 11, 13]. Hypermaps in the Conder's lists are determined up to isomorphism and up to geometric dualities. Two hypermaps are isomorphic if and only if the images of the three generators of Γ are simultaneously conjugated in G . In view of the previous discussion, this happens if and only if the corresponding vectors of the generators are equivalent with respect to the vertical action. The correspondence between the items in Conder's lists and the equivalence classes is not perfect. While Conder lists the hypermaps up to dualities and up to taking mirror image, the corresponding actions may or may not be equivalent. For example, there are four isomorphism classes of chiral maps of genus 7 that split into two pairs of the form: a map and its mirror image. The corresponding actions have the signatures $(0; 2, 6, 9)$ and $(0; 2, 7, 7)$. In the first case, the action of the group $\text{Aut}^+(\Gamma)$ is trivial over the set of isomorphism classes of the maps, and thus the two respective maps correspond with two classes of topological equivalence. In the second case, the action of $\text{Aut}^+(\Gamma)$ on the isomorphism classes is non-trivial and the two chiral maps of type $\{7, 7\}$, distinguished just by orientation, belong to the same equivalence class.

The following observation is well-known; see Broughton [7].

Corollary 4.3. *If G is abelian, then the vertical action is always trivial. Consequently, $\text{Aut}^+(\Gamma)$ acts on $\text{Epi}_0(\Gamma, G)$ as the group of $\mathcal{P}$ -invariant permutations that permute the indices of the corresponding vectors.*

Proof. Since G is abelian, the permutations $\alpha_{s,t}$ are trivial. Furthermore, permutations γ_i just swaps the generators y_i and y_{i+1} of G . $\square$

Example 4.4. In the previous text, we have seen examples where the vertical action on $\text{Epi}_0(\Gamma, G)$ has been trivial or equal to the action of the group of inner automorphisms. In what follows, for each genus greater than 2, we give examples of actions, where the pure braid group action is essential.

Let $D_{2n} = \langle x, y \mid x^n = y^2 = (xy)^2 = 1 \rangle$ be a presentation of the dihedral group of order $2n$. Let Γ be the planar Fuchsian group with signature $(0; 2, 2, n, n)$. One can easily check that the vectors $(x^{-2}y, y, x, x)$ and (y, y, x^{-1}, x) determine smooth epimorphisms $\Gamma \rightarrow D_{2n}$. By the Riemann existence theorem (Theorem 2.1) the corresponding actions of D_{2n} exist. Using the Riemann-Hurwitz equation, we deduce that D_{2n} acts on a surface of genus $n - 1$. Clearly, no automorphism of D_{2n} takes the subsequence (x, x) onto (x^{-1}, x) ; therefore, the corresponding epimorphisms cannot be $\text{Aut}(G)$ -equivalent. Observe that the horizontal action $\langle \gamma_1, \gamma_3 \rangle$, defined in (4.2), preserves the subsequence (x, x) . Thus the two epimorphisms are not equivalent with respect to horizontal action. On the other hand, $\alpha_{2,3}(x^{-2}y, y, x, x) = (x^{-2}y, x^{-2}y, x^{-1}, x)$ and $(x^{-2}y, x^{-2}y, x^{-1}, x)^x = (y, y, x^{-1}, x)$. By Theorem 4.1, the two actions are equivalent.

The corresponding discrete actions of dihedral groups of genus ≤ 8 have identifiers O2.8, O3.22, O4.27, O5.44, O6.48, O7.91, O8.67, see census [22].

5. AUTOMORPHISMS OF NON-PLANAR FUCHSIAN GROUPS

In what follows, we introduce several families of automorphisms of a Fuchsian group Γ . They are defined by the images of generators of Γ with presentation (2.1), each of the missing generators in the expression of an automorphism is fixed.

Automorphisms $\mathcal{H}_j$:

$$(5.1) \quad \begin{aligned} \mathcal{H}_i: x_i &\mapsto x_i x_{i+1} x_i^{-1}, & x_{i+1} &\mapsto x_i, \\ &\text{for } i = 1, \dots, r-1, \text{ and } |x_i| = |x_{i+1}| \end{aligned}$$

These automorphisms come from the action of braid group. It is straightforward to verify that $\mathcal{H}_i$ is an automorphism for every admissible i . They apply if $r \geq 2$.

Automorphisms $\mathcal{B}_j$:

$$(5.2) \quad \begin{aligned} \mathcal{B}_j: a_j &\mapsto a_{j+1}, & b_j &\mapsto b_{j+1}, & a_{j+1} &\mapsto d_{j+1}^{-1} a_j d_{j+1}, \\ &b_{j+1} &\mapsto d_{j+1}^{-1} b_j d_{j+1}, \\ &\text{where } d_{j+1} = a_{j+1} b_{j+1} a_{j+1}^{-1} b_{j+1}^{-1}, \\ &\text{for } j \in \{1, \dots, \gamma-1\}. \end{aligned}$$

This family of automorphisms is introduced by Harvey in [18, p. 396-397]. They apply if $\gamma \geq 2$.

Automorphisms $\mathcal{U}_j$:

$$(5.3) \quad \begin{aligned} \mathcal{U}_1: a_1 &\mapsto a_1 b_1 \\ \mathcal{U}_2: a_1 &\mapsto a_1 b_1, & b_1 &\mapsto a_1^{-1} \\ \mathcal{U}_3: x_i &\mapsto a_2 x_i a_2^{-1}, \text{ for } i = 1, \dots, r, \\ &a_1 \mapsto a_2 a_1, & a_2 &\mapsto b_1 a_2 b_1^{-1}, & b_2 &\mapsto a_2 b_2 a_2^{-1} b_1^{-1}, \\ &a_j \mapsto a_2 a_j a_2^{-1}, & b_j &\mapsto a_2 b_j a_2^{-1} \text{ for } j = 3, \dots, \gamma \\ \mathcal{U}_4: x_r &\mapsto a_1^{-1} x_r a_1, & a_1 &\mapsto a_1^{-1} x_r^{-1} a_1 x_r a_1, & b_1 &\mapsto b_1 a_1^{-1} x_r a_1 \\ \mathcal{U}_4^*: x_1 &\mapsto b_\gamma^{-1} x_1 b_\gamma, & a_\gamma &\mapsto a_\gamma b_\gamma^{-1} x_1^{-1} b_\gamma, & b_\gamma &\mapsto b_\gamma^{-1} x_1 b_\gamma x_1^{-1} b_\gamma. \end{aligned}$$

Harvey referred the automorphisms $\mathcal{U}_i$, $i = 1, 2, 3, 4$, in [18, p. 396-397]. We derived the additional automorphism $\mathcal{U}_4^*$, a twin to $\mathcal{U}_4$. The automorphisms $\mathcal{U}_1$ and $\mathcal{U}_2$ apply if $\gamma \geq 1$. The automorphisms $\mathcal{U}_3$, $\mathcal{U}_4$ and $\mathcal{U}_4^*$ require $\gamma \geq 2$. The group $\langle \mathcal{U}_1, \mathcal{U}_2 \rangle \cong \text{Aut}^+(\pi(\mathcal{S}_1)) \cong \text{Aut}^+(\Gamma(1; -))$. Taking into account that $\pi(\mathcal{S}_1) \cong \mathbb{Z} \times \mathbb{Z}$ one can easily identify $\text{Aut}^+(\pi(\mathcal{S}_1))$ with the group of 2-dimensional unimodular matrices with determinant 1. The automorphisms $\mathcal{U}_i$, and the automorphisms $\mathcal{B}_j$, $j = 1, \dots, \gamma-1$, were used in [18, Theorem 14] to prove that there is just one class of actions of a cyclic group provided the associated Fuchsian group is a surface group. Note that the expression of $\mathcal{U}_3$ in [18, p.396] is incorrect, however, it does not effect the proof of [18, Theorem 14].

Automorphisms $\mathcal{C}_j$:

$$\begin{aligned}
(5.4) \quad & \mathcal{C}_1: a_1 \mapsto a_1 b_1^{-1} \\
& \mathcal{C}_2: b_1 \mapsto b_1 a_1 \\
& \mathcal{C}_3: a_1 \mapsto a_1 b_1^{-1} a_2 b_2 a_2^{-1}, \quad a_2 \mapsto a_2 b_2^{-1} a_2 b_1 a_2, \\
& \quad b_1 \mapsto a_2 b_2^{-1} a_2^{-1} b_1 a_2 b_2 a_2^{-1} \\
& \mathcal{C}_4: a_2 \mapsto a_2 b_2^{-1} \\
& \mathcal{C}_5: b_2 \mapsto b_2 a_2.
\end{aligned}$$

The automorphisms $\mathcal{C}_i$, $i = 1, \dots, 5$ come from McCool [32, p. 458]. The automorphisms $\mathcal{C}_i \cdot \text{Inn}(\Gamma)$, $i = 1, \dots, 5$ generate the outer automorphism group of the fundamental group Γ of a double torus. They form a basis of a set of generators of outer automorphism groups of surface groups for surfaces of genera $g \geq 2$. More precisely, we have the following

Theorem 5.1. *Let Γ be the surface group with signature $(\gamma; -)$, $\gamma > 1$. Then the automorphism group $\text{Aut}^+(\Gamma) = \langle \{\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3, \mathcal{R}\} \cup \text{Inn}(\Gamma) \rangle$, where $\mathcal{C}_i$ for $i = 1, 2, 3$ are defined by (5.4) and $\mathcal{R}$ is defined by (5.5).*

Proof. By [32, p.458] the set $\{\mathcal{R}^j \mathcal{C}_i \mathcal{R}^{-j} \cdot \text{Inn}(\Gamma)\}$, where $i = 1, 2, 3$, and $j = 0, \dots, \gamma - 1$ forms a generating set of $\text{Out}^+(\Gamma)$. $\square$

$$\begin{aligned}
(5.5) \quad & \mathcal{R}: a_j \mapsto a_{j+1}, \quad b_j \mapsto b_{j+1}, \text{ for } j = 1, \dots, \gamma - 1, \\
& a_\gamma \mapsto a_1, \quad b_\gamma \mapsto b_1.
\end{aligned}$$

Consequently, in the case when the investigated finite group is an image of a surface group under a smooth epimorphism, we have complete information to derive the classes of topological equivalence.

6. ALGORITHM

Assume a finite group $H = \langle A \rangle$, $A = \{a_1, a_2, \dots, a_m\}$, acts on a set X . The action graph $\mathcal{G} = (X; A)$ is a directed (multi)graph with set of vertices X and (x_i, x_j) is a (directed) edge of $\mathcal{G}$ if there exists $a_\ell \in A$ such that $x_j = a_\ell \cdot x_i$. Every finite action graph determines an equivalence on X defined by the connectivity components. In particular, let (Γ, G) be a g -admissible pair of groups. Let X be the set of $\text{Aut}(G)$ -classes of smooth epimorphisms $\Gamma \rightarrow G$, and $A \subset \text{Aut}^+(\Gamma)$ be a finite set of automorphisms. Denote by $[\eta]$ the $\text{Aut}(G)$ -class containing a smooth epimorphism $\eta : \Gamma \rightarrow G$. Then we have an action of $\langle A \rangle$ defined as follows: $\alpha[\eta_1] = [\eta_2]$, for $\alpha \in \langle A \rangle < \text{Aut}^+(\Gamma)$ if and only if $[\eta_2] = [\eta_1 \alpha]$. If $\langle A \cup \text{Inn}(\Gamma) \rangle = \text{Aut}^+(\Gamma)$ the equivalence relation defined by the action graph $(X; A)$ coincides with the topological equivalence. The latter holds if and only if the set $\{\alpha \cdot \text{Inn}(\Gamma) \mid \alpha \in A\}$ generates $\text{Out}^+(\Gamma)$.

Algorithm 1.**Input:** the genus, $g \geq 2$,**Output:** List of finite group actions of genus g distinguished up to topological equivalence. For signatures with $\gamma = 0$ or $r = 0$, the list is exact. Otherwise, the resulting list determines a refinement of the topological equivalence.**Step 1** Numerically solve Riemann-Hurwitz equation; find all tuples of the form

$$\mathbf{rh} := \langle g, \gamma, ord, (m_1, m_2, \dots, m_r) \rangle$$

such that $0 \leq \gamma \leq g$, $1 \leq ord \leq 84(g-1)$, $r \leq 2g+2$, m_i divides ord for $i = 1, \dots, r$ and $2g-2 = ord \cdot (2\gamma-2 + \sum_{i=1}^r (1-1/m_i))$;

Step 2 For every numerical solution $\mathbf{rh}$ construct the presentation (2.1) of the Fuchsian group $\Gamma = F(\gamma; m_1, \dots, m_r)$;**Step 3** a) If the library of small groups [1] contains groups of order ord , for every group G of order ord construct all homomorphisms $\eta: \Gamma \rightarrow G$ given by images of generators of Γ :

$$\vec{\eta} := (\eta(x_1), \dots, \eta(x_r), \eta(a_1), \dots, \eta(a_\gamma), \eta(b_1), \dots, \eta(b_\gamma));$$

(hom<> in MAGMA [4] or GroupHomomorphismByImages in GAP [15]);

b) either, construct all low-index normal subgroups of Γ (of index ord); for every subgroup $N \trianglelefteq \Gamma$ in the list set $\eta: \Gamma \rightarrow \Gamma/N$

(LowIndexNormalSubgroups in MAGMA [4] or GQuotients in GAP [15]);

Step 4 For every homomorphism η , check whether elements of $\vec{\eta}$ satisfy the relations of Γ and whether $|\eta(\Gamma)| = ord$. In this way we create a list $\mathcal{L}$ of representatives of the classes of the $\text{Aut}(G)$ -equivalence, where $G = \eta(\Gamma)$ is fixed and Γ is determined by $\mathbf{rh}$; a list $\mathcal{L}$ of tuples of the form $\langle \mathbf{rh}, \eta(\Gamma), \vec{\eta} \rangle$;**Step 5** If $\mathcal{L}$ contains a unique element $\langle \mathbf{rh}, \eta(\Gamma), \vec{\eta} \rangle$, append the tuple to the census;**Step 6** If $|\mathcal{L}| > 1$, construct an action multigraph $(\mathcal{L}; A)$ on the set of representatives $\mathcal{L}$, where $A \subset \text{Aut}^+(\Gamma)$ is chosen as follows:a) if $\gamma = 0$, then

$$A = \{\mathcal{A}_{s,t} \mid 1 \leq s < t \leq r\} \cup \{\mathcal{H}_i \mid i \in \{1, \dots, r-1\} \setminus \{r_1, \dots, r_\ell\}\},$$

see (3.2), (4.2) and (5.1);

b) if Γ is a surface group set $A = \{\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3, \mathcal{R}\}$, see (5.2) and (5.5),

c) otherwise, set

$$A = \{\mathcal{U}_1, \mathcal{U}_2, \mathcal{U}_3, \mathcal{U}_4, \mathcal{U}_4^*\} \cup \{\mathcal{B}_j \mid j = 1, \dots, \gamma-1\} \\ \cup \{\mathcal{H}_i \mid i \in \{1, \dots, r-1\} \setminus \{r_1, \dots, r_\ell\}\},$$

see (5.3), (5.2), and (5.1);

Step 7 Compute the connectivity components of the multigraph $(\mathcal{L}; A)$ and for each component store one representative.

7. COMPLETENESS OF THE CLASSIFICATION FOR $2 \leq g \leq 9$

For any genus $g \geq 2$, Theorem 4.1 and Theorem 5.1 prove that Algorithm 1 produces a complete set of representatives of classes of topological equivalence of finite groups actions on $\mathcal{S}_g$ with planar signatures and with signatures of the form $(\gamma; -)$. If both $r > 0$ and $\gamma > 0$, the algorithm determines orbits of the action of $\text{Aut}^+(G) \times H$ on $\text{Epi}_o(\Gamma, G)$ where $H \leq \text{Aut}^+(\Gamma)$ is a subgroup defined in Step 6(c) of Algorithm 1.

Theorem 7.1. *With the above notation, for $2 \leq g \leq 8$ and for every g -admissible pair (Γ, G) of groups, the orbits of $\text{Aut}^+(G) \times H$ on $\text{Epi}_o(\Gamma, G)$ determine the classes of the topological equivalence.*

Proof. Assume that Theorem 7.1 does not hold. Then there exists a g -admissible pair (Γ, G) , such that $\text{Aut}^+(G) \times \text{Aut}^+(\Gamma)$ has fewer orbits than $\text{Aut}^+(G) \times H$. This can only happen, if $\gamma > 0$, $r > 0$ and the number of orbits of $\text{Aut}^+(G) \times H$ on $\text{Epi}_o(\Gamma, G)$ is > 1 . Checking the data produced by an implementation of Algorithm 1 we identify seven such pairs (Γ, G) for $g \leq 8$, see Table 2. In all cases except O7.39 the group G is abelian. Employing Lemma 2.4, one can easily check that the vectors in Table 2 represent pairwise different classes of the topological equivalence for abelian G . In case O7.39, assume (to the contrary) that the two smooth epimorphisms represent the same equivalence class. Applying Lemma 2.5 we get

$$N_G(\langle \eta_1(x_1), \eta_1(x_2), \eta_1(x_3) \rangle) = N_G(\langle xz, z^2, xz^{-1} \rangle) \cong C_2 \times C_2,$$

and

$$N_G(\langle \eta_2(x_1), \eta_2(x_2), \eta_2(x_3) \rangle) = N_G(\langle z^2, z^2, z^2 \rangle) \cong C_2,$$

a contradiction. □

Table 2: Multiple group actions with $\gamma > 0$ and $r > 0$, $2 \leq g \leq 8$

O5.11	$(1; 2^4)$ $G \cong C_2 \times C_2$
Presentation	$G = \langle x, y \mid x^2 = y^2 = [x, y] = 1 \rangle$
Actions	$(x, x, y, y, 1, 1)$ $(x, x, x, x, y, 1)$
O5.25	$(1; 2^2)$ $G \cong C_4 \times C_2$
Presentation	$\langle x \mid x^4 = y^2 = [x, y] = 1 \rangle$
Actions	$(y, y, x, 1)$ (x^2, x^2, y, x)
O7.9	$(1; 3^6)$ $G \cong C_3$
Presentation	$G = \langle x \mid x^3 = 1 \rangle$
Actions	$(x, x, x, x, x, 1, 1)$ $(x^{-1}, x^{-1}, x^{-1}, x, x, x, 1, 1)$
O7.14	$(1; 2^6)$ $G \cong C_2 \times C_2$
Presentation	$G = \langle x, y, z \mid x^2 = y^2 = [x, y] = xyz = 1 \rangle$
Actions	$(x, x, y, y, y, y, 1, 1)$ $(x, x, y, y, z, z, 1, 1)$
Continued on next page	

Table 2: Multiple group actions with $\gamma > 0$ and $r > 0$, $2 \leq g \leq 8$

	$(x, x, x, x, x, x, y, 1)$	
O7.16	$(1; 4^4)$	$G \cong C_4$
Presentation	$G = \langle x \mid x^4 = 1 \rangle$	
Actions	$(x, x, x, x, 1, 1)$ $(x^{-1}, x^{-1}, x, x, 1, 1)$	
O7.39	$(1; 2^3)$	$G \cong D_8$
Presentation	$G = \langle x, z \mid x^2 = z^4 = (xz)^2 = 1 \rangle$	
Actions	$(xz, z^2, xz^{-1}, x, 1)$ $(z^2, z^2, z^2, xz^{-1}, x)$	
O8.12	$(1; 2^7)$	$G \cong C_2 \times C_2$
Presentation	$G = \langle x, y, z \mid x^2 = y^2 = [x, y] = xyz = 1 \rangle$	
Actions	$(x, y, y, y, z, z, z, 1, 1)$ $(x, y, z, z, z, z, z, 1, 1)$	

Theorem 7.2. *With the above notation, for $g = 9$ and for every g -admissible pair (Γ, G) of groups, except O9.109, the orbits of $\text{Aut}^+(\Gamma) \times H$ on $\text{Epi}_o(\Gamma, G)$ determine the classes of the topological equivalence. There are at most two classes of topological equivalence in case O9.109.*

Proof. Assume to the contrary that Theorem 7.2 does not hold. Then there exists a 9-admissible pair (Γ, G) , such that $\text{Aut}^+(\Gamma) \times \text{Aut}^+(\Gamma)$ has fewer orbits than $\text{Aut}^+(\Gamma) \times H$. This can only happen, if $\gamma > 0$, $r > 0$ and the number of orbits of $\text{Aut}^+(\Gamma) \times H$ on $\text{Epi}_o(\Gamma, G)$ is greater than one. Checking the data produced by an implementation of Algorithm 1 we identified 13 such pairs (Γ, G) , listed in Table 3.

In cases O9.10, O9.15, O9.18, O9.20, O9.26, O9.52, and O9.54 the group G is abelian. Employing Lemma 2.4, it can be checked that in each of the listed cases the corresponding vectors in Table 3 represent pairwise different classes of the topological equivalence for abelian G .

In what follows we shall discuss the remaining cases:

Case O9.50: Since G is abelian, Lemma 2.4 distinguishes O9.50.1 from the other two actions. Now we shall distinguish O.50.2 from O.50.3. Denote by η_1 and η_2 , the epimorphisms $\Gamma \rightarrow \langle x \rangle \times \langle y \rangle \cong C_4 \times C_2$, determined by the vectors $(x^2, x^2, y, y, x, 1)$ and $(yx^2, yx^2, y, y, x, 1)$, respectively. Assume, to the contrary, that $\eta_1 \sim \eta_2$. By Theorem 2.2 there exist $\alpha \in \text{Aut}^+(\Gamma)$ and $a \in \text{Aut}(G)$ such that $\eta_1 \alpha|_{\{x_1, x_2, x_3, x_4\}} = a \eta_2|_{\{x_1, x_2, x_3, x_4\}}$. Since G is abelian, on the left-hand side we get a vector that is a re-arrangement of (x^2, x^2, y, y) . It follows that either $a(y) = y$ and $a(yx^2) = x^2$, or $a(y) = x^2$ and $a(yx^2) = y$. There are exactly three involutions in G , namely y , x^2 , and yx^2 , arranged in the orbits $\{x^2\}$ and $\{y, yx^2\}$ in the action of $\text{Aut}(G)$. It follows that $a^{-1}(x^2) = x^2$, a contradiction with $a^{-1}(x^2) \in \{y, yx^2\}$.

Case O9.51: For each of the three actions we compute the normaliser of the subgroup generated by the images of the elliptic elements by the corresponding epimorphisms $\Gamma \rightarrow G$. We get $N_G(\langle x, y \rangle) \cong D_8$, $N_G(\langle x \rangle) \cong C_2$, and $N_G(\langle (xy)^2, x \rangle) \cong C_2 \times C_2$, respectively. By Lemma 2.5, the actions are pairwise non-equivalent.

Case O9.69: By Table 3, there are two actions to consider. Denote by η_1 and η_2 , the respective epimorphisms $\Gamma \rightarrow \langle x, y \mid x^5 = y^2 = (xy)^2 = 1 \rangle \cong D_{10}$, determined by the vectors $(x, x^{-1}, y, 1)$ and (x, x^{-2}, x^{-2}, y) . Assume to the contrary $\eta_1 \sim \eta_2$. By Theorem 2.2 there exist $\alpha \in \text{Aut}^+(\Gamma)$ and $a \in \text{Aut}(G)$ such that $a\eta_1\alpha|_{\{x_1, x_2\}} = \eta_2|_{\{x_1, x_2\}}$. By [40, Theorem 8], every automorphism $\alpha \in \text{Aut}^+(\Gamma)$ takes an elliptic element to a conjugate of an elliptic element. Taking into the account $G \cong D_{10}$ we get:

$$(x, x^{-2}) \in \{(a(x), a(x^{-1})), (a(x^{-1}), a(x)), (a(x), a(x)), (a(x^{-1}), a(x^{-1}))\}.$$

Since x is element of order five, $x \neq x^{-1}$. Thus $(x, x^{-2}) \neq (a(x), a(x))$, and $(x, x^{-2}) \neq (a(x^{-1}), a(x^{-1}))$. It follows that either $(a(x), a(x^{-1})) = (x, x^{-2})$ or $(a(x^{-1}), a(x)) = (x, x^{-2})$. Clearly, $a(x) = x^i$ for $i \in \{1, 2, 3, 4\}$. We conclude that $(x^{\pm i}, x^{\mp i}) = (x, x^{-2})$, which is impossible.

Case O9.102: For the two actions we compute the normaliser of the subgroup generated by the images of the elliptic elements by the corresponding epimorphisms $\Gamma \rightarrow G$. We get $|N_G(\langle y \rangle)| = 8$, while $|N_G(\langle (yx)^2, x^2 \rangle)| = 16$. By Lemma 2.5, the actions are non-equivalent.

Case O9.104: Denote by η_1 and η_2 , the epimorphisms $\Gamma \rightarrow \langle x \rangle \times \langle y \rangle \cong C_8 \times C_2$, determined by the vectors $(y, y, x, 1)$ and (x^4, x^4, y, x) , respectively. Assume that $\eta_1 \sim \eta_2$. By Theorem 2.2 there exist $\alpha \in \text{Aut}^+(\Gamma)$ and $a \in \text{Aut}(G)$ such that $\eta_1\alpha|_{\{x_1, x_2\}} = a\eta_2|_{\{x_1, x_2\}}$. Since G is abelian, on the left-hand side we get the vector (y, y) , while on the right-hand side we get $(a(x^4), a(x^4))$. It follows that $a(x^4) = y$. There are exactly three involutions in G , namely y , x^4 , and yx^4 , arranged in the orbits $\{x^4\}$ and $\{y, yx^4\}$ in the action of $\text{Aut}(G)$. Similarly as above, $x^4 = a^{-1}(y)$ implying $y = x^4$, a contradiction. $\square$

Table 3: Multiple group actions with $\gamma > 0$ and $r > 0$, $g = 9$

O9.10	$(1; 3^8)$	$G \cong C_3$
Presentation	$G = \langle x \mid x^3 = 1 \rangle$	
Actions	O9.10.1 $(x, x^{-1}, x^{-1}, x^{-1}, x^{-1}, x^{-1}, x^{-1}, 1, 1)$ O9.10.2 $(x, x, x, x, x^{-1}, x^{-1}, x^{-1}, x^{-1}, 1, 1)$	
O9.15	$(2; 2^4)$	$G \cong C_2 \times C_2$
Presentation	$G = \langle x, y \mid x^2 = y^2 = 1, [x, y] = 1 \rangle$	
Actions	O9.15.1 $(x, x, y, y, 1, 1, 1, 1)$ O9.15.2 $(x, x, x, x, y, 1, 1, 1)$	
O9.18	$(1; 2^8)$	$G \cong C_2 \times C_2$
Presentation	$G = \langle x, y, z \mid x^2 = y^2 = z^2 = 1, [x, y] = 1, xyz = 1 \rangle$	
Actions	O9.18.1 $(x, x, y, y, y, y, y, 1, 1)$ O9.18.2 $(x, x, x, x, y, y, y, y, 1, 1)$ O9.18.3 $(z, z, x, x, y, y, y, y, 1, 1)$ O9.18.4 $(x, x, x, x, x, x, x, y, 1)$	
O9.20	$(1; 2^2, 4^4)$	$G \cong C_4$
Presentation	$G = \langle x \mid x^4 = 1 \rangle$	
Actions	O9.20.1 $(x^2, x^2, x, x, x, x, 1, 1)$ O9.20.2 $(x^2, x^2, x^{-1}, x^{-1}, x, x, 1, 1)$	
Continued on next page		

Table 3: Multiple group actions with $\gamma > 0$ and $r > 0$, $g = 9$

O9.26	$(1; 5^4)$	$G \cong C_5$
Presentation	$G = \langle x \mid x^5 = 1 \rangle$	
Actions	O9.26.1 $(x^3, x^{-1}, x^{-1}, x^{-1}, 1, 1)$ O9.26.2 $(x, x, x^{-1}, x^{-1}, 1, 1)$ O9.26.3 $(x^2, x^3, x, x^{-1}, 1, 1)$	
O9.50	$(1; 2^4)$	$G \cong C_4 \times C_2$
Presentation	$G = \langle x, y \mid x^4 = y^2 = 1, [x, y] = 1 \rangle$	
Actions	O9.50.1 $(y, y, y, y, x, 1)$ O9.50.2 $(x^2, x^2, y, y, x, 1)$ O9.50.3 $(yx^2, yx^2, y, y, x, 1)$	
O9.51	$(1; 2^4)$	$G \cong D_8$
Presentation	$G = \langle x, y \mid x^2 = y^2 = (xy)^4 = 1 \rangle$	
Actions	O9.51.1 $(x, x, y, y, 1, 1)$ O9.51.2 $(x, x, x, x, y, 1)$ O9.51.3 $((xy)^2, (xy)^2, x, x, y, 1)$	
O9.52	$(1; 2^4)$	$G \cong C_2 \times C_2 \times C_2$
Presentation	$G = \langle x, y, z \mid x^2 = y^2 = z^2 = 1, [x, y] = [x, z] = [y, z] = 1 \rangle$	
Actions	O9.52.1 $(xyz, x, y, z, 1, 1)$ O9.52.2 $(x, x, y, y, z, 1)$ O9.52.3 (x, x, x, x, y, z)	
O9.54	$(1; 2, 4^2)$	$G \cong C_4 \times C_2$
Presentation	$G = \langle x, y \mid x^4 = y^2 = 1, [x, y] = 1 \rangle$	
Actions	O9.54.1 $(y, x, yx^{-1}, 1, 1)$ O9.54.2 $(x^2, x, x, yx^{-1}, 1)$	
O9.69	$(1; 5^2)$	$G \cong D_{10}$
Presentation	$G = \langle x, y \mid x^5 = y^2 = (xy)^2 = 1 \rangle$	
Actions	O9.69.1 $(x, x^{-1}, y, 1)$ O9.69.2 (x, x^{-2}, x^{-2}, y)	
O9.102	$(1; 2^2)$	$G \cong (C_4 \times C_2) \rtimes C_2$
Presentation	$G = \langle x, y, z \mid x^4 = y^2 = 1, [x, y] = 1, (zx)^2 = (zy)^2 = 1 \rangle$	
Actions	O9.102.1 $(y, y, x, 1)$ O9.102.2 $((yx)^2, x^2, y, x)$	
O9.104	$(1; 2^2)$	$G \cong C_8 \times C_2$
Presentation	$G = \langle x, y \mid x^8 = y^2 = 1, [x^2, y] = 1, (xy)^4 = 1 \rangle$	
Actions	O9.104.1 $(y, y, x, 1)$ O9.104.2 (x^4, x^4, y, x)	
O9.109	$(1; 2^2)$	$G \cong C_2 \times D_8$
Presentation	$G = \langle x, y, z, w \mid x^2 = y^2 = z^2 = w^2 = 1, [x, y] = [y, z] = [x, z] = 1, (zw)^4 = 1 \rangle$	
Actions	O9.109.1 (x, y, z, w) O9.109.2 (wx, yw, z, w)	

8. CLASSIFICATION OF FINITE GROUPS OF SMALL GENERA

The list of groups of genus at most 9 is attached in the Appendix. Full data containing representatives of each class of topological equivalence in JSON format are available at <https://www.savbb.sk/~karabas/science/discactions.html>

For every genus g , $2 \leq g \leq 9$, the rows correspond to g -admissible pairs (Γ, G) . Each row consists of the entries with the following meaning:

- **Ref:** a unique identifier of the form `0g.integer`, it allows us to find the representatives of the classes of topological equivalence in the database, the identifier comes from the census at webpage [22]
- **Signature:** identifies the Fuchsian group Γ ,
- **SMG id:** Identifier of the group G in the library of small groups [1, 4, 15],
- **Structure:** Structure description of G (computed by GAP [15]),
- **epi:** $= |\text{Epi}_0(\Gamma, G)|$, the number of smooth epimorphisms $\Gamma \rightarrow G$
- **autg:** the number of orbits of the action of $\text{Aut}(G)$ on $\text{Epi}_0(\Gamma, G)$, note that $\text{autg} = \text{epi}/|\text{Aut}(G)|$,
- **top:** the number of classes of topological equivalence.

The data were obtained by two independent implementations of Algorithm 1 in MAGMA [4]. The correctness of the outputs has been checked in two ways. First, all multiple actions of g -admissible pairs for $2 \leq g \leq 9$ were checked manually. Secondly, we have compared the obtained data with the published lists of group actions. Independent comparison of outputs was used to check for cyclic actions. For cyclic groups the function enumerating the number of smooth epimorphisms is known, see [29, 33].

We shall discuss our findings below. By the phrase “an admissible pair of groups (Γ, G) has a multiple action on a surface $\mathcal{S}_g$ ” we mean that there are at least two classes of topological equivalence of the actions of G on $\mathcal{S}_g$ with signature determining Γ .

g=2. There are 21 admissible pairs (Γ, G) , each defines a unique equivalence class with respect to the topological equivalence. In particular, $\sim_{\text{top}} = \sim_{\mathcal{H}^1}$ for actions of genus 2. Our list of classes coincides with the one derived by Broughton in [7].

g=3. There are 49 admissible pairs (Γ, G) giving rise to 55 classes of topological equivalence. Four pairs give rise to two non-equivalent actions, namely, O3.9, O3.11, O3.16 and O.23. The pair $(\Gamma(0; 2^2, 4^2), C_4 \times C_2)$ gives rise to three non-equivalent actions. In the list derived by Broughton [7, p.255] one admissible pair, namely, $(\Gamma(0; 2, 3, 12), \text{SL}(2, 3) : C_2)$ is missing (O3.47 in our notation). It gives rise to one equivalence class. There is also apparent misprint in the signature of item 3.ag [7, p.255]. The correct signature should be $(0; 3, 3, 7)$, O3.37 in our list. In all other cases our data fits with the data derived by Broughton. Comparison with [5, p.91] shows that $\sim_{\text{top}} = \sim_{\mathcal{H}^1}$ provided we are restricted to actions of genus 3.

g=4. There are 64 admissible pairs (Γ, G) giving rise to 75 classes of topological equivalence. Our list of actions fit with the one derived by Bogopolski in [3]. Genus four is the smallest genus, where the equivalences $\sim_{\text{top}}$ and $\sim_{\mathcal{H}^1}$ do not coincide. Breuer in [5, p.91] identified 73 classes with respect to $\sim_{\mathcal{H}^1}$. The difference between $\sim_{\text{top}}$ and $\sim_{\mathcal{H}^1}$ is explained by Broughton in [8, p.4] as follows: “The smallest genus example where the Abelian differentials fail to distinguish topologically distinct actions occurs in genus 4

with $G = C_5$ and 4 branch points.” In our list the corresponding item is O4.13, where the pair $(\Gamma(0; 5^4), C_5)$ gives rise to three actions of genus four that are not distinguished by $\sim_{\mathcal{H}^1}$. Kimura in [24] presented another list of actions of genus four, distinguished up to relation $\sim_{\mathcal{H}^1}$. Breuer [5, p.90] noted that one class is doubled, but another one is missing in the Kimura’s list.

g=5. There are 93 admissible pairs (Γ, G) giving rise to 118 classes of topological equivalence. Breuer [5, p.90] identifies 116 classes of $\sim_{\mathcal{H}^1}$. Hence some of the classes of topological equivalence are not distinguished by $\sim_{\mathcal{H}^1}$. The number of $\sim_{\mathcal{H}^1}$ classes fits with the one derived in [26]. Maximum multiplicity of actions of topological equivalence is five and it is achieved by the pair $(\Gamma(0; 2^3, 4^2), C_4 \times C_2)$ identified as O5.30 in our list. The largest order of a group with multiple action is 48. In particular, a group of order 48 (identified as O5.82), acts with signature $(0; 3, 4, 4)$ in two different ways. In the list derived by Conder [11] there are two non-isomorphic regular hypermaps corresponding to these actions, (RPH5.3 and RPH5.4).

g=6. There are 87 admissible pairs (Γ, G) giving rise to 108 classes of topological equivalence. Breuer [5, p.90] identifies 105 classes of $\sim_{\mathcal{H}^1}$. Maximal multiplicity of the action is four, achieved by the actions of cyclic group C_7 with signature $(0; 7^4)$, referenced O6.26. The largest group with multiple action is the cyclic group C_{15} acting with signature $(0; 5, 15, 15)$. The two corresponding regular hypermaps have reference numbers RPH6.12 and RPH6.13 in [11].

g=7. There are 148 admissible pairs (Γ, G) giving rise to 210 classes of topological equivalence. Breuer [5, p.90] identifies 208 classes of $\sim_{\mathcal{H}^1}$. Maximal multiplicity is eight achieved by the actions of $C_4 \times C_2$ with signature $(0; 2^4, 4^2)$, with the identifier O7.48. The largest group with multiple action has order 54 acting with signature $(0; 2, 6, 9)$. It gives rise to two non-equivalent actions. The corresponding regular map is chiral, therefore it appears in two non-isomorphic enantiomers, reference C7.1 in Conder’s list [14].

g=8. There are 108 admissible pairs (Γ, G) giving rise to 150 classes of topological equivalence. Breuer [5, p.90] identifies 141 classes of $\sim_{\mathcal{H}^1}$. Maximal multiplicity is six achieved by the actions of C_{10} , referenced O8.44. The largest group with multiple action is $\text{PSL}(3, 2): C_2$ of order 336 acting with signature $(0; 2, 3, 8)$. It gives rise to two non-equivalent actions. The corresponding regular maps have references R8.1 and R8.2 in Conder’s list [13].

g > 8. Computations of classes of topological equivalence for higher genera and verification of outputs are still running (in August 2023). For most recent results see [22]. For example, we have found that our list of admissible pairs of groups diverges from the Breuer’s list [5, p.91]. In particular, we got different numbers of admissible pairs (Γ, G) for genera 9, 13, 17. Our numbers are 270, 454, 746, which do not fit with corresponding values in [5, p.91]. For instance, comparing our data with these produced by Breuer we have discovered that the pairs $(\Gamma(1; 2^2), C_4 \times C_4)$ and $(\Gamma(1; 2^2), C_4 \times C_2^2)$ are missing the list of 9-admissible pairs in Breuer’s data (personal communication). The same problem also appears in on-line database [35].

Table 4 summarises the above discussion, **Adm**, **H1*** and **top** denote the number of admissible pairs (Γ, G) , the number of $\sim_{\mathcal{H}^1}$ classes and the number of $\sim_{top}$ classes, respectively.

Table 4: The numbers of finite group actions on Riemann surfaces of genera $2 \leq g \leq 9$

genus	Adm	H1	top
2	21	21	21
3	49	55	55
4	64	73	75
5	93	116	118
6	87	105	108
7	148	208	210
8	108	141	150
9	270	428 [†]	441

ACKNOWLEDGEMENT

The three authors were supported by the grant GACR 20-15576S. The first and the second authors were supported by the grant APVV-19-0308 of the Slovak Research and Development Agency and by the grant VEGA 2/0078/20. The authors are very grateful to Robert Jajcay and to the anonymous referee(s) for their useful comments.

REFERENCES

- [1] H. U. Besche, B. Eick, and E. O'Brien. The Small Groups library. http://www.icm.tu-bs.de/ag_algebra/software/small/, 2014. seen August 23, 2022.
- [2] J. S. Birman. *Braids, Links and Mapping Class Groups*. Princeton University Press Princeton, N.J, 1974. ISBN 0691081492.
- [3] O. V. Bogopolski. Classification of actions of finite groups on orientable surface of genus four. *Sib. Adv. Math*, 7:9–38, 1997.
- [4] W. Bosma, J. Cannon, and C. Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3-4):235–265, 1997. ISSN 0747-7171. <https://doi.org/10.1006/jsco.1996.0125>.
- [5] T. Breuer. *Characters and automorphism groups of compact Riemann surfaces*. Number 280. Cambridge University Press, 2000.
- [6] M. R. Bridson and K. Vogtmann. Automorphism groups of free groups, surface groups and free abelian groups. [arXiv:math/0507612](https://arxiv.org/abs/math/0507612) [math.GR], 2005.
- [7] S. A. Broughton. Classifying finite group actions on surfaces of low genus. *Journal of Pure and Applied Algebra*, 69(3):233–270, 1991.

*this value corresponds to the number in column ‘Orb’ in [5, Table 13]

[†]this number is not correct in [5], see the discussion above

- [8] S. A. Broughton. Equivalence of finite group actions on Riemann surfaces and algebraic curves. *Automorphisms of Riemann Surfaces, Subgroups of Mapping Class Groups and Related Topics*, 776:89, 2022.
- [9] S. A. Broughton, J. Paulhus, and A. Wootton. Future directions in automorphisms of surfaces, graphs, and other related topics. *Automorphisms of Riemann Surfaces, Subgroups of Mapping Class Groups and Related Topics*, 776:37, 2022.
- [10] W.-L. Chow. On the algebraical braid group. *Annals of Mathematics*, 49(3):654–658, 1948. ISSN 0003486X. URL <http://www.jstor.org/stable/1969050>.
- [11] M. D. E. Conder. Orientable proper regular hypermaps of genus 2 to 101, 2006. URL <https://www.math.auckland.ac.nz/~conder/OrientableProperHypermaps101.txt>. Accessed: 2023-08-08.
- [12] M. D. E. Conder. Chiral proper orientably-regular hypermaps of genus 2 to 101, 2006. URL <https://www.math.auckland.ac.nz/~conder/ChiralProperHypermaps101.txt>. Accessed: 2023-08-08.
- [13] M. D. E. Conder. Regular orientable maps of genus 2 to 301, 2012. URL <https://www.math.auckland.ac.nz/~conder/RegularOrientableMaps301.txt>. seen August 23, 2022.
- [14] M. D. E. Conder. Chiral rotary maps of genus 2 to 301, 2013. URL <http://www.math.auckland.ac.nz/~conder/ChiralMaps301.txt>. Accessed: 2023-08-08.
- [15] GAP. *GAP – Groups, Algorithms, and Programming, Version 4.12.2*. The GAP Group, 2022. URL <https://www.gap-system.org>.
- [16] S. Gervais. A finite presentation of the mapping class group of a punctured surface. *Topology*, 40(4):703–725, 2001.
- [17] W. J. Harvey. Cyclic groups of automorphisms of a compact Riemann surface. *The Quarterly Journal of Mathematics*, 17(1):86–97, 1966.
- [18] W. J. Harvey. On branch loci in Teichmüller space. *Transactions of the American Mathematical Society*, 153:387–399, 1971.
- [19] A. Hatcher and W. Thurston. A presentation for the mapping class group of a closed orientable surface. *Topology*, 19(3):221–237, 1980.
- [20] S. P. Humphries. Generators for the mapping class group. In *Topology of low-dimensional manifolds*, pages 44–47. Springer, 1979.
- [21] G. A. Jones and D. Singerman. *Complex functions: an algebraic and geometric viewpoint*. Cambridge University Press, 1987.
- [22] J. Karabáš, R. Nedela, and M. Skyvová. Discrete group actions on orientable surfaces. URL <https://www.savbb.sk/~karabas/science/discactions.html>. Accessed: 2023-08-08.
- [23] J. Karabáš, R. Nedela, and M. Skyvová. Computing equivalence classes of finite group actions on orientable surfaces. *Journal of Pure and Applied Algebra*, 228(6):107578, 2024.
- [24] H. Kimura. Classification of automorphism groups, up to topological equivalence, of compact Riemann surfaces of genus 4. *Journal of Algebra*, 264(1):26–54, 2003. ISSN 0021-8693. [https://doi.org/10.1016/S0021-8693\(03\)00138-8](https://doi.org/10.1016/S0021-8693(03)00138-8). URL <https://www.sciencedirect.com/science/article/pii/S0021869303001388>.

- [25] M. Korkmaz. Generating the surface mapping class group by two elements. *Transactions of the American Mathematical Society*, 357(8):3299–3310, 2005.
- [26] A. Kuribayashi and H. Kimura. Automorphism groups of compact Riemann surfaces of genus five. *Journal of Algebra*, 134(1):80–103, 1990. ISSN 0021-8693. [https://doi.org/10.1016/0021-8693\(90\)90212-7](https://doi.org/10.1016/0021-8693(90)90212-7). URL <https://www.sciencedirect.com/science/article/pii/0021869390902127>.
- [27] C. Labruere and L. Paris. Presentations for the punctured mapping class groups in terms of artin groups. *Algebraic & Geometric Topology*, 1(1):73–114, 2001.
- [28] W. B. R. Lickorish. A finite set of generators for the homeotopy group of a 2-manifold. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 60, pages 769–778. Cambridge University Press, 1964.
- [29] V. A. Liskovets. A multivariate arithmetic function of combinatorial and topological significance. *Integers*, 10(1), 2010. <https://doi.org/10.1515/integ.2010.012>.
- [30] E. K. Lloyd. Riemann surface transformation groups. *Journal of Combinatorial Theory, Ser. A*, 13:17–27, 1972.
- [31] J. McCool. Some finitely presented subgroups of the automorphism group of a free group. *Journal of Algebra*, 35(1-3):205–213, 1975.
- [32] J. McCool. Generating the mapping class group. *Publications Mathematiques*, 40:457–468, 1996.
- [33] A. Mednykh and R. Nedela. Enumeration of unrooted maps of a given genus. *Journal of Combinatorial Theory, Series B*, 96(5):706–729, 2006.
- [34] J. Nielsen et al. Untersuchungen zur topologie der geschlossenen zweiseitigen flächen. *Acta Mathematica*, 50:189–358, 1927.
- [35] J. Paulhus. LMFDB: Families of higher genus curves with automorphisms. URL <https://www.lmfdb.org/HigherGenus/C/Aut/>. Accessed: 2023-08-08.
- [36] J. Paulhus. A database of group actions on Riemann surfaces. In *International Conference on Birational Geometry, Kaehler-Einstein Metrics and Degenerations*, pages 693–708. Springer, 2019.
- [37] M. Stukow. A finite presentation for the mapping class group of a nonorientable surface with Dehn twists and one crosscap slide as generators. *Journal of Pure and Applied Algebra*, 218(12):2226–2239, 2014.
- [38] A. O. B. M. Tap. Automorphisms of Fuchsian groups of genus zero. *Pertanika*, (1):115–123, 1988.
- [39] B. Wajnryb. A simple presentation for the mapping class group of an orientable surface. *Israel Journal of Mathematics*, 45(2):157–174, 1983.
- [40] H. Zieschang. Über automorphismen ebener diskontinuierlicher gruppen. *Mathematische Annalen*, 166(2):148–167, 1966.
- [41] H. Zieschang, E. Vogt, and H.-D. Coldewey. *Surfaces and planar discontinuous groups*, volume 835 of *Lecture Notes in Mathematics*. Springer, 2006.

APPENDIX A. CENSUS OF ACTIONS OF FINITE GROUPS
ON RIEMANN SURFACES OF GENERA 2 TO 9

Table 5: Census of finite group actions on Riemann surfaces of genus $g = 2$

Ref	Signature	SMG id	Structure	epi	autg	top
O2.1	$(2; -)$	$\langle 1, 1 \rangle$	1	1	1	1
O2.2	$(1; 2, 2)$	$\langle 2, 1 \rangle$	C_2	4	4	1
O2.3	$(0; 2^6)$	$\langle 2, 1 \rangle$	C_2	1	1	1
O2.4	$(0; 3, 3, 3, 3)$	$\langle 3, 1 \rangle$	C_3	6	3	1
O2.5	$(0; 2^5)$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	60	10	1
O2.6	$(0; 2, 2, 4, 4)$	$\langle 4, 1 \rangle$	C_4	2	1	1
O2.7	$(0; 5, 5, 5)$	$\langle 5, 1 \rangle$	C_5	12	3	1
O2.8	$(0; 2, 2, 3, 3)$	$\langle 6, 1 \rangle$	S_3	12	2	1
O2.9	$(0; 2, 2, 3, 3)$	$\langle 6, 2 \rangle$	C_6	2	1	1
O2.10	$(0; 3, 6, 6)$	$\langle 6, 2 \rangle$	C_6	2	1	1
O2.11	$(0; 2, 2, 2, 4)$	$\langle 8, 3 \rangle$	D_8	24	3	1
O2.12	$(0; 4, 4, 4)$	$\langle 8, 4 \rangle$	Q_8	24	1	1
O2.13	$(0; 2, 8, 8)$	$\langle 8, 1 \rangle$	C_8	4	1	1
O2.14	$(0; 2, 5, 10)$	$\langle 10, 2 \rangle$	C_{10}	4	1	1
O2.15	$(0; 2, 2, 2, 3)$	$\langle 12, 4 \rangle$	D_{12}	36	3	1
O2.16	$(0; 3, 4, 4)$	$\langle 12, 1 \rangle$	$C_3 : C_4$	12	1	1
O2.17	$(0; 2, 6, 6)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	12	1	1
O2.18	$(0; 2, 4, 8)$	$\langle 16, 8 \rangle$	QD_{16}	16	1	1
O2.19	$(0; 3, 3, 4)$	$\langle 24, 3 \rangle$	$SL(2, 3)$	24	1	1
O2.20	$(0; 2, 4, 6)$	$\langle 24, 8 \rangle$	$(C_6 \times C_2) : C_2$	24	1	1
O2.21	$(0; 2, 3, 8)$	$\langle 48, 29 \rangle$	$GL(2, 3)$	48	1	1

Table 6: Census of finite group actions on Riemann surfaces of genus $g = 3$

Ref	Signature	SMG id	Structure	epi	autg	top
O3.1	$(3; -)$	$\langle 1, 1 \rangle$	1	1	1	1
O3.2	$(2; -)$	$\langle 2, 1 \rangle$	C_2	15	15	1
O3.3	$(1; 2, 2, 2, 2)$	$\langle 2, 1 \rangle$	C_2	4	4	1
O3.4	$(0; 2^8)$	$\langle 2, 1 \rangle$	C_2	1	1	1
O3.5	$(1; 3, 3)$	$\langle 3, 1 \rangle$	C_3	18	9	1
O3.6	$(0; 3^5)$	$\langle 3, 1 \rangle$	C_3	10	5	1
O3.7	$(1; 2, 2)$	$\langle 4, 1 \rangle$	C_4	12	6	1
O3.8	$(1; 2, 2)$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	36	6	1
O3.9	$(0; 2^6)$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	180	30	2
O3.10	$(0; 2^3, 4^2)$	$\langle 4, 1 \rangle$	C_4	2	1	1
O3.11	$(0; 4, 4, 4, 4)$	$\langle 4, 1 \rangle$	C_4	8	4	2
O3.12	$(1; 3)$	$\langle 6, 1 \rangle$	S_3	18	3	1
O3.13	$(0; 2^4, 3)$	$\langle 6, 1 \rangle$	S_3	54	9	1

Continued on next page

Table 6 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O3.14	(0; 2, 3, 3, 6)	$\langle 6, 2 \rangle$	C_6	2	1	1
O3.15	(0; 2, 2, 6, 6)	$\langle 6, 2 \rangle$	C_6	2	1	1
O3.16	(0; 7, 7, 7)	$\langle 7, 1 \rangle$	C_7	30	5	2
O3.17	(1; 2)	$\langle 8, 3 \rangle$	D_8	24	3	1
O3.18	(1; 2)	$\langle 8, 4 \rangle$	Q_8	24	1	1
O3.19	(0; 2^5)	$\langle 8, 3 \rangle$	D_8	240	30	1
O3.20	(0; 2^5)	$\langle 8, 5 \rangle$	$C_2 \times C_2 \times C_2$	1680	10	1
O3.21	(0; 2, 2, 4, 4)	$\langle 8, 2 \rangle$	$C_4 \times C_2$	32	4	3
O3.22	(0; 2, 2, 4, 4)	$\langle 8, 3 \rangle$	D_8	16	2	1
O3.23	(0; 4, 8, 8)	$\langle 8, 1 \rangle$	C_8	8	2	2
O3.24	(0; 3, 9, 9)	$\langle 9, 1 \rangle$	C_9	12	2	1
O3.25	(0; 2, 2, 3, 3)	$\langle 12, 3 \rangle$	A_4	72	3	1
O3.26	(0; 2, 2, 2, 6)	$\langle 12, 4 \rangle$	D_{12}	36	3	1
O3.27	(0; 4, 4, 6)	$\langle 12, 1 \rangle$	$C_3 : C_4$	12	1	1
O3.28	(0; 3, 4, 12)	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O3.29	(0; 2, 12, 12)	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O3.30	(0; 2, 7, 14)	$\langle 14, 2 \rangle$	C_{14}	6	1	1
O3.31	(0; 2, 2, 2, 4)	$\langle 16, 11 \rangle$	$C_2 \times D_8$	192	3	1
O3.32	(0; 2, 2, 2, 4)	$\langle 16, 13 \rangle$	$(C_4 \times C_2) : C_2$	48	1	1
O3.33	(0; 4, 4, 4)	$\langle 16, 2 \rangle$	$C_4 \times C_4$	96	1	1
O3.34	(0; 4, 4, 4)	$\langle 16, 4 \rangle$	$C_4 : C_4$	96	3	1
O3.35	(0; 2, 8, 8)	$\langle 16, 5 \rangle$	$C_8 \times C_2$	16	1	1
O3.36	(0; 2, 8, 8)	$\langle 16, 6 \rangle$	$C_8 : C_2$	16	1	1
O3.37	(0; 3, 3, 7)	$\langle 21, 1 \rangle$	$C_7 : C_3$	84	2	1
O3.38	(0; 2, 2, 2, 3)	$\langle 24, 12 \rangle$	S_4	216	9	1
O3.39	(0; 3, 4, 4)	$\langle 24, 12 \rangle$	S_4	24	1	1
O3.40	(0; 3, 3, 6)	$\langle 24, 3 \rangle$	$SL(2, 3)$	24	1	1
O3.41	(0; 2, 6, 6)	$\langle 24, 13 \rangle$	$C_2 \times A_4$	24	1	1
O3.42	(0; 2, 4, 12)	$\langle 24, 5 \rangle$	$C_4 \times S_3$	24	1	1
O3.43	(0; 2, 4, 8)	$\langle 32, 9 \rangle$	$(C_8 \times C_2) : C_2$	64	1	1
O3.44	(0; 2, 4, 8)	$\langle 32, 11 \rangle$	$(C_4 \times C_4) : C_2$	32	1	1
O3.45	(0; 3, 3, 4)	$\langle 48, 3 \rangle$	$(C_4 \times C_4) : C_3$	384	1	1
O3.46	(0; 2, 4, 6)	$\langle 48, 48 \rangle$	$C_2 \times S_4$	48	1	1
O3.47	(0; 2, 3, 12)	$\langle 48, 33 \rangle$	$SL(2, 3) : C_2$	48	1	1
O3.48	(0; 2, 3, 8)	$\langle 96, 64 \rangle$	$((C_4 \times C_4) : C_3) : C_2$	192	1	1
O3.49	(0; 2, 3, 7)	$\langle 168, 42 \rangle$	$PSL(3, 2)$	336	1	1

Table 7: Census of finite group actions on Riemann surfaces of genus $g = 4$

Ref	Signature	SMG id	Structure	epi	autg	top
O4.1	$(4; -)$	$\langle 1, 1 \rangle$	1	1	1	1
O4.2	$(2; 2, 2)$	$\langle 2, 1 \rangle$	C_2	16	16	1
O4.3	$(1; 2^6)$	$\langle 2, 1 \rangle$	C_2	4	4	1

Continued on next page

Table 7 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O4.4	$(0; 2^{10})$	$\langle 2, 1 \rangle$	C_2	1	1	1
O4.5	$(2; -)$	$\langle 3, 1 \rangle$	C_3	80	40	1
O4.6	$(1; 3, 3, 3)$	$\langle 3, 1 \rangle$	C_3	18	9	1
O4.7	$(0; 3^6)$	$\langle 3, 1 \rangle$	C_3	22	11	2
O4.8	$(1; 2, 2, 2)$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	96	16	1
O4.9	$(1; 4, 4)$	$\langle 4, 1 \rangle$	C_4	32	16	1
O4.10	$(0; 2^7)$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	546	91	2
O4.11	$(0; 2^4, 4^2)$	$\langle 4, 1 \rangle$	C_4	2	1	1
O4.12	$(0; 2, 4^4)$	$\langle 4, 1 \rangle$	C_4	8	4	1
O4.13	$(0; 5, 5, 5, 5)$	$\langle 5, 1 \rangle$	C_5	52	13	3
O4.14	$(1; 2, 2)$	$\langle 6, 1 \rangle$	S_3	96	16	1
O4.15	$(1; 2, 2)$	$\langle 6, 2 \rangle$	C_6	32	16	1
O4.16	$(0; 2^6)$	$\langle 6, 1 \rangle$	S_3	240	40	1
O4.17	$(0; 2^2, 3^3)$	$\langle 6, 1 \rangle$	S_3	24	4	1
O4.18	$(0; 2^2, 3^3)$	$\langle 6, 2 \rangle$	C_6	2	1	1
O4.19	$(0; 2^3, 3, 6)$	$\langle 6, 2 \rangle$	C_6	2	1	1
O4.20	$(0; 3, 3, 6, 6)$	$\langle 6, 2 \rangle$	C_6	6	3	2
O4.21	$(0; 2, 6, 6, 6)$	$\langle 6, 2 \rangle$	C_6	2	1	1
O4.22	$(0; 2^4, 4)$	$\langle 8, 3 \rangle$	D_8	176	22	2
O4.23	$(0; 2, 4, 4, 4)$	$\langle 8, 4 \rangle$	Q_8	24	1	1
O4.24	$(0; 2, 2, 8, 8)$	$\langle 8, 1 \rangle$	C_8	4	1	1
O4.25	$(0; 3, 3, 3, 3)$	$\langle 9, 2 \rangle$	$C_3 \times C_3$	432	9	2
O4.26	$(0; 9, 9, 9)$	$\langle 9, 1 \rangle$	C_9	18	3	1
O4.27	$(0; 2, 2, 5, 5)$	$\langle 10, 1 \rangle$	D_{10}	80	4	2
O4.28	$(0; 2, 2, 5, 5)$	$\langle 10, 2 \rangle$	C_{10}	4	1	1
O4.29	$(0; 5, 10, 10)$	$\langle 10, 2 \rangle$	C_{10}	12	3	2
O4.30	$(1; 2)$	$\langle 12, 3 \rangle$	A_4	96	4	1
O4.31	$(0; 2^5)$	$\langle 12, 4 \rangle$	D_{12}	960	80	1
O4.32	$(0; 2, 3, 3, 3)$	$\langle 12, 3 \rangle$	A_4	96	4	1
O4.33	$(0; 2, 2, 3, 6)$	$\langle 12, 4 \rangle$	D_{12}	24	2	1
O4.34	$(0; 2, 2, 3, 6)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	12	1	1
O4.35	$(0; 6, 6, 6)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	12	1	1
O4.36	$(0; 4, 6, 12)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O4.37	$(0; 3, 12, 12)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O4.38	$(0; 3, 5, 15)$	$\langle 15, 1 \rangle$	C_{15}	8	1	1
O4.39	$(0; 2, 2, 2, 8)$	$\langle 16, 7 \rangle$	D_{16}	96	3	1
O4.40	$(0; 4, 4, 8)$	$\langle 16, 9 \rangle$	Q_{16}	32	1	1
O4.41	$(0; 2, 16, 16)$	$\langle 16, 1 \rangle$	C_{16}	8	1	1
O4.42	$(0; 2, 2, 3, 3)$	$\langle 18, 3 \rangle$	$C_3 \times S_3$	48	4	2
O4.43	$(0; 2, 2, 3, 3)$	$\langle 18, 4 \rangle$	$(C_3 \times C_3) : C_2$	432	1	1
O4.44	$(0; 3, 6, 6)$	$\langle 18, 3 \rangle$	$C_3 \times S_3$	24	2	2
O4.45	$(0; 3, 6, 6)$	$\langle 18, 5 \rangle$	$C_6 \times C_3$	48	1	1
O4.46	$(0; 2, 9, 18)$	$\langle 18, 2 \rangle$	C_{18}	6	1	1
O4.47	$(0; 2, 2, 2, 5)$	$\langle 20, 4 \rangle$	D_{20}	120	3	1
O4.48	$(0; 4, 4, 5)$	$\langle 20, 1 \rangle$	$C_5 : C_4$	40	1	1
O4.49	$(0; 4, 4, 5)$	$\langle 20, 3 \rangle$	$C_5 : C_4$	40	2	1

Continued on next page

Table 7 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O4.50	(0; 2, 10, 10)	$\langle 20, 5 \rangle$	$C_{10} \times C_2$	24	1	1
O4.51	(0; 2, 2, 2, 4)	$\langle 24, 12 \rangle$	S_4	96	4	1
O4.52	(0; 3, 4, 6)	$\langle 24, 3 \rangle$	$SL(2, 3)$	24	1	1
O4.53	(0; 2, 6, 12)	$\langle 24, 10 \rangle$	$C_3 \times D_8$	16	1	1
O4.54	(0; 2, 4, 16)	$\langle 32, 19 \rangle$	QD_{32}	64	1	1
O4.55	(0; 2, 2, 2, 3)	$\langle 36, 10 \rangle$	$S_3 \times S_3$	216	3	1
O4.56	(0; 3, 4, 4)	$\langle 36, 9 \rangle$	$(C_3 \times C_3) : C_4$	144	1	1
O4.57	(0; 3, 3, 6)	$\langle 36, 11 \rangle$	$C_3 \times A_4$	144	1	1
O4.58	(0; 2, 6, 6)	$\langle 36, 10 \rangle$	$S_3 \times S_3$	72	1	1
O4.59	(0; 2, 6, 6)	$\langle 36, 12 \rangle$	$C_6 \times S_3$	48	2	1
O4.60	(0; 2, 4, 10)	$\langle 40, 8 \rangle$	$(C_{10} \times C_2) : C_2$	80	1	1
O4.61	(0; 2, 5, 5)	$\langle 60, 5 \rangle$	A_5	120	1	1
O4.62	(0; 2, 4, 6)	$\langle 72, 40 \rangle$	$(S_3 \times S_3) : C_2$	144	1	1
O4.63	(0; 2, 3, 12)	$\langle 72, 42 \rangle$	$C_3 \times S_4$	48	1	1
O4.64	(0; 2, 4, 5)	$\langle 120, 34 \rangle$	S_5	120	1	1

Table 8: Census of finite group actions on Riemann surfaces of genus $g = 5$

Ref	Signature	SMG id	Structure	epi	autg	top
O5.1	(5; -)	$\langle 1, 1 \rangle$	1	1	1	1
O5.2	(3; -)	$\langle 2, 1 \rangle$	C_2	63	63	1
O5.3	(2; 2, 2, 2, 2)	$\langle 2, 1 \rangle$	C_2	16	16	1
O5.4	(1; 2^8)	$\langle 2, 1 \rangle$	C_2	4	4	1
O5.5	(0; 2^{12})	$\langle 2, 1 \rangle$	C_2	1	1	1
O5.6	(1; 3, 3, 3, 3)	$\langle 3, 1 \rangle$	C_3	54	27	1
O5.7	(0; 3^7)	$\langle 3, 1 \rangle$	C_3	42	21	1
O5.8	(2; -)	$\langle 4, 1 \rangle$	C_4	240	120	1
O5.9	(2; -)	$\langle 4, 2 \rangle$	$C_2 \times C_2$	210	35	2
O5.10	(1; 2, 2, 2, 2)	$\langle 4, 1 \rangle$	C_4	12	6	1
O5.11	(1; 2, 2, 2, 2)	$\langle 4, 2 \rangle$	$C_2 \times C_2$	324	54	2
O5.12	(1; 2, 4, 4)	$\langle 4, 1 \rangle$	C_4	32	16	1
O5.13	(0; 2^8)	$\langle 4, 2 \rangle$	$C_2 \times C_2$	1638	273	3
O5.14	(0; $2^5, 4^2$)	$\langle 4, 1 \rangle$	C_4	2	1	1
O5.15	(0; $2^2, 4^4$)	$\langle 4, 1 \rangle$	C_4	8	4	2
O5.16	(1; 5, 5)	$\langle 5, 1 \rangle$	C_5	100	25	1
O5.17	(1; 3, 3)	$\langle 6, 1 \rangle$	S_3	36	6	1
O5.18	(1; 3, 3)	$\langle 6, 2 \rangle$	C_6	54	27	1
O5.19	(0; $2^4, 3^2$)	$\langle 6, 1 \rangle$	S_3	108	18	1
O5.20	(0; $2^4, 3^2$)	$\langle 6, 2 \rangle$	C_6	2	1	1
O5.21	(0; 2, $3^3, 6$)	$\langle 6, 2 \rangle$	C_6	6	3	1
O5.22	(0; $2^2, 3, 6^2$)	$\langle 6, 2 \rangle$	C_6	2	1	1
O5.23	(0; 6, 6, 6, 6)	$\langle 6, 2 \rangle$	C_6	6	3	1
O5.24	(1; 2, 2)	$\langle 8, 1 \rangle$	C_8	48	12	1

Continued on next page

Table 8 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O5.25	(1; 2, 2)	$\langle 8, 2 \rangle$	$C_4 \times C_2$	120	15	2
O5.26	(1; 2, 2)	$\langle 8, 3 \rangle$	D_8	192	24	1
O5.27	(1; 2, 2)	$\langle 8, 5 \rangle$	$C_2 \times C_2 \times C_2$	168	1	1
O5.28	(0; 2 ⁶)	$\langle 8, 3 \rangle$	D_8	1680	210	2
O5.29	(0; 2 ⁶)	$\langle 8, 5 \rangle$	$C_2 \times C_2 \times C_2$	13440	80	3
O5.30	(0; 2 ³ , 4 ²)	$\langle 8, 2 \rangle$	$C_4 \times C_2$	104	13	5
O5.31	(0; 2 ³ , 4 ²)	$\langle 8, 3 \rangle$	D_8	48	6	1
O5.32	(0; 4, 4, 4, 4)	$\langle 8, 2 \rangle$	$C_4 \times C_2$	48	6	2
O5.33	(0; 4, 4, 4, 4)	$\langle 8, 4 \rangle$	Q_8	144	6	1
O5.34	(0; 2, 4, 8, 8)	$\langle 8, 1 \rangle$	C_8	8	2	2
O5.35	(1; 5)	$\langle 10, 1 \rangle$	D_{10}	60	3	1
O5.36	(0; 2 ⁴ , 5)	$\langle 10, 1 \rangle$	D_{10}	500	25	1
O5.37	(0; 2, 2, 10, 10)	$\langle 10, 2 \rangle$	C_{10}	4	1	1
O5.38	(0; 11, 11, 11)	$\langle 11, 1 \rangle$	C_{11}	90	9	2
O5.39	(1; 3)	$\langle 12, 1 \rangle$	$C_3 : C_4$	72	6	1
O5.40	(1; 3)	$\langle 12, 4 \rangle$	D_{12}	36	3	1
O5.41	(0; 2 ⁴ , 3)	$\langle 12, 4 \rangle$	D_{12}	396	33	2
O5.42	(0; 3, 3, 3, 3)	$\langle 12, 3 \rangle$	A_4	360	15	2
O5.43	(0; 2, 3, 4, 4)	$\langle 12, 1 \rangle$	$C_3 : C_4$	12	1	1
O5.44	(0; 2, 2, 6, 6)	$\langle 12, 4 \rangle$	D_{12}	24	2	1
O5.45	(0; 2, 2, 6, 6)	$\langle 12, 5 \rangle$	$C_6 \times C_2$	36	3	2
O5.46	(0; 6, 12, 12)	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O5.47	(0; 3, 15, 15)	$\langle 15, 1 \rangle$	C_{15}	8	1	1
O5.48	(1; 2)	$\langle 16, 3 \rangle$	$(C_4 \times C_2) : C_2$	96	3	1
O5.49	(1; 2)	$\langle 16, 4 \rangle$	$C_4 : C_4$	96	3	1
O5.50	(1; 2)	$\langle 16, 6 \rangle$	$C_8 : C_2$	96	6	1
O5.51	(0; 2 ⁵)	$\langle 16, 7 \rangle$	D_{16}	1920	60	1
O5.52	(0; 2 ⁵)	$\langle 16, 11 \rangle$	$C_2 \times D_8$	4800	75	2
O5.53	(0; 2 ⁵)	$\langle 16, 14 \rangle$	$C_2 \times C_2 \times C_2 \times C_2$	20160	1	1
O5.54	(0; 2, 2, 4, 4)	$\langle 16, 3 \rangle$	$(C_4 \times C_2) : C_2$	320	10	4
O5.55	(0; 2, 2, 4, 4)	$\langle 16, 8 \rangle$	QD_{16}	64	4	1
O5.56	(0; 2, 2, 4, 4)	$\langle 16, 10 \rangle$	$C_4 \times C_2 \times C_2$	192	1	1
O5.57	(0; 2, 2, 4, 4)	$\langle 16, 11 \rangle$	$C_2 \times D_8$	64	1	1
O5.58	(0; 2, 2, 4, 4)	$\langle 16, 13 \rangle$	$(C_4 \times C_2) : C_2$	96	2	1
O5.59	(0; 4, 8, 8)	$\langle 16, 5 \rangle$	$C_8 \times C_2$	16	1	1
O5.60	(0; 4, 8, 8)	$\langle 16, 6 \rangle$	$C_8 : C_2$	16	1	1
O5.61	(0; 2, 2, 2, 10)	$\langle 20, 4 \rangle$	D_{20}	120	3	1
O5.62	(0; 4, 4, 10)	$\langle 20, 1 \rangle$	$C_5 : C_4$	40	1	1
O5.63	(0; 2, 20, 20)	$\langle 20, 2 \rangle$	C_{20}	8	1	1
O5.64	(0; 2, 11, 22)	$\langle 22, 2 \rangle$	C_{22}	10	1	1
O5.65	(0; 2, 2, 3, 3)	$\langle 24, 12 \rangle$	S_4	144	6	1
O5.66	(0; 2, 2, 3, 3)	$\langle 24, 13 \rangle$	$C_2 \times A_4$	120	5	2
O5.67	(0; 2, 2, 2, 6)	$\langle 24, 8 \rangle$	$(C_6 \times C_2) : C_2$	144	6	1
O5.68	(0; 2, 2, 2, 6)	$\langle 24, 14 \rangle$	$C_2 \times C_2 \times S_3$	432	3	1
O5.69	(0; 4, 4, 6)	$\langle 24, 7 \rangle$	$C_2 \times (C_3 : C_4)$	48	1	1
O5.70	(0; 3, 6, 6)	$\langle 24, 13 \rangle$	$C_2 \times A_4$	24	1	1

Continued on next page

Table 8 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O5.71	$(0; 2, 12, 12)$	$\langle 24, 9 \rangle$	$C_{12} \times C_2$	16	1	1
O5.72	$(0; 2, 6, 15)$	$\langle 30, 2 \rangle$	$C_3 \times D_{10}$	40	1	1
O5.73	$(0; 2, 2, 2, 4)$	$\langle 32, 27 \rangle$	$C_2^4: C_2$	1152	3	1
O5.74	$(0; 2, 2, 2, 4)$	$\langle 32, 28 \rangle$	$(C_4 \times C_2 \times C_2): C_2$	384	3	1
O5.75	$(0; 2, 2, 2, 4)$	$\langle 32, 43 \rangle$	$(C_2 \times D_8): C_2$	384	6	1
O5.76	$(0; 4, 4, 4)$	$\langle 32, 2 \rangle$	$(C_4 \times C_2): C_4$	384	1	1
O5.77	$(0; 4, 4, 4)$	$\langle 32, 6 \rangle$	$((C_4 \times C_2): C_2): C_2$	192	3	1
O5.78	$(0; 2, 8, 8)$	$\langle 32, 5 \rangle$	$(C_8 \times C_2): C_2$	64	1	1
O5.79	$(0; 2, 8, 8)$	$\langle 32, 7 \rangle$	$(C_8: C_2): C_2$	128	1	1
O5.80	$(0; 2, 4, 20)$	$\langle 40, 5 \rangle$	$C_4 \times D_{10}$	80	1	1
O5.81	$(0; 2, 2, 2, 3)$	$\langle 48, 48 \rangle$	$C_2 \times S_4$	432	9	1
O5.82	$(0; 3, 4, 4)$	$\langle 48, 30 \rangle$	$A_4: C_4$	144	3	2
O5.83	$(0; 2, 6, 6)$	$\langle 48, 49 \rangle$	$C_2 \times C_2 \times A_4$	144	1	1
O5.84	$(0; 2, 4, 12)$	$\langle 48, 14 \rangle$	$(C_{12} \times C_2): C_2$	96	1	1
O5.85	$(0; 3, 3, 5)$	$\langle 60, 5 \rangle$	A_5	120	1	1
O5.86	$(0; 2, 4, 8)$	$\langle 64, 8 \rangle$	$((C_8 \times C_2): C_2): C_2$	128	1	1
O5.87	$(0; 2, 4, 8)$	$\langle 64, 32 \rangle$	$((C_8: C_2): C_2): C_2$	128	1	1
O5.88	$(0; 2, 5, 5)$	$\langle 80, 49 \rangle$	$C_2^4: C_5$	960	1	1
O5.89	$(0; 3, 3, 4)$	$\langle 96, 3 \rangle$	$((C_4 \times C_2): C_4): C_3$	384	1	1
O5.90	$(0; 2, 4, 6)$	$\langle 96, 195 \rangle$	$(C_2 \times C_2 \times A_4): C_2$	96	1	1
O5.91	$(0; 2, 3, 10)$	$\langle 120, 35 \rangle$	$C_2 \times A_5$	120	1	1
O5.92	$(0; 2, 4, 5)$	$\langle 160, 234 \rangle$	$(C_2^4: C_5): C_2$	960	1	1
O5.93	$(0; 2, 3, 8)$	$\langle 192, 181 \rangle$	$((C_4 \times C_2): C_4): C_3): C_2$	384	1	1

Table 9: Census of finite group actions on Riemann surfaces of genus $g = 6$

Ref	Signature	SMG id	Structure	epi	autg	top
O6.1	$(6; -)$	$\langle 1, 1 \rangle$	1	1	1	1
O6.2	$(3; 2, 2)$	$\langle 2, 1 \rangle$	C_2	64	64	1
O6.3	$(2; 2^6)$	$\langle 2, 1 \rangle$	C_2	16	16	1
O6.4	$(1; 2^{10})$	$\langle 2, 1 \rangle$	C_2	4	4	1
O6.5	$(0; 2^{14})$	$\langle 2, 1 \rangle$	C_2	1	1	1
O6.6	$(2; 3, 3)$	$\langle 3, 1 \rangle$	C_3	162	81	1
O6.7	$(1; 3^5)$	$\langle 3, 1 \rangle$	C_3	90	45	1
O6.8	$(0; 3^8)$	$\langle 3, 1 \rangle$	C_3	86	43	2
O6.9	$(1; 2^5)$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	960	160	1
O6.10	$(1; 2, 2, 4, 4)$	$\langle 4, 1 \rangle$	C_4	32	16	1
O6.11	$(0; 2^9)$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	4920	820	3
O6.12	$(0; 2^6, 4^2)$	$\langle 4, 1 \rangle$	C_4	2	1	1
O6.13	$(0; 2^3, 4^4)$	$\langle 4, 1 \rangle$	C_4	8	4	1
O6.14	$(0; 4^6)$	$\langle 4, 1 \rangle$	C_4	32	16	2
O6.15	$(2; -)$	$\langle 5, 1 \rangle$	C_5	624	156	1
O6.16	$(0; 5^5)$	$\langle 5, 1 \rangle$	C_5	204	51	3

Continued on next page

Table 9 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O6.17	$(1; 2, 2, 3)$	$\langle 6, 1 \rangle$	S_3	216	36	1
O6.18	$(1; 6, 6)$	$\langle 6, 2 \rangle$	C_6	72	36	1
O6.19	$(0; 2^6, 3)$	$\langle 6, 1 \rangle$	S_3	486	81	1
O6.20	$(0; 2^2, 3^4)$	$\langle 6, 1 \rangle$	S_3	48	8	1
O6.21	$(0; 2^2, 3^4)$	$\langle 6, 2 \rangle$	C_6	6	3	1
O6.22	$(0; 2^3, 3^2, 6)$	$\langle 6, 2 \rangle$	C_6	2	1	1
O6.23	$(0; 2^4, 6^2)$	$\langle 6, 2 \rangle$	C_6	2	1	1
O6.24	$(0; 3^3, 6^2)$	$\langle 6, 2 \rangle$	C_6	10	5	2
O6.25	$(0; 2, 3, 6^3)$	$\langle 6, 2 \rangle$	C_6	6	3	1
O6.26	$(0; 7, 7, 7, 7)$	$\langle 7, 1 \rangle$	C_7	186	31	4
O6.27	$(0; 2^5, 4)$	$\langle 8, 3 \rangle$	D_8	720	90	2
O6.28	$(0; 2^2, 4^3)$	$\langle 8, 3 \rangle$	D_8	32	4	1
O6.29	$(0; 2^2, 4^3)$	$\langle 8, 4 \rangle$	Q_8	24	1	1
O6.30	$(0; 2^3, 8^2)$	$\langle 8, 1 \rangle$	C_8	4	1	1
O6.31	$(0; 4, 4, 8, 8)$	$\langle 8, 1 \rangle$	C_8	16	4	2
O6.32	$(0; 3, 3, 9, 9)$	$\langle 9, 1 \rangle$	C_9	24	4	2
O6.33	$(1; 2, 2)$	$\langle 10, 1 \rangle$	D_{10}	480	24	1
O6.34	$(1; 2, 2)$	$\langle 10, 2 \rangle$	C_{10}	96	24	1
O6.35	$(0; 2^6)$	$\langle 10, 1 \rangle$	D_{10}	3120	156	1
O6.36	$(0; 2, 5, 5, 10)$	$\langle 10, 2 \rangle$	C_{10}	12	3	2
O6.37	$(0; 2^3, 3^2)$	$\langle 12, 3 \rangle$	A_4	216	9	1
O6.38	$(0; 2^3, 3^2)$	$\langle 12, 4 \rangle$	D_{12}	72	6	1
O6.39	$(0; 2^3, 3^2)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	12	1	1
O6.40	$(0; 3, 3, 4, 4)$	$\langle 12, 1 \rangle$	$C_3 : C_4$	24	2	1
O6.41	$(0; 3, 3, 4, 4)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O6.42	$(0; 2^4, 6)$	$\langle 12, 4 \rangle$	D_{12}	504	42	2
O6.43	$(0; 2, 4, 4, 6)$	$\langle 12, 1 \rangle$	$C_3 : C_4$	12	1	1
O6.44	$(0; 2, 3, 6, 6)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	12	1	1
O6.45	$(0; 2, 3, 4, 12)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O6.46	$(0; 2, 2, 12, 12)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O6.47	$(0; 13, 13, 13)$	$\langle 13, 1 \rangle$	C_{13}	132	11	3
O6.48	$(0; 2, 2, 7, 7)$	$\langle 14, 1 \rangle$	D_{14}	252	6	2
O6.49	$(0; 2, 2, 7, 7)$	$\langle 14, 2 \rangle$	C_{14}	6	1	1
O6.50	$(0; 7, 14, 14)$	$\langle 14, 2 \rangle$	C_{14}	30	5	3
O6.51	$(0; 5, 15, 15)$	$\langle 15, 1 \rangle$	C_{15}	24	3	2
O6.52	$(0; 2, 2, 4, 8)$	$\langle 16, 7 \rangle$	D_{16}	64	2	1
O6.53	$(0; 2, 2, 4, 8)$	$\langle 16, 8 \rangle$	QD_{16}	32	2	1
O6.54	$(0; 4, 16, 16)$	$\langle 16, 1 \rangle$	C_{16}	16	2	1
O6.55	$(0; 2, 2, 3, 9)$	$\langle 18, 1 \rangle$	D_{18}	108	2	1
O6.56	$(0; 3, 18, 18)$	$\langle 18, 2 \rangle$	C_{18}	12	2	1
O6.57	$(0; 2^5)$	$\langle 20, 4 \rangle$	D_{20}	4800	120	1
O6.58	$(0; 2, 2, 4, 4)$	$\langle 20, 3 \rangle$	$C_5 : C_4$	240	12	1
O6.59	$(0; 4, 5, 20)$	$\langle 20, 2 \rangle$	C_{20}	8	1	1
O6.60	$(0; 3, 7, 21)$	$\langle 21, 2 \rangle$	C_{21}	12	1	1
O6.61	$(0; 2, 2, 3, 4)$	$\langle 24, 6 \rangle$	D_{24}	48	1	1
O6.62	$(0; 2, 2, 3, 4)$	$\langle 24, 8 \rangle$	$(C_6 \times C_2) : C_2$	48	2	1

Continued on next page

Table 9 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O6.63	(0; 2, 2, 3, 4)	$\langle 24, 12 \rangle$	S_4	144	6	1
O6.64	(0; 4, 6, 6)	$\langle 24, 3 \rangle$	$SL(2, 3)$	24	1	1
O6.65	(0; 4, 6, 6)	$\langle 24, 10 \rangle$	$C_3 \times D_8$	16	1	1
O6.66	(0; 3, 8, 8)	$\langle 24, 1 \rangle$	$C_3 : C_8$	24	1	1
O6.67	(0; 2, 2, 2, 12)	$\langle 24, 6 \rangle$	D_{24}	144	3	1
O6.68	(0; 4, 4, 12)	$\langle 24, 4 \rangle$	$C_3 : Q_8$	48	1	1
O6.69	(0; 2, 24, 24)	$\langle 24, 2 \rangle$	C_{24}	8	1	1
O6.70	(0; 5, 5, 5)	$\langle 25, 2 \rangle$	$C_5 \times C_5$	480	1	1
O6.71	(0; 2, 13, 26)	$\langle 26, 2 \rangle$	C_{26}	12	1	1
O6.72	(0; 2, 2, 2, 7)	$\langle 28, 3 \rangle$	D_{28}	252	3	1
O6.73	(0; 4, 4, 7)	$\langle 28, 1 \rangle$	$C_7 : C_4$	84	1	1
O6.74	(0; 2, 14, 14)	$\langle 28, 4 \rangle$	$C_{14} \times C_2$	36	1	1
O6.75	(0; 2, 10, 15)	$\langle 30, 1 \rangle$	$C_5 \times S_3$	24	1	1
O6.76	(0; 2, 9, 9)	$\langle 36, 3 \rangle$	$(C_2 \times C_2) : C_9$	72	1	1
O6.77	(0; 3, 3, 13)	$\langle 39, 1 \rangle$	$C_{13} : C_3$	312	2	1
O6.78	(0; 2, 6, 8)	$\langle 48, 15 \rangle$	$(C_3 \times D_8) : C_2$	96	1	1
O6.79	(0; 2, 6, 8)	$\langle 48, 29 \rangle$	$GL(2, 3)$	48	1	1
O6.80	(0; 2, 4, 24)	$\langle 48, 6 \rangle$	$C_{24} : C_2$	96	1	1
O6.81	(0; 2, 5, 10)	$\langle 50, 3 \rangle$	$C_5 \times D_{10}$	80	1	1
O6.82	(0; 2, 4, 14)	$\langle 56, 7 \rangle$	$(C_{14} \times C_2) : C_2$	168	1	1
O6.83	(0; 2, 2, 2, 3)	$\langle 60, 5 \rangle$	A_5	1080	9	1
O6.84	(0; 2, 4, 9)	$\langle 72, 15 \rangle$	$((C_2 \times C_2) : C_9) : C_2$	216	1	1
O6.85	(0; 3, 3, 5)	$\langle 75, 2 \rangle$	$(C_5 \times C_5) : C_3$	1200	1	1
O6.86	(0; 2, 4, 6)	$\langle 120, 34 \rangle$	S_5	120	1	1
O6.87	(0; 2, 3, 10)	$\langle 150, 5 \rangle$	$((C_5 \times C_5) : C_3) : C_2$	600	1	1

Table 10: Census of finite group actions on Riemann surfaces of genus $g = 7$

Ref	Signature	SMG id	Structure	epi	autg	top
O7.1	(7; −)	$\langle 1, 1 \rangle$	1	1	1	1
O7.2	(4; −)	$\langle 2, 1 \rangle$	C_2	255	255	1
O7.3	(3; 2, 2, 2, 2)	$\langle 2, 1 \rangle$	C_2	64	64	1
O7.4	(2; 2^8)	$\langle 2, 1 \rangle$	C_2	16	16	1
O7.5	(1; 2^{12})	$\langle 2, 1 \rangle$	C_2	4	4	1
O7.6	(0; 2^{16})	$\langle 2, 1 \rangle$	C_2	1	1	1
O7.7	(3; −)	$\langle 3, 1 \rangle$	C_3	728	364	1
O7.8	(2; 3, 3, 3)	$\langle 3, 1 \rangle$	C_3	162	81	1
O7.9	(1; 3^6)	$\langle 3, 1 \rangle$	C_3	198	99	2
O7.10	(0; 3^9)	$\langle 3, 1 \rangle$	C_3	170	85	2
O7.11	(2; 2, 2)	$\langle 4, 1 \rangle$	C_4	240	120	1
O7.12	(2; 2, 2)	$\langle 4, 2 \rangle$	$C_2 \times C_2$	720	120	1
O7.13	(1; 2^6)	$\langle 4, 1 \rangle$	C_4	12	6	1
O7.14	(1; 2^6)	$\langle 4, 2 \rangle$	$C_2 \times C_2$	2916	486	3

Continued on next page

Table 10 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O7.15	$(1; 2^3, 4^2)$	$\langle 4, 1 \rangle$	C_4	32	16	1
O7.16	$(1; 4, 4, 4, 4)$	$\langle 4, 1 \rangle$	C_4	128	64	2
O7.17	$(0; 2^{10})$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	14760	2460	4
O7.18	$(0; 2^7, 4^2)$	$\langle 4, 1 \rangle$	C_4	2	1	1
O7.19	$(0; 2^4, 4^4)$	$\langle 4, 1 \rangle$	C_4	8	4	2
O7.20	$(0; 2, 4^6)$	$\langle 4, 1 \rangle$	C_4	32	16	2
O7.21	$(1; 5, 5, 5)$	$\langle 5, 1 \rangle$	C_5	300	75	1
O7.22	$(2; -)$	$\langle 6, 1 \rangle$	S_3	360	60	1
O7.23	$(2; -)$	$\langle 6, 2 \rangle$	C_6	1200	600	1
O7.24	$(1; 2, 2, 2, 2)$	$\langle 6, 1 \rangle$	S_3	960	160	1
O7.25	$(1; 2, 2, 2, 2)$	$\langle 6, 2 \rangle$	C_6	32	16	1
O7.26	$(1; 3, 3, 3)$	$\langle 6, 1 \rangle$	S_3	72	12	1
O7.27	$(1; 3, 3, 3)$	$\langle 6, 2 \rangle$	C_6	54	27	1
O7.28	$(1; 2, 3, 6)$	$\langle 6, 2 \rangle$	C_6	72	36	1
O7.29	$(0; 2^8)$	$\langle 6, 1 \rangle$	S_3	2184	364	1
O7.30	$(0; 2^4, 3^3)$	$\langle 6, 1 \rangle$	S_3	216	36	1
O7.31	$(0; 2^4, 3^3)$	$\langle 6, 2 \rangle$	C_6	2	1	1
O7.32	$(0; 2^5, 3, 6)$	$\langle 6, 2 \rangle$	C_6	2	1	1
O7.33	$(0; 2, 3^4, 6)$	$\langle 6, 2 \rangle$	C_6	10	5	2
O7.34	$(0; 2^2, 3^2, 6^2)$	$\langle 6, 2 \rangle$	C_6	6	3	2
O7.35	$(0; 2^3, 6^3)$	$\langle 6, 2 \rangle$	C_6	2	1	1
O7.36	$(0; 3, 6^4)$	$\langle 6, 2 \rangle$	C_6	10	5	2
O7.37	$(1; 7, 7)$	$\langle 7, 1 \rangle$	C_7	294	49	1
O7.38	$(1; 2, 2, 2)$	$\langle 8, 2 \rangle$	$C_4 \times C_2$	288	36	1
O7.39	$(1; 2, 2, 2)$	$\langle 8, 3 \rangle$	D_8	600	75	2
O7.40	$(1; 2, 2, 2)$	$\langle 8, 4 \rangle$	Q_8	24	1	1
O7.41	$(1; 2, 2, 2)$	$\langle 8, 5 \rangle$	$C_2 \times C_2 \times C_2$	2016	12	1
O7.42	$(1; 4, 4)$	$\langle 8, 1 \rangle$	C_8	96	24	1
O7.43	$(1; 4, 4)$	$\langle 8, 2 \rangle$	$C_4 \times C_2$	192	24	1
O7.44	$(1; 4, 4)$	$\langle 8, 3 \rangle$	D_8	96	12	1
O7.45	$(1; 4, 4)$	$\langle 8, 4 \rangle$	Q_8	288	12	1
O7.46	$(0; 2^7)$	$\langle 8, 3 \rangle$	D_8	8400	1050	2
O7.47	$(0; 2^7)$	$\langle 8, 5 \rangle$	$C_2 \times C_2 \times C_2$	99120	590	4
O7.48	$(0; 2^4, 4^2)$	$\langle 8, 2 \rangle$	$C_4 \times C_2$	320	40	8
O7.49	$(0; 2^4, 4^2)$	$\langle 8, 3 \rangle$	D_8	352	44	3
O7.50	$(0; 2, 4^4)$	$\langle 8, 2 \rangle$	$C_4 \times C_2$	176	22	3
O7.51	$(0; 2, 4^4)$	$\langle 8, 4 \rangle$	Q_8	144	6	1
O7.52	$(0; 2^2, 4, 8^2)$	$\langle 8, 1 \rangle$	C_8	8	2	2
O7.53	$(0; 8, 8, 8, 8)$	$\langle 8, 1 \rangle$	C_8	64	16	4
O7.54	$(1; 3, 3)$	$\langle 9, 1 \rangle$	C_9	144	24	1
O7.55	$(1; 3, 3)$	$\langle 9, 2 \rangle$	$C_3 \times C_3$	576	12	1
O7.56	$(0; 3^5)$	$\langle 9, 2 \rangle$	$C_3 \times C_3$	3600	75	4
O7.57	$(0; 3, 9, 9, 9)$	$\langle 9, 1 \rangle$	C_9	36	6	3
O7.58	$(0; 2^3, 5, 10)$	$\langle 10, 2 \rangle$	C_{10}	4	1	1
O7.59	$(0; 2, 10, 10, 10)$	$\langle 10, 2 \rangle$	C_{10}	12	3	1
O7.60	$(1; 2, 2)$	$\langle 12, 2 \rangle$	C_{12}	96	24	1

Continued on next page

Table 10 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O7.61	$(1; 2, 2)$	$\langle 12, 3 \rangle$	A_4	288	12	1
O7.62	$(1; 2, 2)$	$\langle 12, 4 \rangle$	D_{12}	576	48	1
O7.63	$(1; 2, 2)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	288	24	1
O7.64	$(0; 2^6)$	$\langle 12, 4 \rangle$	D_{12}	10080	840	3
O7.65	$(0; 2^2, 3^3)$	$\langle 12, 3 \rangle$	A_4	288	12	1
O7.66	$(0; 4, 4, 4, 4)$	$\langle 12, 1 \rangle$	$C_3 : C_4$	192	16	2
O7.67	$(0; 2^3, 3, 6)$	$\langle 12, 4 \rangle$	D_{12}	72	6	1
O7.68	$(0; 2^3, 3, 6)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	36	3	1
O7.69	$(0; 3, 4, 4, 6)$	$\langle 12, 1 \rangle$	$C_3 : C_4$	24	2	1
O7.70	$(0; 3, 4, 4, 6)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O7.71	$(0; 2, 6, 6, 6)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	36	3	1
O7.72	$(0; 3, 3, 4, 12)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O7.73	$(0; 2, 4, 6, 12)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O7.74	$(0; 2, 3, 12, 12)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O7.75	$(1; 7)$	$\langle 14, 1 \rangle$	D_{14}	126	3	1
O7.76	$(0; 2^4, 7)$	$\langle 14, 1 \rangle$	D_{14}	2058	49	1
O7.77	$(0; 2, 2, 14, 14)$	$\langle 14, 2 \rangle$	C_{14}	6	1	1
O7.78	$(0; 15, 15, 15)$	$\langle 15, 1 \rangle$	C_{15}	24	3	1
O7.79	$(1; 4)$	$\langle 16, 7 \rangle$	D_{16}	96	3	1
O7.80	$(1; 4)$	$\langle 16, 8 \rangle$	QD_{16}	96	6	1
O7.81	$(1; 4)$	$\langle 16, 9 \rangle$	Q_{16}	96	3	1
O7.82	$(0; 2^4, 4)$	$\langle 16, 7 \rangle$	D_{16}	768	24	1
O7.83	$(0; 2^4, 4)$	$\langle 16, 11 \rangle$	$C_2 \times D_8$	3072	48	4
O7.84	$(0; 2^4, 4)$	$\langle 16, 13 \rangle$	$(C_4 \times C_2) : C_2$	768	16	2
O7.85	$(0; 2, 4, 4, 4)$	$\langle 16, 2 \rangle$	$C_4 \times C_4$	288	3	1
O7.86	$(0; 2, 4, 4, 4)$	$\langle 16, 4 \rangle$	$C_4 : C_4$	288	9	3
O7.87	$(0; 2, 4, 4, 4)$	$\langle 16, 12 \rangle$	$C_2 \times Q_8$	192	1	1
O7.88	$(0; 2, 4, 4, 4)$	$\langle 16, 13 \rangle$	$(C_4 \times C_2) : C_2$	144	3	1
O7.89	$(0; 2, 2, 8, 8)$	$\langle 16, 5 \rangle$	$C_8 \times C_2$	64	4	3
O7.90	$(0; 2, 2, 8, 8)$	$\langle 16, 6 \rangle$	$C_8 : C_2$	64	4	2
O7.91	$(0; 2, 2, 8, 8)$	$\langle 16, 7 \rangle$	D_{16}	128	4	2
O7.92	$(0; 2, 2, 8, 8)$	$\langle 16, 8 \rangle$	QD_{16}	64	4	2
O7.93	$(0; 8, 16, 16)$	$\langle 16, 1 \rangle$	C_{16}	32	4	3
O7.94	$(1; 3)$	$\langle 18, 3 \rangle$	$C_3 \times S_3$	144	12	1
O7.95	$(0; 2^4, 3)$	$\langle 18, 1 \rangle$	D_{18}	1296	24	1
O7.96	$(0; 2^4, 3)$	$\langle 18, 4 \rangle$	$(C_3 \times C_3) : C_2$	5184	12	1
O7.97	$(0; 2, 3, 3, 6)$	$\langle 18, 3 \rangle$	$C_3 \times S_3$	120	10	4
O7.98	$(0; 2, 3, 3, 6)$	$\langle 18, 5 \rangle$	$C_6 \times C_3$	48	1	1
O7.99	$(0; 2, 2, 6, 6)$	$\langle 18, 3 \rangle$	$C_3 \times S_3$	48	4	1
O7.100	$(0; 6, 9, 18)$	$\langle 18, 2 \rangle$	C_{18}	12	2	2
O7.101	$(0; 4, 10, 20)$	$\langle 20, 2 \rangle$	C_{20}	8	1	1
O7.102	$(0; 3, 21, 21)$	$\langle 21, 2 \rangle$	C_{21}	12	1	1
O7.103	$(1; 2)$	$\langle 24, 10 \rangle$	$C_3 \times D_8$	192	12	1
O7.104	$(1; 2)$	$\langle 24, 11 \rangle$	$C_3 \times Q_8$	192	4	1
O7.105	$(1; 2)$	$\langle 24, 13 \rangle$	$C_2 \times A_4$	288	12	1
O7.106	$(0; 2^5)$	$\langle 24, 6 \rangle$	D_{24}	5760	120	1

Continued on next page

Table 10 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O7.107	$(0; 2^5)$	$\langle 24, 12 \rangle$	S_4	1440	60	1
O7.108	$(0; 2^5)$	$\langle 24, 14 \rangle$	$C_2 \times C_2 \times S_3$	17280	120	1
O7.109	$(0; 2, 3, 3, 3)$	$\langle 24, 3 \rangle$	$SL(2, 3)$	24	1	1
O7.110	$(0; 2, 2, 4, 4)$	$\langle 24, 5 \rangle$	$C_4 \times S_3$	192	8	2
O7.111	$(0; 2, 2, 4, 4)$	$\langle 24, 8 \rangle$	$(C_6 \times C_2): C_2$	192	8	1
O7.112	$(0; 2, 2, 4, 4)$	$\langle 24, 12 \rangle$	S_4	96	4	1
O7.113	$(0; 2, 2, 3, 6)$	$\langle 24, 13 \rangle$	$C_2 \times A_4$	192	8	2
O7.114	$(0; 6, 6, 6)$	$\langle 24, 3 \rangle$	$SL(2, 3)$	24	1	1
O7.115	$(0; 4, 6, 12)$	$\langle 24, 9 \rangle$	$C_{12} \times C_2$	16	1	1
O7.116	$(0; 3, 8, 24)$	$\langle 24, 2 \rangle$	C_{24}	8	1	1
O7.117	$(0; 3, 9, 9)$	$\langle 27, 2 \rangle$	$C_9 \times C_3$	108	1	1
O7.118	$(0; 3, 9, 9)$	$\langle 27, 4 \rangle$	$C_9: C_3$	108	2	2
O7.119	$(0; 2, 2, 2, 14)$	$\langle 28, 3 \rangle$	D_{28}	252	3	1
O7.120	$(0; 4, 4, 14)$	$\langle 28, 1 \rangle$	$C_7: C_4$	84	1	1
O7.121	$(0; 2, 28, 28)$	$\langle 28, 2 \rangle$	C_{28}	12	1	1
O7.122	$(0; 2, 15, 30)$	$\langle 30, 4 \rangle$	C_{30}	8	1	1
O7.123	$(0; 2, 2, 2, 8)$	$\langle 32, 39 \rangle$	$C_2 \times D_{16}$	768	3	1
O7.124	$(0; 2, 2, 2, 8)$	$\langle 32, 42 \rangle$	$(C_8 \times C_2): C_2$	192	3	1
O7.125	$(0; 2, 2, 2, 8)$	$\langle 32, 43 \rangle$	$(C_2 \times D_8): C_2$	384	6	1
O7.126	$(0; 4, 4, 8)$	$\langle 32, 10 \rangle$	$Q_8: C_4$	128	2	1
O7.127	$(0; 4, 4, 8)$	$\langle 32, 11 \rangle$	$(C_4 \times C_4): C_2$	64	2	1
O7.128	$(0; 4, 4, 8)$	$\langle 32, 13 \rangle$	$C_8: C_4$	128	1	1
O7.129	$(0; 4, 4, 8)$	$\langle 32, 14 \rangle$	$C_8: C_4$	128	1	1
O7.130	$(0; 2, 16, 16)$	$\langle 32, 16 \rangle$	$C_{16} \times C_2$	32	1	1
O7.131	$(0; 2, 16, 16)$	$\langle 32, 17 \rangle$	$C_{16}: C_2$	32	1	1
O7.132	$(0; 2, 2, 2, 6)$	$\langle 36, 10 \rangle$	$S_3 \times S_3$	864	12	1
O7.133	$(0; 3, 4, 12)$	$\langle 36, 6 \rangle$	$C_3 \times (C_3: C_4)$	24	1	1
O7.134	$(0; 2, 6, 21)$	$\langle 42, 4 \rangle$	$C_3 \times D_{14}$	84	1	1
O7.135	$(0; 2, 2, 2, 4)$	$\langle 48, 38 \rangle$	$D_8 \times S_3$	1152	12	1
O7.136	$(0; 2, 2, 2, 4)$	$\langle 48, 41 \rangle$	$(C_4 \times S_3): C_2$	1152	4	1
O7.137	$(0; 2, 2, 2, 4)$	$\langle 48, 48 \rangle$	$C_2 \times S_4$	576	12	1
O7.138	$(0; 3, 4, 6)$	$\langle 48, 32 \rangle$	$C_2 \times SL(2, 3)$	48	1	1
O7.139	$(0; 2, 6, 12)$	$\langle 48, 33 \rangle$	$SL(2, 3): C_2$	48	1	1
O7.140	$(0; 2, 6, 9)$	$\langle 54, 3 \rangle$	$C_3 \times D_{18}$	108	1	1
O7.141	$(0; 2, 6, 9)$	$\langle 54, 6 \rangle$	$(C_9: C_3): C_2$	108	2	2
O7.142	$(0; 2, 7, 7)$	$\langle 56, 11 \rangle$	$(C_2 \times C_2 \times C_2): C_7$	336	2	1
O7.143	$(0; 2, 4, 28)$	$\langle 56, 4 \rangle$	$C_4 \times D_{14}$	168	1	1
O7.144	$(0; 2, 4, 16)$	$\langle 64, 38 \rangle$	$(C_{16} \times C_2): C_2$	256	1	1
O7.145	$(0; 2, 4, 16)$	$\langle 64, 41 \rangle$	$(C_{16}: C_2): C_2$	128	1	1
O7.146	$(0; 3, 3, 6)$	$\langle 72, 25 \rangle$	$C_3 \times SL(2, 3)$	144	1	1
O7.147	$(0; 2, 3, 12)$	$\langle 144, 127 \rangle$	$(C_3 \times SL(2, 3)): C_2$	288	1	1
O7.148	$(0; 2, 3, 7)$	$\langle 504, 156 \rangle$	$PSL(2, 8)$	1512	1	1

Table 11: Census of finite group actions on Riemann surfaces of genus $g = 8$

Ref	Signature	SMG id	Structure	epi	autg	top
O8.1	$(8; -)$	$\langle 1, 1 \rangle$	1	1	1	1
O8.2	$(4; 2, 2)$	$\langle 2, 1 \rangle$	C_2	256	256	1
O8.3	$(3; 2^6)$	$\langle 2, 1 \rangle$	C_2	64	64	1
O8.4	$(2; 2^{10})$	$\langle 2, 1 \rangle$	C_2	16	16	1
O8.5	$(1; 2^{14})$	$\langle 2, 1 \rangle$	C_2	4	4	1
O8.6	$(0; 2^{18})$	$\langle 2, 1 \rangle$	C_2	1	1	1
O8.7	$(2; 3, 3, 3, 3)$	$\langle 3, 1 \rangle$	C_3	486	243	1
O8.8	$(1; 3^7)$	$\langle 3, 1 \rangle$	C_3	378	189	1
O8.9	$(0; 3^{10})$	$\langle 3, 1 \rangle$	C_3	342	171	2
O8.10	$(2; 2, 2, 2)$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	1536	256	1
O8.11	$(2; 4, 4)$	$\langle 4, 1 \rangle$	C_4	512	256	1
O8.12	$(1; 2^7)$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	8736	1456	2
O8.13	$(1; 2^4, 4^2)$	$\langle 4, 1 \rangle$	C_4	32	16	1
O8.14	$(1; 2, 4^4)$	$\langle 4, 1 \rangle$	C_4	128	64	1
O8.15	$(0; 2^{11})$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	44286	7381	4
O8.16	$(0; 2^8, 4^2)$	$\langle 4, 1 \rangle$	C_4	2	1	1
O8.17	$(0; 2^5, 4^4)$	$\langle 4, 1 \rangle$	C_4	8	4	1
O8.18	$(0; 2^2, 4^6)$	$\langle 4, 1 \rangle$	C_4	32	16	2
O8.19	$(0; 5^6)$	$\langle 5, 1 \rangle$	C_5	820	205	5
O8.20	$(1; 2, 2, 3, 3)$	$\langle 6, 1 \rangle$	S_3	432	72	1
O8.21	$(1; 2, 2, 3, 3)$	$\langle 6, 2 \rangle$	C_6	72	36	1
O8.22	$(1; 3, 6, 6)$	$\langle 6, 2 \rangle$	C_6	72	36	1
O8.23	$(0; 2^6, 3^2)$	$\langle 6, 1 \rangle$	S_3	972	162	1
O8.24	$(0; 2^6, 3^2)$	$\langle 6, 2 \rangle$	C_6	2	1	1
O8.25	$(0; 2^2, 3^5)$	$\langle 6, 1 \rangle$	S_3	96	16	1
O8.26	$(0; 2^2, 3^5)$	$\langle 6, 2 \rangle$	C_6	10	5	1
O8.27	$(0; 2^3, 3^3, 6)$	$\langle 6, 2 \rangle$	C_6	6	3	1
O8.28	$(0; 2^4, 3, 6^2)$	$\langle 6, 2 \rangle$	C_6	2	1	1
O8.29	$(0; 3^4, 6^2)$	$\langle 6, 2 \rangle$	C_6	22	11	3
O8.30	$(0; 2, 3^2, 6^3)$	$\langle 6, 2 \rangle$	C_6	10	5	2
O8.31	$(0; 2^2, 6^4)$	$\langle 6, 2 \rangle$	C_6	6	3	1
O8.32	$(2; -)$	$\langle 7, 1 \rangle$	C_7	2400	400	1
O8.33	$(1; 2, 2, 4)$	$\langle 8, 3 \rangle$	D_8	512	64	1
O8.34	$(1; 8, 8)$	$\langle 8, 1 \rangle$	C_8	256	64	1
O8.35	$(0; 2^6, 4)$	$\langle 8, 3 \rangle$	D_8	4088	511	4
O8.36	$(0; 2^3, 4^3)$	$\langle 8, 3 \rangle$	D_8	96	12	1
O8.37	$(0; 2^3, 4^3)$	$\langle 8, 4 \rangle$	Q_8	24	1	1
O8.38	$(0; 4^5)$	$\langle 8, 4 \rangle$	Q_8	960	40	1
O8.39	$(0; 2^4, 8^2)$	$\langle 8, 1 \rangle$	C_8	4	1	1
O8.40	$(0; 2, 4^2, 8^2)$	$\langle 8, 1 \rangle$	C_8	16	4	2
O8.41	$(0; 9, 9, 9, 9)$	$\langle 9, 1 \rangle$	C_9	162	27	3
O8.42	$(0; 2^2, 5^3)$	$\langle 10, 1 \rangle$	D_{10}	320	16	2
O8.43	$(0; 2^2, 5^3)$	$\langle 10, 2 \rangle$	C_{10}	12	3	1
O8.44	$(0; 5, 5, 10, 10)$	$\langle 10, 2 \rangle$	C_{10}	52	13	6

Continued on next page

Table 11 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O8.45	$(0; 2^5, 3)$	$\langle 12, 4 \rangle$	D_{12}	2280	190	2
O8.46	$(0; 2, 3^4)$	$\langle 12, 3 \rangle$	A_4	1152	48	1
O8.47	$(0; 2^2, 3, 4^2)$	$\langle 12, 1 \rangle$	$C_3 : C_4$	12	1	1
O8.48	$(0; 2^2, 3^2, 6)$	$\langle 12, 4 \rangle$	D_{12}	48	4	1
O8.49	$(0; 2^2, 3^2, 6)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	12	1	1
O8.50	$(0; 2^3, 6^2)$	$\langle 12, 4 \rangle$	D_{12}	72	6	1
O8.51	$(0; 2^3, 6^2)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	120	10	3
O8.52	$(0; 4, 4, 6, 6)$	$\langle 12, 1 \rangle$	$C_3 : C_4$	24	2	1
O8.53	$(0; 4, 4, 6, 6)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O8.54	$(0; 3, 6, 6, 6)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	36	3	1
O8.55	$(0; 3, 4, 6, 12)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O8.56	$(0; 3, 3, 12, 12)$	$\langle 12, 2 \rangle$	C_{12}	12	3	2
O8.57	$(0; 2, 6, 12, 12)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O8.58	$(1; 2, 2)$	$\langle 14, 1 \rangle$	D_{14}	1344	32	1
O8.59	$(1; 2, 2)$	$\langle 14, 2 \rangle$	C_{14}	192	32	1
O8.60	$(0; 2^6)$	$\langle 14, 1 \rangle$	D_{14}	16800	400	1
O8.61	$(0; 3, 3, 5, 5)$	$\langle 15, 1 \rangle$	C_{15}	8	1	1
O8.62	$(0; 2^4, 8)$	$\langle 16, 7 \rangle$	D_{16}	2240	70	2
O8.63	$(0; 2, 4, 4, 8)$	$\langle 16, 8 \rangle$	QD_{16}	64	4	1
O8.64	$(0; 2, 4, 4, 8)$	$\langle 16, 9 \rangle$	Q_{16}	32	1	1
O8.65	$(0; 2, 2, 16, 16)$	$\langle 16, 1 \rangle$	C_{16}	8	1	1
O8.66	$(0; 17, 17, 17)$	$\langle 17, 1 \rangle$	C_{17}	240	15	3
O8.67	$(0; 2, 2, 9, 9)$	$\langle 18, 1 \rangle$	D_{18}	324	6	2
O8.68	$(0; 2, 2, 9, 9)$	$\langle 18, 2 \rangle$	C_{18}	6	1	1
O8.69	$(0; 9, 18, 18)$	$\langle 18, 2 \rangle$	C_{18}	18	3	2
O8.70	$(0; 2, 2, 5, 10)$	$\langle 20, 4 \rangle$	D_{20}	160	4	2
O8.71	$(0; 2, 2, 5, 10)$	$\langle 20, 5 \rangle$	$C_{10} \times C_2$	24	1	1
O8.72	$(0; 10, 10, 10)$	$\langle 20, 5 \rangle$	$C_{10} \times C_2$	72	3	1
O8.73	$(0; 5, 20, 20)$	$\langle 20, 2 \rangle$	C_{20}	24	3	2
O8.74	$(0; 3, 3, 3, 3)$	$\langle 21, 1 \rangle$	$C_7 : C_3$	2016	48	1
O8.75	$(0; 2, 3, 3, 4)$	$\langle 24, 3 \rangle$	$SL(2, 3)$	24	1	1
O8.76	$(0; 2, 3, 3, 4)$	$\langle 24, 12 \rangle$	S_4	192	8	1
O8.77	$(0; 2, 2, 4, 6)$	$\langle 24, 6 \rangle$	D_{24}	48	1	1
O8.78	$(0; 2, 2, 4, 6)$	$\langle 24, 8 \rangle$	$(C_6 \times C_2) : C_2$	96	4	2
O8.79	$(0; 6, 8, 8)$	$\langle 24, 1 \rangle$	$C_3 : C_8$	24	1	1
O8.80	$(0; 2, 2, 3, 12)$	$\langle 24, 6 \rangle$	D_{24}	96	2	1
O8.81	$(0; 2, 2, 3, 12)$	$\langle 24, 10 \rangle$	$C_3 \times D_8$	16	1	1
O8.82	$(0; 6, 6, 12)$	$\langle 24, 10 \rangle$	$C_3 \times D_8$	16	1	1
O8.83	$(0; 4, 12, 12)$	$\langle 24, 11 \rangle$	$C_3 \times Q_8$	48	1	1
O8.84	$(0; 3, 24, 24)$	$\langle 24, 2 \rangle$	C_{24}	8	1	1
O8.85	$(0; 2^5)$	$\langle 28, 3 \rangle$	D_{28}	13440	160	1
O8.86	$(0; 2, 2, 3, 5)$	$\langle 30, 3 \rangle$	D_{30}	120	1	1
O8.87	$(0; 5, 6, 6)$	$\langle 30, 2 \rangle$	$C_3 \times D_{10}$	40	1	1
O8.88	$(0; 3, 10, 10)$	$\langle 30, 1 \rangle$	$C_5 \times S_3$	24	1	1
O8.89	$(0; 2, 2, 2, 16)$	$\langle 32, 18 \rangle$	D_{32}	384	3	1
O8.90	$(0; 4, 4, 16)$	$\langle 32, 20 \rangle$	Q_{32}	128	1	1

Continued on next page

Table 11 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O8.91	$(0; 2, 32, 32)$	$\langle 32, 1 \rangle$	C_{32}	16	1	1
O8.92	$(0; 2, 17, 34)$	$\langle 34, 2 \rangle$	C_{34}	16	1	1
O8.93	$(0; 2, 2, 2, 9)$	$\langle 36, 4 \rangle$	D_{36}	324	3	1
O8.94	$(0; 4, 4, 9)$	$\langle 36, 1 \rangle$	$C_9 : C_4$	108	1	1
O8.95	$(0; 2, 18, 18)$	$\langle 36, 5 \rangle$	$C_{18} \times C_2$	36	1	1
O8.96	$(0; 2, 10, 20)$	$\langle 40, 10 \rangle$	$C_5 \times D_8$	32	1	1
O8.97	$(0; 2, 2, 3, 3)$	$\langle 42, 1 \rangle$	$(C_7 : C_3) : C_2$	672	16	1
O8.98	$(0; 3, 6, 6)$	$\langle 42, 1 \rangle$	$(C_7 : C_3) : C_2$	84	2	2
O8.99	$(0; 3, 6, 6)$	$\langle 42, 2 \rangle$	$C_2 \times (C_7 : C_3)$	84	2	2
O8.100	$(0; 3, 4, 8)$	$\langle 48, 28 \rangle$	$C_2.S_4 = SL(2, 3).C_2$	48	1	1
O8.101	$(0; 2, 8, 12)$	$\langle 48, 17 \rangle$	$(C_3 \times Q_8) : C_2$	96	1	1
O8.102	$(0; 2, 6, 24)$	$\langle 48, 25 \rangle$	$C_3 \times D_{16}$	64	1	1
O8.103	$(0; 2, 6, 10)$	$\langle 60, 8 \rangle$	$S_3 \times D_{10}$	120	1	1
O8.104	$(0; 2, 4, 32)$	$\langle 64, 53 \rangle$	QD_{64}	256	1	1
O8.105	$(0; 2, 4, 18)$	$\langle 72, 8 \rangle$	$(C_{18} \times C_2) : C_2$	216	1	1
O8.106	$(0; 2, 6, 6)$	$\langle 84, 7 \rangle$	$C_2 \times ((C_7 : C_3) : C_2)$	336	4	2
O8.107	$(0; 3, 3, 4)$	$\langle 168, 42 \rangle$	$PSL(3, 2)$	672	2	2
O8.108	$(0; 2, 3, 8)$	$\langle 336, 208 \rangle$	$PSL(3, 2) : C_2$	672	2	2

Table 12: Census of finite group actions on Riemann surfaces of genus $g = 9$

Ref	Signature	SMG id	Structure	epi	autg	top
O9.1	$(9; -)$	$\langle 1, 1 \rangle$	1	1	1	1
O9.2	$(5; -)$	$\langle 2, 1 \rangle$	C_2	1023	1023	1
O9.3	$(4; 2, 2, 2, 2)$	$\langle 2, 1 \rangle$	C_2	256	256	1
O9.4	$(3; 2^8)$	$\langle 2, 1 \rangle$	C_2	64	64	1
O9.5	$(2; 2^1 2)$	$\langle 2, 1 \rangle$	C_2	16	16	1
O9.6	$(1; 2^1 6)$	$\langle 2, 1 \rangle$	C_2	4	4	1
O9.7	$(0; 2^2 0)$	$\langle 2, 1 \rangle$	C_2	1	1	1
O9.8	$(3; 3, 3)$	$\langle 3, 1 \rangle$	C_3	1458	729	1
O9.9	$(2; 3^5)$	$\langle 3, 1 \rangle$	C_3	810	405	1
O9.10	$(1; 3^8)$	$\langle 3, 1 \rangle$	C_3	774	387	2
O9.11	$(0; 3^1 1)$	$\langle 3, 1 \rangle$	C_3	682	341	2
O9.12	$(3; -)$	$\langle 4, 1 \rangle$	C_4	4032	2016	1
O9.13	$(3; -)$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	3906	651	2
O9.14	$(2; 2, 2, 2, 2)$	$\langle 4, 1 \rangle$	C_4	240	120	1
O9.15	$(2; 2, 2, 2, 2)$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	5328	888	2
O9.16	$(2; 2, 4, 4)$	$\langle 4, 1 \rangle$	C_4	512	256	1
O9.17	$(1; 2^8)$	$\langle 4, 1 \rangle$	C_4	12	6	1
O9.18	$(1; 2^8)$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	26244	4374	4
O9.19	$(1; 2^5, 4^2)$	$\langle 4, 1 \rangle$	C_4	32	16	1
O9.20	$(1; 2^2, 4^4)$	$\langle 4, 1 \rangle$	C_4	128	64	2
O9.21	$(0; 2^1 2)$	$\langle 4, 2 \rangle$	$C_2 \times C_2$	132858	22143	6

Continued on next page

Table 12 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O9.22	$(0; 2^9, 4^2)$	$\langle 4, 1 \rangle$	C_4	2	1	1
O9.23	$(0; 2^6, 4^4)$	$\langle 4, 1 \rangle$	C_4	8	4	2
O9.24	$(0; 2^3, 4^6)$	$\langle 4, 1 \rangle$	C_4	32	16	2
O9.25	$(0; 4^8)$	$\langle 4, 1 \rangle$	C_4	128	64	3
O9.26	$(1; 5, 5, 5, 5)$	$\langle 5, 1 \rangle$	C_5	1300	325	3
O9.27	$(2; 3)$	$\langle 6, 1 \rangle$	S_3	810	135	1
O9.28	$(1; 2^4, 3)$	$\langle 6, 1 \rangle$	S_3	1944	324	1
O9.29	$(1; 3, 3, 3, 3)$	$\langle 6, 1 \rangle$	S_3	144	24	1
O9.30	$(1; 3, 3, 3, 3)$	$\langle 6, 2 \rangle$	C_6	162	81	1
O9.31	$(1; 2, 3, 3, 6)$	$\langle 6, 2 \rangle$	C_6	72	36	1
O9.32	$(1; 2, 2, 6, 6)$	$\langle 6, 2 \rangle$	C_6	72	36	1
O9.33	$(0; 2^8, 3)$	$\langle 6, 1 \rangle$	S_3	4374	729	1
O9.34	$(0; 2^4, 3^4)$	$\langle 6, 1 \rangle$	S_3	432	72	1
O9.35	$(0; 2^4, 3^4)$	$\langle 6, 2 \rangle$	C_6	6	3	1
O9.36	$(0; 2^5, 3^2, 6)$	$\langle 6, 2 \rangle$	C_6	2	1	1
O9.37	$(0; 2, 3^5, 6)$	$\langle 6, 2 \rangle$	C_6	22	11	2
O9.38	$(0; 2^6, 6^2)$	$\langle 6, 2 \rangle$	C_6	2	1	1
O9.39	$(0; 2^2, 3^3, 6^2)$	$\langle 6, 2 \rangle$	C_6	10	5	2
O9.40	$(0; 2^3, 3, 6^3)$	$\langle 6, 2 \rangle$	C_6	6	3	1
O9.41	$(0; 3^2, 6^4)$	$\langle 6, 2 \rangle$	C_6	22	11	3
O9.42	$(0; 2, 6^5)$	$\langle 6, 2 \rangle$	C_6	10	5	1
O9.43	$(0; 7^5)$	$\langle 7, 1 \rangle$	C_7	1110	185	6
O9.44	$(2; -)$	$\langle 8, 1 \rangle$	C_8	3840	960	1
O9.45	$(2; -)$	$\langle 8, 2 \rangle$	$C_4 \times C_2$	3360	420	2
O9.46	$(2; -)$	$\langle 8, 3 \rangle$	D_8	1440	180	1
O9.47	$(2; -)$	$\langle 8, 4 \rangle$	Q_8	1440	60	1
O9.48	$(2; -)$	$\langle 8, 5 \rangle$	$C_2 \times C_2 \times C_2$	2520	15	1
O9.49	$(1; 2, 2, 2, 2)$	$\langle 8, 1 \rangle$	C_8	48	12	1
O9.50	$(1; 2, 2, 2, 2)$	$\langle 8, 2 \rangle$	$C_4 \times C_2$	984	123	4
O9.51	$(1; 2, 2, 2, 2)$	$\langle 8, 3 \rangle$	D_8	4992	624	3
O9.52	$(1; 2, 2, 2, 2)$	$\langle 8, 5 \rangle$	$C_2 \times C_2 \times C_2$	16968	101	3
O9.53	$(1; 2, 4, 4)$	$\langle 8, 1 \rangle$	C_8	96	24	1
O9.54	$(1; 2, 4, 4)$	$\langle 8, 2 \rangle$	$C_4 \times C_2$	704	88	2
O9.55	$(1; 2, 4, 4)$	$\langle 8, 3 \rangle$	D_8	96	12	1
O9.56	$(1; 2, 4, 4)$	$\langle 8, 4 \rangle$	Q_8	288	12	1
O9.57	$(0; 2^8)$	$\langle 8, 3 \rangle$	D_8	46368	5796	4
O9.58	$(0; 2^8)$	$\langle 8, 5 \rangle$	$C_2 \times C_2 \times C_2$	709128	4221	8
O9.59	$(0; 2^5, 4^2)$	$\langle 8, 2 \rangle$	$C_4 \times C_2$	968	121	11
O9.60	$(0; 2^5, 4^2)$	$\langle 8, 3 \rangle$	D_8	1440	180	3
O9.61	$(0; 2^2, 4^4)$	$\langle 8, 2 \rangle$	$C_4 \times C_2$	560	70	11
O9.62	$(0; 2^2, 4^4)$	$\langle 8, 3 \rangle$	D_8	64	8	1
O9.63	$(0; 2^2, 4^4)$	$\langle 8, 4 \rangle$	Q_8	144	6	1
O9.64	$(0; 2^3, 4, 8^2)$	$\langle 8, 1 \rangle$	C_8	8	2	2
O9.65	$(0; 4^3, 8^2)$	$\langle 8, 1 \rangle$	C_8	32	8	4
O9.66	$(0; 2, 8^4)$	$\langle 8, 1 \rangle$	C_8	64	16	3
O9.67	$(1; 9, 9)$	$\langle 9, 1 \rangle$	C_9	486	81	1

Continued on next page

Table 12 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O9.68	$(0; 3^3, 9^2)$	$\langle 9, 1 \rangle$	C_9	48	8	2
O9.69	$(1; 5, 5)$	$\langle 10, 1 \rangle$	D_{10}	240	12	2
O9.70	$(1; 5, 5)$	$\langle 10, 2 \rangle$	C_{10}	300	75	1
O9.71	$(0; 2^4, 5^2)$	$\langle 10, 1 \rangle$	D_{10}	2000	100	2
O9.72	$(0; 2^4, 5^2)$	$\langle 10, 2 \rangle$	C_{10}	4	1	1
O9.73	$(0; 2^2, 5, 10^2)$	$\langle 10, 2 \rangle$	C_{10}	12	3	2
O9.74	$(0; 10, 10, 10, 10)$	$\langle 10, 2 \rangle$	C_{10}	52	13	3
O9.75	$(1; 3, 3)$	$\langle 12, 1 \rangle$	$C_3: C_4$	144	12	1
O9.76	$(1; 3, 3)$	$\langle 12, 2 \rangle$	C_{12}	216	54	1
O9.77	$(1; 3, 3)$	$\langle 12, 3 \rangle$	A_4	1080	45	1
O9.78	$(1; 3, 3)$	$\langle 12, 4 \rangle$	D_{12}	72	6	1
O9.79	$(1; 3, 3)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	108	9	1
O9.80	$(1; 2, 6)$	$\langle 12, 1 \rangle$	$C_3: C_4$	72	6	1
O9.81	$(1; 2, 6)$	$\langle 12, 4 \rangle$	D_{12}	72	6	1
O9.82	$(0; 2^4, 3^2)$	$\langle 12, 3 \rangle$	A_4	648	27	1
O9.83	$(0; 2^4, 3^2)$	$\langle 12, 4 \rangle$	D_{12}	792	66	2
O9.84	$(0; 2^4, 3^2)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	36	3	1
O9.85	$(0; 3^5)$	$\langle 12, 3 \rangle$	A_4	2520	105	2
O9.86	$(0; 2, 3^2, 4^2)$	$\langle 12, 1 \rangle$	$C_3: C_4$	24	2	1
O9.87	$(0; 2, 3^2, 4^2)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O9.88	$(0; 2^5, 6)$	$\langle 12, 4 \rangle$	D_{12}	2280	190	3
O9.89	$(0; 2^2, 4^2, 6)$	$\langle 12, 1 \rangle$	$C_3: C_4$	12	1	1
O9.90	$(0; 2^2, 3, 6^2)$	$\langle 12, 4 \rangle$	D_{12}	48	4	1
O9.91	$(0; 2^2, 3, 6^2)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	36	3	2
O9.92	$(0; 6, 6, 6, 6)$	$\langle 12, 5 \rangle$	$C_6 \times C_2$	108	9	2
O9.93	$(0; 2^2, 3, 4, 12)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O9.94	$(0; 4, 6, 6, 12)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O9.95	$(0; 2^3, 12^2)$	$\langle 12, 2 \rangle$	C_{12}	4	1	1
O9.96	$(0; 4, 4, 12, 12)$	$\langle 12, 2 \rangle$	C_{12}	16	4	3
O9.97	$(0; 3, 6, 12, 12)$	$\langle 12, 2 \rangle$	C_{12}	12	3	2
O9.98	$(0; 2, 7, 7, 14)$	$\langle 14, 2 \rangle$	C_{14}	30	5	3
O9.99	$(0; 3, 3, 5, 15)$	$\langle 15, 1 \rangle$	C_{15}	8	1	1
O9.100	$(1; 2, 2)$	$\langle 16, 1 \rangle$	C_{16}	192	24	1
O9.101	$(1; 2, 2)$	$\langle 16, 2 \rangle$	$C_4 \times C_4$	288	3	1
O9.102	$(1; 2, 2)$	$\langle 16, 3 \rangle$	$(C_4 \times C_2): C_2$	960	30	2
O9.103	$(1; 2, 2)$	$\langle 16, 4 \rangle$	$C_4: C_4$	192	6	1
O9.104	$(1; 2, 2)$	$\langle 16, 5 \rangle$	$C_8 \times C_2$	480	30	2
O9.105	$(1; 2, 2)$	$\langle 16, 6 \rangle$	$C_8: C_2$	384	24	1
O9.106	$(1; 2, 2)$	$\langle 16, 7 \rangle$	D_{16}	1536	48	1
O9.107	$(1; 2, 2)$	$\langle 16, 8 \rangle$	$QD16$	768	48	1
O9.108	$(1; 2, 2)$	$\langle 16, 10 \rangle$	$C_4 \times C_2 \times C_2$	576	3	1
O9.109	$(1; 2, 2)$	$\langle 16, 11 \rangle$	$C_2 \times D_8$	960	15	≤ 2
O9.110	$(1; 2, 2)$	$\langle 16, 12 \rangle$	$C_2 \times Q_8$	192	1	1
O9.111	$(1; 2, 2)$	$\langle 16, 13 \rangle$	$(C_4 \times C_2): C_2$	576	12	1
O9.112	$(0; 2^6)$	$\langle 16, 7 \rangle$	D_{16}	36480	1140	2
O9.113	$(0; 2^6)$	$\langle 16, 11 \rangle$	$C_2 \times D_8$	75840	1185	8

Continued on next page

Table 12 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O9.114	$(0; 2^6)$	$\langle 16, 13 \rangle$	$(C_4 \times C_2): C_2$	2880	60	1
O9.115	$(0; 2^6)$	$\langle 16, 14 \rangle$	$C_2 \times C_2 \times C_2 \times C_2$	504000	25	2
O9.116	$(0; 2^3, 4^2)$	$\langle 16, 3 \rangle$	$(C_4 \times C_2): C_2$	2528	79	12
O9.117	$(0; 2^3, 4^2)$	$\langle 16, 8 \rangle$	QD16	192	12	1
O9.118	$(0; 2^3, 4^2)$	$\langle 16, 10 \rangle$	$C_4 \times C_2 \times C_2$	2112	11	5
O9.119	$(0; 2^3, 4^2)$	$\langle 16, 11 \rangle$	$C_2 \times D_8$	960	15	3
O9.120	$(0; 2^3, 4^2)$	$\langle 16, 13 \rangle$	$(C_4 \times C_2): C_2$	864	18	2
O9.121	$(0; 4, 4, 4, 4)$	$\langle 16, 2 \rangle$	$C_4 \times C_4$	1152	12	3
O9.122	$(0; 4, 4, 4, 4)$	$\langle 16, 3 \rangle$	$(C_4 \times C_2): C_2$	384	12	2
O9.123	$(0; 4, 4, 4, 4)$	$\langle 16, 4 \rangle$	$C_4: C_4$	1152	36	4
O9.124	$(0; 4, 4, 4, 4)$	$\langle 16, 9 \rangle$	Q_{16}	384	12	1
O9.125	$(0; 4, 4, 4, 4)$	$\langle 16, 10 \rangle$	$C_4 \times C_2 \times C_2$	192	1	1
O9.126	$(0; 4, 4, 4, 4)$	$\langle 16, 12 \rangle$	$C_2 \times Q_8$	576	3	1
O9.127	$(0; 2, 4, 8, 8)$	$\langle 16, 5 \rangle$	$C_8 \times C_2$	80	5	5
O9.128	$(0; 2, 4, 8, 8)$	$\langle 16, 6 \rangle$	$C_8: C_2$	80	5	3
O9.129	$(1; 9)$	$\langle 18, 1 \rangle$	D_{18}	162	3	1
O9.130	$(0; 2^4, 9)$	$\langle 18, 1 \rangle$	D_{18}	4374	81	1
O9.131	$(0; 2, 2, 18, 18)$	$\langle 18, 2 \rangle$	C_{18}	6	1	1
O9.132	$(0; 19, 19, 19)$	$\langle 19, 1 \rangle$	C_{19}	306	17	4
O9.133	$(1; 5)$	$\langle 20, 1 \rangle$	$C_5: C_4$	240	6	1
O9.134	$(1; 5)$	$\langle 20, 3 \rangle$	$C_5: C_4$	240	12	1
O9.135	$(1; 5)$	$\langle 20, 4 \rangle$	D_{20}	120	3	1
O9.136	$(0; 2^4, 5)$	$\langle 20, 4 \rangle$	D_{20}	3240	81	2
O9.137	$(0; 2, 4, 4, 5)$	$\langle 20, 1 \rangle$	$C_5: C_4$	40	1	1
O9.138	$(0; 2, 4, 4, 5)$	$\langle 20, 3 \rangle$	$C_5: C_4$	200	10	2
O9.139	$(0; 2, 2, 10, 10)$	$\langle 20, 4 \rangle$	D_{20}	160	4	2
O9.140	$(0; 2, 2, 10, 10)$	$\langle 20, 5 \rangle$	$C_{10} \times C_2$	72	3	2
O9.141	$(0; 10, 20, 20)$	$\langle 20, 2 \rangle$	C_{20}	24	3	2
O9.142	$(0; 7, 21, 21)$	$\langle 21, 2 \rangle$	C_{21}	60	5	3
O9.143	$(1; 3)$	$\langle 24, 1 \rangle$	$C_3: C_8$	288	12	1
O9.144	$(1; 3)$	$\langle 24, 5 \rangle$	$C_4 \times S_3$	144	6	1
O9.145	$(1; 3)$	$\langle 24, 7 \rangle$	$C_2 \times (C_3: C_4)$	144	3	1
O9.146	$(1; 3)$	$\langle 24, 12 \rangle$	S_4	216	9	1
O9.147	$(0; 2^4, 3)$	$\langle 24, 6 \rangle$	D_{24}	2592	54	1
O9.148	$(0; 2^4, 3)$	$\langle 24, 8 \rangle$	$(C_6 \times C_2): C_2$	288	12	1
O9.149	$(0; 2^4, 3)$	$\langle 24, 12 \rangle$	S_4	1944	81	2
O9.150	$(0; 2^4, 3)$	$\langle 24, 14 \rangle$	$C_2 \times C_2 \times S_3$	2160	15	2
O9.151	$(0; 3, 3, 3, 3)$	$\langle 24, 3 \rangle$	$SL(2, 3)$	216	9	1
O9.152	$(0; 2, 3, 4, 4)$	$\langle 24, 5 \rangle$	$C_4 \times S_3$	48	2	1
O9.153	$(0; 2, 3, 4, 4)$	$\langle 24, 7 \rangle$	$C_2 \times (C_3: C_4)$	48	1	1
O9.154	$(0; 2, 3, 4, 4)$	$\langle 24, 12 \rangle$	S_4	72	3	1
O9.155	$(0; 2, 3, 3, 6)$	$\langle 24, 13 \rangle$	$C_2 \times A_4$	120	5	2
O9.156	$(0; 2, 2, 6, 6)$	$\langle 24, 8 \rangle$	$(C_6 \times C_2): C_2$	96	4	2
O9.157	$(0; 2, 2, 6, 6)$	$\langle 24, 10 \rangle$	$C_3 \times D_8$	96	6	2
O9.158	$(0; 2, 2, 6, 6)$	$\langle 24, 13 \rangle$	$C_2 \times A_4$	192	8	3
O9.159	$(0; 2, 2, 6, 6)$	$\langle 24, 14 \rangle$	$C_2 \times C_2 \times S_3$	288	2	1

Continued on next page

Table 12 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O9.160	(0; 2, 2, 6, 6)	$\langle 24, 15 \rangle$	$C_6 \times C_2 \times C_2$	336	1	1
O9.161	(0; 2, 2, 4, 12)	$\langle 24, 5 \rangle$	$C_4 \times S_3$	96	4	3
O9.162	(0; 2, 2, 4, 12)	$\langle 24, 6 \rangle$	D_{24}	96	2	1
O9.163	(0; 8, 8, 12)	$\langle 24, 1 \rangle$	$C_3 : C_8$	48	2	2
O9.164	(0; 6, 12, 12)	$\langle 24, 9 \rangle$	$C_{12} \times C_2$	16	1	1
O9.165	(0; 6, 8, 24)	$\langle 24, 2 \rangle$	C_{24}	8	1	1
O9.166	(0; 4, 24, 24)	$\langle 24, 2 \rangle$	C_{24}	16	2	2
O9.167	(0; 3, 27, 27)	$\langle 27, 1 \rangle$	C_{27}	36	2	1
O9.168	(0; 4, 7, 28)	$\langle 28, 2 \rangle$	C_{28}	12	1	1
O9.169	(0; 3, 10, 30)	$\langle 30, 4 \rangle$	C_{30}	8	1	1
O9.170	(1; 2)	$\langle 32, 2 \rangle$	$(C_4 \times C_2) : C_4$	384	1	1
O9.171	(1; 2)	$\langle 32, 4 \rangle$	$C_8 : C_4$	384	3	1
O9.172	(1; 2)	$\langle 32, 5 \rangle$	$(C_8 \times C_2) : C_2$	384	6	1
O9.173	(1; 2)	$\langle 32, 6 \rangle$	$((C_4 \times C_2) : C_2) : C_2$	384	6	1
O9.174	(1; 2)	$\langle 32, 7 \rangle$	$(C_8 : C_2) : C_2$	384	3	1
O9.175	(1; 2)	$\langle 32, 8 \rangle$	$C_2 \cdot ((C_4 \times C_2) : C_2)$	384	3	1
O9.176	(1; 2)	$\langle 32, 12 \rangle$	$C_4 : C_8$	384	6	1
O9.177	(1; 2)	$\langle 32, 17 \rangle$	$C_{16} : C_2$	384	12	1
O9.178	(0; 2 ⁵)	$\langle 32, 18 \rangle$	D_{32}	15360	120	1
O9.179	(0; 2 ⁵)	$\langle 32, 27 \rangle$	$(C_2 \times C_2 \times C_2 \times C_2) : C_2$	15360	40	1
O9.180	(0; 2 ⁵)	$\langle 32, 34 \rangle$	$(C_4 \times C_4) : C_2$	23040	15	1
O9.181	(0; 2 ⁵)	$\langle 32, 39 \rangle$	$C_2 \times D_{16}$	38400	150	2
O9.182	(0; 2 ⁵)	$\langle 32, 43 \rangle$	$(C_2 \times D_8) : C_2$	7680	120	1
O9.183	(0; 2 ⁵)	$\langle 32, 46 \rangle$	$C_2 \times C_2 \times D_8$	46080	15	1
O9.184	(0; 2 ⁵)	$\langle 32, 49 \rangle$	$(C_2 \times D_8) : C_2$	17280	15	1
O9.185	(0; 2 ⁵)	$\langle 32, 50 \rangle$	$(C_2 \times Q_8) : C_2$	1920	1	1
O9.186	(0; 2, 2, 4, 4)	$\langle 32, 6 \rangle$	$((C_4 \times C_2) : C_2) : C_2$	1152	18	4
O9.187	(0; 2, 2, 4, 4)	$\langle 32, 9 \rangle$	$(C_8 \times C_2) : C_2$	512	8	2
O9.188	(0; 2, 2, 4, 4)	$\langle 32, 11 \rangle$	$(C_4 \times C_4) : C_2$	128	4	2
O9.189	(0; 2, 2, 4, 4)	$\langle 32, 19 \rangle$	QD_{32}	512	8	1
O9.190	(0; 2, 2, 4, 4)	$\langle 32, 22 \rangle$	$C_2 \times ((C_4 \times C_2) : C_2)$	1536	3	2
O9.191	(0; 2, 2, 4, 4)	$\langle 32, 25 \rangle$	$C_4 \times D_8$	512	4	2
O9.192	(0; 2, 2, 4, 4)	$\langle 32, 27 \rangle$	$(C_2 \times C_2 \times C_2 \times C_2) : C_2$	768	2	1
O9.193	(0; 2, 2, 4, 4)	$\langle 32, 28 \rangle$	$(C_4 \times C_2 \times C_2) : C_2$	768	6	2
O9.194	(0; 2, 2, 4, 4)	$\langle 32, 30 \rangle$	$(C_4 \times C_2 \times C_2) : C_2$	512	4	1
O9.195	(0; 2, 2, 4, 4)	$\langle 32, 31 \rangle$	$(C_4 \times C_4) : C_2$	512	2	2
O9.196	(0; 2, 2, 4, 4)	$\langle 32, 34 \rangle$	$(C_4 \times C_4) : C_2$	1536	1	1
O9.197	(0; 2, 2, 4, 4)	$\langle 32, 40 \rangle$	$C_2 \times QD_{16}$	256	2	1
O9.198	(0; 2, 2, 4, 4)	$\langle 32, 42 \rangle$	$(C_8 \times C_2) : C_2$	256	4	1
O9.199	(0; 4, 8, 8)	$\langle 32, 3 \rangle$	$C_8 \times C_4$	128	1	1
O9.200	(0; 4, 8, 8)	$\langle 32, 4 \rangle$	$C_8 : C_4$	128	1	1
O9.201	(0; 4, 8, 8)	$\langle 32, 5 \rangle$	$(C_8 \times C_2) : C_2$	64	1	1
O9.202	(0; 4, 8, 8)	$\langle 32, 8 \rangle$	$C_2 \cdot ((C_4 \times C_2) : C_2)$	128	1	1
O9.203	(0; 4, 8, 8)	$\langle 32, 12 \rangle$	$C_4 : C_8$	128	2	2
O9.204	(0; 2, 2, 2, 18)	$\langle 36, 4 \rangle$	D_{36}	324	3	1
O9.205	(0; 4, 4, 18)	$\langle 36, 1 \rangle$	$C_9 : C_4$	108	1	1

Continued on next page

Table 12 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O9.206	(0; 2, 36, 36)	$\langle 36, 2 \rangle$	C_{36}	12	1	1
O9.207	(0; 2, 19, 38)	$\langle 38, 2 \rangle$	C_{38}	18	1	1
O9.208	(0; 2, 2, 2, 10)	$\langle 40, 8 \rangle$	$(C_{10} \times C_2): C_2$	480	6	1
O9.209	(0; 2, 2, 2, 10)	$\langle 40, 13 \rangle$	$C_2 \times C_2 \times D_{10}$	1440	3	1
O9.210	(0; 4, 4, 10)	$\langle 40, 7 \rangle$	$C_2 \times (C_5: C_4)$	160	1	1
O9.211	(0; 4, 4, 10)	$\langle 40, 12 \rangle$	$C_2 \times (C_5: C_4)$	80	2	1
O9.212	(0; 2, 20, 20)	$\langle 40, 9 \rangle$	$C_{20} \times C_2$	32	1	1
O9.213	(0; 2, 14, 21)	$\langle 42, 3 \rangle$	$C_7 \times S_3$	36	1	1
O9.214	(0; 2, 2, 3, 3)	$\langle 48, 29 \rangle$	$GL(2, 3)$	288	6	1
O9.215	(0; 2, 2, 3, 3)	$\langle 48, 33 \rangle$	$SL(2, 3): C_2$	144	3	1
O9.216	(0; 2, 2, 3, 3)	$\langle 48, 50 \rangle$	$(C_2 \times C_2 \times C_2 \times C_2): C_3$	5760	1	1
O9.217	(0; 2, 2, 2, 6)	$\langle 48, 15 \rangle$	$(C_3 \times D_8): C_2$	1152	12	1
O9.218	(0; 2, 2, 2, 6)	$\langle 48, 38 \rangle$	$D_8 \times S_3$	576	6	1
O9.219	(0; 2, 2, 2, 6)	$\langle 48, 43 \rangle$	$C_2 \times ((C_6 \times C_2): C_2)$	576	3	1
O9.220	(0; 2, 2, 2, 6)	$\langle 48, 48 \rangle$	$C_2 \times S_4$	864	18	2
O9.221	(0; 4, 4, 6)	$\langle 48, 19 \rangle$	$(C_2 \times (C_3: C_4)): C_2$	192	1	1
O9.222	(0; 4, 4, 6)	$\langle 48, 30 \rangle$	$A_4: C_4$	144	3	2
O9.223	(0; 4, 4, 6)	$\langle 48, 48 \rangle$	$C_2 \times S_4$	48	1	1
O9.224	(0; 3, 6, 6)	$\langle 48, 32 \rangle$	$C_2 \times SL(2, 3)$	48	1	1
O9.225	(0; 3, 4, 12)	$\langle 48, 31 \rangle$	$C_4 \times A_4$	48	1	1
O9.226	(0; 2, 12, 12)	$\langle 48, 21 \rangle$	$C_3 \times ((C_4 \times C_2): C_2)$	64	1	1
O9.227	(0; 2, 12, 12)	$\langle 48, 31 \rangle$	$C_4 \times A_4$	96	2	2
O9.228	(0; 2, 8, 24)	$\langle 48, 4 \rangle$	$C_8 \times S_3$	48	1	1
O9.229	(0; 2, 8, 24)	$\langle 48, 5 \rangle$	$C_{24}: C_2$	48	1	1
O9.230	(0; 3, 3, 19)	$\langle 57, 1 \rangle$	$C_{19}: C_3$	684	2	1
O9.231	(0; 3, 5, 5)	$\langle 60, 5 \rangle$	A_5	240	2	2
O9.232	(0; 2, 2, 2, 4)	$\langle 64, 73 \rangle$	$(C_2 \times C_2 \times D_8): C_2$	3072	1	1
O9.233	(0; 2, 2, 2, 4)	$\langle 64, 128 \rangle$	$(C_2 \times C_2 \times D_8): C_2$	3072	6	1
O9.234	(0; 2, 2, 2, 4)	$\langle 64, 134 \rangle$	$((C_4 \times C_4): C_2): C_2$	768	3	1
O9.235	(0; 2, 2, 2, 4)	$\langle 64, 135 \rangle$	$((C_4 \times C_4): C_2): C_2$	768	3	1
O9.236	(0; 2, 2, 2, 4)	$\langle 64, 138 \rangle$	$((C_4 \times C_2): C_2): C_2): C_2$	2304	6	1
O9.237	(0; 2, 2, 2, 4)	$\langle 64, 140 \rangle$	$(C_4 \times D_8): C_2$	3072	6	1
O9.238	(0; 2, 2, 2, 4)	$\langle 64, 177 \rangle$	$(C_2 \times D_{16}): C_2$	3072	3	1
O9.239	(0; 2, 2, 2, 4)	$\langle 64, 190 \rangle$	$(C_2 \times D_{16}): C_2$	3072	12	1
O9.240	(0; 4, 4, 4)	$\langle 64, 23 \rangle$	$(C_4 \times C_2 \times C_2): C_4$	1536	3	1
O9.241	(0; 4, 4, 4)	$\langle 64, 34 \rangle$	$((C_4 \times C_2): C_2): C_2): C_2$	768	3	1
O9.242	(0; 4, 4, 4)	$\langle 64, 35 \rangle$	$(C_4 \times C_4): C_4$	1536	6	2
O9.243	(0; 2, 8, 8)	$\langle 64, 4 \rangle$	$((C_8 \times C_2): C_2): C_2$	256	1	1
O9.244	(0; 2, 8, 8)	$\langle 64, 6 \rangle$	$(C_8 \times C_4): C_2$	256	2	1
O9.245	(0; 2, 8, 8)	$\langle 64, 10 \rangle$	$(C_8: C_4): C_2$	256	1	1
O9.246	(0; 2, 8, 8)	$\langle 64, 12 \rangle$	$(C_4: C_8): C_2$	512	1	1
O9.247	(0; 2, 8, 8)	$\langle 64, 36 \rangle$	$(C_2 \cdot ((C_4 \times C_2): C_2))$	256	1	1
O9.248	(0; 2, 4, 36)	$\langle 72, 5 \rangle$	$C_4 \times D_{18}$	216	1	1
O9.249	(0; 2, 4, 20)	$\langle 80, 14 \rangle$	$(C_{20} \times C_2): C_2$	320	1	1
O9.250	(0; 2, 2, 2, 3)	$\langle 96, 193 \rangle$	$(SL(2, 3): C_2): C_2$	1728	9	1
O9.251	(0; 2, 2, 2, 3)	$\langle 96, 227 \rangle$	$(C_2^4: C_3): C_2$	1728	3	1

Continued on next page

Table 12 – continued from previous page

Ref	Signature	SMG id	Structure	epi	autg	top
O9.252	(0; 3, 4, 4)	$\langle 96, 67 \rangle$	$SL(2, 3): C_4$	96	1	1
O9.253	(0; 3, 4, 4)	$\langle 96, 227 \rangle$	$(C_2^4: C_3): C_2$	576	1	1
O9.254	(0; 3, 3, 6)	$\langle 96, 3 \rangle$	$((C_4 \times C_2): C_4): C_3$	384	1	1
O9.255	(0; 2, 6, 6)	$\langle 96, 70 \rangle$	$(C_2^4: C_3): C_2$	384	1	1
O9.256	(0; 2, 6, 6)	$\langle 96, 202 \rangle$	$(C_2 \times SL(2, 3)): C_2$	192	1	1
O9.257	(0; 2, 4, 12)	$\langle 96, 13 \rangle$	$(C_3 \times ((C_4 \times C_2): C_2)): C_2$	192	1	1
O9.258	(0; 2, 4, 12)	$\langle 96, 186 \rangle$	$C_4 \times S_4$	96	1	1
O9.259	(0; 2, 4, 12)	$\langle 96, 187 \rangle$	$(C_2 \times S_4): C_2$	192	1	1
O9.260	(0; 2, 5, 6)	$\langle 120, 34 \rangle$	S_5	120	1	1
O9.261	(0; 2, 5, 6)	$\langle 120, 35 \rangle$	$C_2 \times A_5$	120	1	1
O9.262	(0; 2, 4, 8)	$\langle 128, 75 \rangle$	$((C_4 \times C_2 \times C_2): C_4): C_2$	512	1	1
O9.263	(0; 2, 4, 8)	$\langle 128, 134 \rangle$	$((C_4: C_8): C_2): C_2$	256	1	1
O9.264	(0; 2, 4, 8)	$\langle 128, 136 \rangle$	$((C_8: C_4): C_2): C_2$	256	1	1
O9.265	(0; 2, 4, 8)	$\langle 128, 138 \rangle$	$((C_4 \times C_4): C_4): C_2$	512	1	1
O9.266	(0; 2, 5, 5)	$\langle 160, 199 \rangle$	$((C_2 \times Q_8): C_2): C_5$	320	1	1
O9.267	(0; 2, 4, 6)	$\langle 192, 955 \rangle$	$((C_2^4: C_3): C_2): C_2$	384	1	1
O9.268	(0; 2, 4, 6)	$\langle 192, 990 \rangle$	$((SL(2, 3): C_2): C_2): C_2$	384	1	1
O9.269	(0; 2, 3, 12)	$\langle 192, 194 \rangle$	$((C_4 \times C_2): C_4): C_3): C_2$	768	1	1
O9.270	(0; 2, 4, 5)	$\langle 320, 1582 \rangle$	$((C_2 \times Q_8): C_2): C_5): C_2$	640	1	1

Email address, J. Karabáš: jan.karabas@umb.sk

Email address, R. Nedela: nedela@savbb.sk

Email address, M. Skyvová: mskyvova@ntis.zcu.cz

(J. Karabáš) DEPARTMENT OF COMPUTER SCIENCE, FACULTY OF NATURAL SCIENCES, MATEJ BEL UNIVERSITY, TAJOVSKÉHO 40, 97401 BANSKÁ BYSTRICA, SLOVAKIA

(J. Karabáš, R. Nedela) MATHEMATICAL INSTITUTE OF SLOVAK ACADEMY OF SCIENCES, ĎUMBIERSKA 1, 97411 BANSKÁ BYSTRICA, SLOVAKIA

(J. Karabáš, R. Nedela, M. Skyvová) NTIS, FACULTY OF APPLIED SCIENCES, TECHNICKÁ 8, 30601 PLZEŇ, CZECH REPUBLIC