

Quantum gravity as a communication resource

Richard Howl,^{1,2,3,*} Ali Akil,^{4,5} Hlér Kristjánsson,^{1,2,3} Xiaobin Zhao,² and Giulio Chiribella^{2,1,3,6}

¹*Quantum Group, Department of Computer Science, University of Oxford,
Wolfson Building, Parks Road, Oxford, OX1 3QD, United Kingdom*

²*QICI Quantum Information and Computation Initiative, Department of Computer Science,
The University of Hong Kong, Pokfulam Road, Hong Kong*

³*HKU-Oxford Joint Laboratory for Quantum Information and Computation*

⁴*Department of Physics, Southern University of Science and Technology, Shenzhen 518055, China*

⁵*Department of Physics, The Hong Kong University of Science and Technology, Hong Kong, China*

⁶*Perimeter Institute for Theoretical Physics, 31 Caroline Street North, Waterloo, Ontario, Canada*

(Dated: March 14, 2022)

It has been theorized that when a quantum communication protocol takes place near a black hole, the spacetime structure induced by the black hole causes an inescapable and fundamental degradation in the protocol's performance compared to if the protocol took place in flat spacetime. This is due to quantum information beyond the event horizon being inaccessible, introducing noise and degrading the entanglement resources of the protocol. However, despite black holes being a place where we expect quantum gravity to be integral, it has been assumed in these results that the black hole is a classical object with a classical spacetime. We show that when the quantum nature of a black hole and its spacetime are taken into account, their quantum properties can be used as resources to allay the degradation of entanglement caused by the event horizon, and thus improve the performance of quantum communication protocols near black holes. Investigating the resourceful nature of quantum gravity could be useful in better understanding the fundamental features of quantum gravity, just as the resourcefulness of quantum theory has revealed new insights into its foundations.

One of, if not the, most important unresolved puzzles of science is how gravity and quantum theory are unified. Many theories have been proposed for the unification, the most prominent of which, string theory and loop quantum gravity, originated several decades ago. Since then, quantum theory has gone through a “second revolution” [1], where it was understood that fundamental quantum properties, such as entanglement, can be used as resources to overcome fundamental constraints of classical physics, leading to technological improvements in areas such as computation and cryptography. This second revolution has led to a new understanding of quantum theory as a theory of information [2–9]. Is this new perception of quantum theory also important for understanding quantum gravity?

An initial approach to this question gave rise to the idea of a quantum gravity computer [10], where the expected inherent indefinite causal structure of a quantum theory of gravity could be used as resource for computation [10, 11], overcoming previously considered performance limitations of standard quantum computation where causal order is fixed a priori. This has been instrumental in the development of the theory of quantum physics with indefinite casual orders [12, 13], with new concepts, such as the quantum switch [12], where the causal order of two channels is determined by a quantum degree of freedom. This is an example of a quantum supermap, a higher order process that maps quantum channels into quantum channels [14]. Once causal order becomes indefinite, higher-order processes appear

to become as important as first-order processes, namely quantum channels, which map density matrices to density matrices [12, 13].

Here, we examine whether quantum gravity can be used as a resource for quantum communication. For this, we consider a system where quantum gravity is expected to play an integral role, a black hole, which is a place where classical general relativity appears to breakdown: it predicts a singularity that would have infinite spacetime curvature. Describing a black hole classically is known to lead to a fundamental decoherence of quantum systems and a degradation of quantum communication protocols that take place near the black hole's event horizon [15–17]. This is connected to other results on how classical gravitational fields can decohere quantum systems, leading to speculation that gravity could be responsible for our world appearing classical at macroscopic scales [15, 18–22], see also [23–25] for earlier work on gravity addressing the measurement problem of quantum theory. All these works considered a classical theory of gravity. We now consider a quantum gravity scenario and show that the quantum nature of a black hole spacetime can be used as a resource to allay the (previously considered fundamental) decoherence of quantum systems and thus also allay the degradation of nearby quantum communication protocols. In particular, we consider a black hole in a superposition of two alternative masses, which induces a superposition of channels acting on the quantum states in the vicinity of the black hole. The superposition of channels is a higher-order process transforming the existing quantum channels into a new quantum channel [26], which may act more favourably in certain quantum information protocols.

* richard.howl@cs.ox.ac.uk

Inherent decoherence from classical black holes.—Here, we review quantum communication near a classical black hole [16, 17]. This is analogous to quantum communication when one or two of the parties are accelerating in flat spacetime [16, 27, 28], which formed important early works in the field of relativistic quantum information [22, 29–31].

Consider two parties, Alice and Rob that share a scalar fermionic (Grassmann) quantum field near the event horizon of a Schwarzschild black hole. For free-falling, inertial observers, the appropriate coordinates are Minkowski. Here, the quantization of the fermionic field is straightforward, resulting in particle \hat{a}_k , \hat{a}_k^\dagger and anti-particle \hat{b}_k , \hat{b}_k^\dagger annihilation and creation operators that are associated with positive (particle) and negative (anti-particle) frequency, monochromatic plane wave mode solutions in Minkowski coordinates $u_k^\pm \propto e^{\pm i k_\mu x^\mu}$ [32], where \mathbf{k} and k_μ are the three- and four-momentum of the mode, with $\mu = 1, \dots, 4$, and we are using $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$. The Minkowski vacuum is defined by the absence of any mode excitations in an inertial frame $|0\rangle_M := \Pi_{\mathbf{k}, \mathbf{k}'} |0_{\mathbf{k}}\rangle_M^+ |0_{\mathbf{k}'}\rangle_M^-$, where $\{+, -\}$ indicate the particle and antiparticle vacua, with $\hat{a}_{\mathbf{k}} |0_{\mathbf{k}}\rangle_M^+ = 0$ and $\hat{b}_{\mathbf{k}} |0_{\mathbf{k}}\rangle_M^- = 0$.

In contrast to a free-falling observer, an observer standing just above the event horizon can consider themselves in flat spacetime but accelerating. Mathematically this is due to the Schwarzschild spacetime metric approximating a Rindler spacetime metric near the event horizon [17], with Rindler coordinates being the natural ones for accelerating observers in flat spacetime [33, 34]. Two different sets of Rindler coordinates are necessary for covering flat spacetime, which define two Rindler regions \mathcal{L} and \mathcal{R} that are causally disconnected from each other and admit a separate quantization procedure for a quantum field [35]. We can expand a scalar fermionic field in terms of Minkowski modes u_k^\pm and, independently, in terms of left and right Rindler modes $u_k^{\pm\mathcal{L}}$, $u_k^{\pm\mathcal{R}}$. The Minkowski and full set of Rindler modes are, therefore, related by a change of basis and in such a way that the vacua of the two spaces differ [35]. This is the Unruh effect, where an accelerating observer will see an inertial vacuum filled with thermal particles [36–38].

There exists an infinite number of orthonormal bases that define the Minkowski vacuum state $|0\rangle_M$. In particular, any complete set of modes made out of independent linear combinations of positive frequency Minkowski modes will define the same vacuum. This means that we can analytically continue the left and right Rindler modes to construct positive energy Minkowski modes as linear combinations of Rindler modes. The resulting modes are called Unruh modes $\nu_k^{\pm\mathcal{L}}$, $\nu_k^{\pm\mathcal{R}}$ that together span the Minkowski modes and have the vacuum $|0\rangle_M$. The Unruh modes of Minkowski space and Rindler modes of Rindler space are simply related: the change of basis is performed

by a two-mode squeezing unitary [28]:

$$\hat{U}_{\mathbf{k}}(r) = e^{r(\hat{c}_{\mathbf{k}}^{\mathcal{R}} \hat{d}_{-\mathbf{k}}^{\mathcal{L}} e^{i\phi} + \hat{c}_{\mathbf{k}}^{\mathcal{R}\dagger} \hat{d}_{-\mathbf{k}}^{\mathcal{L}\dagger} e^{-i\phi})}, \quad (1)$$

such that $\mathbf{B}_{\mathbf{k}} := \hat{U}_{\mathbf{k}} \mathbf{C}_{\mathbf{k}} \hat{U}_{\mathbf{k}}^\dagger$, where $\mathbf{B}_{\mathbf{k}} = (\hat{a}_{\mathbf{k}}, \hat{b}_{-\mathbf{k}}^\dagger)^T$, $\mathbf{C}_{\mathbf{k}} = (\hat{c}_{\mathbf{k}}^{\mathcal{R}}, \hat{d}_{-\mathbf{k}}^{\mathcal{L}\dagger})^T$, and ϕ is a phase that is often neglected on the grounds of being physically irrelevant [27][39]. Here, $\hat{a}_{\mathbf{k}}^{\mathcal{R}}$ is the right Unruh particle annihilation operator for mode \mathbf{k} , $\hat{b}_{-\mathbf{k}}^{\mathcal{R}}$ is the right Unruh anti-particle annihilation operator, $\hat{c}_{\mathbf{k}}^{\mathcal{R}}$ is the annihilation Rindler particle operator in region \mathcal{R} , $\hat{d}_{-\mathbf{k}}^{\mathcal{L}}$ is the annihilation Rindler anti-particle operator in region \mathcal{L} , and r related to the acceleration of the Rindler observer. Taking the observer to be near the event horizon of a black hole and using the equivalence of the Rindler and Schwarzschild spacetimes in this region, r is found through [16, 17]:

$$\tan r = e^{-\hbar\pi\sqrt{f_0}k^0/\kappa}, \quad (2)$$

with $f_0 := 1 - 2m/R_0$ and $\kappa := 1/(4m)$, where m is the mass of the black hole and R_0 the distance of Rob from the centre of the black hole. We are working in units $G = c = 1$, and so $2m$ is also the radius R_s of the event horizon of the black hole. The requirement that Rob is close to the event horizon is expressed by $(R_0 - R_s) \ll R_s$. There is also an analogous expression to (1) for the transformation between the left Unruh operator and Rindler operators.

The spacetime of a black hole can be represented by a Penrose diagram (see Figure 1), consisting of four regions, I, II, III and IV, with regions I and II being outside the black hole and regions III and IV representing being beyond the respective event horizons of each region. Near the event horizon, regions I and II play the role of \mathcal{R} and \mathcal{L} of Rindler space [17, 40, 41].

Now consider that Alice and Rob share a bipartite quantum state that is written as maximally entangled for observers in free fall:

$$|\psi\rangle_{AR} = (|0_{\mathbf{k}_A}\rangle_A^+ |0_{\mathbf{k}_R}\rangle_R^+ + |1_{\mathbf{k}_A}\rangle_A^+ |1_{\mathbf{k}_R}\rangle_R^+) / \sqrt{2}, \quad (3)$$

where $|0_{\mathbf{k}_A, \mathbf{k}_R}\rangle_{A,R}^+$ represents the Minkowski vacuum state in modes $\mathbf{k}_A, \mathbf{k}_R$ for Alice and Rob, $|1_{\mathbf{k}_A}\rangle_A^+ = \hat{a}_{\mathbf{k}_A}^\dagger |0\rangle_A^+$ is a one-particle Minkowski state for Alice in mode \mathbf{k}_A , and $|1_{\mathbf{k}_R}\rangle_R^+ \equiv \hat{a}_{\mathbf{k}_A}^{R\dagger} |0\rangle_R^+$ is a one-particle right Unruh state for Rob in mode \mathbf{k}_R . All other modes are assumed to be in vacua. We could have also chosen the left Unruh state $|1_{\mathbf{k}_R}\rangle_R^+$, but this would just be equivalent to stating that Rob is then in region \mathcal{L} instead of \mathcal{R} [17]. We could have also chosen an arbitrary combination of left and right Unruh modes, which is considered in [42], and taking $|1_{\mathbf{k}_R}\rangle_R^+$ to be equivalent to a one-particle Minkowski state is referred to as the single-mode approximation [27, 42]. Here, as in [17], we do not apply such an approximation as we are interested in fundamental issues rather than a realistic experiment. We, therefore, assume that Rob has access to Unruh modes. Another possibility is that Rob builds a Minkowski wavepacket involving a superposition of general Unruh modes constructed




Figure 1. Penrose diagram of a Schwarzschild black hole. Rob is stationary above the event horizon and constrained to region I. Alice is free-falling into the black hole. Rob in region I is causally disconnected from region II. Alice and Rob are assumed to share a scalar fermionic field and could be involved in a quantum teleportation protocol. For example, as considered in the main text, Alice could attempt to teleport a qubit state $|\Psi\rangle = \alpha|0_{\mathbf{k}_A}\rangle^+ + \beta|1_{\mathbf{k}_A}\rangle^+$ to Rob. In this case, although Alice's signals never reach Rob once she crosses the horizon, from Rob's perspective, if Alice is continuously sending information, for all his eternity he will receive signals from Alice. However, due to the presence of the event horizon, the fidelity of the teleportation will be degraded compared to teleportation with two inertial observers. We could have also chosen the teleportation protocol such that Rob sends his signal to Alice. However, in this case, the receiver, Alice, will eventually be destroyed by the black hole.

in such way that the single-mode approximation is valid [42]. As we are only interested in fundamental issues, we have also only assumed simplistic global modes for Alice and Rob. We do not expect the fundamental picture to change if local modes had been used instead [43].

We now consider that Alice is free-falling just above the event horizon [44], while Rob is stationary a small distance above the horizon. Alice, being a free-fall observer, will define her vacuum and excited states to be that of Minkowski space. Rob, however, will define his vacuum to be the Rindler (or equivalently Boulware) vacuum [17]. That is, Rob will apply the transformation (1) to describe his state in the Rindler basis. However, since Rob is assumed in region I of the black hole spacetime and has no access to II, we must also trace out the Rindler modes of II that form a separate Hilbert space to that of region I. The state $|\psi\rangle_{AR}$ is thus acted upon by the following quantum channel

$$\xi(\hat{\rho}_\psi) = \text{Tr}_{-\mathbf{k}_R} \left[\hat{U}_{AR}(r) (\hat{\rho}_\psi \otimes \hat{\rho}_{-\mathbf{k}_R}) \hat{U}_{AR}^\dagger(r) \right], \quad (4)$$

where $\rho_\psi := |\psi\rangle_{AR} \langle\psi|$, $\hat{U}_{AR}(r) := I_A \otimes \hat{U}_{\mathbf{k}_R}(r)$, $\hat{\rho}_{-\mathbf{k}_R} := |0_{-\mathbf{k}_R}\rangle^- \langle 0_{-\mathbf{k}_R}|$ and $\text{Tr}_{-\mathbf{k}_R}$ means that we trace over mode $-\mathbf{k}_R$ in regions I and II. This is assuming that Alice and Rob are only interested in the \mathbf{k}_A and \mathbf{k}_R modes, i.e. that their detectors are only sensitive to these modes.

The channel (4) is analogous to a fermionic Gaussian amplification channel, which can be decomposed in terms of Kraus operators: $\xi(\hat{\rho}_\psi) = \sum_n \hat{M}_n \hat{\rho}_\psi \hat{M}_n^\dagger$, with $\sum_n \hat{M}_n^\dagger \hat{M}_n = I$. There are only two such Kraus operators [45], which are presented in Appendix A.

On completion of the channel, the maximally-entangled state ρ_ψ becomes [28]:

$$\xi(\rho_\psi) = \frac{1}{2} \begin{pmatrix} \cos^2 r & 0 & 0 & \cos r \\ 0 & \sin^2 r & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \cos r & 0 & 0 & 1 \end{pmatrix}, \quad (5)$$

with the state in the Hilbert space $\mathcal{H}_A \otimes \mathcal{H}_R^I$, where the former is Alice's Hilbert space and the latter is Rob's Hilbert space over region I. Measuring the entanglement of this state using the negativity measure $\mathcal{N}[\rho] := (||\rho^{\Gamma_A}||_1 - 1)/2$, the entanglement is found to be $\mathcal{N}[\xi(\rho_\psi)] = \cos^2 r/2$ [28]. That is, the inertial maximally entangled state has become a non-maximally entangled mixed state for Rob standing next to the event horizon of a black hole. The entanglement resource is thus degraded and this will affect quantum communication between the two parties. For example, if Alice wants to teleport some state $|\Psi\rangle = \alpha|0_{\mathbf{k}_A}\rangle^+ + \beta|1_{\mathbf{k}_A}\rangle^+$ to Rob, with α and β any complex numbers that normalize the state, then the best we can expect Rob to recover at the end of the protocol is $|\Psi\rangle = \alpha|0_{\mathbf{k}_R}^I\rangle^+ + \beta|1_{\mathbf{k}_R}^I\rangle^+$, where $|0_{\mathbf{k}_R}^I\rangle^+$ and $|1_{\mathbf{k}_R}^I\rangle^+$ are the vacuum and single particle states of region I [27, 28]. If Rob is in free fall like Alice, then this protocol is perfect when using their shared maximally entangled state ρ_ψ . However, if Rob sits just above the event horizon of the black hole, and they still choose to perform the protocol using the original ρ_ψ , then the fidelity of the protocol necessarily decreases since the entanglement of ρ_ψ is degraded [17, 27, 28].

Using quantum gravity as a resource.—We have seen how the presence of a black hole can cause inherent degradation of entanglement and thus also quantum communication protocols taking place near the black hole. However, the above treatment assumed a classical description of a black hole and its spacetime. If we were instead to treat the black hole and its spacetime quantum mechanically, does this same degradation of entanglement still always occur? We will now show that the quantum structure of spacetime can be used as a resource to allay this degradation.

We assume that the black hole is now in a superposition of masses m_1 and m_2 . For example, Rob could hold a quantum machine that accesses a control qubit such that if the control is in state $|0\rangle_c$ it does nothing, while if $|1\rangle_c$ it drops a mass $m = m_2 - m_1$ into the black hole. We describe this situation in more detail in Appendix B. A superposition of masses of black holes can also occur naturally due to black hole evaporation. However, for simplicity we ignore black hole evaporation by either assuming a supermassive black hole or that the black hole is in equilibrium such that the amount of energy lost due to




Figure 2. Quantum communication protocol using a quantum black hole. The black line indicates the black hole state. Before time period t_1 , the black hole is in a single mass state $|0\rangle_m$, and the control is in the superposition state $(|0\rangle_c + |1\rangle_c)/\sqrt{2}$. The black hole then moves into an entangled state with the control: if the control is $|0\rangle_c$ the black hole remains in mass state $|0\rangle_c$, whereas if the control is $|1\rangle_c$, the black hole is placed in mass state $|1\rangle_m$. During this time, Alice and Rob share an entangled state. Rob then moves his state to a higher energy mode and accesses the control - the black hole is then subsequently always placed in the state $|1\rangle_m$ irrespective of the state of the control. Finally, Rob measures the control and then performs a quantum communication with the shared entangled state protocol. For example, he could perform quantum teleportation to teleport Alice's state.

evaporation is compensated by the energy from objects falling into the black hole.

With the black hole in a superposition of two masses, if we accept that gravity must be quantized, then this would imply that spacetime is also in a superposition and we have a superposition of spacetime event horizons. Assuming that Alice is always in free fall (she never turns on her rocket engine), and Rob is always standing above the event horizon (his rocket automatically adjusts to the gravitational field it is in), Alice and Rob's state $|\psi\rangle_{AR}$ of (3) would undergo a superposition of two channels with squeezing factors r_1 and r_2 defined by (2) with m replaced by m_1 and m_2 respectively, and become entangled with the black hole and control [46]:

$$|\Psi\rangle_{ARmc} := \frac{1}{\sqrt{2}} \left(\hat{U}_{AR}(r_1) |0\rangle_m |0\rangle_c + \hat{U}_{AR}(r_2) |1\rangle_m |1\rangle_c \right) |\psi\rangle_{AR}^0, \quad (6)$$

where $|0\rangle_m$ and $|1\rangle_m$ respectively denote the two mass states of the black hole with masses m_1 and m_2 , and $|\psi\rangle_{AR}^0 := |\psi\rangle_{AR} \otimes |0_{-\mathbf{k}_R}\rangle^-$. Here, we have ignored all other modes $\mathbf{k} \neq \mathbf{k}_A, \mathbf{k}_R$ for simplicity as they are not relevant to the protocol outlined below. Tracing over

mode $-\mathbf{k}_R$, we have:

$$\hat{\rho}_{ARmc} = \frac{1}{2} \left[\xi_{11}(\hat{\rho}_\psi) \otimes |00\rangle_{mc} {}_c\langle 00| + \xi_{22}(\hat{\rho}_\psi) \otimes |11\rangle_{mc} {}_c\langle 11| + \xi_{12}(\hat{\rho}_\psi) \otimes |00\rangle_{mc} {}_c\langle 11| + \xi_{21}(\hat{\rho}_\psi) \otimes |11\rangle_{mc} {}_c\langle 00| \right], \quad (7)$$

where

$$\xi_{ij}(\hat{\rho}_\psi) := \text{Tr}_{-\mathbf{k}_R} \left[\hat{U}_{AR}(r_i) (\hat{\rho}_\psi \otimes \hat{\rho}_{-\mathbf{k}_R}) \hat{U}_{AR}^\dagger(r_j) \right],$$

with $i, j \in \{1, 2\}$.

We could now assume that Rob is able to perform a projective measurement on the black hole and control in the basis $\{|+\rangle_m, |-\rangle_m\}$ and $\{|+\rangle_c, |-\rangle_c\}$, where $|+\rangle := (|0\rangle + |1\rangle)/\sqrt{2}$ and $|-\rangle := (|0\rangle - |1\rangle)/\sqrt{2}$. However, we do not know the true theory of quantum gravity, so it is unclear what the physical implementation of such a measurement on the black hole would be, although if the superposed masses are close in value and we assume coherent states, a parity measurement would be approximate, which is considered in Appendix C. To overcome this, we instead assume that Rob's machine shifts his state to a higher energy mode, and then accesses the control again. This time, if the control is in state $|0\rangle_c$ the machine drops a mass m into the black hole, whereas if it is in $|1\rangle_c$ the machine does nothing. The reduced state $\hat{\rho}_{ARmc}$ is then $|1\rangle_m {}_m\langle 1| \otimes \hat{\rho}_{ARc}$, where $\hat{\rho}_{ARc}$ is

$$\hat{\rho}_{ARc} = \frac{1}{2} \left[\xi_1(\hat{\rho}_\psi) \otimes |0\rangle_c {}_c\langle 0| + \xi_2(\hat{\rho}_\psi) \otimes |1\rangle_c {}_c\langle 1| + \xi_{12}(\hat{\rho}_\psi) \otimes |0\rangle_c {}_c\langle 1| + \xi_{21}(\hat{\rho}_\psi) \otimes |1\rangle_c {}_c\langle 0| \right], \quad (8)$$

such that the black hole becomes disentangled from Rob and Alice's systems. Here, Alice and Rob's reduced state remains unaffected since high-energy modes are approximately invariant to the channel (4), as $r \approx 0$. Rather than using a high-energy mode, Rob could also have moved his state to helicity modes of a bosonic field [47][48].

Rob now measures the control in the $\{+, -\}$ basis. The posteriori joint states of Alice and Rob would then be $\hat{\rho}_{AR}^+ := {}_c\langle + | \hat{\rho}_{ARc} | + \rangle_c / \text{Tr}[_c\langle + | \hat{\rho}_{ARc} | + \rangle_c]$ and $\hat{\rho}_{AR}^- := {}_c\langle - | \hat{\rho}_{ARc} | - \rangle_c / \text{Tr}[_c\langle - | \hat{\rho}_{ARc} | - \rangle_c]$, which are presented in Appendix D. It is found that $\hat{\rho}_{AR}^+$ is a separable state such that it is now possible for the entanglement between Alice and Rob's systems to completely decohere, and this is independent of the values of r_1, r_2, ϕ_1 and ϕ_2 . This is in stark contrast to the classical case considered above, where there is always entanglement between Alice and Rob's systems.

The average entanglement that Rob would obtain after measuring the control and keeping the results of his measurements is, after many measurements, $p_+ \mathcal{N}[\rho_+]$, where




Figure 3. Density plot of percentage difference between $\mathcal{N}_{Av}[\rho]$ and $\mathcal{N}[\rho_{Av}]$ for r_1 and r_2 from 0 to $\pi/4$. White through yellow and red to black indicates small to larger percentage differences.

p_+ is the probability that Rob measures $|+\rangle_c$. This is found to be

$$\mathcal{N}_{Av}[\rho] = \frac{1}{16} \left(-(\sin r_1 + \sin r_2)^2 + \sqrt{16(\cos r_1 + \cos r_2)^2 + (\sin r_1 + \sin r_2)^4} \right). \quad (9)$$

This is to be compared to the entanglement obtained if there were instead a classical mixture of black holes with masses m_1 and m_2 , which is the same as that obtained when tracing out the control:

$$\mathcal{N}[\rho_{Av}] = \frac{1}{8} \left(-(\sin^2 r_1 + \sin^2 r_2) + \sqrt{4(\cos r_1 + \cos r_2)^2 + (\sin^2 r_1 + \sin^2 r_2)^2} \right). \quad (10)$$

The average entanglement obtained from the quantum superposition of channels $\mathcal{N}_{Av}[\rho]$ is always greater than $\mathcal{N}[\rho_{Av}]$, which is illustrated in Figure 3. This occurs despite ρ_- being separable as the chance of obtaining this state is always extremely low. In Appendix E, we show that the coherent information is also always greater for the quantum superposition of channels compared to a classical mixture.

Quantum gravity can, therefore, in principle, be used as a resource to allay the degradation of entanglement brought about by a black hole spacetime. This then improves the fidelity of quantum teleportation near a black hole [27, 28] as discussed above, or other quantum communication protocols between Alice and Rob.

Discussion.—We have shown how a superposition of black hole states with different masses could be used as a

resource for quantum communication near the black hole. This performance gain comes from quantum gravity naturally setting up a superposition of quantum channels, a higher-order process which is known to generate advantages [49, 50], including in quantum communication [26, 49, 51, 52]. As seen with quantum gravity computers, it also demonstrates that higher-order processes appear to take on an important and natural role in quantum gravity.

A quantum theory of gravity comes with coherence and entanglement and so will always have these resources in comparison to a classical theory of gravity. However, in most cases these resources will be outweighed by the quantum resource of the other fundamental interactions since gravity is very weak. Here, we have considered a situation in which a classical theory of gravity appears to result in a fundamental decoherence that cannot be eradicated when quantum information is stored in a fermionic or scalar field, and shown that quantizing gravity no longer makes this level of decoherence fundamental, illustrating that quantum gravity can be used as a fundamental resource.

Rather than Rob using the superposition of black hole masses to aid in his quantum communication with Alice, he could also use it to provide evidence of the quantum nature of gravity. In a quantum theory of gravity, placing a black hole in a superposition of masses should result in a superposition of event horizons and thus a superposition of Hawking channels, which we have shown can result in less decoherence than if there were instead a classical mixture of Hawking channels. The latter would be expected in a stochastic but fundamentally classical theory of gravity, and so the level of decoherence Rob experiences in his protocol can be used to evidence the quantum nature of gravity. This is connected to very recent work on how a particle detector coupled to a black hole in a superposition of masses can exhibit signatures of quantum-gravitational effects, such as signatory peaks dependent on the mass ratio of the superposed black hole [53].

ACKNOWLEDGMENTS

This work was supported by the Hong Kong Research Grant Council through grants 17326616 and 17300918, and through the Senior Research Fellowship Scheme via SRFS2021-7S02, and by the Croucher Foundation. Research at the Perimeter Institute is supported by the Government of Canada through the Department of Innovation, Science and Economic Development Canada and by the Province of Ontario through the Ministry of Research, Innovation and Science. R.H. and G.C. also acknowledge support of Grant No. 61466 from the John Templeton Foundation as part of the QISS project. The opinions expressed in this publication are those of the authors and do not necessarily reflect the views of the John Templeton Foundation. A.A. thanks G.C. for hosting at

the University of Hong Kong. H.K. acknowledges the UK Engineering and Physical Sciences Research Council (EPSRC) through grant EP/R513295/1.

Appendix A: Kraus decomposition of classical gravity channel

The channel (4), which we refer to as the Hawking channel due to its relationship with Hawking radiation, can be written in terms of Kraus operators:

$$\xi(\hat{\rho}_\psi) = \text{Tr}_{-k_R} \left[\hat{U}_{AR}(r) (\hat{\rho}_\psi \otimes \hat{\rho}_{-k_R}) \hat{U}_{AR}^\dagger(r) \right] \quad (\text{A1})$$

$$\equiv \sum_n \hat{M}_n \hat{\rho}_\psi \hat{M}_n^\dagger, \quad (\text{A2})$$

where $\hat{M}_n := -\langle n_{-k_R} | \hat{U}_{AR} | 0_{-k_R} \rangle^- =: I \otimes \hat{N}_n$, with $\hat{N}_n := -\langle n_{-k_R} | \hat{U}_{k_R} | 0_{-k_R} \rangle^-$. Due to the Pauli exclusion principle, n can only be 0 or 1, resulting in the Kraus operators (see also [45] for a derivation using the Choi-matrix representation of a quantum channel):

$$M_0 = \begin{pmatrix} c(r) & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & c(r) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (\text{A3})$$

$$M_1 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ e^{-i\phi} s(r) & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & e^{-i\phi} s(r) & 0 \end{pmatrix}, \quad (\text{A4})$$

where $c(r) := \cos r$ and $s(r) = \sin r$. On completion of the channel, the maximally-entangled state ρ_ψ becomes:

$$\xi(\rho_\psi) = \frac{1}{2} \begin{pmatrix} c(r)^2 & 0 & 0 & c(r) \\ 0 & s(r)^2 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ c(r) & 0 & 0 & 1 \end{pmatrix}. \quad (\text{A5})$$

Measuring the entanglement of this state using the negativity measure, the entanglement is found to be $\mathcal{N}[\xi(\rho_\psi)] = c(r)^2/2$ [28].

Appendix B: A black hole in a superposition of masses

The most natural way that a black hole could be in a superposition of masses is if the original matter that collapsed to form the black hole were itself in a superposition of masses and then this superposition remains as the black hole is formed. This updates the classical no hair theorem of black holes since, in theory, the quantum phase of the superposition also characterizes the black hole.

Black hole evaporation where the black hole loses mass as it emits Hawking radiation also naturally leads to an indefinite mass for the black hole since it becomes

entangled with the emitted radiation [54]. However, here we have ignored black hole evaporation for simplicity as in previous studies on how classical black holes lead to a degradation in quantum communication [16, 17].

Another possibility is if Rob dropped an object that is in a superposition of energies or masses into the black hole. Since Rob is standing above the event horizon, according to general relativity, if he were to drop a machine that is programmed to transmit information in light signals back to Rob about the distance it has covered, then Rob would in theory never receive a signal saying that the machine has actually hit the horizon, instead he forever receives signals with greater and greater wavelengths about just how close the machine is to the horizon. This is due to the red shifting from the gravitational field. He therefore would in theory only see the machine becoming more and more smeared over the horizon and never actually dropping into it and disappearing [55]. However, that is assuming Rob has access to an unrealistic detector that can pick up any wavelength and can sit around for all of time. Instead, with a physical detector, at some point the detector will just stop seeing a signal since it cannot access all wavelengths, and then to all practical purposes Rob will think that the machine has fallen into the hole and will see the event horizon appear to grow because of this. That is, the black hole's mass will appear to have increased once he no longer receives signals from the machine, which is likely to be a very short time if he is standing very close to the event horizon. This is further illustrated by considering Rob to also have access to a gravimeter which would detect the machine getting closer and closer and becoming more and more smeared over the horizon, until, due to Gauss's law, it cannot tell the difference between the mass being smeared over the spherical horizon or being at the singularity. Therefore, dropping an object with a superposition of energies into a black hole will, to all practical purposes, place the black hole in a superposition of masses.

Appendix C: Projective measurements of a black hole

In the main text we applied a protocol such that the black hole is put into a superposition of masses using a control qubit, and then placed in a definite mass state before a projective measurement on the control in the $\{|+\rangle, |-\rangle\}$ basis. Another option would be to leave the mass of the black hole indefinite and measure it instead in the $\{|+\rangle, |-\rangle\}$ basis. This would have the advantage of Rob not needing to protect his state while he places the black hole in a definite mass state. In theory, it would also allow for the possibility of not needing a control, assuming that the black hole is pre-prepared in a superposition of masses using some different means.

Without knowing the true quantum theory of gravity, it is not possible to state whether a projective measurement in the $\{|+\rangle, |-\rangle\}$ basis exists for a quantum black

hole. However, even in standard quantum theory it is notoriously difficult to perform a projective measurement to determine whether a state is in a macroscopic superposition. Assuming that a black hole is well-described by a coherent state, then a superposition of black hole masses could be described by a superposition of coherent states: $|\alpha\rangle + |\beta\rangle$. One possible method would be to first perform the displacement operator $\hat{D}(-\alpha/2 - \beta/2)$ and then perform a parity operator measurement. If the measurement outcome is +1 then the initial state must have been $|\alpha\rangle + |\beta\rangle$. However, here the displacement operator would have to be performed very quickly since this will change the state of the black hole and thus the Hawking channel.

Assuming that the black hole can be measured in the $\{|+\rangle, |-\rangle\}$ basis, we would also need to disentangle all the modes $\mathbf{k} \neq \mathbf{k}_A, \mathbf{k}_R$ of the quantum field from \mathbf{k}_A and \mathbf{k}_R . This could be achieved by, for example, Rob initially populating all the low-energy modes. Then, due to the Pauli exclusion principle, these modes would not change when Rob stands next to the superposed event horizons of the black hole, and, we can also approximate the unpopulated high-energy modes as unchanging since the strength of the Hawking channel drops off quickly with momentum.

Appendix D: Superposition of Hawking channels

The posteriori joint states of Alice and Rob $\hat{\rho}_{AR}^+$ and $\hat{\rho}_{AR}^-$ after Rob measures the control are:

$$\hat{\rho}_{AR}^+ = \frac{\xi_{11}(\hat{\rho}_\psi) + \xi_{22}(\hat{\rho}_\psi) + \xi_{12}(\hat{\rho}_\psi) + \xi_{22}(\hat{\rho}_\psi)}{\text{Tr}[\xi_1(\hat{\rho}_\psi) + \xi_2(\hat{\rho}_\psi) + \xi_{12}(\hat{\rho}_\psi) + \xi_{22}(\hat{\rho}_\psi)]}, \quad (\text{D1})$$

$$\hat{\rho}_{AR}^- = \frac{\xi_{11}(\hat{\rho}_\psi) + \xi_{22}(\hat{\rho}_\psi) - \xi_{12}(\hat{\rho}_\psi) - \xi_{22}(\hat{\rho}_\psi)}{\text{Tr}[\xi_1(\hat{\rho}_\psi) + \xi_2(\hat{\rho}_\psi) - \xi_{12}(\hat{\rho}_\psi) - \xi_{22}(\hat{\rho}_\psi)]}, \quad (\text{D2})$$

where, using the Kraus decomposition (A1), we can write

$$\xi_{ij}(\hat{\rho}_\psi) = \sum_{n=0}^{n=1} \hat{M}_{in} \hat{\rho}_\psi \hat{M}_{jn}^\dagger, \quad (\text{D3})$$

with

$$M_{i0} = \begin{pmatrix} c(r_i) & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & c(r_i) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (\text{D4})$$

$$M_{i1} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ e^{-i\phi} s(r_i) & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & e^{-i\phi} s(r_i) & 0 \end{pmatrix}, \quad (\text{D5})$$




Figure 4. Density plot of percentage difference between $\mathcal{N}_{Av}[\rho]$ and \mathcal{N}_{Av}^C for r_1 and r_2 from 0 to $\pi/4$. White through yellow and red to black indicates small to larger percentage differences.

and $i \in \{1, 2\}$. The posterior states can then be written as:

$$\hat{\rho}_{AR}^+ = \frac{1}{2A} \begin{pmatrix} (\sum_{i=1}^{i=2} c(r_i))^2 & 0 & 0 & 2 \sum_{i=1}^{i=2} c(r_i) \\ 0 & (\sum_{i=1}^{i=2} s(r_i))^2 - 4B & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 2 \sum_{i=1}^{i=2} c(r_i) & 0 & 0 & 4 \end{pmatrix} \quad (\text{D6})$$

where:

$$A := 3 + c(r_1)c(r_2) + c(\phi_1 - \phi_2)s(r_1)s(r_2), \quad (\text{D7})$$

$$B := s((\phi_1 - \phi_2)/2)^2 s(r_1)s(r_2), \quad (\text{D8})$$

and

$$\hat{\rho}_{AR}^- = \frac{1}{2C} \begin{pmatrix} c(r_1) - c(r_2)^2 & 0 & 0 & 0 \\ 0 & (s(r_1) - s(r_2))^2 + 4B & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad (\text{D9})$$

with:

$$C := 1 - c(r_1)c(r_2) - c(\phi_1 - \phi_2)s(r_1)s(r_2). \quad (\text{D10})$$

Note that $\hat{\rho}_{AR}^-$ is a separable state $\frac{1}{2C} |0\rangle\langle 0| \otimes [(c(r_1) - c(r_2))^2 |0\rangle\langle 0| + ((s(r_1) - s(r_2))^2 + 4B) |1\rangle\langle 1|]$. The negativity of $\hat{\rho}_{AR}^+$ when $\phi_1 = \phi_2$ is

$$\mathcal{N}[\hat{\rho}_{AR}^+] = \left[-[s(r_1) + s(r_2)]^2 + \sqrt{16[c(r_1) + c(r_2)]^2 + [s(r_1) + s(r_2)]^4} \right] / 4[3 + c(r_1 - r_2)]. \quad (\text{D11})$$

The average entanglement that Rob would obtain after measuring the control and keeping the results of his measurements is $\mathcal{N}_{Av}[\hat{\rho}] = p_+ \mathcal{N}[\hat{\rho}_{AR}^+]$, where p_+ is the probability that Rob measures $|+\rangle_c$:

$$p_+ = \text{Tr}[\xi_1(\hat{\rho}_\psi) + \xi_2(\hat{\rho}_\psi) + \xi_{12}(\hat{\rho}_\psi) + \xi_{21}(\hat{\rho}_\psi)]/4 \quad (\text{D12})$$

$$= A/4. \quad (\text{D13})$$

In contrast, the probability that Rob measures $|-\rangle_c$ is:

$$p_- = \text{Tr}[\xi_1(\hat{\rho}_\psi) + \xi_2(\hat{\rho}_\psi) - \xi_{12}(\hat{\rho}_\psi) - \xi_{21}(\hat{\rho}_\psi)]/4 \quad (\text{D14})$$

$$= C/4. \quad (\text{D15})$$

Ignoring the phase as is conventional, or setting $\phi_1 = \phi_2$, the average entanglement $\mathcal{N}_{Av}[\hat{\rho}] := p_+ \mathcal{N}[\hat{\rho}_{AR}^+]$ is

$$\mathcal{N}_{Av}[\hat{\rho}] = \frac{1}{16} \left[-[s(r_1) + s(r_2)]^2 + \sqrt{16[c(r_1) + c(r_2)]^2 + [s(r_1) + s(r_2)]^4} \right]. \quad (\text{D16})$$

This is to be contrasted with the entanglement generated from an equal ‘classical’ mixture of the channels, which has the state:

$$\hat{\rho}_{Av} = \frac{1}{2} [\xi_{11}(\hat{\rho}_\psi) + \xi_{22}(\hat{\rho}_\psi)] \quad (\text{D17})$$

$$= \frac{1}{4} \begin{pmatrix} \sum_{i=1}^{i=2} c(r_i)^2 & 0 & 0 & \sum_{i=1}^{i=2} c(r_i) \\ 0 & \sum_{i=1}^{i=2} s(r_i)^2 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \sum_{i=1}^{i=2} c(r_i) & 0 & 0 & 2 \end{pmatrix}, \quad (\text{D18})$$

and negativity:

$$\mathcal{N}[\hat{\rho}_{Av}] = \frac{1}{8} \left[-[s(r_1)^2 + s(r_2)^2] + \sqrt{4[c(r_1) + c(r_2)]^2 + [s(r_1)^2 + s(r_2)^2]^2} \right]. \quad (\text{D19})$$

This is always smaller than the average entanglement obtained from the quantum superposition of channels $\mathcal{N}[\hat{\rho}_{AR}^+]$, which is illustrated in Figure 3.

Rather than using $\mathcal{N}[\hat{\rho}_{Av}]$, another possibility is to characterize the classical entanglement by

$$\mathcal{N}_{Av}^C := \frac{1}{2} \mathcal{N}[\hat{\rho}_1] + \frac{1}{2} \mathcal{N}[\hat{\rho}_2] \quad (\text{D20})$$

$$= \frac{1}{4} [c(r_1)^2 + c(r_2)^2], \quad (\text{D21})$$

where $\hat{\rho}_1$ and $\hat{\rho}_2$ are the density operators obtained for the Hawking channels with r_1 and r_2 . Since the negativity is not a linear function, $\mathcal{N}_{Av}^C \neq \mathcal{N}[\hat{\rho}_{Av}]$ even though $\hat{\rho}_{Av} = \hat{\rho}_1/2 + \hat{\rho}_2/2$. However, negativity is convex, such that $\sum_i p_i \mathcal{N}[\hat{\rho}_i] \geq \mathcal{N}[\sum_i p_i \hat{\rho}_i]$. Despite this, the entanglement from the quantum superposition of channels is still greater than the classical mixture even by this alternative measure. That is, $\mathcal{N}_{Av}[\hat{\rho}] > \mathcal{N}_{Av}^C$ always, as illustrated by Figure 4.




Figure 5. Density plot of the difference between the coherent information of a superposition of Hawking channels and a classical mixture of Hawking channels for r_1 and r_2 from 0 to $\pi/4$. White through yellow and red to black indicates small to larger differences.

Appendix E: Coherent information

The ability of Alice and Rob to communicate a qubit between each other is quantified by the quantum capacity of the communication channel between them. The quantum capacity of any channel is lower bounded by the coherent information of the channel, which in turn is lower bounded by the coherent information of the output state resulting from the action of the channel on one half of the maximally entangled state. In the scenario considered in the main text, the effective channel described by (8) (arising from the superposition of two fermionic Gaussian amplification channels with squeezing factors r_1, r_2 , respectively) can be thought of as a communication channel between Alice and Rob, where half of the maximally entangled state is sent from Alice to Rob, with Alice keeping the other half herself. Therefore, the coherent information of the output state of (8) gives a lower bound to the quantum capacity of the communication channel available between Alice and Rob.

Similarly, one can calculate the coherent information of the output state arising from a classical mixture of two fermionic Gaussian amplification channels with squeezing factors r_1, r_2 . The difference in the coherent information between the superposition case and the classical mixture case can be used to quantify the effect of a superposition of black hole masses on quantum communication.

The coherent information of a bipartite quantum state $\hat{\rho}_{AB}$ is given by $I(A, B)_\rho = S(B)_\rho - S(AB)_\rho$, where $S(AB)_\rho$ is the von Neumann entropy of $\hat{\rho}_{AB}$ and $S(B)_\rho$ is the von Neumann entropy of the reduced state $\hat{\rho}_B := \text{Tr}_A \hat{\rho}_{AB}$.

Figure 5 shows a density plot of the difference in the

coherent information between the output state of the effective channel (8), arising from a superposition of two channels, and the corresponding output state for a classical mixture of the two channels, for r_1 and r_2 between 0 and $\pi/4$. We find that this difference is always positive, showing that the superposition of channels increases the ability to communicate quantum information between the two relativistic agents.

Appendix F: Superposition of phase channels

Above we considered a superposition of Hawking channels where there are different values of squeezing parameter r but the same phase ϕ . Usually the phase is considered to be physically irrelevant but with a quantum theory of spacetime, this may no longer be the case. Therefore, here we consider a superposition of the Hawking channels that have the same squeezing parameter but different phases. Since the phase is irrelevant in the classical mixture of channels, this also allows us to compare the resource of a superposition of channels to that of just one channel.

Taking a superposition of channels with the same squeezing parameter r but opposite phases $\phi_1 = \phi_2 + \pi$, and assuming a similar protocol to that assumed for a superposition of squeezing parameters, the posterior states after measuring the control in a $\{|+\rangle_c, |-\rangle_c\}$ basis are

$$\hat{\rho}_{AR}^+ := \frac{1}{1 + \cos^2 r} \begin{pmatrix} \cos^2 r & 0 & 0 & \cos r \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \cos r & 0 & 0 & 1 \end{pmatrix}, \quad (\text{F1})$$

and

$$\hat{\rho}_{AR}^- = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad (\text{F2})$$

$$= |10\rangle_{AR} {}_{AR}\langle 10|. \quad (\text{F3})$$

The state $\hat{\rho}_{AR}^+$ has entanglement:

$$\mathcal{N}[\hat{\rho}_{AR}^+] = |\cos r|/(1 + \cos^2 r). \quad (\text{F4})$$

Since we obtain zero entanglement when Rob obtains $|-\rangle_c$, the average entanglement is $p_+ \mathcal{N}[\hat{\rho}_{AR}^+]$, where p_+ is the same as (D13), which in this case is $p_+ = (1 + \cos^2 r)/2$ since $r := r_1 = r_2$. The average entanglement is then:

$$\mathcal{N}_{Av}[\hat{\rho}] = |\cos r|/2. \quad (\text{F5})$$

This is to be compared to the entanglement obtained with the classical gravity case of a classical mixture of the phase channels, which are equivalent, and therefore there is just one channel for which the entanglement is $\cos^2 r/2$. The entanglement for a superposition of channels is clearly always greater than the entanglement from the classical case.




Figure 6. The blue solid line is the average Negativity of the final state of a superposition of channels with opposite phases $\mathcal{N}_{Av}[\rho]$ as a function of squeezing parameter r , while, for comparison, the red dashed line is the negativity of a single channel $\mathcal{N}[\rho] = \cos^2 r/2$ with fixed phase.

However, it is not clear how a superposition of phases could be achieved and what the physical meaning of this would be. Usually the phase of the squeezing operation is neglected and considered to be unphysical [15, 28]. One speculative possibility is that Rob could send a quantum object that has a superposition of phases into a black hole, for example the cat state $|\psi\rangle_m = (|\alpha\rangle + |\alpha e^{-i\pi}\rangle)/\sqrt{2 + 2\exp(-2\alpha^2)}$, where $|\alpha\rangle$ is a coherent state. Another option might be to consider the quantum spin of a black hole.

Appendix G: Global mode assumption

In the quantum communication protocol between Alice and Rob, we only considered global modes and have not considered the physical implementation of Alice and Rob's detectors. In studies of how the classical gravitational field of a black hole and the acceleration of observers causes quantum decoherence, the detectors are generally considered to be Unruh-DeWitt detectors [56], which are pointlike detectors with two internal energy levels that are weakly coupled to a quantum matter field through a monopole interaction. Recent works have considered Unruh-DeWitt detectors in a superposition of different accelerated trajectories in Minkowski spacetime [57–61], finding, for example, that the state of the detector excitations is not in general a convex mixture of the thermal spectrum expected for a classical mixture of accelerations. Most recently, a single particle detector in the presence of a black hole in a superposition of masses has been considered, with similar results [53]. This is related to the work discussed here. However, in the protocol used in this work, at the end of the protocol the black hole is brought back to a definite mass state before a detector would be used. There is, therefore, no need to consider the complications induced by a superposition of Unruh-DeWitt detectors in the protocol considered in the main text.

Localized wave-packets rather than global modes have

been considered for investigations of entanglement degradation near classical black holes, with the same conclusions as in the global case [17]. We, therefore, expect that utilizing localized wave-packets instead of global modes would not entirely negate the alleviation of degradation facilitated by quantum gravity.

Appendix H: Protecting Rob's state

In the protocol in the main text Rob must protect his state while his quantum machine accesses the control to determine whether a mass should be dropped into the black hole or not. This is done so that the black hole becomes disentangled from Rob's state, and he then only needs to measure the control. The way this is achieved in the main text is to move Rob's state to a high energy mode, e.g. using a 90° beam splitter, since the strength of the Hawking channel drops off quickly with momentum. However, there are other ways by which Rob could protect his state. For example, Rob could, in theory, store his state in helicity modes of a spin-1 field since these modes are unaffected by the Hawking channel [62].

Appendix I: A black hole in a superposition of locations

In aiding quantum communication between Alice and Rob, we could have used a superposition of locations of the black hole rather than black hole masses, such that Rob is at a different distance from the event horizon in both cases, although always still close to the event horizon. Exactly the same result as above would be obtained with just r_1 and r_2 following (2) but with different values of R_0 . This illustrates that the same gain would be obtained also if Rob's detectors were placed in a superposition of accelerations, without reference to quantum gravity. This, however, does not diminish the fact that quantum gravity can be used, in principle, as a resource to allay decoherence expected with classical black holes and their classical gravitational fields [16, 17].

There are also attempts to keep gravity classical even when the matter source is described quantum mechanically. In this case, when a black hole is put in a superposition of masses its gravitational field is not in a superposition of states as it remains a classical entity. For example, in fundamental semi-classical gravity, the gravitational field would be that generated by a black hole with a mass which is the expectation of the superposed mass state [63–66]. We do not consider this possibility here.

A black hole that is in a superposition of macroscopic locations has been considered several times before although not in the context of Hawking channels and as a resource for quantum communication (see e.g. [67, 68]). Possible ways for the black hole to be in a superposition of locations include the collapsing matter being in

a location of two positions, and Rob possibly firing an object with a superposition of momenta into the black hole such that the back-reaction of the black hole also places it in a superposition of two momenta. Perhaps then Rob could fire another object with a superposition of momenta to make the black hole stationary in both superposition channels but now with distinct locations.

-
- [1] A. G. J. MacFarlane, J. P. Dowling, and G. J. Milburn, *Philosophical Transactions of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences* **361**, 1655 (2003).
- [2] L. Hardy, arXiv e-prints, quant-ph/0101012 (2001), arXiv:quant-ph/0101012 [quant-ph].
- [3] L. Hardy, (2011), arXiv:1104.2066 [quant-ph].
- [4] L. Hardy, “Operational structures as a foundation for probabilistic theories,” (2009).
- [5] G. Chiribella, G. M. D’Ariano, and P. Perinotti, *Phys. Rev. A* **81**, 062348 (2010).
- [6] Borivoje Dakic and Časlav Brukner, in *Deep Beauty: Understanding the Quantum World Through Mathematical Innovation* (Cambridge University Press).
- [7] L. Masanes and M. P. Müller, *New Journal of Physics* **13**, 063001 (2011).
- [8] J. Barrett, arXiv preprint quant-ph/0508211 (2005).
- [9] G. Chiribella, G. M. D’Ariano, and P. Perinotti, *Phys. Rev. A* **84**, 012311 (2011).
- [10] L. Hardy, “Quantum gravity computers: On the theory of computation with indefinite causal structure,” in *Quantum Reality, Relativistic Causality, and Closing the Epistemic Circle: Essays in Honour of Abner Shimony* (Springer Netherlands, Dordrecht, 2009) pp. 379–401.
- [11] M. Araújo, F. Costa, and i. c. v. Brukner, *Phys. Rev. Lett.* **113**, 250402 (2014).
- [12] G. Chiribella, G. M. D’Ariano, P. Perinotti, and B. Valiron, *Phys. Rev. A* **88**, 022318 (2013), arXiv:0912.0195.
- [13] O. Oreshkov, F. Costa, and Č. Brukner, *Nature communications* **3**, 1 (2012), arXiv:1105.4464.
- [14] G. Chiribella, G. M. D’Ariano, and P. Perinotti, *EPL (Europhysics Letters)* **83**, 30004 (2008).
- [15] M. P. Blencowe, *Phys. Rev. Lett.* **111**, 021302 (2013).
- [16] I. Fuentes-Schuller and R. B. Mann, *Phys. Rev. Lett.* **95**, 120404 (2005).
- [17] E. Martín-Martínez, L. J. Garay, and J. León, *Phys. Rev. D* **82**, 064006 (2010).
- [18] T. Oniga and C. H.-T. Wang, *Phys. Rev. D* **93**, 044027 (2016).
- [19] C. Anastopoulos and B. L. Hu, *Classical and Quantum Gravity* **30**, 165007 (2013).
- [20] I. Pikovski, M. Zych, F. Costa, and Č. Brukner, *Nature Physics* **11**, 668 (2015).
- [21] A. Bassi, A. Großardt, and H. Ulbricht, *Classical and Quantum Gravity* **34**, 193002 (2017).
- [22] R. Howl, L. Hackermüller, D. E. Bruschi, and I. Fuentes, *Advances in Physics: X* **3**, 1383184 (2018), <https://doi.org/10.1080/23746149.2017.1383184>.
- [23] L. Diósi, *Physics Letters A* **120**, 377 (1987).
- [24] R. Penrose, *General relativity and gravitation* **28**, 581 (1996).
- [25] R. Howl, R. Penrose, and I. Fuentes, *New Journal of Physics* **21**, 043047 (2019).
- [26] G. Chiribella and H. Kristjánsson, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* **475**, 20180903 (2019).
- [27] P. M. Alsing and G. J. Milburn, *Phys. Rev. Lett.* **91**, 180404 (2003).
- [28] P. M. Alsing, I. Fuentes-Schuller, R. B. Mann, and T. E. Tessier, *Phys. Rev. A* **74**, 032326 (2006).
- [29] A. Peres and D. R. Terno, *Rev. Mod. Phys.* **76**, 93 (2004).
- [30] T. C. Ralph and T. G. Downes, *Contemporary Physics* **53**, 1 (2012), <https://doi.org/10.1080/00107514.2011.640146>.
- [31] P. M. Alsing and I. Fuentes, *Classical and Quantum Gravity* **29**, 224001 (2012).
- [32] F. Mandl and G. Shaw, *Quantum field theory* (Wiley, 1984).
- [33] W. Rindler, *Phys. Rev.* **119**, 2082 (1960).
- [34] W. Rindler, *American Journal of Physics* **34**, 1174 (1966).
- [35] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, 1982).
- [36] S. A. Fulling, *Phys. Rev. D* **7**, 2850 (1973).
- [37] P. C. W. Davies, *Journal of Physics A: Mathematical and General* **8**, 609 (1975).
- [38] W. G. Unruh, *Phys. Rev. D* **14**, 870 (1976).
- [39] It is possible that with a quantum theory of spacetime curvature, there is a physical interpretation of the phase. In Appendix F, we show that, if it were physical, the phase can be used to allay degradation of entanglement.
- [40] K. S. Thorne, C. W. Misner, and J. A. Wheeler, *Gravitation* (Freeman, 2000).
- [41] D. Tong, .
- [42] D. E. Bruschi, J. Louko, E. Martín-Martínez, A. Dragan, and I. Fuentes, *Phys. Rev. A* **82**, 042332 (2010).
- [43] See Appendix G.
- [44] We follow [17] in considering Alice to be free falling just above the event horizon, and assume that she does not experience Hawking radiation. Alternatively, we could also consider Alice to be free falling but very far from the horizon, in which case the Hawking radiation she would experience would be negligible.
- [45] S. Banerjee, A. K. Alok, S. Omkar, and R. Srikanth, *Journal of High Energy Physics* **2017**, 1 (2017).
- [46] Here, for simplicity, we have chosen the relative phase between the two unitaries $\hat{U}_{AR}(r_1)$ and $\hat{U}_{AR}(r_2)$ to be zero.
- [47] See Appendix H.
- [48] Of course, if Rob had chosen to use high-energy modes in the first place then any quantum communication would not be as degraded. However, by not doing this he is able to demonstrate the resourcefulness of quantum gravity. Alternatively, he could have measured the black hole in the $\{|+\rangle, |-\rangle\}$ basis, in which case Rob does not even need his quantum machine and the control if he already knows that the black hole is in a superposition of masses. In this case there is no need for Rob to then protect his state by using a higher-energy mode, or other modes of a different field, or extremal black holes. This is discussed further in Appendix C where it is found that the other modes of the field become relevant now.
- [49] N. Gisin, N. Linden, S. Massar, and S. Popescu, *Phys. Rev. A* **72**, 012338 (2005).
- [50] M. Araújo, A. Feix, F. Costa, and Č. Brukner, *New Journal of Physics* **16**, 093026 (2014).
- [51] A. A. Abbott, J. Wechs, D. Horsman, M. Mhalla, and C. Branciard, *Quantum* **4**, 333 (2020).
- [52] H. Kristjánsson, G. Chiribella, S. Salek, D. Ebler, and M. Wilson, *New Journal of Physics* **22**, 073014 (2020).
- [53] J. Foo, C. S. Arabaci, M. Zych, and R. B. Mann, (2021),

- arXiv:2111.13315 [gr-qc].
- [54] A. Akil, O. Dahlsten, and L. Modesto, *New Journal of Physics* **23**, 113011 (2021).
 - [55] L. Susskind and J. Lindesay, *An Introduction to Black Holes, Information and the String Theory Revolution* (WORLD SCIENTIFIC, 2004) <https://www.worldscientific.com/doi/pdf/10.1142/5689>.
 - [56] B. DeWitt, *General Relativity: an Einstein Centenary Survey* (Cambridge University Press, 1980).
 - [57] L. C. Barbado, E. Castro-Ruiz, L. Apadula, and i. c. v. Brukner, *Phys. Rev. D* **102**, 045002 (2020).
 - [58] R. Parentani, *Nuclear Physics B* **454**, 227 (1995).
 - [59] E. Martín-Martínez, I. Fuentes, and R. B. Mann, *Phys. Rev. Lett.* **107**, 131301 (2011).
 - [60] N. Stritzelberger and A. Kempf, *Phys. Rev. D* **101**, 036007 (2020).
 - [61] J. Foo, S. Onoe, and M. Zych, *Phys. Rev. D* **102**, 085013 (2020).
 - [62] Y. Ling, S. He, W. Qiu, and H. Zhang, *Journal of Physics A: Mathematical and Theoretical* **40**, 9025 (2007).
 - [63] L. Rosenfeld, *Nuclear Physics* **40**, 353 (1963).
 - [64] C. Möller, *Proceedings, Les théories relativistes de la gravitation: actes du colloque international (Relativistic Theories of Gravitation): Royaumont, France, June 21-27, 1959*, *Colloques Internationaux CNRS* **91**, 15 (1962).
 - [65] T. Kibble, *Communications in Mathematical Physics* **64**, 73 (1978).
 - [66] T. W. B. Kibble and S. Randjbar-Daemi, *Journal of Physics A: Mathematical and General* **13**, 141 (1980).
 - [67] Dimić, Aleksandra and Milivojević, Marko and Gočanin, Dragoljub and Möller, Natália S. and Brukner, Časlav, *Frontiers in Physics* **8**, 470 (2020).
 - [68] A. Arrasmith, A. Albrecht, and W. H. Zurek, *Nature communications* **10**, 1 (2019).