

WEAK DIAMOND, WEAK PROJECTIVITY, AND TRANSFINITE EXTENSIONS OF SIMPLE ARTINIAN RINGS

JAN TRLIFAJ

ABSTRACT. We apply set-theoretic methods to study projective modules and their generalizations over transfinite extensions of simple artinian rings R . We prove that if R is small, then the Weak Diamond implies that projectivity of an arbitrary module can be tested at the layer epimorphisms of R .

The classic Baer's Criterion, saying that a module M is injective, iff it is R -injective, is a basic tool of the structure theory of injective modules over an arbitrary ring R . However, unless R is a perfect ring, there are no criteria available for the dual case, that is, for testing projectivity using a set of epimorphisms, [10].

For many non-perfect rings, it can be proved that there exist small (e.g., countably generated) non-projective R -projective modules (e.g., when R is commutative noetherian of Krull dimension ≥ 1). However, for each cardinal κ , there exists a non-right perfect ring R_κ such that all $\leq \kappa$ -generated R_κ -projective modules are projective, [12]. This is the best one can achieve in ZFC, because it is consistent with ZFC + GCH that if R is not right perfect, then there always exist (large) R -projective modules that are not projective, cf. [1].

Consistency (and hence independence) of the coincidence of R -projectivity and projectivity for certain commutative non-noetherian rings was proved in [11]. This answered in the positive a question from [1, 2.8], and clarified the set-theoretic status of an old problem by Carl Faith [6, p.175]. The consistency result was extended in [12] to further classes of rings that are finite Loewy length extensions of simple artinian rings. The set theoretic tool used in [11] and [12] was Jensen's Diamond.

The goal of the present paper is twofold: to enhance the algebraic tools to cover infinite Loewy length extensions of simple artinian rings, and to weaken the set-theoretic assumptions used in the proofs. In Theorem 3.2 below, we show that the Weak Diamond Principle Φ and CH are sufficient to prove coincidence of the classes of all weak R -projective, R -projective, and projective modules, in the case when R has cardinality at most \aleph_1 , its Loewy length is countable, and each proper layer of R is countably generated. That is, in this case, Φ and CH imply that the projectivity of a module M is equivalent to the factorization of all morphisms from M with finitely generated images through the layer epimorphisms of R . The latter are just the canonical projections $\pi_\alpha : S_{\alpha+1} \rightarrow S_{\alpha+1}/S_\alpha$ ($\alpha < \sigma$), where $(S_\alpha \mid \alpha \leq \sigma + 1)$ is the socle sequence of R .

For basic notions and facts needed from ring and module theory, we refer to [2] and [8]; our references for set-theoretic homological algebra are [4] and [7].

Date: March 14, 2022.

2010 Mathematics Subject Classification. Primary: 16D40, 03E35. Secondary: 16E50, 16D60, 16D70, 03E45, 18G05.

Key words and phrases. weakly projective module, Weak Diamond Principle, von Neumann regular ring, semiartinian ring.

Research supported by GAČR 20-13778S.

1. WEAK PROJECTIVITY

We start by recalling the classic notion of relative projectivity from [2, §16]:

Definition 1.1. Let R be a ring and M, N be modules. Then M is N -projective provided that for each submodule P of N , each homomorphism $f : M \rightarrow N/P$ has a factorization through the canonical projection $N \rightarrow N/P$. That is, the functor $\text{Hom}_R(M, -)$ is exact on all short exact sequences whose middle term is N .

The following Lemma is well-known (see [2, 16.12 and 16.14]):

Lemma 1.2. *The class of all modules N such that M is N -projective is closed under submodules, homomorphic images, and finite direct sums.*

In particular, each finitely generated R -projective module is projective.

Let R be a right semiartinian ring with the right socle sequence $(S_\alpha \mid \alpha \leq \sigma + 1)$. We will call $\sigma + 1$ the *Loewy length* of R . To avoid trivialities, we will tacitly assume that $\sigma > 0$, that is, that R is not completely reducible.

For each $\alpha \leq \sigma$, we will call the completely reducible module $L_\alpha = S_{\alpha+1}/S_\alpha$ the α th layer of R . The layers L_α for $0 < \alpha < \sigma$ are called *proper*. For each $\alpha \leq \sigma$, the canonical epimorphism $\pi_\alpha : S_{\alpha+1} \rightarrow L_\alpha$ is the α th layer epimorphism of R .

In this setting, the following definition of weak R -projectivity was introduced in [12, 3.6]

Definition 1.3. A module M is called *weakly R -projective* provided that for each $0 < \alpha \leq \sigma$, each $f \in \text{Hom}_R(M, L_\alpha)$ with a finitely generated image has a factorization through the α th layer epimorphism π_α .

Note that by Lemma 1.2, each R -projective module is weakly R -projective. The following easy observation [12, 2.3] shows that weak R -projectivity can equivalently be stated in a stronger form:

Lemma 1.4. *Let R be a right semiartinian ring with the right socle sequence $(S_\alpha \mid \alpha \leq \sigma + 1)$. Let M be a module. Then the following are equivalent:*

- (1) M is weakly R -projective,
- (2) For each $\alpha \leq \sigma$, each submodule K such that $S_\alpha \subseteq K \subseteq S_{\alpha+1}$ and each $f \in \text{Hom}_R(M, S_{\alpha+1}/K)$ with a finitely generated image, there exists $g \in \text{Hom}_R(M, S_{\alpha+1})$ such that $f = \pi_K g$, where $\pi_K : S_{\alpha+1} \rightarrow S_{\alpha+1}/K$ is the canonical projection.

In the proof of the next lemma, we will verify N -projectivity of a finitely generated module M by a recursive procedure using a filtration of N .

Here, we call a chain $(N_\alpha \mid \alpha \leq \tau)$ of submodules of a module N a *filtration* of N , provided that τ is an ordinal, $N_0 = 0$, $N_\alpha \subseteq N_{\alpha+1}$ for each $\alpha < \tau$, $N_\alpha = \bigcup_{\beta < \alpha} N_\beta$ for each limit ordinal $\alpha \leq \tau$, and $N_\tau = N$.

Lemma 1.5. *Let R be a ring and N be a module with a filtration $\mathcal{N} = (N_\alpha \mid \alpha \leq \tau)$.*

Let M be a finitely generated module such that for each $\alpha < \tau$, each submodule K such that $N_\alpha \subseteq K \subseteq N_{\alpha+1}$ and each $f \in \text{Hom}_R(M, N_{\alpha+1}/K)$, there exists $g \in \text{Hom}_R(M, N_{\alpha+1})$ such that $f = \pi_K g$, where $\pi_K : N_{\alpha+1} \rightarrow N_{\alpha+1}/K$ is the canonical projection.

Then M is N -projective.

Proof. We claim that M is N_α -projective for each $\alpha \leq \tau$. If so, then for $\alpha = \tau$, we get that M is N -projective.

The claim will be proved by induction on α . There is nothing to prove for $\alpha = 0$.

Assume that the claim is true for some $\alpha < \tau$. Let K be a submodule of $N_{\alpha+1}$. Let $\pi_K : N_{\alpha+1} \rightarrow N_{\alpha+1}/K$, $\rho : N_{\alpha+1} \rightarrow N_{\alpha+1}/(N_\alpha + K)$, $\eta : N_{\alpha+1}/K \rightarrow$

$N_{\alpha+1}/(N_\alpha + K)$, and $\theta : N_\alpha \rightarrow N_\alpha/(N_\alpha \cap K)$ denote the canonical projections. Also, let $\iota_\alpha : N_\alpha/(N_\alpha \cap K) \rightarrow (N_\alpha + K)/K$ be the canonical isomorphism (given by the assignment $\iota_\alpha(x + N_\alpha \cap K) = x + K$).

These homomorphisms fit in the following commutative diagram with exact rows (where, except for the zero maps, all the unnamed single arrows are inclusions):

$$\begin{array}{ccccccccc}
0 & \longrightarrow & N_\alpha + K & \longrightarrow & N_{\alpha+1} & \xrightarrow{\rho} & N_{\alpha+1}/(N_\alpha + K) & \longrightarrow & 0 \\
& & \uparrow & & \parallel & & \eta \uparrow & & \\
0 & \longrightarrow & K & \longrightarrow & N_{\alpha+1} & \xrightarrow{\pi_K} & N_{\alpha+1}/K & \longrightarrow & 0 \\
& & \parallel & & \uparrow & & \uparrow & & \\
0 & \longrightarrow & K & \longrightarrow & N_\alpha + K & \xrightarrow{\pi_K \uparrow} & (N_\alpha + K)/K & \longrightarrow & 0 \\
& & \uparrow & & \uparrow & & \iota_\alpha \uparrow & & \\
0 & \longrightarrow & N_\alpha \cap K & \longrightarrow & N_\alpha & \xrightarrow{\theta} & N_\alpha/(N_\alpha \cap K) & \longrightarrow & 0
\end{array}$$

Let $f \in \text{Hom}_R(M, N_{\alpha+1}/K)$. We have to show that f factorizes through π_K .

By our assumption on M , there exists $g \in \text{Hom}_R(M, N_{\alpha+1})$ such that $\rho g = \eta f$. Since $\rho = \eta \pi_K$, $\eta(f - \pi_K g) = 0$. It follows that $\delta = f - \pi_K g$ maps M into $\ker(\eta) = (N_\alpha + K)/K$, whence $\iota_\alpha^{-1} \delta \in \text{Hom}_R(M, N_\alpha/(N_\alpha \cap K))$. By the inductive premise, there exists $\epsilon : M \rightarrow N_\alpha$ such that $\iota_\alpha \theta \epsilon = \delta$. As $\iota_\alpha \theta = \pi_K \uparrow (N_\alpha + K)$, we conclude that $f = \pi_K g + \delta = \pi_K g + \pi_K \epsilon = \pi_K(g + \epsilon)$, which is the desired factorization of f through π_K .

Let $\alpha \leq \tau$ be a limit ordinal. Let K be a submodule of $N_\alpha = \bigcup_{\beta < \alpha} N_\beta$ and $f \in \text{Hom}_R(M, N_\alpha/K)$. Since M is finitely generated, there exists $\beta < \alpha$ such that $f : M \rightarrow (N_\beta + K)/K$. Let $\iota_\beta : N_\beta/(N_\beta \cap K) \rightarrow (N_\beta + K)/K$ be the canonical isomorphism, and $\theta : N_\beta \rightarrow N_\beta/(N_\beta \cap K)$ the canonical projection. Then $\pi_K \uparrow (N_\beta + K) = \iota_\beta \theta$, and we have the following commutative diagram with exact rows (again, except for the zero maps, all the unnamed single arrows are inclusions):

$$\begin{array}{ccccccccc}
0 & \longrightarrow & K & \longrightarrow & N_\alpha & \xrightarrow{\pi_K} & N_\alpha/K & \longrightarrow & 0 \\
& & \parallel & & \uparrow & & \uparrow & & \\
0 & \longrightarrow & K & \longrightarrow & N_\beta + K & \xrightarrow{\pi_K \uparrow} & (N_\beta + K)/K & \longrightarrow & 0 \\
& & \uparrow & & \uparrow & & \iota_\beta \uparrow & & \\
0 & \longrightarrow & N_\beta \cap K & \longrightarrow & N_\beta & \xrightarrow{\theta} & N_\beta/(N_\beta \cap K) & \longrightarrow & 0
\end{array}$$

By the inductive premise for β , there exists $g \in \text{Hom}_R(M, N_\beta)$ such that $\theta g = \iota_\beta^{-1} f$. Then $f = \iota_\beta \theta g = \pi_K g$, which yields the desired factorization of f through π_K . \square

The results above imply that weak R -projectivity is sufficient to guarantee projectivity for each finitely generated module:

Corollary 1.6. *Let R be any right semiartinian ring. Let M be a finitely generated weakly R -projective module. Then M is projective.*

Proof. Since weak projectivity can be expressed in the stronger form of Lemma 1.4, we can apply Lemma 1.5 for $N = R$ and \mathcal{N} = the right socle sequence of R . Thus, M is R -projective, and by Lemma 1.2, M is projective. \square

The question of the coincidence of weak R -projectivity, R -projectivity, and projectivity for infinitely generated modules over semiartinian rings, that is, whether projectivity can be tested at the layer epimorphisms of R , is much more delicate. We will see that for some semiartinian rings, it is actually independent of ZFC + GCH.

On the one hand, there is the following set-theoretic barrier for all non-right perfect rings R , given by Shelah's Uniformization Principle (SUP). This principle is consistent with ZFC + GCH (see [4, XIII.1.5] or [5, §2] for more details):

Lemma 1.7. *Assume SUP. Let R be a non-right perfect ring. Let κ be a singular cardinal of cofinality ω such that $\text{card}(R) < \kappa$. Then there exists a κ^+ generated module M of projective dimension equal to 1, such that $\text{Ext}_R^1(M, I) = 0$ for each right ideal I of R . In particular, M is R -projective, but not projective.*

Proof. See [1, 2.4 and 2.5]. □

On the other hand, there are different extensions of ZFC + GCH where weak R -projectivity, R -projectivity, and projectivity coincide for suitable semiartinian rings. These rings will be studied in the next section.

2. TRANSFINITE EXTENSIONS OF SIMPLE ARTINIAN RINGS

Recall that a ring R is *von Neumann regular*, if for each $r \in R$ there exists $s \in R$ such that $rsr = r$. We refer to [8] for properties of von Neumann regular rings. A ring R is said to have (right) *primitive factors artinian*, or (right) *pfa* for short, if R/P is (right) artinian for each (right) primitive ideal of R .

If R is von Neumann regular, then minimal right (left) ideals of R correspond to primitive idempotents of R , so the right and left socle sequences of R coincide. In particular, R is right semiartinian, iff R is left semiartinian. Similarly, by [8, 6.2], a von Neumann regular ring has right pfa, iff it has left pfa. The latter condition can equivalently be stated as a property of the module category $\text{Mod-}R$: each homogenous completely reducible module is injective, see [8, 6.28].

Let R be a semiartinian von Neumann regular ring with pfa, and $(S_\alpha \mid \alpha \leq \sigma + 1)$ be its socle sequence. (Notice that $\sigma > 0$ implies that R is not right perfect, because the Jacobson radical of any von Neumann regular ring is 0; thus, Lemma 1.7 applies here.)

By the following theorem from [9], semiartinian von Neumann regular rings with pfa can be viewed as transfinite extensions of full matrix rings over skew-fields (i.e., of simple artinian rings):

Theorem 2.1. *Let R be a right semiartinian ring and $(S_\alpha \mid \alpha \leq \sigma + 1)$ be the right socle sequence of R . The following conditions are equivalent:*

- (1) R is von Neumann regular with pfa.
- (2) for each $\alpha \leq \sigma$ there are a cardinal λ_α , positive integers $n_{\alpha\beta}$ ($\beta < \lambda_\alpha$) and skew-fields $K_{\alpha\beta}$ ($\beta < \lambda_\alpha$), such that $L_\alpha \cong \bigoplus_{\beta < \lambda_\alpha} M_{n_{\alpha\beta}}(K_{\alpha\beta})$, as rings without unit. Moreover, λ_α is infinite iff $\alpha < \sigma$.

The pre-image of $M_{n_{\alpha\beta}}(K_{\alpha\beta})$ in the isomorphism φ_α coincides with the β th homogenous component of $\text{Soc}(R/S_\alpha)$, and it is a finitely generated as right R/S_α -module for all $\beta < \lambda_\alpha$.

The structure of the rings characterized by Theorem 2.1 can be depicted as follows:

L_σ	$M_{n_{\sigma 0}}(\mathbf{K}_{\sigma 0}) \oplus \dots \oplus M_{n_{\sigma, \lambda_\sigma - 1}}(\mathbf{K}_{\sigma, \lambda_\sigma - 1})$	
...	...	
L_α	$M_{n_{\alpha 0}}(\mathbf{K}_{\alpha 0}) \oplus \dots \oplus M_{n_{\alpha \beta}}(\mathbf{K}_{\alpha \beta}) \oplus \dots$	$\beta < \lambda_\alpha$
...	...	
L_1	$M_{n_{10}}(\mathbf{K}_{10}) \oplus \dots \oplus M_{n_{1\beta}}(\mathbf{K}_{1\beta}) \oplus \dots$	$\beta < \lambda_1$
$L_0 = \text{Soc}(R)$	$M_{n_{00}}(\mathbf{K}_{00}) \oplus \dots \oplus M_{n_{0\beta}}(\mathbf{K}_{0\beta}) \oplus \dots$	$\beta < \lambda_0$

Here, $\sigma + 1$ is the Loewy length of R . The rows in this picture represent the layers of R ; λ_α is the number of homogenous components in the α th layer $L_\alpha = S_{\alpha+1}/S_\alpha$ for each $\alpha \leq \sigma$ (this number is infinite except for $\alpha = \sigma$).

$n_{\alpha\beta}$ is the (finite) rank of the β th homogenous component of L_α , and $K_{\alpha\beta}$ is the endomorphism ring (skew-field) of each simple module in the β th homogenous component of L_α (for all $\alpha \leq \sigma$, $\beta < \lambda_\alpha$).

In particular, σ , λ_α ($\alpha \leq \sigma$), $n_{\alpha\beta}$, and $K_{\alpha\beta}$ ($\alpha \leq \sigma$, $\beta < \lambda_\alpha$) are invariants of the ring R .

Lemma 2.2. *Let R be a semiartinian von Neumann regular ring with pfa. Then the following hold:*

- (1) *The class of all weakly R -projective modules is closed under submodules.*
- (2) *All countably generated weakly R -projective modules are projective.*

Proof. (1) This follows from the fact [8, 6.28] that all homogenous semisimple modules are injective, whence so are all finitely generated semisimple modules, and thus all finitely generated submodules of each of the layers L_α ($\alpha < \sigma$).

(2) Let C be a countably generated weakly R -projective module. So $C = \bigcup_{n < \omega} F_n$ where $(F_n \mid n < \omega)$ is a chain of finitely generated submodules of C . By part (1), each F_n is weakly R -projective, hence projective by Corollary 1.6. Since R is von Neumann regular, F_n is a direct summand in F_{n+1} for each $n < \omega$, whence C is projective. \square

From Lemma 2.2(1), we see that if the classes of all weak R -projective and projective modules coincide, then R is a right hereditary ring. What that means in our setting is partially clarified in the next lemma:

Lemma 2.3. *Let R be a semiartinian von Neumann regular ring with pfa. Consider the following two conditions:*

- (1) *R is (left and right) hereditary.*
- (2) *σ is a countable ordinal, and all proper layers of R are countably generated (i.e., λ_α is countable for each $0 < \alpha < \sigma$).*

Then (2) implies (1). Moreover, if λ_0 is countable, then (1) and (2) are equivalent.

Proof. That (2) implies (1) was proved in [12, 3.10].

Assume λ_0 is countable and R is right hereditary. Since all the homogenous components of $\text{Soc}(R)$ are injective, and hence finitely generated, $\text{Soc}(R)$ is countably generated.

Assume (2) fails. If σ is uncountable, then $\sigma = \tau + n$ for an uncountable limit ordinal τ and some $n < \omega$, whence $I = S_\tau$ is not countably generated. If λ_α is uncountable for some ordinal $0 < \alpha < \sigma$, then the α th layer L_α is not countably generated, and the same is true of $I = S_{\alpha+1}$.

Since projective modules over von Neumann regular rings are isomorphic to direct sums of cyclic modules generated by idempotents of R , in either case there is a cardinal $\kappa > \aleph_0$ such that $I \cong \bigoplus_{\gamma < \kappa} e_\gamma R$ where $e_\gamma^2 = e_\gamma \in R$ for each $\gamma < \kappa$. Since R is semiartinian, $\text{Soc}(I) \cong \bigoplus_{\gamma < \kappa} \text{Soc}(e_\gamma R)$ is a direct sum of uncountably many non-zero completely reducible modules, in contradiction with $\text{Soc}(I)$ being a direct summand in the countably generated module $\text{Soc}(R)$. \square

Notice that Lemma 2.3 implies that all semiartinian von Neumann regular rings with pfa of Loewy length 2 are hereditary. The simplest such example is the K -subalgebra R_1 of all eventually constant sequences in K^ω for a field K , studied in [11]. The corresponding picture for R_1 is as follows:

$$\begin{array}{|c|} \hline \mathbf{L}_1 = \mathbf{K} \\ \hline \mathbf{L}_0 = \mathbf{K}^{(\omega)} \\ \hline \end{array} \quad \begin{array}{|c|} \hline \mathbf{K} \\ \hline \mathbf{K} \oplus \dots \oplus \mathbf{K} \oplus \dots \dots \\ \hline \end{array}$$

However, semiartinian von Neumann regular rings with pfa of Loewy length 3 need not be hereditary: for an example, take a set S of cardinality ω_1 consisting of almost disjoint infinite subsets of ω , and consider the K -subalgebra R' of K^ω generated (as a K -linear space) by $B \cup C \cup \{1\}$ where 1 is the unit of K^ω , B is the canonical basis of $K^{(\omega)}$ and C is the set of characteristic functions of all the sets in S . By Lemma 2.3, R' is not hereditary. The corresponding picture for R' is

$$\begin{array}{|c|} \hline \mathbf{L}_2 = \mathbf{K} \\ \hline \mathbf{L}_1 = \mathbf{K}^{(\omega_1)} \\ \hline \mathbf{L}_0 = \mathbf{K}^{(\omega)} \\ \hline \end{array} \quad \begin{array}{|c|} \hline \mathbf{K} \\ \hline \mathbf{K} \oplus \dots \oplus \mathbf{K} \oplus \dots \dots \dots \\ \hline \mathbf{K} \oplus \dots \oplus \mathbf{K} \oplus \dots \\ \hline \end{array}$$

The following recursive construction shows that hereditary semiartinian von Neumann regular rings with pfa of Loewy length $\alpha + 1$ do exist for each countable ordinal $\alpha > 0$. The induction step makes use of a construction of semiartinian von Neumann regular rings from [3, 2.4]:

Example 2.4. Let K be a field. By induction on $0 < \alpha < \omega_1$, we will construct semiartinian von Neumann regular K -algebras with pfa, R_α , of Loewy length $\alpha + 1$ such that R_α has countably generated layers, together with K -algebra embeddings $\nu_\alpha : R_\alpha \rightarrow K^\omega$, and for each $0 < \beta < \alpha$, non-unital K -algebra monomorphisms $f_{\alpha\beta} : R_\beta \rightarrow R_\alpha$ and $g_{\alpha\beta} : K^\omega \rightarrow K^\omega$ such that the squares

$$\begin{array}{ccc} R_\beta & \xrightarrow{f_{\alpha\beta}} & R_\alpha \\ \nu_\beta \downarrow & & \nu_\alpha \downarrow \\ K^\omega & \xrightarrow{g_{\alpha\beta}} & K^\omega \end{array}$$

are commutative, $g_{\alpha\beta}$ splits, and the complement of $\text{Im}(g_{\alpha\beta})$ in K^ω is isomorphic to K^ω .

For $\alpha = 1$, we let R_1 be the K -algebra of all eventually constant sequences of elements of K mentioned above. In particular, R_1 is a K -subalgebra of K^ω of Loewy length 2, and ν_1 is defined as the inclusion of R_1 into K^ω .

The induction step is modeled on [3, 2.4]: Assume the construction is done up to some $0 < \alpha < \aleph_1$. Let $I = \bigoplus_{i < \omega} R_\alpha$. Then $\iota_\alpha = \bigoplus_{i < \omega} \nu_\alpha$ embeds I into $D = \bigoplus_{i < \omega} K^\omega \subseteq K^{\omega^\omega}$. As in [3], ω^ω denotes the ordinal exponentiation, so $\omega^\omega = \sup_{i < \omega} \omega^i$ and $\omega^{i+1} = \omega^i \times \omega$ for each $0 < i < \omega$. Note that $\iota_\alpha(I)$ is an ideal of the K -algebra K^{ω^ω} , and $S_{\alpha+1} = \iota_\alpha(I) \oplus 1.K$ is a K -subalgebra of K^{ω^ω} .

Since the ordinal ω^ω is countable, there is a K -algebra isomorphism $\psi : K^{\omega^\omega} \rightarrow K^\omega$. Let $R_{\alpha+1} = \psi(S_{\alpha+1})$, and let $\nu_{\alpha+1}$ denote the inclusion of $R_{\alpha+1}$ into K^ω .

Let μ_α be the embedding of the first copy of K^ω in D composed with the inclusion $D \subseteq K^{\omega^\omega}$. Notice that μ_α is a split non-unital embedding of K -algebras, and so is $\psi\mu_\alpha$. In fact, $\psi\mu_\alpha(K^\omega)$ is a direct summand in K^ω with a complement isomorphic to K^ω . Moreover, $\mu_\alpha \upharpoonright R_\alpha$ is a non-unital K -algebra embedding of R_α into $S_{\alpha+1}$.

We have the following commutative diagram, where the vertical maps are K -algebra embeddings, and the horizontal ones are non-unital K -algebra embeddings:

$$\begin{array}{ccccc} R_\alpha & \xrightarrow{\iota_\alpha \upharpoonright R_\alpha} & S_{\alpha+1} & \xrightarrow{\psi \upharpoonright S_{\alpha+1}} & R_{\alpha+1} \\ \nu_\alpha \downarrow & & \subseteq \downarrow & & \nu_{\alpha+1} \downarrow \\ K^\omega & \xrightarrow{\mu_\alpha} & K^{\omega^\omega} & \xrightarrow{\psi} & K^\omega \end{array}$$

We let $f_{\alpha+1,\alpha} = \psi \iota_\alpha \upharpoonright R_\alpha$, $g_{\alpha+1,\alpha} = \psi \mu_\alpha$, and for each $0 < \beta < \alpha$, $f_{\alpha+1,\beta} = f_{\alpha+1,\alpha} f_{\alpha\beta}$ and $g_{\alpha+1,\beta} = g_{\alpha+1,\alpha} g_{\alpha\beta}$.

By [3, 2.4], the Loewy length of $R_{\alpha+1}$ is $\alpha + 2$. Moreover, for each $\beta \leq \alpha$, the β th layer of $R_{\alpha+1}$ is countably generated, while its $(\alpha + 1)$ th layer, $R_{\alpha+1}/\psi \iota_\alpha(I)$, is isomorphic to K .

If $\alpha < \omega_1$ is a limit ordinal, then $\alpha = \sup_{n < \omega} \beta_n$ for a strictly increasing chain of countable ordinals $(\beta_n \mid n < \omega)$. We fix one such chain with $\beta_0 > 0$. By the induction hypothesis, we have the following commutative diagram, where ν_{β_n} ($n < \omega$) are K -algebra embeddings, f_{β_{n+1},β_n} , g_{β_{n+1},β_n} ($n < \omega$) are non-unital K -algebra embeddings, g_{β_{n+1},β_n} splits, and the complement of $\text{Im}(g_n)$ in K^ω is isomorphic to K^ω ($n < \omega$):

$$\begin{array}{ccccccc} \dots & \longrightarrow & R_{\beta_n} & \xrightarrow{f_{\beta_{n+1},\beta_n}} & R_{\beta_{n+1}} & \longrightarrow & \dots \\ & & \nu_{\beta_n} \downarrow & & \nu_{\beta_{n+1}} \downarrow & & \\ \dots & \longrightarrow & K^\omega & \xrightarrow{g_{\beta_{n+1},\beta_n}} & K^\omega & \longrightarrow & \dots \end{array}$$

Let $T_\alpha = \varinjlim_{n < \omega} R_{\beta_n}$. Since the direct limit of the split embeddings in the bottom row is isomorphic to $\bigoplus_{n < \omega} K^\omega$, we obtain a non-unital K -algebra embedding $\varinjlim_{n < \omega} \nu_{\beta_n} : T_\alpha \rightarrow \bigoplus_{n < \omega} K^\omega$ which can be extended to a K -algebra homomorphism $\nu_\alpha : R_\alpha \rightarrow K^\omega$ by the same procedure as in the induction step above.

The direct limit of the non-unital K -algebra embeddings in the top row yields $f_{\alpha,\beta_n} : R_{\beta_n} \hookrightarrow T_\alpha \hookrightarrow R_\alpha$ for each $n < \omega$. Moreover, we have the split non-unital K -algebra embeddings $g_{\alpha,\beta_n} : K^\omega \rightarrow K^\omega$ that make the corresponding squares for β_n and α commute.

Since for each $\beta < \alpha$, there exists $n < \omega$ such that $\beta < \beta_n$, we can define $f_{\alpha\beta} = f_{\alpha,\beta_n} f_{\beta_n,\beta}$ and $g_{\alpha\beta} = g_{\alpha,\beta_n} g_{\beta_n,\beta}$. Then also the squares for β and α commute, by the induction premise. Finally, R_α is a semiartinian von Neumann regular ring with pfa of Loewy length $\alpha + 1$, all of whose layers are countably generated.

By Lemma 2.3, R_α is hereditary for each $0 < \alpha < \omega_1$. The corresponding picture for R_α is as follows:

$L_\alpha = \mathbf{K}$			\mathbf{K}		
...
$L_\beta = \mathbf{K}^{(\omega)}$	$\mathbf{K} \oplus$...	$\oplus \mathbf{K} \oplus$
...
$L_1 = \mathbf{K}^{(\omega)}$	$\mathbf{K} \oplus$...	$\oplus \mathbf{K} \oplus$
$L_0 = \mathbf{K}^{(\omega)}$	$\mathbf{K} \oplus$...	$\oplus \mathbf{K} \oplus$

Finally, we recall that in the hereditary setting of Lemma 2.3(2), there is a single short exact sequence that tests for weak R -projectivity of any module M :

Lemma 2.5. *Let R be a semiartinian von Neumann regular ring with pfa. Assume σ is a countable ordinal and all proper layers of R are countably generated.*

Then there exist a module B which is a countable direct product of certain ideals of R , an injective module N , and an epimorphism $\pi : B \rightarrow N$, such that the following are equivalent for a module M :

- (1) M is weakly R -projective.
- (2) The homomorphism $\text{Hom}_R(M, \pi) : \text{Hom}_R(M, B) \rightarrow \text{Hom}_R(M, N)$ is surjective.

Proof. This follows by [12, 4.2]. □

Remark 2.6. By [12, 4.1], the epimorphism π is the product of restrictions of the layer epimorphisms π_α ($\alpha \leq \sigma$) to the right ideals $N_{\alpha, F}$, where F runs over all finite subsets of λ_α , $S_\alpha \subseteq N_{\alpha, F} \subseteq S_{\alpha+1}$, and $N_{\alpha, F}/S_\alpha \cong \bigoplus_{\beta \in F} M_{n_{\alpha\beta}}(K_{\alpha\beta})$ is an injective module, cf. Theorem 2.1.

In particular, [12, 4.2] implies that if M is not weakly R -projective, then there exist an $\alpha \leq \sigma$ and a finite subset F of λ_α such that the homomorphism $\text{Hom}_R(M, \pi_\alpha \upharpoonright N_{\alpha, F}) : \text{Hom}_R(M, N_{\alpha, F}) \rightarrow \text{Hom}_R(M, N_{\alpha, F}/S_\alpha)$ is not surjective.

3. WEAK DIAMOND AND WEAK PROJECTIVITY

By Lemma 2.2(2), if R is a semiartinian von Neumann regular ring with pfa, then the notions of a weak R -projective, R -projective, and projective module coincide for any countably generated module. In contrast with Lemma 1.7, we will show that in the extension of ZFC + CH where the prediction principle Φ (Weak Diamond) holds, these notions coincide for *arbitrary* modules, provided that condition (2) of Lemma 2.3 holds (whence R is hereditary) and $\text{card}(R) \leq \aleph_1$.

To simplify our notation, we introduce the following definition:

Definition 3.1. Let R be a semiartinian von Neumann regular ring with pfa. Then R is *small*, if $\text{card}(R) \leq \aleph_1$, σ is a countable ordinal, and all proper layers of R are countably generated.

Notice that our notion of smallness is more general than the one in [12, Definition 4.3] which required the ordinal σ to be finite rather than countable.

Before introducing the Weak Diamond Principle, we need to recall several basic set-theoretic notions:

Let κ be a regular uncountable cardinal. Let A be a set of cardinality $\leq \kappa$. An increasing continuous chain, $(A_\gamma \mid \gamma < \kappa)$, consisting of subsets of A of cardinality $< \kappa$, is a κ -*filtration of the set* A in case its union is A . If A is moreover a module, we will also use the term κ -*filtration of the module* A , which has the additional assumptions that all the A_γ ($\gamma < \kappa$) are submodules of A , and $A_0 = 0$.

A subset $C \subseteq \kappa$ is a *club* in κ , if C is unbounded (i.e., $\sup C = \kappa$) and closed (i.e. for each $D \subseteq C$, if $s = \sup D < \kappa$ then $s \in C$). A subset $E \subseteq \kappa$ is *stationary* in κ , if $E \cap C \neq \emptyset$ for each club C in κ .

Now, we can introduce the Weak Diamond Principle, Φ . We will use it in the following form presented in [4, Lemma VI.1.7] and [7, Theorem 18.12]:

(Φ) Let κ be a regular uncountable cardinal and E a stationary subset in κ . Let A and B be sets of cardinality $\leq \kappa$. Let $(A_\gamma \mid \gamma < \kappa)$ be a κ -filtration of A , and $(B_\gamma \mid \gamma < \kappa)$ a κ -filtration of B . For each $\gamma \in E$, let $c_\gamma : A_\gamma B_\gamma \rightarrow 2$. Then there exists a function $c : E \rightarrow 2$, such that for each $x \in A B$, the set $E(x) = \{\gamma \in E \mid x \upharpoonright A_\gamma \in A_\gamma B_\gamma \text{ and } c(\gamma) = c_\gamma(x \upharpoonright A_\gamma)\}$ is stationary in κ .

The Weak Diamond Φ is easily seen to be a consequence the better known (and stronger) Jensen's Diamond \diamond , which in turn is a consequence of Gödel's Axiom of Constructibility, and hence is consistent with ZFC + GCH. We refer to [4, §VI.1] and [7, §18] for more details.

We arrive at the promised generalization of [12, 4.4] requiring only the Weak Diamond (rather than the stronger Jensen's Diamond), and allowing for arbitrary countable Loewy length of R :

Theorem 3.2. *Assume CH + Φ . Let R be small. Then all weakly R -projective modules, and hence all R -projective modules, are projective.*

Proof. Let M be a weakly R -projective module. By induction on the minimal number of generators, κ , of M , we will prove that M is projective. For $\kappa \leq \aleph_0$, the result follows by Lemma 2.2(2). If κ is a singular cardinal, then we use the fact that the class of all weakly R -projective modules is closed under submodules (Lemma 2.2(1)) and apply Shelah's Singular Compactness Theorem (e.g., in the version from [7, 7.9], see also [4, XII.1.14]).

Assume that κ is a regular uncountable cardinal. Let $A = \{m_\gamma \mid \gamma < \kappa\}$ be a minimal set of R -generators of M . For each $\gamma < \kappa$, let $A_\gamma = \{m_\delta \mid \delta < \gamma\}$. Let M_γ be the submodule of M generated by A_γ . By the inductive premise, M_γ is projective, and $\mathcal{M} = (M_\gamma \mid \gamma < \kappa)$ is a κ -filtration of the module M . Possibly skipping some terms of \mathcal{M} , we can w.l.o.g. assume that \mathcal{M} has the following property for each $\gamma < \kappa$: if M_δ/M_γ is not weakly R -projective for some $\gamma < \delta < \kappa$, then also $M_{\gamma+1}/M_\gamma$ is not weakly R -projective.

(†) Let E be the set of all $\gamma < \kappa$ such that $M_{\gamma+1}/M_\gamma$ is not weakly R -projective. Also, let $\pi : B \rightarrow N$ be the epimorphism from Lemma 2.5. By that Lemma, for each $\gamma \in E$, we can choose an $h_\gamma \in \text{Hom}_R(M_{\gamma+1}/M_\gamma, N)$ such that h_γ does not factorize through π .

We claim that E is not stationary in κ . Assume this claim is not true. Note that CH implies $\text{card}(B) \leq \aleph_1 \leq \kappa$, so we can fix a κ -filtration of the set B , $(B_\gamma \mid \gamma < \kappa)$.

Let $\gamma < \kappa$. For each $g \in \text{Hom}_R(M_\gamma, N)$, we choose $g^+ \in \text{Hom}_R(M_{\gamma+1}, N)$ such that $g^+ \upharpoonright M_\gamma = g$. This is possible because N is an injective module. Further, for each $f \in \text{Hom}_R(M_\gamma, B)$, we choose $f^e \in \text{Hom}_R(M_{\gamma+1}, B)$ such that $\pi f^e = (\pi f)^+$. This is possible since $M_{\gamma+1}$ is projective. Notice that $\pi(f^e \upharpoonright M_\gamma) = (\pi f)^+ \upharpoonright M_\gamma = \pi f$, so $\delta_f := f^e \upharpoonright M_\gamma - f \in \text{Hom}_R(M_\gamma, K)$ where $K = \text{Ker}(\pi)$.

For each $\gamma \in E$, we define $c_\gamma : {}^{A_\gamma}B_\gamma \rightarrow 2$ as follows: If $x : A_\gamma \rightarrow B_\gamma$ is a restriction of a (necessarily unique) morphism $f \in \text{Hom}_R(M_\gamma, B)$ such that the morphism $\delta_f = f^e \upharpoonright M_\gamma - f$ can be extended to a morphism from $\text{Hom}_R(M_{\gamma+1}, K)$, then we put $c_\gamma(x) = 1$. Otherwise, we let $c_\gamma(x) = 0$.

In this setting, Φ yields a function $c : E \rightarrow 2$ such that for each $x \in {}^A B$, the set $E(x) = \{\gamma \in E \mid x \upharpoonright A_\gamma \in {}^{A_\gamma}B_\gamma \text{ and } c(\gamma) = c_\gamma(x \upharpoonright A_\gamma)\}$ is stationary in κ . We will use c to define a morphism $g \in \text{Hom}_R(M, N)$ as follows:

By induction on $\gamma < \kappa$, we define a sequence $(g_\gamma \mid \gamma < \kappa)$ such that $g_\gamma \in \text{Hom}_R(M_\gamma, N)$. First, $g_0 = 0$. If $\gamma < \kappa$, and g_γ is already defined, we distinguish two cases:

- (I) $\gamma \notin E$ or $c(\gamma) = 0$. In this case, we put $g_{\gamma+1} = (g_\gamma)^+$.
- (II) $\gamma \in E$ and $c(\gamma) = 1$. In this case, we let $g_{\gamma+1} = (g_\gamma)^+ + h_\gamma \rho_\gamma$, where $\rho_\gamma : M_{\gamma+1} \rightarrow M_{\gamma+1}/M_\gamma$ is the canonical projection modulo M_γ .

Notice that in both cases $g_{\gamma+1} \upharpoonright M_\gamma = g_\gamma$. We let $g_\gamma = \bigcup_{\delta < \gamma} g_\delta$ in case $\gamma < \kappa$ is a limit ordinal. Then $g = \bigcup_{\gamma < \kappa} g_\gamma \in \text{Hom}_R(M, N)$.

Since M is weakly R -projective, there exists $f \in \text{Hom}_R(M, B)$ such that $g = \pi f$. By Φ , the set $E(f \upharpoonright A) = \{\gamma \in E \mid f \upharpoonright A_\gamma \in {}^{A_\gamma}B_\gamma \text{ and } c(\gamma) = c_\gamma(f \upharpoonright A_\gamma)\}$ is stationary in κ . Let $\gamma \in E(f \upharpoonright A)$.

Assume that $c(\gamma) = 0$. Then we are in case (I), so $\pi(f \upharpoonright M_{\gamma+1}) = g_{\gamma+1} = (g_\gamma)^+ = (\pi f \upharpoonright M_\gamma)^+ = \pi(f \upharpoonright M_\gamma)^e$. Then the morphism $(f \upharpoonright M_\gamma)^e - f \upharpoonright M_{\gamma+1} \in \text{Hom}_R(M_{\gamma+1}, K)$ is an extension of $\delta_{f \upharpoonright M_\gamma} = (f \upharpoonright M_\gamma)^e \upharpoonright M_\gamma - f \upharpoonright M_\gamma$ to $M_{\gamma+1}$, in contradiction with $c_\gamma(f \upharpoonright A_\gamma) = c(\gamma) = 0$.

So necessarily $c(\gamma) = 1$, and we are in case (II), whence $g_{\gamma+1} = (g_\gamma)^+ + h_\gamma \rho_\gamma$. As $c_\gamma(f \upharpoonright A_\gamma) = c(\gamma) = 1$, the morphism $\delta_{f \upharpoonright M_\gamma} = (f \upharpoonright M_\gamma)^e \upharpoonright M_\gamma - f \upharpoonright M_\gamma$ can be extended to a morphism $\Delta_{f \upharpoonright M_\gamma} \in \text{Hom}_R(M_{\gamma+1}, K)$.

Again, $g_{\gamma+1} = \pi(f \upharpoonright M_{\gamma+1})$ and $(g_\gamma)^+ = \pi(f \upharpoonright M_\gamma)^e$, so

$$h_\gamma \rho_\gamma = \pi(f \upharpoonright M_{\gamma+1} - (f \upharpoonright M_\gamma)^e) = \pi u_\gamma,$$

where $u_\gamma = f \upharpoonright M_{\gamma+1} - (f \upharpoonright M_\gamma)^e + \Delta_{f \upharpoonright M_\gamma} \in \text{Hom}_R(M_{\gamma+1}, B)$. However, $u_\gamma \upharpoonright M_\gamma = f \upharpoonright M_\gamma - (f \upharpoonright M_\gamma)^e \upharpoonright M_\gamma + \delta_{f \upharpoonright M_\gamma} = 0$. Hence u_γ factorizes through ρ_γ by some $v_\gamma \in \text{Hom}_R(M_{\gamma+1}/M_\gamma, B)$, that is, $u_\gamma = v_\gamma \rho_\gamma$.

Thus $h_\gamma \rho_\gamma = \pi v_\gamma \rho_\gamma$. Since ρ_γ is surjective, $h_\gamma = \pi v_\gamma$, in contradiction with our choice of h_γ .

This proves our claim about the set E . So there is a club C in κ such that $C \cap E = \emptyset$. Let $z : \kappa \rightarrow \kappa$ be a strictly increasing continuous function whose image is C . For each $\gamma < \kappa$, let $N_\gamma = M_{z(\gamma)}$. Then $(N_\gamma \mid \gamma < \kappa)$ is a κ -filtration of the module M such that $N_{\gamma+1}/N_\gamma$ is weakly R -projective for each $\gamma < \kappa$. By the inductive premise, $N_{\gamma+1}/N_\gamma$ is projective, hence $N_{\gamma+1} = N_\gamma \oplus P_\gamma$ for a projective module P_γ , for each $\gamma < \kappa$. We conclude that $M = N_0 \oplus \bigoplus_{\gamma < \kappa} P_\gamma$ is projective. This finishes the inductive step for the case when κ is a regular uncountable cardinal. \square

Remark 3.3. As observed by Jan Šároch, Remark 2.6 makes it possible to prove Theorem 3.2 without the assumption of CH, i.e., assuming only Φ . (That is indeed a weaker assumption, since unlike \diamond , the Φ does not imply CH.)

The only modification needed concerns the set E of all $\gamma < \kappa$ such that $M_{\gamma+1}/M_\gamma$ is not weakly R -projective, and the morphisms h_γ ($\gamma \in E$) defined in part (†) of the proof of Theorem 3.2:

For each $\gamma \in E$, Remark 2.6 yields an $\alpha_\gamma \leq \sigma$, a finite subset F_γ of λ_{α_γ} , and an $h_\gamma \in \text{Hom}_R(M_{\gamma+1}/M_\gamma, N_{\alpha_\gamma, F_\gamma}/S_\alpha)$ which does not factorize through $\pi_\alpha \upharpoonright N_{\alpha_\gamma, F_\gamma}$. For each $\alpha \leq \sigma$ and each finite subset F of λ_α , let $E_{\alpha, F} = \{\gamma \in E \mid \alpha_\gamma = \alpha \text{ \& } F_\gamma = F\}$. Then $E = \bigcup_{\alpha, F} E_{\alpha, F}$. Notice that since R is small, the set of all such pairs (α, F) is countable.

Thus, if E is stationary, then so is one of the $E_{\alpha, F}$ (see e.g. [4, II.4.3]), say $E_{\alpha', F'}$. The proof of a contradiction with the stationarity of E then proceeds as that of Theorem 3.2 in the parts following (†), but for $B = N_{\alpha', F'}$, $N = N_{\alpha', F'}/S_{\alpha'}$, $\pi = \pi_{\alpha'} \upharpoonright N_{\alpha', F'}$, and $E = E_{\alpha', F'}$. The point is that in this setting, the cardinality of B is $\leq \text{card}(R) \leq \aleph_1$ even without the assumption of CH.

Combining Lemma 1.7 and Theorem 3.2, we obtain

Corollary 3.4. *Let R be small. Then the assertion ‘All weakly R -projective modules are projective’ is independent of ZFC + GCH.*

Acknowledgment 3.5. The author thanks Kateřina Fuková for valuable comments on an earlier draft of this paper.

REFERENCES

- [1] H. Alhilali, Y. Ibrahim, G. Puninski, M. Yousif, *When R is a testing module for projectivity?*, J. Algebra 484(2017), 198-206.
- [2] F. W. Anderson, K. R. Fuller, *Rings and Categories of Modules*, 2nd ed., GTM 13, Springer, New York 1992.
- [3] P. C. Eklof, K. R. Goodearl, J. Trlifaj, *Dually slender modules and steady rings*, Forum Math. 9(1997), 61 – 74.
- [4] P.C. Eklof, A.H. Mekler, *Almost free modules*, Revised ed., North-Holland, New York 2002.
- [5] P. C. Eklof, S. Shelah, *On Whitehead modules*, J. Algebra 142(1991), 492 – 510.
- [6] C. Faith, *Algebra II. Ring Theory*, GMW 191, Springer-Verlag, Berlin 1976.
- [7] R. Göbel, J. Trlifaj, *Approximations and Endomorphism Algebras of Modules*, 2nd revised and extended ed., GEM 41, W. de Gruyter, Berlin 2012.
- [8] K.R. Goodearl, *Von Neumann Regular Rings*, 2nd ed., Krieger Publ. Co., Malabar 1991.
- [9] P. Růžička, J. Trlifaj, J. Žemlička, *Criteria of steadiness*, Abelian Groups, Module Theory, and Topology, M.Dekker, New York 1998, 359-371.
- [10] J. Šároch, J. Trlifaj, *Test sets for factorization properties of modules*, Rend. Sem. Mat. Univ. Padova 144(2020), 217-238.
- [11] J. Trlifaj, *Faith's problem on R -projectivity is undecidable*, Proc. Amer. Math. Soc. 147(2019), 497-504.
- [12] J. Trlifaj, *The dual Baer criterion for non-perfect rings*, Forum Math. 32(2020), 663-672.

CHARLES UNIVERSITY, FACULTY OF MATHEMATICS AND PHYSICS, DEPARTMENT OF ALGEBRA,
 SOKOLOVSKÁ 83, 186 75 PRAGUE 8, CZECH REPUBLIC
Email address: trlifaj@karlin.mff.cuni.cz