HALF-ISOMORPHISMS OF AUTOMORPHIC LOOPS

MARIA DE LOURDES MERLINI GIULIANI

GILIARD SOUZA DOS ANJOS

Centro de Matemática, Computação e Cognição Universidade Federal do ABC Av. dos Estados, 5001, 09210-580 Santo André - SP, Brazil maria.giuliani@ufabc.edu.br

Instituto de Matemática e Estatística Universidade de São Paulo Rua do Matão, 1010, 05508-090 São Paulo - SP, Brazil giliard.anjos@unesp.br

Abstract

Automorphic loops are loops in which all inner mappings are automorphisms. This variety of loops includes groups and commutative Moufang loops. A half-isomorphism $f: G \longrightarrow K$ between multiplicative systems G and K is a bijection from G onto K such that $f(ab) \in \{f(a)f(b), f(b)f(a)\}$ for any $a, b \in G$. A half-isomorphism is trivial when it is either an isomorphism or an anti-isomorphism. Consider the class of automorphic loops such that the equation $x \cdot (x \cdot y) = (y \cdot x) \cdot x$ is equivalent to $x \cdot y = y \cdot x$. Here we show that this class of loops includes automorphic loops of odd order and uniquely 2-divisible. Furthermore, we prove that every half-isomorphism between loops in that class is trivial.

Keywords: half-isomorphism, half-automorphism, automorphic loop. Mathematics Subject Classification: 20N05.

1 Introduction

A loop (L, .) is a set L with a binary operation (\cdot) such that for each $a, b \in L$ there exist unique elements $x, y \in L$ such that $a \cdot x = b$ and $y \cdot a = b$, and there is an identity element 1 such that $1 \cdot x = x = x \cdot 1$ for every $x \in L$. The operation \cdot is denoted by juxtaposition for the sake of convenience, so (L, \cdot) is denoted by L and $x \cdot y$ by xy.

For any $a \in L$, the right translation and left translation by a are the bijections $R_a : L \to L$; $x \mapsto x \cdot a$ and $L_a : L \to L$; $x \mapsto a \cdot x$ respectively. These bijections generate the multiplication group $Mlt(L) = \langle R_a, L_a; a \in L \rangle$. The inner mapping group, denoted by Inn(L), is the subgroup of Mlt(L) containing the elements that stabilize the identity element. Automorphic loops (or A-loops in Bruck's notation [2]) are loops in which every inner mapping is an automorphism, or Inn(L) is a subgroup of Aut(L), where Aut(L) is the automorphism group of L. It is well known that automorphic loops are power associative, i.e., each element generates a cyclic group.

Let (L, *) and (L', \cdot) be loops. A bijection $f: L \longrightarrow L'$ is called a half-isomorphism if

 $f(x * y) \in \{f(x) \cdot f(y), f(y) \cdot f(x)\}, \text{ for any } x, y \in L.$

A half-automorphism and a half-homomorphism are defined in a similar manner. We say that a half-isomorphism is *trivial* when it is either an isomorphism or an anti-isomorphism. Likewise, a half-isomorphism is *nontrivial* when it is neither an isomorphism nor an anti-isomorphism (In [9], the word "proper" is used instead).

All half-isomorphisms are trivial for groups, as was shown by Scott [16], but not for loops. For example, in the case of the variety of Moufang loops, there exist loops of even order with nontrivial half-isomorphism [3, 12] as well as loops of odd order in which all half-isomorphisms are trivial [4].

These and more recent studies of half-isomorphisms in different classes of loops have shown that the behavior of these functions is rather unpredictable. For example, investigating dihedral automorphic loops, which is a class of automorphic loops of even order, we proved that there exist half-isomorphisms which are neither isomorphisms nor anti-isomorphisms [10]. Another case is the case of the automorphic loops of odd order of the class called Lie automorphic loops for which we proved that all half-automorphisms are trivial [11].

In the current paper, we prove that if Q and Q' are automorphic loops, and in Q' the equation $x \cdot (x \cdot y) = (y \cdot x) \cdot x$ is equivalent to the equation $x \cdot y = y \cdot x$, then every half-isomorphism from Q onto Q' is trivial. The conjecture, presented in [11], that every half-isomorphism between automorphic loops of odd order is trivial, is a corollary of this result.

2 Preliminaries

Part of the investigations for this work (section 3) were assisted by Prover9 [13], an automated computational tool for equational reasoning. However, the raw output is usually overcomplicated and here we present only "humanized" versions on these proofs. Also, the LOOPS package [14] for GAP [5] was used.

In this section, the required definitions and basic results are stated. For general facts about loop theory we recommend [1, 15].

Let L be a loop and $x, y \in L$. Consider the following bijections:

$$R_{(x,y)} = R_x R_y R_{(xy)}^{-1},\tag{1}$$

$$L_{(x,y)} = L_x L_y L_{(yx)}^{-1}, (2)$$

$$T_x = R_x L_x^{-1}. (3)$$

It is well-known that the inner mapping group of L is generated by these bijections, that is:

$$Inn(L) = \langle R_{(x,y)}, L_{(x,y)}, T_x \mid x, y \in L \rangle.$$

The following lemma is straightforward.

Lemma 2.1. Let L be a loop, $x, y, z \in L$ and $\varphi \in Aut(L)$. Then (a) $(y)L_x^{-1}\varphi = (y)\varphi L_{(x)\varphi}^{-1}$, (b) $(y)R_x^{-1}\varphi = (y)\varphi R_{(x)\varphi}^{-1}$, (c) $x(y)L_x^{-1} = y$, (d) $(y)R_x^{-1}x = y$, (e) xy = yx if and only if $(y)T_x = y$.

Let S be a non empty subset of L and $K = \{H \leq L \mid S \subset H\}$. The subloop of L generated by S, denoted by $\langle S \rangle$, is the following subloop of L:

$$\langle S \rangle = \bigcap_{H \in K} H.$$

Proposition 2.2. ([2, Thereom 2.4]) Let Q be an automorphic loop. Then

- (a) Every maximal commutative subset of Q is a subloop of Q.
- (b) Every maximal associative subset of Q is a subgroup of Q.

(c) Q is power associative, that is, $\langle x \rangle$ is associative, for every $x \in Q$.

If Q is non-commutative (non-associative), then by Zorn's Lemma we have that every commutative (associative) subset of Q is contained in at least one maximal commutative (associative) subset of Q. Then the next result is a direct consequence of Proposition 2.2.

Corollary 2.3. Let Q be an automorphic loop and S be a non empty subset of Q. If S is commutative (associative), then $\langle S \rangle$ is commutative (associative).

Proposition 2.4. ([2, Theorem 2.6]) Let Q be an automorphic loop. Then, for all $x, y, z \in Q$ and $m, n, r \in \mathbb{Z}$: (a) $x^m(x^ny) = x^n(x^my)$, (b) $(yx^m)x^n = (yx^n)x^m$, (c) $(x^my)x^n = x^m(yx^n)$,

(d) x((yx)(zx)) = ((xy)(xz))x.

As a direct consequence of the items (a), (b) and (c) of the above Proposition, we have the following result.

Corollary 2.5. Let Q be an automorphic loop and $x \in Q$. Then $\langle L_{x^m}, R_{x^n} | m, n \in \mathbb{Z} \rangle$ is an abelian group. In particular, we have for all $m \in \mathbb{Z}$:

$$T_x^m = R_x^m L_x^{-m} = L_x^{-m} R_x^m.$$
(4)

Remark 2.6. We will omit the parenthesis when occur one of the items (a), (b) and (c) of Proposition 2.4. For example, we will write only $x^{-1}yx$ instead of $(x^{-1}y)x$ or $x^{-1}(yx)$.

Proposition 2.7. ([7, Theorem 7.5]) Every automorphic loop has the antiautomorphic inverse property (AAIP), that is, the following identity holds:

$$(xy)^{-1} = y^{-1}x^{-1} \tag{AAIP}$$

The next result is a consequence of the (AAIP).

Corollary 2.8. If Q is an automorphic loop and $x, y \in Q$, then

$$((y)L_x^{-1})^{-1} = (y^{-1})R_{x^{-1}}^{-1}.$$
(5)

Proposition 2.9. ([8, Lemma 2.7]) In an automorphic loop Q, the following holds, for all $x \in Q$:

$$T_x^{-1} = T_{x^{-1}}. (6)$$

Define $\gamma: Q \to Q$ by $(x)\gamma = x^2$. We say that Q is uniquely 2-divisible if γ is a bijection. In this case, we define $x^{1/2} = (x)\gamma^{-1}$, for every $x \in Q$. Note that, if $\varphi \in Aut(Q)$ and $x \in Q$, then

$$(x^{1/2})\varphi = ((x)\varphi)^{1/2}$$
(7)

Proposition 2.10. ([8, Corollary 4.7]) A finite automorphic loop is uniquely 2-divisible if and only if it has odd order.

A half-isomorphism $f: L \to L'$ is called *special* if the inverse mapping $f^{-1}: L' \to L$ is also a half-isomorphism. Every trivial half-isomorphism is special. The next example presents a half-isomorphism that is not special.

Example 2.11. Let $L = \{1, 2, ..., 7\}$ and consider the following Cayley tables:

*	1	2	3	4	5	6	7		1	2	3	4	5	6	7
1	1	2	3	4	5	6	7	1	1	2	3	4	5	6	7
2	2	3	4	5	6	7	1	2	2	3	7	5	6	1	4
3	3	4	5	6	7	1	2	3	3	4	5	6	7	2	1
4	4	5	6	7	1	2	3	4	4	5	6	7	1	3	2
5	5	6	7	1	2	3	4	5	5	6	4	1	2	7	3
6	6	7	1	2	3	4	5	6	6	7	1	2	3	4	5
7	7	1	2	3	4	5	6	7	7	1	2	3	4	5	6

One can see that $C_7 = (L, *)$ is the cyclic group of order 7 and $L' = (L, \cdot)$ is a non-associative loop. Defining $f: C_7 \to L'$; f(x) = x, then it is not difficult to see that f is a half-isomorphism. Furthermore,

$$f(3*2) = 4 = f(3) \cdot f(2) \neq f(2) \cdot f(3) \quad \text{and} \quad f(3*6) = 1 = f(6) \cdot f(3) \neq f(3) \cdot f(6),$$

then f is a nontrivial half-isomorphism. Now note that $f^{-1}(2 \cdot 3) = 7 \notin \{2 * 3, 3 * 2\}$, and so f^{-1} is not a half-isomorphism.

Proposition 2.12. ([11, Theorem 2.5]) Let L and L' be loops and $f : L \to L'$ be a halfisomorphism. Then the following statements are equivalent: (a) f is special. (b) $\{f(m+n)\} = \{f(n), f(n)\} = \{f(n), f(n)\}$ for any $n \in L$

(b) $\{f(x * y), f(y * x)\} = \{f(x) \cdot f(y), f(y) \cdot f(x)\}$ for any $x, y \in L$.

(c) For all $x, y \in L$ such that x * y = y * x, we have $f(x) \cdot f(y) = f(y) \cdot f(x)$.

Proposition 2.13. Let L and L' be power associative loops and $f: L \to L'$ be a half-isomorphism. Then $f(x^n) = f(x)^n$, for all $x \in L$ and $n \in \mathbb{Z}$.

Proof. By definition, we have $f(1_L) = f(1_L * 1_L) = f(1_L) \cdot f(1_L)$, and then $f(1_L) = 1_{L'}$. Let $x \in L$ and $n \in \mathbb{N}^*$. Suppose that $f(x^n) = f(x)^n$. Since L and L' are power associative, we have that

$$f(x^{n+1}) = f(x^n * x) \in \{f(x)^n \cdot f(x), f(x) \cdot f(x)^n\} = \{f(x)^{n+1}\}.$$

Then $f(x^m) = f(x)^m$, for all $m \in \mathbb{N}$. Now, for $n \in \mathbb{N}$, we have that

$$1_{L'} = f(x^n * x^{-n}) \in \{f(x)^n \cdot f(x^{-n}), f(x^{-n}) \cdot f(x)^n\},\$$

and then $f(x^{-n}) = f(x)^{-n}$ since L' is power associative.

3 Some results about automorphic loops

In this section, we will prove that for every uniquely 2-divisible automorphic loop the equation $x \cdot (x \cdot y) = (y \cdot x) \cdot x$ is equivalent to the equation $x \cdot y = y \cdot x$. First, we need to prove some identities for automorphic loops.

Lemma 3.1. In an automorphic loop Q, the following identities hold:

$$(xy)^2 = (x \cdot (y)R_{x^{-1}}^{-1})y \tag{8}$$

$$(xy)^2 = x \cdot (xy)L_{y^{-1}}^{-1} \tag{9}$$

$$(xyx^{-1})^2 = (xy)(yx^{-1}) \tag{10}$$

$$(x^2)R_y^{-1} = (x)R_{((x)R_y^{-1})^{-1}}^{-1}$$
(11)

$$(yx)x^{-1} = x^{-1}(xy)$$
(12)
$$x^{2} - (x)B^{-1} \cdot (x)L^{-1}$$
(13)

$$x^{2} = (x)R_{y}^{-1} \cdot (x)L_{y^{-1}}^{-1}$$
(13)

$$(y^2 x)x^{-1} = (yx)(x^{-1}y) \tag{14}$$

$$(y^2x)x^{-1} = (yx^{-1})(xy). (15)$$

Proof. (i) Consider $\varphi = R_y^{-1}R_{x^{-1}}L_{xy}$. We have that $(1)\varphi = (xy)(y^{-1}x^{-1}) = 1$, where we used (AAIP) in the second equality, and then $\varphi \in Aut(Q)$. Furthermore, $(xy)\varphi = (xy)(xx^{-1}) = xy$, and so $((xy)^2)\varphi = (xy)^2$. Thus

$$((xy)^2)R_y^{-1}R_{x^{-1}}L_{xy} = (xy)L_{xy},$$

and hence $(xy)^2 = (y)L_x R_{x^{-1}}^{-1} R_y$. Since $R_{x^{-1}}^{-1} L_x = L_x R_{x^{-1}}^{-1}$ (Corollary 2.5), we get (8). (ii) Consider $\varphi = L_x^{-1} L_{y^{-1}} L_{xy}$. We have that $\varphi(1) = (xy)(y^{-1}x^{-1}) = 1$, where we use (AAIP) in the second equality, and then $\varphi \in Aut(Q)$. Furthermore, $\varphi(xy) = (xy)(y^{-1}y) = xy$, and so $((xy)^2)\varphi = (xy)^2$. Thus

$$((xy)^2)L_x^{-1}L_{y^{-1}}L_{xy} = (xy)L_{xy}$$

and hence $(xy)^2 = (xy)L_{y^{-1}}^{-1}L_x$ and we have (9). (iii) Putting $y = vx^{-1}$ in (8), we get

$$(xvx^{-1})^2 = (x(vx^{-1})R_{x^{-1}}^{-1})(vx^{-1})$$

and then (10) follows from the fact that $(vx^{-1})R_{x^{-1}}^{-1} = v$. (iv) Putting $x = (u)R_y^{-1}$ in (8), we get

$$((u)R_y^{-1}y)^2 = ((u)R_y^{-1}(y)R_{((u)R_y^{-1})^{-1}}^{-1})y$$

By Lemma 2.1 (d), we have $u = (u)R_y^{-1}y$, and then

$$(u^2)R_y^{-1} = (u)R_y^{-1}(y)R_{((u)R_y^{-1})^{-1}}^{-1},$$

and using Lemma 2.1 (d) again and Proposition 2.4 we get

$$(u^{2})R_{y}^{-1}((u)R_{y}^{-1})^{-1} = (u)R_{y}^{-1}(y)R_{((u)R_{y}^{-1})^{-1}}^{-1}((u)R_{y}^{-1})^{-1} = (u)R_{y}^{-1}y = u$$

and hence we have (11).

(v) By (6), $(y)T_x = (y)T_{x^{-1}}^{-1}$, and then $(y)R_xL_x^{-1} = (y)L_{x^{-1}}R_{x^{-1}}^{-1}$. Thus $(y)R_xL_x^{-1}R_{x^{-1}} = (y)L_{x^{-1}}$ and hence $(y)R_xR_{x^{-1}}L_x^{-1} = (y)L_{x^{-1}}$ by Corollary 2.5. It follows that $(y)R_xR_{x^{-1}} = (y)L_{x^{-1}}L_x = (y)L_xL_{x^{-1}}$, where we used Corollary 2.5 in the second equality, and therefore we have (12).

(vi) Putting $x = (u)R_u^{-1}$ in (9), we get

$$((u)R_y^{-1}y)^2 = (u)R_y^{-1} \cdot ((u)R_y^{-1}y)L_{y^{-1}}^{-1},$$

and then (13) follows from the fact that $(u)R_y^{-1}y = u$ (Lemma 2.1 (d)). (vii) Let $u, v \in Q$ and $\phi = R_u R_{u^{-1}}$. It is clear that $(1)\phi = 1$, and then $\phi \in Aut(Q)$. Thus $(v^2)\phi = ((v)\phi)^2$, and we have $((vu)u^{-1})^2 = (v^2u)u^{-1}$. By (13), we have

$$(v^{2}u)u^{-1} = ((vu)u^{-1})R_{z}^{-1} \cdot ((vu)u^{-1})L_{z^{-1}}^{-1}, \text{ for all } z \in Q.$$

$$(16)$$

(vii.1) Putting $z = u^{-1}$ in (16), we get

$$(v^{2}u)u^{-1} = ((vu)u^{-1})R_{u^{-1}}^{-1} \cdot ((vu)u^{-1})L_{u}^{-1} = (vu) \cdot ((vu)u^{-1})L_{u}^{-1}$$

Using (12) and Proposition 2.4 (a) we get $((vu)u^{-1})L_u^{-1} = (u^{-1}(uv))L_u^{-1} = (u(u^{-1}v))L_u^{-1} = u^{-1}v$. Therefore $(v^2u)u^{-1} = (vu)(u^{-1}v)$ and we have (14). (vii.2) Putting z = u in (16), we get

$$(v^2u)u^{-1} = ((vu)u^{-1})R_u^{-1} \cdot ((vu)u^{-1})L_{u^{-1}}^{-1}.$$

Using Proposition 2.4 (b) we get $((vu)u^{-1})R_u^{-1} = ((vu^{-1})u)R_u^{-1} = vu^{-1}$. Using (12) we get $((vu)u^{-1})L_u^{-1} = (u^{-1}(uv))L_u^{-1} = uv$. Therefore $(v^2u)u^{-1} = (vu^{-1})(uv)$ and we have (15).

Corollary 3.2. Let Q be an automorphic loop and $x, y \in Q$. Consider that one of the following hold:

(a) $x^{-1}(xy^2) = (xy)(x^{-1}y)$ or (b) $x^{-1}(xy^2) = (x^{-1}y)(xy)$. Then xy = yx.

Proof. (a) Using (12) we get $(y^2x)x^{-1} = (xy)(x^{-1}y)$, and then $(xy)(x^{-1}y) = (yx)(x^{-1}y)$ by (14). Hence xy = yx.

(b) Using (12) we get $(y^2x)x^{-1} = (x^{-1}y)(xy)$, and then $(x^{-1}y)(xy) = (yx^{-1})(xy)$ by (15). Thus $x^{-1}y = yx^{-1}$ and from Corollary 2.3 we get xy = yx.

Lemma 3.3. In an automorphic loop Q, the following identities hold:

(

$$((x)R_{((x)R_y^{-1})^{-1}}^{-1})^{-1} \cdot (x)R_y^{-1} = ((x)R_y^{-1})^{-1} \cdot y^{-1}$$
(17)

$$(((x)R_{xy}^{-1})^{-1}R_{x^{-1}}^{-1})^{-1} \cdot x = (x)R_{yx}^{-1}$$
(18)

$$(x \cdot (y)R_z^{-1})z)R_y^{-1} = (xz)R_y^{-1} \cdot ((z)R_y^{-1})^{-1}$$
(19)

$$(xy)R_z^{-1} \cdot ((y)R_z^{-1})^{-1} = ((x)R_z^{-1} \cdot ((y)R_z^{-1})^{-1})R_y$$
(20)

$$x^{-1}((x)R_{(y)T_x^{-1}}^{-1} \cdot ((y)R_x^{-1}R_{(y)T_x^{-1}}^{-1})^{-1}) = (x)R_{(y)T_x^{-1}}^{-1} \cdot ((y)R_{(y)T_x^{-1}}^{-1})^{-1}$$
(21)

$$((x)R_{xy}^{-1})^{-1}R_{(y)R_{xy}^{-1}\cdot((x)R_{xy}^{-1})^{-1}}^{-1} = x.$$
(22)

Proof. (i) We have that

$$\begin{aligned} ((x)R_y^{-1})^{-1} \cdot (x)R_{((x)R_y^{-1})^{-1}}^{-1} \cdot ((x)R_y^{-1})^{-1} &= ((x)R_y^{-1})^{-1} \cdot x \\ &= ((x)R_y^{-1})^{-1}((x)R_y^{-1} \cdot y) \\ \stackrel{(12)}{=} & (y \cdot (x)R_y^{-1})((x)R_y^{-1})^{-1}. \end{aligned}$$

Then $((x)R_y^{-1})^{-1} \cdot (x)R_{((x)R_y^{-1})^{-1}}^{-1} = y \cdot (x)R_y^{-1}$, and applying (AAIP) we obtain (17).

(ii) By Lemma 2.1 (b), we have that $(x)R_{xy}^{-1}T_x = (x)T_xR_{(xy)T_x}^{-1}$. Since $(x)T_x = x$ and $(xy)T_x = yx$, we get $(x)R_{xy}^{-1}T_x = (x)R_{yx}^{-1}$. By (4), $T_x = L_x^{-1}R_x$, and then $(x)R_{xy}^{-1}L_x^{-1}R_x = (x)R_{yx}^{-1}$. Using (5) in the left side of this equation, we get

$$(((x)R_{xy}^{-1})^{-1}R_{x^{-1}}^{-1})^{-1}R_x = (x)R_{yx}^{-1}.$$

Hence we have (18).

(iii) Note that $R_{((y)R_z^{-1},z)} = R_{(y)R_z^{-1}}R_zR_y^{-1}$ since $y = (y)R_z^{-1} \cdot z$. Applying this automorphism to $x = (x)R_{(y)R_z^{-1}}^{-1} \cdot (y)R_z^{-1}$, we get

Then we have (19).

(iv) Applying the automorphism $R_{((z)R_y^{-1},y)} = R_{(z)R_y^{-1}}R_yR_z^{-1}$ to $x = (x)R_y^{-1} \cdot y$, we get

$$(x)R_{(z)R_y^{-1}}R_yR_z^{-1} = ((x)R_y^{-1})R_{(z)R_y^{-1}}R_yR_z^{-1} \cdot (y)R_{(z)R_y^{-1}}R_yR_z^{-1}$$
(23)

By (19), we have

$$(x)R_{(z)R_y^{-1}}R_yR_z^{-1} = (xy)R_z^{-1} \cdot ((y)R_z^{-1})^{-1} \text{ and } \\ ((x)R_y^{-1})R_{(z)R_y^{-1}}R_yR_z^{-1} = ((x)R_y^{-1} \cdot y)R_z^{-1} \cdot ((y)R_z^{-1})^{-1} = (x)R_z^{-1} \cdot ((y)R_z^{-1})^{-1}.$$

Furthermore,

$$(y)R_{(z)R_y^{-1}}R_yR_z^{-1} = (y \cdot (z)R_y^{-1} \cdot y)R_z^{-1} = (yz)R_z^{-1} = y.$$

Hence (20) follows from (23).

(v) From (20) we have

$$((u)R_v^{-1}\cdot ((x)R_v^{-1})^{-1})x = (ux)R_v^{-1}\cdot ((x)R_v^{-1})^{-1},$$

and applying (AAIP) we get

$$x^{-1}((x)R_v^{-1} \cdot ((u)R_v^{-1})^{-1}) = (x)R_v^{-1} \cdot ((ux)R_v^{-1})^{-1}.$$

Putting $u = (y)R_x^{-1}$ and $v = (y)T_x^{-1}$ in this equation, we obtain (21).

(vi) From (19) we have $((u \cdot (y)R_x^{-1})x)R_y^{-1} = (ux)R_y^{-1} \cdot ((x)R_y^{-1})^{-1}$. Putting $u = ((y)R_x^{-1})^{-1}$ in this equation, we get

$$(x)R_y^{-1} = (((y)R_x^{-1})^{-1} \cdot x)R_y^{-1} \cdot ((x)R_y^{-1})^{-1}$$

$$\begin{array}{ll} \stackrel{(20)}{\Longrightarrow} & (x)R_y^{-1} = (((y)R_x^{-1})^{-1}R_y^{-1} \cdot ((x)R_y^{-1})^{-1})R_x \\ \stackrel{(x)R_y^{-1}R_x^{-1}}{=} ((y)R_x^{-1})^{-1}R_y^{-1} \cdot ((x)R_y^{-1})^{-1} \\ \stackrel{(x)R_y^{-1}R_x^{-1}}{=} ((x)R_y^{-1})^{-1} \\$$

Putting x = wy in this equation and noticing that $(wy)R_y^{-1} = w$, we have

$$w = ((w)R_{wy}^{-1})^{-1}R_{(((y)R_{wy}^{-1})^{-1}R_{y}^{-1})^{-1}}^{-1}.$$
(24)

From (20) we have $(wy)R_{wy}^{-1} \cdot ((y)R_{wy}^{-1})^{-1} = ((w)R_{wy}^{-1} \cdot ((y)R_{wy}^{-1})^{-1})R_y$. Since $(wy)R_{wy}^{-1} = 1$, it follows that

$$((y)R_{wy}^{-1})^{-1}R_y^{-1} = (w)R_{wy}^{-1} \cdot ((y)R_{wy}^{-1})^{-1},$$

and using (AAIP) we get

$$(((y)R_{wy}^{-1})^{-1}R_y^{-1})^{-1} = (y)R_{wy}^{-1} \cdot ((w)R_{wy}^{-1})^{-1}.$$

Putting the above equation in (24), we obtain (22).

Proposition 3.4. In an automorphic loop Q, the following identity holds:

$$(x)R_{(x)T_y^{-1}}^{-1} \cdot y = (y)T_x^{-1}$$
(25)

Proof. From (17) we have $((u)R_{((u)R_v^{-1})^{-1}}^{-1} \cdot (u)R_v^{-1} = ((u)R_v^{-1})^{-1} \cdot v^{-1}$. If $u = (x)R_{xy}^{-1}$ and $v = (y)R_{xy}^{-1} \cdot ((x)R_{xy}^{-1})^{-1}$, then $(u)R_v^{-1} = x$ by (22), and we get

$$(((x)R_{xy}^{-1})^{-1}R_{x^{-1}}^{-1})^{-1} \cdot x = x^{-1} \cdot ((y)R_{xy}^{-1} \cdot ((x)R_{xy}^{-1})^{-1})^{-1}$$

Using (18) in the left side and (AAIP) in the right side of this equation, we obtain

$$(x)R_{yx}^{-1} = x^{-1}((x)R_{xy}^{-1} \cdot ((y)R_{xy}^{-1})^{-1}).$$

Putting w = yx in the above equation and noticing that $y = (w)R_x^{-1}$, we get

$$(x)R_w^{-1} = x^{-1}((x)R_{x\cdot(w)R_x^{-1}}^{-1}\cdot((w)R_x^{-1}R_{x\cdot(w)R_x^{-1}}^{-1})^{-1}).$$

Since $(w)T_x^{-1} = x \cdot (w)R_x^{-1}$, from (21) we have

$$(x)R_w^{-1} = (x)R_{(w)T_x^{-1}}^{-1} \cdot ((w)R_{(w)T_x^{-1}}^{-1})^{-1},$$

and then

$$x = ((x)R_{(w)T_x^{-1}}^{-1} \cdot ((w)R_{(w)T_x^{-1}}^{-1})^{-1})R_w.$$
(26)

Putting $z = (w)T_x^{-1}$ in (20), we have

$$(xw)R_{(w)T_x^{-1}}^{-1} \cdot ((w)R_{(w)T_x^{-1}}^{-1})^{-1} = ((x)R_{(w)T_x^{-1}}^{-1} \cdot ((w)R_{(w)T_x^{-1}}^{-1})^{-1})R_w,$$

and putting this in (26), we obtain

$$x = (xw)R_{(w)T_x^{-1}}^{-1} \cdot ((w)R_{(w)T_x^{-1}}^{-1})^{-1}$$

$$\begin{array}{lll} \underset{(x)}{\overset{(x)}{\longrightarrow}}{\longrightarrow} & x = ((xw)T_{x}R_{w}^{-1} \cdot ((w)T_{x}R_{w}^{-1})^{-1})T_{x}^{-1} \\ \underset{(xw)}{\overset{(x)}{\longrightarrow}}{\longrightarrow} & x = (xw)T_{x}R_{w}^{-1} \cdot ((w)T_{x}R_{w}^{-1})^{-1} \\ \underset{(xw)}{\overset{(xw)}{\longrightarrow}}{\longrightarrow} & x = (wx)R_{w}^{-1} \cdot ((w)T_{x}R_{w}^{-1})^{-1} \\ \underset{(xv)}{\longrightarrow} & x = (x)T_{w}^{-1} \cdot ((w)T_{x}R_{w}^{-1})^{-1} \\ \underset{(x)}{\longrightarrow} & (x)L_{(x)T_{w}^{-1}}^{-1} = ((w)T_{x}R_{w}^{-1})^{-1} \\ \underset{(x)}{\overset{(5)}{\longrightarrow}} & (x^{-1})R_{(x^{-1})T_{w}^{-1}}^{-1} = (w)T_{x}R_{w}^{-1} \\ \underset{(x^{-1})}{\longrightarrow} & (x^{-1})R_{(x^{-1})T_{w}^{-1}}^{-1}R_{w} = (w)T_{x} \\ \underset{(x)}{\overset{(6)}{\longrightarrow}} & (x^{-1})R_{(x^{-1})T_{w}^{-1}}^{-1}R_{w} = (w)T_{x^{-1}} \end{array} \right$$

Let Q be an automorphic loop. For $x, y \in Q$, consider the following condition:

$$x(xy) = (yx)x \text{ if and only if } xy = yx.$$
(27)

By (4), we have $T_x^2 = R_x^2 L_x^{-2}$, and then the above condition can be rewritten as:

$$(y)T_x^2 = y$$
 if and only if $(y)T_x = y.$ (28)

Remark 3.5. In flexible loops (and therefore in automorphic loops), xy = yx implies that x(xy) = (yx)x.

As a consequence of (28) and Corollary 2.3, we have the following result.

Lemma 3.6. Let Q be an automorphic loop satisfying (27) and let $x, y \in Q$. If $(y)T_x^2 = y$, then $\langle x, y \rangle$ is commutative.

Corollary 3.7. Let Q be an automorphic loop satisfying (27) and let $x, y \in Q$. If $xyx^{-1} = x^{-1}yx$, then $\langle x, y \rangle$ is commutative.

Proof. By assumption, $(y)R_{x^{-1}}L_x = (y)R_xL_{x^{-1}}$, and then $(y)R_{x^{-1}}L_{x^{-1}}^{-1} = (y)R_xL_x^{-1}$ by Corollary 2.5. Thus $(y)T_{x^{-1}} = (y)T_x$, and hence $(y)T_x^2 = y$ by (6). Therefore the claim follows from Lemma 3.6.

Theorem 3.8. Let Q be an automorphic loop and $x, y \in Q$. Then

$$x(xy) = (yx)x$$
 if and only if $x^2y = yx^2$.

Proof. We have that

Corollary 3.9. If Q is an uniquely 2-divisible automorphic loop, then it satisfies (27).

Proof. Let $x, y \in Q$ be such that x(xy) = (yx)x. By Theorem 3.8, we have that $x^2y = yx^2$, and then $(x^2)T_y = x^2$. Since $T_y \in Aut(Q)$, it follows that $(x)T_y = x$ by (7).

The next result is immediate from Proposition 2.10 and Corollary 3.9.

Corollary 3.10. Every automorphic loop of odd order satisfies (27).

It is clear that every commutative automorphic loop satisfies (27) and the direct product of two automorphic loops that satisfy (27) also satisfies this condition. Then there are also non-commutative automorphic loops of even whose satisfy (27).

4 Main results

In this section, (Q, *) and (Q', \cdot) are automorphic loops, and $f : (Q, *) \to (Q', \cdot)$ is a halfisomorphism. Furthermore, we consider that Q' satisfies (27), that is, for every $x, y \in Q'$ we have

 $x \cdot (x \cdot y) = (y \cdot x) \cdot x$ if and only if $x \cdot y = y \cdot x$.

Lemma 4.1. Let $x, y \in Q$ be such that $f(x * y) = f(x) \cdot f(y)$ and $f(y * x^{-1}) = f(x^{-1}) \cdot f(y)$. Then $f(x) \cdot f(y) = f(y) \cdot f(x)$.

Proof. By Proposition 2.13, $f(x^{-1}) = f(x)^{-1}$. Then using the definition of half-isomorphisms:

$$\begin{split} f((x*y)*x^{-1}) &\in U = \{f(x) \cdot f(y) \cdot f(x)^{-1}, f(x)^{-1} \cdot (f(x) \cdot f(y))\}, \\ f(x*(y*x^{-1})) &\in V = \{f(x) \cdot (f(x)^{-1} \cdot f(y)), f(x)^{-1} \cdot f(y) \cdot f(x)\}. \end{split}$$

Since $(x * y) * x^{-1} = x * (y * x^{-1})$, we have that $U \cap V \neq \emptyset$. If $f(x) \cdot f(y) \cdot f(x)^{-1} = f(x) \cdot (f(x)^{-1} \cdot f(y))$, then $f(y) \cdot f(x)^{-1} = f(x)^{-1} \cdot f(y)$, and so $f(x) \cdot f(y) = f(y) \cdot f(x)$ by Corollary 2.3. When $f(x) \cdot f(y) \cdot f(x)^{-1} = f(x)^{-1} \cdot f(y) \cdot f(x)$ we have that $f(x) \cdot f(y) = f(y) \cdot f(x)$ by Corollary 3.7. Now consider that $f(x * y * x^{-1}) = f(x)^{-1} \cdot (f(x) \cdot f(y))$. By Lemma 3.1 (b), we have that $(x * y * x^{-1})^2 = (x * y) * (y * x^{-1})$, and then

$$f((x * y * x^{-1})^2) \in \{(f(x) \cdot f(y)) \cdot (f(x)^{-1} \cdot f(y)), (f(x)^{-1} \cdot f(y)) \cdot (f(x) \cdot f(y))\}$$
(29)

Note that $\varphi = L_{f(x)}L_{f(x)^{-1}} \in Aut(Q')$. Then

$$f((x * y * x^{-1})^2) = (f(x * y * x^{-1}))^2 = ((f(y))\varphi)^2 = (f(y)^2)\varphi = f(x)^{-1} \cdot (f(x) \cdot f(y)^2),$$

and putting this in (29), we get

$$\begin{aligned} f(x)^{-1} \cdot (f(x) \cdot f(y)^2) &= (f(x) \cdot f(y)) \cdot (f(x)^{-1} \cdot f(y)) \text{ or } \\ f(x)^{-1} \cdot (f(x) \cdot f(y)^2) &= (f(x)^{-1} \cdot f(y)) \cdot (f(x) \cdot f(y)). \end{aligned}$$

By Corollary 3.2, we get that $f(x) \cdot f(y) = f(y) \cdot f(x)$ in both cases.

Proposition 4.2. *f* is a special half-isomorphism.

Proof. Let $x, y \in Q$ be such that x * y = y * x. It is suffices to prove that $f(x) \cdot f(y) = f(y) \cdot f(x)$ by Proposition 2.12.

The case $f(x*y) = f(y*x) = f(x) \cdot f(y)$ is symmetric to the case $f(y*x) = f(y) \cdot f(x)$, so we only prove the first one. If $f(y * x^{-1}) = f(x^{-1}) \cdot f(y)$, then $f(x) \cdot f(y) = f(y) \cdot f(x)$ by Lemma 4.1. Now consider that $f(y * x^{-1}) = f(y) \cdot f(x^{-1})$. By Corollary 2.3, we have that $\langle x, y \rangle$ is commutative, and then $y * x^{-1} = x^{-1} * y$ and $f(x^{-1} * y) = f(y) \cdot f(x^{-1})$. Since $f(x * y) = f(x) \cdot f(y)$, using (AAIP) and Proposition 2.13 we get

$$f(y^{-1} * x^{-1}) = (f(x * y))^{-1} = f(y)^{-1} \cdot f(x)^{-1} = f(y^{-1}) \cdot f(x^{-1}).$$

By Lemma 4.1, we have that $f(y^{-1}) \cdot f(x^{-1}) = f(x^{-1}) \cdot f(y^{-1})$, and then using (AAIP) and Proposition 2.13 we get that $f(x) \cdot f(y) = f(y) \cdot f(x)$.

Since f is special, by Proposition 2.12 we have, for all $x, y \in Q$:

If
$$x * y = y * x$$
, then $f(x) \cdot f(y) = f(y) \cdot f(x)$, (30)

If
$$f(x * y) = f(x) \cdot f(y)$$
, then $f(y * x) = f(y) \cdot f(x)$, (31)

If
$$f(x * y) = f(y) \cdot f(x)$$
, then $f(y * x) = f(x) \cdot f(y)$. (32)

Let L, L' be flexible loops. A mapping f from L into L' is called a semi-homomorphism if f(xyx) = f(x)f(y)f(x), for all $x, y \in L$.

Proposition 4.3. *f* is a semi-homomorphism.

Proof. Let $x, y \in Q$. We have two cases: (i) $f(x * y) = f(x) \cdot f(y)$. Since f is special, we have $f(y * x) = f(y) \cdot f(x)$ by (31). Then:

$$\begin{aligned} &f((x * y) * x) \in \{f(x) \cdot f(y) \cdot f(x), f(x) \cdot (f(x) \cdot f(y))\}, \\ &f(x * (y * x)) \in \{f(x) \cdot f(y) \cdot f(x), (f(y) \cdot f(x)) \cdot f(x)\}. \end{aligned}$$

Since Q is flexible, either $f(x * y * x) = f(x) \cdot f(y) \cdot f(x)$ or

$$f(x * y * x) = f(x) \cdot (f(x) \cdot f(y)) = (f(y) \cdot f(x)) \cdot f(x).$$
(33)

In the second case, we get $f(x) \cdot f(y) = f(y) \cdot f(x)$ since Q' satisfies (27). Putting this in (33), we get $f(x * y * x) = f(x) \cdot f(y) \cdot f(x)$.

(ii) $f(x * y) = f(y) \cdot f(x)$. This case is analogous to (i).

Proposition 4.4. *Q* satisfies (27).

Proof. Let $x, y \in Q$ be such that x * (x * y) = (y * x) * x. We have two cases: (i) $f(x * y) = f(x) \cdot f(y)$. By Proposition 4.3, we have that

$$f((x * y) * x) = (f(x) \cdot f(y)) \cdot f(x) = f(x) \cdot (f(y) \cdot f(x)) = f(x * (y * x)).$$

Since f is special, we get that $f(x*(x*y)) = f(x) \cdot (f(x) \cdot f(y))$ and $f((y*x)*x) = (f(y) \cdot f(x)) \cdot f(x)$ by (31). Since x * (x * y) = (y * x) * x, we get $f(x) \cdot (f(x) \cdot f(y)) = (f(y) \cdot f(x)) \cdot f(x)$, and then $f(x) \cdot f(y) = f(y) \cdot f(x)$ since Q' satisfies (27). Therefore x * y = y * x. (ii) $f(x * y) = f(y) \cdot f(x)$. This case is analogous to (i). **Definition 4.5.** ([6]) Let $x, y, z \in Q$. The triple (x, y, z) is called a *GG-triple* of f if (i) $f(x * y) = f(x) \cdot f(y) \neq f(y) \cdot f(x)$, (ii) $f(x * z) = f(z) \cdot f(x) \neq f(x) \cdot f(z)$.

Remark 4.6. In Example 2.11, (3, 2, 6) is a GG-triple.

The next lemma is a well-known result about loops. A proof of it can be found in [10, Lemma 1].

Lemma 4.7. Let L be a loop and H, K be subloops of L. If $L = H \cup K$, then either H = L or K = L.

Let S be a non empty subset of a loop L. The *commutant* of S is the set

$$C_L(S) = \{ x \in L \mid xy = yx, \forall y \in S \}.$$

If L is an automorphic loop, then $C_L(S)$ is a subloop of L [8, Proposition 2.10].

Proposition 4.8. Let Q_1 and Q_2 be automorphic loops and $\varphi : Q_1 \to Q_2$ be a special halfisomorphism. If φ has no GG-triple, then φ is trivial.

Proof. Let $A = \{x \in Q_1 | \varphi(x * u) = \varphi(x) \cdot \varphi(u), \forall u \in Q_1\}$ and $B = \{y \in Q_1 | \varphi(y * u) = \varphi(u) \cdot \varphi(y), \forall u \in Q_1\}$. Since φ has no GG-triple, then $Q_1 = A \cup B$.

For $x \in A$ and $y \in B$, we have by definition that $\varphi(x * y) = \varphi(x) \cdot \varphi(y) = \varphi(y * x)$, and then x * y = y * x. Thus $A \subset C_{Q_1}(y)$ and $B \subset C_{Q_1}(x)$, for all $x \in A$ and $y \in B$. Since $Q_1 = A \cup B$, it follows by Lemma 4.7 that, for every $(x, y) \in (A, B)$, either $Q_1 = C_{Q_1}(x)$ or $Q_1 = C_{Q_1}(y)$.

If there exists $x \in A$ such that $C_{Q_1}(x) \neq Q_1$, then $C_{Q_1}(y) = Q_1$, for all $y \in B$, and so $\varphi(y * u) = \varphi(u) \cdot \varphi(y) = \varphi(y) \cdot \varphi(u)$, for all $u \in Q_1$ and $y \in B$, since φ is special. Thus $B \subset A$, and we get that $Q_1 = A$. Hence φ is an isomorphism. Similarly, when $C_{Q_1}(x) = Q_1$, for all $x \in A$, we get that $A \subset B$, and then $Q_1 = B$ and φ is an anti-isomorphism. \Box

For $x \in Q$, define the mapping $\phi_x = R_x L_{x^{-1}}$. Then $(u)\phi_x = x^{-1} * u * x = u^x$, for all $u \in Q$. Since $(1)\phi_x = 1$, we have that $\phi_x \in Aut(Q)$, and then, for all $u, v \in Q$:

$$(u*v)^x = u^x * v^x \tag{34}$$

Lemma 4.9. Let $x, y \in Q$.

(a) If x * y = y * x, then $f(y^x) = f(y)^{f(x)} = f(y)^{f(x)^{-1}}$. (b) If $x * y \neq y * x$ and $f(x * y) = f(x) \cdot f(y)$, then $f(y^x) = f(y)^{f(x)}$ and $f(y^{x^{-1}}) = f(y)^{f(x)^{-1}}$. (c) If $x * y \neq y * x$ and $f(x * y) = f(y) \cdot f(x)$, then $f(y^x) = f(y)^{f(x)^{-1}}$ and $f(y^{x^{-1}}) = f(y)^{f(x)}$.

Proof. Using the definition of half-isomorphisms, we get

$$f(x^{-1} * (x * y)) \in U = \{f(x)^{-1} \cdot f(x * y), f(x * y) \cdot f(x)^{-1}\},$$
(35)

$$f(x * (x^{-1} * y)) \in V = \{f(x) \cdot f(x^{-1} * y), f(x^{-1} * y) \cdot f(x)\}.$$
(36)

Note that $x^{-1} * (x * y) = x * (x^{-1} * y)$, and then $U \cap V \neq \emptyset$.

(a) Since f is special, we have that $f(x * y) = f(x) \cdot f(y) = f(y) \cdot f(x)$. Furthermore, $\langle x, y \rangle$ and $\langle f(x), f(y) \rangle$ are commutative by Corollary 2.3. Hence $f(x)^{-1} \cdot f(x * y) = f(x * y) \cdot f(x)^{-1}$ and by (35):

$$f(x^{-1} * y * x) = f(x^{-1} * (x * y)) = f(x)^{-1} \cdot f(y) \cdot f(x) = f(x) \cdot (f(x)^{-1} \cdot f(y)) = f(x) \cdot f(y) \cdot f(x)^{-1}.$$

(b) Since f is special, we have that $f(y * x^{-1}) = f(y) \cdot f(x)^{-1}$ by Lemma 4.1 and (30), and then $f(x^{-1} * y) = f(x)^{-1} \cdot f(y)$ by (31). Putting this in (35) and (36), we get

$$\begin{split} f(x^{-1}*(x*y)) &\in U = \{f(x)^{-1} \cdot (f(x) \cdot f(y)), f(x) \cdot f(y) \cdot f(x)^{-1}\}, \\ f(x*(x^{-1}*y)) &\in V = \{f(x) \cdot (f(x)^{-1} \cdot f(y)), f(x)^{-1} \cdot f(y) \cdot f(x)\}. \end{split}$$

Since $f(x) \cdot f(y) \neq f(y) \cdot f(x)$, we have that $f(x)^{-1} \cdot (f(x) \cdot f(y)) \neq f(x)^{-1} \cdot f(y) \cdot f(x)$ and $f(x) \cdot f(y) \cdot f(x)^{-1} \neq f(x) \cdot (f(x)^{-1} \cdot f(y))$. Furthermore, $f(x) \cdot f(y) \cdot f(x)^{-1} \neq f(x)^{-1} \cdot f(y) \cdot f(x)$ by Corollary 3.7. Thus

$$f(x * (x^{-1} * y)) = f(x) \cdot (f(x)^{-1} \cdot f(y)) = f(x)^{-1} \cdot (f(x) \cdot f(y)) = f(x^{-1} * (x * y))$$

and hence $f(y^x) = f(y)^{f(x)}$ and $f(y^{x^{-1}}) = f(y)^{f(x)^{-1}}$ by (31). (c) Using (AAIP), we get $f(y^{-1}x^{-1}) = f(x)^{-1} \cdot f(y)^{-1}$. Since f is special, we have $f(y) \cdot f(x)^{-1} \neq f(x)^{-1} \cdot f(y)$ by (30) and Corollary 2.3, and then $f(x^{-1} * y) = f(y) \cdot f(x)^{-1}$ by Lemma 4.1. Putting this in (35) and (36), we get

$$\begin{split} f(x^{-1}*(x*y)) &\in U = \{f(x)^{-1} \cdot f(y) \cdot f(x), (f(y) \cdot f(x)) \cdot f(x)^{-1}\}, \\ f(x*(x^{-1}*y)) &\in V = \{f(x) \cdot f(y) \cdot f(x)^{-1}, (f(y) \cdot f(x)^{-1}) \cdot f(x)\}. \end{split}$$

Since $f(x) \cdot f(y) \neq f(y) \cdot f(x)$, we have that $f(x)^{-1} \cdot f(y) \cdot f(x) \neq (f(y) \cdot f(x)^{-1}) \cdot f(x)$ and $(f(y) \cdot f(x)) \cdot f(x)^{-1} \neq f(x) \cdot f(y) \cdot f(x)^{-1}$. Furthermore, $f(x)^{-1} \cdot f(y) \cdot f(x) \neq f(x) \cdot f(y) \cdot f(x)^{-1}$ by Corollary 3.7. Thus

$$f(x * (x^{-1} * y)) = (f(y) \cdot f(x)^{-1}) \cdot f(x) = (f(y) \cdot f(x)) \cdot f(x)^{-1} = f(x^{-1} * (x * y)),$$

and hence $f(y^x) = f(y)^{f(x)^{-1}}$ and $f(y^{x^{-1}}) = f(y)^{f(x)}$ by (32).

Theorem 4.10. Let (Q, *) and (Q', \cdot) be automorphic loops and $f : (Q, *) \to (Q', \cdot)$ be a halfisomorphism. If (Q', \cdot) satisfies (27), then f is trivial.

Proof. Suppose by contradiction that f has a GG-triple (x, y, z). Then $x * y \neq y * x$, $x * z \neq z * x$, $f(x * y) = f(x) \cdot f(y)$ and $f(x * z) = f(z) \cdot f(x)$. By Lemma 4.9, we have that $f(y^x) = f(y)^{f(x)}$, $f(y^{x^{-1}}) = f(y)^{f(x)^{-1}}$, $f(z^x) = f(z)^{f(x)^{-1}}$ and $f(z^{x^{-1}}) = f(z)^{f(x)}$. Using (34), we get

$$f(y^{x}) \cdot f(z^{x^{-1}}) \cdot f(y^{x}) = (f(y) \cdot f(z) \cdot f(y))^{f(x)} \text{ and } f(y^{x^{-1}}) \cdot f(z^{x}) \cdot f(y^{x^{-1}}) = (f(y) \cdot f(z) \cdot f(y))^{f(x)^{-1}}.$$

Since f is a semi-homomophism, it follows that

$$f(y^{x} * z^{x^{-1}} * y^{x}) = (f(y * z * y))^{f(x)},$$
(37)

$$f(y^{x^{-1}} * z^x * y^{x^{-1}}) = (f(y * z * y))^{f(x)^{-1}}.$$
(38)

If $f(x * (y * z * y)) = f(x) \cdot f(y * z * y)$, then $f(y^x * z^x * y^x) = f((y * z * y)^x) = (f(y * z * y))^{f(x)}$ by (34) and Lemma 4.9, and so $y^x * z^x * y^x = y^x * z^{x^{-1}} * y^x$ by (37). Thus $z^x = z^{x^{-1}}$, and hence x * z = z * x by Proposition 4.4 and Corollary 3.7, which is a contradiction. When $f(x * (y * z * y)) = f(y * z * y) \cdot f(x)$ we have that $f(y^{x^{-1}} * z^{x^{-1}} * y^{x^{-1}}) = f((y * z * y)^{x^{-1}}) = (f(y * z * y))^{f(x)^{-1}}$ by (34) and Lemma 4.9, and so $y^{x^{-1}} * z^{x^{-1}} * y^{x^{-1}} = y^{x^{-1}} * z^x * y^{x^{-1}}$ by (38). Thus $z^x = z^{x^{-1}}$, and hence x * z = z * x by Proposition 4.4 and Corollary 3.7, which is a contradiction.

Therefore f has no GG-triple and the claim follows by Proposition 4.8.

As direct consequences of Theorem 4.10 and Corollaries 3.9 and 3.10, we have the following results.

Corollary 4.11. Every half-isomorphism between uniquely 2-divisible automorphic loops is trivial.

Corollary 4.12. Every half-isomorphism between automorphic loops of odd order is trivial.

5 Open problems

We proved that every half-isomorphism between loops in the class of automorphic loops satisfying (27) is special, as was demonstrated earlier for automorphic Moufang loops [6, 9]. Examples of nontrivial half-isomorphisms between automorphic loops exist (cf. [10]), but so far all of them are special. We propose the conjecture that this can be generalized to all automorphic loops:

Conjecture 5.1. Every half-isomorphism between automorphic loops is special.

References

- [1] R.H. Bruck, A Survey of Binary Systems, Springer, 1971.
- [2] R.H. Bruck, L.J. Paige, Loops whose inner mappings are automorphisms, Ann. Math., 63, (1956), 308–323.
- [3] S. Gagola III, M.L. Merlini Giuliani, On half-automorphisms of certain Moufang Loops with even order, Journal of Algebra, 386, (2013), 131–141.
- [4] S. Gagola III, M.L. Merlini Giuliani, Half-isomorphisms of Moufang loops of odd order, Journal of Algebra and its Application, 11, no. 5 (2012), 194-199.
- [5] The GAP Group, GAP Groups, Algorithms, and Programming, Version 4.10.1, 2019. http://www.gap-system.org
- [6] A. Grishkov, M.L. Merlini Giuliani, M. Rasskazova, L. Sabinina, Half-Isomorphisms of finite Automorphic Moufang Loops, Communications in Algebra, 44, (2016), 4252–4261.
- [7] K. W. Johnson, M.K. Kinyon, G.P. Nagy, and P. Vojtěchovský, Searching for small simple automorphic loops, LMS J. Comput. Math., 14, (2011), 200–213,
- [8] M. K. Kinyon, K. Kunen, J.D. Phillips, P. Vojtěchovský. The structure of automorphic loops. Trans. Amer. Math. Soc., 368, (2016), 8901–8927.
- M. Kinyon, I. Stuhl, P. Vojtěchovský, *Half-isomorphisms of Moufang loops*, Journal of Algebra, 450, (2016), 152–161.
- [10] M.L. Merlini Giuliani, G. Souza dos Anjos, Half-isomorphisms of dihedral automorphic loops, Communications in Algebra, 48, (2020), no. 3, 1150–1162.
- [11] M.L. Merlini Giuliani, G. Souza dos Anjos, Lie automorphic loops under halfautomorphisms, Journal of Algebra and its Application, 19, (2020), no. 11, 2050221.
- [12] M.L. Merlini Giuliani, P. Plaumann, L. Sabinina, *Half-automorphisms of Cayley-Dickson loops*, In: Falcone, G., ed., Lie Groups, Differential Equations, and Geometry. Cham: Springer, (2017), 109–125.

- [13] W. McCune, Prover9 and Mace4, http://www.cs.unm.edu/~mccune/prover9, 2005-2010.
- [14] G.P. Nagy, P. Vojtěchovský. LOOPS: Computing with quasigroups and loops in GAP, version 3.4.0, package for GAP, https://cs.du.edu/~petr/loops/
- [15] H.O. Pflugfelder, Quasigroups and Loops: Introduction, Sigma Series in Pure Math., 7, Heldermann, 1990.
- [16] W.R. Scott, Half-homomorphisms of groups, Proc. Amer. Math. Soc., 8, (1957), 1141– 1144.