VARIATIONAL INEQUALITIES ON UNBOUNDED DOMAINS FOR ZERO-SUM SINGULAR-CONTROLLER VS. STOPPER GAMES

ANDREA BOVO, TIZIANO DE ANGELIS, AND ELENA ISSOGLIO

ABSTRACT. We study a class of zero-sum games between a singular-controller and a stopper over finite-time horizon. The underlying process is a multi-dimensional (locally non-degenerate) controlled stochastic differential equation (SDE) evolving in an unbounded domain. We prove that such games admit a value and provide an optimal strategy for the stopper. The value of the game is shown to be the maximal solution, in a suitable Sobolev class, of a variational inequality of 'minmax' type with obstacle constraint and gradient constraint. Although the variational inequality and the game are solved on an unbounded domain we do not require boundedness of either the coefficients of the controlled SDE or of the cost functions in the game.

1. INTRODUCTION

We consider a class of zero-sum games on a finite-time horizon [0,T] between a controller and a stopper. The underlying stochastic dynamics $X^{[n,\nu]}$ is given by a multi-dimensional, singularly controlled, stochastic differential equation of the form

(1.1)
$$dX_t^{[n,\nu]} = b(X_t^{[n,\nu]})dt + \sigma(X_t^{[n,\nu]})dW_t + n_t d\nu_t$$

where W is a multi-dimensional Brownian motion and the control pair $(n_t, \nu_t)_{t \in [0,T]}$ is given by a unitary vector $n_t(\omega) \in \mathbb{R}^d$ and a real-valued, right-continuous, increasing process $\nu_t(\omega)$. The stopper decides when the game ends and receives from the controller an amount depending on a running payoff h(t, x), a terminal payoff g(t, x) and a cost f(t, x) per unit of control exerted, see Section 3 for details. Using a blend of analytical and probabilistic techniques we prove that the game admits a value v which is the maximal solution in a suitable Sobolev space of the variational inequalities:

(1.2)
$$\min\left\{\max\left\{\partial_t u + \mathcal{L}u - ru + h, g - u\right\}, f - |\nabla u|_d\right\} = 0,\\ \max\left\{\min\left\{\partial_t u + \mathcal{L}u - ru + h, f - |\nabla u|_d\right\}, g - u\right\} = 0, \end{cases}$$

a.e. in $[0,T) \times \mathbb{R}^d$ with terminal condition u(T,x) = g(T,x). Here, \mathcal{L} is the infinitesimal generator of the uncontrolled SDE, $r \geq 0$ is a constant discount rate for the game's payoff and $|\cdot|_d$ is the Euclidean norm in \mathbb{R}^d . We also provide an optimal stopping rule for the stopper, which provides a best response against *any* admissible control for the controller.

The two variational problems in (1.2) have not received much attention in the literature and they pose a number of challenges. The first obvious one is that swapping the order of 'min' and 'max' is non-trivial and it relates in some sense to proving the equivalence between the so-called upper and lower value of the game. Secondly, a solution of the variational problem is subject to two hard constraints: an obstacle constraint $u \ge g$ and a gradient constraint $|\nabla u|_d \le f$. Thirdly, we solve the problem on an *unbounded domain* but without imposing boundedness of the coefficients of the SDE or of the payoff functions, and without requiring uniform ellipticity of the matrix $\sigma\sigma^{\top}$ in the whole space. These seem important technical improvements even when compared to variational inequalities on unbounded domains for singular control problems (e.g., Chow et al. [11], Soner and

Date: October 31, 2023.

²⁰²⁰ Mathematics Subject Classification. 91A05, 91A15, 60G40, 93E20, 49J40, 35K58.

Key words and phrases. zero-sum stochastic games, singular control, optimal stopping, controlled diffusions, variational inequalities, obstacle problems, gradient constraints, penalisation methods, unbounded domains.

Shreve [34, 35], Menaldi and Taksar [33] and Zhu [40]) or optimal stopping games (e.g., Bensoussan and Friedman [5], Friedman [18] and Stettner [36]).

Our method of proof builds upon penalisation techniques that address simultaneously the two hard constraints in (1.2). We find bounds on the Sobolev norm of the solution of the penalised (semilinear) PDE problem, uniformly with respect to the penalisation parameters, thanks to analytical techniques rooted in early work by Evans [14] and new probabilistic tricks developed ad-hoc in our framework. Indeed, it turns out that the co-existence of two hard constraints in (1.2), the 'min-max' structure of the problem, its parabolic nature and unboundedness of the domain make the use of purely analytical ideas as in Evans [14] not sufficient to provide the necessary bounds (see also the references given in the previous paragraph and more recent work by Hynd [25] and Kelbert and Moreno-Franco [28], for comparison). In the process of obtaining our main result (Theorem 3.3) we also contribute a detailed proof of the existence and uniqueness of the solution for the penalised problem (Theorem 4.11 for bounded domain and Theorem 5.3+Proposition 5.4 for unbounded domain), which hopefully will serve as a useful reference for future work in the field. Finally, the existence of an optimal stopping time τ_* is interesting in its own right as it may enable free-boundary techniques for the study of the optimal strategy of the controller. Thanks to the structure of τ_* the game can be turned into a singular control problem with absorption along the boundary of the (unknown) contact set $\{v = q\}$. Then it may be possible to use ideas from, e.g., De Angelis and Ekström [12] to determine the optimal control and a saddle point in the game.

The probabilistic study of so-called controller-stopper zero-sum games originates from work by Maitra and Sudderth [32]. A 'gambler' selects the distribution of a discrete-time process $(X_n)_{n \in \mathbb{N}}$ and a stopper ends the game at a stopping time $\tau \in \mathbb{N}$ of her choosing and pays an amount $u(X_{\tau})$ to the gambler, for some bounded function u. The problem was then cast in a continuoustime framework by Karatzas and Sudderth [26] who consider one-dimensional Itô diffusions whose drift and diffusion coefficients are chosen by the controller. They obtain (almost explicit) optimal strategies for both players using methods based on the general theory of one-dimensional linear diffusions. We as diffusion coefficient is controlled, and finds that the game admits a value which is not continuously differentiable as function of the initial state of the process.

Prior to [32], Bensoussan and Friedman [4] solved, via penalisation methods, nonlinear variational inequalities linked to zero-sum games in which each player uses both a control and a stopping time.

Following those early contributions, the literature on zero-sum (and to some extent also nonzerosum) controller-stopper games has grown steadily. A wide variety of methods has been deployed spanning, for example, martingale theory (Karatzas and Zamfirescu [27]), backward stochastic differential equations (e.g., Hamadène [21], Choukroun et al. [10]) and solution of variational problems via viscosity theory (e.g., Bayraktar and Huang [2] and Bayraktar and Young [3]). A common denominator of those papers is that the controller uses so-called 'classical' controls, i.e., progressively measurable maps $(t, \omega) \mapsto \alpha_t(\omega)$ that enter the drift and diffusion coefficient, b and σ of the controlled SDE in the form

$$\mathrm{d}X_t^\alpha = b(X_t^\alpha, \alpha_t)\mathrm{d}t + \sigma(X_t^\alpha, \alpha_t)\mathrm{d}W_t.$$

From an analytical point of view, those games are connected to Hamilton-Jacobi-Bellman (HJB) equations with obstacle constraint but without gradient constraint, hence different from (1.2).

Much less attention instead has been devoted to the study of games in which the controller can adopt singular controls as in (1.1). The resulting dynamics is 'singular' in the sense that the mapping $t \mapsto \nu_t(\omega)$ need not be absolutely continuous with respect to the Lebesgue measure (nor even continuous at all). A notable contribution to this strand of the literature was given by Hernandez-Hernandez et al. [23] who consider a zero-sum game in which $X^{[n,\nu]}$ is real-valued. They provide a general verification theorem and explicit optimal strategies for both players in some examples. They also show that the value function of the game need not be smooth if the stopping payoff is not continuously differentiable. The methods in [23] rely crucially on the one-dimensional set-up that allows to solve the variational problem with ordinary differential equations and an educated guess on the structure of the optimal strategies. Here instead we develop a general theory for multi-dimensional state-dynamics that leads us to consider PDEs. The introduction of [23] also explains numerous important applications of zero-sum games like the ones we consider here. Such applications include models for a central bank controlling exchange rates up to the time of a possible political veto and models for the control of inflation. We refer the interested reader to the Introduction of [23] for more insight on applications.

Our paper is organised as follows. We set out the notation in Section 2. Then, in Section 3 we cast our problem, state our main assumptions and the main result (Theorem 3.3). In Section 4 we introduce and solve a penalised problem on bounded domain. In Section 5 we let the domain become unbounded and solve the corresponding penalised problem. Finally, in Section 6 we prove our main result. A technical appendix completes the paper.

2. NOTATION

Fix $d, d' \in \mathbb{N}$, $d \leq d'$ and $T \in (0, \infty)$. Given $u \in \mathbb{R}^d$ we let $|u|_d$ be its Euclidean norm. For vectors $u, v \in \mathbb{R}^d$ their scalar product is denoted by $\langle u, v \rangle$. Given a matrix $M \in \mathbb{R}^{d \times d'}$, with entries M_{ij} , $i = 1, \ldots d, j = 1, \ldots d'$, we denote its norm by

$$|M|_{d \times d'} \coloneqq \left(\sum_{i=1}^{d} \sum_{j=1}^{d'} M_{ij}^2\right)^{1/2}$$

and, if d = d', we let $\operatorname{tr}(M) \coloneqq \sum_{i=1}^{d} M_{ii}$. The *d*-dimensional *open* ball centred in 0 with radius *m* is denoted by B_m and the state space in our problem is going to be

$$\mathbb{R}^{d+1}_{0,T} := [0,T] \times \mathbb{R}^d.$$

Finally, given a bounded set A we denote by \overline{A} its closure.

For a smooth function $f : \mathbb{R}_{0,T}^{d+1} \to \mathbb{R}$ we denote its partial derivatives by $\partial_t f$, $\partial_{x_i} f$, $\partial_{tx_j} f$, $\partial_{x_ix_j} f$, for $i, j = 1, \ldots d$. We will also use $f_t = \partial_t f$, $f_{x_i} = \partial_{x_i} f$, $f_{tx_i} = \partial_{tx_i} f$ and $f_{x_ix_j} = \partial_{x_ix_j} f$ to simplify long expressions. By ∇f we intend the spatial gradient, i.e., $\nabla f = (\partial_{x_1} f, \ldots \partial_{x_d} f)$, and by $D^2 f$ the spatial Hessian matrix with entries $\partial_{x_ix_j} f$ for $i, j = 1, \ldots d$. As usual $C_c^{\infty}(\mathbb{R}_{0,T}^{d+1})$ is the space of functions with compact support and infinitely many continuous derivatives. Continuous functions on a domain D are denoted by C(D). For an open bounded set $\mathcal{O} \subset \mathbb{R}_{0,T}^{d+1}$ we let $C^0(\overline{\mathcal{O}})$ be the space of continuous functions $f : \overline{\mathcal{O}} \to \mathbb{R}$ equipped with the supremum norm

(2.1)
$$\|f\|_{C^0(\overline{\mathcal{O}})} \coloneqq \sup_{(t,x)\in\overline{\mathcal{O}}} |f(t,x)|$$

Analogously, $C^0(\mathbb{R}^{d+1}_{0,T})$ is the space of bounded and continuous functions $f: \mathbb{R}^{d+1}_{0,T} \to \mathbb{R}$ equipped with the norm $\|f\|_{\infty} \coloneqq \|f\|_{C^0(\mathbb{R}^{d+1}_{0,T})}$ as in (2.1) but with $\overline{\mathcal{O}}$ replaced by $\mathbb{R}^{d+1}_{0,T}$.

For bounded $\mathcal{O} \subset \mathbb{R}^{d+1}_{0,T}$, we consider the following function spaces:

- $C^{0,1}(\overline{\mathcal{O}})$ be the class of continuous functions with $\partial_{x_i} f \in C(\overline{\mathcal{O}})$ for $i = 1, \ldots, d$;
- $C^{1,2}(\overline{\mathcal{O}})$ be the class of continuous functions with $\partial_t f, \partial_{x_i} f, \partial_{x_i x_j} f \in C(\overline{\mathcal{O}})$ for $i, j = 1, \dots, d$;
- $C^{1,3}(\overline{\mathcal{O}})$ be the class of continuous functions with

 $\partial_t f, \, \partial_{x_i} f, \, \partial_{x_i x_j} f, \, \partial_{x_i x_j x_k} f, \, \partial_{t x_i} f \in C(\overline{\mathcal{O}})$

for $i, j, k = 1, \ldots, d$ (notice the mixed derivatives $\partial_{tx_i} f$).

The above definitions extend obviously to continuously differentiable functions on $\mathbb{R}^{d+1}_{0,T}$.

Let $d(z_1, z_2) = (|t - s| + |x - y|_d^2)^{\frac{1}{2}}$ be the parabolic distance between points $z_1 = (t, x)$ and $z_2 = (s, y)$ in $\mathbb{R}^{d+1}_{0,T}$. For a fixed $\alpha \in (0, 1)$ and a continuous function $f : \overline{\mathcal{O}} \to \mathbb{R}$ we set [19, p. 61]

$$\|f\|_{C^{\alpha}(\overline{\mathcal{O}})} \coloneqq \|f\|_{C^{0}(\overline{\mathcal{O}})} + \sup_{\substack{z_{1}, z_{2} \in \overline{\mathcal{O}} \\ z_{1} \neq z_{2}}} \frac{|f(z_{1}) - f(z_{2})|}{\mathrm{d}^{\alpha}(z_{1}, z_{2})}$$

We say that $f \in C^{\alpha}(\overline{\mathcal{O}})$ if $f \in C^{0}(\overline{\mathcal{O}})$ and $||f||_{C^{\alpha}(\overline{\mathcal{O}})} < \infty$. We work with the following norms, defined for sufficiently smooth functions f:

$$\begin{split} \|f\|_{C^{0,1,\alpha}(\overline{\mathcal{O}})} &\coloneqq \|f\|_{C^{\alpha}(\overline{\mathcal{O}})} + \sum_{i=1}^{d} \|\partial_{x_{i}}f\|_{C^{\alpha}(\overline{\mathcal{O}})}; \\ \|f\|_{C^{1,2,\alpha}(\overline{\mathcal{O}})} &\coloneqq \|f\|_{C^{0,1,\alpha}(\overline{\mathcal{O}})} + \|\partial_{t}f\|_{C^{\alpha}(\overline{\mathcal{O}})} + \sum_{i,j=1}^{d} \|\partial_{x_{i}x_{j}}f\|_{C^{\alpha}(\overline{\mathcal{O}})}; \\ \|f\|_{C^{1,3,\alpha}(\overline{\mathcal{O}})} &\coloneqq \|f\|_{C^{1,2,\alpha}(\overline{\mathcal{O}})} + \sum_{i=1}^{d} \|\partial_{tx_{i}}f\|_{C^{\alpha}(\overline{\mathcal{O}})} + \sum_{i,j,k=1}^{d} \|\partial_{x_{i}x_{j}x_{k}}f\|_{C^{\alpha}(\overline{\mathcal{O}})}; \end{split}$$

For $(j,k) \in \{(0,0); (0,1); (1,2); (1,3)\}$ and bounded \mathcal{O} let us define

$$C^{j,k,\alpha}(\overline{\mathcal{O}}) \coloneqq \left\{ f \in C^{j,k}(\overline{\mathcal{O}}) \middle| \, \|f\|_{C^{j,k,\alpha}(\overline{\mathcal{O}})} < \infty \right\},$$

$$C^{j,k,\alpha}_{\ell oc}(\mathbb{R}^{d+1}_{0,T}) \coloneqq \left\{ f \in C^{j,k}(\mathbb{R}^{d+1}_{0,T}) \middle| \, f \in C^{j,k,\alpha}(\overline{\mathcal{O}}) \text{ for all bounded } \mathcal{O} \subset \mathbb{R}^{d+1}_{0,T} \right\}.$$

For B and B' open balls in \mathbb{R}^d and $S \in [0,T)$, let $\mathcal{O}_B := [0,T) \times B$ and $\mathcal{O}_{S,B'} := [0,S) \times B'$. We denote $C_{Loc}^{j,k,\alpha}(\mathcal{O}_B)$ the class of functions $f \in C(\overline{\mathcal{O}}_B)$ such that $f \in C^{j,k,\alpha}(\overline{\mathcal{O}}_{S,B'})$ for all S < T and B' such that $\overline{B'} \subset B$. Finally, we let

$$C_{Loc}^{j,k,\alpha}(\mathbb{R}^{d+1}_{0,T}) \coloneqq \left\{ f \in C(\mathbb{R}^{d+1}_{0,T}) \mid f \in C_{Loc}^{j,k,\alpha}(\mathcal{O}_B) \text{ for all open balls } B \subset \mathbb{R}^d \right\}.$$

Notice that, the derivatives of functions in $C_{loc}^{j,k,\alpha}(\mathbb{R}^{d+1}_{0,T})$ are Hölder continuous on $\overline{\mathcal{O}}_B$ for any ball $B \subset \mathbb{R}^d$. Instead, the derivatives of functions in $C_{Loc}^{j,k,\alpha}(\mathcal{O}_B)$ need not be continuous along the parabolic boundary of \mathcal{O}_B and derivatives of functions in $C_{Loc}^{j,k,\alpha}(\mathbb{R}^{d+1}_{0,T})$ may be discontinuous at T. To simplify long formulae, sometimes we use the notations:

$$\|\nabla f\|_{C^{0}(\overline{\mathcal{O}})} := \left(\sum_{i=1}^{d} \|\partial_{x_{i}}f\|_{C^{0}(\overline{\mathcal{O}})}^{2}\right)^{\frac{1}{2}} \text{ and } \|D^{2}f\|_{C^{0}(\overline{\mathcal{O}})} := \left(\sum_{i,j=1}^{d} \|\partial_{x_{i}x_{j}}f\|_{C^{0}(\overline{\mathcal{O}})}^{2}\right)^{\frac{1}{2}}.$$

For $p \in [1, \infty]$ we say that $f \in L^p_{\ell oc}(\mathbb{R}^{d+1}_{0,T})$ if $f \in L^p(\mathcal{O})$ for any bounded $\mathcal{O} \subset \mathbb{R}^{d+1}_{0,T}$. Denote by $\mathcal{D}^{1,2}_{\mathcal{O}}$ the class of functions $f \in L^p_{\ell oc}(\mathbb{R}^{d+1}_{0,T})$ whose partial derivatives $\partial_t f$, $\partial_{x_i} f$, $\partial_{x_i x_j} f$ exist in the weak sense on \mathcal{O} , for $i, j = 1, \ldots d$, and let

$$\|f\|_{W^{1,2,p}(\mathcal{O})} \coloneqq \|f\|_{L^{p}(\mathcal{O})} + \|\partial_{t}f\|_{L^{p}(\mathcal{O})} + \sum_{i=1}^{d} \|\partial_{x_{i}}f\|_{L^{p}(\mathcal{O})} + \sum_{i,j=1}^{d} \|\partial_{x_{i}x_{j}}f\|_{L^{p}(\mathcal{O})}.$$

Then we define $W^{1,2,p}(\mathcal{O}) \coloneqq \{ f \in \mathcal{D}_{\mathcal{O}}^{1,2} \mid ||f||_{W^{1,2,p}(\mathcal{O})} < \infty \}$ and

$$W_{\ell oc}^{1,2,p}(\mathbb{R}^{d+1}_{0,T}) \coloneqq \left\{ f \in L^p_{\ell oc}(\mathbb{R}^{d+1}_{0,T}) \middle| f \in W^{1,2,p}(\mathcal{O}), \, \forall \mathcal{O} \subseteq \mathbb{R}^{d+1}_{0,T}, \mathcal{O} \text{ bounded} \right\}.$$

For $\alpha = 1 - \frac{d+2}{p}$ and p > d+2, and for any bounded $\mathcal{O} \subset \mathbb{R}^{d+1}_{0,T}$, we have (2.2) $C^{1,2}(\overline{\mathcal{O}}) \subset W^{1,2,p}(\mathcal{O}) \hookrightarrow C^{0,1,\alpha}(\overline{\mathcal{O}}),$ where the first inclusion is obvious and the second one is a compact Sobolev embedding ([16, eq. (E.9)] or [30, Exercise 10.1.14]).

3. Setting and Main Results

Let $(\Omega, \mathcal{F}, \mathsf{P})$ be a complete probability space, $\mathbb{F} = (\mathcal{F}_s)_{s \in [0,\infty)}$ a right-continuous filtration completed by the P-null sets and $(W_s)_{s \in [0,\infty)}$ a \mathbb{F} -adapted, d'-dimensional Brownian motion. Fix $T \in (0,\infty)$. For $t \in [0,T]$, we denote

$$\mathcal{T}_t := \{ \tau \mid \tau \text{ is } \mathbb{F} \text{-stopping time with } \tau \in [0, T-t], \text{ P-a.s.} \}$$

and we let \mathcal{A}_t be the class of processes

$$\mathcal{A}_{t} \coloneqq \left\{ (n,\nu) \middle| \begin{array}{l} (n_{s})_{s \in [0,\infty)} \text{ is progressively measurable, } \mathbb{R}^{d} \text{-valued,} \\ \text{with } |n_{s}|_{d} = 1, \text{ P-a.s. for all } s \in [0,\infty); \\ (\nu_{s})_{s \in [0,\infty)} \text{ is } \mathbb{F} \text{-adapted, real-valued, non-decreasing and} \\ \text{right-continuous with } \nu_{0-} = 0, \text{ P-a.s., and } \mathbb{E}[|\nu_{T-t}|^{2}] < \infty \end{array} \right\}.$$

The notation $\nu_{0-} = 0$ accounts for a possible jump of ν at time zero. For a given pair $(n, \nu) \in \mathcal{A}_t$ we consider the following (controlled) stochastic differential equation:

(3.1)
$$X_s^{[n,\nu]} = x + \int_0^s b(X_u^{[n,\nu]}) \mathrm{d}u + \int_0^s \sigma(X_u^{[n,\nu]}) \mathrm{d}W_u + \int_{[0,s]} n_u \mathrm{d}\nu_u, \quad 0 \le s \le T - t,$$

where $b : \mathbb{R}^d \to \mathbb{R}^d$ and $\sigma : \mathbb{R}^d \to \mathbb{R}^{d \times d'}$ are continuous functions. For P-a.e. ω , the map $s \mapsto n_s(\omega)$ is Borel-measurable on [0, T] and $s \mapsto \nu_s(\omega)$ defines a measure on [0, T]; thus the Lebesgue-Stieltjes integral $\int_{[0,s]} n_u(\omega) d\nu_u(\omega)$ is well-defined for P-a.e. ω . Under our Assumption 3.1 on (b, σ) there is a unique \mathbb{F} -adapted solution of (3.1) by, e.g., [29, Thm. 2.5.7]. We denote

$$\mathsf{P}_x(\ \cdot\) = \mathsf{P}\big(\ \cdot\ \big|X_{0-}^{[n,\nu]} = x\big) \quad \text{and} \quad \mathsf{E}_x\big[\ \cdot\ \big] = \mathsf{E}\big[\ \cdot\ \big|X_{0-}^{[n,\nu]} = x\big].$$

We study a class of 2-player zero-sum games (ZSGs) between a (singular) controller and a stopper. The stopper picks a stopping time $\tau \in \mathcal{T}_t$ and the controller chooses a pair $(n, \nu) \in \mathcal{A}_t$. At time τ the game ends and the controller pays to the stopper a random payoff depending on τ and on the path of $X^{[n,\nu]}$ up to time τ . Given continuous functions $f, g, h : \mathbb{R}^{d+1}_{0,T} \to [0,\infty)$, a fixed discount rate $r \geq 0$ and $(t, x) \in \mathbb{R}^{d+1}_{0,T}$, the game's *expected* payoff reads

$$\mathcal{J}_{t,x}(n,\nu,\tau) = \mathsf{E}_x \bigg[e^{-r\tau} g(t+\tau, X^{[n,\nu]}_{\tau}) + \int_0^\tau e^{-rs} h(t+s, X^{[n,\nu]}_s) \,\mathrm{d}s + \int_{[0,\tau]} e^{-rs} f(t+s, X^{[n,\nu]}_s) \,\mathrm{o}\mathrm{d}\nu_s \bigg],$$

where

(3.2)
$$\int_{[0,\tau]} e^{-rs} f(t+s, X_s^{[n,\nu]}) \circ d\nu_s \coloneqq \int_0^\tau e^{-rs} f(t+s, X_s^{[n,\nu]}) d\nu_s^c + \sum_{0 \le s \le \tau} e^{-rs} \int_0^{\Delta\nu_s} f(t+s, X_{s-}^{[n,\nu]} + \lambda n_s) d\lambda.$$

Here ν^c is the continuous part of the process ν in the decomposition $\nu_s = \nu_s^c + \sum_{u \leq s} \Delta \nu_u$, with $\Delta \nu_u = \nu_u - \nu_{u-}$. If f(t, x) = f(t) the integral (3.2) reduces to the standard Lebesgue-Stieltjes integral $\int_{[0,\tau]} f(s) d\nu_s$. In general, (3.2) gives a cost of exerting control that is consistent with the gradient constraint appearing in Hamilton-Jacobi-Bellman (HJB) equations for singular stochastic control (see, e.g., [40]). It can be interpreted as the limit of standard Lebesgue-Stieltjes integrals if each jump of the control ν is approximated by infinitesimally small sequential jumps ([1, Cor. 1]).

The game admits *lower* and *upper* value, defined respectively by

(3.3)
$$\underline{v}(t,x) \coloneqq \sup_{\tau \in \mathcal{T}_t} \inf_{(n,\nu) \in \mathcal{A}_t} \mathcal{J}_{t,x}(n,\nu,\tau) \quad \text{and} \quad \overline{v}(t,x) \coloneqq \inf_{(n,\nu) \in \mathcal{A}_t} \sup_{\tau \in \mathcal{T}_t} \mathcal{J}_{t,x}(n,\nu,\tau),$$

so that $\underline{v}(t,x) \leq \overline{v}(t,x)$. If equality holds then we say that our game admits a value

(3.4)
$$v(t,x) \coloneqq \underline{v}(t,x) = \overline{v}(t,x).$$

For $a(x) := (\sigma \sigma^{\top})(x) \in \mathbb{R}^{d \times d}$, the infinitesimal generator of $X^{[e_1,0]}$ (with e_1 the unit vector with 1 in the first entry) reads

$$(\mathcal{L}\varphi)(x) = \frac{1}{2} \mathrm{tr} \left(a(x) D^2 \varphi(x) \right) + \langle b(x), \nabla \varphi(x) \rangle, \text{ for any } \varphi \in C^2(\mathbb{R}^d).$$

By density arguments the linear operator \mathcal{L} admits a unique extension $\overline{\mathcal{L}}$ to $W^{2,p}_{\ell oc}(\mathbb{R}^d)$ and, with a slight abuse of notation, we set $\overline{\mathcal{L}} = \mathcal{L}$.

A heuristic use of the dynamic programming principle, suggests that the value of the game vshould be solution of a free boundary problem of the following form:

Problem A. Fix p > d+2. Find a function $u \in W^{1,2,p}_{\ell oc}(\mathbb{R}^{d+1}_{0,T})$ such that, letting

$$\mathcal{I} \coloneqq \left\{ (t,x) \in \mathbb{R}^{d+1}_{0,T} \big| \, |\nabla u(t,x)|_d < f(t,x) \right\} \quad \text{and} \quad \mathcal{C} \coloneqq \left\{ (t,x) \in \mathbb{R}^{d+1}_{0,T} \big| \, u(t,x) > g(t,x) \right\},$$

u satisfies:

(3.5)
$$\begin{cases} (\partial_t u + \mathcal{L}u - ru)(t, x) = -h(t, x), & \text{for all } (t, x) \in \mathcal{C} \cap \mathcal{I}; \\ (\partial_t u + \mathcal{L}u - ru)(t, x) \ge -h(t, x), & \text{for a.e. } (t, x) \in \mathcal{C}; \\ (\partial_t u + \mathcal{L}u - ru)(t, x) \le -h(t, x), & \text{for a.e. } (t, x) \in \mathcal{I}; \\ u(t, x) \ge g(t, x), & \text{for all } (t, x) \in \mathbb{R}^{d+1}_{0,T}; \\ |\nabla u(t, x)|_d \le f(t, x), & \text{for all } (t, x) \in \mathbb{R}^{d+1}_{0,T}; \\ u(T, x) = g(T, x), & \text{for all } x \in \mathbb{R}^d, \end{cases}$$

with $|u(t,x)| \leq c(1+|x|_d^2)$ for all $(t,x) \in \mathbb{R}^{d+1}_{0,T}$ and a suitable c > 0.

Notice that the conditions $u \ge g$ and $|\nabla u|_d \le f$ hold for all (t, x) because of the embedding (2.2). Thus, the two sets \mathcal{I} and \mathcal{C} are open in $[0,T) \times \mathbb{R}^d$. It is not hard to check that a function $u \in W^{1,2,p}_{\ell oc}(\mathbb{R}^{d+1}_{0,T})$ solves Problem A if and only if it solves the variational inequalities in (1.2) a.e. on $\mathbb{R}^{d+1}_{0,T}$ with quadratic growth. Next we give assumptions under which we obtain our main result (Theorem 3.3).

Assumption 3.1 (Controlled SDE). The functions b and σ are continuously differentiable and locally Lipschitz on \mathbb{R}^d . Moreover, there is $D_1 > 0$ such that

$$|b(x)|_d + |\sigma(x)|_{d \times d'} \le D_1(1 + |x|_d), \text{ for all } x \in \mathbb{R}^d.$$

Recalling $a = \sigma \sigma^{\top}$, for any bounded set $B \subset \mathbb{R}^d$ there is $\theta_B > 0$ such that $a(\cdot)$ is locally elliptic:

(3.6)
$$\langle \zeta, a(x)\zeta \rangle \ge \theta_B |\zeta|_d^2 \quad \text{for any } \zeta \in \mathbb{R}^d \text{ and all } x \in \overline{B}.$$

Assumption 3.2 (Functions f, g, h). For the functions $f, g, h : \mathbb{R}^{d+1}_{0,T} \to [0, \infty)$ the following hold:

- (i) $f^2, g \in C^{1,2,\alpha}_{\ell oc}(\mathbb{R}^{d+1}_{0,T})$ and $h \in C^{0,1,\alpha}_{\ell oc}(\mathbb{R}^{d+1}_{0,T})$ for some $\alpha \in (0,1)$; (ii) $t \mapsto f(t,x)$ is non-increasing for each $x \in \mathbb{R}^d$ and $0 \le f(t,x) \le c(1+|x|^p_d)$ for some c, p > 0; (iii) there is $K_0 \in (0,\infty)$ such that for all $0 \le s < t \le T$ and all $x \in \mathbb{R}^{d+1}_{0,T}$

(3.7)
$$h(t,x) - h(s,x) \le K_0(t-s) \quad and \quad g(t,x) - g(s,x) \le K_0(t-s);$$

(iv) there is $K_1 \in (0, \infty)$ such that

(3.8)
$$0 \le g(t,x) + h(t,x) \le K_1(1+|x|_d^2), \quad for \ (t,x) \in \mathbb{R}^{d+1}_{0,T};$$

(v) f and g are such that

(3.9)
$$|\nabla g(t,x)|_d \le f(t,x), \quad \text{for all } (t,x) \in \mathbb{R}^{d+1}_{0,T};$$

(vi) there is $K_2 \in (0, \infty)$ such that

(3.10)
$$\Theta(t,x) \coloneqq \left(h + \partial_t g + \mathcal{L}g - rg\right)(t,x) \ge -K_2, \quad \text{for all } (t,x) \in \mathbb{R}^{d+1}_{0,T}.$$

Condition (3.7) is immediately satisfied if h and q are time-homogeneous (as it is often the case in investment problems, see, e.g., [8], [9], [13]) or if the maps $t \mapsto (h(t,x), g(t,x))$ are nonincreasing for all $x \in \mathbb{R}^d$. Otherwise, that condition amounts to setting a maximum growth rate on $t \mapsto (h(t,x), q(t,x))$ as time increases. Condition (3.10) guarantees that there is no region in the state space where the controller (minimiser) can push the process and obtain arbitrarily large (negative) running gains. Condition (3.9) is sufficient to guarantee that the stopping payoff satisfies the gradient constraint in (3.5) and therefore, from a probabilistic point of view, the stopper can stop at any point in the state-space. On a more technical level, we use (3.9) to obtain crucial bounds on the spatial gradient of the solution of a penalised problem introduced below (cf. proof of Lemma 5.8) and on its time-derivative (cf. proof of Lemma 5.7, where we use the lower bound on the Hamiltonian (4.5) implied by (3.9). Finally, it will be shown in Lemma 6.4 that (3.9) implies that the controller should never exert a jump at the time the stopper ends the game. Here we are not concerned with the construction of equilibria and Lemma 6.4 is not needed for the rest of our analysis. However, we notice that the lemma points to potential difficulties in the construction of equilibria without condition (3.9). Intuitively, when (3.9) is violated, the controller could make arbitrarily large (negative) gains by performing a large jump at the time the stopper decides to end the game. This can be seen by choosing $n_{\tau} = -\nabla g / |\nabla g|_d$ in the third line of (6.9), so that the controller gains $|\nabla g|_d - f > 0$ per unit of control exerted.

The next theorem is the main result of the paper and its proof is distilled in the following sections through a number of technical results and estimates.

Theorem 3.3. The game described above admits a value (i.e., (3.4) holds) and the value function v of the game is the maximal solution to Problem A. Moreover, for any given $(t, x) \in \mathbb{R}^{d+1}_{0,T}$ and any admissible control $(n, \nu) \in \mathcal{A}_t$, the stopping time $\tau_* = \tau_*(t, x; n, \nu) \in \mathcal{T}_t$ defined under P_x as

(3.11)
$$\tau_* \coloneqq \inf \left\{ s \ge 0 \, \middle| \, v(t+s, X_s^{[n,\nu]}) = g(t+s, X_s^{[n,\nu]}) \right\} \wedge (T-t),$$

is optimal for the stopper. That is

$$v(t,x) = \inf_{(n,\nu)\in\mathcal{A}_t} \mathcal{J}_{t,x}(n,\nu,\tau_*(n,\nu)).$$

Remark 3.4. The optimal stopping rule τ_* is in so-called feedback form, because it only depends on the sample paths of the controlled dynamics $X^{[n,\nu]}$. This means that it characterises a best response against any choice of the control (n,ν) available to the controller.

Remark 3.5. Uniqueness of the solution to Problem A remains an open question. Methods used in, e.g., [35], do not apply due to the presence of obstacle and gradient constraints. Existence of an optimal control pair $(n^*, \nu^*) \in \mathcal{A}_t$ is also subtle and cannot be addressed in the generality of our setting. Even in standard singular control problems (not games) abstract existence results rely on compactness arguments in the space of increasing processes under more stringent assumptions (e.g., convexity/concavity or independence from the state dynamics) on the functions f, g, h, b and σ (see, e.g., [7], [22], [31], [37]).

4. PENALISED PROBLEM AND A PRIORI ESTIMATES

In this section we first introduce a class of penalised problems and illustrate their connection with a class of ZSGs of control (Section 4.1). Then we provide important *a priori* estimates on the

growth and gradient of the solution of such penalised problems (Sections 4.2 and 4.3) and, finally, we prove existence and uniqueness of the solution (Section 4.4).

4.1. A penalised problem. For technical reasons related to solvability of the penalised problem and the probabilistic representation of its solution we choose to work on a sequence of bounded domains $(\mathcal{O}_m)_{m\in\mathbb{N}}\subset\mathbb{R}^{d+1}_{0,T}$. Recall that $B_m\subset\mathbb{R}^d$ is the open ball of radius m centred in the origin and set $\mathcal{O}_m := [0,T) \times B_m$ with parabolic boundary $\partial_P \mathcal{O}_m = ([0,T) \times \partial B_m) \cup (\{T\} \times \overline{B}_m)$.

Let $(\xi_m)_{m\in\mathbb{N}}\subset C_c^{\infty}(\mathbb{R}^d)$ be such that for each $m\in\mathbb{N}$ we have:

- (i) $0 \le \xi_m \le 1$ on \mathbb{R}^d , with $\xi_m = 1$ on B_m and $\xi_m = 0$ on $\mathbb{R}^d \setminus B_{m+1}$;
- (ii) there is $C_0 > 0$ independent of $m \in \mathbb{N}$ such that

(4.1)
$$|\nabla \xi_m|_d^2 \le C_0 \xi_m \text{ on } \mathbb{R}^d.$$

An example of such functions is provided in Appendix A.1 for completeness. We define

$$g_m(t,x) \coloneqq \xi_{m-1}(x)g(t,x)$$
 and $h_m(t,x) \coloneqq \xi_{m-1}(x)h(t,x)$, for $(t,x) \in \mathbb{R}^{d+1}_{0,T}$.

Clearly $g_m = h_m = 0$ on $\mathbb{R}^{d+1}_{0,T} \setminus \mathcal{O}_m$ while $g_m = g$ and $h_m = h$ on \mathcal{O}_{m-1} . We also define a version f_m of the function f so that $f_m = f$ on \mathcal{O}_{m-1} and the condition

(4.2)
$$|\nabla g_m(t,x)|_d \le f_m(t,x), \quad \text{for } (t,x) \in \overline{\mathcal{O}}_m$$

is preserved. For $(t, x) \in \mathbb{R}^{d+1}_{0,T}$ we let

$$f_m(t,x) \coloneqq \left(f^2(t,x) + \|g\|_{C^0(\overline{\mathcal{O}}_m)}^2 |\nabla\xi_{m-1}(x)|_d^2 + 2(g\xi_{m-1}\langle\nabla\xi_{m-1},\nabla g\rangle)(t,x) \right)^{\frac{1}{2}}$$

and notice that on $\overline{\mathcal{O}}_m$

$$|\nabla g_m|_d^2 = \sum_{i=1}^d \left(\xi_{m-1}\partial_{x_i}g + g\partial_{x_i}\xi_{m-1}\right)^2 = \xi_{m-1}^2 |\nabla g|_d^2 + g^2 |\nabla \xi_{m-1}|_d^2 + 2\xi_{m-1}g\langle \nabla \xi_{m-1}, \nabla g \rangle \le f_m^2,$$

where the inequality follows by the assumption $|\nabla g|_d \leq f$ and $|\xi_m| \leq 1$. Since $\nabla \xi_{m-1} = \mathbf{0}$ on B_{m-1} , we have $f_m = f$ on \mathcal{O}_{m-1} . Notice that $f_m^2 \in C^{0,1,\alpha}(\overline{\mathcal{O}}_m)$ by Assumption 3.2 but it does not vanish on the boundary of \mathcal{O}_m . By construction, it is clear that

$$g_m \to g$$
, $h_m \to h$ and $f_m \to f$, as $m \to \infty$,

uniformly on any compact $\mathcal{K} \subset \mathbb{R}^{d+1}_{0,T}$.

Let us now state the penalised problem. Fix $(\varepsilon, \delta) \in (0, 1)^2$ and $m \in \mathbb{N}$. Let $\psi_{\varepsilon} \in C^2(\mathbb{R})$ be a non-negative, convex function such that $\psi_{\varepsilon}(y) = 0$ for $y \leq 0$, $\psi_{\varepsilon}(y) > 0$ for y > 0, $\psi'_{\varepsilon} \geq 0$ and $\psi_{\varepsilon}(y) = \frac{y-\varepsilon}{\varepsilon}$ for $y \geq 2\varepsilon$. We also denote $(y)^+ := \max\{0, y\}$ for $y \in \mathbb{R}$.

Problem B. Find $u = u_m^{\varepsilon,\delta}$ with $u \in C^{1,2,\alpha}(\overline{\mathcal{O}}_m)$, for $\alpha \in (0,1)$ as in Assumption 3.2, that solves:

(4.3)
$$\begin{cases} \partial_t u + \mathcal{L}u - ru = -h_m - \frac{1}{\delta} (g_m - u)^+ + \psi_{\varepsilon} \left(|\nabla u|_d^2 - f_m^2 \right), & \text{on } \mathcal{O}_m, \\ u(t, x) = g_m(t, x), & (t, x) \in \partial_P \mathcal{O}_m \end{cases}$$

There are two useful probabilistic interpretations of a solution to Problem B, which we are going to illustrate next. Given $t \in [0, T]$, define the control classes

$$\mathcal{A}_{t}^{\circ} \coloneqq \left\{ (n,\nu) \middle| \begin{array}{l} (n_{s})_{s \in [0,\infty)} \text{ is progressively measurable, } \mathbb{R}^{d} \text{-valued,} \\ \text{with } |n_{s}|_{d} = 1, \text{ P-a.s. for all } s \in [0,\infty); \\ (\nu_{s})_{s \in [0,\infty)} \text{ is } \mathbb{F} \text{-adapted, real-valued, non-decreasing and} \\ \text{absolutely continuous in time, P-a.s., with } \mathbb{E}[|\nu_{T-t}|^{2}] < \infty \end{array} \right\}$$

and

$$\mathcal{T}_t^{\delta} \coloneqq \left\{ w \middle| \begin{array}{l} (w_s)_{s \in [0,\infty)} \text{ is progressively measurable,} \\ \text{with } 0 \le w_s \le \frac{1}{\delta}, \text{ P-a.s. for all } s \in [0, T-t] \end{array} \right\}.$$

It is obvious that $\mathcal{A}_t^{\circ} \subset \mathcal{A}_t$. For $(t, x) \in \mathbb{R}_{0,T}^{d+1}$ and $y \in \mathbb{R}^d$ we define the Hamiltonian

(4.4)
$$H_m^{\varepsilon}(t,x,y) \coloneqq \sup_{p \in \mathbb{R}^d} \left\{ \langle y, p \rangle - \psi_{\varepsilon} \left(|p|_d^2 - f_m^2(t,x) \right) \right\}$$

The function H_m^{ε} is non-negative (pick p = 0). Thanks to (4.2), choosing $p = -\nabla g_m(t, x)$ we have $H_m^{\varepsilon}(t,x,y) \ge -\langle y, \nabla g_m(t,x) \rangle, \quad \text{for all } (t,x,y) \in [0,T] \times \mathbb{R}^d \times \mathbb{R}^d.$ (4.5)

For any admissible pair $(n, \nu) \in \mathcal{A}_t^{\circ}$, we consider the controlled dynamics

$$X_{s}^{[n,\nu]} = x + \int_{0}^{s} \left[b(X_{u}^{[n,\nu]}) + n_{u}\dot{\nu}_{u} \right] \mathrm{d}u + \int_{0}^{s} \sigma(X_{u}^{[n,\nu]}) \mathrm{d}W_{u}, \quad \text{for } 0 \le s \le T - t,$$

and define $\rho_m = \rho_{\mathcal{O}_m}(t, x; n, \nu)$ as

(4.6)
$$\rho_m = \inf\{s \ge 0 \mid X_s^{[n,\nu]} \notin B_m\} \land (T-t).$$

Instead, for $w \in \mathcal{T}_t^{\delta}$, we introduce a controlled discount factor $R_s^w \coloneqq \exp(-\int_0^s [r+w_{\lambda}] d\lambda)$. For $(t,x) \in \overline{\mathcal{O}}_m$ and a triple $[(n,\nu),w] \in \mathcal{A}_t^{\circ} \times \mathcal{T}_t^{\delta}$ let us consider an expected payoff:

(4.7)
$$\mathcal{J}_{t,x}^{\varepsilon,\delta,m}(n,\nu,w) = \mathsf{E}_x \bigg[\int_0^{\rho_m} R_s^w \big[h_m(\cdot) + w_s g_m(\cdot) + H_m^{\varepsilon}(\cdot, n_s \dot{\nu}_s) \big] (t+s, X_s^{[n,\nu]}) \, \mathrm{d}s + R_{\rho_m}^w g_m(t+\rho_m, X_{\rho_m}^{[n,\nu]}) \bigg].$$

The associated upper and lower value read, respectively,

$$\overline{v}_{m}^{\varepsilon,\delta}(t,x) = \inf_{(n,\nu)\in\mathcal{A}_{t}^{\circ}} \sup_{w\in\mathcal{T}_{t}^{\delta}} \mathcal{J}_{t,x}^{\varepsilon,\delta,m}(n,\nu,w) \quad \text{and} \quad \underline{v}_{m}^{\varepsilon,\delta}(t,x) = \sup_{w\in\mathcal{T}_{t}^{\delta}} \inf_{(n,\nu)\in\mathcal{A}_{t}^{\circ}} \mathcal{J}_{t,x}^{\varepsilon,\delta,m}(n,\nu,w)$$

so that $\underline{v}_m^{\varepsilon,\delta} \leq \overline{v}_m^{\varepsilon,\delta}$. A solution of Problem B coincides with the value function of this ZSG.

Proposition 4.1. Let $u_m^{\varepsilon,\delta}$ be a solution of Problem *B*. Then

(4.8)
$$u_m^{\varepsilon,\delta}(t,x) = \overline{v}_m^{\varepsilon,\delta}(t,x) = \underline{v}_m^{\varepsilon,\delta}(t,x), \quad \text{for all } (t,x) \in \overline{\mathcal{O}}_m.$$

Proof. For simplicity denote $u = u_m^{\varepsilon,\delta}$. By definition $\overline{v}_m^{\varepsilon,\delta} = \underline{v}_m^{\varepsilon,\delta} = u = g_m$ on $\partial_P \mathcal{O}_m$. Fix $(t,x) \in \mathcal{O}_m$ and an arbitrary triple $[(n,\nu),w] \in \mathcal{A}_t^{\circ} \times \mathcal{T}_t^{\delta}$. Applying Dynkin's formula to $R_{\rho_m}^w u(t+\rho_m, X_{\rho_m}^{[n,\nu]})$ and using that u solves (4.3) we have

(4.9)
$$u(t,x) = \mathsf{E}_{x} \left[R^{w}_{\rho_{m}} g_{m}(t+\rho_{m}, X^{[n,\nu]}_{\rho_{m}}) + \int_{0}^{\rho_{m}} R^{w}_{s} h_{m}(t+s, X^{[n,\nu]}_{s}) \,\mathrm{d}s \right. \\ \left. + \int_{0}^{\rho_{m}} R^{w}_{s} \left[\frac{1}{\delta} \left(g_{m} - u \right)^{+} - \psi_{\varepsilon} \left(|\nabla u|^{2}_{d} - f^{2}_{m} \right) \right] (t+s, X^{[n,\nu]}_{s}) \,\mathrm{d}s \right. \\ \left. + \int_{0}^{\rho_{m}} R^{w}_{s} \left[w_{s} u(t+s, X^{[n,\nu]}_{s}) - \left\langle n_{s} \dot{\nu}_{s}, \nabla u(t+s, X^{[n,\nu]}_{s}) \right\rangle \right] \,\mathrm{d}s \right]$$

By definition of the Hamiltonian we have (choosing $p = -\nabla u(t + s, X_s^{[n,\nu]})$ in (4.4)) $(4.10) \quad -\left\langle n_s \dot{\nu}_s, \nabla u(t+s, X_s^{[n,\nu]}) \right\rangle - \psi_{\varepsilon} \left(|\nabla u|_d^2 - f_m^2 \right) (t+s, X_s^{[n,\nu]}) \le H_m^{\varepsilon}(t+s, X_s^{[n,\nu]}, n_s \dot{\nu}_s).$ Moreover, choosing $w = w^* \in \mathcal{T}_t^{\delta}$ defined as

(4.11)
$$w_s^* \coloneqq \begin{cases} 0 & \text{if } u(t+s, X_s^{[n,\nu]}) > g_m(t+s, X_s^{[n,\nu]}), \\ \frac{1}{\delta} & \text{if } u(t+s, X_s^{[n,\nu]}) \le g_m(t+s, X_s^{[n,\nu]}), \end{cases}$$

we also have

(4.12)
$$\frac{1}{\delta} (g_m - u)^+ (t + s, X_s^{[n,\nu]}) + w_s^* u(t + s, X_s^{[n,\nu]}) = w_s^* g_m(t + s, X_s^{[n,\nu]}).$$

Then, plugging (4.10) and (4.12) into (4.9) we arrive at $u(t,x) \leq \mathcal{J}_{t,x}^{\varepsilon,\delta,m}(n,\nu,w^*)$. Since the pair (n, ν) was arbitrary, then we have

$$u(t,x) \leq \inf_{(n,\nu)\in\mathcal{A}_t^o} \mathcal{J}_{t,x}^{\varepsilon,\delta,m}(n,\nu,w^*),$$

and therefore $u(t,x) \leq \underline{v}_m^{\varepsilon,\delta}(t,x)$. Next we are going to prove that $u \geq \overline{v}_m^{\varepsilon,\delta}$. For any $w \in \mathcal{T}_t^{\delta}$ it is not hard to see that

(4.13)
$$\frac{1}{\delta} (g_m - u)^+ (t + s, X_s^{[n,\nu]}) + w_s u(t + s, X_s^{[n,\nu]}) \ge w_s g_m(t + s, X_s^{[n,\nu]}),$$

since $0 \le w_s \le \frac{1}{\delta}$ for all $s \in [0, T - t]$. Assume, it is possible to find a pair $(n^*, \nu^*) \in \mathcal{A}_t^\circ$ such that, putting $X^* = X^{[n^*, \nu^*]}$ we obtain

(4.14)
$$-\langle n_s^* \dot{\nu}_s^*, \nabla u(t+s, X_s^*) \rangle - \psi_{\varepsilon} \left(|\nabla u|_d^2 - f_m^2 \right) (t+s, X_s^*) = H_m^{\varepsilon} (t+s, X_s^*, n_s^* \dot{\nu}_s^*).$$

Then, plugging (4.13) and (4.14) into (4.9), we have $u(t,x) \geq \mathcal{J}_{t,x}^{\varepsilon,\delta,m}(n^*,\nu^*,w)$. Since $w \in \mathcal{T}_t^{\delta}$ is arbitrary we have

$$u(t,x) \ge \sup_{w \in \mathcal{T}_t^{\delta}} \mathcal{J}_{t,x}^{\varepsilon,\delta,m}(n^*,\nu^*,w) \ge \overline{v}_m^{\varepsilon,\delta}(t,x).$$

It remains to find the pair (n^*, ν^*) . By concavity, the supremum in $H^{\varepsilon}_m(t, x, y)$ is uniquely attained at a point $p = p(t, x, y) \in \mathbb{R}^d$ identified by the first-order condition $y = 2\psi'_{\varepsilon}(|p|^2_d - f^2_m(t, x))p$. Taking

(4.15)
$$n_s^* \coloneqq \begin{cases} -\frac{\nabla u_m^{\varepsilon,\delta}(t+s,X_s^*)}{|\nabla u_m^{\varepsilon,\delta}(t+s,X_s^*)|_d}, & \text{if } \nabla u_m^{\varepsilon,\delta}(t+s,X_s^*) \neq \mathbf{0}, \\ \text{any unit vector}, & \text{if } \nabla u_m^{\varepsilon,\delta}(t+s,X_s^*) = \mathbf{0}, \\ \dot{\nu}_s^* \coloneqq 2\psi_{\varepsilon}' \left(|\nabla u_m^{\varepsilon,\delta}(t+s,X_s^*)|_d^2 - f_m^2(t+s,X_s^*) \right) |\nabla u_m^{\varepsilon,\delta}(t+s,X_s^*)|_d, \end{cases}$$

with

(

$$(4.16) X_{s\wedge\rho_m}^* = x + \int_0^{s\wedge\rho_m} \left[b(X_\lambda^*) - 2\left(\psi_\varepsilon'(|\nabla u_m^{\varepsilon,\delta}|_d^2 - f_m^2)\nabla u_m^{\varepsilon,\delta}\right)(t+\lambda, X_\lambda^*) \right] \mathrm{d}\lambda + \int_0^{s\wedge\rho_m} \sigma(X_\lambda^*) \mathrm{d}W_\lambda$$

for $s \in [0, T-t]$, we have that (4.14) holds (we restored the notation $u_m^{\varepsilon,\delta}$ for future reference). It remains to check that $(X^*_{s \wedge \rho_m})_{s \in [0,T]}$ is actually well-defined.

Since $u_m^{\varepsilon,\delta} \in C^{1,2,\alpha}(\overline{\mathcal{O}}_m)$ and $\psi_{\varepsilon} \in C^2(\mathbb{R})$, then both the drift and diffusion coefficients of the controlled SDE are Lipschitz in space (recall Assumption 3.1). Hence (4.16) admits a unique strong solution. Moreover, both n^* and ν^* are progressively measurable and since $0 \le \psi'_{\varepsilon} \le 1/\varepsilon$ then also $0 \leq \dot{\nu}_s^* \leq 2\varepsilon^{-1} \|\nabla u_m^{\varepsilon,\delta}\|_{C^0(\overline{\mathcal{O}}_m)} < \infty.$ Therefore $(n^*, \nu^*) \in \mathcal{A}_t^{\circ}$.

Remark 4.2. From the proof of Proposition 4.1 we see that w^* defined in (4.11) is optimal for the maximiser and (n^*, ν^*) defined in (4.15) is optimal for the minimiser in the ZSG with payoff (4.7).

We also show that $u_m^{\varepsilon,\delta}$ is the value function of a control problem with a recursive structure.

Proposition 4.3. Let $u_m^{\varepsilon,\delta}$ be a solution of Problem *B*. Then, for $(t,x) \in \overline{\mathcal{O}}_m$,

$$u_m^{\varepsilon,\delta}(t,x) = \inf_{(n,\nu)\in\mathcal{A}_t^{\circ}} \mathsf{E}_x \left[e^{-(r+\delta^{-1})\rho_m} g_m(t+\rho_m, X_{\rho_m}^{[n,\nu]}) + \int_0^{\rho_m} e^{-(r+\delta^{-1})s} \left[h_m + \frac{1}{\delta} (g_m \vee u_m^{\varepsilon,\delta}) + H_m^{\varepsilon}(\cdot, n_s \dot{\nu}_s) \right] (t+s, X_s^{[n,\nu]}) \,\mathrm{d}s \right],$$

$$4.17)$$

and the pair (n^*, ν^*) from (4.15) is optimal.

Proof. For simplicity denote $u = u_m^{\varepsilon,\delta}$. Since $\frac{1}{\delta}(g_m - u)^+ + \frac{1}{\delta}u = \frac{1}{\delta}g_m \vee u$, then taking $w \equiv \frac{1}{\delta}$ in (4.9) and using (4.10) we get

$$\begin{split} u(t,x) &\leq \inf_{(n,\nu)\in\mathcal{A}_t^{\diamond}} \mathsf{E}_x \bigg[e^{-(r+\delta^{-1})\rho_m} g_m(t+\rho_m, X_{\rho_m}^{[n,\nu]}) \\ &+ \int_0^{\rho_m} e^{-(r+\delta^{-1})s} \big[h_m + \frac{1}{\delta} g_m \vee u + H_m^{\varepsilon}(\cdot, n_s \dot{\nu}_s) \big](t+s, X_s^{[n,\nu]}) \,\mathrm{d}s \bigg]. \end{split}$$

The equality is obtained by substituting in (4.9) the controls $w \equiv \frac{1}{\delta}$ and (n^*, ν^*) defined in (4.15). Recalling the notation $X^* = X^{[n^*, \nu^*]}$ and (4.14) we obtain (4.17) and optimality of (n^*, ν^*) .

From the probabilistic representation of $u_m^{\varepsilon,\delta}$ in (4.8) we establish uniqueness in Problem B. Corollary 4.4. There is at most one solution to Problem B.

4.2. Quadratic growth and stability. Here we establish growth and stability results for $u_m^{\varepsilon,\delta}$.

Lemma 4.5. Let $u_m^{\varepsilon,\delta}$ be a solution of Problem *B*. Then, there is a constant $K_3 > 0$ independent of ε, δ, m such that

(4.18)
$$0 \le u_m^{\varepsilon,\delta}(t,x) \le K_3(1+|x|_d^2), \quad \text{for all } (t,x) \in \overline{\mathcal{O}}_m$$

Proof. Since f_m , g_m and h_m are non-negative, then by (4.8) we get $u_m^{\varepsilon,\delta} \ge 0$. For the upper bound we pick the control pair $(n,\nu) = (e_1,0)$ (with e_1 the unit vector with 1 in the first coordinate) in (4.7) and notice that $H_m^{\varepsilon}(t+s, X_s^{[e_1,0]}, 0) = 0$. Then

$$\begin{split} u_m^{\varepsilon,\delta}(t,x) &\leq \sup_{w \in \mathcal{T}^{\delta}} \mathsf{E}_x \bigg[\int_0^{\rho_m} R_s^w \big[h_m + w_s g_m \big] (t+s, X_s^{[e_1,0]}) \,\mathrm{d}s + R_{\rho_m}^w g_m(t+\rho_m, X_{\rho_m}^{[e_1,0]}) \bigg] \\ &\leq K_1 \sup_{w \in \mathcal{T}^{\delta}} \mathsf{E}_x \bigg[\bigg(1 + \sup_{0 \leq s \leq T-t} e^{-rs} \big| X_s^{[e_1,0]} \big|_d^2 \bigg) \Big(1 + \int_0^{\rho_m} e^{-\int_0^s w_\lambda \mathrm{d}\lambda} \big[1 + w_s \big] \,\mathrm{d}s \Big) \bigg], \end{split}$$

where the inequality is using the quadratic growth of h_m and g_m (see (3.8)) and the definition of R^w . For P-a.e. ω we have the simple bound

$$\int_0^{\rho_m(\omega)} e^{-\int_0^s w_\lambda(\omega) \mathrm{d}\lambda} \left[1 + w_s(\omega)\right] \mathrm{d}s \le 1 + T.$$

Therefore

$$u_m^{\varepsilon,\delta}(t,x) \le K_1(2+T)\mathsf{E}_x \left[1 + \sup_{0 \le s \le T-t} e^{-rs} \left| X_s^{[e_1,0]} \right|_d^2 \right] \le K_3(1+|x|_d^2),$$

by standard estimates for SDEs with coefficients with linear growth ([29, Cor. 2.5.10]) and the constant $K_3 > 0$ depends only on T, D_1 and K_1 in Assumptions 3.1 and 3.2.

Remark 4.6. In particular, (4.18) implies that for any $m \ge m_0 \in \mathbb{N}$ and $\varepsilon, \delta \in (0, 1)$

$$\left\|u_m^{\varepsilon,\delta}\right\|_{C^0(\overline{\mathcal{O}}_{m_0})} \le K_3(1+|m_0|^2) \rightleftharpoons M_1(m_0).$$

The next result relies upon standard PDE arguments. Its proof is in Appendix for completeness. **Lemma 4.7.** Let $u_m^{\varepsilon,\delta}$ be a solution of Problem B. Let $u^n \in C^{\infty}(\overline{\mathcal{O}}_m)$ and $\chi_n \in C^{\infty}(\mathbb{R})$ be such that $\chi_n(0) = 0, \ \chi'_n \geq 0, \ (\chi_n)_{n \in \mathbb{N}}$ are equi-Lipschitz, and

$$\|u^n - u_m^{\varepsilon,\delta}\|_{C^{1,2,\gamma}(\overline{\mathcal{O}}_m)} + \|\chi_n - (\cdot)^+\|_{C^0(\mathbb{R})} \le \frac{1}{n}, \quad n \in \mathbb{N},$$

for some $\gamma \in (0, \alpha)$. Then, there exists a unique solution $w^n \in C^{1,2,\alpha}(\overline{\mathcal{O}}_m)$ of

(4.19)
$$\begin{cases} \partial_t w^n + \mathcal{L}w^n - rw^n = -h_m - \frac{1}{\delta}\chi_n \left(g_m - u_m^{\varepsilon,\delta}\right) + \psi_{\varepsilon}(|\nabla u^n|_d^2 - f_m^2 - \frac{1}{n}), & \text{on } \mathcal{O}_m, \\ w^n(t,x) = g_m(t,x), & (t,x) \in \partial_P \mathcal{O}_m. \end{cases}$$

Moreover, $w^n \in C^{1,3,\alpha}_{Loc}(\mathcal{O}_m)$ and $w^n \to u^{\varepsilon,\delta}_m$ in $C^{1,2,\beta}(\overline{\mathcal{O}}_m)$ as $n \to \infty$, for all $\beta \in (0,\alpha)$.

4.3. Gradient bounds. Our next goal is to find a bound for the norm of the gradient of $u_m^{\varepsilon,\delta}$ uniformly in ε, δ . We start by considering an estimate on the parabolic boundary $\partial_P \mathcal{O}_m$ that will be later used to bound $u_m^{\varepsilon,\delta}$ on the whole $\overline{\mathcal{O}}_m$.

Lemma 4.8. Let $u_m^{\varepsilon,\delta}$ be a solution of Problem *B*. Then, there is $M_2 = M_2(m) > 0$ such that (4.20) $\sup_{(t,x)\in\partial_P\mathcal{O}_m} |\nabla u_m^{\varepsilon,\delta}(t,x)|_d \leq M_2, \quad \text{for all } \varepsilon,\delta \in (0,1).$

Proof. For simplicity we denote $u = u_m^{\varepsilon,\delta}$. If t = T we have $u(T, x) = g_m(T, x)$ for $x \in \overline{B}_m$ and the bound is trivial because $\nabla u(T, x) = \nabla g_m(T, x)$.

Next, let
$$t \in [0, T)$$
. Notice that $u|_{\partial B_m} = g_m|_{\partial B_m} = 0$. Fix $x \in B_m$ and $y \in \partial B_m$. Then
(4.21) $0 \le u(t, x) - u(t, y) = u(t, x) - g_m(t, y) \le \|\nabla g_m\|_{C^0(\overline{\mathcal{O}}_m)} |x - y|_d + u(t, x) - g_m(t, x).$

For arbitrary $(n, \nu) \in \mathcal{A}_t^{\circ}$ and $w \in \mathcal{T}_t^{\delta}$, Dynkin's formula gives

$$g_m(t,x) = \mathsf{E}_x \bigg[R^w_{\rho_m} g_m(t+\rho_m, X^{[n,\nu]}_{\rho_m}) - \int_0^{\rho_m} R^w_s \langle n_s \dot{\nu}_s, \nabla g_m(t+s, X^{[n,\nu]}_s) \rangle \,\mathrm{d}s (4.22) \qquad \qquad - \int_0^{\rho_m} R^w_s \big[(\partial_t g_m + \mathcal{L} g_m - rg_m)(t+s, X^{[n,\nu]}_s) - w_s g_m(t+s, X^{[n,\nu]}_s) \big] \,\mathrm{d}s \bigg].$$

Then, setting $\Theta_m = \partial_t g_m + \mathcal{L} g_m - rg_m + h_m$ and recalling (4.8), we can write

$$\begin{split} u(t,x) &- g_m(t,x) \\ &= \inf_{(n,\nu)\in\mathcal{A}^{\diamond}_t} \sup_{w\in\mathcal{T}^{\delta}_t} \mathsf{E}_x \bigg[\int_0^{\rho_m} R^w_s \Big(\big[\Theta_m + \langle n_s \dot{\nu}_s, \nabla g_m \rangle \big] + H^{\varepsilon}_m(\cdot, n_s \dot{\nu}_s) \Big)(t+s, X^{[n,\nu]}_s) \,\mathrm{d}s \bigg]. \end{split}$$

Picking $(n, \nu) = (e_1, 0)$ (with e_1 the unit vector with 1 in the first coordinate) and recalling that $H_m^{\varepsilon}(\cdot, 0) = 0$ and $\rho_m = \rho_{\mathcal{O}_m}$ we obtain the upper bound

$$u(t,x) - g_m(t,x) \le \sup_{w \in \mathcal{T}_t^{\delta}} \mathsf{E}_x \left[\int_0^{\rho_m} R_s^w \Theta_m(t+s, X_s^{[e_1,0]}) \,\mathrm{d}s \right] \le \left\| \Theta_m \right\|_{C^0(\overline{\mathcal{O}}_m)} \mathsf{E}_x\left[\rho_{\mathcal{O}_m}\right].$$

Combining the latter with (4.21) we obtain

$$0 \le u(t,x) - u(t,y) \le \|\nabla g_m\|_{C^0(\overline{\mathcal{O}}_m)} |x - y|_d + \|\Theta_m\|_{C^0(\overline{\mathcal{O}}_m)} \mathsf{E}_x\left[\rho_{\mathcal{O}_m}\right].$$

Since $\rho_{\mathcal{O}_m} = \rho_{\mathcal{O}_m}(t, x; e_1, 0)$ is associated to the control pair $(n, \nu) = (e_1, 0)$, then

$$\rho_{\mathcal{O}_m} = \inf\{s \ge 0 \mid X_s^{[e_1,0]} \notin B_m\} \land (T-t) =: \tau_m \land (T-t),$$

and clearly $\mathsf{E}_x[\rho_{\mathcal{O}_m}] \leq \mathsf{E}_x[\tau_m] =: \pi(x)$. It is well-known that $\pi \in C^2(\overline{B}_m)$ and it solves

$$\mathcal{L}\pi(x) = -1$$
 for $x \in B_m$ with $\pi(x) = 0$ for $x \in \partial B_m$,

by uniform ellipticity of \mathcal{L} on B_m (see [20, Thm. 6.14]). That is sufficient to conclude

$$\mathsf{E}_x\left[\rho_{\mathcal{O}_m}\right] \le \pi(x) = \pi(x) - \pi(y) \le L_{\pi,m} |x - y|_d,$$

for some constant $L_{\pi,m} > 0$ depending only on the coefficients of \mathcal{L} and the radius m. Then, for all $t \in [0,T)$ we have $0 \le u(t,x) - u(t,y) \le M_2 |x-y|_d$, with $M_2 = \|\nabla g_m\|_{C^0(\overline{\mathcal{O}}_m)} + \|\Theta_m\|_{C^0(\overline{\mathcal{O}}_m)} L_{\pi,m}$. This implies (4.20) because $u \in C^{1,2,\alpha}(\overline{\mathcal{O}}_m)$.

Using Lemma 4.8 we can also provide a bound on $|\nabla u_m^{\varepsilon,\delta}|_d$ in the whole domain $\overline{\mathcal{O}}_m$. It is useful to recall that a function $\varphi \in C^{1,2}(\overline{\mathcal{O}}_m)$ attaining a maximum at a point $(t_0, x_0) \in \mathcal{O}_m$ also satisfies

(4.23)
$$\mathcal{L}\varphi(t_0, x_0) + \partial_t \varphi(t_0, x_0) \le 0,$$

by the maximum principle (see [19, Lemma 2.1]). Since $(t_0, x_0) \in \mathcal{O}_m$, then $\nabla \varphi(t_0, x_0) = \mathbf{0}$ and, when $t_0 \in (0, T)$, also $\partial_t \varphi(t_0, x_0) = 0$.

Proposition 4.9. Let $u_m^{\varepsilon,\delta}$ be a solution of Problem *B*. Then, there is $M_3 = M_3(m)$ such that $\sup_{(t,x)\in\overline{\mathcal{O}}_m} |\nabla u_m^{\varepsilon,\delta}(t,x)|_d \le M_3, \quad \text{for all } \varepsilon, \delta \in (0,1).$ (4.24)

Proof. This proof refines and extends arguments from [40, Lemma A.2] (see also [24, Lemma 3.4] and [28, Lemma 2.8]). For simplicity we denote $u = u_m^{\varepsilon,\delta}$. Let $\lambda_m \in (0,\infty)$ be a constant depending on *m* but independent of ε, δ , which will be chosen later. Let $v^{\lambda} \in C^{0,1,\alpha}(\overline{\mathcal{O}}_m)$ be defined as

$$v^{\lambda}(t,x) \coloneqq |\nabla u(t,x)|_d^2 - \lambda u(t,x)$$

for some $\lambda \in (0, \lambda_m]$. Recalling $M_1 = M_1(m)$ from Remark 4.6 we have for any $\lambda \in (0, \lambda_m]$

(4.25)
$$\sup_{(t,x)\in\overline{\mathcal{O}}_m} |\nabla u(t,x)|_d^2 \le \sup_{(t,x)\in\overline{\mathcal{O}}_m} v^{\lambda}(t,x) + \lambda_m M_1(m)$$

Let $(t^{\lambda}, x^{\lambda}) \in \overline{\mathcal{O}}_m$ be a maximum point for v^{λ} . Two situations may arise: either $(t^{\lambda}, x^{\lambda}) \in \mathcal{O}_m$ or $(t^{\lambda}, x^{\lambda}) \in \partial_P \mathcal{O}_m$. If $(t^{\lambda}, x^{\lambda}) \in \partial_P \mathcal{O}_m$, then by Lemmas 4.5 and 4.8 we have

$$v^{\lambda}(t^{\lambda}, x^{\lambda}) \leq |\nabla u(t^{\lambda}, x^{\lambda})|_{d}^{2} \leq M_{2}^{2} \implies \sup_{(t,x)\in\overline{\mathcal{O}}_{m}} |\nabla u(t,x)|_{d}^{2} \leq M_{2}^{2} + \lambda_{m}M_{1},$$

where the implication follows by (4.25). Thus, if

(4.26)
$$\Lambda_m := \{\lambda \in (0, \lambda_m] : \text{ there exists a maximiser of } v^\lambda \text{ in } \partial_P \mathcal{O}_m\} \neq \emptyset$$

it is sufficient to pick $\lambda \in \Lambda_m$ and (4.24) holds.

Let us now assume $\Lambda_m = \emptyset$ (i.e., $(t^{\lambda}, x^{\lambda}) \in \mathcal{O}_m$ for all $\lambda \in (0, \lambda_m]$). With no loss of generality:

(4.27)
$$|\nabla u(t^{\lambda}, x^{\lambda})|_{d} > 1 \quad \text{and} \quad \langle \nabla u(t^{\lambda}, x^{\lambda}), \nabla g_{m}(t^{\lambda}, x^{\lambda}) \rangle - |\nabla u(t^{\lambda}, x^{\lambda})|_{d}^{2} < 0.$$

If either condition fails, then (4.24) trivially holds. Likewise, we assume

(4.28)
$$\psi_{\varepsilon}' \left(|\nabla u(t^{\lambda}, x^{\lambda})|_d^2 - f_m^2(t^{\lambda}, x^{\lambda}) \right) > 1$$

because ψ_{ε} is convex and $\psi'_{\varepsilon}(r) = \frac{1}{\varepsilon} > 1$ for $r \ge 2\varepsilon$. So, if (4.28) fails, it must be $|\nabla u(t^{\lambda}, x^{\lambda})|_d^2 < \varepsilon$ $f_m^2(t^{\lambda}, x^{\lambda}) + 2\varepsilon$ and (4.24) holds because f_m is bounded on $\overline{\mathcal{O}}_m$.

We would like to compute $\partial_t v + \mathcal{L} v$ but the term containing $(\cdot)^+$ in the PDE for u is not continuously differentiable and, therefore, it is not clear that $|\nabla u|_d^2$ admits classical derivatives. That is why we resort to an approximation procedure. Let u^n , χ_n and w^n be defined as in Lemma 4.7. Recall that $w^n \in C^{1,3,\alpha}_{Loc}(\mathcal{O}_m) \cap C^{1,2,\alpha}(\overline{\mathcal{O}}_m)$ and $w^n \to u$ in $C^{1,2,\gamma}(\overline{\mathcal{O}}_m)$ for all $\gamma \in (0,\alpha)$. Define

$$v^{\lambda,n}(t,x) \coloneqq |\nabla w^n(t,x)|_d^2 - \lambda u(t,x),$$

so that $v^{\lambda,n} \in C^{1,2,\alpha}_{Loc}(\mathcal{O}_m) \cap C^{0,1,\alpha}(\overline{\mathcal{O}}_m)$. Clearly, $v^{\lambda,n} \to v^{\lambda}$ uniformly on $\overline{\mathcal{O}}_m$. Let $(t_n^{\lambda}, x_n^{\lambda})_{n \in \mathbb{N}}$ be a sequence with $(t_n^{\lambda}, x_n^{\lambda}) \in \arg \max_{\overline{\mathcal{O}}_m} v^{\lambda,n}$ for $n \in \mathbb{N}$. Since $\overline{\mathcal{O}}_m$ is compact, the sequence admits a subsequence $(t_{n_k}^{\lambda}, x_{n_k}^{\lambda})_{k \in \mathbb{N}}$ converging to some $(\tilde{t}, \tilde{x}) \in \overline{\mathcal{O}}_m$. It is not hard to show that $(\tilde{t}, \tilde{x}) \in \arg \max_{\overline{\mathcal{O}}_m} v^{\lambda}$ (we provide the full argument in Appendix for completeness). Then, with no loss of generality we can assume $(t^{\lambda}, x^{\lambda}) = (\tilde{t}, \tilde{x})$. That implies that we can choose $(t_{n_k}^{\lambda}, x_{n_k}^{\lambda})_{k \in \mathbb{N}} \subset \mathcal{O}_m$. More precisely, by continuity of v^{λ} , for any $\eta > 0$ there exist bounded open sets $U_{\lambda,\eta} \subset \mathcal{O}_m$ and $V_{\lambda,\eta} \subset B_m$ such that $(t^{\lambda}, x^{\lambda}) \in U_{\lambda,\eta} \cup (\{0\} \times V_{\lambda,\eta})$ and $v^{\lambda}(t,x) > v^{\lambda}(t^{\lambda}, x^{\lambda}) - \eta$ for all $(t,x) \in U_{\lambda,\eta} \cup (\{0\} \times V_{\lambda,\eta})$ (by convention, if $t^{\lambda} \neq 0$ we take $V_{\lambda,\eta} = \emptyset$). Moreover, for $k \in \mathbb{N}$ sufficiently large we have $(t_{n_i}^{\lambda}, x_{n_i}^{\lambda})_{j \geq k} \subset U_{\lambda,\eta} \cup (\{0\} \times V_{\lambda,\eta}).$

From now on we simply relabel our subsequence by $(t_n^{\lambda}, x_n^{\lambda})_{n \in \mathbb{N}}$ with a slight abuse of notation. Since $(t_n^{\lambda}, x_n^{\lambda})$ is a maximum point of $v^{\lambda,n}$ from the maximum principle (see (4.23)) we get

(4.29)
$$\mathcal{L}v^{\lambda,n}(t_n^{\lambda}, x_n^{\lambda}) + \partial_t v^{\lambda,n}(t_n^{\lambda}, x_n^{\lambda}) \le 0.$$

Next we compute explicitly all terms in (4.29) and to simplify notation we drop the argument $(t_n^{\lambda}, x_n^{\lambda})$. Denoting $\partial_t v = v_t, \ \partial_{x_i} v = v_{x_i}$ and $\partial_{x_i x_j} v = v_{x_i x_j}$, we obtain

$$v_t^{\lambda,n} = 2 \langle \nabla w^n, \nabla w_t^n \rangle - \lambda u_t;$$

$$(4.30) \qquad v_{x_i}^{\lambda,n} = 2 \langle \nabla w^n, \nabla w_{x_i}^n \rangle - \lambda u_{x_i}, \qquad 1 \le i \le d;$$

$$v_{x_i x_j}^{\lambda,n} = 2 \langle \nabla w_{x_i}^n, \nabla w_{x_j}^n \rangle + 2 \langle \nabla w^n, \nabla w_{x_i x_j}^n \rangle - \lambda u_{x_i x_j}, \qquad 1 \le i, j \le d.$$

Substituting in (4.29) gives:

(4.31)
$$0 \ge \sum_{i,j=1}^{d} a_{ij} \langle \nabla w_{x_i}^n, \nabla w_{x_j}^n \rangle + 2 \sum_{k=1}^{d} w_{x_k}^n (\partial_t w_{x_k}^n + \mathcal{L} w_{x_k}^n) - \lambda (\partial_t u + \mathcal{L} u).$$

From uniform ellipticity (3.6) on B_m and denoting $\theta_{B_m} = \theta$ we have

(4.32)
$$\sum_{i,j=1}^{d} a_{ij} \langle \nabla w_{x_i}^n, \nabla w_{x_j}^n \rangle = \sum_{k=1}^{d} \sum_{i,j=1}^{d} a_{ij} w_{x_k x_i}^n w_{x_k x_j}^n \ge \sum_{k=1}^{d} \theta |\nabla w_{x_k}^n|_d^2 \ge \theta |D^2 w^n|_{d \times d}^2.$$

To study the second term in (4.31) we introduce the differential operator \mathcal{L}_{x_k} :

(4.33)
$$(\mathcal{L}_{x_k}\varphi)(x) = \frac{1}{2} \operatorname{tr} \left(a_{x_k}(x) D^2 \varphi(x) \right) + \langle b_{x_k}(x), \nabla \varphi(x) \rangle, \quad \text{for } \varphi \in C^2(\mathbb{R}^d),$$

where $a_{x_k} \in \mathbb{R}^{d \times d}$ is the matrix with entries $(\partial_{x_k} a_{ij})_{i,j=1}^d$ and $b_{x_k} \in \mathbb{R}^d$ the vector with entries $(\partial_{x_k} b_i)_{i=1}^d$. Differentiating with respect to x_k the PDE in (4.19) and rearranging terms we get

(4.34)
$$\partial_t w_{x_k}^n + \mathcal{L} w_{x_k}^n = r w_{x_k}^n + \psi_{\varepsilon}' (\bar{\zeta}_n) \cdot (|\nabla u^n|_d^2 - f_m^2)_{x_k} \\ - \frac{1}{\delta} \chi_n' (g_m - u) \cdot (g_m - u)_{x_k} - \partial_{x_k} h_m - \mathcal{L}_{x_k} w^n,$$

where we set $\bar{\zeta}_n \coloneqq (|\nabla u^n|_d^2 - f_m^2)(t_n^{\lambda}, x_n^{\lambda}) - \frac{1}{n}$ for the argument of ψ_{ε}' . In the third term of (4.31) we substitute (4.3) and, combining with (4.32) and (4.34), we obtain

$$0 \ge \theta |D^2 w^n|_{d\times d}^2 + 2\Big[r|\nabla w^n|_d^2 + \psi_{\varepsilon}'(\bar{\zeta}_n)\langle \nabla w^n, \nabla(|\nabla u^n|_d^2 - f_m^2)\rangle - \frac{1}{\delta}\chi_n'(g_m - u)\langle \nabla w^n, \nabla(g_m - u)\rangle (4.35) - \langle \nabla w^n, \nabla h_m \rangle - \sum_{k=1}^d w_{x_k}^n \mathcal{L}_{x_k} w^n \Big] - \lambda \big(ru + \psi_{\varepsilon}(\zeta_n) - \frac{1}{\delta}(g_m - u)^+ - h_m\big),$$

where we set $\zeta_n := (|\nabla u|_d^2 - f_m^2)(t_n^\lambda, x_n^\lambda)$ for the argument of ψ_{ε} . Let us denote $\hat{w}^n = u - w^n$. Then $\|\hat{w}^n\|_{C^{1,2,\gamma}(\overline{\mathcal{O}}_m)} \to 0$ as $n \to \infty$ because $w^n \to u$ in $C^{1,2,\gamma}(\overline{\mathcal{O}}_m)$, for all $\gamma \in (0, \alpha)$. We claim that

$$(4.36) \qquad 2\Big[r|\nabla w^{n}|_{d}^{2} - \frac{1}{\delta}\chi_{n}'(g_{m} - u)\langle\nabla w^{n}, \nabla(g_{m} - u)\rangle - \langle\nabla w^{n}, \nabla h_{m}\rangle - \sum_{k=1}^{d} w_{x_{k}}^{n}\mathcal{L}_{x_{k}}w^{n}\Big] \\ -\lambda\Big(ru + \psi_{\varepsilon}(\zeta_{n}) - \frac{1}{\delta}(g_{m} - u)^{+} - h_{m}\Big) \\ \geq -C_{1}|\nabla u|_{d}^{2} - \theta|D^{2}w^{n}|_{d\times d}^{2} - C_{2} - \lambda rM_{1} - \lambda\psi_{\varepsilon}'(\bar{\zeta}_{n})|\nabla u|_{d}^{2} - R_{n},$$

for M_1 as in Remark 4.6, constants $C_1 = C_1(m) > 0$, $C_2 = C_2(m) > 0$ depending only on m and with

$$0 \le R_n \le \kappa_{\delta,m} \Big(\|\hat{w}^n\|_{C^{0,1,\gamma}(\overline{\mathcal{O}}_m)} + \lambda \|\psi_{\varepsilon}'(\zeta_n) - \psi_{\varepsilon}'(\bar{\zeta}_n)\|_{C^0(\overline{\mathcal{O}}_m)} \Big)$$

where $\kappa_{\delta,m} > 0$ depends on δ , $\|\nabla u\|_{C^0(\overline{\mathcal{O}}_m)}$ and $\|\nabla g_m\|_{C^0(\overline{\mathcal{O}}_m)}$. Clearly $R_n \to 0$ as $n \to \infty$. For the ease of exposition the proof of (4.36) is given separately at the end of this proof.

Plugging (4.36) into (4.35) we obtain

$$0 \ge 2\psi_{\varepsilon}'(\bar{\zeta}_n)\langle \nabla w^n, \nabla(|\nabla u^n|_d^2 - f_m^2)\rangle - C_1|\nabla u|_d^2 - C_2 - \lambda r M_1 - \lambda \psi_{\varepsilon}'(\bar{\zeta}_n)|\nabla u|_d^2 - R_n.$$

By (4.28), $\psi'_{\varepsilon}(\bar{\zeta}_n) \geq 1$ for large n. Then, multiplying both sides of the inequality by -1 we obtain

$$0 \le \psi_{\varepsilon}'(\bar{\zeta}_n) \left(\lambda |\nabla u|_d^2 - 2\langle \nabla w^n, \nabla (|\nabla u^n|_d^2 - f_m^2) \rangle + C_1 |\nabla u|_d^2 + C_2 + \lambda r M_1 + R_n\right)$$

That implies

(4.37)
$$0 \le (C_1 + \lambda) |\nabla u|_d^2 - 2\langle \nabla w^n, \nabla (|\nabla u^n|_d^2 - f_m^2) \rangle + C_2 + \lambda r M_1 + R_n.$$

We claim that

(4.38)
$$-2\langle \nabla w^n, \nabla (|\nabla u^n|_d^2 - f_m^2) \rangle \le -2\lambda |\nabla u|_d^2 + 2|\nabla u|_d |\nabla f_m^2|_d + \tilde{R}_n,$$

where $\tilde{R}_n = \tilde{R}_n(m,\varepsilon,\delta)$ goes to zero as $n \to \infty$. Again, for ease of exposition the proof of (4.38) is given separately at end of this proof, after the one for (4.36). Thus (4.37) becomes

$$0 \le (C_1 - \lambda) |\nabla u|_d^2 + 2 |\nabla u|_d |\nabla f_m^2|_d + C_2 + \lambda r M_1 + R_n + \tilde{R}_n.$$

By definition of f_m it is not hard to verify that $|\nabla f_m^2|_d \leq C_3$ for a constant $C_3 = C_3(m) > 0$ independent of ε and δ . Then from the inequality above we obtain

$$|(\lambda - C_1)|\nabla u|_d^2 \le 2C_3|\nabla u|_d + C_2 + \lambda r M_1 + R_n + \tilde{R}_n.$$

Choosing $\lambda = \overline{\lambda} \coloneqq C_1 + 1$ and recalling our shorthand notation $\nabla u = \nabla u(t_n^{\lambda}, x_n^{\lambda})$, we have

(4.39)
$$|\nabla u(t_n^{\bar{\lambda}}, x_n^{\bar{\lambda}})|_d^2 \le 2C_3 |\nabla u(t_n^{\bar{\lambda}}, x_n^{\bar{\lambda}})|_d + C_2 + \bar{\lambda} r M_1 + R_n + \tilde{R}_n.$$

Thanks to (4.27) we have $|\nabla u(t^{\bar{\lambda}}, x^{\bar{\lambda}})|_d > 1$. Thus, with no loss of generality we can assume that $|\nabla u(t, x)|_d \geq 1$ for all $(t, x) \in \overline{U}_{\bar{\lambda},\eta} \cup (\{0\} \times \overline{V}_{\bar{\lambda},\eta})$ and recall also that $(t_n^{\bar{\lambda}}, x_n^{\bar{\lambda}})_{n \in \mathbb{N}} \subset U_{\bar{\lambda},\eta} \cup (\{0\} \times V_{\bar{\lambda},\eta})$. Thus, dividing by $|\nabla u(t_n^{\bar{\lambda}}, x_n^{\bar{\lambda}})|_d$ in (4.39) we get

(4.40)
$$|\nabla u(t_n^{\bar{\lambda}}, x_n^{\bar{\lambda}})|_d \le 2C_3 + C_2 + \bar{\lambda}rM_1 + R_n + \tilde{R}_n$$

From (4.40) and the definition of $U_{\bar{\lambda},\eta} \cup (\{0\} \times V_{\bar{\lambda},\eta})$

$$\sup_{(t,x)\in\overline{\mathcal{O}}_m} v^{\bar{\lambda}}(t,x) = v^{\bar{\lambda}}(t^{\bar{\lambda}},x^{\bar{\lambda}}) \le v(t_n^{\bar{\lambda}},x_n^{\bar{\lambda}}) + \eta \le (2C_3 + C_2 + \bar{\lambda}rM_1 + R_n + \tilde{R}_n)^2 + \eta$$

Letting $n \to \infty$ we obtain

$$\sup_{(t,x)\in\overline{\mathcal{O}}_m} v^{\bar{\lambda}}(t,x) \le (2C_3 + C_2 + \bar{\lambda}rM_1)^2 + \eta.$$

By the arbitrariness of η and since (4.25) holds for any $\lambda \in (0, \lambda_m]$, taking $\lambda_m = \overline{\lambda}$ we have

$$\sup_{(t,x)\in\overline{\mathcal{O}}_m} |\nabla u(t,x)|_d^2 \le (2C_3 + C_2 + \bar{\lambda}rM_1)^2 + \bar{\lambda}M_1$$

Hence, the proposition holds with $M_3 = ((2C_3 + C_2 + \bar{\lambda}rM_1)^2 + \bar{\lambda}M_1)^{1/2}$ independent of ε and δ . \Box **Proof of** (4.36). Recalling $\hat{w}^n = u - w^n$ we first notice

(4.41)
$$|\nabla w^{n}|_{d}^{2} = \left(|\nabla u|_{d}^{2} - 2\langle \nabla u, \nabla \hat{w}^{n} \rangle + |\nabla \hat{w}^{n}|_{d}^{2} \right)$$
$$\leq |\nabla u|_{d}^{2} + \|\nabla \hat{w}^{n}\|_{C^{0}(\overline{\mathcal{O}}_{m})} \left(2\|\nabla u\|_{C^{0}(\overline{\mathcal{O}}_{m})} + \|\nabla \hat{w}^{n}\|_{C^{0}(\overline{\mathcal{O}}_{m})} \right).$$

The first term on the left-hand side of (4.36) is positive. For the second one, notice

$$\begin{split} \langle \nabla w^n, \nabla (g_m - u) \rangle = & \langle \nabla (u - \hat{w}^n), \nabla (g_m - u) \rangle \\ = & \langle \nabla u, \nabla g_m \rangle - |\nabla u|_d^2 - \langle \nabla \hat{w}^n, \nabla (g_m - u) \rangle \\ \leq & \langle \nabla \hat{w}^n, \nabla (u - g_m) \rangle \leq \| \hat{w}^n \|_{C^{0,1,\gamma}(\overline{\mathcal{O}}_m)} (\| \nabla u \|_{C^0(\overline{\mathcal{O}}_m)} + \| \nabla g_m \|_{C^0(\overline{\mathcal{O}}_m)}), \end{split}$$

where the first inequality holds by (4.27) in $U_{\lambda,\eta} \cup (\{0\} \times V_{\lambda,\eta})$. Since $0 \le \chi'_n \le 2$, then (4.42) $\frac{1}{\delta}\chi'_n(g_m - u)\langle \nabla w^n, \nabla (g_m - u)\rangle \le \frac{2}{\delta} \|\hat{w}^n\|_{C^{0,1,\gamma}(\overline{\mathcal{O}}_m)}(\|\nabla u\|_{C^0(\overline{\mathcal{O}}_m)} + \|\nabla g_m\|_{C^0(\overline{\mathcal{O}}_m)}),$

and this term can be collected into R_n in (4.36).

Let us look at the third and fourth term on the left-hand side of (4.36). For the former we have

(4.43)
$$\langle \nabla w^n, \nabla h_m \rangle \le |\nabla w^n|_d |\nabla h_m|_d \le \frac{1}{2} |\nabla w^n|_d^2 + \frac{1}{2} |\nabla h_m|_d^2.$$

For the latter, recalling (4.33) we have

(4.44)
$$\sum_{k=1}^{d} w_{x_k}^n \mathcal{L}_{x_k} w^n = \frac{1}{2} \sum_{i,j=1}^{d} \langle \nabla w^n, \nabla a_{ij} \rangle w_{x_i x_j}^n + \sum_{i=1}^{d} \langle \nabla w^n, \nabla b_i \rangle w_{x_i}^n$$
$$\leq \frac{d^2}{2} A_m |\nabla w^n|_d |D^2 w^n|_{d \times d} + dA_m |\nabla w^n|_d^2,$$

where we used Cauchy-Schwarz inequality and set

(4.45)
$$A_m \coloneqq \max_{i,j} \left(\left\| \nabla a_{ij} \right\|_{C^0(\overline{B}_m)} + \left\| \nabla b_i \right\|_{C^0(\overline{B}_m)} \right).$$

Using that $d^2 A_m |\nabla w^n|_d |D^2 w^n|_{d \times d} \leq \theta^{-1} d^4 A_m^2 |\nabla w^n|_d^2 + \theta |D^2 w^n|_{d \times d}^2$, with $\theta = \theta_{B_m}$ as in (3.6), and combining (4.43), (4.44) and (4.41), we have

$$(4.46) \quad \langle \nabla w^{n}, \nabla h_{m} \rangle + \sum_{k=1}^{d} w_{x_{k}}^{n} \mathcal{L}_{x_{k}} w^{n} \leq \frac{1}{2} C_{1} |\nabla w^{n}|_{d}^{2} + \frac{1}{2} \theta |D^{2} w^{n}|_{d \times d}^{2} + \frac{1}{2} C_{2}$$

$$\leq \frac{1}{2} \Big[C_{1} |\nabla u|_{d}^{2} + \theta |D^{2} w^{n}|_{d \times d}^{2} + C_{2} + C_{1} \|\nabla \hat{w}^{n}\|_{C^{0}(\overline{\mathcal{O}}_{m})} \Big(2 \|\nabla u\|_{C^{0}(\overline{\mathcal{O}}_{m})} + \|\nabla \hat{w}^{n}\|_{C^{0}(\overline{\mathcal{O}}_{m})} \Big) \Big],$$

with

(4.47)
$$C_1 = C_1(m) \coloneqq 1 + d^4 A_m^2 \theta^{-1} + 2dA_m$$
 and $C_2 = C_2(m) \coloneqq \|\nabla h_m\|_{C^0(\overline{\mathcal{O}}_m)}^2$

The expression involving $\|\nabla \hat{w}^n\|_{C^0}$ can be collected into R_n in (4.36).

It remains to find an upper bound for $\lambda \left(ru + \psi_{\varepsilon}(\zeta_n) - \frac{1}{\delta}(g_m - u)^+ - h_m \right)$. Since $h_m \ge 0$ and $(g_m - u)^+ \ge 0$ and taking $M_1 = M_1(m) > 0$ as in Remark 4.6 we have

(4.48)
$$\lambda \left(ru + \psi_{\varepsilon}(\zeta_n) - \frac{1}{\delta}(g_m - u)^+ - h_m \right) \le \lambda \left(rM_1 + \psi_{\varepsilon}(\zeta_n) \right).$$

By convexity of ψ_{ε} and since $\psi_{\varepsilon}(0) = 0$, we have

$$(4.49) \quad \psi_{\varepsilon}(\zeta_{n}) \leq \psi_{\varepsilon}'(\zeta_{n})(|\nabla u|_{d}^{2} - f_{m}^{2}) \leq \psi_{\varepsilon}'(\zeta_{n})|\nabla u|_{d}^{2} \leq \psi_{\varepsilon}'(\bar{\zeta}_{n})|\nabla u|_{d}^{2} + \left|\psi_{\varepsilon}'(\zeta_{n}) - \psi_{\varepsilon}'(\bar{\zeta}_{n})\right||\nabla u|_{d}^{2} \\ \leq \psi_{\varepsilon}'(\bar{\zeta}_{n})|\nabla u|_{d}^{2} + \left\|\psi_{\varepsilon}'(\zeta_{n}) - \psi_{\varepsilon}'(\bar{\zeta}_{n})\right\|_{C^{0}(\overline{\mathcal{O}}_{m})} \|\nabla u\|_{C^{0}(\overline{\mathcal{O}}_{m})}^{2}.$$

Combining (4.42), (4.46), (4.48) and (4.49) we obtain (4.36).

Proof of (4.38). By Cauchy-Schwarz inequality and recalling that $\hat{w}^n = u - w^n$ we have

$$(4.50) \qquad -2\langle \nabla w^{n}, \nabla (|\nabla u^{n}|_{d}^{2} - f_{m}^{2})\rangle \\ \leq -2\langle \nabla w^{n}, \nabla |\nabla u^{n}|_{d}^{2}\rangle + 2|\nabla w^{n}|_{d}|\nabla f_{m}^{2}|_{d} \\ \leq -2\langle \nabla w^{n}, \nabla |\nabla u^{n}|_{d}^{2}\rangle + 2|\nabla u|_{d}|\nabla f_{m}^{2}|_{d} + 2\|\nabla \hat{w}^{n}\|_{C^{0}(\overline{\mathcal{O}}_{m})}\|\nabla f_{m}^{2}\|_{C^{0}(\overline{\mathcal{O}}_{m})}.$$

The first term on the right-hand side of (4.50) can be written as

$$\left\langle \nabla w^n, \nabla |\nabla u^n|_d^2 \right\rangle = \left\langle \nabla u, \nabla |\nabla u^n|_d^2 \right\rangle - \left\langle \nabla \hat{w}^n, \nabla |\nabla u^n|_d^2 \right\rangle.$$

The k-th entry of the vector $\nabla |\nabla u^n|_d^2$ reads $(|\nabla u^n|_d^2)_{x_k} = 2\langle \nabla u^n, \nabla u_{x_k}^n \rangle$ and therefore

$$\begin{split} \left| \nabla |\nabla u^{n}|_{d}^{2} \right|_{d}^{2} &\leq 4 \sum_{k=1}^{d} \left| \langle \nabla u^{n}, \nabla u_{x_{k}}^{n} \rangle \right|^{2} \leq 4 \sum_{k=1}^{d} |\nabla u^{n}|_{d}^{2} |\nabla u_{x_{k}}^{n}|_{d}^{2} \\ &\leq 4 |\nabla u^{n}|_{d}^{2} |D^{2}u^{n}|_{d \times d}^{2} \leq 2 \Big(\|\nabla u^{n}\|_{C^{0}(\overline{\mathcal{O}}_{m})}^{4} + \|D^{2}u^{n}\|_{C^{0}(\overline{\mathcal{O}}_{m})}^{4} \Big). \end{split}$$

Thus, using $\sqrt{a^2 + b^2} \le |a| + |b|$, we have

(4.51)
$$\langle \nabla w^n, \nabla | \nabla u^n |_d^2 \rangle \ge \langle \nabla u, \nabla | \nabla u^n |_d^2 \rangle - \sqrt{2} \| \nabla \hat{w}^n \|_{C^0(\overline{\mathcal{O}}_m)} \Big(\| \nabla u^n \|_{C^0(\overline{\mathcal{O}}_m)}^2 + \| D^2 u^n \|_{C^0(\overline{\mathcal{O}}_m)}^2 \Big).$$

For the term $\langle \nabla u, \nabla | \nabla u^n |_d^2 \rangle$ we argue as follows:

$$(4.52) \begin{aligned} u_{x_k} \partial_{x_k} |\nabla u^n|_d^2 &= 2u_{x_k} \langle \nabla u^n, \nabla u_{x_k}^n \rangle \\ &= 2u_{x_k} \langle \nabla u^n - \nabla u, \nabla u_{x_k}^n \rangle + 2u_{x_k} \langle \nabla u, \nabla u_{x_k}^n - \nabla u_{x_k} \rangle + 2u_{x_k} \langle \nabla u, \nabla u_{x_k} \rangle \\ &\geq -2 \|u^n - u\|_{C^{1,2,\gamma}(\overline{\mathcal{O}}_m)} \|\nabla u\|_{C^0(\overline{\mathcal{O}}_m)} \left(\|D^2 u^n\|_{C^0(\overline{\mathcal{O}}_m)} + \|\nabla u\|_{C^0(\overline{\mathcal{O}}_m)} \right) \\ &+ 2u_{x_k} \langle \nabla u, \nabla u_{x_k} \rangle. \end{aligned}$$

For the last term above, recall that all expressions are evaluated at $(t_n^{\lambda}, x_n^{\lambda})$, which is a stationary point for $v^{\lambda,n}$ in the spatial coordinates. Thus, $2\langle \nabla w^n, \nabla w_{x_i}^n \rangle = \lambda u_{x_i}$ by (4.30) and we get

$$2u_{x_k} \langle \nabla u, \nabla u_{x_k} \rangle = 2u_{x_k} \langle \nabla (w^n + \hat{w}^n), \nabla (w^n_{x_k} + \hat{w}^n_{x_k}) \rangle$$

$$(4.53) = 2u_{x_k} \langle \nabla w^n, \nabla w^n_{x_k} \rangle + 2u_{x_k} \langle \nabla w^n, \nabla \hat{w}^n_{x_k} \rangle + 2u_{x_k} \langle \nabla \hat{w}^n, \nabla (w^n_{x_k} + \hat{w}^n_{x_k}) \rangle$$

$$\geq \lambda u^2_{x_k} - 2 \| \hat{w}^n \|_{C^{1,2,\gamma}(\overline{\mathcal{O}}_m)} \| \nabla u \|_{C^0(\overline{\mathcal{O}}_m)} \left(\| \nabla w^n \|_{C^0(\overline{\mathcal{O}}_m)} + \| D^2 u \|_{C^0(\overline{\mathcal{O}}_m)} \right).$$

Summing over all k's in (4.52) we have

$$\langle \nabla u, \nabla | \nabla u^n |_d^2 \rangle \geq \lambda | \nabla u |_d^2 - 2d \| \hat{w}^n \|_{C^{1,2,\gamma}(\overline{\mathcal{O}}_m)} \| \nabla u \|_{C^0(\overline{\mathcal{O}}_m)} \left(\| \nabla w^n \|_{C^0(\overline{\mathcal{O}}_m)} + \| D^2 u \|_{C^0(\overline{\mathcal{O}}_m)} \right)$$
$$- 2d \| u^n - u \|_{C^{1,2,\gamma}(\overline{\mathcal{O}}_m)} \| \nabla u \|_{C^0(\overline{\mathcal{O}}_m)} \left(\| D^2 u^n \|_{C^0(\overline{\mathcal{O}}_m)} + \| \nabla u \|_{C^0(\overline{\mathcal{O}}_m)} \right).$$

Plugging this expression back into (4.51) we get

$$(4.54) \qquad \langle \nabla w^{n}, \nabla | \nabla u^{n} |_{d}^{2} \rangle \\ \geq \lambda | \nabla u |_{d}^{2} - 2d \| \hat{w}^{n} \|_{C^{1,2,\gamma}(\overline{\mathcal{O}}_{m})} \| \nabla u \|_{C^{0}(\overline{\mathcal{O}}_{m})} \left(\| \nabla w^{n} \|_{C^{0}(\overline{\mathcal{O}}_{m})} + \| D^{2} u \|_{C^{0}(\overline{\mathcal{O}}_{m})} \right) \\ - 2d \| u^{n} - u \|_{C^{1,2,\gamma}(\overline{\mathcal{O}}_{m})} \| \nabla u \|_{C^{0}(\overline{\mathcal{O}}_{m})} \left(\| D^{2} u^{n} \|_{C^{0}(\overline{\mathcal{O}}_{m})} + \| \nabla u \|_{C^{0}(\overline{\mathcal{O}}_{m})} \right) \\ - \sqrt{2} \| \nabla \hat{w}^{n} \|_{C^{0}(\overline{\mathcal{O}}_{m})} \left(\| \nabla u^{n} \|_{C^{0}(\overline{\mathcal{O}}_{m})}^{2} + \| D^{2} u^{n} \|_{C^{0}(\overline{\mathcal{O}}_{m})}^{2} \right).$$

Since $w^n \to u$ and $u^n \to u$ in $C^{1,2,\gamma}(\overline{\mathcal{O}}_m)$ as $n \to \infty$ for all $\gamma \in (0,\alpha)$ (Lemma 4.7), then all the terms in (4.50) and (4.54) depending on \hat{w}^n and $u^n - u$ can be collected in a remainder $\tilde{R}_n = \tilde{R}_n(m,\varepsilon,\delta)$ that goes to zero as $n \to \infty$. So, (4.50) and (4.54) give us

$$-2\langle \nabla w^n, \nabla (|\nabla u^n|_d^2 - f_m^2) \rangle \le -2\lambda |\nabla u|_d^2 + 2|\nabla u|_d |\nabla f_m^2|_d + \tilde{R}_n,$$

as claimed in (4.38).

4.4. Solution to the penalised problem. In this section we prove that Problem B admits a unique solution. The proof is based on a fixed point argument that requires the a priori estimates derived in Sections 4.2 and 4.3 as well as the next bound.

Proposition 4.10. Let $u_m^{\varepsilon,\delta}$ be a solution of Problem B. For any $\beta \in (0,1)$ there is $M_4 = M_4(m,\varepsilon,\delta,\beta)$ such that

$$(4.55) \|u_m^{\varepsilon,\delta}\|_{C^{0,1,\beta}(\overline{\mathcal{O}}_m)} \le M_4.$$

Proof. Let $u = u_m^{\varepsilon,\delta}$ for simplicity. Then, u can be seen as the unique solution φ of the linear PDE

$$\partial_t \varphi + \mathcal{L} \varphi - r \varphi = -h_m - \frac{1}{\delta} (g_m - u)^+ + \psi_{\varepsilon} (|\nabla u|_d^2 - f_m^2), \text{ on } \mathcal{O}_m$$

with boundary conditions $\varphi(t, x) = 0$ for $x \in \partial B_m$ and $\varphi(T, x) = g_m(T, x)$ for all $x \in B_m$. The theory of strong solutions for linear parabolic PDEs (see [6, Thm. 2.6.5 and Rem. 2.6.4]) gives

$$(4.56) \quad \|u\|_{W^{1,2,p}(\mathcal{O}_m)} \le C\left(\left\|h_m + \frac{1}{\delta}(g_m - u)^+ - \psi_{\varepsilon}(|\nabla u|_d^2 - f_m^2)\right\|_{L^p(\mathcal{O}_m)} + \|g_m\|_{W^{1,2,p}(\mathcal{O}_m)}\right)$$

for any $p \in (1, \infty)$, with C a constant independent of ε and δ . Denoting $|\mathcal{O}_m|$ the volume of the set \mathcal{O}_m , thanks to Proposition 4.9 and $u \ge 0$ we have

$$\|u\|_{W^{1,2,p}(\mathcal{O}_m)} \le C |\mathcal{O}_m|^{\frac{1}{p}} \left(\left\|h_m + \frac{1}{\delta}g_m\right\|_{C^0(\overline{\mathcal{O}}_m)} + \frac{1}{\varepsilon}M_3^2 \right) + C \|g_m\|_{W^{1,2,p}(\mathcal{O}_m)} < \infty,$$

having also used that $\psi_{\varepsilon}(x) \leq \frac{1}{\varepsilon}x$ for all $x \geq 0$.

Since p is arbitrary, then Sobolev embedding (2.2) guarantees that for any $\beta \in (0, 1)$ there exists a constant $M_4 = M_4(m, \varepsilon, \delta, \beta)$ such that (4.55) holds.

Theorem 4.11. There exists a unique solution $u_m^{\varepsilon,\delta}$ of Problem B.

Proof. Uniqueness is by Corollary 4.4. Existence will be proved refining arguments from [28, Prop. 1.2]. Fix $\varepsilon, \delta \in (0, 1)$ and $m \in \mathbb{N}$. Let us define a subset of $C^{0,1,\alpha}(\overline{\mathcal{O}}_m)$ as

$$C^{0,1,\alpha}_{*}(\overline{\mathcal{O}}_{m}) \coloneqq \left\{ \varphi \in C^{0,1,\alpha}(\overline{\mathcal{O}}_{m}) \middle| \begin{array}{l} \varphi = 0 \text{ on } \{T\} \times \partial B_{m} \text{ and} \\ \nabla \varphi = \mathbf{0} \text{ on } \{T\} \times \partial B_{m} \end{array} \right\}$$

The space $C^{0,1,\alpha}_*(\overline{\mathcal{O}}_m)$ is closed under the norm $\|\cdot\|_{C^{0,1,\alpha}(\overline{\mathcal{O}}_m)}$. Hence $(C^{0,1,\alpha}_*(\overline{\mathcal{O}}_m), \|\cdot\|_{C^{0,1,\alpha}(\overline{\mathcal{O}}_m)})$ is a Banach space. For $C^{1,2,\alpha}_*(\overline{\mathcal{O}}_m) \coloneqq (C^{1,2,\alpha} \cap C^{0,1,\alpha}_*)(\overline{\mathcal{O}}_m)$, also $(C^{1,2,\alpha}_*(\overline{\mathcal{O}}_m), \|\cdot\|_{C^{1,2,\alpha}(\overline{\mathcal{O}}_m)})$ is a Banach space.

Given $\varphi \in C^{0,1,\alpha}_*(\overline{\mathcal{O}}_m)$ we consider the linear partial differential equation for $w = w^{\varphi}$:

(4.57)
$$\begin{cases} \partial_t w + \mathcal{L}w - rw = -h_m - \frac{1}{\delta} \left(g_m - \varphi \right)^+ + \psi_{\varepsilon} \left(|\nabla \varphi|_d^2 - f_m^2 \right), & \text{on } \mathcal{O}_m, \\ w(t, x) = g_m(t, x), & (t, x) \in \partial_P \mathcal{O}_m. \end{cases}$$

For $x \in \partial B_m$ the compatibility condition

$$\lim_{s\uparrow T} (\partial_t g_m + \mathcal{L}g_m - rg_m)(s, x) = \left[-h_m - \frac{1}{\delta} \left(g_m - \varphi \right)^+ + \psi_{\varepsilon} \left(|\nabla \varphi|_d^2 - f_m^2 \right) \right](T, x),$$

holds, with both sides of the expression being equal to zero. Indeed, on the left-hand side, properties of the cut-off function $\xi_{m-1} \in C_c^{\infty}(B_m)$ guarantee $g_m = \partial_{x_i}g_m = \partial_{x_i,x_j}g_m = 0$ on $[0,T] \times \partial B_m$, hence also $\partial_t g_m = 0$. On the right-hand side of the equation, we use that $h_m = \varphi = \partial_{x_i}\varphi = 0$ on $\{T\} \times \partial B_m$. Therefore (4.57) admits a unique solution in $C^{1,2,\alpha}(\overline{\mathcal{O}}_m)$ by [19, Thm. 3.3.7]. The boundary condition of the PDE implies w = 0 and $\nabla w = \mathbf{0}$ on $\{T\} \times \partial B_m$. Hence $w \in C_*^{1,2,\alpha}(\overline{\mathcal{O}}_m)$.

Define the operator $\Gamma : C^{0,1,\alpha}_*(\overline{\mathcal{O}}_m) \to C^{0,1,\alpha}_*(\overline{\mathcal{O}}_m)$ that maps φ to the solution of the PDE (4.57), i.e., $\Gamma[\varphi] = w^{\varphi}$. Next, we show that Γ has a fixed point by Schaefer's fixed point theorem (stated in Appendix for completeness). So Problem B admits a solution. We have $\Gamma[\varphi] \in C^{1,2,\alpha}_*(\overline{\mathcal{O}}_m) \Longrightarrow \Gamma[\varphi] \in C^{0,1,\beta}_*(\overline{\mathcal{O}}_m)$ for all $\beta \in (0,1)$ by (2.2). We must prove that Γ is continuous and compact in $C^{0,1,\alpha}_*(\overline{\mathcal{O}}_m)$. Consider a sequence $(\varphi_n)_{n\in\mathbb{N}} \subset C^{0,1,\alpha}_*(\overline{\mathcal{O}}_m)$ such that $\varphi_n \to \varphi$ in $C^{0,1,\alpha}_*(\overline{\mathcal{O}}_m)$. Let $F : \mathbb{R} \times \mathbb{R}^d \to \mathbb{R}$ be defined as

$$F(q,p) \coloneqq \psi_{\varepsilon} \left(|p|_d^2 - f_m^2 \right) - \frac{1}{\delta} (g_m - q)^+$$

Clearly $|\varphi|$ and $|\nabla \varphi|_d$ are bounded on $\overline{\mathcal{O}}_m$. Since F is locally Lipschitz, then standard estimates for Hölder norms allow to prove $F(\varphi_n, \nabla \varphi_n) \to F(\varphi, \nabla \varphi)$ in $C^{\gamma}(\overline{\mathcal{O}}_m)$ as $n \to \infty$ for any $\gamma \in (0, \alpha)$. By Lemma A.1 in Appendix, $\Gamma[\varphi_n] \to \Gamma[\varphi]$ in $C^{1,2,\gamma'}_*(\overline{\mathcal{O}}_m)$ for any $\gamma' \in (0,\gamma)$, as $n \to \infty$. Sobolev embedding (2.2) implies $\Gamma[\varphi_n] \to \Gamma[\varphi]$ in $C^{0,1,\beta}_*(\overline{\mathcal{O}}_m)$ for any $\beta \in (0,1)$, hence continuity of Γ .

For compactness, notice that Γ maps bounded sets of $C^{0,1,\alpha}_*(\overline{\mathcal{O}}_m)$ into bounded sets of $C^{1,2,\alpha}_*(\overline{\mathcal{O}}_m)$ by [19, Thms. 3.2.6 and 3.3.7]. Since bounded sets in $C^{1,2,\alpha}_*(\overline{\mathcal{O}}_m)$ are bounded in $C^{0,1,\beta}_*(\overline{\mathcal{O}}_m)$ for all $\beta \in (0,1)$ (see (2.2)), then $C^{1,2,\alpha}_*(\overline{\mathcal{O}}_m) \hookrightarrow C^{0,1,\alpha}_*(\overline{\mathcal{O}}_m)$ is a compact embedding.

It remains to prove that the set

$$\mathcal{B} \coloneqq \left\{ \varphi \in C^{0,1,\alpha}_*(\overline{\mathcal{O}}_m) \,\middle|\, \rho \Gamma[\varphi] = \varphi \text{ for some } \rho \in [0,1] \right\}$$

is bounded in $C^{0,1,\alpha}_*(\overline{\mathcal{O}}_m)$. If $\rho = 0$, then $\varphi = 0$. If $\rho\Gamma[\varphi] = \varphi$ for some $\rho \in (0,1]$, then φ satisfies

(4.58)
$$\begin{cases} \partial_t \varphi + \mathcal{L}\varphi - r\varphi = \rho \left[-h_m - \frac{1}{\delta} \left(g_m - \varphi \right)^+ + \psi_{\varepsilon} \left(|\nabla \varphi|_d^2 - f_m^2 \right) \right], & \text{on } \mathcal{O}_m, \\ \varphi = \rho g_m, & \text{on } \partial_P \mathcal{O}_m \end{cases}$$

The PDE in (4.58) is the same as the one in (4.3) but with h_m , ψ_{ε} , g_m , $\frac{1}{\delta}$ replaced by ρh_m , $\rho \psi_{\varepsilon}$, ρg_m , $\frac{\rho}{\delta}$. Then, all the results from this section and Proposition 4.10 apply to φ . In particular, $\|\varphi\|_{C^{0,1,\alpha}(\overline{\mathcal{O}}_m)} \leq M_4$ uniformly for all $\rho \in (0,1]$, with M_4 as in Proposition 4.10. Finally, Schaefer's fixed point theorem guarantees existence of the solution of Problem B, for every triple (ε, δ, m) .

5. Penalised Problem on Unbounded Domain and Further Estimates

We refine our a priori estimates on the solution of Problem B. First we shall make all bounds independent of m so that we can construct a solution to a penalised problem on unbounded domain as $m \to \infty$. Then we shall find bounds independent of ε and δ so as to pass to the limit for $\varepsilon, \delta \downarrow 0$ and obtain a solution to Problem A.

5.1. Estimates independent of m. First we bound $\nabla u_m^{\varepsilon,\delta}$ independently of m on each compact.

Proposition 5.1. Fix $m_0 \in \mathbb{N}$ and $q \ge m_0 + 2$. Let $u_q^{\varepsilon, \delta}$ be the unique solution of Problem B on \mathcal{O}_q . Then, there is $N_1 = N_1(m_0)$ independent of ε , δ and q, such that

(5.1)
$$\sup_{(t,x)\in\overline{\mathcal{O}}_{m_0}} |\nabla u_q^{\varepsilon,\delta}(t,x)|_d \le N_1.$$

Proof. For notational simplicity set $u = u_q^{\varepsilon,\delta}$, $\xi = \xi_{m_0}$ and $\|\cdot\|_0 = \|\cdot\|_{C^0(\overline{\mathcal{O}}_{m_0+1})}$. Since $q \ge m_0 + 2$, we have $f_q = f$, $g_q = g$ and $h_q = h$ on $\overline{\mathcal{O}}_{m_0+1}$. Let $\lambda_0 \in (0, \infty)$ be a constant depending on m_0 but independent of ε, δ, q , which will be chosen later. Let $v^{\lambda} \in C^{0,1,\alpha}(\overline{\mathcal{O}}_{m_0+1})$ be defined as

(5.2)
$$v^{\lambda}(t,x) \coloneqq \xi(x) \left| \nabla u(t,x) \right|_{d}^{2} - \lambda u(t,x), \quad \text{for } (t,x) \in \overline{\mathcal{O}}_{m_{0}+1}$$

for some $\lambda \in (0, \lambda_0]$. We will use later that

(5.3)
$$\sup_{(t,x)\in\overline{\mathcal{O}}_{m_0}} |\nabla u(t,x)|_d^2 \le \sup_{(t,x)\in\overline{\mathcal{O}}_{m_0+1}} \xi(x) |\nabla u(t,x)|_d^2 \le \sup_{(t,x)\in\overline{\mathcal{O}}_{m_0+1}} v^{\lambda}(t,x) + \lambda_0 ||u||_0,$$

where we also notice that $||u||_0 \le M_1$ with $M_1 = M_1(m_0+1)$ as in Remark 4.6.

Let $(t^{\lambda}, x^{\lambda}) \in \overline{\mathcal{O}}_{m_0+1}$ be a maximum point for v^{λ} . If $x^{\lambda} \in \partial B_{m_0+1}$ then $\xi(x^{\lambda}) = 0$ and $v(t^{\lambda}, x^{\lambda}) = -\lambda u(t^{\lambda}, x^{\lambda}) \leq 0$. If $t^{\lambda} = T$, then $v^{\lambda}(T, x^{\lambda}) \leq \xi(x^{\lambda}) |\nabla g(T, x^{\lambda})|_d^2 \leq ||f||_0^2$ (see Assumption 3.2). Thus,

in both cases (5.1) holds with $N_1 \geq (\|f\|_0^2 + \lambda_0 M_1)^{1/2}$. Defining Λ_{m_0+1} as in (4.26) but with λ_0 instead of λ_m , if $\Lambda_{m_0+1} \neq \emptyset$ the bound holds taking $\lambda \in \Lambda_{m_0+1}$. It remains to consider the case $\Lambda_{m_0+1} = \emptyset$, so that $(t^{\lambda}, x^{\lambda}) \in \mathcal{O}_{m_0+1}$ for all $\lambda \in (0, \lambda_0]$.

As in the proof of Proposition 4.9 we use the smooth approximation w^n of u, obtained from (4.19). Analogously, let us define $v^{\lambda,n} := \xi |\nabla w^n|_d^2 - \lambda u$ in $\overline{\mathcal{O}}_{m_0+1}$ and let $(t_n^\lambda, x_n^\lambda)_{n \in \mathbb{N}}$ be a sequence converging to (t^λ, x^λ) with $(t_n^\lambda, x_n^\lambda) \in \arg \max_{\overline{\mathcal{O}}_{m_0+1}} v^{\lambda,n}$. For any $\eta > 0$ there exists a neighbourhood $U_{\lambda,\eta} \cup (\{0\} \times V_{\lambda,\eta})$ of (t^λ, x^λ) such that

(5.4)
$$v^{\lambda}(t,x) > v^{\lambda}(t^{\lambda},x^{\lambda}) - \eta, \quad \text{for all } (t,x) \in U_{\lambda,\eta} \cup \left(\{0\} \times V_{\lambda,\eta}\right)$$

and $(t_n^{\lambda}, x_n^{\lambda}) \in U_{\lambda,\eta} \cup (\{0\} \times V_{\lambda,\eta})$ for sufficiently large *n*'s (with the convention $V_{\lambda,\eta} = \emptyset$ if $t^{\lambda} \neq 0$). Taking derivatives of $v^{\lambda,n}$ we have, for $1 \leq i, j \leq d$,

$$\begin{split} v_t^{\lambda,n} &= 2\xi \langle \nabla w^n, \nabla w_t^n \rangle - \lambda u_t; \\ v_{x_i}^{\lambda,n} &= 2\xi \langle \nabla w^n, \nabla w_{x_i}^n \rangle - \lambda u_{x_i} + \xi_{x_i} |\nabla w^n|_d^2; \\ v_{x_i x_j}^{\lambda,n} &= 2\xi \langle \nabla w_{x_i}^n, \nabla w_{x_j}^n \rangle + 2\xi \langle \nabla w^n, \nabla w_{x_i x_j}^n \rangle \\ &- \lambda u_{x_i x_j} + \xi_{x_i x_j} |\nabla w^n|_d^2 + 2 \left(\xi_{x_i} \langle \nabla w^n, \nabla w_{x_j}^n \rangle + \xi_{x_j} \langle \nabla w^n, \nabla w_{x_i}^n \rangle \right). \end{split}$$

Since $(t_n^{\lambda}, x_n^{\lambda}) \in \mathcal{O}_{m_0+1}$ then (4.23) gives $(\partial_t v^{\lambda,n} + \mathcal{L}v^{\lambda,n})(t_n^{\lambda}, x_n^{\lambda}) \leq 0$. Substituting the expressions for the derivatives of $v^{\lambda,n}$, some tedious but straightforward calculations and symmetry of a_{ij} give

(5.5)
$$0 \ge 2\xi \langle \nabla w^n, (\partial_t + \mathcal{L})(\nabla w^n) \rangle - \lambda (\partial_t u + \mathcal{L}u) + \xi \sum_{i,j=1}^d a_{ij} \langle \nabla w^n_{x_i}, \nabla w^n_{x_j} \rangle$$
$$+ (\mathcal{L}\xi) |\nabla w^n|_d^2 + 2 \sum_{i,j=1}^d a_{ij} \xi_{x_i} \langle \nabla w^n, \nabla w^n_{x_j} \rangle,$$

where $(\partial_t + \mathcal{L})(\nabla w^n)$ denotes the vector with entries $(\partial_t + \mathcal{L})w_{x_k}^n$ for $k = 1, \ldots, d$. Here we omit the dependence on $(t_n^{\lambda}, x_n^{\lambda})$ for notational convenience.

The expressions for $(\partial_t + \mathcal{L})w_{x_k}^n$ and $(\partial_t + \mathcal{L})u$ are the same as in (4.34) and (4.3), respectively. A lower bound for $\sum a_{ij} \langle \nabla w_{x_i}^n, \nabla w_{x_j}^n \rangle$ was also obtained in (4.32) but with θ therein replaced by $\theta = \theta_{B_{m_0+1}}$. Thus, from (5.5) we get

$$(5.6) \qquad 0 \ge \xi \theta |D^2 w^n|_{d \times d}^2 + 2\xi \Big[r |\nabla w^n|^2 + \psi_{\varepsilon}'(\bar{\zeta}_n) \langle \nabla w^n, \nabla (|\nabla u^n|_d^2 - f^2) \rangle - \frac{1}{\delta} \chi_n'(g-u) \langle \nabla w^n, \nabla (g-u) \rangle - \langle \nabla w^n, \nabla h \rangle - \sum_{k=1}^d w_{x_k}^n \mathcal{L}_{x_k} w^n \Big] - \lambda \Big(ru + \psi_{\varepsilon}(\zeta_n) - h - \frac{1}{\delta}(g-u)^+ \Big) + (\mathcal{L}\xi) |\nabla w^n|_d^2 + 2 \sum_{i,j=1}^d a_{ij} \xi_{x_i} \langle \nabla w^n, \nabla w_{x_j}^n \rangle,$$

where $\bar{\zeta}_n = (|\nabla u^n|_d^2 - f^2)(t_n^{\lambda}, x_n^{\lambda}) - \frac{1}{n}$ and $\zeta_n = (|\nabla u|_d^2 - f^2)(t_n^{\lambda}, x_n^{\lambda})$. Up to a factor ξ , the expression above is the analogue of (4.35) but with two additional terms. As in (4.36) we have

(5.7)
$$2\xi \Big[r |\nabla w^n|_d^2 - \frac{1}{\delta} \chi'_n(g-u) \langle \nabla w^n, \nabla (g-u) \rangle - \langle \nabla w^n, \nabla h \rangle - \sum_{k=1}^d w^n_{x_k} \mathcal{L}_{x_k} w^n \Big] \\ -\lambda \Big(ru + \psi_{\varepsilon}(\zeta_n) - \frac{1}{\delta}(g-u)^+ - h \Big) \\ \ge -C_1 \xi |\nabla u|_d^2 - \frac{\theta}{2} \xi |D^2 w^n|_{d\times d}^2 - \xi C_2 - \lambda r M_1 - \lambda \psi'_{\varepsilon}(\bar{\zeta}_n) |\nabla u|_d^2 - R_n,$$

where $R_n \to 0$ as $n \to \infty$ uniformly on $\overline{\mathcal{O}}_{m_0+1}$. We notice that, differently to (4.36), we have a factor 1/2 multiplying $|D^2 w^n|_{d\times d}^2$. That of course is obtained by adjusting the constant $C_1 = (1 + 2d^4 A_{m_0+1}^2 \theta^{-1} + 2dA_{m_0+1})$ (see (4.47)) with A_{m_0+1} as in (4.45) and $C_2 = \|\nabla h\|_0$.

The last term on the right-hand side of (5.6) can be easily bounded. Set $\bar{a}_0 := \max_{i,j} ||a_{ij}||_0$, which is finite by continuity of a_{ij} , and recall that $|\nabla \xi|_d^2 \leq C_0 \xi$ by (4.1). Then

(5.8)
$$2\sum_{i,j=1}^{d} a_{ij}\xi_{x_i} \langle \nabla w^n, \nabla w_{x_j}^n \rangle \ge -2\bar{a}_0 d \sum_{j=1}^{d} |\nabla \xi|_d |\nabla w^n|_d |\nabla w_{x_j}^n|_d \\\ge -2\bar{a}_0 d^2 \sqrt{C_0 \xi} |\nabla w^n|_d |D^2 w^n|_{d \times d} \ge -C_3 |\nabla w^n|_d^2 - \xi \frac{\theta}{2} |D^2 w^n|_{d \times d}^2,$$

where the final inequality is by $|ab| \leq pa^2 + \frac{b^2}{p}$ with $a = 2\bar{a}_0 d^2 \sqrt{C_0} |\nabla w^n|_d$, $b = \sqrt{\xi} |D^2 w^n|_{d \times d}$ and $p = 2/\theta$, and setting $C_3 = 8\theta^{-1}\bar{a}_0^2 d^4 C_0$. It is also easy to check that $||\mathcal{L}\xi||_0 \leq \kappa$ for some $\kappa = \kappa(||b||_0, ||\sigma||_0) > 0$, because the derivatives of ξ are bounded independently of m_0 .

Plugging (5.7) and (5.8) into (5.6) and setting $C_4 = C_3 + \kappa$ we have

(5.9)
$$0 \ge 2\xi \psi_{\varepsilon}'(\bar{\zeta}_n) \langle \nabla w^n, \nabla \left(|\nabla u^n|_d^2 - f^2 \right) \rangle - \xi C_1 |\nabla u|_d^2 - C_4 |\nabla w^n|_d^2 - \lambda \psi_{\varepsilon}'(\bar{\zeta}_n) |\nabla u|_d^2 - \xi C_2 - \lambda r M_1 - R_n.$$

Using (4.41) we have $\xi C_1 |\nabla u|_d^2 + C_4 |\nabla w^n|_d^2 \leq C_5 |\nabla u|_d^2 + \tilde{R}_n$, where $\tilde{R}_n \to 0$ as $n \to \infty$, uniformly on $\overline{\mathcal{O}}_{m_0+1}$ and $C_5 = C_1 + C_4$. It remains to find a lower bound for $\xi \langle \nabla w^n, \nabla (|\nabla u^n|_d^2 - f^2) \rangle$. By the same arguments as in (4.50), the bounds in (4.51) and (4.52) continue to hold, up to the inclusion of the multiplicative factor ξ . We have an additional term in the final expression in (4.53), because now $v_{x_k}^{\lambda,n} = 0$ gives $2\xi \langle \nabla w^n, \nabla w_{x_k}^n \rangle = \lambda u_{x_k} - \xi_{x_k} |\nabla w^n|_d^2$. So, the extra term appearing in (4.53) reads $-u_{x_k} \xi_{x_k} |\nabla w^n|_d^2$ and, similarly to (4.41), we get

$$-u_{x_k}\xi_{x_k}|\nabla w^n|_d^2 \ge -u_{x_k}\xi_{x_k}|\nabla u|_d^2 - |u_{x_k}||\xi_{x_k}||\nabla \hat{w}^n||_0 \left(2\|\nabla u\|_0 + \|\nabla \hat{w}^n\|_0\right)$$

In summary, we have

(5.10)
$$\xi \langle \nabla w^n, \nabla \left(|\nabla u^n|_d^2 - f^2 \right) \rangle \ge \lambda |\nabla u|_d^2 - \xi |\nabla u|_d |\nabla f^2|_d - |\nabla u|_d^3 |\nabla \xi|_d - \xi \hat{R}_n,$$

where $\dot{R}_n \to 0$ as $n \to \infty$, uniformly on $\overline{\mathcal{O}}_{m_0+1}$.

Substituting (5.10) into (5.9) and grouping together all terms that vanish as $n \to \infty$ we obtain

$$0 \ge 2\psi_{\varepsilon}'(\bar{\zeta}_n) \left[\frac{\lambda}{2} |\nabla u|_d^2 - \xi |\nabla u|_d |\nabla f^2|_d - |\nabla u|_d^3 |\nabla \xi|_d\right] - C_5 |\nabla u|_d^2 - C_2 - \lambda r M_1 - \bar{R}_n$$

where $\bar{R}_n \to 0$ as $n \to \infty$. Using that $\psi'_{\varepsilon}(\bar{\zeta}_n) \ge 1$ by the analogue of (4.28), for sufficiently large n, and multiplying the above expression by -1 we arrive at

$$0 \le \psi_{\varepsilon}'(\bar{\zeta}_n) \Big[-\lambda |\nabla u|_d^2 + 2\xi |\nabla u|_d |\nabla f^2|_d + 2|\nabla u|_d^3 |\nabla \xi|_d + C_5 |\nabla u|_d^2 + C_2 + \lambda r M_1 + \bar{R}_n \Big].$$

Using $2\xi |\nabla u|_d |\nabla f^2|_d \leq |\nabla u|_d^2 + |\nabla f^2|_d^2$ and $|\nabla f^2|_d \leq ||\nabla f^2||_0$, the above inequality leads to

$$(\lambda - 1 - 2|\nabla \xi|_d |\nabla u|_d - C_5) |\nabla u|_d^2 \le ||\nabla f^2||_0^2 + C_2 + \lambda r M_1 + \bar{R}_n.$$

Then, recalling that $|\nabla \xi|_d^2 \leq C_0 \xi$ (see (4.1)) and setting $\bar{\lambda} = 2 + 2\sqrt{C_0} \left\|\sqrt{\xi}|\nabla u|_d\right\|_0 + C_5$, we obtain

(5.11)
$$|\nabla u|_d^2(t_n^{\bar{\lambda}}, x_n^{\bar{\lambda}}) \le ||\nabla f||_0^2 + C_2 + \bar{\lambda}rM_1 + \bar{R}_n$$

The parameter $\bar{\lambda}$ is bounded from above independently of ∇u as follows. Let c > 0 be a constant that varies from one expression to the next, independent of $\bar{\lambda}$, ε , δ , but depending on d and the $C^0(\overline{\mathcal{O}}_{m_0+1})$ -norms of b, σ, g, h, f^2 , and their spatial gradient. From (5.2), (5.3) and (5.4), we obtain

$$\left\|\xi|\nabla u|_{d}^{2}\right\|_{0} \leq |\nabla u(t_{n}^{\bar{\lambda}}, x_{n}^{\bar{\lambda}})|_{d}^{2} + \bar{\lambda}M_{1} + \eta \leq c(1 + \bar{\lambda} + \bar{R}_{n}) + \eta \leq c(1 + \left\|\sqrt{\xi}|\nabla u|_{d}\right\|_{0} + \bar{R}_{n}) + \eta,$$

where the second inequality uses (5.11) and the final one the definition of $\overline{\lambda}$. Since $\|\xi|\nabla u\|_d^2\|_0 = \|\sqrt{\xi}|\nabla u\|_d\|_0^2$, then $\|\sqrt{\xi}|\nabla u\|_d\|_0 \le \max\{1, c(1+\overline{R}_n)\} + \eta$. As $n \uparrow \infty$ and $\eta \downarrow 0$ we get (5.1) from (5.3) and (5.11), choosing $\lambda_0 = 2 + 2(1+c)\sqrt{C_0} + C_5$, thanks to the bound on $\overline{\lambda}$.

The bound in Proposition 4.10 depends on m. The next result instead provides a uniform bound on any compact $\overline{\mathcal{O}}_{m_0}$ for $m_0 < m$. This can be achieved thanks to (5.1).

Proposition 5.2. Fix $m_0 \in \mathbb{N}$ and $q \ge m_0 + 3$ and let $u_q^{\varepsilon,\delta}$ be the unique solution of Problem *B* on \mathcal{O}_q . For any $p \in (d+2,\infty)$ and $\beta = 1 - (d+2)/p$ there is $N_2 = N_2(m_0,\varepsilon,\delta,p)$ such that

(5.12)
$$\|u_q^{\varepsilon,\delta}\|_{W^{1,2,p}(\mathcal{O}_{m_0})} + \|u_q^{\varepsilon,\delta}\|_{C^{0,1,\beta}(\overline{\mathcal{O}}_{m_0})} \le N_2$$

Proof. Define $\varphi(t, x) \coloneqq \xi_{m_0}(x) u_q^{\varepsilon,\delta}(t, x)$. Since $u_q^{\varepsilon,\delta}$ solves (4.3) and $f_q = f$, $g_q = g$ and $h_q = h$ on \mathcal{O}_{m_0+1} , then φ solves

$$\partial_t \varphi + \mathcal{L} \varphi - r \varphi = \xi_{m_0} \left[-h - \frac{1}{\delta} (g - u_q^{\varepsilon, \delta})^+ + \psi_{\varepsilon} \left(|\nabla u_q^{\varepsilon, \delta}|_d^2 - f^2 \right) \right] + Q, \quad \text{on } \mathcal{O}_{m_0 + 1}$$

where $Q(t,x) = u_q^{\varepsilon,\delta}(t,x)(\mathcal{L}\xi_{m_0})(x) + 2\langle a(x)\nabla\xi_{m_0}(x), \nabla u_q^{\varepsilon,\delta}(t,x)\rangle$, and with boundary conditions $\varphi(t,x) = 0$ for $x \in \partial B_{m_0+1}$ and $\varphi(T,x) = \xi_{m_0}(x)g(T,x)$ for $x \in B_{m_0+1}$. As in (4.56) we have

(5.13)
$$\|\varphi\|_{W^{1,2,p}(\mathcal{O}_{m_0+1})}$$

 $\leq C\Big(\|\xi_{m_0}[h+\frac{1}{\delta}(g-u_q^{\varepsilon,\delta})^+-\psi_{\varepsilon}(|\nabla u_q^{\varepsilon,\delta}|_d^2-f^2)]+Q\|_{L^p(\mathcal{O}_{m_0+1})}+\|\xi_{m_0}g\|_{W^{1,2,p}(\mathcal{O}_{m_0+1})}\Big)$

for any $p \in (1, \infty)$ and with C > 0 independent of ε , δ and q. Denoting $|\mathcal{O}_{m_0+1}|$ the volume of \mathcal{O}_{m_0+1} and using Proposition 5.1 we obtain

$$\begin{aligned} \|u_q^{\varepsilon,\delta}\|_{W^{1,2,p}(\mathcal{O}_{m_0})} &\leq \|\varphi\|_{W^{1,2,p}(\mathcal{O}_{m_0+1})} \\ &\leq C |\mathcal{O}_{m_0}|^{\frac{1}{p}} \left(\|h + \frac{1}{\delta}g + Q\|_{C^0(\overline{\mathcal{O}}_{m_0+1})} + \frac{1}{\varepsilon}(N_1)^2 \right) + C \|\xi_{m_0}g\|_{W^{1,2,p}(\mathcal{O}_{m_0+1})}, \end{aligned}$$

where the first inequality is due to $u_q^{\varepsilon,\delta} = \varphi$ on \mathcal{O}_{m_0} . Since Q is bounded on $\overline{\mathcal{O}}_{m_0+1}$ independently of q, the estimate above and Sobolev embedding (2.2) give us (5.12).

5.2. Penalised problem on unbounded domain. Combining the results obtained so far we can prove existence and uniqueness of the solution to a penalised problem on $\mathbb{R}^{d+1}_{0,T}$.

Problem C. Find $u = u^{\varepsilon,\delta}$ with $u^{\varepsilon,\delta} \in (C^{1,2,\alpha}_{Loc} \cap W^{1,2,p}_{\ell oc})(\mathbb{R}^{d+1}_{0,T})$, for any $p \in (1,\infty)$ and $\alpha \in (0,1)$ as in Assumption 3.2, that solves:

(5.14)
$$\begin{cases} \partial_t u + \mathcal{L}u - ru = -h - \frac{1}{\delta} (g - u)^+ + \psi_{\varepsilon} \left(|\nabla u|_d^2 - f^2 \right), & \text{on } [0, T) \times \mathbb{R}^d, \\ u(T, x) = g(T, x), & \text{for all } x \in \mathbb{R}^d. \end{cases}$$

Theorem 5.3. There exists a solution $u^{\varepsilon,\delta}$ of Problem C.

Proof. Fix n and take m > n + 3. From Proposition 5.2 we know that for any $\beta \in (0, 1)$ and $p \in (1, \infty)$, the norms $\|u_m^{\varepsilon,\delta}\|_{W^{1,2,p}(\mathcal{O}_n)}$ and $\|u_m^{\varepsilon,\delta}\|_{C^{0,1,\beta}(\overline{\mathcal{O}}_n)}$ are bounded by a constant independent of m. By weak compactness in $W^{1,2,p}$ and Ascoli-Arzelá's theorem we can then extract a sequence $(u_{m_k}^{\varepsilon,\delta})_{k\in\mathbb{N}}$ and there exists a function $u^{\varepsilon,\delta;n} \in W^{1,2,p}(\mathcal{O}_n)$ (both possibly depending on the choice of \mathcal{O}_n) such that, as $k \to \infty$ (and $m_k^n \to \infty$), we obtain

(5.15)
$$\begin{aligned} u_{m_k^{\varepsilon,\delta}}^{\varepsilon,\delta} &\to u^{\varepsilon,\delta;n} \quad \text{and} \quad \nabla u_{m_k^{\varepsilon,\delta}}^{\varepsilon,\delta} \to \nabla u^{\varepsilon,\delta;n} \quad \text{in } C^{\alpha}(\overline{\mathcal{O}}_n), \\ \partial_t u_{m_k^{\varepsilon,\delta}}^{\varepsilon,\delta} &\to \partial_t u^{\varepsilon,\delta;n} \quad \text{and} \quad D^2 u_{m_k^{\varepsilon,\delta}}^{\varepsilon,\delta} \to D^2 u^{\varepsilon,\delta;n} \quad \text{weakly in } L^p(\mathcal{O}_n). \end{aligned}$$

Since the sequence $(u_{m_k^n}^{\varepsilon,\delta})_{k\in\mathbb{N}}$ is bounded also in the $W^{1,2,p}(\mathcal{O}_{n+1})$ -norm (perhaps by a larger constant), then up to selecting a further subsequence we have convergence as in (5.15) but with n replaced by n + 1. Therefore $u^{\varepsilon,\delta;n} = u^{\varepsilon,\delta;n+1}$ on \mathcal{O}_n . Using that $\overline{\mathcal{O}}_n \uparrow \mathbb{R}_{0,T}^{d+1}$ as $n \to \infty$ and iterating the extraction of further subsequences (if needed), we can uniquely define a limit function $u^{\varepsilon,\delta} \in (C_{\ell oc}^{0,1,\alpha} \cap W_{\ell oc}^{1,2,p})(\mathbb{R}_{0,T}^{d+1}).$

Fix n and take m > n. Multiply the PDE solved by $u_m^{\varepsilon,\delta}$ (see (4.3)) by a test function supported on $\overline{\mathcal{O}}_n$. Then passing to the limit along the subsequence constructed above, it is standard procedure to show that $u^{\varepsilon,\delta}$ satisfies the first equation in (5.14) in the a.e. sense on \mathcal{O}_n thanks to locally uniform convergence on compacts of $u_m^{\varepsilon,\delta}$ and $\nabla u_m^{\varepsilon,\delta}$ and the weak convergence of $\partial_t u_m^{\varepsilon,\delta}$ and $D^2 u_m^{\varepsilon,\delta}$. Since, this can be done for any \mathcal{O}_n and $u^{\varepsilon,\delta}(T,\cdot) = g(T,\cdot)$, then $u^{\varepsilon,\delta}$ solves (5.14) in the a.e. sense (it is a strong solution). It now remains to prove it is actually a classical solution.

Fix an arbitrary open bounded domain $\mathcal{O} \subset \mathbb{R}^{d+1}_{0,T}$ with smooth parabolic boundary $\partial_P \mathcal{O}$. Let $v \in C^{1,2,\alpha}_{Loc}(\mathcal{O})$ be the unique classical solution of the boundary value problem

$$\begin{cases} \partial_t v + \mathcal{L}v - rv = -h - \frac{1}{\delta} \left(g - u^{\varepsilon, \delta} \right)^+ + \psi_{\varepsilon} \left(|\nabla u^{\varepsilon, \delta}|_d^2 - f^2 \right), & \text{on } \mathcal{O}, \\ v(t, x) = u^{\varepsilon, \delta}(t, x), & \text{for } (t, x) \in \partial_P \mathcal{O} \end{cases}$$

Existence and uniqueness of such v is guaranteed by [19, Thm. 3.4.9] because

$$-h - \frac{1}{\delta} (g - u^{\varepsilon, \delta})^+ + \psi_{\varepsilon} (|\nabla u^{\varepsilon, \delta}|_d^2 - f^2) \in C^{\alpha}(\overline{\mathcal{O}})$$

and \mathcal{L} is uniformly elliptic on $\overline{\mathcal{O}}$ with continuously differentiable coefficients (Assumption 3.1). Since v is also a strong solution, then $v - u^{\varepsilon,\delta} \in W^{1,2,p}(\mathcal{O})$ is a strong solution of $\partial_t w + \mathcal{L}w - rw = 0$ in \mathcal{O} with w = 0 on $\partial_P \mathcal{O}$. It follows that $\|v - u^{\varepsilon,\delta}\|_{W^{1,2,p}(\mathcal{O})} = 0$ by the same estimate as in (4.56). By arbitrariness of \mathcal{O} we can choose a $C_{Loc}^{1,2,\alpha}$ -representative of $u^{\varepsilon,\delta}$, as claimed. \Box

We now give a probabilistic representation for $u^{\varepsilon,\delta}$ analogue of (4.8) but on unbounded domain. For $(n,\nu) \in \mathcal{A}_t^{\circ}$ and $w \in \mathcal{T}_t^{\delta}$ let us denote by $\mathcal{J}_{t,x}^{\varepsilon,\delta}(n,\nu,w)$ a payoff analogue of (4.7) but with ρ_m , g_m , h_m replaced by T - t, g, h, respectively, and with the Hamiltonian H_m^{ε} replaced by

(5.16)
$$H^{\varepsilon}(t,x,y) \coloneqq \sup_{p \in \mathbb{R}^d} \left\{ \langle y, p \rangle - \psi_{\varepsilon} \left(|p|_d^2 - f^2(t,x) \right) \right\}.$$

Notice that

(5.17)
$$t \mapsto H^{\varepsilon}(t, x, y)$$
 is non-increasing for all $(x, y) \in \mathbb{R}^d \times \mathbb{R}^d$

because $t \mapsto f(t, x)$ is non-increasing by Assumption 3.2. Moreover, taking $p = \varepsilon y/2$ in H^{ε} gives

(5.18)
$$H^{\varepsilon}(t,x,y) \ge \frac{\varepsilon}{2} |y|_d^2 - \psi_{\varepsilon} \left(\frac{\varepsilon^2}{4} |y|_d^2 - f^2(t,x)\right) \ge \frac{\varepsilon}{2} |y|_d^2 - \psi_{\varepsilon} \left(\frac{\varepsilon^2}{4} |y|_d^2\right) \ge \frac{\varepsilon}{4} |y|_d^2$$

Proposition 5.4. Let $u^{\varepsilon,\delta}$ be a solution of Problem C. Then

(5.19)
$$u^{\varepsilon,\delta}(t,x) = \inf_{(n,\nu)\in\mathcal{A}_t^\circ} \sup_{w\in\mathcal{T}_t^\delta} \mathcal{J}_{t,x}^{\varepsilon,\delta}(n,\nu,w) = \sup_{w\in\mathcal{T}_t^\delta} \inf_{(n,\nu)\in\mathcal{A}_t^\circ} \mathcal{J}_{t,x}^{\varepsilon,\delta}(n,\nu,w).$$

Proof. Fix $[(n,\nu),w] \in \mathcal{A}_t^{\circ} \times \mathcal{T}_t^{\delta}$. An application of Dynkin's formula to $R_{\rho_m}^w u^{\varepsilon,\delta}(t+\rho_m, X_{\rho_m}^{[n,\nu]})$ combined with (5.14) gives (recall $u^{\varepsilon,\delta} \in (W_{\ell oc}^{1,2,p} \cap C_{\ell oc}^{0,1,\beta})(\mathbb{R}_{0,T}^{d+1}))$

$$(5.20) \qquad u^{\varepsilon,\delta}(t,x) = \mathsf{E}_{x} \left[R^{w}_{\rho_{m}} u^{\varepsilon,\delta} \left(t + \rho_{m}, X^{[n,\nu]}_{\rho_{m}} \right) \right. \\ \left. + \int_{0}^{\rho_{m}} R^{w}_{s} \left(h + \frac{1}{\delta} \left(g - u^{\varepsilon,\delta} \right)^{+} - \psi_{\varepsilon} \left(|\nabla u^{\varepsilon,\delta}|^{2}_{d} - f^{2} \right) \right) \left(t + s, X^{[n,\nu]}_{s} \right) \, \mathrm{d}s \right. \\ \left. + \int_{0}^{\rho_{m}} R^{w}_{s} \left(w_{s} u^{\varepsilon,\delta} \left(t + s, X^{[n,\nu]}_{s} \right) - \left\langle n_{s} \dot{\nu}_{s}, \nabla u^{\varepsilon,\delta} \left(t + s, X^{[n,\nu]}_{s} \right) \right\rangle \right) \, \mathrm{d}s \right] .$$

By definition of the Hamiltonian H^{ε} in (5.16) we have

(5.21)
$$u^{\varepsilon,\delta}(t,x) \leq \mathsf{E}_x \Big[R^w_{\rho_m} u^{\varepsilon,\delta} \big(t + \rho_m, X^{[n,\nu]}_{\rho_m} \big) \Big] \\ + \mathsf{E}_x \Big[\int_0^{\rho_m} R^w_s \Big(h + \frac{1}{\delta} \big(g - u^{\varepsilon,\delta}\big)^+ + w_s u^{\varepsilon,\delta} + H^{\varepsilon}(\cdot, n_s \dot{\nu}_s) \Big) \big(t + s, X^{[n,\nu]}_s \big) \, \mathrm{d}s \Big].$$

Letting $m \uparrow \infty$ we have $\rho_m \uparrow T - t$, P_x -a.s. We can take the limit inside the second expectation by monotone convergence as all the terms under the integral are positive. By Lemma 4.5 the term under the first expectation has quadratic growth in $X^{[n,\nu]}$. Thanks to standard estimates for SDEs (see [29, Thm. 2.5.10]) there is a constant c > 0 independent of m such that

$$\mathsf{E}_{x} \left[\sup_{0 \le s \le T-t} |X_{s}^{[n,\nu]}|_{d}^{2} \right] \le c \left(1 + |x|_{d}^{2} + \mathsf{E}_{x}[|\nu_{T-t}|^{2}] \right)$$

Then, dominated convergence and $u^{\varepsilon,\delta}(T, \cdot) = g(T, \cdot)$ give us

By arguments as in the proof of Proposition 4.1, with $w^* \in \mathcal{T}_t^{\delta}$ defined as in (4.11) but with u and g_m replaced by $u^{\varepsilon,\delta}$ and g, respectively, we obtain $u^{\varepsilon,\delta}(t,x) \leq \mathcal{J}_{t,x}^{\varepsilon,\delta}(n,\nu,w^*)$. Therefore

(5.22)
$$u^{\varepsilon,\delta}(t,x) \le \sup_{w \in \mathcal{T}_t^{\delta}} \inf_{(n,\nu) \in \mathcal{A}_t^{\circ}} \mathcal{J}_{t,x}^{\varepsilon,\delta}(n,\nu,w)$$

As in Proposition 4.1, for the reverse inequality we set $X^* = X^{[n^*,\nu^*]}$ and denote

(5.23)
$$n_{s}^{*} \coloneqq \begin{cases} -\frac{\nabla u^{\varepsilon,\delta}(t+s,X_{s}^{*})}{|\nabla u^{\varepsilon,\delta}(t+s,X_{s}^{*})|_{d}}, & \text{if } \nabla u^{\varepsilon,\delta}(t+s,X_{s}^{*}) \neq \mathbf{0}, \\ \text{any unit vector}, & \text{if } \nabla u^{\varepsilon,\delta}(t+s,X_{s}^{*}) = \mathbf{0}, \\ \dot{\nu}_{s}^{*} \coloneqq 2\psi_{\varepsilon}^{\prime} \left(|\nabla u^{\varepsilon,\delta}(t+s,X_{s}^{*})|_{d}^{2} - f^{2}(t+s,X_{s}^{*}) \right) |\nabla u^{\varepsilon,\delta}(t+s,X_{s}^{*})|_{d}. \end{cases}$$

We claim here and will prove later that $(n^*, \nu^*) \in \mathcal{A}_t^{\circ}$ and the SDE for X^* admits a unique nonexploding strong solution. For (n^*, ν^*) equality holds in (5.21). As $m \uparrow \infty$ Fatou's lemma gives $u^{\varepsilon,\delta}(t,x) \geq \mathcal{J}_{t,x}(n^*, \nu^*, w)$, hence

(5.24)
$$u^{\varepsilon,\delta}(t,x) \ge \inf_{(n,\nu)\in\mathcal{A}_t^\circ} \sup_{w\in\mathcal{T}_t^\delta} \mathcal{J}_{t,x}(n,\nu,w).$$

Combining (5.22) and (5.24) we conclude.

It remains to check that $(n^*, \nu^*) \in \mathcal{A}_t^{\circ}$ and X^* is non-exploding. We use an argument from [34, Lemma 13.7]. Let $\zeta_m = \inf\{s \ge 0 : |X_s^*|_d \ge m\}$. On the random time-interval $[0, \zeta_m \land (T-t)]$ the process X^* is well-defined and the pair (n^*, ν^*) is adapted because $u^{\varepsilon,\delta} \in C^{1,2,\alpha}_{Loc}(\mathcal{O}_m) \cap C^{0,1,\alpha}(\overline{\mathcal{O}}_m)$. Notice that $\zeta_k \le \zeta_{k+1}$ and it may occur $\zeta_\infty := \lim_{k \to \infty} \zeta_k < T-t$ with positive probability. Moreover $\rho_m = \zeta_m \land (T-t)$ in (5.20) and let us take $w \equiv 0$ therein. By construction, for $s \in [0, \zeta_m \land (T-t)]$

$$-\left\langle n_s^* \dot{\nu^*}_s, \nabla u^{\varepsilon,\delta}(t+s, X_s^*) \right\rangle - \psi_{\varepsilon} \left(|\nabla u^{\varepsilon,\delta}|_d^2 - f^2 \right) (t+s, X_s^*) = H^{\varepsilon}(t+s, X_s^*, n_s^* \dot{\nu^*}_s).$$

Then, by positivity of all remaining terms in (5.20)

$$u^{\varepsilon,\delta}(t,x) \ge \mathsf{E}_x\Big[\int_0^{\zeta_m \wedge (T-t)} e^{-rs} H^{\varepsilon}(t+s, X_s^*, n_s^* \dot{\nu}_s^*) \mathrm{d}s\Big]$$

By positivity of H^{ε} and monotone convergence, we can let $m \uparrow \infty$ and preserve the inequality while the integral in time extends to $\zeta_{\infty} \land (T-t)$. Combining with (5.18) and Lemma 4.5 we have

(5.25)
$$\frac{\varepsilon}{4}\mathsf{E}_x\Big[\int_0^{\zeta_\infty\wedge(T-t)} e^{-rs} |\dot{\nu}_s^*|^2 \,\mathrm{d}s\Big] \le u^{\varepsilon,\delta}(t,x) \le K_3(1+|x|_d^2).$$

Since $\dot{\nu}^* \ge 0$ then $s \mapsto \nu_s^*$ is non-decreasing and

$$|\nu_{\zeta_m \wedge (T-t)}^*|^2 = 2 \int_0^{\zeta_m \wedge (T-t)} \nu_s^* \dot{\nu}_s^* \, \mathrm{d}s \le \int_0^{\zeta_m \wedge (T-t)} |\nu_s^*|^2 \, \mathrm{d}s + \int_0^{\zeta_m \wedge (T-t)} |\dot{\nu}_s^*|^2 \, \mathrm{d}s,$$

where we used $2ab \leq a^2 + b^2$. By Gronwall's lemma and taking expectations we obtain

$$\mathsf{E}_{x}\big[|\nu_{\zeta_{m}\wedge(T-t)}^{*}|^{2}\big] \leq e^{T}\mathsf{E}_{x}\Big[\int_{0}^{\zeta_{m}\wedge(T-t)}|\dot{\nu}_{s}^{*}|^{2}\,\mathrm{d}s\Big] \leq e^{T(1+r)}\mathsf{E}_{x}\Big[\int_{0}^{\zeta_{m}\wedge(T-t)}e^{-rs}|\dot{\nu}_{s}^{*}|^{2}\,\mathrm{d}s\Big].$$

Combining with (5.25) and letting $m \to \infty$, Fatou's lemma gives us

(5.26)
$$\mathsf{E}_{x}\left[|\nu_{\zeta_{\infty}\wedge(T-t)}^{*}|^{2}\right] \leq 4e^{T(1+r)}\varepsilon^{-1}K_{3}(1+|x|_{d}^{2})$$

Linear growth of (b, σ) and well-posedness of $X^*_{s \wedge \zeta_m}$ give, by Markov inequality and standard bounds,

$$\begin{aligned} \mathsf{P}_{x}(\zeta_{m} < T - t) \leq & \frac{1}{m^{2}} \mathsf{E}_{x} \Big[\sup_{s \in [0, \zeta_{m} \land (T - t)]} |X_{s}^{*}|_{d}^{2} \Big] \\ \leq & \frac{C}{m^{2}} \Big(1 + |x|_{d}^{2} + \mathsf{E}_{x} \big[\big| \nu_{\zeta_{m} \land (T - t)}^{*} \big|^{2} \big] \Big) \leq \frac{C}{m^{2}} c(\varepsilon) (1 + |x|_{d}^{2}), \end{aligned}$$

where C > 0 depends only on T and D_1 from Assumption 3.1, and $c(\varepsilon)$ depends on the constants from (5.26). Since $\zeta_m \uparrow \zeta_\infty$, then $\mathsf{P}_x(\zeta_m < T - t) \downarrow \mathsf{P}_x(\zeta_\infty \leq T - t)$ as $m \to \infty$ and by taking limits in the expression above we conclude $\mathsf{P}_x(\zeta_\infty \leq T - t) = 0$. Thus, X_s^* is well-defined for all $s \in [0, T - t]$ and $\mathsf{E}_x[|\nu_{T-t}^*|^2] < \infty$, by (5.26) implying $(n^*, \nu^*) \in \mathcal{A}_t^\circ$ as claimed. \Box

Proposition 5.4 implies that Problem C admits a *unique* solution and that the triple $[(n^*, \nu^*), w^*]$ is optimal in (5.19). By arguments as in the proof of Proposition 4.3 we also obtain the next result.

Proposition 5.5. Let $u^{\varepsilon,\delta}$ be the unique solution of Problem C. Then

$$\begin{split} u^{\varepsilon,\delta}(t,x) &= \inf_{(n,\nu)\in\mathcal{A}_t^\circ} \mathsf{E}_x \bigg[e^{-(r+\delta^{-1})(T-t)} g(T, X_{T-t}^{[n,\nu]}) \\ &+ \int_0^{T-t} e^{-(r+\delta^{-1})s} \big[h + \frac{1}{\delta}g \vee u^{\varepsilon,\delta} + H^{\varepsilon}(\,\cdot\,, n_s \dot{\nu}_s) \big](t+s, X_s^{[n,\nu]}) \,\mathrm{d}s \bigg], \end{split}$$

and the pair (n^*, ν^*) from (5.23) is optimal.

5.3. Refined estimates independent of ε and δ . Here we develop bounds for the penalty terms in the PDE of Problem C which are independent of ε and δ .

Lemma 5.6. For K_2 as in (3.10) we have

(5.27)
$$\frac{1}{\delta} \left\| \left(g - u^{\varepsilon, \delta} \right)^+ \right\|_{\infty} \le K_2.$$

Proof. For any $(n, \nu) \in \mathcal{A}_t^{\circ}$, the function g has the same probabilistic representation as in (4.22) but with g_m , ρ_m and w replaced by g, T - t and $1/\delta$, respectively. Combining that with the expression for $u^{\varepsilon,\delta}$ in Proposition 5.5, and recalling Θ defined in (3.10), we get

As in (4.5),
$$[\langle n_s \dot{\nu}_s, \nabla g \rangle + H^{\varepsilon}(\cdot, n_s \dot{\nu}_s)](t+s, X_s^{[n,\nu]}) \ge 0$$
, and observing that $g \lor u^{\varepsilon,\delta} - g \ge 0$ we get
 $u^{\varepsilon,\delta}(t,x) - g(t,x) \ge \inf_{(n,\nu)\in\mathcal{A}_t^{\circ}} \mathsf{E}_x \left[\int_0^{T-t} e^{-(r+\delta^{-1})s} \Theta(t+s, X_s^{[n,\nu]}) \, \mathrm{d}s \right] \ge -K_2 \frac{\delta}{r\delta+1},$

where K_2 was defined in (3.10). The above implies $\frac{1}{\delta}(g(t,x) - u^{\varepsilon,\delta}(t,x))^+ \leq K_2$ as needed. \Box

Next we give an upper bound on $\partial_t u^{\varepsilon,\delta}$. In the lemma below we understand

$$\partial_t u^{\varepsilon,\delta}(T,x) \coloneqq \lim_{s \to 0} \frac{u^{\varepsilon,\delta}(T,x) - u^{\varepsilon,\delta}(T-s,x)}{s}.$$

Lemma 5.7. There is $K_4 > 0$ only depending on K_0 and K_2 from Assumption 3.2 such that (5.28) $\partial_t u^{\varepsilon,\delta}(t,x) \leq K_4, \quad \text{for } (t,x) \in \mathbb{R}^{d+1}_{0,T}.$

Proof. Let $u = u^{\varepsilon,\delta}$ for simplicity and take $T \ge t_2 > t_1 \ge 0$. Let $w^{(2)} \in \mathcal{T}_{t_2}^{\delta}$ be optimal for the value function $u(t_2, x)$ and let $(n^{(1)}, \nu^{(1)}) \in \mathcal{A}_{t_1}^{\circ}$ be optimal for the value function $u(t_1, x)$. Set $X^{(1)} = X^{[n^{(1)}, \nu^{(1)}]}$ and notice that $w_s^{(1)} \coloneqq w_s^{(2)} \mathbb{1}_{\{s \le T - t_2\}}$ lies in $\mathcal{T}_{t_1}^{\delta}$ and $(n^{(1)}, \nu^{(1)})$ restricted to $[0, T - t_2]$ lies in $\mathcal{A}_{t_2}^{\circ}$. To simplify notation let us also set $[\Delta_{t_2, t_1} g](s, x) = g(t_2 + s, x) - g(t_1 + s, x)$ and analogously for $[\Delta_{t_2, t_1} h](s, x)$ and $[\Delta_{t_2, t_1} H^{\varepsilon}](s, x, y)$. Then

$$u(t_{2},x) - u(t_{1},x) \leq \mathcal{J}_{t_{2},x}^{\varepsilon,\delta}(n^{(1)},\nu^{(1)},w^{(2)}) - \mathcal{J}_{t_{1},x}^{\varepsilon,\delta}(n^{(1)},\nu^{(1)},w^{(1)})$$

$$\leq \mathsf{E}_{x} \bigg[R_{T-t_{2}}^{w^{(2)}}g(T,X_{T-t_{2}}^{(1)}) - R_{T-t_{1}}^{w^{(1)}}g(T,X_{T-t_{1}}^{(1)}) - \int_{T-t_{2}}^{T-t_{1}} R_{s}^{w^{(1)}} \Big(h + H^{\varepsilon}(\cdot,n_{s}^{(1)}\dot{\nu}_{s}^{(1)})\Big) (t_{1} + s,X_{s}^{(1)}) \,\mathrm{d}s \bigg]$$

$$(5.29) \qquad + \int_{0}^{T-t_{2}} R_{s}^{w^{(2)}} \Big([\Delta_{t_{2},t_{1}}h] + [\Delta_{t_{2},t_{1}}H^{\varepsilon}](\cdot,n_{s}^{(1)}\dot{\nu}_{s}^{(1)}) + w_{s}^{(2)}[\Delta_{t_{2},t_{1}}g] \Big) (s,X_{s}^{(1)}) \,\mathrm{d}s \bigg].$$

By (5.17) the Hamiltonian H^{ε} is non-increasing in time, so $[\Delta_{t_2,t_1}H^{\varepsilon}] \leq 0$. Next, we apply Dynkin's formula to $R_{T-t_1}^{w^{(1)}}g(T, X_{T-t_1}^{(1)})$ on the time interval $[T - t_2, T - t_1]$, to obtain

$$\mathsf{E}_{x} \Big[R_{T-t_{1}}^{w^{(1)}} g(T, X_{T-t_{1}}^{(1)}) \Big] = \mathsf{E}_{x} \Big[R_{T-t_{2}}^{w^{(1)}} g(T - (t_{2} - t_{1}), X_{T-t_{2}}^{(1)}) \\ + \int_{T-t_{2}}^{T-t_{1}} R_{s}^{w^{(1)}} \big(\partial_{t}g + \mathcal{L}g - rg + \langle \nabla g, n_{s}^{(1)} \dot{\nu}_{s}^{(1)} \rangle \big) (t_{1} + s, X_{s}^{(1)}) \mathrm{d}s \Big]$$

$$(5.30)$$

Let us plug (5.30) into (5.29), recall that $R_{T-t_2}^{w^{(1)}} = R_{T-t_2}^{w^{(2)}}$ and use (4.5) with f, g replacing f_m, g_m : $u(t_2, x) - u(t_1, x)$

$$\leq \mathsf{E}_{x} \bigg[R_{T-t_{2}}^{w^{(2)}} \left(g(T, X_{T-t_{2}}^{(1)}) - g(T - (t_{2} - t_{1}), X_{T-t_{2}}^{(1)}) \right) - \int_{T-t_{2}}^{T-t_{1}} R_{s}^{w^{(1)}} \Theta(t_{1} + s, X_{s}^{(1)}) \, \mathrm{d}s \\ + \int_{0}^{T-t_{2}} R_{s}^{w^{(2)}} \Big([\Delta_{t_{2},t_{1}}h](s, X_{s}^{(1)}) + w_{s}^{(2)} [\Delta_{t_{2},t_{1}}g](s, X_{s}^{(1)}) \Big) \, \mathrm{d}s \bigg],$$

where we recall $\Theta = \partial_t g + \mathcal{L}g - rg + h$. Thanks to condition (3.7) on h and g and (3.10) on Θ

$$u(t_2, x) - u(t_1, x) \le (K_0(1+T) + K_2)(t_2 - t_1)$$

by evaluating explicitly $\int_0^{T-t_2} w_s^{(2)} R_s^{w^{(2)}} ds$. Then, the claim holds with $K_4 = K_0(1+T) + K_2$. \Box

Our next goal is to find a uniform bound for the penalty term involving ψ_{ε} . Here we are inspired by estimates obtained under different assumptions in [40] (see also [24] and [28]).

Lemma 5.8. There is $M_5 = M_5(m)$, independent of ε and δ , such that (5.31) $\|\psi_{\varepsilon}(|\nabla u^{\varepsilon,\delta}|^2_d - f^2)\|_{C^0(\overline{\mathbb{C}}_{-1})} \leq M_5.$ *Proof.* For notational simplicity we set $u = u^{\varepsilon,\delta}$ and $\xi = \xi_m$. Let

(5.32)
$$v(t,x) \coloneqq \xi(x)\psi_{\varepsilon}\left(|\nabla u(t,x)|_d^2 - f^2(t,x)\right), \quad \text{for } (t,x) \in \overline{\mathcal{O}}_{m+1}.$$

Since v is continuous, then it attains a maximum on $\overline{\mathcal{O}}_{m+1}$. If such maximum is attained at a point $(t^*, x^*) \in \partial_P \mathcal{O}_{m+1}$ then $v(t^*, x^*) = 0$ because either $x^* \in \partial B_{m+1}$ and $\xi(x^*) = 0$ or $t^* = T$ and $|\nabla u(t^*, x^*)|_d = |\nabla g(T, x^*)|_d \leq f(T, x^*)$ by (3.9). Thus, suppose the maximum is attained in \mathcal{O}_{m+1} .

We argue similarly to the proof of Proposition 4.9. For any $\eta > 0$ there exists a neighbourhood $U_{\eta} \cup (\{0\} \times V_{\eta})$ of (t^*, x^*) such that $v(t, x) > v(t^*, x^*) - \eta$ for all $(t, x) \in U_{\eta} \cup (\{0\} \times V_{\eta})$. With no loss of generality, there is S < T and B an open ball with $\overline{B} \subset B_{m+1}$, so that $U_{\eta} \cup (\{0\} \times V_{\eta}) \subset \mathcal{O}_{S,B}$, where $\mathcal{O}_{S,B} = [0,S) \times B$. Let w^n be the solution of a PDE as in (5.14) but with $\nabla u^{\varepsilon,\delta}$ and $(\cdot)^+$ on the right-hand side of that equation replaced by smooth approximations ∇u^n and χ_n , and the function f^2 in the argument of ψ_{ε} replaced by $f^2 + \frac{1}{n}$. By arguments analogous to those in Lemma 4.7, $w^n \in C^{1,3,\alpha}_{Loc}(\mathcal{O}_{m+1})$ and $w^n \to u$ in $C^{1,2,\beta}(\overline{\mathcal{O}}_{S,B})$ and in $(C^{0,1,\beta} \cap W^{1,2,p})(\overline{\mathcal{O}}_m)$ as $n \to \infty$ for all $\beta \in (0, \alpha)$ (see Remark A.2). Define

$$v^{n}(t,x) \coloneqq \xi(x)\psi_{\varepsilon}\left(|\nabla w^{n}(t,x)|_{d}^{2} - f^{2}(t,x) - \frac{1}{n}\right), \quad \text{for } (t,x) \in \overline{\mathcal{O}}_{m+1}.$$

We have that v^n belongs to $C_{Loc}^{1,2,\alpha}(\mathcal{O}_{m+1}) \cap C^{0,1,\alpha}(\overline{\mathcal{O}}_{m+1})$, it is non-negative and it is equal to zero for $x \in \partial B_{m+1}$. Moreover $v^n \to v$ in $C^{0,1,\gamma}(\overline{\mathcal{O}}_{m+1})$ for all $\gamma \in (0,\alpha)$.

Let $(t_n^*, x_n^*)_{n \in \mathbb{N}}$ be such that $(t_n^*, x_n^*) \in \arg \max_{\overline{\mathcal{O}}_{m+1}} v^n$ and, with no loss of generality, assume $(t_n^*, x_n^*) \to (t^*, x^*)$. We also assume $|\nabla g(t^*, x^*)|_d - |\nabla u(t^*, x^*)|_d < 0$, as otherwise $f(t^*, x^*) \geq |\nabla g(t^*, x^*)|_d \geq |\nabla u(t^*, x^*)|_d$ (cf. (3.9)) implies $0 \leq v(t, x) \leq v(t^*, x^*) = 0$. By uniform convergence of ∇w^n to ∇u , we can also assume $|\nabla g|_d - |\nabla w^n|_d \leq 0$ on $U_\eta \cup (\{0\} \times V_\eta)$ for all $n \in \mathbb{N}$. We denote $\overline{\zeta_n} := (|\nabla w^n|_d^2 - f^2 - \frac{1}{n})(t_n^*, x_n^*)$ and taking derivatives of v^n we obtain

$$v_{t}^{n} = \xi \psi_{\varepsilon}'(\bar{\zeta}_{n}) \left(|\nabla w^{n}|_{d}^{2} - f^{2} \right)_{t}$$

$$v_{x_{i}}^{n} = \xi_{x_{i}} \psi_{\varepsilon}(\bar{\zeta}_{n}) + \xi \psi_{\varepsilon}'(\bar{\zeta}_{n}) \left(|\nabla w^{n}|_{d}^{2} - f^{2} \right)_{x_{i}}$$

$$(5.33) \qquad v_{x_{i}x_{j}}^{n} = \xi_{x_{i}x_{j}} \psi_{\varepsilon}(\bar{\zeta}_{n}) + \psi_{\varepsilon}'(\bar{\zeta}_{n}) \left(\xi_{x_{i}} \left(|\nabla w^{n}|_{d}^{2} - f^{2} \right)_{x_{j}} + \xi_{x_{j}} \left(|\nabla w^{n}|_{d}^{2} - f^{2} \right)_{x_{i}} \right)$$

$$+ \xi \psi_{\varepsilon}''(\bar{\zeta}_{n}) \left(|\nabla w^{n}|_{d}^{2} - f^{2} \right)_{x_{i}} \left(|\nabla w^{n}|_{d}^{2} - f^{2} \right)_{x_{j}}$$

$$+ \xi \psi_{\varepsilon}'(\bar{\zeta}_{n}) \left(2 \langle \nabla w_{x_{j}}^{n}, \nabla w_{x_{i}}^{n} \rangle + 2 \langle \nabla w^{n}, \nabla w_{x_{i}x_{j}}^{n} \rangle - \left(f^{2} \right)_{x_{i}x_{j}} \right).$$

By (4.23) we have $0 \ge (\partial_t v^n + \mathcal{L}v^n)(t_n^*, x_n^*)$. Since (t_n^*, x_n^*) is fixed, we omit it from the calculations that follow, for notational simplicity. Then using (5.33) and symmetry of a_{ij}

$$(5.34) \qquad 0 \ge (\mathcal{L}\xi)\psi_{\varepsilon}(\bar{\zeta}_{n}) - \xi\psi_{\varepsilon}'(\bar{\zeta}_{n})\left(\partial_{t}(f^{2}) + \mathcal{L}(f^{2})\right) + 2\xi\psi_{\varepsilon}'(\bar{\zeta}_{n})\left\langle\nabla w^{n}, (\partial_{t} + \mathcal{L})(\nabla w^{n})\right\rangle \\ + \frac{1}{2}\xi\psi_{\varepsilon}''(\bar{\zeta}_{n})\left\langle a\nabla(|\nabla w^{n}|_{d}^{2} - f^{2}), \nabla(|\nabla w^{n}|_{d}^{2} - f^{2})\right\rangle \\ + \psi_{\varepsilon}'(\bar{\zeta}_{n})\left\langle a\nabla\xi, \nabla(|\nabla w^{n}|_{d}^{2} - f^{2})\right\rangle + \xi\psi_{\varepsilon}'(\bar{\zeta}_{n})\sum_{i,j=1}^{d}a_{ij}\left\langle\nabla w_{x_{i}}^{n}, \nabla w_{x_{j}}^{n}\right\rangle,$$

where $(\partial_t + \mathcal{L})(\nabla w^n)$ is the vector with entries $(\partial_t + \mathcal{L})w_{x_k}^n$ for $k = 1, \ldots, d$. Recall that ψ_{ε} is non-decreasing and convex, so $\psi'_{\varepsilon}, \psi''_{\varepsilon} \ge 0$. Then, the last term on the right-hand side above is bounded from below by $\xi \psi'_{\varepsilon}(\bar{\zeta}_n) \theta |D^2 w^n|^2_{d \times d}$ as in (4.32) with θ equal to $\theta_{B_{m+1}}$. Uniform ellipticity (3.6) on B_{m+1} also gives

$$\frac{1}{2}\xi\psi_{\varepsilon}''(\bar{\zeta}_n)\big\langle a\nabla(|\nabla w^n|_d^2 - f^2), \nabla(|\nabla w^n|_d^2 - f^2)\big\rangle \ge 0.$$

Set $\bar{a}_m \coloneqq \max_{i,j} \|a_{ij}\|_{C^0(\overline{B}_{m+1})}$ and recall $|\nabla \xi|_d^2 \le C_0 \xi$ (see (4.1)). Then

$$\langle a\nabla\xi, \nabla(|\nabla w^{n}|_{d}^{2} - f^{2}) \rangle \geq -\bar{a}_{m}d^{2}|\nabla\xi|_{d} \Big(2|\nabla w^{n}|_{d}|D^{2}w^{n}|_{d\times d} + |\nabla f^{2}|_{d} \Big)$$

$$\geq -\xi \frac{\theta}{4}|D^{2}w^{n}|_{d\times d}^{2} - \frac{16}{\theta}\bar{a}_{m}^{2}d^{4}C_{0}|\nabla w^{n}|_{d}^{2} - \bar{a}_{m}d^{2}\sqrt{C_{0}\xi}|\nabla f^{2}|_{d},$$

where we used $|ab| \leq pa^2 + b^2/p$ with $p = \frac{4}{\theta}$, $b = \sqrt{\xi} |D^2 w^n|_{d \times d}$ and $a = 2\bar{a}_m d^2 \sqrt{C_0} |\nabla w^n|_d$. Since $\nabla w^n \to \nabla u$ uniformly on $\overline{\mathcal{O}}_{m+1}$, then by Proposition 5.1 we can assume $|\nabla w^n|_d \leq 1 + N_1$ and obtain

$$\left\langle a\nabla\xi, \nabla(|\nabla w^n|_d^2 - f^2) \right\rangle \ge -\xi\frac{\theta}{4}|D^2w^n|_{d\times d}^2 - C_1$$

with $C_1 = 16d^4 \bar{a}_m^2 C_0 \theta^{-1} (1+N_1)^2 + \bar{a}_m d^2 \sqrt{C_0 \xi} \|\nabla f^2\|_{C^0(\overline{\mathcal{O}}_{m+1})}$. Since $\mathcal{L}\xi$ and $(\partial_t + \mathcal{L})f^2$ are continuous on $\overline{\mathcal{O}}_{m+1}$ we have $|\mathcal{L}\xi| + |(\partial_t + \mathcal{L})f^2| \leq C_2$ on $\overline{\mathcal{O}}_{m+1}$. Similarly to the first inequality in (4.49), $\psi_{\varepsilon}(\bar{\zeta}_n) \leq \psi'_{\varepsilon}(\bar{\zeta}_n) |\nabla w^n|_d^2 \leq \psi'_{\varepsilon}(\bar{\zeta}_n) (1+N_1)^2$. Thus

$$\psi_{\varepsilon}(\bar{\zeta}_n)|\mathcal{L}\xi| \le \psi_{\varepsilon}'(\bar{\zeta}_n)(1+N_1)^2 C_2 =: \psi_{\varepsilon}'(\bar{\zeta}_n)C_3,$$

where $C_3 = C_3(m) > 0$.

We claim that, for any $\lambda > 0$ there are constants $C_4 = C_4(m) > 0$ and $\kappa_{\delta,m} > 0$ such that

(5.35)
$$\xi \langle \nabla w^n, (\partial_t + \mathcal{L})(\nabla w^n) \rangle \ge -\frac{\theta}{8} \xi |D^2 w^n|_{d \times d}^2 - \xi \lambda \theta \psi_{\varepsilon}^2(\zeta_n) - C_4(1 + \lambda^{-1}) - \kappa_{\delta,m} R_n,$$

where $\zeta_n \coloneqq (|\nabla u^n|_d^2 - f^2 - \frac{1}{n})(t_n^*, x_n^*)$ and R_n is independent of (t_n^*, x_n^*) and such that $R_n \to 0$ as $n \to \infty$. The claim is proven separately at the end of this proof, for the sake of readability. Plugging all the above estimates into (5.34) and factoring out $\psi_{\varepsilon}'(\bar{\zeta}_n) \ge 1$ gives us

$$0 \le -\frac{\theta}{2}\xi |D^2 w^n|_{d\times d}^2 + 2\lambda\theta\xi\psi_{\varepsilon}^2(\zeta_n) + C_1 + C_2 + C_3 + 2C_4(1+\lambda^{-1}) + 2\kappa_{\delta,m}R_n.$$

Letting $C_5 = C_5(m) > 0$ be a suitable constant the expression simplifies to

$$\frac{\theta}{2}\xi|D^2w^n|^2_{d\times d} \le 2\lambda\theta\xi\psi^2_\varepsilon(\zeta_n) + C_5(1+\lambda^{-1}) + 2\kappa_{\delta,m}R_n$$

We want to bound $\xi |D^2 w^n|_{d \times d}$ by $\xi \psi_{\varepsilon}(\zeta_n)$. So we multiply both sides of the inequality above by ξ , take square root and use $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$ for $a, b \geq 0$ and $|\xi| \leq 1$. That gives

(5.36)
$$\xi |D^2 w^n|_{d \times d} \le 2\sqrt{\lambda} \xi \psi_{\varepsilon}(\zeta_n) + \sqrt{2\theta^{-1}C_5(1+\lambda^{-1})} + \sqrt{4\theta^{-1}\kappa_{\delta,m}R_n}.$$

Recall that w^n solves

$$\partial_t w^n + \mathcal{L} w^n - r w^n = \psi_{\varepsilon}(\zeta_n) - h - \frac{1}{\delta} \chi_n(g-u), \quad \text{on } [0,T) \times \mathbb{R}^d$$

Multiplying by ξ we can express $\xi\psi_{\varepsilon}(\zeta_n)$ in terms of the remaining functions in the equation above. Since $(t_n^*, x_n^*) \in \mathcal{O}_{S,B}$, $w^n \to u$ in $C^{1,2,\beta}(\overline{\mathcal{O}}_{S,B})$ and $\chi_n(g-u) \to (g-u)^+$ uniformly on compacts, we can assume with no loss of generality that on $\overline{\mathcal{O}}_{S,B}$ the following hold: $\frac{1}{\delta}\chi_n(g-u) \leq (1+K_2)$ by (5.27), $|\nabla w^n|_d \leq (1+N_1)$ by (5.1), $\partial_t w^n \leq \frac{1}{2} + K_4$ by (5.28) and $rw^n \geq -\frac{1}{2}$ because $u \geq 0$. The coefficients a and b in \mathcal{L} are bounded on \overline{B}_{m+1} by a constant A_{m+1} (slightly abusing notation). Thus,

$$\xi\psi_{\varepsilon}(\zeta_{n}) = \xi\partial_{t}w^{n} - \xi rw^{n} + \xi\mathcal{L}w^{n} + \xi h + \xi\frac{1}{\delta}\chi_{n}(g-u)$$

$$\leq 1 + K_{4} + A_{m+1}(1+N_{1}) + \frac{1}{2}A_{m+1}\xi|D^{2}w^{n}|_{d\times d} + ||h||_{C^{0}(\overline{\mathcal{O}}_{m+1})} + (1+K_{2}).$$

Substituting (5.36) and grouping together the constants we obtain, for some $C_6 = C_6(m) > 0$,

$$(1 - \sqrt{\lambda}A_{m+1})\xi\psi_{\varepsilon}(\zeta_n) \leq C_6\sqrt{1 + \lambda^{-1} + \kappa_{\delta,m}R_n}$$

Then, choosing $\lambda = (4A_{m+1}^2)^{-1}$ and recalling that all expressions are evaluated at (t_n^*, x_n^*) we obtain

$$\xi(x_n^*)\psi_{\varepsilon}\left(|\nabla u^n(t_n^*, x_n^*)|_d^2 - f^2(t_n^*, x_n^*) - \frac{1}{n}\right) \le 2C_6\sqrt{1 + 4A_{m+1}^2 + \kappa_{\delta,m}R_n}$$

Taking limits as $n \to \infty$, using that $(t_n^*, x_n^*) \to (t^*, x^*)$, $R_n \to 0$ and $\nabla u^n \to \nabla u$ (uniformly on compacts), we have

$$\xi(x^*)\psi_{\varepsilon}\left(|\nabla u(t^*,x^*)|_d^2 - f^2(t^*,x^*)\right) \le 2C_6\sqrt{1 + 4A_{m+1}^2} =: M_5.$$

Recalling the definition of v in (5.32) we can conclude:

$$\left\|\psi_{\varepsilon}\left(|\nabla u|_{d}^{2}-f^{2}\right)\right\|_{C^{0}(\overline{\mathcal{O}}_{m})} \leq \sup_{(t,x)\in\overline{\mathcal{O}}_{m+1}}v(t,x) = v(t^{*},x^{*}) \leq M_{5}.$$

Proof of (5.35). Recall that $u = u^{\varepsilon,\delta}$ and that w^n solves

$$\partial_t w^n + \mathcal{L} w^n - r w^n = -h - \frac{1}{\delta} \chi_n \left(g - u^{\varepsilon, \delta} \right) + \psi_{\varepsilon} (|\nabla u^n|_d^2 - f^2 - \frac{1}{n}), \quad \text{on } [0, T) \times \mathbb{R}^d.$$

Differentiating with respect to x_k , multiplying by ξ and evaluating at (t_n^*, x_n^*) we get (5.37) $\xi(\partial_t w_{x_k}^n + \mathcal{L} w_{x_k}^n) = -\xi \mathcal{L}_{x_k} w^n + \xi r w_{x_k}^n - \xi h_{x_k} - \xi \frac{1}{\lambda} \chi'_n (g-u)(g-u)_{x_k}$

(5.37)
$$\xi(\partial_t w_{x_k}^* + \mathcal{L} w_{x_k}^n) = -\xi \mathcal{L}_{x_k} w^* + \xi r w_{x_k}^* - \xi h_{x_k} - \xi \frac{1}{\delta} \chi_n^* (g-u)(g-v) + \xi \psi_{\varepsilon}'(\zeta_n) (|\nabla u^n|_d^2 - f^2 - \frac{1}{n})_{x_k},$$

where $\zeta_n = (|\nabla u^n|_d^2 - f^2 - \frac{1}{n})(t_n^*, x_n^*)$. We subtract $v_{x_k}^n = \xi_{x_k}\psi_{\varepsilon}(\bar{\zeta}_n) + \xi\psi'_{\varepsilon}(\bar{\zeta}_n)(|\nabla w^n|_d^2 - f^2)_{x_k}$ from both sides of (5.37), and we add and subtract $\xi_{x_k}\psi_{\varepsilon}(\zeta_n)$ on the right-hand side of (5.37). Then

(5.38)
$$\xi(\partial_t w_{x_k}^n + \mathcal{L} w_{x_k}^n) - v_{x_k}^n = -\xi \mathcal{L}_{x_k} w^n + \xi r w_{x_k}^n - \xi h_{x_k} - \xi \frac{1}{\delta} \chi_n' (g-u) (g-u)_{x_k} - \xi_{x_k} \psi_{\varepsilon}(\zeta_n) + P_{n,k},$$

where

$$P_{n,k} = \xi_{x_k} \Big(\psi_{\varepsilon}(\zeta_n) - \psi_{\varepsilon}(\bar{\zeta}_n) \Big) + \xi \Big(\psi_{\varepsilon}'(\zeta_n) \big(|\nabla u^n|_d^2 - f^2 \big)_{x_k} - \psi_{\varepsilon}'(\bar{\zeta}_n) \big(|\nabla w^n|_d^2 - f^2 \big)_{x_k} \Big).$$

Recall that $(t_n^*, x_n^*) \in \mathcal{O}_{S,B}$ is a stationary point for v^n in the spatial coordinates, so $v_{x_k}^n = 0$ in (5.38) for each $1 \leq k \leq d$. Then

$$\xi \langle \nabla w^n, (\partial_t + \mathcal{L})(\nabla w^n) \rangle = \xi \left(-\sum_{k=1}^d w_{x_k}^n \mathcal{L}_{x_k} w^n + r |\nabla w^n|_d^2 - \langle \nabla w^n, \nabla h + \frac{1}{\delta} \chi'_n (g - u) \nabla (g - u) \rangle \right)$$

$$(5.39) \qquad -\psi_{\varepsilon}(\zeta_n) \langle \nabla w^n, \nabla \xi \rangle + \sum_{k=1}^d w_{x_k}^n P_{n,k}.$$

Since $(t_n^*, x_n^*) \in \mathcal{O}_{S,B}$, then denoting $\|\cdot\|_{S,B} = \|\cdot\|_{C^0(\overline{\mathcal{O}}_{S,B})}$ we have

$$\sum_{k=1}^{a} w_{x_{k}}^{n} P_{n,k} \geq - \|\psi_{\varepsilon}(\zeta_{n}) - \psi_{\varepsilon}(\bar{\zeta}_{n})\|_{S,B} \|\nabla w^{n}\|_{S,B} \|\nabla \xi\|_{S,B} - \|\psi_{\varepsilon}'(\zeta_{n}) - \psi_{\varepsilon}'(\bar{\zeta}_{n})\|_{S,B} \|\nabla w^{n}\|_{S,B} \|\nabla \left(|\nabla w^{n}|_{d}^{2} - f^{2}\right)\|_{S,B} - \|\psi_{\varepsilon}'(\zeta_{n})\|_{S,B} \|\nabla w^{n}\|_{S,B} \|\nabla (|\nabla u^{n}|_{d}^{2} - |\nabla w^{n}|_{d}^{2})\|_{S,B} =: \tilde{R}_{n},$$

where $\tilde{R}_n \to 0$ as $n \to \infty$ thanks to $C^{1,2,\beta}(\overline{\mathcal{O}}_{S,B})$ -convergence of w^n and u^n to u, for $\beta \in (0,\alpha)$. By Cauchy-Schwarz inequality, recalling that $0 \leq \chi'_n(\cdot) \leq 2$ and using arguments as in (4.42) we have

$$\begin{split} \langle \nabla w^n, \nabla h \rangle &\leq \|\nabla w^n\|_{C^0(\overline{\mathcal{O}}_{m+1})} \|\nabla h\|_{C^0(\overline{\mathcal{O}}_{m+1})} \leq (N_1+1) \|\nabla h\|_{C^0(\overline{\mathcal{O}}_{m+1})}, \\ \frac{1}{\delta} \chi'_n(g-u) \langle \nabla w^n, \nabla (g-u) \rangle &\leq \frac{2}{\delta} \|\nabla \hat{w}^n\|_{S,B} \big(\|\nabla g\|_{C^0(\overline{\mathcal{O}}_{m+1})} + \|\nabla u\|_{C^0(\overline{\mathcal{O}}_{m+1})} \big) =: \kappa_{\delta,m} \tilde{R}'_n, \end{split}$$

where $N_1 = N_1(m+1)$ is as in Proposition 5.1 and $\hat{w}^n = u - w^n$. Recall that $\|\nabla \hat{w}^n\|_{S,B} \to 0$ as $n \to \infty$, therefore $\tilde{R}'_n \to 0$ too. For the penultimate term on the right-hand side of (5.39), recalling $|\nabla \xi|_d \leq \sqrt{C_0\xi}$ (see (4.1)), we have

$$\psi_{\varepsilon}(\zeta_n) \langle \nabla w^n, \nabla \xi \rangle \leq |\nabla w^n|_d |\nabla \xi|_d \psi_{\varepsilon}(\zeta_n) \leq (N_1 + 1) \sqrt{C_0 \xi} \psi_{\varepsilon}(\zeta_n) \leq \frac{(N_1 + 1)^2 C_0}{\lambda \theta} + \xi \lambda \theta \psi_{\varepsilon}^2(\zeta_n),$$

where in the last inequality we used $ab \leq \frac{a^2}{p} + pb^2$ with $a = (N_1 + 1)\sqrt{C_0}$, $b = \sqrt{\xi}\psi_{\varepsilon}(\zeta_n)$, and $p = \lambda\theta$, with λ a constant to be chosen later and $\theta = \theta_{B_{m+1}}$ as in (3.6). For the first term on the right-hand side of (5.39) we argue as in (4.44) and obtain

$$\sum_{k=1}^{d} w_{x_k}^n \mathcal{L}_{x_k} w^n \le \frac{\theta}{8} |D^2 w^n|_{d \times d}^2 + C_1 (N_1 + 1)^2,$$

where $C_1 \coloneqq 8d^4 A_{m+1}^2 \theta^{-1} + 2dA_{m+1}$, the constant A_{m+1} is defined as in (4.45) and, differently from (4.46), we use $ab \leq pa^2 + \frac{b^2}{p}$ with $p = \frac{\theta}{8}$, $a = |D^2 w^n|_{d \times d}$ and $b = \frac{d^2}{2}A_{m+1}|\nabla w^n|_d$. Combining these bounds we get

$$\xi \langle \nabla w^n, (\partial_t + \mathcal{L})(\nabla w^n) \rangle \ge -\xi \frac{\theta}{8} |D^2 w^n|_{d \times d}^2 - \xi \lambda \theta \psi_{\varepsilon}^2(\zeta_n) - C_4(1 + \lambda^{-1}) - \kappa_{\delta,m} R_n,$$

where we define $C_4 \coloneqq C_1(N_1+1)^2 + (N_1+1)^2 C_0 \theta^{-1} + (N_1+1) \|\nabla h\|_{C^0(\overline{\mathcal{O}}_{m+1})}$ and we collect \tilde{R}_n and $\kappa_{\delta,m} \tilde{R}'_n$ in $\kappa_{\delta,m} R_n$ with an abuse of notation.

The bounds on the penalty terms in the PDE for $u^{\varepsilon,\delta}$ enable the next estimate.

Theorem 5.9. For any $p \in (1, \infty)$, there is $M_6 = M_6(m, p)$ such that

(5.40)
$$\|u^{\varepsilon,\delta}\|_{W^{1,2,p}(\mathcal{O}_m)} \le M_6, \quad \text{for all } \varepsilon, \delta \in (0,1).$$

Proof. The proof repeats the exact same arguments as in the proof of Proposition 5.2 but applied to $\varphi = \xi_{m+1} u^{\varepsilon,\delta}$ rather than to $\varphi = \xi_{m_0} u_q^{\varepsilon,\delta}$. In addition, we use Lemmas 5.6 and 5.8 to obtain the upper bound for $\|\varphi\|_{W^{1,2,p}(\mathcal{O}_{m+1})}$ as in (5.13), which is therefore independent of ε, δ .

6. The Variational Inequality

In this section, we finally prove our main result, i.e., Theorem 3.3. First we prove that Problem A admits a solution (Theorem 6.1), then we prove that such solution is the value function of our game (Theorem 6.2) and it is the maximal solution for Problem A.

Theorem 6.1. There exists a solution u of Problem A.

Proof. Let $(\varepsilon_k)_{k\in\mathbb{N}}$ be a decreasing sequence with $\varepsilon_k \to 0$. Fix $k, m \in \mathbb{N}$. Thanks to Theorem 5.9 and the compact embedding of $W^{1,2,p}(\mathcal{O}_m)$ into $C^{0,1,\beta}(\overline{\mathcal{O}}_m)$ for $\beta = 1 - (d+2)/p$, we can extract a sequence $(u^{\varepsilon_k,\delta_{k,j}^m})_{j\in\mathbb{N}}$ converging to a limit $u^{\varepsilon_k;[m]}$ (possibly depending on \mathcal{O}_m) as $j \to \infty$, in the following sense:

(6.1)
$$\begin{aligned} u^{\varepsilon_k,\delta_{k,j}^m} &\to u^{\varepsilon_k;[m]} \quad \text{and} \quad \nabla u^{\varepsilon_k,\delta_{k,j}^m} \to \nabla u^{\varepsilon_k;[m]} \quad \text{in } C^{\alpha}(\overline{\mathcal{O}}_m), \\ \partial_t u^{\varepsilon_k,\delta_{k,j}^m} &\to \partial_t u^{\varepsilon_k;[m]} \quad \text{and} \quad D^2 u^{\varepsilon_k,\delta_{k,j}^m} \to D^2 u^{\varepsilon_k;[m]} \quad \text{weakly in } L^p(\mathcal{O}_m). \end{aligned}$$

Up to selecting further subsequences (if needed), we find analogous limits on $\mathcal{O}_{m+1} \subset \mathcal{O}_{m+2} \subset \ldots$ so that $u^{\varepsilon_k;[m]} = u^{\varepsilon_k;[m+1]}$ on $\overline{\mathcal{O}}_m$, $u^{\varepsilon_k;[m+1]} = u^{\varepsilon_k;[m+2]}$ on $\overline{\mathcal{O}}_{m+1}$ and so on. Since $\overline{\mathcal{O}}_m \uparrow \mathbb{R}^{d+1}_{0,T}$ as $m \to \infty$, iterating this procedure we can define a limit function u^{ε_k} on $\mathbb{R}^{d+1}_{0,T}$.

The sequence $(u^{\varepsilon_k})_{k\in\mathbb{N}}$ satisfies the same bound as in (5.40). Therefore, by the same argument as above we can extract a further converging subsequence, which we denote still by $(u^{\varepsilon_k})_{k\in\mathbb{N}}$ with an abuse of notation. That is, there is a function u on $\mathbb{R}^{d+1}_{0,T}$ such that for any $m \in \mathbb{N}$

$$u^{\varepsilon_k} \to u \quad \text{and} \quad \nabla u^{\varepsilon_k} \to \nabla u \quad \text{in } C^{\alpha}(\overline{\mathcal{O}}_m),$$

 $\partial_t u^{\varepsilon_k} \to \partial_t u \quad \text{and} \quad D^2 u^{\varepsilon_k} \to D^2 u \quad \text{weakly in } L^p(\mathcal{O}_m).$

Finally, we can extract a diagonal subsequence $(u^{\varepsilon_i,\delta_i})_{i\in\mathbb{N}}$ that converges to u locally on $\mathbb{R}^{d+1}_{0,T}$ in the sense above as $(\varepsilon_i, \delta_i) \to 0$, simultaneously.

Next we prove that the limit function u is solution of Problem A. By construction, $u \in W^{1,2,p}_{\ell oc}(\mathbb{R}^{d+1}_{0,T})$. Thanks to (5.27), (5.31) and $C^{0,1,\alpha}$ -convergence on compacts, in the limit as $\varepsilon, \delta \to 0$ we obtain

$$g(t,x) - u(t,x) \le 0$$
 and $|\nabla u(t,x)|_d - f(t,x) \le 0$, for all $(t,x) \in \mathbb{R}^{d+1}_{0,T}$.

Fix $(\bar{t}, \bar{x}) \in \mathcal{C}$, i.e., $u(\bar{t}, \bar{x}) > g(\bar{t}, \bar{x})$. By continuity of u and g there is an open neighbourhood \mathcal{O} of (\bar{t}, \bar{x}) such that u(t, x) > g(t, x) for all $(t, x) \in \mathcal{O}$. Uniform convergence on compacts of $u^{\varepsilon_i, \delta_i}$ to u also guarantees that $u^{\varepsilon_i, \delta_i} > g$ on \mathcal{O} , for sufficiently large *i*'s. Then, for large *i*'s (5.14) reads

$$\partial_t u^{\varepsilon_i,\delta_i} + \mathcal{L} u^{\varepsilon_i,\delta_i} - r u^{\varepsilon_i,\delta_i} = -h + \psi_{\varepsilon} \left(|\nabla u^{\varepsilon_i,\delta_i}|_d^2 - f^2 \right) \ge -h, \quad \text{on } \mathcal{O}.$$

Multiplying the equation above by $\phi \in C_c^{\infty}(\mathcal{O}), \phi \geq 0$ and letting $i \to \infty$ we obtain the second equation in (3.5). Analogously, let $(\bar{t}, \bar{x}) \in \mathcal{I}$, i.e., $|\nabla u(\bar{t}, \bar{x})|_d < f(\bar{t}, \bar{x})$. Then, by continuity of ∇u and uniform convergence on compacts of $\nabla u^{\varepsilon_i, \delta_i} \to \nabla u$ we find an open neighbourhood \mathcal{O} such that $|\nabla u|_d < f$ on \mathcal{O} and $|\nabla u^{\varepsilon_i, \delta_i}|_d < f$ on \mathcal{O} for sufficiently large *i*. In such neighbourhood (5.14) reads

$$\partial_t u^{\varepsilon_i,\delta_i} + \mathcal{L} u^{\varepsilon_i,\delta_i} - r u^{\varepsilon_i,\delta_i} = -h - \frac{1}{\delta} (g - u^{\varepsilon_i,\delta_i})^+ \le -h,$$

and by the same argument as above, using test functions, we can pass to the limit and obtain the third equation in (3.5). The case in which $(\bar{t}, \bar{x}) \in \mathcal{I} \cap \mathcal{C}$ is now obvious and the first equation in (3.5) also holds for all (t, x) by standard PDE theory (e.g., as in the proof of Theorem 5.3). Finally, the terminal condition is trivially satisfied since $u^{\varepsilon,\delta}(T, x) = g(T, x)$ for all $(\varepsilon, \delta) \in (0, 1)^2$.

Notice that u has at most quadratic growth by Lemma 4.5.

We are now in the position to prove our main result.

Theorem 6.2. The game in (3.3) admits a value v which is also the maximal solution of Problem A. Moreover, τ_* defined in (3.11) is optimal for the stopper.

Proof. Fix $(t, x) \in \mathbb{R}^{d+1}_{0,T}$, let $[(n, \nu), \tau] \in \mathcal{A}_t \times \mathcal{T}_t$ and redefine ρ_m as in (4.6) but with càdlàg controls rather than continuous ones. By regularity of $u^{\varepsilon,\delta}$ (Theorem 5.3), letting $\rho_m^k = \rho_m \wedge (T - t - k^{-1})^+$ Itô's formula applies to $e^{-r(\tau \wedge \rho_m^k)} u^{\varepsilon,\delta}(t + \tau \wedge \rho_m^k, X^{[n,\nu]}_{\tau \wedge \rho_m^k})$. Using that $u^{\varepsilon,\delta}$ solves Problem C, taking expectations and letting $k \uparrow \infty$ we obtain

$$(6.2) \qquad u^{\varepsilon,\delta}(t,x) = \mathsf{E}_{x} \left[e^{-r(\tau \wedge \rho_{m})} u^{\varepsilon,\delta} (t + \tau \wedge \rho_{m}, X_{\tau \wedge \rho_{m}}^{[n,\nu]}) \right. \\ \left. + \int_{0}^{\tau \wedge \rho_{m}} e^{-rs} \left[h + \frac{1}{\delta} \left(g - u^{\varepsilon,\delta} \right)^{+} - \psi_{\varepsilon} \left(|\nabla u^{\varepsilon,\delta}|_{d}^{2} - f^{2} \right) \right] \left(t + s, X_{s-}^{[n,\nu]} \right) \mathrm{d}s \right. \\ \left. - \int_{0}^{\tau \wedge \rho_{m}} e^{-rs} \langle \nabla u^{\varepsilon,\delta} (t + s, X_{s-}^{[n,\nu]}), n_{s} \rangle \, \mathrm{d}\nu_{s}^{c} \right. \\ \left. - \sum_{0 \le s \le \tau \wedge \rho_{m}} e^{-rs} \int_{0}^{\Delta \nu_{s}} \langle \nabla u^{\varepsilon,\delta} (t + s, X_{s-}^{[n,\nu]} + \lambda n_{s}), n_{s} \rangle \, \mathrm{d}\lambda \right].$$

We want to take limits as $\varepsilon, \delta \to 0$ and pass the limits under expectations. To do that we notice that $X_{s-}^{[n,\nu]} \in \overline{B}_m$ for all $s \in [0, \rho_m]$, P_x -a.s. Then the terms under the integral with respect to 'ds' are bounded thanks to Assumption 3.2, Lemma 5.6 and Lemma 5.8. Since $\nabla u^{\varepsilon,\delta}$ is also bounded by $N_1(m)$ (Proposition 5.1 with m_0 therein replaced by m), the integrals with respect to the control are bounded by $N_1(m)\nu_{T-t}$, which is square integrable by definition of \mathcal{A}_t . This reasoning does not apply to the final jump in the sum, because that jump could bring the process $X^{[n,\nu]}$ outside the ball B_m if $\rho_m \leq \tau$. It is therefore convenient to isolate that term from the sum and rewrite

(6.3)
$$u^{\varepsilon,\delta}(t+\tau\wedge\rho_m, X^{[n,\nu]}_{\tau\wedge\rho_m}) - \int_0^{\Delta\nu_{\tau\wedge\rho_m}} \langle \nabla u^{\varepsilon,\delta}(t+\tau\wedge\rho_m, X^{[n,\nu]}_{\tau\wedge\rho_m-} + \lambda n_{\tau\wedge\rho_m}), n_{\tau\wedge\rho_m} \rangle \, \mathrm{d}\lambda$$
$$= u^{\varepsilon,\delta}(t+\tau\wedge\rho_m, X^{[n,\nu]}_{\tau\wedge\rho_m-}).$$

Finally, we recall that $u^{\varepsilon,\delta}$ has quadratic growth by Lemma 4.5 and notice that

(6.4)
$$\mathsf{E}_{x}\left[\sup_{0\leq s\leq T-t}\left|X_{s}^{[n,\nu]}\right|_{d}^{2}\right]\leq c\left(1+|x|_{d}^{2}+\mathsf{E}_{x}[|\nu_{T-t}|^{2}]\right),$$

by standard estimates for SDEs [29, Thm. 2.5.10], with c > 0 independent of m, ε , δ . Then we are allowed to use dominated convergence and it remains to evaluate the limit.

For P_x -a.e. $\omega \in \Omega$ there is a compact $\mathcal{K}_\omega \subset \mathbb{R}^d$ such that $X_s^{[n,\nu]}(\omega) \in \mathcal{K}_\omega$ for all $s \in [0, \rho_m(\omega)]$, by right-continuity of the process and the fact that ν is square integrable. Then, uniform convergence of $(u^{\varepsilon,\delta}, \nabla u^{\varepsilon,\delta})$ to $(u, \nabla u)$ on compacts implies

$$\lim_{\varepsilon,\delta\to 0} \langle \nabla u^{\varepsilon,\delta}(t+s, X_{s-}^{[n,\nu]}), n_s \rangle(\omega) = \langle \nabla u(t+s, X_{s-}^{[n,\nu]}), n_s \rangle(\omega),$$
$$\lim_{\varepsilon,\delta\to 0} \langle \nabla u^{\varepsilon,\delta}(t+s, X_{s-}^{[n,\nu]} + \lambda n_s), n_s \rangle(\omega) = \langle \nabla u(t+s, X_{s-}^{[n,\nu]} + \lambda n_s), n_s \rangle(\omega).$$

for all $s \in [0, \rho_m(\omega)]$ and all $\lambda \in [0, \Delta \nu_s(\omega)]$, for P_x -a.e. $\omega \in \Omega$.

Let us now choose $\tau = \tau_* = \inf\{s \ge 0 \mid u(t+s, X_s^{[n,\nu]}) \le g(t+s, X_s^{[n,\nu]})\}$ (notice that $\tau_* = \tau_*(n,\nu)$). Fix $\omega \in \Omega$ outside of a P_x -null set and recall that $s \mapsto \nu_s(\omega)$ has at most countably many jumps on any bounded interval. Then, for a.e. $0 \le s < \tau_*(\omega) \land \rho_m(\omega)$ there is $q_{s,\omega} > 0$ such that

$$(u-g)(t+s, X_{s-}^{[n,\nu]}(\omega)) = (u-g)(t+s, X_s^{[n,\nu]}(\omega)) \ge q_{s,\omega}$$

Therefore, the pointwise convergence of $u^{\varepsilon,\delta}$ to u implies that for a.e. $0 \le s < \tau_*(\omega) \land \rho_m(\omega)$

$$\limsup_{\varepsilon,\delta\to 0} \frac{1}{\delta} \left(g - u^{\varepsilon,\delta}\right)^+ \left(t + s, X_{s-}^{[n,\nu]}(\omega)\right) = 0$$

Using the observations above, combined with (6.3), dominated convergence and $\psi_{\varepsilon} \geq 0$ we obtain

$$\begin{split} u(t,x) \leq \mathsf{E}_x \bigg[e^{-r(\tau_* \wedge \rho_m)} u(t+\tau_* \wedge \rho_m, X_{\tau_* \wedge \rho_m}^{[n,\nu]}) + \int_0^{\tau_* \wedge \rho_m} e^{-rs} h(t+s, X_s^{[n,\nu]}) \,\mathrm{d}s \\ &- \int_0^{\tau_* \wedge \rho_m} e^{-rs} \langle \nabla u(t+s, X_{s-}^{[n,\nu]}), n_s \rangle \,\mathrm{d}\nu_s^c - \sum_{0 \leq s \leq \tau_* \wedge \rho_m} e^{-rs} \int_0^{\Delta \nu_s} \langle \nabla u(t+s, X_{s-}^{[n,\nu]} + \lambda n_s), n_s \rangle \,\mathrm{d}\lambda \bigg]. \end{split}$$

By $|\nabla u|_d \leq f$ and the definition of $\tau_* \wedge \rho_m$ we have

$$\begin{split} u(t,x) &\leq \mathsf{E}_x \left[e^{-r\tau_*} g(t+\tau_*, X^{[n,\nu]}_{\tau_*}) \mathbb{1}_{\{\tau_* \leq \rho_m\}} + e^{-r\rho_m} u(t+\rho_m, X^{[n,\nu]}_{\rho_m}) \mathbb{1}_{\{\tau_* > \rho_m\}} \right] \\ &+ \mathsf{E}_x \bigg[\int_0^{\tau_* \wedge \rho_m} e^{-rs} h(t+s, X^{[n,\nu]}_s) \,\mathrm{d}s + \int_{[0,\tau_* \wedge \rho_m]} e^{-rs} f(t+s, X^{[n,\nu]}_s) \circ \,\mathrm{d}\nu_s \bigg]. \end{split}$$

Now we let $m \to \infty$. Clearly $\rho_m \uparrow T - t$, P_x -a.s. by (6.4). Since the bound in (6.4) is independent of m and functions u and g have at most quadratic growth we can apply the dominated convergence theorem to pass the limit inside the first expectation. We can also take the limit inside the second expectation by monotone convergence as all terms under the integral are non-negative. For P_x -a.e. $\omega \in \Omega$ we have $X_s^{[n,\nu]}(\omega) \in \mathcal{K}_\omega$ for all $s \in [0, T - t]$. Then

$$\lim_{m \to \infty} e^{-r\rho_m(\omega)} u \big(t + \rho_m(\omega), X_{\rho_m}^{[n,\nu]}(\omega) \big) \mathbb{1}_{\{\tau_* > \rho_m\}}(\omega) = e^{-r(T-t)} u \big(T, X_{T-t}^{[n,\nu]}(\omega) \big) \mathbb{1}_{\{\tau_* \ge T-t\}}(\omega) = e^{-r(T-t)} g \big(T, X_{T-t}^{[n,\nu]}(\omega) \big) \mathbb{1}_{\{\tau_* = T-t\}}(\omega), \quad \mathsf{P}_x\text{-a.e. } \omega \in \Omega,$$

because $\tau_*(\omega) \leq T - t$ and u is uniformly continuous on $[0,T] \times \mathcal{K}_{\omega}$. Hence, for $m \to \infty$ we obtain (6.5) $u(t,x) \leq \mathcal{J}_{t,x}(n,\nu,\tau_*).$

By arbitrariness of $(n, \nu) \in \mathcal{A}_t$ and sub-optimality of τ_* we have $u(t, x) \leq \underline{v}(t, x)$ by definition of lower value.

Next we prove $u \geq \overline{v}$. Since $\frac{1}{\delta}(g - u^{\varepsilon,\delta})^+ \geq 0$ that term can de dropped from (6.2) to obtain a lower bound for $u^{\varepsilon,\delta}$. We let $\delta \to 0$ in (6.2) along the sequence constructed in (6.1) while keeping ε fixed. As above, dominated convergence applies, and thanks to $u^{\varepsilon} \geq g$ (Lemma 5.6) we obtain

$$u^{\varepsilon}(t,x) \geq \mathsf{E}_{x} \left[e^{-r(\tau \wedge \rho_{m})} g(t+\tau \wedge \rho_{m}, X_{\tau \wedge \rho_{m}}^{[n,\nu]}) + \int_{0}^{\tau \wedge \rho_{m}} e^{-rs} \left[h - \psi_{\varepsilon} \left(|\nabla u^{\varepsilon}|_{d}^{2} - f^{2} \right) \right] \left(t+s, X_{s-}^{[n,\nu]} \right) \mathrm{d}s \right]$$

$$(6.6) \qquad -\int_{0}^{\tau \wedge \rho_{m}} e^{-rs} \langle \nabla u^{\varepsilon}(t+s, X_{s-}^{[n,\nu]}), n_{s} \rangle \mathrm{d}\nu_{s}^{c} - \sum_{0 \leq s \leq \tau \wedge \rho_{m}} e^{-rs} \int_{0}^{\Delta \nu_{s}} \langle \nabla u^{\varepsilon}(t+s, X_{s-}^{[n,\nu]} + n_{s}\lambda), n_{s} \rangle \mathrm{d}\lambda \right].$$

We can now choose a control pair $(n, \nu) = (n^{\varepsilon}, \nu^{\varepsilon})$ defined as in (5.23) but with $u^{\varepsilon,\delta}$ therein replaced by u^{ε} . Although $\nabla u^{\varepsilon}(t, \cdot)$ is not Lipschitz, it can be shown by standard localisation procedure and the use of [38, Thm. 1] that the associated controlled SDE admits a unique, non-exploding, strong solution $X^{\varepsilon} = X^{[n^{\varepsilon},\nu^{\varepsilon}]}$ on [0, T-t] (the proof is given in Appendix A.4 for completeness).

By construction, the pair $(n^{\varepsilon}, \nu^{\varepsilon})$ satisfies

(6.7)
$$-\left\langle n_s^{\varepsilon} \dot{\nu}_s^{\varepsilon}, \nabla u^{\varepsilon}(t+s, X_s^{\varepsilon}) \right\rangle - \psi_{\varepsilon} \left(|\nabla u^{\varepsilon}|_d^2 - f^2 \right) \left(t+s, X_s^{\varepsilon} \right) = H^{\varepsilon} \left(t+s, X_s^{\varepsilon}, n_s^{\varepsilon} \dot{\nu}_s^{\varepsilon} \right)$$

with H^{ε} as in (5.16). Then, from (6.6) we obtain

$$u^{\varepsilon}(t,x) \geq \mathsf{E}_{x}\bigg[e^{-r(\tau \wedge \rho_{m})}g(t+\tau \wedge \rho_{m}, X^{\varepsilon}_{\tau \wedge \rho_{m}}) + \int_{0}^{\tau \wedge \rho_{m}} e^{-rs}\big(h + H^{\varepsilon}(\cdot, n^{\varepsilon}_{s}\dot{\nu}^{\varepsilon}_{s})\big)(t+s, X^{\varepsilon}_{s})\mathrm{d}s\bigg].$$

Taking $p = f(t+s, X_s^{\varepsilon})n_s^{\varepsilon}$ in the Hamiltonian, letting $m \to \infty$ and using Fatou's lemma, we obtain

$$u^{\varepsilon}(t,x) \geq \mathsf{E}_x \bigg[e^{-r\tau} g\big(t+\tau, X^{\varepsilon}_{\tau}\big) + \int_0^{\tau} e^{-rs} h(t+s, X^{\varepsilon}_s) \mathrm{d}s + \int_{[0,\tau]} e^{-rs} f(t+s, X^{\varepsilon}_s) \circ \mathrm{d}\nu^{\varepsilon}_s \bigg],$$

where we notice that ν^{ε} is absolutely continuous so that the final integral is obvious.

By arbitrariness of τ we can take supremum over all stopping times. Since $(n^{\varepsilon}, \nu^{\varepsilon}) \in \mathcal{A}_t$, we can also take infimum over all admissible controls and continue with the same direction of inequalities. That is, $u^{\varepsilon}(t, x) \geq \overline{v}(t, x)$. Finally, letting $\varepsilon \to 0$ we obtain $u(t, x) \geq \overline{v}(t, x)$, as needed. Since $u \leq \underline{v}$ was proven above, we conclude $u = v = \overline{v} = \underline{v}$.

Now that we know that u = v is the value of the game, (6.5) also yields optimality of τ_* . Notice that, in particular, $\tau_* = \tau_*(n, \nu)$ is best response against any admissible pair (n, ν) .

It remains to prove that v is indeed the maximal solution of Problem A. Let w be another solution of Problem A. By standard mollification procedure we can construct a sequence $(w_k)_{k\in\mathbb{N}} \subset C_c^{\infty}(\mathbb{R}^{d+1}_{0,T})$ such that $w_k \to w$ and $\nabla w_k \to \nabla w$ uniformly on compact sets, with $\partial_t w_k \to \partial_t w$ and $D^2 w_k \to D^2 w$ strongly in $L^p_{\elloc}(\mathbb{R}^{d+1}_{0,T})$ for all $p \in [1, \infty)$, as $k \to \infty$. For $\eta > 0$, set

$$\mathcal{C}_w^{\eta} = \{(t, x) \in \mathbb{R}^{d+1}_{0, T} : w(t, x) > g(t, x) + \eta\}.$$

For fixed $\eta > 0$ and $m \in \mathbb{N}$, by the same argument as in the proof of [17, Thm. 4.1, Ch. VIII]

(6.8)
$$\liminf_{k \to \infty} \inf_{(t,x) \in \mathcal{O}_m \cap \mathcal{C}_w^{\eta}} (\partial_t w_k + \mathcal{L} w_k - r w_k + h)(t,x) \ge 0.$$

Pick an arbitrary pair $(n, \nu) \in \mathcal{A}_t$ and denote

$$\zeta_{\eta} = \inf\{s \ge 0 | w(t+s, X_s^{[n,\nu]}) \le g(t+s, X_s^{[n,\nu]}) + \eta\} \land (T-t)$$

By an application of Itô's formula to $e^{-rs}w_k(t+s, X_s^{[n,\nu]})$ we obtain

$$\begin{split} w_k(t,x) = \mathsf{E}_x \bigg[e^{-r(\zeta_\eta \wedge \rho_m)} w_k(t + \zeta_\eta \wedge \rho_m, X_{\zeta_\eta \wedge \rho_m}^{[n,\nu]}) - \int_0^{\zeta_\eta \wedge \rho_m} e^{-rs} (\partial_t w_k + \mathcal{L}w_k - rw_k)(t + s, X_s^{[n,\nu]}) \mathrm{d}s \\ - \int_0^{\zeta_\eta \wedge \rho_m} e^{-rs} \langle \nabla w_k(t + s, X_{s-}^{[n,\nu]}), n_s \rangle \, \mathrm{d}\nu_s^c - \sum_{s \leq \zeta_\eta \wedge \rho_m} e^{-rs} \int_0^{\Delta \nu_s} \langle \nabla w_k(t + s, X_{s-}^{[n,\nu]} + \lambda n_s), n_s \rangle \, \mathrm{d}\lambda \bigg] \end{split}$$

Letting $k \to \infty$, dominated convergence (up to possibly selecting a subsequence) and reverse Fatou's lemma (justified by (6.8)) allow us to pass the limit under expectation. Then, exploiting the uniform convergence of $(w_k, \nabla w_k)$ to $(w, \nabla w)$ on $\overline{\mathcal{O}_m \cap \mathcal{C}_w^{\eta}}$, (6.8), the definition of $\zeta_{\eta} \wedge \rho_m$ and the fact that $|\nabla w|_d \leq f$ we have

$$\begin{split} w(t,x) &\leq \eta + \mathsf{E}_x \bigg[e^{-r\zeta_{\eta}} g(t+\zeta_{\eta}, X_{\zeta_{\eta}}^{[n,\nu]}) \mathbb{1}_{\{\zeta_{\eta} \leq \rho_m\}} + e^{-r\rho_m} w(t+\rho_m, X_{\rho_m}^{[n,\nu]}) \mathbb{1}_{\{\zeta_{\eta} > \rho_m\}} \\ &+ \int_0^{\zeta_{\eta} \wedge \rho_m} e^{-rs} h(t+s, X_s^{[n,\nu]}) \mathrm{d}s + \int_0^{\zeta_{\eta} \wedge \rho_m} e^{-rs} f(t+s, X_s^{[n,\nu]}) \circ \mathrm{d}\nu_s \bigg]. \end{split}$$

Finally, letting $m \to \infty$, the same arguments that lead to (6.5) give $w(t,x) \leq \eta + \mathcal{J}_{t,x}(n,\nu,\zeta_{\eta})$. Hence, $w(t,x) \leq \eta + v(t,x)$ and letting $\eta \to 0$ we conclude.

Remark 6.3. It is worth noticing that the proof above can be repeated verbatim if we replace \mathcal{A}_t with \mathcal{A}_t° everywhere. Thus we conclude that

$$v(t,x) = \inf_{(n,\nu)\in\mathcal{A}_t^\circ} \sup_{\tau\in\mathcal{T}_t} \mathcal{J}_{t,x}(n,\nu,\tau) = \sup_{\tau\in\mathcal{T}_t} \inf_{(n,\nu)\in\mathcal{A}_t^\circ} \mathcal{J}_{t,x}(n,\nu,\tau).$$

That is, the game with absolutely continuous controls admits the same value as the game with singular controls. It is however expected, but it will not be proven here, that an optimal control cannot be found in \mathcal{A}°_{t} whereas it should be possible to find one in \mathcal{A}_{t} in some cases.

We make a simple final observation concerning the nature of the optimal control in our game.

Lemma 6.4. Let $(t, x) \in \mathbb{R}^{d+1}_{0,T}$. For any $\tau \in \mathcal{T}_t$ we have

$$\inf_{(n,\nu)\in\mathcal{A}_t}\mathcal{J}_{t,x}(n,\nu,\tau) = \inf_{(n,\nu)\in\mathcal{A}_t^\tau}\mathcal{J}_{t,x}(n,\nu,\tau),$$

where $\mathcal{A}_{t}^{\tau} := \{ (n, \nu) \in \mathcal{A}_{t} \mid \nu_{\tau} = \nu_{\tau-}, \mathsf{P}_{x} - a.s. \}.$

Proof. In the expression of $\mathcal{J}_{t,x}(n,\nu,\tau)$, for any triple $[(n,\nu),\tau] \in \mathcal{A}_t \times \mathcal{T}_t$ we have

$$e^{-r\tau}g(t+\tau, X_{\tau}^{[n,\nu]}) + \int_{[0,\tau]} e^{-rs}f(t+s, X_{s}^{[n,\nu]}) \circ d\nu_{s}$$

$$= e^{-r\tau}g(t+\tau, X_{\tau-}^{[n,\nu]}) + \int_{[0,\tau)} e^{-rs}f(t+s, X_{s}^{[n,\nu]}) \circ d\nu_{s}$$

$$+ e^{-r\tau}\int_{0}^{\Delta\nu_{\tau}} (\langle \nabla g, n_{\tau} \rangle + f)(t+\tau, X_{\tau-}^{[n,\nu]} + \lambda n_{\tau}) d\lambda$$

$$\geq e^{-r\tau}g(t+\tau, X_{\tau-}^{[n,\nu]}) + \int_{[0,\tau)} e^{-rs}f(t+s, X_{s}^{[n,\nu]}) \circ d\nu_{s},$$

where the final inequality is due to (3.9). Therefore, the controller attains a lower payoff by avoiding a jump of the control at time τ . That concludes the proof.

(

7. Short Remark about Methodology

The methodology in our paper combines analytical and probabilistic estimates. The former are predominant in particular in Proposition 4.9, Proposition 5.1, Lemma 5.8, which yield bounds on the spatial gradient of the solution of the penalised problem (first on bounded domains and then on unbounded domains), uniformly with respect to the penalisation parameters and the cut-off functions. Those analytical estimates are rather convoluted and do not necessarily offer a clear intuition for generalisations to different setups.

Probabilistic techniques instead are used in Lemma 4.5 and Lemma 4.8 to obtain, respectively, growth estimates on the solution of the penalised problem and bounds on its gradient on $\partial_P \mathcal{O}_m$. In Lemma 5.6 and Lemma 5.7 quick probabilistic arguments applied to the solution $u^{\varepsilon,\delta}$ of the penalised problem on unbounded domain yield a uniform bound on $\delta^{-1}(g - u^{\varepsilon,\delta})^+$ and an upper bound for the time derivative $\partial_t u^{\varepsilon,\delta}$. Finally, Theorem 6.2 is proven using probabilistic representations of the functions $u^{\varepsilon,\delta}$ and $u^{\varepsilon} = \lim_{\delta \to 0} u^{\varepsilon,\delta}$ combined with limiting arguments. That approach is needed because an application of Itô's formula (or generalisations thereof) to $u(t, X_t^{[n,\nu]})$, for generic (n,ν) , is not possible due to the lack of sufficient regularity of the maximal solution u of Problem C.

It seems to us that proofs obtained via probabilistic methods are generally easy to interpret. It would be interesting to obtain fully probabilitic arguments also for the bounds on the spatial gradient, which we currently obtain via analytical methods. This, however, appears a very challenging task and we leave it for future work.

APPENDIX A.

A.1. Cut-off functions. Here we give a construction of the functions in (4.1). Let $\xi : \mathbb{R} \to [0,1]$ be defined as

$$\xi(z) \coloneqq \begin{cases} 1, & z \le 0, \\ 0, & z \ge 1, \\ \exp\left(\frac{1}{z-1}\right) / \left[\exp\left(\frac{1}{z-1}\right) + \exp\left(-\frac{1}{z}\right) \right], & 0 < z < 1, \end{cases}$$

and set $\xi_m(x) = \xi(|x|_d - m)$ for $x \in \mathbb{R}^d$. Then $\xi_m \in C_c^{\infty}(\mathbb{R}^d)$, $0 \le \xi_m \le 1$, $\xi_m = 1$ on \overline{B}_m and $\xi_m = 0$ on $\mathbb{R}^d \setminus B_{m+1}$. It is clear that $\nabla \xi_m = \mathbf{0}$ on B_m and $\mathbb{R}^d \setminus B_{m+1}$. It can also be checked that for $x \in B_{m+1} \setminus B_m$

$$\partial_{x_k}\xi_m(x) = \frac{x_k}{|x|_d}\xi'(|x|_d - m), \quad \text{for } k = 1, \dots d,$$

and therefore $|\nabla \xi_m(x)|_d^2 = [\xi'(|x|_d - m)]^2$. Since $\xi'(z) = -\xi(z)(1 - \xi(z))(\frac{1}{(z-1)^2} + \frac{1}{z^2})$, then $|\nabla \xi_m(x)|_d^2 \le C_0 \xi_m(x)$ for all $x \in \mathbb{R}^d$, for a suitable $C_0 > 0$ independent of m.

A.2. **Proof of Lemma 4.7.** For existence and uniqueness of the solution to (4.19) we invoke [19, Thm. 3.3.7]. Indeed the smoothing of $u_m^{\varepsilon,\delta}$ and $(\cdot)^+$ guarantees that

(A.1)
$$h_m + \frac{1}{\delta}\chi_n(g_m - u_m^{\varepsilon,\delta}) - \psi_\varepsilon \left(|\nabla u^n|_d^2 - f_m^2 - \frac{1}{n}\right) \in C^{0,1,\alpha}(\overline{\mathcal{O}}_m)$$

Moreover, the compatibility condition

$$\lim_{s\uparrow T} (\partial_t g_m + \mathcal{L}g_m - rg_m)(s, x) = \left[-h_m - \frac{1}{\delta}\chi_n(g_m - u_m^{\varepsilon,\delta}) + \psi_\varepsilon \left(|\nabla u^n|_d^2 - f_m^2 - \frac{1}{n} \right) \right](T, x), \text{ for } x \in \partial B_m,$$

holds with both sides of the equation equal to zero. Indeed, given that $\xi_{m-1} \in C_c^{\infty}(B_m)$ we have $g_m = \partial_{x_i x_j} g_m = 0$ on $[0, T] \times \partial B_m$. Moreover, $g_m = 0$ on $[0, T] \times \partial B_m$ also implies $\partial_t g_m = 0$ on $[0, T] \times \partial B_m$. So the left-hand side of the equation is equal to zero. On the right-hand side, for $x \in \partial B_m$ we have $h_m(T, x) = g_m(T, x) = u_m^{\varepsilon, \delta}(T, x) = 0$ and

$$|\nabla u^n(T,x)|_d^2 \le |\nabla u_m^{\varepsilon,\delta}(T,x)|_d^2 + \frac{1}{n} = |\nabla g_m(T,x)|_d^2 + \frac{1}{n} \le f_m^2(T,x) + \frac{1}{n},$$

by uniform convergence of ∇u^n to $\nabla u_m^{\varepsilon,\delta}$ and (4.2). The compatibility condition follows upon recalling $\chi_n(0) = 0$ and $\psi_{\varepsilon}(z) = 0$ for $z \leq 0$.

The fact that $w^n \in C^{1,3,\alpha}_{Loc}(\mathcal{O}_m)$ is also consequence of (A.1) and standard interior estimates for PDEs [19, Thm. 3.5.11+Cor. 3.5.1]. Instead, the convergence result $w^n \to u_m^{\varepsilon,\delta}$ in $C^{1,2,\beta}(\overline{\mathcal{O}}_m)$, as $n \to \infty$, for $\beta \in (0, \alpha)$, is a special case of Lemma A.1.

Lemma A.1. Let $F : \mathbb{R} \times \mathbb{R}^d \to \mathbb{R}$ be a Lipschitz continuous function. Fix $\phi, \varphi \in C^{0,1,\alpha}(\overline{\mathcal{O}}_m)$ and let u be a solution in $C^{1,2,\alpha}(\overline{\mathcal{O}}_m)$ of

(A.2)
$$\begin{cases} \partial_t u + \mathcal{L}u - ru = -h_m + F(\phi, \nabla \varphi), & \text{on } \mathcal{O}_m, \\ u(t, x) = g_m(t, x), & (t, x) \in \partial_P \mathcal{O}_m \end{cases}$$

Let $(\phi_n)_{n\in\mathbb{N}}, (\varphi_n)_{n\in\mathbb{N}} \subseteq C^{0,1,\alpha}(\overline{\mathcal{O}}_m)$ be such that $\phi_n \to \phi$ and $\varphi_n \to \varphi$ in $C^{0,1,\gamma}(\overline{\mathcal{O}}_m)$ as $n \to \infty$ for all $\gamma \in (0, \alpha)$. Let $(F_n)_{n\in\mathbb{N}}$ be equi-Lipschitz continuous functions $F_n : \mathbb{R} \times \mathbb{R}^d \to \mathbb{R}$ such that $F_n \to F$ in $C^0_{loc}(\mathbb{R}^{1+d})$. Finally, denote by u_n a solution to (A.2) in $C^{1,2,\alpha}(\overline{\mathcal{O}}_m)$ with $F_n(\phi_n, \nabla \varphi_n)$ instead of $F(\phi, \nabla \varphi)$.

Then, up to possibly selecting a subsequence,

(A.3)
$$\lim_{n \to \infty} \|u_n - u\|_{C^{1,2,\gamma}(\overline{\mathcal{O}}_m)} = 0, \quad \text{for all } \gamma \in (0,\alpha).$$

If
$$(\varphi_n)_{n\in\mathbb{N}}\subseteq C^{0,2,\alpha}(\overline{\mathcal{O}}_m)$$
 and $(F_n)_{n\in\mathbb{N}}\subseteq C^{1,\alpha}(\mathbb{R}^{1+d})$, then $(u_n)_{n\in\mathbb{N}}\subseteq C^{1,3,\alpha}_{Loc}(\mathcal{O}_m)\cap C^{1,2,\alpha}(\overline{\mathcal{O}}_m)$.

Proof. Define $\hat{u}_n \coloneqq u - u_n$. Then \hat{u}_n solves

$$\begin{cases} \partial_t \hat{u}_n + \mathcal{L}\hat{u}_n - r\hat{u}_n = F(\phi, \nabla \varphi) - F_n(\phi_n, \nabla \varphi_n), & \text{on } \mathcal{O}_m, \\ \hat{u}_n(t, x) = 0, & (t, x) \in \partial_P \mathcal{O}_m. \end{cases}$$

By [19, Thm. 3.2.6] we have the estimate

(A.4)
$$\|\hat{u}_n\|_{C^{1,2,\gamma}(\overline{\mathcal{O}}_m)} \le K \|F(\phi,\nabla\varphi) - F_n(\phi_n,\nabla\varphi_n)\|_{C^{0,0,\gamma}(\overline{\mathcal{O}}_m)},$$

for a constant K > 0 independent of n. Notice that by equi-Lipschitz continuity, the sequence $(F_n)_{n \in \mathbb{N}}$ is compact in any $C^{\beta}(\overline{U})$ for $\beta \in (0,1)$ and bounded set $U \subset \mathbb{R}^{d+1}$. Thanks to the convergence of the functions ϕ_n, φ_n and F_n we have that, up to possibly selecting a subsequence, $F_n(\phi_n, \nabla \varphi_n) \to F(\phi, \nabla \varphi)$ in $C^{\gamma}(\overline{\mathcal{O}}_m)$ as $n \to \infty$ for all $\gamma \in (0, \alpha)$. Thus, (A.3) holds.

If we also assume that $(\varphi_n)_{n\in\mathbb{N}} \subseteq C^{0,2,\alpha}(\overline{\mathcal{O}}_m)$ and $(F_n)_{n\in\mathbb{N}} \subseteq C^{1,\alpha}(\mathbb{R}^{1+d})$, it turns out that $F_n(\phi_n,\varphi_n) \in C^{0,1,\alpha}(\overline{\mathcal{O}}_m)$ and since the coefficients of \mathcal{L} are continuously differentiable then $u_n \in C^{1,3,\alpha}_{Loc}(\mathcal{O}_m)$ for all n by [19, Thm. 3.5.11+Cor. 3.5.1].

Remark A.2. Thanks to [6, Thm. 2.6.5 and Rem. 2.6.4] the bound (A.4) can be replaced by

$$\|\hat{u}_n\|_{W^{1,2,p}(\mathcal{O}_m)} \le K \|F(\phi, \nabla\varphi) - F_n(\phi_n, \nabla\varphi_n)\|_{L^p(\mathcal{O}_m)}, \quad p \in (1,\infty)$$

Hence, stability of solutions of (A.2) also holds in $W^{1,2,p}(\mathcal{O}_m)$, i.e., $\lim_n u_n = u$ in $W^{1,2,p}(\mathcal{O}_m)$.

A.3. Convergence of the sequence $(t_{n_k}^{\lambda}, x_{n_k}^{\lambda})_{k \in \mathbb{N}}$ in Proposition 4.9. Here we prove that $(\tilde{t}, \tilde{x}) \in \arg \max_{\overline{\mathcal{O}}_m} v^{\lambda}$. Arguing by contradiction let us assume $(\tilde{t}, \tilde{x}) \notin \arg \max_{\overline{\mathcal{O}}_m} v^{\lambda}$. Then there exists a $\epsilon > 0$ such that $v^{\lambda}(\tilde{t}, \tilde{x}) \leq \max_{\overline{\mathcal{O}}_m} v^{\lambda} - \epsilon$ and so there exists a neighbourhood U_{ϵ} of (\tilde{t}, \tilde{x}) such that $v^{\lambda}(t, x) \leq \max_{\overline{\mathcal{O}}_m} v^{\lambda} - \frac{\epsilon}{2}$ for all $(t, x) \in \overline{U}_{\epsilon}$. For all sufficiently large k's we also have $(t_{n_k}^{\lambda}, x_{n_k}^{\lambda}) \in U_{\epsilon}$ and by uniform convergence

(A.5)
$$|v^{\lambda,n_k} - v^{\lambda}|(t,x) \le \frac{\epsilon}{4}, \quad \text{for } (t,x) \in \overline{\mathcal{O}}_m$$

Hence

(A.6)
$$\max_{(t,x)\in\overline{\mathcal{O}}_m} v^{\lambda,n_k}(t,x) = v^{\lambda,n_k}(t_{n_k}^{\lambda},x_{n_k}^{\lambda}) \le v^{\lambda}(t_{n_k}^{\lambda},x_{n_k}^{\lambda}) + \frac{\epsilon}{4} \le \max_{(t,x)\in\overline{\mathcal{O}}_m} v^{\lambda}(t,x) - \frac{\epsilon}{4},$$

where the first equality is by definition of $(t_{n_k}^{\lambda}, x_{n_k}^{\lambda})$, the first inequality by (A.5) and the final inequality follows by $(t_{n_k}^{\lambda}, x_{n_k}^{\lambda}) \in \overline{U}_{\epsilon}$.

With no loss of generality we can assume v^{λ} and $v^{\lambda,n}$ be positive. Otherwise we apply our argument to $\tilde{v}^{\lambda} = v^{\lambda} - \min_{\overline{\mathcal{O}}_m} v^{\lambda} + 1$ and the associated sequence $\tilde{v}^{\lambda,n} = v^{\lambda,n} - \min_{\overline{\mathcal{O}}_m} v^{\lambda} + 1$. By triangular inequality and positivity of v^{λ} and $v^{\lambda,n}$ we have

$$\max_{\overline{\mathcal{O}}_m} |v^{\lambda} - v^{\lambda, n_k}| \ge \max_{\overline{\mathcal{O}}_m} |v^{\lambda}| - \max_{\overline{\mathcal{O}}_m} |v^{\lambda, n_k}| = \max_{\overline{\mathcal{O}}_m} v^{\lambda} - \max_{\overline{\mathcal{O}}_m} v^{\lambda, n_k} \ge \frac{\epsilon}{4},$$

for all k's sufficiently large, where the inequality is due to (A.6). This contradicts uniform convergence and therefore $(\tilde{t}, \tilde{x}) \in \arg \max_{\overline{\mathcal{O}}_m} v^{\lambda}$ as claimed.

A.4. Existence of X^{ε} in the proof of Theorem 6.2. For $m \in \mathbb{N}$ let (b^m, σ^m) be functions $\mathbb{R}^d \to \mathbb{R}^d \times \mathbb{R}^{d \times d'}$ that are equal to (b, σ) on B_m and extend continuously to be constant on $\mathbb{R}^d \setminus B_m$. Similarly, $\alpha^m : \mathbb{R}^{d+1}_{0,T} \to \mathbb{R}^d$ is defined as

$$\alpha^m(t,x) = -2\psi_{\varepsilon}'\left(|\nabla u^{\varepsilon}(t,x)|_d^2 - f^2(t,x)\right)\nabla u^{\varepsilon}(t,x), \quad \text{on } B_m,$$

and extended continuously to be constant on $\mathbb{R}^d \setminus B_m$. Since $u^{\varepsilon} \in C^{0,1,\alpha}_{\ell oc}(\mathbb{R}^{d+1}_{0,T})$, then thanks to [38, Thm. 1] there exists a unique strong solution of

$$X_{s}^{m} = x + \int_{0}^{s} \left(b^{m}(X_{u}^{m}) + \alpha^{m}(t+u, X_{u}^{m}) \right) \mathrm{d}u + \int_{0}^{s} \sigma^{m}(X_{u}^{m}) \mathrm{d}W_{u}, \quad s \in [0, T-t].$$

Notice that $X^m = X^{m;t}$ depends on t via the time-inhomogeneous drift α^m but we omit it for simplicity. Notice also that here we should understand $n_s^m \dot{\nu}_s^m = \alpha^m (t+s, X_s^m)$ and $X^m = X^{m;[n^m,\nu^m]}$. Letting $\zeta_{m,k} = \inf\{s \ge 0 : |X_s^m|_d \ge k\}$, for any $m \ge k$ we have $X_{s \land \zeta_{m,k}}^m = X_{s \land \zeta_{k,k}}^k$ for all $s \in [0, T-t]$, P_x -a.s. (i.e., the two processes are indistinguishable). Thus, setting $X_s^{\varepsilon}(\omega) := X_s^k(\omega)$ for $s < \zeta_{k,k}(\omega)$ and denoting $\tau_k = \inf\{s \ge 0 : |X_s^{\varepsilon}|_d \ge k\}$ it is clear that, by uniqueness of strong solutions and the definition of the pair $(n^{\varepsilon}, \nu^{\varepsilon})$, the process X^{ε} satisfies

$$X_{s\wedge\tau_k}^{\varepsilon} = x + \int_0^{s\wedge\tau_k} \left(b(X_u^{\varepsilon}) + n_u^{\varepsilon} \dot{\nu}_u^{\varepsilon} \right) \mathrm{d}u + \int_0^{s\wedge\tau_k} \sigma(X_u^{\varepsilon}) \mathrm{d}W_u, \quad s \in [0, T-t].$$

By continuity of paths $\tau_k \leq \tau_{k+1}$ and $\tau_{\infty} := \lim_{k \to \infty} \tau_k$ is well-defined. Moreover, ν^{ε} satisfies the same bound as in (5.26) thanks to (6.7). Therefore, linear growth of the coefficients of the SDE and the same arguments as those at the end of the proof of Proposition 5.4 imply that $\mathsf{P}_x(\tau_{\infty} \leq T-t) = 0$. Then X^{ε} is well-defined on [0, T-t].

A.5. Shaefer's fixed point theorem.

Theorem A.3 (Thm. 9.2.4 in [15]). Suppose $T : \mathcal{D} \to \mathcal{D}$ is a continuous and compact mapping on a Banach space \mathcal{D} . Assume further that the set

$$\{f \in \mathcal{D} \mid f = \rho T[f] \text{ for some } \rho \in [0,1]\}$$

is bounded. Then T has a fixed point.

References

- L.H.R. Alvarez. Singular stochastic control in the presence of a state-dependent yield structure. *Stochastic Process.* Appl., 86(2):323–343, 2000.
- [2] E. Bayraktar and Y.-J. Huang. On the multidimensional controller-and-stopper games. SIAM J. Control Optim., 51(2):1263–1297, 2013.
- [3] E. Bayraktar and V.R. Young. Proving regularity of the minimal probability of ruin via a game of stopping and control. *Finance Stoch.*, 15(4):785–818, 2011.
- [4] A. Bensoussan and A. Friedman. Nonlinear variational inequalities and differential games with stopping times. J. Funct. Anal., 16(3):305–352, 1974.

- [5] A. Bensoussan and A. Friedman. Nonzero-sum stochastic differential games with stopping times and free boundary problems. Trans. Amer. Math. Soc., 231(2):275–327, 1977.
- [6] A. Bensoussan and J.-L. Lions. Applications of variational inequalities in stochastic control, volume 12 of Studies in Mathematics and its Applications. North-Holland, Amsterdam, New York, Oxford, 1982.
- [7] A. Budhiraja and K. Ross. Existence of optimal controls for singular control problems with state constraints. Ann. Appl. Probab., 16(4):2235-2255, 2006.
- [8] M.B. Chiarolla and U.G. Haussmann. Optimal control of inflation: A central bank problem. SIAM J. Control Optim., 36(3):1099–1132, 1998.
- M.B. Chiarolla and U.G. Haussmann. On a stochastic, irreversible investment problem. SIAM J. Control Optim., 48(2):438–462, 2009.
- [10] S. Choukroun, A. Cosso, and H. Pham. Reflected BSDEs with nonpositive jumps, and controller-and-stopper games. *Stochastic Process. Appl.*, 125(2):597–633, 2015.
- [11] P.-L. Chow, J.-L. Menaldi, and M. Robin. Additive control of stochastic linear systems with finite horizon. SIAM J. Control Optim., 23(6):858–899, 1985.
- [12] T. De Angelis and E. Ekström. The dividend problem with a finite horizon. Ann. Appl. Probab., 27(6):3525–3546, 2017.
- [13] T. De Angelis and G. Ferrari. A stochastic partially reversible investment problem on a finite time-horizon: Free-boundary analysis. *Stochastic Process. Appl.*, 124(12):4080–4119, 2014.
- [14] L.C. Evans. A second order elliptic equation with gradient constraint. Comm. Partial Differential Equations, 4(5):555–572, 1979.
- [15] L.C. Evans. Partial differential equations. American Mathematical Society, Providence, R.I., 2010.
- [16] W.H. Fleming and R.W. Rishel. Deterministic and stochastic optimal control, volume 1 of Applications of Mathematics. Springer-Verlag, New York Inc, 1986.
- [17] W.H. Fleming and H.M. Soner. Controlled Markov processes and viscosity solutions, volume 25. Springer Science & Business Media, 2006.
- [18] A. Friedman. Stochastic games and variational inequalities. Arch. Ration. Mech. Anal., 51(5):321–346, 1973.
- [19] A. Friedman. Partial differential equations of parabolic type. Courier Dover Publications, 2008.
- [20] D. Gilbarg and N.S. Trudinger. Elliptic partial differential equations of second order. Springer-Verlag, Berlin, Heidelberg, 2001.
- [21] S. Hamadène. Mixed zero-sum stochastic differential game and American game options. SIAM J. Control Optim., 45(2):496–518, 2006.
- [22] U.G. Haussmann and W. Suo. Singular optimal stochastic controls I: Existence. SIAM J. Control Optim., 33(3):916–936, 1995.
- [23] D. Hernandez-Hernandez, R.S. Simon, and M. Zervos. A zero-sum game between a singular stochastic controller and a discretionary stopper. Ann. Appl. Probab., 25(1):46 – 80, 2015.
- [24] R. Hynd. Partial differential equations with gradient constraints arising in the optimal control of singular stochastic processes. PhD thesis, UC Berkeley, 2010.
- [25] R. Hynd. Analysis of Hamilton-Jacobi-Bellman equations arising in stochastic singular control. ESAIM Control Optim. Calc. Var., 19(1):112–128, 2013.
- [26] I. Karatzas and W.D. Sudderth. The controller-and-stopper game for a linear diffusion. Ann. Probab., 29(3):1111– 1127, 2001.
- [27] I. Karatzas and I.-M. Zamfirescu. Martingale approach to stochastic differential games of control and stopping. Ann. Probab., 36(4):1495–1527, 2008.
- [28] M. Kelbert and H.A. Moreno-Franco. HJB equations with gradient constraint associated with controlled jumpdiffusion processes. SIAM J. Control Optim., 57(3):2185–2213, 2019.
- [29] N.V. Krylov. Controlled diffusion processes, volume 14 of Stochastic Modelling and Applied Probability. Springer, New York, NY, 1980. Reprint: Springer-Verlag Berlin Heidelberg, 2009.
- [30] N.V. Krylov. Lectures on elliptic and parabolic equations in Sobolev spaces, volume 96. American Mathematical Soc., 2008.
- [31] J. Li and G. Zitkovic. Existence, characterization, and approximation in the generalized monotone-follower problem. SIAM J. Control Optim., 55(1):94–118, 2017.
- [32] A.P. Maitra and W.D. Sudderth. The gambler and the stopper. Lecture Notes-Monograph Series, 30:191–208, 1996.
- [33] J.L. Menaldi and M.I. Taksar. Optimal correction problem of a multidimensional stochastic system. Automatica J. IFAC, 25(2):223–232, 1989.
- [34] H.M. Soner and S.E. Shreve. Regularity of the value function for a two-dimensional singular stochastic control problem. SIAM J. Control Optim., 27(4):876–907, 1989.
- [35] H.M. Soner and S.E. Shreve. A free boundary problem related to singular stochastic control: the parabolic case. Comm. Partial Differential Equations, 16(2-3):373–424, 1991.

- [36] L. Stettner. Penalty method for finite horizon stopping problems. SIAM J. Control Optim., 49(3):1078–1099, 2011.
- [37] M.I. Taksar. Skorohod problems with nonsmooth boundary conditions. J. Comput. Appl. Math., 40(2):233–251, 1992.
- [38] A.Y. Veretennikov. On strong solutions and explicit formulas for solutions of stochastic integral equations. Mat. Sb., 39(3):387–403, 1981.
- [39] A. Weerasinghe. A controller and a stopper game with degenerate variance control. Electron. Commun. Probab., 11:89–99, 2006.
- [40] H. Zhu. Generalized solution in singular stochastic control: the nondegenerate problem. *Appl. Math. Optim.*, 25(3):225–245, 1992.

A. BOVO: SCHOOL OF MANAGEMENT AND ECONOMICS, DEPT. ESOMAS, UNIVERSITY OF TORINO, CORSO UNIONE SOVIETICA, 218 BIS, 10134, TORINO, ITALY.

Email address: andrea.bovo@unito.it

T. DE ANGELIS: SCHOOL OF MANAGEMENT AND ECONOMICS, DEPT. ESOMAS, UNIVERSITY OF TORINO, CORSO UNIONE SOVIETICA, 218 BIS, 10134, TORINO, ITALY; COLLEGIO CARLO ALBERTO, PIAZZA ARBARELLO 8, 10122, TORINO, ITALY.

Email address: tiziano.deangelis@unito.it

E. Issoglio: Dept. of Mathematics "G. Peano", University of Torino, Via Carlo Alberto, 10, 10123, Torino, Italy.

Email address: elena.issoglio@unito.it