

Improved Tail Estimates for the Distribution of Quadratic Weyl Sums

Francesco Cellarosi^{*†} Jory Griffin[‡] Tariq Osman[§].

February 7, 2023

Abstract

We consider quadratic Weyl sums $S_N(x; c, \alpha) = \sum_{n=1}^N \exp \{2\pi i ((\frac{1}{2}n^2 + cn)x + \alpha n)\}$ for $c = \alpha = 0$ (the rational case) or $(c, \alpha) \notin \mathbb{Q}^2$ (the irrational case), where $x \in \mathbb{R}$ is randomly distributed according to a probability measure absolutely continuous with respect to the Lebesgue measure. The limiting distribution in the complex plane of $\frac{1}{\sqrt{N}} S_N(x; c, \alpha)$ as $N \rightarrow \infty$ was described in [17] (respectively [6]) in the rational (resp. irrational) case. According to the limiting distribution, the probability of landing outside a ball of radius R is known to be asymptotic to $\frac{4 \log 2}{\pi^2} R^{-4}(1 + o(1))$ in the rational case and to $\frac{6}{\pi^2} R^{-6}(1 + O(R^{-12/31}))$ in the irrational case, as $R \rightarrow \infty$. In this work we refine the technique of [6] to improve the known tail estimates to $\frac{4 \log 2}{\pi^2} R^{-4}(1 + O_\varepsilon(R^{-2+\varepsilon}))$ and $\frac{6}{\pi^2} R^{-6}(1 + O_\varepsilon(R^{-2+\varepsilon}))$ for every $\varepsilon > 0$. In the rational case, we rely on the equidistribution of a rational horocycle lift to a torus bundle over the unit tangent bundle to the classical modular surface. All the constants implied by the O_ε -notations are made explicit.

Contents

1	Introduction	3
2	Preliminaries	8
2.1	The universal Jacobi group and $\mathrm{ASL}(2, \mathbb{R})$	9
2.2	Integral transforms on $L^2(\mathbb{R})$	10
2.3	The Schrödinger-Weil representation of \tilde{G} on $L^2(\mathbb{R})$	10
2.4	Jacobi theta functions	11
2.5	Invariance of $\Theta_{f_1} \overline{\Theta_{f_2}}$	12
2.6	Geodesic and horocycle flows	15

^{*}Department of Mathematics and Statistics. Queen's University. Kingston, ON, Canada.

[†]Corresponding author: fc19@queensu.ca

[‡]School of Mathematics. University of Bristol. Bristol, U.K.

[§]Department of Mathematics. Brandeis University. Waltham, MA, U.S.A.

3	Limit Theorems	15
3.1	Equidistribution in the smooth rational case	15
3.2	Equidistribution in the smooth irrational case	17
3.3	Limit theorems for indicators	17
4	Growth in the cusp and tail bounds for $\Theta_{f_2} \overline{\Theta_{f_2}}$	22
4.1	Tail bound in the smooth case	23
4.1.1	Tail bound in the smooth rational case	23
4.1.2	Tail bound in the smooth irrational case	26
5	Dynamical approximation of the indicators	28
5.1	Smooth trapezoidal approximation of indicators	28
5.2	Defining Θ_{χ_s} almost everywhere	30
5.3	Expansion of the product $\Theta_{\chi} \overline{\Theta_{\chi_b}}$	31
6	Estimating κ_{η} for $1 < \eta \leq 2$	32
6.1	Divergence of expected displacement for harmonic quantum oscillator with piecewise initial data	36
7	Tail approximations	38
7.1	Rational case	38
7.2	Irrational case	42
8	Proof of the main theorems	46
8.1	Rational case	46
8.2	Irrational case	53
8.3	Final discussion on the explicit constants	57
A	Bounds for the zeta function	59
	References	61

1 Introduction

Set $e(x) := e^{2\pi i x}$ and consider quadratic Weyl sums of the form

$$S_N(x) = S_N(x; c, \alpha) = \sum_{n=1}^N e\left(\left(\frac{1}{2}n^2 + cn\right)x + \alpha n\right), \quad (1.1)$$

where N is a positive integer, and x, c and α are real. We assume that c and α are fixed, and that x in (1.1) is randomly distributed on \mathbb{R} according to a probability measure λ , absolutely continuous with respect to the Lebesgue measure on \mathbb{R} . Sums (1.1) can be thought of as the N -th partial sum of a particular sequence of strongly dependent random variables on the unit circle, or as the position after N steps of a deterministic walk in \mathbb{C} with a random seed x . It is natural to ask how the parameters (c, α) affect the distribution of (1.1), after suitable rescaling, as $N \rightarrow \infty$. The limiting distribution on the complex plane of $\frac{S_N}{\sqrt{N}}$ as $N \rightarrow \infty$ has been studied by Marklof [17] when $c = \alpha = 0$ and by Cellarosi and Marklof [6] when $(c, \alpha) \notin \mathbb{Q}^2$. We shall refer to the two cases as *rational* and *irrational*, respectively, see also Remark 1.4 below. In both cases the limiting distribution of $\frac{S_N}{\sqrt{N}}$ is not Gaussian, as can be seen by the following heavy tailed asymptotics.

- Rational case: if $c = \alpha = 0$, then

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \frac{|S_N(x; 0, 0)|}{\sqrt{N}} > R \right\} = \frac{4 \log 2}{\pi^2} \frac{1}{R^4} (1 + o(1)) \quad (1.2)$$

as $R \rightarrow \infty$, see [17].

- Irrational case: if $(c, \alpha) \notin \mathbb{Q}^2$, then

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \frac{|S_N(x; c, \alpha)|}{\sqrt{N}} > R \right\} = \frac{6}{\pi^2} \frac{1}{R^6} \left(1 + O\left(R^{-\frac{12}{31}}\right) \right) \quad (1.3)$$

as $R \rightarrow \infty$, see [6].

More generally, the convergence of $t \mapsto \frac{1}{\sqrt{N}} S_{[tN]}$ as $N \rightarrow \infty$ to a random process has been shown by Cellarosi [5] and Cellarosi and Marklof [6]. In this paper we are able to improve the tail estimates (1.2)-(1.3) of [17] and [6] to the following ‘optimal’ ones (see Remark 1.8 below).

Theorem 1.1. *For all $0 < \varepsilon \leq 1$*

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \frac{|S_N(x; 0, 0)|}{\sqrt{N}} > R \right\} = \frac{4 \log 2}{\pi^2 R^4} (1 + O_\varepsilon(R^{-2+\varepsilon})) \quad (1.4)$$

as $R \rightarrow \infty$.

Theorem 1.2. *Let $(c, \alpha) \notin \mathbb{Q}^2$. Then for all $0 < \varepsilon \leq 1$*

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \frac{|S_N(x; c, \alpha)|}{\sqrt{N}} > R \right\} = \frac{6}{\pi^2 R^6} (1 + O_\varepsilon(R^{-2+\varepsilon})) \quad (1.5)$$

as $R \rightarrow \infty$.

The recent literature includes several examples of heavy-tailed limiting distribution arising from the study of problems in number theory (e.g. the distribution of visible lattice points by Boca, Cobeli and Zaharescu [2], the gaps distribution between Farey fractions satisfying divisibility constraints by Boca, Heersink and Spiegelhalter [3], the distribution of short Gauss sums by Demirci Akarsu [8], the celebrated result by Elkies and McMullen on the gaps in the sequence $(\sqrt{n} \bmod 1)_{n \geq 1}$ [13], and the distribution of Frobenius numbers by Marklof [20]), mathematical physics (e.g. the distribution of free-path lengths in various Lorentz gas models in the small scatterer limit, see the works of Boca and Zaharescu [4], Marklof and Strömbergsson [21]-[22]-[23], and Nándori, Szász and Varjú [24]), and ergodic theory (e.g. the gap distribution between the slopes of saddle connections in flat surfaces as in the works of Athreya, Chaika and Lelièvre [1], of Uyanik and Work [26], as well as the deviations of ergodic averages for toral rotations by Dolgopyat and Fayad [10]-[11]).

Theorems 1.1 and 1.2 address the size of fluctuations about the tail of the limiting distribution of rescaled quadratic Weyl sums. Our results can also be interpreted as refined large deviations estimates. In Figure 1 we illustrate how the distribution of $\frac{|S_N|}{\sqrt{N}}$ approaches a different limiting distribution as $N \rightarrow \infty$ in the rational and irrational cases. The predicted tail asymptotics for the limiting distributions are shown in Figure 2, along with the fluctuations predicted by Theorems 1.1 and 1.2.

Remark 1.3. The slightly more general sums

$$\sum_{n=1}^N e\left(\left(\frac{1}{2}n^2 + c_1 n + c_0\right)x + \alpha n\right) \quad (1.6)$$

are considered in [6]. In this work we are only concerned in the absolute value of such sums, which does not depend on c_0 , and hence we simplify the notation by setting $c_0 = 0$ and $c_1 = c$.

Remark 1.4. The remaining cases not covered by Theorems 1.1 and 1.2 correspond to $(c, \alpha) \in \mathbb{Q}^2 \setminus \{(0, 0)\}$. For *all* rational pairs $(c, \alpha) \in \mathbb{Q}^2$ we have an asymptotic tail formula similar to (1.4), namely of the form

$$\frac{\mathcal{T}(c, \alpha)}{R^4} (1 + O_\varepsilon(R^{-2+\varepsilon})), \quad (1.7)$$

where $\mathcal{T}(c, \alpha) \geq 0$ for all $(c, \alpha) \in \mathbb{Q}^2$ and $\mathcal{T}(0, 0) = \frac{4 \log 2}{\pi^2}$. To find a formula for $\mathcal{T}(c, \alpha)$ in the general rational case, one can use a combination of number-theoretical and group-theoretical ideas which are independent of the techniques used here to obtain the error




Figure 1: Histograms of the empirical distribution of $\frac{1}{\sqrt{N}}|S_N(x; c, \alpha)|$ for 10^6 uniformly sampled values of $x \in [0, 1]$. We use $c = \alpha = 0$ (top row), $(c, \alpha) = (0, \sqrt{2})$ (bottom row), $N = 100$ (left column), $N = 500$ (middle column), and $N = 1000$ (right column).

term $O_\varepsilon(R^{-2+\varepsilon})$ in (1.4) and (1.7). We therefore defer the determination of $\mathcal{T}(c, \alpha)$ to a separate work [7] and here we only consider the rational case $c = \alpha = 0$ as in Theorem 1.1. It is nevertheless worthwhile to mention that there exist infinitely many rational pairs (c, α) for which $\mathcal{T}(c, \alpha) = 0$. For such parameters, such as $(c, \alpha) = (\frac{1}{2(2k-1)}, \frac{1}{2(2k-1)})$ with $k \in \mathbb{N}$, the limiting distribution of $\frac{S_N}{\sqrt{N}}$ is therefore compactly supported. Other examples of compactly supported limiting distributions can be found in the work of Demirci Akarsu and Marklof on incomplete Gauss sums [9], and of Kowalski and Sawin on Kloosterman and Birch sums [15].

In this paper we actually obtain precise tail estimates for the limiting distribution of products of sums of the form (1.1), i.e. $\lim_{N \rightarrow \infty} \lambda \{x \in \mathbb{R} : |\frac{1}{N}S_{[aN]}(x; c, \alpha)\overline{S_{[bN]}(x; c, \alpha)}| > R^2\}$ where, up to relabelling, $0 < a \leq b$. Furthermore, without loss of generality, we may assume that $1 = a \leq b$ since

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{N}S_{[aN]}(x; c, \alpha)\overline{S_{[bN]}(x; c, \alpha)} \right| > R^2 \right\} = \quad (1.8)$$

$$= \lim_{M \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{M}S_M(x; c, \alpha)\overline{S_{[\frac{b}{a}M]}(x; c, \alpha)} \right| > (\sqrt{a}R)^2 \right\}. \quad (1.9)$$

The following two theorems are the main result of this paper.




Figure 2: We sample 5×10^6 points according to (an approximation of) the limiting distribution in the rational case (top row) and in the irrational case (bottom row), see Remark 3.10. The empirical distributions corresponding to our samples are shown in the left panels along with the predicted tail density, obtained by differentiating $1 - \frac{4 \log 2}{\pi^2} R^{-4}$ in the rational case and $1 - \frac{6}{\pi^2} R^{-6}$ in the irrational case, with respect to R . In the right panels we illustrate Theorems 1.1-1.2 for $\varepsilon = \frac{1}{2}$ and plot $(1 - \frac{4 \log 2}{\pi^2} R^{-4} - \text{cdf}(R)) R^{5.5}$ in the rational case, and $(1 - \frac{6}{\pi^2} R^{-6} - \text{cdf}(R)) R^{7.5}$ in the irrational case, where $\text{cdf}(R)$ denotes the empirical distribution function. Our results show the existence of a uniform horizontal band within which these graphs are confined. Given our sample size, we expect to see $\frac{4 \log 2}{\pi^2} 16^{-4} \times 5 \times 10^6 \approx 21$ values larger than 16 in the rational case and $\frac{6}{\pi^2} 7^{-6} \times 5 \times 10^6 \approx 26$ values larger than 7 in the irrational case. This is the reason why the ranges of R shown in the two cases end at 16 and 7, respectively. In this way, the accuracy of these graphs is comparable. The bin size in the two histograms on the left is the same.

Theorem 1.5. *Let $b \geq 1$ and $0 < \varepsilon \leq 1$. There exists a constant $R^{\text{rat}}(b, \varepsilon) > 0$ such that for all $R > R^{\text{rat}}(b, \varepsilon)$ we have*

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{N} S_N(x; 0, 0) \overline{S_{[bN]}(x; 0, 0)} \right| > R^2 \right\} = \frac{2D_{\text{rat}}(b)}{\pi^2 R^4} (1 + O_\varepsilon(R^{-2+\varepsilon})), \quad (1.10)$$

where

$$D_{\text{rat}}(b) = \begin{cases} 2 \log 2, & \text{if } b = 1, \\ 2b \coth^{-1}(b) + \frac{1}{2} \log(b^2 - 1) + \frac{b^2}{2} \log(1 - \frac{1}{b^2}), & \text{if } b > 1. \end{cases} \quad (1.11)$$

Theorem 1.6. *Let $(c, \alpha) \notin \mathbb{Q}^2$, $b \geq 1$, and $0 < \varepsilon \leq 1$. There exists a constant $R^{\text{irr}}(b, \varepsilon) > 0$ such that for all $R > R^{\text{irr}}(b, \varepsilon)$ we have*

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{N} S_N(x; c, \alpha) \overline{S_{[bN]}(x; c, \alpha)} \right| > R^2 \right\} = \frac{2D_{\text{irr}}(b)}{\pi^2 R^6} (1 + O_\varepsilon(R^{-2+\varepsilon})), \quad (1.12)$$

where $D_{\text{irr}}(b)$ is given in (8.57) and $D_{\text{irr}}(1) = 3$.

In this paper, we aim to make all the constants implied by the O -notations explicit and simple, often at the cost of making such constants bigger (see also Section 8.3). Explicit versions of Theorems 1.5 and 1.6 are given by Theorems 8.4 and 8.7, respectively. By taking $b = 1$ we obtain Theorems 1.1 and 1.2 as particular cases of Theorems 1.5 and 1.6. Their explicit versions are given in Corollaries 8.5 and 8.8, respectively. To obtain Theorems 1.5 and 1.6, we first prove a more general tail estimate for products of Weyl sums of the form

$$S_N(x; c, \alpha; f) = \sum_{n \in \mathbb{Z}} f\left(\frac{n}{N}\right) e\left(\left(\frac{1}{2}n^2 + cn\right)x + n\alpha\right), \quad (1.13)$$

where $f : \mathbb{R} \rightarrow \mathbb{R}$ is bounded and of sufficient decay at $\pm\infty$ so that the series (1.13) is absolutely convergent. Note that (1.13) reduces to (1.1) when $f = \mathbf{1}_{(0,1]}$. Assuming a certain degree of decay and regularity for two functions f_1 and f_2 , we can accurately study the tails of the limiting distribution of the product $\frac{1}{N} S_N(x; c, \alpha; f_1) \overline{S_N(x; c, \alpha; f_2)}$ as $N \rightarrow \infty$, where x is distributed on \mathbb{R} according to the probability measure λ . Using the uniform decay of a family of Fourier-like transforms, we define the regularity class $\mathcal{S}_\eta(\mathbb{R})$, see Section 2.4. We then prove the following

Theorem 1.7. *Let $\eta > 1$ and let $f_1, f_2 \in \mathcal{S}_\eta(\mathbb{R})$. Then*

$$\begin{aligned} \lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{N} S_N(x; c, \alpha; f_1) \overline{S_N(x; c, \alpha; f_2)} \right| > R^2 \right\} = \\ = \begin{cases} \frac{2D_{\text{rat}}(f_1, f_2)}{\pi^2 R^4} (1 + O(R^{-2\eta})), & \text{if } c = \alpha = 0, \\ \frac{2D_{\text{irr}}(f_1, f_2)}{\pi^2 R^6} (1 + O(R^{-2\eta})), & \text{if } (c, \alpha) \notin \mathbb{Q}^2, \end{cases} \end{aligned} \quad (1.14)$$

where $D_{\text{rat}}(f_1, f_2)$ and $D_{\text{irr}}(f_1, f_2)$ are given in (4.10) and (4.37), respectively. The constants implied by the O -notations in (1.14) depend explicitly on η, f_1, f_2 .

Both instances of Theorem 1.7 are a combination of a dynamical statement and an analytical tail estimate. The dynamical ingredient is the equidistribution of horocycle lifts under the ‘stretching’ action of the geodesic flow in a suitably defined 5-dimensional homogeneous space $\Gamma \backslash G$. In the rational case, the equidistribution takes place on a codimension-2 submanifold (see Theorem 3.2, extending Theorem 5.2 in [17]), while in the irrational case, already studied in [6], horocycle lifts equidistribute in the whole space according to its normalised Haar measure (see Theorem 3.3). Therefore, the dynamics singles out a limiting probability measure on $\Gamma \backslash G$, which is different in the two cases we consider, and we are left to study the push-forward of such probability measures via a product of two Jacobi theta functions $\Theta_{f_1} \overline{\Theta_{f_2}} : \Gamma \backslash G \rightarrow \mathbb{C}$. The corresponding probability measures on \mathbb{C} are the ones whose tail behaviour we address in Lemmata 4.2 and 4.3.

Remark 1.8. The assumption $\eta > 1$ in Theorem 1.7 is crucial, and hence Theorems 1.5-1.6, in which $\eta = 1$, do not follow directly. Consequently, the error terms in Theorems 1.5-1.6 are ‘optimal’, in the sense that improving the statements to include $\varepsilon = 0$ is not possible. Following the strategy of [6], we use a careful smoothing procedure and a dynamically defined partition of unity to, in effect, increase the regularity η . This then allows us to derive (1.10) and (1.12) from (1.14). By improving a key estimate of [6], in which the case $\eta = 2$ is considered, and instead pushing $\eta \rightarrow 1^+$, we achieve the improved error terms in (1.10) and (1.12).

The above-mentioned improvement follows from our Theorem 6.4, which we also use to illustrate the divergence of the expected displacement of a quantum harmonic oscillator as the initial condition approaches a piecewise C^1 function with finitely many jump discontinuities.

2 Preliminaries

In this section we define the Jacobi theta function Θ_f , which plays a central role in our analysis. We will see how the quantity $\frac{1}{N} S_N(x; c, \alpha; f_1) \overline{S_N(x; c, \alpha; f_2)}$ can be seen as the product of two theta functions $\Theta_{f_1} \overline{\Theta_{f_2}}$ evaluated at certain points in the Lie group $G = \text{SL}(2, \mathbb{R}) \ltimes \mathbb{R}^2$. We also introduce *two* measures on G . The first one is the Haar measure μ , which is classical and is used in the analysis of the irrational case $(c, \alpha) \notin \mathbb{Q}^2$. The second one, μ^0 , is the product of the Haar measure on $\text{SL}(2, \mathbb{R})$ with an atomic measure on \mathbb{R}^2 and is used when addressing the rational case $(c, \alpha) = (0, 0)$. The analysis of all other rational cases (see Remark 1.4) requires similar measures $\mu^{(c, \alpha)}$. We also introduce a lattice $\Gamma < G$ under which $\Theta_{f_1} \overline{\Theta_{f_2}}$ is invariant.

2.1 The universal Jacobi group and $\text{ASL}(2, \mathbb{R})$

Let $\mathfrak{H} := \{w \in \mathbb{C} : \text{Im}(w) > 0\}$ denote the upper half plane. For each $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}(2, \mathbb{R})$ and $z \in \mathfrak{H}$, we let $Mz = \frac{az+b}{cz+d}$ and $\epsilon_M(z) = (cz+d)/|cz+d|$. Let $\widetilde{\text{SL}}(2, \mathbb{R})$ denote the universal cover of $\text{SL}(2, \mathbb{R})$, i.e.

$$\widetilde{\text{SL}}(2, \mathbb{R}) := \{[M, \beta_M] : M \in \text{SL}(2, \mathbb{R}), \beta_M \text{ a continuous function on } \mathfrak{H} \text{ s.t. } e^{i\beta_M(z)} = \epsilon_M(z)\}. \quad (2.1)$$

The group law on $\widetilde{\text{SL}}(2, \mathbb{R})$ is given by

$$[M_1, \beta'_{M_1}][M_2, \beta''_{M_2}] = [M_1 M_2, \beta'''_{M_1 M_2}], \quad \beta'''_{M_1 M_2}(z) = \beta'_{M_1}(M_2 z) + \beta''_{M_2}(z), \quad (2.2)$$

$$[M, \beta_M]^{-1} = [M^{-1}, \beta'_{M^{-1}}], \quad \beta'_{M^{-1}}(z) = -\beta_M(M^{-1}z). \quad (2.3)$$

We identify $\widetilde{\text{SL}}(2, \mathbb{R})$ with $\mathfrak{H} \times \mathbb{R}$ via $[M, \beta_M] \mapsto (z, \phi) = (Mi, \beta_M(i))$ and we have an action of $\widetilde{\text{SL}}(2, \mathbb{R})$ on $\mathfrak{H} \times \mathbb{R}$ given by

$$[M, \beta_M](z, \phi) = (Mz, \phi + \beta_M(z)). \quad (2.4)$$

Let $\omega : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$ be the standard symplectic form $\omega(\xi, \xi') = xy' - yx'$, where $\xi = \begin{pmatrix} x \\ y \end{pmatrix}, \xi' = \begin{pmatrix} x' \\ y' \end{pmatrix}$. The Heisenberg group $\mathbb{H}(\mathbb{R})$ is defined as $\mathbb{R}^2 \times \mathbb{R}$ with group law

$$(\xi, t)(\xi', t') = (\xi + \xi', t + t' + \frac{1}{2}\omega(\xi, \xi')). \quad (2.5)$$

We consider the *universal Jacobi group* $\tilde{G} = \widetilde{\text{SL}}(2, \mathbb{R}) \ltimes \mathbb{H}(\mathbb{R})$, with group law

$$([M, \beta_M]; \xi, \zeta)([M', \beta'_{M'}]; \xi', \zeta') = ([MM', \beta''_{MM'}]; \xi + M\xi', \zeta + \zeta' + \frac{1}{2}\omega(\xi, g\xi')), \quad (2.6)$$

where $\beta''_{MM'}(z) = \beta_M(M'z) + \beta'_{M'}(z)$ as in (2.2). Identifying \tilde{G} with $\mathfrak{H} \times \mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}$, the Haar measure $\tilde{\mu}$ on \tilde{G} is given in coordinates $\tilde{g} = (x + iy, \phi; (\xi_1, \xi_2), \zeta)$ by $d\tilde{\mu}(\tilde{g}) = \frac{1}{y^2} dx dy d\phi d\xi_1 d\xi_2 d\zeta$. Later, we will focus our attention on the special affine group $G := \text{ASL}(2, \mathbb{R}) = \text{SL}(2, \mathbb{R}) \ltimes \mathbb{R}^2$, which is the range of the homomorphism

$$\pi : \tilde{G} \rightarrow G, \quad \pi([M, \beta_M]; \xi, \zeta) = (M, \xi) \in G \quad (2.7)$$

and hence

$$G \cong \tilde{G} / \{([I, 2\pi k]; \mathbf{0}, \zeta) : k \in \mathbb{Z}, \zeta \in \mathbb{R}\}. \quad (2.8)$$

The group law on G is

$$(M; \xi)(M'; \xi') = (MM'; \xi + M\xi') \quad (2.9)$$

and by identifying G with $\mathfrak{H} \times [0, 2\pi) \times \mathbb{R}^2$ the Haar measure μ on G is given in coordinates $g = (x + iy, \phi; (\xi_1, \xi_2))$, up to a multiplicative factor, by

$$d\mu(g) = \frac{3}{\pi^2} \frac{dx dy d\phi d\xi_1 d\xi_2}{y^2}. \quad (2.10)$$

We will also need the measure μ^0 on G , given by

$$d\mu^0(g) = \frac{1}{\pi^2} \frac{dx dy d\phi}{y^2} d\nu, \quad \nu = \sum_{\mathbf{k} \in \mathbb{Z}^2} \left(\delta_{\begin{pmatrix} 0 \\ 0 \end{pmatrix} + \mathbf{k}} + \delta_{\begin{pmatrix} 1/2 \\ 0 \end{pmatrix} + \mathbf{k}} + \delta_{\begin{pmatrix} 0 \\ 1/2 \end{pmatrix} + \mathbf{k}} \right), \quad (2.11)$$

which projects to the Haar measure on $\mathrm{SL}(2, \mathbb{R}) = \mathfrak{H} \times [0, 2\pi)$ and to an atomic measure on \mathbb{R}^2 . The reason for the normalising factors $\frac{3}{\pi^2}$ and $\frac{1}{\pi^2}$ in (2.10) and (2.11) will be clear in Section 2.5.

2.2 Integral transforms on $L^2(\mathbb{R})$

For $f \in L^2(\mathbb{R})$ and $\phi \in \mathbb{R}$, let us define the operator $\mathcal{I}_\phi : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ as

$$[\mathcal{I}_\phi f](w) = \begin{cases} f(w), & \text{if } \phi \equiv 0 \pmod{2\pi}, \\ f(-w), & \text{if } \phi \equiv \pi \pmod{2\pi}, \\ |\sin \phi|^{-1/2} \int_{\mathbb{R}} e\left(\frac{\frac{1}{2}(w^2 + w'^2) \cos \phi - ww'}{\sin \phi}\right) f(w') dw', & \text{if } \phi \not\equiv 0 \pmod{\pi}. \end{cases} \quad (2.12)$$

We will see in Section 6 how to interpret $w \mapsto [\mathcal{I}_\phi f](w)$ as the solution at time ϕ to a quantum harmonic oscillator with initial condition f . Let us also introduce the function $\phi \mapsto \sigma_\phi$, given by

$$\sigma_\phi := \begin{cases} 2\nu, & \text{if } \phi = \nu\pi, \nu \in \mathbb{Z}; \\ 2\nu + 1, & \text{if } \nu\pi < \phi < (\nu + 1)\pi, \nu \in \mathbb{Z}. \end{cases} \quad (2.13)$$

Finally, define the operator $\mathcal{F}_\phi : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ as

$$[\mathcal{F}_\phi f](w) = e(-\sigma_\phi/8) [\mathcal{I}_\phi f](w) \quad (2.14)$$

The reason for the factor $e(-\sigma_\phi/8)$ in (2.14) is that $\lim_{\phi \rightarrow 0^\pm} [\mathcal{I}_\phi f](w) = e(\pm \frac{1}{8}) f(w)$ for $f \in \mathcal{S}(\mathbb{R})$. The 1-parameter family $(\mathcal{F}_\phi)_{\phi \in \mathbb{R}}$ is a continuous group of unitary operators on $L^2(\mathbb{R})$. The particular case $\mathcal{F}_{\frac{\pi}{2}}$ coincides with the classical unitary Fourier transform. Note that $\phi \mapsto \mathcal{F}_\phi$ is 4π -periodic and that $\overline{\mathcal{F}_\phi f} = \mathcal{F}_{-\phi} \bar{f}$. For a more compact notation, unless explicitly stated, we will write f_ϕ instead of $\mathcal{F}_\phi f$.

2.3 The Schrödinger-Weil representation of \tilde{G} on $L^2(\mathbb{R})$

By Iwasawa decomposition, every $M \in \mathrm{SL}(2, \mathbb{R})$ can be written uniquely as

$$M = n_x a_y k_\phi, \quad (2.15)$$

where

$$n_x = \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}, \quad a_y = \begin{pmatrix} y^{1/2} & 0 \\ 0 & y^{-1/2} \end{pmatrix}, \quad k_\phi = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix}, \quad (2.16)$$

and $z = x + iy \in \mathfrak{H}$, $\phi \in [0, 2\pi)$. This allows us to parametrize $\mathrm{SL}(2, \mathbb{R})$ by $\mathfrak{H} \times [0, 2\pi)$. We will use the shorthand $(z, \phi) = n_x a_y k_\phi$. Recall the identification of $\widetilde{\mathrm{SL}}(2, \mathbb{R})$ with $\mathfrak{H} \times \mathbb{R}$. We can extend (2.15) to a decomposition of $\widetilde{\mathrm{SL}}(2, \mathbb{R})$: for every $\tilde{M} = [M, \beta_M] \in \widetilde{\mathrm{SL}}(2, \mathbb{R})$ we have

$$\tilde{M} = [M, \beta_M] = \tilde{n}_x \tilde{a}_y \tilde{k}_\phi = [n_x, 0][a_y, 0][k_\phi, \beta_{k_\phi}]. \quad (2.17)$$

The classical (projective) *Shale-Weil representation* of $\mathrm{SL}(2, \mathbb{R})$ can be lifted to a genuine representation of $\widetilde{\mathrm{SL}}(2, \mathbb{R})$ using (2.17), see [16]: for $f \in L^2(\mathbb{R})$ let

$$[R(\tilde{n}_x)f](w) = e\left(\frac{1}{2}w^2x\right) f(w), \quad (2.18)$$

$$[R(\tilde{a}_y)f](w) = y^{1/4} f(y^{1/2}w), \quad (2.19)$$

$$[R(\tilde{k}_\phi)f](w) = f_\phi(w). \quad (2.20)$$

Recalling (2.5), elements of $\mathbb{H}(\mathbb{R})$ can be decomposed as

$$\left(\begin{pmatrix} x \\ y \end{pmatrix}, \zeta\right) = \left(\begin{pmatrix} x \\ 0 \end{pmatrix}, 0\right) \left(\begin{pmatrix} 0 \\ y \end{pmatrix}, 0\right) \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \zeta - \frac{xy}{2}\right) \quad (2.21)$$

and this allows us to define the *Schrödinger representation* of $\mathbb{H}(\mathbb{R})$ on $L^2(\mathbb{R})$ as

$$[W\left(\begin{pmatrix} x \\ 0 \end{pmatrix}, 0\right)f](w) = e(xw) f(w), \quad (2.22)$$

$$[W\left(\begin{pmatrix} 0 \\ y \end{pmatrix}, 0\right)f](w) = f(w - y), \quad (2.23)$$

$$[W\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \zeta\right)f](w) = e(\zeta) f(w). \quad (2.24)$$

Combining the two above representations, we can define the *Schrödinger-Weil representation* of the universal Jacobi group. Recalling the identification of \tilde{G} with $\mathfrak{H} \times \mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}$, we set

$$\tilde{R}(z, \phi; \boldsymbol{\xi}, \zeta) = W(\boldsymbol{\xi}, \zeta) R(z, \phi), \quad (2.25)$$

for $(z, \phi; \boldsymbol{\xi}, \zeta) \in \tilde{G}$.

2.4 Jacobi theta functions

We consider the space of functions $f : \mathbb{R} \rightarrow \mathbb{R}$ for which f_ϕ defined in (2.14) has a certain decay at infinity, uniformly in ϕ : denote

$$\kappa_\eta(f) = \sup_{w, \phi} |f_\phi(w)| (1 + |w|^2)^{\frac{\eta}{2}}. \quad (2.26)$$

and define

$$\mathcal{S}_\eta(\mathbb{R}) := \{f : \mathbb{R} \rightarrow \mathbb{R} : \kappa_\eta(f) < \infty\}, \quad (2.27)$$

see [19]. These function spaces play a key role in our analysis when $\eta > 1$. We stress that the indicator functions $\mathbf{1}_{(0,s]}$ and $\chi_s = \mathbf{1}_{(0,s)}$ belong to \mathcal{S}_1 but not to \mathcal{S}_η for any $\eta > 1$. In Section 5, we introduce approximations $\chi_s^{(J)}$ of χ_s which belong to \mathcal{S}_η with $\eta > 1$ and it is crucial to estimate $\kappa_\eta(\chi_s^{(J)})$. Most of the analysis in [6] deals with the case $\eta = 2$; here we see how to obtain the optimal tail bounds (1.4) and (1.5) by pushing $\eta \rightarrow 1^+$.

We can now use the representation (2.25) to define the most important function in our investigation. For $(z, \phi; \boldsymbol{\xi}, \zeta) \in \tilde{G}$ and $f \in \mathcal{S}_\eta(\mathbb{R})$, with $\eta > 1$, define the *Jacobi theta function* as

$$\Theta_f(z, \phi; \boldsymbol{\xi}, \zeta) = \sum_{n \in \mathbb{Z}} [\tilde{R}(z, \phi; \boldsymbol{\xi}, \zeta) f](n) = \quad (2.28)$$

$$= y^{1/4} e(\zeta - \frac{1}{2} \xi_1 \xi_2) \sum_{n \in \mathbb{Z}} f_\phi((n - \xi_2) y^{1/2}) e(\frac{1}{2} (n - \xi_2)^2 x + n \xi_1), \quad (2.29)$$

where $z = x + iy$, and $\boldsymbol{\xi} = \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix}$. The reason for introducing theta functions is that we can rewrite (1.13), after normalization, as

$$N^{-\frac{1}{2}} S_N(x; c, \alpha; f) = \Theta_f(x + iN^{-2}, 0; (\alpha_0^{+cx}), 0). \quad (2.30)$$

2.5 Invariance of $\Theta_{f_1} \overline{\Theta_{f_2}}$

It is shown in [18] that if $f \in \mathcal{S}_\eta$ with $\eta > 1$, then that the function $\Theta_f : \tilde{G} \rightarrow \mathbb{C}$ is invariant under the left action by the lattice $\tilde{\Gamma} = \langle \tilde{\gamma}_1, \tilde{\gamma}_2, \tilde{\gamma}_3, \tilde{\gamma}_4, \tilde{\gamma}_5 \rangle$, where

$$\tilde{\gamma}_1 = \left(\left[\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \arg \right]; \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \frac{1}{8} \right) = \left(i, \frac{\pi}{2}; \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \frac{1}{8} \right), \quad (2.31)$$

$$\tilde{\gamma}_2 = \left(\left[\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, 0 \right]; \left(\begin{pmatrix} 1/2 \\ 0 \end{pmatrix}, 0 \right) = \left(1 + i, 0; \begin{pmatrix} 1/2 \\ 0 \end{pmatrix}, 0 \right), \quad (2.32)$$

$$\tilde{\gamma}_3 = \left(\left[\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, 0 \right]; \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, 0 \right) = \left(i, 0; \begin{pmatrix} 1 \\ 0 \end{pmatrix}, 0 \right), \quad (2.33)$$

$$\tilde{\gamma}_4 = \left(\left[\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, 0 \right]; \left(\begin{pmatrix} 0 \\ 1 \end{pmatrix}, 0 \right) = \left(i, 0; \begin{pmatrix} 0 \\ 1 \end{pmatrix}, 0 \right), \quad (2.34)$$

$$\tilde{\gamma}_5 = \left(\left[\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, 0 \right]; \left(\begin{pmatrix} 0 \\ 1 \end{pmatrix}, 0 \right) = \left(i, 0; \begin{pmatrix} 0 \\ 0 \end{pmatrix}, 1 \right). \quad (2.35)$$

Now, let us fix two functions $f_1, f_2 \in \mathcal{S}_\eta$ with $\eta > 1$ and consider the product $\Theta_{f_1} \overline{\Theta_{f_2}}$. Since $\Theta_{f_1} \overline{\Theta_{f_2}}(z, \phi; \boldsymbol{\xi}, \zeta)$ does not depend on ζ and depends only on the value of $\phi \bmod 2\pi$, recalling (2.8), we consider $\Theta_{f_1} \overline{\Theta_{f_2}}$ as a function on G . This function is therefore invariant

under the left action by the group $\Gamma = \pi(\tilde{\Gamma}) = \langle \gamma_1, \gamma_2, \gamma_3, \gamma_4 \rangle$, where $\gamma_i = \pi(\tilde{\gamma}_i)$ for $1 \leq i \leq 4$. Therefore $\Theta_{f_1} \overline{\Theta_{f_2}}$ is well-defined on the quotient $\Gamma \backslash G$, which is a non-compact 5-dimensional manifold with finite volume. A fundamental domain for the action of Γ on G is

$$\mathcal{F}_\Gamma = \left\{ (z, \phi; \boldsymbol{\xi}) \in \mathcal{F}_{\text{SL}(2, \mathbb{Z})} \times [0, \pi) \times \left(-\frac{1}{2}, \frac{1}{2}\right]^2 \right\}, \quad (2.36)$$

where

$$\mathcal{F}_{\text{SL}(2, \mathbb{Z})} = \left\{ z \in \mathfrak{H} : |z| > 1, -\frac{1}{2} \leq \text{Re}(z) < \frac{1}{2} \right\} \cup \left\{ z \in \mathfrak{H} : |z| = 1, \text{Re}(z) \leq 0 \right\}. \quad (2.37)$$

Note that \mathcal{F}_Γ can be seen as a torus bundle over the unit tangent bundle to the classical modular surface.

Lemma 2.1. *The measure μ^0 is Γ -invariant.*

Proof. Let $A = A_H \times A_R$ be a measurable rectangle, where A_H is a Borel subset of $\mathfrak{H} \times [0, 2\pi)$ and A_R is a Borel subset of \mathbb{R}^2 . Suppose that A has finite μ^0 -measure. Let us show that $\mu^0(\gamma^{-1}A) = \mu^0(A)$ for every $\gamma \in \Gamma$. The statement of the lemma follows from this fact. Since A is a measurable rectangle, $\mathbf{1}_A(x + iy, \phi; \boldsymbol{\xi}) = \mathbf{1}_{A_H}(x + iy, \phi) \mathbf{1}_{A_R}(\boldsymbol{\xi})$. Let $\gamma = (M; \mathbf{v})$. We have

$$\mu^0(\gamma^{-1}A) = \int_G \mathbf{1}_{\gamma^{-1}A}(x + iy, \phi; \boldsymbol{\xi}) d\mu^0 = \int_G \mathbf{1}_A(\gamma(x + iy, \phi; \boldsymbol{\xi})) d\mu^0 \quad (2.38)$$

$$= \int_G \mathbf{1}_{A_H}(M(x + iy, \phi)) \mathbf{1}_{A_R}(\gamma \cdot \boldsymbol{\xi}) d\mu^0 \quad (2.39)$$

$$= \int_{\mathbb{R}^2} \mathbf{1}_{A_R}(\gamma \cdot \boldsymbol{\xi}) d\nu \int_0^\pi \int_0^\infty \int_{-\infty}^\infty \mathbf{1}_{A_H}(M(x + iy, \phi)) \frac{dx dy d\phi}{y^2} \quad (2.40)$$

Note that the action of M on $\text{SL}(2, \mathbb{R})$ preserves the measure $\frac{dx dy d\phi}{y^2}$. Moreover, using the generators of Γ , we see that the action of each $\gamma \in \Gamma$ is a bijection on the support of ν , i.e.

$$\bigcup_{\mathbf{k} \in \mathbb{Z}^2} \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} + \mathbf{k}, \begin{pmatrix} 1/2 \\ 0 \end{pmatrix} + \mathbf{k}, \begin{pmatrix} 0 \\ 1/2 \end{pmatrix} + \mathbf{k} \right\}. \quad (2.41)$$

Since ν is atomic with equal weights for its atoms, we obtain $\gamma_*^{-1}\nu = \nu$. Therefore (2.40) implies $\mu^0(\gamma^{-1}A) = \mu^0(A)$. \square

The projection of the measure μ (respectively μ^0) from G to $\Gamma \backslash G$ will be denoted by $\mu_{\Gamma \backslash G}$ (respectively $\mu_{\Gamma \backslash G}^0$). These projection measures are well-defined thanks to the Γ -invariance of the Haar measure μ and to Lemma 2.1. Our choice of normalisation in (2.10) and (2.11) yields $\mu_{\Gamma \backslash G}(\mathcal{F}_\Gamma) = \mu_{\Gamma \backslash G}^0(\mathcal{F}_\Gamma) = 1$ and therefore $\mu_{\Gamma \backslash G}$ and $\mu_{\Gamma \backslash G}^0$ are both probability measures on $\Gamma \backslash G$.

Remark 2.2. It is not hard to numerically sample points $(x, y, \phi; (\xi_1)) \in \mathcal{F}_\Gamma$ according to $\mu_{\Gamma \setminus G}^0$ or $\mu_{\Gamma \setminus G}$, assuming that we have a generator of random numbers uniformly distributed on $[0, 1]$, such as `RandomReal[]` in the Mathematica[©] software. The only nontrivial step is to generate pairs of real numbers $(x, y) \in \mathcal{F}_{\text{SL}(2, \mathbb{Z})}$ distributed according to the joint density $\frac{3}{\pi y^2} \mathbf{1}_{\mathcal{F}_{\text{SL}(2, \mathbb{Z})}}(x, y)$. The marginal density of $x \in [-\frac{1}{2}, \frac{1}{2}]$ is $\int_{\sqrt{1-x^2}}^{\infty} \frac{3}{\pi y^2} dy = \frac{3}{\pi \sqrt{1-x^2}}$ and hence its distribution function is $[-\frac{1}{2}, \frac{1}{2}] \ni t \mapsto \frac{1}{2} + \frac{3 \arcsin(t)}{\pi} \in [0, 1]$. If x_0 is uniformly distributed on $[0, 1]$, then evaluating the inverse distribution function $[0, 1] \ni s \mapsto \sin(\frac{\pi s}{3} - \frac{\pi}{6}) \in [-\frac{1}{2}, \frac{1}{2}]$ at $s = x_0$ yields a random number distributed according to the marginal density of x . Since the conditional density of y given x is $\frac{3}{\pi y^2} \mathbf{1}_{\mathcal{F}_{\text{SL}(2, \mathbb{Z})}}(x, y) / \frac{3}{\pi \sqrt{1-x^2}} = \frac{\sqrt{1-x^2}}{y^2} \mathbf{1}_{\mathcal{F}_{\text{SL}(2, \mathbb{Z})}}(x, y)$, we find the conditional distribution function of y given x as $[\sqrt{1-x^2}, \infty) \ni t \mapsto 1 - \frac{\sqrt{1-x^2}}{t} \in [0, 1]$. If y_0 is uniformly distributed on $[0, 1]$, then evaluating the inverse conditional distribution function $[0, 1] \ni s \mapsto \frac{\sqrt{1-x^2}}{1-s} \in [\sqrt{1-x^2}, \infty)$ at $s = y_0$ yields a random number distributed according to the conditional density of y given x . In summary, if $(x_0, y_0) = (\text{RandomReal}[], \text{RandomReal}[])$ is a pair of independent uniformly distributed numbers on $[0, 1]$ and we define $x = \sin(\frac{\pi x_0}{3} - \frac{\pi}{6})$ and $y = \frac{\sqrt{1-x^2}}{1-y_0}$, then (x, y) is distributed according to the joint density $\frac{3}{\pi y^2} \mathbf{1}_{\mathcal{F}_{\text{SL}(2, \mathbb{Z})}}(x, y)$, see Figure 3. Generating the remaining




Figure 3: A sample of 5×10^4 points $(x, y) \in \mathcal{F}_{\text{SL}(2, \mathbb{Z})}$ according to the probability density $\frac{3}{\pi y^2} \mathbf{1}_{\mathcal{F}_{\text{SL}(2, \mathbb{Z})}}(x, y)$ as described in Remark 2.2. About 32% of the points sampled lie above the line $y = 3$ and are not shown.

three variables (ϕ, ξ_1, ξ_2) , independently from (x, y) , is easy. In the rational case, if $(\phi_0, \xi_0) = (\text{RandomReal}[], \text{RandomReal}[])$, then we take $\phi = \pi \phi_0$ and we define $(\xi_1, \xi_2) =$

$(0, 0)$ if $\xi_0 \in [0, \frac{1}{3})$, $(\xi_1, \xi_2) = (\frac{1}{2}, 0)$ if $\xi_0 \in [\frac{1}{3}, \frac{2}{3})$, and $(\xi_1, \xi_2) = (0, \frac{1}{2})$ if $\xi_0 \in [\frac{2}{3}, 1]$. In the irrational case, we take $(\phi, \xi_1, \xi_2) = (\pi \text{RandomReal}[], \text{RandomReal}[], \text{RandomReal}[])$.

2.6 Geodesic and horocycle flows

The group G naturally acts on itself by multiplication. Let us consider the action by right-multiplication by elements of the 1-parameter groups $\{\Phi^t : t \in \mathbb{R}\}$ (the *geodesic flow*) and $\{\Psi^x : x \in \mathbb{R}\}$ (the *horocycle flow*), where

$$\Phi^t = \left(a_{e^{-t}}; \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right) = \left(\begin{pmatrix} e^{-t/2} & 0 \\ 0 & e^{t/2} \end{pmatrix}; \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right), \quad (2.42)$$

$$\Psi^x = \left(n_x; \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right) = \left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}; \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right). \quad (2.43)$$

We will also consider the right-multiplication by (2.42) and (2.43) as flows on $\Gamma \backslash G$. In Section 5 we will also use the flows $\tilde{\Phi}^t = (\tilde{a}_{e^{-t}}; \begin{pmatrix} 0 \\ 0 \end{pmatrix}, 0) = \left(\left[\begin{pmatrix} e^{-t/2} & 0 \\ 0 & e^{t/2} \end{pmatrix}, 0 \right]; \begin{pmatrix} 0 \\ 0 \end{pmatrix}, 0 \right)$ and $\tilde{\Psi}^t = (\tilde{n}_x; \begin{pmatrix} 0 \\ 0 \end{pmatrix}, 0) = \left(\left[\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}, 0 \right]; \begin{pmatrix} 0 \\ 0 \end{pmatrix}, 0 \right)$ on \tilde{G} and on $\tilde{\Gamma} \backslash \tilde{G}$.

3 Limit Theorems

In this section we discuss some fundamental dynamical results that will allow us to take $N \rightarrow \infty$ in our main theorems. Such limit theorems rely on the equidistribution of horocycle lifts in the homogeneous space $\Gamma \backslash G$ under the action of the geodesic flow.

3.1 Equidistribution in the smooth rational case

Theorem 3.1 (Limit Theorem, rational case). *Let $F : \Gamma \backslash G \rightarrow \mathbb{R}$ be a bounded continuous function, and λ a Borel probability measure on \mathbb{R} absolutely continuous with respect to the Lebesgue measure. Then*

$$\lim_{t \rightarrow \infty} \int_{\mathbb{R}} F(\Gamma(I; \mathbf{0}) \Psi^u \Phi^t) d\lambda(u) = \int_{\Gamma \backslash G} F d\mu_{\Gamma \backslash G}^{\mathbf{0}},$$

where $\mu_{\Gamma \backslash G}^{\mathbf{0}}$ is the probability measure defined in (2.11).

Proof. Consider the action of Γ on $\mathbb{T}^2 = \mathbb{R}^2 / \mathbb{Z}^2$ via affine transformations, i.e. $(M, \mathbf{v})(\boldsymbol{\xi} + \mathbb{Z}^2) = (\mathbf{v} + M\boldsymbol{\xi} + \mathbb{Z}^2)$, and let $\Sigma = \text{Stab}_{\Gamma}(\mathbf{0} + \mathbb{Z}^2)$. Using the generators of Γ it is not hard to see that $\Sigma = \Gamma_{\theta} \ltimes \mathbb{Z}^2$, where $\Gamma_{\theta} = \langle \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix} \rangle$ is the so-called *theta group*. Note that the orbit of $\mathbf{0} + \mathbb{Z}^2$ under Γ consists of three points in $\mathbb{T}^2 = (-\frac{1}{2}, \frac{1}{2}]^2$, namely $S = \text{Orb}_{\Gamma}(\mathbf{0} + \mathbb{Z}^2) = \{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1/2 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1/2 \end{pmatrix} \} + \mathbb{Z}^2$. By the orbit-stabiliser theorem we have $[\Gamma : \Sigma] = 3$. A fundamental domain \mathcal{F}_{Σ} for the action of Σ on G is therefore given by three copies of (2.36). More explicitly, $\mathcal{F}_{\Sigma} = \mathcal{F}_{\Gamma} \cup \rho_1 \mathcal{F}_{\Gamma} + \rho_2 \mathcal{F}_{\Gamma}$, where $\rho_1 = \gamma_3^{-1} \gamma_2 = \left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}; \begin{pmatrix} -1/2 \\ 0 \end{pmatrix} \right)$ and $\rho_2 = \gamma_2 \gamma_1 = \left(\begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}; \begin{pmatrix} 1/2 \\ 0 \end{pmatrix} \right)$. Let $A \subset \text{SL}(2, \mathbb{Z}) \backslash \text{SL}(2, \mathbb{R}) = \mathcal{F}_{\text{SL}(2, \mathbb{Z})} \times [0, \pi)$. We have

- (i) $(z, \phi; \xi) \in A \times \left\{ \begin{pmatrix} 1/2 \\ 0 \end{pmatrix} \right\}$ if and only if $\rho_1(z, \phi; \xi) \in \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} A \times \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$ and
- (ii) $(z, \phi; \xi) \in A \times \left\{ \begin{pmatrix} 0 \\ 1/2 \end{pmatrix} \right\}$ if and only if $\rho_2(z, \phi; \xi) \in \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix} A \times \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}$.

If A is a set whose boundary has zero $\mu_{\text{SL}(2, \mathbb{Z}) \backslash \text{SL}(2, \mathbb{R})}$ -measure, by Sarnak's theorem on equidistribution of long horocycles [25], we obtain

$$\lim_{t \rightarrow \infty} \int_{\mathbb{R}} \mathbf{1}_{A \times \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right\}} (\Gamma(I; \mathbf{0}) \Psi^x \Phi^t) d\lambda(x) = \lim_{y \rightarrow 0} \int_{\mathbb{R}} \mathbf{1}_A (\Gamma_{\theta} n_x a_y) d\lambda(x) = \quad (3.1)$$

$$= \mu_{\Gamma_{\theta} \backslash \text{SL}(2, \mathbb{R})}(A) = \frac{1}{3} \mu_{\text{SL}(2, \mathbb{Z}) \backslash \text{SL}(2, \mathbb{R})}(A) \quad (3.2)$$

Similarly, using the fact that Möbius transformations are isometries on quotients of $\text{SL}(2, \mathbb{R})$,

$$\lim_{t \rightarrow \infty} \int_{\mathbb{R}} \mathbf{1}_{A \times \left\{ \begin{pmatrix} 1/2 \\ 0 \end{pmatrix} \right\}} (\Gamma(I; \mathbf{0}) \Psi^x \Phi^t) d\lambda(x) = \lim_{y \rightarrow 0} \int_{\mathbb{R}} \mathbf{1}_{\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} A} (\Gamma_{\theta} n_x a_y) d\lambda(x) = \quad (3.3)$$

$$= \mu_{\Gamma_{\theta} \backslash \text{SL}(2, \mathbb{R})} \left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} A \right) = \frac{1}{3} \mu_{\text{SL}(2, \mathbb{Z}) \backslash \text{SL}(2, \mathbb{R})}(A), \quad (3.4)$$

$$\lim_{t \rightarrow \infty} \int_{\mathbb{R}} \mathbf{1}_{A \times \left\{ \begin{pmatrix} 0 \\ 1/2 \end{pmatrix} \right\}} (\Gamma(I; \mathbf{0}) \Psi^x \Phi^t) d\lambda(x) = \lim_{y \rightarrow 0} \int_{\mathbb{R}} \mathbf{1}_{\begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix} A} (\Gamma_{\theta} n_x a_y) d\lambda(x) = \quad (3.5)$$

$$= \mu_{\Gamma_{\theta} \backslash \text{SL}(2, \mathbb{R})} \left(\begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix} A \right) = \frac{1}{3} \mu_{\text{SL}(2, \mathbb{Z}) \backslash \text{SL}(2, \mathbb{R})}(A). \quad (3.6)$$

The limits (3.2)-(3.4)-(3.6), along with a standard approximation argument, imply that the limiting distribution, as $t \rightarrow \infty$, of $\Gamma(I; \mathbf{0}) \Psi^x \Phi^t$ (where $x \in \mathbb{R}$ is randomly distributed according to λ) is given by the product probability measure $\mu_{\text{SL}(2, \mathbb{Z}) \backslash \text{SL}(2, \mathbb{R})} \times \frac{1}{3} \left(\delta_{\begin{pmatrix} 0 \\ 0 \end{pmatrix}} + \delta_{\begin{pmatrix} 1/2 \\ 0 \end{pmatrix}} + \delta_{\begin{pmatrix} 0 \\ 1/2 \end{pmatrix}} \right) = \mu_{\Gamma \backslash G}^{\mathbf{0}}$. \square

Now we consider $\mathcal{A} \subset \mathbb{C}$ and aim to replace F in the previous theorem by the indicator $\mathbf{1}_{(\Theta_{f_1} \overline{\Theta_{f_2}})^{-1}(\mathcal{A})}$, which is bounded but not continuous. This is done in the following

Theorem 3.2. *Assume λ is any Borel probability measure on \mathbb{R} absolutely continuous with respect to Lebesgue measure. Then, for any $f_1, f_2 \in \mathcal{S}_{\eta}(\mathbb{R})$ with $\eta > 1$, we have*

$$\lim_{t \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \Theta_{f_1} \overline{\Theta_{f_2}} (\Gamma(I; \mathbf{0}) \Psi^u \Phi^t) \in \mathcal{A} \right\} = \mu_{\Gamma \backslash G}^{\mathbf{0}} ((\Theta_{f_1} \overline{\Theta_{f_2}})^{-1}(\mathcal{A})) \quad (3.7)$$

whenever $\mathcal{A} \subset \mathbb{C}$ is a measurable set whose boundary is a null set with respect to the push-forward measure $(\Theta_{f_1} \overline{\Theta_{f_2}})_* \mu_{\Gamma \backslash G}^{\mathbf{0}}$.

Proof. The regularity assumption on f_1, f_2 implies that $\Theta_{f_1} \overline{\Theta_{f_2}} : \Gamma \backslash G \rightarrow \mathbb{C}$ defined via (2.29) is a continuous function. Theorem (3.1) implies that for every bounded continuous $B : \mathbb{C} \rightarrow \mathbb{R}$, we have

$$\lim_{t \rightarrow \infty} \int_{\mathbb{R}} B(\Theta_{f_1} \overline{\Theta_{f_2}} (\Gamma(I; \mathbf{0}) \Psi^u \Phi^t)) d\lambda(u) = \int_{\Gamma \backslash G} B \circ (\Theta_{f_1} \overline{\Theta_{f_2}}) d\mu_{\Gamma \backslash G}^{\mathbf{0}}. \quad (3.8)$$

The Portmanteau theorem from probability theory (see, e.g., Theorem 11.1.1 in [12]) allows us to replace B in (3.8) by $\mathbf{1}_{\mathcal{A}}$, yielding (3.7). \square

3.2 Equidistribution in the smooth irrational case

While Theorem 3.1 is used in the special rational case $\alpha = c = 0$, the following theorem addresses the irrational case $(\alpha, c) \notin \mathbb{Q}^2$.

Theorem 3.3 (Limit Theorem, irrational case). *Assume λ is any Borel probability measure on \mathbb{R} absolutely continuous with respect to Lebesgue measure. Then for any $(\alpha, c) \notin \mathbb{Q}^2$ and any $f_1, f_2 \in \mathcal{S}_\eta(\mathbb{R})$ with $\eta > 1$, we have*

$$\lim_{t \rightarrow \infty} \lambda \{u \in \mathbb{R} : \Theta_{f_1} \overline{\Theta_{f_2}} (\Gamma(I; (\alpha + cu)_0) \Psi^u \Phi^t) \in \mathcal{A}\} = \mu_{\Gamma \setminus G}((\Theta_{f_1} \overline{\Theta_{f_2}})^{-1}(\mathcal{A})). \quad (3.9)$$

whenever $\mathcal{A} \subset \mathbb{C}$ is a measurable set whose boundary has measure zero with respect to the push-forward measure $(\Theta_{f_1} \overline{\Theta_{f_2}})_* \mu_{\Gamma \setminus G}$.

Proof. Because of the regularity assumption on f_1 and f_2 , this theorem follows immediately from Corollary 4.3 in [6]. \square

3.3 Limit theorems for indicators

We now extend the limit Theorems 3.2 and 3.3 to the case when f_1 and f_2 are indicator functions (which do not belong in \mathcal{S}_η for $\eta > 1$). Here we write $\chi = \mathbf{1}_{(0,1)}$ and $\chi_b = \mathbf{1}_{(0,b)}$. Although $\frac{1}{N} S_N(x; c, \alpha) \overline{S_{[bN]}(x; c, \alpha)} = \Theta_{f_1} \overline{\Theta_{f_2}}(x + iN^{-2}, 0; (\alpha + cx)_0, 0)$ when $f_1 = \mathbf{1}_{(0,1]}$ and $f_2 = \mathbf{1}_{(0,b]}$, here we use indicators of open intervals. The arguments used to prove Theorems 3.4-3.5 below will work with either kind of indicators. We will see in Section 5 how indicators of open intervals are a natural choice. However, replacing one kind of indicator by another will have no effect in the limit, see Corollaries 3.11 and 3.12. We also note that in this case the function $\Theta_\chi \overline{\Theta_{\chi_b}}$ is not continuous and in fact is only defined $\mu_{\Gamma \setminus G}$ -almost everywhere (as well as $\mu_{\Gamma \setminus G}^0$ -almost everywhere as we will see in Section 5.2) on $\Gamma \setminus G$. It follows from Corollary 2.4 in [6] that $\Theta_f \in L^4(\Gamma \setminus G, \mu_{\Gamma \setminus G})$ for every $f \in L^2(\mathbb{R})$ and hence $\Theta_\chi \overline{\Theta_{\chi_b}}$ is a well defined element of $L^2(\Gamma \setminus G, \mu_{\Gamma \setminus G})$.

Theorem 3.4 (Limit theorem, rational case, for indicators). *Assume λ is any Borel probability measure on \mathbb{R} absolutely continuous with respect to Lebesgue measure. Then*

$$\lim_{t \rightarrow \infty} \lambda \{u \in \mathbb{R} : \Theta_\chi \overline{\Theta_{\chi_b}} (\Gamma(I; \mathbf{0}) \Psi^u \Phi^t) \in \mathcal{A}\} = \mu_{\Gamma \setminus G}^0((\Theta_\chi \overline{\Theta_{\chi_b}})^{-1}(\mathcal{A})). \quad (3.10)$$

whenever $\mathcal{A} \subset \mathbb{C}$ is a measurable set whose boundary has measure zero with respect to the push-forward measure $(\Theta_\chi \overline{\Theta_{\chi_b}})_* \mu_{\Gamma \setminus G}^0$.

Theorem 3.5 (Limit theorem, irrational case, for indicators). *Assume λ is any Borel probability measure on \mathbb{R} absolutely continuous with respect to Lebesgue measure. Then for any $(\alpha, c) \in \mathbb{R}^2 \setminus \mathbb{Q}^2$*

$$\lim_{t \rightarrow \infty} \lambda \{u \in \mathbb{R} : \Theta_\chi \overline{\Theta_{\chi_b}} (\Gamma(I; (\alpha + cu)_0) \Psi^u \Phi^t) \in \mathcal{A}\} = \mu_{\Gamma \setminus G}((\Theta_\chi \overline{\Theta_{\chi_b}})^{-1}(\mathcal{A})). \quad (3.11)$$

whenever $\mathcal{A} \subset \mathbb{C}$ is a measurable set whose boundary has measure zero with respect to the push-forward measure $(\Theta_\chi \overline{\Theta_{\chi_b}})_* \mu_{\Gamma \setminus G}$.

The approximation arguments needed to prove Theorems 3.4–3.5 are given in Lemmata 3.6–3.9 below, which are similar to Lemmata 4.6–4.9 in [6].

Lemma 3.6. *Let f_1, f_2, h be compactly supported, real-valued, Riemann-integrable functions, and assume $h \geq 0$. Then for all $\alpha, c \in \mathbb{R}$, we have*

$$\limsup_{t \rightarrow \infty} \int_{\mathbb{R}} |\Theta_{f_1} \overline{\Theta_{f_2}}((I, (\alpha + cu)) \Psi^u \Phi^t)| h(u) du \leq 2 \|f_1\|_2 \|f_2\|_2 \|h\|_1. \quad (3.12)$$

Proof. Using the Cauchy-Schwarz inequality and Lemma 4.6 in [6], we obtain

$$\left| \limsup_{t \rightarrow \infty} \int_{\mathbb{R}} |\Theta_{f_1} \overline{\Theta_{f_2}}((I, (\alpha + cu)) \Psi^u \Phi^t)| h(u) du \right|^2 \leq \quad (3.13)$$

$$\leq \limsup_{t \rightarrow \infty} \int_{\mathbb{R}} |\Theta_{f_1}((I, (\alpha + cu)) \Psi^u \Phi^t)|^2 h(u) du \int_{\mathbb{R}} |\Theta_{f_2}((I, (\alpha + cu)) \Psi^u \Phi^t)|^2 h(u) du \leq \quad (3.14)$$

$$\leq 4 \|f_1\|_2^2 \|f_2\|_2^2 \|h\|_1^2. \quad (3.15)$$

□

Lemma 3.7. *Let f_1, f_2 be compactly supported, real-valued, Riemann-integrable functions on \mathbb{R} , and let λ be a Borel probability measure on \mathbb{R} absolutely continuous with respect to Lebesgue measure. Then for all $\alpha, c \in \mathbb{R}$ and every $K > 0$ we have*

$$\limsup_{t \rightarrow \infty} \lambda(\{u \in \mathbb{R} : |\Theta_{f_1} \overline{\Theta_{f_2}}((I, (\alpha + cu)) \Psi^u \Phi^t)| > K\}) \leq \frac{2 \|f_1\|_2 \|f_2\|_2}{K}. \quad (3.16)$$

Proof. As in the proof of Lemma 4.7 in [6], let $\lambda' \in L^1(\mathbb{R})$ be the density of λ and let m_h be the positive measure whose density is $h \in C_c(\mathbb{R})$. Markov inequality and Lemma 3.6 yield

$$\lambda(\{u \in \mathbb{R} : |\Theta_{f_1} \overline{\Theta_{f_2}}((I, (\alpha + cu)) \Psi^u \Phi^t)| > K\}) \leq \quad (3.17)$$

$$\leq m_h(\{u \in \mathbb{R} : |\Theta_{f_1} \overline{\Theta_{f_2}}((I, (\alpha + cu)) \Psi^u \Phi^t)| > K\}) + \|\lambda' - h\|_1 \leq \quad (3.18)$$

$$\leq \frac{2 \|f_1\|_2 \|f_2\|_2 \|h\|_1}{K} + \|\lambda' - h\|_1. \quad (3.19)$$

Finally, using the density of $C_c(\mathbb{R})$ in $L^1(\mathbb{R})$ and the continuity of the L^1 -norm, (3.19) implies (3.16) □

Lemma 3.8. *Let f_1, f_2 be compactly supported, real-valued, Riemann-integrable functions on \mathbb{R} , and let λ be a Borel probability measure on \mathbb{R} absolutely continuous with respect to Lebesgue measure. Then for all $\alpha, c \in \mathbb{R}$ and every $\varepsilon > 0$ there exists $K_\varepsilon > 0$ such that*

$$\limsup_{t \rightarrow \infty} \lambda(\{u \in \mathbb{R} : |\Theta_{f_1} \overline{\Theta_{f_2}}((I, (\alpha + cu)) \Psi^u \Phi^t)| > K_\varepsilon\}) \leq \varepsilon \|f_1\|_2 \|f_2\|_2. \quad (3.20)$$

Proof. The statement follows from Lemma 3.7 by taking $K_\varepsilon = 2\varepsilon^{-1}$. \square

Lemma 3.9. *Let f_1, f_2 be compactly supported, real-valued, Riemann-integrable functions on \mathbb{R} , and let λ be a Borel probability measure on \mathbb{R} absolutely continuous with respect to Lebesgue measure. Let $\eta > 1$. Then for all $\alpha, c \in \mathbb{R}$ and every $\varepsilon, \delta > 0$ there exist compactly supported functions $\tilde{f}_1, \tilde{f}_2 \in \mathcal{S}_\eta(\mathbb{R})$ such that*

$$\limsup_{t \rightarrow \infty} \lambda(\{u \in \mathbb{R} : |\Theta_{f_1} \overline{\Theta_{f_2}}((I; (\alpha + cu)_0^+) \Psi^u \Phi^t) - \Theta_{\tilde{f}_1} \overline{\Theta_{\tilde{f}_2}}((I; (\alpha + cu)_0^+) \Psi^u \Phi^t)| > \delta\}) < \varepsilon. \quad (3.21)$$

Proof. Fix $\varepsilon, \delta > 0$. Since f_1, f_2 are compactly supported and $\mathcal{S}_\eta(\mathbb{R})$ is dense in $L^2(\mathbb{R})$, we can find compactly supported $\tilde{f}_1, \tilde{f}_2 \in \mathcal{S}_\eta(\mathbb{R})$ such that

$$\frac{4}{\delta} \left(\|f_1\|_2 \|f_2 - \tilde{f}_2\|_2 + \|f_1 - \tilde{f}_1\|_2 \|\tilde{f}_2\|_2 \right) \leq \varepsilon. \quad (3.22)$$

Note that $f \mapsto \Theta_f$ is linear and hence we can write

$$\Theta_{f_1} \overline{\Theta_{f_2}} - \Theta_{\tilde{f}_1} \overline{\Theta_{\tilde{f}_2}} = \Theta_{f_1} \overline{\Theta_{f_2 - \tilde{f}_2}} + \Theta_{f_1 - \tilde{f}_1} \overline{\Theta_{\tilde{f}_2}}. \quad (3.23)$$

Using (3.23) we obtain the union bound

$$\lambda(\{u \in \mathbb{R} : |\Theta_{f_1} \overline{\Theta_{f_2}}((I; (\alpha + cu)_0^+) \Psi^u \Phi^t) - \Theta_{\tilde{f}_1} \overline{\Theta_{\tilde{f}_2}}((I; (\alpha + cu)_0^+) \Psi^u \Phi^t)| > \delta\}) \leq \quad (3.24)$$

$$\leq \lambda(\{u \in \mathbb{R} : |\Theta_{f_1} \overline{\Theta_{f_2 - \tilde{f}_2}}((I; (\alpha + cu)_0^+) \Psi^u \Phi^t)| > \frac{\delta}{2}\}) + \quad (3.25)$$

$$+ \lambda(\{u \in \mathbb{R} : |\Theta_{f_1 - \tilde{f}_1} \overline{\Theta_{\tilde{f}_2}}((I; (\alpha + cu)_0^+) \Psi^u \Phi^t)| > \frac{\delta}{2}\}). \quad (3.26)$$

Now we apply Lemma 3.7 three times with $K = \frac{\delta}{2}$ to the limsup of the expressions in (3.25)-(3.26), thus obtaining (3.22) as an upper bound. \square

We can now proceed to the proof of Theorems 3.4 and 3.5. Since the Lemmata 3.6–3.9 above hold regardless of the values of α and c , we combine the proofs into one.

Proof of Theorems 3.4 and 3.5. We follow the strategy of the proof of Theorem 4.4 in [6]. Note that (3.10)–(3.11) are equivalent to the following statement: for every bounded continuous function $F : \mathbb{C} \rightarrow \mathbb{R}$, we have

$$\lim_{t \rightarrow \infty} \int_{\mathbb{R}} F(\Theta_\chi \overline{\Theta_{\chi_b}}(\Gamma(I; (\alpha + cu)_0^+) \Psi^u \Phi^t)) d\lambda(u) = \begin{cases} \int_{\Gamma \setminus G} F \circ (\Theta_\chi \overline{\Theta_{\chi_b}}) d\mu_{\Gamma \setminus G}^0 & \text{if } \alpha = c = 0, \\ \int_{\Gamma \setminus G} F \circ (\Theta_\chi \overline{\Theta_{\chi_b}}) d\mu_{\Gamma \setminus G} & \text{if } (\alpha, c) \notin \mathbb{Q}^2. \end{cases} \quad (3.27)$$

Lemma 3.8 and the Helly-Prokhorov theorem imply that for every sequence $(t_j)_{j \geq 1}$ such that $t_j \rightarrow \infty$ as $j \rightarrow \infty$ there exists subsequence $(t_{j_\ell})_{\ell \geq 1}$ and a probability measure ρ on \mathbb{C} such that for every bounded continuous function $F : \mathbb{C} \rightarrow \mathbb{R}$ we have

$$\lim_{\ell \rightarrow \infty} \int_{\mathbb{R}} F(\Theta_\chi \overline{\Theta_{\chi_b}} (\Gamma(I; (\alpha + cu)_0) \Psi^u \Phi^{t_{j_\ell}})) d\lambda(u) = \int_{\mathbb{C}} F d\rho. \quad (3.28)$$

We want to make sure that the limit along the original sequence

$$I(\chi, \chi_b) = \lim_{j \rightarrow \infty} \int_{\mathbb{R}} F(\Theta_\chi \overline{\Theta_{\chi_b}} (\Gamma(I; (\alpha + cu)_0) \Psi^u \Phi^{t_j})) d\lambda(u) \quad (3.29)$$

exists and does not depend on the sequence $(t_j)_{j \geq 1}$. To this end, suppose that $F \in C_c^\infty(\mathbb{C})$. By Lemma 3.9, for every $\varepsilon, \delta > 0$, there exist compactly supported functions $f_1, f_2 \in \mathcal{S}_\eta$, $\eta > 1$, such that

$$\int_{\mathbb{R}} |F(\Theta_\chi \overline{\Theta_{\chi_b}} (\Gamma(I; (\alpha + cu)_0) \Psi^u \Phi^{t_j})) - F(\Theta_{f_1} \overline{\Theta_{f_2}} (\Gamma(I; (\alpha + cu)_0) \Psi^u \Phi^{t_j}))| d\lambda(u) \leq \quad (3.30)$$

$$\leq L \int_{\mathbb{R}} |\Theta_\chi \overline{\Theta_{\chi_b}} (\Gamma(I; (\alpha + cu)_0) \Psi^u \Phi^{t_j}) - \Theta_{f_1} \overline{\Theta_{f_2}} (\Gamma(I; (\alpha + cu)_0) \Psi^u \Phi^{t_j})| d\lambda(u) \leq \quad (3.31)$$

$$\leq L(\varepsilon + \delta), \quad (3.32)$$

where L is the Lipschitz constant of F . Observe that the limit

$$I(f_1, f_2) = \lim_{j \rightarrow \infty} \int_{\mathbb{R}} F(\Theta_{f_1} \overline{\Theta_{f_2}} (\Gamma(I; (\alpha + cu)_0) \Psi^u \Phi^{t_j})) d\lambda(u) \quad (3.33)$$

exists by either Theorem 3.2 if $\alpha = c = 0$ or Theorem 3.3 if $(\alpha, c) \notin \mathbb{Q}^2$. This fact, together with the bound (3.32), imply that the sequence

$$\left(\int_{\mathbb{R}} F(\Theta_\chi \overline{\Theta_{\chi_b}} (\Gamma(I; (\alpha + cu)_0) \Psi^u \Phi^{t_j})) d\lambda(u) \right)_{j \geq 1} \quad (3.34)$$

is Cauchy and hence the limit (3.29) exists. Note that, as $f_1 \rightarrow \chi$ and $f_2 \rightarrow \chi_b$ in $L^2(\mathbb{R})$ we have from (3.32) that $I(f_1, f_2) \rightarrow I(\chi, \chi_b)$. Therefore

$$I(\chi, \chi_b) = \int_{\mathbb{C}} F d\rho = \begin{cases} \int_{\Gamma \setminus G} F \circ (\Theta_\chi \overline{\Theta_{\chi_b}}) d\mu_{\Gamma \setminus G}^0 & \text{if } \alpha = c = 0, \\ \int_{\Gamma \setminus G} F \circ (\Theta_\chi \overline{\Theta_{\chi_b}}) d\mu_{\Gamma \setminus G} & \text{if } (\alpha, c) \notin \mathbb{Q}^2. \end{cases} \quad (3.35)$$

We have shown that, given α and c , the limiting measure ρ is the same for every sequence $(t_j)_{j \geq 1}$ tending to infinity, and this implies that (3.27) holds. \square

Remark 3.10. In order to numerically sample points in \mathbb{C} whose law is the push forward of $\mu_{\Gamma \setminus G}^0$ or $\mu_{\Gamma \setminus G}$ via $\Theta_\chi \overline{\Theta_{\chi b}}$, it is enough to generate sample points $(x, y, \phi; (\xi_1^1)) \in \mathcal{F}_\Gamma$ as explained in Remark 2.2 and then evaluate the function $\Theta_\chi \overline{\Theta_{\chi b}}$ at those points. The evaluation is numerically challenging since the convergence of the series (2.29) is very delicate when $(f_1, f_2) = (\chi, \chi_b)$. In order to obtain Figure 2 (in which $b = 1$ and hence $|\Theta_\chi|^2 \in \mathbb{R}_{\geq 0}$) we use Mathematica[©] to find a closed form for $\chi_\phi(w)$ (involving special functions **Erf** and **Erfi** that can be evaluated to arbitrary precision by the software) and then approximate $|\Theta_\chi|$ by truncating the sum over $n \in \mathbb{Z}$ in (2.29) to $-100 \leq n \leq 100$.

Corollary 3.11. *Let $\alpha = c = 0$, and $b \geq 1$. Let λ be a probability measure, absolutely continuous with respect to Lebesgue measure. Then for every $b \geq 1$ and $R > 0$ we have that*

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \frac{1}{N} |S_N \overline{S_{[bN]}}(x; 0, 0)| > R^2 \right\} = \mu_{\Gamma \setminus G}^0 \{ |\Theta_\chi \overline{\Theta_{\chi b}}| > R^2 \}. \quad (3.36)$$

Corollary 3.12. *Let $(c, \alpha) \in \mathbb{R}^2 \setminus \mathbb{Q}^2$, and $b \geq 1$. Let λ be a probability measure, absolutely continuous with respect to Lebesgue measure. Then for every $b \geq 1$ and $R > 0$ we have that*

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \frac{1}{N} |S_N \overline{S_{[bN]}}(x; c, \alpha)| > R^2 \right\} = \mu_{\Gamma \setminus G} \{ |\Theta_\chi \overline{\Theta_{\chi b}}| > R^2 \}. \quad (3.37)$$

Proof of Corollaries 3.11 and 3.12. Let $s \in \{1, b\}$. Observe that $S_{[sN]} = S_N^{1(0, s]}$ and $S_N^{\chi s}$ are both finite sums and differ in absolute value by at most 1. Therefore, for any $(c, \alpha) \in \mathbb{R}^2$,

$$\frac{1}{\sqrt{N}} |S_{[sN]}(x; c, \alpha)| = \frac{1}{\sqrt{N}} |S_N^{\chi s}(x; c, \alpha)| + O\left(\frac{1}{\sqrt{N}}\right), \quad (3.38)$$

and so

$$\frac{1}{N} |S_N \overline{S_{[sN]}}(x; c, \alpha)| = \quad (3.39)$$

$$= \frac{1}{N} \left| S_N^\chi \overline{S_N^{\chi s}}(x; c, \alpha) \right| + O\left(\frac{1}{N}\right) + O\left(\frac{1}{\sqrt{N}} \left| \frac{S_N^\chi(x; c, \alpha)}{\sqrt{N}} \right| \right) + O\left(\frac{1}{\sqrt{N}} \left| \frac{S_N^{\chi s}(x; c, \alpha)}{\sqrt{N}} \right| \right). \quad (3.40)$$

Theorems 3.4 and 3.5 and (1.8)-(1.9) imply that $\left| \frac{S_N(x; c, \alpha)}{\sqrt{N}} \right|$ and $\left| \frac{S_N^{\chi s}(x; c, \alpha)}{\sqrt{N}} \right|$ converge in law when $\alpha = c = 0$ and $(c, \alpha) \in \mathbb{R}^2 \setminus \mathbb{Q}^2$. By the Crámer-Slutsky theorem (see [14], Theorem 11.4 therein) we have that the sum of the three O -terms in (3.40) goes to zero in law, as $N \rightarrow \infty$. Therefore, as $|\cdot|$ is continuous, both $\frac{1}{N} |S_N \overline{S_{[sN]}}(x; c, \alpha)|$ and $\frac{1}{N} |S_N^\chi \overline{S_N^{\chi s}}(x; c, \alpha)|$ converge in law to the same limit as $N \rightarrow \infty$. More precisely,

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \frac{1}{N} |S_N \overline{S_{[bN]}}(x; c, \alpha)| \in \mathcal{A} \right\} = \begin{cases} \mu_{\Gamma \setminus G}^0 \{ |\Theta_\chi \overline{\Theta_{\chi b}}| \in \mathcal{A} \} & \text{if } \alpha = c = 0, \\ \mu_{\Gamma \setminus G} \{ |\Theta_\chi \overline{\Theta_{\chi b}}| \in \mathcal{A} \} & \text{if } (c, \alpha) \notin \mathbb{Q}^2 \end{cases} \quad (3.41)$$

for any measurable set $\mathcal{A} \subset \mathbb{R}$ whose boundary is of measure zero with respect to the appropriate limiting measure, namely $|\Theta_\chi \overline{\Theta_{\chi_b}}|_* \mu_{\Gamma \setminus G}^0$ or $|\Theta_\chi \overline{\Theta_{\chi_b}}|_* \mu_{\Gamma \setminus G}$ as the case may be. Taking $\mathcal{A} = (R, \infty)$ gives the result. \square

4 Growth in the cusp and tail bounds for $\Theta_{f_2} \overline{\Theta_{f_2}}$

In this section we focus on the size of $\Theta_{f_1} \overline{\Theta_{f_2}}$ when f_1 and f_2 are of sufficient regularity. The following lemma, which extends Lemma 2.1 in [6], is crucial and will be used later. Recall the definition of $\kappa_\eta(f)$ (2.26).

Lemma 4.1. *Given $\xi_2 \in \mathbb{R}$, write $\xi_2 = k + \theta$, with $k \in \mathbb{Z}$, and $\theta \in (-\frac{1}{2}, \frac{1}{2}]$. Let $\eta > 1$, then, there exist constants $d > 0$ and $C_\eta = d^\eta \zeta(\eta)^2$ such that for every $f_1, f_2 \in \mathcal{S}_\eta(\mathbb{R})$ we have*

$$\left| \Theta_{f_1} \overline{\Theta_{f_2}}(x + iy, \phi, \xi) - y^{\frac{1}{2}} (f_1)_\phi(-\theta y^{\frac{1}{2}}) \overline{(f_2)_\phi(-\theta y^{\frac{1}{2}})} \right| \leq C_\eta \kappa_\eta(f_1) \kappa_\eta(f_2) y^{-\frac{(\eta-1)}{2}} \quad (4.1)$$

for every $y \geq \frac{1}{2}$, and every x, ϕ, ξ . We may take $d = 2^6$.

Proof. The regularity assumption on f_1 and f_2 ensures that the series defining Θ_{f_1} and Θ_{f_2} are absolutely convergent. Therefore we may write

$$\begin{aligned} \Theta_{f_1} \overline{\Theta_{f_2}}(x + iy, \phi, \xi) &= y^{\frac{1}{2}} \sum_{n, m \in \mathbb{Z}} (f_1)_\phi((n - \xi_2)y^{\frac{1}{2}}) \overline{(f_2)_\phi((m - \xi_2)y^{\frac{1}{2}})} \\ &\quad e\left(\frac{1}{2}(n - \xi_2)^2 x + n\xi_1 - \frac{1}{2}(m - \xi_2)^2 x - m\xi_1\right). \end{aligned} \quad (4.2)$$

Since the term $y^{\frac{1}{2}} (f_1)_\phi(-\theta y^{\frac{1}{2}}) \overline{(f_2)_\phi(-\theta y^{\frac{1}{2}})}$ in (4.1) corresponds to the term $n = m = k$, it is enough to prove that the above sum restricted to $(n, m) \in \mathbb{Z}^2 \setminus \{(k, k)\}$ is bounded in absolute value by $C_\eta \kappa_\eta(f_1) \kappa_\eta(f_2) y^{-\frac{(\eta-1)}{2}}$. Indeed, by (2.26),

$$y^{\frac{1}{2}} \sum_{(n, m) \in \mathbb{Z}^2 \setminus \{(k, k)\}} \left| (f_1)_\phi((n - \xi_2)y^{\frac{1}{2}}) \overline{(f_2)_\phi((m - \xi_2)y^{\frac{1}{2}})} \right| \leq \quad (4.3)$$

$$\leq y^{\frac{1}{2}} \sum_{(n, m) \in \mathbb{Z}^2 \setminus \{(k, k)\}} \frac{\kappa_\eta(f_1)}{(1 + (n - \xi_2)^2 y)^{\frac{\eta}{2}}} \frac{\kappa_\eta(f_2)}{(1 + (m - \xi_2)^2 y)^{\frac{\eta}{2}}} = \quad (4.4)$$

$$= y^{\frac{1}{2}} \kappa_\eta(f_1) \kappa_\eta(f_2) \left(y^{-\eta} \left(\sum_{n \in \mathbb{Z} \setminus \{k\}} \frac{1}{(y^{-1} + (n - \xi_2)^2)^{\frac{\eta}{2}}} \right)^2 + \right. \quad (4.5)$$

$$\left. + 2y^{-\frac{\eta}{2}} \frac{1}{(1 + (k - \xi_2)^2 y)^{\frac{\eta}{2}}} \sum_{n \in \mathbb{Z} \setminus \{k\}} \frac{1}{(y^{-1} + (n - \xi_2)^2)^{\frac{\eta}{2}}} \right). \quad (4.6)$$

Now note that the sum appearing in both (4.5) and (4.6) is bounded above by

$$\sum_{\ell=1}^{\infty} \frac{1}{(\ell - \theta)^\eta} + \sum_{j=1}^{\infty} \frac{1}{(j + \theta)^\eta}. \quad (4.7)$$

By setting $\theta = \pm \frac{1}{2}$ we can find an upper bound for (4.7) given by $2\zeta(\eta)(2^\eta - 1)$. We have therefore found an upper bound for the absolute value of the sum (4.2) restricted to $(n, m) \in \mathbb{Z}^2 \setminus \{(k, k)\}$ of the form

$$y^{\frac{1}{2}} \kappa_\eta(f_1) \kappa_\eta(f_2) \left(A_\eta y^{-\eta} + B_\eta y^{-\frac{\eta}{2}} \right), \quad (4.8)$$

where $A_\eta = (2\zeta(\eta)(2^\eta - 1))^2$ and $B_\eta = 4\zeta(\eta)(2^\eta - 1)$. Since $y^{-\eta} \leq 2^{\frac{\eta}{2}} y^{-\frac{\eta}{2}}$ for $y \geq \frac{1}{2}$ and $B_\eta \leq 2^{\frac{\eta}{2}} A_\eta$, we obtain (4.1) with $C_\eta \geq 2^{1+\frac{\eta}{2}} A_\eta$ and the result follows. \square

4.1 Tail bound in the smooth case

We now consider the function $\Theta_{f_1} \overline{\Theta_{f_2}} : \Gamma \backslash G \rightarrow \mathbb{C}$ as *two* possible complex-valued random variables by assuming that its argument is distributed either according to $\mu_{\Gamma \backslash G}^0$ or $\mu_{\Gamma \backslash G}$. In both cases, we use Lemma 4.1 to accurately study the tails of these random variables.

4.1.1 Tail bound in the smooth rational case

Lemma 4.2. *Let $\eta > 1$. Then, for every $f_1, f_2 \in \mathcal{S}_\eta(\mathbb{R})$ and $R^2 > 2^\eta C_\eta \kappa_\eta(f_1) \kappa_\eta(f_2)$ we have*

$$\mu_{\Gamma \backslash G}^0(|\Theta_{f_1} \overline{\Theta_{f_2}}| > R^2) = \frac{2D_{\text{rat}}(f_1, f_2)}{\pi^2 R^4} (1 + O_\eta((\kappa_\eta(f_1) \kappa_\eta(f_2))^\eta R^{-2\eta})), \quad (4.9)$$

where C_η is as in Lemma 4.1,

$$D_{\text{rat}}(f_1, f_2) = \int_0^\pi |(f_1)_\phi(0) (f_2)_\phi(0)|^2 d\phi, \quad (4.10)$$

and the implied constant by O_η -notation in (4.9) can be taken to be $3(2^\eta C_\eta)^\eta$.

Proof. Recall the fundamental domain (2.36) for the action of Γ on G , and define

$$\mathcal{F}_T = \{(x + iy, \phi; \xi) \in \mathfrak{H} \times [0, \pi) \times (-\frac{1}{2}, \frac{1}{2}]^2 : x \in [-1/2, 1/2), y \geq T\} \quad (4.11)$$

so that we have the containments $\mathcal{F}_1 \subseteq \mathcal{F}_\Gamma \subseteq \mathcal{F}_{\frac{1}{2}}$. Set $\tilde{\kappa} = C_\eta \kappa_\eta(f_1) \kappa_\eta(f_2)$. We obtain, by Lemma 4.1,

$$\mu_{\Gamma \backslash G}^0 \left\{ (x + iy, \phi; \xi) \in \mathcal{F}_\Gamma : |\Theta_{f_1} \overline{\Theta_{f_2}}(x + iy, \phi; \xi)| > R^2 \right\} \leq \quad (4.12)$$

$$\leq \mu^0 \left\{ (x + iy, \phi; \xi) \in \mathcal{F}_{\frac{1}{2}} : |\Theta_{f_1} \overline{\Theta_{f_2}}(x + iy, \phi; \xi)| > R^2 \right\} \leq \quad (4.13)$$

$$\leq \mu^0 \left\{ (x + iy, \phi; \xi) \in \mathcal{F}_{\frac{1}{2}} : \left| y^{\frac{1}{2}} (f_1)_\phi(-\theta y^{\frac{1}{2}}) (f_2)_\phi(-\theta y^{\frac{1}{2}}) \right| + \tilde{\kappa} y^{-\frac{\eta-1}{2}} > R^2 \right\}. \quad (4.14)$$

Since $f_1, f_2 \in \mathcal{S}_\eta(\mathbb{R})$, we have

$$\left| (f_1)_\phi(-\theta y^{\frac{1}{2}}) (f_2)_\phi(-\theta y^{\frac{1}{2}}) \right| \leq \frac{\kappa_\eta(f_1) \kappa_\eta(f_2)}{\left(1 + \left| -\theta y^{\frac{1}{2}} \right|^2\right)^\eta} \leq \kappa_\eta(f_1) \kappa_\eta(f_2). \quad (4.15)$$

Therefore, the condition

$$\left| y^{\frac{1}{2}} (f_1)_\phi(-\theta y^{\frac{1}{2}}) (f_2)_\phi(-\theta y^{\frac{1}{2}}) \right| + \tilde{\kappa} y^{-\frac{\eta-1}{2}} > R^2 \quad (4.16)$$

in (4.14) implies

$$\tilde{\kappa} \left(y^{\frac{1}{2}} + y^{-\frac{\eta-1}{2}} \right) > R^2. \quad (4.17)$$

Furthermore, since $y \geq \frac{1}{2}$ for all points $(x + iy, \phi; \xi) \in \mathcal{F}_{\frac{1}{2}}$, we have $y^{-\frac{\eta-1}{2}} \leq 2^{\frac{\eta}{2}} y^{\frac{1}{2}}$ and (4.17) in turn implies

$$y > \frac{R^4}{\tilde{\kappa}^{24\eta}}. \quad (4.18)$$

Therefore (4.16) implies

$$\left| y^{\frac{1}{2}} (f_1)_\phi(-\theta y^{\frac{1}{2}}) (f_2)_\phi(-\theta y^{\frac{1}{2}}) \right| + \tilde{\kappa} \left(\frac{R^4}{\tilde{\kappa}^{24\eta}} \right)^{-\frac{\eta-1}{2}} > R^2. \quad (4.19)$$

Let us introduce the shorthand

$$I_T^{\mathbf{0}}(\Lambda) = \mu^{\mathbf{0}} \left\{ (x + iy, \phi; \xi) \in \mathcal{F}_T : \left| y^{\frac{1}{2}} (f_1)_\phi(-\theta y^{\frac{1}{2}}) (f_2)_\phi(-\theta y^{\frac{1}{2}}) \right| > \Lambda \right\}. \quad (4.20)$$

Then, (4.14) and (4.19) yield the upper bound

$$\mu_{\Gamma \setminus G}^{\mathbf{0}}(|\Theta_{f_1} \overline{\Theta_{f_2}}| > R^2) \leq I_{\frac{1}{2}}^{\mathbf{0}} \left(R^2 - \tilde{\kappa} \left(\frac{2^\eta \tilde{\kappa}}{R^2} \right)^{\eta-1} \right). \quad (4.21)$$

Similarly, we obtain the lower bound

$$\mu_{\Gamma \setminus G}^{\mathbf{0}}(|\Theta_{f_1} \overline{\Theta_{f_2}}| > R^2) \geq I_1^{\mathbf{0}} \left(R^2 + \tilde{\kappa} \left(\frac{2^\eta \tilde{\kappa}}{R^2} \right)^{\eta-1} \right). \quad (4.22)$$

Using the fact that $f_1, f_2 \in \mathcal{S}_\eta(\mathbb{R})$, with $\eta > 1$, we see that for fixed $\theta \neq 0$ we have $\lim_{y \rightarrow \infty} y^{\frac{1}{2}} (f_1)_\phi(-\theta y^{\frac{1}{2}}) (f_2)_\phi(-\theta y^{\frac{1}{2}}) = 0$. Thus, in order for the inequality in (4.20) to hold for large Λ , we must have that $|\theta y^{\frac{1}{2}}| = O(1)$ as $y \rightarrow \infty$, i.e. $\theta = O(y^{-\frac{1}{2}})$. In particular, if θ is fixed, then $\theta = 0$ (i.e. $\xi_2 \in \mathbb{Z}$). Recall (2.11) and note that integrating the indicator of the event $\{|y^{\frac{1}{2}} (f_1)_\phi(-\theta y^{\frac{1}{2}}) (f_2)_\phi(-\theta y^{\frac{1}{2}})| > \Lambda\}$ against the measure $\mu^{\mathbf{0}}$ on \mathcal{F}_T forces $\xi \in \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1/2 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1/2 \end{pmatrix} \right\}$. More precisely, $\xi_2 = \theta = 0$ with probability $\frac{2}{3}$ and $\xi_2 = \theta = \frac{1}{2}$

with probability $\frac{1}{3}$. The terms with $\xi_2 = \theta = \frac{1}{2}$ do not contribute for Λ large. Indeed, note that if $\xi_2 = \theta = \frac{1}{2}$, then we have the estimates

$$\left| \frac{y^{\frac{1}{2}}(f_1)_\phi(-\frac{1}{2}y^{\frac{1}{2}})(f_2)_\phi(-\frac{1}{2}y^{\frac{1}{2}})}{\kappa_\eta(f_1)\kappa_\eta(f_2)} \right| \leq \frac{y^{\frac{1}{2}}}{(1+\frac{1}{4}y)^\eta} < 4^\eta y^{\frac{1}{2}-\eta} \leq 2^{3\eta} \quad (4.23)$$

for $y \geq \frac{1}{2}$. Therefore, if we assume

$$\Lambda \geq 2^{3\eta} \kappa_\eta(f_1) \kappa_\eta(f_2), \quad (4.24)$$

then we obtain

$$I_T^0(\Lambda) = \frac{3}{\pi^2} \frac{2}{3} \int_{-\frac{1}{2}}^{\frac{1}{2}} \int_0^\pi \int_{y > \max\{T, |(f_1)_\phi(0)(f_2)_\phi(0)|^{-2}\Lambda^2\}} \frac{1}{y^2} dy d\phi dx. \quad (4.25)$$

Note that (4.24) implies that $\Lambda \geq \kappa_\eta(f_1) \kappa_\eta(f_2)$ and hence $|(f_1)_\phi(0)(f_2)_\phi(0)|^2 \leq \frac{\Lambda^2}{T}$ for $T \in \{\frac{1}{2}, 1\}$. In this case (4.25) can be rewritten as

$$I_T^0(\Lambda) = \frac{2}{\pi^2} \int_{-\frac{1}{2}}^{\frac{1}{2}} \int_0^\pi \int_{|(f_1)_\phi(0)(f_2)_\phi(0)|^{-2}\Lambda^2}^\infty \frac{1}{y^2} dy d\phi dx = \quad (4.26)$$

$$= \frac{2}{\pi^2} \frac{1}{\Lambda^2} \int_0^\pi |(f_1)_\phi(0)(f_2)_\phi(0)|^2 d\phi. \quad (4.27)$$

We now use the inequality (valid for $0 \leq u \leq \frac{1}{2}$)

$$\left| 1 - \frac{1}{(1 \mp u)^2} \right| \leq 6u, \quad (4.28)$$

and hence

$$\frac{1}{(1 \mp u)^2} = 1 + O(u) \quad (4.29)$$

where the implied constant can be taken to be 6. Hence, with

$$\Lambda = R^2 \mp \tilde{\kappa} \left(\frac{2^\eta \tilde{\kappa}}{R^2} \right)^{\eta-1}, \quad (4.30)$$

we obtain

$$\frac{1}{\Lambda^2} = \frac{1}{\left(R^2 \mp \tilde{\kappa} \left(\frac{2^\eta \tilde{\kappa}}{R^2} \right)^{\eta-1} \right)^2} = \frac{1}{R^4} \frac{1}{\left(1 \mp \left(\frac{2^{\eta-1} \tilde{\kappa}}{R^2} \right)^\eta \right)^2} = \frac{1}{R^4} (1 + O(2^{\eta(\eta-1)} \tilde{\kappa}^\eta R^{-2\eta})) \quad (4.31)$$

provided $2^{\eta(\eta-1)}\tilde{\kappa}^\eta R^{-2\eta} < \frac{1}{2}$. This is implied by the inequality

$$R^2 > 2^\eta \tilde{\kappa}. \quad (4.32)$$

Therefore, the assumption (4.32) and the bound (4.31) allow us to write

$$\frac{1}{\Lambda^2} = \frac{1}{R^4} \left(1 + O_\eta((\kappa_\eta(f_1)\kappa_\eta(f_2))^\eta R^{-2\eta}) \right), \quad (4.33)$$

where the constant implied by the O_η -notation can be taken to be $3(2^\eta C_\eta)^\eta$. Finally we notice that (4.32) also guarantees that (4.24) holds. In fact, we obtain

$$\Lambda = R^2 \mp \tilde{\kappa} \left(\frac{2^\eta \tilde{\kappa}}{R^2} \right)^{\eta-1} > \tilde{\kappa} (2^\eta - 1) \geq \tilde{\kappa}. \quad (4.34)$$

Therefore, using the fact that $C_\eta \geq 2^{6\eta}$, we see that (4.32) implies (4.24) and hence in view of (4.27) and (4.33) we see

$$I_T^0(\Lambda) = \frac{2}{\pi^2} D_{\text{rat}}(f_1, f_2) R^{-4} \left(1 + O_\eta((\kappa_\eta(f_1)\kappa_\eta(f_2))^\eta R^{-2\eta}) \right) \quad (4.35)$$

for (T, Λ) either equal to $(\frac{1}{2}, R^2 - \tilde{\kappa}(\frac{2^\eta \tilde{\kappa}}{R^2})^{\eta-1})$ or to $(1, R^2 + \tilde{\kappa}(\frac{2^\eta \tilde{\kappa}}{R^2})^{\eta-1})$. Finally, the bounds (4.21) and (4.22) imply (4.9). \square

Combining Theorem 3.2 and Lemma 4.2, we get the rational case of Theorem 1.7.

4.1.2 Tail bound in the smooth irrational case

Lemma 4.3. *Let $\eta > 1$. Then for every $f_1, f_2 \in \mathcal{S}_\eta(\mathbb{R})$ and $R^2 > 2^\eta C_\eta \kappa_\eta(f_1)\kappa_\eta(f_2)$ we have*

$$\mu_{\Gamma \setminus G}(|\Theta_{f_1} \overline{\Theta_{f_2}}| > R^2) = \frac{2D_{\text{irr}}(f_1, f_2)}{\pi^2 R^6} \left(1 + O_\eta((\kappa_\eta(f_1)\kappa_\eta(f_2))^\eta R^{-2\eta}) \right), \quad (4.36)$$

where

$$D_{\text{irr}}(f_1, f_2) = \int_0^\pi \int_{-\infty}^\infty |(f_1)_\phi(w) (f_2)_\phi(w)|^3 \, d\phi \, dw \quad (4.37)$$

and the constant implied by the O_η -notation in (4.36) can be taken to be $7(2^\eta C_\eta)^\eta$.

Proof. Proceeding as in the proof of Lemma 4.2, if we set

$$I_T(\Lambda) = \mu \left\{ (x + iy, \phi; \boldsymbol{\xi}) \in \mathcal{F}_T : \left| y^{\frac{1}{2}} (f_1)_\phi(-\theta y^{\frac{1}{2}}) (f_2)_\phi(-\theta y^{\frac{1}{2}}) \right| > \Lambda \right\}, \quad (4.38)$$

then we obtain the bounds

$$I_1 \left(R^2 + \tilde{\kappa} \left(\frac{2^\eta \tilde{\kappa}}{R^2} \right)^{\eta-1} \right) \leq \mu_{\Gamma \setminus G}(|\Theta_{f_1} \overline{\Theta_{f_2}}| > R^2) \leq I_{\frac{1}{2}} \left(R^2 - \tilde{\kappa} \left(\frac{2^\eta \tilde{\kappa}}{R^2} \right)^{\eta-1} \right). \quad (4.39)$$

If we assume that

$$\Lambda \geq 2\kappa_\eta(f_1)\kappa_\eta(f_2), \quad (4.40)$$

then $|(f_1)_\phi(-\theta y^{\frac{1}{2}})(f_2)_\phi(-\theta y^{\frac{1}{2}})|^2 \leq \frac{\Lambda^2}{T}$ for $T \in \{\frac{1}{2}, 1\}$ and we obtain

$$I_T(\Lambda) = \frac{3}{\pi^2} \int_{-\frac{1}{2}}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{\frac{1}{2}} \int_0^\pi \iint \frac{1}{y^2} dy d\theta d\phi d\xi_1 dx. \quad (4.41)$$

$$\left\{ -\frac{1}{2} < \theta \leq \frac{1}{2}, y > |(f_1)_\phi(-\theta y^{\frac{1}{2}})(f_2)_\phi(-\theta y^{\frac{1}{2}})|^{-2} \Lambda^2 \right\}$$

Now we use the change of variables $w = -\theta y^{1/2}$ and notice that

$$-\frac{1}{2} < \theta < 0, \quad y > |(f_1)_\phi(-\theta y^{\frac{1}{2}})(f_2)_\phi(-\theta y^{\frac{1}{2}})|^{-2} \Lambda^2 \quad (4.42)$$

$$\iff 0 < w < \infty, \quad -|w| |(f_1)_\phi(w)(f_2)_\phi(w)| \Lambda^{-1} < \theta < 0 \quad (4.43)$$

(since (4.40) implies that $-|w| |(f_1)_\phi(w)(f_2)_\phi(w)| \Lambda^{-1} > -\frac{1}{2}$) and similarly

$$0 < \theta < \frac{1}{2}, \quad y > |(f_1)_\phi(-\theta y^{\frac{1}{2}})(f_2)_\phi(-\theta y^{\frac{1}{2}})|^{-2} \Lambda^2 \quad (4.44)$$

$$\iff -\infty < w < 0, \quad 0 < \theta < |w| |(f_1)_\phi(w)(f_2)_\phi(w)| \Lambda^{-1}. \quad (4.45)$$

Therefore, since $\frac{1}{y^2} dy = \frac{2\theta^2}{|w|^3} dw$, (4.41) becomes

$$I_T(\Lambda) = \frac{6}{\pi^2} \int_0^\pi \left(\int_0^\infty \frac{1}{|w|^3} \int_{-|w| |(f_1)_\phi(w)(f_2)_\phi(w)| \Lambda^{-1}}^0 \theta^2 d\theta dw + \int_{-\infty}^0 \frac{1}{|w|^3} \int_0^{|w| |(f_1)_\phi(w)(f_2)_\phi(w)| \Lambda^{-1}} \theta^2 d\theta dw \right) d\phi = \quad (4.46)$$

$$= \frac{2}{\pi^2} \frac{1}{\Lambda^3} \int_0^\pi \int_{-\infty}^\infty |(f_1)_\phi(w)(f_2)_\phi(w)|^3 dw d\phi. \quad (4.47)$$

Now we use the inequality (valid for $0 \leq u \leq \frac{1}{2}$)

$$\left| 1 - \frac{1}{(1 \mp u)^3} \right| \leq 14u \quad (4.48)$$

and hence

$$\frac{1}{(1 \mp u)^3} = 1 + O(u) \quad (4.49)$$

where the implied constant can be taken to be 14. Hence, with

$$\Lambda = R^2 \mp \tilde{\kappa} \left(\frac{2^\eta \tilde{\kappa}}{R^2} \right)^{\eta-1} \quad (4.50)$$

we obtain

$$\frac{1}{\Lambda^3} = \frac{1}{R^6} (1 + O(2^{\eta(\eta-1)} \tilde{\kappa}^\eta R^{-2\eta})) \quad (4.51)$$

provided $2^{\eta(\eta-1)}\tilde{\kappa}^\eta R^{-2\eta} < \frac{1}{2}$. This is implied by the inequality

$$R^2 > 2^\eta \tilde{\kappa}. \quad (4.52)$$

Therefore, assumption (4.52) and the bound (4.51) allow us to write

$$\frac{1}{\Lambda^3} = \frac{1}{R^6} \left(1 + O_\eta((\kappa_\eta(f_1)\kappa_\eta(f_2))^\eta R^{-2\eta}) \right), \quad (4.53)$$

where the constant implied by the O_η -notation can be taken to be $7(2^\eta C_\eta)^\eta$. As in the rational case, we notice that (4.52) implies (4.40) and hence in view of (4.47) and (4.53) we obtain

$$I_T(\Lambda) = \frac{2}{\pi^2} D_{\text{irr}}(f_1, f_2) R^{-6} \left(1 + O_\eta((\kappa_\eta(f_1)\kappa_\eta(f_2))^\eta R^{-2\eta}) \right) \quad (4.54)$$

for (T, Λ) either equal to $(\frac{1}{2}, R^2 - \tilde{\kappa}(\frac{2^\eta \tilde{\kappa}}{R^2})^{\eta-1})$ or to $(1, R^2 + \tilde{\kappa}(\frac{2^\eta \tilde{\kappa}}{R^2})^{\eta-1})$. The bounds (4.39) allow us to conclude the proof. \square

Combining Theorem 3.3 and Lemma 4.3, we get the irrational case of Theorem 1.7.

5 Dynamical approximation of the indicators

Let $\chi_s = \mathbf{1}_{(0,s)}$ and $\chi = \chi_1$. Our goal is to accurately estimate $\mu_{\Gamma \setminus G}^0((\Theta_\chi \overline{\Theta_{\chi_b}})^{-1}(\mathcal{A}))$ and $\mu_{\Gamma \setminus G}((\Theta_\chi \overline{\Theta_{\chi_b}})^{-1}(\mathcal{A}))$ from (3.10) and (3.11) when \mathcal{A} is the complement of a closed ball of large radius R . Since $\chi, \chi_b \notin \mathcal{S}_\eta(\mathbb{R})$ with $\eta > 1$, we cannot apply Lemmata 4.2 and 4.3 directly. Instead, we will apply them to $\mu_{\Gamma \setminus G}^0((\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}})^{-1}(B_R^c))$ and $\mu_{\Gamma \setminus G}((\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}})^{-1}(B_R^c))$, where $\chi^{(J)}, \chi_b^{(J)}$ are smooth approximations of χ, χ_b belonging in $\mathcal{S}_\eta(\mathbb{R})$ with $\eta > 1$. We now illustrate the construction of such approximations as in [6].

5.1 Smooth trapezoidal approximation of indicators

Let $a \leq b$ and $\varepsilon, \delta \geq 0$. We introduce a piecewise quadratic function that is identically zero outside $(a - \varepsilon, b + \delta)$ and identically one within (a, b) . We set

$$T_{a,b}^{\varepsilon,\delta}(w) = \begin{cases} 0 & w \leq a - \varepsilon \\ \frac{2}{\varepsilon^2}(w - (a - \varepsilon))^2 & a - \varepsilon < w \leq a - \frac{\varepsilon}{2} \\ 1 - \frac{2}{\varepsilon^2}(w - a)^2 & a - \frac{\varepsilon}{2} < w \leq a \\ 1 & a < w \leq b \\ 1 - \frac{2}{\delta^2}(w - b)^2 & b < w \leq b + \frac{\delta}{2} \\ \frac{2}{\delta^2}(w - (b + \delta))^2 & b + \frac{\delta}{2} < w \leq b + \delta \\ 0 & w > b + \delta. \end{cases} \quad (5.1)$$

We have the following

Lemma 5.1 ([6], Lemma 3.1 therein). *Let $\varepsilon, \delta > 0$. Then $T_{a,b}^{\varepsilon,\delta} \in \mathcal{S}_2(\mathbb{R})$.*

As a consequence, $T_{a,b}^{\varepsilon,\delta} \in \mathcal{S}_\eta(\mathbb{R})$ for every $1 < \eta \leq 2$. Using piecewise functions consisting of higher degree polynomials, one could construct analogues of $T_{a,b}^{\varepsilon,\delta}$ that belong to $\mathcal{S}_\eta(\mathbb{R})$ for larger η . This, however, is not necessary in our analysis since, as we shall see in Section 8, values of η closer to 1 will yield better power savings in the error terms. Note that Lemma (5.1) does not apply to χ_s . Let $\Delta = T_{\frac{1}{3},\frac{1}{3}}^{\frac{1}{6},\frac{1}{3}}$. We have the following partition of unity:

$$\chi(w) = \sum_{j=0}^{\infty} \Delta(2^j w) + \sum_{j=0}^{\infty} \Delta(2^j(1 - w)). \quad (5.2)$$

It is crucial to rewrite (5.2) dynamically, using the geodesic flow $\tilde{\Phi}$ from Section 2.6. Using (2.19), (2.23), and (2.25), we get the identities

$$[\tilde{R}(\tilde{\Phi}^{(-2 \log 2)^j})f](w) = 2^{\frac{j}{2}} f(2^j w), \quad (5.3)$$

$$[\tilde{R}(\tilde{\gamma}_4 \tilde{\Phi}^{(-2 \log 2)^j})f](w) = 2^{\frac{j}{2}} f(2^j(w - 1)), \quad (5.4)$$

where $\tilde{\gamma}_4$ is as in (2.34). We then rewrite (5.2) as

$$\chi(w) = \sum_{j=0}^{\infty} 2^{-\frac{j}{2}} [\tilde{R}(\tilde{\Phi}^{(-2 \log 2)^j})\Delta](w) + \sum_{j=0}^{\infty} 2^{-\frac{j}{2}} [\tilde{R}(\tilde{\gamma}_4 \tilde{\Phi}^{(-2 \log 2)^j})\Delta_-](w). \quad (5.5)$$

where $\Delta_-(w) = \Delta(-w)$. We observe that each sum in (5.5) is a weighted sum along the backward orbit of the geodesic flow. The flow renormalizes each sum, mapping one term to the next and providing the exponential weights. Furthermore, the action of the geodesic flow $\tilde{\Phi}$ allows us to obtain indicators χ_s from χ by rescaling, namely

$$\chi_s = \sqrt{s} \tilde{R}(\tilde{\Phi}^{2 \log s}) \chi. \quad (5.6)$$

Therefore, using the group law (2.6), we obtain

$$\chi_s(w) = \sqrt{s} \left(\sum_{j=0}^{\infty} 2^{-\frac{j}{2}} [\tilde{R}(\tilde{\Phi}^{2\log(\frac{s}{2^j})})\Delta](w) + \sum_{j=0}^{\infty} 2^{-\frac{j}{2}} [\tilde{R}(\tilde{\gamma}_4\tilde{\Phi}^{2\log(\frac{s}{2^j})})\Delta_-](w) \right) \quad (5.7)$$

$$= \chi_{s,L}(w) + \chi_{s,R}(w), \quad (5.8)$$

where $\chi_{s,L} = T_{0,\frac{s}{3}}^{0,\frac{s}{3}}$ and $\chi_{s,R} = T_{\frac{2s}{3},s}^{\frac{s}{3},0}$. A natural way to construct an approximation $\chi_s^{(J)}$ of χ_s that is dynamically compatible with the action of the geodesic flow is to simply truncate the sums in (5.7) to $0 \leq j \leq J-1$ for some $J \geq 1$. This gives a smooth trapezoid function that equals $T_{\frac{s}{3 \cdot 2^{J-1}}, \frac{s}{3 \cdot 2^{J-1}}}^{\frac{s}{3}, \frac{s}{3}}$ and can be written as

$$\chi_s^{(J)}(w) = \sqrt{s} \left(\sum_{j=0}^{J-1} 2^{-\frac{j}{2}} [\tilde{R}(\tilde{\Phi}^{2\log(\frac{s}{2^j})})\Delta](w) + \sum_{j=0}^{J-1} 2^{-\frac{j}{2}} [\tilde{R}(\tilde{\gamma}_4\tilde{\Phi}^{2\log(\frac{s}{2^j})})\Delta_-](w) \right) \quad (5.9)$$

$$= \chi_{s,L}^{(J)}(w) + \chi_{s,R}^{(J)}(w), \quad (5.10)$$

where $\chi_{s,L}^{(J)} = T_{\frac{s}{3 \cdot 2^{J-1}}, \frac{s}{3}}^{\frac{s}{3}, \frac{s}{3}}$ and $\chi_{s,R}^{(J)} = T_{\frac{2s}{3}, \frac{s}{3 \cdot 2^{J-1}}}^{\frac{s}{3}, \frac{s}{3}}$, respectively. When $s = 1$, we will drop s from the notation and simply write χ , χ_L , χ_R , $\chi^{(J)}$, $\chi_L^{(J)}$, and $\chi_R^{(J)}$.

5.2 Defining Θ_{χ_s} almost everywhere

By linearity, (2.28), and (5.7) we may write

$$\Theta_{\chi_s}(\tilde{g}) := \sqrt{s} \left(\sum_{j=0}^{\infty} 2^{-\frac{j}{2}} \Theta_{\Delta}(\tilde{g}\tilde{\Phi}^{2\log(\frac{s}{2^j})}) + \sum_{j=0}^{\infty} 2^{-\frac{j}{2}} \Theta_{\Delta_-}(\tilde{g}\tilde{\gamma}_4\tilde{\Phi}^{2\log(\frac{s}{2^j})}) \right). \quad (5.11)$$

Since $\chi_s \notin \mathcal{S}_{\eta}$ for $\eta > 1$ this theta function is not defined for all $\tilde{g} \in \tilde{G}$. On the other hand, for $J \geq 1$ we have

$$\Theta_{\chi_s^{(J)}}(\tilde{g}) = \mathcal{L}_s^{(J)}(\tilde{g}) + \mathcal{R}_s^{(J)}(\tilde{g}\tilde{\gamma}_4^s) \quad (5.12)$$

where

$$\mathcal{L}_s^{(J)}(\tilde{g}) = \Theta_{\chi_{s,L}^{(J)}}(\tilde{g}) = \sqrt{s} \sum_{j=0}^{J-1} 2^{-\frac{j}{2}} \Theta_{\Delta}(\tilde{g}\tilde{\Phi}^{2\log(\frac{s}{2^j})}), \quad (5.13)$$

$$\mathcal{R}_s^{(J)}(\tilde{g}) = \Theta_{\chi_{s,R}^{(J)}}(\tilde{g}) = \sqrt{s} \sum_{j=0}^{J-1} 2^{-\frac{j}{2}} \Theta_{\Delta_-}(\tilde{g}\tilde{\gamma}_4\tilde{\Phi}^{2\log(\frac{s}{2^j})}). \quad (5.14)$$

Note that these theta functions for the smoothed approximations are well defined on all of \tilde{G} by Lemma 5.1. By splitting the summation in (5.11) into blocks of length J as in the proof of Lemma 3.14 in [6], we can write

$$\Theta_{\chi_s}(\tilde{g}) = \sum_{k=0}^{\infty} 2^{-k\frac{J}{2}} \left(\mathcal{L}_s^{(J)}(\tilde{g}\tilde{\Phi}^{-(2\log 2)kJ}) + \mathcal{R}_s^{(J)}(\tilde{g}\tilde{\gamma}_4\tilde{\Phi}^{-(2\log 2)kJ}) \right). \quad (5.15)$$

The term $k = 0$ in (5.15) corresponds to (5.12) and will provide the main contribution to our estimates, while the other terms, which are obtained by the renormalising action of the backward geodesic flow, will contribute to the error term.

As in Lemma 4.2 and Lemma 4.3, we need to consider the push forward of the measures $\mu_{\Gamma \backslash G}^0$ and $\mu_{\Gamma \backslash G}$ via the function $\Theta_\chi \overline{\Theta_{\chi_b}}$, and hence it is crucial that latter is well defined on a set of full $\mu_{\Gamma \backslash G}^0$ and $\mu_{\Gamma \backslash G}$ -measure. When $s = 1$, Theorem 3.10 in [6] shows that there is a $\tilde{\Gamma}$ -invariant set $\tilde{W} \subset \tilde{G}$ (explicitly defined by means of a Diophantine condition) of full $\tilde{\mu}$ -measure such that the series in (5.11) are absolutely convergent for all $\tilde{g} \in \tilde{W}$. Furthermore, the $\tilde{\mu}$ -a.e. defined function $\Theta_\chi(\tilde{g})$ is $\tilde{\Gamma}$ -invariant and hence we can write it as $\Theta_\chi(\tilde{\Gamma}\tilde{g})$. All other values of $s > 0$ can be handled in the same way using (5.6). To describe the set \tilde{W} we use a set of coordinates on \tilde{G} different from $(x + iy, \phi; \xi, \zeta) \in \mathfrak{H} \times \mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}$ on \tilde{G} . We consider the flow $\tilde{\Phi}^t$ together with its stable, unstable, and neutral manifolds. Letting $\tilde{n}_+(v, \rho) := ([(\frac{1}{0} \ v), 0]; (\frac{\rho}{0}), 0)$ and $\tilde{n}_-(u, \sigma) := ([(\frac{1}{u} \ 0), \arg(u \cdot + 1)]; (\frac{0}{\sigma}), 0)$ parametrise the unstable and stable manifolds respectively, we can write any $\tilde{g} \in \tilde{G}$ as

$$\tilde{g} = \tilde{n}_+(v, \rho)\tilde{n}_-(u, \sigma)\tilde{\Phi}^t([I, 0]; (\frac{0}{0}), \tau) \quad (5.16)$$

with $v, \rho, u, \sigma, t, \tau \in \mathbb{R}$. The set \tilde{W} of full $\tilde{\mu}$ -measure consists of all elements of the form (5.16) such that $v + \frac{1}{u}$ is Diophantine of type $\kappa > 1$. Note that the projection $W = \pi(\tilde{W}) \subset G$ via (2.7) is of full μ -measure and Γ -invariant. If we write (5.16) with respect to our usual coordinates on $\tilde{\Gamma} \backslash \tilde{G}$, we obtain

$$\tilde{\Gamma}n_+(v, \rho)n_-(u, \sigma)\tilde{\Phi}^t([I, 0]; (\frac{0}{0}), \tau) = \tilde{\Gamma}(n_x a_y k_\phi; (\frac{\xi_1}{\xi_2}), \zeta) \quad (5.17)$$

where $x = v + \frac{u}{u^2 + e^{2t}}$, $y = \frac{e^t}{u^2 + e^{2t}}$, $\xi_1 = \rho + v\sigma$, $\xi_2 = \sigma$, $\zeta = \tau$, and ϕ is determined by $\cos \phi = \frac{\text{sgn}(u)e^t}{\sqrt{u^2 + e^{2t}}}$ and $\sin \phi = \frac{\text{sgn}(u)u}{\sqrt{u^2 + e^{2t}}}$. In particular, since ξ_1 and ξ_2 do not depend on u , we have that for every $\xi \in \mathbb{R}^2$ there exists a set $U_\xi \subset \widetilde{\text{SL}}(2, \mathbb{R})$ of full $\mu_{\widetilde{\text{SL}}(2, \mathbb{R})}$ -measure such that $\Theta_{\chi_s}(x + iy, \phi; \xi, \zeta)$ is well-defined for every $(x + iy, \phi) \in U_\xi$, every $\zeta \in \mathbb{R}$, and every $s > 0$. For $\xi \in \mathbb{R}^2$, we let $\tilde{W}_\xi = \bigcup_{\tilde{\gamma} \in \tilde{\Gamma}} \tilde{\gamma}(U_\xi \times \{\xi\} \times \mathbb{R})$ and consider its projection

$W_\xi := \pi(\tilde{W}_\xi)$, which is Γ -invariant since $\pi(\tilde{\Gamma}) = \Gamma$. In particular, W_0 is Γ -invariant and is of full μ^0 -measure and hence ΓW_0 is of full $\mu_{\Gamma \backslash G}^0$ -measure. Therefore, since $\Theta_\chi(\tilde{g})$ and $\Theta_{\chi_b}(\tilde{g})$ are simultaneously defined (and $\tilde{\Gamma}$ -invariant) for all $\tilde{g} \in \tilde{W}_0$ and the product $\Theta_\chi(\tilde{g})\overline{\Theta_{\chi_b}(\tilde{g})}$ only depends on $g = \pi(\tilde{g})$, then $\Theta_\chi \overline{\Theta_{\chi_b}}(\Gamma g)$ is defined for all Γg in the full $\mu_{\Gamma \backslash G}^0$ -measure set ΓW_0 .

5.3 Expansion of the product $\Theta_\chi \overline{\Theta_{\chi_b}}$

Let us now use (5.15) to expand $\Theta_\chi \overline{\Theta_{\chi_b}}(g)$. Depending on whether we are considering the irrational case or the rational case, we consider $g \in W$ or $g \in W_0$ and we consider

$\tilde{g} \in \tilde{W}$ or $\tilde{g} \in \tilde{W}_0$ such that $\pi(\tilde{g}) = g$. Then, we obtain

$$\Theta_\chi \overline{\Theta_{\chi_b}}(g) = \Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}(g) + \quad (5.18)$$

$$+ \sum_{k+\ell>0} 2^{-\frac{(k+\ell)J}{2}} \mathcal{L}_1^{(J)}(\tilde{g}\tilde{\Phi}^{-(2\log 2)kJ}) \overline{\mathcal{L}_b^{(J)}(\tilde{g}\tilde{\Phi}^{-(2\log 2)\ell J})} + \quad (5.19)$$

$$+ \sum_{k+\ell>0} 2^{-\frac{(k+\ell)J}{2}} \mathcal{L}_1^{(J)}(\tilde{g}\tilde{\Phi}^{-(2\log 2)kJ}) \overline{\mathcal{R}_b^{(J)}(\tilde{g}\tilde{\gamma}_4\tilde{\Phi}^{-(2\log 2)\ell J})} + \quad (5.20)$$

$$+ \sum_{k+\ell>0} 2^{-\frac{(k+\ell)J}{2}} \mathcal{R}_1^{(J)}(\tilde{g}\tilde{\Phi}^{-(2\log 2)kJ}) \overline{\mathcal{L}_b^{(J)}(\tilde{g}\tilde{\Phi}^{-(2\log 2)\ell J})} + \quad (5.21)$$

$$+ \sum_{k+\ell>0} 2^{-\frac{(k+\ell)J}{2}} \mathcal{R}_1^{(J)}(\tilde{g}\tilde{\Phi}^{-(2\log 2)kJ}) \overline{\mathcal{R}_b^{(J)}(\tilde{g}\tilde{\gamma}_4\tilde{\Phi}^{-(2\log 2)\ell J})}. \quad (5.22)$$

Note that each term in (5.19) only depends on $g = \pi(\tilde{g})$ since we can once again use the renormalisation given by the geodesic flow to write

$$\mathcal{L}_1^{(J)}(\tilde{g}\tilde{\Phi}^{-(2\log 2)kJ}) \overline{\mathcal{L}_b^{(J)}(\tilde{g}\tilde{\Phi}^{-(2\log 2)\ell J})} = \Theta_{\chi_{1,L}^{(J)}}(\tilde{g}\tilde{\Phi}^{-(2\log 2)kJ}) \overline{\Theta_{\chi_{b,L}^{(J)}}(\tilde{g}\tilde{\Phi}^{-(2\log 2)\ell J})} = \quad (5.23)$$

$$= \Theta_{\chi_{1,L}^{(J,k)}}(\tilde{g}) \overline{\Theta_{\chi_{b,L}^{(J,\ell)}}(\tilde{g})} = \Theta_{\chi_{1,L}^{(J,k)}} \overline{\Theta_{\chi_{b,L}^{(J,\ell)}}}(g), \quad (5.24)$$

where $\chi_{s,L}^{(J,m)} = R(\tilde{\Phi}^{-(2\log 2)mJ})\chi_{s,L}^{(J)}$ for $(s, m) \in \{(1, k), (b, \ell)\}$. A similar argument holds for all other summands in (5.20)–(5.22). We now rewrite (5.18)–(5.22) as

$$\Theta_\chi \overline{\Theta_{\chi_b}}(g) = \Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}(g) + \sum_{k+\ell>0} 2^{-\frac{(k+\ell)J}{2}} \mathcal{L}\overline{\mathcal{L}}_{k,\ell}^{(J)}(g) + \sum_{k+\ell>0} 2^{-\frac{(k+\ell)J}{2}} \mathcal{L}\overline{\mathcal{R}}_{k,\ell}^{(J)}(g) + \quad (5.25)$$

$$+ \sum_{k+\ell>0} 2^{-\frac{(k+\ell)J}{2}} \mathcal{R}\overline{\mathcal{L}}_{k,\ell}^{(J)}(g) + \sum_{k+\ell>0} 2^{-\frac{(k+\ell)J}{2}} \mathcal{R}\overline{\mathcal{R}}_{k,\ell}^{(J)}(g), \quad (5.26)$$

with the obvious notations, where we have dropped the explicit dependence on b .

6 Estimating κ_η for $1 < \eta \leq 2$

Recall the function $\chi_{s,L}^{(J)}$ introduced in Section 5.1 and used, along with its mirror image $\chi_{s,L}^{(J)}(-\cdot)$, in Section 5.2 to define $\mathcal{L}_s^{(J)}$ and $\mathcal{R}_s^{(J)}$. These functions are also implicitly used in (5.25)–(5.26). It is crucial for us to estimate $\kappa_\eta(\chi_s^{(J)})$, $\kappa_\eta(\chi_{s,L}^{(J)})$, and $\kappa_\eta(\chi_{s,L}^{(J)}(-\cdot))$, extending the case $\eta = 2$ is considered in [6].

In this section we will write $f(t, x) = [\mathcal{J}_t f](x)$, where \mathcal{J}_t is the integral operator defined in (2.12).

Lemma 6.1. *The function $f(t, x) = [\mathcal{J}_t f](x)$ satisfies the Schrödinger equation*

$$\begin{cases} \frac{i}{2\pi} \partial_t f(t, x) = \frac{1}{2} \left(-\frac{1}{4\pi^2} \frac{d^2}{dx^2} + x^2 \right) f(t, x). \\ f(0, x) = f(x). \end{cases} \quad (6.1)$$

Proof. The fact that $f(0, x) = f(x)$ is part of the definition (2.12). A direct computation yields

$$\begin{aligned} \frac{i}{2\pi} \partial_t f(t, x) = & |\sin(t)|^{-\frac{1}{2}} \int_{\mathbb{R}} e\left(\frac{1}{2}(x^2 + w^2) \cot(t) - xw \csc(t)\right) \\ & \left(-\frac{i}{4\pi} \cot(t) + \frac{1}{2}(x^2 + w^2) \csc^2(t) - xw \cos(t) \csc^2(t)\right) f(w) dw \end{aligned} \quad (6.2)$$

and

$$\begin{aligned} \frac{1}{2} \left(-\frac{1}{4\pi^2} \frac{d^2}{dx^2} + x^2\right) f(t, x) = & |\sin(t)|^{-\frac{1}{2}} \int_{\mathbb{R}} e\left(\frac{1}{2}(x^2 + w^2) \cot(t) - xw \csc(t)\right) \\ & \left(\frac{1}{2}x^2 \cot^2(t) + \frac{1}{2}w^2 \csc^2(t) - xw \cos(t) \csc^2(t) - \frac{i \cot(t)}{4\pi} + \frac{1}{2}x^2\right) f(w) dw. \end{aligned} \quad (6.3)$$

The result then follows from the identity $1 + \cot^2(t) = \csc^2(t)$. \square

The following lemma can be seen as the consequence of a ‘coarse’ stationary phase argument, and is very important in our bounds for κ_η .

Lemma 6.2 ([6], Lemma 3.3 therein). *Let f be a compactly supported function with bounded variation. Let $f(t, x)$ be the solution to (6.1) with initial condition f . Then*

$$|f(t, x)| \leq \|f\|_{\text{sp}} \quad (6.4)$$

uniformly in t , where $\|f\|_{\text{sp}} := \max\{2\|f\|_1, 3(\|f\|_\infty + V(f))\}$ and $V(f)$ is the total variation of f . In particular, if f is bounded and piecewise C^1 , then $V(f) = \|f'\|_1$.

Remark 6.3. We can bound the norm in (6.4) for the functions $\chi_s, \chi_s^{(J)}, \chi_{s,L}, \chi_{s,L}^{(J)}$ introduced in Section 5.2 as follows. Since $\|T_{a,b}^{\varepsilon,\delta}\|_1 = b - a + \frac{\varepsilon+\delta}{2}$, $\|T_{a,b}^{\varepsilon,\delta}\|_\infty = 1$, and $V(T_{a,b}^{\varepsilon,\delta}) = 2$, we get

$$\|\chi_s\|_{\text{sp}} = \max\{2s, 9\}, \quad (6.5)$$

$$\|\chi_s^{(J)}\|_{\text{sp}} = \max\{2(1 - 2^{-J})s, 9\} \leq \max\{2s, 9\}, \quad (6.6)$$

$$\|\chi_{s,L}\|_{\text{sp}} = \max\{s, 9\}, \quad (6.7)$$

$$\|\chi_{s,L}^{(J)}\|_{\text{sp}} = \max\{(1 - 2^{-J})s, 9\} \leq \max\{s, 9\}. \quad (6.8)$$

The following theorem will be used to estimate $\kappa_\eta(\chi_s^{(J)})$ and $\kappa_\eta(\chi_{s,L}^{(J)})$. It is in this theorem that we make the assumption that $1 < \eta \leq 2$ for the first time.

Theorem 6.4. *Let f be a bounded, compactly supported C^1 function such that $\|f\|_1, \|f'\|_1$, and $\|f''\|_1$ are all finite. Let $f(t, x)$ be the solution to (6.1) with initial condition f . Define $h^{p,q}(w) = w^p f^{(q)}(w)$. Then for all $1 < \eta \leq 2$, $b \geq 1$ and all $x, t \in \mathbb{R}$ we have*

$$\begin{aligned} |f(t, x)|(1 + |x|^2)^{\frac{\eta}{2}} \leq & 2^{\frac{\eta}{2}} \max \left\{ \|h^{0,0}\|_{\text{sp}}, b^{\eta-1} (\|h^{0,1}\|_1 + \|h^{1,0}\|_{\text{sp}}), \right. \\ & \left. b^{\eta-2} (\|h^{1,1}\|_1 + \|h^{0,2}\|_1 + \|h^{0,0}\|_1 + \|h^{2,0}\|_{\text{sp}}) \right\}. \end{aligned} \quad (6.9)$$

Proof. We first use the bound coming from applying Lemma 6.2 directly to f . For $|x| < 1$ and all $t \in \mathbb{R}$ we get

$$|(1 + |x|^2)^{\frac{\eta}{2}} f(t, x)| \leq 2^{\frac{\eta}{2}} \|f\|_{\text{sp}}. \quad (6.10)$$

Let $b \geq 1$ and assume that $1 \leq |x| < b$. Using integration by parts we get

$$f(t, x) = |\sin(t)|^{-\frac{1}{2}} \int_{\mathbb{R}} e(\frac{1}{2}(x^2 + w^2) \cot(t)) f(w) e(-xw \csc(t)) dw = \quad (6.11)$$

$$= \left[-|\sin(t)|^{-\frac{1}{2}} \frac{\sin(t)}{2\pi i x} e(\frac{1}{2}(x^2 + w^2) \cot(t)) f(w) \right]_{-\infty}^{\infty} + \quad (6.12)$$

$$|\sin(t)|^{-\frac{1}{2}} \frac{\sin(t)}{2\pi i x} \int_{\mathbb{R}} e(\frac{1}{2}(x^2 + w^2) \cot(t) - xw \csc(t)) [f'(w) + 2\pi i \cot(t) w f(w)] dw. \quad (6.13)$$

The boundary term in (6.12) is zero since f is compactly supported. The integral in (6.13) splits into two pieces by linearity. The first half is bounded in absolute value trivially by $\|f'\|_1$. To bound the absolute value of the second half of the integral (together with the prefactor $|\sin(t)|^{-\frac{1}{2}}$), we apply Lemma 6.2 to the function $h^{1,0}$. We obtain

$$\left| (1 + |x|^2)^{\frac{\eta}{2}} f(t, x) \right| \leq \frac{(1 + |x|^2)^{\frac{\eta}{2}}}{|x|} (\|h^{0,1}\|_1 + \|h^{1,0}\|_{\text{sp}}). \quad (6.14)$$

Since $|x| \geq 1$, we can write $(1 + |x|^2)^{\frac{\eta}{2}} |x|^{-1} \leq 2^{\frac{\eta}{2}} |x|^{\eta-1}$. Therefore, since $\eta > 1$ and $1 \leq |x| < b$, we have

$$\left| (1 + |x|^2)^{\frac{\eta}{2}} f(t, x) \right| \leq 2^{\frac{\eta}{2}} b^{\eta-1} (\|h^{0,1}\|_1 + \|h^{1,0}\|_{\text{sp}}), \quad (6.15)$$

for all in $t \in \mathbb{R}$. In the remaining range $|x| \geq b$ we start from (6.12)–(6.13) and perform one more integration by parts. We get

$$f(t, x) = \quad (6.16)$$

$$= |\sin(t)|^{-\frac{1}{2}} \left(\frac{\sin(t)}{2\pi i x} \right)^2 \int_{\mathbb{R}} e(\frac{1}{2}(x^2 + w^2) \cot(t) - xw \csc(t)) \times \quad (6.17)$$

$$\times (4\pi i \cot(t) w f'(w) + f''(w) + 2\pi i \cot(t) f(w) - 4\pi^2 \cot^2(t) w^2 f(w)) dw. \quad (6.18)$$

We split (6.17)–(6.18) into four integrals by linearity. The first one yields the bound $\frac{1}{\pi} |\sin(t)|^{\frac{1}{2}} |\cos(t)| \|h^{1,1}\|_1$, the second the bound $\frac{1}{4\pi^2} |\sin(t)|^{\frac{3}{2}} \|f''\|_1$, and the third the bound $\frac{1}{2\pi} |\sin(t)|^{\frac{1}{2}} |\cos(t)| \|f\|_1$. The fourth integral is dealt with by applying Lemma 6.2 to the function $h^{2,0}$ in order to avoid the possible divergence for t close to an integer multiple of π . We thus obtain

$$\left| (1 + |x|^2)^{\frac{\eta}{2}} f(t, x) \right| \leq \frac{(1 + |x|^2)^{\frac{\eta}{2}}}{|x|^2} (\|h^{1,1}\|_1 + \|h^{0,2}\|_1 + \|h^{0,0}\|_1 + \|h^{2,0}\|_{\text{sp}}). \quad (6.19)$$

Since $|x| \geq b \geq 1$ and $\eta \leq 2$, we have

$$\left| (1 + |x|^2)^{\frac{\eta}{2}} f(t, x) \right| \leq 2^{\frac{\eta}{2}} b^{\eta-2} (\|h^{1,1}\|_1 + \|h^{0,2}\|_1 + \|h^{0,0}\|_1 + \|h^{2,0}\|_{\text{sp}}) \quad (6.20)$$

for all $t \in \mathbb{R}$. Combining (6.10), (6.15) and (6.20), we obtain (6.9). \square

We now extend Lemma 3.1 in [6], in which the case $\eta = 2$ is considered, by proving the following

Proposition 6.5. *Fix $s \geq 1$, and $J \geq 1$ as above. Then for $1 < \eta \leq 2$ we have*

$$\kappa_\eta(\chi_s^{(J)}) \leq K(s) 2^{(\eta-1)J} \quad \text{and} \quad \kappa_\eta(\chi_{s,L}^{(J)}(\pm \cdot)) \leq K_L(s) 2^{(\eta-1)J}, \quad (6.21)$$

where $K(s)$ and $K_L(s)$ are given explicitly in (6.28) and (6.35) respectively.

Proof. Let $f = \chi_s^{(J)} = T^{\frac{s}{6 \cdot 2^{J-1}}, \frac{s}{6 \cdot 2^{J-1}}}$. Let us estimate the norms featured in (6.9). Using the notation $h^{p,q}$ as in Theorem 6.4 and the definition (5.1) directly (see also Remark 6.3), we obtain $\|h^{0,0}\|_{\text{sp}} \leq \max\{2s, 9\}$, $\|(h^{0,0})'\|_1 = \|h^{0,1}\|_1 = 2$, $\|h^{1,0}\|_1 = \frac{1}{2}s^2(1 - 2^{-J}) \leq \frac{s^2}{2}$, $\|h^{1,0}\|_\infty \leq s$, $\|h^{1,1}\|_1 = s$, $\|(h^{1,0})'\|_1 \leq \|h^{0,0}\|_1 + \|h^{1,1}\|_1 \leq 2s$, $\|h^{1,0}\|_{\text{sp}} \leq \max\{s^2, 9s\}$, $\|h^{0,2}\|_1 = \frac{24 \cdot 2^J}{s} \leq 24 \cdot 2^J$, $\|h^{2,0}\|_1 \leq \frac{s^3}{3}$, $\|h^{2,0}\|_\infty \leq s^2$, $\|h^{2,1}\|_1 \leq s^2$, $\|(h^{2,0})'\|_1 \leq 2\|h^{1,0}\|_1 + \|h^{2,1}\|_1 \leq 2s^2$, and $\|h^{2,0}\|_{\text{sp}} \leq \max\{\frac{2}{3}s^3, 9s^2\}$. Using the definition (2.26) and Theorem 6.4 with $b = 2^J$, we obtain

$$\sup_{\phi, w} |f(\phi, w)|(1 + |w|^2)^{\frac{\eta}{2}} \leq \quad (6.22)$$

$$\leq 2^{\frac{\eta}{2}} \max \left\{ \max\{2s, 9\}, \right. \quad (6.23)$$

$$\left. 2^{(\eta-1)J} (2 + \max\{s^2, 9s\}), \right. \quad (6.24)$$

$$\left. 2^{(\eta-2)J} (s + 24 \cdot 2^J + s + \max\{\frac{2}{3}s^3, 9s^2\}) \right\} \leq \quad (6.25)$$

$$\leq 2^{\frac{\eta}{2}} 2^{(\eta-1)J} \max \left\{ 2s, 9, 2 + \max\{9s, s^2\}, s + 24 + \max\left\{\frac{9}{2}s^2, \frac{1}{3}s^3\right\} \right\} \leq \quad (6.26)$$

$$\leq K(s) 2^{(\eta-1)J}, \quad (6.27)$$

where

$$K(s) = 48 + 2s + \max\left\{9s^2, \frac{2}{3}s^3\right\}. \quad (6.28)$$

We can repeat the same argument letting $f = \chi_{s,L}^{(J)} = T^{\frac{s}{6 \cdot 2^{J-1}}, \frac{s}{3 \cdot 2^{J-1}}}$ instead. Also in this case the norms featured in (6.9) can be estimated directly using (5.1), although the computations are rather tedious. We have $\|h^{0,0}\|_{\text{sp}} \leq \max\{s, 9\}$, $\|(h^{0,0})'\|_1 = \|h^{0,1}\|_1 = 2$, $\|h^{1,0}\|_1 = \frac{55}{432}s^2(1 - 2^{-2J}) \leq \frac{55}{432}s^2$, $\|h^{1,0}\|_\infty \leq \frac{2}{3}s$, $\|h^{1,1}\|_1 = \frac{1}{2}s(1 + 2^{-J}) \leq \frac{3}{4}s$, $\|(h^{1,0})'\|_1 \leq \|h^{0,0}\|_1 + \|h^{1,1}\|_1 \leq \frac{5}{4}s$, $\|h^{1,0}\|_{\text{sp}} \leq \max\{\frac{55}{216}s^2, \frac{23}{4}s\}$, $\|h^{0,2}\|_1 = \frac{12}{s}(1 + 2^J) \leq 18 \cdot 2^J$, $\|h^{2,0}\|_1 = \frac{19}{432}s^3(1 - 2^{3J}) \leq \frac{19}{432}s^3$, $\|h^{2,0}\|_\infty \leq \frac{4}{9}s^2$, $\|h^{2,1}\|_1 = \frac{55}{216}s^2(1 + 2^{-2J}) \leq \frac{275}{864}s^2$, $\|(h^{2,0})'\|_1 \leq 2\|h^{1,0}\|_1 + \|h^{2,1}\|_1 \leq \frac{55}{96}s^2$, and $\|h^{2,0}\|_{\text{sp}} \leq \max\{\frac{19}{216}s^3, \frac{293}{96}s^2\}$. Arguing as above, we obtain

$$\sup_{\phi, w} |f(\phi, w)|(1 + |w|^2)^{\frac{\eta}{2}} \leq \quad (6.29)$$

$$\leq 2^{\frac{\eta}{2}} \max \left\{ \max\{s, 9\}, \right. \quad (6.30)$$

$$\left. 2^{(\eta-1)J} \left(2 + \max\left\{ \frac{55}{216}s^2, \frac{23}{4}s \right\} \right), \right. \quad (6.31)$$

$$\left. 2^{(\eta-2)J} \left(\frac{3}{4}s + 18 \cdot 2^J + \frac{s}{2} + \max\left\{ \frac{19}{216}s^3, \frac{293}{96}s^2 \right\} \right) \right\} \leq \quad (6.32)$$

$$\leq 2^{\frac{\eta}{2}} 2^{(\eta-1)J} \max \left\{ s, 9, 2 + \max\left\{ \frac{23}{4}s, \frac{55}{216}s^2 \right\}, 18 + \frac{5}{8}s + \max\left\{ \frac{293}{192}s^2, \frac{19}{432}s^3 \right\} \right\} \leq \quad (6.33)$$

$$\leq K_L(s) 2^{(\eta-1)J}, \quad (6.34)$$

where

$$K_L(s) = 36 + \frac{5}{4}s + \max \left\{ \frac{293}{96}s^2, \frac{19}{216}s^3 \right\}. \quad (6.35)$$

The argument for $\chi_{s,L}^{(J)}(-\cdot)$ is the same as for $\chi_{s,L}^{(J)}$. \square

Remark 6.6. An elementary but tedious analysis shows that for all $s \geq 1$ we have

$$2^{\frac{1}{2}} \leq \frac{K(s)}{K_L(s)} \leq 2^3. \quad (6.36)$$

6.1 Divergence of expected displacement for harmonic quantum oscillator with piecewise initial data

We illustrate here an application of Theorem 6.4. The equation (6.1) describes a 1-dimensional quantum harmonic oscillator. If we interpret $x \mapsto |f(t, x)|^2$ as a probability density, then we can consider the m -th moment of the displacement of the quantum particle at time t , i.e. $\langle x^m \rangle = \int_{\mathbb{R}} x^m |f(t, x)|^2 dx$. It is well known that the regularity of

the initial condition f determines the decay of $f(t, \cdot)$. If, for instance, $f \in \mathcal{S}(\mathbb{R})$, then $f(t, \cdot) \in \mathcal{S}(\mathbb{R})$ and hence all moments $\langle x^m \rangle$ are finite. However, if the initial condition is only piecewise C^1 with finite jumps (such as $f = \chi$), then $\langle x \rangle$ diverges because $|f(t, x)|^2$ decays too slowly as $|x| \rightarrow \infty$. In Theorem 6.8 below we consider smooth approximations to such initial data, and we precisely quantify the rate at which the divergence in the expected displacement occurs as the approximation increases.

For the rest of this section, f_ϵ will not denote $\mathcal{F}_\epsilon f$, and we will instead use it to denote a sequence of functions approaching f in the suitable sense as $\epsilon \rightarrow 0^+$.

Definition 6.7. A sequence $(f_\epsilon)_{1 \geq \epsilon > 0}$ of C^1 functions converging in $L^1(\mathbb{R})$ to a piecewise C^1 function $f_0 \in C_{\text{pw}}^1$ as $\epsilon \rightarrow 0^+$ is said to *converge with bounded gradient* if there exists a constant $C > 0$ such that for all $\epsilon > 0$ we have $\|f_\epsilon\|_1 \leq C$, $\|f'_\epsilon\|_1 \leq C$, and $\|f''_\epsilon\|_1 \leq C\epsilon^{-1}$. That is to say, the derivative of the functions f_ϵ will diverge as $\epsilon \rightarrow 0$ to approximate the piecewise function f_0 , but such divergence must occur in proportionally small intervals.

This definition can be motivated as follows. Let $f_0 \in C_{\text{pw}}^1$ be piecewise C^1 with finitely many jump discontinuities at the points x_j , and construct a sequence $f_\epsilon(x) = f_0(x) + \sum_j g_j((x - x_j)/\epsilon)$ where the g_j 's are compactly supported functions in $[-1, 1]$ chosen in order to make f_ϵ differentiable. By direct computation one can check that $f_\epsilon \rightarrow f_0$ in L^1 with bounded gradient as $\epsilon \rightarrow 0^+$.

Theorem 6.8. *Let $f_0 \in C_{\text{pw}}^1$ be a bounded, compactly supported function with finitely many jump discontinuities. Let $(f_\epsilon)_{\epsilon>0}$ be a sequence of bounded, compactly supported C^1 functions converging to f_0 with bounded gradient. Let $f_\epsilon(t, x)$ be the solution to (6.1) with initial condition f_ϵ . Then for all $1 < \eta \leq 2$ and all sufficiently small ϵ there exists a constant $C' > 0$ such that*

$$|f_\epsilon(t, x)| \leq C'(1 + |x|^2)^{-\frac{\eta}{2}} \epsilon^{-(\eta-1)} \quad (6.37)$$

uniformly for all t .

To illustrate the above theorem, fix $1 < \eta \leq 2$. As in (2.26)-(2.27), we think of η as a regularity parameter. Observe that (6.37) means that $f_\epsilon \in \mathcal{S}_\eta(\mathbb{R})$ for every $\epsilon > 0$ and provides a bound $\kappa_\eta(f_\epsilon) = O(\epsilon^{1-\eta})$. This also implies that for every fixed $\epsilon > 0$ the probability density $x \mapsto |f_\epsilon(t, x)|^2$ decays sufficiently fast as $|x| \rightarrow \infty$ so that the expected displacement $\int_{\mathbb{R}} x |f_\epsilon(t, x)|^2 dx$ is finite (and in fact uniformly bounded for all times). On the other hand, as mentioned above, the integral $\int_{\mathbb{R}} x |f_0(t, x)|^2 dx$ diverges for all $t > 0$ due to the lack of regularity of f_0 . These two contrasting behaviours are reconciled by the theorem: the smaller $\epsilon > 0$ is, the closer f_ϵ is to f_0 , and (6.37) shows how the uniform bound for the expected displacement blows up to infinity as $\epsilon \rightarrow 0^+$.

Proof of Theorem 6.8. In order to apply Theorem 6.4 to the function $f = f_\epsilon$, we need to consider the various norms featured in (6.9). We have

$$\|h^{0,0}\|_{\text{sp}} = \max\{2\|f_\epsilon\|_1, 3(\|f_\epsilon\|_\infty + \|f'_\epsilon\|_1)\} \leq K_1 \quad (6.38)$$

with $K_1 = 3(M + C)$, C as in Definition 6.7 and $M = \sup_{0 < \epsilon < 1} \|f_\epsilon\|_\infty$. Also,

$$\|h^{0,1}\|_1 + \|h^{1,0}\|_{\text{sp}} \leq C + \max\{2AC, 3(AM + C + AC)\} \leq K_2 \quad (6.39)$$

where $K_2 = 4C + 3A(C + M)$ and A is such that $\text{supp } f_\epsilon \subset [-A, A]$ for all $0 < \epsilon < 1$. Finally,

$$\|h^{1,1}\|_1 + \|h^{0,2}\|_1 + \|h^{0,0}\|_1 + \|h^{2,0}\|_{\text{sp}} \quad (6.40)$$

$$\leq AC + C\epsilon^{-1} + C + \max\{2A^2C, 3(A^2M + 2AC + A^2C)\} \leq \epsilon^{-1}K_3 \quad (6.41)$$

with $K_3 = 2C + 3A^2(C + M) + 7AC$. Using $b = \epsilon^{-1}$ in Theorem 6.4, we obtain Theorem 6.37 with $C' = 2^{\frac{\eta}{2}} \max\{K_1, K_2, K_3\}$. \square

7 Tail approximations

Recall (5.25)–(5.26), in which we have approximated $\Theta_\chi \overline{\Theta_{\chi_b}}(g)$ by $\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}(g)$. Let $\mu = \mu_{\Gamma \setminus G}^0$ in the rational case and $\mu = \mu_{\Gamma \setminus G}$ in the irrational case. In this section:

- (i) we first estimate how much the tail probability $\mu(|\Theta_\chi \overline{\Theta_{\chi_b}}| > R^2)$ differs from $\mu(|\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}| > R^2(1 \pm \delta)^2)$ for suitable choices of parameters $J \geq 4$ and $\delta > 0$ (see Lemmata 7.1 and 7.3),
- (ii) then we give an asymptotic for $\mu(|\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}| > R^2(1 \pm \delta)^2)$ which includes explicit error terms (see Lemmata 7.2 and 7.4).

7.1 Rational case

Lemma 7.1. *Fix $1 < \eta \leq 2$, $b \geq 1$, and let $K_L(b)$ as in Proposition 6.5. Then for all $J \geq 4$, $R^2 > 64 C_\eta K_L(b)^2 2^{2(\eta-1)J}$ and $2^{-\frac{J-1}{2}} < \delta < \frac{1}{2}$ we have*

$$\mu_{\Gamma \setminus G}^0(|\Theta_\chi \overline{\Theta_{\chi_b}}| > R^2) - \mu_{\Gamma \setminus G}^0(|\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}| > R^2(1 - \delta)^2) = O\left(\frac{1}{R^4 \delta^2 2^J}\right) \quad \text{and} \quad (7.1)$$

$$\mu_{\Gamma \setminus G}^0(|\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}| > R^2(1 + \delta)^2) - \mu_{\Gamma \setminus G}^0(|\Theta_\chi \overline{\Theta_{\chi_b}}| > R^2) = O\left(\frac{1}{R^4 \delta^2 2^J}\right), \quad (7.2)$$

where the constants implied by the O -notation in (7.1)–(7.2) can both be taken to be $2^{17} \|\chi_{b,L}\|_{\text{sp}}^4$ (see (5.8), Lemma 6.2, and (6.7)).

Proof. Set $S = \{(k, \ell) \in \mathbb{Z}_{\geq 0}^2 : (k, \ell) \neq (0, 0)\}$. Note that $1 = (1 - \delta)^2 + 4 \frac{(1 - \delta)^2}{4} \sum_{(k, \ell) \in S} \delta^{k+\ell}$.

We use (5.25)–(5.26) and a union bound to obtain

$$\mu_{\Gamma \setminus G}^0(|\Theta_\chi \overline{\Theta_{\chi_b}}| > R^2) - \mu_{\Gamma \setminus G}^0(|\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}| > R^2(1 - \delta)^2) \leq \quad (7.3)$$

$$\leq \sum_{(k, \ell) \in S} \mu_{\Gamma \setminus G}^0\left(2^{-\frac{(k+\ell)J}{2}} |\mathcal{L} \overline{\mathcal{L}}_{k, \ell}^{(J)}| > R^2 \frac{(1 - \delta)^2}{4} \delta^{k+\ell}\right) + \quad (7.4)$$

$$+ \sum_{(k, \ell) \in S} \mu_{\Gamma \setminus G}^0\left(2^{-\frac{(k+\ell)J}{2}} |\mathcal{L} \overline{\mathcal{R}}_{k, \ell}^{(J)}| > R^2 \frac{(1 - \delta)^2}{4} \delta^{k+\ell}\right) + \quad (7.5)$$

$$+ \sum_{(k, \ell) \in S} \mu_{\Gamma \setminus G}^0\left(2^{-\frac{(k+\ell)J}{2}} |\mathcal{R} \overline{\mathcal{L}}_{k, \ell}^{(J)}| > R^2 \frac{(1 - \delta)^2}{4} \delta^{k+\ell}\right) + \quad (7.6)$$

$$+ \sum_{(k, \ell) \in S} \mu_{\Gamma \setminus G}^0\left(2^{-\frac{(k+\ell)J}{2}} |\mathcal{R} \overline{\mathcal{R}}_{k, \ell}^{(J)}| > R^2 \frac{(1 - \delta)^2}{4} \delta^{k+\ell}\right). \quad (7.7)$$

We focus on estimating the sum in (7.4), since (7.5)–(7.7) are similar. First observe that

another union bound and the invariance of the measure imply, for every $(k, \ell) \in S$,

$$\mu_{\Gamma \setminus G}^{\mathbf{0}} \left(2^{-\frac{(k+\ell)J}{2}} |\mathcal{L}\bar{\mathcal{L}}_{k,\ell}^{(J)}| > R^2 \frac{(1-\delta)^2}{4} \delta^{k+\ell} \right) \leq \quad (7.8)$$

$$\leq \mu_{\Gamma \setminus G}^{\mathbf{0}} \left\{ g \in \mathcal{F}_{\Gamma} : \left| \Theta_{\chi_L^{(J)}}(\tilde{g} \tilde{\Phi}^{-(2 \log 2)kJ}) \right| > R^{\frac{1-\delta}{2}} (\delta 2^{\frac{J}{2}})^{\frac{k+\ell}{2}} \right\} + \quad (7.9)$$

$$+ \mu_{\Gamma \setminus G}^{\mathbf{0}} \left\{ g \in \mathcal{F}_{\Gamma} : \left| \Theta_{\chi_{b,L}^{(J)}}(\tilde{g} \tilde{\Phi}^{-(2 \log 2)\ell J}) \right| > R^{\frac{1-\delta}{2}} (\delta 2^{\frac{J}{2}})^{\frac{k+\ell}{2}} \right\} = \quad (7.10)$$

$$= \mu_{\Gamma \setminus G}^{\mathbf{0}} \left\{ g \in \mathcal{F}_{\Gamma} : \left| \Theta_{\chi_L^{(J)}}(\tilde{g}) \right| > R^{\frac{1-\delta}{2}} (\delta 2^{\frac{J}{2}})^{\frac{k+\ell}{2}} \right\} + \quad (7.11)$$

$$+ \mu_{\Gamma \setminus G}^{\mathbf{0}} \left\{ g \in \mathcal{F}_{\Gamma} : \left| \Theta_{\chi_{b,L}^{(J)}}(\tilde{g}) \right| > R^{\frac{1-\delta}{2}} (\delta 2^{\frac{J}{2}})^{\frac{k+\ell}{2}} \right\}. \quad (7.12)$$

In (7.9)–(7.12) use \tilde{g} such that $\pi(\tilde{g}) = g$ as discussed in Section 5.3. To apply Lemma 4.2 to (7.11) and (7.12) we need

$$R^{\frac{1-\delta}{2}} (\delta 2^{\frac{J}{2}})^{\frac{k+\ell}{2}} > 2^{\frac{\eta}{2}} C_{\eta}^{\frac{1}{2}} \max\{\kappa_{\eta}(\chi_L^{(J)}), \kappa_{\eta}(\chi_{b,L}^{(J)})\}. \quad (7.13)$$

By Proposition 6.5 we can write $\kappa_{\eta}(\chi_{s,L}^{(J)}) \leq K_L(s) 2^{(\eta-1)J}$ for $s \in \{1, b\}$ and thus, since $\eta \leq 2$ and $\delta < \frac{1}{2}$, (7.13) is implied by the inequality

$$R > 8 C_{\eta}^{\frac{1}{2}} K_L(s) 2^{(\eta-1)J}. \quad (7.14)$$

Under this assumption, Lemma 4.2 gives

$$\mu_{\Gamma \setminus G}^{\mathbf{0}} \left\{ g \in \mathcal{F}_{\Gamma} : \left| \Theta_{\chi_L^{(J)}}(\tilde{g}) \right| > R^{\frac{1-\delta}{2}} (\delta 2^{\frac{J}{2}})^{\frac{k+\ell}{2}} \right\} = \frac{\frac{2}{\pi^2} D_{\text{rat}}(\chi_L^{(J)}, \chi_L^{(J)})}{R^4 \frac{(1-\delta)^4}{16} (\delta^2 2^J)^{k+\ell}} (1 + \mathcal{E}), \quad (7.15)$$

where $|\mathcal{E}| \leq 3(2^{\eta} C_{\eta})^{\eta} \kappa_{\eta}(\chi_L^{(J)})^{2\eta} \left(R^{\frac{1-\delta}{2}} (\delta 2^{\frac{J}{2}})^{\frac{k+\ell}{2}} \right)^{-2\eta}$. By (7.13) we obtain $|\mathcal{E}| \leq 3$, and by Lemma 6.2 we have

$$D_{\text{rat}}(\chi_L^{(J)}, \chi_L^{(J)}) = \int_0^{\pi} \left| \left(\chi_L^{(J)} \right)_{\phi}(0) \right|^4 d\phi \leq \pi \|\chi_L^{(J)}\|_{\text{sp}}^4. \quad (7.16)$$

Since $\delta < \frac{1}{2}$ we have $(1-\delta)^4 > 2^{-4}$, and thus (7.15) can be bounded above by

$$\frac{2^{11} \|\chi_L^{(J)}\|_{\text{sp}}^4}{\pi R^4 (\delta^2 2^J)^{k+\ell}}. \quad (7.17)$$

An identical argument allows us to bound the quantity in (7.12) from above by

$$\frac{2^{11} \|\chi_{b,L}^{(J)}\|_{\text{sp}}^4}{\pi R^4 (\delta^2 2^J)^{k+\ell}}. \quad (7.18)$$

Now observe that for $q > 2$ we have $\sum_{(k,\ell) \in S} q^{-(k+\ell)} = \frac{1}{q} \left(\frac{2q^2 - q}{(q-1)^2} \right) \leq \frac{6}{q}$. This bound can be used for $q = \delta^2 2^J$ since $\delta > 2^{-\frac{J-1}{2}}$ by hypothesis. Therefore, since $\|\chi_L^{(J)}\|_{\text{sp}} < \|\chi_{b,L}^{(J)}\|_{\text{sp}}$, the sum in (7.4) is bounded above by

$$\frac{2^{13} \|\chi_{b,L}^{(J)}\|_{\text{sp}}^4}{R^4 \delta^2 2^J}. \quad (7.19)$$

We can obtain the same upper bound for each of the sums in (7.5)–(7.7) and therefore obtain (7.1) with an implied constant equal to $2^{15} \|\chi_{b,L}^{(J)}\|_{\text{sp}}^4$. To prove (7.2), we use a slightly different union bound, arising from the identity

$$(1 + \delta)^2 = 1 + (1 - \delta)^2 \left(\sum_{k \geq 0} \sum_{\ell \geq 1} \delta^{k+\ell} + \sum_{k \geq 1} \sum_{\ell \geq 0} \delta^{k+\ell} + \sum_{k \geq 1} \sum_{\ell \geq 1} \delta^{k+\ell} \right) = \quad (7.20)$$

$$= 1 + 4 \frac{(1 - \delta)^2}{4} \sum_{(k,\ell) \in S} c_{k,\ell} \delta^{k+\ell}, \quad (7.21)$$

where $c_{k,\ell} = 1$ if $k = 0$ or $\ell = 0$ and $c_{k,\ell} = 3$ otherwise. Now (5.25)–(5.26) and a union bound yield

$$\mu_{\Gamma \setminus G}^0 (|\Theta_\chi \overline{\Theta_{\chi_b}}| > R^2 (1 + \delta)^2) - \mu_{\Gamma \setminus G}^0 (|\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}| > R^2) \leq \quad (7.22)$$

$$\leq \sum_{(k,\ell) \in S} \mu_{\Gamma \setminus G}^0 \left(2^{-\frac{(k+\ell)J}{2}} |\mathcal{L} \overline{\mathcal{L}}_{k,\ell}^{(J)}| > c_{k,\ell} R^2 \frac{(1-\delta)^2}{4} \delta^{k+\ell} \right) + \quad (7.23)$$

$$+ \sum_{(k,\ell) \in S} \mu_{\Gamma \setminus G}^0 \left(2^{-\frac{(k+\ell)J}{2}} |\mathcal{L} \overline{\mathcal{R}}_{k,\ell}^{(J)}| > c_{k,\ell} R^2 \frac{(1-\delta)^2}{4} \delta^{k+\ell} \right) + \quad (7.24)$$

$$+ \sum_{(k,\ell) \in S} \mu_{\Gamma \setminus G}^0 \left(2^{-\frac{(k+\ell)J}{2}} |\mathcal{R} \overline{\mathcal{L}}_{k,\ell}^{(J)}| > c_{k,\ell} R^2 \frac{(1-\delta)^2}{4} \delta^{k+\ell} \right) + \quad (7.25)$$

$$+ \sum_{(k,\ell) \in S} \mu_{\Gamma \setminus G}^0 \left(2^{-\frac{(k+\ell)J}{2}} |\mathcal{R} \overline{\mathcal{R}}_{k,\ell}^{(J)}| > c_{k,\ell} R^2 \frac{(1-\delta)^2}{4} \delta^{k+\ell} \right). \quad (7.26)$$

Arguing as before, the sum in (7.23) can be bounded above by

$$\frac{2^{15} \|\chi_{b,L}^{(J)}\|_{\text{sp}}^4}{R^4 \delta^2 2^J}, \quad (7.27)$$

and similarly for (7.24)–(7.26). Using $\|\chi_{b,L}^{(J)}\|_{\text{sp}} \leq \|\chi_{b,L}\|_{\text{sp}}$ (see Remark 6.3) we obtain (7.2) with an implied constant equal to $2^{17} \|\chi_{b,L}\|_{\text{sp}}^4$. \square

Lemma 7.2. *Fix $1 < \eta \leq 2$, $b \geq 1$, and let $K(b)$ be as in Proposition 6.5. Then for all $J \geq 1$, $R^2 > 16 C_\eta K(b)^2 2^{2(\eta-1)J}$ and $0 < \delta < \frac{1}{2}$ we have*

$$\begin{aligned} & \mu_{\Gamma \setminus G}^0 \left\{ |\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}| > R^2 (1 \pm \delta)^2 \right\} = \\ & = \frac{2D_{\text{rat}}(\chi, \chi_b)}{\pi^2 R^4} (1 + O(\delta)) (1 + O_\eta(2^{2\eta(\eta-1)J} R^{-2\eta})) (1 + O(2^{-J})), \end{aligned} \quad (7.28)$$

where the constants implied by the O -notations can be taken to be 30 , $2^{10}C_\eta^2K(b)^4$, and $2^5b\|\chi_b\|_{\text{sp}}^3$ (see (6.5)) respectively.

Proof. In order to apply Lemma 4.2 we require

$$R^2(1 \pm \delta)^2 > 2^\eta C_\eta \kappa_\eta(\chi^{(J)}) \kappa_\eta(\chi_b^{(J)}). \quad (7.29)$$

By Proposition 6.5 we have $\kappa_\eta(\chi^{(J)}) \kappa_\eta(\chi_b^{(J)}) \leq K(1)K(b)2^{2(\eta-1)J}$. Since $\eta \leq 2$, $\delta < \frac{1}{2}$ and $K(1) \leq K(b)$, (7.29) is implied by the inequality

$$R^2 > 16 C_\eta K(b)^2 2^{2(\eta-1)J}. \quad (7.30)$$

We obtain

$$\mu_{\Gamma \setminus G}^0 \left\{ |\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}| > R^2(1 \pm \delta)^2 \right\} = \frac{\frac{2}{\pi^2} D_{\text{rat}}(\chi^{(J)}, \chi_b^{(J)})}{R^4(1 \pm \delta)^4} (1 + \mathcal{E}_1), \quad (7.31)$$

where

$$|\mathcal{E}_1| \leq 3(2^\eta C_\eta \kappa_\eta(\chi^{(J)}) \kappa_\eta(\chi_b^{(J)}))^\eta (R(1 \pm \delta))^{-2\eta} \quad (7.32)$$

$$\leq 3(2^{\eta+2} C_\eta K(b)^2)^\eta 2^{2\eta(\eta-1)J} R^{-2\eta} \quad (7.33)$$

$$\leq 2^{10} C_\eta^2 K(b)^4 2^{2\eta(\eta-1)J} R^{-2\eta}. \quad (7.34)$$

This yields the second implied constant in the statement of the Lemma. For the first implied constant, note that for $0 < \delta < \frac{1}{2}$ we have

$$\frac{1}{(1 \pm \delta)^4} \leq \frac{1}{(1 - \delta)^4} = (1 + \mathcal{E}_2), \quad |\mathcal{E}_2| \leq 30\delta. \quad (7.35)$$

The third implied constant is more laborious. We have the estimate

$$\left| D_{\text{rat}}(\chi, \chi_b) - D_{\text{rat}}(\chi^{(J)}, \chi_b^{(J)}) \right| \leq \quad (7.36)$$

$$\leq \int_0^\pi \left| |\chi_\phi(0) (\chi_b)_\phi(0)|^2 - |(\chi^{(J)})_\phi(0) (\chi_b^{(J)})_\phi(0)|^2 \right| d\phi \leq. \quad (7.37)$$

By the triangle inequality this is bounded above by

$$\int_0^\pi \left| |\chi_\phi(0) (\chi_b)_\phi(0)|^2 - |(\chi^{(J)})_\phi(0) (\chi_b)_\phi(0)|^2 \right| d\phi + \quad (7.38)$$

$$+ \int_0^\pi \left| |(\chi^{(J)})_\phi(0) (\chi_b)_\phi(0)|^2 - |(\chi^{(J)})_\phi(0) (\chi_b^{(J)})_\phi(0)|^2 \right| d\phi \leq. \quad (7.39)$$

Now apply Lemma 6.2 to $|(\chi_b)_\phi(0)|$ in the first term, and $|(\chi^{(J)})_\phi(0)|$ in the second term to obtain

$$\left| D_{\text{rat}}(\chi, \chi_b) - D_{\text{rat}}(\chi^{(J)}, \chi_b^{(J)}) \right| \leq \|\chi_b\|_{\text{sp}}^2 \int_0^\pi \left| |\chi_\phi(0)|^2 - |(\chi^{(J)})_\phi(0)|^2 \right| d\phi \quad (7.40)$$

$$+ \|\chi^{(J)}\|_{\text{sp}}^2 \int_0^\pi \left| |(\chi_b)_\phi(0)|^2 - |(\chi_b^{(J)})_\phi(0)|^2 \right| d\phi. \quad (7.41)$$

Factorising the integrands, applying Lemma 6.2 twice more, and using the reverse triangle inequality gives

$$\left| D_{\text{rat}}(\chi, \chi_b) - D_{\text{rat}}(\chi^{(J)}, \chi_b^{(J)}) \right| \leq \quad (7.42)$$

$$\leq \|\chi_b\|_{\text{sp}}^2 (\|\chi\|_{\text{sp}} + \|\chi^{(J)}\|_{\text{sp}}) \int_0^\pi |\chi_\phi(0) - (\chi^{(J)})_\phi(0)| d\phi \quad (7.43)$$

$$+ \|\chi^{(J)}\|_{\text{sp}}^2 (\|\chi_b\|_{\text{sp}} + \|\chi_b^{(J)}\|_{\text{sp}}) \int_0^\pi |(\chi_b)_\phi(0) - (\chi_b^{(J)})_\phi(0)| d\phi. \quad (7.44)$$

Furthermore, since $\|\chi_s - \chi_s^{(J)}\|_1 = s2^{-J}$, note that

$$\int_0^\pi |(\chi_s)_\phi(0) - (\chi_s^{(J)})_\phi(0)| d\phi = \int_0^\pi |(\chi_s - \chi_s^{(J)})_\phi(0)| d\phi \leq \quad (7.45)$$

$$\leq \int_0^\pi \frac{d\phi}{\sqrt{\sin(\phi)}} \int_{\mathbb{R}} |(\chi_s - \chi_s^{(J)})(v)| dv = \frac{\sqrt{2}\pi^{\frac{3}{2}}s}{\Gamma(\frac{3}{4})^2} 2^{-J}. \quad (7.46)$$

By Remark 6.3, each of the norms is bounded above by $\|\chi_b\|_{\text{sp}}$ and we obtain

$$\left| D_{\text{rat}}(\chi, \chi_b) - D_{\text{rat}}(\chi^{(J)}, \chi_b^{(J)}) \right| \leq 2^5 \|\chi_b\|_{\text{sp}}^3 b 2^{-J}. \quad (7.47)$$

Combining (7.31) with (7.34), (7.35) and (7.47) gives the result. \square

7.2 Irrational case

Lemma 7.3. Fix $1 < \eta \leq 2$, $b \geq 1$ and let $K_L(b)$ be as in Proposition 6.5. Then for all $J \geq 4$, $R^2 > 64C_\eta K_L(b)^2 2^{2(\eta-1)J}$ and $2^{-\frac{J-1}{2}} < \delta < \frac{1}{2}$ we have

$$\mu_{\Gamma \setminus G}(|\Theta_\chi \overline{\Theta_{\chi_b}}| > R^2) - \mu(|\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}| > R^2(1-\delta)^2) = O\left(\frac{1}{R^6 \delta^3 2^{\frac{3}{2}J}}\right) \text{ and} \quad (7.48)$$

$$\mu_{\Gamma \setminus G}(|\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}| > R^2(1+\delta)^2) - \mu(|\Theta_\chi \overline{\Theta_{\chi_b}}| > R^2) = O\left(\frac{1}{R^6 \delta^3 2^{\frac{3}{2}J}}\right), \quad (7.49)$$

where the constants implied by the O -notation in (7.1)-(7.2) can both be taken to be $2^{22} \|\chi_{b,L}\|_{\text{sp}}^5$ (see (6.7)).

Proof. We follow the same strategy as the proof of Lemma 7.1 and we omit several details. Set $S = \{(k, \ell) \in \mathbb{Z}_{\geq 0}^2 : (k, \ell) \neq (0, 0)\}$. We use (5.25)–(5.26) and a union bound to obtain

$$\mu_{\Gamma \setminus G}(|\Theta_{\chi} \overline{\Theta_{\chi_b}}| > R^2) - \mu_{\Gamma \setminus G}(|\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}| > R^2(1-\delta)^2) \leq \quad (7.50)$$

$$\leq \sum_{(k, \ell) \in S} \mu_{\Gamma \setminus G} \left(2^{-\frac{(k+\ell)J}{2}} |\mathcal{L} \overline{\mathcal{L}}_{k, \ell}^{(J)}| > R^2 \frac{(1-\delta)^2}{4} \delta^{k+\ell} \right) + \quad (7.51)$$

$$+ \sum_{(k, \ell) \in S} \mu_{\Gamma \setminus G} \left(2^{-\frac{(k+\ell)J}{2}} |\mathcal{L} \overline{\mathcal{R}}_{k, \ell}^{(J)}| > R^2 \frac{(1-\delta)^2}{4} \delta^{k+\ell} \right) + \quad (7.52)$$

$$+ \sum_{(k, \ell) \in S} \mu_{\Gamma \setminus G} \left(2^{-\frac{(k+\ell)J}{2}} |\mathcal{R} \overline{\mathcal{L}}_{k, \ell}^{(J)}| > R^2 \frac{(1-\delta)^2}{4} \delta^{k+\ell} \right) + \quad (7.53)$$

$$+ \sum_{(k, \ell) \in S} \mu_{\Gamma \setminus G} \left(2^{-\frac{(k+\ell)J}{2}} |\mathcal{R} \overline{\mathcal{R}}_{k, \ell}^{(J)}| > R^2 \frac{(1-\delta)^2}{4} \delta^{k+\ell} \right). \quad (7.54)$$

Each term of the sum in (7.51) can be estimated as

$$\mu_{\Gamma \setminus G} \left(2^{-\frac{(k+\ell)J}{2}} |\mathcal{L} \overline{\mathcal{L}}_{k, \ell}^{(J)}| > R^2 \frac{(1-\delta)^2}{4} \delta^{k+\ell} \right) \leq \quad (7.55)$$

$$\leq \mu_{\Gamma \setminus G} \left\{ g \in \mathcal{F}_{\Gamma} : \left| \Theta_{\chi_L^{(J)}}(\tilde{g}) \right| > R^{\frac{1-\delta}{2}} (\delta 2^{\frac{J}{2}})^{\frac{k+\ell}{2}} \right\} + \quad (7.56)$$

$$+ \mu_{\Gamma \setminus G} \left\{ g \in \mathcal{F}_{\Gamma} : \left| \Theta_{\chi_{b,L}^{(J)}}(\tilde{g}) \right| > R^{\frac{1-\delta}{2}} (\delta 2^{\frac{J}{2}})^{\frac{k+\ell}{2}} \right\}. \quad (7.57)$$

In (7.56)–(7.57) we use \tilde{g} such that $\pi(\tilde{g}) = g$ as discussed in Section 5.3. To apply Lemma 4.3 to each of these terms we need

$$R^{\frac{1-\delta}{2}} (\delta 2^{\frac{J}{2}})^{\frac{k+\ell}{2}} > 2^{\frac{\eta}{2}} C_{\eta}^{\frac{1}{2}} \max\{\kappa_{\eta}(\chi_L^{(J)}), \kappa_{\eta}(\chi_{b,L}^{(J)})\}. \quad (7.58)$$

By Proposition 6.5, and since $\eta \leq 2$ this is implied by the assumption

$$R > 8C_{\eta}^{\frac{1}{2}} K_L(b) 2^{(\eta-1)J}. \quad (7.59)$$

Under this assumption, Lemma 4.3 gives

$$\mu_{\Gamma \setminus G} \left\{ g \in \mathcal{F}_{\Gamma} : \left| \Theta_{\chi_L^{(J)}}(\tilde{g}) \right| > R^{\frac{1-\delta}{2}} (\delta 2^{\frac{J}{2}})^{\frac{k+\ell}{2}} \right\} = \frac{2}{\pi^2} D_{\text{irr}}(\chi_L^{(J)}, \chi_L^{(J)}) \frac{1}{R^6 \frac{(1-\delta)^6}{64} (\delta^3 2^{\frac{3}{2}J})^{k+\ell}} (1 + \mathcal{E}), \quad (7.60)$$

where $|\mathcal{E}| \leq 7$. Note that $\delta < \frac{1}{2}$ implies $(1-\delta)^{-6} \leq 64$. Additionally, Lemma 6.2 and the unitarity of $f \mapsto f_{\phi}$ yields

$$D_{\text{irr}}(\chi_L^{(J)}, \chi_L^{(J)}) = \int_0^{\pi} \int_{-\infty}^{\infty} \left| \left(\chi_L^{(J)} \right)_{\phi}(w) \right|^6 dw d\phi \leq \|\chi_L^{(J)}\|_{\text{sp}}^4 \int_0^{\pi} \left\| \left(\chi_L^{(J)} \right)_{\phi} \right\|_2^2 d\phi = \quad (7.61)$$

$$= \pi \|\chi_L^{(J)}\|_{\text{sp}}^4 \|\chi_L^{(J)}\|_2^2. \quad (7.62)$$

Since $|\chi_L^{(J)}(w)| \leq 1$, we have $\|\chi_L^{(J)}\|_2^2 \leq \|\chi_L^{(J)}\|_1 \leq \frac{1}{2}\|\chi_L^{(J)}\|_{\text{sp}}$, and thus

$$\mu_{\Gamma \setminus G} \left\{ g \in \mathcal{F}_\Gamma : \left| \Theta_{\chi_L^{(J)}}(\tilde{g}) \right| > R^{\frac{1-\delta}{2}} (\delta 2^{\frac{J}{2}})^{\frac{k+\ell}{2}} \right\} < \frac{2^{14} \|\chi_L^{(J)}\|_{\text{sp}}^5}{R^6 (\delta^3 2^{\frac{3}{2}J})^{k+\ell}}. \quad (7.63)$$

An identical argument allows us to bound the quantity in (7.57) from above by $\frac{2^{14} \|\chi_{b,L}^{(J)}\|_{\text{sp}}^5}{R^6 (\delta^3 2^{\frac{3}{2}J})^{k+\ell}}$.

Now observe that for $q > 2^{\frac{3}{2}}$ we have $\sum_{(k,\ell) \in S} q^{-(k+\ell)} \leq \frac{4}{q}$. This bound can be used for $q = \delta^3 2^{\frac{3}{2}J}$ since $\delta > 2^{-\frac{J-1}{2}}$ by hypothesis. Therefore, the sum in (7.51) is bounded above by

$$\frac{2^{15} \|\chi_{b,L}^{(J)}\|_{\text{sp}}^5}{R^6} \sum_{(k,\ell) \in S} \frac{1}{(\delta^3 2^{\frac{3}{2}J})^{k+\ell}} \leq \frac{2^{17} \|\chi_{b,L}^{(J)}\|_{\text{sp}}^5}{R^6 \delta^3 2^{\frac{3}{2}J}}. \quad (7.64)$$

We can obtain the same upper bound for each of the sums in (7.52)–(7.54) and therefore obtain (7.48) with an implied constant equal to $2^{19} \|\chi_{b,L}^{(J)}\|_{\text{sp}}^5$. To prove (7.49), we can use a slightly different union bound as in the proof of (7.2). Combined with the upper bound $\|\chi_{b,L}^{(J)}\|_{\text{sp}} \leq \|\chi_{b,L}\|_{\text{sp}}$, this yields (7.49) with an implied constant equal to $2^{21} \|\chi_{b,L}\|_{\text{sp}}^5$. \square

Lemma 7.4. *Fix $1 < \eta \leq 2$, $b \geq 1$, and let $K(b)$ be as in Proposition 6.5. Suppose that $R^2 > 16C_\eta K(b)^2 2^{2(\eta-1)J}$ and $0 < \delta < \frac{1}{2}$. Then*

$$\begin{aligned} \mu_{\Gamma \setminus G} \left\{ |\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}| > R^2 (1 \pm \delta)^2 \right\} &= \\ &= \frac{2D_{\text{irr}}(\chi, \chi_b)}{\pi^2 R^6} (1 + O(\delta)) \left(1 + O_\eta(2^{2\eta(\eta-1)J} R^{-2\eta}) \right) \left(1 + O(2^{-\frac{J}{2}}) \right), \end{aligned} \quad (7.65)$$

where the constants implied by the O -notations can be taken to be 126 , $2^{11} C_\eta^2 K(b)^4$, and $8\sqrt{b} \|\chi_b\|_{\text{sp}}^5$ (see (6.5)) respectively.

Proof. We follow the strategy of Lemma 3.16 in [6], but keep track of all implied constants. The argument is similar to the proof of Lemma 7.2 and omit several details. In order to apply Lemma 4.3 we require

$$R^2 (1 \pm \delta)^2 > 2^\eta C_\eta \kappa_\eta(\chi^{(J)}) \kappa_\eta(\chi_b^{(J)}). \quad (7.66)$$

Applying Proposition 6.5, and using $\eta \leq 2$ and $\delta < \frac{1}{2}$, this is implied by the assumption

$$R^2 > 16C_\eta K(b)^2 2^{2(\eta-1)J}. \quad (7.67)$$

Applying Lemma 4.3 we obtain

$$\mu_{\Gamma \setminus G} \left\{ |\Theta_{\chi^{(J)}} \overline{\Theta_{\chi_b^{(J)}}}| > R^2 (1 \pm \delta)^2 \right\} = \frac{\frac{2}{\pi^2} D_{\text{irr}}(\chi^{(J)}, \chi_b^{(J)})}{R^6 (1 \pm \delta)^6} (1 + \mathcal{E}_1), \quad (7.68)$$

where

$$|\mathcal{E}_1| \leq 7(2^\eta C_\eta \kappa_\eta(\chi^{(J)}) \kappa_\eta(\chi_b^{(J)}))^\eta (R(1 \pm \delta))^{-2\eta} \quad (7.69)$$

$$\leq 7(2^{\eta+2} C_\eta K(b)^2)^\eta 2^{2\eta(\eta-1)J} R^{-2\eta} \quad (7.70)$$

$$\leq 2^{11} C_\eta^2 K(b)^4 \quad (7.71)$$

This yields the second implied constant in the statement of the Lemma. For the first implied constant, note that for $0 < \delta < \frac{1}{2}$ we have

$$\frac{1}{(1 \pm \delta)^6} \leq \frac{1}{(1 - \delta)^6} = (1 + \mathcal{E}_2), \quad |\mathcal{E}_2| \leq 126\delta. \quad (7.72)$$

For the third implied constant, note that

$$\left| D_{\text{irr}}(\chi, \chi_b) - D_{\text{irr}}(\chi^{(J)}, \chi_b^{(J)}) \right| \leq \quad (7.73)$$

$$\leq \int_0^\pi \left| \int_{-\infty}^\infty \left(|\chi_\phi(w)(\chi_b)_\phi(w)|^3 - |(\chi^{(J)})_\phi(w)(\chi_b^{(J)})_\phi(w)|^3 \right) dw \right| d\phi = \quad (7.74)$$

$$= \int_0^\pi \left| \left\| (\chi_\phi(\chi_b)_\phi)^3 \right\|_1 - \left\| ((\chi^{(J)})_\phi(\chi_b^{(J)})_\phi)^3 \right\|_1 \right| d\phi \leq \quad (7.75)$$

$$\leq \int_0^\pi \left\| (\chi_\phi(\chi_b)_\phi)^3 - ((\chi^{(J)})_\phi(\chi_b^{(J)})_\phi)^3 \right\|_1 d\phi, \quad (7.76)$$

where in (7.76) we have used the reverse triangle inequality for the 1-norm. Now, using the identity $x^3 - y^3 = (x - y)(x^2 + xy + y^2)$ and the bound

$$\begin{aligned} & \left| (\chi_\phi(w)(\chi_b)_\phi(w))^2 + \chi_\phi(w)(\chi_b)_\phi(w)(\chi^{(J)})_\phi(w)(\chi_b^{(J)})_\phi(w) + ((\chi^{(J)})_\phi(w)(\chi_b^{(J)})_\phi(w))^2 \right| \leq \\ & \leq 3\|\chi_b\|_{\text{sp}}^4 \end{aligned} \quad (7.77)$$

coming from Lemma 6.2 and Remark 6.3, we see that (7.76) implies

$$\left| D_{\text{irr}}(\chi, \chi_b) - D_{\text{irr}}(\chi^{(J)}, \chi_b^{(J)}) \right| \leq \quad (7.78)$$

$$\leq 3\|\chi_b\|_{\text{sp}}^4 \int_0^\pi \left\| \chi_\phi(\chi_b)_\phi - (\chi^{(J)})_\phi(\chi_b^{(J)})_\phi \right\|_1 d\phi. \quad (7.79)$$

Now we add and subtract $\chi_\phi(\chi_b^{(J)})_\phi$ in (7.79) and use the triangle inequality, followed by the Cauchy-Schwarz inequality, to obtain

$$\left| D_{\text{irr}}(\chi, \chi_b) - D_{\text{irr}}(\chi^{(J)}, \chi_b^{(J)}) \right| \leq \quad (7.80)$$

$$\leq 3\|\chi_b\|_{\text{sp}}^4 \int_0^\pi \left(\|\chi_\phi\|_2 \|(\chi_b)_\phi - (\chi_b^{(J)})_\phi\|_2 + \|(\chi_b^{(J)})_\phi\|_2 \|\chi_\phi - (\chi^{(J)})_\phi\|_2 \right) d\phi. \quad (7.81)$$

The unitarity of the Shale-Weil representation and the bounds $\|\chi_\phi\|_2^2 \leq \frac{1}{2}\|\chi_b\|_{\text{sp}}, \|(\chi_b^{(J)})_\phi\|_2^2 \leq \frac{1}{2}\|\chi_b\|_{\text{sp}}$, and $\|\chi_s - \chi_s^{(J)}\|_2 \leq 3^{-\frac{1}{2}}2^{-\frac{J}{2}}\sqrt{s}$ imply¹

$$\left| D_{\text{irr}}(\chi, \chi_b) - D_{\text{irr}}(\chi^{(J)}, \chi_b^{(J)}) \right| \leq 8\sqrt{b}\|\chi_b\|_{\text{sp}}^5 2^{-\frac{J}{2}}. \quad (7.82)$$

Combining (7.68) with (7.71), (7.72) and (7.82) gives the result. \square

8 Proof of the main theorems

In this section, we combine the tail bounds in Lemmata 4.2 and 4.3 with the tail approximations from Section 7 to prove Theorems 8.3 and 8.6, which are analogue to Theorem 1.7 but hold for $(f_1, f_2) = (\chi, \chi_b)$ and have explicit, η -dependent power savings, namely $O_\eta(R^{-\frac{2\eta}{6\eta(\eta-1)+1}})$ and $O_\eta(R^{-\frac{6\eta}{16\eta(\eta-1)+3}})$, respectively. We then take $\eta \rightarrow 1^+$ to obtain Theorems 8.4 and 8.7, which are explicit versions of Theorems 1.5 and 1.6, respectively.

8.1 Rational case

It is convenient to express $D_{\text{rat}}(\chi, \chi_b)$ from Lemma 7.2 in closed form. Using the definition (4.10) we have

$$D_{\text{rat}}(\chi, \chi_b) = \int_0^\pi |\chi_\phi(0)|^2 |(\chi_b)_\phi(0)|^2 d\phi. \quad (8.1)$$

Theorem 8.1. *Let $b \geq 1$. Then*

$$D_{\text{rat}}(\chi, \chi_b) = \begin{cases} 2 \log 2 & \text{if } b = 1; \\ 2b \coth^{-1}(b) + \frac{1}{2} \log(b^2 - 1) + \frac{b^2}{2} \log(1 - \frac{1}{b^2}) & \text{if } b > 1. \end{cases} \quad (8.2)$$

Proof. Note that for $0 < \phi < \pi$ we have

$$|f_\phi(0)|^2 = \frac{1}{\sin(\phi)} \left| \int_{-\infty}^{\infty} e\left(v^2 \frac{\cot(\phi)}{2}\right) f(v) dv \right|^2 \quad (8.3)$$

and the change of variables $x = \frac{\cot(\phi)}{2}$ in (8.1) yields

$$D_{\text{rat}}(\chi, \chi_b) = 2 \int_{-\infty}^{\infty} \left| \int_0^1 e(v^2 x) dv \right|^2 \left| \int_0^b e(w^2 x) dw \right|^2 dx. \quad (8.4)$$

¹Note that $\|\chi_s - \chi_s^{(J)}\|_2$ is erroneously claimed to be $O(2^{-J})$ in [6] but this does not affect the proof of Theorem 3.13 therein. In the same way, if we replaced $O(2^{-\frac{J}{2}})$ in (7.65) by $O(2^{-J})$, we would get the same optimal power saving in Theorem 8.6, see β^* in (8.67).

By performing the three successive changes of variables $w' = w/b$, $x' = bx$, and $v' = v/b$, we obtain

$$D_{\text{rat}}(\chi, \chi_b) = b^2 D_{\text{rat}}(\chi, \chi_{1/b}) \quad (8.5)$$

Therefore it is enough to compute $D_{\text{rat}}(\chi, \chi_c)$ with $0 < c \leq 1$. Observing that the integrand is even with respect to the variable x , we can rewrite it as

$$D_{\text{rat}}(\chi, \chi_c) = 4 \int_0^\infty p(x) q\left(\frac{1}{x}\right) \frac{dx}{x}, \quad (8.6)$$

where

$$p(x) = \int_0^1 \int_0^1 e(x(v_1^2 - v_2^2)) dv_1 dv_2 \quad \text{and} \quad q(x) = \frac{1}{x} \int_0^c \int_0^c e\left(\frac{1}{x}(w_1^2 - w_2^2)\right) dw_1 dw_2. \quad (8.7)$$

The integral in (8.6) can be seen as a Mellin convolution (evaluated at $z = 1$, see also (8.13)) over the multiplicative group $\mathbb{R}_{>0}$. Let us compute the Mellin transforms of p and q respectively. Let $0 < \text{Re}(s) < 1$. We use Fubini's theorem to change the order of integration and integrate with respect to x first and then with respect to the two remaining variables ($dv_1 dv_2$ and $dw_1 dw_2$ respectively). We obtain

$$P(s) = \int_0^\infty x^{s-1} p(x) dx = \int_0^1 \int_0^1 \int_0^\infty x^{s-1} e(x(v_1^2 - v_2^2)) dx dv_1 dv_2 = \quad (8.8)$$

$$= \frac{\pi^{\frac{3}{2}-s} \csc\left(\frac{\pi s}{2}\right)}{2^{s+2}(1-s) \Gamma\left(\frac{3}{2}-s\right)} \quad (8.9)$$

and

$$Q(s) = \int_0^\infty x^{s-1} q(x) dx = \int_0^c \int_0^c \int_0^\infty x^{s-2} e(x^{-1}(w_1^2 - w_2^2)) dx dw_1 dw_2 = \quad (8.10)$$

$$= \frac{c^{2s} \pi^{\frac{1}{2}+s} \sec\left(\frac{\pi s}{2}\right)}{2^{3-s} s \Gamma\left(\frac{1}{2}+s\right)}. \quad (8.11)$$

We use the double angle formula for sine and the identity $\Gamma\left(\frac{3}{2}-s\right) \Gamma\left(\frac{1}{2}+s\right) = \frac{1}{2}\pi(1-2s) \sec(\pi s)$ to simplify

$$P(s)Q(s) = \frac{\pi c^{2s}}{8} \frac{\cot(\pi s)}{s(s-1)(2s-1)}. \quad (8.12)$$

Now we apply the Mellin inversion formula

$$\int_0^\infty p(x) q\left(\frac{z}{x}\right) \frac{dx}{x} = \int_{\frac{1}{2}-i\infty}^{\frac{1}{2}+i\infty} P(s)Q(s) z^{-s} ds \quad (8.13)$$

in the particular case $z = 1$. Note that for $s \rightarrow \frac{1}{2}$ we have $\cot(\pi s) = -\pi \left(s - \frac{1}{2}\right) + O((s - \frac{1}{2})^3)$ and therefore $s = \frac{1}{2}$ is not a pole of $P(s)Q(s)$. The poles of $\cot(\pi s)$ are the integers and $\text{Res}(\cot(\pi s), s = n) = \frac{1}{\pi}$ for every $n \in \mathbb{Z}$. Therefore

$$\text{Res}(P(s)Q(s), s = n) = \begin{cases} \frac{-3c^2 + 2c^2 \log(c)}{8} & \text{if } n = 1, \\ \frac{c^{2n}}{8n(n-1)(2n-1)} = \frac{c^{2n}}{8} \left(\frac{1}{n} - \frac{2}{n-\frac{1}{2}} + \frac{1}{n-1} \right) & \text{if } n \geq 2. \end{cases} \quad (8.14)$$

By the residue's theorem, we have

$$D_{\text{rat}}(\chi, \chi_c) = \frac{4}{2\pi i} \int_{\frac{1}{2}-i\infty}^{\frac{1}{2}+i\infty} P(s)Q(s)ds = -4 \sum_{n \geq 1} \text{Res}(P(s)Q(s), s = n) = \quad (8.15)$$

$$= -\frac{1}{2} \left(-3c^2 + 2c^2 \log(c) + \sum_{n=2}^{\infty} c^{2n} \left(\frac{1}{n} - \frac{2}{n-\frac{1}{2}} + \frac{1}{n-1} \right) \right). \quad (8.16)$$

Now, if $c = 1$, we can use the digamma function $\psi(x) = \frac{\Gamma'(x)}{\Gamma(x)}$, which satisfies $\psi(x+1) = \psi(x) + 1/x$, to write

$$\sum_{n=2}^N \left(\frac{1}{n} + \frac{1}{n-1} \right) = 2\psi(N) + 2\gamma - 1 + \frac{1}{N} \quad \text{and} \quad \sum_{n=2}^N \frac{1}{n-\frac{1}{2}} = \psi\left(N + \frac{1}{2}\right) - \psi\left(\frac{3}{2}\right). \quad (8.17)$$

Since $\psi(N + \frac{1}{2}) - \psi(N) = \frac{1}{2N} + O(N^{-2})$ as $N \rightarrow \infty$ and $\psi(\frac{1}{2}) = -2\log 2 - \gamma$, we have

$$D_{\text{rat}}(\chi, \chi) = -\frac{1}{2} \left(-3 + \lim_{N \rightarrow \infty} \sum_{n=2}^N \left(\frac{1}{n} - \frac{2}{n-\frac{1}{2}} + \frac{1}{n-1} \right) \right) = \quad (8.18)$$

$$= -\frac{1}{2} \left(-4 + 2\gamma + 2\psi\left(\frac{3}{2}\right) \right) = 2\log 2. \quad (8.19)$$

This proves the first part of (8.2). On the other hand, for $0 < c < 1$, we can use the series

$$-\log(1-x) = \sum_{k=1}^{\infty} \frac{x^k}{k} \quad \text{and} \quad \tanh^{-1}(x) = \sum_{k=0}^{\infty} \frac{x^{2k+1}}{2k+1} \quad (8.20)$$

to simplify

$$D_{\text{rat}}(\chi, \chi_c) = -\frac{1}{2} \left(-3c^2 + 2c^2 \log(c) + \sum_{n=2}^{\infty} c^{2n} \left(\frac{1}{n} - \frac{2}{n - \frac{1}{2}} + \frac{1}{n-1} \right) \right) = \quad (8.21)$$

$$= -\frac{1}{2} \left(-3c^2 + 2c^2 \log(c) - c^2 - \log(1 - c^2) + 4c^2 - 4c \tanh^{-1}(c) - c^2 \log(1 - c^2) \right) = \quad (8.22)$$

$$= 2c \tanh^{-1}(c) - c^2 \log(c) + \frac{1 + c^2}{2} \log(1 - c^2). \quad (8.23)$$

Using (8.23) with $c = b^{-1}$ and (8.5), we obtain the second part of (8.2). \square

Remark 8.2. The function $b \mapsto D_{\text{rat}}(\chi, \chi_b)$ is once but not twice continuously differentiable at $b = 1$. Its first derivative at 1 equals $2 \log 2$, while its second derivative equals $\log(1 - b^{-2})$ for $b > 1$. Moreover, $D_{\text{rat}}(\chi, \chi_b) = \frac{3+2\log b}{2} + O(b^{-2})$ as $b \rightarrow \infty$.

Theorem 8.3. Let λ be a probability measure on \mathbb{R} which is absolutely continuous with respect to the Lebesgue measure. Let $1 < \eta \leq 2$ and $b \geq 1$. Set $R_0^{\text{rat}} = R_0^{\text{rat}}(b, \eta)$ as in (8.45). Then for all $R > R_0^{\text{rat}}$ we have

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{N} S_N(x; 0, 0) \overline{S_{[bN]}(x; 0, 0)} \right| > R^2 \right\} = \frac{2D_{\text{rat}}(\chi, \chi_b)}{\pi^2 R^4} \left(1 + O_{\eta} \left(R^{-\frac{2\eta}{6\eta(\eta-1)+1}} \right) \right), \quad (8.24)$$

where the constant implied by the O_{η} -notation can be taken to be as in (8.46)

Proof. Corollary 3.11 gives

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{N} S_N(x; 0, 0) \overline{S_{[bN]}(x; 0, 0)} \right| > R^2 \right\} = \mu_{\Gamma \setminus G}^0 (|\Theta_{\chi} \overline{\Theta_{\chi b}}| > R^2). \quad (8.25)$$

Let us now write $J = \alpha \log_2(R)$ and $\delta = R^{-\beta}$, for some $\alpha, \beta > 0$. We then rewrite the assumptions of Lemma 7.1 as

$$R^2 > 64K_L(b)^2 C_{\eta} R^{2(\eta-1)\alpha}, \quad \sqrt{2} R^{-\frac{\alpha}{2}} < R^{-\beta} < \frac{1}{2}, \quad R^{\alpha} \geq 16, \quad (8.26)$$

and those of Lemma 7.2 as

$$R^2 > 16K(b)^2 C_{\eta} R^{2(\eta-1)\alpha}, \quad 0 < R^{-\beta} < \frac{1}{2}, \quad R^{\alpha} \geq 2. \quad (8.27)$$

In order for the first inequality of (8.26) to hold for all sufficiently large R , we need $(\eta - 1)\alpha < 1$. The same is true for the first inequality of (8.27). Similarly, the second inequality of (8.26) implies $\beta < \frac{\alpha}{2}$. By the lower bound in Remark 6.6, we have $K_L(b)^2 \leq \frac{1}{2} K(b)^2$. Therefore if we assume that

$$R^2 > 32K(b)^2 C_{\eta} R^{2(\eta-1)\alpha}, \quad R^{\alpha} > 16, \quad R^{\beta} > 2, \quad R^{\frac{\alpha}{2}-\beta} > \sqrt{2}, \quad (8.28)$$

then (8.26) and (8.27) hold and we can apply both Lemma 7.1 and Lemma 7.2. We obtain

$$\mu_{\Gamma \setminus G}^0(|\Theta_\chi \overline{\Theta_{\chi_b}}| > R^2) = \quad (8.29)$$

$$= \frac{2D_{\text{rat}}(\chi, \chi_b)}{\pi^2 R^4} (1 + O(R^{-\beta})) (1 + O_\eta(R^{-2\eta+2\eta(\eta-1)\alpha})) (1 + O(R^{-\alpha})) + O(R^{-\alpha+2\beta-4}), \quad (8.30)$$

where $D_{\text{rat}}(\chi, \chi_b)$ is given explicitly in (8.2). Now we choose R sufficiently large so that, in addition to satisfying (8.28), each of the first three O -terms in (8.30) is bounded above in absolute value by 1. This means, recalling the implicit constants in Lemma 7.2,

$$R > \max \left\{ (32K(b)^2 C_\eta)^{\frac{1}{2-2(\eta-1)\alpha}}, 16^{\frac{1}{\alpha}}, 2^{\frac{1}{\beta}}, 2^{\frac{1}{\alpha-2\beta}}, 30^{\frac{1}{\beta}}, \right. \quad (8.31)$$

$$\left. (32K(b)^2 C_\eta)^{\frac{1}{\eta-\eta(\eta-1)\alpha}}, (32b\|\chi_b\|_{\text{sp}}^3)^{\frac{1}{\alpha}} \right\}. \quad (8.32)$$

Observing that $\frac{1}{2-2(\eta-1)\alpha} \leq \frac{1}{\eta-\eta(\eta-1)\alpha}$ for $1 < \eta \leq 2$ and $0 < \alpha < \frac{1}{\eta-1}$ and that, by (6.5), $32b\|\chi_b\|_{\text{sp}}^3 \geq 32 \cdot 9^2 \geq 16$, we can rewrite the above condition on R as

$$R > R_0^{\text{rat}}(\alpha, \beta; b, \eta) := \max \left\{ (32K(b)^2 C_\eta)^{\frac{1}{\eta-\eta(\eta-1)\alpha}}, 30^{\frac{1}{\beta}}, 2^{\frac{1}{\alpha-2\beta}}, (32b\|\chi_b\|_{\text{sp}}^3)^{\frac{1}{\alpha}} \right\}. \quad (8.33)$$

Under the assumption (8.33), combining the contribution of the error terms in (8.30), we can write²

$$\mu_{\Gamma \setminus G}^0(|\Theta_\chi \overline{\Theta_{\chi_b}}| > R^2) = \frac{2D_{\text{rat}}(\chi, \chi_b)}{\pi^2 R^4} (1 + O_\eta(R^{-\beta} + R^{-2\eta+2\eta(\eta-1)\alpha} + R^{-\alpha} + R^{-\alpha+2\beta})), \quad (8.34)$$

where the implied constant in the O_η -notation in (8.34) can be taken as

$$\max \left\{ 120, 2^{10}K(b)^4 C_\eta^2, 2^6 b \|\chi_b\|_{\text{sp}}^3, \frac{2^{17} \|\chi_{b,L}\|_{\text{sp}}^4}{\frac{2}{\pi^2} D_{\text{rat}}(\chi, \chi_b)} \right\}. \quad (8.35)$$

Using the trivial bounds $K(b) \geq 2^5$ and $C_\eta \geq 2^6$, we see that $2^{10}K(b)^4 C_\eta^2 \geq 2^{42} \geq 120$ and hence the first term in 8.35 can be dropped. Using the explicit formulæ (6.28) and (6.5) for $K(b)$ and $\|\chi_b\|_{\text{sp}}$ respectively, we can verify that for $b \geq 1$ the bound $K(b)^4 \geq 2^{14}b\|\chi_b\|_{\text{sp}}^3$ holds. Therefore $2^{10}K(b)^4 C_\eta^2 \geq 2^{22}K(b)^4 \geq 2^{36}b\|\chi_b\|_{\text{sp}}^3$ and hence the third term in 8.35 can be dropped. Finally, using the explicit formula (6.7) for $\|\chi_{b,L}\|_{\text{sp}}$, we can check that the bound $K(b)^4 \geq 2^{10}\|\chi_{b,L}\|_{\text{sp}}^4$ holds for every $b \geq 1$. This fact, along with the lower bound $D_{\text{rat}}(\chi, \chi_b) \geq 2 \log(2)$ (see Theorem 8.1), yield $2^{10}K(b)^4 C_\eta^2 \geq 2^{22}K(b)^4 \geq 2^{32}\|\chi_{b,L}\|_{\text{sp}}^4 \geq \frac{\pi^2 2^{16}}{2 \log(2)} \|\chi_{b,L}\|_{\text{sp}}^4 \geq \pi^2 2^{16} \frac{\|\chi_{b,L}\|_{\text{sp}}^4}{D_{\text{rat}}(\chi, \chi_b)}$ and hence the

²The transition from (8.30) to (8.34) uses the fact that $A(1+B)(1+C)(1+D) + E \leq A(1+4B+C+2D+\frac{E}{A})$ provided $A, E > 0$ and $0 < B, C, D < 1$.

last term in 8.35 can also be dropped. We have shown that the implied constant in the O_η -notation in (8.34) can be taken as

$$P_0^{\text{rat}}(b, \eta) = 2^{10} K(b)^4 C_\eta^2. \quad (8.36)$$

The best power saving in (8.34) comes from choosing (α^*, β^*) in the region $\{(\alpha, \beta) : 0 < \alpha < \frac{1}{\eta-1}, 0 < \beta < \frac{\alpha}{2}\}$ in order to maximise $m(\alpha, \beta) = \min\{\alpha, \beta, 2\eta - 2\eta(\eta-1)\alpha, \alpha - 2\beta\} = \min\{\beta, 2\eta - 2\eta(\eta-1)\alpha, \alpha - 2\beta\}$. That is

$$(\alpha^*, \beta^*) = \left(\frac{6\eta}{6\eta(\eta-1) + 1}, \frac{2\eta}{6\eta(\eta-1) + 1} \right), \quad (8.37)$$

at which $m(\alpha^*, \beta^*) = \beta^*$. Therefore, combining (8.25), (8.30), (8.33), (8.34), and (8.37), we obtain that for every $R > R_0^{\text{rat}}(\alpha^*, \beta^*; b, \eta)$ we have

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{N} S_N(x; 0, 0) \overline{S_{[bN]}(x; 0, 0)} \right| > R^2 \right\} = \frac{2D_{\text{rat}}(\chi, \chi_b)}{\pi^2 R^4} \left(1 + O_\eta \left(R^{-\frac{2\eta}{6\eta(\eta-1)+1}} \right) \right), \quad (8.38)$$

where the implicit constant in the O_η -notation in (8.38) can be taken as $4P_0^{\text{rat}}(b, \eta)$. Note that

$$\frac{1}{\eta - \eta(\eta-1)\alpha^*} = \frac{1}{\eta} + 6(\eta-1), \quad (8.39)$$

$$\frac{1}{\beta^*} = \frac{1}{\alpha^* - 2\beta^*} = \frac{1}{2\eta} + 3(\eta-1), \quad (8.40)$$

$$\frac{1}{\alpha^*} = \frac{1}{6\eta} + (\eta-1). \quad (8.41)$$

We claim that

$$R_0^{\text{rat}}(\alpha^*, \beta^*, \eta) = (32K(b)^2 C_\eta)^{\frac{1}{\eta} + 6(\eta-1)}. \quad (8.42)$$

From (8.40) we immediately have $30^{\frac{1}{\beta^*}} \geq 2^{\frac{1}{\alpha^* - 2\beta^*}}$. Then we see by (8.39)-(8.40) that

$$(32K(b)^2 C_\eta)^{\frac{1}{\eta - \eta(\eta-1)\alpha^*}} = (2^{10} K(b)^4 C_\eta^2)^{\frac{1}{2\eta} + 3(\eta-1)} \geq 30^{\frac{1}{2\eta} + 3(\eta-1)} = 30^{\frac{1}{\beta^*}} \quad (8.43)$$

because $2^{10} K(b)^4 C_\eta^2 \geq 2^{10} K(1)^4 2^{12} \geq 2^{45} \geq 30$. Finally, (8.41) and the explicit bound $K(b)^2 \geq 2^{10} b^{\frac{1}{6}} \|\chi_b\|_{\text{sp}}^{\frac{1}{2}}$ imply

$$(32K(b)^2 C_\eta)^{\frac{1}{\eta - \eta(\eta-1)\alpha^*}} = (32K(b)^2 C_\eta)^{\frac{1}{\eta} + 6(\eta-1)} \geq (2^{\frac{5}{6}} b^{\frac{1}{6}} \|\chi_b\|_{\text{sp}}^{\frac{1}{2}})^{\frac{1}{\eta} + 6(\eta-1)} = (32b \|\chi_b\|_{\text{sp}}^3)^{\frac{1}{\alpha^*}} \quad (8.44)$$

since $2^5 K(b)^2 C_\eta \geq 2^{11} K(b)^2 \geq 2^{21} b^{\frac{1}{6}} \|\chi_b\|_{\text{sp}}^{\frac{1}{2}} \geq 2^{\frac{5}{6}} b^{\frac{1}{6}} \|\chi_b\|_{\text{sp}}^{\frac{1}{2}}$. This concludes the proof of the claim (8.42). Now, if we set

$$R_0^{\text{rat}}(b, \eta) := (32K(b)^2 C_\eta)^{\frac{1}{\eta} + 6(\eta-1)}, \quad (8.45)$$

then for every $R > R_0^{\text{rat}}(b, \eta)$ we have (8.38) with an implicit constant of

$$P^{\text{rat}}(b, \eta) := 2^{12} K(b)^4 C_\eta^2 \quad (8.46)$$

and the theorem is proven. \square

Theorem 8.4 (Explicit version of Theorem 1.5). *Let λ be a probability measure on \mathbb{R} which is absolutely continuous with respect to the Lebesgue measure. Let $b \geq 1$ and $0 < \varepsilon \leq 1$, and set*

$$R^{\text{rat}}(b, \varepsilon) = 2^{38} K(b)^4 \varepsilon^{-2} \quad (8.47)$$

where $K(b)$ is given in (6.28). Then for all $R > R^{\text{rat}}(b, \varepsilon)$ we have

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{N} S_N(x; 0, 0) \overline{S_{\lfloor bN \rfloor}(x; 0, 0)} \right| > R^2 \right\} = \frac{2D_{\text{rat}}(b)}{\pi^2 R^4} (1 + O_\varepsilon(R^{-2+\varepsilon})), \quad (8.48)$$

where $D_{\text{rat}}(b) = D_{\text{rat}}(\chi, \chi_b)$ is given in (8.2), and the constant implied by the O_ε -notation in (8.48) can be taken to be

$$2^{42} K(b)^4 \varepsilon^{-4}. \quad (8.49)$$

Proof. Note that the exponent in (8.24) satisfies

$$\lim_{\eta \rightarrow 1^+} \frac{2\eta}{6\eta(\eta-1)+1} = 2^-.$$

This means that the closer $\eta > 1$ is to 1, the better the power saving in Theorem 8.3 is. If we set $\frac{2\eta}{6\eta(\eta-1)+1} = 2 - \varepsilon$ for some $\varepsilon > 0$, then $\varepsilon \mapsto \eta(\varepsilon) = \frac{7-3\varepsilon+\sqrt{3\varepsilon^2-18\varepsilon+25}}{6(2-\varepsilon)}$ maps $(0, 1]$ to $(1, \frac{4+\sqrt{10}}{6}]$ monotonically. Considering the Taylor expansion of $\eta(\varepsilon)$ near $\varepsilon = 0$ we find the lower bound $\eta(\varepsilon) > 1 + \frac{\varepsilon}{10}$.

We aim to find a simple upper bound for $R_0^{\text{rat}}(b, \eta(\varepsilon))$. If the function $\eta \mapsto R_0^{\text{rat}}(b, \eta)$ were decreasing, then we could get an upper bound by replacing $R_0^{\text{rat}}(b, \eta(\varepsilon))$ with $R_0^{\text{rat}}(b, 1 + \frac{\varepsilon}{10})$. However, we only have monotonicity near $\eta = 1$ and not on the whole interval $(1, \frac{4+\sqrt{10}}{6}]$. To address this issue, let us find an upper bound for $R_0^{\text{rat}}(b, \eta)$ which is decreasing as a function of η in the desired range. For the rest of this proof we will assume that $1 < \eta \leq \frac{4+\sqrt{10}}{6}$. We have

$$\zeta(\eta) \leq \frac{\sqrt{2}}{\eta-1} \quad (8.50)$$

(see Corollary A.4 in Appendix A). Recall that $C_\eta = 2^{6\eta} \zeta(\eta)^2$. We use (8.45), (8.50), the fact that $\frac{1}{\eta} + 6(\eta-1) \leq 2$, and the bound $(\eta-1)^{-2(\frac{1}{\eta}+6(\eta-1))} \leq 2^5(\eta-1)^{-2}$ to obtain

$$R_0^{\text{rat}}(b, \eta) \leq \left(\frac{2^{6+6\eta} K(b)^2}{(\eta-1)^2} \right)^{\frac{1}{\eta}+6(\eta-1)} \leq \frac{2^{20+2\sqrt{10}} K(b)^4}{(\eta-1)^{2(\frac{1}{\eta}+6(\eta-1))}} \leq \frac{2^{25+2\sqrt{10}} K(b)^4}{(\eta-1)^2}. \quad (8.51)$$

Therefore, since the right-most upper bound in (8.51) is decreasing in η ,

$$R_0^{\text{rat}}(b, \eta(\varepsilon)) \leq \frac{2^{25+2\sqrt{10}} K(b)^4}{(\frac{\varepsilon}{10})^2} \leq 2^{38} K(b)^4 \varepsilon^{-2} =: R^{\text{rat}}(b, \varepsilon). \quad (8.52)$$

Therefore, if $R > R^{\text{rat}}(b, \varepsilon)$, then by Theorem 8.3 we obtain

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{N} S_N(x; 0, 0) \overline{S_{[bN]}(x; 0, 0)} \right| > R^2 \right\} = \frac{2D_{\text{rat}}(\chi, \chi_b)}{\pi^2 R^4} (1 + O_\varepsilon(R^{-2+\varepsilon})). \quad (8.53)$$

The constant implied by the O_ε -notation in (8.53) is given by $P^{\text{rat}}(b, \eta(\varepsilon))$ using (8.46) and can be bounded above as follows:

$$P^{\text{rat}}(b, \eta(\varepsilon)) \leq 2^{22+2\sqrt{10}} K(b)^4 (\frac{\varepsilon}{10})^{-4} \leq 2^{42} K(b)^4 \varepsilon^{-4}. \quad (8.54)$$

□

By taking $b = 1$ in Theorem 8.4 and bounding $K(1)^4 = 59^4 \leq 2^{24}$, we obtain the following

Corollary 8.5 (Explicit version of Theorem 1.1). *Let λ be a probability measure on \mathbb{R} which is absolutely continuous with respect to the Lebesgue measure. Let $0 < \varepsilon \leq 1$ and set $R^{\text{rat}}(\varepsilon) = 2^{62} \varepsilon^{-2}$. Then for all $R > R^{\text{rat}}(\varepsilon)$ we have*

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{\sqrt{N}} S_N(x; 0, 0) \right| > R \right\} = \frac{4 \log 2}{\pi^2 R^4} (1 + O_\varepsilon(R^{-2+\varepsilon})), \quad (8.55)$$

the constant implied by the O_ε -notation in (8.55) can be taken to be $2^{66} \varepsilon^{-4}$.

8.2 Irrational case

Theorem 8.6. *Let λ be a probability measure on \mathbb{R} which is absolutely continuous with respect to the Lebesgue measure. Let $(c, \alpha) \notin \mathbb{Q}^2$, $1 < \eta \leq 2$, and $b \geq 1$. Set $R_0^{\text{irr}} = R_0^{\text{irr}}(b, \eta)$ as in (8.77). Then for all $R > R_0^{\text{irr}}$ we have*

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{N} S_N(x; c, \alpha) \overline{S_{[bN]}(x; c, \alpha)} \right| > R^2 \right\} = \frac{2D_{\text{irr}}(\chi, \chi_b)}{\pi^2 R^6} \left(1 + O_\eta \left(R^{-\frac{6\eta}{16\eta(\eta-1)+3}} \right) \right), \quad (8.56)$$

where

$$D_{\text{irr}}(\chi, \chi_b) = \int_{-\infty}^{\infty} \int_0^\pi |\chi_\phi(w) (\chi_b)_\phi(w)|^3 d\phi dw \quad (8.57)$$

and the constant implied by the O_η -notation can be taken to be as in (8.78)

Proof. Corollary 3.12 gives

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{N} S_N(x; c, \alpha) \overline{S_{[bN]}(x; c, \alpha)} \right| > R^2 \right\} = \mu_{\Gamma \setminus G} (|\Theta_\chi \overline{\Theta_{\chi_b}}| > R^2). \quad (8.58)$$

We argue as in the proof of Theorem 8.3 and write $J = \alpha \log_2(R)$, and $\delta = R^{-\beta}$, for some $\alpha, \beta > 0$. The assumptions of Lemmata 7.3 and 7.4 read as (8.26) and (8.27) respectively, i.e. they are the same as in the rational case. Therefore, if we assume (8.28), then we can apply both Lemma 7.3 and Lemma 7.4.

$$\mu_{\Gamma \setminus G} (|\Theta_\chi \overline{\Theta_{\chi_b}}| > R^2) = \quad (8.59)$$

$$= \frac{2D_{\text{irr}}(\chi, \chi_b)}{\pi^2 R^6} (1 + O(R^{-\beta})) (1 + O_\eta(R^{-2\eta+2\eta(\eta-1)\alpha})) (1 + O(R^{-\frac{\alpha}{2}})) + O\left(R^{-\frac{3}{2}\alpha+3\beta-6}\right). \quad (8.60)$$

Now we choose R sufficiently large so that, in addition to satisfying (8.28), each of the first three O -terms in (8.60) is bounded above in absolute value by 1. This means, recalling the implicit constants in Lemma 7.4,

$$R > \max \left\{ (32K(b)^2 C_\eta)^{\frac{1}{2-2(\eta-1)\alpha}}, 16^{\frac{1}{\alpha}}, 2^{\frac{1}{\beta}}, 2^{\frac{1}{\alpha-2\beta}}, 126^{\frac{1}{\beta}}, \right. \quad (8.61)$$

$$\left. (2^{\frac{11}{2}} K(b)^2 C_\eta)^{\frac{1}{\eta-\eta(\eta-1)\alpha}}, (2^6 b \|\chi_b\|_{\text{sp}}^{10})^{\frac{1}{\alpha}} \right\} \quad (8.62)$$

$$= \max \left\{ (2^{\frac{11}{2}} K(b)^2 C_\eta)^{\frac{1}{\eta-\eta(\eta-1)\alpha}}, 126^{\frac{1}{\beta}}, 2^{\frac{1}{\alpha-2\beta}}, (2^6 b \|\chi_b\|_{\text{sp}}^{10})^{\frac{1}{\alpha}} \right\} =: R_0^{\text{irr}}(\alpha, \beta; b, \eta). \quad (8.63)$$

Under the assumption $R > R_0^{\text{irr}}(\alpha, \beta; b, \eta)$, combining the contribution of the error terms in (8.60), we can write

$$\mu_{\Gamma \setminus G} (|\Theta_\chi \overline{\Theta_{\chi_b}}| > R^2) = \frac{2D_{\text{irr}}(\chi, \chi_b)}{\pi^2 R^6} \left(1 + O_\eta \left(R^{-\beta} + R^{-2\eta+2\eta(\eta-1)\alpha} + R^{-\frac{\alpha}{2}} + R^{-\frac{3}{2}\alpha+3\beta} \right) \right), \quad (8.64)$$

where the implied constant in the O_η -notation in (8.34) can be taken as

$$\max \left\{ 504, 2^{11} K(b)^4 C_\eta^2, 16\sqrt{b} \|\chi_b\|_{\text{sp}}^5, \frac{2^{22} \|\chi_{b,L}\|_{\text{sp}}^5}{\pi^2 D_{\text{irr}}(\chi, \chi_b)} \right\}. \quad (8.65)$$

Since $2^{11} K(b)^4 C_\eta^2 \geq 2^{46} \geq 504$, the first term in (8.65) can be dropped. Since $K(b)^4 \geq 2^7 \sqrt{b} \|\chi_b\|_{\text{sp}}^5$ for every $b \geq 1$, we have $2^{11} K(b)^4 C_\eta^2 \geq 2^{23} K(b)^4 \geq 2^{30} \sqrt{b} \|\chi_b\|_{\text{sp}}^5$ and the third term in (8.65) can be dropped. We conjecture that $D_{\text{irr}}(\chi, \chi_b) \geq D_{\text{irr}}(\chi, \chi) = 3$. If that were the case, then we could use the bound $K(b)^4 \geq 2^7 \|\chi_{b,L}\|_{\text{sp}}^5$ to obtain $2^{11} K(b)^4 C_\eta^2 \geq 2^{23} K(b)^4 \geq 2^{30} \|\chi_{b,L}\|_{\text{sp}}^5 \geq \frac{\pi^2 2^{21}}{3} \|\chi_{b,L}\|_{\text{sp}}^5 \geq \pi^2 2^{21} \frac{\|\chi_{b,L}\|_{\text{sp}}^5}{D_{\text{irr}}(\chi, \chi_b)}$ and we would also drop the last term of (8.65). However, since we do not have an explicit lower bound for $D_{\text{irr}}(\chi, \chi_b)$,

we avoid the comparison between the second and the last term of (8.65). Therefore, we take the implied constant in the O_η -notation in (8.64) as

$$P_0^{\text{irr}}(b, \eta) = \max \left\{ 2^{11} K(b)^4 C_\eta^2, \frac{\pi^2 2^{21} \|\chi_{b,L}\|_{\text{sp}}^5}{D_{\text{irr}}(\chi, \chi_b)} \right\}. \quad (8.66)$$

Note that, refining the argument above, the second term of (8.66) can be dropped as long as we have the lower bound $D_{\text{irr}}(\chi, \chi_b) \geq 2^{-9} \pi^2$. The best power saving in (8.64) comes from choosing (α^*, β^*) in the region $\{(\alpha, \beta) : 0 < \alpha < \frac{1}{\eta-1}, 0 < \beta < \frac{\alpha}{2}\}$ in order to maximise $m(\alpha, \beta) = \min\{\frac{\alpha}{2}, \beta, 2\eta - 2\eta(\eta-1)\alpha, \frac{3}{2}\alpha - 3\beta\} = \min\{\beta, 2\eta - 2\eta(\eta-1)\alpha, \frac{3}{2}\alpha - 3\beta\}$. That is

$$(\alpha^*, \beta^*) = \left(\frac{16\eta}{16\eta(\eta-1) + 3}, \frac{6\eta}{16\eta(\eta-1) + 3} \right) \quad (8.67)$$

at which $m(\alpha^*, \beta^*) = \beta^*$. Therefore, combining (8.58), (8.60), (8.63), (8.64), and (8.67), we obtain that for every $R > R_0^{\text{irr}}(\alpha^*, \beta^*, \eta)$ we have

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{N} S_N(x; c, \alpha) \overline{S_{\lfloor bN \rfloor}(x; c, \alpha)} \right| > R^2 \right\} = \frac{2D_{\text{irr}}(\chi, \chi_b)}{\pi^2 R^6} \left(1 + O_\eta \left(R^{-\frac{6\eta}{16\eta(\eta-1)+3}} \right) \right), \quad (8.68)$$

where the implicit constant in the O_η -notation in (8.68) can be taken as $4P_0^{\text{irr}}(b, \eta)$. Note that

$$\frac{1}{\eta - \eta(\eta-1)\alpha^*} = \frac{1}{\eta} + \frac{16}{3}(\eta-1), \quad (8.69)$$

$$\frac{1}{\beta^*} = \frac{1}{2\eta} + \frac{8}{3}(\eta-1), \quad (8.70)$$

$$\frac{1}{\alpha^* - 2\beta^*} = \frac{3}{4\eta} + 4(\eta-1), \quad (8.71)$$

$$\frac{1}{\alpha^*} = \frac{3}{16\eta} + (\eta-1). \quad (8.72)$$

We claim that

$$R_0^{\text{irr}}(\alpha^*, \beta^*; b, \eta) = (2^{\frac{11}{2}} K(b)^2 C_\eta)^{\frac{1}{\eta} + \frac{16}{3}(\eta-1)}. \quad (8.73)$$

To see this, first use (8.70)-(8.71) to see that $126^{\frac{1}{\beta^*}} = (126^{\frac{2}{3}})^{\frac{3}{4\eta} + 4(\eta-1)} \geq 2^{\frac{3}{4\eta} + 4(\eta-1)} = 2^{\frac{1}{\alpha^* - 2\beta^*}}$. Then (8.69)-(8.70) to obtain

$$(2^{\frac{11}{2}} K(b)^2 C_\eta)^{\frac{1}{\eta - \eta(\eta-1)\alpha^*}} = (2^{11} K(b)^4 C_\eta^2)^{\frac{1}{2\eta} + \frac{8}{3}(\eta-1)} \geq 126^{\frac{1}{2\eta} + \frac{8}{3}(\eta-1)} \geq 126^{\frac{1}{\beta^*}} \quad (8.74)$$

because $2^{11} K(b)^4 C_\eta^2 \geq 2^{46} \geq 126$. Finally, (8.72) and the explicit bound $K(b)^2 \geq 2^5 b^{\frac{3}{16}} \|\chi_b\|_{\text{sp}}^{\frac{15}{8}}$ yield

$$(2^{\frac{11}{2}} K(b)^2 C_\eta)^{\frac{1}{\eta - \eta(\eta-1)\alpha^*}} = (2^{\frac{11}{2}} K(b)^2 C_\eta)^{\frac{1}{\eta} + \frac{16}{3}(\eta-1)} \quad (8.75)$$

$$\geq (2^{\frac{9}{8}} b^{\frac{3}{16}} \|\chi_b\|_{\text{sp}}^{\frac{15}{8}})^{\frac{1}{\eta} + \frac{16}{3}(\eta-1)} = (2^6 b \|\chi_b\|_{\text{sp}}^{10})^{\frac{1}{\alpha^*}} \quad (8.76)$$

since $2^{\frac{11}{2}} K(b)^2 C_\eta \geq 2^{\frac{23}{2}} K(b)^2 \geq 2^{\frac{33}{2}} b^{\frac{3}{16}} \|\chi_b\|_{\text{sp}}^{\frac{15}{8}} \geq 2^{\frac{9}{8}} b^{\frac{3}{16}} \|\chi_b\|_{\text{sp}}^{\frac{15}{8}}$. This concludes the proof of the claim (8.73). Therefore, if we set

$$R_0^{\text{irr}}(b, \eta) = (2^{\frac{11}{2}} K(b)^2 C_\eta)^{\frac{1}{\eta} + \frac{16}{3}(\eta-1)}, \quad (8.77)$$

then for every $R > R_0^{\text{irr}}(b, \eta)$ we have (8.68) with an implicit constant of

$$P^{\text{irr}}(b, \eta) := \max \left\{ 2^{13} K(b)^4 C_\eta^2, \frac{\pi^2 2^{24} \|\chi_{b,L}\|_{\text{sp}}^5}{D_{\text{irr}}(\chi, \chi_b)} \right\}. \quad (8.78)$$

and the theorem is proven. \square

Theorem 8.7 (Explicit version of Theorem 1.6). *Let λ be a probability measure on \mathbb{R} which is absolutely continuous with respect to the Lebesgue measure. Let $(c, \alpha) \notin \mathbb{Q}^2$, $b \geq 1$ and $0 < \varepsilon \leq 1$. Set*

$$R^{\text{irr}}(b, \varepsilon) = 2^{39} K(b)^4 \varepsilon^{-2} \quad (8.79)$$

where $K(b)$ is given in (6.28). Then for all $R > R_b^{\text{irr}}(\varepsilon)$ we have

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{N} S_N(x; c, \alpha) \overline{S_{\lfloor bN \rfloor}(x; c, \alpha)} \right| > R^2 \right\} = \frac{2D_{\text{irr}}(b)}{\pi^2 R^6} (1 + O_\varepsilon(R^{-2+\varepsilon})), \quad (8.80)$$

where $D_{\text{irr}}(b) = D_{\text{irr}}(\chi, \chi_b)$ is given in (8.57), and the constant implied by the O_ε -notation in (8.80) can be taken to be

$$\max \{ 2^{41} K(b)^4 \varepsilon^{-4}, 2^{28} \max\{b^5, 3^{10}\} D_{\text{irr}}(b)^{-1} \}. \quad (8.81)$$

Proof. Note that the exponent in (8.56) satisfies

$$\lim_{\eta \rightarrow 1^+} \frac{6\eta}{16\eta(\eta-1) + 3} = 2^-.$$

This means that the closer $\eta > 1$ is to 1, the better the power saving in Theorem 8.6 is. If we set $\frac{6\eta}{16\eta(\eta-1) + 3} = 2 - \varepsilon$, then $\varepsilon \mapsto \eta(\varepsilon) = \frac{19-8\varepsilon+\sqrt{16\varepsilon^2-112\varepsilon+169}}{16(2-\varepsilon)}$ maps $(0, 1]$ to $(1, \frac{11+\sqrt{73}}{16}]$ monotonically. Hence we assume that $1 < \eta \leq \frac{11+\sqrt{73}}{16}$. We aim to employ the lower bound $\eta(\varepsilon) > 1 + \frac{3}{26}\varepsilon$. Note that, by Corollary A.4, (8.50) holds in this case. We use (8.77), along with the fact that $\frac{1}{\eta} + \frac{16}{3}(\eta-1) \leq 2$ and the bound $(\eta-1)^{-2(\frac{1}{\eta} + \frac{16}{3}(\eta-1))} \leq 2^5(s-1)^{-2}$ to obtain

$$R_0^{\text{irr}}(b, \eta) \leq \left(\frac{2^{\frac{13}{2}+6\eta} K(b)^2}{(\eta-1)^2} \right)^{\frac{1}{\eta} + \frac{16}{3}(\eta-1)} \leq \frac{2^{\frac{85}{4} + \frac{3}{4}\sqrt{73}} K(b)^4}{(\eta-1)^{2(\frac{1}{\eta} + \frac{16}{3}(\eta-1))}} \leq \frac{2^{\frac{105}{4} + \frac{3}{4}\sqrt{73}} K(b)^4}{(\eta-1)^2}. \quad (8.82)$$

By monotonicity we have

$$R_0^{\text{irr}}(b, \eta(\varepsilon)) \leq \frac{2^{\frac{105}{4} + \frac{3}{4}\sqrt{73}} K(b)^4}{(\frac{3}{26}\varepsilon)^2} \leq 2^{39} K(b)^4 \varepsilon^{-2} =: R^{\text{irr}}(b, \varepsilon). \quad (8.83)$$

Therefore, if $R > R^{\text{irr}}(b, \varepsilon)$, then by Theorem 8.6 we obtain

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{N} S_N(x; c, \alpha) \overline{S_{[bN]}(x; c, \alpha)} \right| > R^2 \right\} = \frac{2D_{\text{irr}}(\chi, \chi_b)}{\pi^2 R^6} (1 + O_\eta(R^{-2+\varepsilon})) \quad (8.84)$$

where the constant implied by the O_ε -notation is given by $P^{\text{irr}}(b, \eta(\varepsilon))$ using (8.78). The bounds $2^{11} K(b)^4 C_{\eta(\varepsilon)}^2 \leq 2^{\frac{85}{4} + \frac{3}{4}\sqrt{73}} K(b)^4 (\eta(\varepsilon) - 1)^{-4} \leq 2^{41} K(b)^4 \varepsilon^{-4}$ and (6.7) imply that the constant implied by the O_ε -notation in (8.80) can be taken as (8.81) \square

If we take $b = 1$ in Theorem 8.7 and, use the fact that $D_{\text{irr}}(1) = 3$ (see (3.121) in [6]), we obtain the following

Corollary 8.8 (Explicit version of Theorem 1.2). *Let λ be a probability measure on \mathbb{R} which is absolutely continuous with respect to the Lebesgue measure. Let $(c, \alpha) \notin \mathbb{Q}^2$ and $0 < \varepsilon \leq 1$. Set $R^{\text{irr}}(\varepsilon) = 2^{63} \varepsilon^{-2}$. Then for all $R > R^{\text{irr}}(\varepsilon)$ we have*

$$\lim_{N \rightarrow \infty} \lambda \left\{ x \in \mathbb{R} : \left| \frac{1}{\sqrt{N}} S_N(x; c, \alpha) \right| > R \right\} = \frac{6}{\pi^2 R^6} (1 + O_\varepsilon(R^{-2+\varepsilon})), \quad (8.85)$$

the constant implied by the O_ε -notation in (8.85) can be taken to be $2^{65} \varepsilon^{-4}$.

8.3 Final discussion on the explicit constants

As pointed out in the Introduction, we put a lot of effort into finding explicit, simple-looking constants for the implied O_ε -notation in Theorems 1.1-1.2 and in their generalisations 1.5-1.6. These constants are given in Corollaries 8.5 and 8.8, and in (8.47), (8.49), (8.79), (8.81). We do not claim that the dependence of these constants upon b and ε is optimal. One can in principle make these constants smaller by improving (8.45), (8.46), (8.77), and (8.78). This task would involve fairly elementary estimates, but would also be extremely lengthy and we purposefully avoided it. For the enthusiastic reader who seeks to optimise the constants, we list below some steps in which improvements can be made, and highlight some of the effects of such potential changes.

1. To improve the constant $C_\eta = 2^{6\eta} \zeta(\eta)^2$ in Lemma 4.1:

- (a) Bound (4.7) above by $\zeta(\eta)(2^\eta - 1) + \zeta(\eta, \frac{3}{2})$, where $\zeta(\eta, a)$ is the Hurwitz zeta function $\zeta(\eta, a) = \sum_{n=0}^{\infty} (n+a)^{-\eta}$.
- (b) Replace A_η by $A'_\eta = (\zeta(\eta)(2^\eta - 1) + \zeta(\eta, \frac{3}{2}))^2$ and B_η by $B'_\eta = 2(\zeta(\eta)(2^\eta - 1) + \zeta(\eta, \frac{3}{2}))$ in (4.8). These new constants satisfy the inequality $B'_\eta \leq \frac{7}{20} A'_\eta$.
- (c) Replace C_η by $A'_\eta(2^{\frac{\eta}{2}} + \frac{7}{20})$. Near $\eta = 1$, the improved constant would be smaller than C_η by at least a factor of 9.

2. With the revised definition of C_η discussed above, the statement of Lemma 4.2 would still hold, but its proof would need to be changed. In fact, since C_η enters the definition of $\tilde{\kappa}$, one has to show that (4.32) still implies (4.24). Albeit true, this would be more laborious to prove. Similarly, Lemma 4.3 would still hold. Its proof would not require major changes as it would still be easy to see that (4.52) implies (4.40).
3. The lower bound $C_\eta \geq 2^6$, which we use in the proof of Theorems 8.3 and 8.6, would still hold but its proof would require some effort.
4. The constants $K(s)$ and $K_L(s)$ in the key Proposition 6.5 could be improved by avoiding the bound $2^{\frac{\eta}{2}} \leq 2$ for $1 < \eta \leq 2$ in (6.27) and (6.34). This would complicate the statements and the arguments in Sections 7-8 by introducing dependence of such constants upon η .
5. In the proof of Lemma 7.2, in addition to improving C_η as discussed in point 1 above, we could avoid the use of the inequality $K(1) \leq K(b)$ and replace (7.30) with $R^2 > 16 \cdot C_\eta K(1) K(b) 2^{2(\eta-1)J} = 16 \cdot 59 \cdot C_\eta K(b) 2^{2(\eta-1)J}$. The improvement would be especially beneficial for large values of b , since $K(b) \sim \frac{2}{3}b^3$ as $b \rightarrow \infty$. Moreover, restricting $1 < \eta \leq \eta_0$ (see point 8a below), one could replace the factor 16 with $2^{2+\eta_0}$. In the same way, we could lower the implied constant $2^{10}C_\eta^2 K(b)^4$ by replacing (7.34) with $3 \cdot 2^{\eta_0^2+2\eta_0} C_\eta^{\eta_0} K(b)^{2\eta_0} 2^{2\eta(\eta-1)J} R^{-2\eta}$. Similar considerations hold for Lemma 7.4.
6. In Lemma 7.1 the power 2 of $K_L(b)$ could not be reduced due to (7.13). On the other hand, assuming $1 < \eta \leq \eta_0$ (see point 8a below), the factor 64 in the assumption of the lemma could be lowered to $2^{\frac{7}{2}+\eta_0}$. The same holds for Lemma 7.3.
7. If we carried out the improvements described in points 5-6 above, then we would need to modify the proof of Theorem 8.3. Specifically, we would replace the first inequality of (8.26) by $R^2 > 2^{\frac{7}{2}+\eta_0} K_L(b)^2 C_\eta R^{2(\eta-1)\alpha}$ and the first inequality of (8.27) by $R^2 > 59 \cdot 2^{2+\eta_0} K(b) C_\eta 2^{2(\eta-1)\alpha}$. Consequently, because of the different powers of $K_L(b)$ and $K(b)$, Remark 6.6 could not be used and the sufficient condition (8.28) for (8.26)-(8.27) would have to be changed. In turn, this change would affect (8.31), (8.33), (8.42), and (8.45). The lowering of the implied constants $2^{10}C_\eta^2 K(b)^4$ in Lemma 7.2 would affect (8.32), (8.33), (8.42), (8.45), but also (8.35), (8.36), and (8.46). Ultimately the term $(K(b)^2 C_\eta)^{\frac{1}{\eta}+6(\eta-1)}$ in $R_0^{\text{rat}}(n, \eta)$ could be replaced by its square root. The term $K(b)^4 C_\eta^2$ in (8.35) could be replaced by $K(b)^\eta C_\eta^\eta$ but (8.36) would have to be written as the maximum of two terms (one depending on b and η and one only dependent on b). Similar considerations apply to Theorem 8.6, ultimately affecting (8.77) and (8.78).

8. Reducing the interval $0 < \varepsilon \leq 1$ in Theorem 8.4 to $0 < \varepsilon \leq \varepsilon_0$ would improve the constant (8.47) (and hence the constant $R^{\text{rat}}(\varepsilon)$ in Corollary 8.5) as follows:

- (a) It would reduce the interval $1 < \eta \leq \frac{4+\sqrt{10}}{6}$ to $1 < \eta \leq \eta_0 = \frac{7-3\varepsilon_0+\sqrt{3\varepsilon_0^2-18\varepsilon_0+25}}{6(2-\varepsilon_0)}$ in the proof of the theorem.
- (b) The constant $\sqrt{2}$ in Corollary A.4 (and hence in (8.50)) could be lowered and hence improve the first inequality in (8.51).
- (c) For sufficiently small ε_0 we would gain the monotonicity of the function $\eta \mapsto R_0^{\text{rat}}(b, \eta)$ (or its improved version, as described in point 7 above) and of the second term in (8.51) (i.e. the upper bound of (8.45) obtained using (8.50)) on the interval $(1, \eta_0]$.
- (d) Even if we did not alter (8.45), the bound $\frac{1}{\eta} + 6(\eta - 1) \leq 2$, used in (8.51), could be replaced by $\frac{1}{\eta} + 6(\eta - 1) \leq \frac{2}{2-\varepsilon_0}$ and consequently reduce the exponent to which $K(b)^2$ is raised from 2 to slightly above 1. Therefore $K(b)^4$ in (8.47) could be replaced by $K(b)^{\frac{4}{2-\varepsilon_0}}$. The dependence of (8.47) upon ε would be unchanged. If we did alter (8.45) as described in points 5-7 above, then the dependence of (8.47) upon b and ε would be $K(b)^{\frac{2}{2-\varepsilon_0}} \varepsilon^{-1}$ (up to a multiplicative constant).

9. The improvement of the constant (8.49) is a little more subtle. The reason is that, as outlined in point 7 above, (8.36) would be written as the maximum of two terms. The term involving b and η would include $K(b)$ raised to a power slightly above 1 (improving the power 4 in (8.49)) and ε raised to a power slightly below -2 (improving the power -4). We could improve the constant $2^{66} \varepsilon^{-4}$ in Corollary 8.5 accordingly.

10. Arguing as in points 8-9, we could also improve the constants (8.79) and (8.81) in Theorem 8.7 and those in Corollary 8.8.

A Bounds for the zeta function

The constant $C_\eta = 2^{6\eta} \zeta(\eta)^2$ in Lemmata 4.1 and 4.3 involves the Riemann zeta function and is used throughout Sections 4, 7, and 8 of the paper. In the proofs of Theorems 8.4 and 8.7 we use the upper bound (8.50), which we prove in this appendix.

For $\eta > 1$ let $\zeta_a(\eta) = \sum_{n=1}^{\infty} (-1)^{n-1} n^{-\eta}$ be the alternating zeta function (also known as the Dirichlet eta function).

Lemma A.1. *For $\eta > 1$ we have*

$$\zeta_a(\eta) \left(\frac{1}{\log(2)(\eta-1)} + \frac{1}{2} \right) \leq \zeta(\eta) \leq \zeta_a(\eta) \left(\frac{1}{\log(2)(\eta-1)} + \frac{1}{2} + \frac{\log(2)}{12}(\eta-1) \right). \quad (\text{A.1})$$

Proof. The identity $\zeta(\eta) = \zeta_a(\eta)(1 - 2^{1-\eta})^{-1}$ is classical. Truncating the Laurent series $(1 - 2^{1-\eta})^{-1}$ near $\eta = 1$ to the first order we have $\frac{1}{\log(2)(\eta-1)} + \frac{1}{2} + \frac{\log(2)}{12}(\eta - 1)$. This is an upper bound for $(1 - 2^{1-\eta})^{-1}$ if $\eta > 1$. In fact, an application of l'Hôpital's rule gives

$$\lim_{\eta \rightarrow 1} \frac{1}{\log(2)(\eta - 1)} + \frac{1}{2} + \frac{\log(2)}{12}(\eta - 1) - (1 - 2^{1-\eta})^{-1} = 0 \quad (\text{A.2})$$

and we can check that for $\eta > 1$

$$\frac{d}{d\eta} \left(\frac{1}{\log(2)(\eta - 1)} + \frac{1}{2} + \frac{\log(2)}{12}(\eta - 1) - (1 - 2^{1-\eta})^{-1} \right) > 0. \quad (\text{A.3})$$

This yields the upper bound in (A.1). A similar argument applied to $(1 - 2^{1-\eta})^{-1} - (\frac{1}{\log(2)(\eta-1)} + \frac{1}{2})$ yields the lower bound. \square

Corollary A.2. *For every $\eta_0 > 1$ there is a constant $c(\eta_0) > 1$ such that if $1 < \eta \leq \eta_0$ we have $\frac{1}{\eta-1} < \zeta(\eta) \leq \frac{c(\eta_0)}{\eta-1}$.*

Proof. Note that functions $\eta \mapsto \zeta_a(\eta)$ and $\eta \mapsto \frac{1}{\log(2)} + \frac{1}{2}(\eta - 1) + \frac{\log(2)}{12}(\eta - 1)^2$ are both positive and increasing for $\eta > 1$. Therefore, for $\eta \in (1, \eta_0]$, their product is maximised at $\eta = \eta_0$. Similarly, since $\frac{1}{\log(2)} + \frac{1}{2}(\eta - 1)$ is bounded below by its value at $\eta = 1$ and $\zeta_a(1) = \log(2)$ (this is the alternating harmonic series), we have $\zeta_a(\eta) \left(\frac{1}{\log(2)(\eta-1)} + \frac{1}{2} \right) > 1$. The statement then follows from Lemma A.1 with

$$c(\eta_0) = \zeta_a(\eta_0) \left(\frac{1}{\log(2)} + \frac{1}{2}(\eta_0 - 1) + \frac{\log(2)}{12}(\eta_0 - 1)^2 \right). \quad (\text{A.4})$$

\square

Remark A.3. Note that the lower bound in Corollary A.2 is only meaningful for $1 < \eta \leq 2$ since for $\eta > 2$ it is worse than the trivial bound $\zeta(\eta) > 1$.

Corollary A.4. *For $1 < \eta \leq \frac{5}{4}$, we have $\frac{1}{\eta-1} < \zeta(\eta) \leq \frac{\sqrt{2}}{\eta-1}$.*

Proof. The statement follows immediately from Corollary A.2 since, using (A.4), it is easy to verify that $1.1487793 \approx c(\frac{5}{4}) < \sqrt{2}$. \square

Note that Corollary A.4 can be used in the proofs of Theorems 8.4 and 8.7 since they use the ranges $1 < \eta \leq \frac{4+\sqrt{10}}{6}$ and $1 < \eta \leq \frac{11+\sqrt{73}}{16}$ respectively and $\frac{4+\sqrt{10}}{6}, \frac{11+\sqrt{73}}{16} \leq \frac{5}{4}$.

Acknowledgements

The first and third authors acknowledge the support from the NSERC Discovery Grant “Statistical and Number-Theoretical Aspects of Dynamical Systems”. The authors thank the anonymous referee for the useful corrections they suggested.

References

- [1] J. S. Athreya, J. Chaika, and S. Lelièvre. The gap distribution of slopes on the golden L. In *Recent trends in ergodic theory and dynamical systems*, volume 631 of *Contemp. Math.*, pages 47–62. Amer. Math. Soc., Providence, RI, 2015.
- [2] F. P. Boca, C. Cobeli, and A. Zaharescu. Distribution of lattice points visible from the origin. *Comm. Math. Phys.*, 213(2):433–470, 2000.
- [3] F. P. Boca, B. Heersink, and P. Spiegelhalter. Gap distribution of Farey fractions under some divisibility constraints. *Integers*, 13:Paper No. A44, 15, 2013.
- [4] F. P. Boca and A. Zaharescu. The distribution of the free path lengths in the periodic two-dimensional Lorentz gas in the small-scatterer limit. *Comm. Math. Phys.*, 269(2):425–471, 2007.
- [5] F. Cellarosi. Limiting curlicue measures for theta sums. *Ann. Inst. Henri Poincaré Probab. Stat.*, 47(2):466–497, 2011.
- [6] F. Cellarosi and J. Marklof. Quadratic weyl sums, automorphic functions and invariance principles. *Proceedings of the London Mathematical Society*, 113(6):775–828, 2016.
- [7] F. Cellarosi and T. Osman. Heavy tailed and compactly supported distributions of quadratic Weyl sums with rational parameters. *arXiv:2210.09838*.
- [8] E. Demirci Akarsu. Short incomplete Gauss sums and rational points on metaplectic horocycles. *Int. J. Number Theory*, 10(6):1553–1576, 2014.
- [9] E. Demirci Akarsu and J. Marklof. The value distribution of incomplete Gauss sums. *Mathematika*, 59(2):381–398, 2013.
- [10] D. Dolgopyat and B. Fayad. Deviations of ergodic sums for toral translations I. Convex bodies. *Geom. Funct. Anal.*, 24(1):85–115, 2014.
- [11] D. Dolgopyat and B. Fayad. Deviations of ergodic sums for toral translations II. Boxes. *Publ. Math. Inst. Hautes Études Sci.*, 132:293–352, 2020.
- [12] R. M. Dudley. *Real Analysis and Probability*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2 edition, 2002.
- [13] N. D. Elkies and C. T. McMullen. Gaps in $\sqrt{n} \bmod 1$ and ergodic theory. *Duke Math. J.*, 123(1):95–139, 2004.
- [14] Allan Gut. *Probability: a graduate course*. Springer Texts in Statistics. Springer, New York, second edition, 2013.

- [15] E. Kowalski and W.F. Sawin. Kloosterman paths and the shape of exponential sums. *Compos. Math.*, 152(7):1489–1516, 2016.
- [16] G. Lion and M. Vergne. *The Weil representation, Maslov index and theta series*, volume 6 of *Progress in Mathematics*. Birkhäuser, Boston, Mass., 1980.
- [17] J. Marklof. Limit theorems for theta sums. *Duke Math. J.*, 97(1):127–153, 1999.
- [18] J. Marklof. Almost modular functions and the distribution of n^2x modulo one. *Int. Math. Res. Not.*, (39):2131–2151, 2003.
- [19] J. Marklof. Spectral theta series of operators with periodic bicharacteristic flow. *Ann. Inst. Fourier (Grenoble)*, 57(7):2401–2427, 2007. Festival Yves Colin de Verdière.
- [20] J. Marklof. The asymptotic distribution of Frobenius numbers. *Invent. Math.*, 181(1):179–207, 2010.
- [21] J. Marklof and A. Strömbergsson. The distribution of free path lengths in the periodic Lorentz gas and related lattice point problems. *Ann. of Math. (2)*, 172(3):1949–2033, 2010.
- [22] J. Marklof and A. Strömbergsson. The periodic Lorentz gas in the Boltzmann-Grad limit: asymptotic estimates. *Geom. Funct. Anal.*, 21(3):560–647, 2011.
- [23] J. Marklof and A. Strömbergsson. Power-law distributions for the free path length in Lorentz gases. *J. Stat. Phys.*, 155(6):1072–1086, 2014.
- [24] P. Nándori, D. Szász, and T. Varjú. Tail asymptotics of free path lengths for the periodic Lorentz process: on Dettmann’s geometric conjectures. *Comm. Math. Phys.*, 331(1):111–137, 2014.
- [25] P. Sarnak. Asymptotic behavior of periodic orbits of the horocycle flow and Eisenstein series. *Comm. Pure Appl. Math.*, 34(6):719–739, 1981.
- [26] C. Uyanik and G. Work. The distribution of gaps for saddle connections on the octagon. *Int. Math. Res. Not. IMRN*, (18):5569–5602, 2016.