BREDON AND FARRELL-JONES HOMOLOGY OF ARTIN GROUPS OF DIHEDRAL TYPE

YAGO ANTOLÍN AND RAMÓN FLORES

ABSTRACT. For Artin groups of dihedral type, we compute the Bredon homology groups of the classifying space for the family of virtually cyclic subgroups with coefficients in the K-theory of a group ring, and the Farrell-Jones homology groups of their classifying spaces with coefficients in regular rings.

1. Introduction

Let G be a group, R a ring and $n \in \mathbb{Z}$. In [18], Farrell-Jones stated the existence of an assembly map:

$$H_n^G(\underline{\underline{E}}G, \mathbf{K}(R)) \to K_n(RG),$$

where $\underline{\underline{E}}G$ is the classifying space of G with respect to the family of virtually cyclic groups, $H_*^G(-, \overline{\mathbf{K}}(R))$ is the G-homology theory defined in Section 1 of [18] and $K_n(RG)$ stands for the n-th group of algebraic K-theory of the group ring RG. The Farrell-Jones conjecture predicts that the assembly map is an isomorphism. The conjecture has been verified for a big family of groups (see [28] for an excellent survey), and no counterexample has been found so far. The philosophy in this context is, for a group for which the assembly map is known to be an isomorphism, to perform computations in the topological side in order to extract information about the algebraic K-theory of the group ring. It is remarkable that the latter are extremely difficult to compute, and at the same time encode fundamental invariants of manifolds, including obstructions to the existence of cobordisms and information about groups of pseudoisotopies.

The left-hand side of the conjecture can be approached by means of a G-equivariant version of the Atiyah-Hirzebruch spectral sequence, which converges to the Farrell-Jones K-homology, and whose E_2 -page is the Bredon homology of the classifying space $\underline{E}G$ with coefficients in the K-theory of the group rings of the virtually cyclic subgroups of G. This strategy has been used in the past with great success in the context of the Baum-Connes Conjecture; however, as the level of complexity increases when passing from the family of finite groups to the bigger family of virtually cyclic groups, there are not a lot of examples of explicit computations in the literature in the Farrell-Jones context ([5], [29], [41]). We remark here that the knowledge of the Bredon homology groups has always been of independent interest, because of the prominent role that they play in equivariant homotopy and the dimension theory of groups. In this sense,

⁰Mathematics Subject Classification (2020): Primary: 19E20; Secondary: 19A31, 19B28, 20F36. The second author was supported by grant PID2020-117971GB-C21 of the Spanish Ministery of Science and Innovation, and grant FQM-213 of the Junta de Andalucía.

the methods of the present paper may be useful in a more general context, switching the K-theoretic coefficients to a general coefficient module.

The computation of the Farrell-Jones K-homology groups appearing in the left-hand side of the conjecture can be undertaken in this way for any group G and any coefficient ring R. This is the content of Section 4 of this paper for G a dihedral Artin group. Moreover, with additional hypotheses over G and R, it is sometimes possible to simplify the input of the Atiyah-Hirzebruch spectral sequence. For example, if G is torsion-free and R is regular, a splitting result of Lück-Steimle [30] permits the use of the ordinary homology of G in the E_2 -page instead of the Bredon homology, which usually makes computations quite easier. We will take profit of these tools in the cases in which the additive abelian group of the coefficient ring is cyclic in Section 6.

This paper is part of a program aimed to understand equivariant and K-theoretic invariants for Artin groups. These groups are ubiquitious in geometric group theory, as they comprehend many different subfamilies of groups (braid groups, right-angled, spherical type, extra-large, free groups...), provide (counter)examples for many interesting phenomena, and at the same time they are not well understood from a global point of view: it is not even known if every Artin group is torsion-free. See [38] for a good survey. In this context, Azzali et al. have recently computed explicitly both sides of Baum-Connes for pure braid groups [4]; in joint work of the second author with J. González-Meneses [19] it is computed the minimal dimension of a model of $\underline{E}G$ for braid groups; J. Aramayona et al. have published two papers ([1], [2]) about virtually-cyclic dimensions of mapping class groups (which include un particular certain Artin groups).

In the present paper we consider the case of Artin groups of dihedral type (see Definition 3.1), which are one-relator, torsion-free and two-generated; moreover, they have small geometric dimension and its group-theoretic structure is quite well understood. These features makes them strongly appropriate for computations, and in fact have been recently studied from different angles, as growth series [34], systems of equations [11] or geodesics [46]. Moreover, these groups are free-by-cyclic, and the Farrell-Jones conjecture has been recently verified for this class of groups [7]. Hence, our computations in the left-hand side can be read in terms of algebraic K-theory of RG.

Our main results about these groups exploit the two points of view exposed above, i.e. computation of Bredon homology and description of the Farrell-Jones K-homology making use of the simplified version of Atiyah-Hirzebruch spectral sequence. Concerning the former, our main result is the following:

Theorem 4.17. Let A_n be an Artin group of dihedral type and R a ring. Then we have the following:

$$\begin{aligned} &(1) \ \ H_i^{vc}(A_n,K_q(R[-])) = \{0\} \ \text{for} \ i \geq 4. \\ &(2) \ \ H_3^{vc}(A_n,K_q(R[-])) = \begin{cases} \bigoplus_{[H] \neq [Z(A_n)]} K_q(R) & n \ \text{odd} \\ \ker g_2^2 & n \ \text{even.} \end{cases} \\ &(3) \ \ H_2^{vc}(A_n,K_q(R[-])) = \ker g_2^1. \\ &(4) \ \ H_1^{vc}(A_n,K_q(R[-])) = \operatorname{coker} g_2^1 = \begin{cases} (\bigoplus_{[H] \neq [Z(A_n)]} N_q^{[H]}) \oplus T_1(K_q(R)) \oplus T_2(K_q(R)) & n \ \text{odd} \\ \bigoplus_{[H]} N_q^{[H]} & n \ \text{even.} \end{cases}$$

(5)
$$H_0^{vc}(A_n, K_q(R[-])) = \operatorname{coker} g_2^0 = \begin{cases} (\bigoplus_{[H] \neq [Z(A_n)]} N_q^{[H]}) \oplus K_q(R) \oplus \overline{C}(K_q(R)) & n \text{ odd} \\ (\bigoplus_{[H]} N_q^{[H]}) \oplus K_q(R) & n \text{ even.} \end{cases}$$

Here g_i^j stands for a homomorphism in the Degrijse-Petrosyan exact sequence [14, Section 7], [H] for the commensurability class of a non-trivial cyclic subgroup H, and $N_q^{[H]}$, $T_i(K_q(R))$ and $\overline{C}(K_q(R))$ for groups that depend on H and the Bass-Heller-Swan decomposition of $K_q(R[\mathbb{Z}])$. See Section 4 for details.

As an example, we note that the previous theorem, together with the fact that dihedral Artin groups satisfy the Farrell-Jones conjecture provides the following information about the K-theory of the group ring.

Corollary 4.18. Let A_n , n > 2 be an Artin group of dihedral type. Let R be a regular ring. Then $K_0(RA_n) = K_0(R)$.

The main tool used in the proof of Theorem 4.17 is an exact sequence in Bredon homology [14], which is in turn the Mayer-Vietoris sequence associated to the push-out that defines the Lück-Weiermann model for $\underline{E}G$ ([31], see also Section 2 below). The knowledge about the group-theoretic structure of the groups includes a complete understanding of the commensurators, which is crucial in the computations. Using the theorem, we are able to describe with precision the Bredon homology of A_n with coefficients in the K-theory of several rings, as \mathbb{Z} , $\mathbb{Z}/2$ or some group rings of finite groups (see Section 5).

When the coefficient ring is regular, it is more efficient to use the Atiyah-Hirzebruch sequence whose E^2 -page is given by ordinary homology. This method is used in Section 6, and allows us to describe Farrell-Jones K-homology groups of A_n with coefficients in $\mathbf{K}(\mathbb{Z})$ and $\mathbf{K}(\mathbb{F}_r)$ (where \mathbb{F}_r is the field of r elements and r is prime), with a good degree of sharpness when n is odd. For example, up to a few extension problems, the groups $H^{A_{2n-1}}(\underline{\mathbb{E}} A_{2n-1}, \mathbf{K}(\mathbb{Z}))$ are computed in Proposition 6.2, while Proposition 6.6, Proposition 6.7 and Corollary 6.8 identify the groups $H^{A_n}(\underline{\mathbb{E}} A_n, \mathbf{K}(\mathbb{F}_r))$. When $R = \mathbb{Z}$ and n even the information is less complete.

We finish this introduction by summarizing the contents of the different sections of this paper. In Section 2 we recall the main definitions about classifying spaces for families and Bredon homology, with special emphasis in the Lück-Weiermann model and its associated Mayer-Vietoris sequence. In Section 3 the main properties of Artin groups that will be used on the rest of the paper are studied. In Section 4, we carefully analyze the homomorphisms in the exact sequence and prove Theorem 4.17, which in Section 5 is applied to describe different concrete examples. Finally, in Section 6 we compute the Farrell-Jones K-homology over regular rings.

2. Preliminaries

 $\langle \mathtt{Sect:Prelim} \rangle$

In this section we state some notions of G-equivariant homotopy that will frequently appear in the rest of the paper. The exposition will be sketchy and very focused to our goals; the reader interested in a thorough treatment of the subject is referred to [16] for the theory of G-CW-complexes and actions on them, to [26] for the theory of classifying spaces and to the first part of [36] for Bredon homology.

 $?\langle \texttt{Sect:classify} \rangle ?$

2.1. Classifying spaces for families. In this section we will briefly recall the notion of classifying space for a family of subgroups, which is the central object in the topological side of the Isomorphism Conjectures. Then we will review Lück-Weiermann model and the definition of commensurator, which will be crucial in our computations.

Definition 2.1. Let G be a discrete group, and \mathcal{F} be a family of subgroups of G closed under passing to subgroups and conjugation. A G-CW-complex X is a classifying space for the family \mathcal{F} if for every $H \in \mathcal{F}$ the fixed-point set X^H is contractible, and empty otherwise.

The classifying space for the family \mathcal{F} is usually denoted by $E_{\mathcal{F}}G$. Moreover, two models for $E_{\mathcal{F}}G$ are G-homotopy equivalent. A point is always a model for $E_{\mathcal{F}}G$ if $G \in \mathcal{F}$, and the closeness under subgroups implies that $E_{\mathcal{F}}G$ is always a contractible space.

If there is a family \mathcal{F} of subgroups of G with the previous closeness properties and a subgroup $H \leq G$, we denote by $\mathcal{F} \cap H$ the family whose elements are the intersections $F \cap H$, with $F \in \mathcal{F}$. The family $\mathcal{F} \cap H$ of subgroups of H is again closed under H-conjugation and taking subgroups. In these conditions the action of H over $E_{\mathcal{F}}G$ by restriction turns $E_{\mathcal{F}}G$ into a model for $E_{\mathcal{F} \cap H}H$.

The most important families of subgroups in this context are the trivial family $\mathcal{F}_{\{1\}}$, the family \mathcal{F}_{Fin} of finite groups and the family \mathcal{F}_{vc} of virtually cyclic groups of G; the classifying spaces for these families are respectively denoted by EG, $\underline{E}G$ and $\underline{E}G$. Observe that $\mathcal{F}_{\{1\}} \subseteq \mathcal{F}_{Fin} \subseteq \mathcal{F}_{vc}$, and that $\mathcal{F}_{\{1\}} = \mathcal{F}_{Fin}$ if and only if G is torsion-free. It is also a standard argument to show that torsion-free virtually cyclic groups are cyclic (see for example Lemma 3.2 in [33]). Then, for torsion-free groups, \mathcal{F}_{vc} is the set of cyclic subgroups.

From now on we will describe the model of $\underline{E}G$ developed by Lück-Weiermann in [31], for the special families we are interested (the construction is indeed more general). Given a group G, and two subgroups H and K, we consider the equivalence relation generated by $H \sim K$ if $H \cap K$ has finite index in both H and K. Observe that if H is in $\mathcal{F}_{vc} \setminus \mathcal{F}_{Fin}$ then $H \sim K$ if and only if K is virtually cyclic and $H \cap K$ is infinite. Also remark that if $H \in \mathcal{F}_{Fin}$, then $H \sim K$ if and only if $K \in \mathcal{F}_{Fin}$. The equivalence class of H will be denoted by [H]. This equivalence relation is preserved by conjugation, it can be defined $g^{-1}[H]g$ as $[g^{-1}Hg]$ for any $H \leqslant G$ and for any $g \in G$.

Now we can define the notion of *commensurator*, central in this model and in our paper:

 $?\langle defn:comm \rangle ?$

Definition 2.2. Given an equivalence class [H] of the relation \sim , the *commensurator* of [H] in G is defined as the subgroup

$$\text{Comm}_G[H] = \{ g \in G \mid g^{-1}[H]g = [H] \}.$$

In [31], it is also defined a family of subgroups of $Comm_G[H]$ as:

$$\mathcal{F}[H] := \{ K < \text{Comm}_G[H] \mid K \sim H \text{ or } |K| < \infty \}.$$

If H is virtually cyclic, it is easy to check that this family is closed under taking subgroups and conjugation in $Comm_G[H]$. We remark that the commensurator of [H] is sometimes denoted by $N_G[H]$ in the literature.

Now we have all the ingredients needed for building Lück-Weiermann model:

 $\langle \text{maintheorem} \rangle$ Theorem 2.3. ([31], Theorem 2.3) We denote by I a complete set of representatives of the G-orbits (under conjugation) of equivalence classes [H] of infinite virtually cyclic subgroups of G, and we choose, for every $[H] \in I$, models for the classifying spaces $E \operatorname{Comm}_G[H]$ and $E_{\mathcal{F}[H]}\operatorname{Comm}_G[H]$. We also choose a model for $\underline{E}G$. Consider the G-pushout:

$$\coprod_{[H] \in I} G \times_{\operatorname{Comm}_G[H]} \underline{E} \operatorname{Comm}_G[H] \xrightarrow{i} \underline{E} G$$

$$\downarrow \coprod_{[H] \in I} id_G \times_{\operatorname{Comm}_G[H]} f_{[H]} \qquad \downarrow$$

$$\coprod_{[H] \in I} G \times_{\operatorname{Comm}_G[H]} E_{\mathcal{F}[H]} \operatorname{Comm}_G[H] \xrightarrow{i} \underline{E} G$$

where $f_{[H]}$ is a cellular $Comm_G[H]$ -map for every $[H] \in I$ and i is the inclusion. In these conditions, X is a model for $\underline{E}G$.

In practice, this theorem implies that the existence of good models for the proper classifying space of the commensurators and G, and also of the classifying spaces with respect to the families $\mathcal{F}[H]$ will lead to the knowledge of good models for $\underline{E}G$. Moreover, the push-out implies the existence of a long exact sequence in Bredon homology, and dimensional consequences that we will analyze in next section.

? (Sect:Bredon)?

2.2. Bredon homology. In this subsection we will briefly review the main definitions concerning Bredon homology. We follow the topological concise approach from [40], which we use in our computations.

Consider a discrete group G, \mathcal{F} a family of groups which is closed under conjugation and taking subgroups. Let $O_{\mathcal{F}}(G)$ be the *orbit category* whose objects are the homogeneous spaces G/K, $K \subset G$ with $K \in \mathcal{F}$, and whose morphisms are the G-equivariant maps. Then a left Bredon module N over $O_{\mathcal{F}}(G)$ is a covariant functor

$$N: O_{\mathcal{F}}(G) \to \mathbf{Ab}$$
,

where \mathbf{Ab} is the category of abelian groups.

Let N be a left Bredon module and X a G-CW-complex, and assume that all the stabilizers of the G-action belong to the family \mathcal{F} . Then the Bredon chain complex $(C_n^{\mathcal{F}}(X,N),\Phi_n)$ can be defined in the following way. For every $d \geq 0$, consider a set $\{e_i^d\}_{i \in I}$ of representatives of orbits of d-cells in X, and denote by $\operatorname{Stab}(e_i^d)$ the stabilizer of e_i^d . Then we define the n-th group of Bredon chains as $C_n^{\mathcal{F}}(X,N) = \bigoplus_{i \in I} N(G/\operatorname{Stab}(e_i^d)).$

Consider now a (d-1)-face of e_i^d , which can be given as ge for a certain (d-1)-cell e. Then we have an inclusion of stabilizers g^{-1} Stab $(e_i^d)g \subseteq \text{Stab}(e)$. As g^{-1} Stab $(e_i^d)g$ and $stab(e_i^d)$ are isomorphic, the previous inclusion induces an equivariant G-map $f: G/\operatorname{Stab}(e_i^d) \to G/\operatorname{Stab}(e)$. In turn, as N is a functor, we have an induced homomorphism $N(f): N(G/\operatorname{Stab}(e_i^d)) \to$ $N(G/\operatorname{Stab}(e))$. Taking into account that the boundary of e_i^d can be written as $\partial e_i^d = \sum_{j=1}^n g_j e_j^{d-1}$ for certain $g_j \in G$ and using linear extension to all representatives of equivariant d-cells, we obtain a differential $\Phi_d \colon C_d^{\mathcal{F}}(X,N) \to C_{d-1}^{\mathcal{F}}(X,N)$ for every d > 0. So we have the following definition:

Definition 2.4. The homology groups of the chain complex $(C_i^{\mathcal{F}}(X, N), \Phi_i)$ will be denoted by $H_i^{\mathcal{F}}(X, N)$ and called *Bredon homology groups* of X with coefficients in N with respect to the family \mathcal{F} .

We define $H_i^{\mathcal{F}}(G, N)$, the Bredon homology groups of G with coefficients in N with respect to the family \mathcal{F} as $H_i^{\mathcal{F}}(E_{\mathcal{F}}G, M)$.

These groups are preserved under G-equivariant homotopy equivalence.

Notation. When \mathcal{F} is the family of finite groups, we use indistinctly the notations $\underline{E}G$ or $E_{\mathcal{F}}G$. If \mathcal{F} is the family that only contains the trivial group, the superindex in the Bredon homology will be supressed, as it is ordinary homology in this case. It is worth noticing that there is an algebraic definition of $H_*^{\mathcal{F}}(G,M)$, however there is an isomorphism $H_*^{\mathcal{F}}(G,M) \simeq H_*^{\mathcal{F}}(E_{\mathcal{F}}G,M)$ between the algebraic and the topological definitions of Bredon homology [36, page 15]. To not overload the paper with unnecessary notation, we commonly denote this homology groups by $H_*^{\mathcal{F}}(G,M)$, although as said above, we mainly deal with the topological definition.

The cyclic group of order n will be denoted by C_n . When we want to consider its ring structure we might use \mathbb{Z}/\mathbf{n} . When n is prime, we might use \mathbb{F}_n to emphasize its field structure.

3. Artin groups of dihedral type

(Sect:Artin)

In this section we present the main features of the Artin groups of dihedral type that we will need in the remaining of the paper. We start with the definition of the groups:

(Defn:Artindih)

Definition 3.1. Let $n \ge 1$. By $\operatorname{prod}(x, y; n)$ we denote the word of length n that alternates x and y and starts with x. For example, $\operatorname{prod}(x, y; 3) = xyx$ and $\operatorname{prod}(x, y; 4) = xyxy$. With this notation, a dihedral Artin group of type n is the group

$$A_n = \langle a, b \mid \operatorname{prod}(a, b; n) = \operatorname{prod}(b, a; n) \rangle$$
.

The name "dihedral" comes from the associated Coxeter group,

$$\langle a, b \mid a^2 = b^2 = 1, \operatorname{prod}(a, b; n) = \operatorname{prod}(b, a; n) \rangle$$

which is the dihedral group of order 2n. Dihedral Artin groups are torsion-free, even more $A_n \cong F_{n-1} \rtimes \mathbb{Z}$, where F_k is a free group of rank k. To see this, one can check that the kernel of $A_n \to \mathbb{Z}$, $a, b \mapsto 1$ is free on rank n-1. In particular, Dihedral Artin groups satisfy the Farrell-Jones conjecture [7].

We are interested on understanding the commensurators of the virtually cyclic subgroups of A_n , for that we will use that A_n is also a central extension of a virtually free group.

As indicated above an important ingredient of our calculations will be the description of some commensurators inside these Artin groups of subgroups from the family of virtually cyclic groups (which in this case turn to be just cyclic groups). We start by observing that for any virtually cyclic subgroup H of a group G, and any $h \in H$ of infinite order, we have that

$$\operatorname{Comm}_G[H] = \operatorname{Comm}_G(\langle h \rangle) = \{ g \in G \mid gh^m g^{-1} = h^n \text{ for some } n, m \in \mathbb{Z} - \{0\} \}.$$

We will be interested in computing commensurators up to isomorphism, and we will use the fact that commensurators of conjugated subgroups are conjugated. Let us denote by π the

natural projection map $\pi: A_n \to \overline{A_n} := A_n/Z(A_n)$. Then, for every $g \in A_n$,

$$\pi(\operatorname{Comm}_{A_n}[\langle g \rangle]) \leqslant \operatorname{Comm}_{\overline{A_n}}[\langle \pi(g) \rangle],$$

and

$$Z(A_n) \leqslant \operatorname{Comm}_{A_n}[\langle g \rangle].$$

We will prove the following.

 $\mathtt{em} : \mathtt{commensurators}
angle$

Lemma 3.2. Let A_n be a dihedral Artin group and $g \in A_n$ of infinite order. Then

- (1) If $\langle g \rangle \cap Z(A_n) \neq \{1\}$ then $Comm_{A_n}[\langle g \rangle] = A_n$ and $Z(A_n) \in [H]$,
- (2) If $\langle g \rangle \cap Z(A_n) = \{1\}$ then $Comm_{A_n}[\langle g \rangle] \cong \mathbb{Z}^2$ and there is $\langle g' \rangle \in [\langle g \rangle]$ that is a direct factor of $Comm_{A_n}[\langle g \rangle]$.

Proof. It is a well-known fact that the centers of the dihedral Artin groups have different shape depending on the parity of n. Hence, we should divide the study of the commensurators of their cyclic subgroups in two subcases, even and odd.

Case n even: In the group

$$A_{2n} = \langle a, b \mid \operatorname{prod}(a, b; 2n) = \operatorname{prod}(b, a; 2n) \rangle$$
,

it is known that $Z(A_{2n}) = \langle (ab)^n \rangle$. Moreover, $A_{2n} \cong \langle x, y \mid x^{-1}y^nx = y^n \rangle$ via the isomorphism defined by $x \to b$, $y \mapsto ba$. Therefore $A_{2n}/Z(A_{2n}) = \langle x, y \mid x^{-1}y^nx = y^n, y^n = 1 \rangle = \langle \overline{x}, \overline{y} \mid \overline{y}^n = 1 \rangle \cong C_{\infty} * C_n$.

Since $\overline{A_{2n}}$ is hyperbolic (even more, virtually free), the commensurator of an infinite order element is virtually cyclic. (See for example [3, Theorem 2] bearing in mind that infinite cyclic subgroups of hyperbolic groups are quasi-convex).

Suppose first that $\pi(g)$ has infinite order. Then $\operatorname{Comm}_{\overline{A_{2n}}}[\langle \pi(g) \rangle]$ is infinite and virtually cyclic and hence $\pi(\operatorname{Comm}_{A_{2n}}[\langle g \rangle])$ is infinite and virtually cyclic. Since all infinite virtually cyclic subgroups of $\overline{A_{2n}}$ are infinite cyclic we have that $\pi(\operatorname{Comm}_{A_{2n}}[\langle g \rangle])$ is infinite cyclic. Therefore, $\operatorname{Comm}_{A_{2n}}[\langle g \rangle]$ is a central extension of $\mathbb Z$ by $\mathbb Z$ and hence $\operatorname{Comm}_{A_n}[\langle g \rangle] \cong \mathbb Z^2$. We can take g' as a pre-image of a generator of $\pi(\operatorname{Comm}_{A_{2n}}[\langle g \rangle])$.

Suppose now that $\pi(g)$ has finite order. Thus $\langle g \rangle \cap Z(A_{2n})$ is non-trivial and infinite, and thus $[\langle g \rangle] = [Z(A_{2n})]$ and $\operatorname{Comm}_{A_{2n}}[\langle g \rangle] = \operatorname{Comm}_{A_{2n}}Z(A_n) = A_{2n}$.

Case n odd: In the group

$$A_{2n+1} = \langle a, b \mid \operatorname{prod}(a, b; 2n+1) = \operatorname{prod}(b, a; 2n+1) \rangle$$

we have that $Z(A_{2n+1}) = \langle (ab)^{2n+1} \rangle$. Moreover, $A_{2n+1} \cong \langle x, y \mid xy^n = y^{n+1}x^{-1} \rangle$ via the isomorphism defined by $x \to b$, $y \mapsto ab$. Therefore

$$A_{2n+1}/Z(A_{2n+1}) = \langle x, y \mid xy^n x = y^{n+1}, y^{2n+1} = 1 \rangle = \langle x, y \mid (xy^n)^2 = 1 = y^{2n+1} \rangle$$
$$= \langle \overline{z}, \overline{y} \mid \overline{z}^2 = 1 = \overline{y}^{2n+1} \rangle \cong C_2 * C_{2n+1},$$

where \overline{z} denotes the class of $xy^nZ(A_{2n+1})$.

Since $\overline{A_{2n+1}}$ is hyperbolic and infinite virtually cyclic subgroups of $\overline{A_{2n+1}}$ are infinite cyclic we get, arguing as above, that if $\pi(g)$ has infinite order then $\operatorname{Comm}_{A_{2n+1}}[\langle g \rangle] \cong \mathbb{Z}^2$.

Suppose now that $\pi(g)$ has finite order. Arguing as above, $\operatorname{Comm}_{A_{2n+1}}[\langle g \rangle] = \operatorname{Comm}_{A_{2n+1}}[Z(A_{2n+1})] = A_{2n+1}$.

We recall the ordinary homology of these groups, which will be important in the remaining sections of the paper.

 $\langle \mathtt{OrdHom} \rangle$

Proposition 3.3. For $n \geq 2$.

If n is even, we have
$$H_0(A_n) = \mathbb{Z}$$
, $H_1(A_n) = \mathbb{Z} \oplus \mathbb{Z}$, $H_2(A_n) = \mathbb{Z}$ and $H_i(A_n) = 0$ for $i > 2$.
If n is odd, we have $H_0(A_n) = \mathbb{Z}$, $H_1(A_n) = \mathbb{Z}$ and $H_i(A_n) = 0$ for $i > 1$.

Proof. As dihedral Artin groups are one-relator groups, the Cayley complex (associated to the one-relator presentation) gives a 2-dimensional model for $K(A_n, 1)$ (see [32] or [10] for more general Artin groups). As A_n is not free, this implies that $\operatorname{gd} A_n = \operatorname{cd} A_n = 2$ for every n, and then in particular $H_i(A_n) = 0$ for $i \geq 3$. Moreover, as $K(A_n, 1)$ is connected, $H_0(A_n) = \mathbb{Z}$ for every n. The formulae for H_1 follow from performing abelianization to the groups. Finally, the results for the Schur multiplier H_2 are a consequence of Theorem 3.1 in [12].

4. Bredon homology of Artin groups of dihedral type

(Sect:BredonArtin)

In this section we start with the computation of homological invariants of the Artin groups of dihedral type, which is the main goal of this paper. The computation of the Farrell-Jones homology in the case of a general ring is usually very complicated. A powerful tool to undertake this computation is the G-equivariant version of the Atiyah-Hirzebruch spectral sequence, which is a spectral sequence of the 1st and 4th quadrant. In our case of interest, the E_2 -page of this sequence is given by Bredon homology of the classifying space $\underline{E}G$, and the E_{∞} -page encodes the Farrell-Jones homology. Let us make the last statement more precise.

Let G be a discrete group, \mathcal{F} its family of virtually cyclic subgroups, R a ring. For every $q \in \mathbb{Z}$, denote by $K_q(R[-])$ the covariant module over the orbit category $O_{\mathcal{F}}(G)$ that sends every (left) coset G/H to the K-theory group $K_q(R[H])$. In these conditions, the E_2 -page of the Atiyah-Hirzebruch spectral sequence that we will use is defined as $E_2^{p,q} = H_p^{\mathcal{F}}(\underline{\underline{E}}G, K_q(R[-]))$, for every $p \geq 0$ and $q \in \mathbb{Z}$, and converges to $H_{p+q}^G(\underline{\underline{E}}G, \mathbf{K}(R))$ the (p+q)-th group of Farrell-Jones homology of $\underline{\underline{E}}G$ with coefficients in R. An excellent source for more information about this sequence is [28].

Although the path to the computation is very clear, in general it is difficult to obtain explicit formulae for the Bredon homology of these classifying spaces. Different reasons for this are the complexity of the models for $\underline{E}G$ and the fact that the exact values of $K_q(R[H])$ are only known for very special instances $\overline{of} R$ and H. In fact, taking $R = \mathbb{Z}$ and H the trivial group, the groups $K_q(\mathbb{Z})$ are not completely listed, as their value in some cases depend on the solution of the Vandiver conjecture, which remains unsolved. See [48].

In this section we show that for Artin groups of dihedral type it is possible to describe to some extent the aforementioned Bredon homology groups, with coefficients in $K_q(R[-])$ for a general ring R (see Theorem 4.17); of course, the result strongly depends on the concrete shape of the K-theory groups of the group rings involved. The key to our computations is the description of the commensurators of the virtually cyclic subgroups of the groups A_n (see previous section), and mainly the Mayer-Vietoris sequence associated to the push-out of Lück-Weiermann model (Theorem 2.3), that was explicitly stated by Degrijse-Petrosyan in Section 7 of [14], for the cohomological case. We offer here the homological version, which is the one needed in our context. To avoid confusions, we denote Bredon homology with respect to the family of virtually cyclic groups as H_*^{vc} from now on.

Proposition 4.1. ([14, Proposition 7.1], see also [31, Theorem 2.3]) Let M be a left Bredon $\langle \text{mainMV} \rangle$ module over $O_{\mathcal{F}_{vc}}(A_n)$. There is an exact sequence:

$$\dots \to H^{vc}_{i+1}(A_n, M) \overset{g_1^{i+1}}{\to} \bigoplus_{[H] \in I} H_i^{Fin \cap \operatorname{Comm}_{A_n}[H]}(\operatorname{Comm}_{A_n}[H], M) \overset{g_2^i}{\to}$$

$$\stackrel{g_2^i}{\to} (\bigoplus_{[H] \in I} H_i^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], M)) \oplus H_i^{Fin}(A_n, M) \stackrel{g_3^i}{\to} H_i^{vc}(A_n, M) \to \dots$$

Before we undertake our computations of the homology groups, we will compute their geometric dimension gd with respect to the family of virtually cyclic groups.

? $\langle gd \rangle$? **Proposition 4.2.** The geometric dimension of A_n with respect of the family of virtually cyclic groups is 3.

Proof. As seen in Lemma 3.2, $A_n = \langle a, b | \operatorname{prod}(a, b; n) = \operatorname{prod}(b, a; n) \rangle$, contains subgroups isomorphic to $\mathbb{Z} \oplus \mathbb{Z}$ (for example $\operatorname{Comm}_{A_n}(\langle a \rangle)$). According to Example 5.21 in [31], this implies that $\operatorname{\underline{gd}} A_n \geq 3$.

On the other hand, as the Artin groups A_n are one-relator, Corollary 3 in [13] implies that $\underline{gd} A_n \leq 3$, and then $\underline{gd} A_n = 3$.

As stated, the strategy to compute $H_i^{vc}(A_n, M)$ goes through describing the different elements of the exact sequence of Proposition 4.1. We first understand the terms $H_i^{\mathcal{F}[H]}$ on Subsection 4.1. Then, in Subsection 4.2, we give a more concrete description of the different homology groups when we take coefficients in the K-theory. Finally, in Subsection 4.3 we study the homomorphisms g_1 and g_2 of the exact sequence of Proposition 4.1 and prove our main theorems.

- $\langle \mathtt{HiF[H]} \rangle$
 - 4.1. Computing the homology of the commensurators. Let us start the main calculations of this section. It is clear from the previous Mayer-Vietoris sequence that the computations of the ordinary homology of the commensurators (which includes the homology of A_n , as the center of A_n is virtually cyclic) and the homology of the commensurators with coefficients in $\mathcal{F}[H]$ will give us valuable information about the Bredon homology of A_n with respect to the family of virtually cyclic subgroups, so we will perform these calculations in the sequel. We start with the ordinary homology, which is straightforward and depends on the shape of the commensurators:
 - If $\operatorname{Comm}_{A_n}[H] \simeq \mathbb{Z} \oplus \mathbb{Z}$, we have $H_0(\operatorname{Comm}_{A_n}[H]) = H_2(\operatorname{Comm}_{A_n}[H]) = \mathbb{Z}$, $H_1(\operatorname{Comm}_{A_n}[H]) = \mathbb{Z} \oplus \mathbb{Z}$, $H_i(\operatorname{Comm}_{A_n}[H]) = 0$ for i > 2.
 - If $\operatorname{Comm}_{A_n}[H] \simeq A_n$, see Proposition 3.3.

We concentrate now in the case of $\mathcal{F}[H]$, which we recall is the family of subgroups of $\operatorname{Comm}_{A_n}[H]$ that are either finite or commensurable with H. Concretely, we intend to compute $H_i^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], M)$ for every non-trivial cyclic subgroup H of A_n . We have two cases, either $H \cap Z(A_n)$ is trivial or not. We will use the following Convention throughout the paper without explictly referring to it.

Convention 4.3. For [H] a class of infinite cyclic subgroups, we can always take a representative H that is normal in $Comm_{A_n}[H]$.

If $H \cap Z(A_n)$ is trivial, then Lemma 3.2 states that $H \subseteq \operatorname{Comm}_{A_n}[H]$. Moreover, by Lemma 3.2 we assume that H is a direct factor of $\operatorname{Comm}_{A_n}[H]$.

If $H \cap Z(A_n)$ is non-trivial, then $[H] = [Z(A_n)]$ and we will assume in this case that we choose $Z(A_n)$ as the representative of [H], and thus $H \subseteq \text{Comm}_{A_n}[H]$ again.

With this convention, the projection π : $Comm_{A_n}[H] \to Comm_{A_n}[H]/H$ is well-defined and $Comm_{A_n}[H]/H$ is isomorphic to C_{∞} , $C_{\infty} * C_n$ or $C_2 * C_{2n+1}$.

Let now M be a module over the orbit category of $\operatorname{Comm}_{A_n}[H]$ with respect to $\mathcal{F}[H]$, and $\pi^{-1}M$ the induced module over the orbit category of $\operatorname{Comm}_{A_n}[H]/H$ with respect to the family of finite groups. That is, for $K \leq \operatorname{Comm}_{A_n}[H]/H$ finite it is defined

$$\pi^{-1}M((\operatorname{Comm}_{An}[H]/H)/K) := M(\operatorname{Comm}_{An}[H]/\pi^{-1}(K)).$$

Observe that this assignation gives rise to a natural transformation of functors $\pi^{-1}M \to M$. In the next result, which is essentially [14, Lemma 4.2], but stated for homology instead of cohomology, we will see that this natural transformation induces an isomorphism in Bredon homology. It will be a powerful tool in our computations.

 $\langle induced \rangle$

Proposition 4.4. Let H be an infinite cyclic subgroup of A_n normal in its commensurator. For every $n \geq 0$, and every module M over the orbit category of $Comm_{A_n}[H]$ there is an isomorphism

$$H_i^{Fin}(\operatorname{Comm}_{A_n}[H]/H, \pi^{-1}M) \simeq H_i^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], M).$$

Moreover, every model for $\underline{\mathrm{E}}(\mathrm{Comm}_{A_n}[H]/H)$ is a model for $E_{\mathcal{F}[H]}\mathrm{Comm}_{A_n}H$, with the action induced by the quotient map $\mathrm{Comm}_{A_n}[H] \to \mathrm{Comm}_{A_n}[H]/H$.

Proof. The argument here is taken from the proof of [14, Lemma 4.2]. We write it here to make our paper more self-contained.

The projection π : $\operatorname{Comm}_{A_n}[H] \to \operatorname{Comm}_{A_n}[H]/H$ maps the family $\mathcal{F}[H]$ onto the family Fin of finite subgroups of the quotient $\operatorname{Comm}_{A_n}[H]/H$. Moreover, the pre-image $\pi^{-1}(K)$ for any finite group K of $\operatorname{Comm}_{A_n}[H]/H$ lies in $\mathcal{F}[H]$. Therefore, $\underline{\operatorname{E}}\operatorname{Comm}_{A_n}[H]/H$ is a model for $E_{\mathcal{F}[H]}\operatorname{Comm}_{A_n}[H]$, with the action induced by the projection π .

Consider now the spectral sequence associated to the short exact sequence $1 \to H \to \operatorname{Comm}_{A_n}[H] \xrightarrow{\pi} \operatorname{Comm}_{A_n}[H]/H \to 1$ for homology [35]. For every module M over the orbit category of $\operatorname{Comm}_{A_n}[H]$ we have

$$E_2^{p,q}(M) = H_p^{Fin}(\mathrm{Comm}_{A_n}[H]/H, H_q^{\mathcal{F}[H] \cap \pi^{-1}(-)}(\pi^{-1}(-), M)),$$

which converges to $E_{\infty}^{p,q}(M) = H_{p+q}^{\mathcal{F}[H]}(\mathrm{Comm}_{A_n}[H], M)$.

Observe that $E_2^{p,q}$ is trivial for $q \geq 1$, as for every finite subgroup $K < \text{Comm}_{A_n}[H]/H$, $\pi^{-1}(K)$ belongs to the family $\mathcal{F}[H] \cap \pi^{-1}(K)$, and then $H_q^{\mathcal{F}[H] \cap \pi^{-1}(-)}(\pi^{-1}(-), M)$ is zero. Thus, as $E_2^{i,0}(M) = H_i^{Fin}(\text{Comm}_{A_n}[H]/H, \pi^{-1}M)$ in the 0-th row of the sequence, we have

$$H_i^{Fin}(\operatorname{Comm}_{A_n}[H]/H, \pi^{-1}M) \simeq H_i^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], M).$$

for every $i \geq 0$. We are done.

The family of finite subgroups is easier to deal with than the family $\mathcal{F}[H]$, taking account of the previous proposition, the next step is to construct the corresponding classifying spaces for proper actions of the commensurators in A_n modulo the corresponding subgroups. We have the following:

- If $\operatorname{Comm}_{A_n}[H] \simeq \mathbb{Z} \oplus \mathbb{Z}$, then there is a representative of the class [H] that can be identified with one of these copies of \mathbb{Z} ; hence, we may assume that the inclusion $H \hookrightarrow \operatorname{Comm}_{A_n}[H] = \mathbb{Z} \oplus \mathbb{Z}$ is the inclusion of the first factor, and so $\operatorname{Comm}_{A_n}[H]/H$ is isomorphic to \mathbb{Z} . Then a model for $\underline{E}(\operatorname{Comm}_{A_n}[H]/H)$ is the straight line, and the action is by shifting.
- If $\operatorname{Comm}_{A_n}[H] \simeq A_n$, then $Z(A_n)$ is a representative of the class [H]. We have seen that $\operatorname{Comm}_{A_n}[H]/H = A_n/Z(A_n)$ is an amalgamated product of two cyclic groups, depending its concrete shape on the parity of n. In this case a tree model for $\underline{\operatorname{E}}(\operatorname{Comm}_{A_n}[H]/H)$ can be explicitly constructed.

If n is even, we have that $A_n/Z(A_n) \cong C_\infty * C_n$. Denote $C_\infty * C_n$ by $\overline{A_n}$. Let s be a generator of C_∞ . Then Bass-Serre theorem guarantees that the graph with vertex set $\overline{A_n}/C_n$, edge set $\overline{A_n}$ and incidence maps $\iota(g) = gC_\infty$ and $\tau(g) = gsC_\infty$ is a $\overline{A_n}$ -equivariant oriented tree.

If n is odd, we have that $A_n/Z(A_n) \cong C_2 * C_n$. Denote $C_2 * C_n$ by $\overline{A_n}$. Then Bass-Serre theorem guarantees that the graph with vertex set $\overline{A_n}/C_2 \sqcup \overline{A_n}/C_n$, edge set $\overline{A_n}$ and incidence maps $\iota(g) = gC_2$ and $\tau(g) = gC_n$ is a $\overline{A_n}$ -equivariant oriented tree.

Note that in both cases, the isotropy groups are the finite subgroups of $\overline{A_n}$ and they fix exactly a vertex; and therefore these are models for $\underline{E} \overline{A_n}$.

As the classifying spaces of the commensurators that we have described are all 1-dimensional, we will use the following result of Mislin:

Lemma 4.5 ([36], Lemma 3.14). Suppose that for a family \mathcal{F} of subgroups of a group G there is a tree model T for $E_{\mathcal{F}}G$. Let S_e be the stabilizer of the edge $e \in T$ and S_v the stabilizer of a vertex. Let N be a coefficient module. Then $H_i^{\mathcal{F}}(G,N) = 0$ for i > 1 and there is an exact sequence:

$$0 \to H_1^{\mathcal{F}}(G, N) \to \bigoplus_{[e]} N(G/S_e) \to \bigoplus_{[v]} N(G/S_v) \to H_0^{\mathcal{F}}(G, N) \to 0,$$

where [e] and [v] run over the G-orbits of edges and vertices of T, respectively.

 $\langle \text{diff} \rangle$ Remark 4.6. It is interesting to remark that the middle map $\bigoplus_{[e]} N(G/S_e) \to \bigoplus_{[v]} N(G/S_v)$ is induced by the (formal) border operation defined by $\partial[e] = [v_1] - [v_2]$, being v_1 and v_2 vertices of a representative e of [e]. The tree is assumed to be oriented.

In particular we obtain:

Corollary 4.7. For every $n \geq 2$, $i \geq 2$, $H \leqslant A_n$ infinite cyclic and every module M over the orbit category with respect to the family $\mathcal{F}[H]$, we have $H_i^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], M) = 0$.

Proof. Recall that by Proposition 4.4 $H_i^{\mathcal{F}[H]}(\mathrm{Comm}_{A_n}[H], M) \simeq H_i^{Fin}(\mathrm{Comm}_{A_n}[H]/H, \pi^{-1}M)$. As $\underline{\mathrm{E}}\,\mathrm{Comm}_{A_n}[H]/H$ is 1-dimensional, the result follows by Lemma 4.5.

So it remains to compute $H_0^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], M)$ and $H_1^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], M)$. We assume here a general coefficient module M, but the reader may keep in mind that our case of interest is $M(A_n/-) = K_q(R[-]), q \in \mathbb{Z}$.

 $\langle homology0and1 \rangle$

Proposition 4.8. Let $A_n = \langle a, b \mid \operatorname{prod}(a, b; n) = \operatorname{prod}(b, a; n) \rangle$ be a dihedral Artin group, let $H \leq A_n$ be an infinite cyclic group that is normal in its commensurator, let $\pi \colon \operatorname{Comm}_{A_n}[H] \to \operatorname{Comm}_{A_n}[H]/H$ be the natural projection. Let M be a module over the orbit category of $\operatorname{Comm}_{A_n}[H]$. Then

(i) If $H \cap Z(A_n)$ is trivial, then

$$H_0^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], M) = H_1^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], M) = M(\operatorname{Comm}_{A_n}[H]/H).$$

(ii) If $H = Z(A_n)$, n = 2k + 1 and if $f: M(A_n/Z(A_n)) \to M(A_n/\langle b(ab)^k \rangle) \oplus M(A_n/\langle ab \rangle)$ is induced by the natural projections, we have

$$H_0^{\mathcal{F}[H]}(A_n,M) = \operatorname{coker} f \quad \text{ and } \quad H_1^{\mathcal{F}[H]}(A_n,M) = \ker f.$$

(iii) If $H = Z(A_n)$ and n = 2k, we have

$$H_0^{\mathcal{F}[H]}(A_n, M) = M(A_n/\langle ab \rangle)$$
 and $H_1^{\mathcal{F}[H]}(A_n, M) = M(A_n/Z(A_n)).$

Note that in cases (ii) and (iii), $A_n = \text{Comm}_{A_n}[H]$.

Proof. In the case $H \cap Z(A_n)$ is trivial, the classifying space $\underline{E} \operatorname{Comm}_{A_n}[H]/H$ is the real line, there is one orbit of edges and one orbit of vertices, and the action is free. Then by Remark 4.6, the central map in that exact sequence is trivial and by Lemma 4.5,

$$H_1^{Fin}(\text{Comm}_{A_n}[H]/H, \pi^{-1}M) = \pi^{-1}M((\text{Comm}_{A_n}[H]/H)/\{1\})$$

and

$$H_0^{Fin}(\text{Comm}_{A_n}[H]/H, \pi^{-1}M) = \pi^{-1}M((\text{Comm}_{A_n}[H]/H)/\{1\}).$$

Using Proposition 4.4, case (i) follows.

Let us discuss the case $H = Z(A_n)$. Now $\operatorname{Comm}_{A_n}[H]/H = A_n/Z(A_n)$ is an amalgamated product of two finite cyclic groups or of a finite cyclic and an infinite cyclic group, depending on the parity of n. In both cases, there is a tree model for the classifying space for proper actions of the commutator modulo H. Denote $\operatorname{Comm}_{A_n}[H]/H$ by $\overline{A_n}$ for brevity.

We consider first the case n=2k+1. Here $\overline{A_n}=S*L$, with $S=\langle b(ab)^kZ(A_n)\rangle\cong C_2$ and $L=\langle abZ(A_n)\rangle\cong C_n$. The Bass-Serre tree has two orbits of vertices, with stabilizers conjugated to S or L, and one free orbit of edges. Then, by Lemma 4.5

$$(1) ? \underline{\mathsf{MVodd}}? \qquad H_1^{Fin}(\overline{A_n}, \pi^{-1}M) \hookrightarrow \pi^{-1}M(\overline{A_n}/\{1\}) \xrightarrow{f}$$

$$\stackrel{f}{\to} \pi^{-1}M(\overline{A_n}/S) \oplus \pi^{-1}M(\overline{A_n}/L) \twoheadrightarrow H_0^{Fin}(\overline{A_n}, \pi^{-1}M).$$

Hence, we obtain that

$$H_0^{Fin}(\overline{A_n}, \pi^{-1}M) = (\pi^{-1}M(\overline{A_n}/S) \oplus \pi^{-1}M(\overline{A_n}/L))/\text{Im } f$$

and

$$H_1^{Fin}(\overline{A_n}, \pi^{-1}M) = \text{Ker } f.$$

Observe that the two components of f are induced by the images of the projections $\overline{A_n}/\{1\}$ $\overline{A_n}/S$ and $\overline{A_n}/\{1\} \to \overline{A_n}/L$ by the functor $\pi^{-1}M$. The case (ii) follows by applying Proposition

Consider now the case n=2k. In this situation $\overline{A_n}=A_n/Z(A_n)$ is a free product G=S*L, with $S = \langle bZ(A_n) \rangle \simeq C_{\infty}$ and $L = \langle abZ(A_n) \rangle \simeq C_n$. We denote by s the element $bZ(A_n)$. The Bass-Serre tree for this group has one orbit of vertices with stabilizers conjugate to L and one orbit of edges with trivial stabilizers.

The exact sequence of Proposition 4.5 has the following form:

$$(2) ? \underline{\mathsf{MVeven}} ? H_1^{Fin}(\overline{A_n}, \pi^{-1}M) \hookrightarrow M(\overline{A_n}/\{1\}) \xrightarrow{f} \pi^{-1}M(\overline{A_n}/L) \twoheadrightarrow H_0^{Fin}(\overline{A_n}, \pi^{-1}M).$$

The function f is induced by $g\{1\} \in A_{2n}/\{1\} \mapsto gsL - gL$ by the functor M. And hence f is

Hence, we deduce that

$$H_1^{Fin}(\overline{A_n}, \pi^{-1}M) = \pi^{-1}M(\overline{A_n}/\{1\})$$

and

$$H_0^{Fin}(\overline{A_n}, \pi^{-1}M) = \pi^{-1}M(\overline{A_n}/L).$$

The result now follows by applying Proposition 4.4.

Observe that taking into account that $\overline{A_{2n}}$ is one-relator, this computation agrees with the result of Corollary 3.23 in [36], in the case of $M = R_{\mathbb{C}}$, the complex representation ring.

 $\langle \mathtt{Ktheory} \rangle$

4.2. Coefficients in K-theory groups. From the point of view of Farrell-Jones conjecture, the case of interest is the coefficients in the K-theory $K_q(R)$ of the group ring. We now describe all the homology groups of the exact sequence of Proposition 4.1 with coefficients in this module, except the groups $H^{vc}(A_n, K_q(R[-]))$ which will be studied in the next section.

We begin recalling the following definition.

?(defn:regular)?

Definition 4.9. A regular ring is a commutative noetherian ring such that in the localization at every prime ideal, the Krull dimension of the maximal ideal is equal to the cardinal of a minimal set of generators.

Examples of regular rings include fields (of dimension zero) and Dedekind domains. If R is regular then so is the polynomial ring R[x], with dimension one greater than that of R.

Before stating our results, we also need to recall the Bass-Heller-Swan decomposition in K-theory, which permits to decompose the K-theory of $R[\mathbb{Z}]$. For thorough approaches to algebraic K-theory the reader is referred to [6], [50] or [27].

?\(\text{BHS}\)? **Theorem 4.10** ([43], Theorem 9.8). Given a ring R and $q \in \mathbb{Z}$, there exists an isomorphism

$$K_q(R[\mathbb{Z}]) \simeq K_q(R) \oplus K_{q-1}(R) \oplus NK_q(R) \oplus NK_q(R),$$

which is natural in the ring R.

The additional terms $NK_q(R)$ are called the Nil-terms, and $NK_n(R)$ is defined as the kernel of the homomorphism induced in K_q by the homomorphism $R[t] \to R$ which sends t to 1. These terms vanish for a regular ring R, see [43, Section 9].

In our computations it will be important to understand the endomorphism ind_n of $K_q(\mathbb{Z})$ induced by multiplication by n in \mathbb{Z} . The references for the sequel are [21, Section 2], [45] and

According to Bass-Heller-Swan-decomposition, ind_n can be seen as a homomorphism

$$ind_n: K_q(R) \oplus K_{q-1}(R) \oplus NK_q(R) \oplus NK_q(R) \to K_q(R) \oplus K_{q-1}(R) \oplus NK_q(R) \oplus NK_q(R).$$

Here, by naturality of the decomposition, the image of $ind_n|_{K_q(R)}$ lies inside $K_q(R)$, the image of $ind_n|_{K_{q-1}(R)}$ lies inside $K_{q-1}(R)$ and analogously for the Nil-terms. Now, the restriction of ind_n to $K_q(R)$ is the identity and the restriction of ind_n to $K_{q-1}(R)$ is multiplication by n. By Farrell's trick [17], ind_n admits a transfer res_n such that $res_n \circ ind_n$ is multiplication by n. This implies the following:

 $\langle \mathtt{kernel} \rangle$

Proposition 4.11. In the previous notation, the kernel of ind_n is isomorphic to a direct sum $T_1(K_q(R)) \oplus T_2(K_q(R))$, where $T_1(K_q(R))$ is the n-torsion subgroup of $K_{q-1}(R)$ and $T_2(K_q(R))$ is a subgroup of the n-torsion subgroup of $NK_a(R) \oplus NK_a(R)$.

It should be remarked that the restriction of $ind_n|_{NK_q(R)}$ (sometimes call the Frobenius map in the literature) is related to the action of big Witt vectors in $NK_q(R)$, and is hard to describe in general. Anyhow, the previous proposition will be important when computing the Bredon homology of Artin groups of dihedral type.

The following propositions particularize our previous results when the coefficient module is K-theory.

For the sake of clarity, we maintain the notation from Proposition 4.1, although we are aware that sometimes can be found redundant.

(Anordinary)

Proposition 4.12. Let us consider an Artin group A_n of dihedral type, $q \in \mathbb{Z}$, R a ring. Then we have:

Proof. As A_n is torsion-free, Bredon homology with respect to the family of finite groups is in fact ordinary homology. The proposition is then obtained by applying Universal Coefficient Theorem [22, Theorem 3A.3] to the ordinary homology groups of A_n (Proposition 3.3), taking into account that the latter are torsion-free and that the K-theory groups are abelian.

We consider now the Bredon homology of the commensurators.

Proposition 4.13. Let us consider an Artin group A_n of dihedral type, $q \in \mathbb{Z}$, R a ring. Let $Comm_{A_n}[H]$ be the commensurator of a virtually cyclic group in A_n . Then:

• If
$$Comm_{A_n}[H] = \mathbb{Z} \oplus \mathbb{Z}$$
, then

$$H_i^{Fin\cap \operatorname{Comm}_{A_n}[H]}(\operatorname{Comm}_{A_n}[H], K_q(R[-])) = \begin{cases} K_q(R) & \text{if } i = 0, 2 \\ K_q(R) \oplus K_q(R) & \text{if } i = 1 \\ 0 & \text{otherwise.} \end{cases}$$

• If $Comm_{A_n}[H] = A_n$, then

$$H_i^{Fin\cap \operatorname{Comm}_{A_n}[H]}(\operatorname{Comm}_{A_n}[H],K_q(R[-]))=H_i^{Fin}(A_n,K_q(R))$$

and this case was described in the previous proposition.

Proof. Given a commensurator $\operatorname{Comm}_{A_n}[H]$, its Bredon homology with respect to the family $Fin \cap \operatorname{Comm}_{A_n}[H]$ is its ordinary homology. Then, as in the previous proposition, the result follows by applying the Universal Coefficient Theorem to the ordinary homology groups of the commensurators.

We now undertake the remaining case. In item (3) we follow the notation of Proposition 4.11. Moreover, we use the following notation.

 $\langle \mathtt{not} \colon \mathtt{CKR} \rangle$

Notation 4.14. For n odd, we denote by $C(K_q(R))$ the cokernel of the homomorphism

$$\tilde{f} \colon K_q(R[\mathbb{Z}]) \to K_q(R[\mathbb{Z}]) \oplus K_q(R[\mathbb{Z}])$$

induced by $f: R[\mathbb{Z}] \to R[\mathbb{Z}] \oplus R[\mathbb{Z}]$ induced in the first component by multiplication by 2 and in the second by multiplication by n = 2k + 1.

By Bass-Heller-Swan decomposition, $C(K_q(R))$ this is a quotient of $K_q(R) \oplus K_q(R) \oplus (\oplus_{i=1}^2 K_{q-1}(R)) \oplus (\oplus_{i=1}^4 NK_q(R))$. Moreover, \tilde{f} restricts to the component corresponding to $K_q(R)$ of the Bass-Heller-Swan decomposition of $K_q(R[\mathbb{Z}])$ to the diagonal map to $K_q(R) \oplus K_q(R)$, a component of $K_q(R[\mathbb{Z}]) \oplus K_q(R[\mathbb{Z}])$. Thus, $C(K_q(R))$ can be viewed as a quotient of $K_q(R) \oplus (\oplus_{i=1}^2 K_{q-1}(R)) \oplus (\oplus_{i=1}^4 NK_q(R))$ that can be identified in many cases, see Section 5.

(FHcomm)

Proposition 4.15. Let A_n be an Artin group of dihedral type, $q \in \mathbb{Z}$, R a ring. Consider for every non-trivial virtually cyclic $H \leq A_n$ and the family $\mathcal{F}[H]$. Then:

(1) If $i \geq 2$, then

$$H_i^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], K_q(R[-])) = 0.$$

(2) If $H \cap Z(A_n)$ is trivial, then

$$H_0^{\mathcal{F}[H]}(\text{Comm}_{A_n}[H], K_q(R[-])) = H_1^{\mathcal{F}[H]}(\text{Comm}_{A_n}[H], K_q(R[-])) = K_q(R[\mathbb{Z}]).$$

(3) If n is odd and $Comm_{A_n}[H] = A_n$, then

$$H_0^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], K_q(R[-])) = C(K_q(R))$$

ana

$$H_1^{\mathcal{F}[H]}(\text{Comm}_{A_n}[H], K_q(R[-])) = T_1(K_q(R)) \oplus T_2(K_q(R)).$$

(4) If n is even and $Comm_{A_n}[H] = A_n$, then

$$H_0^{\mathcal{F}[H]}(\text{Comm}_{A_n}[H], K_q(R[-])) = H_1^{\mathcal{F}[H]}(\text{Comm}_{A_n}[H], K_q(R[-])) = K_q(R[\mathbb{Z}]).$$

Proof. We will check every item separately.

- (1) It follows straightforward from Corollary 4.7.
- (2) The claim follows from item (i) in Proposition 4.8, taking into account that H is infinite cyclic an then $K_q(R[H]) = K_q(R[\mathbb{Z}])$.
- (3) Let n = 2k + 1. Consider the homomorphism $f: M(A_n/Z(A_n)) \to M(A_n/\langle b(ab)^k \rangle) \oplus M(A_n/\langle ab \rangle)$ defined in item (2) of Proposition 4.8. Taking $M(A_n/-) = K_q(R[-])$, we obtain a homomorphism $f_K: K_q(R[Z(A_n)]) \to K_q(R[\langle b(ab)^k \rangle]) \oplus K_q(R[\langle ab \rangle])$. Observe that the two components of f_K are respectively induced by the inclusions $Z(A_n) \hookrightarrow \langle b(ab)^k \rangle$ and $Z(A_n) \hookrightarrow \langle ab \rangle$, which are both inclusions $\mathbb{Z} \hookrightarrow \mathbb{Z}$ given respectively by multiplication by 2 and multiplication by n. Then, f_K can be seen as the homomorphism $K_q(R[\mathbb{Z}]) \to K_q(R[\mathbb{Z}]) \oplus K_q(R[\mathbb{Z}])$ induced by each multiplication in the corresponding component. According to Proposition 4.11, the kernel of the first component of this homomorphism is equal to $T_1(K_q(R)) \oplus T_2(K_q(R))$, while its cokernel is $C(K_q(R))$ by definition. The result now follows from item (2) in Proposition 4.8.
- (4) In this case $A_n = \operatorname{Comm}_{A_n}[H]$. The claim follows from item (3) in Proposition 4.8, taking into account that H is infinite cyclic and then $K_q(R[H]) = K_q(R[\mathbb{Z}])$.

\(\lambda\) morphism\(\rangle\)

4.3. Understanding the homomorphisms of Proposition 4.1. In our way to describe the Bredon homology of A_n with respect to the family of virtually cyclic groups, we should describe to some extent the homomorphisms that appear in the Mayer-Vietoris sequence of Proposition 4.1.

In the following we will use without explicit mention the previous three propositions, which identify the terms of the exact sequence. We also maintain the name of the homomorphisms in the sequence. Note that the superscript of the homomorphism g_i^k specifies the degree of the homology in the source of g_i^k . Moreover, when we need to refer to the j-th component of g_i^k , we will write g_{ij}^k . For example, g_{22}^1 is the second component of the homomorphism g_2^1 defined over the first homology group.

To prove our results, we need to analyze in detail the following homomorphism

$$(3) [g2] \qquad \bigoplus_{[H]} H_i^{Fin \cap \operatorname{Comm}_{A_n}[H]} (\operatorname{Comm}_{A_n}[H], M) \stackrel{g_2^i}{\to}$$

$$\stackrel{g_2^i}{\to} \left(\bigoplus_{[H]} H_i^{\mathcal{F}[H]} (\operatorname{Comm}_{A_n}[H], M) \right) \oplus H_i^{Fin}(A_n, M).$$

The homomorphism g_{21}^i is induced by the vertical left arrow $\sqcup_{[H]\in I} \mathrm{id}_{A_n} \times_{\mathrm{Comm}_{A_n}[H]} f_{[H]}$ of Lück-Weiermann push-out (4) below (see also Theorem 2.3). In turn, the homomorphism g_{22}^i is induced by the inclusion in the upper horizontal arrow of the push-out (4)

$$(4) \begin{tabular}{l} \hline (4) \begin{tabular}{l} \hline \end{tabular} (4) \begin{tabular}{l} \hline \end{$$

We will decompose the homomorphism g_2^i of Equation (3) into homomorphisms $g_{2[H]}^i$ which are the restriction of g_2^i to the factor of the domain corresponding to [H]. We will write

$$(5) [\underline{\mathtt{eq}} \underline{g}_{[H]}^i \colon H_i^{Fin \cap \mathrm{Comm}_{A_n}[H]}(\mathrm{Comm}_{A_n}[H], M) \to H_i^{\mathcal{F}[H]}(\mathrm{Comm}_{A_n}[H], M) \oplus H_i^{Fin}(A_n, M).$$

Note that we are making a slight abuse of notation, as the real codomain of $g_{2[H]}^i$ is the same as that of g_2^i , but we have chosen to write only the subgroup where the image of $g_{2[H]}^i$ lies. Moreover, when needed, we will further decompose $g_{2[H]}^i$ into $g_{21[H]}^i \oplus g_{22[H]}^i$ indicating the different factors of the image of $g_{2[H]}^i$. Observe that this notation is coherent with the previous one.

As all the commensurators are torsion-free, $H_i^{Fin\cap \operatorname{Comm}_{A_n}[H]}(\operatorname{Comm}_{A_n}[H], M)$ is ordinary homology with coefficients in $M(\operatorname{Comm}_{A_n}[H]/\{1\})$.

We now analyze the homomorphism g_2^i on the case $M = K_q(R[-])$. Before that, we introduce the following notation.

Notation 4.16. We will denote $N_q^{[H]} = K_{q-1}(R) \oplus NK_q(R) \oplus NK_q(R)$, where the superindex means that this group is associated to a concrete commensurability class [H].

We now will give more detailed descriptions of the homomorphism g_2^i and the cokernels introduced above.

- 4.3.1. The homomorphism g_2^2 when n odd: In this case we have that the codomain of g_2^2 is $(\bigoplus H_2^{\mathcal{F}[H]}(\mathrm{Comm}_{A_n}[H], M)) \oplus H_2^{Fin}(A_n, M) = \{0\}$, and therefore for all commensurability classes of infinite cyclic subgroups [H] the homomorphisms $g_{2[H]}^2$ are trivial.
 - 4.3.2. **The homomorphism** g_2^2 **when** n **even:** In this case we have that the codomain of g_2^2 is $(\bigoplus H_2^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], M)) \oplus H_2^{Fin}(A_n, M) = (\bigoplus_{[H]} \{0\}) \oplus K_q(R)$. Moreover, by Proposition 4.12 and Proposition 4.13, the domain of all $g_{2[H]}^2$ are the same, and we have

$$g_{2[H]}^2: K_q(R) \to \{0\} \oplus K_q(R).$$

Note that $g_{22[H]}^2$ is induced by the inclusion given by the upper arrow in the push-out. When $H = Z(A_n)$, we have that $Comm_{A_n}[H] = A_n$ and therefore this inclusion is the identity. Since $g_{22[Z(A_n)]}^2$ is surjective, we have that g_2^2 is surjective.

 $\langle g^{21} \rangle$ 4.3.3. The homomorphism g_2^1 when n is odd: Before we describe this case, we need to make some considerations. Recall that we assume that H is normal in $\operatorname{Comm}_{A_n}[H]$. Observe that in the models described in Section 4.1 for $\underline{E}(\operatorname{Comm}_{A_n}[H]/H)$, the stabilizers of the edges are trivial for every H. This means that if we consider the space $\underline{E}(\operatorname{Comm}_{A_n}[H]/H)$ as a model for $E_{\mathcal{F}[H]}\operatorname{Comm}_{A_n}[H]$ (see Proposition 4.4) the stabilizers of the edges are always isomorphic to H. On the other hand, in any model of $\underline{E}\operatorname{Comm}_{A_n}[H]$ the stabilizer of the edges should be trivial, as the action of $\operatorname{Comm}_{A_n}[H]$ is free. Then, the homomorphism induced by $f_{[H]}:\underline{E}\operatorname{Comm}_{A_n}[H]\to E_{\mathcal{F}[H]}\operatorname{Comm}_{A_n}[H]$ in the first chain group of the Bredon complex with coefficients in a module M takes every copy of $M(\operatorname{Comm}_{A_n}[H]/1)$ to a copy of $M(\operatorname{Comm}_{A_n}[H]/H)$ with the homomorphism induced by the inclusion of the trivial

group in H. In particular, if $M = K_q(R[-])$ for some q, the corresponding homomorphism $K_q(R) \to K_q(R[\mathbb{Z}])$ is given by the inclusion of $K_q(R)$ in the corresponding piece of the Bass-Heller-Swan decomposition of $K_q(R[\mathbb{Z}])$. This fact will be very useful in the sequel.

In the following we use the notation of Proposition 4.11 when needed. We have that

$$g_{2[Z(A_n)]}^1 \colon K_q(R) \to T_1(K_q(R)) \oplus T_2(K_q(R)) \oplus K_q(R)$$

and for H nontrivial, $[H] \neq [Z(A_n)]$

$$g^1_{2[H]} \colon K_q(R) \oplus K_q(R) \to K_q(R[\mathbb{Z}]) \oplus K_q(R).$$

Note that there are infinitely many commensurability classes of infinity cyclic subgroups different from the $[Z(A_n)]$ and hence infinitely many terms of this kind. We now examine the cases $[H] \neq [Z(A_n)]$ and $[H] = [Z(A_n)]$ separately.

If $[H] = [Z(A_n)]$ then we have $Comm_{A_n}[H] = A_n$.

In order to describe $g_{21[Z(A_n)]}^1$, consider the homomorphism

$$C_1^{Fin}(\underline{E}\operatorname{Comm}_{A_n}[H], K_q) \to C_1^{\mathcal{F}[H]}(E_{\mathcal{F}[H]}\operatorname{Comm}_{A_n}[H], K_q)$$

at the level of Bredon chains that induces $g_{21[Z(A_n)]}^1$ in homology. According to the previous considerations, this homomorphism is given by the inclusion $i: K_q(R) \hookrightarrow K_q(R[H]) = K_q(R[\mathbb{Z}])$ given by Bass-Heller-Swan decomposition. But by item (3) of Proposition 4.15, the image of this homomorphism is trivial in $H_1^{\mathcal{F}[H]}(E_{\mathcal{F}[H]}\operatorname{Comm}_{A_n}[H], K_q)$, and hence $g_{21[Z(A_n)]}^1$ is also trivial. In turn, $g_{22[Z(A_n)]}^1$ is the identity.

If $[H] \neq [Z(A_n)]$ then $\operatorname{Comm}_{A_n}[H] = \mathbb{Z}^2$, and $\operatorname{Comm}_{A_n}[H]/H$ is isomorphic to \mathbb{Z} . In this case $g^1_{21[H]}$ is given by a homomorphism $K_q(R) \oplus K_q(R) \to K_q(R[\mathbb{Z}])$, where we assume that the first component of the domain corresponds to H and the second to $Z(A_n)$. As the homomorphism $g^1_{21[H]}$ is induced in homology by the quotient homomorphism $\operatorname{Comm}_{A_n}[H] \to \operatorname{Comm}_{A_n}[H]/H$, the previous results imply that the first component of $g^1_{21[H]}$ is trivial, while the second, which corresponds to the center, identifies the copy of $K_q(R)$ in the Bass-Heller-Swan decomposition of $K_q(R[\mathbb{Z}])$.

On the other hand, $g_{22[H]}^1: (\mathbb{Z} \oplus \mathbb{Z}) \otimes K_q(R) \to \mathbb{Z} \otimes K_q(R)$ is defined by the abelianization of the inclusion $H_1(\operatorname{Comm}_{A_n}[H]) \to H_1(A_n)$ in the first component of the tensor product and by the identity in the second.

Now since the image of $g_{2[Z(A_n)]}^1$ is precisely given by the copy of $K_q(R)$ that corresponds to $H_1(A_n, K_q)$, the previous computations imply that the cokernel of g_2^1 is then equal to $\bigoplus_{[H]\neq[Z(A_n)]} N_q^{[H]}) \oplus T_1(K_q(R)) \oplus T_2(K_q(R))$.

4.3.4. The homomorphism g_2^1 when n is even: This homomomorphism is defined in equation (5) and, according to Propositions 4.12, 4.13 and 4.15, its first component is given by

$$g_{2[H]}^1 \colon K_q(R) \oplus K_q(R) \to K_q(R[\mathbb{Z}]) \oplus (K_q(R) \oplus K_q(R))$$

for every commensurability class [H]. Let us describe this component with more detail.

We consider first the case $[H] \neq [Z(A_n)]$. Here, the same argument as in the odd case proves that the component of $g^1_{21[H]}$ given by the center is the inclusion $K_q(R) \hookrightarrow K_q(R[\mathbb{Z}])$ via Bass-Heller-Swan decomposition, and the other component is trivial.

Also when $[H] \neq [Z(A_n)]$ the homomorphism $g_{22[H]}^1$ is identified (via Proposition 4.13 and the Universal Coefficient Theorem) with a homomorphism $(\mathbb{Z} \oplus \mathbb{Z}) \otimes K_q(R) \to (\mathbb{Z} \oplus \mathbb{Z}) \otimes K_q(R)$, which comes, as above, from tensoring with $K_q(R)$ the homomomorphism $H_1(\text{Comm}_{A_n}[H]) \to H_1(A_n)$ given by abelianization of the inclusion of the commensurator in A_n .

Now we consider $[H] = [Z(A_n)]$. As in the previous case, the homomorphism $g_{21[H]}^1$ can be described as:

$$H_1^{Fin}(A_n, K_q(-)) \to H_1^{F[H]}(A_n, \pi^{-1}K_q(-)) \cong H_1^{Fin}(A_n/Z(A_n), \pi^{-1}K_q(-)).$$

Now recall from Proposition 4.8 that

$$H_1^{Fin}(A_n/Z(A_n), \pi^{-1}K_q(-)) = \pi^{-1}K_q(R[\{1\}]) = K_q(R[Z(A_n)]) = K_q(R[Z]),$$

where $Z(A_n)$ is interpreted here as the stabilizer of the unique A_n -class of edges in the model of $E_{\mathcal{F}[H]}A_n$ described in Section 4.1. On the other hand, the two copies of $K_q(R)$ in $H_1(A_n, K_q(R))$ come from taking values of the module $K_q(R[-])$ on the trivial group, interpreted as the stabilizer of two different A_n -classes of edges in a model of EA_n . Again taking into account our previous considerations about stabilizers, the two components of $g^1_{21[Z(A_n)]}: K_q(R) \oplus K_q(R) \to K_q(R[\mathbb{Z}])$ induce inclusion of the $K_0(R)$ via Bass-Heller-Swan decomposition. On the other hand, it is clear that $g^1_{22[Z(A_n)]}$ is the identity.

As $H_1(A_n)$ is a free abelian group of rank 2 generated by the images of a and b under abelianization, we may assume that the two copies of $K_q(R)$ in the image of $g_{2[H]}^1$ correspond respectively to these two copies of \mathbb{Z} , after tensoring with $K_q(R)$. Now observe that if $H = \langle a \rangle$, the image of the restriction of $g_{22[H]}^1$ to the homology of its commensurator is exactly the first of the two copies of $K_q(R)$, while the image of the restriction of $g_{22[H]}^1$ to the homology of the commensurator of $H = \langle b \rangle$ is the other copy. As the restrictions of $g_{21[H]}^1$ to the homology of these commensurators are trivial, we obtain that $\{0\} \oplus K_q(R) \oplus K_q(R)$ lies in the image of $g_{2[H]}^1$. In fact, the description of $g_{21[H]}^1$ for $H = Z(A_n)$ implies that the copy of $K_q(R)$ inside $K_q(R[\mathbb{Z}])$ is also in the image, and now it is easy to conclude that the image of $g_{2[H]}^1$ is in fact $\bigoplus_{I=1}^n K_q(R) \oplus K_q(R) \oplus K_q(R)$, corresponding the big direct sum to the copies of $K_q(R)$ included in each copy of $K_q(R[\mathbb{Z}])$, and the remaining two copies corresponding to the ordinary homology of A_n . In particular, we have that coker $g_2^1 = \bigoplus_{I=1}^n N_q^{[H]}$.

4.3.5. The homomorphism g_2^0 for every n: Similar considerations to those of the beginning of Section 4.3.3 hold here. If x is a vertex of $\underline{E}\mathrm{Comm}_{A_n}[H]$ such that the stabilizer of $f_{[H]}(x)$ is infinite cyclic (here $f_{[H]}$ is the function of (4)), then the induced map $K_q(R) \to K_q(R[\mathbb{Z}])$ induces the injection in the correspondent component of Bass-Heller-Swan decomposition. This is always the case except when n is odd, $H = Z(A_n)$ and $f_{[H]}(x)$ has the shape gC_{∞} if we consider $\underline{E}(\mathrm{Comm}_{A_n}[H]/H)$ as a $\mathrm{Comm}_{A_n}[H]/H$ -complex. This will be enough to describe g_2^0 to the extent we need.

We begin describing $g_{2[H]}^0$ according to the different prossibilities of [H] and n.

When $H \neq Z(A_n)$, the results of Propositions 4.12, 4.13 and 4.15 imply that $g_{2[H]}^0$ is defined in the following way:

$$g_{2[H]}^0 \colon K_q(R) \to K_q(R[\mathbb{Z}]) \oplus K_q(R).$$

By the previous considerations, the homomorphism $g_{21[H]}^0$ identifies $K_q(R)$ as the corresponding direct summand of $K_q(R[\mathbb{Z}])$ in the Bass-Heller-Swan decomposition, while $g_{22[H]}^0$ is induced by the inclusion $\operatorname{Comm}_{A_n}[H] \hookrightarrow A_n$, counts the number of connected components of the classifying space, and then is the identity.

When n is even and $H = Z(A_n)$, we have that $H_0^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], K_q(-)) = K_q(R[\mathbb{Z}])$ by Proposition 4.15, and this copy of \mathbb{Z} corresponds to a stabilizer which is isomorphic to C_n , a cyclic group of order n. Then, by the previous considerations, the homomorphism $g_{2[H]}^0$ behaves as in the case n even and $H \neq Z(A_n)$.

In the case n odd and $H = Z(A_n)$, again by Proposition 4.15 we have that $H_0^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], K_q(-)) = C(K_q(R))$, where $C(K_q(R))$ was defined in Notation 4.14. Then $g_{21[H]}^0$ is defined as a homomorphism

$$g_{2[H]}^0 \colon K_q(R) \to C(K_q(R)) \oplus K_q(R).$$

Remark from the definition of $C(K_q(R))$ that this group is the direct sum of $K_q(R)$ with quotients of $\bigoplus_{i=1}^2 K_{q-1}(R)$ and $\bigoplus_{i=1}^4 NK_q(R)$, and that this copy of $K_q(R)$ comes from the identification of the respective copies of $K_q(R)$ that appear in the Bass-Heller-Swan decomposition of $K_q(R[\langle b(ab)^k \rangle])$ and $K_q(R[\langle ab \rangle])$. These copies correspond respectively to the stabilizers C_2 and C_{2k+1} in $\underline{E}(\operatorname{Comm}_{A_n}[H]/H)$ (as a model for the $\operatorname{Comm}_{A_n}[H]/H$ -action). Then, the previous considerations about stabilizers imply that $g^0_{21[H]}$ maps this $K_0(R)$ isomorphically to the mentioned copy of itself inside $C(K_q(R))$. In turn, $g^0_{22[H]}$ is again the identity, arguing as in the even case.

With the previous information, we can describe $\operatorname{coker} g_2^0$.

Now if n is even, observe that the source of g_2^0 is $\bigoplus_{[H]} K_q(R)$ and the codomain is $(\bigoplus_{[H]} K_q(R[\mathbb{Z}])) \oplus K_q(R)$. Moreover, the image of every $g_{2[H]}^0$ consists in a copy of $K_q(R)$ inside $K_q(R[\mathbb{Z}])$ (again the copy that appears in the Bass-Heller-Swan decomposition), and the copy of $K_q(R)$ that corresponds to $H_0^{Fin}(A_n, M)$, which is fixed for any choice of H. Hence, the cokernel of the homomorphism g_2^0 is isomorphic to the quotient of a direct sum of copies of $K_q(R[\mathbb{Z}])$ (indexed by [H]) by the identification of all the copies $K_q(R)$ which are the images of the homomorphisms $g_{22[H]}^0$. Observe that the copies of $K_q(R[\mathbb{Z}])$ are "glued" by the copy of $K_q(R)$ that corresponds to $H_0^{Fin}(A_n, M)$, and hence we have

$$\operatorname{coker} g_2^0 = (\bigoplus_{[H]} N_q^{[H]}) \oplus K_q(R).$$

When n is odd, we only need to take into account that in the case of $H = Z(A_n)$ the role of $K_q(R[\mathbb{Z}])$ in the codomain is played by $C(K_q(R))$. Then, using the same argument as in the previous case and denoting by $\overline{C}(K_q(R))$ the quotient of $C(K_q(R))$ under the copy of $K_q(R)$ in the Bass-Heller-Swan decomposition, we obtain that:

$$\operatorname{coker} g_2^0 = (\bigoplus_{[H] \neq [Z(A_n)]} N_q^{[H]}) \oplus K_q(R) \oplus \overline{C}(K_q(R)).$$

Observe that g_2^0 is a monomorphism for every n, as the inclusions of $K_q(R)$ in $K_q(R[\mathbb{Z}])$ and $C(K_q(R))$ are so.

Now we can describe the Bredon homology of A_n with respect to the family of virtually cyclic groups. Observe that the (co)kernels of the statement have been previously described, and that the groups $N_q^{[H]}$ were defined in Section 4.3.1.

 $\langle \mathtt{Thm} : \mathtt{Bredon} \rangle$

Theorem 4.17. Let A_n be an Artin group of dihedral type. In the previous notation, we have the following:

(1)
$$H_i^{vc}(A_n, K_q(R[-])) = \{0\} \text{ for } i \ge 4.$$

(2) $H_3^{vc}(A_n, K_q(R[-])) = \begin{cases} \bigoplus_{[H] \ne [Z(A_n)]} K_q(R) & n \text{ odd} \\ \ker g_2^2 & n \text{ even.} \end{cases}$
(3) $H_2^{vc}(A_n, K_q(R[-])) = \ker g_2^1.$

$$(3) \ H_2^{vc}(A_n, K_q(R[-])) = \ker g_2^1.$$

$$(4) \ H_1^{vc}(A_n, K_q(R[-])) = \operatorname{coker} g_2^1 = \begin{cases} (\bigoplus_{[H] \neq [Z(A_n)]} N_q^{[H]}) \oplus T_1(K_q(R)) \oplus T_2(K_q(R)) & n \ odd \\ \bigoplus_{[H]} N_q^{[H]} & n \ even \end{cases}$$

$$(5) \ H_0^{vc}(A_n, K_q(R[-])) = \operatorname{coker} g_2^0 = \begin{cases} (\bigoplus_{[H] \neq [Z(A_n)]} N_q^{[H]}) \oplus K_q(R) \oplus \overline{C}(K_qR) & n \text{ odd} \\ (\bigoplus_{[H]} N_q^{[H]}) \oplus K_q(R) & n \text{ even.} \end{cases}$$

Proof. The proof is based on the sequence of Proposition 4.1 and the previous homological computations. We will check the five items in a separate way.

- (1) By Proposition 4.12, Proposition 4.13 and Proposition 4.15 the sequence of Proposition 4.1 is identically trivial to the left of g_3^3 .
- (2) As the exact sequence of Proposition 4.1 is identically trivial to the left of g_3^3 , the map g_3^3 is trivial and by exactness, $H_3^{vc}(A_n, K_q(R[-]))$ is isomorphic to the kernel of g_2^2 . This completes the even case.

For the case n odd, we claim that g_1^3 is an isomorphism. Indeed, from Proposition 4.12 and Proposition 4.15, the term $\bigoplus H_2^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], M)) \oplus H_2^{Fin}(A_n, M) = \{0\}$ and in particular $\operatorname{Im}(g_2^2) = \{0\}$. Since g_1^3 is injective, the claim follows. By Proposition 4.13, the specific description of the odd case follows.

(3) There are two different arguments depending if n is odd or even.

For n odd, as the term $\bigoplus H_2^{\mathcal{F}[H]}(\operatorname{Comm}_{A_n}[H], M) \oplus H_2^{Fin}(A_n, M)$ is trivial, the statement is a direct consequence of the exactness of the sequence of Proposition 4.1.

For n even, we have seen in the discussion above about g_2^2 that this map is surjective, and hence g_3^2 is the trivial map. This implies that $H_2^{vc}(A_n, K_q(R[-]))$ is the image of g_1^2 , or equivalently, the kernel of g_2^1 .

- (4) The previous description of the homomorphisms in the Mayer-Vietoris sequence proves that g_2^0 is a monomorphism, and hence g_3^1 is surjective and $H_1^{vc}(A_n, K_q(R[-]))$ the cokernel of g_2^1 , which has been described above in terms of the summands $N_q^{[H]}$.
- (5) Since the Mayer-Vietoris sequence ends at $H_0^{vc}(A_n, K_q(R[-]))$ this term is equal to the image of g_3^0 which is equal to the cokernel of g_2^0 that was described above.

As said in the introduction, this theorem opens the door to concrete computations of Bredon homology of A_n with respect to the family of virtually cyclic groups, provided there is available information about the K-theory of the coefficient ring. In particular, when R is a regular ring the absence of Nil-terms and negative K-theory groups make the calculations easier and more precise. For instance, we have

 $\langle cor: K_0 \rangle$ Corollary 4.18. Let A_n , n > 2 be an Artin group of dihedral type. Let R be a regular ring. Then $K_0(RA_n) = K_0(R)$.

Proof. As R is regular, $K_i(R)$ vanishes for i < 0 and also the Nil-Terms of the Bass-Heller-Swan decomposition. In particular, by Theorem 4.17, $H_0^{vc}(A_n, K_0(R[-])) = K_0(R)$. Moreover, as $K_i(R)$ vanishes for i < 0 we see by Theorem 4.17, that $H_j^{vc}(A_n, K_i(R[-])) = \{0\}$ for i < 0. Then the E_2 -page of the Atiyah-Hirzebruch spectral sequence is concentrated in the non-negative part of the 0th, 1st, 2nd and 3rd columns, and $E_{\infty}^{0,0} = E_2^{0,0} = K_0(R)$.

In particular, this implies that every finitely dominated CW-complex whose fundamental group is A_n has the homotopy type of a finite CW-complex.

In the following section we compute these Bredon homology groups for different choices of the ring, including some non-regular ones.

5. Computations of $H_i^{vc}(A_n, K_q(R[-]))$ for several coefficient rings

(Sect:concrete)

In this section we use Theorem 4.17 to describe $H_i^{vc}(A_n, K_q(R[-]))$ for some instances of the ring R, both regular $(\mathbb{Z}, \mathbb{F}_q)$ and non-regular $(\mathbb{Z}[\mathbb{Z}/2], \mathbb{Z}[\mathbb{Z}/2 \times \mathbb{Z}/2]$ and $\mathbb{Z}[\mathbb{Z}/4])$. We recall that all these groups give information about the E^2 -term of the corresponding Atiyah spectral sequence. We point out that in the regular cases many groups can be computed, because the K-theory of \mathbb{Z} is nearly known and the K-theory of \mathbb{F}_q is known (see the corresponding examples below). In the non-regular framework, by contrast, it is very difficult to find concrete descriptions of $K_q(R)$ for q > 1, or of the corresponding Nil-terms.

Recall that the groups $H_i^{vc}(A_n, K_0(R[-]))$ are trivial for $i \geq 4$ and any ring R.

Example 5.1. First we compute the Bredon homology of A_n with respect to the family of virtually cyclic subgroups, taking as coefficients $K_q(\mathbb{Z}[-])$, for q = 0, 1, 2.

We need in our computations the lower algebraic K-theory groups of the integers. The groups $K_q(\mathbb{Z})$ are known for all $q \leq 7$ and all $q \geq 8$ such that $q \not\equiv 0 \mod 4$, see [48, page 2]. For example, the first values of $K_q(\mathbb{Z})$ are given in the following table:

\overline{q}	< 0	0	1	2	3	4	5	6	7
$K_q(\mathbb{Z})$	0	\mathbb{Z}	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/48$	0	\mathbb{Z}	0	$\mathbb{Z}/240$

Observe that, as \mathbb{Z} is regular, the Nil-terms in the Bass-Heller-Swan decomposition are trivial, and then for every $i \in \mathbb{N}$, $K_i(\mathbb{Z}[\mathbb{Z}]) = K_i(\mathbb{Z}) \oplus K_{i-1}(\mathbb{Z})$ and $N_q^{[H]} = K_{q-1}(\mathbb{Z})$. Moreover, also by regularity, $K_i(\mathbb{Z}) = 0$ if $i \leq -1$.

Taking into account of all these considerations, the previous theorem implies the following. For q=0, we have:

- (1) $H_3^{vc}(A_n, K_0(\mathbb{Z}[-])) \simeq H_2^{vc}(A_n, K_0(\mathbb{Z}[-])) \simeq \bigoplus_{\aleph_0} \mathbb{Z}.$ (2) $H_1^{vc}(A_n, K_0(\mathbb{Z}[-])) = 0.$

(3) $H_0^{vc}(A_n, K_0(\mathbb{Z}[-])) = \mathbb{Z}.$

Now for q=1,

- $(1) \ H_3^{vc}(A_n, K_1(\mathbb{Z}[-])) \simeq H_2^{vc}(A_n, K_1(\mathbb{Z}[-])) \simeq \bigoplus_{\aleph_0} \mathbb{Z}/2.$ $(2) \ H_1^{vc}(A_n, K_1(\mathbb{Z}[-])) = \bigoplus_{\aleph_0} \mathbb{Z}.$ $(3) \ H_0^{vc}(A_n, K_1(\mathbb{Z}[-])) = (\bigoplus_{\aleph_0} \mathbb{Z}) \oplus \mathbb{Z}/2.$

And finally, for q=2, all these groups are isomorphic to $\bigoplus_{\aleph_0} \mathbb{Z}/2$.

Let us briefly explain these computations. We consider first q = 0. For $H_3^{vc}(A_n, K_0(\mathbb{Z}[-]))$, the odd case is immediate. In the even case, $H_3^{vc}(A_n, K_1(\mathbb{Z}[-]))$ is the kernel of a homomorphism $\bigoplus_{\aleph_0} \mathbb{Z} \to \mathbb{Z}$, and then isomorphic to $\bigoplus_{\aleph_0} \mathbb{Z}$. This easy argument will be frequently used in the examples of this section without express mention.

We now deal with $H_2^{vc}(A_n, K_0(\mathbb{Z}[-]))$. We only deal with the odd case, the even one is very similar. First observe that the kernel of g_2^1 is isomorphic to $\bigoplus_{\aleph_0} \mathbb{Z}$. More precisely, taking account the description of Section 4.3.3 and the values of the $K_i(\mathbb{Z})$, g_2^1 (for n odd) is defined in the following way:

$$g_2^1: (\bigoplus_{[H] \neq [Z(A_n)]} \mathbb{Z}^2) \oplus \mathbb{Z} \to (\bigoplus_{[H] \neq [Z(A_n)]} \mathbb{Z}) \oplus \mathbb{Z}.$$

Here the kernel of $g_{21[H]}^1$ is isomorphic to \mathbb{Z} for every $[H] \neq [Z(A_n)]$. As there is an infinite number of such commensurators, we conclude that the kernel should be also infinite, and then $H_2^{vc}(A_n, K_1(\mathbb{Z}[-])) = \bigoplus_{\aleph_0} \mathbb{Z}.$

Observe that, as $K_{-1}(\mathbb{Z}) = 0$ and \mathbb{Z} is regular, $N_0^{[H]} = T_i(K_0\mathbb{Z}) = \overline{C}(K_0\mathbb{Z}) = 0$. Now the values of $H_1^{vc}(A_n, K_0(\mathbb{Z}[-]))$ and $H_0^{vc}(A_n, K_0(\mathbb{Z}[-]))$ are easily deduced from items (4) and (5) of Theorem 4.17.

Now take q=1. For $H_3^{vc}(A_n,K_1(\mathbb{Z}[-]))$ and $H_2^{vc}(A_n,K_1(\mathbb{Z}[-]))$ it is argued as in the previous case, taking into account that every subgroup of an \mathbb{F}_2 -vector space is again an \mathbb{F}_2 vector space. Now observe that $N_1^{[H]} = K_0(\mathbb{Z}) = \mathbb{Z}$ by the Bass-Heller-Swan decomposition, and regularity and the fact that $K_0(\mathbb{Z}) = \mathbb{Z}$ is torsion-free imply that $T_1(K_1\mathbb{Z}) = 0$, $T_2(K_1\mathbb{Z}) = 0$ 0 and $\overline{C}(K_1\mathbb{Z}) = \mathbb{Z}$. The values of $H_1^{vc}(A_n, K_1(\mathbb{Z}[-]))$ and $H_0^{vc}(A_n, K_1(\mathbb{Z}[-]))$ follow again from items (4) and (5) of Theorem 4.17.

Finally, for q=2, the values of the homology are immediately implied by regularity and the fact that $K_2(\mathbb{Z}) = K_1(\mathbb{Z}) = \mathbb{Z}/2$.

Example 5.2. Now we will compute the groups $H_i^{vc}(A_n, K_q(\mathbb{F}_2[-]))$, for $0 \le q \le 3$. First, the following table (see [39]) includes the algebraic K-groups that are necessary in our computations:

q	< 0	0	1	2	3
$K_q(\mathbb{F}_2)$	0	\mathbb{Z}	0	0	$\mathbb{Z}/3$

As \mathbb{F}_2 is regular, the same considerations about the Nil-terms and the negative K-groups apply also in this case. Hence, again as a direct consequence of Theorem 4.17 we have the following results. For q=0 and every $i\in\mathbb{N},\ H_i^{vc}(A_n,K_0(\mathbb{F}_2[-]))=H_i^{vc}(A_n,K_0(\mathbb{Z}[-])),$ because $K_0(\mathbb{Z}) = K_0(\mathbb{F}_2)$ and then $K_0(\mathbb{F}_2[-]) = K_0(\mathbb{Z}[-])$ because both have the same Bass-Heller-Swan decomposition.

For q = 1, we have $H_3^{vc}(A_n, K_1(\mathbb{F}_2[-])) = H_2^{vc}(A_n, K_1(\mathbb{F}_2[-])) = 0$, as $K_1(\mathbb{F}_2) = 0$. On the other hand, as $N_1^{[H]} = \mathbb{Z}$ for every H, $H_1^{vc}(A_n, K_1(\mathbb{F}_2[-])) = H_0^{vc}(A_n, K_1(\mathbb{F}_2[-])) = \bigoplus_{\aleph_0} \mathbb{Z}$.

For q=2, the triviality of $K_2(\mathbb{F}_2)$ and $K_1(\mathbb{F}_2)$ implies the triviality of $H_i^{vc}(A_n, K_2(\mathbb{F}_2[-]))$ for every i.

Finally, for q=3, we have $H_2^{vc}(A_n,K_3(\mathbb{F}_2[-])) \simeq H_3^{vc}(A_n,K_3(\mathbb{F}_2[-])) = \bigoplus_{\aleph_0} \mathbb{Z}/3$. As the groups $N_3^{[H]}$ are trivial and $K_2(\mathbb{F}_2)$ is so, we obtain that $H_1^{vc}(A_n,K_3(\mathbb{F}_2[-])) = \mathbb{Z}/3$ is trivial, and $H_0^{vc}(A_n,K_3(\mathbb{F}_2[-])) = \mathbb{Z}/3$.

We follow with some non-regular examples, namely the rings $\mathbb{Z}[\mathbb{Z}/2]$, $\mathbb{Z}[\mathbb{Z}/2 \times \mathbb{Z}/2]$ and $\mathbb{Z}[\mathbb{Z}/4]$, which we will respectively denote by R_1 , R_2 and R_3 . We will compute the groups $H_i^{vc}(A_n, K_q(R_j[-]))$, for $0 \le i \le 3$, q = 0, 1 and $1 \le j \le 3$. In order to do this we will need the values of their lower algebraic K-theory groups, as well as the Nil groups. All these groups are displayed in the following table:

	K_1	K_0	K_{-1}	NK_0	NK_1
$\mathbb{Z}[\mathbb{Z}/2]$	$(\mathbb{Z}/2)^2$	\mathbb{Z}	0	0	0
$\mathbb{Z}[\mathbb{Z}/2 imes \mathbb{Z}/2]$	$(\mathbb{Z}/2)^3$	\mathbb{Z}	\mathbb{Z}^r	$igoplus_{leph_0} \mathbb{Z}/2$	$igoplus_{leph_0} \mathbb{Z}/2$
$\mathbb{Z}[\mathbb{Z}/4]$	$\mathbb{Z}/2 imes\mathbb{Z}/4$	\mathbb{Z}	\mathbb{Z}^s	$igoplus_{leph_0} \mathbb{Z}/2$	$igoplus_{leph_0} \mathbb{Z}/2$

Let us briefly explain the values of the table. By work of Oliver [37, Theorem 14.1-2], the kernels $SK_1(R_j)$ of the determinant maps are trivial, and hence (see for example [44]) $K_1(R_j)$ is the group of units of R_j . Now the values of K_1 follow from a theorem of Higman ([24], see also [42, II.4.1]). In turn, by [9, Proposition 6], the reduced K_0 of these three rings is trivial, and then [27, Lemma 2.18] implies that $K_0(R_j) = \mathbb{Z}$ for every j. The values of the third column follow from work of Carter [8, Theorem 1], being the figures r and s positive integers that depend on the Schur indexes. Finally, the Nil-terms of the two last columns were computed by Weibel in [49].

We are now ready to describe the homology of the dihedral Artin groups with respect to the family of virtually cyclic groups, referred to the K-theory K_q of these group rings, q = 0, 1. As before, our main tool is Theorem 4.17.

Example 5.3. We start with $R_1 = \mathbb{Z}[\mathbb{Z}/2]$. When q = 0, $H_3^{vc}(A_n, K_0(R_1[-])) = H_2^{vc}(A_n, K_0(R_1[-])) = \bigoplus_{\aleph_0} \mathbb{Z}$. Taking into account that $N_0^{[H]} = 0$ for every H (because $K_{-1}(R_1)$ is trivial, and also the Nil-terms), we obtain that $H_1^{vc}(A_n, K_0(R_1[-])) = 0$. On the other hand, $H_0^{vc}(A_n, K_0(R_1[-])) = \mathbb{Z}$ too.

If q = 1, $H_3^{vc}(A_n, K_1(R_1[-])) = H_2^{vc}(A_n, K_1(R_1[-])) = \bigoplus_{\aleph_0} \mathbb{Z}/2$. Moreover, as \mathbb{Z} is torsion-free and then the correspondent $T_1(K_1(R_1))$ is trivial, we have $H_1^{vc}(A_n, K_1(R_1[-])) = \bigoplus_{\aleph_0} \mathbb{Z}$. Finally, taking into account that $C(K_1(R_1)) = \mathbb{Z}$ in this case, $H_0^{vc}(A_n, K_1(R_1[-])) = \mathbb{Z}/2 \oplus \mathbb{Z}$

Example 5.4. We continue by considering $R_2 = \mathbb{Z}[\mathbb{Z}/2 \times \mathbb{Z}/2]$. Now $H_3^{vc}(A_n, K_0(R_2[-])) = H_2^{vc}(A_n, K_0(R_2[-])) = \bigoplus_{\aleph_0} \mathbb{Z}$, exactly as in the previous example. We have $N_0^{[H]} = K_{-1}(R_2) \oplus NK_0(R_2) \oplus NK_0(R_2) = \mathbb{Z}^r \oplus (\bigoplus_{\aleph_0} \mathbb{Z}/2)$ and then $H_1^{vc}(A_n, K_0(R_2[-])) = (\bigoplus_{\aleph_0} \mathbb{Z}) \oplus (\bigoplus_{\aleph_0} \mathbb{Z}/2)$. Observe that the extra term $T_2(K_0(R_2))$ is an \mathbb{F}_2 -vector space of at most countable dimension, and then it is included in the previous direct sum. Finally, $H_0^{vc}(A_n, K_0(R_2[-])) = (\bigoplus_{\aleph_0} \mathbb{Z}) \oplus$

 $(\bigoplus_{\aleph_0} \mathbb{Z}/2)$ as in the previous case, taking into account that $\overline{C}(K_0(R_2))$ is a direct sum of free abelian groups and an \mathbb{F}_2 -vector space, both of at most countable dimension.

When q = 1, $H_3^{vc}(A_n, K_1(R_2[-])) = H_2^{vc}(A_n, K_0(R_2[-])) = \bigoplus_{\aleph_0} \mathbb{Z}/2$, because $K_1(R_2) = (\mathbb{Z}/2)^3$. Now $N_q^{[H]} = \mathbb{Z} \oplus \bigoplus_{\aleph_0} \mathbb{Z}/2$, and then by similar reasons to the previous case, $H_1^{vc}(A_n, K_1(R_2[-])) = (\bigoplus_{\aleph_0} \mathbb{Z}) \oplus (\bigoplus_{\aleph_0} \mathbb{Z}/2)$. Similarly $H_0^{vc}(A_n, K_1(R_2[-])) = (\bigoplus_{\aleph_0} \mathbb{Z}) \oplus (\bigoplus_{\aleph_0} \mathbb{Z}/2)$.

Example 5.5. To conclude, we consider $R_2 = \mathbb{Z}[\mathbb{Z}/4]$. As the groups K_0 , NK_0 and NK_1 are all isomorphic as abelian groups to their counterparts in the previous example and K_{-1} is also free abelian, $H_3^{vc}(A_n, K_0(R_3[-])) = H_2^{vc}(A_n, K_0(R_3[-])) = \bigoplus_{\aleph_0} \mathbb{Z}, H_1^{vc}(A_n, K_0(R_3[-])) = (\bigoplus_{\aleph_0} \mathbb{Z}) \oplus (\bigoplus_{\aleph_0} \mathbb{Z}/2)$, and $H_0^{vc}(A_n, K_0(R_3[-])) = (\bigoplus_{\aleph_0} \mathbb{Z}) \oplus (\bigoplus_{\aleph_0} \mathbb{Z}/2)$.

For q=1, it is clear that $H_3^{vc}(A_n,K_1(R_3[-]))=(\bigoplus_{\aleph_0}\mathbb{Z}/2)\oplus(\bigoplus_{\aleph_0}\mathbb{Z}/4)$. Now, the analysis of g_2^1 in Section 4.3.3 guarantees that $H_2^{vc}(A_n,K_1(R_3[-]))$ should contain a subgroup isomorphic to $(\bigoplus_{\aleph_0}\mathbb{Z}/2)\oplus(\bigoplus_{\aleph_0}\mathbb{Z}/4)$, and then be isomorphic to it, as the source of g_2^1 is also isomorphic to $(\bigoplus_{\aleph_0}\mathbb{Z}/2)\oplus(\bigoplus_{\aleph_0}\mathbb{Z}/4)$. We have $N_1^{[H]}=K_0(R_3)\oplus NK_0(R_2)\oplus NK_0(R_2)=\mathbb{Z}\oplus(\bigoplus_{\aleph_0}\mathbb{Z}/2)$. By an analogous reasoning to the case of R_2 , $H_1^{vc}(A_n,K_1(R_3[-]))=(\bigoplus_{\aleph_0}\mathbb{Z})\oplus(\bigoplus_{\aleph_0}\mathbb{Z}/2)$. Finally, $H_0^{vc}(A_n,K_1(R_3[-]))=(\bigoplus_{\aleph_0}\mathbb{Z})\oplus(\bigoplus_{\aleph_0}\mathbb{Z}/2)\oplus\mathbb{Z}/4$, corresponding the extra term to $K_1(R_3)$.

Remark 5.6. Observe that the proposed method permits computations of the Bredon homology (with respect to the family of virtually cyclic groups) of A_n with respect to the K-theory of any ring for which there is some knowledge of the algebraic K-groups. We remark that the computation of the lower algebraic K-theory groups is a hot topic nowadays (see for example [25], [23] or [20]), so it seems possible that good knowledge about the E^2 -term of the Atiyah-Hirzebruch spectral sequence is achieved in these cases, even for more general families of Artin groups.

According to Theorem 4.17, the Atiyah-Hirzebruch sequence has four columns. This indicates that the sequence should collapse in the page E_4 , and hence this page should provide the knowledge of Farrell-Jones groups. The analysis of the differentials, as well as the computation of the E_3 and E_4 -pages, seems a difficult and interesting future line of research.

6. Farrell-Jones homology of Artin groups of dihedral type with coefficients in regular rings

⟨Sect:Farrell⟩

In this last section we will prove that an Atiyah-Hirzebruch spectral sequence whose E_2 page is given by ordinary homology leads to the description of (some) Farrell-Jones homology
groups of A_n , assuming coefficients in \mathbb{Z} or \mathbb{F}_r with r prime.

In our approach to $H^G(\underline{\underline{E}}G, \mathbf{K}(\mathbb{Z}))$, it is crucial Theorem 0.1 in [30], that establishes a splitting

$$H^G(\underline{E}G, \mathbf{K}(R)) \simeq H^G(\underline{E}G, \mathbf{K}(R)) \oplus H^G(\underline{E}G, \underline{E}G, \mathbf{K}(R)),$$

for every group G and ring R. When the ring R is regular and G is torsion-free, the second term of the direct sum vanishes [28, Proposition 2.6]. Likewise, when G is torsion-free $\underline{E}G = EG$, the classical universal space for principal G-bundles, and then $H^G(\underline{E}G, \mathbf{K}(R)) = H(BG, \mathbf{K}(R))$,

being the latter ordinary homology. As said above, we are interested in the case $G = A_n$, which is torsion-free, and R a ring whose underlying group is cyclic. Our goal here will be to compute $H^{A_n}(\underline{E} A_n, \mathbf{K}(R))$.

6.1. Homology with coefficients in $K_q(\mathbb{Z})$. From now on, we undertake the computation of $H(A_n, \mathbf{K}(\mathbb{Z}))$. Consider first n odd. By Proposition 4.12 and Universal Coefficient Theorem, there are only two non-trivial columns in $E_{p,q}^2 = H_p(BG, K_q(\mathbb{Z}))$, corresponding to p = 0 and p = 1. The spectral sequence then collapses in the E_2 -term, and we obtain the following:

Proposition 6.1. For every $q \in \mathbb{Z}$ and every odd $n \geq 2$, there exists an extension

$$K_q(\mathbb{Z}) \to H_q(BA_n, \mathbf{K}(\mathbb{Z})) \to K_{q-1}(\mathbb{Z}).$$

Proof. It is a direct consequence of the definition of the edge homomorphisms in the spectral sequence. \Box

Now, the K-theory groups of \mathbb{Z} described in Example 5.1 and the previous proposition imply the following result.

⟨Prop:FJodd⟩

Proposition 6.2. If $n \geq 2$ is odd, we have the following:

- $H_q(BA_n, \mathbf{K}(\mathbb{Z})) = 0$ if q < 0.
- $H_0(BA_n, \mathbf{K}(\mathbb{Z})) = \mathbb{Z}$.
- $H_1(BA_n, \mathbf{K}(\mathbb{Z})) = \mathbb{Z} \oplus \mathbb{Z}/2$.
- $H_2(BA_n, \mathbf{K}(\mathbb{Z}))$ is a group of order 4.
- $H_3(BA_n, \mathbf{K}(\mathbb{Z}))$ is isomorphic to $\mathbb{Z}/\mathbf{96}$ or to $\mathbb{Z}/\mathbf{2} \times \mathbb{Z}/\mathbf{48}$.
- $H_4(BA_n, \mathbf{K}(\mathbb{Z})) = \mathbb{Z}/48$.
- $H_5(A_n, \mathbf{K}(\mathbb{Z})) = \mathbb{Z}$.
- $H_6(A_n, \mathbf{K}(\mathbb{Z})) = \mathbb{Z}$.
- $H_7(A_n, \mathbf{K}(\mathbb{Z})) = \mathbb{Z}/240$.

It is possible to approach many more groups taking as input the algebraic K-theory of \mathbb{Z} . For example, for a > 1, the groups $H_{8a}(BA_n, \mathbf{K}(R))$ are always isomorphic to a product $\mathbb{Z} \times \mathbb{Z}/\mathbf{k}$, and the number k can always be computed in terms of the Bernoulli numbers. Moreover the groups $H_{4a}(BA_n, \mathbf{K}(\mathbb{Z}))$ and $H_{4a+1}(BA_n, \mathbf{K}(\mathbb{Z}))$ depend on a solution of the Vandiver conjecture, which is still open. See Section 9 of [48] for details.

Remark 6.3. Recall that, by [7], Farrell-Jones holds for A_n , so the previous results also compute $K_i(\mathbb{Z}A_n)$.

When n is even, the computation gets harder. In this case, following similar arguments to the odd case, the non-trivial homology groups of A_n with coefficients in $K_q(\mathbb{Z})$ are $H_0(A_n, K_q(\mathbb{Z})) = H_2(A_n, K_q(\mathbb{Z})) = K_q(\mathbb{Z})$, and $H_1(A_n, K_q(\mathbb{Z})) = K_q(\mathbb{Z}) \oplus K_q(\mathbb{Z})$. Then, the Atiyah-Hirzebruch spectral sequence has three non-trivial columns, corresponding to p = 0, 1, 2 in $E_{p,q}^2 = H_p(BG, K_q(\mathbb{Z}))$. But now it is possible that non-trivial differentials exist in the page E^3 , and the sequence will collapse in E^3 , but perhaps not before. Let us collect the information we have about the E_3 -page:

Proposition 6.4. For every $q \in \mathbb{Z}$, we have $E_{0,q}^3 = \operatorname{Ker} d_3 \subseteq K_q(\mathbb{Z})$, $E_{1,q}^3 = K_q(\mathbb{Z}) \oplus K_q(\mathbb{Z})$, $E_{2,q}^3 = K_q(\mathbb{Z})/\operatorname{Im} d_3$ and, $E_{p,q}^3 = 0$ otherwise.

Proof. The result follows from dimensional reasons from the definition of the spectral sequence, and the ordinary homology of A_n with coefficients in $K_q(\mathbb{Z})$.

As said above, $E_{p,q}^3 = E_{p,q}^{\infty}$, and we may say something about the homology groups.

Proposition 6.5. There always exist surjective homomorphisms $H_q(BA_n, \mathbf{K}(\mathbb{Z})) \to K_{q-2}(\mathbb{Z})/Im\ d_3$, for every $q \in \mathbb{Z}$. Moreover, the kernel B_q of this homomorphism can be described by means of a central extension $Ker\ d_3 \to B_q \to K_{q-1}(\mathbb{Z}) \oplus K_{q-1}(\mathbb{Z})$.

Proof. It is a consequence of the previous proposition and the structure of the filtration in the spectral sequence, see for example Section 3 in [15].

6.2. Homology with coefficients in $K_q(\mathbb{F}_r)$. In the following let r be a prime number. Unlike the previous case, the algebraic K-theory groups of the field \mathbb{F}_r are computed [39]. If q is negative or q is even, $K_q(\mathbb{Z}/\mathbf{r}) = 0$. If q = 0, $K_0(\mathbb{F}_r) = \mathbb{Z}$, and if q = 2k - 1 $(k \ge 1)$, $K_q(\mathbb{F}_r) = \mathbb{Z}/(\mathbf{r}^k - 1)$.

As before, we consider first the case n odd, in which the E^2 -page of the Atiyah-Hirzebruch spectral sequence has only two non-trivial columns that correspond to p = 0, 1. Moreover, according with the previous data and Proposition 4.12, we have

$$\begin{cases} E_{0,0}^2 = E_{0,1}^2 = \mathbb{Z} \\ E_{2k-1,0}^2 = E_{2k-1,1}^2 = \mathbb{Z}/(\mathbf{r}^{\mathbf{k}} - \mathbf{1}) & \text{for } k \ge 1 \\ E_{p,q}^2 = 0 & \text{otherwise.} \end{cases}$$

For dimensional reasons there are no differentials in the spectral sequence, and hence $E_{p,q}^2 = E_{p,q}^{\infty}$. Moreover, in the diagonal p+q=s there is only one possible nonzero entry, so we have: $\langle \text{Fqodd} \rangle$ **Proposition 6.6.** Given a prime r and a natural number $s \geq 0$, we have

$$H_s(BA_n, \mathbf{K}(\mathbb{F}_r)) = \begin{cases} \mathbb{Z} & \text{if } s = 0\\ \mathbb{Z}/(\mathbf{r}^{\mathbf{k}} - \mathbf{1}) & \text{if } s = 2k - 1, \ k \ge 0\\ \mathbb{Z}/(\mathbf{r}^{\mathbf{k}} - \mathbf{1}) & \text{if } s = 2k, \ k \ge 1. \end{cases}$$

Proof. It follows straightforward from the previous remarks and the fact that this Atiyah-Hirzebruch spectral sequence converges to Farrell-Jones K-homology, i.e $H_p(A_n, K_q(\mathbb{F}_r)) \Rightarrow H_{p+q}(BA_n, \mathbf{K}(\mathbb{F}_r))$.

Assume now n even. Now the E^2 -page of the spectral sequence has three non-trivial columns, corresponding to q = 0, 1, 2. According again to Proposition 4.12 and the aforementioned values of $K_q(\mathbb{F}_r)$, we have:

$$\begin{cases} E_{0,0}^2 = E_{0,2}^2 = \mathbb{Z} \\ E_{0,1}^2 = \mathbb{Z} \\ E_{2k-1,0}^2 = E_{2k-1,2}^2 = \mathbb{Z}/(\mathbf{r^k-1}) & \text{for } k \geq 1 \\ E_{2k-1,1}^2 = (\mathbb{Z}/(\mathbf{r^k-1}))^2 & \text{for } k \geq 1 \\ E_{p,q}^2 = 0 & \text{otherwise.} \end{cases}$$

Again by dimensional reasons there are no differentials in the spectral sequence, which collapses in the E^2 -term. Then the Farrell-Jones K-homology groups of A_n fit in the following extensions:

 $\langle Feven \rangle$

Proposition 6.7. Given a prime r and a natural number $s \geq 0$, we have

$$= \begin{cases} H_s(BA_n, \mathbf{K}(\mathbb{F}_r)) = \mathbb{Z} & \text{if } s = 0 \\ H_s(BA_n, \mathbf{K}(\mathbb{F}_r)) = \mathbb{Z}/(\mathbf{r}^s - \mathbf{1}) \times \mathbb{Z} & \text{if } s = 1 \\ \mathbb{Z}/(r^s - 1) \to H_s(BA_n, \mathbf{K}(\mathbb{F}_r)) \to (\mathbb{Z}/(\mathbf{r}^{s-1} - \mathbf{1}))^2 & \text{if } s = 2k - 1, \ k > 1 \\ (\mathbb{Z}/(\mathbf{r}^{s-1} - \mathbf{1}))^2 \to H_s(BA_n, \mathbf{K}(\mathbb{F}_r)) \to \mathbb{Z}/(\mathbf{r}^{s-2} - \mathbf{1}) & \text{if } s = 2k, \ k \ge 1. \end{cases}$$

Proof. Observe that in every line p+q=s there are exactly two entries of $E^2=E^{\infty}$ which are non-trivial, and are given by the correspondent groups of K-theory of \mathbb{F}_q . The extensions are of course given by the edge homomorphisms. The extension that identifies $H_1(BA_n, \mathbf{K}(\mathbb{F}_r))$ splits because there are no non-splitting (abelian) extensions of \mathbb{Z}/\mathbf{m} by \mathbb{Z} , for any natural $m \geq 2$. We finish by observing again that the spectral sequence converges to Farrell-Jones K-homology.

⟨cor:nonsplit⟩

Corollary 6.8. In the notation of the previous proposition, if r = 2, all the previous extensions split.

Proof. It is enough to remark that $2^r - 1$ and $2^{r+1} - 1$ are coprimes for r > 1, and then there are no non-trivial extensions of $\mathbb{Z}/(2^{r+1} - 1)$ by $\mathbb{Z}/(2^r - 1)$.

Remark 6.9. It is important to highlight again that Proposition 6.6 and Proposition 6.7 do compute the algebraic K-theory of $\mathbb{F}_r A_n$, because Farrell-Jones conjecture holds for A_n .

Questions. It is an interesting and difficult problem to investigate how to generalize our techniques to more general families of Artin groups. It would be also good to know which statements are also valid for the bigger family of torsion-free one-relator groups.

Acknowledgments.

We warmly thank D. Juan-Pineda, W. Lück, N. Petrosyan, V. Srinivasan and J. Stienstra for their useful comments.

References

- AJT18 [1] J. Aramayona, D. Juan-Pineda and A. Trujillo-Negrete, On the virtually cyclic dimension of mapping class groups of punctured spheres, Algebr. Geom. Topol. 18 (2018), no. 4, 2471—2495.
- Arma [2] J. Aramayona and C. Martínez-Pérez, The proper geometric dimension of the mapping class group, Algebr. Geom. Topol. 14 (2014), no. 1, 217—227.
- [3] G.N. Arzhantseva, On quasiconvex subgroups of word hyperbolic groups, Geom. Dedicata, 87 (2001), 191–208.
- ABGRW [4] S Azzali, S.L. Browne, M.P. Gomez-Aparicio, L.C. Ruth and H. Wang K-homology and K-theory of pure braid groups, arXiv:2105.07848
- BJV14 [5] N. Bárcenas, D. Juan-Pineda and M. Velásquez, Bredon cohomology, K-theory and K-homology of pullbacks of groups, arXiv:1311.3138.
- Bas68 [6] H. Bass, Algebraic K-theory, W. A. Benjamin, Inc., New York-Amsterdam, 1968.
- BFW21 [7] M. Bestvina. K. Fujiwara and D. Wigglesworth, Farrell-Jones conjecture for free-by-cyclic groups, arXiv:1906.00069.pdf

- Car80 [8] D.W. Carter, Lower K-theory of finite groups, Comm. Algebra 8 (1980), 1927-1937.
- Cas73 [9] P. Cassou-Noguès, Classes d'idéaux de l'algèbre d'un groupe abélien, C. R. Acad. Sci. Paris Ser. A-B, 276:A973-A975, 1973.
- [CMW04] [10] R. Charney, J. Meier and K. Whitlessey, Bestvina's normal form complex and the homology of Garside groups, Geom. Dedicata 105 (2004), 171-–188.
- [CHR20] [11] L. Ciobanu, D. Holt and S. Rees, Equations in groups that are virtually direct products J. Algebra 545 (2020), 88–99.
- [CEO9] [12] M. Clancy and G. Ellis. Homology of some Artin and twisted Artin groups, J. K-Theory 6 (2009), 171–196.
- Deg16 [13] D. Degrijse. A cohomological characterization of locally virtually cyclic groups, Adv. Math. 305 (2017), 935–952.
 - [DP] [14] D. Degrijse and N. Petrosyan, Geometric dimension of groups for the family of virtually cyclic subgroups, J. Topol. 7 3 (2014), 697–726.
 - [DR][15] A Díaz-Ramos Spectral sequences via examples, The Graduate Journal of Mathematics, Mediterranean Institute for the Mathematical Sciences (MIMS), Volume 2, Issue 1 (2017), 10–28.
- TDieck [16] T. tom Dieck, Transformation groups Vol. 8. Walter de Gruyter, 1987.
- [Far77][17] T. Farrell, The nonfiniteness of Nil, Proc. Amer. Math. Soc. 65 (1977), 215—216.
- FaJo93 [18] T. Farrell and L. Jones, Isomorphism conjectures in algebraic K-theory, J. Amer. Math. Soc., 6 (1993), 249—297.
- [F1Go18] [19] R. Flores and J. González-Meneses, Classifying spaces for the family of virtually cyclic subgroups of braid groups, Int. Math. Res. Notices. 2020 (2020), 1575–1600.
- GJM18 [20] J. Guaschi, D. Juan-Pineda and S. Millán-López, The lower algebraic K-theory of virtually cyclic subgroups of the braid groups of the sphere and of $\mathbb{Z}[B_4(S^2)]$, Springer Briefs in Mathematics. Springer, Cham, 2018.
- [HaLu12] [21] I. Hambleton and W. Lück, Induction and computation of Bass Nil groups for finite groups, Pure Appl. Math. Q. 8 (2012), 199–219.
- [122] A. Hatcher, Algebraic topology, Cambridge University Press, 2002.
- [HiJu2021] [23] C. Hidber and D. Juan-Pineda, The algebraic K-theory of the group ring of the Klein bottle group, Topol. Appl. 293 (2021), 8 pages.
 - [Hig40][24] G. Higman, The units of group rings, Proc. London Math. Soc. 46 (1940), 231–248.
 - LaOro7 [25] J. Lafont and I. Ortiz, Relative hyperbolicity, classifying spaces, and lower algebraic K-theory, Topology 46 (2007), 527–553.
 - Luc05 [26] W. Lück, Survey on classifying spaces for families of subgroups, in Infinite groups: geometric, combinatorial and dynamical aspects, 269—322, Progr. Math., 248, Birkhäuser, Basel, 2005.
 - Luc21 [27] W. Lück, Conjectures in K and L-theory, in preparation, https://www.him.uni-bonn.de/luck/data/ic.pdf
 - [28] W. Lück and H. Reich, The Baum-Connes and the Farrell-Jones conjectures in K- and L-theory, Handbook of K-theory, volume 2, pp. 703–842, Springer, 2005.
 - Luro14 [29] W. Lück and D. Rosenthal, On the K- and L-theory of hyperbolic and virtually finitely generated abelian groups, Forum Math. 26 (2014), no. 5, 1565—1609.
 - [LS16] [30] W. Lück and W. Steimle, Splitting the relative assembly map, Nil-terms and involutions, Ann. K-theory 1 (2016) 339–377.
 - [IW] [31] W. Lück and M. Weiermann, On the classifying space of the family of virtually cyclic subgroups, Pure Appl. Math. Q. 8 2 (2012), 497–555.
 - Lyndon [32] R. Lyndon, Cohomology theory of groups with a single defining relation, Ann. of Math. 52 (1950), 650–665.
 - Mac96 [33] D. Macpherson, Permutation groups whose subgroups have just finitely many orbits, in Ordered groups and infinite permutation groups, pp. 221–230, Kluwer Academic Publishers, 1996.
 - MaMa06 [34] J. Mairesse and F. Mathéus, Growth series for Artin groups of dihedral type Internat. J. Algebra Comput. 16 (2006), 1087–1107.
 - Mar02 [35] C. Martínez-Pérez, A spectral sequence in Bredon (co)homology, J. Pure Appl. Algebra 176 (2002), 161–173.
 - MiVaO3 [36] G. Mislin and A. Valette, Proper group actions and the Baum-Connes conjecture, Advanced courses in Mathematics, no. 55, CRM Barcelona, 2003.
 - [01188][37] R. Oliver, Whitehead groups of finite groups. Cambridge University Press, Cambridge, 1988.
 - Par09 [38] L. Paris, Braid groups and Artin groups, Handbook of Teichmüller theory. Vol. II, pp. 389—451, IRMA Lect. Math. Theor. Phys., 13, Eur. Math. Soc., Zürich, 2009.

[Qui73] [39] D. Quillen, Finite generation of the groups Ki of rings of algebraic integers, in Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972), pages 179–198. Lecture Notes in Math., vol. 341. Springer-Verlag, Berlin, 1973.

San08 [40] R. Sánchez-García, Bredon homology and equivariant K-homology of SL(3, Z), J. Pure Appl. Algebra, 212 (2008), 1046–1059.

[SaVe18] [41] L. Sánchez-Saldaña and M. Velásquez, The algebraic and topological K-theory of the Hilbert modular group, Homology Homotopy Appl. 20 (2018), no. 2, 377—402.

She78 [42] S.K. Shehgal Topics in group rings, Dekkar, New York, 1978.

Sri91 [43] V. Srinivas, Algebraic K-theory, Birkhäuser Boston Inc., Boston, MA, 1991.

Ste78 [44] M.R. Stein, Whitehead groups of finite groups, Bull. Amer. Math. Soc. 84 (1978), 201–212.

[Sti82] [45] J. Stienstra, Operations in the higher K-theory of endomorphisms, Current trends in algebraic topology, Part 1 (London, Ont., 1981), CMS Conf. Proc., 2, Amer. Math. Soc., Providence, R.I., 1982, 59–115.

Wal09 [46] C. Walsh, Busemann points of Artin groups of dihedral type, Internat. J. Algebra Comput. 19 (2009), 891–910.

Wei81 [47] C. Weibel, Mayer-Vietoris sequences and module structures on NK, Algebraic K-theory, Evanston 1980 (Proc. Conf., Northwestern Univ., Evanston, Ill., 1980), Lecture Notes in Math., 854, Springer, Berlin, 1981, 466—493.

[48] C. Weibel, Algebraic K-theory of rings of integers in local and global fields, Handbook of K-theory, volume 2, pp. 339–390, Springer, 2005.

[Wei09] [49] C. Weibel, NK_0 and NK_1 of the groups C_4 and D_4 , Comment. Math. Helv. 84 (2009), 339–349.

Wei13 [50] C. Weibel, The K-book: an introduction to algebraic K-theory, Graduate Studies in Mathematics, 145 (2013), American Mathematical Society.

Universidad Autónoma de Madrid, Departamento de Matemáticas, Campus de Cantoblanco, 28049, Madrid (Spain)

Email address, Yago Antolín: yago.anpi@gmail.com

Universidad de Sevilla, Departmento de Geometría y Topología, c/ Tarfia, s/n 41080, Seville (Spain)

Email address, Ramón Flores: ramonjflores@us.es