LOCAL SOLUBILITY FOR A FAMILY OF QUADRICS OVER A SPLIT QUADRIC SURFACE

TIM BROWNING, JULIAN LYCZAK, AND ROMAN SARAPIN

ABSTRACT. We study the density of everywhere locally soluble diagonal quadric surfaces, parameterised by rational points that lie on a split quadric surface.

1. Introduction

A great deal of recent activity has been directed at the question of local solubility for families of varieties. Let X and Y be smooth, proper and geometrically irreducible varieties over \mathbb{Q} , equipped with a dominant morphism $\pi\colon X\to Y$, with geometrically integral generic fibre. Suppose that Y is Fano and equipped with an anticanonical height function $H\colon Y(\mathbb{Q})\to \mathbb{R}$. This short note is concerned with the behaviour of the counting function

$$N_{\text{loc}}(\pi, B) = \# \left\{ y \in Y(\mathbb{Q}) \cap \pi(X(\mathbf{A}_{\mathbb{Q}})) \colon H(y) \le B \right\},\,$$

as $B \to \infty$, where $\mathbf{A}_{\mathbb{Q}}$ denotes the ring of adèles. We shall consider an explicit example, with $Y \subset \mathbb{P}^3$ a split quadric surface, which calls into question our expectations about the behaviour of $N_{\text{loc}}(\pi, B)$.

The quantity $N_{\text{loc}}(\pi, B)$ measures the density of fibres of π which are everywhere locally soluble, and it has received the most attention when Y is projective space. For example, it follows from work of Poonen and Voloch [10, Thm. 3.6] that

$$N_{\text{loc}}(\pi, B) \sim c_{d,n} B, \tag{1.1}$$

as $B \to \infty$, for an explicit constant $c_{d,n} > 0$, when $\pi: X \to \mathbb{P}^N$ is the family of degree d hypersurfaces in \mathbb{P}^n over \mathbb{Q} , with $(d,n) \neq (2,2)$ and $N = \binom{n+d}{d} - 1$. Moreover, when $Y = \mathbb{P}^n$, work of Loughran and Smeets [7] shows that

$$N_{\text{loc}}(\pi, B) \ll \frac{B}{(\log B)^{\Delta(\pi)}},$$
 (1.2)

for a certain quantity $\Delta(\pi) \geq 0$, whose definition will be recalled in §2 and which can be extended to arbitrary fibrations $\pi \colon X \to Y$. For $Y = \mathbb{P}^n$, it is conjectured in [7] that the upper bound (1.2) should be sharp whenever the fibre $\pi^{-1}(D)$ above each codimension one point D in \mathbb{P}^n contains a component of multiplicity one.

It is natural to ask what happens when Y is a Fano variety over \mathbb{Q} that is not projective space. We shall assume that $Y(\mathbb{Q})$ is Zariski dense in Y and that

Date: March 15, 2022.

2010 Mathematics Subject Classification. 14G05 (11N36, 14D10).

Y conforms to the version of the Manin conjecture [2] without accumulating subvarieties, so that

$$\#\{y \in Y(\mathbb{Q}) : H(y) \le B\} \sim c_Y B(\log B)^{\rho_Y - 1}$$

as $B \to \infty$, where $c_Y > 0$ is a constant and ρ_Y is the rank of the Picard group of Y. Assuming that Y is a Fano variety over \mathbb{Q} without accumulating subvarieties, a naive combination of the Manin conjecture with the upper bound (1.2) might lead us to suppose that

$$N_{\text{loc}}(\pi, B) \sim \frac{c_{\pi} B(\log B)^{\rho_Y - 1}}{(\log B)^{\Delta(\pi)}},\tag{1.3}$$

as $B \to \infty$, for a suitable constant $c_{\pi} > 0$. This is compatible with (1.1), since in this case $\Delta(\pi) = 0$ and $\rho_{\mathbb{P}^N} = 1$. It also compatible with an upper bound of Browning and Loughran [1, Thm. 1.10], which concerns the case where $Y \subset \mathbb{P}^n$ is a smooth quadric over \mathbb{Q} of dimension at least 3, since then $\rho_Y = 1$.

We study an explicit example that addresses a case where the variety Y has Picard number $\rho_Y > 1$ and which contradicts the expectation (1.3). Let $Y \subset \mathbb{P}^3$ be the split quadric surface

$$y_0 y_1 = y_2 y_3. (1.4)$$

Clearly Y is smooth and $\rho_Y = 2$. Let $X \subset \mathbb{P}^3 \times \mathbb{P}^3$ be the variety defined by the pair of equations

$$y_0 x_0^2 + y_1 x_1^2 + y_2 x_2^2 + y_3 x_3^2 = 0$$

$$y_0 y_1 = y_2 y_3.$$
 (1.5)

Then X is equipped with a dominant morphism $\pi\colon X\to Y$ whose generic fibre is geometrically irreducible. We shall see in §2 that the total space X is singular and so we let $\widetilde{X}\to X$ be a desingularisation. Taking $\widetilde{\pi}$ to be the composition with $\pi\colon X\to Y$, we shall prove the following bounds for the associated counting function $N_{\text{loc}}(\widetilde{\pi},B)$ in §3.

Theorem 1.1. We have $B \ll N_{loc}(\widetilde{\pi}, B) \ll B$.

Since the Hasse principle holds for quadratic forms, we deduce from Theorem 1.1 that the same upper and lower bounds hold for the quantity $N_{\text{glob}}(\tilde{\pi}, B)$, counting the number of $y \in Y(\mathbb{Q}) \cap \pi(\tilde{X}(\mathbb{Q}))$ such that $H(y) \leq B$.

The next result, proved in §2, shows that our example is not compatible with the naive expectation (1.3), since $\rho_Y = 2$.

Theorem 1.2. We have $\Delta(\widetilde{\pi}) = 2$.

Just as thin sets have been found to have a large influence on modern formulations of the Manin conjecture, as discussed by Peyre [8], so we expect that the

naive expectation (1.3) should hold after the removal of appropriate thin sets from $Y(\mathbb{Q})$. In fact, for our example (1.5), we conjecture that

$$\#\left\{y \in Y(\mathbb{Q}) \cap \pi(\widetilde{X}(\mathbf{A}_{\mathbb{Q}})) \colon H(y) \le B, \ -y_0 y_2 \ne \square, \ -y_0 y_3 \ne \square\right\} \sim c \frac{B}{\log B},$$

for an appropriate constant c > 0. It seems plausible that the character sum method developed by Friedlander and Iwaniec [4] will prove useful for tackling this counting problem, although the hyperboloid height conditions that arise from parameterising the points in $Y(\mathbb{Q})$ will undoubtedly complicate the argument.

In this note we have focused attention on the quadric surface bundle $X \to Y$, where X is given by (1.5). One can also study the conic bundle $\eta: W \to Y$, where $W \subset \mathbb{P}^2 \times \mathbb{P}^3$ is given by

$$y_0 x_0^2 + y_1 x_1^2 + y_2 x_2^2 = 0$$
$$y_0 y_1 = y_2 y_3.$$

This variety is singular and so we consider the morphism $\widetilde{\eta} \colon \widetilde{W} \to Y$, where $\widetilde{\eta}$ is the composition of a desingularisation $\widetilde{W} \to W$ with η . The methods of this paper carry over to this case with little adjustment and can be used to prove that $B \ll N_{\text{loc}}(\widetilde{\eta}, B) \ll B$. Moreover, one also has $\Delta(\widetilde{\eta}) = 2$ in this case, providing a further example that counters the expectation (1.3), but one which we believe can be repaired via the removal of appropriate thin sets. (In fact, for $i \in \{0, 1\}$, the fibre of η above the line $y_i = y_2 = 0$ has multiplicity two in W, whereas the corresponding fibre of $\widetilde{\eta}$ admits an additional component of multiplicity one.)

Acknowledgement. While working on this paper the first author was supported by FWF grant P 32428-N35. This paper was written while the third author was an intern at IST.

2. Geometric input

We begin by extending the definition of $\Delta(\pi)$ from [7] to general fibrations. Let X and Y be arbitrary proper and geometrically irreducible varieties over \mathbb{Q} , equipped with a dominant morphism $\pi \colon X \to Y$, with geometrically integral generic fibre. Assume that Y is smooth and associate the residue field $\kappa(D)$ to any codimension one point $D \in Y^{(1)}$. Let S_D be the set of geometrically irreducible components of $\pi^{-1}(D)$ and choose a finite group Γ_D through which the action of $\operatorname{Gal}(\overline{\kappa(D)}/\kappa(D))$ on S_D factors. Then $\Delta(\pi)$ is defined to be

$$\Delta(\pi) = \sum_{D \in Y^{(1)}} (1 - \delta_D(\pi)), \qquad (2.1)$$

where

$$\delta_D(\pi) = \frac{\#\{\sigma \in \Gamma_D \colon \sigma \text{ acts with a fixed point on } S_D\}}{\#\Gamma_D}.$$

Henceforth, we take $Y \subset \mathbb{P}^3$ to be the split quadric (1.4) and $X \subset \mathbb{P}^3 \times \mathbb{P}^3$ to be the variety (1.5). Consider the fibration $\pi \colon X \to Y$. There are precisely 4 codimension one points in Y which produce reducible fibres. These are the four lines

$$D_{i,j} = \{y_i = y_j = 0\}, \text{ for } i \in \{0,1\}, j \in \{2,3\}.$$
 (2.2)

Above each of these the fibre of π is split by a quadratic extension, so that $\delta_{D_{i,j}}(\pi) = \frac{1}{2}$. It follows that $\Delta(\pi) = 4 \cdot \frac{1}{2} = 2$ in (2.1).

However, the Jacobian of X is

$$J = \begin{pmatrix} x_0^2 & x_1^2 & x_2^2 & x_3^2 & 2y_0x_0 & 2y_1x_1 & 2y_2x_2 & 2y_3x_3 \\ y_1 & y_0 & -y_3 & -y_2 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

Thus X is singular, with singular locus supported on the union of subschemes

$$\Sigma_{i,j} = V(y_i, y_j, x_{i'}, x_{j'}, y_{j'}x_i^2 + y_{i'}x_j^2), \quad \text{for } \{i, i'\} = \{0, 1\} \text{ and } \{j, j'\} = \{2, 3\},$$

where we note that a choice of i and j uniquely determines i' and j'. In fact, away from the intersection of two such subschemes, the singular locus Σ is precisely the union of these four reduced subschemes. Let $\widetilde{X} \to X$ be a desingularisation of X and let $\widetilde{\pi}$ be the composition with $\pi \colon X \to Y$. The main aim of this section is to prove that

$$\Delta(\widetilde{\pi}) = 2, \tag{2.3}$$

as claimed in Theorem 1.2.

We begin by establishing the following geometric lemma that will be useful in our treatment.

Lemma 2.1. Let C and C' be two integral curves smooth and projective over a field k. Let us write κ and κ' for their function fields. Consider a proper morphism

$$\rho: W \to C \times C'$$
.

If $W \times C'_{\kappa}$ is smooth over κ and $W \times C_{\kappa'}$ is smooth over κ' , then the image of the singular locus of W/k under ρ has dimension 0.

Proof. Consider the composite morphism

$$W \to C \times_k C' \to C$$

which has a smooth generic fibre by assumption. By spreading out [9, Thm. 3.2.1], we see that $W_U \to U$ is smooth for a dense open subset $U \subset C$. We conclude that $W_U \to U \to \operatorname{Spec} k$ is smooth. Similarly, we have a $V \subset C'$ such that $W_V \to V$ is smooth. We conclude that the image of the smooth locus of $W \to \operatorname{Spec} k$ under ρ contains both $U \times C'$ and $C \times V$.

It suffices to prove (2.3) for a single desingularisation.

Definition 2.2. Let $X' \to X$ be the blowup in the union of the closed subschemes $\Sigma_{i,j}$. Let $\widetilde{X} \to X'$ be a fixed strong desingularisation.

We have seen that the image of the singular locus Σ under $X \to Y$ is onedimensional and supported on the union of lines $D_{i,j} \subset Y$ in (2.2). Let $\eta_{i,j}$ be the generic point of the line $D_{i,j}$. We will prove that the image of the singular locus of X' under $\pi' \colon X' \to Y$ has lower dimension.

Lemma 2.3. The image of the singular locus Σ' of X' under $X' \to Y$ is (at most) zero-dimensional. The fibre $X'_{\eta_{i,j}}$ has four geometrically irreducible components, of which two make up the strict transform of $X_{\eta_{i,j}}$ and two come from the exceptional divisor of the blowup $X' \to X$. These components are conjugated in pairs.

Proof. We identify Y with $\mathbb{P}^1 \times \mathbb{P}^1$, where the projective lines have coordinates $u = \frac{y_i}{y_j} = \frac{y_{j'}}{y_{i'}}$ and $t = \frac{y_i}{y_{j'}} = \frac{y_j}{y_{i'}}$, respectively. By Lemma 2.1 and symmetry we only need to show that $X' \times_Y \mathbb{P}^1_{\mathbb{Q}(u)}$ is smooth over $\mathbb{P}^1_{\mathbb{Q}(u)}$. Again, by symmetry, we can restrict to the affine opens where $t \neq \infty$.

Since blowing up commutes with pulling back along flat morphisms we find that $X' \times_Y \mathbb{A}^1_{\mathbb{Q}(u)}$ and the blowup of $X \times_Y \mathbb{A}^1_{\mathbb{Q}(u)}$ in the pullback of Σ are naturally isomorphic over $\mathbb{A}^1_{\mathbb{Q}(u)}$. The three subschemes $\Sigma_{i',j}$, $\Sigma_{i,j'}$ and $\Sigma_{i',j'}$ pull back to the empty scheme under $X \times_Y \mathbb{A}^1_{\mathbb{Q}(u)} \to X$ and it will therefore suffice to compute the blowup of $X \times_Y \mathbb{A}^1_{\mathbb{Q}(u)}$ in the pullback of $\Sigma_{i,j}$.

We see that $X \times_Y \mathbb{A}^1_{\mathbb{Q}(u)} \subset \mathbb{P}^3 \times_{\mathbb{Q}} \mathbb{A}^1_{\mathbb{Q}(u)}$ is given by $utx_i^2 + x_{i'}^2 + tx_j^2 + ux_{j'}^2 = 0$. The pullback of $\Sigma_{i,j}$ equals $V(t, x_{i'}, x_{j'}, ux_i^2 + x_j^2)$, which is contained in the affine open given by $x_i \neq 0$. By abuse of notation we will use the same notation for the variables on this affine open.

We can compute the blowup Z of $\{ut + x_{i'}^2 + tx_j^2 + ux_{j'}^2 = 0\} \subset \mathbb{A}^3 \times_{\mathbb{Q}} \mathbb{A}^1_{\mathbb{Q}(u)}$ in the subscheme $V(t, x_{i'}, x_{j'}, u + x_j^2)$ explicitly. We find

$$Z = \operatorname{Proj} \mathbb{Q}(u)[t][x_{i'}, x_j, x_{j'}][A, B, C, D]/I$$

where the grading comes from A, B, C and D, and I is the ideal generated by the 2×2 -minors of

$$\begin{pmatrix} A & B & C & D \\ t & x_{i'} & x_{j'} & u + x_j^2 \end{pmatrix},$$

together with the polynomials

$$ut + x_{i'}^2 + tx_j^2 + ux_{j'}^2$$
, $(x_{i'}^2 + ux_{j'}^2)B + tx_{i'}D$, $ux_{j'}C + x_{i'}B + tD$, $AD + B^2 + uC^2$.

One can now directly check that $Z \to \operatorname{Spec} \mathbb{Q}(u)$ is indeed smooth. Thus the first statement follows from Lemma 2.1.

For the second part we note that the fibre of $Z \to \mathbb{A}^1_{\mathbb{Q}(u)}$ over t = 0 is an affine open of $X'_{\eta_{i,j}}$. It consists of the strict transform of $X_{\eta_{i,j}}$, which we already know to consist of two conjugate components. Any other component comes from the exceptional divisor, which we can compute from the equations above to be

$$\text{Proj } \mathbb{Q}(u)[x_j][A, B, C, D]/(u + x_i^2, AD + B^2 + uC^2).$$

Hence, $X'_{\eta_{i,j}}$ has four geometrical components which are conjugated in pairs. \square

It follows from this result $\delta_{D_{i,j}}(\pi') = \frac{1}{2}$. Moreover, the singular locus of X' does not meet the fibres $X'_{\eta_{i,j}}$ over the generic points $\eta_{i,j}$ of the $D_{i,j}$. By using a strong desingularisation, which was proven to exist by Hironaka [5], we can conclude that $\widetilde{X}_{\eta_{i,j}} \stackrel{\cong}{\to} X'_{\eta_{i,j}}$ over Y. It therefore follows that $\delta_{D_{i,j}}(\widetilde{\pi}) = \frac{1}{2}$, which thereby completes the proof of (2.3), via (2.1).

3. Analytic input

In this section we prove Theorem 1.1. We note that the map $\widetilde{X} \to X$ is an isomorphism outside the locus of $y \in \mathbb{P}^3$ such that $y_0 \dots y_3 = 0$. Let $|\mathbf{y}| = \max_{0 \le i \le 3} |y_i|$ if $y \in \mathbb{P}^3(\mathbb{Q})$ is represented by a vector $\mathbf{y} \in \mathbb{Z}^4_{\text{prim}}$. The anticanonical height function on $Y(\mathbb{Q})$ is then $H(y) = |\mathbf{y}|^2$. Since there are $O(\sqrt{B})$ choices of $y \in \mathbb{P}^3(\mathbb{Q})$ of height at most B for which $y_0 \dots y_3 = 0$, it follows that

$$N_{\text{loc}}(\widetilde{\pi}, B) = N_{\text{loc}}(\pi, B) + O(\sqrt{B}),$$

where $\pi\colon X\to Y$. For a given vector \mathbf{y} , let $X_{\mathbf{y}}\subset\mathbb{P}^3$ be the diagonal quadric

$$y_0 x_0^2 + y_1 x_1^2 + y_2 x_2^2 + y_3 x_3^2 = 0.$$

Then

$$N_{\text{loc}}(\pi, B) = \frac{1}{2} \# \left\{ \mathbf{y} \in \mathbb{Z}_{\text{prim}}^4 \colon |\mathbf{y}| \le \sqrt{B}, \ X_{\mathbf{y}}(\mathbf{A}_{\mathbb{Q}}) \ne \emptyset \right\}.$$

In order to prove Theorem 1.1, it will suffice to prove that

$$B \ll N_{\text{loc}}(\pi, B) \ll B. \tag{3.1}$$

To see the lower bound, we take $\mathbf{y} = (u, v, -u, -v)$, for coprime $u, v \in \mathbb{Z}$ such that $|u|, |v| \leq \sqrt{B}$. For such \mathbf{y} we obviously have $(1:1:1:1) \in X_{\mathbf{y}}(\mathbf{A}_{\mathbb{Q}})$. It easily follows that $N_{\text{loc}}(\pi, B) \gg B$.

Remark 3.1. The lower bound for $N_{\text{loc}}(\pi, B)$ arises from points on the antidiagonal line $D \subset Y$ defined by the equations $y_0 + y_2 = y_1 + y_3 = 0$. Consider the fibration $\pi_D \colon X_D \to D$. The reducible fibres occur precisely above the closed points $y_0 = 0$ and $y_1 = 0$ in $D \cong \mathbb{P}^1$. However, the fibres above these points are split and so it follows from (2.1) that $\Delta(\pi_D) = 0$. On the other hand $\rho_D = 1$ and so our lower bound is consistent with (1.3), which would predict that $N_{\text{loc}}(\pi_D, B) \sim cB$ for an appropriate constant c > 0.

It remains to prove the upper bound in (3.1). Let us first dispatch the contribution to $N_{\text{loc}}(\pi, B)$ from \mathbf{y} for which $y_0y_1y_2y_3 = 0$. But then at least two components of \mathbf{y} must vanish and the fibre over any such point has an obvious rational point. There are $\frac{4}{\zeta(2)}B + O(\sqrt{B})$ vectors $(a, b) \in \mathbb{Z}_{\text{prim}}^2$ with $|a|, |b| \leq B$.

We denote by $N^{\circ}(B)$ the contribution from \mathbf{y} for which $y_0y_1y_2y_3 \neq 0$. Then our work so far shows that

$$N_{\text{loc}}(\pi, B) = N^{\circ}(B) + \frac{8}{\zeta(2)}B + O(\sqrt{B}).$$
 (3.2)

For each $\mathbf{y} \in \mathbb{Z}_{\neq 0}^4$ with $y_0 y_1 = y_2 y_3$, there exist $\mathbf{t} = (t_0, t_1, t_2, t_3) \in \mathbb{Z}_{\neq 0}^4$ such that $y_0 = t_0 t_2$, $y_1 = t_1 t_3$, $y_2 = t_0 t_3$, $y_3 = t_1 t_2$.

In this way it follows that

$$N^{\circ}(B) \ll \# \left\{ \mathbf{t} \in \mathbb{Z}_{\neq 0}^{4} \colon \begin{array}{l} |t_{0}t_{2}|, |t_{0}t_{3}|, |t_{1}t_{2}|, |t_{1}t_{3}| \leq \sqrt{B} \\ X_{t_{0}t_{2}, t_{1}t_{3}, t_{0}t_{3}, t_{1}t_{2}}(\mathbf{A}_{\mathbb{Q}}) \neq \emptyset \end{array} \right\}. \tag{3.3}$$

We shall begin by studying a related counting function in which only square-free integers appear, before later deducing our final estimate for $N^{\circ}(B)$.

3.1. Square-free coefficients. For given $U_0, \ldots, U_3 > 0$, let

$$\mathscr{U} = \left\{ \mathbf{u} \in \mathbb{Z}^4 : U_i/2 < |u_i| \le U_i \text{ for } 0 \le i \le 3 \right\}.$$

In this section we shall be concerned with estimating

$$N^*(U_0, \dots, U_3) = \# \left\{ \mathbf{u} \in \mathscr{U} : \begin{array}{l} X_{u_0 u_2, u_1 u_3, u_0 u_3, u_1 u_2}(\mathbf{A}_{\mathbb{Q}}) \neq \emptyset \\ \mu^2(u_0) \cdots \mu^2(u_3) = 1 \end{array} \right\}, \tag{3.4}$$

where μ is the Möbius function. We shall prove the upper bound

$$N^*(U_0, \dots, U_3) \ll \frac{U_0 U_1 U_2 U_3}{(1 + \log U_{\min})^2},$$
 (3.5)

if $U_{\min} = \min\{U_0, \dots, U_3\} \ge 1$.

The upper bound (3.5) is trivial if $1 \leq U_{\min} < 2$ and so we may proceed under the assumption that $U_{\min} \geq 2$. Our primary tool will be the following multidimensional form of the large sieve inequality, as worked out by Kowalski [6, Thm. 4.1].

Lemma 3.1. Let

$$X = \{(n_1, \dots, n_r) \in \mathbb{Z}^r \colon -N_j \le n_j \le N_j \text{ for } 1 \le j \le r\}$$

and let $\Omega_p \subset \mathbb{F}_p^r$ for all p. Then

$$\#\{x \in X \colon x \mod p \notin \Omega_p \ \forall p \le L\} \ll \frac{\prod_{i=1}^r (N_i + L^2)}{F(L)},$$

where

$$F(L) = \sum_{n < L} \mu^2(n) \prod_{p \mid n} \frac{|\Omega_p|}{p^r - |\Omega_p|}.$$

Define
$$\Omega_p = \bigcup_{i=0}^{3} \Omega_{p,i}$$
, where
$$\Omega_{p,0} = \{ (r_0, r_1, r_2, r_3) \in \mathbb{F}_p^4 \colon r_0 = 0, \ r_1 r_2 r_3 \neq 0, \ -r_2 r_3 \notin \mathbb{F}_p^{\times,2} \},$$

$$\Omega_{p,1} = \{ (r_0, r_1, r_2, r_3) \in \mathbb{F}_p^4 \colon r_1 = 0, \ r_0 r_2 r_3 \neq 0, \ -r_2 r_3 \notin \mathbb{F}_p^{\times,2} \},$$

$$\Omega_{p,2} = \{ (r_0, r_1, r_2, r_3) \in \mathbb{F}_p^4 \colon r_2 = 0, \ r_0 r_1 r_3 \neq 0, \ -r_0 r_1 \notin \mathbb{F}_p^{\times,2} \},$$

$$\Omega_{p,3} = \{ (r_0, r_1, r_2, r_3) \in \mathbb{F}_p^4 \colon r_3 = 0, \ r_0 r_1 r_2 \neq 0, \ -r_0 r_1 \notin \mathbb{F}_p^{\times,2} \}.$$

Let **u** be a vector counted by $N^*(U_0, \ldots, U_3)$. The following result shows that **u** mod $p \notin \Omega_p$ for any prime p.

Lemma 3.2. Let p be a prime and let $\mathbf{u} \in \mathbb{Z}^4$ be such that $\mu^2(u_0) \cdots \mu^2(u_3) = 1$. If $\mathbf{u} \mod p \in \Omega_p$ then $X_{u_0u_2,u_1u_3,u_0u_3,u_1u_2}(\mathbb{Q}_p) = \emptyset$.

Proof. This is standard but we include a full proof for the sake of completeness. Assume without loss of generality that $\mathbf{u} \mod p \in \Omega_{p,0}$. We suppose for a contradiction that there exists a solution $(x_0 \colon x_1 \colon x_2 \colon x_3)$ over \mathbb{Q}_p to the equation

$$u_0 u_2 x_0^2 + u_1 u_3 x_1^2 + u_0 u_3 x_2^2 + u_1 u_2 x_3^2 = 0.$$

We can assume that $(x_0, x_1, x_2, x_3) \in \mathbb{Z}_p^4$ is primitive. Since $p \mid u_0$, it follows that $u_1u_3x_1^2 + u_1u_2x_3^2 \equiv 0 \mod p$. But $p \nmid u_1$ and so

$$u_3 x_1^2 + u_2 x_3^2 \equiv 0 \mod p$$
.

But $-u_2u_3$ is a non-square modulo p and so $x_1 \equiv x_3 \equiv 0 \mod p$. Write $x_1 = px_1'$, $x_3 = px_3'$ and $u_0 = pu_0'$, where $p \nmid u_0'$ since u_0 is square-free. Then it follows that

$$u_0'u_2x_0^2 + pu_1u_3x_1'^2 + u_0'u_3x_2^2 + pu_1u_2x_3'^2 = 0,$$

whence $u_2x_0^2 + u_3x_2^2 \equiv 0 \mod p$. But this implies that $x_0 \equiv x_2 \equiv 0 \mod p$, which contradicts the primitivity of (x_0, x_1, x_2, x_3) .

In order to apply the large sieve we need to calculate the size of Ω_p . But for p > 2 we clearly have

$$|\Omega_{p,i}| = (p-1)^2 \cdot \frac{p-1}{2} = \frac{(p-1)^3}{2},$$

for $0 \le i \le 3$. Hence it follows that

$$|\Omega_p| = 2p^3 \left(1 - \frac{1}{p}\right)^3. \tag{3.6}$$

Lemma 3.2 implies that

$$N^*(U_0,\ldots,U_3) \le \# \{\mathbf{u} \in \mathscr{U} : \mathbf{u} \bmod p \notin \Omega_p \ \forall p \le L\},$$

for any $L \geq 1$. On appealing to Lemma 3.1, we conclude that

$$N^*(U_0, \dots, U_3) \ll \frac{\prod_{i=0}^3 (U_i + L^2)}{F(L)},$$
 (3.7)

where

$$F(L) = \sum_{n \le L} \mu^2(n) \prod_{p|n} \frac{|\Omega_p|}{p^4 - |\Omega_p|}.$$

The following result gives a lower bound for this quantity.

Lemma 3.3. We have $F(L) \gg (\log L)^2$, for any $L \geq 2$.

Proof. It follows from (3.6) that

$$\frac{|\Omega_p|}{p^4 - |\Omega_p|} = \frac{2p^3(1 - \frac{1}{p})^3}{p^4(1 - \frac{2}{p}(1 - \frac{1}{p})^3)}$$
$$\ge \frac{2}{p} \left(1 - \frac{1}{p}\right)^3.$$

Hence

$$F(L) \ge \sum_{n \le L} \frac{\mu^2(n)\tau(n)\phi^*(n)^3}{n},$$

where $\tau(n)$ is the divisor function and

$$\phi^*(n) = \prod_{p|n} \left(1 - \frac{1}{p}\right).$$

We now apply a standard result on the asymptotic evaluation of multiplicative arithmetic functions supported on square-free integers, as supplied by Friedlander and Iwaniec [3, Thm. A.5]. Let $g(n) = \mu^2(n)\tau(n)\phi^*(n)^3/n$. Then

$$\sum_{p \le x} g(p) \log p = 2 \sum_{p \le x} \frac{\log p}{p} + O(1) = 2 \log x + O(1),$$

by Mertens' theorem. Moreover, the conditions (A.16) and (A.17) of [3, Thm. A.5] are easily verified. Hence, on taking k=2 in this result, it follows that

$$\sum_{n \le L} g(n) \gg (\log L)^2,$$

which thereby establishes the lemma.

Let $U_{\min} = \min\{U_0, \dots, U_3\}$ and recall that we are assuming that $U_{\min} \geq 2$. Inserting Lemma 3.3 into (3.7) and taking $L = \sqrt{U_{\min}}$, we finally arrive at the upper bound (3.5).

3.2. **Deduction of the general case.** We now turn to the estimation of $N^{\circ}(B)$, starting from (3.3). Pick T_0, \ldots, T_3 such that

$$T_0, \dots, T_3 \ge 1$$
 and $T_0 T_2, T_0 T_3, T_1 T_2, T_1 T_3 \le 4\sqrt{B}$. (3.8)

We shall examine the contribution $N(T_0, ..., T_3)$, say, from those $\mathbf{t} \in \mathbb{Z}_{\neq 0}^4$ for which $T_i/2 < |t_i| \le T_i$, for $0 \le i \le 3$. Then

$$N^{\circ}(B) \ll \sum_{T_0,\dots,T_3} N(T_0,\dots,T_3),$$
 (3.9)

where T_0, \ldots, T_3 run over powers of 2 subject to the inequalities in (3.8).

Each non-zero integer t_i admits a factorisation $u_i m_i^2$ for square-free $u_i \in \mathbb{Z}$ and $m_i \in \mathbb{Z}_{>0}$. Moreover,

$$X_{t_0t_2,t_1t_3,t_0t_3,t_1t_2}(\mathbf{A}_{\mathbb{Q}}) \neq \emptyset \iff X_{u_0u_2,u_1u_3,u_0u_3,u_1u_2}(\mathbf{A}_{\mathbb{Q}}) \neq \emptyset.$$

Hence it follows that

$$N(T_0, \dots, T_3) \le \sum_{m_0 \le \sqrt{T_0}} \dots \sum_{m_3 \le \sqrt{T_3}} N^* \left(\frac{T_0}{m_0^2}, \dots, \frac{T_3}{m_3^2} \right),$$

in the notation of (3.4). On appealing to (3.5), we deduce that

$$N(T_0, \dots, T_3) \ll \sum_{m_0 \leq \sqrt{T_0}} \dots \sum_{m_3 \leq \sqrt{T_3}} \sum_{j \in \{0, \dots, 3\}} \frac{T_0 T_1 T_2 T_3}{m_0^2 m_1^2 m_2^2 m_3^2 (1 + \log(T_j / m_j^2))^2}$$

$$\ll T_0 T_1 T_2 T_3 \sum_{j \in \{0, \dots, 3\}} \sum_{m \leq \sqrt{T_j}} \frac{1}{m^2 (1 + \log(T_j / m^2))^2}$$

$$\ll \frac{T_0 T_1 T_2 T_3}{(1 + \log T_{\min})^2},$$

where $T_{\min} = \min\{T_0, \dots, T_3\}$. Returning to (3.9), we finally conclude that

$$N^{\circ}(B) \ll \sum_{T_0, \dots, T_3} \frac{T_0 T_1 T_2 T_3}{(1 + \log T_{\min})^2},$$

where T_0, \ldots, T_3 run over powers of 2 subject to the inequalities in (3.8).

By symmetry, we can suppose without loss of generality that $T_0 \leq T_1$ and $T_2 \leq T_3$. Summing first over $T_1 \leq 4\sqrt{B}/T_3$, we see that

$$\sum_{T_0, \dots, T_3} \frac{T_0 T_1 T_2 T_3}{(1 + \log T_{\min})^2} \ll \sqrt{B} \sum_{T_0, T_2, T_3} \frac{T_0 T_2}{(1 + \log \min\{T_0, T_2\})^2}.$$

Suppose first that $T_0 \leq T_2$. Then we sum over $T_2 \leq T_3$ and then $T_3 \leq 4\sqrt{B}/T_0$ to get

$$\sum_{T_0,\dots,T_3} \frac{T_0 T_1 T_2 T_3}{(1 + \log T_{\min})^2} \ll \sqrt{B} \sum_{T_0,T_3} \frac{T_0 T_3}{(1 + \log T_0)^2}$$

$$\ll B \sum_{T_0} \frac{1}{(1 + \log T_0)^2},$$

where the sum is over $T_0 = 2^j$ with $1 \le 2^j \le 2B^{1/4}$. This sum is convergent, whence this case contributes O(B). Alternatively, if $T_0 > T_2$, then we sum over

 $T_0 \leq 4\sqrt{B}/T_3$ and then $T_3 \geq T_2$ to get

$$\sum_{T_0, \dots, T_3} \frac{T_0 T_1 T_2 T_3}{(1 + \log T_{\min})^2} \ll \sqrt{B} \sum_{T_0, T_2, T_3} \frac{T_0 T_2}{(1 + \log T_2)^2}$$

$$\ll B \sum_{T_2, T_3} \frac{T_2}{T_3 (1 + \log T_2)^2}$$

$$\ll B \sum_{T_2} \frac{1}{(1 + \log T_2)^2}.$$

This also makes the satisfactory contribution O(B).

Hence we have shown that $N^{\circ}(B) = O(B)$. Once inserted into (3.2), this therefore completes the proof of the upper bound in (3.1).

References

- [1] T.D. Browning and D. Loughran, Sieving rational points on varieties. *Trans. Amer. Math. Soc.* **371** (2019), 5757–5785.
- [2] J. Franke, Y.I. Manin and Y. Tschinkel, Rational points of bounded height on Fano varieties. *Invent. Math.* **95** (1989), 421–435.
- [3] J.B. Friedlander and H. Iwaniec, Opera de cribro. American Math. Soc., 2010.
- [4] J.B. Friedlander and H. Iwaniec, Ternary quadratic forms with rational zeros. *J. Théor. Nombres Bordeaux* **22** (2010), 97–113.
- [5] H. Hironaka, Resolution of singularities of an algebraic variety over a field of characteristic zero. I & II. Annals Math. **79** (1964), 109–203 & 205–326.
- [6] E. Kowalski, The large sieve and its applications. Cambridge University Press, 2009.
- [7] D. Loughran and A. Smeets, Fibrations with few rational points. Geom. Funct. Anal. 26 (2016), 1449–1482.
- [8] E. Peyre, Liberté et accumulation. Documenta Math. 22 (2017), 1615–1659.
- [9] B. Poonen, Rational Points on Varieties. Volume 186 of Graduate Studies in Mathematics.
 American Mathematical Society, Providence, RI, 2017.
- [10] B. Poonen and J.F. Voloch, Random Diophantine equations. Arithmetic of higher-dimensional algebraic varieties (Palo Alto, CA, 2002), 175–184, Progr. Math. 226, Birkhäuser, 2004.

IST Austria, Am Campus 1, 3400 Klosterneuburg, Austria

Email address: tdb@ist.ac.at
Email address: jlyczak@ist.ac.at

Email address: romansarapin13@gmail.com